# Siegel $p^2$ Vectors for Representations of GSp(4)

Jonathan Cohen

Abstract. Let F be a p-adic field and  $(\pi, V)$  an irreducible complex representation of  $G = \operatorname{GSp}(4, F)$  with trivial central character. Let  $\operatorname{Si}(\mathfrak{p}^2) \subset G$  denote the Siegel congruence subgroup of level  $\mathfrak{p}^2$  and  $u \in N_G(\operatorname{Si}(\mathfrak{p}^2))$  the Atkin-Lehner element. We compute the dimension of the space of  $\operatorname{Si}(\mathfrak{p}^2)$ -fixed vectors in V as well as the signatures of the involutions  $\pi(u)$  acting on these spaces.

# 1 Introduction

This paper is concerned with a dimension counting problem in *p*-adic representation theory for the group  $\operatorname{GSp}(4)$ . Part of the motivation comes from the following problem in the classical theory of Siegel modular forms. If  $\Gamma \subset \operatorname{Sp}(4, \mathbb{Q})$  is a congruence subgroup, then the dimension of the space of cusp forms of level  $\Gamma$  (and integer weight k, say) is not known in general. For example, if

$$\Gamma = \begin{bmatrix} \mathbb{Z} & 4\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & \mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & 4\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ 4\mathbb{Z} & 4\mathbb{Z} & 4\mathbb{Z} & \mathbb{Z} \end{bmatrix} \cap \operatorname{Sp}(4, \mathbb{Z})$$

is the "Klingen congruence subgroup of level 4" then the associated dimensions were only recently computed in [RSY2022]. The method by which this was achieved required, as one of its several inputs, the dimensions of spaces of fixed vectors in all irreducible smooth representations of  $GSp(4, \mathbb{Q}_2)$  for the subgroups

$$\mathrm{Kl}(4) = \begin{bmatrix} \mathbb{Z}_2 & \mathbb{Z}_2 & \mathbb{Z}_2 & \mathbb{Z}_2 \\ 4\mathbb{Z}_2 & \mathbb{Z}_2 & \mathbb{Z}_2 & \mathbb{Z}_2 \\ 4\mathbb{Z}_2 & \mathbb{Z}_2 & \mathbb{Z}_2 & \mathbb{Z}_2 \\ 4\mathbb{Z}_2 & 4\mathbb{Z}_2 & 4\mathbb{Z}_2 & \mathbb{Z} \end{bmatrix} \cap \mathrm{GSp}(4, \mathbb{Z}_2).$$

These dimensions, and more, had been computed in [Yi2021]. We remark that the different "shape" of the subgroups in question is an artifact of different conventions for alternating forms in the classical and representation-theoretic contexts.

If one hopes to use the approach taken in [RSY2022] for other congruence subgroups, then a necessary component is the determination of dimensions of spaces of fixed vectors in all irreducible smooth representations  $(\pi, V)$  of  $\text{GSp}(4, \mathbb{Q}_p)$  for appropriate local subgroups. In this paper we carry out this local computation for the "Siegel congruence subgroups of level  $p^{2n}$ given by

$$\operatorname{Si}(p^2) = \begin{bmatrix} \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ p^2 \mathbb{Z}_p & p^2 \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ p^2 \mathbb{Z}_p & p^2 \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \end{bmatrix} \cap \operatorname{GSp}(4, \mathbb{Z}_p).$$

#### 2 NOTATION

If we let

$$u_{2} := \begin{bmatrix} & 1 & \\ & & -1 \\ p^{2} & & \\ & -p^{2} & \end{bmatrix} \in N_{G}(\operatorname{Si}(p^{2}))$$

then  $u_2$  acts by an involution on the space of  $\operatorname{Si}(p^2)$ -fixed vectors in any representation of  $\operatorname{GSp}(4, \mathbb{Q}_p)$  with trivial central character. We compute the signature of this action. Our main result is stated in Theorem 3.1. We remark that the case p = 2 may be found in [RSY2022], and the case where  $(\pi, V)$  has Iwahori-invariant vectors is handled in [Ro2006].

Our situation is in one sense easier than the corresponding Klingen subgroup because  $\operatorname{Si}(p^2)$  contains a conjugate of the pro-unipotent radical of  $\operatorname{GSp}(4, \mathbb{Z}_p)$ , and the work of [Ros2018] then permits us to reduce to a finite computation employing the character tables of [Sh1982] and [En1972]. Since there are some errors in these character tables, we include a fully corrected table (when p = 2) since there does not appear to be one published elsewhere; see tables 8 and 9. When q is odd the only errors we have found in [Sh1982] are that the three entries in the  $\tau_2$  column for the conjugacy classes of type C should all be negated.

We thank Ralf Schmidt for many useful conversations in preparing this article.

# 2 Notation

Let  $\mathbb{F}_q$  be the field with q elements, where q is a power of a prime number p. If q is odd let  $\xi$  denote a fixed non-square in  $\mathbb{F}_q^{\times}$ , and  $\alpha_0 : \mathbb{F}_q^{\times} \to \{\pm 1\}$  the nontrivial quadratic character.

Let F be a field. Throughout we fix the symplectic form

$$J = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}.$$
 (1)

We let GSp(4) be the algebraic group whose F-points are

$$\operatorname{GSp}(4,F) = \{g \in \operatorname{GL}(4,F) \mid {}^{t}gJg = \mu(g)J \text{ for some } \mu(g) \in F^{\times}\}.$$
(2)

The kernel of the multiplier homomorphism  $\mu : \operatorname{GSp}(4, F) \to F^{\times}$  is the symplectic group  $\operatorname{Sp}(4, F)$ . The center of  $\operatorname{GSp}(4, F)$  is

$$Z(F) = \left\{ \begin{bmatrix} a & a \\ & a \\ & a \end{bmatrix} \mid a \in F^{\times} \right\}$$
(3)

If F is a non-archimedean local field of characteristic zero, we let  $\mathfrak{o}$  be its ring of integers,  $\mathfrak{p}$  the maximal ideal of  $\mathfrak{o}$ , and  $\varpi$  a generator of  $\mathfrak{p}$ . We denote by q the cardinality of the residue class field  $\mathfrak{o}/\mathfrak{p}$ . We further define the maximal compact subgroup

$$K = \mathrm{GSp}(4, \mathfrak{o}),\tag{4}$$

its prounipotent radical

$$K^{+} = \ker(K \to \operatorname{GSp}(4, \mathfrak{o}/\mathfrak{p})), \tag{5}$$

and for  $n \geq 1$ , the Siegel congruence subgroups

$$\operatorname{Si}(\mathfrak{p}^n) = K \cap \begin{bmatrix} \mathfrak{o} & \mathfrak{o} & \mathfrak{o} & \mathfrak{o} \\ \mathfrak{o} & \mathfrak{o} & \mathfrak{o} & \mathfrak{o} \\ \mathfrak{p}^n & \mathfrak{p}^n & \mathfrak{o} & \mathfrak{o} \\ \mathfrak{p}^n & \mathfrak{p}^n & \mathfrak{o} & \mathfrak{o} \end{bmatrix}.$$
(6)

We also define the Atkin-Lehner element  $u_n \in N_G(Si(\mathfrak{p}^n))$  by

$$u_n = \begin{bmatrix} 1 & & \\ & -1 \\ & & -\varpi^n \end{bmatrix}.$$
 (7)

Observe that  $u_n^2 = \varpi^n I_4$ .

Let  $(\pi, V)$  be a smooth complex representation of  $G := \operatorname{GSp}(4, F)$  where F is a p-adic field. For a compact open subgroup  $C \subset G$  we define

$$V^C := \{ v \in V \mid \pi(c)v = v \text{ for all } c \in C \}.$$

$$(8)$$

For an irreducible representation  $\rho$  of any group, we write  $\omega_{\rho}$  for its central character.

### 3 Main result and proof

We now state and prove the main result of this article.

**3.1 Theorem.** Let  $(\pi, V)$  be an irreducible smooth complex representation of G with trivial central character. The integers dim  $V^{\text{Si}(\mathfrak{p}^2)}$  and the signatures of  $\pi(u_2)$  are given in Table 4 when q is even and in tables 6 and 7 when q is odd.

*Proof.* Define the two groups H, M and the element u by

$$H = K \cap \begin{bmatrix} \begin{smallmatrix} \mathfrak{o} & \mathfrak{o} & \mathfrak{p} & \mathfrak{p} \\ \mathfrak{o} & \mathfrak{o} & \mathfrak{p} & \mathfrak{p} \\ \mathfrak{p} & \mathfrak{p} & \mathfrak{o} & \mathfrak{o} \end{bmatrix} = \begin{bmatrix} \varpi & \varpi & \\ & 1 & \\ & & 1 \end{bmatrix} \operatorname{Si}(\mathfrak{p}^2) \begin{bmatrix} \varpi^{-1} & \varpi^{-1} & \\ & & 1 & \\ & & & 1 \end{bmatrix}$$
(9)

$$M = H/K^{+} = \left\{ \begin{bmatrix} a & b \\ c & d \\ & \lambda a & -\lambda b \\ & -\lambda c & \lambda d \end{bmatrix} \mid \lambda, ad - bc \in \mathbb{F}_{q}^{\times} \right\}$$
(10)

$$u = \begin{bmatrix} 1 & & \\ & -1 \\ & & -1 \end{bmatrix} = \varpi^{-1} I_4 \begin{bmatrix} \varpi & & \\ & & 1 \\ & & & 1 \end{bmatrix} u_2 \begin{bmatrix} \varpi^{-1} & & \\ & & \pi^{-1} \\ & & & 1 \end{bmatrix} \in N_G(H)$$
(11)

Observe that  $M \cong \operatorname{GL}(2, \mathbb{F}_q) \times \mathbb{F}_q^{\times}$ . If q is even then  $\operatorname{GSp}(4, \mathbb{F}_q) = Z(\mathbb{F}_q) \times \operatorname{Sp}(4, \mathbb{F}_q)$  and  $M = Z(\mathbb{F}_q) \times (M \cap \operatorname{Sp}(4, \mathbb{F}_q))$ . Note  $u \in \operatorname{Sp}(4, F) \cap K$ . Let  $s(\pi)$  denote the signature of  $\pi(u_2)$  acting on  $V^{\operatorname{Si}(\mathfrak{p}^2)}$ , or equivalently, of  $\pi(u)$  acting on  $V^H$ . Let  $\chi$  denote the trace character of  $V^{K^+}$  as a  $K/K^+ = \operatorname{GSp}(4, \mathbb{F}_q)$ -representation. We have

$$\dim V^{\operatorname{Si}(\mathfrak{p}^2)} = \dim V^H$$
  
=  $\dim((V^{K^+})^M)$   
=  $|M|^{-1} \sum_{m \in M} \chi(m)$  (12)

$$s(\pi) = |M|^{-1} \sum_{m \in M} \chi(mu)$$
 (13)

where we identify u with its image in  $K/K^+$ .

For each  $(\pi, V)$ , the corresponding  $\chi$  is given in table 3 of [Ros2018] in terms of the virtual characters listed in [Sh1982] and [En1972]. In tables 8 and 9 are corrected versions of the latter character tables.

**Proof of dimension formulas.** We must determine the distribution of conjugacy classes of  $GSp(4, \mathbb{F}_q)$  in M.

Table 1: The representatives of the conjugacy classes of M, with the sizes of the M-conjugacy class and the labels of the corresponding classes in  $GSp(4, \mathbb{F}_q)$ , q odd, as listed in [Sh1982]. Some rows have representative given in a quadratic extension of  $\mathbb{F}_q$ . In the last column we also indicate the equivalences that occur over M, for purposes of counting multiplicities.

g	$[M:C_M(g)]$	conditions	$K/K^+$ -conjugacy class
	1	$a\in \mathbb{F}_q^\times$	$A_0(a)$
$\begin{bmatrix} a & 1 \\ a & \\ & a & 1 \\ & & a \end{bmatrix}$	$q^2 - 1$	$a \in \mathbb{F}_q^{\times}$	$A_{21}(a)$
$\left[\begin{array}{c}a\\-a\\-a\\a\end{array}\right]$	$q^2 + q$	$a \in \mathbb{F}_q^{\times}$	$B_0(a) = B_0(-a)$
$\left[ egin{array}{cc} a & & \ -a & & \ & -a & \ & & a \end{array}  ight]$	$q^2 - q$	$a^q = -a$	$C_0(a) = C_0(-a)$
$\begin{bmatrix} a & & \\ & a & \\ & & b \end{bmatrix}$	1	$a, b \in \mathbb{F}_q^{\times}, \ a \neq b$	$D_0(a,b)$
$\begin{bmatrix} a \\ b \\ a \\ b \end{bmatrix}$	$q^2 + q$	$a, b \in \mathbb{F}_q^{\times}, \ a \neq b$	$D_0(a,b) = D_0(b,a)$
$\begin{bmatrix} a & 1 \\ a \\ b & 1 \\ b & b \end{bmatrix}$	$q^2 - 1$	$a,b\in \mathbb{F}_q^{\times},a\neq b$	$D_1(a,b)$
$\begin{bmatrix} ac \\ c \\ c \\ ca^{-1} \end{bmatrix}$	$q^2 + q$	$a,c\in \mathbb{F}_q^{\times},a\neq \pm 1$	$E_0(c,a)$
$\begin{bmatrix} a & & \\ & a^{q} & \\ & & a \\ & & & a^{q} \end{bmatrix}$	$q^2 - q$	$a^q \neq a$	$F_0(a) = F_0(a^q)$
$\begin{bmatrix} a_1 & & \\ & a_2 & \\ & & a_3 \\ & & & a_4 \end{bmatrix}$	$q^2 + q$	$\begin{array}{c} a_i \in \mathbb{F}_q^{\times}, \; a_1 a_4 = a_2 a_3, \\ a_i \neq a_j \; \text{if} \; i \neq j \end{array}$	$H_0(a_1, a_2, a_3, a_4) = H_0(a_2, a_1, a_4, a_3)$
$\begin{bmatrix} a & a^q & & \\ & ba^{-q} & \\ & & ba^{-1} \end{bmatrix}$	$q^2 - q$	$\begin{array}{c} a^{q} \neq a, \ b \in \mathbb{F}_{q}^{\times} \\ b \neq a^{2}, a^{q+1} \end{array}$	$I_0(a,b) = I_0(a^q,b)$

Table 2: The first eight rows give the conjugacy classes of  $M \cap \operatorname{Sp}(4, \mathbb{F}_q)$  with the sizes of the M-conjugacy class and the labels of the corresponding classes in  $\operatorname{Sp}(4, \mathbb{F}_q)$ , q even, as listed in [En1972]. Some elements have representatives given in a quadratic extension of  $\mathbb{F}_q$ . In particular  $\theta$  denotes an element of order  $q^2 - 1$  in  $\mathbb{F}_{q^2}$ , with  $\eta = \theta^{q-1}$  and  $\gamma = \theta^{q+1}$ . The last four rows give representatives of the  $\operatorname{Sp}(4, q)$ -conjugacy classes of  $Mu \cap \operatorname{Sp}(4, q)$ . In the last column we also indicate the equivalences that occur over M, for purposes of counting multiplicities.  $g = \lfloor [M: C_M(q)] \rfloor$  conditions  $\lfloor K/K^+$ -conjugacy class

g	$[M:C_M(g)]$	conditions	$K/K^+$ -conjugacy cla
	1		$A_1$
	$q^2 - 1$		$A_{31}$
$\begin{bmatrix} \gamma^i & & & \\ & \gamma^j & & \\ & & \gamma^{-j} & \\ & & & \gamma^i \end{bmatrix}$	$q^2 + q$	$\gamma^i,  \gamma^j,  \gamma^{i\pm j} \neq 1$	$B_1(i,j) = B_1(j,i)$
$\begin{bmatrix} \theta^{i} & & & \\ & \theta^{qi} & \\ & & \theta^{-qi} & \\ & & \theta^{-i} \end{bmatrix}$	$q^2 - q$	$\theta^{i(q\pm 1)} \neq 1$	$B_2(i) = B_2(qi)$
$\begin{bmatrix} 1 & & & \\ & \gamma^i & \\ & & \gamma^{-i} & \\ & & & 1 \end{bmatrix}$	$q^2 + q$	$1 \le i \le q-2$	$C_1(i)$
$\left \begin{array}{c}\gamma^{i}\\\gamma^{-i}\\\gamma^{i}\\\gamma^{-i}\end{array}\right $	$q^2 + q$	$1 \le i \le q-2$	$C_2(i) = C_2(-i)$
$\begin{bmatrix} \eta^i & & & \\ & \eta^{-i} & \\ & & \eta^i & \\ & & & n^{-i} \end{bmatrix}$	$q^2 - q$	$1 \leq i \leq q$	$C_4(i) = C_4(-i)$
$\begin{bmatrix} \gamma^i & 1 & & \\ & \gamma^i & \\ & & \gamma^{-i} & 1 \\ & & & \gamma^{-i} \end{bmatrix}$	$q^2 - 1$	$1 \le i \le q-2$	$D_2(i)$
	-		$A_{31}$
	-		$A_{32}$
$\begin{bmatrix} \gamma^{i} & \gamma^{i} \\ \gamma^{-i} & \gamma^{-i} \\ \gamma^{i} & \gamma^{-i} \\ \gamma^{i} & \gamma^{-i} \end{bmatrix}$	-	$1 \le i \le q-2$	$D_2(i) = D_2(-i)$
$\begin{bmatrix} \eta^{i} & \eta^{i} & \\ \eta^{-i} & \eta^{-i} \\ \eta^{i} & \\ & n^{-i} \end{bmatrix}$	_	$1 \le i \le q$	$D_4(i) = D_4(-i)$
L '/ J	1		

From tables 1 and 2 and the triviality of the central character of  $\pi$  we can conclude the following formulas: when q is even we have

$$(q^{2}-1)(q^{2}-q)\dim V^{H} = \chi(A_{1}) + (q^{2}-1)\chi(A_{31}) + \frac{q(q+1)}{2} \sum_{\substack{i,j=1\\i\neq j,q-1-j}}^{q-2} \chi(B_{1}(i,j)) + \frac{q(q-1)}{2} \sum_{\substack{i=1\\(q\pm1)\nmid i}}^{q^{2}-2} \chi(B_{2}(i)) + q(q+1) \sum_{i=1}^{q-2} \chi(C_{1}(i)) + \frac{q^{2}+q+2}{2} \sum_{i=1}^{q-2} \chi(C_{2}(i)) + \frac{q(q-1)}{2} \sum_{i=1}^{q} \chi(C_{4}(i)) + (q^{2}-1) \sum_{i=1}^{q-2} \chi(D_{2}(i)).$$
(14)

and when q is odd we have

$$(q^{2}-1)(q^{2}-q)\dim V^{H} = \chi(A_{0}) + (q^{2}-1)\chi(A_{21}) + \frac{q(q+1)}{2}\chi(B_{0}) + \frac{q(q-1)}{2}\chi(C_{0}(\xi^{1/2})) + \frac{q^{2}+q+2}{2} \sum_{1\neq b\in \mathbb{F}_{q}^{\times}} \chi(D_{0}(1,b)) + (q^{2}-1) \sum_{1\neq b\in \mathbb{F}_{q}^{\times}} \chi(D_{1}(1,b)) + q(q+1) \sum_{\pm 1\neq a\in \mathbb{F}_{q}^{\times}} \chi(E_{0}(1,a)) + \frac{q(q-1)}{2} \sum_{a\in \mathbb{F}_{q}} \chi(F_{0}(a+\xi^{1/2})) + \frac{q(q+1)}{2} \sum_{\substack{a,b\in \mathbb{F}_{q}^{\times}\setminus\{1\}\\b\neq a,\ a^{2}}} \chi(H_{0}(1,a,ba^{-1},b)) + \frac{q(q-1)}{2} \sum_{\pm 1\neq b\in \mathbb{F}_{q}^{\times},} \chi(I_{0}(\xi^{1/2},-b\xi)) + \frac{q(q-1)}{2} \sum_{\substack{a\in \mathbb{F}_{q}^{\times}\\1\neq b\in \mathbb{F}_{q}^{\times}}} \chi(I_{0}(a+\xi^{1/2},b(a^{2}-\xi))$$
(15)

In tables 3 and 5 we list the evaluations of the expressions in the right side of (14) and (15) for all of the virtual characters constructed in [En1972] and [Sh1982]. Coupled with table 3 of [Ros2018], one obtains the Si( $\mathfrak{p}^2$ )-columns of tables 4, 6, and 7.

**Proof of signature formulas.** We must determine the distribution of conjugacy classes of  $GSp(4, \mathbb{F}_q)$  in Mu = uM.

Suppose first that q is even. Elements in uM have the form  $\begin{bmatrix} \chi & \lambda X \end{bmatrix}$  for  $X \in GL(2, \mathbb{F}_q)$  and  $\lambda \in \mathbb{F}_q^{\times}$ . The equalities

$$\begin{bmatrix} \lambda^{-1/2}I \\ I \end{bmatrix} \begin{bmatrix} \chi & \lambda X \end{bmatrix} \begin{bmatrix} \lambda^{1/2}I \\ I \end{bmatrix} = \lambda^{1/2} \cdot \begin{bmatrix} \chi & X \end{bmatrix}$$
$$\begin{bmatrix} I \\ I & I \end{bmatrix} \begin{bmatrix} \chi & X \end{bmatrix} \begin{bmatrix} I \\ I & I \end{bmatrix} = \begin{bmatrix} X & X \\ X \end{bmatrix}$$

together show that

$$\sum_{m \in M} \chi(mu) = (q-1) \sum_{X \in \operatorname{GL}(2,\mathbb{F}_q)} \chi(\begin{bmatrix} X & X \\ X \end{bmatrix}) = (q-1)^2 \sum_{X \in \operatorname{SL}(2,\mathbb{F}_q)} \chi(\begin{bmatrix} X & X \\ X \end{bmatrix}).$$

Conjugating by an appropriate  $\begin{bmatrix} B \\ B \end{bmatrix} \in M$  allows us to assume that X is in Jordan form. From the last four rows in table 2, and the orders of centralizers of elements of  $GL(2, \mathbb{F}_q)$ , we obtain

$$q(q^{2}-1)s(\pi) = \chi(A_{31}) + (q^{2}-1)\chi(A_{32}) + \frac{q(q+1)}{2}\sum_{i=1}^{q-2}\chi(D_{2}(i)) + \frac{q(q-1)}{2}\sum_{i=1}^{q}\chi(D_{4}(i)).$$
(16)

Finally, suppose q is odd. Unlike the even q case, not every element of uM is conjugate to one which is block upper triangular. Define the following subgroups and elements of M:

$$T = \left\{ \begin{bmatrix} 1 & d \\ & c \\ & & cd \end{bmatrix} \mid c, d \in \mathbb{F}_q^{\times} \right\}, \qquad Z = Z(\mathbb{F}_q), \qquad N = \left\{ \begin{bmatrix} 1 & b \\ & 1 & -b \\ & & 1 \end{bmatrix} \mid b \in \mathbb{F}_q \right\},$$
$$t_{c,d} = \begin{bmatrix} 1 & d \\ & c \\ & & cd \end{bmatrix}, \qquad x = \begin{bmatrix} 1 & 1 \\ & 1 & -1 \\ & & 1 \end{bmatrix}, \qquad w = \begin{bmatrix} 1 & 1 \\ & & 1 \end{bmatrix}.$$

Clearly T and u normalize N and  $\mu(w) = 1$ . Using the Bruhat decomposition, every element in M may be written uniquely in the form ztn or in the form ztnwn' for  $z \in Z$ ,  $t \in T$ ,  $n, n' \in N$ . Since  $ztnwn'u = ztnwu(u^{-1}n'u)$  is conjugate to  $zt(t^{-1}u^{-1}n'utn)wu$ , we obtain

$$\sum_{m \in M} \chi(mu) = (q-1) \sum_{t \in T, n \in N} \left( \chi(tnu) + q\chi(tnwu) \right)$$
(17)

Note that  $\mu(t_{c,d}nu) = \mu(t_{c,d}nwu) = cd$ . The reader can readily verify the following assertions:

- i) We have  $t_{a^2c,d}nu$  is T-conjugate to  $aI_4 \cdot t_{c,d}nu$  and  $t_{a^2c,d}nwu$  is T-conjugate to  $aI_4 \cdot t_{c,d}nwu$ ,
- ii) If  $1 \neq n \in N$  has corresponding parameter  $b \neq 0$  then  $t_{c,d}nu$  is *T*-conjugate to  $t_{c,d}xu$  and  $t_{c,b^2d}nwu$  is *T*-conjugate to  $bI_4 \cdot t_{c,d}xwu$ ,
- iii) If  $d \neq 1$  then  $t_{c,d}nu$  is N-conjugate to  $t_{c,d}u$ .

Thus

$$\sum_{m \in M} \chi(mu) = \frac{(q-1)^2}{2} \sum_{c \in \{1,\xi\}, d \in \mathbb{F}_q^{\times}, n \in N} (\chi(t_{c,d}nu) + q\chi(t_{c,d}nwu))$$
$$= \frac{(q-1)^2}{2} \sum_{c \in \{1,\xi\}, d \in \mathbb{F}_q^{\times}} \left( \chi(t_{c,d}u) + (q-1)\chi(t_{c,d}xu) + q \sum_{n \in N} \chi(t_{c,d}nwu) \right)$$
$$= \frac{(q-1)^2}{2} \sum_{c \in \{1,\xi\}} \left( \chi(t_{c,1}u) + (q-1)\chi(t_{c,1}xu) + q \sum_{1 \neq d \in \mathbb{F}_q^{\times}} \chi(t_{c,d}u) \right)$$

$$+q\sum_{d\in\mathbb{F}_q^{\times}} \left(\chi(t_{c,d}wu) + (q-1)\chi(t_{c,d}xwu)\right)\right).$$
(18)

We now record the following facts, referring to [Sh1982] for the notation for conjugacy classes:

$$t_{c,d}u \text{ is in a conjugacy class of type} \begin{cases} B_0(1) & \text{if } c = d = 1\\ C_0(\xi^{1/2}) & \text{if } c = \xi, d = 1\\ D_0(1,-1) & \text{if } c = 1, d = -1\\ F_0(\xi^{1/2}) & \text{if } c = \xi, d = -1\\ H_0(1,-1,-d,d) & \text{if } c = 1, d \neq \pm 1\\ I_0(\xi^{1/2},d\xi) & \text{if } c = \xi, d \neq \pm 1 \end{cases}$$
(19)  
$$t_{c,1}xu \text{ is in a conjugacy class of type} \begin{cases} B_{31}(1) & \text{if } c = 1, -1 \in \mathbb{F}_q^{\times 2}\\ B_{32}(1) & \text{if } c = 1, -1 \notin \mathbb{F}_q^{\times 2}\\ B_{32}(1) & \text{if } c = 1, -1 \notin \mathbb{F}_q^{\times 2}\\ C_{11}(\xi^{1/2}) & \text{if } c = \xi, -1 \notin \mathbb{F}_q^{\times 2},\\ C_{12}(\xi^{1/2}) & \text{if } c = \xi, -1 \in \mathbb{F}_q^{\times 2} \end{cases}$$
(20)

$$t_{c,d}wu \text{ is in a conjugacy class of type} \begin{cases} D_0((-cd)^{1/2}, -(-cd)^{1/2}) & \text{if } -cd \in \mathbb{F}_q^{\times 2} \\ F_0((-cd)^{1/2}) & \text{if } -cd \notin \mathbb{F}_q^{\times 2} \end{cases}.$$
(21)

For the last elements, observe that  $t_{c,d}xwu$  has eigenvalues  $(-c)^{1/2}\left(\frac{\pm 1\pm (1+4d)^{1/2}}{2}\right)$ . In particular there are four distinct eigenvalues unless d = -1/4. It is not hard to show that

$$t_{c,-1/4}xwu \text{ is in a conjugacy class of type} \begin{cases} B_{31}(1/2) & \text{if } -c, -1 \in \mathbb{F}_q^{\times 2}, \\ B_{32}(1/2) & \text{if } -c \in \mathbb{F}_q^{\times 2}, -1 \notin \mathbb{F}_q^{\times 2} \\ C_{11}(\xi^{1/2}/2) & \text{if } -c, -1 \notin \mathbb{F}_q^{\times 2}, \\ C_{12}(\xi^{1/2}/2) & \text{if } -c \notin \mathbb{F}_q^{\times 2}, -1 \in \mathbb{F}_q^{\times 2}. \end{cases}$$
(22)

Note that  $\{-1, -\xi\}$  is a set of representatives for  $\mathbb{F}_q^{\times}/\mathbb{F}_q^{\times 2}$ . Finally we consider  $t_{c,d}xwu$  with  $d \neq -1/4$ . For  $r \in \mathbb{F}_{q^2}^{\times}$  we have  $r^2 \in \mathbb{F}_q$  if and only if  $r^{q-1} \in \{\pm 1\}$ , with the sign determined by whether  $r \in \mathbb{F}_q^{\times}$  or not. Thus

$$\sum_{d \in \mathbb{F}_q^{\times}, d \neq -1/4} \chi(t_{c,d} x w u) = \sum_{d \in \mathbb{F}_q^{\times}, d \neq -1} \chi(t_{c,d/4} x w u)$$
(23)

$$= \frac{1}{2} \sum_{\substack{r \in \mathbb{F}_{q^2}^{\times} \setminus \{\pm 1\} \\ r^{q-1} = \pm 1}} \chi(t_{c, \frac{r^2 - 1}{4}} nwu).$$
(24)

The eigenvalues of  $t_{c,\frac{r^2-1}{4}}nwu$  are  $(-c)^{1/2}\left(\frac{\pm 1\pm r}{2}\right)$ . It is not hard to show that the conjugacy class of  $t_{c,\frac{r^2-1}{4}}xwu$  is of type

#### 4 TABLES

$$\begin{cases} (-c)^{1/2} \cdot H_0(\frac{1+r}{2}, \frac{-1-r}{2}, \frac{-1+r}{2}, \frac{1-r}{2}) & \text{if } -c \in \mathbb{F}_q^{\times 2}, r \in \mathbb{F}_q^{\times} \\ (-c)^{1/2} L_0(\frac{1+r}{2}, \frac{-1+r}{2}) & \text{if } -c \in \mathbb{F}_q^{\times 2}, r \notin \mathbb{F}_q^{\times} \\ I_0((-c)^{1/2}(\frac{1+r}{2}), \frac{c(r^2-1)}{4}) & \text{if } -c \notin \mathbb{F}_q^{\times 2}, r \in \mathbb{F}_q^{\times} \\ I_0((-c)^{1/2}(\frac{1+r}{2}), \frac{c(r^2-1)}{4}) & \text{if } -c \notin \mathbb{F}_q^{\times 2}, r \notin \mathbb{F}_q^{\times} \end{cases}$$
(25)

Here the conjugacy class  $I_0(a, b)$  denotes the one with an eigenvalue  $a \notin \mathbb{F}_q$  and similitude  $b \neq a^2, a^{q+1}$ , and  $L_0(a, b)$  denotes a conjugacy class with eigenvalues  $a, b \notin \mathbb{F}_q$  and  $a \neq b^q$ . Putting everything together and using the triviality of the central character, we obtain

$$2q(q^{2}-1)s(\pi) = \chi(B_{0}) + \chi(C_{0}(\xi^{1/2})) + (q^{2}-1)\left(\chi(B_{31})\left(\frac{1+\alpha_{0}(-1)}{2}\right) + \chi(B_{32})\left(\frac{1-\alpha_{0}(-1)}{2}\right)\right) + (q^{2}-1)\left(\chi(C_{11}(\xi^{1/2}))\left(\frac{1-\alpha_{0}(-1)}{2}\right) + \chi(C_{12}(\xi^{1/2}))\left(\frac{1+\alpha_{0}(-1)}{2}\right)\right) + q^{2}\left(\chi(D_{0}(1,-1)) + \chi(F_{0}(\xi^{1/2}))\right) + \frac{q(q+1)}{2}\sum_{\pm 1\neq a\in\mathbb{F}_{q}^{\times}}\left(\chi(H_{0}(1,-1,-a,a)) + \chi(I_{0}(\xi^{1/2},\xi a))\right) + \frac{q(q-1)}{2}\sum_{a\in\mathbb{F}_{q}^{\times}}\left(\chi(I_{0}(1+a\xi^{1/2},-1+a^{2}\xi)) + \chi(L_{0}(1+a\xi^{1/2},-1-a\xi^{1/2}))\right).$$
(26)

In tables 3 and 5 we list the evaluations of the expressions in the right side of (16) and (26) for all of the virtual characters given in the tables of [En1972] and [Sh1982]. Coupled with table 3 of [Ros2018], one obtains the *s*-columns of tables 4, 6, and 7.

**3.2 Corollary.** Let  $(\pi, V)$  be an irreducible representation of G with trivial central character Then dim  $V^{\text{Si}(\mathfrak{p}^2)} \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 12\}.$ 

#### 4 Tables

The irreducible characters of GSp(4, q), with q even, are classified in [En1972], and those with q odd are handled in [Sh1982]. We refer to their papers for the definition of the virtual characters that appear in the first column of tables 3 and 5. In table 3 we have listed the parameters for each character. In table 5, when q is odd, the parameters are as follows:  $\lambda, \mu, \nu : \mathbb{F}_q^{\times} \to \mathbb{C}^{\times}$ ,  $\Lambda : \mathbb{F}_{q^2}^{\times} \to \mathbb{C}^{\times}, \omega : \mathbb{F}_{q^2}^{\times}[q+1] \to \mathbb{C}^{\times}, \Theta : \mathbb{F}_{q^2}^{\times}[(q-1)(q^2+1)] \to \mathbb{C}^{\times}$ , and  $\lambda' : \mathbb{F}_{q^2}^{\times}[2(q-1)] \to \mathbb{C}^{\times}$ . The character  $\omega_0 : \mathbb{F}_{q^2}^{\times}[q+1] \to \{\pm 1\}$  is the nontrivial quadratic character, and  $N : \mathbb{F}_{q^2}^{\times} \to \mathbb{F}_q^{\times}$  denotes the norm map. The  $\delta_{-,-}$  denotes the Kronecker delta.

To allow for reducible representations of  $GSp(4, \mathbb{F}_q)$ , we have not restricted the parameters defining these characters. For example, the unramified parabolically induced representations Vof GSp(4, F) will have reducible parahoric restriction  $V^{K^+}$  as a K-representation. Some choices of parameters can result in non-genuine representations, hence the possibility of negative entries in the M columns of tables 3 and 5.

The irreducible nonsupercuspidal representations of  $\operatorname{GSp}(4, F)$  were classified in [Sa1993]. The families they construct constitute the representations given in families I, II,..., XI in tables 4, 6 and 7. There are also three families of supercuspidals that appear in these tables in the last rows. The characters  $\chi_i, \chi, \sigma : F^{\times} \to \mathbb{C}^{\times}$  and quadratic characters  $\xi : F^{\times} \to \{\pm 1\}$  are all tamely ramified. We write  $a(\sigma) \in \{0, 1\}$  for the conductor of a tamely ramified character  $\sigma$ . The letter  $\rho$  indicates a depth zero supercuspidal representation of  $\operatorname{GL}(2, F)$ , to which one may associate a parameter  $\Lambda$  as above with  $\Lambda^{q-1} \neq 1$ , and  $\omega_{\Lambda} := \Lambda^{q-1}$ .

In table 8 and 9,  $\zeta_n$  denotes a complex primitive *n*th root of unity.

$\sigma$	parameters	M	s
$\theta_0$		1	1
$ heta_1$		3	1
$\theta_2$		3	1
$ heta_3$		0	0
$ heta_4$		2	0
$ heta_5$		0	0
$\chi_1(k,l)$	$k, l \in \mathbb{Z}/(q-1)\mathbb{Z}$	$2 + 2(\delta_{k,0} + \delta_{l,0})^2 + \delta_{k,l} + \delta_{k,-l}$	$2 + \delta_{k,l} + \delta_{k,-l}$
$\chi_2(k)$	$k \in \mathbb{Z}/(q^2 - 1)\mathbb{Z}$	$\delta_{(q+1)k,0} - \delta_{(q-1)k,0} - 2\delta_{k,0}$	$-\delta_{(q+1)k,0} - \delta_{(q-1)k,0}$
$\chi_3(k,l)$	$k \in \mathbb{Z}/(q-1)\mathbb{Z}$ $l \in \mathbb{Z}/(q+1)/\mathbb{Z}$	$2 + 2\delta_{k,0}$	0
$\chi_4(k,l)$	$k, l \in \mathbb{Z}/(q+1)/\mathbb{Z}$	$2 - \delta_{k,l} - \delta_{k,-l}$	$-2 + \delta_{k,l} + \delta_{k,-l}$
$\chi_5(k)$	$k \in \mathbb{Z}/(q^2 + 1)/\mathbb{Z}$	0	0
$\chi_6(k)$	$k \in \mathbb{Z}/(q-1)\mathbb{Z}$	$2 + 5\delta_{k,0}$	$2 + \delta_{k,0}$
$\chi_7(k)$	$k \in \mathbb{Z}/(q-1)\mathbb{Z}$	$1 + 3\delta_{k,0}$	$1 + \delta_{k,0}$
$\chi_8(k)$	$k \in \mathbb{Z}/(q+1)\mathbb{Z}$	$-\delta_{k,0}$	$-\delta_{k,0}$
$\chi_9(k)$	$k \in \mathbb{Z}/(q+1)/\mathbb{Z}$	$1 + \delta_{k,0}$	$1 - \delta_{k,0}$
$\chi_{10}(k)$	$k \in \mathbb{Z}/(q-1)\mathbb{Z}$	$1+4\delta_{k,0}$	1
$\chi_{11}(k)$	$k \in \mathbb{Z}/(q-1)\mathbb{Z}$	$3 + 5\delta_{k,0}$	$1 + \delta_{k,0}$
$\chi_{12}(k)$	$k \in \mathbb{Z}/(q+1)\mathbb{Z}$	$1 - 2\delta_{k,0}$	-1
$\chi_{13}(k)$	$k \in \mathbb{Z}/(q+1)\mathbb{Z}$	$3-\delta_{k,0}$	$-1 + \delta_{k,0}$

Table 3: The "dimensions" of M-fixed vectors and signature s of the involution u for q even.

Type	$\pi$	data	$\operatorname{Si}(\mathfrak{p}^2)$	s
Ι	$\chi_1  imes \chi_2  times \sigma$	$a(\chi_1) = a(\chi_2) = 0$	12	4
		$a(\chi_1) + a(\chi_2) = 1$	4	2
		$a(\chi_i) = a(\sigma) + a(\chi_i \sigma) = 1$	3	3
		$a(\chi_i) = a(\sigma) = a(\chi_i \sigma) = 1$	2	2
IIa	$\chi \mathrm{St}_{\mathrm{GL}(2)} \rtimes \sigma$	$a(\chi) = 0$	5	1
		$a(\chi) = 1$	1	1
IIb	$\chi 1_{\mathrm{GL}(2)} \rtimes \sigma$	$a(\chi) = 0$	7	3
		$a(\chi) = 1$	2	2
IIIa	$\chi \rtimes \sigma \operatorname{St}_{\operatorname{GL}(2)}$	$a(\chi) = 0$	8	2
		$a(\chi) = 1$	3	1
IIIb	$\chi \rtimes \sigma 1_{\mathrm{GL}(2)}$	$a(\chi) = 0$	4	2
		$a(\chi) = 1$	1	1
IVa	$\sigma \mathrm{St}_G$	$a(\sigma) = 0$	2	0
IVb	$L(\nu^2, \nu^{-1}\sigma \operatorname{St}_{\operatorname{GSp}(2)})$	$a(\sigma) = 0$	6	2
IVc	$L(\nu^{3/2} \mathrm{St}_{\mathrm{GL}(2)}, \nu^{-3/2}\sigma)$	$a(\sigma) = 0$	3	1
IVd	$\sigma 1_G$	$a(\sigma) = 0$	1	1
Va	$\delta([\xi,\nu\xi],\nu^{-1/2}\sigma)$	$a(\sigma) = 0$	2	0
Vb	$L(\nu^{1/2}\xi St_{GL(2)}, \nu^{-1/2}\sigma)$	$a(\sigma) = 0$	3	1
Vc	$L(\nu^{1/2}\xi \operatorname{St}_{\operatorname{GL}(2)}, \nu^{-1/2}\xi\sigma)$	$a(\sigma) = 0$	3	1
Vd	$L(\nu\xi,\xi\rtimes\nu^{-1/2}\sigma)$	$a(\sigma) = 0$	4	2
VIa	$ au(S,  u^{-1/2}\sigma)$	$a(\sigma) = 0$	5	1
VIb	$ au(T, u^{-1/2}\sigma)$	$a(\sigma) = 0$	3	1
VIc	$L(\nu^{1/2} \mathrm{St}_{\mathrm{GL}(2)}, \nu^{-1/2}\sigma)$	$a(\sigma) = 0$	0	0
VId	$L(\nu, 1_{F^{\times}} \rtimes \nu^{-1/2}\sigma)$	$a(\sigma) = 0$	4	2
VII	$\chi  times  ho$	$a(\chi) = 0$	4	0
		$a(\chi) = 1$	2	0
VIIIa	au(S, ho)		3	-1
VIIIb	au(T, ho)		1	1
IXa	$\delta(\nu\xi,\nu^{-1/2}\rho)$		3	-1
IXb	$L( u\xi, u^{-1/2} ho)$		1	1
Х	$\rho\rtimes\sigma$	$a(\sigma) = 0$	1	-1
		$a(\sigma) = 1$	0	0
XIa	$\delta(\nu^{1/2}\rho,\nu^{-1/2}\sigma)$	$a(\sigma) = 0$	1	-1
XIb	$L(\nu^{1/2}\rho,\nu^{-1/2}\sigma)$	$a(\sigma) = 0$	0	0
s.c.	$\operatorname{c-Ind}_{ZK}^G \chi_5(k)$		0	0
s.c.	$\operatorname{c-Ind}_{ZK}^G \chi_4(k,l)$		2	-2
s.c.	$\operatorname{c-Ind}_{ZK}^G \theta_5$		0	0

Table 4: Si( $\mathfrak{p}^2$ )-fixed vector dimensions and Atkin-Lehner signatures, q even.

$\sigma$	$\omega_{\sigma}$		S
$X_1(\lambda,\mu,\nu)$	$\lambda\mu\nu^2$	$1 + \lambda(-1) + \delta_{\nu,1} + \delta_{\lambda\nu,1} + 2(\delta_{\lambda,1} + \delta_{\mu,1}) + 4\delta_{\lambda,1}\delta_{\nu,1}$	$\nu(-1)(1 + \delta_{\lambda\nu,1}) + \lambda\nu(-1)(1 + \delta_{\nu,1})$
$X_2(\Lambda,  u)$	$\nu^2 \cdot \Lambda _{\mathbb{F}_q^{\times}}$	$1 - \Lambda(\xi^{1/2})\nu(\xi) + \delta_{\nu,1} - \delta_{\Lambda\nu\circ N,1} - 2\delta_{\nu,1}\delta_{\Lambda,1}$	$\nu(-1) \left( 1 - \Lambda(\xi^{1/2})\nu(\xi)(1 + \delta_{\nu,1}) - \delta_{\Lambda\nu \circ N,1} \right)$
$X_3(\Lambda, \nu)$	$ \nu \cdot \Lambda _{\mathbb{F}_{a}^{\times}}$	$1 + \nu(-1) + 2\delta_{\nu,1}$	0
$X_4(\Theta)$	$\Theta _{\mathbb{F}_{a}^{\times}}$	$1 - \Theta(\xi^{1/2})$	0
$X_5(\Lambda,\omega)$	$\Lambda _{\mathbb{F}_q^{\times}}$	$1+\omega(-1)-\delta_{\Lambda,1}-\delta_{\Lambda\widetilde\omega,1}$	$-\Lambda(\xi^{1/2})(1+\omega(-1))+\omega(-1)(\delta_{\Lambda,1}+\delta_{\Lambda\widetilde{\omega},1})$
$\chi_1(\lambda, u)$	$\lambda  u^2$	$1+3\delta_{\nu,1}$	$ u(-1) + \delta_{\nu,1} $
$\chi_2(\lambda, u)$	$\lambda \nu^2$	$3 + 3\delta_{\nu,1} + 2\delta_{\nu^2,1}$	$ u(-1) + \delta_{\nu,1} $
$\chi_3(\lambda, u)$	$\lambda^2 \nu^2$	$\delta_{\lambda\nu,1}(1+\lambda(-1)) + 2\delta_{\lambda,1} + 3\delta_{\lambda,1}\delta_{\nu,1}$	$\delta_{\lambda\nu,1}(1+\lambda(-1))+\delta_{\lambda,1}\delta_{\nu,1}$
$\chi_4(\lambda, u)$	$\lambda^2 \nu^2$	$1 + \lambda(-1)\delta_{\lambda\nu,\alpha_0} + \delta_{\nu,1} + 2\delta_{\lambda,1} + \delta_{\lambda,1}\delta_{\nu,1}$	$\nu(-1) + \alpha_0(-1)(1+\delta_{\nu,1})\delta_{\lambda\nu,\alpha_0}$
$\chi_5(\omega, u)$	$ u^2 $	$\delta_{ u,1}(1-\omega(-1)-\delta_{\omega,1})$	$\delta_{\nu,1}(1-\omega(-1)-\delta_{\omega,1})$
$\chi_6(\omega, u)$	$ u^2$	$1 + \delta_{\nu,\alpha_0}(\omega(-1) - \delta_{\omega,\omega_0}) - 2\delta_{\nu,1}\delta_{\omega,1}$	$-\omega(-1)\nu(-\xi) + \nu(-1)\delta_{\nu,\alpha_0}(1-\delta_{\omega,\omega_0})$
$\chi_7(\Lambda)$	$\Lambda _{\mathbb{F}_q^{\times}}$	$1 + \delta_{\Lambda,1}$	$\Lambda(\xi^{1/2}) - \delta_{\Lambda,1}$
$\chi_8(\Lambda)$	$\Lambda _{\mathbb{F}_q^{\times}}$	$3-\delta_{\Lambda,1}$	$-\Lambda(\xi^{1/2}) + \delta_{\Lambda,1}$
$ au_1$	1	0	0
$ au_2( u)$	$ u^2$	$\delta_{\nu,1}(1+\alpha_0(-1))$	$\delta_{\nu,1}(1+\alpha_0(-1))$
$ au_3$	1	1	$\alpha_0(-1)$
$ au_4(\lambda')$	$\lambda' _{\mathbb{F}_a^{\times}}$	0	0
$ au_5(\lambda')$	$\lambda' _{\mathbb{F}_q^{\times}}$	0	0
$ heta_1( u)$	$\nu^2$	$1+2\delta_{\nu,1}$	u(-1)
$ heta_2( u)$	$ u^2 $	0	0
$ heta_3( u)$	$ u^2$	$1+2\delta_{\nu,1}$	$-\nu(-1) + 2\delta_{\nu,1}$
$ heta_4( u)$	$ u^2$	0	0
$ heta_5( u)$	$ u^2$	$3 - \delta_{ u,1}$	$ u(-1) - \delta_{\nu,1} $
$ heta_0( u)$	$ u^2 $	$\delta_{ u,1}$	$\delta_{ u,1}$

Table 5: The "dimensions" of M-fixed vectors and signature s of the involution u, for q odd. All representations are assumed to have trivial central character.

	Table 6: Si( $\mathfrak{p}^2$ )-fixed vector dimensions and Atkin-Lehner signatures, q odd.							
Type	$\pi$	$\omega_{\pi}$	conditions	$\operatorname{Si}(\mathfrak{p}^2)$	s			
Ι	$\chi_1  imes \chi_2  times \sigma$	$\chi_1\chi_2\sigma^2$	$a(\sigma) = a(\chi_i) = 0$	12	4			
			$a(\sigma) = 1, \ a(\chi_i) = 0$	6	$2\sigma(-1)$			
			$a(\sigma) = a(\chi_1) + a(\chi_2) = 1$	4	$2\sigma(-1)$			
			$a(\chi_i) = a(\sigma) + a(\chi_i \sigma) = 1$	$2 + \chi_1(-1)$	$1 + 2\chi_1(-1)$			
			$a(\chi_i) = a(\sigma) = a(\chi_i \sigma) = 1$	$1 + \chi_1(-1)$	$(1 + \chi_1(-1))\sigma(-1)$			
IIa	$\chi St_{GL(2)} \rtimes \sigma$	$\chi^2 \sigma^2$	$a(\sigma) = a(\chi) = 0$	5	1			
			$a(\sigma) = 0,  a(\chi) = 1$	$2 + \chi(-1)$	$1 + 2\chi(-1)$			
			$a(\sigma) = 1,  a(\chi) = 0$	4	$2\sigma(-1)$			
			$a(\sigma) = a(\chi) = 1, a(\sigma\chi) = 0$	1	$\sigma(-1)$			
			$a(\sigma) = a(\chi) = a(\sigma\chi) = 1$	$1 + \chi(-1)$	$(1+\chi(-1))\sigma(-1)$			
IIb	$\chi 1_{\mathrm{GL}(2)} \rtimes \sigma$	$\chi^2 \sigma^2$	$a(\sigma) = a(\chi) = 0$	7	3			
			$a(\sigma) = 0, \ a(\chi) = 1$	0	0			
			$a(\sigma) = 1, \ a(\chi) = 0$	2	0			
			$a(\sigma) = a(\chi) = 1, a(\sigma\chi) = 0$	$1 + \chi(-1)$	$1 + \chi(-1)$			
	~	0	$a(\sigma) = a(\chi) = a(\sigma\chi) = 1$	0	0			
IIIa	$\chi \rtimes \sigma \mathrm{St}$	$\chi \sigma^2$	$a(\sigma) = a(\chi) = 0$	8	2			
			$a(\sigma) = 1, a(\chi) = 0$	5	$\sigma(-1)$			
		2	$a(\sigma) = a(\chi) = 1$	3	$\sigma(-1)$			
IIIb	$\chi \rtimes \sigma 1$	$\chi \sigma^2$	$a(\sigma) = 0$	4	2			
<b>TT</b> 7	C.	9	$a(\sigma) = 1$	1	$\sigma(-1)$			
IVa	$\sigma \mathrm{St}_G$	$\sigma^2$	$a(\sigma) = 0$	2	( 1)			
TX 71	$\tau(2) = 1 \alpha$	9	$a(\sigma) = 1$	3	$\sigma(-1)$			
IVD	$L(\nu^2, \nu^{-1}\sigma \operatorname{St}_{\operatorname{GSp}(2)})$	$\sigma^2$	$a(\sigma) = 0$	6	2			
<b>TT</b> 7	$\tau$ ( 3/2 g) $-3/2$ )	9	$a(\sigma) = 1$	2	0			
IVC	$L(\nu^{\sigma/2}\operatorname{St}_{\operatorname{GL}(2)},\nu^{-\sigma/2}\sigma)$	$\sigma^2$	$a(\sigma) = 0$	3				
TT 7 1	1	2	$a(\sigma) = 1$	1	$\sigma(-1)$			
IVd	$\sigma \iota_G$	$\sigma^2$	$a(\sigma) = 0$	1				
			$a(\sigma) = 1$	U	0			

# TABLES

Type	$\pi$	$\omega_{\pi}$	conditions	$\operatorname{Si}(\mathfrak{p}^2)$	S
Va	$\delta([\xi, \nu\xi], \nu^{-1/2}\sigma)$	$\sigma^2$	$a(\xi) = a(\sigma) = 0$	2	0
			$a(\xi) = 0, \ a(\sigma) = 1$	3	$\sigma(-1)$
			$a(\xi) = 1$	1	$\xi(-1)$
Vb	$L(\nu^{1/2}\xi St_{GL(2)}, \nu^{-1/2}\sigma)$	$\sigma^2$	$a(\xi) = a(\sigma) = 0$	3	1
			$a(\xi) = 0, \ a(\sigma) = 1$	1	$\sigma(-1)$
			$a(\xi) = 1, \ a(\sigma) = 0$	$1 + \xi(-1)$	$1 + \xi(-1)$
			$a(\xi) = a(\sigma) = 1$	0	0
Vc	$L(\nu^{1/2}\xi St_{GL(2)}, \nu^{-1/2}\xi\sigma)$	$\sigma^2$	$a(\xi) = a(\sigma) = 0$	3	1
			$a(\xi) = 0, \ a(\sigma) = 1$	1	$\sigma(-1)$
			$a(\xi) = 1 \ a(\sigma) = 0$	0	0
			$a(\xi) = a(\sigma) = 1$	$1 + \sigma(-1)$	$1 + \sigma(-1)$
Vd	$L(\nu\xi,\xi \rtimes \nu^{-1/2}\sigma)$	$\sigma^2$	$a(\xi) = a(\sigma) = 0$	4	2
			$a(\xi) = 0, \ a(\sigma) = 1$	1	$\sigma(-1)$
			$a(\xi) = 1$	0	0
VIa	$ au(S,  u^{-1/2}\sigma)$	$\sigma^2$	$a(\sigma) = 0$	5	1
			$a(\sigma) = 1$	4	$2\sigma(-1)$
VIb	$\tau(T, \nu^{-1/2}\sigma)$	$\sigma^2$	$a(\sigma) = 0$	3	1
			$a(\sigma) = 1$	1	$-\sigma(-1)$
VIc	$L(\nu^{1/2} \mathrm{St}_{\mathrm{GL}(2)}, \nu^{-1/2}\sigma)$	$\sigma^2$		0	0
VId	$L(\nu, 1_{E^{\times}} \rtimes \nu^{-1/2}\sigma)$		$a(\sigma) = 0$	4	2
			$a(\sigma) = 1$	1	$\sigma(-1)$
VII	$\chi  times  ho$	$\chi\omega_{o}$	$a(\chi) = 0$	4	0
		, e , r	$a(\chi) = 1$	$1 + \chi(-1)$	0
VIIIa	au(S, ho)	$\omega_{ ho}$	· - /	3	$-\Lambda(t)$
VIIIb	au(T, ho)	$\omega_{ ho}$		1	$\Lambda(t)$
IXa	$\delta(\nu\xi,\nu^{-1/2} ho)$	$\xi \omega_{ ho}$	$a(\xi) = 0$	3	$-\Lambda(t)$
		- /	$a(\xi) = 1$	0	0
IXb	$L(\nu\xi, \nu^{-1/2}\rho)$	$\xi\omega_{ ho}$	$a(\xi) = 0$	1	$-\Lambda(t)$
		,	$a(\xi) = 1$	0	0
Х	$\rho \rtimes \sigma$	$\omega_{ ho}\sigma^2$	$a(\sigma) = 0$	$2 - \Lambda(t)$	$1 - 2\Lambda(t)$
			$a(\sigma) = 1$	$1 - \Lambda \sigma^2(t)$	$\sigma(-1)(1 - \Lambda \sigma^2(t))$
XIa	$\delta( u^{1/2} ho, u^{-1/2}\sigma)$	$\sigma^2$	$a(\sigma) = 0$	1	$-\omega_{\Lambda}(-1)$
		$\sigma^2$	$a(\sigma) = 1$	$1 + \omega_{\Lambda}(-1)$	$\sigma(-1)(1+\omega_{\Lambda}(-1))$
XIb	$L(.\nu^{1/2}\rho,\nu^{-1/2}\sigma)$	$\sigma^2$	$a(\sigma) = 0$	$1-\omega_{\Lambda}(-1)$	$1-\omega_{\Lambda}(-1)$
	· · · / /	$\sigma^2$	$a(\sigma) = 1$	0	0
s.c.	$\operatorname{c-Ind}_{ZK}^G X_4(\Theta)$	$\widetilde{\Theta}$		$1 - \Theta(t)$	0
s.c.	c-Ind $_{ZK}^{G}X_5(\Lambda,\omega)$	$\widetilde{\Lambda}$		$1 + \omega(-1)$	$-\Lambda(t)(1+\omega(-1))$
s.c.	$\operatorname{c-Ind}_{\mathcal{Z}V}^{G}\theta_2$	1		0	
	$\Delta K^{-2}$	-	I		

Table 7: Si( $\mathfrak{p}^2$ )-fixed vector dimensions and Atkin-Lehner signatures, q odd, continued. Here  $t \in \mathbb{F}_{q^2}^{\times}$  denotes any element with  $t^{q-1} = -1$ .

Class	$\theta_0$	$\theta_1$	$\theta_2$	$\theta_3$	$\theta_4$	$\theta_5$	$\chi_1(k,l)$	$\chi_2(k)$	$\chi_3(k,l)$	$\chi_4(k,l)$
$A_1$	1	$q(q+1)^2/2$	$q(q^2+1)/2$	$q(q^2+1)/2$	$q^4$	$q(q-1)^2/2$	$(q+1)^2(q^2+1)$	$q^4 - 1$	$q^4 - 1$	$(q-1)^2(q^2+1)$
$A_2$	1	q(q+1)/2	-q(q-1)/2	q(q+1)/2		-q(q-1)/2	$(q+1)^2$	$q^2 - 1$	$-q^2 - 1$	$(q-1)^2$
A <sub>31</sub>	1	q(q+1)/2	q(q+1)/2	-q(q-1)/2		-q(q-1)/2	$(q+1)^2$	$-q^2 - 1$	$q^2 - 1$	$(q-1)^2$
A <sub>32</sub>	1	$\frac{q/2}{a/2}$	$\frac{q/2}{-a/2}$	$\frac{q/2}{-a/2}$		$\frac{q/2}{a/2}$	2q + 1 1	-1	-1	-2q + 1
$A_{42}$	1	-q/2	$\frac{q}{2}$	$\frac{q}{2}$		-q/2	1	-1	-1	1
		1/	17	17		17				
$B_1(i,j)$	1	2	1	1	1		$\alpha_{ik}\alpha_{jl} + \alpha_{il}\alpha_{jk}$			
$B_2(i)$	1		1	-1	-1			$-\theta_{ik} - \theta_{qik}$	2	
$B_3(i,j)$	1		-1	1	-1	0			$-\alpha_{ik}\beta_{jl}$	00100
$B_4(i, j)$ $B_7(i)$	1	_1	-1	-1	1	-2				$\rho_{ik}\rho_{jl}+\rho_{il}\rho_{jk}$
$D_{5}(i)$	1	Ť			1	1				
$C_1(i)$	1	q+1	q	1	q		$(q+1)(\alpha_{ik}+\alpha_{il})$		$(q-1)\alpha_{ik}$	
$C_2(i)$	1	q+1	1	q	q		$(q+1)\alpha_{ik}\alpha_{il}$	$(q-1)\alpha_{ik}$		
$C_3(i)$	1		-1	q	-q	q-1		(	$-(q+1)\beta_{il}$	$-(q-1)(\beta_{ik}+\beta_{il})$
$C_4(i)$	1		q	-1	-q	q-1		$-(q+1)\beta_{ik}$		$-(q-1)\beta_{ik}\beta_{il}$
$D_1(i)$	1	1		1			$\alpha_{21} + \alpha_{22}$		$-\alpha$	
$D_{2}(i)$	1	1	1	Ĩ			$\alpha_{ik} + \alpha_{il}$ $\alpha_{ik} \alpha_{il}$	$-\alpha_{ik}$	$\alpha_{ik}$	
$D_3(i)$	1		-1			-1	<i>en * 11</i>	in	$-\beta_{il}$	$\beta_{ik} + \beta_{il}$
$D_4(i)$	1			-1		-1		$-\beta_{ik}$		$\beta_{ik}\beta_{il}$

Table 8: character table for Sp(4, q), q even;  $\alpha_i = \zeta_{q-1}^i + \zeta_{q-1}^{-i}, \beta_i = \zeta_{q+1}^i + \zeta_{q+1}^{-i}, \theta_i = \zeta_{q^2-1}^i + \zeta_{q^2-1}^{-i}$ 

Table 9: character table for Sp(4, q), q even, continued;  $\tau_i = \zeta_{q^2+1}^i + \zeta_{q^2+1}^{-i}$ 

Class	$\chi_5(k)$	$\chi_6(k)$	$\chi_7(k)$	$\chi_8(k)$	$\chi_9(k)$	$\chi_{10}(k)$	$\chi_{11}(k)$	$\chi_{12}(k)$	$\chi_{13}(k)$	
$A_1$	$(q^2 - 1)^2$	$(q+1)(q^2+1)$	$(q+1)(q^2+1)$	$(q-1)(q^2+1)$	$(q-1)(q^2+1)$	$q(q+1)(q^2+1)$	$q(q+1)(q^2+1)$	$q(q-1)(q^2+1)$	$q(q-1)(q^2+1)$	
$A_2$	$-q^2 + 1$	q + 1	$q^2 + q + 1$	q-1	$-q^2 + q - 1$	q(q + 1)	q	q(q - 1)	-q	
$A_{31}$	$-q^2 + 1$	$q^2 + q + 1$	q + 1	$-q^2 + q - 1$	q-1	q	q(q + 1)	-q	q(q - 1)	
$A_{32}$	1	q + 1	q + 1	q-1	q-1	q	q	-q	-q	
$A_{41}$	1	1	1	-1	-1					
$A_{42}$	1	1	1	-1	-1					
- ( )										
$B_1(i,j)$		$\alpha_{ik}\alpha_{jk}$	$\alpha_{ik} + \alpha_{jk}$			$\alpha_{ik}\alpha_{jk}$	$\alpha_{ik} + \alpha_{jk}$			
$B_2(i)$		$\alpha_{ik}$		$-\beta_{ik}$	0	$-\alpha_{ik}$		$-\beta_{ik}$		
$B_3(i,j)$			$\alpha_{ik}$	0.0	$-\beta_{jk}$		$-\alpha_{ik}$	0.0	$-\beta_{jk}$	
$B_4(i,j)$				$-\beta_{ik}\beta_{jk}$	$-\beta_{ik} - \beta_{jk}$			$\beta_{ik}\beta_{jk}$	$\beta_{ik} + \beta_{jk}$	
$B_5(i)$	$\tau_{ik} + \tau_{qik}$									
$\alpha$ (i)		( 1)	. 1 .		1	( 11)			1	
$C_1(i)$		$(q+1)\alpha_{ik}$	$q + 1 + \alpha_{ik}$	- 1	q-1	$(q+1)\alpha_{ik}$	$q + 1 + q\alpha_{ik}$	- 1	q - 1	
$C_2(i)$		$q + 1 + \alpha_{2ik}$	$(q+1)\alpha_{ik}$	q = 1	a 1 8.	$q + 1 + q\alpha_{2ik}$	$(q+1)\alpha_{ik}$	q-1	$a + 1 = a\beta$	
$C_3(i)$		$\alpha \perp 1$	q + 1	$(q-1)\beta_{ik}$	$q = 1 - \beta_{ik}$	a 1	-q - 1	$-(q-1)\rho_{ik}$	$-q + 1 - q\rho_{ik}$	
$C_4(i)$		q + 1		$q = 1 - \rho_{2ik}$	$(q-1)\rho_{ik}$	-q - 1		$-q+1-qp_{2ik}$	$-(q-1)\rho_{ik}$	
$D_1(i)$		0	$1 \pm \alpha v$		_1	0	1		_1	
$D_1(i)$ $D_2(i)$		$1 \pm \alpha_{0ik}$	$1 + \alpha_{ik}$	-1	1	1	0.1	-1		
$D_2(i)$ $D_2(i)$		1   Ct21k	1		$-1 - \beta_{ii}$	1	_1	ß.,	1	
$D_3(i)$ $D_4(i)$		1	1	$-1 - \beta_{av}$	$-\beta_{ik}$	_1		Pik	л В.,	
$\nu_{4}(\iota)$		1		1 P2ik	$\rho_{ik}$	1		1	Pik	

### REFERENCES

# References

Dabbaghian-Abdoly, Vahid. "Characters of some finite groups of Lie type with a [Da2007] restriction containing a linear character once." Journal of Algebra 309.2 (2007): 543-558. Enomoto, Hikoe. "The characters of the finite symplectic group  $Sp(4,q), q = 2^{f}$ ." [En1972] Osaka J. Math 9.1 (1972): 75-94. [Iw1965] Iwahori, Nagayoshi, and Hideya Matsumoto. "On some Bruhat decomposition and the structure of the Hecke rings of p-adic Chevalley groups." Publications Mathématiques de l'IHÉS 25 (1965): 5-48. [J2022] Johnson-Leung, Jennifer, Brooks Roberts, and Ralf Schmidt. Stable Klingen vectors and Paramodular Newforms. Springer, 2023. [L2010] Lust, Jaime. Verifying depth-zero supercuspidal L-packets for inner forms of GSp(4). Diss. UC San Diego, 2010. [Ro2006] Roberts, Brooks, and Ralf Schmidt. "A decomposition of the spaces  $S_k(\Gamma_0(N))$ in degree 2 and the construction of hypercuspidal modular forms." Proceedings of the 9th Autumn Workshop on Number Theory, Hakuba, Japan, 2006. [Ro2007] Roberts, Brooks, and Ralf Schmidt. Local newforms for GSp(4). Vol. 1918. Springer Science and Business Media, 2007. [Ros2016] Rösner, Mirko. Parahoric restriction for GSp(4) and the inner cohomology of Siegel modular threefolds. Heidelberg University PhD Thesis, 2016. Available at https://www.mathi.uni-heidelberg.de/fg-sga Rösner, Mirko. "Parahoric restriction for GSp(4)." Algebras and Representation [Ros2018] Theory 21 (2018): 145-161. [RSY2022] Roy, Manami, Ralf Schmidt, and Shaoyun Yi. "Dimension formulas for Siegel modular forms of level 4." Mathematika 69.3 (2023): 795-840. Sally, Paul, and Marko Tadic. "Induced representations and classifications for [Sa1993] GSp(2, F) and Sp(2, F)." Mém. Soc. Math. France (NS) 52 (1993): 75-133. [Sh1982] Shinoda, Ken-ichi. "The characters of the finite conformal symplectic group, CSp(4,q)." Communications in Algebra 10.13 (1982): 1369-1419. Yi, Shaoyun. "Klingen  $\mathfrak{p}^2$  vectors for GSp(4)." The Ramanujan Journal 54, no. [Yi2021] 3 (2021): 511-554.