

EXTENSIONS OF THE BLOCH - PÓLYA THEOREM ON THE NUMBER OF REAL ZEROS OF POLYNOMIALS (II)

TAMÁS ERDÉLYI

ABSTRACT. We prove that there are absolute constants $c_1 > 0$ and $c_2 > 0$ such that for every

$$a_0, a_1, \dots, a_n \in [1, M], \quad 1 \leq M \leq \exp(c_2 n),$$

there are

$$b_0, b_1, \dots, b_n \in \{-1, 0, 1\}$$

such that the polynomial $P(z) = \sum_{j=0}^n b_j a_j z^j$ has at least

$$c_1 \left(\frac{n}{\log(4M)} \right)^{1/2}$$

distinct sign changes in $I_a := (1-2a, 1-a)$, where $a := \left(\frac{\log(4M)}{n} \right)^{1/2} \leq 1/3$. This improves and extends earlier results of Bloch and Pólya and Erdélyi, and, as a special case, recaptures a special case of a more general recent result of Jacob and Nazarov.

1. INTRODUCTION

In this note c and c_j will denote suitable positive absolute constants. Let \mathcal{F}_n denote the set of polynomials of degree at most n with coefficients from $\{-1, 0, 1\}$. Let \mathcal{L}_n denote the set of polynomials of degree n with coefficients from $\{-1, 1\}$. Let \mathcal{A}_n denote the set of polynomials of degree at most n with coefficients from $\{0, 1\}$. Updating what the authors write in [BEK99] and [Er08] we have the following. The study of the location of zeros of the above classes of polynomials starts with Bloch and Pólya [BP32]. They prove that the average number of real zeros of a polynomial from \mathcal{F}_n is at most $cn^{1/2}$. They also prove that if $n \geq 3$ then a polynomial from \mathcal{F}_n cannot have more than

$$\frac{cn \log \log n}{\log n}$$

Key words and phrases. location of zeros, polynomials with constrained coefficients.
2020 Mathematics Subject Classifications. 26C10, 11C08

real zeros. This quite weak result appears to be the first on this subject. Schur [Sch33] and by different methods Szegő [Sze34] and Erdős and Turán [ET50] improve this to $c(n \log n)^{1/2}$ (see also [BE95]). (Their results are more general, but in this specialization not sharp.) Our Theorem 4.1 in [BEK99] gives the right upper bound of $cn^{1/2}$ for the number of real zeros of polynomials from a much larger class, namely for all polynomials P of the form

$$P(x) = \sum_{j=0}^n a_j x^j, \quad |a_0| = |a_n| = 1, \quad |a_j| \leq 1, \quad a_j \in \mathbb{C}.$$

Schur [Sch33] claims that Schmidt gives a version of part of this theorem. However, it does not appear in the reference he gives, namely [Sch32], and we have not been able to trace it to any other source. In [BE97] we prove that $\eta n^{1/2}$ is an upper bound for the number of zeros of polynomials P of the form

$$P(x) = \sum_{j=0}^n a_j x^j, \quad |a_0| = 1, \quad |a_j| \leq 1, \quad a_j \in \mathbb{C},$$

inside any polygon with vertices on the unit circle, where the constant η depends only on the polygon. Bloch and Pólya [BP32] also proved that there are polynomials $P \in \mathcal{F}_n$ with at least

$$\frac{cn^{1/4}}{(\log n)^{1/2}}$$

sign changes on the real line \mathbb{R} . Schur [Sch33] claims they do it for polynomials with coefficients only from $\{-1, 1\}$, but this appears to be incorrect.

In a seminal paper Littlewood and Offord [LO39] prove that the number of real roots of a polynomial $P \in \mathcal{L}_n$, on average, lies between

$$\frac{c_1 \log n}{\log \log \log n} \quad \text{and} \quad c_2 (\log n)^2,$$

and it is proved by Boyd [Bo97] that every $P \in \mathcal{L}_n$ has at most $\frac{c(\log n)^2}{\log \log n}$ zeros at 1 (in the sense of multiplicity).

Kac [Ka48] prove that the expected number of real roots of a polynomial of degree n with random uniformly distributed independent coefficients is asymptotically $(2/\pi) \log n$.

In [Er08] we prove that there are absolute constants $c_1 > 0$ and $c_2 > 0$ such that for every

$$a_0, a_1, \dots, a_n \in [1, M], \quad 1 \leq M \leq \exp(c_2 n^{1/4}),$$

there are $b_0, b_1, \dots, b_n \in \{-1, 0, 1\}$ such that $P(z) = \sum_{j=0}^n b_j a_j z^j$ has at least $c_2 n^{1/4}$ distinct

sign changes in $(0, 1)$. This improves and extends the earlier result of Bloch and Pólya [BP32] and, as a special case, it recaptures a special case

Jacob and Nazarov [JN24] prove quite a general theorem including that there are polynomials $P \in \mathcal{L}_n$ having at least $cn^{1/2}$ distinct zeros in $[0, 1]$; and there are polynomials $0 \neq P \in \mathcal{A}_n$ for which the polynomials Q defined by $Q(x) := P(-x)$ have at least $cn^{1/2}$ distinct zeros in $[0, 1]$.

2. NEW RESULT

Theorem 2.1. *There are absolute constants $c_1 > 0$ and $c_2 > 0$ such that for every*

$$a_0, a_1, \dots, a_n \in [1, M], \quad 1 \leq M \leq \exp(c_2 n),$$

there are

$$b_0, b_1, \dots, b_n \in \{-1, 0, 1\}$$

such that the polynomial P of the form $P(z) = \sum_{j=0}^n b_j a_j z^j$ has at least

$$c_1 \left(\frac{n}{\log(4M)} \right)^{1/2}$$

distinct sign changes in $I_a := (1 - 2a, 1 - a)$, where $a := \left(\frac{\log(4M)}{n} \right)^{1/2} \leq 1/3$.

This improves and extends earlier results of Bloch and Pólya [BP32] and Erdélyi [Er08], and recaptures a special case of a more general result of Jacob and Nazarov [JN24].

3. LEMMAS

We use the notation $\log^- z := \max\{0, -\log z\}$ and $\log^+ z := \max\{0, \log z\}$. The geometric mean $L_0([a, b])$ of a continuous function P on a finite interval $[a, b] \subset \mathbb{R}$ is

$$\|P\|_{L_0([a, b])} := \exp \left(\frac{1}{b-a} \int_a^b \log |P(x)| dx \right).$$

In [JN24] Jacob and Nazarov showed the following.

Lemma 3.1. *Let $a \in (0, 1/3)$ and $I_a := (1 - 2a, 1 - a)$. Then*

$$\int_{I_a} \log^- P(x) dx \leq c_3 \log(4M)$$

for every polynomial P of the form

$$(3.1) \quad P(x) = \sum_{j=0}^n b_j a_j x^j, \quad a_j \in [1, M], \quad b_j \in \{-1, 0, 1\}, \quad b_0 = \pm 1.$$

Moreover $c_3 := 9\pi/\sqrt{2}$ is a suitable choice.

As a consequence we have the following.

Lemma 3.2. *Let $a \in (0, 1/3)$ and $I_a := (1 - 2a, 1 - a)$. Then*

$$\|P\|_{L_0(I_a)} \geq \exp\left(\frac{-c_3 \log(4M)}{a}\right)$$

for every polynomial P of the form (3.1).

We will need the version of Lemma 3.2 below in the proof of Theorem 2.1.

Lemma 3.3. *Let $a \in (0, 1/3)$, $I_a := (1 - 2a, 1 - a)$, and $0 \leq k \leq n$. Then*

$$\|R\|_{L_0(I_a)} \geq \exp\left(\frac{-c_3 \log(4M)}{a}\right)$$

for every polynomial R of the form

$$(3.2) \quad R(x) = \sum_{j=k}^n b_j a_j x^{j-k}, \quad a_j \in [1, M], \quad b_j \in \{-1, 0, 1\}, \quad b_k = \pm 1.$$

4. PROOFS

Proof of Lemma 3.2. Using Lemma 3.1 we have

$$\begin{aligned} \|P\|_{L_0(I_a)} &= \exp\left(\frac{1}{a} \int_{I_a} \log^+ P(x) dx - \frac{1}{a} \int_{I_a} \log^- P(x) dx\right) \\ &\geq \exp\left(\frac{-1}{a} \int_{I_a} \log^- P(x) dx\right) \\ &\geq \exp\left(\frac{-c_3 \log(4M)}{a}\right). \end{aligned}$$

□

Proof of Theorem 2.1. We modify the outline given by Nazarov in MathOverflow where $a_j = 1$ for each $j = 0, 1, \dots, n$, is assumed. Let $a \in (0, 1/3)$ and $I_a := (1 - 2a, 1 - a)$. Associated with a polynomial P we define the polynomial $\tilde{P}(x) := P(1 - a - ax)$. The function $f(x) := 1 - a - ax$ maps the interval $(0, 1)$ onto the interval $(1 - 2a, 1 - a)$, hence the polynomial P of the form

$$P(x) = \sum_{j=0}^n a_j x^j, \quad a_j \in [1, M],$$

satisfies

$$0 \leq \tilde{P}(x) < (n + 1)M, \quad x \in [0, 1].$$

For every fixed $a_0, a_1, \dots, a_n \in [1, M]$ there are 2^{n+1} different polynomials P of the form

$$(4.1) \quad P(x) = \sum_{j=0}^n d_j a_j x^j, \quad d_j \in \{0, 1\}.$$

By the pigeon hole principle there are two different polynomials P_1 and P_2 of the form (4.1) such that the polynomial $0 \neq P := P_1 - P_2$ is of the form (3.1) and

$$(4.2) \quad \left| \int_0^1 \tilde{P}(x) x^j dx \right| \leq (n+1)M2^{-n/m}, \quad j = 0, 1, \dots, m-1.$$

We fix this P and $b_0, b_1, \dots, b_n \in \{-1, 0, 1\}$ in the rest of the proof. There is a $0 \leq k \leq n$ such that $b_0 = b_1 = \dots = b_{k-1} = 0$ and $b_k = \pm 1$. We have $P(x) = x^k R(x)$, where R is a polynomial of the form (3.2) and

$$x^k \geq (1 - 2a)^n \geq \exp(-4an), \quad x \in I_a, \quad a \in (0, 1/3).$$

Combining this with Lemma 3.3 gives

$$\|P\|_{L_0(I_a)} \geq \exp\left(\frac{-c_3 \log(4M)}{a} - 4an\right).$$

Now we choose $a := \left(\frac{\log(4M)}{n}\right)^{1/2} \leq 1/3$ to get

$$(4.3) \quad \|\tilde{P}\|_{L_0([0,1])} = \|P\|_{L_0(I_a)} \geq \exp(-c_4 \log(4M)n^{1/2}).$$

Assume that P has exactly $m-1$ sign changes in I_a . Then \tilde{P} has exactly $m-1 \geq 0$ sign changes in $(0, 1)$. We denote the sign changes of \tilde{P} in $(0, 1)$ by x_1, x_2, \dots, x_{m-1} . We define

$$Q(x) := \prod_{j=1}^{m-1} (x - x_j)$$

(the empty product is defined to be 1). Then $\tilde{P}Q$ does not change sign in $(0, 1)$. Observe that

$$\|x - x_j\|_{L_0([0,1])} \geq c_5, \quad j = 1, 2, \dots, m-1,$$

hence

$$(4.4) \quad \|Q\|_{L_0([0,1])} \geq \exp(-c_6 m).$$

Combining (4.3), (4.4), the multiplicative property of the geometric mean, the comparison between the $L_0([0, 1])$ and $L_1([0, 1])$ norms, the fact that $\tilde{P}Q$ does not change sign, (4.2), and the fact that the sum of the absolute values of the coefficients of Q is at most 2^m , we obtain

$$\begin{aligned} \exp(-c_4 \log(4M)n^{1/2} - c_6 m) &\leq \|\tilde{P}Q\|_{L_0([0,1])} \leq \|\tilde{P}Q\|_{L_1([0,1])} = \left| \int_0^1 (\tilde{P}Q)(x) dx \right| \\ &\leq 2^m (n+1)M2^{-n/m}. \end{aligned}$$

Taking the log of both sides above and choosing

$$m := \left\lfloor \frac{c_7 n^{1/2}}{\log(4M)} \right\rfloor$$

with a sufficiently small absolute constant $c_7 > 0$, we get a contradiction. \square

5. ACKNOWLEDGMENT

The author thanks Fedor Nazarov for the discussions on the topic and for presenting the main idea of the proof of Theorem 2.1 on MathOverflow.

REFERENCES

- BP32. A. Bloch and G. Pólya, *On the roots of certain algebraic equations*, Proc. London Math. Soc. **33** (1932), 102–114.
- BE95. P. Borwein and T. Erdélyi, *Polynomials and Polynomial Inequalities*, Springer-Verlag, New York, 1995.
- BE97. P. Borwein and T. Erdélyi, *On the zeros of polynomials with restricted coefficients*, Illinois J. Math. **41** (1997), 667–675.
- BEK99. P. Borwein, T. Erdélyi, and G. Kós, *Littlewood-type problems on $[0, 1]$* , Proc. London Math. Soc. **79** (1999), 22–46.
- Bo97. D.W. Boyd, *On a problem of Byrne’s concerning polynomials with restricted coefficients*, Math. Comp. **66** (1997), 1697–1703.
- Er08. T. Erdélyi, *Extensions of the Bloch-Pólya theorem on the number of real zeros of polynomials*, Journal de théorie des nombres de Bordeaux **20**, no. **2** (2008), 281–287.
- ET50. P. Erdős and P. Turán, *On the distribution of roots of polynomials*, Ann. Math. **57** (1950), 105–119.
- JN24. M. Jacob and F. Nazarov, *The Newman algorithm for constructing polynomials with restricted coefficients and many real roots*, <https://arxiv.org/pdf/2404.07971>.
- Ka48. M. Kac, *On the average number of real roots of a random algebraic equation (II)*, Proc. London Math. Soc. **50** (1948), 390–408.
- LO39. J.E. Littlewood and A.C. Offord, *On the number of real roots of a random algebraic equation. II*, Proc. Cam. Phil. Soc. **35**, 133–148.
- Sch32. E. Schmidt, *Über algebraische Gleichungen vom Pólya-Bloch-Typos*, Sitz. Preuss. Akad. Wiss., Phys.-Math. Kl. (1932), 321.
- Sch33. I. Schur, *Untersuchungen über algebraische Gleichungen.*, Sitz. Preuss. Akad. Wiss., Phys.-Math. Kl. (1933), 403–428.
- Sze34. G. Szegő, *Bemerkungen zu einem Satz von E. Schmidt über algebraische Gleichungen.*, Sitz. Preuss. Akad. Wiss., Phys.-Math. Kl. (1934), 86–98.

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843
E-mail address: `terdelyi@math.tamu.edu`