

A SHORT NONSTANDARD PROOF OF THE SPECTRAL THEOREM FOR UNBOUNDED SELF-ADJOINT OPERATORS

TAKASHI MATSUNAGA

ABSTRACT. By nonstandard analysis, a very short and elementary proof of the Spectral Theorem for unbounded self-adjoint operators is given.

1. INTRODUCTION

The Spectral Theorem for unbounded self-adjoint operators is one of the most fundamental theorems in functional analysis. The proofs in standard mathematics are long and not straightforward.

On the other hand, Luxemburg offered a nonstandard proof of the Spectral Theorem for bounded self-adjoint operators in [4], while Moore gave another nonstandard proof of the theorem for bounded self-adjoint operators via the nonstandard hull construction in [5]. Yamashita and Ozawa [7] gave three equivalent definitions of the nonstandard hull of internal bounded (possibly unbounded) self-adjoint operators. Recently, Goldbring presented a nonstandard proof of the theorem for unbounded self-adjoint operators using the projection-valued Loeb measure in [2] inspired by Raab's [6] work.

Suggested by the papers cited above, we give a very short and elementary nonstandard proof of the Spectral Theorem for unbounded self-adjoint operators.

2. PRELIMINARIES

Our framework of nonstandard analysis is presented in Davis [1]. In this framework, we have countable saturation and there exists a *-finite dimensional linear space that includes a given linear space. Throughout this note, we follow the arguments and notations of Moore [5] unless otherwise stated.

Let T be a (possibly unbounded) self-adjoint operator on a dense subspace $D(T)$ of a complex Hilbert space H and let S be a *-finite dimensional subspace of ${}^*D(T)$ such that $D(T) \subseteq \hat{S}$, where \hat{S} denotes the nonstandard hull of S . \hat{S} is a Hilbert space by countable saturation. Since $D(T)$ is dense in H , we can assume $H \subseteq \hat{S}$. Let $\pi : \text{fin}(S) \rightarrow \hat{S}$ be the natural projection, where $\text{fin}(S) = \{x \in S \mid {}^*\|x\| \text{ is finite}\}$. If A is a *-linear operator on S such that ${}^*\|A\|$ is finite, one can define the nonstandard hull \hat{A} of A on \hat{S} by setting $\pi(Ax) = \hat{A}(\pi(x))$. Let T_S denote the restriction of $P_S T$ to S , where P_S is the internal projection of *H to S .

Since T_S is a *-finite dimensional self-adjoint operator, the internal spectral family $F(\lambda)(\lambda \in {}^*\mathbb{R})$ exists by transferring the finite dimensional Spectral Theorem in [3]. For $\lambda \in \mathbb{R}$ and $n \in \mathbb{N}$, $\hat{F}(\lambda + 1/n)$ are internal non-increasing projections.

2020 Mathematics Subject Classification. Primary nonstandard functional analysis 46S20; Secondary linear symmetric and selfadjoint operators (unbounded) 47B25.

Hence we can define $E(\lambda) = \lim_{n \rightarrow \infty} \hat{F}(\lambda + 1/n)$. Thus we obtain a family of right-continuous non-decreasing projections $E(\lambda)$ on \hat{S} .

Lemma 2.1. *If $K \in {}^*\mathbb{R}$ is positive infinite, for $x \in D(T)$*

$$\begin{aligned} \pi\left({}^*\int_{|\lambda| \geq K} dF(\lambda)x\right) &= 0, & \pi\left({}^*\int_{|\lambda| \geq K} \lambda dF(\lambda)x\right) &= 0, \\ \lim_{\lambda \rightarrow -\infty} E(\lambda)x &= 0, & \lim_{\lambda \rightarrow \infty} E(\lambda)x &= x. \end{aligned}$$

Proof. Since $F(\lambda)$ is the internal spectral family of T_S , for $x \in D(T)$

$${}^*\|T_S x\|^2 = {}^*\int_{-\infty}^{\infty} \lambda^2 d(F(\lambda)x, x) \geq K^2 {}^*\int_{|\lambda| \geq K} d(F(\lambda)x, x)$$

is finite, we obtain

$${}^*\int_{|\lambda| \geq K} d(F(\lambda)x, x) \simeq 0, \quad (\#)$$

which leads to the first equality. For the second equality, using the Cauchy-Schwartz inequality, one can get

$$\left({}^*\int_{|\lambda| \geq K} \lambda d(F(\lambda)x, x)\right)^2 \leq \left({}^*\int_{|\lambda| \geq K} \lambda^2 d(F(\lambda)x, x)\right) \left({}^*\int_{|\lambda| \geq K} d(F(\lambda)x, x)\right).$$

Since again ${}^*\|T_S x\|^2 \geq {}^*\int_{|\lambda| \geq K} \lambda^2 d(F(\lambda)x, x)$ is finite, applying (#) we have ${}^*\int_{|\lambda| \geq K} \lambda d(F(\lambda)x, x) \simeq 0$. Using the Polarization Identity, it follows that for $y \in D(T)$

$$|{}^*\int_{|\lambda| \geq K} \lambda d(F(\lambda)x, y)| \simeq 0$$

to obtain the second equality. The third and fourth equalities easily follow by definition. \square

Lemma 2.2. *If $E(\lambda)x$ is continuous in λ at $a \in \mathbb{R}$, for $x \in D(T)$*

$$\pi(F(a)x) = E(a)x.$$

Proof. Since $E(\lambda)x$ is continuous at $a \in \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \pi\left(F\left(a + \frac{1}{n}\right)x\right) = \lim_{n \rightarrow \infty} \pi\left(F\left(a - \frac{1}{n}\right)x\right) = E(a)x.$$

On the other hand, we have

$${}^*\|F\left(a + \frac{1}{n}\right)x - F\left(a - \frac{1}{n}\right)x\|^2 = {}^*\|F\left(a + \frac{1}{n}\right)x - F(a)x\|^2 + {}^*\|F(a)x - F\left(a - \frac{1}{n}\right)x\|^2$$

because $F(\lambda)$ is an internal spectral family. From these equalities, the conclusion follows. \square

Lemma 2.3. (the Representation Lemma) *If $E(\lambda)x$ is continuous in λ at $a < b \in \mathbb{R}$, for $x \in D(T)$*

$$\pi\left({}^*\int_a^b \lambda dF(\lambda)x\right) = \int_a^b \lambda dE(\lambda)x.$$

Proof. Let $E(\lambda)x$ is continuous in λ at $a = a_0 < a_1 < \dots < a_n = b \in \mathbb{R}$. Using Lemma 2.2 to obtain the first and last equalities and noting that $F(\lambda)$ is an internal spectral family to get the third and fourth equalities, we have

$$\begin{aligned}
& \|\pi(* \int_a^b \lambda dF(\lambda)x) - \sum_{k=1}^n a_k((E(a_k) - E(a_{k-1}))x)\|^2 \\
&= \|\sum_{k=1}^n \pi(* \int_{a_{k-1}}^{a_k} \lambda dF(\lambda)x) - \sum_{k=1}^n \pi(a_k((F(a_k) - F(a_{k-1}))x))\|^2 \\
&= \|\sum_{k=1}^n \pi(* \int_{a_{k-1}}^{a_k} (\lambda - a_k)dF(\lambda)x)\|^2 = \sum_{k=1}^n \|\pi(* \int_{a_{k-1}}^{a_k} (\lambda - a_k)dF(\lambda)x)\|^2 \\
&\leq \sum_{k=1}^n \|\pi(* \int_{a_{k-1}}^{a_k} (a_{k-1} - a_k)dF(\lambda)x)\|^2 \leq \max_k (a_k - a_{k-1})^2 \sum_{k=1}^n \|\pi((F(a_k) - F(a_{k-1}))x)\|^2 \\
&= \max_k (a_k - a_{k-1})^2 \|\pi((F(b) - F(a))x)\|^2 = \max_k (a_k - a_{k-1})^2 \|((E(b) - E(a))x)\|^2.
\end{aligned}$$

Since $E(\lambda)x$ is right-continuous in λ , one can choose a_k 's such that $\max_k (a_k - a_{k-1})^2 < \epsilon$ for any $\epsilon > 0$. Letting $\epsilon \downarrow 0$, the desired result follows. \square

3. A NONSTANDARD PROOF THE SPECTRAL THEOREM

Lemma 3.1. For $x \in D(T)$

$$Tx = \pi(T_S x) = \pi(* \int_{-\infty}^{\infty} \lambda dF(\lambda)x) = \int_{-\infty}^{\infty} \lambda dE(\lambda)x.$$

Proof. The first equality is by construction. The second equality is just the spectral resolution of the *-finite dimensional self-adjoint operator T_S . For the third equality, applying Lemmas 2.1 and 2.3, let $a \rightarrow -\infty$ and $b \rightarrow \infty$. \square

Lemma 3.2. (the Operational Calculus) Suppose that $z \in \mathbb{C} \setminus \mathbb{R}$, then for $x \in H$

$$\int_{-\infty}^{\infty} \frac{1}{\lambda - z} dE(\lambda)x = (T - z)^{-1}x \in D(T).$$

Proof. Since T is self-adjoint, there exists $y \in D(T)$ such that $(T - z)y = x$. Using the previous lemma, we have

$$\begin{aligned}
& \int_{-\infty}^{\infty} \frac{1}{\lambda - z} dE_\lambda(\lambda)(T - z)y = \int_{-\infty}^{\infty} \frac{1}{\lambda - z} d_\lambda E(\lambda) \int_{-\infty}^{\infty} (\mu - z) d_\mu E(\mu)y \\
&= \int_{-\infty}^{\infty} \frac{1}{\lambda - z} d_\lambda \int_{-\infty}^{\infty} (\mu - z) d_\mu E(\lambda)E(\mu)y = \int_{-\infty}^{\infty} \frac{1}{\lambda - z} d_\lambda \int_{-\infty}^{\lambda} (\mu - z) d_\mu E(\mu)y \\
&= \int_{-\infty}^{\infty} \frac{1}{\lambda - z} (\lambda - z) d_\lambda E(\lambda)y = \int_{-\infty}^{\infty} d_\lambda E(\lambda)y = y.
\end{aligned}$$

This leads to the conclusion. \square

Lemma 3.3. $E(\lambda)$ is a spectral family on H .

Proof. Recall an elementary formula

$$\lim_{\epsilon \downarrow 0} \frac{1}{2\pi i} \int_a^b \left(\frac{1}{\lambda - \mu - i\epsilon} - \frac{1}{\lambda - \mu + i\epsilon} \right) d\mu = 0 (\lambda < a, \lambda > b), \quad 1 (a < \lambda < b).$$

From this formula, if $E(\lambda)x$ is continuous in λ at $a < b \in \mathbb{R}$, using the previous lemma to get the fourth equality and the fifth set membership, we have for $x \in D(T)$

$$\begin{aligned} (E(b) - E(a))x &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} \lim_{\epsilon \downarrow 0} \int_a^b \left(\frac{1}{\lambda - \mu - i\epsilon} - \frac{1}{\lambda - \mu + i\epsilon} \right) d\mu dE(\lambda)x \\ &= \lim_{\epsilon \downarrow 0} \frac{1}{2\pi i} \int_{-\infty}^{\infty} \int_a^b \left(\frac{1}{\lambda - \mu - i\epsilon} - \frac{1}{\lambda - \mu + i\epsilon} \right) d\mu dE(\lambda)x \\ &= \lim_{\epsilon \downarrow 0} \frac{1}{2\pi i} \int_a^b \int_{-\infty}^{\infty} \left(\frac{1}{\lambda - \mu - i\epsilon} - \frac{1}{\lambda - \mu + i\epsilon} \right) dE(\lambda) d\mu x \\ &= \lim_{\epsilon \downarrow 0} \frac{1}{2\pi i} \int_a^b \left((T - \mu - i\epsilon)^{-1} - (T - \mu + i\epsilon)^{-1} \right) d\mu x \in H, \end{aligned}$$

which completes the proof because $E(\lambda)$ is a right-continuous non-decreasing family of projections with $\lim_{\lambda \rightarrow -\infty} E(\lambda)x = 0$ and $\lim_{\lambda \rightarrow \infty} E(\lambda)x = x$ for $x \in D(T)$ (lemma 2.1). \square

The lemmas above lead to

Theorem 3.4. For $x \in D(T)$

$$Tx = \int_{-\infty}^{\infty} \lambda dE(\lambda)x,$$

where $E(\lambda)$ is a spectral family on H .

4. CONCLUDING REMARKS

For those who are familiar with nonstandard analysis, this is a very short and elementary nonstandard proof of the Spectral Theorem for unbounded self-adjoint operators.

REFERENCES

- [1] M. Davis, *Applied nonstandard analysis*, John Wiley & Sons (1977), Dover Publications (2005).
- [2] I. Goldbring, *A nonstandard proof of the spectral theorem for unbounded self-adjoint operators*, *Expositiones Mathematicae*, **39** (2021), 590-603.
- [3] P.R. Halmos, *Finite-dimensional vector spaces, 2nd edition*, Van Nostrand (1958), Dover Publications (2017).
- [4] W.A.J. Luxemburg, *On some concurrent binary relations occurring in analysis*, *Studies in Logic and the Foundations of Mathematics*, **69** (1972), 85-100.
- [5] L.C. Moore, Jr., *Hyperfinite extensions of bounded operators on a separable Hilbert space*, *Transactions of the American Mathematical Society*, **218** (1976), 285-295.
- [6] A. Raab, *An approach to nonstandard quantum mechanics*, *Journal of Mathematical Physics*, **45**(2004), 4791-4809.
- [7] H Yamashita and M Ozawa, *Nonstandard representations of unbounded self-adjoint operators*, *RIMS Kokyuroku*, **1186**(2001), 119-124.

DEPARTMENT OF MEDICAL INFORMATICS, OSAKA INTERNATIONAL CANCER INSTITUTE, 3-1-69 OTEMAE, CHUO, OSAKA CITY, 541-8567, JAPAN

Email address: matsunaga-ta@nifty.com