

# NONLOCAL LIOUVILLE THEOREMS WITH GRADIENT NONLINEARITY

ANUP BISWAS, ALEXANDER QUAAS, AND ERWIN TOPP

**ABSTRACT.** In this article we consider a large family of nonlinear nonlocal equations involving gradient nonlinearity and provide a unified approach, based on the Ishii-Lions type technique, to establish Liouville property of the solutions. We also answer an open problem raised by [19]. Some applications to regularity issues are also studied.

## 1. INTRODUCTION

The main goal of this article is to study Liouville property of nonlinear nonlocal equations of the form

$$-\mathcal{I}u + H(u, \nabla u) = 0 \quad \text{in } \mathbb{R}^n,$$

where the Hamiltonian  $H(r, p)$  are models of the forms (I)  $|p|^m$  for  $m > 1$ , (II)  $-r^q|p|^m$  and (III)  $r^q|p|^m$ , and  $\mathcal{I}$  is a Pucci type nonlinear operator of order  $2s$ ,  $s \in (0, 1)$ . Liouville property of the solutions to various nonlinear partial differential equations has been a central topic of research not only because of its own importance, but also due to its applications in the regularity theory (see [11, 35, 36, 38]), a priori bounds [6] etc. This particular article is inspired from the recent survey of Cirant and Goffi [19] where the authors provide a comprehensive account on the the development on various Liouville theorems for classical (or local) elliptic equations. The nonlocal analogue of most of these Liouville theorems remained open because of several difficulties stemming from the nonlocal nature of the equations and unavailability of appropriate gradient bound, also known as Bernstein estimate. In this article we propose a unified approach to prove Liouville property for a large class of nonlinear nonlocal operators including gradient nonlinearity of type I, II and III above.

For  $H = 0$ , the above Liouville property is quite well studied and have also been used to establish interior regularity of solutions [21, 35, 36, 38]. The case  $H(u, \nabla u) = -u^q$  is known as the Lane-Emden equation and has been extensively studied in literature, see [9, 17, 34, 42] and references therein. Recently, Barrios and Del Pezzo [7] study Liouville property of positive solutions to the equations  $(-\Delta)^s u + |\nabla u|^m = \lambda u^q$  in the exterior domains. The authors rely on the maximum principles to obtain an appropriate decay estimate of the solutions at infinity which helps them to establish the required results. This particular idea in the local case goes back to the work of Alarcón et. al. [2], see also the earlier results by Cutri and Leoni [20] and Armstrong and Sirakov in [4]; and in the context of nonlocal equation see Felmer and second author [24] and the references therein. We should also mention the work of Alibaud et. al.[3], Berger and Schilling [13], Fall and Weth [22] which study Liouville property for a general linear integro-differential operator with constant coefficients. In fact, [3, 13] provides a necessary and sufficient condition for the Liouville property to hold in the class of bounded solutions.

To understand the difficulty in treating problems considered in this article, let us focus on the model I. That is, the Hamiltonian is of the form  $H(u, \nabla u) = |\nabla u|^m$ . For the classical elliptic operator, that is  $\mathcal{I} = \Delta$ , Liouville property is obtained from the gradient estimate. More precisely,

---

2010 *Mathematics Subject Classification.* Primary: 35J60, 35B53.

*Key words and phrases.* Lipschitz regularity, Bernstein estimate, nonexistence, regularity, Hamilton-Jacobi equations.

since  $v(x) := R^{-\frac{m-2}{m-1}}u(Rx)$ ,  $R > 0$ , satisfies the same equation, from the gradient estimate it follows that

$$|\nabla u(x)| \leq CR^{-\frac{1}{m-1}} \quad x \in B_{R/2}.$$

Letting  $R \rightarrow \infty$ , we see that  $u = \text{const}$ . This idea dates back to the work of Peletier and Serrin [33], Lions [32]. Later similar gradient estimate is used to obtain Liouville property for model II, see Bidaut-Véron, García-Huidobro and Véron [14], Filipucci, Pucci and Souplet [25]. Therefore, gradient estimate plays a key role in establishing the Liouville property. Surprisingly, this Bernstein type gradient estimate remained a challenging problem for the nonlocal operators (see [15, 28]). Very recently, the first and third author in [10] obtained a gradient estimate which seems to be far from being optimal and only works for model I.

Our method in this article is based on Ishii-Lions type argument which was introduced by Ishii and Lions in [31] to obtain Hölder regularity of classical viscosity solutions. Recall that, Ishii-Lions argument is applied in a ball of fixed radius and with an appropriate test function with a large Hölder constant. Recently, this argument has also been employed by Barles et. al. [12] to establish Hölder regularity for a large class of nonlinear nonlocal equations. We apply a similar Ishii-Lions method but in large balls with a small Hölder constant and then making use of the ellipticity, we establish the results. Interestingly, this technique works for a large class of nonlinear equations and therefore, produces a unified approach in obtaining Liouville property. As can be seen from the proofs below, similar technique can also be applied to nonlocal operators with kernels of variable order and more general integro-differential operators with non-degenerate second order elliptic term. In the later part of this article we also apply Liouville theorems to establish certain regularity results. For instance, in Theorem 4.1 we treat a nonlinear nonlocal equation corresponding to the fractional order  $s = \frac{1}{2}$  and show that the viscosity solutions are  $C^\gamma$  for any  $\gamma \in (0, 1)$ . This extends the result of Schwab and Silvestre [37] which obtains  $C^\alpha$  regularity for some  $\alpha \in (0, 1)$ .

The rest of the article is organized as follows: in the next Section we introduce our model and state all our main results on Liouville property. Section 3 is devoted to the proof of these results. Finally, in Section 4 we provide two applications of our Liouville theorems to obtain some regularity results.

**1.1. Setting and main results.** Given ellipticity constants  $0 < \lambda \leq \Lambda$ , we denote by  $\mathcal{K}_s = \mathcal{K}_s(\lambda, \Lambda)$  the family of nonnegative, measurable kernels  $K : \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{R}$  for which the following condition holds

$$\frac{\lambda}{|z|^{n+2s}} \leq K(z) \leq \frac{\Lambda}{|z|^{n+2s}}, \quad z \neq 0. \quad (1.1)$$

Note that we do not assume  $K$  to be symmetric with respect to 0.

For each  $K \in \mathcal{K}_s$ , we denote

$$L_K u(x) = \int_{\mathbb{R}^d} [u(x+z) - u(x) - 1_{\{2s \in [1,2)\}} 1_B(z) \nabla u(x) \cdot z] K(z) dz,$$

which is well-defined for adequately regular functions  $u : \mathbb{R}^n \rightarrow \mathbb{R}$ . Here  $B$  denotes the unit ball in  $\mathbb{R}^n$  around 0. Let  $\mathcal{I}$  be a translation invariant, positively 1-homogeneous, nonlocal operator which is elliptic with respect to the class  $\mathcal{K}_s$ , that is,

$$\begin{aligned} \mathcal{I}u_h(x) &= \mathcal{I}u(x+h) \quad \text{for all } x, h \in \mathbb{R}^n, \quad \text{where } u_h := u(\cdot + h), \\ \inf_{K \in \mathcal{K}_s} (L_K u(x) - L_K v(x)) &\leq \mathcal{I}u(x) - \mathcal{I}v(x) \leq \sup_{K \in \mathcal{K}_s} (L_K u(x) - L_K v(x)) \end{aligned} \quad (1.2)$$

whenever  $u, v \in C^{1,1}(x) \cap L^1(\omega_s)$  where  $\omega_s(z) = (1 + |z|)^{-n-2s}$ . Note that for  $u \in C^{1,1}(x) \cap L^1(\omega_s)$  and  $v \in C^{1,1}(y) \cap L^1(\omega_s)$ , letting  $\tilde{v} = v(\cdot + y - x)$ , we also have from (1.2) that

$$\inf_{K \in \mathcal{K}_s} (L_K u(x) - L_K v(y)) \leq \mathcal{I}u(x) - \mathcal{I}v(y) \leq \sup_{K \in \mathcal{K}_s} (L_K u(x) - L_K v(y)). \quad (1.3)$$

We also denote by  $\mathcal{K}_{\text{sym}} = \mathcal{K}_{\text{sym}}(\lambda, \Lambda)$  the subclass of kernels in  $\mathcal{K}_s$  which are symmetric, that is,  $K(z) = K(-z)$  for all  $z \neq 0$ . Some of our results would use a translation invariant, positively 1-homogeneous, nonlocal operator  $\mathcal{I}_{\text{sym}}$  which is elliptic with respect to the class  $\mathcal{K}_{\text{sym}}$ .

Now we are ready to state our first main result.

**Theorem 1.1.** *Let  $s \in (\frac{1}{2}, 1)$  and  $H : \mathbb{R}^n \rightarrow \mathbb{R}$  be a continuous function satisfying the following: for every  $\varepsilon, L > 0$  there exists  $C = C(\varepsilon, L)$  satisfying*

$$|H(p) - H(q)| \leq C |p - q| \quad (1.4)$$

for all  $\varepsilon \leq |p|, |q| \leq L$ . Consider a viscosity solution  $u$  to

$$-\mathcal{I}u + H(\nabla u) = 0 \quad \text{in } \mathbb{R}^n. \quad (1.5)$$

If for some  $\gamma \in [0, \frac{1}{2s})$  and  $M > 0$  we have

$$|u(x)| \leq M(1 + |x|)^\gamma \quad x \in \mathbb{R}^n, \quad (1.6)$$

then  $u$  is necessarily a constant.

Furthermore, for  $s \in (0, \frac{1}{2}]$  the same conclusion holds, provided  $\gamma \in [0, 2s)$  and one of the following holds:

(i)  $u$  is uniformly continuous in  $\mathbb{R}^n$  and  $H$  is locally Lipschitz in the sense that for every  $L > 0$  there exists  $C = C(L)$  satisfying

$$|H(p) - H(q)| \leq C |p - q| \quad \text{for all } |p|, |q| \leq L. \quad (1.7)$$

(ii)  $H$  is globally Lipschitz.

**Remark 1.1.** *It is interesting to note that we do not require  $H$  to be locally Lipschitz for (1.4) to hold. Also, under the stated conditions of Theorem 1.1, (1.5) does not have any solution unless  $H(0) = 0$ . Again, for  $s > 1/2$ , we can not take the growth exponent  $\gamma$  to be 1. For instance, if we consider the equation*

$$(-\Delta)^s u + |e \cdot \nabla u|^m = 0 \quad \text{in } \mathbb{R}^n, \quad m > 0$$

for some unit vector  $e$ , then  $u(x) = e^\perp \cdot x$  is a non-zero solution for any unit vector  $e^\perp$  satisfying  $e^\perp \cdot e = 0$ .

Theorem 1.1 deals with a quite general Hamiltonian in the sense that we do not impose any sign or growth conditions on  $H$ . If we impose a slightly restrictive condition on  $H$  in Theorem 1.1, the Liouville property holds for a larger class of functions, as stated in the following theorem

**Theorem 1.2.** *Let  $s > 1/2$  and  $H : \mathbb{R}^n \rightarrow \mathbb{R}$  be a locally Lipschitz function satisfying*

$$|H(p) - H(q)| \leq C_H (|p|^{m-1} + |q|^{m-1}) |p - q| \quad \text{for all } p, q \in \mathbb{R}^n, \quad (1.8)$$

where  $m > 1$  and  $C_H > 0$ . Let  $\mathcal{I}_{\text{sym}}$  be a translation invariant, positively 1-homogeneous operator which is elliptic with respect to the set of all symmetric kernels  $\mathcal{K}_{\text{sym}}$ . Consider the equation

$$-\mathcal{I}_{\text{sym}}u + H(\nabla u) = 0 \quad \text{in } \mathbb{R}^n.$$

Then there is no non-constant solution  $u$  satisfying

$$|u(x)| \leq M(1 + |x|)^\gamma, \quad x \in \mathbb{R}^n$$

for some  $\gamma < \max\{\frac{m-2s}{m-1}, \frac{1}{2s}\}$ .

It is also interesting to ask whether there exists a solution that is comparable to  $|x|^\gamma$  near infinity. Our next result gives an answer to this question.

**Theorem 1.3.** *Suppose that  $u$  is a non-trivial solution to*

$$-\mathcal{I}_{\text{sym}}u + |\nabla u|^m = 0 \quad \text{in } \mathbb{R}^n \quad (1.9)$$

for  $m > 1$  and  $s > \frac{1}{2}$ , that satisfies

$$|u(x)| \leq M(1 + |x|)^\gamma, \quad \text{for some } \gamma < 2s \text{ and all } x \in \mathbb{R}^n, \quad (1.10)$$

$$\limsup_{|x| \rightarrow \infty} \frac{|u(x)|}{|x|^\gamma} > 0. \quad (1.11)$$

Then  $\gamma \leq \max\{0, \frac{m-2s}{m-1}\}$ .

Now we start with our second model. In [25], Filipucci, Pucci and Souplet consider the equation  $\Delta u + u^q |\nabla u|^m = 0$  in  $\mathbb{R}^n$  for  $m > 2, q > 0$ , and it is shown that any positive bounded classical solution to this equation is a constant. One of the main ingredients of their proof is a Bernstein type estimate on  $u^{\frac{q+m-1}{m-2}}$  which is used to establish that  $u^{q+1} |\nabla u|^{m-2}$  is bounded in  $\mathbb{R}^n$ . Interestingly, our proof of Theorem 1.1 can be modified suitably to get our next result which extends the Liouville property [25, Theorem 1.1] to the nonlocal setting.

**Theorem 1.4.** *Suppose that  $s > \frac{1}{2}$ . Then there exists no bounded positive viscosity solution to*

$$-\mathcal{I}u = u^q |\nabla u|^m \quad \text{in } \mathbb{R}^n,$$

for  $m > 2s, q > 0$ , unless  $u$  is a constant.

Our next result considers equations with coercive Hamiltonian, that is, model (III). In case of local operators, such problems are studied in [23, 26, 27]. More precisely, these works deal with a general family of nonlinear variational operators of  $p$ -Laplacian type. In particular, Filipucci, Pucci and Ragoli [26] show that the equation

$$\Delta_p u \geq u^q |\nabla u|^m \quad \text{in } \mathbb{R}^n, \quad \text{with } 0 \leq m < p - 1,$$

has only trivial  $C^1$  solutions if and only if  $q + 1 > p - m$ . Farina and Serrin [23] study Liouville property for similar equations without imposing any sign condition on the  $C^1$  solutions. In this article we prove the following Liouville property

**Theorem 1.5.** *Let  $s \geq 1/2$  and  $u$  be a non-negative viscosity solution to*

$$-\mathcal{I}u + u^q |\nabla u|^m = 0 \quad \text{in } \mathbb{R}^n, \quad (1.12)$$

for some  $q \geq 0, m \geq 0, q + m > 1$ , and

$$|u(x)| \leq M(1 + |x|)^\gamma$$

for some  $\gamma < 1$ . Then  $u$  is necessarily a constant. In particular, in view of Theorem 1.3, there exists no non-negative solution to (1.9) satisfying (1.10)-(1.11), other than constants.

Theorems 1.2, 1.3 and 1.5 partly answers the open problem mentioned in [19, Open problem 6.6].

The proofs of Filipucci, Pucci and Ragoli [26] rely on the construction of appropriate radial solution by the method of ode and then application of maximum principles whereas the arguments of Farina and Serrin [23] use variational techniques. None of this methods seems to be useful in proving Theorem 1.5.

**Remark 1.2.** *The conclusion of Theorem 1.5 also holds for a more general equation of the form*

$$-\mathcal{I}u + u^q |\nabla u|^m = u^{-\delta} \quad \text{in } \mathbb{R}^n,$$

for some  $\delta \geq -1$  and with the same conditions on  $q$  and  $m$ . Note that for  $\delta \in [-1, 0)$  and  $q = 0$ , this produces Liouville property not covered by [7].

The technique developed in this article can also deal with equations with singular nonlinearity. In fact, we could prove the following Liouville property

**Theorem 1.6.** *Let  $f : (0, \infty) \rightarrow \mathbb{R}$  be a continuous non-increasing function. Then there exists no non-constant positive viscosity solution to the equation*

$$-\mathcal{I}_{\text{sym}}u = f(u) \quad \text{in } \mathbb{R}^n, \quad (1.13)$$

provided

$$|u(x)| \leq C(1 + |x|)^\gamma \quad (1.14)$$

and one of the following holds

- (i)  $\gamma < 1 \wedge 2s$ ;
- (ii)  $s > 1/2$  and  $f$  is a  $C^1$  convex function satisfying

$$\limsup_{R \rightarrow \infty} R^{2s - (2-\beta)\gamma} |f'(R^\gamma)|^{\frac{2s-1}{2s}} = \infty, \quad (1.15)$$

for some  $\beta \in (0, 1)$ .

Here  $\mathcal{I}_{\text{sym}}$  is a translation invariant, positively 1-homogeneous operator which is elliptic with respect to the set of all symmetric kernels  $\mathcal{K}_{\text{sym}}$ .

The following corollary is immediate from the above theorem.

**Corollary 1.7.** *The equation  $(-\Delta)^s u = u^{-p}$ ,  $p > 0$ , does not have a solution satisfying (1.14) provided one of the following holds.*

- (i)  $2s \leq 1$  and  $\gamma < 2s$ ;
- (ii)  $2s > 1$  and either  $\gamma < 1$  or

$$2s > \gamma \left[ 1 + (p+1) \frac{2s}{2s-1} \right].$$

When  $\mathcal{I}_{\text{sym}}$  is replaced by the Laplacian operator, the proof follows by taking a logarithmic transformation on  $u$  and then applying the Bernstein estimate (see [41]). This method does not apply to the nonlocal setting as explained before in the introduction.

**Notations:**  $\kappa, \kappa_1, \dots, C, C_1, \dots$  are arbitrary constants whose value may change from line to line.  $B_r(x)$  will denote the ball of radius  $r$  around  $x$ . For  $x = 0$ , we simply denote this ball by  $B_r$  whereas the unit ball around 0 is denoted by  $B$ .

## 2. GENERAL STRATEGY AND TECHNICAL LEMMAS.

In this section we sketch the central idea for proving our results. We start with the notion of solution, for which we introduce a general equation, allowing the discussion in a general context.

Let  $\Omega \subseteq \mathbb{R}^n$ ,  $U \subseteq \mathbb{R}$  be open sets,  $H \in C(\mathbb{R}^n \times U \times \mathbb{R}^n)$  and  $\mathcal{I}$  be a nonlocal operator as mentioned in the introduction. We consider Hamilton-Jacobi equation of the form

$$-\mathcal{I}u + H(x, u, \nabla u) = 0 \quad \text{in } \Omega. \quad (2.1)$$

By a solution to this equation we would mean viscosity solution which we recall for convenience.

**Definition 2.1.** *We say  $u : \mathbb{R}^n \rightarrow U$  measurable,  $u \in L^1(\omega_s)$ , upper semicontinuous (lower semicontinuous) in  $\bar{\Omega}$ , is a viscosity subsolution (supersolution) to (2.1) in  $\Omega$  if, for each  $x_0 \in \Omega$  and  $\varphi \in C^2(B_r(x_0))$  for some  $r > 0$  satisfying  $u(x_0) = \varphi(x_0)$  and  $u \leq \varphi$  (resp.  $u \geq \varphi$ ) in  $B_r(x_0)$ , we have*

$$-\mathcal{I}v(x_0) + H(x_0, u(x_0), \nabla \varphi(x_0)) \leq (\text{resp. } \geq) f(x_0),$$

where

$$v(y) = \begin{cases} \varphi(y) & \text{for } y \in B_r(x_0), \\ u(y) & \text{elsewhere.} \end{cases} \quad (2.2)$$

We say that  $u$  is a viscosity solution to problem (2.1) in  $\Omega$  if it is both sub and supersolution in the above sense.

In the remaining part of the paper, we refer  $v$  in (2.2) as a test function for  $u$  with smooth representative  $\varphi$  at  $x_0$ .

Related to viscosity evaluations, we notice that in the case of subsolutions  $u$ , it is enough if  $x_0$  is a point of maximum of  $u - \varphi$  in  $B_r(x_0)$ . This follows from the observation that if we have two functions  $v_1, v_2$  such that  $v_2 = v_1 + c$  in  $B_r(x_0)$  for some  $c \in \mathbb{R}$ , then from (1.2), we have

$$\mathcal{I}v_1(x_0) = \mathcal{I}v_2(x_0).$$

This will allow us to use a relaxed version of viscosity subsolution. The same observation can be made for supersolutions, and hence for solutions.

**2.1. General Strategy.** Since the proofs involve technical calculations, for the ease of reading we first explain the common strategy to prove Theorems 1.1-1.5, and Theorem 1.6. These arguments are inspired from the celebrated Ishii-Lions method [31, Section VII] and have been adapted to the nonlocal setting following the ideas of Barles et. al. [12].

We consider a function  $\varphi \in C([0, +\infty))$ ,  $\varphi(0) = 0$ , so that  $\varphi$  is smooth, increasing and concave in some interval  $(0, a_0)$  for some  $a_0 > 0$ . This function  $\varphi$  could be understood as a modulus of continuity. For  $R > 0$ , we also require a function  $\tilde{\chi}_R : \mathbb{R}^n \rightarrow \mathbb{R}$ , smooth and nonnegative with  $\tilde{\chi}_R = 0$  in  $B_R$ , such that

$$\limsup_{|x| \rightarrow +\infty} \frac{|u(x)|}{\tilde{\chi}_R(x)} < 1. \quad (2.3)$$

This function plays the role of the localization. With these ingredients, take  $\delta \in (0, 1)$  and define the function  $\Phi : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  as

$$\Phi(x, y) = u(x) - u(y) - \delta\varphi(|x - y|) - \tilde{\chi}_R(x) - \tilde{\chi}_R(y), \quad x, y \in \mathbb{R}^n.$$

Our key point is to prove that for all  $\delta \in (0, 1)$  small enough, there exists  $R_0 = R_0(\delta) > 1$  large enough, such that for all  $R \geq R_0$  we have

$$\sup_{\mathbb{R}^n \times \mathbb{R}^n} \Phi(x, y) \leq 0. \quad (2.4)$$

Once this inequality is established, it can be easily seen that  $u$  is locally constant. In fact, for each compact set  $K \subset \mathbb{R}^n$  and each  $\delta$ , we can find  $R \geq R_0$  large enough so that  $K \subset B_R$ , implying that, for each  $x, y \in K$  we have

$$u(x) - u(y) - \delta\varphi(|x - y|) \leq \sup_{\mathbb{R}^n \times \mathbb{R}^n} \Phi(x, y) \leq 0.$$

In particular,

$$|u(x) - u(y)| \leq \delta\varphi(|x - y|) \quad \text{for all } x, y \in K,$$

and since  $\delta$  is arbitrary, we conclude that  $u$  is locally constant, and hence constant in  $\mathbb{R}^n$ .

The proof of (2.4) is carried out by contradiction, that is, assuming that:

$$\text{for each } \delta > 0 \text{ small and } R \text{ large enough, we have } \sup_{\mathbb{R}^n \times \mathbb{R}^n} \Phi(x, y) > 0. \quad (2.5)$$

By the choice of  $\chi_R$ , the suprema is attained at some point  $(\bar{x}, \bar{y}) \in \mathbb{R}^n \times \mathbb{R}^n$ , and since  $\varphi, \chi_R$  are nonnegative, we also have  $\bar{x} \neq \bar{y}$ . We denote

$$\phi(x, y) = \delta\varphi(|x - y|) + \tilde{\chi}_R(x) + \tilde{\chi}_R(y), \quad x, y \in \mathbb{R}^n.$$

Therefore,

$$\begin{cases} \bar{x} \text{ is a point of global maximum of } x \mapsto u(x) - (u(\bar{y}) + \phi(x, \bar{y})), \\ \bar{y} \text{ is a point of global minimum of } y \mapsto u(y) - (u(\bar{x}) - \phi(\bar{x}, y)), \end{cases} \quad (2.6)$$

and since  $\bar{x} \neq \bar{y}$ , we can test the equation at  $\bar{x}$  treating  $u$  as a subsolution, and at  $\bar{y}$  treating  $u$  as a supersolution. This leads us to the set of inequalities

$$-\mathcal{I}u(\bar{x}) + H(\bar{x}, u(\bar{x}), \nabla u(\bar{x})) \leq 0,$$

$$-\mathcal{I}u(\bar{y}) + H(\bar{y}, u(\bar{y}), \nabla u(\bar{y})) \geq 0,$$

understood in the viscosity sense. Subtracting these inequalities, we get that

$$I \leq I_H,$$

where

$$\begin{aligned} I &= -\mathcal{I}u(\bar{x}) + \mathcal{I}u(\bar{y}), \\ I_H &= -H(\bar{x}, u(\bar{x}), \nabla u(\bar{x})) + H(\bar{y}, u(\bar{y}), \nabla u(\bar{y})). \end{aligned}$$

We show that for suitable choice of  $\delta$  and  $R$  we must have  $I_H \ll I$  which contradicts the above inequality and therefore, (2.5) does not hold.

From this point, we need to consider the particularities of each problem to estimate both the terms  $I$  and  $I_H$ . Though already treated in the literature, the estimates for the nonlocal terms are less standard and because of this we provide next some technical lemmas that are going to be useful for our proofs.

**2.2. Technical Lemmas.** This section consists of three lemmas which are used repeatedly in our proofs. We need a few notations to state these lemmas. For  $K \in \mathcal{K}_s$ ,  $A \subset \mathbb{R}^n$  measurable,  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  measurable, and  $x, \xi \in \mathbb{R}^n$ , we denote

$$L_K[A](f, x, \xi) = \int_A (f(x+z) - f(x) - \mathbf{1}_B \xi \cdot z) K(z) dz,$$

whenever the integral makes sense. This is the case, for instance, when  $f \in C^2$  in a neighborhood of  $x$ ,  $\xi = \nabla f(x)$ , and  $f \in L^1(\omega_s)$ . In that case, we simply write

$$L_K[A]f(x) = L_K[A](f, x, \nabla f(x)).$$

For technical reasons, we also consider the notation

$$\tilde{L}_K[A](f, x, \xi) = \int_A (f(x+z) - f(x) - \xi \cdot z) K(z) dz, \quad (2.7)$$

that is, when the compensator term acts in the whole domain of integration. As before, when  $f$  is smooth we write

$$\tilde{L}_K[A]f(x) = \tilde{L}_K[A](f, x, \nabla f(x)).$$

The first one, Lemma 2.2, encodes the contribution coming from the ellipticity of the operator  $\mathcal{I}$ .

**Lemma 2.2.** *Let  $\bar{a} \in \mathbb{R}^n$ ,  $\bar{a} \neq 0$ , and for  $\eta_0, \delta_0 \in (0, 1)$ , consider the cone*

$$\mathcal{C} = \mathcal{C}_{\delta_0, \eta_0} := \{z \in B_{\delta_0|\bar{a}|} : |\bar{a} \cdot z| \geq (1 - \eta_0)|\bar{a}||z|\}. \quad (2.8)$$

*Assume  $\varphi \in C([0, +\infty))$  with  $\varphi \in C^2(0, |\bar{a}|(1 + \delta_0))$  increasing and concave, and denote  $\tilde{\varphi}(x) = \varphi(|x|)$  for  $x \in \mathbb{R}^n$ . Then, for each  $K$  we have*

$$\tilde{L}_K[\mathcal{C}](\tilde{\varphi}, \bar{a}) \leq \frac{1}{2} \int_{\mathcal{C}} \sup_{0 \leq t \leq 1} \left\{ (1 - \tilde{\eta}^2) \frac{\varphi'(|\bar{a} + tz|)}{(1 - \delta_0)|\bar{a}|} + \tilde{\eta}^2 \varphi''(|\bar{a} + tz|) \right\} |z|^2 K(z) dz,$$

where

$$\tilde{\eta} = \frac{1 - \eta_0 - \delta_0}{1 + \delta_0}.$$

*Proof.* Since  $z \mapsto \varphi(|\bar{a} + z|)$  is  $C^2$  in the cone  $\mathcal{C}$ , by Taylor's expansion we have

$$\begin{aligned} & \varphi(|\bar{a} + z|) - \varphi(|\bar{a}|) - z \cdot \varphi'(|\bar{a}|) \frac{\bar{a}}{|\bar{a}|} \\ & \leq \frac{1}{2} \sup_{0 \leq t \leq 1} \left\{ \frac{\varphi'(|\bar{a} + tz|)}{|\bar{a} + tz|} (|z|^2 - \langle \widehat{\bar{a} + tz}, z \rangle^2) + \varphi''(\bar{a} + tz) \langle \widehat{\bar{a} + tz}, z \rangle^2 \right\}, \end{aligned}$$

for all  $z \in B_{\delta_0|\bar{a}|}$ , where  $\hat{\zeta}$  denotes the unit vector along  $\zeta \neq 0$ . Thus, for each  $K \in \mathcal{K}_s$ , we obtain

$$\tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a}) \leq \frac{1}{2} \int_{\mathcal{C}} \sup_{0 \leq t \leq 1} \left\{ \frac{\varphi'(|\bar{a} + tz|)}{|\bar{a} + tz|} (|z|^2 - \langle \widehat{\bar{a} + tz}, z \rangle^2) + \varphi''(\bar{a} + tz) \langle \widehat{\bar{a} + tz}, z \rangle^2 \right\} K(z) dz. \quad (2.9)$$

For  $z \in \mathcal{C}_{\delta_0, \eta_0}$  and  $|t| \leq 1$  we have

$$|\langle \bar{a} + tz, z \rangle| \geq (1 - \eta_0 - \delta_0)|\bar{a}||z|,$$

and

$$(1 - \delta_0)|\bar{a}| \leq |\bar{a} + tz| \leq (1 + \delta_0)|\bar{a}|. \quad (2.10)$$

Using these inequalities in (2.9) we get the result.  $\square$

We require the following estimate for the nonlocal operator evaluated on the localization function.

**Lemma 2.3.** *Let  $\chi_0 : [0, +\infty) \rightarrow [0, 1]$  be a smooth, nondecreasing function satisfying  $\chi_0(t) = 0$  if  $t < 1/8$ ,  $\chi_0(t) = 1$  if  $t \geq 1/4$ . For  $\alpha \in [0, 2s)$  and  $R > 1$ , define*

$$\chi(x) = \chi_R(x) = \chi_0(|x|/R)(1 + |x|^2)^{\alpha/2}, \quad x \in \mathbb{R}^n. \quad (2.11)$$

Then, there exists a constant  $C$  such that for each  $A > 1$  and  $K \in \mathcal{K}_s$ , we have

$$|L_K \chi(x)| \leq \begin{cases} CA^\alpha R^{\alpha-1} & \text{for } s \in (1/2, 1), \\ CA^\alpha R^{\alpha-1} \log R & \text{for } s = 1/2, \\ CA^\alpha R^{\alpha-2s} & \text{for } s \in (0, 1/2), \end{cases} \quad \text{for all } |x| \leq AR,$$

and if  $K \in \mathcal{K}_{\text{sym}}$ , then we have

$$|L_K \chi(x)| \leq CA^\alpha R^{\alpha-2s}, \quad \text{for all } |x| \leq AR.$$

*Proof.* Denoting  $\psi(x) = (1 + |x|^2)^{\frac{\alpha}{2}}$ , we see that

$$\begin{aligned} \partial_{x_i} \chi(x) &= R^{-1} \chi'_0(|x|/R) \psi(x) \frac{x_i}{|x|} + \chi_0(|x|/R) \partial_{x_i} \psi(x), \\ \partial_{x_i x_j}^2 \chi(x) &= (R^{-2} \chi''_0(|x|/R) \frac{x_i x_j}{|x|^2} + R^{-1} \chi'_0(|x|/R) (\delta_{ij} |x|^{-1} - x_i x_j |x|^{-3})) \psi(x) \\ &\quad + R^{-1} \chi'_0(|x|/R) \left( \frac{x_i}{|x|} \partial_{x_j} \psi(x) + \frac{x_j}{|x|} \partial_{x_i} \psi(x) \right) + \chi_0(|x|/R) \partial_{x_i x_j}^2 \psi(x). \end{aligned}$$

Now we estimate  $L_K \chi(x)$  by dividing the domain in parts. Using that  $\chi_0(|x|/R) = 0$  for  $|x| \leq R/8$ ,  $\chi'_0, \chi''_0$  are uniformly bounded, and  $\alpha < 2s < 2$ , there exists  $\kappa > 0$ , not depending on  $R$  and  $A$ , such that

$$\|D^2 \chi\|_{L^\infty(\mathbb{R}^n)} \leq \kappa R^{\alpha-2},$$

giving us, for all  $K \in \mathcal{K}_s$ , that

$$|L_K[B]\chi(x)| \leq \kappa_1 \Lambda R^{\alpha-2},$$

for some  $\kappa_1 > 0$ , not dependent on  $R$  and  $A$ . Also, this estimate is uniform in  $x$ .

Now consider  $|x| \leq AR$ . There exists  $C > 0$  (not dependent on  $R$  and  $A$ ) such that

$$\|\nabla \chi\|_{L^\infty(B_R(x))} \leq CA^\alpha R^{\alpha-1},$$

which gives us

$$|L[B_R \setminus B]\chi(x)| \leq CA^\alpha R^{\alpha-1} \int_{B_R \setminus B} |z| K(z) dz \leq \begin{cases} C\Lambda A^\alpha R^{\alpha-1} & \text{for } s > 1/2, \\ C\Lambda A^\alpha R^{\alpha-1} \log R & \text{for } s = 1/2, \\ C\Lambda A^\alpha R^{\alpha-2s} & \text{for } s < 1/2. \end{cases}$$

for all  $K \in \mathcal{K}_s$ . Finally, there exists  $C > 0$ , not depending on  $A, R$ , such that

$$|L[B_R^c]\chi(x)| \leq C \int_{B_R^c} ((AR)^\alpha + |z|^\alpha) K(z) dz \leq C\Lambda A^\alpha R^{\alpha-2s}$$



Thus, gathering the above estimates, we conclude the result for  $K \in \mathcal{K}_s$ .

When  $K \in \mathcal{K}_{\text{sym}}$ , we can write

$$L_K \chi(x) = \tilde{L}_K[B_R] \chi(x) + L_K[B_R^c] \chi(x),$$

and using the estimate for  $D^2 \chi$ , we get

$$|\tilde{L}_K[B_R] \chi(x)| \leq CR^{\alpha-2} \int_{B_R} |z|^2 K(z) dz \leq CR^{\alpha-2s},$$

for some  $C > 0$ , not depending on  $R$  and  $A$ . Again, for the integral on  $B_R^c$ , we use the same estimate as in the previous case. This concludes the proof.  $\square$

**Lemma 2.4.** *Let  $u_1, u_2 \in C(\mathbb{R}^n) \cap L^1(\omega_s)$ ,  $\varphi$  as in Lemma 2.2, and  $\chi_1, \chi_2 \in C^2(\mathbb{R}^n) \cap L^1(\omega_s)$ . Assume there exists  $(\bar{x}, \bar{y}) \in \mathbb{R}^n$  with  $\bar{x} \neq \bar{y}$ , global maximum point of the function*

$$(x, y) \mapsto u_1(x) - u_2(y) - \phi(x, y), \quad x, y \in \mathbb{R}^n,$$

where  $\phi(x, y) = \varphi(|x - y|) - \chi_1(x) - \chi_2(y)$ .

Denote  $v_1$  the test function for  $u_1$  with smooth representative  $\phi(\cdot, \bar{y})$  at  $\bar{x}$ , and  $v_2$  the test function of  $u_2$  with smooth representative  $-\phi(\bar{x}, \cdot)$  at  $\bar{y}$ . Then, setting  $\bar{a} = \bar{x} - \bar{y}$  and considering  $\mathcal{C}$  as in (2.8) for some  $\delta_0, \eta_0 \in (0, 1)$ , the following inequality holds

$$\begin{aligned} -\mathcal{I}v_1(\bar{x}) + \mathcal{I}v_2(\bar{y}) &\geq -2 \sup_{K \in \mathcal{K}_s} \{\tilde{L}_K[\mathcal{C}] \tilde{\varphi}(\bar{a})\} - \sup_{K \in \mathcal{K}_s} \{L_K[\mathcal{C} \cup B_r^c] \chi_1(\bar{x})\} - \sup_{K \in \mathcal{K}_s} \{L_K[\mathcal{C} \cup B_r^c] \chi_2(\bar{y})\} \\ &\quad - \sup_{K \in \mathcal{K}_s} |L_K[\mathcal{C}^c \cap B_r] v_1(\bar{x})| - \sup_{K \in \mathcal{K}_s} |L_K[\mathcal{C}^c \cap B_r] v_2(\bar{y})|, \end{aligned}$$

where,  $\tilde{\varphi}(x) = \varphi(|x|)$  as in Lemma 2.2,  $\tilde{L}_K$  is given by (2.7), and  $B_r$  is the ball given by (2.2). In particular,

$$-\mathcal{I}v_1(\bar{x}) + \mathcal{I}v_2(\bar{y}) \geq -2 \sup_{K \in \mathcal{K}_s} \{\tilde{L}_K[\mathcal{C}] \tilde{\varphi}(\bar{a})\} - \sup_{K \in \mathcal{K}_s} \{L_K \chi_1(\bar{x})\} - \sup_{K \in \mathcal{K}_s} \{L_K \chi_2(\bar{y})\} + \Theta(r),$$

for some function  $\Theta(r)$ , dependent on  $\phi, \chi_1, \chi_2$ , satisfying  $\lim_{r \rightarrow 0} \Theta(r) = 0$ .

The same result holds for  $\mathcal{I}_{\text{sym}}$ , elliptic with respect to the family  $\mathcal{K}_{\text{sym}}$ , replacing  $\mathcal{K}_s$  by  $\mathcal{K}_{\text{sym}}$  in the above inequalities.

*Proof.* Denote  $\hat{a} = \frac{1}{|\bar{a}|} \bar{a}$  and  $\Phi(x, y) = u_1(x) - u_2(y) - \phi(x, y)$  for  $x, y \in \mathbb{R}^n$ .

Fix  $K \in \mathcal{K}_s$ . Using the fact that

$$\Phi(\bar{x} + z, \bar{y}) \leq \Phi(\bar{x}, \bar{y}) \Rightarrow u_1(\bar{x} + z) - u_2(\bar{x}) \leq \phi(|\bar{x} + z - \bar{y}|) - \phi(|\bar{x} - \bar{y}|),$$

we have

$$L_K[\mathcal{C}] v_1(\bar{x}) \leq L_K[\mathcal{C}] \phi(\cdot, \bar{y})(\bar{x}). \quad (2.12)$$

Similarly, the relation  $\Phi(\bar{x}, \bar{y} + z) \leq \Phi(\bar{x}, \bar{y})$  would lead to

$$L_K[\mathcal{C}] v_2(\bar{y}) \geq L_K[\mathcal{C}] (-\phi(\bar{x}, \cdot))(\bar{y}). \quad (2.13)$$

Since

$$\nabla_x \phi(\bar{x}, \bar{y}) = \varphi'(|\bar{a}|) \hat{a} + \nabla \chi_1(\bar{x}) \quad \text{and} \quad -\nabla_y \phi(\bar{x}, \bar{y}) = \varphi'(|\bar{a}|) \hat{a} - \nabla \chi_2(\bar{y}),$$

we have that

$$\begin{aligned} L_K[\mathcal{C}] \phi(\cdot, \bar{y})(\bar{x}) &= \int_{\mathcal{C}} [\varphi(|\bar{a} + z|) - \varphi(|\bar{a}|) - 1_B(z) \varphi'(|\bar{a}|) \hat{a} \cdot z] K(z) dz + L_K[\mathcal{C}] \chi_1(\bar{x}), \\ L_K[\mathcal{C}] (-\phi(\bar{x}, \cdot))(\bar{y}) &= - \int_{\mathcal{C}} [\varphi(|\bar{a} - z|) - \varphi(|\bar{a}|) + 1_B(z) \varphi'(|\bar{a}|) \hat{a} \cdot z] K(z) dz - L_K[\mathcal{C}] \chi_2(\bar{y}). \end{aligned}$$

Using (2.12) and (2.13), we conclude that

$$\begin{aligned} L_K[\mathcal{C}]v_1(\bar{x}) - L_K[\mathcal{C}]v_2(\bar{y}) &\leq \int_{\mathcal{C}} [\varphi(|\bar{a} + z|) - \varphi(|\bar{a}|) - 1_B(z)\varphi'(|\bar{a}|)\hat{a} \cdot z]K(z)dz \\ &\quad + \int_{\mathcal{C}} [\varphi(|\bar{a} - z|) - \varphi(|\bar{a}|) + 1_B(z)\varphi'(|\bar{a}|)\hat{a} \cdot z]K(z)dz \\ &\quad + L_K[\mathcal{C}]\chi_1(\bar{x}) + L_K[\mathcal{C}]\chi_2(\bar{y}). \end{aligned}$$

Writing  $K^*(z) = K(-z)$  for  $z \neq 0$ , we see that

$$L_K[\mathcal{C}]v_1(\bar{x}) - L_K[\mathcal{C}]v_2(\bar{y}) \leq \tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a}) + \tilde{L}_{K^*}[\mathcal{C}]\tilde{\varphi}(\bar{a}) + L_K[\mathcal{C}]\chi_1(\bar{x}) + L_K[\mathcal{C}]\chi_2(\bar{y}).$$

Notice that if  $K \in \mathcal{K}_s$ , then  $K^* \in \mathcal{K}_s$  too, and if  $K \in \mathcal{K}_{\text{sym}}$ , then  $K^* = K$ . Also, note that  $\Phi(\bar{x} + z, \bar{y} + z) \leq \Phi(\bar{x}, \bar{y})$  for all  $z$  implies

$$(u_1(\bar{x} + z) - u_1(\bar{x})) - (u_2(\bar{y} + z) - u_2(\bar{y})) \leq (\chi_1(\bar{x} + z) - \chi_1(\bar{x})) + (\chi_2(\bar{y} + z) - \chi_2(\bar{y})),$$

from which we readily have

$$L_K[\mathcal{C}^c]v_1(\bar{x}) - L_K[\mathcal{C}^c]v_2(\bar{y}) \leq L_K[\mathcal{C}^c \cap B_r^c]\chi_1(\bar{x}) + L_K[\mathcal{C}^c \cap B_r^c]\chi_2(\bar{y}) + |L_K[\mathcal{C}^c \cap B_r]v_1(\bar{x})| + |L_K[\mathcal{C}^c \cap B_r]v_2(\bar{y})|.$$

Gathering the above inequalities, and using the ellipticity of  $\mathcal{I}$  (see (1.3)), we see that

$$\begin{aligned} -\mathcal{I}v_1(\bar{x}) + \mathcal{I}v_2(\bar{y}) &\geq - \sup_{K \in \mathcal{K}_s} \{ \tilde{L}_K v_1(\bar{x}) - \tilde{L}_K v_2(\bar{y}) \} \\ &\geq -2 \sup_{K \in \mathcal{K}_s} \{ \tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a}) \} - \sup_{K \in \mathcal{K}_s} \{ L_K[\mathcal{C} \cup B_r^c]\chi_1(\bar{x}) \} - \sup_{K \in \mathcal{K}_s} \{ L_K[\mathcal{C} \cup B_r^c]\chi_2(\bar{y}) \} \\ &\quad - \sup_{K \in \mathcal{K}_s} |L_K[\mathcal{C}^c \cap B_r]v_1(\bar{x})| - \sup_{K \in \mathcal{K}_s} |L_K[\mathcal{C}^c \cap B_r]v_2(\bar{y})|, \end{aligned}$$

which concludes the proof for  $\mathcal{K}_s$ . The proof for  $\mathcal{K}_{\text{sym}}$  is a straightforward adaptation.  $\square$

### 3. PROOFS OF MAIN RESULTS

In this section we prove our main results. As mentioned before, the central idea of our proofs relies on the Ishii-Lions method [31, Section VII] which was introduced to prove Hölder regularity of classical viscosity solutions. We begin with the proof of Theorem 1.1 and most of the other proofs (other than Theorems 1.5 and 1.3) will be variation of this particular one.

*Proof of Theorem 1.1.* To simplify some computations, up to modifying  $M$ , we will use (1.6) as

$$|u(x)| \leq M|x|^\gamma \quad \text{for } |x| > 1 \quad \text{and} \quad \max_{|x| \leq 1} |u(x)| \leq M.$$

We consider the different cases separately.

**1. Case  $s \in (1/2, 1)$ :** We divide the proof in four steps.

*Step 1: Setting the General Strategy.* For  $\theta \in (0, 1)$  to be fixed later and  $R > 4$ , we define  $\varphi = \varphi_R : [0, +\infty) \rightarrow [0, +\infty)$  as

$$\varphi(t) = t - \frac{1}{4R^\theta}t^{1+\theta}, \quad t \in [0, R], \tag{3.1}$$

and extended as a constant function equal to  $\varphi(R)$  on  $[R, +\infty)$ . Take  $\alpha \in (\gamma, 2s)$  and consider the function  $\chi = \chi_R$  as in (2.11) above. Let

$$\tilde{\chi}_R(x) = M2^{1+2\alpha}R^{\gamma-\alpha}\chi(x), \quad x \in \mathbb{R}^n.$$

Since  $\tilde{\chi}_R(x) \geq M2^{1+2\alpha}R^{\gamma-\alpha}|x|^\alpha$  for  $|x| > R/4$  and  $\alpha > \gamma$ , (2.3) holds.

Now consider the function

$$\Phi(x, y) := u(x) - u(y) - \delta\varphi(|x - y|) - (\tilde{\chi}_R(x) + \tilde{\chi}_R(y)), \quad x, y \in \mathbb{R}^n,$$

and as mention in the Section 2.1, our aim is to prove that (2.4) holds. By contradiction, we assume (2.5) holds, namely, there exists  $\delta > 0$  and a sequence  $R \rightarrow +\infty$  such that

$$\sup_{x,y \in \mathbb{R}^n} \Phi(x, y) > 0.$$

By our choice of localization function  $\tilde{\chi}$ , the above supremum is attained at some point  $(\bar{x}, \bar{y}) \in \mathbb{R}^n \times \mathbb{R}^n$ , and also  $\bar{x} \neq \bar{y}$ . From now on, we denote

$$\bar{a} = \bar{x} - \bar{y}.$$

Notice that for  $|x| > \frac{R}{4}$  we can write

$$\begin{aligned} |u(x)| - M2^{1+2\alpha}R^{\gamma-\alpha}\chi(x) &\leq M|x|^\gamma - M2^{1+2\alpha}R^{\gamma-\alpha}|x|^\alpha \\ &\leq |x|^\gamma M(1 - 2^{1+2\alpha}R^{\gamma-\alpha}|x|^{\alpha-\gamma}) \\ &\leq |x|^\gamma M(1 - 2^{1+2\gamma}) \leq -(R/4)^\gamma M, \end{aligned}$$

Thus, for  $|x| > R/4$  and  $y \in \mathbb{R}^n$ , we have

$$\Phi(x, y) \leq -(R/4)^\gamma M - u(y) - \tilde{\chi}_R(y) \leq 0.$$

A similar analysis can be performed for  $|y| > R/4$ , giving us  $\Phi(x, y) \leq 0$  for  $|y| > R/4$ ,  $x \in \mathbb{R}^n$ . Therefore, we must have

$$|\bar{x}|, |\bar{y}| \leq R/4.$$

Denoting

$$\phi(x, y) = \delta\varphi(|x - y|) + (\tilde{\chi}_R(x) + \tilde{\chi}_R(y)), \quad (x, y) \in \mathbb{R}^n \times \mathbb{R}^n,$$

we have  $x \mapsto \phi(x, \bar{y})$  is a test function for  $u$  at  $\bar{x}$ , and  $y \mapsto -\phi(\bar{x}, y)$  is a test function for  $u$  at  $\bar{y}$  (c.f. (2.6)). Then, denoting  $v_1$  the test function of  $u$  with smooth representative  $\phi(\cdot, \bar{y})$  at  $\bar{x}$ , and  $v_2$  the test function of  $u$  with smooth representative  $-\phi(\bar{x}, \cdot)$  at  $\bar{y}$ , (c.f. (2.2)), we have

$$\begin{aligned} -\mathcal{I}v_1(\bar{x}) + H(\nabla_x \phi(\bar{x}, \bar{y})) &\leq 0, \\ -\mathcal{I}v_2(\bar{y}) + H(-\nabla_y \phi(\bar{x}, \bar{y})) &\geq 0. \end{aligned}$$

Then, subtracting both the inequalities, we arrive at

$$I \leq I_H, \tag{3.2}$$

where

$$\begin{aligned} I &:= -\mathcal{I}v_1(\bar{x}) + \mathcal{I}v_2(\bar{y}), \\ I_H &:= -H(\nabla_x \phi(\bar{x}, \bar{y})) + H(-\nabla_y \phi(\bar{x}, \bar{y})). \end{aligned}$$

Now we estimate  $I$  and  $I_H$  separately.

*Step 2: Estimate of  $I_H$ .* A simple computation shows that for  $t \leq R$  we have

$$\frac{1}{2} \leq (1 - \frac{1+\theta}{4R^\theta}t^\theta) = \varphi'(t) \leq 1, \quad \varphi''(t) = -\frac{(1+\theta)\theta}{4R^\theta}t^{\theta-1}. \tag{3.3}$$

Furthermore, there exists a constant  $\kappa > 0$ , independent of  $R$ , such that

$$|\nabla\chi(\bar{y})|, |\nabla\chi(\bar{x})| \leq \kappa(1 + R^{\alpha-1}),$$

giving us

$$|\nabla\tilde{\chi}_R(\bar{x})|, |\nabla\tilde{\chi}_R(\bar{y})| \leq \kappa M(R^{\gamma-1} + R^{\gamma-\alpha}),$$

for some constant  $\kappa > 0$ .

Since  $\gamma < 1 \wedge \alpha$ , the l.h.s. of the above display tends to 0 as  $R \rightarrow \infty$ , and therefore, combining with (3.3) we see that

$$\frac{\delta}{8} \leq |\nabla_x \phi(\bar{x}, \bar{y})|, |\nabla_y \phi(\bar{x}, \bar{y})| \leq 1 + 2\kappa M,$$

for all large  $R$ . Thus we can apply (1.4) with the local Lipschitz constant  $C_\delta$ , depending on  $\delta$ , from which we arrive at

$$I_H \leq C_\delta(R^{\gamma-1} + R^{\gamma-\alpha}), \quad (3.4)$$

for all large  $R$ .

*Step 3: Estimate of  $I$ .* For convenience, we recall the cone

$$\mathcal{C} = \mathcal{C}_{\delta_0, \eta_0} := \{z \in B_{\delta_0|\bar{a}|} : |\bar{a} \cdot z| \geq (1 - \eta_0)|\bar{a}||z|\} \subset B_{\frac{R}{2}}$$

for  $\delta_0, \eta_0 \in (0, 1)$  to be fixed, Using Lemma 2.4, we have

$$\begin{aligned} I &\geq -2\delta \sup_{K \in \mathcal{K}_s} \{\tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a})\} - \sup_{K \in \mathcal{K}_s} \{L_K\tilde{\chi}_R(\bar{x})\} - \sup_{K \in \mathcal{K}_s} \{L_K\tilde{\chi}_R(\bar{y})\} + \Theta(r) \\ &=: -2\delta I_1 - I_2 - I_3 + \Theta(r), \end{aligned} \quad (3.5)$$

where  $\Theta(r) \rightarrow 0$  as  $r \rightarrow 0$ , for each fixed  $\delta$  and  $R$ . Now it remains to estimate  $I_1, I_2$  and  $I_3$  above. Using Lemma 2.3, obtain at once that

$$I_2, I_3 \leq CMR^{\gamma-1}. \quad (3.6)$$

Thus we deal with  $I_1$ . For this, we start by noticing that

$$\frac{\delta}{2}|\bar{a}| \leq \delta\varphi(|\bar{a}|) \leq u(\bar{x}) - u(\bar{y}) \leq 2M(R^\gamma + 1),$$

from which, we also get that

$$|\bar{a}| \leq \frac{CM}{\delta}R^\gamma, \quad (3.7)$$

for some constant  $C$ .

Now, we choose

$$\eta_0 = \tau \frac{|\bar{a}|^\theta}{R^\theta}, \quad \delta_0 = \tau \frac{|\bar{a}|^\theta}{R^\theta} < 1, \quad (3.8)$$

for some  $\tau \in (0, 1)$  small enough so that

$$\frac{1 - \tilde{\eta}^2}{1 - \delta_0} < 4(\eta_0 + 2\delta_0) \quad \text{and} \quad \tilde{\eta}^2 \geq \frac{1}{2}.$$

Then, using (2.10), (3.3) together with Lemma 2.2, we have

$$\tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a}) \leq \frac{1}{2} \frac{|\bar{a}|^{\theta-1}}{R^\theta} \sup_{K \in \mathcal{K}} \int_{\mathcal{C}} [12\tau - \frac{\theta(\theta+1)}{8}(1 + \delta_0)^{\theta-1}] |z|^2 K(z) dz.$$

We can choose  $\tau$  depending on  $\theta$  to conclude the existence of  $c_\theta \in (0, 1)$  such that

$$\tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a}) \leq -c_\theta \lambda \frac{|\bar{a}|^{\theta-1}}{R^\theta} \int_{\mathcal{C}} |z|^2 \frac{dz}{|z|^{n+2s}}.$$

To estimate the integral on the cone we use [12, Example 1] to obtain

$$\tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a}) \leq -\tilde{c}_\theta \lambda \frac{|\bar{a}|^{\theta-1}}{R^\theta} \eta_0^{\frac{n-1}{2}} (|\bar{a}|\delta_0)^{2-2s},$$

and from definition of  $\delta_0, \eta_0$  this leads to

$$\begin{aligned} \tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a}) &\leq -\tilde{c}_\theta \tau^{\frac{n-1}{2}+2-2s} \lambda \frac{|a|^{\theta-1}}{R^\theta} (|\bar{a}|/R)^{\theta \frac{d-1}{2}} (|\bar{a}|^{1+\theta} R^{-\theta})^{2-2s} \\ &= -\tilde{c}_\theta \tau^{\frac{n-1}{2}+2-2s} \lambda R^{-\theta((n+1)/2+2-2s)} |\bar{a}|^{\theta((n+1)/2+2-2s)+1-2s} \\ &= -\tilde{c}_\theta \tau^{\frac{n-1}{2}+2-2s} \lambda R^{-\theta\beta} |\bar{a}|^{\theta\beta+1-2s}, \end{aligned}$$

where  $\beta = (n+1)/2 + 2 - 2s$ . Since  $2s > 1$ , we now set  $\theta \in (0, \frac{2s-1}{\beta})$  so that the exponent of  $|\bar{a}|$  becomes negative. By definition of  $I_1$  and (3.7) we then obtain

$$2\delta I_1 = 2\delta \sup_K \{ \tilde{L}_K[C] \tilde{\varphi}(\bar{a}) \} \leq -\kappa_3 \lambda \delta^{2s-\theta\beta} R^{\theta\beta(\gamma-1)+\gamma(1-2s)}, \quad (3.9)$$

for some constant  $\kappa_3$  not dependent on  $\delta$  and  $R$ . Combining this estimate together with (3.6) and replacing them into (3.5), we conclude that

$$I \geq \kappa_4 \delta^{2s-\theta\beta} R^{\theta\beta(\gamma-1)+\gamma(1-2s)} - CM R^{\gamma-1} + \Theta(r). \quad (3.10)$$

for some constant  $\kappa_4$ , not dependent on  $R$  and  $\delta$ .

*Step 4: Conclusion.* Placing this last estimate for  $I$  and (3.4) into (3.2) and letting  $r \rightarrow 0$ , we obtain

$$\kappa_4 \delta^{2s-\theta\beta} R^{\theta\beta(\gamma-1)+\gamma(1-2s)} \leq C(R^{\gamma-\alpha} + R^{\gamma-1}) \quad (3.11)$$

for some constant  $\kappa_4, C$ , not dependent on  $R$ , and for all large  $R$ .

Now we can set  $\alpha > 1$  so that  $\alpha - \gamma > 1 - \gamma$ , and  $\theta \in (0, \frac{2s-1}{\beta})$  small enough to satisfy

$$\theta\beta(\gamma-1) + \gamma(1-2s) > \gamma-1.$$

This can be done because  $2s\gamma < 1$ . Then, taking  $R \rightarrow +\infty$  we arrive at a contradiction to (3.11), confirming (2.4) holds. As explained in our General Strategy (see Section 2.1), this implies  $u$  is locally constant, and therefore the result.

**2. Case  $s \in (0, 1/2]$ :** We start with (i). Suppose that  $u$  is not constant and for some  $\eta > 0$  there exist points  $\tilde{x}, \tilde{y}$  such that

$$u(\tilde{x}) - u(\tilde{y}) > \eta.$$

As before, for  $\delta \in (0, 1)$  and  $R > 4$  we let  $\tilde{\chi}_R(x) = M2^{1+2\alpha} R^{\gamma-\alpha} \chi(x)$  with  $\chi$  as in (2.11) with  $\alpha \in (\gamma, 2s)$ . We also denote  $\varphi(t) = t^\beta$  for  $\beta \in (\gamma, 2s)$  fixed. We consider the function

$$\Phi(x, y) = u(x) - u(y) - \delta\varphi(|x-y|) - (\tilde{\chi}_R(x) + \tilde{\chi}_R(y)), \quad x, y \in \mathbb{R}^n. \quad (3.12)$$

Let us fix  $R$  large enough so that  $|\tilde{x}| + |\tilde{y}| < R/8$ . Set  $\hat{\delta} = \hat{\delta}(\eta, \beta)$  small enough so that

$$u(\tilde{x}) - u(\tilde{y}) - \delta\varphi(|\tilde{x} - \tilde{y}|) > \eta/2 \quad \text{for all } \delta \in (0, \hat{\delta}).$$

Using the definition of  $\chi_0$  we note that

$$\sup_{\mathbb{R}^n \times \mathbb{R}^n} \Phi \geq \sup_{B_R \times B_R} \Phi \geq u(\tilde{x}) - u(\tilde{y}) - \delta|\tilde{x} - \tilde{y}|^\beta > \eta/2,$$

and since (2.3) holds, the supremum is attained at some  $(\bar{x}, \bar{y}) \in \mathbb{R}^n \times \mathbb{R}^n$ . Denote  $\bar{a} = \bar{x} - \bar{y}$ . Let  $\omega$  be the modulus of continuity of  $u$  and since  $u$  is not a constant, we necessarily have  $\omega(t) > 0$  for  $t > 0$ . By maximality, we see that

$$\eta/2 \leq \Phi(\bar{x}, \bar{y}) \leq u(\bar{x}) - u(\bar{y}) \leq \omega(|\bar{a}|) \Rightarrow |\bar{a}| \geq \omega^{-1}(\eta/2).$$

In particular,  $|\bar{a}|$  is bounded from below, independent of  $R$  and  $\delta$ . Hence  $\varphi'(|\bar{a}|) \leq \beta|\omega^{-1}(\eta)|^{\beta-1}$ .

Now, letting  $\phi(x, y) = \delta\varphi(|x-y|) + \tilde{\chi}_R(x) + \tilde{\chi}_R(y)$ , and denoting  $v_1$  the test function for  $u$  with smooth representative  $\phi(\cdot, \bar{y})$  at  $\bar{x}$ , and  $v_2$  the test function for  $u$  with smooth representative  $-\phi(\bar{x}, \cdot)$  at  $\bar{y}$ , we use the viscosity inequalities and arrive at (3.2), the notation being the same as before, just the assumptions need to be specified.

First, by (1.7), similarly as in (3.4), we get

$$I_H \leq C_\eta R^{\gamma-\alpha},$$

for some constant  $C_\eta > 0$  that depends on  $\eta$  but not on  $\delta$  nor  $R$ .

Using Lemmas 2.3 and 2.4, we obtain

$$I \geq \begin{cases} -2\delta \sup_{K \in \mathcal{K}_s} \{ \tilde{L}_K \tilde{\varphi}(\bar{a}) \} + \Theta(r) - CR^{\gamma-2s} & \text{for } s \in (0, 1/2), \\ -2\delta \sup_{K \in \mathcal{K}_s} \{ \tilde{L}_K \tilde{\varphi}(\bar{a}) \} + \Theta(r) - CR^{\gamma-1} \log R & \text{for } s = 1/2, \end{cases}$$

for some  $C > 0$ , not dependent on  $\eta, \delta$  and  $R$ , and  $\Theta(r) \rightarrow 0$  as  $r \rightarrow 0$ .

Now, using Lemma 2.2 with  $\varphi(t) = t^\beta$  and  $\delta_0 = \eta_0 = \tau$  suitably small depending on  $\beta$  (see (3.9)), we obtain

$$\sup_{K \in \mathcal{K}_s} \{\tilde{L}_K \tilde{\varphi}(\bar{a})\} \leq -\kappa \lambda |\bar{a}|^{\beta-2s} \leq -\kappa_1 \delta^{\frac{2s-\beta}{\beta}} R^{\frac{\gamma}{\beta}(\beta-2s)},$$

where the last inequality follows from the fact  $\delta\varphi(|\bar{a}|) \leq 2M(R^\gamma + 1)$  (see (3.7)) and  $\kappa, \kappa_1$  are suitable constants. Combining the above estimates, letting  $r \rightarrow 0$ , we arrive at

$$\kappa_2 \delta^{\frac{2s}{\beta}} R^{\frac{\gamma}{\beta}(\beta-2s)} \leq \kappa_3 R^{\gamma-\alpha},$$

for some constants  $\kappa_2, \kappa_3$ , independent of  $R$  and  $\delta$ , for all large  $R$ . Fixing  $\beta \in (\gamma, 2s)$  we can choose  $\alpha \in (\gamma, 2s)$  to satisfy  $2s\gamma < \alpha\beta \Leftrightarrow \gamma(2s - \beta) < \beta(\alpha - \gamma)$ . Then the above inequality can not hold for all large  $R$ , leading to a contradiction. Hence  $u$  must be a constant.

Now we consider (ii). We again work with the same coupling function  $\Phi$  given by (3.12), and show that  $\Phi \leq 0$  in  $\mathbb{R}^n \times \mathbb{R}^n$  for all large  $R$ . Suppose, to the contrary, that for some  $\delta \in (0, 1)$  and large  $R$  we have  $\sup \Phi > 0$ . Then we proceed as before to arrive at (3.2). Lower bound for  $I$  can be computed from (i) above. The only difference is the estimate of  $I_H$ , where we use the (uniform) Lipschitz property of  $H$ . In particular,

$$\begin{aligned} I_H &= -H(\nabla_x \phi(\bar{x}, \bar{y})) + H(-\nabla_y \phi(\bar{x}, \bar{y})) \\ &\leq H_{\text{Lip}}(|\nabla \tilde{\chi}_R(\bar{x})| + |\nabla \tilde{\chi}_R(\bar{y})|) \\ &\leq CR^{\gamma-\alpha}, \end{aligned}$$

for some constant  $C$ , where  $H_{\text{Lip}}$  denotes the Lipschitz constant of  $H$ . Now the remaining part of the proof can be completed as in (i).  $\square$

*Proof of Theorem 1.2.* In view of Theorem 1.1 we only need to consider the case  $\frac{1}{2s} \leq \gamma < \frac{m-2s}{m-1}$ . In particular, we have

$$m > 2s + 1$$

in this case.

Let  $\varphi$  as in (3.1) and  $\chi$  as in (2.11) with  $\alpha \in (\gamma, 2s)$ . Denote  $\tilde{\chi}_R(x) = M2^{1+2\alpha}R^{\gamma-\alpha}\chi(x)$ , and consider

$$\Phi(x, y) = u(x) - u(y) - \delta\varphi(|x - y|) - (\tilde{\chi}_R(x) + \tilde{\chi}_R(y)) \quad x, y \in \mathbb{R}^n.$$

We pick  $\epsilon_0 \in (0, 1)$ , to be chosen later, and set

$$R = R(\delta) = \delta^{-\frac{1}{(1-\gamma)\epsilon_0}} \quad \text{for } \delta > 0, \quad (3.13)$$

The result follows by proving the existence of  $\delta_0 > 0$  such that

$$\Phi_\delta := \sup_{\mathbb{R}^n \times \mathbb{R}^n} \Phi(x, y) \leq 0 \quad \text{for all } \delta \leq \delta_0.$$

Suppose, to the contrary, that for some sequence  $\delta \rightarrow 0^+$  we have  $\Phi_\delta > 0$ . Since (2.3) holds for the choice of  $\tilde{\chi}_R$ , the supremum of  $\Phi_\delta$  is finite and attained at some point  $(\bar{x}, \bar{y}) \in \mathbb{R}^n \times \mathbb{R}^n$ , that is  $\Phi(\bar{x}, \bar{y}) = \Phi_\delta > 0$ . As before, we have  $\bar{x} \neq \bar{y}$ , and  $|\bar{x}|, |\bar{y}| \leq R/4$ . From now on, we denote

$$\bar{a} = \bar{x} - \bar{y},$$

and  $\phi(x, y) = \delta\varphi(|x - y|) + \tilde{\chi}_R(x) + \tilde{\chi}_R(y)$ .

Then, denoting  $v_1$  the test function of  $u$  with smooth representative  $\phi(\cdot, \bar{y})$  at  $\bar{x}$ , and  $v_2$  the test function of  $u$  with smooth representative  $-\phi(\bar{x}, \cdot)$  at  $\bar{y}$ , (c.f (2.2)), we can write

$$-\mathcal{I}v_1(\bar{x}) + H(\nabla_x \phi(\bar{x}, \bar{y})) \leq 0; \quad -\mathcal{I}v_2(\bar{y}) + H(-\nabla_y \phi(\bar{x}, \bar{y})) \geq 0.$$

We subtract these inequalities to arrive at

$$I \leq I_H,$$

with

$$I = -\mathcal{I}v_1(\bar{x}) + \mathcal{I}v_2(\bar{x}); \quad I_H = H(-\nabla_y \phi(\bar{x}, \bar{y})) - H(\nabla_x \phi(\bar{x}, \bar{y})).$$

As before, we estimate each term. For  $I_H$ , we see that

$$|\nabla \tilde{\chi}_R(\bar{y})|, |\nabla \tilde{\chi}_R(\bar{x})| \leq \kappa MR^{\gamma-1}$$

for some  $\kappa > 0$ , due to the fact  $\chi_0(|x|/R) = 0$  for  $|x| \leq R/8$ . Hence, from (1.8), we get

$$\begin{aligned} I_H &\leq C_H \left( |\nabla_x \phi(\bar{x}, \bar{y})|^{m-1} + |\nabla_y \phi(\bar{x}, \bar{y})|^{m-1} \right) |\nabla \tilde{\chi}_R(\bar{x}) + \nabla \tilde{\chi}_R(\bar{y})| \\ &\leq \kappa C_H MR^{\gamma-1} (\delta^{m-1} + R^{(\gamma-1)(m-1)}), \end{aligned}$$

from which, by the choice of  $R(\delta)$  in (3.13) we arrive at

$$I_H \leq \kappa C_H M \delta^{m-1 + \frac{1}{\epsilon_0}},$$

for some  $\kappa > 0$ , just depending on  $n$ .

For  $I$ , using Lemma 2.4, we have

$$\begin{aligned} I &\geq -2\delta \sup_{K \in \mathcal{K}_{\text{sym}}} \{ \tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a}) \} - \sup_{K \in \mathcal{K}_{\text{sym}}} \{ L_K \tilde{\chi}_R(\bar{x}) \} - \sup_{K \in \mathcal{K}_{\text{sym}}} \{ L_K \tilde{\chi}_R(\bar{y}) \} + \Theta(r) \\ &=: -2\delta I_1 - I_2 - I_3 + \Theta(r), \end{aligned}$$

for some  $\Theta(r) \rightarrow 0$  as  $r \rightarrow 0$ . Using Lemma 2.2, with the same choice for  $\delta_0, \eta_0$  in (3.8), we arrive at the same estimate (3.9), namely,

$$-2\delta I_1 \geq \kappa_1 \delta^{2s - \theta\beta} R^{\theta\beta(\gamma-1) + \gamma(1-2s)} = \kappa_1 \delta^{2s + \frac{\gamma(2s-1)}{(1-\gamma)\epsilon_0} + \theta\beta(\frac{1}{\epsilon_0} - 1)},$$

where  $\beta = (n+1)/2 + 2 - 2s > 0$  and  $\kappa_1$  depends on  $n, s, \theta$  but not on  $\delta$  and  $R$ .

On the other hand, since we are considering symmetric kernels, by Lemma 2.3 and the definition of  $\tilde{\chi}_R$  we have

$$I_2, I_3 \leq CR^{\gamma-2s} = C\delta^{\frac{2s-\gamma}{(1-\gamma)\epsilon_0}}.$$

This gives a lower bound for  $I$ . Gathering these estimates and the upper bound of  $I_H$ , and replacing them into the inequality  $I \leq I_H$ , we arrive at

$$\kappa_1 \delta^{2s + \frac{\gamma(2s-1)}{(1-\gamma)\epsilon_0} + \theta\beta(\frac{1}{\epsilon_0} - 1)} \leq \bar{\kappa} \left( \delta^{m-1 + \frac{1}{\epsilon_0}} + \delta^{\frac{2s-\gamma}{(1-\gamma)\epsilon_0}} \right), \quad (3.14)$$

after letting  $r \rightarrow 0$ . Since  $\epsilon_0 < 1$ , we have

$$2s + \frac{\gamma(2s-1)}{(1-\gamma)\epsilon_0} < \frac{2s-\gamma}{(1-\gamma)\epsilon_0}.$$

On the other hand, since  $\gamma \mapsto \gamma(1-\gamma)^{-1}$  is strictly increasing in  $(0, 1)$  and  $\gamma < \frac{m-2s}{m-1}$ , we have

$$1 - \frac{\gamma(2s-1)}{1-\gamma} > 2s + 1 - m.$$

Thus, taking  $\epsilon_0 \in (0, 1)$  close enough to 1, we get that

$$2s - m + 1 < \frac{1}{\epsilon_0} - \frac{\gamma(2s-1)}{(1-\gamma)\epsilon_0}.$$

Summarizing, if we let

$$\tau_1 = 2s + \frac{\gamma(2s-1)}{(1-\gamma)\epsilon_0}; \quad \tau_2 = m - 1 + \frac{1}{\epsilon_0}; \quad \tau_3 = \frac{2s-\gamma}{(1-\gamma)\epsilon_0},$$

from the choice of  $\epsilon_0$  we have  $0 < \tau_1 < \tau_2$  and  $\tau_1 < \tau_3$ , and inequality (3.14) reads

$$\kappa_1 \delta^{\tau_1 + \theta\beta(1 - \frac{1}{\epsilon_0})} \leq \bar{\kappa}(\delta^{\tau_2} + \delta^{\tau_3}).$$

Now fix  $\epsilon_0$  and take  $\theta$  small enough so that  $\tau_1 + \theta\beta(1 - \epsilon_0^{-1}) < \tau_2, \tau_3$  (this would modify  $\kappa_1$  but this is not going to play a role), and then letting  $\delta \rightarrow 0^+$  we arrive at a contradiction. As before, this implies that  $u$  is constant.  $\square$

Now we come to the proof of Theorem 1.4.

*Proof of Theorem 1.4.* In this case, we denote  $M = \|u\|_\infty$ . Recalling  $\varphi$  as in (3.1), we consider  $\chi$  as in (2.11) with  $\alpha = 0$ , denote  $\tilde{\chi}_R(x) = 2M\chi(x)$ , and the function

$$\Phi(x, y) = u(x) - u(y) - \delta\varphi(|x - y|) - (\tilde{\chi}_R(x) + \tilde{\chi}_R(y)), \quad x, y \in \mathbb{R}^n.$$

We also write

$$\phi(x, y) = \delta\varphi(|x - y|) + (\tilde{\chi}_R(x) + \tilde{\chi}_R(y)), \quad x, y \in \mathbb{R}^n.$$

We fix  $\epsilon_0 \in (0, (2s)^{-1})$  and let

$$R = R(\delta) = \delta^{-1/\epsilon_0} > \delta^{-1}.$$

Following our general strategy in Section 2.1, we claim that there exists  $\delta_0 > 0$  such that

$$\Phi_\delta := \sup_{\mathbb{R}^n \times \mathbb{R}^n} \Phi(x, y) \leq 0 \quad \text{for all } \delta \leq \delta_0.$$

Arguing by contradiction, we assume  $\Phi_\delta > 0$ . Since (2.3) holds, there exist  $\bar{x}, \bar{y} \in \mathbb{R}^n, \bar{x} \neq \bar{y}$ , such that  $\Phi(\bar{x}, \bar{y}) = \Phi_\delta$ . These points are such that  $|\bar{x}|, |\bar{y}| \leq R/4$ .

Denoting  $v_1$  the test function for  $u$  with smooth representative  $\phi(\cdot, \bar{y})$  at  $\bar{x}$ , and  $v_2$  the test function for  $u$  with smooth representative  $-\phi(\bar{x}, \cdot)$  at  $\bar{y}$ , (c.f. (2.2)), we can write

$$-\mathcal{I}v_1(\bar{x}) \leq u(\bar{x})^q |\nabla_x \phi(\bar{x}, \bar{y})|^m; \quad -\mathcal{I}v_2(\bar{y}) \geq u(\bar{y})^q |-\nabla_y \phi(\bar{x}, \bar{y})|^m.$$

Subtract both the inequalities to arrive at

$$I \leq I_H, \tag{3.15}$$

with

$$I = -\mathcal{I}v_1(\bar{x}) + \mathcal{I}v_2(\bar{x}); \quad I_H = u(\bar{x})^q |\nabla_x \phi(\bar{x}, \bar{y})|^m - u(\bar{y})^q |\nabla_y \phi(\bar{x}, \bar{y})|^m.$$

To compute  $I_H$ , we first note that

$$|\nabla \tilde{\chi}_R(\bar{x})|, |\nabla \tilde{\chi}_R(\bar{y})| \leq CMR^{-1},$$

for some  $C$ . Thus, denoting  $\bar{a} = \bar{x} - \bar{y}$  and  $\hat{a} = \bar{a}/|\bar{a}|$ , using (3.3), we obtain

$$\begin{aligned} I_H &= u(\bar{x})^q |\delta\varphi'(|\bar{a}|)\hat{a} + \nabla \tilde{\chi}_R(\bar{x})|^m - u(\bar{y})^q |\delta\varphi'(|\bar{a}|)\hat{a} - \nabla \tilde{\chi}_R(\bar{y})|^m \\ &= [u(\bar{x})^q - u(\bar{y})^q] |\delta\varphi'(|\bar{a}|)\hat{a} + \nabla \tilde{\chi}_R(\bar{x})|^m \\ &\quad + u(\bar{y})^q (|\delta\varphi'(|\bar{a}|)\hat{a} + \nabla \tilde{\chi}_R(\bar{x})|^m - |\delta\varphi'(|\bar{a}|)\hat{a} - \nabla \tilde{\chi}_R(\bar{y})|^m) \\ &\leq \kappa_2 [\|u\|_\infty^q (\delta^m + R^{-m} + (\delta^{m-1} + R^{-m+1})R^{-1})] \\ &\leq \kappa_3 \delta^m, \end{aligned}$$

where  $\kappa_3$  depends on  $\|u\|_\infty$ , and the last inequality follows from the fact that  $R^{-1} < \delta$ .

For  $I$ , using Lemma 2.4, we see that

$$\begin{aligned} I &\geq -2\delta \sup_{K \in \mathcal{K}_s} \{\tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a})\} - \sup_{K \in \mathcal{K}_s} \{L_K \tilde{\chi}_R(\bar{x})\} - \sup_{K \in \mathcal{K}_s} \{L_K \tilde{\chi}_R(\bar{y})\} + \Theta(r) \\ &=: -2\delta I_1 - I_2 - I_3 + \Theta(r), \end{aligned}$$

for some  $\Theta(r) \rightarrow 0$  as  $r \rightarrow 0$ , and in the same way as in (3.10), we conclude that

$$I \geq \kappa_4 \delta^{2s - \theta\beta} R^{-\theta\beta} - CMR^{-1} = \kappa_4 \delta^{2s + \theta\beta(\epsilon_0^{-1} - 1)} - CM\delta^{1/\epsilon_0} + \Theta(r).$$



Gathering these estimates in (3.15) and letting  $r \rightarrow 0$ , reveals

$$\kappa_4 \delta^{2s+\theta\beta(\varepsilon_0^{-1}-1)} \leq \kappa_5 (\delta^m + \delta^{\frac{1}{\varepsilon_0}}). \quad (3.16)$$

Fix  $\theta$  small, depending of  $\varepsilon_0$ , so that

$$2s + \theta\beta(\varepsilon_0^{-1} - 1) < \min\{m, 1/\varepsilon_0\}.$$

With this choice of  $\theta$ , it is easily seen that (3.16) can not hold for all small  $\delta$ . Hence  $\Phi_\delta \leq 0$ , and rest of the proof follows as before.  $\square$

Now we come to the proof of Theorem 1.5. This particular proof is different from the previous ones in the sense that the coercivity structure of the Hamiltonian becomes crucial for our argument.

*Proof of Theorem 1.5.* We may assume  $u > 0$ . Otherwise, if  $u(x_0) = 0$  for some  $x_0$ , we can apply strong maximum principle to obtain  $u \equiv 0$ , proving the theorem. Consider  $\beta, \tilde{\gamma}$  satisfying

$$\gamma < \tilde{\gamma} < \beta < 1.$$

Later we shall further choose  $\tilde{\gamma}, \beta$  in this range.

For  $\mu \in (0, 1)$ , define  $\bar{u} = \mu u$ . It is direct to see that  $\bar{u}$  solves the problem

$$\mathcal{I}\bar{u} - \mu^{1-q-m}\bar{u}^q |\nabla \bar{u}|^m = 0 \quad \text{in } \mathbb{R}^n, \quad (3.17)$$

in the viscosity sense. The proof follows the general strategy (see Section 2.1) with subtle modifications. In correspondence with (2.4), the aim is to prove the following

**Claim A.** *There exists  $\delta_1 \in (0, 1)$  such that, for all  $\mu \in (1 - \delta_1, 1)$  and  $\delta \in (0, \delta_1)$ , there exists  $\alpha_0 > 0$ , dependent on  $\delta, \mu$ , such that*

$$\Phi(x, y) := \mu u(x) - u(y) - \delta |x - y|^\beta - \alpha(1 + |x|^2)^{\tilde{\gamma}/2} \leq 0 \quad \text{in } \mathbb{R}^n \times \mathbb{R}^n,$$

for all  $\alpha \leq \alpha_0$ .

As before, this concludes the result, since by fixing two points  $x, y \in \mathbb{R}^n$ , and taking  $\alpha \rightarrow 0, \mu \rightarrow 1$  and  $\delta \rightarrow 0$  respectively, we arrive at  $u(x) - u(y) \leq 0$ , and since the argument is symmetric in  $x, y$ , we conclude that  $u$  is constant in  $\mathbb{R}^n$ .

Now we prove Claim A. Suppose on the contrary that that for some  $\mu < 1$  close to 1, and  $\delta, \alpha > 0$  close to zero, we have

$$\sup_{x, y} \Phi(x, y) > 0. \quad (3.18)$$

We remark that the contradiction argument allows us to take  $\alpha$  far smaller than  $\delta$  and  $1 - \mu$ . For the ease of notation we denote  $\psi(x) = (1 + |x|^2)^{\tilde{\gamma}/2}$ . Notice that by the growth condition imposed on  $u$  we have

$$\limsup_{|x| \rightarrow \infty} \frac{u(x)}{\alpha \psi(x)} < 1,$$

and therefore, we can find a number  $k$  such that

$$|x| \geq k \Rightarrow \Phi(x, y) \leq \mu u(x) - \alpha \psi(x) < 0.$$

On the other hand, since  $\gamma < \beta$ ,

$$\max_{|x| \leq k} \Phi(x, y) \leq \max_{|x| \leq k} (\mu u(x) - \alpha \psi(x)) - u(y) - \delta \min_{|x| \leq k} |x - y|^\beta \rightarrow -\infty,$$

as  $|y| \rightarrow \infty$ . Hence, a maximizer  $(\bar{x}, \bar{y}) \in \mathbb{R}^n \times \mathbb{R}^n$  for (3.18) exists.

Denote

$$\bar{a} = \bar{x} - \bar{y}.$$

Note that since  $u > 0$  we have  $\bar{u} \leq u$  in  $\mathbb{R}^n$ . This, together with  $\Phi(\bar{x}, \bar{y}) > 0$  imply

$$0 < \bar{u}(\bar{x}) - u(\bar{y}) \leq u(\bar{x}) - u(\bar{y}),$$

which gives  $\bar{a} = \bar{x} - \bar{y} \neq 0$ .

Again, since  $\Phi(\bar{x}, \bar{y}) > 0$ , it follows that

$$\frac{\alpha}{2^{\tilde{\gamma}/2}}(1 + |\bar{x}|)^{\tilde{\gamma}} \leq \alpha\psi(\bar{x}) \leq \mu u(\bar{x}) \leq M(1 + |\bar{x}|)^\gamma,$$

which in turn, gives

$$|\bar{x}| < 1 + |\bar{x}| \leq C_1 \alpha^{-\frac{1}{\tilde{\gamma}-\gamma}},$$

for some  $C_1 > 0$  depending on  $M$ . Moreover, using the same inequality, we also have that

$$\delta|\bar{a}|^\beta \leq \bar{u}(\bar{x}) \leq \mu M(1 + |\bar{x}|)^\gamma \leq M(C_1)^\gamma \alpha^{-\frac{\gamma}{\tilde{\gamma}-\gamma}}. \quad (3.19)$$

These bounds will be useful for our calculations below.

Denote  $\varphi(t) = t^\beta$ ,  $\phi(x, y) = \delta\varphi(|x-y|) + \alpha\psi(x)$ , and denote  $v_1$  the test function for  $\bar{u}$  with smooth representative  $\phi(\cdot, \bar{y})$  at  $\bar{x}$ , and  $v_2$  the test function for  $u$  with smooth representative  $-\phi(\bar{x}, \cdot)$  at  $\bar{y}$ . Using the equations for  $u$  and  $\bar{u}$  (recall  $\bar{u}$  solves (3.17)), and subtracting, we have

$$\tilde{I} + \tilde{I}_H \leq 0, \quad (3.20)$$

where

$$\begin{aligned} \tilde{I} &= -\mathcal{I}v_1(\bar{x}) + \mathcal{I}v_2(\bar{y}), \\ \tilde{I}_H &= \mu^{1-q-m}\bar{u}^q(\bar{x})|\delta\beta|\bar{a}^{\beta-1}\hat{a} + \alpha|\nabla\psi(\bar{x})|^m - u^q(\bar{y})|\delta\beta|\bar{a}^{\beta-1}\hat{a}^m, \end{aligned}$$

For  $\tilde{I}$ , using Lemma 2.4 we have

$$\begin{aligned} \tilde{I} &\geq -2\delta \sup_{K \in \mathcal{K}_s} \{\tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a})\} - \alpha \sup_{K \in \mathcal{K}_s} \{L_K\psi(\bar{x})\} + \Theta(r) \\ &=: -2\delta I_1 - \alpha I_2 + \Theta(r), \end{aligned}$$

where  $\Theta(r) \rightarrow 0$  as  $r \rightarrow 0$ . A simple modification of Lemma 2.3 implies that

$$I_2 \leq C\Lambda, \quad (3.21)$$

for some  $C > 0$ , not dependent on  $\delta$  and  $\alpha$ . More precisely, we note that

$$\sup_{x \in \mathbb{R}^n} [ |D^2\psi(x)| + |\nabla\psi(x)| ] \leq \kappa_2$$

for some  $\kappa_2 > 0$ . Thus

$$\sup_{K \in \mathcal{K}} L_K[B_1](\psi, x, \nabla\psi(x)) \leq \kappa_3.$$

Again, since  $x \mapsto (1 + |x|^2)^{\frac{1}{2}}$  is 1-Lipschitz, we have

$$(1 + |x - z|^2)^{\frac{\tilde{\gamma}}{2}} - (1 + |x|^2)^{\frac{\tilde{\gamma}}{2}} \leq [(1 + |x|^2)^{\frac{1}{2}} + |z|]^{\tilde{\gamma}} - (1 + |x|^2)^{\frac{\tilde{\gamma}}{2}} \leq |z|^{\tilde{\gamma}}.$$

Thus for  $s \geq 1/2$ , we get

$$\sup_{K \in \mathcal{K}} \int_{|z| \geq 1} ((1 + |x - z|^2)^{\frac{\tilde{\gamma}}{2}} - (1 + |x|^2)^{\frac{\tilde{\gamma}}{2}}) K(z) dz \leq \Lambda \int_{|z| \geq 1} |z|^{\tilde{\gamma}} \frac{1}{|z|^{n+2s}} dz \leq \kappa_4.$$

This gives us (3.21).

On the other hand, using Lemma 2.2, we can take  $\delta_0, \eta_0$  small enough (not depending on  $\bar{a}$ ,  $\delta$  and  $\alpha$ ) to obtain, for all  $K \in \mathcal{K}_s$ , that

$$\tilde{L}_K[\mathcal{C}]\tilde{\varphi}(\bar{a}) \leq -c|\bar{a}|^{\beta-2} \int_{\mathcal{C}} |z|^2 K(z) dz \leq -c\lambda|\bar{a}|^{\beta-2} \int_{\mathcal{C}} |z|^{2-n-2s} dz,$$

for some  $c > 0$ . Using the estimates on the cone in [12, Example 1], we arrive at

$$I_1 \leq -c\lambda|\bar{a}|^{\beta-2s}.$$

Combining the above estimates, we conclude that

$$\tilde{I} \geq \kappa_1\delta|\bar{a}|^{\beta-2s} - C\Lambda\alpha + \Theta(r), \quad (3.22)$$

where the constants  $\kappa_1, C$  do not depend on  $\alpha, \delta, \mu$ .

Now we compute the term  $\tilde{I}_H$ . We write

$$\tilde{I}_H = (\mu^{1-q-m} - 1) \underbrace{\bar{u}^q(\bar{x}) |\delta \beta \bar{a}|^{\beta-1} \hat{a} + \alpha \nabla \psi(\bar{x})^m}_{\Xi_1} + \underbrace{\left( \bar{u}^q(\bar{x}) |\delta \beta \bar{a}|^{\beta-1} \hat{a} + \alpha \nabla \psi(\bar{x})^m - u(\bar{y}) \delta^m \beta^m |\bar{a}|^{m(\beta-1)} \right)}_{\Xi_2}.$$

Choose  $\beta$  close to 1 so that

$$\max \left\{ \frac{(1-\beta)\gamma}{(\tilde{\gamma}-\gamma)\beta}, \frac{m(1-\beta)\gamma}{(\tilde{\gamma}-\gamma)\beta} \right\} < 1. \quad (3.23)$$

Using (3.19), we have

$$|\bar{a}|^{1-\beta} \leq \kappa_6 \delta^{\frac{\beta-1}{\beta}} \alpha^{-\frac{\gamma(1-\beta)}{(\tilde{\gamma}-\gamma)\beta}} \Rightarrow \alpha |\bar{a}|^{1-\beta} \leq \kappa_6 \delta^{\frac{\beta-1}{\beta}} \alpha^{1-\frac{\gamma(1-\beta)}{(\tilde{\gamma}-\gamma)\beta}}, \quad (3.24)$$

for some  $\kappa_6 > 0$ , not dependent on  $\delta$  and  $\alpha$ . By the choice of  $\beta$  in (3.23), we have the exponent of  $\alpha$  positive. Thus, since  $\nabla \psi(\bar{x})$  is bounded with an upper bound independent of  $\delta$  and  $\alpha$ , for  $\delta > 0$  we can find  $\alpha$  small enough so that

$$\begin{aligned} \Xi_1 &= \bar{u}(\bar{x})^q (\delta \beta |\bar{a}|^{\beta-1})^m \left| \hat{a} + \delta^{-1} \beta^{-1} \alpha |\bar{a}|^{1-\beta} \nabla \psi(\bar{x}) \right|^m \\ &\geq \frac{1}{2} \bar{u}(\bar{x})^q \delta^m \beta^m |\bar{a}|^{m(\beta-1)}, \end{aligned}$$

where we use the fact that  $m > 0$  and  $|\hat{a}| = 1$ . Similarly, for  $\Xi_2$ , using that  $\bar{u}(\bar{x}) > u(\bar{y})$  and  $m > 0$ , and taking  $\alpha$  small in terms of  $\delta$ , we see that

$$\begin{aligned} \Xi_2 &\geq u(\bar{y})^q (\delta \beta |\bar{a}|^{\beta-1})^m \left( \left| \hat{a} + \delta^{-1} \beta^{-1} \alpha |\bar{a}|^{1-\beta} \nabla \psi(\bar{x}) \right|^m - 1 \right) \\ &\geq -\kappa u(\bar{y})^p \alpha \delta^{m-1} \beta^{m-1} |\bar{a}|^{(\beta-1)(m-1)}, \end{aligned}$$

for some constant  $\kappa > 0$  depending on  $m$  and  $\|\nabla \psi\|_\infty$ , but not on  $\delta$  and  $\alpha$ , where the last inequality follows from the Lipschitz property of the map  $z \mapsto |\hat{a} + z|^m$  in  $\{|z| \leq 1/2\}$ .

Thus, using the above estimates for  $\Xi_1, \Xi_2$  and that  $\bar{u}(\bar{x}) > u(\bar{y})$ , we see that

$$\begin{aligned} \tilde{I}_H &\geq \bar{u}(\bar{x})^q \delta^m \beta^m |\bar{a}|^{m(\beta-1)} \left( \frac{1}{2} (\mu^{1-q-m} - 1) - \kappa \delta^{-1} \beta^{-1} \alpha |\bar{a}|^{1-\beta} \right) \\ &\geq \frac{(\mu^{1-q-m} - 1)}{4} \bar{u}(\bar{x})^q \delta^m \beta^m |\bar{a}|^{m(\beta-1)} \end{aligned}$$

for  $\alpha$  small enough dependent on  $\mu$  and  $\delta$ .

Combining the above estimate with (3.22) in (3.20), letting  $r \rightarrow 0$ , we see that for all  $\alpha$  small enough in terms of  $\delta$  and  $\mu$ , we have

$$\kappa_1 \delta |\bar{a}|^{\beta-2s} + \frac{(\mu^{1-q-m} - 1)}{4} \bar{u}(\bar{x})^q \delta^m |\bar{a}|^{m(\beta-1)} \leq C_1 \Lambda \alpha. \quad (3.25)$$

Now we can have one of the following two situations.

**Case 1.** There exists a subsequence  $\alpha_k \rightarrow 0$  and a constant  $\kappa_9$  such that  $|\bar{a}| = |\bar{a}(\alpha_k)| \leq \kappa_9$  for all  $k$ . Since  $\beta < 2s$ , the left most term in (3.25) remains positive whereas the r.h.s. goes to zero. This is a contradiction.

**Case 2.** There exists a subsequence  $\alpha_k \rightarrow 0$  such that  $|\bar{a}_k| = |\bar{x}_k - \bar{y}_k| \rightarrow \infty$  as  $k \rightarrow \infty$ . From  $\Phi(\bar{x}_k, \bar{y}_k) > 0$  we see that

$$\delta |\bar{a}_k|^\beta \leq \bar{u}(\bar{x}_k) \Rightarrow \bar{u}(\bar{x}_k)^q \geq \delta^q |\bar{a}_k|^{q\beta} > \kappa > 0,$$

for some  $\kappa > 0$ , as  $k \rightarrow \infty$ . Therefore, for all large  $k$ , dropping the first term in the l.h.s. of (3.25), and using (3.19) (more specifically (3.24)), we see that

$$\kappa \frac{(\mu^{1-q-m} - 1)}{4} \delta^m \leq \kappa_9 \alpha_k |\bar{a}_n|^{m(1-\beta)} \leq \kappa_{10} \alpha_k^{1 - \frac{m(1-\beta)\gamma}{\beta(\bar{\gamma}-\gamma)}},$$

where the constant  $\kappa_{10}$  does not depend on  $\alpha_k$ . By (3.23), the exponent of  $\alpha_k$  is positive, and therefore, the inequality cannot hold for large  $k$ . Therefore, (3.25) cannot hold for arbitrary small  $\alpha$ , proving Claim A.  $\square$

Next we prove Theorem 1.3

*Proof of Theorem 1.3.* Suppose, on the contrary, that  $\gamma > \max\{0, \frac{m-2s}{m-1}\}$ . Since  $u + c$  also solves (1.9), we assume that  $u(0) = 0$ . Let

$$A(r) := \max_{\bar{B}_r(0)} |u(x)|.$$

In view of (1.10)-(1.11) we have

$$0 < \limsup_{r \rightarrow \infty} \frac{A(r)}{r^\gamma} < \infty.$$

Let  $r_k \rightarrow \infty$  be such that

$$\frac{1}{C_1} r_k^\gamma \leq A(r_k) \leq C_1 r_k^\gamma \quad \text{for all } n \geq 1,$$

for some large constant  $C_1$ . Let us define

$$v_k(0) = 0, \quad v_k(x) = \frac{1}{A(r_k)} u(r_k x) \quad x \in \mathbb{R}^n.$$

Then, from the above estimate and (1.10) it follows that

$$\max_{\bar{B}} |v_k| = 1 \quad \text{and} \quad |v_k(x)| \leq C(1 + |x|)^\gamma,$$

for all  $k \geq 1$  and some constant  $C$ , independent of  $n$ . For each  $k \geq 1$ , let us now define a nonlocal operator

$$[\mathcal{I}_k \phi(\cdot)](x) = [r_k^{2s} \mathcal{I} \phi(r_k^{-1} \cdot)](r_k x).$$

Then  $\mathcal{I}_k$  is elliptic with respect to the family  $\mathcal{K}_{\text{sym}}$ . That is,

$$\inf_{K \in \mathcal{K}_{\text{sym}}} (L_K u(x) - L_K v(x)) \leq \mathcal{I}_n u(x) - \mathcal{I}_k v(x) \leq \sup_{K \in \mathcal{K}_{\text{sym}}} (L_K u(x) - L_K v(x)),$$

for all  $u, v \in C^{1,1}(x) \cap L^1(\omega_s)$ . From (1.9) we see that

$$\mathcal{I}^k v_k - \left[ A(r_k) r_k^{-\frac{m-2s}{m-1}} \right]^{m-1} |\nabla v_k|^m = 0 \quad \text{in } \mathbb{R}^n. \quad (3.26)$$

**Claim:** The family  $\{v_k\}$  locally equicontinuous in  $\mathbb{R}^n$ .

Let us first complete the proof assuming the claim. Since  $\{v_k\}$  locally equicontinuous, we can extract a subsequence  $v_{k_\ell} \rightarrow v$ , uniformly on compacts, where  $v \in C(\mathbb{R}^n)$  satisfying

$$v(0) = 0 \quad \max_{\bar{B}} |v| = 1 \quad \text{and} \quad |v(x)| \leq C(1 + |x|)^\gamma.$$

Moreover, since

$$A(r_k) r_k^{-\frac{m-2s}{m-1}} \rightarrow \infty \quad \text{as } n \rightarrow \infty,$$

using the stability property of viscosity solution and (3.26), we have

$$-|\nabla v| = 0 \quad \text{in } \mathbb{R}^n$$

in the viscosity sense. That is, for any smooth test function  $\varphi$  touching  $v$  from above at a point  $x$  we must have  $|\nabla\varphi(x)| = 0$ . It is easily seen that  $v$  must be constant, and therefore,  $v \equiv 0$ . This contradicts the fact that  $\max_{\bar{B}} |v| = 1$ . Hence  $\gamma \leq \max\{0, \frac{m-2s}{m-1}\}$ .

*Proof of the claim.* The proof follows from the argument of [10, Lemma 3.6] and we provide only a sketch here for convenience. Fix  $\ell > 0$  and define

$$\Theta = 2 \sup_k \sup_{B_{\ell+1}} |v_k| < \infty.$$

Let  $\psi : \mathbb{R}^n \rightarrow [0, 1]$  is a smooth cut-off function satisfying  $\psi = 0$  in  $B_\ell$  and  $\psi = 1$  in  $B_{\ell+\frac{1}{2}}^c$ . To prove the claim it is enough to show that for each  $\eta \in (0, 1)$  and  $\varepsilon \in (0, 1)$ , there exists  $L > 0$  satisfying

$$v_k(x) - v_k(y) \leq L|x - y|^\eta + 2\Theta(1 - \varepsilon) + \Theta\psi(x) \quad \text{for all } x, y \in B_{\ell+1}, n \geq 1. \quad (3.27)$$

To show (3.27) we consider the function

$$\Phi(x, y) = \varepsilon v_k(x) - v_k(y) - L|x - y|^\eta - \Theta\psi(x),$$

and assume on the contrary that

$$\sup_{B_{\ell+1} \times B_{\ell+1}} \Phi > 0,$$

for all large  $L$  satisfying  $L > 2^{1+2\eta}\Theta$ . Let  $(\bar{x}, \bar{y}) \in B_{\ell+1} \times B_{\ell+1}$  be a point where the supremum is attained. Also, we only need to analyze the case where  $\bar{x} \neq \bar{y}$ . Since  $\Phi(\bar{x}, \bar{y}) \geq \Phi(\bar{x}, \bar{x})$  we obtain  $L|\bar{x} - \bar{y}|^\eta \leq \Theta$ , implying  $|\bar{x} - \bar{y}| < 1/4$ . Also, let

$$p = L\eta|\bar{x} - \bar{y}|^{\eta-2}(\bar{x} - \bar{y}), \quad q = \nabla\psi(\bar{x}).$$

It is important for us to observe here

$$|p| = L\eta|\bar{x} - \bar{y}|^{\eta-1} \geq \eta\Theta^{\frac{\eta-1}{\eta}} L^{1/\eta} \rightarrow \infty,$$

as  $L \rightarrow \infty$ . Denoting

$$\varrho_k = \left[ A(r_k) r_k^{-\frac{m-2s}{m-1}} \right]^{m-1},$$

we observe from (3.26) that

$$\mathcal{I}^k(\varepsilon v_k) - \varrho_k \varepsilon^{1-m} |\nabla(\varepsilon v_k)|^m = 0 \quad \text{in } \mathbb{R}^n.$$

Now applying sub and super-solution inequality and then subtracting the inequalities we obtain the estimate

$$\begin{aligned} \varrho_k \varepsilon^{1-m} |p + \Theta q|^m - \varrho_k |p|^m &\leq \underbrace{\sup_{K \in \mathcal{K}_{\text{sym}}} \{L_K[B_\delta](\phi(\cdot, \bar{y}), \bar{x}, p + \Theta q) - L_K[B_\delta](\phi(\bar{x}, \cdot), \bar{y}, p)\}}_{:=A_1} \\ &+ \sup_{K \in \mathcal{K}_{\text{sym}}} \{L_K[B_\delta^c](\varepsilon v_k, \bar{x}, p + \Theta q) - L_K[B_\delta^c](v_k, \bar{y}, p)\}, \end{aligned} \quad (3.28)$$

for all  $\delta > 0$  small, where  $\phi(x, y) := L|x - y|^\eta + \Theta\psi(x)$  and

$$L_K[D](f, x, \xi) = \int_D (f(x+z) - f(x) - 1_{B_1}(z)z \cdot \xi) K(z) dz.$$

Using the maximality of  $\Phi$  at  $(\bar{x}, \bar{y})$ , it is standard to show that

$$\sup_{K \in \mathcal{K}_{\text{sym}}} \{L_K[B_\delta^c \cap B_{\frac{1}{4}}](\varepsilon v_n, \bar{x}, p + \Theta q) - L_K[B_\delta^c \cap B_{\frac{1}{4}}](v_n, \bar{y}, p)\} \leq \kappa\Theta,$$

for some constant  $\kappa$ . Since  $z \mapsto |z|$  is integrable with respect to  $\omega_s$ , it follows that

$$\sup_{K \in \mathcal{K}_{\text{sym}}} \{L_K[B_{\frac{1}{4}}^c](\varepsilon v_k, \bar{x}, p + \Theta q) - L_K[B_{\frac{1}{4}}^c](v_k, \bar{y}, p)\} \leq \kappa_1(1 + \Theta + |p|),$$

for some constant  $\kappa_1$ . Since  $A_1 \rightarrow 0$  as  $\delta \rightarrow 0$ , the rhs of (3.28) is dominated by  $\kappa\Theta + \kappa_1(1 + \Theta + |p|)$ . Note that for some constants  $\kappa_2, \kappa_3$  we have

$$\begin{aligned} \varepsilon^{1-m}|p + \Theta q|^m - |p|^m &\geq (1 - \varepsilon)(m - 1)|p + \Theta q|^m - \kappa_2\Theta|q|(|p|^{m-1} + |\theta q|^{m-1}) \\ &\geq \frac{1}{2}(1 - \varepsilon)(m - 1)|p|^m - \kappa_3 \end{aligned}$$

Combining these estimates in (3.28) we get

$$\frac{\varrho k}{2}(1 - \varepsilon)(m - 1)|p|^m - \kappa_3 \leq \kappa\Theta + \kappa_1(1 + \Theta + |p|).$$

Since  $|p| \rightarrow \infty$  as  $L \rightarrow \infty$  and  $m > 1$ , we obtain a contradiction from the above estimate. This proves the claim.  $\square$

We finish this section by proving Theorem 1.6

*Proof of Theorem 1.6.* We show that if there is a solution  $u$  to (1.13) then it must be positive constant. It would lead to a contradiction since  $f$  does not vanish on  $(0, \infty)$ . As before (see Theorem 1.1(i)), we consider

$$\Phi(x, y) := u(x) - u(y) - \delta\varphi(|x - y|) - M2^{1+2\alpha}R^{\gamma-\alpha}(\chi(x) + \chi(y)).$$

where  $\varphi(t) = t^\beta$  for some  $\beta \in (\gamma, 2s \wedge 1)$ . In this case, (3.2) will be replaced by

$$I \leq f(u(\bar{x})) - f(u(\bar{y})), \quad (3.29)$$

where

$$I := -\mathcal{I}v_1(\bar{x}) + \mathcal{I}v_2(\bar{y}),$$

and  $v_1, v_2$  are defined as in Theorem 1.1. Since

$$u(\bar{x}) - u(\bar{y}) \geq \delta\varphi(|\bar{x} - \bar{y}|) > 0,$$

we have  $f(u(\bar{x})) - f(u(\bar{y})) \leq 0$ . In view of Lemma 2.3 we obtain (see Theorem 1.1(i))

$$I \geq \kappa_1\delta^{\frac{2s}{\beta}}R^{\frac{\gamma}{\beta}(\beta-2s)} + \Theta(r) - CR^{\gamma-2s},$$

where  $\Theta(r) \rightarrow 0$  as  $r \rightarrow 0$ . Letting  $r \rightarrow 0$  we get

$$\kappa_1\delta^{\frac{2s}{\beta}}R^{\frac{\gamma}{\beta}(\beta-2s)} \leq CR^{\gamma-2s}, \quad (3.30)$$

where the constants  $C, \kappa_1$  does not depend on  $R$ . Since  $\gamma(2s - \beta) < \beta(2s - \gamma)$ , (3.30) can not hold for large  $R$ . Thus  $\Phi \leq 0$  for all large  $R$ , proving  $u$  to be a constant. This completes the proof of (i).

For the second part, we fix  $\beta$  as given by (1.15). Now we estimate  $f(u(\bar{x})) - f(u(\bar{y}))$  as follows: using the convexity of  $f$  we get

$$f(u(\bar{x})) - f(u(\bar{y})) \leq -f'(u(\bar{x}))(u(\bar{y}) - u(\bar{x})).$$

Since  $u(\bar{x}) - u(\bar{y}) \geq \delta\varphi(|\bar{x} - \bar{y}|) = \delta|a|^\beta$ , we have

$$f(u(\bar{x})) - f(u(\bar{y})) \leq f'(u(\bar{x}))\delta|a|^\beta,$$

using the fact  $f' \leq 0$ . Since  $f'$  is increasing, using the fact that  $u(\bar{x}) \leq CR^\gamma$ , (3.29) would give us

$$\kappa_1|a|^{\beta-2s} + \delta|a|^\beta|f'(CR^\gamma)| = \kappa_1|a|^{2s-\beta} - \delta|a|^\beta f'(CR^\gamma) \leq C_1R^{\gamma-2s}.$$

Applying Young's inequality with  $p = 2s$  and  $p' = \frac{2s}{2s-1}$ , to the left hand side, we note that

$$\kappa_2|a|^{\frac{\beta-2s}{2s} + \frac{\beta(2s-1)}{2s}}|f'(CR^\gamma)|^{\frac{2s-1}{2s}} \leq C_1R^{\gamma-2s},$$

for some constant  $\kappa_2$ . Since  $\delta|a|^\beta \leq CR^\gamma$ , we get  $|a|^{\beta-1} \geq \kappa_3R^{(\beta-1)\gamma}$ , implying

$$R^{2s-(2-\beta)\gamma}|f'(CR^\gamma)|^{\frac{2s-1}{2s}} \leq C_2$$

for some constant  $C_2$ , independent of  $R$ . This contradicts (1.15) for a sequence of  $R$  tending to infinity. Rest of the proof follows as before.  $\square$

#### 4. APPLICATIONS OF LIOUVILLE PROPERTY TO REGULARITY

In this section we showcase application of above Liouville properties in regularity estimates.

We start with interior regularity estimates for HJ equations with *critical* diffusion, in the sense that the nonlocal operator has a diffusion comparable to the one of the square root of the fractional Laplacian  $\sqrt{-\Delta}$ , that is, when  $s = 1/2$ .

In this setting, we can mention the contributions by Silvestre [40, 39], where the author considers operators of the form  $\sqrt{-\Delta}u + b(x) \cdot \nabla u = f$ . Assuming  $b, f \in L^\infty$ , it is shown in [39] that  $u \in C^\alpha$  for some  $\alpha \in (0, 1)$  whereas [40] establishes  $C^{1,\alpha}$  regularity of the solution assuming  $C^\alpha$  regularity of  $b$  and  $f$ . Later, Schwab and Silvestre [37] prove  $C^\alpha$  regularity of solutions of (4.1) for some  $\alpha$ , dependent on  $\lambda, \Lambda, C_2$  and  $n$ . In [18, Theorem 3.1], the authors consider Pucci type nonlinear equation with Hölder continuous coefficients and obtain Lipschitz regularity of the solutions. It should be noted that [18] deals with non-symmetric kernels. Theorem 4.1 improves [39, 40, 37] by proving  $C^\gamma$  regularity for any  $\gamma \in (0, 1)$  and complements the regularity result of [18] as we do not impose any regularity hypothesis on the coefficients other than continuity.

To state our next result we introduce the following state dependent operator corresponding to  $s = \frac{1}{2}$ .

$$\mathcal{J}u(x) = \inf_{a \in \mathcal{A}} \sup_{b \in \mathcal{B}} \int_{\mathbb{R}^d} (u(x+z) - u(x) - 1_B(z)z \cdot \nabla u(x)) \frac{k_{ab}(x, z)}{|z|^{n+1}} dz,$$

where  $\mathcal{A}, \mathcal{B}$  are some index sets. We impose the following condition on the kernels  $k_{ab}$ :

**(K)** *The family  $k_{ab}$  is symmetric in the sense that  $k_{ab}(z) = k_{ab}(-z)$ .  $x \mapsto k_{ab}(x, z)$  continuous in  $B_1$ , uniformly with respect to  $a, b, z$ . That is, for any compact set  $\mathcal{K} \subset B_1$ , there exists a modulus of continuity function  $\varrho$  satisfying*

$$|k_{ab}(x_1, z) - k_{ab}(x_2, z)| \leq \varrho(|x_1 - x_2|) \quad \text{for all } x_1, x_2 \in \mathcal{K}, a \in \mathcal{A}, b \in \mathcal{B}, z \in \mathbb{R}^n.$$

Furthermore, for some constants  $0 < \lambda \leq \Lambda$  we have  $\lambda \leq k_{ab} \leq \Lambda$  for all  $(a, b) \in \mathcal{A} \times \mathcal{B}$ .

The next theorem is a consequence of 1.1.

**Theorem 4.1.** *Consider the operator  $\mathcal{J}$  satisfying (K). Suppose that  $H : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  be a continuous function satisfying the following:*

(i) *There exist constants  $C_1, C_2$  satisfying*

$$|H(x, p)| \leq C_1 + C_2|p| \quad \text{for all } (x, p) \in B_1 \times \mathbb{R}^n;$$

(ii) *There exists a  $\check{H}$ , Lipschitz in  $p$ , such that*

$$\lim_{t \rightarrow 0^+} tH(x, \frac{1}{t}p) = \check{H}(x, p),$$

*uniformly on compact subsets of  $\mathbb{R}^n \times \mathbb{R}^n$ .*

Then for any  $\gamma \in (0, 1)$ , the viscosity solution  $u$  to

$$-\mathcal{J}u + H(x, u) = f \quad \text{in } B_1 \tag{4.1}$$

is in  $C^\gamma(B_{\frac{1}{2}})$ , and the  $C^\gamma$  norm depends only on  $\lambda, \Lambda, n, \gamma, \sup_{B_1} |u|, \|u\|_{L^1(\omega_{1/2})}, \|f\|_\infty$  and modulus of continuity  $\varrho$ .

*Proof.* We follow the idea of Serra [38] (see also [11, section 5]). First of all, consider a smooth cut-off function  $\xi : \mathbb{R}^n \rightarrow [0, 1]$  such that  $\xi(y) = 1$  for  $|y| \leq \frac{7}{8}$  and  $\xi(y) = 0$  for  $|y| \geq 1$ . Letting  $v = \xi u$  we see that

$$f - C\Lambda \|u\|_{L^1(\omega_s)} \leq -\mathcal{J}v + H(x, v) \leq f + C\Lambda \|u\|_{L^1(\omega_s)} \quad \text{in } B_{\frac{3}{4}}, \tag{4.2}$$

for some constant  $C$ . Dividing both side by  $\tau := 1 + \|u\|_{L^\infty(B_1)} + \|u\|_{L^1(\omega_{1/2})} + \|f\|_{L^\infty(B_1)} > 1$ , and replacing  $v$  by  $\frac{v}{\tau}$  and  $H$  by  $\frac{1}{\tau}H(x, \tau p)$  (which also satisfy the above condition with the same  $C_1, C_2$ ) we may assume that  $\|u\|_{L^\infty(B_1)} + \|u\|_{L^1(\omega_{1/2})} + \|f\|_{L^\infty(B_1)} \leq 1$ .

Set  $\gamma \in (0, 1)$ . We prove the theorem by contradiction. Suppose that there exists a family of nonempty index sets  $\mathcal{A}_k, \mathcal{B}_k$  (with kernels satisfying the continuity assumption **(K)** uniformly in  $k$ ), and a sequence of  $u_k, f_k$  solving

$$-\mathcal{J}u_k + H(x, \nabla u_k) = f_k \quad \text{in } B_1$$

such that

$$\lim_{k \rightarrow \infty} \sup_{x, y \in B_{\frac{1}{2}}} \frac{|u_k(x) - u_k(y)|}{|x - y|^\gamma} = \infty.$$

Let  $v_k = \xi u_k$ , where  $\xi$  is defined above, and we consider the equation (4.2) satisfied by  $v_k$ . As in [38, Lemma 4.3], the above is equivalent to

$$\sup_k \sup_{r > 0} \sup_{z \in B_{\frac{1}{2}}} r^{-\gamma} \|v_k(\cdot) - v_k(z)\|_{L^\infty(B_r(z))} = +\infty. \quad (4.3)$$

For  $r > 0$ , define

$$\Theta(r) = \sup_k \sup_{r_1 \geq r} \sup_{z \in B_{\frac{1}{2}}} r_1^{-\gamma} \|v_k - v_k(z)\|_{L^\infty(B_{r_1}(z))},$$

which is nondecreasing as  $r \rightarrow 0$ , and  $\Theta(r) \rightarrow +\infty$  as  $r \rightarrow 0$ .

Now choose a sequence  $(z_m, k_m, r_m) \in B_{1/2} \times \mathbb{N} \times (0, 1)$  with  $r_m \rightarrow 0$  such that

$$r_m^{-\gamma} \|v_{k_m} - v_{k_m}(z_m)\|_{L^\infty(B_{r_m}(z_m))} > \frac{\Theta(r_m)}{2},$$

and define

$$\tilde{v}_m(y) = \frac{v_{k_m}(z_m + r_m y) - v_{k_m}(z_m)}{r_m^\gamma \Theta(r_m)}.$$

Then, the following can be easily verified

$$\tilde{v}_m(0) = 0, \quad \|\tilde{v}_m\|_{L^\infty(B_1)} \geq 1/2, \quad |\tilde{v}_m(y)| \leq \kappa(1 + |y|)^\gamma, \quad y \in \mathbb{R}^n, \quad (4.4)$$

for some constant  $\kappa$ , independent of  $m$ . The first two facts are direct from the definition and the choice of  $r_m$ . For the last inequality, by the monotonicity property of  $\Theta$  it follows that

$$\|\tilde{v}_m\|_{L^\infty(B_R)} \leq R^\gamma \frac{\Theta(Rr_m)}{\Theta(r_m)} \leq R^\gamma.$$

Moreover, defining

$$-\mathcal{J}_m \phi = \inf_{a \in \mathcal{A}_{k_m}} \sup_{b \in \mathcal{B}_{k_m}} \int_{\mathbb{R}^n} (\phi(x+z) - \phi(x) - 1_B(z)z \cdot \nabla \phi(x)) \frac{k_{ab}(z_m + r_m x, r_m z)}{|z|^{n+1}} dz,$$

we see from (4.2) that

$$\left| -\mathcal{J}_m v_m + \frac{r_m^{1-\gamma}}{\Theta(r_m)} H(z_m + r_m x, r_m^{\gamma-1} \Theta(r_m) \nabla v_m) \right| \leq \frac{r_m^{1-\gamma}}{\Theta(r_m)} |f(z_m + r_m x)| + C \Lambda \frac{r_m^{1-\gamma}}{\Theta(r_m)} \quad \text{in } B_{\frac{1}{2r_m}}.$$

Due to compactness, we shall assume without any loss of generality that  $z_m \rightarrow z_\circ \in \bar{B}_{\frac{1}{2}}$ , as  $m \rightarrow \infty$ . One can also extract a subsequence  $m_k$  such that

$$\mathcal{J}_{m_k} \rightarrow \tilde{\mathcal{J}}$$



weakly (see [11, Lemma 5.4]), as  $m_k \rightarrow \infty$ , where  $\tilde{\mathcal{J}}$  is a translation invariant, positively 1-homogeneous operator which is elliptic with respect to  $\mathcal{K}_{\text{sym}}$ . Since

$$\left| \frac{r_m^{1-\gamma}}{\Theta(r_m)} H(z_m + r_m x, r_m^{\gamma-1} \Theta(r_m) \nabla v_m) \right| \leq C_1 + C_2 |\nabla v_m|,$$

applying [37, Theorem 7.2] we see that  $\{\tilde{v}_m\}$  is locally Hölder continuous, uniformly in  $m$ . Thus, we can extract a subsequence of  $\tilde{v}_k$  converging to  $w$  so that

$$-\tilde{\mathcal{J}}w + \check{H}(z_o, \nabla w) = 0 \quad \text{in } \mathbb{R}^n,$$

and, due to (4.4), we also have

$$w(0) = 0, \quad \|w\|_{L^\infty(B_1)} \geq 1/2, \quad |w(y)| \leq \kappa(1 + |y|)^\gamma. \quad (4.5)$$

Applying Theorem 1.1, we see that  $w$  is a constant, but this contradicts the first two criterion in (4.5). Hence (4.3) can not hold. This completes the proof.  $\square$

Our next result is an application of Theorem 1.6. To understand the result let us consider the following equation

$$-\mathcal{I}_{\text{sym}} u = u^{-\delta} \quad \text{in } \Omega, \quad u > 0 \quad \text{in } \Omega, \quad \text{and} \quad u = 0 \quad \text{in } \Omega^c,$$

in a smooth bounded domain  $\Omega$ . Besides the existence-uniqueness of the solutions, the regularity of the solutions up to the boundary has been of special interest. For the classical case of Laplacian we refer to [29, 30]. Similar problem for the nonlocal operator is considered recently in [1, 8] for  $-\mathcal{I}_{\text{sym}} u = (-\Delta)^s$ . More precisely, it is shown in [1] that  $u \in C^s(\bar{\Omega})$  for  $\delta < 1$  and  $u \in C^{\frac{2s}{\delta+1}}(\bar{\Omega})$  for  $\delta > 1$ . The proof techniques relies on the Green function representation of the solutions and sharp boundary regularity of the Green function. As can be seen from the proofs of [1, 29], the boundary regularity of  $u$  is *dictated* by the boundary regularity of the torsion function or the principal eigenfunction. Interestingly, for the operator  $\mathcal{I}_{\text{sym}}$  one may not expect a sharp boundary behaviour of the Dirichlet solution as point out by Ros-Oton and Serra in [35]. Our next result provides some insight on the regularity of the solution compare with the "torsion problem" when  $\delta$  is large.

**Theorem 4.2.** *Let  $u$  be a positive viscosity solution to*

$$-\mathcal{I}_{\text{sym}} u + 1_{(1,2)}(s)H(\nabla u) = \frac{k(x)}{u^\delta} \quad \text{in } \Omega, \quad u > 0 \quad \text{in } \Omega, \quad \text{and} \quad u = 0 \quad \text{in } \Omega^c,$$

for some  $\delta > 0$  where  $k \in C(\bar{\Omega})$  and  $k > 0$  in  $\bar{\Omega}$ .  $H$  is positively 1-homogeneous and locally Lipschitz. Let  $\psi$  be the solution to the torsion problem

$$-\mathcal{I}_{\text{sym}} \psi + 1_{(1,2)}(s)H(\nabla \psi) = 1 \quad \text{in } \Omega, \quad \psi > 0 \quad \text{in } \Omega, \quad \text{and} \quad u = 0 \quad \text{in } \Omega^c,$$

and for some  $0 < \beta < 1 \wedge 2s$  we have

$$(\text{dist}(x, \Omega^c))^\beta \leq C\psi(x) \quad \text{in } \Omega,$$

for some constant  $C$ . If  $\beta(1 + \delta) > 2s$ , then we have

$$(\text{dist}(x, \Omega^c))^\beta \leq C_1 u(x), \quad \text{and} \quad \sup_{x \in \Omega} \frac{u(x)}{(\text{dist}(x, \Omega^c))^\beta} = \infty.$$

In particular,  $u \notin C^\beta(\bar{\Omega})$ .

*Proof.* To simplify the notation we write  $1_{(1,2)}(s)H(p) = H_s(p)$ . The first part follows from comparison principle. More precisely, let  $k \geq c_k$  in  $\bar{\Omega}$ , for some positive constant  $c_k$ . Then

$$-\mathcal{I}_{\text{sym}} u + H_s(\nabla u) \geq \frac{c_k}{\|u\|_\infty^\delta} \quad \text{in } \Omega.$$

By comparison principle [5], we have

$$\frac{c_k}{\|u\|_\infty^\delta} \psi \leq u \quad \text{in } \mathbb{R}^d.$$

This gives the first part. Now we consider the second part. Suppose, on the contrary, that

$$\sup_{x \in \Omega} \frac{u(x)}{(\text{dist}(x, \Omega^c))^\beta} = \kappa \in (0, \infty). \quad (4.6)$$

In particular, we have from (4.6) that

$$u(x) \leq \kappa (\text{dist}(x, \Omega^c))^\beta.$$

Consider a sequence of points  $\{x_n\}$  in  $\Omega$  approaching the boundary. Let  $r_n = \text{dist}(x_n, \Omega^c)$ . Set  $\gamma = \frac{\beta(1+\delta)}{2s} \in (1, \infty)$  and define

$$w_n(y) = \frac{1}{r_n^\beta} u(x_n + r_n^\gamma y), \quad y \in \mathbb{R}^d.$$

We claim that

$$|w_n(y)| \leq \kappa_1 (1 + |y|^\beta) \quad \text{for } y \in \mathbb{R}^d, \quad (4.7)$$

for some  $\kappa_1$ , independent of  $n$ . Note that, for  $|y|^\beta \geq \|u\|_\infty$ , we have  $u(x_n + y) \leq |y|^\beta$ . Again, for  $|y|^\beta \leq \|u\|_\infty$ , we have

$$\text{dist}(x_n + y, \Omega^c) \leq \text{dist}(x_n, \Omega^c) + |y| \leq r_n + |y|.$$

Applying (4.6), we have  $u(x_n + y) \leq \kappa (r_n + |y|)^\beta \leq \kappa (r_n^\beta + |y|^\beta)$ . Since  $\beta < 1 < \gamma$ , (4.7) follows.

On the other hand, since  $2s\gamma = \beta(1 + \delta)$ ,

$$\begin{aligned} -\mathcal{I}_{\text{sym}} w_n + r_n^{(2s-1)\gamma} H_s(\nabla w_n) &= r_n^{2s\gamma - \beta - \delta} \frac{k(x_n + r_n^\gamma y)}{(w_k(y))^\delta} \\ &= \left[ \frac{r_n^{\beta\delta}}{(\text{dist}(x_n + r_n^\gamma y))^{\beta\delta}} \right] k(x_n + r_n^\gamma y) \left[ \frac{(\text{dist}(x_n + r_n^\gamma y)^\beta}{u(x_n + r_n^\gamma y)} \right]^\delta. \end{aligned}$$

Since from the first part  $w_k$  is uniformly bounded from below on every compact set  $K$  for large  $n$ , we have  $|k(x_n + r_n^\gamma y)/w_k^\delta|$  bounded. Also,

$$\left| \frac{\text{dist}(x_n + r_n^\gamma y)}{r_n} - 1 \right| = \left| \frac{\text{dist}(x_n + r_n^\gamma y) - \text{dist}(x_n, \Omega^c)}{r_n} \right| \leq r_n^{\gamma-1} |y| \rightarrow 0,$$

uniformly on compacts, as  $n \rightarrow \infty$ . Thus, using (4.7) and standard regularity theory [16, 37], we see that  $\{w_n\}$  is locally Hölder continuous, uniformly in  $n$ . Let  $w_{n_k} \rightarrow w$ , along some sub-sequence. It is evident that  $w > 0$  and  $|w| \leq \kappa_1 (1 + |y|^\beta)$ . Furthermore, from the stability property of viscosity solution, we obtain

$$-\mathcal{I}_{\text{sym}} w = \frac{c}{w^\delta} \quad \text{in } \mathbb{R}^d,$$

where  $c = \lim_{n_k \rightarrow \infty} k(x_{n_k}) > 0$ , which we can assume to exist or chose a further subsequence. Now applying Theorem 1.6 we see that such  $w$  can not exist. Hence (4.6) cannot hold. This completes the proof.  $\square$

**Acknowledgement.** Part of this project was done during a visit of A.B. and E.T. at Departamento de Matemáticas (DM) of Universidad Técnica Federico Santa María at Valparaíso. The kind hospitality of the department is acknowledged. This research of A.B. is partly supported by a SwarnaJayanti fellowship SB/SJF/2020-21/03. A.Q. was partially supported by Fondecyt Grant 1231585. E.T. was supported by Fondecyt Grant 1201897, and CNPq Grants 408169 and 306022.

## REFERENCES

- [1] Adimurthi, J. Giacomoni, and S. Santra. Positive solutions to a fractional equation with singular nonlinearity, *J. Differential Equations* 265, no.4, 1191–1226, 2018
- [2] S. Alarcón, J. García-Melián, and A. Quaas. Nonexistence of positive supersolutions to some nonlinear elliptic problems, *J. Math. Pures Appl.* (9), 99(5):618–634, 2013
- [3] N. Alibaud, F. del Teso, J. Endal, and E. R. Jakobsen. The Liouville theorem and linear operators satisfying the maximum principle, *J. Math. Pures Appl.* (9) 142 , 229–242, 2020
- [4] N. Armstrong, S. Boyan, Nonexistence of positive supersolutions of elliptic equations via the maximum principle, *Communications in Partial Differential Equations.* 36 (2011), 2011–2047.
- [5] G. Barles, E. Chasseigne, and C. Imbert. On the Dirichlet problem for second-order elliptic integro-differential equations. *Indiana Univ. Math. J.* 57(1), 213–246, 2008
- [6] B. Barrios, L. Del Pezzo, J. García-Melián, and A. Quaas. A priori bounds and existence of solutions for some nonlocal elliptic problems, *Rev. Mat. Iberoam.* 34, 195–220, 2018
- [7] B. Barrios and L. M. Del Pezzo. Study of the existence of supersolutions for nonlocal equations with gradient terms, *Milan J. Math.* 88, no.2, 267–294, 2020
- [8] B. Barrios, I. De Bonis, M. Medina, and I. Peral. Semilinear problems for the fractional laplacian with a singular nonlinearity, *Open Math.*, 13, 390–407, 2015
- [9] A. Biswas. Liouville type results for systems of equations involving fractional Laplacian in exterior domains, *Nonlinearity* 32, 2246–2268, 2019
- [10] A. Biswas and E. Topp. Nonlocal ergodic control problem in  $\mathbb{R}^d$ , appeared online in *Math. Annalen*
- [11] A. Biswas and S. Khan. Existence-Uniqueness for nonlinear integro-differential equations with drift in  $\mathbb{R}^d$ , *SIAM J. Math. Anal.* 55 (5), 4378–4409, 2023
- [12] G. Barles, E. Chasseigne, A. Ciomaga, and C. Imbert. Lipschitz regularity of solutions for mixed integro-differential equations, *J. Differential Equations*, 252(11):6012–6060, 2012
- [13] D. Berger and R. L. Schilling. On the Liouville and strong Liouville properties for a class of non-local operators, *Math. Scand.* 128(2): 365–388, 2022
- [14] M-F. Bidaut-Véron, M. García-Huidobro, and L. Véron. Estimates of solutions of elliptic equations with a source reaction term involving the product of the function and its gradient, *Duke Math. J.* 168, no.8, 1487–1537, 2019
- [15] X. Cabré, S. Dipierro, and E. Valdinoci. The Bernstein technique for integro-differential equations, *Arch. Ration. Mech. Anal.*, 243(3):1597–1652, 2022
- [16] L. Caffarelli and L. Silvestre. Regularity theory for fully nonlinear integro-differential equations, *Comm. Pure Appl. Math.*, 62(5):597–638, 2009
- [17] W. Chen and C. Li. An integral system and the Lane-Emden conjecture, *Discret. Contin. Dyn. Syst.* 24, 1167–1184, 2009
- [18] A. Ciomaga, D. Ghilli, and E. Topp. Periodic homogenization for weakly elliptic Hamilton-Jacobi- Bellman equations with critical fractional diffusion, *Comm. Partial Differential Equations*, 47(1):1-38, 2022
- [19] M. Cirant and A. Goffi. On the Liouville property for fully nonlinear equations with superlinear first-order terms, "Proceedings of the Conference on Geometric and Functional Inequalities and Recent Topics in Nonlinear PDEs", *Contemporary Mathematics, American Mathematical Society*, 781 (2023)
- [20] A. Cutri and F. Leoni. On the Liouville property for fully nonlinear equations, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 17 219–245, 2000
- [21] M. M. Fall. Entire s-harmonic functions are affine, *Proc. Amer. Math. Soc.* 144, no.6, 2587–2592, 2016
- [22] M. M. Fall and T. Weth. Liouville theorems for a general class of nonlocal operators, *Potential Anal.* 45, no.1, 187–200, 2016
- [23] A. Farina and J. Serrin, Entire solutions of completely coercive quasilinear elliptic equations II, *J. Differential Equations* 250, no. 12, 4409–4436, 2011
- [24] P. Felmer, A. Quaas, Fundamental solutions and Liouville type theorems for nonlinear integral operators, *Adv. Math.* 226 (2011), 2712–2738.
- [25] R. Filippucci, P. Pucci, and P. Souplet. A Liouville-Type Theorem for an Elliptic Equation with Superquadratic Growth in the Gradient, *Adv. Nonlinear Stud.* 20(2): 245–251, 2020
- [26] R. Filippucci, P. Pucci, and M. Rigoli. Nonlinear weighted p-Laplacian elliptic inequalities with gradient terms, *Commun. Contemp. Math.* 12 ,501–535, 2010
- [27] R. Filippucci. Nonexistence of positive weak solutions of elliptic inequalities, *Nonlinear Anal.* ,2903–2916, 2009
- [28] A. Goffi. A priori Lipschitz estimates for nonlinear equations with mixed local and nonlocal diffusion via the adjoint-Bernstein method, *Boll Unione Mat Ital*, 2022. <https://doi.org/10.1007/s40574-022-00340-w>
- [29] C. Gui and F. Lin. Regularity of an elliptic problem with a singular nonlinearity, *Proc. Roy. Soc. Edinburgh, Sec. A*, 123, 1021–1029, 1993

- [30] A. C. Lazer and P. J. McKenna. On a singular nonlinear elliptic boundary-value problem, *Proc. Amer. Math. Soc.*, 111, 721–730, 1991
- [31] H. Ishii and P. L. Lions. Viscosity solutions of fully non-linear second-order elliptic partial differential equations, *J. Differential Equations*, 83, No.1, 26–78, 1990
- [32] P. L. Lions. Quelques remarques sur les problèmes elliptiques quasilineaires du second ordre, *J. Anal. Math.* 45, 234–254, 1985
- [33] L. A. Peletier and J. Serrin. Gradient bounds and Liouville theorems for quasilinear elliptic equations, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* 5, no. 1, 65–104, 1978
- [34] A. Quaas and A. Xia. A Liouville type theorem for Lane-Emden systems involving the fractional Laplacian, *Nonlinearity* 29, 2279–2297, 2016
- [35] X. Ros-Oton and J. Serra. Boundary regularity for fully nonlinear integro-differential equations, *Duke Math. J.* 165, no.11, 2079–2154, 2016
- [36] X. Ros-Oton and J. Serra. Regularity theory for general stable operators, *J. Differential Equations* 260, no.12, 8675–8715, 2016
- [37] R. W. Schwab and L. Silvestre. Regularity for parabolic integro-differential equations with very irregular kernels, *Anal. PDE*, 9(3):727–772, 2016
- [38] J. Serra. Regularity for fully nonlinear nonlocal parabolic equations with rough kernels, *Calc. Var. Partial Differential Equations*, 54(1):615–629, 2015
- [39] Luis Silvestre. On the differentiability of the solution to an equation with drift and fractional diffusion, *Indiana University Mathematics Journal* Vol. 61, No. 2, 557–584, 2012
- [40] L. Silvestre. Hölder estimates for advection fractional-diffusion equations, *Annali della Scuola Normale Superiore di Pisa. Classe di Scienze*, 2012
- [41] Y. Yang. Gradient Estimates for the Equation  $\Delta u + cu^{-\alpha} = 0$  on Riemannian Manifolds, *Acta Mathematica Sinica*, Vol. 26, 1177–1182, 2010
- [42] X. Yu. Liouville type theorems for integral equations and integral systems, *Calc. Var. Partial Differential Equations* 46, 75–95, 2013

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH-PUNE, DR. HOMI BHABHA ROAD, PASHAN, PUNE 411008. INDIA EMAIL: [anup@iiserpune.ac.in](mailto:anup@iiserpune.ac.in)

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA CASILLA V-110, AVDA. ESPAÑA, 1680 – VALPARAÍSO, CHILE. EMAIL: [alexander.quaas@usm.cl](mailto:alexander.quaas@usm.cl)

INSTITUTO DE MATEMÁTICAS, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, RIO DE JANEIRO - RJ, 21941-909, BRAZIL; AND DEPARTAMENTO DE MATEMÁTICA Y C.C., UNIVERSIDAD DE SANTIAGO DE CHILE, CASILLA 307, SANTIAGO, CHILE. EMAIL: [etopp@im.ufrj.br](mailto:etopp@im.ufrj.br); [erwin.topp@usach.cl](mailto:erwin.topp@usach.cl).