

# ON ALGEBRAIC DEGREES OF INVERTED KLOOSTERMAN SUMS

XIN LIN AND DAQING WAN

ABSTRACT. The study of  $n$ -dimensional inverted Kloosterman sums was suggested by Katz (1995) who handled the case when  $n = 1$  from complex point of view. For general  $n \geq 1$ , the  $n$ -dimensional inverted Kloosterman sums were studied from both complex and  $p$ -adic point of view in our previous paper. In this note, we study the algebraic degree of the inverted  $n$ -dimensional Kloosterman sum as an algebraic integer.

## 1. INTRODUCTION

Let  $\mathbb{F}_q$  be the finite field of  $q$  elements with characteristic  $p$  and let  $\zeta_p$  be a fixed primitive  $p$ -th root of unit in  $\mathbb{C}$ . Let  $\psi : \mathbb{F}_q \rightarrow \mathbb{C}^*$  be the nontrivial additive character defined by

$$\psi(x) = \zeta_p^{\text{Tr}(x)},$$

where  $\text{Tr}$  is the trace map from  $\mathbb{F}_q$  to  $\mathbb{F}_p$ . For  $b \in \mathbb{F}_q^*$  and integer  $n \geq 1$ , recall that the classical  $n$ -dimensional Kloosterman sum is defined by

$$K_n(q, b) = \sum_{x_1 \cdots x_{n+1} = b, x_i \in \mathbb{F}_q^*} \psi(x_1 + \cdots + x_{n+1}) = \sum_{x_i \in \mathbb{F}_q^*} \psi\left(x_1 + \cdots + x_n + \frac{b}{x_1 \cdots x_n}\right).$$

For each fixed  $n$ , this is a family of exponential sums, parametrized by the one parameter  $b \in \mathbb{F}_q^*$ . This sum has been studied extensively in the literature. Deligne's deep theorem [Del80] gives the following sharp estimate.

$$|K_n(q, b)| \leq (n+1)\sqrt{q}^n.$$

Sperber [Spe80] determined the  $p$ -adic slopes for the L-function of the Kloosterman sums. These are local results on the number  $K_n(q, b)$ .

A global arithmetic problem is to view  $K_n(q, b)$  as an algebraic integer in the  $p$ -th cyclotomic field  $\mathbb{Q}(\zeta_p)$  and study its degree  $\deg K_n(q, b)$  over  $\mathbb{Q}$ . This degree problem turns out to be rather intricate and remains far from well-understood. A simple result from [Wan95] says that if  $\text{Tr}(b) \neq 0$ , then  $\deg K_n(q, b)$  is equal to its maximal possible value  $(p-1)/(n+1, p-1)$ . In particular, this settles the degree problem when  $q = p$ . If  $\text{Tr}(b) = 0$ , then  $\deg K_n(q, b)$  can be quite subtle. Motivated by [Wan95], the degree of a generalized form of  $K_n(q, b)$  is presented by Yang [Yan24] recently.

Katz [Kat95] raised the question of studying the following *inverted  $n$ -dimensional Kloosterman sum* defined by

$$IK_n(q, b) = \sum_{\substack{x_1 \cdots x_{n+1} = b, x_i \in \mathbb{F}_q^* \\ x_1 + \cdots + x_{n+1} \neq 0}} \psi\left(\frac{1}{x_1 + \cdots + x_{n+1}}\right).$$

Namely,

$$IK_n(q, b) = \sum_{\substack{x_i \in \mathbb{F}_q^* \\ x_1 + \cdots + \frac{b}{x_1 \cdots x_n} \neq 0}} \psi\left(\frac{1}{x_1 + \cdots + x_n + \frac{b}{x_1 \cdots x_n}}\right).$$

When  $n = 1$ , Katz [Kat95] obtained a sharp upper bound for  $IK_1(q, b)$ . This result along with paper [Eva95, Ang96] motivates the study of Ramanujan graphs. As we shall see, this inverted sum

2020 *Mathematics Subject Classification.* 11T23, 11L05.

*Key words and phrases.* Inverted Kloosterman sums, Exponential sums, Finite field, Algebraic degree.

turns out to be quite interesting in the sense that several new features occur. First, as a complex number, one does not have the expected square root cancellation estimate because there is a non-trivial main term. For any positive integer  $n$ , the sum  $IK_n(q, b)$  is estimated recently in our previous paper [LW24]. Two estimates were obtained. The first estimate is elementary. The second estimate is much deeper but assumes that  $p$  does not divide  $n + 1$ . It is based on the theorem of Denef-Loeser [DL91] for toric exponential sums which in turn depends on Deligne's theorem on the Weil conjectures. We state the two estimates below.

**Theorem 1.1.** *Notations as above. We have*

$$\left| IK_n(q, b) + \frac{(q-1)^n}{q} \right| \leq q^{\frac{n+1}{2}}.$$

If  $p \nmid n + 1$ , we further have

$$\left| IK_n(q, b) + \frac{(q-1)^n - (-1)^n(q+1)}{q} \right| \leq 2nq^{\frac{n}{2}}.$$

In this note, we view the inverted Kloosterman sum  $IK_n(q, b)$  as an algebraic integer in the  $p$ -th cyclotomic field  $\mathbb{Q}(\zeta_p)$  and study its degree as an algebraic integer. This is again quite subtle in general. As easily shown later,  $\deg IK_n(q, b)$  always divides  $(p-1)/(n+1, p-1)$ . The main result of this note is to prove that the equality holds in the case of prime field  $\mathbb{F}_p$ .

**Theorem 1.2.** *Notations as above. For all  $b \in \mathbb{F}_p^*$ , we have*

$$\deg IK_n(p, b) = \frac{p-1}{(n+1, p-1)}.$$

Our method is  $p$ -adic in nature. We first derive an elementary  $p$ -adic formula for  $IK_n(q, b)$  in terms of Gauss sums. Then, we apply the classical Stickelberger theorem to identify the  $p$ -adic main terms that would hopefully distinguish the Galois conjugates. This is not always possible for general  $q$ , but we show it works if  $q = p$ .

For the classical Kloosterman sum  $K_n(q, b)$ , all the Hodge numbers are 1. However, for the inverted Kloosterman sum  $IK_n(q, b)$ , all the Hodge numbers are 2, except for two of them which are 1. This perhaps explains why the inverted sum  $IK_n(q, b)$  is more complicated than the classical sum  $K_n(q, b)$ .

We conclude this introduction with some further research problems. The first natural problem is to prove degree results for the inverted Kloosterman sum  $IK_n(q, b)$  when  $q$  is a proper power of  $p$ . In this paper, we only treated the case  $q = p$ . It would be interesting to prove some results when  $q$  is a proper power of  $p$ . The second problem is to prove degree results for other classical exponential sums which have been studied from both complex and  $p$ -adic point of view, such as the exponential sums in [Wan04, Wan21, FW21, Li21, CL22, LC22, YZ22, ZF14].

## 2. A $p$ -ADIC FORMULA

We view the inverted Kloosterman sum as an algebraic integer in the  $p$ -adic field  $\mathbb{C}_p$ . For this purpose, it suffices to take  $\zeta_p$  to be a primitive  $p$ -th root of unity in  $\mathbb{C}_p$ . Then, the character  $\psi : \mathbb{F}_q \rightarrow \mathbb{C}_p^*$  becomes a nontrivial additive  $p$ -adic character. Recall that for integer  $n \geq 1$  and  $b \in \mathbb{F}_q^*$ , the inverted  $n$ -dimensional Kloosterman sum is defined to be

$$IK_n(q, b) = \sum_{\substack{x_1 \cdots x_{n+1} = b, \\ x_i \in \mathbb{F}_q^* \\ x_1 + \cdots + x_{n+1} \neq 0}} \psi \left( \frac{1}{x_1 + \cdots + x_{n+1}} \right).$$

The notation  $\chi : \mathbb{F}_q^* \rightarrow \mathbb{C}_p^*$  denotes a multiplicative  $p$ -adic character. By the orthogonality of characters, we have

$$IK_n(q, b) = \frac{1}{q(q-1)} \sum_{\lambda, x_i \in \mathbb{F}_q^*} \sum_{u \in \mathbb{F}_q} \psi(u(x_1 + \cdots + x_{n+1} - \lambda)) \psi \left( \frac{1}{\lambda} \right) \sum_{\chi} \chi \left( \frac{x_1 \cdots x_{n+1}}{b} \right)$$

$$\begin{aligned}
&= \frac{1}{q(q-1)} \sum_{\lambda \in \mathbb{F}_q^*} \sum_{x_i \in \mathbb{F}_q^*} \psi\left(\frac{1}{\lambda}\right) \sum_{\chi} \chi\left(\frac{x_1 \cdots x_{n+1}}{b}\right) \\
&\quad + \frac{1}{q(q-1)} \sum_{\lambda \in \mathbb{F}_q^*} \sum_{x_i \in \mathbb{F}_q^*} \sum_{u \in \mathbb{F}_q^*} \psi(u(x_1 + \cdots + x_{n+1} - \lambda)) \psi\left(\frac{1}{\lambda}\right) \sum_{\chi} \chi\left(\frac{x_1 \cdots x_{n+1}}{b}\right) \\
(2.1) \quad &:= S_1 + S_2.
\end{aligned}$$

Then

$$\begin{aligned}
S_1 &= \frac{1}{q(q-1)} \sum_{\lambda \in \mathbb{F}_q^*} \sum_{\chi} \chi^{-1}(b) \psi\left(\frac{1}{\lambda}\right) \sum_{x_i \in \mathbb{F}_q^*} \chi(x_1) \cdots \chi(x_{n+1}) \\
&= \frac{1}{q(q-1)} \sum_{\lambda \in \mathbb{F}_q^*} \psi\left(\frac{1}{\lambda}\right) \sum_{\chi} \chi^{-1}(b) \left( \sum_{x \in \mathbb{F}_q^*} \chi(x) \right)^{n+1} \\
(2.2) \quad &= -\frac{(q-1)^n}{q}.
\end{aligned}$$

Define the Gauss sum  $G(\chi)$  by

$$G(\chi) = \sum_{x \in \mathbb{F}_q^*} \chi(x) \psi(x).$$

Note that if  $\chi$  is trivial, then  $G(\chi) = -1$ . We have

$$\begin{aligned}
S_2 &= \frac{1}{q(q-1)} \sum_{\lambda, u \in \mathbb{F}_q^*} \sum_{\chi} \chi^{-1}(b) \sum_{x_i \in \mathbb{F}_q^*} \chi(x_1) \psi(ux_1) \cdots \chi(x_{n+1}) \psi(ux_{n+1}) \psi(-u\lambda) \psi\left(\frac{1}{\lambda}\right) \\
&= \frac{1}{q(q-1)} \sum_{\lambda, u \in \mathbb{F}_q^*} \sum_{\chi} \chi^{-1}(b) \chi^{-(n+1)}(u) \psi(-u\lambda) \psi\left(\frac{1}{\lambda}\right) G(\chi)^{n+1} \\
&= \frac{1}{q(q-1)} \sum_{\chi} \chi^{-1}(b) \left( \sum_{\lambda \in \mathbb{F}_q^*} \chi^{-(n+1)}\left(-\frac{1}{\lambda}\right) \psi\left(\frac{1}{\lambda}\right) \right) G(\chi^{-(n+1)}) G(\chi)^{n+1} \\
&= \frac{1}{q(q-1)} \sum_{\chi} \chi^{-1}(b) \chi^{n+1}(-1) G(\chi^{-(n+1)})^2 G(\chi)^{n+1} \\
&= \frac{1}{q(q-1)} \sum_{\chi} \chi^{n+1}(-1) \chi^{-1}(b) G(\chi^{-(n+1)})^2 G(\chi)^{n+1} \\
(2.3) \quad &= \frac{1}{q(q-1)} \left( (-1)^{n+1} + \sum_{\chi \neq 1} \chi^{n+1}(-1) \chi^{-1}(b) G(\chi^{-(n+1)})^2 G(\chi)^{n+1} \right).
\end{aligned}$$

It follows that

$$(2.4) \quad IK_n(q, b) = \frac{1}{q(q-1)} \left( -(q-1)^{n+1} + (-1)^{n+1} + \sum_{\chi \neq 1} \chi^{n+1}(-1) \chi^{-1}(b) G(\chi^{-(n+1)})^2 G(\chi)^{n+1} \right).$$

### 3. PROOF OF THEOREM

We now turn to the proof of our theorem on the degree of the inverted Kloosterman sum. At this point, we do not assume that  $q = p$  yet.

The Galois group of  $\mathbb{Q}(\zeta_p)$  over  $\mathbb{Q}$  is isomorphic to the cyclic group  $\mathbb{F}_p^*$  of order  $p - 1$ . For  $a \in \mathbb{F}_p^*$ , let  $\sigma_a$  denote the automorphism such that  $\sigma_a(\zeta_p) = \zeta_p^a$ . One checks that

$$\sigma_a(IK_n(q, b)) = \sum_{\substack{x_1 \cdots x_{n+1} = b, x_i \in \mathbb{F}_q^* \\ x_1 + \cdots + x_{n+1} \neq 0}} \psi \left( \frac{a}{x_1 + \cdots + x_{n+1}} \right) = IK_n(q, ba^{-(n+1)}).$$

Let  $H = \{a \in \mathbb{F}_p^* \mid a^{(n+1, p-1)} = 1\}$ . This is a cyclic subgroup of order  $(n+1, p-1)$  in  $\mathbb{F}_p^*$ . It is clear that  $H$  acts trivially on  $IK_n(q, b)$ , namely,  $IK_n(q, b)$  is in the fixed field  $\mathbb{Q}(\zeta_p)^H$ . By Galois theory, we deduce that  $\deg IK_n(q, b)$  divides  $(p-1)/(n+1, p-1)$ , giving the desired upper bound for  $\deg IK_n(q, b)$ . Non-trivial lower bound is harder to obtain. One needs to show that the set  $\{IK_n(q, ba^{-(n+1)}) : a \in \mathbb{F}_p^*\}$  contains many different elements. This can be difficult to do in general. Our strategy is to find a  $p$ -adic asymptotic formula for  $IK_n(q, ba^{-(n+1)})$  and shows that its main term contains many different elements as  $a$  varies in  $\mathbb{F}_p^*$ . This will produce a non-trivial lower bound.

For  $a \in \mathbb{F}_p^*$ , equation (2.4) implies

$$(3.1) \quad IK_n(q, b) - IK_n(q, ba^{-(n+1)}) = \frac{1}{q(q-1)} \left( \sum_{\chi \neq 1} \chi^{n+1}(-1)(\chi^{-1}(b) - \chi^{-1}(ba^{-(n+1)}))G(\chi^{-(n+1)})^2 G(\chi)^{n+1} \right).$$

We would like to know when this difference of Galois conjugates is non-zero. For this purpose, we first analyze each term when the non-trivial character  $\chi$  varies. If  $\chi^{n+1} = 1$ , then

$$\chi^{-1}(ba^{-(n+1)}) = \chi^{-1}(b)\chi^{n+1}(a) = \chi^{-1}(b).$$

This shows that all terms in equation (3.1) with  $\chi^{n+1} = 1$  are zero, and hence they cannot distinguish the Galois conjugates of  $IK_n(q, b)$ . One needs to find other terms with  $\chi^{n+1} \neq 1$  and with minimal  $p$ -adic valuation. It is not clear how to identify such  $p$ -adic main terms in general, but the Stickelberger theorem is useful in this regard.

Let  $\pi$  be the  $(p-1)$ th root of  $-p$  such that  $\pi \equiv \psi(1) - 1 \pmod{\pi^2}$ . Then one has  $\text{ord}_p(\pi) = 1/(p-1)$ . Let  $\omega : \mathbb{F}_q^* \rightarrow \mathbb{C}_p^*$  be the Teichmüller character. This is a primitive character of order  $q-1$ . Any character  $\chi : \mathbb{F}_q^* \rightarrow \mathbb{C}_p^*$  can be written as  $\omega^{-m}$  for a unique integer  $0 \leq m \leq q-2$ . The character  $\chi^{n+1}$  is non-trivial if and only if  $(n+1)m$  is not divisible by  $q-1$ . The Stickelberger theorem says that

$$(3.2) \quad v_p(G(\omega^{-m})) = \frac{\sigma(m)}{p-1} \quad \text{or} \quad v_\pi(G(\omega^{-m})) = \sigma(m),$$

where  $\sigma(m)$  is the sum of  $p$ -digits in the base  $p$  expansion of  $m$ , and  $v_p(z)$  denotes the  $p$ -adic valuation of  $z$  normalized such that  $v_p(p) = 1$ .

We now assume that  $q = p$  and denote  $IK_n(p, b)$  as  $IK_n(b)$ . Then,  $\chi = \omega^{-m}$  for a unique integer  $0 \leq m \leq p-2$  and the Stickelber theorem simplifies to

$$v_p(G(\omega^{-m})) = \frac{\sigma(m)}{p-1} = \frac{m}{p-1}.$$

We only need to consider those  $0 \leq m \leq p-2$  such that  $(n+1)m$  is not divisible by  $p-1$ . In this case, equation (3.1) shows that

$$(3.3) \quad p(p-1) \left( IK_n(b) - IK_n(ba^{-(n+1)}) \right) = \left( \sum_{\substack{1 \leq m \leq p-2 \\ (p-1) \nmid (n+1)m}} (-1)^{m(n+1)} \omega^m(b)(1 - \omega^m(a^{-(n+1)}))G(\omega^{m(n+1)})^2 G(\omega^{-m})^{n+1} \right).$$

If  $n + 1$  is divisible by  $p - 1$ , it is clear that  $IK_n(b) = IK_n(ba^{-(n+1)})$  for all  $a \in \mathbb{F}_p^*$  and thus  $\deg IK_n(b) = 1 = (p - 1)/(n + 1, p - 1)$ . The theorem is trivially true. In the following, we assume that  $n + 1$  is not divisible by  $p - 1$  and  $1 \leq m \leq p - 2$  with  $(p - 1) \nmid (n + 1)m$ . By the Stickelberger theorem, the  $p$ -adic valuation

$$V(m) := v_p(G(\omega^{m(n+1)})^2 G(\omega^{-m})^{n+1}) = \frac{(n+1)m + 2\{-(n+1)m\}_{p-1}}{p-1},$$

where  $\{x\}_{p-1}$  denotes the smallest non-negative residue of  $x$  modulo  $(p - 1)$ . We shall find all the terms with lower  $p$ -adic valuations in different cases.

*Case I:*  $p - 1 > n + 1$

We first consider the case  $p - 1 > n + 1$  and there is at least one term on the right side of equation (3.3), say  $m = 1$ . Since  $1 \leq m \leq p - 2$  and  $(n + 1)m$  is not divisible by  $p - 1$ , there is a unique integer  $0 \leq i \leq n$  such that

$$(p - 1)i < (n + 1)m < (p - 1)(i + 1).$$

It follows that

$$V(m) = \frac{(n+1)m + 2((p-1)(i+1) - (n+1)m)}{p-1} = \frac{2(p-1)(i+1) - (n+1)m}{p-1}$$

and

$$(3.4) \quad i + 1 < V(m) = \frac{(p-1)(i+1) + (p-1)(i+1) - (n+1)m}{p-1} < i + 2.$$

In order for  $V(m)$  to be minimal, it is necessary to have  $i = 0$ . In this case,  $m$  satisfies

$$0 < (n + 1)m < p - 1.$$

Write

$$p - 1 = k + (n + 1)h, \quad 1 \leq k \leq n + 1.$$

Then,  $m$  can take all the values in  $\{1, 2, \dots, h\}$  and one checks that

$$V(m) = 1 + \frac{k + (n + 1)(h - m)}{p - 1}.$$

These valuations are distinct as  $m$  varies in  $\{1, 2, \dots, h\}$ . In fact,

$$1 < V(h) < V(h - 1) < \dots < V(1) < 2.$$

Since  $(n + 1)m$  lies strictly between 0 and  $p - 1$ , it follows that  $(n + 1)m$  is not divisible by  $p - 1$ . This implies that for each  $1 \leq m \leq h$ , the character  $\omega^{-m(n+1)}$  is non-trivial. Reducing equation (3.3), we obtain

$$(3.5) \quad p(p - 1) \left( IK_n(b) - IK_n(ba^{-(n+1)}) \right) \equiv \sum_{1 \leq m \leq h} (1 - \omega^m(a^{-(n+1)})) A_m(b) p^{V(m)} \pmod{p^2},$$

where each  $A_m(b)$  is a unit in  $\mathbb{Z}_p[\pi]$ . For  $1 \leq m \leq h$ , it is clear that

$$1 - \omega^m(a^{-(n+1)}) \equiv 1 - a^{-m(n+1)} \pmod{p}.$$

Let  $a \in \mathbb{F}_p^*$  such that  $a^{(n+1, p-1)} \neq 1$ . The coefficient  $1 - a^{-m(n+1)}$  is not zero in  $\mathbb{F}_p$  for  $m = 1$ . Let  $m^*$  be the largest integer in the interval  $[1, h]$  such that  $1 - a^{-m(n+1)}$  is not zero in  $\mathbb{F}_p$ . Then,

$$\begin{aligned} & p(p - 1) \left( IK_n(b) - IK_n(ba^{-(n+1)}) \right) \\ & \equiv (1 - \omega^{m^*}(a^{-(n+1)})) A_{m^*}(b) p^{V(m^*)} \pmod{p^{V(m^*) + \frac{1}{p-1}}} \\ & \not\equiv 0 \pmod{p^{V(m^*) + \frac{1}{p-1}}}. \end{aligned}$$

This shows that  $IK_n(b) - IK_n(ba^{-(n+1)}) \neq 0$ . In summary, for  $a \in \mathbb{F}_p^*$ , we have proved that  $IK_n(b) - IK_n(ba^{-(n+1)}) = 0$  if and only if  $a^{(n+1, p-1)} = 1$ . It follows that  $\deg IK_n(b) = (p-1)/(n+1, p-1)$ .

*Case II:*  $\frac{1}{2}(n+1) < p-1 < n+1$

In this case, there is a unique integer  $1 \leq i \leq n-1$  such that

$$(p-1)i < (n+1)m < (p-1)(i+1).$$

Since we assume  $p-1 < n+1 < 2(p-1)$ , the integer  $m$  is uniquely determined by  $i$  and  $i$  is also bounded by  $m$ . As shown in equation (3.4), the minimal  $V(m)$  corresponds to  $i = 1$  and one has

$$(p-1) < (n+1)m < 2(p-1),$$

which implies  $m = 1$  and

$$2 < V(1) = 2 + \frac{2(p-1) - (n+1)}{p-1} < 3.$$

For  $m \geq 2$ , the restriction  $(p-1)i < (n+1)m < (p-1)(i+1)$  implies that  $i \geq 2$  and thus  $V(m) > 3$ . Combining with equation (3.3), one obtains

$$p(p-1) \left( IK_n(b) - IK_n(ba^{-(n+1)}) \right) \equiv \left( 1 - \omega \left( a^{-(n+1)} \right) \right) p^{V(1)} C_1(b) \pmod{p^3},$$

where  $C_1(b)$  is a unit in  $\mathbb{Z}_p[\pi]$ . Then for  $a \in \mathbb{F}_p^*$ , we have proved that  $IK_n(b) = IK_n(ba^{-(n+1)})$  if and only if  $a^{(n+1, p-1)} = 1$ .

*Case III:*  $p-1 < \frac{1}{2}(n+1)$

By the Stickelberger theorem, the  $p$ -adic valuation

$$(3.6) \quad W(m) := v_\pi(G(\omega^{m(n+1)})^2 G(\omega^{-m})^{n+1}) = (n+1)m + 2\{-(n+1)m\}_{p-1}.$$

We shall determine when  $W(m)$  reaches its minimum as  $1 \leq m \leq p-2$ . Denote  $h = \{n+1\}_{p-1}$ . Since we assumed that  $n+1$  is not divisible by  $p-1$ , one has  $1 \leq h \leq p-2$ . For  $m = 1$ , it can be deduced from equation (3.6) that  $W(1) = n+1 + 2(p-1-h)$ . For  $2 \leq m \leq p-2$ , there is a unique integer  $i \geq 4$  such that

$$(p-1)i < (n+1)m < (p-1)(i+1).$$

Then one has

$$(3.7) \quad W(m) - W(1) = (m-1)(n+1) - 2(p-1) + 2((i+1)(p-1) - m(n+1)) + 2h.$$

Since we assume  $n+1 > 2(p-1)$  in this case, equation (3.7) implies that  $W(m) > W(1) + 4$  for  $2 \leq m \leq p-2$ . Combining with equation (3.3), one obtains

$$p(p-1) \left( IK_n(b) - IK_n(ba^{-(n+1)}) \right) \equiv \left( 1 - \omega \left( a^{-(n+1)} \right) \right) \pi^{W(1)} C_2(b) \pmod{\pi^{W(1)+1}},$$

where  $C_2(b)$  is a unit in  $\mathbb{Z}_p[\pi]$ . Then for  $a \in \mathbb{F}_p^*$ , we have proved that  $IK_n(b) = IK_n(ba^{-(n+1)})$  if and only if  $a^{(n+1, p-1)} = 1$ . The theorem is proved.

## REFERENCES

- [Ang96] Jeff Angel, *Finite upper half planes over finite fields*, Finite Fields Appl. **2** (1996), no. 1, 62–86. MR 1371720
- [CL22] Chao Chen and Xin Lin, *L-functions of certain exponential sums over finite fields*, Math. Z. **300** (2022), no. 2, 1851–1871. MR 4363799
- [Del80] Pierre Deligne, *La conjecture de Weil. II*, Inst. Hautes Études Sci. Publ. Math. (1980), no. 52, 137–252. MR 601520
- [DL91] J. Denef and F. Loeser, *Weights of exponential sums, intersection cohomology, and Newton polyhedra*, Invent. Math. **106** (1991), no. 1, 275–294. MR 1128216
- [Eva95] Ronald Evans, *Spherical functions for finite upper half planes with characteristic 2*, Finite Fields Appl. **1** (1995), no. 3, 376–394. MR 1341954

- [FW21] Lei Fu and Daqing Wan, *On Katz's  $(A, B)$ -exponential sums*, *Q. J. Math.* **72** (2021), no. 3, 773–793. MR 4310299
- [Kat95] Nicholas M. Katz, *A note on exponential sums*, *Finite Fields Appl.* **1** (1995), no. 3, 395–398. MR 1341955
- [LC22] Xin Lin and Chao Chen, *L-functions of certain exponential sums over finite fields II*, *J. Number Theory* **241** (2022), 198–220. MR 4472439
- [Li21] Jiyou Li, *Newton polygons of L-functions associated to Deligne polynomials*, *Finite Fields Appl.* **75** (2021), Paper No. 101880, 10. MR 4272552
- [LW24] Xin Lin and Daqing Wan, *On inverted kloosterman sums over finite fields*, *Math. Z.* **306** (2024), 60.
- [Spe80] Steven Sperber, *Congruence properties of the hyper-Kloosterman sum*, *Compositio Math.* **40** (1980), no. 1, 3–33. MR 558257
- [Wan95] Daqing Wan, *Minimal polynomials and distinctness of Kloosterman sums*, *Finite Fields Appl.* **1** (1995), 189–203. MR 1337743
- [Wan04] ———, *Variation of  $p$ -adic Newton polygons for L-functions of exponential sums*, *Asian J. Math.* **8** (2004), no. 3, 427–472. MR 2129244
- [Wan21] ———, *Exponential sums over finite fields*, *J. Syst. Sci. Complex.* **34** (2021), no. 4, 1225–1278. MR 4298049
- [Yan24] Liping Yang, *Degrees of generalized kloosterman sums*, *Forum Math.* (2024), <http://doi.org/10.1515/forum-2023-0295>.
- [YZ22] Liping Yang and Hao Zhang, *Generic Newton polygons for L-functions of  $(A, B)$ -exponential sums*, *Finite Fields Appl.* **78** (2022), 101980. MR 4349885
- [ZF14] Jun Zhang and Weiduan Feng, *On L-functions of certain exponential sums*, *Finite Fields Appl.* **26** (2014), 7–31. MR 3151355

DEPARTMENT OF MATHEMATICS, SHANGHAI MARITIME UNIVERSITY, SHANGHAI 201306, PR CHINA.  
*Email address:* xlin1126@hotmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CA 92697-3875 USA.  
*Email address:* dwan@math.uci.edu