

A higher index on finite-volume locally symmetric spaces

Hao Guo,^{*} Peter Hochs[†] and Hang Wang[‡]

July 24, 2024

Abstract

Let G be a connected, real semisimple Lie group. Let $K < G$ be maximal compact, and let $\Gamma < G$ be discrete and such that $\Gamma \backslash G$ has finite volume. If the real rank of G is 1 and Γ is torsion-free, then Barbasch and Moscovici obtained an index theorem for Dirac operators on the locally symmetric space $\Gamma \backslash G/K$. We obtain a higher version of this, by constructing an index of Dirac operators on G/K in the K -theory of an algebra on which the conjugation-invariant terms in Barbasch and Moscovici's index theorem define continuous traces. The resulting index theorems also apply when Γ has torsion. The cases of these index theorems for traces defined by semisimple orbital integrals extend to Song and Tang's higher orbital integrals, and yield nonzero and computable results even when $\text{rank}(G) > \text{rank}(K)$, or the real rank of G is larger than 1.

Contents

1	Introduction	2
2	A higher index	5
2.1	An algebra of exponentially decaying kernels	5
2.2	Construction of the index	9

^{*}Tsinghua University, haoguo@mail.tsinghua.edu.cn

[†]Radboud University, p.hochs@math.ru.nl

[‡]East China Normal University, wanghang@math.ecnu.edu.cn

3	Index theorems	11
3.1	Geometric traces	11
3.2	Spectral traces	14
3.3	Relation with the classical index	16
3.4	Index theorems: semisimple and non-semisimple cases	17
4	Exponentially decaying kernels	22
4.1	The algebra $\mathcal{A}_S(E)$	22
4.2	Suprema of Schwartz kernels	24
4.3	Decay behaviour of Schwartz kernels	28
5	Traces on a Schwartz algebra	31
5.1	Schwartz spaces	31
5.2	Geometric traces on $\mathcal{S}(G)$	33
5.3	Spectral traces on $\mathcal{S}(G)$	36
6	Contributions from semisimple elements	38
6.1	A holomorphic fixed-point formula	38
6.2	A fixed-point set	41
6.3	A normal bundle	43
6.4	A fixed-point formula on G/T	45
6.5	Higher orbital integrals	48
6.6	Proofs of the index theorems	49
7	Roe algebras and von Neumann algebras	50
7.1	Roe algebras	50
7.2	The equivariant index for cocompact actions	52
7.3	Von Neumann algebras and the von Neumann trace	54

1 Introduction

Consider a connected, real semisimple Lie group G , a maximal compact subgroup $K < G$ and a discrete subgroup $\Gamma < G$. *Locally symmetric spaces* of the form $X = \Gamma \backslash G/K$ come up in many places in mathematics, and include, for example, all hyperbolic manifolds, several moduli spaces, and the space $\mathrm{SL}(2, \mathbb{Z}) \backslash \mathrm{SL}(2, \mathbb{R}) / \mathrm{SO}(2)$ on which modular forms are defined. Spaces X that arise in this way are sometimes compact, but in many relevant cases (such as the example mentioned) they only have finite volume. Because of the relevance of these spaces, it is a worthwhile question how to construct and compute topological invariants for them.

Index theory is a useful tool for constructing and computing invariants of manifolds. This works particularly well if Γ is torsion-free and $\Gamma \backslash G$ is compact, so that X is a compact, smooth manifold. Then elliptic differential operators on X have well-defined Atiyah–Singer indices. And by considering the action by Γ on G/K rather than just its quotient, one can construct a higher index of Γ -equivariant elliptic operators [7, 24, 39], with values in $K_*(C^*(\Gamma))$. Pairings of this index with various cyclic cocycles were computed by Connes and Moscovici [14, 15].

If X is smooth and has finite volume, but is not necessarily compact, then Moscovici [29] constructed an index of elliptic operators on X . This was computed for Dirac operators by Barbasch and Moscovici [6] in the case where the real rank of G is 1. That computation was based on a version of Selberg’s trace formula for real rank 1 groups due to Osborne and Warner [30, 37]. The result is an expression for the index of Dirac operators as a sum of various contributions, all but finitely many of which are given by semisimple orbital integrals. In fact, if X is compact, then one only needs these semisimple contributions, so the non-semisimple contributions reflect the noncompactness of X . Some of these non-semisimple contributions are defined by conjugation-invariant functionals, which can be expected to define traces on suitable convolution algebras. The sum of the other contributions is conjugation-invariant as well, even though the individual terms are not.

Our goal here is to unify these constructions and computations, and obtain a higher index theorem on finite-volume locally symmetric spaces if G has real rank 1. More precisely, we aim to construct a K -theoretic index from which

- the classical index can be recovered,
- if G has real rank 1, then the conjugation-invariant parts of Barbasch and Moscovici’s expression for the classical index on X can be extracted and computed individually, and
- the usual Γ -equivariant index can be recovered if X is compact.

To achieve this aim, we show that

1. every G -equivariant, self-adjoint, elliptic, first-order differential operator on G/K has a well-defined index in $K_*(\mathcal{A}(G))$, where $\mathcal{A}(G)$ is an analogue of Casselman’s Schwartz algebra [12];
2. this index can be represented by a standard idempotent involving heat operators [15];

3. the semisimple orbital integrals and the conjugation-invariant non-semisimple components of Barbasch and Moscovici's index theorem define continuous traces on $\mathcal{A}(G)$.

There are several commonly used convolution algebras on G besides Casselman's Schwartz algebra on which semisimple orbital integrals define continuous traces. These include $C_c^\infty(G)$ and Harish-Chandra's L^1 - or L^2 -Schwartz algebras. We use Casselman's algebra because it is large enough so that the first two points above are true (the second is not true for $C_c^\infty(G)$), and small enough so that the third point is true (which is not true for some non-semisimple contributions if one uses Harish-Chandra's Schwartz algebras). See also Remark 5.3.

Because of the second point above, applying the traces from the third point to the index in the first point leads to expressions involving heat operators, which were computed in [6]. For traces not defined by semisimple orbital integrals, we then simply apply the computations in [6]. For semisimple orbital integrals, we give an independent index-theoretic computation. This may be of independent interest, and it also has the advantages that

- it generalises to the higher orbital integrals from [35], which yield nonzero results even when $\text{rank}(G) > \text{rank}(K)$, in which case the classical index vanishes;
- our computations also apply when Γ has torsion, so that X is an orbifold, and we obtain nonzero contributions from nontrivial elliptic elements;
- the index theorems for semisimple orbital integrals apply even when G does not have real rank 1.

(We note, however, that the computations for semisimple orbital integrals are more about the action by G on G/K than about the group Γ .)

Our higher index on G/K is constructed as a special case of a general higher index of elliptic operators equivariant with respect to proper, co-compact group actions. This takes values in the K -theory of an algebra of exponentially fast decaying kernels. The more general index may be useful in a wider context, for pairing traces or cyclic cocycles with higher indices.

At the end of the paper, we relate our index to the widely used index in terms of Roe algebras, and to an index involving a von Neumann algebra. From the latter, one can recover the L^2 -index. In the special case where X is compact, we use the Roe algebra index to recover the Γ -equivariant index from our higher index.

Acknowledgements

PH was supported by the Netherlands Organisation for Scientific Research NWO, through ENW-M grants OCENW.M.21.176 and OCENW.M.23.063. HW was supported by the grants 23JC1401900, NSFC 12271165 and in part by Science and Technology Commission of Shanghai Municipality (No. 22DZ2229014).

2 A higher index

2.1 An algebra of exponentially decaying kernels

Consider a Riemannian manifold X , with Riemannian distance d and Riemannian density μ .

Definition 2.1. The manifold X has *uniformly exponential growth* if there are $a_0, C > 0$ such that for all $x \in X$,

$$\int_X e^{-a_0 d(x, x')} d\mu(x') \leq C.$$

Example 2.2. Suppose that a group H acts isometrically on X , such that X/H is compact. Let $x_0 \in X$ and $a_0 > 0$, and suppose that

$$\int_X e^{-a_0 d(x_0, x')} d\mu(x')$$

converges. Let $r > 0$ be such that $H \cdot B_r(x_0) = X$. Then for all $x \in X$, write $x = hx_1$, for $h \in H$ and $x_1 \in B_r(x_0)$. Then by H -invariance of d and μ and the triangle inequality,

$$\int_X e^{-a_0 d(x, x')} d\mu(x') = \int_X e^{-a_0 d(x_1, x')} d\mu(x') \leq e^{a_0 r} \int_X e^{-a_0 d(x_0, x')} d\mu(x').$$

So X has uniformly exponential growth. This example includes the case where $X = G/K$, for a connected Lie group G , compact subgroup $K < G$, and a G -invariant Riemannian metric on X .

We suppose from now on that X has uniformly exponential growth. We consider a Hermitian vector bundle $E \rightarrow X$. A section κ of the exterior tensor product $E \boxtimes E^* \rightarrow X \times X$ has *finite propagation* if there is an $r > 0$ such that for all $x, x' \in X$ with $d(x, x') > r$, we have $\kappa(x, x') = 0$. If two

L^∞ -sections κ, λ of $E \boxtimes E^*$ have finite propagation, then their composition $\kappa\lambda$ is well-defined as an L^∞ -section of $E \boxtimes E^*$ with finite propagation, by

$$(\kappa\lambda)(x, x') = \int_X \kappa(x, x'')\lambda(x'', x') d\mu(x''), \quad (2.1)$$

for all $x, x' \in X$.

Fix a vector space S of differential operators on $E \boxtimes E^*$ such that

1. S contains the identity operator;
2. for all $l \in \mathbb{Z}_{\geq 0}$, the set of order l differential operators in S is a finite-dimensional linear subspace of S ;
3. for all $P \in S$, there are $Q, R \in S$ such that for all $\kappa, \lambda \in \Gamma^\infty(E \boxtimes E^*)$ with finite propagation,

$$P(\kappa\lambda) = (Q\kappa)(R\lambda). \quad (2.2)$$

Example 2.3. Let G be a Lie group, and let V be a finite-dimensional, complex vector space. Consider the trivial vector bundle $E = G \times V \rightarrow G$. Let R be the right regular representation of the universal enveloping algebra $\mathcal{U}(\mathfrak{g})$ on $C^\infty(G)$. For $P_1, P_2 \in \mathcal{U}(\mathfrak{g})$, let the operator $R(P_1) \otimes R(P_2)$ on $\Gamma^\infty(E \boxtimes E^*) = C^\infty(G \times G) \otimes \text{End}(V)$ be defined by the action by $R(P_1)$ on the first copy of G , and the action by $R(P_2)$ on the second copy of G . Let

$$S := \{R(P_1) \otimes R(P_2); P_1, P_2 \in \mathcal{U}(\mathfrak{g})\}.$$

Then S satisfies the three conditions listed above. For the third, if $P = R(P_1) \otimes R(P_2)$, then we may take $Q = R(P_1) \otimes 1$ and $R = 1 \otimes R(P_2)$.

Let $\Gamma_{S, \text{fp}}^\infty(E \boxtimes E^*)$ be the space of smooth sections κ of $E \boxtimes E^* \rightarrow X \times X$ with finite propagation, such that $P\kappa$ is bounded for all $P \in S$.

Definition 2.4. For $a > 0$ and $P \in S$, the seminorm $\|\cdot\|_{a, P}$ on $\Gamma_{S, \text{fp}}^\infty(E \boxtimes E^*)$ is given by

$$\|\kappa\|_{a, P} = \sup_{x, x' \in X} e^{ad(x, x')} \|(P\kappa)(x, x')\|, \quad (2.3)$$

for $\kappa \in \Gamma_{S, \text{fp}}^\infty(E \boxtimes E^*)$. We denote the completion of $\Gamma_{S, \text{fp}}^\infty(E \boxtimes E^*)$ in the seminorms $\|\cdot\|_{a, P}$, for $a > 0$ and $P \in S$, by $\mathcal{A}_S(E)$.

Lemma 2.5. *The space $\mathcal{A}_S(E)$ is a Fréchet algebra under the composition (2.1).*

Lemma 2.5 is proved in Subsection 4.1.

Now let H be a Lie group (possibly with infinitely many connected components, so H may be discrete), acting isometrically on X and such that X/H is compact. Suppose that E is an H -equivariant vector bundle, and that the action preserves the Hermitian metric on E . Consider the action by $H \times H$ on $\Gamma^\infty(E \boxtimes E^*)$ given by

$$((h, h') \cdot \kappa)(x, x') := h\kappa(h^{-1}x, h'^{-1}x')h'^{-1}, \quad (2.4)$$

for $h, h' \in H$, $x, x' \in X$ and $\kappa \in \Gamma^\infty(E \boxtimes E^*)$. Suppose that the operators in S commute with this action. Unless specified otherwise, by the action of H on $\Gamma^\infty(E \boxtimes E^*)$, we mean the restriction of the action (2.4) to the diagonal. (But note the notation (7.7).)

Lemma 2.6. *The action by H on $\Gamma^\infty(E \boxtimes E^*)$ restricts to an isometric action on $\mathcal{A}_S(E)$. The fixed-point set $\mathcal{A}_S(E)^H$ of this action is a Fréchet algebra.*

Lemma 2.6 is proved in Subsection 4.1.

We will use slight generalisations of the algebras $\mathcal{A}_S(E)$ and $\mathcal{A}_S(E)^H$. Let Y be another Riemannian manifold with uniformly exponential growth. Let $\varphi: Y \rightarrow X$ be a smooth map. If $\kappa \in \Gamma^\infty(E \boxtimes E^*)$, then we define $\varphi^*\kappa \in \Gamma^\infty(\varphi^*E \boxtimes \varphi^*E^*)$ by

$$(\varphi^*\kappa)(y, y') = \kappa(\varphi(y), \varphi(y')) \in E_{\varphi(y)} \otimes E_{\varphi(y')}^* = (\varphi^*E \boxtimes \varphi^*E^*)_{(y, y')}, \quad (2.5)$$

for $y, y' \in Y$. We assume that φ is

1. surjective;
2. a *quasi-isometry* in the sense that there are $A, B > 0$ such that for all $y, y' \in Y$,

$$A^{-1}d_Y(y, y') - B \leq d_X(\varphi(y), \varphi(y')) \leq Ad_Y(y, y') + B, \quad (2.6)$$

where d_X is the Riemannian distance on X , and d_Y is the one on Y ;

3. an *algebra homomorphism* in the sense that for all $\kappa, \lambda \in \Gamma_{\text{fp}}^\infty(E \boxtimes E^*)$, we have

$$\varphi^*(\kappa\lambda) = (\varphi^*\kappa)(\varphi^*\lambda).$$

In the third point, we note that the second point implies that $(\varphi^*\kappa)$ and $(\varphi^*\lambda)$ have finite propagation, so their composition is defined.

Let S be a vector space of differential operators as above, but on the vector bundle $\varphi^*E \rightarrow Y$ rather than on E itself. Let $\Gamma_{S,\varphi,\text{fp}}^\infty(E \boxtimes E^*)$ be the space of smooth sections κ of $E \boxtimes E^* \rightarrow X \times X$ with finite propagation, such that $P\varphi^*\kappa$ is bounded for all $P \in S$.

Definition 2.7. For $a > 0$ and $P \in S$, the seminorm $\|\cdot\|_{a,P}$ on $\Gamma_{S,\varphi,\text{fp}}^\infty(E \boxtimes E^*)$ is given by

$$\|\kappa\|_{a,P} = \|\varphi^*\kappa\|_{a,P} \quad (2.7)$$

for $\kappa \in \Gamma_{S,\text{fp}}^\infty(E \boxtimes E^*)$, where the right hand side is defined as in (2.3). We denote the completion of $\Gamma_{S,\varphi,\text{fp}}^\infty(E \boxtimes E^*)$ in the seminorms $\|\cdot\|_{a,P}$, for $a > 0$ and $P \in S$, by $\mathcal{A}_{S,\varphi}(E)$.

In this definition, note that surjectivity of φ implies injectivity of φ^* . Hence if all seminorms (2.7) are zero, then in particular $\|\kappa\|_{1,1} = 0$, so $\kappa = 0$.

By the construction and assumptions, the expression (2.5) defines an isometric algebra homomorphism

$$\varphi^* : \mathcal{A}_{S,\varphi}(E) \xrightarrow{\cong} \mathcal{A}_S(\varphi^*E). \quad (2.8)$$

Hence $\mathcal{A}_{S,\varphi}(E)$ is a Fréchet algebra by Lemma 2.5.

If the Lie group H also acts on Y , and φ is H -equivariant, then φ^*E naturally becomes an H -equivariant vector bundle. The algebra isomorphism 2.8 is now H -equivariant, so $\mathcal{A}_{S,\varphi}(E)^H \subset \mathcal{A}_{S,\varphi}(E)$ is a Fréchet algebra by Lemma 2.6, isomorphic to $\mathcal{A}_S(\varphi^*E)^H$ via (2.8).

Example 2.8. If $Y = X$ and $\varphi = \text{id}_X$ is the identity map, then $\mathcal{A}_{S,\text{id}_X}(E) = \mathcal{A}_S(E)$ and $\mathcal{A}_{S,\text{id}_X}(E)^H = \mathcal{A}_S(E)^H$. So Definition 2.7 is a generalisation of Definition 2.4.

Example 2.9. Consider the setting of Example 2.3. Let $K < G$ be a compact subgroup, and suppose that V carries a unitary representation of K . Let $X = G/K$, and let $E = G \times_K V \rightarrow X$ be the quotient of $G \times V$ by the K -action given by

$$k \cdot (g, v) = (gk^{-1}, k \cdot v),$$

for $k \in K$, $g \in G$ and $v \in V$. Let $q: G \rightarrow X$ be the quotient map. Then $q^*E = G \times V$, and q satisfies all the assumptions on the map φ above. For the third, we choose a left Haar measure dg on G , and define the measure $d(gK)$ on G/K by

$$\int_{G/K} f(gK) d(gK) = \int_G (q^*f)(g) dg \quad (2.9)$$

for all $f \in C_c(G/K)$. Then for all $\kappa, \lambda \in \Gamma_{\text{fp}}^\infty(E \boxtimes E^*)$, and all $g, g' \in G$,

$$\begin{aligned} q^*(\kappa\lambda)(g, g') &= \int_{G/K} \kappa(gK, g''K) \lambda(g''K, g'K) d(gK) \\ &= \int_G \kappa(gK, g''K) \lambda(g''K, g'K) dg = (q^*\kappa)(q^*\lambda)(g, g'). \end{aligned}$$

Consider the actions by G on X , E and G induced by left multiplication. Then the operators $R(P_1) \otimes R(P_2)$ in Example 2.3 are $G \times G$ -equivariant for all $P_1, P_2 \in \mathcal{U}(\mathfrak{g})$. And q is equivariant for this action, so we obtain the Fréchet algebra $\mathcal{A}_{S,q}(E)^G$.

Note that the operators $R(P_1) \otimes R(P_2)$ do not preserve the subspace

$$\Gamma^\infty(E \boxtimes E^*) = (C^\infty(G \times G) \otimes \text{End}(V))^{K \times K} \subset C^\infty(G \times G) \otimes \text{End}(V),$$

which is why the algebra $\mathcal{A}_{S,q}(E)^G$ is used in this case.

Some motivation for the use of the algebra $\mathcal{A}_{S,\varphi}(E)^H$, in particular in the setting of Example 2.9, is given in Remark 5.3.

2.2 Construction of the index

Let $\hat{B}(\mathbb{R})$ be the completion of $C_c^\infty(\mathbb{R})$ in the seminorms defined by

$$\|\psi\|_{a,k} := \sup_{\xi \in \mathbb{R}} e^{a|\xi|} |\psi^{(k)}(\xi)|, \quad (2.10)$$

for $\psi \in C_c^\infty(\mathbb{R})$, $a > 0$ and $k \in \mathbb{Z}_{\geq 0}$. Then $\hat{B}(\mathbb{R})$ lies inside the space of Schwartz functions, hence so does its inverse Fourier transform $B(\mathbb{R})$. We equip the latter with the seminorms given by

$$\|f\|_{a,k} := \|\hat{f}\|_{a,k} \quad (2.11)$$

for $f \in B(\mathbb{R})$, $a > 0$ and $k \in \mathbb{Z}_{\geq 0}$.

Lemma 2.10. *The space $B(\mathbb{R})$ is a Fréchet algebra with respect to pointwise multiplication.*

This lemma is proved in Subsection 4.1.

Consider the setting of Definition 2.7 in the equivariant case, i.e. including the actions by H . We will denote the Schwartz kernel of an operator T by κ_T . Let D be an H -equivariant, first-order, elliptic, self-adjoint differential operator on E .

Proposition 2.11. *For all $f \in B(\mathbb{R})$, the Schwartz kernel $\kappa_{f(D)}$ of the operator $f(D)$ lies in $\mathcal{A}_{S,\varphi}(E)^H$. In fact, the map $f \mapsto \kappa_{f(D)}$ is a continuous, injective $*$ -homomorphism from $B(\mathbb{R})$ to $\mathcal{A}_{S,\varphi}(E)^H$.*

This proposition is proved at the end of Subsection 4.3.

Suppose that the conditions of Proposition 2.11 hold. Then this proposition yields a continuous, injective $*$ -homomorphism

$$j: B(\mathbb{R}) \rightarrow \mathcal{A}_{S,\varphi}(E)^H. \quad (2.12)$$

Let $B(D) \subset \mathcal{A}_{S,\varphi}(E)^H$ be the image of j ; we denote the inclusion map $B(D) \hookrightarrow \mathcal{A}_{S,\varphi}(E)^H$ by j as well. We view $B(D)$ as a Fréchet algebra, with the topology for which j defines a homeomorphism from $B(\mathbb{R})$ to $B(D)$. Let $\mathcal{M}(B(D))$ be the multiplier algebra of $B(D)$. This is the algebra of all continuous linear maps $T: B(D) \rightarrow B(D)$ such that for all $f_1, f_2 \in B(D)$, we have $T(f_1 f_2) = (T f_1) f_2 = f_1 T f_2$. Because $B(D)$ is commutative and has a bounded approximate identity, $\mathcal{M}(B(D))$ is a Fréchet algebra containing $B(D)$ as an ideal [1].

Consider the maps

$$K_1(\mathcal{M}(B(D))/B(D)) \xrightarrow{\partial} K_0(B(D)) \xrightarrow{j^*} K_0(\mathcal{A}_{S,\varphi}(E)^H), \quad (2.13)$$

where the first map is a boundary map in the long exact sequence associated to the ideal $B(D) \subset \mathcal{M}(B(D))$.

Suppose that E has a $\mathbb{Z}/2\mathbb{Z}$ grading. Suppose that D is odd with respect to this grading. We denote the restrictions of D to even- and odd-graded sections by D^+ and D^- , respectively.

Theorem 2.12. *The operator D lies in $\mathcal{M}(B(D))$. It is invertible in $\mathcal{M}(B(D))$ modulo $B(D)$. The image of the resulting class in $K_1(\mathcal{M}(B(D))/B(D))$ under the map (2.13) equals*

$$\left[\begin{pmatrix} e^{-tD^- D^+} & e^{-\frac{t}{2}D^- D^+} \frac{1 - e^{-tD^- D^+}}{D^- D^+} D^+ \\ e^{-\frac{t}{2}D^+ D^-} D^+ & 1 - e^{-tD^+ D^-} \end{pmatrix} - \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right] \in K_0(\mathcal{A}_{S,\varphi}(E)^H). \quad (2.14)$$

This theorem is proved at the end of Subsection 4.3.

Definition 2.13. The H -equivariant *higher index* of D with respect to S and φ is the class (2.14). It is denoted by $\text{index}_{S,H}(D) \in K_0(\mathcal{A}_{S,\varphi}(E)^H)$.

Remark 2.14. We use the algebra $B(D)$ to define the higher index of D , because D may not be a multiplier of the larger algebra $\mathcal{A}_{S,\varphi}(E)^H$, and it may not have an inverse modulo $\mathcal{A}_{S,\varphi}(E)^H$ in the multiplier algebra of $\mathcal{A}_{S,\varphi}(E)^H$. See also Remark 4.6.

3 Index theorems

In this section, let G be a noncompact, connected, real semisimple Lie group, with finite centre. (Later, G will sometimes be assumed to have real rank 1.) Fix a Haar measure dg on G . Let $K < G$ be maximal compact, and let dk be the Haar measure on K giving it unit volume. Let $X = G/K$, equipped with a G -invariant Riemannian metric defined by an $\text{Ad}(K)$ -invariant inner product on the Lie algebra \mathfrak{g} . The resulting Riemannian distance is denoted by d as before. Let d_G be the Riemannian distance on G defined by the same inner product on \mathfrak{g} .

3.1 Geometric traces

We will use some ‘geometric’ traces on an algebra of matrix-valued functions on G , defined by integrals over subsets of G . We also use some ‘spectral’ traces, involving the operator trace, introduced in Subsection 3.2. All these traces correspond to terms in Barbasch and Moscovici’s index theorem [6].

We write L and R for the left and right regular representations of G on smooth functions, respectively. We use the same notation for the corresponding representation $\mathcal{U}(\mathfrak{g})$.

Definition 3.1. For $f \in C_c^\infty(G)$, $a > 0$ and $P_1, P_2 \in \mathcal{U}(\mathfrak{g})$, we write

$$\|f\|_{a, P_1, P_2} := \sup_{g \in G} e^{ad_G(e, g)} |(L(P_1)R(P_2)f)(g)|. \quad (3.1)$$

Let $\mathcal{S}(G)$ be the completion of $C_c^\infty(G)$ in the seminorms $\|\cdot\|_{a, P_1, P_2}$.

Remark 3.2. The algebra $\mathcal{S}(G)$ is a version of Casselman’s Schwartz algebra [12]. We use seminorms defined by left and right derivatives, whereas only left derivatives are used in [12]. We use both left and right derivatives so that $\mathcal{S}(G)$ embeds into Harish-Chandra’s Schwartz spaces, see Lemma 5.2. It is not essential to use both left and right derivatives, because in the end we will always restrict to subalgebras of $\mathcal{S}(G)$ corresponding to a finite number of K -types. Then it is sufficient to only use left or right derivatives to obtain an embedding into Harish-Chandra’s Schwartz spaces, by (a slight extension of) Lemma 18 in [18].

Lemma 3.3. *The space $\mathcal{S}(G)$ is a Fréchet algebra with respect to convolution.*

This lemma is proved in Subsection 5.1.

A relation between the algebra $\mathcal{S}(G)$ and the algebra from Subsection 2.1 will play a central role. Consider the setting of Example 2.9. Then we have the Fréchet algebra $\mathcal{A}_{S,q}(E)^G$. The Fréchet algebra $\mathcal{S}(G) \otimes \text{End}(V)$ is the completion of $C_c^\infty(G) \otimes \text{End}(V)$ in the seminorms given by

$$\|f\|_{a,P_1,P_2}^V := \sup_{g \in G} e^{ad_G(e,g)} \|(L(P_1)R(P_2)f)(g)\|_{\text{End}(V)},$$

for $a > 0$, $P_1, P_2 \in \mathcal{U}(\mathfrak{g})$ and $f \in C_c^\infty(G) \otimes \text{End}(V)$. Consider the action by $K \times K$ on $\mathcal{S}(G) \otimes \text{End}(V)$ given by

$$((k, k') \cdot f)(g) = \pi(k)f(k^{-1}gk')\pi(k')^{-1}, \quad (3.2)$$

for $k, k' \in K$, $f \in \mathcal{S}(G) \otimes \text{End}(V)$ and $g \in G$. Analogously to Lemma 2.6, elements of $K \times K$ act on $\mathcal{S}(G) \otimes \text{End}(V)$ by continuous operators, so that the fixed point set $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$ is a Fréchet algebra.

Consider the fixed-point set $(C^\infty(G) \otimes \text{End}(V))^{K \times K}$ of the action (3.2) by $K \times K$ on $C^\infty(G) \otimes \text{End}(V)$.

Lemma 3.4. *For all $\kappa \in \Gamma^\infty(E \boxtimes E^*)^G$, setting*

$$[e, \Psi(\kappa)(g)v] = \kappa(eK, gK)[g, v], \quad (3.3)$$

for $g \in G$ and $v \in V$, defines a function $\Psi(\kappa) \in (C^\infty(G) \otimes \text{End}(V))^{K \times K}$. This defines a linear isomorphism $\Psi: \Gamma^\infty(E \boxtimes E^*)^G \rightarrow (C^\infty(G) \otimes \text{End}(V))^{K \times K}$. This linear isomorphism restricts to an algebra isomorphism from $\Gamma_{\text{fp}}^\infty(E \boxtimes E^*)^G$ to $(C_c^\infty(G) \otimes \text{End}(V))^{K \times K}$, which in turn extends continuously to an isometric algebra isomorphism from $\mathcal{A}_{S,q}(E)^G$ to $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$.

This lemma is proved in Subsection 5.1.

We will use several continuous traces on the algebra $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$. If τ is a linear functional on $\mathcal{S}(G)$, or on the subspace of $\mathcal{S}(G)$ corresponding to the $K \times K$ -isotypical components in $\text{End}(V)$, then we write

$$\tau^V := \tau \otimes \text{tr}_V: (\mathcal{S}(G) \otimes \text{End}(V))^{K \times K} \rightarrow \mathbb{C}.$$

If H is any group, $H' < H$ a subgroup, and $h \in H$, then we write H'_h for the centraliser of h in H' .

Let $g \in G$ be such that G_g is unimodular. Fix a Haar measure dh on G_g . Because G_g is unimodular, there is a unique G -invariant measure $d(G_g h)$ on $G_g \backslash G$ such that for all $f \in C_c(G)$,

$$\int_G f(h) dh = \int_{G_g \backslash G} \int_{G_g} f(h'h) dh' d(G_g h).$$

For $f \in C^\infty(G)$ such that the following converges, consider the orbital integral

$$\tau_g(f) := \int_{G_g \backslash G} f(h^{-1}gh) d(G_g h). \quad (3.4)$$

Proposition 3.5. *If g is semisimple, then τ_g^V defines a continuous trace on $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$.*

This proposition is proved in Subsection 5.2.

In Theorem 3.17, we will see that semisimple orbital integrals (the traces from Proposition 3.5) map the higher index of the operators we consider to zero if $\text{rank}(G) > \text{rank}(K)$. The higher index itself is generally nonzero in that case, however, and this nonvanishing can be detected by pairing the index with cyclic cocycles defined by the *higher orbital integrals* from [35].

Let $P < G$ be a cuspidal parabolic subgroup, with Langlands decomposition $P = MA'N'$. (We use primes to distinguish the groups A' and N' from the groups A and N in an Iwasawa decomposition $G = KAN$ below.) Let $g \in M$ be semisimple. In Definition 3.3 and Theorem 3.5 in [35], Song and Tang constructed a cyclic cocycle $\Phi_{P,g}$ over Harish-Chandra's Schwartz algebra $\mathcal{C}^2(G)$, whose degree is the dimension of A' . If $\text{rank}(G) = \text{rank}(K)$, then G is a cuspidal parabolic subgroup of itself, and $\Phi_{G,g} = \tau_g$.

Proposition 3.6. *The operator trace on $\text{End}(V)$ and the cyclic cocycle $\Phi_{P,g}$ combine into a cyclic cocycle $\Phi_{P,g}^V$ on $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$.*

This proposition is proved in Subsection 5.2.

We will also use some traces that are not associated to semisimple elements of G . From now on, until the end of Subsection 3.3, we suppose that G has real rank 1. Consider an Iwasawa decomposition $G = KAN$, and a compatible Γ -cuspidal parabolic subgroup $P = MAN$. Then A is one-dimensional by assumption. Let $\lambda \in \mathfrak{a}^*$ be the simple restricted root of $(\mathfrak{g}, \mathfrak{a})$ that determines N . Then we have the restricted root space decomposition $\mathfrak{n} = \mathfrak{n}_\lambda \oplus \mathfrak{n}_{2\lambda}$. With $N_\lambda := \exp(\mathfrak{n}_\lambda)$ and $N_{2\lambda} := \exp(\mathfrak{n}_{2\lambda})$, we have $N = N_\lambda N_{2\lambda}$, where $N_\lambda \cap N_{2\lambda} = \{e\}$.

We equip \mathfrak{n} , and hence \mathfrak{n}_λ and $\mathfrak{n}_{2\lambda}$ with the $\text{Ad}(K)$ -invariant inner product defined by the Killing form and a Cartan involution. Since $N_{2\lambda}$ lies in the centre of N , we have $G_n = M_n N$ for all nontrivial $n \in N_{2\lambda}$, so G_n is unimodular. So the orbital integral $\tau_n(f)$ can be defined on suitable functions.

Lemma 3.7. *For all $n \in N_{2\lambda}$, the orbital integral τ_n^V defines a continuous trace on $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$.*

This lemma is proved in Subsection 5.2.

For $a \in A$ and $\xi \in \mathfrak{a}^*$, we write $a^\xi := e^{\langle \xi, \log a \rangle}$. Let

$$\rho_N := (\dim(\mathfrak{n}_\lambda)/2 + \dim(\mathfrak{n}_{2\lambda}))\lambda. \quad (3.5)$$

In a few places, we will use the fact that the Haar measure dg on G can be decomposed as

$$\int_G f(g) dg = \int_{KAN} f(kan) a^{2\rho_N} dk da dn, \quad (3.6)$$

for all $f \in C_c(G)$, for Haar measures dk on K , da on A and dn on N .

For $f \in C^\infty(G)$ such that the following converges, write

$$\tau_\lambda(f) := \int_N \int_K f(knk^{-1}) dk dn. \quad (3.7)$$

Proposition 3.8. *The integral (3.7) defines a continuous trace τ_λ^V on $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$.*

This proposition is proved in Subsection 5.2.

3.2 Spectral traces

Apart from the geometric traces of Subsection 3.1, we also use some ‘spectral’ traces on $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$.

Let $\Gamma < G$ be a discrete subgroup such that $\Gamma \backslash G$ has finite volume. As in [6, 30, 37], we make the following technical assumption. Let P_1, \dots, P_l be representatives of the Γ -conjugacy classes of Γ -cuspidal parabolic subgroups of G , with Langlands decompositions $P_j = M_j A_j N_j$. Let Z_Γ be the centre of Γ . Then we assume that for all j ,

$$\Gamma \cap P_j = Z_\Gamma \cdot (\Gamma \cap N_j). \quad (3.8)$$

This condition implies that a noncentral element $\gamma \in \Gamma$ is semisimple if and only if its conjugacy class has empty intersection with each of the groups P_j ; see Lemma 5.3 in [37]. That in turn implies that for all semisimple $\gamma \in \Gamma$, the quotient $\Gamma_\gamma \backslash G_\gamma$ has finite volume; see Lemma 5.4 in [37]. In general, a discrete subgroup $\Gamma < G$ such that $\Gamma \backslash G$ has finite volume has a finite-index normal subgroup with the above property. (See page 56 of [37] and Lemma 6.5 in [16].) It is possible to work without this condition, then noncentral semisimple elements of Γ should be replaced by noncentral elements whose conjugacy classes have empty intersections with the groups P_j .

Example 3.9. Let $G = \mathrm{SL}(2, \mathbb{R})$ and $\Gamma = \mathrm{SL}(2, \mathbb{Z})$. There is one Γ -conjugacy class of Γ -cuspidal parabolic subgroups, represented by the subgroup $P < G$ of upper-triangular matrices. In its Langlands decomposition $P = MA'N'$, we have

$$N' = \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}; b \in \mathbb{R} \right\}, \quad \text{so}$$

$$\Gamma \cap N' = \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}; b \in \mathbb{Z} \right\}.$$

And $Z_\Gamma = \{\pm I\}$, so

$$Z_\Gamma \cdot (\Gamma \cap N') = \left\{ \begin{pmatrix} a & b \\ 0 & a \end{pmatrix}; a \in \{\pm 1\}, b \in \mathbb{Z} \right\} = \Gamma \cap P.$$

So (3.8) holds.

Consider the right quasiregular representation R^Γ of G in $L^2(\Gamma \backslash G)$: for all $g, h \in G$ and $\varphi \in L^2(\Gamma \backslash G)$,

$$(R^\Gamma(g)\varphi)(\Gamma h) = \varphi(\Gamma hg).$$

Write $L^2(\Gamma \backslash G) = L_d^2(\Gamma \backslash G) \oplus L_c^2(\Gamma \backslash G)$, where $L_d^2(\Gamma \backslash G)$ decomposes discretely into irreducible representations of G , and $L_c^2(\Gamma \backslash G)$ decomposes continuously. Let R_d^Γ be the restriction of R^Γ to $L_d^2(\Gamma \backslash G)$. It defines a representation of $L^1(G)$ in the usual way. We consider its extension

$$(R_d^\Gamma)^V : (L^1(G) \otimes \mathrm{End}(V))^{K \times K} \rightarrow \mathcal{B}((L_d^2(\Gamma \backslash G) \otimes V)^K).$$

Proposition 3.10. *For all $f \in (\mathcal{S}(G) \otimes \mathrm{End}(V))^{K \times K}$, the operator $(R_d^\Gamma)^V(f)$ is trace-class. The algebra homomorphism*

$$(R_d^\Gamma)^V : (\mathcal{S}(G) \otimes \mathrm{End}(V))^{K \times K} \rightarrow \mathcal{L}^1((L_d^2(\Gamma \backslash G) \otimes V)^K) \quad (3.9)$$

is continuous.

Corollary 3.11. *The map $f \mapsto \mathrm{Tr}(R_d^\Gamma(f))$ defines a continuous trace $(\mathrm{Tr} \circ R_d^\Gamma)^V$ on $(\mathcal{S}(G) \otimes \mathrm{End}(V))^{K \times K}$.*

Proposition 3.10 and Corollary 3.11 are proved in Subsection 5.3.

Let $A(\mathfrak{n}_\lambda)$ be the area of the unit sphere in \mathfrak{n}_λ , and let $C_\lambda(\Gamma)$ and $C_{2\lambda}(\Gamma)$ be as on page 299 of [30]. Fix $n_0 \in N_{2\lambda}$ such that $\|\log(n_0)\| = 1$. For

$f \in C^\infty(G)$ such that $R_d^\Gamma(f)$ is trace-class and the other terms converge, we consider the ‘remainder’

$$\begin{aligned} \tau_{\text{rem}}(f) &:= \text{Tr}(R_d^\Gamma(f)) \\ &- \sum_{(\gamma) \text{ semisimple}} \text{vol}(\Gamma_\gamma \backslash G_\gamma) \tau_\gamma(f) - \frac{C_{2\lambda}(\Gamma)}{2} (\tau_{n_0}(f) + \tau_{n_0^{-1}}(f)) - \frac{C_\lambda(\Gamma)}{A(\mathfrak{n}_\lambda)|\lambda|} \tau_\lambda(f), \end{aligned} \quad (3.10)$$

Here the sum on the right hand side is over the Γ -conjugacy classes (γ) of semisimple elements of Γ .

Remark 3.12. An explicit form of the functional τ_{rem} is given in [30]: see the theorems on pages 293 and 299 in that paper.

Proposition 3.13. *The functional τ_{rem} defines a continuous trace τ_{rem}^V on $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$.*

This proposition is proved at the end of Subsection 5.3.

3.3 Relation with the classical index

We first note that the index of elliptic operators on real rank one, finite-volume locally symmetric spaces studied in [6, 29] can be recovered from the higher index of Definition 2.13. Then we show that in the special case where X is compact, then Γ -equivariant index can be recovered from this higher index as well.

Consider the setting of Example 2.9. Let D be an operator as above Proposition 2.11, on sections of $E \rightarrow X$. Suppose that D is G -equivariant. Then by Theorem 2.12 and Example 2.9, it has an equivariant higher index

$$\text{index}_{\mathcal{S},G}(D) \in K_0(\mathcal{A}_{\mathcal{S},q}(E)^G). \quad (3.11)$$

Using the isomorphism $\Psi: \mathcal{A}_{\mathcal{S},q}(E)^G \rightarrow (\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$ from Lemma 3.4, we can map this index to $K_0((\mathcal{S}(G) \otimes \text{End}(V))^{K \times K})$. Then applying the map induced by the algebra homomorphism $(R_d^\Gamma)^V$ from Proposition 3.10 yields an element of $K_0(\mathcal{L}^1(L_d^2(\Gamma \backslash G) \otimes V)) = \mathbb{Z}$.

Let D^Γ be the elliptic differential operator on $\Gamma \backslash G \times_K V \rightarrow \Gamma \backslash G/K$ induced by D . It has a well-defined index by Theorem 2.1 in [29].

Corollary 3.14. *If G has real rank 1, then under the isomorphism*

$$K_0(\mathcal{L}^1(L_d^2(\Gamma \backslash G) \otimes V)) = \mathbb{Z}, \quad (3.12)$$

we have

$$(R_d^\Gamma)_*^V \circ \Psi_*(\text{index}_{\mathcal{S},G}(D)) = \text{index}(D^\Gamma). \quad (3.13)$$

Proof. The isomorphism (3.12) is given by the operator trace. So by Theorem 2.12, the left hand side of (3.13) equals

$$\mathrm{Tr}(R_d^\Gamma(\Psi(e^{-tD^-D^+}))) - \mathrm{Tr}(R_d^\Gamma(\Psi(e^{-tD^+D^-}))).$$

By Proposition 3.1 in [6], this equals the right hand side of (3.13). \square

Remark 3.15. The number (3.13) is the value of the continuous trace $\mathrm{Tr} \circ (R_d^\Gamma)^V \circ \Psi$ on $\mathrm{index}_{S,G}(D)$.

Remark 3.16. By (1.2.5) in [6], the index (3.13) vanishes if $\mathrm{rank}(G) > \mathrm{rank}(K)$. The higher index (3.11) will generally be nonzero, however. Indeed, by Corollary 7.4, the higher index (3.11) is nonzero whenever $\mathrm{index}_G(D)$ is nonzero. In fact, Bradd [9, 10] showed that the inclusion $\mathcal{S}(G) \hookrightarrow C_r^*(G)$ induces an isomorphism on K -theory, if one restricts to isotypical components corresponding to certain finite sets of K -types. This implies that the higher index (3.11) is nonzero if and only if $\mathrm{index}_G(D)$ is nonzero. And that is true for the operators D_W that we consider in the next subsection.

3.4 Index theorems: semisimple and non-semisimple cases

Next, we use the algebra isomorphism from Lemma 3.4 to apply the traces from Subsections 3.1 and 3.2 to the higher index (3.11) in the case of Dirac operators, and obtain a higher version of the index theorem in [6]. For now we drop the assumption that G has real rank 1, this will be assumed again in the index theorem for non-semisimple contributions, Theorem 3.20.

The index theorems are stated for Dirac operators on G/K used in many places, e.g. [4, 6, 31, 38]. These are defined as follows. Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be a Cartan decomposition. We equip \mathfrak{p} with the K -invariant inner product defined by the Killing form, and use the corresponding G -invariant Riemannian metric on G/K . The representation $\mathrm{Ad}: K \rightarrow \mathrm{SO}(\mathfrak{p})$ lifts to $\widetilde{\mathrm{Ad}}: \widetilde{K} \rightarrow \mathrm{Spin}(\mathfrak{p})$, for a double cover $\widetilde{K} \rightarrow K$. In this way, we view the standard representation $\Delta_{\mathfrak{p}}$ of $\mathrm{Spin}(\mathfrak{p})$ as a representation of \widetilde{K} . Let W be an irreducible representation of \widetilde{K} . Suppose that the representation of \widetilde{K} in $\Delta_{\mathfrak{p}} \otimes W$ descends to a representation of K . This is the case, for example, if $\widetilde{\mathrm{Ad}}$ descends to K , and we take W to be a lift of a representation of K .

The space of smooth sections of $E = G \times_K (\Delta_{\mathfrak{p}} \otimes W) \rightarrow G/K$ is isomorphic to $(C^\infty(G) \otimes \Delta_{\mathfrak{p}} \otimes W)^K$, where K acts on $C^\infty(G)$ via the right regular representation. Let c be the Clifford action by \mathfrak{p} on $\Delta_{\mathfrak{p}}$. Let $\{X_1, \dots, X_r\}$

be an orthonormal basis of \mathfrak{p} . Then we have the Dirac operator

$$D_W := \sum_{j=1}^r R(X_j) \otimes c(X_j) \otimes 1_W \quad (3.14)$$

on $\Gamma^\infty(E) = (C^\infty(G) \otimes \Delta_{\mathfrak{p}} \otimes W)^K$. It is equivariant with respect to the natural action by G on E . We suppose that G/K is even-dimensional. Then The space $\Delta_{\mathfrak{p}}$ has a $\text{Spin}(\mathfrak{p})$ -invariant $\mathbb{Z}/2\mathbb{Z}$ -grading $\Delta_{\mathfrak{p}} = \Delta_{\mathfrak{p}}^+ \oplus \Delta_{\mathfrak{p}}^-$, and the operator D_W is odd with respect to the induced grading on E . Consider its equivariant higher index

$$\text{index}_{S,G}(D_W) \in K_0(\mathcal{A}_{S,q}(E)^G). \quad (3.15)$$

Consider the traces $\tau_\gamma^{\Delta_{\mathfrak{p}} \otimes W}$ from Proposition 3.5, for semisimple $\gamma \in \Gamma$, $(\tau_{n_0} + \tau_{n_0-1})^{\Delta_{\mathfrak{p}} \otimes W}$ from Lemma 3.7, $\tau_\lambda^{\Delta_{\mathfrak{p}} \otimes W}$ from Proposition 3.8 and $T_\lambda^{\Delta_{\mathfrak{p}} \otimes W}$ from Proposition 3.13 on the algebra $(\mathcal{S}(G) \otimes \text{End}(\Delta_{\mathfrak{p}} \otimes W))^{K \times K}$. Via the isomorphism Ψ from Lemma 3.4, these define continuous traces on $\mathcal{A}_{S,q}(E)^G$. These can be applied to (3.15). In fact, higher cyclic cocycles $\Phi_{P,\gamma}^{\Delta_{\mathfrak{p}} \otimes W}$ as in Proposition 3.6, for γ semisimple, can also be applied to the index (3.15) and computed, yielding possibly nonzero results in cases where the trace $\tau_\gamma^{\Delta_{\mathfrak{p}} \otimes W}$ maps (3.15) to zero.

To state index theorems for the resulting numbers, we introduce some notation in the various cases.

Index theorem for semisimple contributions

Let $T < K$ be a maximal torus. If T is a Cartan subgroup of G , so $\text{rank}(G) = \text{rank}(K)$, then we use the following notation. Fix a positive root system $R^+(G, T)$ for (G, T) , and consider the Weyl group $W_K = N_K(T)/T$. Let ρ be half the sum of the roots in $R^+(G, T)$, let ρ_c be half the sum of the roots in $R^+(K, T) = R^+(G, T) \cap R(K, T)$, and let $\rho_n = \rho - \rho_c$.

If $g \in T$, then we write $R^+(G_g, T) = R^+(G, T) \cap R(G_g, T)$, and $W_{K_g} = N_{K_g}(T)/T$. Let ρ_g be half the sum of the roots in $R^+(G_g, T)$.

Let λ be the highest weight of W with respect to the set of compact positive roots $R^+(K, T) \subset R^+(G, T)$. For $\mu \in i\mathfrak{t}^*$, we write

$$d_\mu^{G_g} = \prod_{\alpha \in R^+(G_g, T)} \frac{(\mu, \alpha)}{(\rho_g, \alpha)}. \quad (3.16)$$

(This includes the special case where $g = e$, so $G_g = G$.) For discrete series representation of G_g with Harish-Chandra parameter μ , this is its formal degree.

Theorem 3.17 (Index theorem, semisimple case). *Suppose that $\text{rank}(G) = \text{rank}(K)$.*

(a) *If $\gamma \in \Gamma$ is central in G , then*

$$\tau_\gamma^{\Delta_{\mathfrak{p}} \otimes W} \circ \Psi_*(\text{index}_{S,G}(D_W)) = e^{\lambda - \rho_n}(\gamma) d_{\lambda + \rho_c}^G.$$

(b) *If $\gamma \in \Gamma$ is hyperbolic, then*

$$\tau_\gamma^{\Delta_{\mathfrak{p}} \otimes W} \circ \Psi_*(\text{index}_{S,G}(D_W)) = 0.$$

(c) *If $\gamma \in \Gamma$ is elliptic, then with respect to a compact Cartan subgroup T containing γ ,*

$$\begin{aligned} & \tau_\gamma^{\Delta_{\mathfrak{p}} \otimes W} \circ \Psi_*(\text{index}_{S,G}(D_W)) = \\ & (-1)^{(\dim(G/K) + \dim(G_\gamma/K_\gamma))/2} \frac{\sum_{w \in W_{K_\gamma} \setminus W_K} \det(w) e^{w(\lambda + \rho_c - \rho_\gamma)}(\gamma) d_{w(\lambda + \rho_\gamma - \rho_n)}^{G_\gamma}}{e^{\rho - \rho_\gamma}(\gamma) \prod_{\alpha \in R^+(G,T) \setminus R^+(G_\gamma,T)} (1 - e^\alpha(\gamma))}. \end{aligned}$$

Index theorem for semisimple contributions from higher cocycles

As in Proposition 3.6, consider a maximal cuspidal parabolic subgroup $P = MA'N' < G$, and a semisimple element $g \in M$. If P is a maximal cuspidal parabolic subgroup, then $T < M$, after conjugation if necessary. Then T is a compact Cartan subgroup of M . Fix a positive root system $R^+(M, T) \subset R(M, T)$. If $g \in T$, we write

$$R^+(M_g, T) := R(M_g, T) \cap R^+(M, T).$$

Let ρ^M be half the sum of the elements of $R^+(M, T)$, let ρ_c^M be half the sum of the elements of $R^+(K \cap M, T) := R^+(M, T) \cap R(K \cap M, T)$, and let $\rho_n^M = \rho^M - \rho_c^M$. Let ρ_g^M be half the sum of the roots in $R^+(M_g, T)$.

As in [20], we now assume that the map $\widetilde{\text{Ad}}: \widetilde{K} \rightarrow \text{Spin}(\mathfrak{p})$ descends to K . By Lemma 4.3 in [20], there is a natural $K \cap M$ -equivariant embedding of $\mathfrak{k}/(\mathfrak{k} \cap \mathfrak{m})$ into \mathfrak{p} , via an identification of $\mathfrak{k}/(\mathfrak{k} \cap \mathfrak{m})$ with the orthogonal complement of $(\mathfrak{p} \cap \mathfrak{m}) \oplus \mathfrak{a}$. Hence we obtain the representation $K \cap M \rightarrow \text{Spin}(\mathfrak{k}/(\mathfrak{k} \cap \mathfrak{m})) \rightarrow \text{GL}(\Delta_{\mathfrak{k}/(\mathfrak{k} \cap \mathfrak{m})})$. The spaces $\mathfrak{k}/\mathfrak{t}$ and $(\mathfrak{k} \cap \mathfrak{m})/\mathfrak{t}$ are both even-dimensional, hence so is $\mathfrak{k}/(\mathfrak{k} \cap \mathfrak{m})$. So the representation $\Delta_{\mathfrak{k}/(\mathfrak{k} \cap \mathfrak{m})}$ of $K \cap M$ has a natural $\mathbb{Z}/2\mathbb{Z}$ grading $\Delta_{\mathfrak{k}/(\mathfrak{k} \cap \mathfrak{m})} = \Delta_{\mathfrak{k}/(\mathfrak{k} \cap \mathfrak{m})}^+ \oplus \Delta_{\mathfrak{k}/(\mathfrak{k} \cap \mathfrak{m})}^-$. If we view

$\Delta_{\mathfrak{t}/(\mathfrak{t} \cap \mathfrak{m})}$ as the virtual representation $[\Delta_{\mathfrak{t}/(\mathfrak{t} \cap \mathfrak{m})}^+] - [\Delta_{\mathfrak{t}/(\mathfrak{t} \cap \mathfrak{m})}^-] \in R(K \cap M)$, then some of the multiplicities $m_U \in \mathbb{Z}$ in

$$\Delta_{\mathfrak{t}/(\mathfrak{t} \cap \mathfrak{m})} \otimes W|_{K \cap M} = \bigoplus_{U \in \widehat{K \cap M}} m_U U, \quad (3.17)$$

may be negative.

For each $U \in \widehat{K \cap M}$, let $\lambda_U \in i(\mathfrak{t} \cap \mathfrak{m})^*$ be its highest weight with respect to the positive root system $R^+(K \cap M, T)$. Consider the Weyl groups $W_{K \cap M} = N_{K \cap M}(T)/T$ and $W_{K \cap M_g} = N_{K \cap M_g}(T)/T$. For $\mu \in i(\mathfrak{t} \cap \mathfrak{m})^*$, define $d_\mu^{M_g}$ analogously to (3.16), where the product now runs over $\alpha \in R^+(M_g, T)$. For semisimple $g \in M$, let $\Phi_{P,g}^{\Delta_p \otimes W}$ be as in Proposition 3.6.

Theorem 3.18 (Index theorem, semisimple case for higher cocycles). *If P is not a maximal cuspidal parabolic subgroup of G , then*

$$\langle \Phi_{P,\gamma}^{\Delta_p \otimes W}, \Psi_*(\text{index}_{S,G}(D_W)) \rangle = 0$$

for all semisimple $\gamma \in \Gamma \cap M$. Suppose now that P is a maximal cuspidal parabolic subgroup of G .

(a) *If $\gamma \in \Gamma \cap M$ is central in M , then*

$$\langle \Phi_{P,\gamma}^{\Delta_p \otimes W}, \Psi_*(\text{index}_{S,G}(D_W)) \rangle = \sum_{U \in \widehat{K \cap M}} m_U e^{\lambda_U - \rho_n^M}(\gamma) d_{\lambda_U + \rho_n^M}^M.$$

(b) *If $\gamma \in \Gamma \cap M$ is hyperbolic in M , then*

$$\langle \Phi_{P,\gamma}^{\Delta_p \otimes W}, \Psi_*(\text{index}_{S,G}(D_W)) \rangle = 0.$$

(c) *If $\gamma \in \Gamma \cap M$ is elliptic in M , then with respect to a compact Cartan subgroup $T < M$ containing γ ,*

$$\begin{aligned} \langle \Phi_{P,\gamma}^{\Delta_p \otimes W}, \Psi_*(\text{index}_{S,G}(D_W)) \rangle = & \\ & (-1)^{(\dim(M/(K \cap M)) + \dim(M_\gamma/(K \cap M_\gamma)))/2} \\ & \sum_{U \in \widehat{K \cap M}} m_U \frac{\sum_{w \in W_{K \cap M_\gamma} \setminus W_{K \cap M}} \det(w) e^{w(\lambda_U + \rho_c^M - \rho_\gamma^M)}(\gamma) d_{w(\lambda_U + \rho_\gamma^M - \rho_n^M)}^{M_\gamma}}{e^{\rho^M - \rho_\gamma^M}(\gamma) \prod_{\alpha \in R^+(M,T) \setminus R^+(M_\gamma,T)} (1 - e^\alpha(\gamma))}. \end{aligned}$$

Remark 3.19. If $\text{rank}(G) = \text{rank}(K)$, then G is a maximal cuspidal parabolic subgroup of itself, and $\Phi_{G,g} = \tau_g$ for all semisimple $g \in G$. Hence Theorem 3.18 reduces to Theorem 3.17 in this case.

Index theorem for non-semisimple contributions

To state an index theorem for traces not associated to semisimple elements of Γ , we introduce some further notation. We now assume that G has real rank 1.

Let l be the number of Γ -conjugacy classes of Γ -cuspidal parabolic subgroups of G . Choose representatives $P_j = M_j A_j N_j$ of these conjugacy classes. If $G = \mathrm{SU}(2n, 1)$, then we write

$$C_2(\Gamma) := \sum_{j=1}^l \frac{\mathrm{vol}(\Gamma \cap N_j \backslash N_j)}{\mathrm{vol}(\Gamma \cap (N_j)_{2\lambda} \backslash (N_j)_{2\lambda})} \frac{(2\pi)^{2n}}{(2n)!} B_n,$$

where B_n is the n th Bernoulli number. Furthermore, let $\mathfrak{t} \subset \mathfrak{su}(2n, 1)$ be the diagonal Cartan. Let $Z_0 \in \mathfrak{t}$ be the diagonal matrix with diagonal elements $i, i, \dots, i, -2ni$. Let

$$\varepsilon(R^+(G, T)) = \mathrm{sign} \prod_{\alpha \in R^+(G, T) \backslash R^+(K, T)} \langle \alpha, Z_0 \rangle.$$

For general G , if $R_0^+(G, T)$ is a positive root system chosen as in Section 2.a of [2], then we write

$$\varepsilon(\lambda + \rho_c) := \mathrm{sign} \prod_{\alpha \in R_0^+(G, T)} (\lambda + \rho_c, \alpha).$$

If $N_{2\lambda}$ is trivial, then we choose a vector $\tilde{X} \in \mathfrak{n}_\lambda$ of norm $\frac{\sqrt{2}}{\|\lambda\|}$. If $N_{2\lambda}$ is nontrivial, then we choose a vector $\tilde{X} \in \mathfrak{n}_{2\lambda}$ of norm $\frac{1}{\sqrt{2}\|\lambda\|}$. Write $\tilde{Y} := \theta(\tilde{X})$, where θ is the Cartan involution. For integral $\mu \in i\mathfrak{t}^*$, define $k(\mu) \in \mathbb{Z}$ by the relation

$$e^{\langle \mu, \mathfrak{t}(\tilde{X} - \tilde{Y}) \rangle} = e^{ik(\mu)t}$$

for all $t \in \mathbb{R}$. Recall the choice of $n_0 \in N_{2\lambda}$ above (3.10).

Theorem 3.20 (Index theorem, non-semisimple case). *Suppose that G has real rank 1 and that $\mathrm{rank}(G) = \mathrm{rank}(K)$.*

(a) *If $G \neq \mathrm{SU}(2n, 1)$, then*

$$(\tau_{n_0} + \tau_{n_0^{-1}})^{\Delta_{\mathfrak{p}} \otimes W} \circ \Psi_*(\mathrm{index}_{S, G}(D_W)) = 0.$$

If $G = \mathrm{SU}(2n, 1)$, then

$$\begin{aligned} & (\tau_{n_0} + \tau_{n_0^{-1}})^{\Delta_{\mathfrak{p}} \otimes W} \circ \Psi_*(\mathrm{index}_{S, G}(D_W)) = \\ & \|\lambda\| \frac{2^{3n-3} (2n+1)^n}{\mathrm{area}(S^{4n-1})} C_2(\Gamma) \varepsilon(R^+(G, T)) \dim(W). \end{aligned} \quad (3.18)$$

(b) We have

$$\tau_\lambda^{\Delta_{\mathfrak{p}} \otimes W} \circ \Psi_*(\text{index}_{S,G}(D_W)) = 0.$$

(c) If $\lambda + \rho_c$ is regular, then

$$\begin{aligned} \tau_{\text{rem}}^{\Delta_{\mathfrak{p}} \otimes W} \circ \Psi_*(\text{index}_{S,G}(D_W)) = \\ \frac{(-1)^{\dim(G/K)/2}}{2} l\varepsilon(\lambda + \rho_c) \sum_{w \in W_K} \det(w) k(w(\lambda + \rho_c)). \end{aligned} \quad (3.19)$$

If G/K is real hyperbolic space of dimension at least 4, then

$$\tau_{\text{rem}}^{\Delta_{\mathfrak{p}} \otimes W} \circ \Psi_*(\text{index}_{S,G}(D_W)) = 0.$$

Remark 3.21. Without the assumptions that $\lambda + \rho_c$ is regular or that G/K is real hyperbolic space of dimension at least 4, there is a less explicit version of part (c) of Theorem 3.20. See Propositions 5.6 and 6.5 in [6].

Theorems 3.17, 3.18 and 3.20 are proved in Subsection 6.6. The proof of Theorem 3.20 is taken from [6]. We give an independent, index-theoretic proof of Theorem 3.17 in Section 6. In the setting of Theorem 3.18, the computations from [6] do not apply directly, and we use results from [20] combined with arguments from the proof of Theorem 3.17.

4 Exponentially decaying kernels

4.1 The algebra $\mathcal{A}_S(E)$

Consider the setting of Subsection 2.1.

Proof of Lemma 2.5. If $\kappa \in \Gamma_{S,\text{fp}}(E \boxtimes E^*)$ and $\|\kappa\|_{a,P} = 0$ for all $a > 0$ and $P \in S$, then in particular $\|\kappa\|_{1,1} = 0$, because $1 \in S$. Hence $\kappa = 0$.

Using bases of the finite-dimensional spaces of order l differential operators in S , we can form a countable subset of the set of seminorms (2.3) inducing the same topology as the set of all those seminorms. So the space $\mathcal{A}_S(E)$ is a Fréchet space.

Let a_0 and C be as in Definition 2.1. Let $a \geq a_0$, and $P \in S$. Let $Q, R \in S$ be as in (2.2). Let $\kappa, \lambda \in \Gamma_{S,\text{fp}}^\infty(E \boxtimes E^*)$. Then for all $x, y \in X$,

$$(P(\kappa\lambda))(x, y) = \int_X (Q\kappa)(x, z)(R\lambda)(z, y) d\mu(z),$$

where the differential operators Q and R may be taken inside the integral because the integrand has compact support. The norm of the right hand side is smaller than or equal to

$$\|\kappa\|_{2a,Q}\|\lambda\|_{a,R} \int_X \int_X e^{-2ad(x,z)} e^{-ad(z,y)} d\mu(z). \quad (4.1)$$

And by the triangle inequality,

$$d(z,y) \geq d(x,y) - d(x,z).$$

So the right hand side of (4.1) is at most equal to

$$\|\kappa\|_{2a,Q}\|\lambda\|_{a,R} e^{-ad(x,y)} \int_X e^{-ad(x,z)} d\mu(z) \leq \|\kappa\|_{2a,Q}\|\lambda\|_{a,R} e^{-ad(x,y)} C.$$

Here we used that $a \geq a_0$. We conclude that

$$\|\kappa\lambda\|_{a,P,Q} \leq C\|\kappa\|_{2a,Q}\|\lambda\|_{a,R}.$$

If $a < a_0$, then

$$\|\kappa\lambda\|_{a,P} \leq \|\kappa\lambda\|_{a_0,P,Q} \leq C\|\kappa\|_{2a_0,Q}\|\lambda\|_{a_0,R}.$$

□

Proof of Lemma 2.6. If $\kappa \in \mathcal{A}_S(E)$, $h \in H$, $a > 0$ and $P \in S$, then by H -equivariance of P , and H -invariance of the metric on E ,

$$\|h \cdot \kappa\|_{a,P} = \|\kappa\|_{a,P}.$$

So the action by H on $\Gamma^\infty(E \boxtimes E^*)$ indeed restricts to an isometric action by H on $\mathcal{A}_S(E)$. Because the action is continuous and $\mathcal{A}_S(E)$ is Hausdorff, the fixed point set of every element $h \in H$ is closed. Hence so is the intersection $\mathcal{A}_S(E)^H$ of all these fixed-point sets. □

Proof of Lemma 2.10. We show that $\hat{B}(\mathbb{R})$ is a Fréchet algebra with respect to convolution, which implies the claim. The topology on $\hat{B}(\mathbb{R})$ is given by the countable subset of seminorms $\|\cdot\|_{a,k}$ for $a, k \in \mathbb{Z}_{\geq 0}$, so $\hat{B}(\mathbb{R})$ is a Fréchet space. Let $\psi_1, \psi_2 \in \hat{B}(\mathbb{R})$, $a > 0$ and $k \in \mathbb{Z}_{\geq 0}$. Then

$$\begin{aligned} \|\psi_1 * \psi_2\|_{a,k} &= \sup_{\xi \in \mathbb{R}} e^{a|\xi|} |(\psi_1^{(k)} * \psi_2)(\xi)| \\ &\leq \|\psi_1\|_{2a,k} \|\psi_2\|_{a,0} \sup_{\xi} e^{a|\xi|} \int_{\mathbb{R}} e^{-2a|\eta|} e^{-a|\xi-\eta|} d\eta \\ &\leq \|\psi_1\|_{2a,k} \|\psi_2\|_{a,0} \int_{\mathbb{R}} e^{-a|\eta|} d\eta, \end{aligned}$$

because $|\xi| - |\eta| \leq |\xi - \eta|$ for all $\xi, \eta \in \mathbb{R}$. So convolution is continuous. □

Lemma 4.1. *For all $t > 0$, the function*

$$g_t(x) = \frac{1 - e^{-tx^2}}{x}$$

is a multiplier of $B(\mathbb{R})$.

Proof. We have

$$\widehat{g}_t(\xi) = i\pi \left(\operatorname{erf} \left(\frac{\xi}{2\sqrt{t}} \right) - \operatorname{sgn}(\xi) \right).$$

This function decays faster than any exponential function. Let $a > 0$, and let $C_a > 0$ be such that for all $\xi \in \mathbb{R}$,

$$|\widehat{g}_t(\xi)| \leq C_a e^{-a|\xi|}.$$

Let $\varphi \in C_c^\infty(\mathbb{R})$ and $k \in \mathbb{Z}_{\geq 0}$. Analogously to the proof of Lemma 2.10, we find that

$$\|\varphi * \widehat{g}_t\|_{a,k} \leq \|\varphi\|_{2a,k} C_a \int_{\mathbb{R}} e^{-a|\eta|} d\eta.$$

It follows that convolution by \widehat{g}_t extends to a continuous operator on $\widehat{B}(\mathbb{R})$. Hence pointwise multiplication by g_t is a continuous operator on $B(\mathbb{R})$. \square

4.2 Suprema of Schwartz kernels

Until the end of this section, we use the notation and assumptions in Proposition 2.11.

We will prove and use extensions of Lemmas 3.5 and 3.8 in [13]. Versions of the following fact seem to be well-known, compare e.g. Proposition 5.9 in [11]. We include a proof here because we were not immediately able to find a concise proof in the literature.

Lemma 4.2. *For all $s > \dim(X)/2$, and all $f \in C_c(X)$, the operator $f(1 + D^2)^{-s}$ on $L^2(E)$ is Hilbert–Schmidt.*

Proof. Let $s > 0$. If f is real-valued, then f and $(1 + D^2)^{-s}$ are self-adjoint. Hence $f(1 + D^2)^{-s}$ is Hilbert–Schmidt if and only if $(1 + D^2)^{-s}f$ is. If f is not real-valued, then we reach the same conclusion by splitting f up into real and imaginary parts.

We will show that $(1 + D^2)^{-s}f$ is Hilbert–Schmidt. Using the Mellin transform, we write

$$(1 + D^2)^{-s}f = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{-t} e^{-tD^2} f dt.$$

Let κ_t be the Schwartz kernel of e^{-tD^2} . Because all structures are preserved by the cocompact action by H , the manifold M and the bundle E have bounded geometry. So there is an $a > 0$ such that $C_1 > 0$ such that for all $t \in (0, 1]$ and $x, y \in X$,

$$\|\kappa_t(x, y)\| \leq C_1 t^{-\dim(X)/2} e^{-ad(x,y)^2/t}.$$

See Proposition 5.2 in [11]. Because X has uniformly exponential growth, it follows that the Schwartz kernel of $e^{-tD^2} f$ is square-integrable, so $e^{-tD^2} f$ is Hilbert–Schmidt.

Now for all $t > 1$,

$$\|e^{-tD^2} f\|_{\mathcal{L}^2(L^2(E))} = \|e^{-(t-1)D^2} e^{-D^2} f\|_{\mathcal{L}^2(L^2(E))} \leq \|e^{-D^2} f\|_{\mathcal{L}^2(L^2(E))},$$

because $e^{-(t-1)D^2}$ has operator norm at most 1. And for all $t \leq 1$ and $x, y \in X$, we have $e^{-ad(x,y)^2/t} \leq e^{-ad(x,y)^2}$. So then

$$\|e^{-tD^2} f\|_{\mathcal{L}^2(L^2(E))}^2 \leq C_1^2 t^{-\dim(X)} \int_{X \times X} e^{-2ad(x,x')^2} |f(x')|^2 d\mu(x) d\mu(x').$$

We find that for all $s > \dim(X)/2$,

$$\begin{aligned} \|(1 + D^2)^{-s} f\|_{\mathcal{L}^2(L^2(E))} &\leq \\ C_1 \left(\int_{X \times X} e^{-ad(x,x')^2} f(x') d\mu(x) d\mu(x') \right)^{1/2} &\frac{1}{\Gamma(s)} \int_0^1 t^{s-\dim(X)/2-1} e^{-t} dt \\ &+ \|e^{-D^2} f\|_{\mathcal{L}^2(L^2(E))} \int_1^\infty t^{s-1} e^{-t} dt. \end{aligned}$$

Both terms on the right converge. \square

On a relatively compact open subset of a manifold, Sobolev norms of sections of a vector bundles can be defined unambiguously, up to equivalence. We denote the resulting L^2 -Sobolev space of order k by W^k .

Lemma 4.3. *Let X and Y be smooth manifolds, and $\varphi: Y \rightarrow X$ smooth and proper. Let $E \rightarrow X$ be a vector bundle. Let $U \subset X$ be open and relatively compact. For $s \in \Gamma^\infty(E|_U)$, define $\varphi^* s \in \Gamma^\infty((\varphi^* E)|_{\varphi^{-1}(U)})$ by*

$$\varphi^* s(y) = s(\varphi(y)),$$

for $y \in \varphi^{-1}(U)$. For all $k \in \mathbb{Z}_{\geq 0}$, this defines a bounded map

$$\varphi^*: W^k(E|_U) \rightarrow W^k((\varphi^* E)|_{\varphi^{-1}(U)}).$$

Proof. By relative compactness of U , the statement reduces to the case where U admits local coordinates x^1, \dots, x^m for X and a trivialisation of E . Then with respect to local coordinates y^1, \dots, y^n for Y in an open subset $V \subset \varphi^{-1}(U)$, the chain rule implies that for all multi-indices $\alpha \in \mathbb{Z}_{\geq 0}^n$, there are functions $F_\alpha^\beta \in C^\infty(V)$ (defined in terms of φ and its partial derivatives) such that for all $s \in \Gamma^\infty(E|_U) = C^\infty(U) \otimes \mathbb{C}^r$,

$$\frac{\partial^\alpha(\varphi^*s|_V)}{\partial y^\alpha} = \sum_{\beta \in \mathbb{Z}_{\geq 0}^n; |\beta| \leq |\alpha|} F_\alpha^\beta \frac{\partial^\beta s}{\partial x^\beta} \circ \varphi|_V.$$

Because φ is proper, the set $\varphi^{-1}(U)$ is relatively compact. So it can be covered by finitely many coordinate neighbourhoods V , and the functions F_α^β on these sets are bounded. The Jacobian of φ is bounded on $\varphi^{-1}(U)$, hence the claim follows. \square

Fix integers $p > \dim(X)/2$ and $l \geq 0$. Let $\mathcal{B}_l(L^2(E))^H$ be the space of $T \in \mathcal{B}(L^2(E))^H$ such that for all $j, k \in \mathbb{Z}_{\geq 0}$ with $j+k \leq \dim(X) + 2p + l + 1$, the composition $D^j T D^k$ defines a bounded operator on $L^2(E)$. For $T \in \mathcal{B}_l(L^2(E))^H$ and $r \leq \dim(X) + 2p + l + 1$, we write

$$\|T\|_r := \max \left\{ \|D^j T D^k\|_{\mathcal{B}(L^2(E))}; j, k \in \mathbb{Z}_{\geq 0}, j+k \leq r \right\}.$$

Let S_l be a finite-dimensional vector space of $H \times H$ -equivariant differential operators on $E \boxtimes E^*$ of orders at most l . Let $\varphi: Y \rightarrow X$ be as in Definition 2.7.

Lemma 4.4. *There is a $C > 0$ such that for all $T \in \mathcal{B}_l(L^2(E))^H$ and $P \in S_l$, the section $P(\varphi^* \kappa_T)$ is continuous and bounded, with sup-norm at most $C \|T\|_{\dim(X) + 2p + l + 1}$.*

Proof. We generalise some arguments in the proof of Lemma 3.5 in [13].

Let ε be the grading operator on E , equal to 1 on the even part of E and -1 on the odd part. We write D for the operator on $\Gamma^\infty(E^*)$ corresponding to D via the identification $E^* \cong E$ defined by the metric. Then the operator

$$D \otimes 1 + \varepsilon \otimes D$$

on $\Gamma^\infty(E \boxtimes E^*)$ is elliptic.

Let $f \in C_c(X)$, $n \leq \dim(X) + 1$ and $T \in \mathcal{B}_l(L^2(E))^H$. The operator $f(1 + D^2)^{-p}$ is Hilbert–Schmidt by Lemma 4.2, so

$$\begin{aligned} \|f(D \otimes 1 + \varepsilon \otimes D)^n \kappa_T\|_{L^2(E \boxtimes E^*)} &\leq \sum_{j=0}^n \binom{n}{j} \|f D^j T D^{n-j}\|_{\mathcal{L}^2(L^2(E))} \\ &\leq \|f(1 + D^2)^{-p}\|_{\mathcal{L}^2(L^2(E))} \sum_{j=0}^n \binom{n}{j} \|(1 + D^2)^p D^j T D^{n-j}\|_{\mathcal{B}(L^2(E))}. \end{aligned} \quad (4.2)$$

There is a constant $C_1 > 0$ (depending on f , D and p) such that the right hand side of (4.2) is equal to or less than $C_1 \|T\|_{2p+n}$. (In particular, the section $f(D \otimes 1 + \varepsilon \otimes D)^n \kappa_T$ is indeed square-integrable.)

Now let $U \subset X$ be open and relatively compact. Let $f \in C_c(X)$ be such that $f|_U \equiv 1$. Then by a Sobolev embedding theorem, relative compactness of U , Lemma 4.3 and finite-dimensionality of S_l , there are constants $C_2, C_3 > 0$ (depending on U , D and S_l), such that for all $P \in S_l$, and all $\kappa \in \Gamma^\infty(E \boxtimes E^*)$,

$$\begin{aligned} \sup_{x, x' \in U} \|P(\varphi^* \kappa)(x, x')\| &\leq C_2 \|\varphi^* \kappa\|_{W^{\dim(X)+l+1}(\varphi^* E|_{\varphi^{-1}(U)} \boxtimes \varphi^* E^*|_{\varphi^{-1}(U)})} \\ &\leq C_3 \|\kappa\|_{W^{\dim(X)+l+1}(E|_U \boxtimes E^*|_U)}. \end{aligned}$$

By ellipticity of $D \otimes 1 + \varepsilon \otimes D$, the right hand side is smaller than or equal to

$$C_4 \sup_{n \leq \dim(X)+l+1} \|(D \otimes 1 + \varepsilon \otimes D)^n \kappa\|_{L^2(E|_U \boxtimes E^*|_U)}.$$

for some constant $C_4 > 0$.

In particular, we have for all $P \in S_l$,

$$\begin{aligned} \sup_{x, x' \in U} \|P(\varphi^* \kappa_T)(x, x')\| &\leq C_4 \sup_{n \leq \dim(X)+l+1} \|(D \otimes 1 + \varepsilon \otimes D)^n \kappa_T\|_{L^2(E|_U \boxtimes E^*|_U)} \\ &\leq C_4 \sup_{n \leq \dim(X)+l+1} \|f(D \otimes 1 + \varepsilon \otimes D)^n \kappa_T\|_{L^2(E \boxtimes E^*)} \\ &\leq C_1 C_4 \|T\|_{\dim(X)+2p+l+1}. \end{aligned} \quad (4.3)$$

Now choose U (still relatively compact) such that $H \cdot U = X$. Then for all $P \in S_l$, the facts that P is $H \times H$ -equivariant and φ is H -equivariant imply that

$$\begin{aligned} \sup_{x, x' \in X} \|P(\varphi^* \kappa_T)(x, x')\| &= \sup_{x, x' \in U, h, h' \in H} \|P(\varphi^* \kappa_T)(h^{-1}x, h'^{-1}x')\| \\ &= \sup_{x, x' \in U, h, h' \in H} \|P(\varphi^* \kappa_{h \circ T \circ h'})(x, x')\|. \end{aligned} \quad (4.4)$$

By (4.3), the right hand side is smaller than or equal to

$$C_1 C_2 \|h \circ T \circ h'^{-1}\|_{\dim(X)+2p+l+1} = C_1 C_2 \|T\|_{\dim(X)+2p+l+1}.$$

Here we used H -equivariance of D , and the fact that H acts unitarily on $L^2(E)$. \square

4.3 Decay behaviour of Schwartz kernels

Recall the definition of the Fréchet algebra $B(\mathbb{R})$ in Subsection 2.2. For $f \in B(\mathbb{R})$, $n \in \mathbb{Z}_{\geq 0}$ and $s \geq 0$, write

$$F_{f,n}(s) := \sum_{k=0}^n \int_{\mathbb{R} \setminus [-s,s]} |\hat{f}^{(k)}(\xi)| d\xi.$$

Let c_D be the propagation speed of D , i.e. the maximum of the norm of the principal symbol of D on unit cotangent vectors. (This is finite, and indeed a maximum, because D commutes with the cocompact action by H .) The following estimate is an extension of Lemma 3.8 in [13]. Recall that we fixed an integer $p > \dim(X)/2$.

Proposition 4.5. *Let $n := \dim(X) + 2p + l + 1$, and let $f \in B(\mathbb{R})$.*

- (a) *The section $P(\varphi^* \kappa_{f(D)})$ is continuous and bounded for all $P \in S_l$.*
- (b) *Let A and B be as in (2.6). For all $\mu > 1$, there is a $C > 0$ such that for all $P \in S_l$ and all $y, y' \in Y$,*

$$\|P(\varphi^* \kappa_{f(D)})(y, y')\| \leq C F_{f,n} \left(\frac{d(y, y') - AB}{A\mu c_D} \right). \quad (4.5)$$

Proof. The proof is a modification of the proof of Lemma 3.8 in [13].

The function $\hat{f}^{(n)}$ is a Schwartz function, hence so is the function $x \mapsto x^n f(x)$. In particular, the latter function is bounded. So $f(D) \in \mathcal{B}_l(L^2(E))^H$. Hence Lemma 4.4 implies part (a).

To prove part (b), let $r > 0$ and $\mu > 1$. Let $\varphi \in C^\infty(\mathbb{R})$ be such that $\varphi(\xi) = 1$ if $|\xi| \geq 1$, and $\varphi(\xi) = 0$ if $|\xi| \leq 1/\mu$. Define $\varphi_r \in C^\infty(\mathbb{R})$ by $\varphi_r(\xi) := \varphi(\xi c_D/r)$. Let g_r be the inverse Fourier transform of $\varphi_r \hat{f}$.

As in the proof of Lemma 3.8 in [13], we have for all $j, k \in \mathbb{Z}_{\geq 0}$ with $j + k \leq n$,

$$\|D^j g_r(D) D^k\|_{\mathcal{B}(L^2(E))} \leq \psi(r) F_{f,n}(r/\mu c_D),$$

where

$$\begin{aligned}\psi(r) &:= \frac{1}{2\pi} \max \left\{ \binom{n}{j} \|\varphi_r^{(j)}\|_\infty; j = 0, \dots, n \right\} \\ &= \frac{1}{2\pi} \max \left\{ \binom{n}{j} \left(\frac{c_D}{r}\right)^j; j = 0, \dots, n \right\}.\end{aligned}$$

So ψ is decreasing in r . We find that $g_r(D) \in \mathcal{B}_l(L^2(E))^H$, and Lemma 4.4 implies that there is a $C_1 > 0$ such that for all $r > 0$ and all $P \in S_l$, the section $P(\kappa_{g_r(D)})$ is continuous and bounded, with sup-norm

$$\|P(\varphi^* \kappa_{g_r(D)})\|_\infty \leq C_1 \psi(r) F_{f,n}(r/\mu c_D). \quad (4.6)$$

Using finite-propagation arguments (specifically, Propositions 10.3.1 and 10.3.5 in [19] and their proofs) and the fact that $\hat{g}_r(\xi) = \hat{f}(\xi)$ if $|\xi| \geq r/c_D$, one shows that for all $s, s' \in \Gamma_c^\infty(E)$ whose supports are at least a distance r apart,

$$(s, g_r(D)s')_{L^2(E)} = (s, f(D)s')_{L^2(E)}.$$

So for all $x, x' \in X$ with $d_X(x, x') \geq r$, we have $\kappa_{g_r(D)}(x, x') = \kappa_{f(D)}(x, x')$. If $y, y' \in Y$ and $d_Y(y, y') \geq A(r + B)$, then $d_X(\varphi(\varphi(y)), \varphi(\varphi(y'))) \geq r$, so

$$\varphi^* \kappa_{g_r(D)}(y, y') = \varphi^* \kappa_{f(D)}(y, y').$$

Hence for all $y, y' \in Y$ with $d_Y(y, y') > A(r + B)$, we have

$$P(\varphi^* \kappa_{f(D)})(y, y') = P(\varphi^* \kappa_{g_r(D)})(y, y'). \quad (4.7)$$

Now let $y, y' \in X$, and suppose that $d_Y(y, y') > A(B + 1)$. Then (4.6) and (4.7) imply that for all $r > 1$ such that $A(r + B) < d_Y(y, y')$, and all $P \in S_l$,

$$\begin{aligned}\|P(\varphi^* \kappa_{f(D)})(y, y')\| &= \|P(\varphi^* \kappa_{g_r(D)})(y, y')\| \\ &\leq C_1 \psi(r) F_{f,n}(r/\mu c_D) \leq C_1 \psi(1) F_{f,n}(r/\mu c_D).\end{aligned}$$

Here we used the fact that ψ is decreasing. Because this inequality holds for all $r \in (1, d_Y(y, y')/A - B)$, it extends by continuity to

$$\|P(\varphi^* \kappa_{f(D)})(y, y')\| \leq C_1 \psi(1) F_{f,n} \left(\frac{d(y, y') - AB}{A\mu c_D} \right), \quad (4.8)$$

for all $y, y' \in X$ with $d_Y(y, y') > A(B + 1)$.

To obtain an estimate for $y, y' \in Y$ with $d_Y(y, y') \leq A(B+1)$, we first note that the function

$$(y, y') \mapsto \sup_{P \in S_l} \|P(\varphi^* \kappa_{f(D)})(y, y')\| \quad (4.9)$$

is invariant under the diagonal action by H , because D and φ are H -equivariant and P is $H \times H$ equivariant. Because X/H is compact, and φ is H -equivariant and proper, the quotient Y/H is compact as well. Hence so is the quotient of the set $\{(y, y') \in Y \times Y; d_Y(y, y') \leq A(B+1)\}$ by H . So the H -invariant function (4.9) is bounded on the latter set. Together with the estimate (4.8) for the case where $d(x, y) > A(B+1)$, and the fact that $F_{f,n}$ is a decreasing function, this implies part (b). \square

Proof of Proposition 2.11. Let $f \in B(\mathbb{R})$. Then for all $a, s > 0$, and $n \in \mathbb{Z}_{\geq 0}$,

$$F_{f,n}(s) \leq \frac{2e^{-as}}{a} \sum_{k=0}^n \|f\|_{a,k}. \quad (4.10)$$

By (4.10), Proposition 4.5 (with $\mu = 2$) implies that there are $n \in \mathbb{Z}_{\geq 0}$ and $C > 0$ such that for all $P \in S_l$, and all $a, a' > 0$,

$$\begin{aligned} \|\kappa_{f(D)}\|_{a,P} &\leq C \sup_{y,y' \in Y} e^{ad_Y(y,y')} F_{f,n} \left(\frac{d_Y(y, y') - AB}{2Ac_D} \right) \\ &\leq \frac{2C}{a'} e^{a'B/2c_D} \sup_{y,y' \in Y} e^{(a-a'/2Ac_D)d_Y(y,y')} \sum_{k=0}^n \|f\|_{a',k}. \end{aligned}$$

For $a' = 2Ac_D a$, the right hand side equals

$$\frac{C}{Ac_D a} e^{aAB} \sum_{k=0}^n \|f\|_{2Ac_D a, k}.$$

So the map $f \mapsto \kappa_{f(D)}$ is continuous. It is clearly an injective $*$ -homomorphism. \square

Proof of Theorem 2.12. The identity function $x \mapsto x$ is clearly a multiplier of $B(\mathbb{R})$, so $D \in \mathcal{M}(B(D))$. By Lemma 4.1, the operator

$$R_t := \frac{1 - e^{-tD^2}}{D^2} D \quad (4.11)$$

lies in $\mathcal{M}(B(D))$ as well. And

$$1 - DR_t = 1 - R_t D = e^{-tD^2},$$

which lies in $B(D)$. Using the pseudo-inverse R_t of D , one can show that the image of $[D] \in K_1(\mathcal{M}(B(D))/B(D))$ under (2.13) is (2.14), see page 356 of [15]. \square

Remark 4.6. We use the algebra $B(D)$ to define the higher index of D , because in general the operator (4.11) (or D itself, depending on S) may not be a multiplier of the larger algebra $\mathcal{A}_S(E)$.

5 Traces on a Schwartz algebra

We consider the setting described at the start of Section 3, and initially do not assume that G has real rank 1.

5.1 Schwartz spaces

Proof of Lemma 3.3. Let a_0 and C be as in Definition 2.1, for $X = G$. Let $f_1, f_2 \in C_c^\infty(G)$, $a \geq a_0$ and $P_1, P_2 \in \mathcal{U}(\mathfrak{g})$. Then, as in the proof of Lemma 2.5,

$$\|f_1 * f_2\|_{a, P_1, P_2} = \|L(P_1)f_1 * R(P_2)f_2\|_{a, 1} \leq C\|f_1\|_{2a, P_1, 1}\|f_2\|_{a, 1, P_2}.$$

If $a < a_0$, then

$$\|f_1 * f_2\|_{a, P_1, P_2} \leq \|f_1 * f_2\|_{a_0, P_1, P_2} \leq C\|f_1\|_{2a_0, P_1, 1}\|f_2\|_{a_0, 1, P_2}.$$

So $\mathcal{S}(G)$ is closed under convolution, and convolution is a continuous operation on $\mathcal{S}(G)$. \square

Proof of Lemma 3.4. The following facts can be verified directly from the definitions.

1. For all $\kappa \in \Gamma^\infty(E \boxtimes E^*)^G$, the relation (3.3) indeed yields a well-defined function $\Psi(\kappa) \in (C^\infty(G) \otimes \text{End}(V))^{K \times K}$.
2. The map Ψ is linear.
3. For $f \in (C^\infty(G) \otimes \text{End}(V))^{K \times K}$, define $\kappa_f \in \Gamma^\infty(E \boxtimes E^*)$ by

$$\kappa_f(gK, g'K)[g, v] = [g, f(g^{-1}g')v], \quad (5.1)$$

for $g, g' \in G$ and $v \in V$. This yields a well defined section $\kappa_f \in \Gamma^\infty(E \boxtimes E^*)^G$.

4. The map $f \mapsto \kappa_f$ from the previous point is a two-sided inverse of Ψ .

5. The map Ψ restricts to a bijection from $\Gamma_{\text{fp}}^\infty(E \boxtimes E^*)^G$ to $(C_c^\infty(G) \otimes \text{End}(V))^{K \times K}$.
6. For all $\kappa, \lambda \in \Gamma_{\text{fp}}^\infty(E \boxtimes E^*)^G$, we have $\Psi(\kappa\lambda) = \Psi(\kappa)\Psi(\lambda)$.
7. For all $f \in (C^\infty(G) \otimes \text{End}(V))^{K \times K}$, all $P_1, P_2 \in \mathcal{U}(\mathfrak{g})$ and all $g, g' \in G$,

$$(R(P_1) \otimes R(P_2)q^* \kappa_f)(g, g') = (L(P_1)R(P_2)f)(g^{-1}g').$$

By left G -invariance of the Riemannian distance on G , the last point implies that for all $a > 0$ and $P_1, P_2 \in \mathcal{U}(\mathfrak{g})$,

$$\|\kappa_f\|_{a, R(P_1) \otimes R(P_2)} = \|f\|_{a, P_1, P_2}.$$

So Ψ indeed defines an isometric algebra isomorphism

$$\Psi: \mathcal{A}_{S, q}(E)^G \xrightarrow{\cong} (\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}.$$

□

We will use a relation between $\mathcal{S}(G)$ and Harish-Chandra's L^p -Schwartz spaces. Let Ξ be the matrix coefficient for a unit vector in the representation induced from the trivial representation of a minimal parabolic. Fix $p > 0$. For $f \in C_c^\infty(G)$, $a > 0$ and $P_1, P_2 \in \mathcal{U}(\mathfrak{g})$, we write

$$\|f\|_{\mathcal{C}^p; a, P_1, P_2} := \sup_{g \in G} (1 + d_{G/K}(eK, gK))^a \Xi(x)^{-2/p} |(L(P_1)R(P_2)f)(g)|.$$

Let $\mathcal{C}^p(G)$ be the completion of $C_c^\infty(G)$ in the seminorms $\|\cdot\|_{\mathcal{C}^p; a, P, Q}$.

Remark 5.1. The space $\mathcal{C}^p(G)$ is the L^p -version of Harish-Chandra's Schwartz space. See e.g. pages 161–162 of [6]. For $p = 1$, this reduces to the definition on page 34 of [37]; for equivalence of the seminorms used, see Proposition 7.15 and (7.49)–(7.51) in [25]. The space $\mathcal{C}_\varepsilon^1(G)$ on page 98 of [37] equals $\mathcal{C}^{2/(2+\varepsilon)}(G)$.

Lemma 5.2. *The space $\mathcal{S}(G)$ is contained in $\mathcal{C}^p(G)$ for all $p > 0$, and the inclusion map is continuous.*

Proof. The weight functions in the definition of the seminorms on $\mathcal{C}^p(G)$ can be bounded by functions that increase exponentially in the distance to the identity element. □

Remark 5.3. At this point, let us comment on why we use the algebra $\mathcal{S}(G)$ in the construction of the higher index on G/K . The spaces $\mathcal{C}^1(G)$ and $\mathcal{C}^2(G)$ are in fact Fréchet algebras as well, see pages 49–51 of [37] for the former, and Proposition 12.16 in [25] for the latter. These have two disadvantages for our purposes, however.

1. While the geometric traces from Subsection 3.1 extend continuously to $(\mathcal{C}^2(G) \otimes \text{End}(V))^{K \times K}$, the spectral traces in Subsection 3.2 are only continuous on $(\mathcal{C}^p(G) \otimes \text{End}(V))^{K \times K}$ for $0 < p < 1$, hence not necessarily on the algebras $(\mathcal{C}^1(G) \otimes \text{End}(V))^{K \times K}$ and $(\mathcal{C}^2(G) \otimes \text{End}(V))^{K \times K}$. A related point is that Proposition 3.10, used to recover the classical index from the higher index in Corollary 3.14, is not true if $\mathcal{S}(G)$ is replaced by the algebra $\mathcal{C}^1(G)$ or $\mathcal{C}^2(G)$.
2. It was shown in Lemma 3.8 in [20] that the image of the idempotent in (2.14) under the map Ψ from Lemma 3.4 lies in the unitisation of $(\mathcal{C}^2(G) \otimes \text{End}(V))^{K \times K}$, but the Fourier transform techniques used there do not seem to generalise easily to an argument that it lies in $(\mathcal{C}^p(G) \otimes \text{End}(V))^{K \times K}$ for smaller p . The heat operators on the diagonal in (2.14) were shown to map into $(\mathcal{C}^p(G) \otimes \text{End}(V))^{K \times K}$ for all $p > 0$ in Proposition 2.4 in [6]. An extension of these arguments to the off-diagonal elements in (2.14) may be possible, but in view of the proof of Lemma 2.3 in [6], we expect this to be similar to our proof that they map into $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$.

So two key advantages of the algebra $\mathcal{S}(G)$ over $\mathcal{C}^1(G)$ and $\mathcal{C}^2(G)$ are that

1. the functionals in Subsection 3.2 are continuous traces on $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$, and
2. the idempotent in (2.14) maps into $(\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}$,

as shown in the present paper.

Furthermore, the algebra $\mathcal{A}_{\mathcal{S}, \varphi}(E)^H$, and hence the higher index of Definition 2.13, can be defined for more general spaces than G/K .

5.2 Geometric traces on $\mathcal{S}(G)$

In this subsection, we prove Propositions 3.5, 3.6 and 3.8 and Lemma 3.7. We start with a precise form of the statement that a conjugation-invariant functional is a trace.

Lemma 5.4. *Let H be a unimodular Lie group. Let $\mathcal{A}(H)$ be a locally convex convolution algebra of functions on H , containing $C_c^\infty(H)$ as a dense subalgebra. Suppose that convolution is continuous as a map from $\mathcal{A}(H) \times \mathcal{A}(H)$ to $\mathcal{A}(H)$. Let $\tau: \mathcal{A}(H) \rightarrow \mathbb{C}$ be a conjugation-invariant, continuous linear functional. Suppose that for all $f_1, f_2 \in C_c^\infty(H)$,*

$$\tau(f_1 * f_2) = \int_H \tau(x \mapsto f_1(xy^{-1})f_2(y)) dy. \quad (5.2)$$

Then τ is a trace on $\mathcal{A}(H)$.

Proof. Let $f_1, f_2 \in C_c^\infty(H)$. Then by (5.2) and conjugation-invariance of τ ,

$$\tau(f_1 * f_2) = \int_H \tau(x \mapsto f_1(y^{-1}x))f_2(y) dy.$$

By a substitution $y \mapsto y^{-1}x$ (where we use unimodularity of H) and (5.2), this equals

$$\int_H \tau(x \mapsto f_2(xy^{-1})f_1(y)) dy = \tau(f_2 * f_1).$$

So τ is a trace on $C_c^\infty(H)$. By continuity of convolution and of τ , this implies that τ is a trace on $\mathcal{A}(H)$. \square

Proposition 5.5. *If g is semisimple, then τ_g defines a continuous trace on $\mathcal{S}(G)$.*

Proof. If g is semisimple, then τ_g is a continuous functional on $\mathcal{C}^2(G)$; see Theorem 2.1 in [21], which is a consequence of Theorem 6 in [18]. By Lemma 5.2, τ_g restricts to a continuous functional on $\mathcal{S}(G)$. It is clearly conjugation-invariant, and it has the property (5.2) by the Fubini-Tonelli theorem. Convolution is continuous on $\mathcal{S}(G)$ by Lemma 3.3. So the claim follows by Lemma 5.4. \square

Proposition 5.5 implies Proposition 3.5.

Proof of Proposition 3.6. By Theorem 3.5 in [35], $\Phi_{P,g}$ is a continuous cyclic cocycle on $\mathcal{C}^2(G)$. So the claim follows from Lemma 5.2. \square

For the rest of this section, we assume that G has real rank 1. It is noted on page 299 of [30] that $\tau_{n_0} + \tau_{n_0^{-1}}$ is a continuous linear functional on $\mathcal{C}^2(G)$. In fact, the two terms separately define continuous traces on $\mathcal{S}(G)$.

Lemma 5.6. *For all $n \in N_{2\lambda}$, the orbital integral τ_n defines a continuous trace on $\mathcal{S}(G)$.*

Proof. We first show that τ_n is a continuous linear functional on $\mathcal{S}(G)$. Note that $N < G_n$, because $n \in N_{2\lambda}$. So for all $f \in \mathcal{S}(G)$, (3.6) implies that

$$|\tau_n(f)| \leq \int_{G/N} |f(xnx^{-1})| d(xN) = \int_{KA} |f(kana^{-1}k^{-1})| a^{2\rho_N} dk da.$$

For all $c > 0$, this is smaller than or equal to

$$\|f\|_{c,0} \int_{KA} e^{-cd(e,kana^{-1}k^{-1})} a^{2\rho_N} da \leq \|f\|_{c,0} e^{c \operatorname{diam}(K)} \int_A e^{-cd(e,ana^{-1})} a^{2\rho_N} da \quad (5.3)$$

by the triangle inequality. If $n = \exp(Y)$ for $Y \in \mathfrak{n}_{2\lambda}$, then the distance from e to ana^{-1} in the abelian group $N_{2\lambda}$ is

$$d_{N_{2\lambda}}(1, \exp(\operatorname{Ad}(a)Y)) = \|\operatorname{Ad}(a)Y\| = a^{2\lambda} \|Y\|.$$

This is the same as the distance from e to ana^{-1} in G , because $t \mapsto \exp(tY)$ is a geodesic. So the right hand side of (5.3) is smaller than or equal to

$$\|f\|_{c,0} e^{c \operatorname{diam}(K)} \|Y\| \int_A a^{-2c\lambda} a^{2\rho_N} da.$$

By (3.5), the integral converges if $c > \dim(\mathfrak{n}_\lambda)/2 + \dim(\mathfrak{n}_{2\lambda})$. So τ_n is indeed continuous.

It is immediate that τ_n is conjugation-invariant. And it satisfies (5.2) for compactly supported functions, by the Fubini-Tonelli theorem. So Lemma 5.4 implies the claim. \square

Lemma 5.6 implies Lemma 3.7.

Proposition 5.7. *The integral (3.7) converges for all $f \in \mathcal{S}(G)$, and defines a continuous trace on this algebra.*

Proof. The integral defines a continuous functional on $\mathcal{C}^2(G)$, as noted on page 299 of [30]. So by Lemma 5.2, it is a continuous functional on $\mathcal{S}(G)$.

For the trace property, we use Lemma 5.4. Let us show that τ_λ is conjugation-invariant.¹ Because the action by AN on G by right multiplication is proper and cocompact, there exists a function $\varphi \in C_c(G)$ such that for all $x \in G$,

$$\int_{AN} \varphi(xan) a^{4\rho_N} da dn = 1. \quad (5.4)$$

¹This seems to be known; the functional τ_λ is called an ‘invariant term’ in [6] (and denoted by T_1 there). We include a proof here because we could not immediately find a reference.

The group AN normalises N , and for all $f \in C_c(G)$, and all $a \in A$ and $n' \in N$,

$$\int_N f(an'n(an')^{-1}) dn = a^{2\rho_N} \int_N f(n) dn,$$

where $a^{2\rho_N}$ is a Jacobian. It follows that for any function φ as in (5.4), and all $f \in C_c^\infty(G)$,

$$\begin{aligned} \tau_\lambda(f) &= \int_N \int_{KAN} \varphi(kan') f(knk^{-1}) a^{4\rho_N} dk da dn' dn \\ &= \int_N \int_{KAN} \varphi(kan') f(kan'n(kan')^{-1}) a^{2\rho_N} dk da dn' dn \quad (5.5) \\ &= \int_N \int_G \varphi(g) f(gng^{-1}) dg dn, \end{aligned}$$

by (3.6). For $h \in G$, let $f_h \in C_c^\infty(G)$ be given by $f_h(g) = f(hgh^{-1})$. Then by (5.5) and a substitution $hg \mapsto g$,

$$\tau_\lambda(f_h) = \int_N \int_G (L_h\varphi)(g) f(gng^{-1}) dg dn. \quad (5.6)$$

The function $L_h\varphi$ has the same property (5.4) as φ , so (5.6) equals $\tau_\lambda(f)$.

We have seen that τ_λ is continuous and conjugation-invariant. It satisfies (5.2) by the Fubini–Tonelli theorem, so it is a trace on $\mathcal{S}(G)$ by Lemma 5.4. \square

Proposition 5.7 implies Proposition 3.8.

5.3 Spectral traces on $\mathcal{S}(G)$

In this subsection, we prove Propositions 3.10 and 3.13 and Corollary 3.11.

Proposition 5.8. *Let $F \subset \hat{K}$ be a finite set, and let $\mathcal{C}_F^1(G) \subset \mathcal{C}^1(G)$ be the subspace of isotypical components corresponding to F for the right K -action. For all $f \in \mathcal{C}_F^1(G)$, the operator $R_d^\Gamma(f)$ is trace-class. The linear map*

$$R_d^\Gamma: \mathcal{C}_F^1(G) \rightarrow \mathcal{L}^1(L_d^2(\Gamma \backslash G)) \quad (5.7)$$

is continuous.

Proof. The first claim is Theorem 4.6 in [37].

Let $L_0^2(\Gamma \backslash G) \subset L_d^2(\Gamma \backslash G)$ be the G -invariant subspace of cusp forms. By Theorem 4.3 in [37], the representation R^Γ defines a linear map

$$R_0^\Gamma: \mathcal{C}^1(G) \rightarrow \mathcal{L}^1(L_0^2(\Gamma \backslash G)). \quad (5.8)$$

On pages 39–40 of [37], it is shown that $f \mapsto \text{Tr}(R_0^\Gamma(f))$ is a continuous functional on $\mathcal{C}^1(G)$. In fact, the arguments given there can be strengthened slightly, to show that (5.8) is continuous. Indeed, apart from replacing the left quasiregular representation used in [37] by the right quasiregular representation we use here, the only changes that need to be made are that in the second-last set of displayed equations on page 40 of [37], one should take absolute values of $L_{G/\Gamma}^0(\alpha)$, $(U(\alpha)e_{U,i}, e_{U,i})$ and the integral over G in the third line. Then the bottom equation on page 40 of [37] becomes $\text{tr}(|L_{G/\Gamma}^0(\alpha)|) \leq C\|\Delta_L^{2N}\alpha\|_1$, and continuity of (5.8) follows.

Let $L_0^2(\Gamma \backslash G)^\perp \subset L_d^2(\Gamma \backslash G)$ be the orthogonal complement to $L_0^2(\Gamma \backslash G)$. It is shown on page 32 of [37] that the representation of G on this space is K -finite. Hence the image of the compression

$$(R_d^\Gamma)^\perp: \mathcal{C}_F^1(G) \rightarrow \mathcal{L}^1(L_0^2(\Gamma \backslash G)^\perp) \quad (5.9)$$

of (5.7) is finite-dimensional. Let n_F be its dimension. Because $\mathcal{C}^1(G)$ embeds continuously into $L^1(G)$, we find that for all $f \in \mathcal{C}_F^1(G)$,

$$\text{Tr}(|(R_d^\Gamma)^\perp(f)|) \leq n_F \|(R_d^\Gamma)^\perp(f)\|_{\mathcal{B}(L_d^2(\Gamma \backslash G)^\perp)} \leq n_F \|f\|_{L^1(G)},$$

and (5.9) is also continuous.

Continuity of (5.8) and (5.9) implies continuity of (5.7). \square

Proposition 5.8 and Lemma 5.2 imply Proposition 3.10.

Now consider a finite subset $F \subset \widehat{K \times K}$. Let $\mathcal{S}_F(G) \subset \mathcal{S}(G)$ be the subalgebra of functions in isotypical components corresponding to F , with respect to the left and right regular representations. Lemma 5.2 and Proposition 5.8 have the following consequence. This implies Corollary 3.11.

Corollary 5.9. *The map $f \mapsto \text{Tr}(R_d^\Gamma(f))$ is a continuous trace on $\mathcal{S}_F(G)$.*

Remark 5.10. To prove Corollary 5.9, we could have used the result from [37] that $f \mapsto \text{Tr}(R_0^\Gamma(f))$ is a continuous functional on $\mathcal{C}^1(G)$, rather than the stronger Proposition 5.8. We have included this proposition, because it is also used in Corollary 3.14.

The following result is a consequence of Theorem 8.4 in [37] and the theorem on page 299 of [30] and the comments below it.

Theorem 5.11 (Osborne–Warner). *For all $0 < p < 1$, the expression (3.10) defines a continuous linear functional τ_{rem} on $\mathcal{C}^p(G)$.*

Corollary 5.12. *The functional τ_{rem} from Theorem 5.11 restricts to a continuous trace on $\mathcal{S}(G)$.*

Proof. The restriction of τ_{rem} to $\mathcal{S}(G)$ is continuous by Lemma 3.3 and Theorem 5.11. It is a trace by Propositions 3.5 and 3.8 and Lemma 3.7. \square

Corollary 5.12 implies Proposition 3.13.

6 Contributions from semisimple elements

To prove Theorem 3.17, one can use the computations in Section 4.1 of [6]. Those computations are based on some nontrivial results from representation theory. We give an independent, index-theoretic proof here, based on a generalisation of the main result in [23]. We hope that this approach and some of the intermediate results may be of independent interest. The index-theoretic arguments also extend to a proof of Theorem 3.18.

For a locally compact group H acting properly and cocompactly on a manifold, we will use the standard equivariant index

$$\text{index}_H(D) \in K_0(C_r^*(H)) \tag{6.1}$$

of H -equivariant elliptic operators D . There are many equivalent definitions, see e.g. [7, 24, 39].

6.1 A holomorphic fixed-point formula

The first step is a version of the fixed-point formula in [21] for Dolbeault–Dirac operators. Let X be a complex manifold. Let G be a connected, real semisimple Lie group acting properly and cocompactly on X by holomorphic maps. Let $E \rightarrow M$ be a holomorphic, Hermitian, G -equivariant vector bundle. Let $\bar{\partial}_E$ be the Dolbeault operator on antiholomorphic differential forms twisted by E , and let $\bar{\partial}_E^*$ be its formal adjoint. Then we have the Dolbeault–Dirac operator $\bar{\partial}_E + \bar{\partial}_E^*$.

Let $g \in G$ be semisimple. Because G acts holomorphically on X , the fixed point set $X^g \subset X$ of g is a complex submanifold, and the normal bundle $N \rightarrow X^g$ is a complex vector bundle. (To be precise, the connected components of X^g are complex submanifolds of possibly different dimensions, so the rank of N may jump between these components.) Its complexification decomposes into holomorphic and antiholomorphic parts

as $N \otimes \mathbb{C} = N^{1,0} \oplus N^{0,1}$. Fix a Haar measure dz on the centraliser G_g . Let $\chi^g \in C_c^\infty(X^g)$ be such that for all $x \in X^g$,

$$\int_{G_g} \chi^g(zx) dz = 1.$$

(Here we use compactness of X^g/Z ; see Lemma 2.4 in [21].)

The restriction $E^g \rightarrow X^g$ of E is a holomorphic vector bundle over the complex manifold X^g , so we have the Dolbeault–Dirac operator $\bar{\partial}_{E^g} + \bar{\partial}_{E^g}^*$ on X^g . For every connected component X_j^g of X^g , we write $E_j^g := E|_{X_j^g}$. We similarly have the Dolbeault–Dirac operator $\bar{\partial}_{E_j^g} + \bar{\partial}_{E_j^g}^*$ on X_j^g .

Consider the equivariant indices

$$\begin{aligned} \text{index}_G(\bar{\partial}_E + \bar{\partial}_E^*) &\in K_0(C_r^*(G)); \\ \text{index}_{G_g}(\bar{\partial}_{E_j^g} + \bar{\partial}_{E_j^g}^*) &\in K_0(C_r^*(G_g)). \end{aligned}$$

The orbital integral trace τ_g is a well defined map on $K_0(C_r^*(G))$; see Section 2.1 of [21]. The von Neumann trace τ_e defines a map on $K_0(C_r^*(G_g))$. We write $\overline{g^{\mathbb{Z}}}$ for the closure of the cyclic subgroup of G generated by g .

Theorem 6.1. (a) *If g does not lie in a compact subgroup of G , then $\tau_g(\text{index}_G(\bar{\partial}_E + \bar{\partial}_E^*)) = 0$.*

(b) *If g lies in a compact subgroup of G , then*

$$\tau_g(\text{index}_G(\bar{\partial}_E + \bar{\partial}_E^*)) = \int_{T(X^g)} \chi^g \frac{\text{ch} \left(\left[\sigma_{\bar{\partial}_{E^g} + \bar{\partial}_{E^g}^*} \right] (g) \right) \text{Todd}(T(X^g) \otimes \mathbb{C})}{\text{ch} \left([\wedge N^{0,1}] (g) \right)}.$$

(c) *Suppose, furthermore, that on each connected component X_j^g of X^g , the group element g acts on every fibre of E_j^g as the same scalar $g|_{E_j^g}$. Suppose also that the bundle $N^{0,1}$ is trivial as a $\overline{g^{\mathbb{Z}}}$ -equivariant vector bundle. Then*

$$\tau_g(\text{index}_G(\bar{\partial}_E + \bar{\partial}_E^*)) = \sum_j \frac{g|_{E_j^g}}{\det_{\mathbb{C}}(1 - g|_{N^{0,1}})|_{X_j^g}} \tau_e(\text{index}_{G_g}(\bar{\partial}_{E_j^g} + \bar{\partial}_{E_j^g}^*)).$$

Proof. Part (a) follows from the second statement in Theorem 2.5 in [21]. (This in fact holds for all semisimple $g \in G$ not contained on compact subgroups.)

To prove part (b), we again apply Theorem 2.5 in [21]. This now yields

$$\tau_g(\text{index}_G(\bar{\partial}_E + \bar{\partial}_E^*)) = \int_{T(X^g)} \chi^g \frac{\text{ch} \left(\left[\sigma_{\bar{\partial}_E + \bar{\partial}_E^*} \right] (g) \right) \text{Todd}(T(X^g) \otimes \mathbb{C})}{\text{ch}([\Lambda N \otimes \mathbb{C}](g))}.$$

To be precise, in Theorem 2.5 in [21] it is stated that this equality holds for almost all elliptic elements g not contained in compact subgroups. The result applies to “almost all” such g because of the “finite Gaussian orbital integrals” condition. But this turns out to hold in general, see below Definition 4.3.3 in [8].

We use the decomposition

$$\left[\sigma_{\bar{\partial}_E + \bar{\partial}_E^*} |_{X^g} \right] (g) = [\Lambda(N^{0,1})^*] (g) \otimes \left[\sigma_{\bar{\partial}_{E^g} + \bar{\partial}_{E^g}^*} \right] (g) = [\Lambda N^{1,0}] (g) \otimes \left[\sigma_{\bar{\partial}_{E^g} + \bar{\partial}_{E^g}^*} \right] (g)$$

By multiplicativity of the Chern character,

$$\frac{\text{ch}([\Lambda N^{1,0}](g))}{\text{ch}([\Lambda N \otimes \mathbb{C}](g))} = \frac{1}{\text{ch}([\Lambda N^{0,1}](g))},$$

so (b) follows.

To deduce (c) from (b), we first note that the assumption on the way g acts on $E|_{X^g}$ implies that

$$\left[\sigma_{\bar{\partial}_{E^g} + \bar{\partial}_{E^g}^*} |_{X_j^g} \right] (g) = g|_{E_j^g} \left[\sigma_{\bar{\partial}_{E_j^g} + \bar{\partial}_{E_j^g}^*} \right],$$

for every connected component X_j^g of X^g . And if $N^{0,1}$ is trivial, then for all j , we have $N^{0,1}|_{X_j^g} = X_j^g \times W_j$ for some representation W_j of the compact abelian group $\overline{g^{\mathbb{Z}}}$. (The dimension of X_j^g , and hence of W_j , may depend on j .) Then

$$[\Lambda N^{0,1}|_{X_j^g}] = [X_j^g \times \mathbb{C}] \otimes [\Lambda W_j] \in K_{\overline{g^{\mathbb{Z}}}}^0(X_j^g) \cong K^0(X_j^g) \otimes R(\overline{g^{\mathbb{Z}}}).$$

Here ΛW_j should be interpreted as a graded representation, i.e.

$$[\Lambda W_j] = \left[\Lambda^{\text{even}} W_j \right] - \left[\Lambda^{\text{odd}} W_j \right] \in R(\overline{g^{\mathbb{Z}}}).$$

So

$$\text{ch}([\Lambda N^{0,1}](g)) = \text{tr} \left(g|_{\Lambda^{\text{even}} W_j} \right) - \text{tr} \left(g|_{\Lambda^{\text{odd}} W_j} \right) = \det_{\mathbb{C}}(1-g|_{W_j}) = \det_{\mathbb{C}}(1-g|_{N^{0,1}})|_{X_j^g}.$$

We find that

$$\tau_g(\text{index}_G(\bar{\partial}_E + \bar{\partial}_E^*)) = \sum_j \frac{\text{tr}(g|_E)}{\det_{\mathbb{C}}(1 - g|_{N^{0,1}})} \Big|_{X_j^g} \int_{T(X_j^g)} \chi^g|_{X_j^g} \text{ch} \left(\left[\sigma_{\bar{\partial}_{E_j^g} + \bar{\partial}_{E_j^g}^*} \right] \right) \text{Todd}(T(X_j^g) \otimes \mathbb{C}).$$

By Proposition 4.4 and Theorem 6.12 in [36], the integral on the right equals $\tau_e(\text{index}_{G_g}(\bar{\partial}_{E_j^g} + \bar{\partial}_{E_j^g}^*))$. \square

Example 6.2. If X and G are compact, and X^g is finite, then Theorem 6.1(c) reduces to

$$\text{index}_G(\bar{\partial}_E + \bar{\partial}_E^*)(g) = \sum_{x \in X^g} \frac{g|_{E_x}}{\det_{\mathbb{C}}(1 - g|_{T_x X^{0,1}})} = \sum_{x \in X^g} \frac{g|_{E_x}}{\det_{\mathbb{C}}(1 - g^{-1}|_{T_x X^{1,0}})}.$$

This is consistent with Atiyah and Bott's holomorphic fixed-point theorem, Theorem 4.12 in [3], because in the notation of that paper,

$$\text{index}_G(\bar{\partial}_E + \bar{\partial}_E^*)(g) = L(g^{-1}, g).$$

6.2 A fixed-point set

Until the end of Subsection 6.4, we suppose that $T < G$ is a compact Cartan subgroup, so $\text{rank}(G) = \text{rank}(K)$. We fix a positive root system $R^+(G, T)$ for (G, T) . This determines a G -invariant complex structure on G/T such that $T_{eT}^{1,0}(G/T) = \bigoplus_{\alpha \in R^+(G, T)} \mathfrak{g}_{\alpha}^{\mathbb{C}}$. Let $g \in T$. We will apply Theorem 6.1 to the space $X = G/T$, and we start by analysing the fixed-point set $(G/T)^g$.

Lemma 6.3. *For all $x \in G$ such that $xgx^{-1} \in T$, there is a $y \in N_G(T)$ such that $xgx^{-1} = ygy^{-1}$.*

Proof. Because $xTx^{-1} \subset G_{xgx^{-1}}$, we have two maximal tori T and xTx^{-1} in the connected component $(G_{xgx^{-1}})_0$. These are conjugate in $(G_{xgx^{-1}})_0$. (See e.g. Propositions 6.59 and 6.61 in [26], or use the facts that any two maximal compact subgroups of $(G_{xgx^{-1}})_0$ are conjugate, and so are any two maximal tori inside such a compact subgroup.) Let $h \in (G_{xgx^{-1}})_0$ be such that $hxTx^{-1}h^{-1} = T$. Then $y := hx$ has the desired properties. \square

Recall that we used the Weyl groups $W_K = N_K(T)/T$ and $W_{K_g} = N_{K_g}(T)/T$. Note that $G_g w$ is a right T -invariant subset of G for all $w \in N_G(T)$.

Lemma 6.4. *The fixed-point set $(G/T)^g$ decomposes as*

$$(G/T)^g = \coprod_{w \in W_{K_g} \setminus W_K} G_g w/T, \quad (6.2)$$

where the disjoint union is over representatives $w \in N_K(T)$ of classes in $W_{K_g} \setminus W_K$.

Proof. We first show that

$$(G/T)^g = G_g N_G(T)/T. \quad (6.3)$$

It is immediate that the right hand side is contained in the left hand side. Now let $xT \in (G/T)^g$. Then $x^{-1}gx \in T$. By Lemma 6.3, there is a $y \in N_G(T)$ such that $x^{-1}gx = y^{-1}gy$. So $xy^{-1} \in G_g$, and $xT = (xy^{-1})yT$ lies in the right hand side of (6.3).

Next, we show that

$$(G/T)^g = \coprod_{w \in N_{G_g}(T) \setminus N_G(T)} G_g w/T, \quad (6.4)$$

where the disjoint union is over a set of representatives $w \in N_G(T)$. Indeed, if $y \in N_{G_g}(T) \subset G_g$ and $w \in N_G(T)$, then $G_g y w/T = G_g w/T$. And if $w, w' \in N_G(T)$ and $G_g w/T = G_g w'/T$, then let $y := w'w^{-1} \in N_G(T)$. Then $G_g w/T = G_g y w'/T$, so there are $a \in G_g$ and $b \in T$ such that $awb = yw'$. Hence $a^{-1}y = bw'^{-1} \in T \subset G_g$. So $y \in G_g$, and therefore $y \in N_{G_g}(T)$. So (6.4) follows.

Finally, we have $N_G(T) = N_K(T)$ and $N_{G_g}(T) = N_{K_g}(T)$; see for example Lemma 6.15 in [22]. So the decomposition (6.2) follows from (6.4) and the equality

$$N_{K_g}(T) \setminus N_K(T) = (N_{K_g}(T)/T) \setminus (N_K(T)/T).$$

□

Example 6.5. If $g = e$, then Lemma 6.4 reduces to the equality $G/T = G/T$. At the other extreme, if the powers of g are dense in T , then Lemma 6.4 becomes $(G/T)^T = W_K$; this is Lemma 6.15 in [22].

6.3 A normal bundle

For $w \in N_G(T)$, let J_w be the G -invariant complex structure on G/T such that

$$T_{eT}^{1,0}(G/T) \cong \bigoplus_{\alpha \in R^+(G,T)} \mathfrak{g}_{w\alpha}^{\mathbb{C}}.$$

Then J_e is the complex structure we have considered so far.

Lemma 6.6. *Let $w \in N_G(T)$. The map $R_w: G/T \rightarrow G/T$, given by $R_w(xT) = xwT$ for $x \in G$, is a well-defined, G -equivariant diffeomorphism. It is holomorphic as a map*

$$R_w: (G/T, J_w) \rightarrow (G/T, J_e). \quad (6.5)$$

Proof. The first claim follows directly from the definitions. We show that (6.5) is holomorphic.

Note that $R_w = L_w \circ C_{w^{-1}}$, where L is the standard action by G on G/T , and $C_{w^{-1}} = L_{w^{-1}} \circ R_w$ is induced by conjugation by w^{-1} . The map L_w is holomorphic for any of the complex structures $J_{w'}$, because they are G -invariant. And

$$T_{eT}C_{w^{-1}} = \text{Ad}(w^{-1}): \mathfrak{g}/\mathfrak{t} \rightarrow \mathfrak{g}/\mathfrak{t}.$$

Because $\text{Ad}(w^{-1})\mathfrak{g}_{\alpha}^{\mathbb{C}} = \mathfrak{g}_{w^{-1}\alpha}^{\mathbb{C}}$, this map induces a complex linear map from $(T_{eT}(G/J), J_w)$ to $(T_{eT}(G/J), J_e)$. For general $x \in G$, we have $C_{w^{-1}} = L_{w^{-1}xw} \circ C_{w^{-1}} \circ L_{x^{-1}}$. So

$$T_{xT}C_{w^{-1}} = T_{eT}L_{w^{-1}xw} \circ T_{eT}C_{w^{-1}} \circ T_{xT}L_{x^{-1}}.$$

The three maps on the right hand side are complex-linear, hence so is the map on the left. We find that the map (6.5) is indeed holomorphic. \square

The complex structure J_w , for $w \in N_G(T)$, restricts to the complex sub-bundle $N \subset T(G/T)|_{(G/T)^g}$. We denote this restriction by J_w as well.

Lemma 6.7. *For every $w \in N_G(T)$, the map $R_w: G_g/T \rightarrow G_gw/T$ given by $R_w(xT) = xwT$ for $x \in G_g$, is a well-defined, G_g -equivariant diffeomorphism. The vector bundles*

$$\begin{aligned} (N|_{G_g/T}, J_w) &\rightarrow G_g/T; \\ R_w^*(N|_{G_gw/T}, J_e) &\rightarrow G_g/T \end{aligned} \quad (6.6)$$

are isomorphic as G_g -equivariant, holomorphic vector bundles.

Proof. Let $w \in N_G(T)$. The first claim follows from the first claim in Lemma 6.6.

We write $N(S)$ for the normal bundle of a submanifold $S \subset G/T$. Because the sets $G_g w/T$ are connected components of $(G/T)^g$, we have

$$TR_w(N|_{G_g/T}) = TR_w(N(G_g/T)) = N(R_w(G_g/T)) = N(G_g w/T) = N|_{G_g w/T}.$$

This fact, together with G -equivariance of R_w and the fact that the map (6.5) is holomorphic by Lemma 6.6, implies that the bundles (6.6) are equivariantly isomorphic as holomorphic vector bundles. \square

Lemma 6.8. *The bundle $N^{0,1} \rightarrow (G/T)^g$ is trivial as a G_g -vector bundle.*

Proof. Let $w \in N_K(T)$. We first prove that $(N^{0,1}|_{G_g/T}, J_w) \rightarrow G_g/T$ is G_g -equivariantly trivial. Indeed,

$$N|_{G_g/T} = G_g \times_T \mathfrak{g}/\mathfrak{g}_g,$$

where \mathfrak{g}_g is the Lie algebra of G_g . So

$$(N^{0,1}|_{G_g/T}, J_w) = G_g \times_T ((\mathfrak{g}/\mathfrak{g}_g)^{0,1}, J_w), \quad (6.7)$$

where

$$((\mathfrak{g}/\mathfrak{g}_g)^{0,1}, J_w) = \bigoplus_{\alpha \in R^+(G,T) \setminus R^+(G_g,T)} \mathfrak{g}_{-w\alpha}^{\mathbb{C}}. \quad (6.8)$$

The bundle (6.7) is isomorphic as a G_g -equivariant, complex vector bundle to $G_g/T \times ((\mathfrak{g}/\mathfrak{g}_g)^{0,1}, J_w)$ via the map given by $[x, Y] \mapsto (xY, \text{Ad}(x)Y)$, for $x \in G_g$ and $Y \in ((\mathfrak{g}/\mathfrak{g}_g)^{0,1}, J_w)$. So $(N^{0,1}|_{G_g/T}, J_w) \rightarrow G_g/T$ is equivariantly trivial as claimed.

The claim now follows from Lemmas 6.4 and 6.7. \square

As in Theorem 3.17, let ρ be half the sum of the positive roots in $R^+(G, T)$, and let ρ_g be half the sum of the positive roots in $R^+(G_g, T)$.

Lemma 6.9. *For all $w \in W_K$,*

$$\det_{\mathbb{C}}(1-g|_{N^{0,1}})|_{G_g w/T} = e^{w(\rho_g - \rho)}(g) \det(w)e^{\rho - \rho_g}(g) \prod_{\alpha \in R^+(G,T) \setminus R^+(G_g,T)} (1 - e^{-\alpha}(g)). \quad (6.9)$$

Proof. Let $w \in N_K(T)$. Then by (6.7) and (6.8),

$$\begin{aligned} \det_{\mathbb{C}}(1 - g|_{(N^{0,1}, J_w)})|_{G_g/T} &= \prod_{\alpha \in R^+(G,T) \setminus R^+(G_g,T)} (1 - e^{-w\alpha}(g)) \\ &= e^{w(\rho_g - \rho)}(g) e^{w(\rho - \rho_g)}(g) \prod_{\alpha \in R^+(G,T) \setminus R^+(G_g,T)} (1 - e^{-w\alpha}(g)). \end{aligned}$$

The right hand side with the first exponential factor removed is a ratio of Weyl denominators. This is alternating with respect to w . So the last expression equals the right hand side of (6.9). The claim now follows by Lemma 6.7. \square

6.4 A fixed-point formula on G/T

For an analytically integral $\nu \in \mathfrak{it}^*$, we write \mathbb{C}_ν for the complex numbers equipped with the unitary representation of T with weight ν .

Lemma 6.10. *Let $\mu \in \mathfrak{it}^*$ be integral, and let $w \in N_K(T)$ be such that $w \cdot \mu = \mu \circ \text{Ad}(w)^{-1}$ is integral. Then there is a G_g -equivariant isomorphism of complex line bundles*

$$R_w^*(G_g w/T \times_T \mathbb{C}_\mu) \cong G_g \times_T \mathbb{C}_{w \cdot \mu} \rightarrow G_g/T.$$

Proof. We have

$$R_w^*(G_g w/T \times_T \mathbb{C}_\mu) = \{(xT, [xwt, z]); x \in G_g, t \in T, z \in \mathbb{C}_\mu\}.$$

The map from the right hand side to $G_g \times_T \mathbb{C}_{w \cdot \mu}$ mapping $(xT, [xwt, z])$ in this set to $[x, e^\mu(t)z]$, is a well-defined, G_g -equivariant isomorphism of complex line bundles. \square

We consider a Dirac operator D_W of the form (3.14). Let $\lambda \in \mathfrak{it}^*$ be the highest weight of W with respect to the set of compact positive roots $R^+(K, T) \subset R^+(G, T)$. As in Theorem 3.17, let ρ_c be half the sum of the positive compact roots in $R^+(K, T)$. Let $\rho_n = \rho - \rho_c$. For analytically integral $\nu \in \mathfrak{it}^*$, consider the holomorphic line bundle $L_\nu := G \times_T \mathbb{C}_\nu \rightarrow G/T$. Let $d_\mu^{G_g}$ be as in (3.16).

Lemma 6.11. *The element $\lambda - \rho_n \in \mathfrak{it}^*$ is analytically integral. For all $w \in N_K(T)$,*

$$\tau_e(\text{index}_{G_g}(\bar{\partial}_{L_{\lambda - \rho_n}|_{G_g w/T}} + \bar{\partial}_{L_{\lambda - \rho_n}|_{G_g w/T}}^*)) = (-1)^{\dim(G_g/K_g)/2} d_{w(\lambda + \rho_g - \rho_n)}^{G_g}. \quad (6.10)$$

Proof. The element $\lambda - \rho_n \in \mathfrak{it}^*$ is analytically integral, because $\Delta_{\mathfrak{p}} \otimes W$ defines a representation of K with highest weight $\lambda + \rho_n$.

For $w \in N_K(T)$, let $J_w^{G_g}$ be the G_g -invariant complex structure on G_g/T such that

$$T_{eT}^{1,0}(G_g/T) = \bigoplus_{\alpha \in R^+(G_g, T)} \mathfrak{g}_{w\alpha}^{\mathbb{C}}.$$

Let $w \in N_K(T)$. Then Lemma 6.6 implies that the map R_w restricts to a G_g -equivariant, holomorphic diffeomorphism

$$R_w : (G_g/T, J_w^{G_g}) \rightarrow (G_g w/T, J_e^{G_g}).$$

And Lemma 6.10 implies that

$$R_w^*(L_{\lambda - \rho_n}|_{G_g w/T}) \cong L_{w(\lambda - \rho_n)}^{G_g/T} := G_g \times_T \mathbb{C}_{w(\lambda - \rho_n)} \rightarrow G_g/T$$

as G_g -equivariant, complex line bundles. We find that

$$\text{index}_{G_g}(\bar{\partial}_{L_{\lambda - \rho_n}|_{G_g w/T}} + \bar{\partial}_{L_{\lambda - \rho_n}|_{G_g w/T}}^*) = \text{index}_{G_g}(\bar{\partial}_{L_{w(\lambda - \rho_n)}^{G_g/T}} + \bar{\partial}_{L_{w(\lambda - \rho_n)}^{G_g/T}}^*), \quad (6.11)$$

where the Dolbeault operator on the right is defined with respect to $J_w^{G_g}$.

In the proof of Proposition 5.2 in [21], it is shown that the right hand side of (6.11) equals $(-1)^{\dim(G_g/K_g)/2} \text{D-Ind}_{K_g}^{G_g}[W_{w(\lambda - \rho_n + \rho_{n,g})}^{K_g}]$, where $\text{D-Ind}_{K_g}^{G_g}$ denotes Dirac induction, $\rho_{n,g}$ is half the sum of the noncompact roots in $R^+(G_g, T)$, and $W_{w(\lambda - \rho_n + \rho_{n,g})}^{K_g}$ is the irreducible representation of K_g with highest weight $w(\lambda - \rho_n + \rho_{n,g})$ with respect to the positive root system $wR^+(K, T) \cap R(K_g, T)$.

If $\lambda - \rho_n + \rho_g$ is regular for G_g , then Lemma 5.4 in [21] implies that

$$\text{D-Ind}_{K_g}^{G_g}[W_{w(\lambda - \rho_n + \rho_{n,g})}^{K_g}] = [\pi_{w(\lambda + \rho_g - \rho_n)}^{G_g}],$$

the K -theory class of the discrete series representation of G_g with Harish-Chandra parameter $w(\lambda + \rho_g - \rho_n)$. Hence (6.10) follows, because the value of τ_e on a K -theory class of a discrete series representation is the formal degree of that representation [4, 14, 27].

If $\lambda - \rho_n + \rho_g$ is singular for G_g , then $d_{w(\lambda + \rho_g - \rho_n)}^{G_g} = 0$. Then the Dirac operator whose G_g -equivariant index is $\text{D-Ind}_{K_g}^{G_g}[W_{w(\lambda - \rho_n + \rho_{n,g})}^{K_g}]$ has trivial kernel by Theorem 9.3 in [4]. And $\tau_e(\text{D-Ind}_{K_g}^{G_g}[W_{w(\lambda - \rho_n + \rho_{n,g})}^{K_g}])$ equals the L^2 -index of this Dirac operator (see Proposition 4.4 in [36]), which is defined in terms of the projection onto its kernel, and hence zero. So now the left hand side of (6.10) is zero as well, and (6.10) holds. \square

Theorem 6.12. *Let $g \in G$ be semisimple. Then $\tau_g(\text{index}_G(D_W)) = 0$ if g is hyperbolic. If $\text{rank}(G) = \text{rank}(K)$ and g is elliptic, then*

$$\tau_g(\text{index}_G(D_W)) = (-1)^{(\dim(G/K) + \dim(G_g/K_g))/2} \frac{\sum_{w \in W_{K_g} \setminus W_K} \det(w) e^{w(\lambda + \rho_c - \rho_g)}(g) d_{w(\lambda + \rho_g - \rho_n)}^{G_g}}{e^{\rho - \rho_g}(g) \prod_{\alpha \in R^+(G, T) \setminus R^+(G_g, T)} (1 - e^\alpha(g))}. \quad (6.12)$$

Proof. The case for hyperbolic g follows from Theorem 6.1(a).

Suppose that $\text{rank}(G) = \text{rank}(K)$ and that g is elliptic. The element $\lambda - \rho_n \in \mathfrak{it}^*$ is analytically integral as noted in Lemma 6.11. In the proof of Proposition 5.2 in [21], it is shown that

$$\text{index}_G(D_W) = (-1)^{\dim(G/K)/2} \text{index}_G(\bar{\partial}_{L_{\lambda - \rho_n}} + \bar{\partial}_{L_{\lambda - \rho_n}}^*).$$

(This is a place where we use the assumption that $\text{rank}(G) = \text{rank}(K)$.) By this equality and Lemmas 6.4, 6.8 and 6.9, Theorem 6.1(c) implies that

$$\tau_g(\text{index}_G(D_W)) = (-1)^{\dim(G/K)/2} \frac{\sum_{w \in W_{K_g} \setminus W_K} \det(w) e^{w(\rho - \rho_g)}(g) (\text{tr}(g|_{L_{\lambda - \rho_n}})|_{G_g w/T}) \tau_e(\text{index}_{G_g}(\bar{\partial}_{L_{\lambda - \rho_n}|_{G_g w/T}} + \bar{\partial}_{L_{\lambda - \rho_n}|_{G_g w/T}}^*))}{e^{\rho - \rho_g}(g) \prod_{\alpha \in R^+(G, T) \setminus R^+(G_g, T)} (1 - e^\alpha(g))}.$$

For all $w \in N_K(T)$, Lemma 6.10 implies that

$$\text{tr}(g|_{L_{\lambda - \rho_n}})|_{G_g w/T} = e^{w(\lambda - \rho_n)}(g).$$

So (6.12) follows by Lemma 6.11. \square

Example 6.13. If g is central, then Theorem 6.12 reduces to

$$\tau_g(\text{index}_G(D_W)) = e^{\lambda - \rho_n}(g) d_{\lambda + \rho_c}^G.$$

This is Proposition 4.2 in [6]. If $g = e$, then versions of this equality were obtained in [14, 27]. This case was used in the proof of Theorem 6.12.

Example 6.14. If the powers of g are dense in T , then $G_g = K_g = T$, and for all $w \in W_K$,

$$\begin{aligned} \text{tr}(g|_{L_{\lambda - \rho_n}})|_{G_g w/T} &= e^{w(\lambda - \rho_n)}(g); \\ \tau_e(\text{index}_{G_g}(\bar{\partial}_{L_{\lambda - \rho_n}|_{G_g w/T}} + \bar{\partial}_{L_{\lambda - \rho_n}|_{G_g w/T}}^*)) &= 1. \end{aligned}$$

So Theorem 6.12 reduces to

$$\tau_g(\text{index}_G(D_W)) = (-1)^{\dim(G/K)/2} \frac{\sum_{w \in W_K} \det(w) e^{w(\lambda + \rho_c)}(g)}{e^\rho(g) \prod_{\alpha \in R^+(G, T)} (1 - e^{-\alpha}(g))}.$$

This is Theorem 3.1(a) in [23].

Remark 6.15. If $\text{rank}(G) > \text{rank}(K)$, then Theorem 3.2(a) in [23] implies that the number $\tau_g(\text{index}_G(D_W))$ equals zero for all hyperbolic g and all elliptic g with powers dense in a maximal torus. Then also $\tau_e(\text{index}_G(D_W)) = 0$, as in the proof of Lemma 6.11. We expect that $\tau_g(\text{index}_G(D_W)) = 0$ for all semisimple g in this case, but we have not worked out the details. That would imply that all numbers in Theorem 3.17 equal zero if $\text{rank}(G) > \text{rank}(K)$.

6.5 Higher orbital integrals

We no longer assume that G has a compact Cartan subgroup. We saw in Theorem 6.12 that $\tau_g(\text{index}_G(D_W)) = 0$ if $\text{rank}(G) > \text{rank}(K)$. However, $\text{index}_G(D_W) \in K_0(C_r^*(G))$ is nonzero, and one way to detect its nonvanishing is to pair it with the higher orbital integrals from [35]. We can compute the resulting pairings using the results in [20] to generalise Theorem 6.12. We use notation and assumptions as in Theorem 3.18. As in [20] and in Theorem 3.18, we assume that the map $\widetilde{\text{Ad}}: \widetilde{K} \rightarrow \text{Spin}(\mathfrak{p})$ from Subsection 3.4 descends to K .

Corollary 6.16. *If P is not a maximal cuspidal parabolic subgroup, or $g \in M$ is hyperbolic, then*

$$\langle \Phi_{P,g}, \text{index}_G(D_W) \rangle = 0.$$

If P is a maximal cuspidal parabolic subgroup and $g \in T \subset M$, then

$$\begin{aligned} \langle \Phi_{P,g}, \text{index}_G(D_W) \rangle &= (-1)^{(\dim(M/(K \cap M)) + \dim(M_g/(K \cap M_g)))/2} \\ &\sum_{U \in \widehat{K \cap M}} m_U \frac{\sum_{w \in W_{K \cap M_g} \setminus W_{K \cap M}} \det(w) e^{w(\lambda_U + \rho_c^M - \rho_g^M)}(g) d_{w(\lambda_U + \rho_g^M - \rho_n^M)}^{M_g}}{e^{\rho^M - \rho_g^M}(g) \prod_{\alpha \in R^+(M, T) \setminus R^+(M_g, T)} (1 - e^\alpha(g))}. \end{aligned} \tag{6.13}$$

Proof. The first statement is one conclusion of Theorem 2.1 in [20].

Now suppose that P is a maximal cuspidal parabolic subgroup and $g \in T$. Let $\{X_1, \dots, X_s\}$ be an orthonormal basis of $\mathfrak{p} \cap \mathfrak{m}$. For every finite-dimensional representation Z of $K \cap M$, consider the Dirac operator

$$D_Z^M = \sum_{j=1}^s R(X_j) \otimes c(X_j) \otimes 1_Z$$

on $(C^\infty(M) \otimes \Delta_{\mathfrak{p} \cap \mathfrak{m}} \otimes Z)^{K \cap M}$, analogous to (3.14). Let τ_g^M be the orbital integral with respect to g as an element of M . Then (5.4) and (5.5) in [20] imply that

$$\begin{aligned} \langle \Phi_{P,g}, \text{index}_G(D_W) \rangle &= \\ & \tau_g^M(\text{index}_M(D_{\Delta_{\mathfrak{t}/(\mathfrak{t} \cap \mathfrak{m})}^+}^M \otimes W|_{K \cap M})) - \tau_g^M(\text{index}_M(D_{\Delta_{\mathfrak{t}/(\mathfrak{t} \cap \mathfrak{m})}^-}^M \otimes W|_{K \cap M})). \end{aligned}$$

Hence (6.13) follows by (3.17) and Theorem 6.12. \square

Example 6.17. If $\text{rank}(G) = \text{rank}(K)$, then G is a (maximal) cuspidal parabolic subgroup of itself, and $\Phi_{G,g} = \tau_g$. So (6.13) reduces to (6.12).

Example 6.18. If P is a maximal cuspidal parabolic subgroup and g lies in the centre of M , then analogously to Example 6.13, the equality (6.13) reduces to

$$\langle \Phi_{P,g}, \text{index}_G(D_W) \rangle = \sum_{U \in \widehat{K \cap M}} m_U e^{\lambda_U - \rho_n^M}(g) d_{\lambda_U + \rho_c^M}^M.$$

Example 6.19. If P is a maximal cuspidal parabolic subgroup, $g \in T$ and the powers of g are dense in T , then analogously to Example 6.14, the equality (6.13) reduces to

$$\begin{aligned} \langle \Phi_{P,g}, \text{index}_G(D_W) \rangle &= \\ (-1)^{\dim(M/(K \cap M))/2} \sum_{U \in \widehat{K \cap M}} m_U \frac{\sum_{w \in W_{K \cap M}} \det(w) e^{w(\lambda_U + \rho_c^M)}(g)}{e^{\rho^M}(g) \prod_{\alpha \in R^+(M,T)} (1 - e^{-\alpha}(g))} &= \\ & (-1)^{\dim(M/(K \cap M))/2} \frac{\chi_{\Delta_{\mathfrak{t}/(\mathfrak{t} \cap \mathfrak{m})}}(g) \chi_W(g)}{\chi_{\Delta_{\mathfrak{p} \cap \mathfrak{m}}}(g)}, \end{aligned}$$

where χ denotes the character of a finite-dimensional representation. This is (6.3) in [20].

6.6 Proofs of the index theorems

Let $F \subset \widehat{K \times K}$ be the set of irreducible subrepresentations of $\text{End}(\Delta_{\mathfrak{p}} \otimes W)$. For any continuous trace τ on $\mathcal{S}_F(G)$ (with notation as above Corollary 5.9), Theorem 2.12 implies that

$$\tau^{\Delta_{\mathfrak{p}} \otimes W} \circ \Psi_*(\text{index}_{\mathcal{S}_F(G)}(D_W)) = \tau^{\Delta_{\mathfrak{p}} \otimes W}(\Psi(e^{-tD_W^- D_W^+})) - \tau^{\Delta_{\mathfrak{p}} \otimes W}(\Psi(e^{-tD_W^+ D_W^-})). \quad (6.14)$$

If $g \in G$ is semisimple and $\tau = \tau_g$, then Proposition 3.6 in [21] implies that the right hand side equals $\tau_g(\text{index}_G(D_W))$. So parts (b) and (c) of Theorem 3.17 follow from Theorem 6.12. Part (a) follows by Example 6.13.

Similarly, in the setting of Theorem 3.18, we have

$$\langle \Phi_{M,g}^{\Delta_p \otimes W}, \text{index}_{S,G}(D_W) \rangle = \langle \Phi_{P,g}, \text{index}_G(D_W) \rangle.$$

So Theorem 3.18 follows from Corollary 6.16 and Example 6.18.

To prove Theorem 3.20, we just apply the computations in [6] to the right hand side of (6.14). The case of part (a) of Theorem 3.20 where $G \neq \text{SU}(2n, 1)$, and case (b) of this theorem, are (4.5.3) in [6], which comes from [5]. In the setting of part (a) of Theorem 3.20 for $G = \text{SU}(2n, 1)$, the equality (4.5.6) in [6] states that the right hand side of (6.14) equals the right hand side of (3.18). In the setting of part (c) of Theorem 3.20 with $\lambda + \rho_c$ regular, it is proved on page 194 of [6] that the right hand side of (6.14) equals the right hand side of (3.19). The case of part (c) of Theorem 3.20 for real hyperbolic space of dimension at least 4 is proved on the same page of [6].

7 Roe algebras and von Neumann algebras

We now consider a slightly more general setting than in Section 3.1, and relate the equivariant higher index from Definition 2.13 to the well-known equivariant higher index defined in terms of Roe algebras [39], and to a higher index involving the group von Neumann algebra of Γ . From the latter, we can recover the L^2 -index using the von Neumann trace. We have seen that this, and more, can be recovered from the index of Definition 2.13. (See Theorem 3.17(a).) In the special case where $\Gamma \backslash X$ is compact, we show that the Γ -equivariant index (6.1) can be recovered from the index from Definition 2.13.

7.1 Roe algebras

Suppose that $X = G/K$, for a connected, unimodular Lie group G , and a compact subgroup $K < G$. We write $\mathcal{A}(E) := \mathcal{A}_{\{1\}}(E)$ for the algebra of Definition 2.4, with $S = \{1\}$. For $a > 0$, we write $\|\cdot\|_a := \|\cdot\|_{a,1}$ for the corresponding seminorm on $\mathcal{A}(E)$. Let $a_0, C > 0$ be as in Definition 2.1.

Lemma 7.1. *The action by an element $\kappa \in \mathcal{A}(E)$ on $L^2(E)$ is a bounded operator, and*

$$\|\kappa\|_{B(L^2(E))} \leq C \|\kappa\|_{2a_0}. \quad (7.1)$$

Proof. Let $\kappa \in \Gamma_{\{1\}, \text{fp}}^\infty(E)$. Let $s \in L^2(E)$. Then

$$\begin{aligned} \|\kappa \cdot s\|_{L^2(E)}^2 &\leq \int_X \int_X \int_X \|\kappa(x, x')\| \|\kappa(x, x'')\| \|s(x')\| \|s(x'')\| d\mu(x') d\mu(x'') d\mu(x) \\ &\leq \|\kappa\|_{2a_0}^2 \int_X \int_X \int_X e^{-2a_0(d(x, x') + d(x, x''))} \|s(x')\| \|s(x'')\| d\mu(x') d\mu(x'') d\mu(x) \\ &\leq \|\kappa\|_{2a_0}^2 \int_X \int_X \int_X e^{-a_0(d(x, x') + d(x, x''))} e^{-a_0 d(x', x'')} \|s(x')\| \|s(x'')\| d\mu(x') d\mu(x'') d\mu(x), \end{aligned}$$

by the triangle inequality. Because the integrand is nonnegative, the integral on the right converges if and only if the following integral converges, and then these integrals are equal:

$$\|\kappa\|_{2a_0}^2 \int_X \int_X e^{-a_0 d(x', x'')} \|s(x')\| \|s(x'')\| \left(\int_X e^{-a_0(d(x, x') + d(x, x''))} d\mu(x) \right) d\mu(x') d\mu(x''). \quad (7.2)$$

For fixed $x', x'' \in X$, the Cauchy–Schwartz inequality implies that

$$\int_X e^{-a_0(d(x, x') + d(x, x''))} d\mu(x) \leq C.$$

So (7.2) is bounded above by

$$\|\kappa\|_{2a_0}^2 C \int_X \int_X e^{-a_0 d(x', x'')} \|s(x')\| \|s(x'')\| d\mu(x') d\mu(x'').$$

By G -invariance of d , the latter expression equals

$$\begin{aligned} &\|\kappa\|_{2a_0}^2 C \int_X \int_X e^{-a_0 d(eK, g^{-1}g'K)} \|s(gK)\| \|s(g'K)\| d\mu(gK) d\mu(g'K) \\ &= \|\kappa\|_{2a_0}^2 C \int_X \int_X e^{-a_0 d(eK, g''K)} \|s(gK)\| \|s(gg''K)\| d\mu(gK) d\mu(g''K), \quad (7.3) \end{aligned}$$

where we substituted $g''K = g^{-1}g'K$ and used G -invariance of μ .

Since G -invariant densities on G/K differ by constants, the measure μ equals the measure $d(gK)$ in (2.9) for some Haar measure dg on G . By right invariance of dg and the Cauchy–Schwartz inequality, we have for all $g'' \in G$,

$$\int_X \|s(gK)\| \|s(gg''K)\| d\mu(gK) \leq \|s\|_{L^2}^2.$$

So the right hand side of (7.3) is less than or equal to

$$\|\kappa\|_{2a_0}^2 C \int_X e^{-a_0 d(eK, g''K)} d\mu(g''K) \|s\|_{L^2(E)}^2 \leq \|\kappa\|_{2a_0}^2 C^2 \|s\|_{L^2(E)}^2.$$

So (7.1) follows. We also find that, by the Fubini–Tonelli theorem, all integrals that appeared converge (absolutely). \square

Let $H < G$ be any closed subgroup. Suppose that the action by H on X lifts to E , preserving the Hermitian metric. Let $C^*(X; E)^H$ be the H -equivariant Roe algebra of X defined on the $C_0(X)$ -module $L^2(E)$. This is the closure in $\mathcal{B}(L^2(E))$ of the algebra of operators that are H -equivariant and locally compact, and have finite propagation.

Lemma 7.2. *The algebra $\mathcal{A}(E)^H$ is contained in $C^*(X; E)^H$, and the inclusion map is continuous.*

Proof. The dense subalgebra $\Gamma_{\{1\}, \text{fp}}^\infty(E \boxtimes E^*)^H$ of $\mathcal{A}(E)^H$ consists of H -equivariant, locally compact operators with finite propagation. So the claim follows from Lemma 7.1. \square

Let D be as in Subsection 2.2. It has an H -equivariant higher index [17, 32, 39]

$$\text{index}_{C^*(X; E)^H}(D) \in K_0(C^*(X; E)^H). \quad (7.4)$$

Proposition 7.3. *Under the map $K_0(\mathcal{A}(E)^H) \rightarrow K_0(C^*(X; E)^H)$ induced by the inclusion from Lemma 7.2, the class $\text{index}_{\{1\}, H}(D) \in K_0(\mathcal{A}(E)^H)$ is mapped to $\text{index}_{C^*(X; E)^H}(D)$.*

Proof. The operator (4.11) is a multiplier of $C^*(X; E)^H$; see e.g. Lemma 2.1 in [34]. Hence $\text{index}_{C^*(X; E)^H}(D)$ is also given by (2.14). See also (3.9) in [21], Exercise 12.7.3 in [19] and page 256 of [15]. \square

7.2 The equivariant index for cocompact actions

The index (7.4) is closely related to the equivariant index (6.1). The inclusion map

$$j_{C_r^*(G)} : (\mathcal{S}(G) \otimes \text{End}(V))^{K \times K} \hookrightarrow C_r^*(G) \otimes \text{End}(V)$$

is continuous, because $\mathcal{S}(G)$ is continuously included in $L^1(G)$. It induces

$$(j_{C_r^*(G)})_* : K_0((\mathcal{S}(G) \otimes \text{End}(V))^{K \times K}) \rightarrow K_0(C_r^*(G) \otimes \text{End}(V)) = K_0(C_r^*(G)).$$

Corollary 7.4. *We have*

$$(j_{C_r^*(G)})_* \circ \Psi_*(\text{index}_{\mathcal{S}, G}(D)) = \text{index}_G(D) \in K_0(C_r^*(G)).$$

Proof. By Lemma 7.1, the map Ψ from Lemma 3.4 defines an algebra isomorphism

$$\Psi: C^*(G/K; E)^G \xrightarrow{\cong} (C_r^*(G) \otimes \text{End}(V))^{K \times K}.$$

By Corollary 4.3 in [17] (compare also [33]),

$$\Psi_*(\text{index}_{C^*(G/K; E)^G}(D)) = \text{index}_G(D).$$

As in the proof of Proposition 7.3, $\text{index}_{C^*(G/K; E)^G}(D)$ is given by the idempotents in (2.14). By Lemma 7.2, we have a continuous inclusion maps

$$\mathcal{A}_{S,q}(E)^G \hookrightarrow \mathcal{A}(E)^G \hookrightarrow C^*(G/K; E)^G.$$

By Proposition 7.3, the map on K -theory induced by these inclusions maps $\text{index}_{S,G}(D)$ to $\text{index}_{C^*(G/K; E)^G}(D)$. Because of compatibility between the map Ψ on different algebras and the relevant inclusion maps, the claim follows. \square

Our main interest is in the case where $\Gamma \backslash X$ is noncompact, but has finite volume. It is still a natural question how the higher index from Definition 2.13 is related to the Γ -equivariant index (6.1) in the case where $\Gamma \backslash X$ is compact. So suppose that $\Gamma \backslash X$ is compact. Then we have

$$\text{index}_\Gamma(D) \in K_0(C_r^*(\Gamma)). \quad (7.5)$$

In Proposition 4.10 in [28], a restriction class

$$[\text{Res}_\Gamma^G] \in KK_0(C_r^*(G), C_r^*(\Gamma)) \quad (7.6)$$

is constructed. The Kasparov product with this class from the right defines a restriction map

$$\text{Res}_\Gamma^G: K_0(C_r^*(G)) \rightarrow K_0(C_r^*(\Gamma)).$$

This allows us to recover the index (7.5) from the index of Definition 2.13.

Corollary 7.5. *If $\Gamma \backslash X$ is compact, then*

$$\text{Res}_\Gamma^G \circ (j_{C_r^*(G)})_* \circ \Psi_*(\text{index}_{S,G}(D)) = \text{index}_\Gamma(D) \in K_0(C_r^*(\Gamma)).$$

Proof. By Lemma 4.15 in [28],

$$\text{Res}_\Gamma^G(\text{index}_G(D)) = \text{index}_\Gamma(D).$$

Hence the claim follows by Corollary 7.4. \square

Remark 7.6. In the construction of the class (7.6) in [28], discreteness of Γ is not essential. So there is a straightforward generalisation of Corollary 7.5 to an equality

$$\text{Res}_H^G \circ (j_{C_r^*(G)})_* \circ \Psi_*(\text{index}_{S,G}(D)) = \text{index}_H(D)$$

for any closed subgroup $H < G$ such that $H \backslash G$ is compact.

7.3 Von Neumann algebras and the von Neumann trace

Let $Y \subset X$ be a fundamental domain for the action by Γ , such that the closure of Y equals the closure of its interior. Consider the unitary isomorphism

$$U: L^2(E) \rightarrow l^2(\Gamma) \otimes L^2(E|_Y)$$

given by

$$(Us)(\gamma, x) = \gamma s(\gamma^{-1}x),$$

for $s \in L^2(E)$, $\gamma \in \Gamma$ and $x \in Y$.

As in Subsection 7.1, we write $\mathcal{A}(E) := \mathcal{A}_{\{1\}}(E)$. Let L be the left regular representation of Γ in $l^2(\Gamma)$. If $\kappa \in \mathcal{A}(E)^\Gamma$ and $\gamma \in \Gamma$, then we write $\gamma \circ \kappa = (\gamma, e) \cdot \kappa \in \mathcal{A}(E)$ in terms of the action (2.4), so that

$$(\gamma \circ \kappa)(x, x') = \gamma \kappa(\gamma^{-1}x, x'), \quad (7.7)$$

for $x, x' \in X$.

Lemma 7.7. *Consider a Γ -invariant, measurable section κ of $E \boxtimes E^*$ that is the Schwartz kernel of a bounded operator on $L^2(E)$, which we also denote by κ . Then for all $s, s' \in L^2(E)$,*

$$\sum_{\gamma \in \Gamma} (s, U^{-1} \circ (L_\gamma \otimes (\gamma \circ \kappa)|_{Y \times Y}) \circ U s')_{L^2(E)} = (s, \kappa s')_{L^2(E)}. \quad (7.8)$$

Proof. For $\psi \in l^2(\Gamma) \otimes L^2(E|_Y)$, $\gamma \in \Gamma$ and $x \in Y$,

$$(U^{-1}\psi)(\gamma x) = \gamma \psi(\gamma^{-1}, x).$$

Using this, and writing out definitions, we find that for all $s' \in L^2(E)$, $\gamma, \gamma' \in \Gamma$ and $x \in Y$,

$$(U^{-1} \circ (L_\gamma \otimes (\gamma \circ \kappa)|_{Y \times Y}) \circ U s')(\gamma' x) = \int_Y \gamma' \gamma \kappa(\gamma^{-1}x, y) \gamma^{-1} \gamma'^{-1} s'(\gamma' \gamma y) d\mu(y).$$

By Γ -invariance of κ , the right hand side equals

$$\int_Y \kappa(\gamma'x, \gamma'\gamma y) s'(\gamma'\gamma y) d\mu(y).$$

So the left hand side of (7.8) equals

$$\sum_{\gamma, \gamma' \in \Gamma} \int_Y \int_Y (s(\gamma'x), \kappa(\gamma'x, \gamma'\gamma y) s'(\gamma'\gamma y))_E d\mu(y) d\mu(x),$$

which equals the right hand side of (7.8). \square

Let $A(E)^\Gamma$ be the Banach space of bounded, Γ -equivariant operators on $L^2(E)$ with bounded, measurable Schwartz kernels. For $T \in A(E)^\Gamma$, with Schwartz kernel κ , define

$$\|T\|_{A(E)^\Gamma} := \|T\|_{\mathcal{B}(L^2(E))} + \|\kappa\|_{L^\infty(E \otimes E^*)}.$$

Note that $A(E)^\Gamma$ is not a Banach algebra. But by Lemma 7.1, the algebra $\mathcal{A}(E)^\Gamma$ is continuously included in $A(E)^\Gamma$.

Conjugation by U is an isometric algebra isomorphism from $\mathcal{B}(L^2(E))$ onto $\mathcal{B}(l^2(\Gamma) \otimes L^2(E|_Y))$. It restricts to an injective, continuous linear map

$$j_{\mathcal{N}}: A(E)^\Gamma \rightarrow \mathcal{B}(l^2(\Gamma) \otimes L^2(E|_Y)).$$

This map is an algebra homomorphism on any algebra included in $A(E)^\Gamma$, such as $\mathcal{A}(E)^\Gamma$.

If \mathcal{H} is any Hilbert space, and $A \subset \mathcal{B}(\mathcal{H})$ a C^* -algebra, then we write $\mathcal{N}(\Gamma) \otimes A$ for the algebra of $T \in \mathcal{B}(l^2(\Gamma) \otimes \mathcal{H})$ for which there is a sequence $(T_j)_{j=1}^\infty$ in the algebraic tensor product $\mathbb{C}\Gamma \otimes A$ such that for all $\psi, \psi' \in l^2(\Gamma) \otimes \mathcal{H}$,

$$(\psi, T_j \psi')_{l^2(\Gamma) \otimes \mathcal{H}} \rightarrow (\psi, T \psi')_{l^2(\Gamma) \otimes \mathcal{H}}. \quad (7.9)$$

Here we let $\mathbb{C}\Gamma$ act on $l^2(\Gamma)$ via the left regular representation.

Example 7.8. If $\mathcal{H} = \mathbb{C} = A$, then $\mathcal{N}(\Gamma) \otimes \mathbb{C}$ is the completion of $\mathbb{C}\Gamma$ in the weak topology on operators on $l^2(\Gamma)$, i.e. the von Neumann algebra of Γ .

Proposition 7.9. *If $\Gamma \backslash G/K$ has finite volume, then*

$$j_{\mathcal{N}}(A(E)^\Gamma) \subset \mathcal{N}(\Gamma) \otimes \mathcal{K}(L^2(E|_Y)).$$

Furthermore, the map $j_{\mathcal{N}}: A(E)^\Gamma \rightarrow \mathcal{N}(\Gamma) \otimes \mathcal{K}(L^2(E|_Y))$ is continuous.

Proof. Let $\kappa \in \mathcal{A}(E)^\Gamma$. Then $j_{\mathcal{N}}(\kappa) \in \mathcal{B}(l^2(\Gamma) \otimes L^2(E|_Y))$, and

$$\|j_{\mathcal{N}}(\kappa)\|_{\mathcal{B}(l^2(\Gamma) \otimes L^2(E|_Y))} = \|\kappa\|_{\mathcal{B}(L^2(E))}. \quad (7.10)$$

Now for every $\gamma \in \Gamma$, the operator $(\gamma \circ \kappa)|_{Y \times Y}$ on $L^2(E|_Y)$ is Hilbert–Schmidt, hence compact, because Y has finite volume and κ is bounded. So

$$L_\gamma \otimes (\gamma \circ \kappa)|_{Y \times Y} \in \mathbb{C}\Gamma \otimes \mathcal{K}(L^2(E|_Y)).$$

By Lemma 7.7 and unitarity of U , the sum

$$\sum_{\gamma} L_\gamma \otimes (\gamma \circ \kappa)|_{Y \times Y}$$

converges to $j_{\mathcal{N}}(\kappa)$ in the topology defined by (7.9). So $j_{\mathcal{N}}(\kappa) \in \mathcal{N}(\Gamma) \otimes \mathcal{K}(L^2(E|_Y))$.

Continuity of $j_{\mathcal{N}}$ follows from (7.10): if $\psi, \psi' \in l^2(\Gamma) \otimes L^2(E|_Y)$, then

$$|(\psi, j_{\mathcal{N}}(\kappa)\psi')_{l^2(\Gamma) \otimes L^2(E|_Y)}| \leq \|\kappa\|_{\mathcal{B}(L^2(E))} \|\psi\|_{l^2(\Gamma) \otimes L^2(E|_Y)} \|\psi'\|_{l^2(\Gamma) \otimes L^2(E|_Y)}.$$

□

Suppose that $\Gamma \backslash G/K$ has finite volume. Then by Proposition 7.9, we obtain an algebra homomorphism $j_{\mathcal{N}}: \mathcal{A}(E)^\Gamma \rightarrow \mathcal{N}(\Gamma) \otimes \mathcal{K}(L^2(E|_Y))$. This induces

$$(j_{\mathcal{N}})_*: K_0(\mathcal{A}(E)^\Gamma) \rightarrow K_0(\mathcal{N}(\Gamma) \otimes \mathcal{K}(L^2(E|_Y))) = K_0(\mathcal{N}(\Gamma)).$$

We obtain a link with the index studied in [14]. Let D be as in Subsection 2.2, with $H = \Gamma$. Then we have

$$(j_{\mathcal{N}})_*(\text{index}_{\{1\}, \Gamma}(D)) \in K_0(\mathcal{N}(\Gamma)).$$

On the algebra $\mathcal{N}(\Gamma)$, we have the von Neumann trace τ_e^Γ , given by evaluation at $e \in \Gamma$. We will extend this to a trace on a subalgebra of $\mathcal{N}(\Gamma) \otimes \mathcal{K}(L^2(E|_Y))$, and apply it to $j_{\mathcal{N}}(\text{index}_{\{1\}, \Gamma}(D))$.

Lemma 7.10. *For all $\kappa \in \mathcal{A}(E)^\Gamma$, the integral*

$$\text{Tr}_e(\kappa) := \int_Y \text{tr}(\kappa(x, x)) d\mu(x)$$

converges. This defines a continuous trace Tr_e on $\mathcal{A}(E)^\Gamma$.

Proof. This follows from boundedness of kernels in $\mathcal{A}(E)^\Gamma$ and finite volume of Y . \square

For a bounded, continuous $\kappa_Y \in \Gamma(E|_Y \boxtimes E^*|_Y)$, we write

$$\mathrm{Tr}_Y(\kappa_Y) = \int_Y \mathrm{tr}(\kappa_Y(y, y)) d\mu(y).$$

It is immediate from the definitions that for all $\kappa \in \mathcal{A}(E)^\Gamma$,

$$(\tau_e^\Gamma \otimes \mathrm{Tr}_Y)(j_{\mathcal{N}}(\kappa)) = \mathrm{Tr}_e(\kappa).$$

In particular, Lemma 7.10 implies that $\tau_e^\Gamma \otimes \mathrm{Tr}_Y$ defines a trace on $\mathrm{im}(j_{\mathcal{N}}) \subset \mathcal{N}(\Gamma) \otimes \mathcal{K}(L^2(E|_Y))$, which is continuous in the topology on $\mathrm{im}(j_{\mathcal{N}})$ that makes $j_{\mathcal{N}}$ a homeomorphism. Hence we can apply $\tau_e^\Gamma \otimes \mathrm{Tr}_Y$ to

$$(j_{\mathcal{N}})_*(\mathrm{index}_{\{1\}, \Gamma}(D)) \in K_0(\mathrm{im}(j_{\mathcal{N}})).$$

If D is equivariant with respect to G , rather than just Γ , then $\mathrm{Tr}_e(\mathrm{index}_{\{1\}, \Gamma}(D))$ equals $\mathrm{vol}(\Gamma \backslash G)$ times the L^2 -index of D , with respect to the action by G . This was computed in Theorem 5.2 in [14]. In the setting of Theorem 3.17, we have the following L^2 -index theorem.

Corollary 7.11. *Let D_W be as in (3.14), and $d_{\lambda+\rho_c}^G$ as in (3.16). Then*

$$(\tau_e^\Gamma \otimes \mathrm{Tr}_Y)(j_{\mathcal{N}}(\mathrm{index}_{\{1\}, \Gamma}(D_W))) = \mathrm{vol}(\Gamma \backslash G) d_{\lambda+\rho_c}^G.$$

Proof. Let $\mathcal{A}(G)$ be the completion of $C_c^\infty(G)$ in the seminorms $\|\cdot\|_{a,1,1}$ in (3.1). Then as in Lemma 3.4, we obtain a continuous isomorphism $\Psi: \mathcal{A}(E)^G \rightarrow (\mathcal{A}(G) \otimes \mathrm{End}(V))^{K \times K}$. One can check directly that

$$(\tau_e^\Gamma \otimes \mathrm{Tr}_Y) \circ j_{\mathcal{N}}|_{\mathcal{A}(E)^G} = \mathrm{Tr}_e|_{\mathcal{A}(E)^G} = \mathrm{vol}(\Gamma \backslash G) \tau_e \circ \Psi.$$

Hence the claim follows from case (a) of Theorem 3.17. \square

References

- [1] Z. Alimohammadi and A. Rejali. On the multiplier algebra of Fréchet algebras. *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.*, 81(4):125–132, 2019.
- [2] James Arthur. Some tempered distributions on semisimple groups of real rank one. *Ann. of Math. (2)*, 100:553–584, 1974.

- [3] Michael Atiyah and Raoul Bott. A Lefschetz fixed point formula for elliptic complexes. II. Applications. *Ann. of Math. (2)*, 88:451–491, 1968.
- [4] Michael Atiyah and Wilfried Schmid. A geometric construction of the discrete series for semisimple Lie groups. *Invent. Math.*, 42:1–62, 1977.
- [5] Dan Barbasch. Fourier inversion for unipotent invariant integrals. *Trans. Amer. Math. Soc.*, 249(1):51–83, 1979.
- [6] Dan Barbasch and Henri Moscovici. L^2 -index and the Selberg trace formula. *J. Funct. Anal.*, 53(2):151–201, 1983.
- [7] Paul Baum, Alain Connes, and Nigel Higson. Classifying space for proper actions and K -theory of group C^* -algebras. In *C^* -algebras: 1943–1993 (San Antonio, TX, 1993)*, volume 167 of *Contemp. Math.*, pages 240–291. American Mathematical Society, Providence, RI, 1994.
- [8] Jean-Michel Bismut. *Hypoelliptic Laplacian and orbital integrals*, volume 177 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2011.
- [9] Jacob Bradd. Compatible decomposition of the Casselman algebra and the reduced group C^* -algebra of a real reductive group. ArXiv:2312.11773, 2023.
- [10] Jacob Bradd and Nigel Higson. On Novodvorskii’s theorem and the Oka principle. *Eur. J. Math.*, 7(4):1729–1752, 2021.
- [11] A. L. Carey, V. Gayral, A. Rennie, and F. A. Sukochev. Index theory for locally compact noncommutative geometries. *Mem. Amer. Math. Soc.*, 231(1085):vi+130, 2014.
- [12] William Casselman. Introduction to the Schwartz space of $\Gamma \backslash G$. *Canad. J. Math.*, 41(2):285–320, 1989.
- [13] Xiaoman Chen, Jinmin Wang, Zhizhang Xie, and Guoliang Yu. De-localized eta-invariants, cyclic cohomology and higher rho-invariants. ArXiv:1901.02378, 2019.
- [14] Alain Connes and Henri Moscovici. The L^2 -index theorem for homogeneous spaces of Lie groups. *Ann. of Math. (2)*, 115(2):291–330, 1982.
- [15] Alain Connes and Henri Moscovici. Cyclic cohomology, the Novikov conjecture and hyperbolic groups. *Topology*, 29(3):345–388, 1990.

- [16] H. Garland and M. S. Raghunathan. Fundamental domains for lattices in (R-)rank 1 semisimple Lie groups. *Ann. of Math. (2)*, 92:279–326, 1970.
- [17] Hao Guo, Peter Hochs, and Varghese Mathai. Equivariant Callias index theory via coarse geometry. *Ann. Inst. Fourier (Grenoble)*, 71(6):2387–2430, 2021.
- [18] Harish-Chandra. Discrete series for semisimple Lie groups. II: Explicit determination of the characters. *Acta Math.*, 116:1–111, 1966.
- [19] Nigel Higson and John Roe. *Analytic K-homology*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2000. Oxford Science Publications.
- [20] Peter Hochs, Yanli Song, and Xiang Tang. An index theorem for higher orbital integrals. *Math. Ann.*, 382(1-2):169–202, 2022.
- [21] Peter Hochs and Hang Wang. A fixed point formula and Harish-Chandra’s character formula. *Proc. London Math. Soc.*, 00(3):1–32, 2017.
- [22] Peter Hochs and Hang Wang. A fixed point theorem on noncompact manifolds. *Ann. K-theory*, 2(3):235–286, 2018.
- [23] Peter Hochs and Hang Wang. Orbital integrals and K -theory classes. *Ann. K-Theory*, 4(2):185–209, 2019.
- [24] Gennadi Kasparov. Index of invariant elliptic operators, K -theory and representations of Lie groups. *Dokl. Akad. Nauk SSSR*, 268(3):533–537, 1983.
- [25] Anthony Knapp. *Representation theory of semisimple groups*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, Oxford, 1986.
- [26] Anthony W. Knapp. *Lie groups beyond an introduction*, volume 140 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1996.
- [27] Vincent Lafforgue. Banach KK-theory and the Baum-Connes conjecture. In *Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002)*, pages 795–812. Higher Ed. Press, Beijing, 2002.

- [28] Bram Mesland, Mehmet Haluk Şengün, and Hang Wang. A K -theoretic Selberg trace formula. In *Operator theory, operator algebras and their interactions with geometry and topology—Ronald G. Douglas memorial volume*, volume 278 of *Oper. Theory Adv. Appl.*, pages 403–424. Birkhäuser/Springer, Cham, [2020] ©2020.
- [29] Henri Moscovici. L^2 -index of elliptic operators on locally symmetric spaces of finite volume. In *Operator algebras and K -theory (San Francisco, Calif., 1981)*, volume 10 of *Contemp. Math.*, pages 129–137. Amer. Math. Soc., Providence, R.I., 1982.
- [30] M. Scott Osborne and Garth Warner. Multiplicities of the integrable discrete series: the case of a nonuniform lattice in an R -rank one semisimple group. *J. Functional Analysis*, 30(3):287–310, 1978.
- [31] Rajagopalan Parthasarathy. Dirac operator and the discrete series. *Ann. of Math. (2)*, 96:1–30, 1972.
- [32] John Roe. *Index theory, coarse geometry, and topology of manifolds*, volume 90 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1996.
- [33] John Roe. Comparing analytic assembly maps. *Q. J. Math.*, 53(2):241–248, 2002.
- [34] John Roe. Positive curvature, partial vanishing theorems and coarse indices. *Proc. Edinb. Math. Soc. (2)*, 59(1):223–233, 2016.
- [35] Yanli Song and Xiang Tang. Higher orbit integrals, cyclic cocycles, and K -theory of reduced group C^* -algebra. ArXiv:1910.00175.
- [36] Hang Wang. L^2 -index formula for proper cocompact group actions. *J. Noncommut. Geom.*, 8(2):393–432, 2014.
- [37] Garth Warner. Selberg’s trace formula for nonuniform lattices: the R -rank one case. In *Studies in algebra and number theory*, Adv. in Math. Suppl. Stud., 6, pages 1–142, 1979.
- [38] Antony Wassermann. Une démonstration de la conjecture de Connes-Kasparov pour les groupes de Lie linéaires connexes réductifs. *C. R. Acad. Sci. Paris Sér. I Math.*, 304(18):559–562, 1987.

- [39] Rufus Willett and Guoliang Yu. *Higher index theory*, volume 189 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2020.