

ON NEW TYPES OF CONVEX FUNCTIONS AND THEIR PROPERTIES

M.EMIN ÖZDEMİR

ABSTRACT. In this study, Firstly, we will write two new convex functions for $-1 < n - \alpha \leq 1$ and two new lemmas. Then we will find the relevance of the two new lemmas to Caputo-left-sided derivatives under additional conditions and draw conclusions about them. Secondly, We will give examples that support it. In what follows, we will rewrite the Hermite Hadamard integral inequality based on this new definitions. Furthermore, we will write an elementary inequality valid for value $-1 < n - \alpha \leq 1$.

1. INTRODUCTION

Many important inequalities are established for the class of convex functions, but one of the most important is so-called Hermite–Hadamard’s inequality (or Hadamard’s inequality). This double inequality is stated as follows in literature:

Let $f : I \subseteq R \rightarrow R$ be a convex function and let $a, b \in I$, with $a < b$. The following double inequality;

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2}$$

is known as Hermite- Hadamard’s inequality for convex functions. For many papers connected with (1.1) for different types of convex functions see [1]-[35] the references therein. The paper is organized as follows :

Firstly, we need some representations and preliminary information that appear in the inequalities we will write.

Caputo left-sided derivative

$${}^C D_{a+}^{\alpha} [f](x) = \frac{1}{\Gamma(n-\alpha)} \int_a^x (x-\xi)^{n-\alpha-1} \frac{d^n (f(\xi))}{d\xi^n} dt, \quad x > a$$

Caputo right-sided derivative

$$(1.2) \quad {}^C D_{b-}^{\alpha} [f](x) = \frac{(-1)^n}{\Gamma(n-\alpha)} \int_x^b (\xi-x)^{n-\alpha-1} \frac{d^n (f(\xi))}{d\xi^n} dt, \quad x < b$$

As can be seen, In Caputo, The first calculated the derivative of the integer order and then the integral of the non-integer order.

Second, we need the new definition and some of its implications, which we present below.

2020 *Mathematics Subject Classification.* Primary 26A33, 26A51; Secondary 05A15, 15A18.

Key words and phrases. convexity,

This paper is in final form and no version of it will be submitted for publication elsewhere.

Definition 1. ($(n - \alpha)$ convex) The function $f : I \subseteq [1, \infty) \rightarrow \mathbb{R}^+$ is said to be $(n - \alpha)$ convex-concave, if the following inequality holds:

$$(1.3) \quad f(tx + (1-t)y) \leq t^{n-\alpha}f(x) + \left| \left(t^{n-\alpha} - (1-t)^{n-\alpha} \right) \right| f(y)$$

for $-1 < n - \alpha \leq 1, (0 < n \leq 1, 0 \leq \alpha < 1)$ $t \in (0, 1]$, $x, y \in I$. with $y < x$. If (1.3) reverses, then f is said to be concave on I . Below are some interpretations of the above definition.

a) Since $|t_1^{n-\alpha} - t_2^{n-\alpha}| \leq |t_1 - t_2|^{n-\alpha}$, for $n-\alpha \in (-1, 1]$ and $\forall t_1, t_2 \in [0, 1]$ the definition in(1.3) can be rewritten as

$$\begin{aligned} & f(tx + (1-t)y) \\ & \leq t^{n-\alpha}f(x) + \left| \left(t^{n-\alpha} - (1-t)^{n-\alpha} \right) \right| f(y) \\ & \leq t^{n-\alpha}f(x) + |2t - 1|^{n-\alpha} f(y). \end{aligned}$$

b) Since $n-\alpha \in (-1, 1]$, If we take into account the fact $|2t - 1|^{n-\alpha} \leq t^{n-\alpha}$, for $t \in [\frac{1}{2}, 1]$ we can rewrite the inequality (1.3).

We obtain a new definition written for convex functions:

$$\begin{aligned} f(tx + (1-t)y) & \leq t^{n-\alpha}f(x) + \left| \left(t^{n-\alpha} - (1-t)^{n-\alpha} \right) \right| f(y) \\ & \leq t^{n-\alpha}f(x) + |1 - 2t|^{n-\alpha} f(y) \\ & \leq t^{n-\alpha}f(x) + t^{n-\alpha}f(y) \\ & = t^{n-\alpha} (f(x) + f(y)) \end{aligned}$$

From the first and last terms

$$(1.4) \quad f(tx + (1-t)y) \leq t^{n-\alpha} (f(x) + f(y))$$

c) If we choose $t = \frac{1}{2}$, $n - \alpha = 1$ in (1.4) for $t \in [\frac{1}{2}, 1]$. we obtain

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x) + f(y)}{2}$$

d) If we just choose $t = \frac{1}{2}$ in (1.4)

$$f\left(\frac{x+y}{2}\right) \leq \left(\frac{1}{2}\right)^{n-\alpha} [f(x) + f(y)]$$

e) If we choose $t = \frac{\frac{1}{2}+1}{2}$ in (1.4) for $t \in [\frac{1}{2}, 1]$ we have

$$f\left(\frac{3x+y}{4}\right) \leq \left(\frac{3}{4}\right)^{n-\alpha} [f(x) + f(y)]$$

f) If we choose $t = \frac{\frac{3}{2}+1}{2}$ in (1.4) for $t \in [\frac{1}{2}, 1]$ we have

$$f\left(\frac{7x+6y}{8}\right) \leq \left(\frac{7}{8}\right)^{n-\alpha} [f(x) + f(y)]$$

g) If we choose and $t = \frac{\frac{7}{2}+1}{2}$ in (1.4) for $t \in [\frac{1}{2}, 1]$ we have

$$f\left(\frac{15x+14y}{8}\right) \leq \left(\frac{15}{16}\right)^{n-\alpha} [f(x) + f(y)]$$

If we continue in this sense, we obtain the sequence $\left(\frac{2^k-1}{2^k}\right)^{n-\alpha}$ which converges to 1 for $(n-\alpha; k) \rightarrow (\pm 1, \infty)$

$$\left(\frac{1}{2}\right)^{n-\alpha}, \left(\frac{3}{4}\right)^{n-\alpha}, \dots, \left(\frac{2^k-1}{2^k}\right)^{n-\alpha}, \dots \rightarrow 1 \quad (k = 1, 2, 3, \dots)$$

That is,

$$(1.5) \quad f\left(\frac{\left(\frac{2^k-1}{2^k}\right)x + \frac{1}{2^k}y}{2^k}\right) \leq \left(\frac{2^k-1}{2^k}\right)^{n-\alpha} [f(x) + f(y)]$$

and without losing generality we can write

$$(1.6) \quad f\left(\frac{\left(\frac{2^k-1}{2^k}\right)x + \frac{1}{2^k}y}{2^k}\right) \leq \left(\frac{2^k-1}{2^k}\right)^{n-\alpha} f(x) + \left(\frac{(2^k)^2-1}{2^k}\right)^{n-\alpha} f(y)$$

for $k = 1, 2, 3, \dots$ and $n - \alpha \in (-1, 1]$.

h) if we only integrate both sides of the inequality(1.4) according to t :

$$\frac{1}{x-y} \int_x^y f(u) du \leq \frac{f(x) + f(y)}{n-\alpha+1}; \quad u = tx + (1-t)y$$

i) If we choose $t = 1$ in (1.4) for $t \in [\frac{1}{2}, 1]$ we obtain

$$f(y) \geq 0$$

Additionally, If we take limit of (1.5) we again find the $f(y) \geq 0$ relation for $(n-\alpha; k) \rightarrow (\pm 1, \infty)$.

i) If we choose $n = \frac{1}{2}$, $\alpha = -\frac{1}{2}$ and $t = \frac{1}{2}$ in(1.3) . we obtain Jensen convexity:

$$\begin{aligned} f\left(\frac{x+y}{2}\right) &\leq t^{n-\alpha} f(x) + \left|t^{n-\alpha} - (1-t)^{n-\alpha}\right| f(y) \\ &= \frac{1}{2} f(x) + \left|\left(\left(\frac{1}{2}\right) - \left(1 - \frac{1}{2}\right)\right)\right| f(y) \\ &= \frac{1}{2} f(x) + |0| \cdot f(y) \leq \frac{1}{2} f(x) + \frac{1}{2} f(y) = \frac{f(x) + f(y)}{2} \end{aligned}$$

We used the fact that inequality $|t_1^{n-\alpha} - t_2^{n-\alpha}| \leq |t_1 - t_2|^{n-\alpha}$, for $n - \alpha \in (-1, 1]$, and $\forall t_1, t_2 \in [0, 1]$.

Now, in the light of the informations above, let's rearrange inequality(1.6) as definition.

Definition 2. The function $f : I \subseteq [1, \infty) \rightarrow \mathbb{R}^+$ is said to be $k^{n-\alpha}$ - convex, if the following inequality holds:

$$f\left(\frac{\left(\frac{2^k-1}{2^k}\right)x + \frac{1}{2^k}y}{2^k}\right) \leq \left(\frac{2^k-1}{2^k}\right)^{n-\alpha} f(x) + \left(\frac{(2^k)^2-1}{2^k}\right)^{n-\alpha} f(y)$$

for $k = 1, 2, 3, \dots$ and $n - \alpha \in (-1, 1]$.

Example : The function

$$f(x) = x$$

is an $k^{n-\alpha}$ - convex. Because

$$\begin{aligned} f\left(\frac{\left(\frac{2^k-1}{2^k}\right)x + \frac{1}{2^k}y}{2^k}\right) &= \frac{\left(\frac{2^k-1}{2^k}\right)x + \frac{1}{2^k}y}{2^k} \\ &\leq \left(\frac{2^k-1}{2^k}\right)^{n-\alpha} x + \left(\frac{(2^k)^2-1}{2^k}\right)^{n-\alpha} y \\ &\leq \left(\frac{2^k-1}{2^k}\right)^{n-\alpha} f(x) + \left(\frac{(2^k)^2-1}{2^k}\right)^{n-\alpha} f(y) \end{aligned}$$

In light of the above information, we write new theorems and new lemmas and interpreted their results.

2. HERMITE-HADAMARD INEQUALITY FOR $(n - \alpha)$ CONVEX FUNCTION

Theorem 1. *Let $f : I \subseteq [1, \infty) \rightarrow R^+$ be an $(n - \alpha)$ convex function. If $a < b$ and $f \in L[a, b]$, with $n - \alpha \in (-1, 1]$ then the following Hermite-Hadamard type inequality hold:*

$$(2.1) \quad 2^{n-\alpha-1} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{n - \alpha + 1}$$

Proof. firstly , let be $ta + (1-t)b = x$, for $t \in [0, 1]$

$$\begin{aligned} \frac{1}{b-a} \int_a^b f(x) dx &= \int_0^1 f(ta + (1-t)b) dt \\ &\leq \int_0^1 \left(t^{n-\alpha} f(a) + \left| t^{n-\alpha} - (1-t)^{n-\alpha} \right| f(b) \right) dt \\ &\leq \int_0^1 \left(t^{n-\alpha} f(a) + |1-2t|^{n-\alpha} f(b) \right) dt \\ &= \int_0^1 t^{n-\alpha} f(a) dt + \int_0^1 |1-2t|^{n-\alpha} f(b) dt \\ &= f(a) \int_0^1 t^{n-\alpha} dt + f(b) \int_0^{\frac{1}{2}} (1-2t)^{n-\alpha} dt \\ &\quad + f(b) \int_{\frac{1}{2}}^1 (2t-1)^{n-\alpha} dt \\ &= \frac{f(a) + f(b)}{n - \alpha + 1} \end{aligned}$$

From the first and last terms,we have

$$(2.2) \quad \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{n - \alpha + 1}$$

Here we again used the fact that inequality $|t_1^{n-\alpha} - t_2^{n-\alpha}| \leq |t_1 - t_2|^{n-\alpha}$, for $n - \alpha \in (-1, 1]$ and $\forall t_1, t_2 \in [0, 1]$,secondly,

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) &= f\left(\frac{[ta + (1-t)b] + [(1-t)a + tb]}{2}\right) \\
&= f\left(\frac{1}{2}[ta + (1-t)b] + \frac{1}{2}[(1-t)a + tb]\right) \\
&\leq \frac{1}{2^{n-\alpha}}f(ta + (1-t)b) \\
&\quad + \left|\left(\left(\frac{1}{2}\right)^{n-\alpha} - \left(1 - \frac{1}{2}\right)^{n-\alpha}\right)\right|f((1-t)a + tb) \\
&= \frac{1}{2^{n-\alpha}}f(ta + (1-t)b) + 0.f((1-t)a + tb) \\
&\leq \frac{1}{2^{n-\alpha}}f(ta + (1-t)b) + \frac{1}{2^{n-\alpha}}f((1-t)a + tb)
\end{aligned}$$

From the first and last terms, we have

$$(2.3) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{2^{n-\alpha}}f(ta + (1-t)b) + \frac{1}{2^{n-\alpha}}f((1-t)a + tb)$$

Now, If we integrate both sides of (2.3) over $[0, 1]$. We have

$$(2.4) \quad 2^{n-\alpha-1}f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx$$

Finally. If we combine inequalities (2.2) and 2.4 we get obtain we obtain (2.1).

Here we used fact that $0 < \frac{1}{2^{n-\alpha}}$. \square

Corollary 1. 1. *If we choose n and α so that $n = \frac{1}{k}$ ($k = 1, 2, 3, 4, \dots$), $\alpha = \frac{s-1}{s}$ ($k, s = 1, 2, 3, \dots$) .in (2.1) Then we have*

$$(2.5) \quad 2^{\frac{1}{k} + \frac{1}{s} - 2}f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{\frac{1}{k} + \frac{1}{s}}$$

If we choose $k = 1$ and $s = 1$ and in (2.5) we obtain directly well known Hermit- Hadamard inequalities in (1.1) :

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$$

or If we take limit of the inequality (2.5) for $(n, \alpha) \rightarrow (1, 0)$ we again obtain in (1.1).

If we choose n and α so that $n = \frac{1}{k}$ ($k = 1, 2, 3, \dots$) .and $\alpha = \frac{1}{s}$ ($s = 2, 3, \dots$)in (2.1) Then we have

$$2^{\frac{1}{k} - \frac{1}{s} - 1}f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{\frac{1}{k} - \frac{1}{s} + 1}$$

On the other hand, Since

$(\frac{1}{k} - \frac{1}{s} - 1, \frac{1}{k} - \frac{1}{s} + 1) > (-1, 1)$ for $(k, s) \rightarrow (\infty, \infty)$ we obtain

$$f\left(\frac{a+b}{2}\right) \leq \frac{2}{b-a} \int_a^b f(x) dx \leq 2(f(a) + f(b))$$

Suppose that the function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^+$ is $(n - \alpha)$ -convex. By applying the (2.1) on each of the intervals $[a, \frac{a+b}{2}]$ and $[\frac{a+b}{2}, b]$ we get

$$\begin{aligned} 2^{n-\alpha-1} f\left(\frac{a+b}{2}\right) &\leq 2^{n-\alpha-2} \left(f\left(\frac{3a+b}{4}\right) + f\left(\frac{3b+a}{4}\right) \right) \\ &\leq \frac{1}{b-a} \int_a^b f(x) dx \\ &\leq \frac{1}{n-\alpha+1} \left\{ \frac{f(a)+f(b)}{2} + f\left(\frac{a+b}{2}\right) \right\} \\ &\leq \frac{1}{n-\alpha+1} \left\{ \frac{f(a)+f(b)}{2} \right\} \end{aligned}$$

If the division process is continued, the difference between the image of the arithmetic mean of $[a, b]$ and the arithmetic mean of $[f(a), f(b)]$ will become smaller.

From first and last inequalities we obtain a Jensen type inequality for the $n - \alpha$ convex functions :

$$2^{n-\alpha-1} f\left(\frac{a+b}{2}\right) \leq \frac{1}{n-\alpha+1} \left\{ \frac{f(a)+f(b)}{2} \right\}$$

Theorem 2. let be $f : I \subseteq [1, \infty) \rightarrow \mathbb{R}^+$ a $(n - \alpha)$ convex function with $n - \alpha \in (-1, 1]$, $t \in (0, 1]$, $x, y \in I$. Then the following inequality holds :

$$(2.6) \quad 2^{n-\alpha-1} f\left(\frac{a+b}{2}\right)$$

$$(2.7) \quad \leq \frac{f(a)}{n-\alpha+2} + \frac{f(b)}{2(n-\alpha+1)} \left(1 - (i)^{2(n-\alpha+1)}\right)$$

Here $i^2 = -1$ ($i \in \mathbb{C}$ is complex number)

Proof. If $t = 1/2$ is taken in inequality (1.3),

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \frac{1}{2^{n-\alpha}} f(x) + \left| \left(\left(\frac{1}{2}\right)^{n-\alpha} - \left(1 - \frac{1}{2}\right)^{n-\alpha} \right) \right| f(y) \\ &= \frac{1}{2^{n-\alpha}} f(x) + \left| \left(\left(\frac{1}{2}\right)^{n-\alpha} - \left(1 - \frac{1}{2}\right)^{n-\alpha} \right) \right| f(y) \\ &= \frac{1}{2^{n-\alpha}} f(x) + |0| f(y) \\ &\leq \frac{1}{2^{n-\alpha}} f(x) + \frac{1}{2^{n-\alpha}} f(y) \end{aligned}$$

From the first and last terms, we have

$$2^{n-\alpha} f\left(\frac{a+b}{2}\right) \leq f(x) + f(y)$$

□

Now, for all $n - \alpha \in (-1, 1]$, $\frac{1}{2^{n-\alpha}} > 0$, Now let $x = ta + (1-t)b$, $y = (1-t)a + tb$. Then

$$(2.8) \quad 2^{n-\alpha} f\left(\frac{a+b}{2}\right) \leq f(ta + (1-t)b) + f((1-t)a + tb)$$

Multiplying both sides of (2.8) by t and taking integral over $[0, 1]$ with respect to t with the convexity of f :

$$\begin{aligned} 2^{n-\alpha-1} f\left(\frac{a+b}{2}\right) &\leq \int_0^1 t f(ta + (1-t)b) dt + \int_0^1 t f((1-t)a + tb) dt \\ &= \int_0^1 \left\{ \begin{array}{l} t^{n-\alpha+1} f(a) + t \left| (t^{n-\alpha} - (1-t)^{n-\alpha}) \right| f(b) dt \\ + t \left| ((1-t)^{n-\alpha} - t^{n-\alpha}) \right| f(b) dt \end{array} \right\} \\ &\leq \int_0^1 \left\{ t^{n-\alpha+1} f(a) + t |2t-1|^{n-\alpha} f(b) dt \right\} + \int_0^1 t |1-2t|^{n-\alpha} f(b) dt \\ &= \frac{f(a)}{n-\alpha+2} + f(b) \left\{ \begin{array}{l} \int_0^{\frac{1}{2}} t (2t-1)^{n-\alpha} dt + \int_{\frac{1}{2}}^1 t (1-2t)^{n-\alpha} dt \\ + \int_0^{\frac{1}{2}} t (1-2t)^{n-\alpha} dt + \int_{\frac{1}{2}}^1 t (2t-1)^{n-\alpha} dt \end{array} \right\} \\ &= \frac{f(a)}{n-\alpha+2} + f(b) \left\{ \frac{1}{2(n-\alpha+1)} - \frac{(-1)^{n-\alpha+1}}{2(n-\alpha+1)} \right\} \\ &= \frac{f(a)}{n-\alpha+2} + \frac{f(b)}{2(n-\alpha+1)} (1 - (-1)^{n-\alpha+1}) \end{aligned}$$

which completes the proof.

Here we used integrals

$$\begin{aligned} \int_0^{\frac{1}{2}} t (2t-1)^{n-\alpha} dt &= -\frac{1}{4} \left(\frac{(-1)^{n-\alpha+2}}{n-\alpha+2} + \frac{(-1)^{n-\alpha+1}}{n-\alpha+1} \right) \\ \int_{\frac{1}{2}}^1 t (1-2t)^{n-\alpha} dt &= -\frac{1}{4} \left(\frac{(-1)^{n-\alpha+1}}{n-\alpha+1} - \frac{(-1)^{n-\alpha+2}}{n-\alpha+2} \right) \\ \int_0^{\frac{1}{2}} t (1-2t)^{n-\alpha} dt &= \frac{1}{4} \left(\frac{1}{n-\alpha+1} - \frac{1}{n-\alpha+2} \right) \\ \int_{\frac{1}{2}}^1 t (2t-1)^{n-\alpha} dt &= \frac{1}{4} \left(\frac{1}{n-\alpha+2} + \frac{1}{n-\alpha+1} \right) \end{aligned}$$

i) If we take limit for $(n, \alpha) \rightarrow (1^-, 1^-)$ or $(n, \alpha) \rightarrow (0^+, 0^+)$ in (2.6) we have

$$f\left(\frac{a+b}{2}\right) \leq f(a) + \frac{f(b)}{2}$$

ii) If we choose $n - \alpha = \frac{1}{2}$ in (2.6) we have

$$f\left(\frac{a+b}{2}\right) \leq 2\sqrt{2} \left(\frac{f(a)}{5} + \frac{f(b)}{3} (1 + e^{-\frac{\pi}{2}i}) \right)$$

where $i \in C$ is a complex number,

Now, we will write two new lemmas and new results based on them for $(n - \alpha)$ convex functions.

Lemma 1. Let $f : I \subseteq [1, \infty) \rightarrow R^+$ be a differentiable mapping on (y, x) with $y < x$, $f^{(n)} > 0$. If $f^{(n)} \in L[y, x]$, for $0 < n \leq 1$, $0 \leq \alpha < 1$, $t \in [0, 1]$. The following equality for Caputo right-sided derivative holds:

$$\begin{aligned} & \left[\frac{(-1)^n}{(x-y)^2} \Gamma(2) - \frac{(-1)^n}{(x-y)^{(n-\alpha)}} \Gamma(n-\alpha+2) \right] \left(C_{D_{x-}}^\alpha f \right) (u) \\ &= \int_0^1 t (1-t^{n-\alpha}) f^{(n)}(tx + (1-t)y) dt \end{aligned}$$

Here $\left(C_{D_{x-}}^\alpha f \right) (u)$ is as in(1.2) and $\Gamma(r) = \int_0^\infty e^{-t} t^{r-1} dt$, ($r > 0$) is the Gamma function .

Proof. Since $-1 < n-\alpha \leq 1$, $t \in [0, 1]$ or $n-\alpha \in (0, 1]$ with $y < x$, $f^{(n)} > 0$ we have Firstly , by integrating and setting $u = tx + (1-t)y$

$$\begin{aligned} & \int_0^1 t (1-t^{n-\alpha}) f^{(n)}(tx + (1-t)y) dt \\ &= \int_0^1 (t - t^{n-\alpha+1}) f^{(n)}(tx + (1-t)y) dt \quad , \quad (u = tx + (1-t)y) \\ &= \frac{1}{x-y} \int_y^x \left(\frac{u-y}{x-y} - \left(\frac{u-y}{x-y} \right)^{n-\alpha+1} \right) f^{(n)}(u) du \\ &= \frac{1}{(x-y)^2} \int_y^x (u-y) f^{(n)}(u) du \\ & \quad - \frac{1}{(x-y)^{n-\alpha+2}} \int_y^x (u-y)^{n-\alpha+1} f^{(n)}(u) du \\ &= \frac{(-1)^n}{(x-y)^2} \Gamma(2) \left(C_{D_{x-}}^\alpha f \right) (u) \\ & \quad - \frac{(-1)^n}{(x-y)^{n-\alpha+2}} \Gamma(n-\alpha+2) \left(C_{D_{x-}}^\alpha f \right) (u) \\ &= \frac{(-1)^n}{(x-y)^2} \left[\Gamma(2) - \frac{1}{(x-y)^{n-\alpha}} \Gamma(n-\alpha+2) \right] \left(C_{D_{x-}}^\alpha f \right) (u) \end{aligned}$$

□

Lemma 2. Let $f : I \subseteq [1, \infty) \rightarrow R^+$ be a differentiable mapping on (y, x) with $y < x$. If $f^{(n)} \in L[y, x]$, for $0 < n \leq 1$, $0 \leq \alpha < 1$, $t \in [0, 1]$ The following equality for Caputo right-sided derivative holds:

$$(2.9) \quad \frac{(-1)^n}{(x-y)^{n-\alpha+1}} \left[\Gamma(n-\alpha+1) - \frac{1}{x-y} \Gamma(n-\alpha+1) \right] \left(C_{D_{x-}}^\alpha f \right) (u)$$

$$(2.10) = \int_0^1 t^{n-\alpha} (1-t) f^{(n)}(tx + (1-t)y) dt$$

Proof.

$$\begin{aligned}
& \int_0^1 t^{n-\alpha} (1-t) f^{(n)}(tx + (1-t)y) dt \\
= & \int_0^1 (t^{n-\alpha} - t^{n-\alpha+1}) f^{(n)}(tx + (1-t)y) dt \quad , \quad (u = tx + (1-t)y) \\
= & \frac{1}{x-y} \int_y^x \left(\left(\frac{u-y}{x-y} \right)^{n-\alpha} - \left(\frac{u-y}{x-y} \right)^{n-\alpha+1} \right) f^{(n)}(u) du \\
= & \frac{1}{(x-y)^{n-\alpha+1}} \int_y^x (u-y)^{n-\alpha} f^{(n)}(u) du \\
& - \frac{1}{(x-y)^{n-\alpha+2}} \int_y^x (u-y)^{n-\alpha+1} f^{(n)}(u) du \\
= & \frac{(-1)^n}{(x-y)^{n-\alpha+1}} \Gamma(n-\alpha+1) \left(C_{D_{x-}^\alpha} f \right) (u) \\
& - \frac{(-1)^n}{(x-y)^{n-\alpha+2}} \Gamma(n-\alpha+2) \left(C_{D_{x-}^\alpha} f \right) (u) \\
= & \frac{(-1)^n}{(x-y)^{n-\alpha+1}} \left[\Gamma(n-\alpha+1) - \frac{1}{x-y} \Gamma(n-\alpha+1) \right] \left(C_{D_{x-}^\alpha} f \right) (u)
\end{aligned}$$

□

Corollary 2. *Since $0 < n \leq 1$, $0 \leq \alpha < 1$, $t \in [0, 1]$ for $n - \alpha \in (0, 1]$ we have*

$$\begin{aligned}
t & \leq t^{n-\alpha} \implies 1 - t^{n-\alpha} \leq 1 - t \\
& t(1 - t^{n-\alpha}) f^{(n)}(tx + (1-t)y) \\
& \leq t^{n-\alpha} (1-t) f^{(n)}(tx + (1-t)y) \quad ,
\end{aligned}$$

or

$$(2.11) \quad \int_0^1 t(1 - t^{n-\alpha}) f^{(n)}(tx + (1-t)y) dt$$

$$(2.12) \quad \leq \int_0^1 t^{n-\alpha} (1-t) f^{(n)}(tx + (1-t)y) dt$$

From (2.11), that is

$$\begin{aligned}
& \frac{(-1)^n}{(x-y)^2} \left[\Gamma(2) - \frac{1}{(x-y)^{n-\alpha}} \Gamma(n-\alpha+2) \right] \\
\leq & \frac{(-1)^n}{(x-y)^{n-\alpha+1}} \left[\Gamma(n-\alpha+1) - \frac{1}{x-y} \Gamma(n-\alpha+1) \right]
\end{aligned}$$

or

$$\begin{aligned}
& \frac{1}{(x-y)^2} \left[1 - \frac{1}{(x-y)^{n-\alpha}} \Gamma(n-\alpha+2) \right] \\
\leq & \frac{1}{(x-y)^{n-\alpha+1}} \left[1 - \frac{1}{x-y} \right] \Gamma(n-\alpha+1)
\end{aligned}$$

Theorem 3. $0 \leq [\alpha] < 1$, $t \in [0, 1]$, $n = [\alpha] + 1$, If $f : I \subseteq [1, \infty) \rightarrow \mathbb{R}^+$, $I \subset [0, \infty)$, be a differentiable function on I such that $f \in AC^1 L[y, x]$ with $y < x$. If $|f|$ is $(n - \alpha)$ is convex, The following inequalities holds:

$$\begin{aligned}
& \left| \left[\frac{(-1)^n}{(x-y)^2} \Gamma(2) - \frac{(-1)^n}{(x-y)^{(n-\alpha)}} \Gamma(n-\alpha+2) \right] \left(C_{D_x^-}^\alpha f \right) (u) \right| \\
\leq & |f(x)| \left(\frac{1}{n-\alpha+2} - \frac{1}{2(n-\alpha+1)} \right) \\
(2.13) \quad & + \frac{|f(y)|}{4} \left[\beta(n-\alpha+1, 2) - \frac{\beta(n-\alpha+1, n-\alpha+2)}{2^{n-\alpha+2}} \right] \\
& + \frac{|f(y)|}{4} (-1)^{n-\alpha+1} \left[1 - \frac{1}{2^{n-\alpha+1}} \right] \beta(n-\alpha+1, n-\alpha+2)
\end{aligned}$$

Here is $\beta(a, b) = \int_0^1 t^{x-1} (1-t)^{y-1} dt$, $(x, y > 0)$ and $\Gamma(r) = \int_0^\infty e^{-t} t^{r-1} dt$, $(x > 0)$.

Considering Lemma 1 and $(n - \alpha)$ convexity of $|f|$,

$$\begin{aligned}
& \left| \left[\frac{(-1)^n}{(x-y)^2} \Gamma(2) - \frac{(-1)^n}{(x-y)^{(n-\alpha)}} \Gamma(n-\alpha+2) \right] \left(C_{D_x^-}^\alpha f \right) (u) \right| \\
\leq & \left| \int_0^1 t(1-t^{n-\alpha}) f(tx + (1-t)y) dt \right| \\
& \left| \int_0^1 t(1-t^{n-\alpha}) f(tx + (1-t)y) dt \right| \\
\leq & \int_0^1 |t(1-t^{n-\alpha}) f(tx + (1-t)y)| dt \\
= & \int_0^1 t(1-t^{n-\alpha}) |f(tx + (1-t)y)| dt \\
\leq & \int_0^1 t(1-t^{n-\alpha}) \left\{ t^{n-\alpha} |f(x)| + |1-2t|^{n-\alpha} |f(y)| \right\} dt \\
= & |f(x)| \int_0^1 t^{n-\alpha+1} (1-t^{n-\alpha}) dt + |f(y)| \int_0^1 t(1-t^{n-\alpha}) |1-2t|^{n-\alpha} dt \\
& |f(x)| \int_0^1 t^{n-\alpha+1} (1-t^{n-\alpha}) dt + |f(y)| \\
& \times \left\{ \int_0^{\frac{1}{2}} t(1-t^{n-\alpha}) (1-2t) dt + \int_{\frac{1}{2}}^1 t(1-t^{n-\alpha}) (2t-1) dt \right\} \\
= & |f(x)| \left(\frac{1}{n-\alpha+2} - \frac{1}{2(n-\alpha+1)} \right) \\
& + \frac{|f(y)|}{4} \left[\beta(n-\alpha+1, 2) - \frac{\beta(n-\alpha+1, n-\alpha+2)}{2^{n-\alpha+2}} \right]
\end{aligned}$$

$$+ \frac{|f(y)|}{4} (-1)^{n-\alpha+1} \left[1 - \frac{1}{2^{n-\alpha+1}} \right] \beta(n-\alpha+1, n-\alpha+2)$$

Here

$$\begin{aligned} & \int_0^1 t^{n-\alpha+1} (1-t^{n-\alpha}) dt = \frac{1}{n-\alpha+2} - \frac{1}{2(n-\alpha+1)} \\ & \int_0^1 (t-t^{n-\alpha+1}) |1-2t|^{n-\alpha} |f(y)| dt \\ = & \int_0^{\frac{1}{2}} (t-t^{n-\alpha+1}) (1-2t)^{n-\alpha} dt + \int_{\frac{1}{2}}^1 (t-t^{n-\alpha+1}) (2t-1)^{n-\alpha} dt \\ = & \frac{1}{4} \left(\frac{1}{n-\alpha+2} - \frac{1}{n-\alpha+1} \right) - \frac{1}{2^{n-\alpha+2}} \beta(n-\alpha+1, n-\alpha+2) \\ + & \frac{1}{4} \left(\frac{1}{n-\alpha+2} + \frac{1}{n-\alpha+1} \right) - (-1)^{n-\alpha+1} \beta(n-\alpha+1, n-\alpha+2) \\ & \int_0^1 u^{n-\alpha} (u+1)^{n-\alpha} du = (-1)^{n-\alpha+1} \beta(n-\alpha+1, n-\alpha+2), \end{aligned}$$

Here we used the fact that

$$\begin{aligned} & f(tx + (1-t)y) \\ & \leq t^{n-\alpha} f(x) + \left| (t^{n-\alpha} - (1-t)^{n-\alpha}) \right| f(y) \\ & \leq t^{n-\alpha} f(x) + |2t-1|^{n-\alpha} f(y). \end{aligned}$$

Theorem 4. *Let the conditions of Theorem 3 be satisfied with $|f^n|$ is $(n-\alpha)$ is convex. In this case, the following inequality is true.*

$$\begin{aligned} & \left[\frac{(-1)^n}{(x-y)^2} \Gamma(2) - \frac{(-1)^n}{(x-y)^{(n-\alpha)}} \Gamma(n-\alpha+2) \right] \left(C_{D_x^-}^\alpha f \right) (u) \\ & \leq \left| f^{(n)}(y) \right| (\beta((n-\alpha)+1, (n-\alpha)+2) + \beta(2, 2(n-\alpha)+1)) \\ & \quad + \left| f^{(n)}(x) \right| \beta(2, 2(n-\alpha)+1) \end{aligned}$$

Proof. From Lemma 2 and the the properties of Modulus

$$\begin{aligned} & \left| \frac{(-1)^n}{(x-y)^{n-\alpha+1}} \left[\Gamma(n-\alpha+1) - \frac{1}{x-y} \Gamma(n-\alpha+1) \right] \left(C_{D_x^-}^\alpha f \right) (u) \right| \\ = & \left| \int_0^1 t^{n-\alpha} (1-t) f^{(n)}(tx + (1-t)y) dt \right| \\ & \leq \int_0^1 t^{n-\alpha} (1-t) \left| f^{(n)}(tx + (1-t)y) \right| dt \\ & \leq \left| f^{(n)}(x) \right| \int_0^1 t^{2(n-\alpha)} (1-t) dt \\ & \quad + \left| f^{(n)}(y) \right| \int_0^1 t^{n-\alpha} (1-t) \left\{ t^{n-\alpha} + (1-t)^{n-\alpha} \right\} dt \end{aligned}$$

$$\begin{aligned}
&= \left| f^{(n)}(x) \right| \beta(2, 2(n-\alpha) + 1) \\
&\quad + \left| f^{(n)}(y) \right| \left\{ \int_0^1 t^{2(n-\alpha)} (1-t) dt + \int_0^1 t^{(n-\alpha)} (1-t)^{(n-\alpha+1)} dt \right\} \\
&= \left| f^{(n)}(x) \right| \beta(2, 2(n-\alpha) + 1) \\
&\quad + \left| f^{(n)}(y) \right| (\beta(2, 2(n-\alpha) + 1) + \beta((n-\alpha) + 1, (n-\alpha) + 2))
\end{aligned}$$

□

Theorem 5. For all $n-\alpha \in (-1, 1]$ we obtain a general elementary inequality

$$\frac{1}{2} \leq \frac{n-\alpha+3}{(n-\alpha+1)(n-\alpha+2)}$$

Here $0 < n \leq 1$, $0 \leq \alpha < 1$

Proof. Since $t \leq t^{n-\alpha} \implies 1 - t^{n-\alpha} \leq 1 - t$ we can write inequality, for $n-\alpha \in (-1, 1]$ ve $t \in [0, 1]$.

$$t(1 - t^{n-\alpha}) \leq t^{n-\alpha}(1 - t).$$

Now, if we only integrate both sides of the inequality $t(1 - t^{n-\alpha}) \leq t^{n-\alpha}(1 - t)$ over $[0, 1]$, We obtain the desired inequality:

$$\begin{aligned}
\int_0^1 t(1 - t^{n-\alpha}) dt &\leq \int_0^1 t^{n-\alpha}(1 - t) dt \\
\frac{t^2}{2} - \frac{t^{n-\alpha+1}}{n-\alpha+1} \Big|_0^1 &\leq \frac{t^{n-\alpha+1}}{n-\alpha+1} - \frac{t^{n-\alpha+2}}{n-\alpha+2} \Big|_0^1 \\
\frac{1}{2} &\leq \frac{n-\alpha+3}{(n-\alpha+1)(n-\alpha+2)}
\end{aligned}$$

□

3. CONCLUSION

In fact, higher mathematics does not go beyond the study of numbers or number systems in one way or another (e.g. derivative or integral, etc.). One of the three basic properties of real numbers is the order relation is especially important in optimization theory and nonlinear programming. In this context, the convex functions discussed play a leading role. For example, the H-H integral inequality and its variants, well known in the literature, have been obtained through convex functions and classical inequalities. The results obtained above in such that $(n-\alpha)$ -convex and $f \in AC^1L[y, x]$ with $y < x$ are just a few of the infinite number of upper bounds. Researchers interested in the subject can approach the smallest of the upper limits by writing more specific parameters.

REFERENCES

- [1] Muhamet Emin Özdemir, Çetin Yıldız, Ahmet Ocak Akdemir, and Erhan Set, On some inequalities for s-convex functions and applications, Journal of Inequalities and Applications 2013, 2013:333
- [2] M.K. Bakula, M.E. Ozdemir and J.Pečarić, Hadamard type inequalities for m -convex and (α, m) -convex functions, J. Ineq. Pure Appl. Math. 9(4) (2008) Art. 96.
- [3] U.S. Kirmaci a, M.K. Bakula, M.E. Ozdemir J. Pecaric , Hadamard-type inequalities for s-convex functions, Applied Mathematics and Computation 193 (2007) 26-35.

- [4] M. Emin Özdemir and Alper Ekinici, On (α, β) -Convex Function, Eastern Anatolian Journal of Science, Volume VI, Issue II, 2020, 9-14.
- [5] M. Emin Özdemir, New Refinements of Hadamard Integral Inequality via k-Fractional Integrals for P-Convex Function, Turkish Journal of Science, 6-1, 1-5. ISSN: 2587-0971
- [6] M. Emin Özdemir, Hemen Dutta, Ahmet Ocak Akdemir, New estimations for Hadamard inequality via k-Riemann-Liouville fractional integral operators, Mathematics in Engineering, Science and Aerospace (MESA), Vol 11 No 2, pp. 323-332, 2020. CSP-Cambridge, UK, I&S-Florida, USA, 2020.
- [7] M. Emin Özdemir, Çetin Yıldız, New inequalities for n-time differentiable functions, TBILISI MATHEMATICAL JOURNAL, Volume: 12, Issue: 2, Pages: 1-15,
- [8] M. Emin Özdemir, Saad I. Butt, Bahtiyar Bayraktar, Jamshed Nasir, Several integral inequalities for (α, s, m) -convex functions, AIMS Mathematics, 2020, 5(4): 3906-3921. doi: 10.3934/math.2020253.
- [9] M. Emin Özdemir and Ahmet Ocak Akdemir, New Refinements and Integral Inequalities for Concave Functions, TWMS J. And Eng.. Math . V.9, No.1, Special Issue, 2019, pp. 73-83.
- [10] M. Emin Özdemir, Inequalities On Geometrically Convex Functions, Math Sci. Lett.7, No.3, pp. 145-149 (2018).
- [11] M. Emin Ozdemir, Ahmet Ocak Akdemir, Integral Inequalities for Some Convex Functions, Eastern Anatolian Journal of Science, Volume II, Issue I, 1-6.
- [12] M.E.Ozdemir, M. A. Latif and A.O.Akdemir, On Some Hadamard-Type Inequalities for Product of Two Convex Functions On the Co-ordinates, Turkish Journal of Science, Volume I, Issue I, 41-58, 2016.
- [13] M. Emin Özdemir, Merve Avcı-Ardıç and H. Kavurmacı-Önalın, Hermite-Hadamard type inequalities for s-convex and s-concave functions via fractional integrals, Turkish Journal of Science, Vol 1, Issue 1, pages 28-40, 2016.
- [14] M. Emin Özdemir, Ahmet Ocak Akdemir, Erhan Set, On (h-m) Convexity and Hadamard-Type Inequalities, TJMM, 8 (2016), No. 51-58.
- [15] M. Emin Özdemir and A. Ekinici, Generalized Integral Inequalities for Convex Functions, Mathematical Inequalities and Applications (MIA), Volume 19, Number 4 (2016), 1429-1439 (SCI)
- [16] M. E. Özdemir, Some inequalities for the s-Godunova-Levin type functions, Math Sci, DOI 10.1007/s40096-015-0144-y, 17 March 2015.
- [17] M. E. Özdemir, Merve Avci, Ardic, Some Companions of Ostrowski type inequality for functions whose second derivatives are convex and concave with applications, Arab Journal of Mathematical Sciences, Arb J Math Sci 21(1), 53-56. 2015.
- [18] M. E. Özdemir, Havva Kavurmacı Önalın, Merve Avcı, Ardıç, Hermite Hadamard type Inequalities for -Convex Functions, J. Concrete and Applicable Mathematics, Vol.13, 1-2, 96-107, 2015
- [19] M. Emin Özdemir, Erhan Set and Ahmet Ocak Akdemir, On The (r,s) -Convexity and Some Hadamard-Type Inequalities, J. Applied Functional Analysis, Vol, No' S 1-22, 95-100, 2015, Copyright 2015 Eudoxus Press, LLC.
- [20] M. Emin Ozdemir, On Iyengar- Type Inequalities via Quasi-Convexity and Quasi-Concavity(SCI.) Miskolc Mathematical Notes, vol.15 (2014), No.1, pp. 171-181.
- [21] M. Emin Ozdemir, Çetin Yıldız, Mustafa Gürbüz, A note on geometrically convex functions, Journal of Ineq. And Appl., (SCI.) 2014, 2014:180.
- [22] M. Emin Ozdemir, Mustafa Gürbüz, Çetin Yıldız, Inequalities for mappings whose second derivatives are quasi-convex or h-convex functions, Miskolc Mathematics Notes(SCI.) (2014), vol 15, 2014 NO:2 pp. 635-649.
- [23] M. Emin Özdemir, Havva Kavurmacı and Merve Avcı, Some Integral Inequalities For Convex Functions, Tamkang Journal Of Mathematics, Accepted. Vol. : 44, Number 3., 253-259, Autumn 2013, doi. 10.5556/j.tkjm.44.2013.1016.
- [24] M. Emin Özdemir, Çetin Yıldız, New Inequalities for Hermite- Hadamard and Simpson Type with Applications, Tamkang Journal of Mathematics, 44 (2), Page: 209-216, 2013.
- [25] M., Emin Özdemir, Alper Ekinici and Ahmet Ocak Akdemir, Generalization of Integral Inequalities for Functions whose Second Derivatives are Convex and m-Convex, Miskolc Mathematical Notes, vol.13(2012), No.2, pp.441-457(SCI).
- [26] M., E. Özdemir, A. O. Akdemir, Ç. Yıldız, On Co-ordinated Quasi-Convex Functions, Czechoslovak Mathematical Journal, 62(137) (2012), (2012), 889-900 (SCI).

- [27] M. Emin Özdemir, Havva Kavurmacı, Ahmet Ocak Akdemir, Merve Avcı, Inequalities for Convex and Convex Functions on $\Delta = [a, b] \times [c, d]$, Journal of Inequalities and Applications, 2012, 2012:20, doi:10.1186/1029-242X-2012-20, Published: 1 February 2012. (SCI).
- [28] M. Emin Özdemir, Ahmet Ocak Akdemir and Erhan Set On the Ostrowski-Grüss type inequality for twice differentiable functions, Hacettepe Journal of Mathematics and Statistics, 41(5) (2012), 651-655. (SCI).
- [29] M.,Emin Özdemir, E. Set, M. Alomari, Integral inequalities via several kind of convexity Creative Math. Inf. 20(2011), No.1, 62-73.
- [30] Özdemir, M.,E , Kavurmacı Havva, Set Erhan "Ostrowski's Type Inequalities for (α, m) – Convex Functions" Kyungpook Math . J. 50(2010) , 371-378.
- [31] M.,E.Özdemir, Merve Avcı,Erhan Set On Some Inequalities of Hermite- Hadamard Type Via m- Convexity, Applied Mathematics Letters 23 (2010) 1065-1070.,(SCI).
- [32] Özdemir, M.Emin; Kirmaci, Ugur; Ocak, Rahim; Dönmez, Ali, On automorphic and modular forms in the space of the homogeneous polynomials with degree 2l and applications to the special matrix. (English) (SCI), Appl. Math. Comput. 152, No.3, 897-904 (2004).
- [33] Özdemir, M.Emin, A theorem on mappings with bounded derivatives with applications to quadrature rules and means. (English) ,(SCI), Appl. Math. Comput. 138, No.2-3, 425-434 (2003).
- [34] Özdemir, M.Emin; Kirmaci, Ugur, Two new theorems on mappings uniformly continuous and convex with applications to quadrature rules and means. (English) ,(SCI), Appl. Math. Comput. 143, No.2-3, 269-274 (2003).
- [35] U.S. Kirmaci, M. Klaricic Bakula, M. Emin Özdemir, Josip Pecaric, On Some Inequalities For P-Norms, JIPAM(Journal of Inequalities Pure and Apply Mathematics, vol 9 (2008) Issue 1.ISSN:1443-5756.

Email address: eminozdemir@uludag.edu.tr

Current address: Bursa Uludağ University, Department of Mathematics, and Science Education , Bursa-Türkiye