



UNCOUNTABLE INFINITE EXACT SOLUTIONS TO THE FITZHUGH-NAGUMO MODEL

A PREPRINT

 **Shahid Sultan Ali Ramji***
Department of Mathematics
University of Karachi
shahidsultanali@uok.edu.pk

 **Eddy Kwessi**
Trinity University
Texas, U.S.A.

 **Mujahid Abbas**
Government College University
Lahore, Pakistan

July 24, 2024

ABSTRACT

First time in six decades, uncountable infinite exact solutions of FitzHugh-Nagumo model with diffusion have been found. FitzHugh-Nagumo model is a nonlinear dynamical system applicable to neurosciences, chemical kinetics, cell division, population dynamics, electronics, epidemiology, cardiac physiology and pattern formation. Non-classical symmetry analysis has been carried out to find invariant solutions. In many ways this finding makes the corpus of asymptotics and numerics obsolete. Non singular, physically stable and meaningful solutions could now be found without distorting the actual model or introducing forced conditions.

1 Introduction:

Embedded within the cells of nervous system, exists a composition of a special classification termed neurons. This classification of the neurons is a highly significant aspect acting as the main worker units in the nervous system. The neurons are designed in a very unique way in the communication system of the gland cells and muscles thus, enabling them importantly, to act as carriers of information transmission between them. It is very natural to think that any model for neurons would be complicated. FitzHugh-Nagumo model with diffusion (FHND) arose as a special case of the famed Hodgkin-Huxley (Nobel Prize 1963) model. The pioneering work by Nagumo [6] opened several avenues for research in neuroscience. As a result there exists many different variants of FHND. It can be used to model how voltage of neuron is changing as a function of time and space,

$$\begin{aligned}\frac{\partial u}{\partial t} &= D \frac{\partial^2 u}{\partial x^2} - v + g(u) \\ \frac{\partial v}{\partial t} &= \varepsilon[-\beta v + u],\end{aligned}\tag{1}$$

where $g(u) = u - \frac{u^3}{3}$ and $D, \varepsilon,$ and β are real parameters. The holy grail are the integral surfaces $u(x, t)$ and $v(x, t)$. These quantities are known as voltage potential (fast variable) and recovery variable (slow variable) respectively. It is a well-established neurodynamic model. For decades this model has validated itself against empirical data. A cursory Google search produces more than 200,000 results for FHN. According to Web of Science FHN has 2564 primary articles and 50930 primary citations after 1970. Despite of its importance, this original version of FHND has never been cracked exactly before. Although if we either change $g(u)$ slightly or introduce a diffusion term with Laplacian in the second equation or impose some extra conditions on the model then it becomes very easy to work out numerous exact solutions. In the literature, many approaches have been proposed to study its dynamics. In [11], we proved that an exact discrete scheme exist for FitzHugh-Nagumo model. We also worked out discrete steady states for our version of discrete FitzHugh-Nagumo model. Later we identified the problem to carry out stability analysis of FitzHugh-Nagumo with diffusion. In the the same article we found numerous steady states for the original continuous FitzHugh-Nagumo model with diffusion. Most of steady states found were space dependent. As pointed out linear stability analysis was

*Abdus Salam School of Mathematical Sciences

not a reliable tool. Luckily we were also able to find isolated fixed points. Here in this manuscript we presenting the Jacobian for FitzHugh-Nagumo model also.

Isolated fixed points found in [11]:

$$u(t, x) = \frac{\left(\left(4\sqrt{-\frac{4\beta^3-12\beta^2+12\beta-4}{\beta}}\right)\beta^2\right)^{\frac{1}{3}}}{2\beta} + \frac{2(\beta-1)}{\left(\left(4\sqrt{-\frac{4\beta^3-12\beta^2+12\beta-4}{\beta}}\right)\beta^2\right)^{\frac{1}{3}}}, \quad (2)$$

$$v(t, x) = -\frac{1}{\beta} \left(-\frac{\left(\left(4\sqrt{-\frac{4\beta^3-12\beta^2+12\beta-4}{\beta}}\right)\beta^2\right)^{\frac{1}{3}}}{2\beta} \right) + \frac{1}{\beta} \left(\frac{2(\beta-1)}{\left(\left(4\sqrt{-\frac{4\beta^3-12\beta^2+12\beta-4}{\beta}}\right)\beta^2\right)^{\frac{1}{3}}} \right). \quad (3)$$

$$u(t, x) = \frac{\sqrt{3}\sqrt{\beta-1}}{\sqrt{\beta}}, \quad v(t, x) = \frac{\sqrt{3}\sqrt{\beta-1}}{\beta^{\frac{3}{2}}}. \quad (4)$$

$$u(t, x) = -\frac{\sqrt{3}\sqrt{\beta-1}}{\sqrt{\beta}}, \quad v(t, x) = -\frac{\sqrt{3}\sqrt{\beta-1}}{\beta^{\frac{3}{2}}}. \quad (5)$$

$$u(t, x) = 0, \quad v(t, x) = 0. \quad (6)$$

$$u(t, x) = \frac{\left(\sqrt{-\frac{(\beta-1)^3}{\beta}}\beta^2\right)^{\frac{2}{3}} + \beta^2 - \beta}{\left(\sqrt{-\frac{(\beta-1)^3}{\beta}}\beta^2\right)^{\frac{1}{3}}\beta},$$

$$v(t, x) = -\frac{\left(\sqrt{-\frac{(\beta-1)^3}{\beta}}\beta^2\right)^{\frac{2}{3}} + \beta^2 - \beta}{\left(\sqrt{-\frac{(\beta-1)^3}{\beta}}\beta^2\right)^{\frac{1}{3}}\beta}. \quad (7)$$

Space dependent steady states found in [11]

$$u(t, x) = -\frac{\tanh\left(\frac{\sqrt{2}\sqrt{\frac{\beta-1}{D\beta}}(x+x_0)}{2}\right)\sqrt{3}\sqrt{\beta-1}}{\sqrt{\beta}},$$

$$v(t, x) = -\frac{\tanh\left(\frac{\sqrt{2}\sqrt{\frac{\beta-1}{D\beta}}(x+x_0)}{2}\right)\sqrt{3}\sqrt{\beta-1}}{\beta^{\frac{3}{2}}}. \quad (8)$$

$$u(t, x) = \frac{\tanh\left(\frac{\sqrt{2}\sqrt{\frac{\beta-1}{D\beta}}(x+x_0)}{2}\right)\sqrt{3}\sqrt{\beta-1}}{\sqrt{\beta}},$$

$$v(t, x) = \frac{\tanh\left(\frac{\sqrt{2}\sqrt{\frac{\beta-1}{D\beta}}(x+x_0)}{2}\right)\sqrt{3}\sqrt{\beta-1}}{\beta^{\frac{3}{2}}}. \quad (9)$$

Using Similariry Solutions a verified group invariant steady state solution was obtained.

$$u(x, t) = c_2 \sqrt{6} \sqrt{\frac{\beta - 1}{\beta c_2^2 + 5\beta - 6}}$$

$$\text{JacobiSN} \left(\frac{\left(6c_1 D\beta + \sqrt{6} \sqrt{D\beta(-6 + 5\beta)} x \right) \sqrt{6} \sqrt{\frac{\beta - 1}{\beta c_2^2 + 5\beta - 6}}}{6D\beta}, \frac{c_2 \sqrt{5\beta^2 - 6\beta}}{-6 + 5\beta} \right).$$
(10)

There is limited research available on the similarity analysis of FHND, although some articles focus on the conditional symmetry [12] of the fast equation. Slow variable is always neglected. This research carries out a thorough symmetry analysis for the reaction diffusion system (1). Conditional symmetry analysis has been worked out to find invariant solutions. Yet again this investigation establishes the potential of symmetry methods to unravel non-linearity. First time ever in the sixty two years history of this paradigm infinite exact solutions have been found. Annexure collects some exact results obtained. These results are verified here using the ptest function available in Maple's library.

2 Lie Symmetry Analysis

This part will try to serve as an introduction to lie symmetry analysis. The aim is to use symmetries to solve differential equations. Like any other mathematical persuasion, one needs a definition to proceed. Mostly differential equations are solved by changing the variables involved.

$$x^* = x^*(x, y), \quad y^* = y^*(x, y) \quad (11)$$

Now there are at least two ways to view this transformation. One can either geometrically view XY plane as the domain of this mapping and X^*Y^* plane as the codomain. Or one can superimpose XY and X^*Y^* plane one over another with the common origin. So primarily this mapping maps (x, y) to (x^*, y^*) . As goal is to built continuous groups these transformation depends on an arbitrary parameter ϵ ,

$$x^* = x^*(x, y, \epsilon) \quad y^* = y^*(x, y, \epsilon). \quad (12)$$

Definition of a group requires these elements must locally be closed under composition, must contain the identity element and invertible to posses inverses.

$$x^{**} = x^{**}(x^*, y^*, \epsilon^*) = x^{**}(x, y, \epsilon^{**}) \quad (13)$$

$$x^*(x, y, 0) = x; \quad y^*(x, y, 0) = y. \quad (14)$$

$$\det \begin{vmatrix} \frac{\partial x^*}{\partial x} & \frac{\partial x^*}{\partial y} \\ \frac{\partial y^*}{\partial x} & \frac{\partial y^*}{\partial y} \end{vmatrix} \neq 0, \quad (15)$$

There are ways to define lie symmetry. A very useful way [13] is $[X, A] = \lambda A$. To connect differential equations with the operator A consider an example. Let $y = (x, a^\alpha), \alpha = 1, \dots, n$ be the general solution of a differential equation. For the corresponding partial differential equation $Af = 0$, independent solutions $\phi^\alpha, \alpha = 1, \dots, n$ can be obtained by differentiation of the solution set. Differentiation of solution system n times results in a new system of equations. Generalized version of implicit function theorem guarantees that a^α can be made the subject of this system. That is to say a^α can be written explicitly and appearing left side of the system. These n constants are precisely the level surfaces of n linearly independent solutions to the associated pde $Af = 0$, related to the given n th order ode. In particular the general solution of $y'' + y = 0$ is given by

$$y = (\phi_0^1)^{1/2} \sin(x - \phi_0^2). \quad (16)$$

$$\text{So } y' = (\phi_0^1)^{1/2} \cos(x - \phi_0^2), \quad (17)$$

$$y'' = -(\phi_0^1)^{1/2} \sin(x - \phi_0^2). \quad (18)$$

Dividing (16) by (17)

$$\begin{aligned} \tan(x - \phi_0^2) &= y/y', \\ x - \phi_0^2 &= \tan^{-1}(y/y'), \\ \phi_0^2 &= x - \tan^{-1} y/y'. \end{aligned} \quad (19)$$

Squaring and adding (16) and (17)

$$y^2 + y'^2 = \phi_0^1. \quad (20)$$

(19) and (20) are the two linearly independent solutions of

$$\left(\frac{\partial}{\partial x} + y' \frac{\partial}{\partial y} - y \frac{\partial}{\partial y'}\right)f = 0.$$

In fact they are functionally independent. \triangleleft

Given X , a lie symmetry of A i.e. $[X, A] = \lambda A$. Define

$$\hat{X} = X + \mu(x, y, y', \dots, y^{n-1})A$$

then for some $\hat{\lambda}$, $[\hat{X}, A] = \hat{\lambda}A$.

$$\begin{aligned} [\hat{X}, A] &= [X + \mu A, A] \\ &= [X, A] + [\mu A, A] \\ &= \lambda A + (\mu A)A - A(\mu A) \\ &= \lambda A + (\mu A)A - (A\mu)A - \mu(AA) \\ &= (\lambda - A\mu)A \\ &= \hat{\lambda}A \end{aligned}$$

The question is when \hat{X} is going to be a lie symmetry for A ? As already worked in [13], $\lambda = -A\xi$ and ξ is a function x and y . In order for \hat{X} to be a symmetry μ should be independent of the derivatives $y^{(i)}$. On the contrary let us say μ has the term $y^{(n-1)}$ then the presence of $\frac{\partial}{\partial x}$ in the total differential A will throw the $\hat{\lambda}$ into the world of $y^{(n)}$ and hence \hat{X} will not remain a symmetry. \triangleleft

In order to use lie symmetry understanding of this operator A could be very useful. Let us explore more about it. $\hat{A} = \mu(x, y, \dots, y^{n-1})A$ and A are equivalent as their solutions ϕ^a are same. Let us show that for every symmetry X of A , a commutator \hat{A} with X exists. μ is already a yanked out factor of A so obviously anything that satisfies A necessarily satisfies μA . Proof of the statement resides in the following argument.

$$\begin{aligned} [X, \hat{A}] &= [X, \mu A] = X(\mu A) - (\mu A)X \\ &= (X\mu)A + \mu(XA) - \mu(AX) \\ &= (X\mu)A + \mu(XA - AX) \\ &= (X\mu)A + \mu\lambda A \\ &= (X\mu + \mu\lambda)A \end{aligned}$$

Now $X\mu + \mu\lambda = 0$ is a pde whose solution space is non empty, hence for every lie symmetry commutator exists. \triangleleft

As pointed out earlier pragmatism will be the philosophy for this discourse. Hence the best approach would be exemplification. So let us directly integrate the following first order ordinary differential equations by their respective lie symmetries.

$$\begin{aligned} y' &= \frac{-(y+2x)}{x}, \quad X = \frac{1}{y+2x} \frac{\partial}{\partial x} \\ y' &= f(y)g(x), \quad X = \frac{1}{g(x)} \frac{\partial}{\partial x} \end{aligned}$$

For the former the first integral can directly be computed.

$$\begin{aligned} \phi &= \int \frac{dy + \frac{(y+2x)}{x} dx}{-\left(\frac{1}{y+2x}\right) \left(\frac{-(y+2x)}{x}\right)} \\ &= \int x dx + \left(\frac{(y+2x)}{x}\right) x dx \end{aligned}$$

$$\begin{aligned}
 &= xy + x^2 \\
 y &= \frac{c - x^2}{x}
 \end{aligned}$$

As for the second equation, exploiting given symmetry as an integrating factor, conserved quantity, the first integral can directly be given via the following integral.

$$\begin{aligned}
 \phi &= \int \frac{dy - f(y)(g(x))dx}{\frac{1}{g(x)}f(y)g(x)} \\
 \phi &= - \int \frac{dy}{f(y)} + \int g(x)dx
 \end{aligned}$$

◁

Beside finding solutions of different equations, symmetries could be of great help for order reduction. As an example see

$$2y'y''' - 3y''^2 = 0$$

admits the following symmetry

$$X = a \frac{\partial}{\partial x} + b \frac{\partial}{\partial y} + cy \frac{\partial}{\partial y}.$$

It is not difficult to see that differential equation does not depend on y so the given equation has the structure

$$s^n = \hat{\omega}(t, s', s'', \dots, s^{n-1}).$$

Hence it is already an equation of order $n - 1$ for y' . ◁

For the equation $y''' - yy'' + y'^2 = 0$;

$$X = a \frac{\partial}{\partial x} + b \left(x \frac{\partial}{\partial x} - y \frac{\partial}{\partial y} \right)$$

is a lie point symmetry. If we use translation invariance along x part of this two parameter local group. Then we can reduce one order in the following way.

$$\begin{aligned}
 X &= \frac{\partial}{\partial x}. \\
 t &= y, \quad s = x. \\
 y' &= \left(\frac{ds}{dt} \right)^{-1}, \\
 y'' &= \frac{-s''}{(s')^3}, \\
 \frac{d^3s}{dt^3} &= \frac{1}{t_n + t_y y'} \frac{d \left(\frac{ds^2}{dr^2} \right)}{dx} \\
 &= \frac{1}{y'} \frac{d \left(\frac{-y''}{y'^3} \right)}{dx} = \frac{-y'^3 y''' + 3y''^2 y'^2}{y'^7} \\
 s''' &= \frac{-y' y''' + 3y''^2}{y'^5}, \\
 y''' &= \frac{s'''}{s'^4} + \frac{3s''}{s'^2}, \\
 \frac{s'''}{s'^5} - \frac{3s''^2}{s'^6} &= -y' y''', \\
 y'' &= \frac{3s''^2}{s'^5} - \frac{s'''}{s'^4} = \frac{3s''^2 - s' s'''}{s'^5},
 \end{aligned}$$

$$\begin{aligned} \frac{3s''^2 - s's'''}{s'^5} - t \left(\frac{-s''}{s'^3} \right) + \frac{1}{s'} &= 0, \\ 3s''^2 - s's''' + s''s'^2 t + s'^3 &= 0. \end{aligned}$$

◁

As another example not for order reduction but to actually find a solution. Transform the following symmetry

$$X = [a + b(x - y)] \frac{\partial}{\partial x} + a \frac{\partial}{\partial y}$$

with $b = 0$ to coordinates t, s given in

$$t = y, \quad s = \ln(x - y),$$

and then use the generator to integrate

$$s'' + s'^2 + 1 = 0.$$

$$X = a \frac{\partial}{\partial x} + a \frac{\partial}{\partial y}.$$

$$\begin{aligned} X &= X(t) \frac{\partial}{\partial t} + X(s) \frac{\partial}{\partial s} \\ &= a \frac{\partial}{\partial t} \end{aligned}$$

$$\begin{aligned} \phi &= \int \frac{ds' - (-1 - s'^2) dt}{-a(-1 - s'^2)} \\ &= \frac{1}{a} \int \frac{ds'}{1 + s'^2} + \frac{1}{a} \int dt \\ &= \frac{1}{a} \tan^{-1} s' + \frac{1}{a} t, \\ s' &= \tan(a\phi - t), \\ s &= \int \tan(a\phi - t) + C \end{aligned}$$

◁

When the prolonged generators are intransitive in the solution space of differential equation, the integration process fails. Even with these generators, the action of whose group doesn't cover all solution space, differential equations could be solved merely by algebraic manipulation. When the action of these generators divide the solution space into families of different orbits a linear relationship $\mu_1 X_1 + \mu_2 X_2 + \nu A = 0$ between these operators exists. When $\mu_1 X_1 + \mu_2 X_2 + \nu A = 0$ holds then ν, μ_1 and μ_2 can't be constants. Let us assume that such a relationship holds with constants, then one of it being nonzero makes, say

$$X_2 = \Psi X_1 + \nu A. \quad (21)$$

Here Ψ and ν are constants.

$$\begin{aligned} [A, X_2] &= -\lambda_2 A = (A\Psi)X_1 - \Psi\lambda_1 A + (Av)A \\ \Rightarrow \lambda_2 &= \Psi\lambda_1 \\ \Rightarrow [A, X_2] &= \Psi[A, X_1] \\ \Rightarrow X_2 &= \Psi X_1. \end{aligned}$$

Putting X_2 in (21),

$$\nu A = 0.$$

But this is nonsensical. Similarly taking only one as constant it can be shown that if such a linear relation hold then ν, μ_1 and μ_2 can't be constants. ◁

This discussion about connecting operators become unavoidable when one wants to solve a differential equation with G_3IX . For the time being revert to integration process again. If X_1, X_2 form a lie algebra, $\delta_1 = \xi_1 n_1 - \xi_2 n_1 = 0$ holds then show that $y'' = \omega(x, y, y')$ is transformable into $d^2\hat{s}/dt^2 = 0$, admitting eight symmetries.

Case I.

$$\begin{aligned} [X_1, X_2] &= 0, \quad \delta = 0. \\ s'' &= \hat{\omega}(t). \end{aligned}$$

Lets try

$$\begin{aligned} \hat{s}(t) &= s - \int \int^t \hat{\omega}(t') dt' dt + \phi_0 t + \psi_0, \\ \frac{d\hat{s}(t)}{dt} &= \frac{ds}{dt} - \int \hat{\omega}(t) dt + \phi_0, \\ \frac{d^2\hat{s}(t)}{dt^2} &= \frac{d^2s}{dt^2} - \hat{\omega}(t) = 0. \end{aligned}$$

From Chapter 4 Question 2 [13], the last equation is transformable to trivial second order differential equation and they both admit eight symmetries.

Case II.

$$\begin{aligned} [X_1, X_2] &= X_1, \quad \delta = 0. \\ s'' &= s' \hat{\omega}(t). \end{aligned}$$

Lets try

$$\begin{aligned} \hat{s}(t) &= s - \int \phi_0 e^{\hat{\omega}(t') dt'} dt + \phi_0, \\ \frac{d\hat{s}(t)}{dt} &= \frac{ds}{dt} - \phi_0 e^{\int \hat{\omega}(t) dt}, \\ \frac{d^2\hat{s}(t)}{dt^2} &= \frac{d^2s}{dt^2} - \phi_0 e^{\hat{\omega}(t) dt} \cdot \hat{\omega}(t), \\ \frac{d^2s}{dt^2} &- s' \hat{\omega}(t) = 0. \end{aligned}$$

Again this second order homogeneous linear differential equation admits eight symmetries as it can be transformed to $y'' = 0$. \triangleleft

When a second order ordinary differential equation admits G_2 , two independent first integrals can directly be given via line integrals. Hence the solution can be found by eliminating y' from these two first integrals. Below is an example with an abelian lie algebra.

Solve $y'' = y + x^2$ using $X_1 = e^x \frac{\partial}{\partial y}$, $X_2 = e^{-x} \frac{\partial}{\partial y}$.

$$[X_1, X_2] = \left(e^x \left(\frac{\partial}{\partial y} (e^{-x}) - e^{-x} \frac{\partial}{\partial y} (e^x) \right) \frac{\partial}{\partial y} \right) = 0.$$

$$\begin{aligned} \delta &= 0. \\ \eta'_1 &= \frac{\partial \eta_1}{\partial x} + \eta' \left(\frac{\partial \eta_1}{\partial y} - \frac{\partial \xi_1}{\partial x} \right) - y'^2 \frac{\partial \xi_1}{\partial y} = e^x. \\ \eta'_2 &= -e^{-x}. \end{aligned}$$

$$\begin{aligned}
 \phi &= \int \frac{\begin{vmatrix} dx & dy & dy' \\ 1 & y' & y+x^2 \\ 0 & e^x & e^x \end{vmatrix}}{\begin{vmatrix} 1 & y' & y+x \\ 0 & e^x & e^x \\ 0 & e^{-x} & -e^{-x} \end{vmatrix}} \\
 &= \frac{-1}{2} \int (y'e^x - ye^x - x^2e^x) dx + e^x dy + e^x dy' \\
 &= -\frac{1}{2} \int d(y'e^x) - d(ye^x) - x^2e^x dx \\
 &= -\frac{1}{2} (e^x y') + \frac{1}{2} (e^x y) + \frac{1}{2} \int x^2 e^x dx \\
 &= \frac{1}{2} e^x (y - y') + \frac{1}{2} \left[x^2 e^x - \int 2x e^x dx \right] \\
 &= \frac{1}{2} e^x (y - y') + \frac{1}{2} [x^2 e^x - 2x e^x + 2e^x] \\
 \phi &= \frac{1}{2} e^x (y - y' + x^2 + 2x + 2) \\
 \psi &= -\frac{1}{2} \int \begin{vmatrix} dx & dy & dy' \\ 1 & y' & y+x^2 \\ 0 & e^{-x} & -e^{-x} \end{vmatrix} \\
 &= \frac{-1}{2} \int (-e^{-x} y' - ye^{-x} - x^2 e^{-x}) dx + e^{-x} dy + e^{-x} dy' \\
 &= -\frac{1}{2} \int d(y'e^{-x}) + d(ye^{-x}) - x^2 e^{-x} dx \\
 &= -\frac{1}{2} e^{-x} y' - \frac{1}{2} ye^{-x} + \frac{1}{2} \int x^2 e^{-x} dx \\
 &= -\frac{1}{2} e^{-x} (y + y') + \frac{1}{2} (-x^2 e^{-x} - 2x e^{-x} - 2e^{-x}) \\
 \psi &= \frac{-1}{2} e^{-x} (y + y' + x^2 + 2x + 2), \\
 2e^{-x}\psi &= y - y' + x^2 - 2x + 2, \\
 -2e^x\phi &= y + y' + x^2 + 2x + 2, \\
 y &= \phi e^{-x} - \psi e^x - x^2 - 2.
 \end{aligned}$$

Although Lie's integration procedure has led us to the correct solution but observe that the given differential equation is second order non-homogeneous linear. And simpler and faster methods are already available to find the solution. The problem with the theory of differential equations is that it makes mathematics look more of an art than science. Diverse methods each requiring ingenuity of the solver if not guesses. Lie's method could unifies all techniques. It has the potential to be 'one size fits all' or the 'theory of everything' for differential equations. \triangleleft

To see lie's method of integration using normal form work, consider $3yy'' = 5y'^2$. It admits $X_1 = \frac{\partial}{\partial x}$, $X_2 = x \frac{\partial}{\partial x}$, $X_3 = y \frac{\partial}{\partial y}$. Its algorithmic to integrate given differential equation using three symmetries. The given equation has no direct dependence on x hence X_1 , is a symmetry. In fact its not hard to see X_2 is also a symmetry. For the sake of completeness,

$$\begin{aligned}
 \tilde{x} &= \epsilon x, \\
 \tilde{y} &= y. \\
 \frac{d\tilde{y}}{d\tilde{x}} &= \frac{dy}{d(\epsilon x)} = \frac{1}{\epsilon} \frac{dy}{dx},
 \end{aligned}$$

$$\begin{aligned}\frac{d^2\tilde{y}}{d\tilde{x}^2} &= \frac{d}{d\tilde{x}} \left(\frac{1}{\epsilon} \frac{dy}{dx} \right) = \frac{1}{\epsilon^2} \frac{d^2y}{dx^2}, \\ 3y \frac{1}{\epsilon^2} \frac{d^2y}{dx^2} &= 5 \left(\frac{1}{\epsilon} \frac{dy}{dx} \right)^2 \\ \Rightarrow 3yy'' &= 5y'^2;\end{aligned}$$

X_2 is a symmetry. Now take

$$\begin{aligned}\tilde{x} &= x, \\ \tilde{y} &= \epsilon y. \\ \frac{d\tilde{y}}{d\tilde{x}} &= \epsilon \frac{dy}{dx}, \\ \frac{d^2\tilde{y}}{d\tilde{x}^2} &= \epsilon \frac{d^2y}{dx^2}, \\ 3\epsilon y \epsilon y'' &= 5\epsilon^2 y'^2, \\ \Rightarrow 3yy'' &= 5y'^2;\end{aligned}$$

X_3 is a symmetry. Now take $X_1 = \frac{\partial}{\partial x}$, $X_3 = y \frac{\partial}{\partial y}$. These are commuting operators and for them,

$$\Delta = \begin{vmatrix} 1 & y' & \frac{5y'^2}{3y} \\ 1 & 0 & 0 \\ 0 & y & y' \end{vmatrix} = \frac{2y'^2}{3}.$$

Nonzero determinant means the lie algebra generated by X_1 and X_3 acts transitively in the space of first integrals. ϕ, ψ are now guaranteed.

$$\begin{aligned}\phi &= \int \frac{\begin{vmatrix} dx & dy & dy' \\ 1 & y' & \frac{5y'^2}{3y} \\ 1 & 0 & 0 \end{vmatrix}}{\frac{2y'^2}{3}} \\ &= \int \frac{\frac{5y'^2}{3y} dy - y' dy}{\frac{2y'^2}{3}}, \\ \phi &= \frac{5}{2} \ln |y| - \frac{3}{2} \ln |y'|. \\ \phi_0 &= \frac{y'}{y^{5/3}}. \\ \psi &= \int \frac{\begin{vmatrix} dx & dy & dy' \\ 1 & y' & \frac{5y'}{3y} \\ 0 & y & y' \end{vmatrix}}{\frac{2y'^2}{3}} \\ &= \int \frac{3}{2y'^2} \left[\left(y^2 - \frac{5y'^2}{3} \right) dx - y' dy + y dy' \right] \\ &= \frac{3}{2} y'^2 \left(-\frac{2}{3} 5y'^2 x \right) - \frac{3}{2y'^2} y' y - \frac{3y}{2y'}, \\ \psi_0 &= -5x - \frac{3y}{y'}.\end{aligned}$$

$$\begin{aligned}
 -5xy' - 3y &= y'\psi_0, \\
 y' &= \frac{3y}{\psi_0 + 5x}, \\
 \phi_0 y^{5/3} &= \frac{3y}{\psi_0 + 5x}, \\
 y^{2/3} &= \frac{3}{\phi_0(\psi_0 + 5x)}, \\
 y &= \frac{c_1}{(c_2 + 5x)^{3/2}}.
 \end{aligned}$$

◁

In order to solve differential equations almost always some sort of transformations are involved. At the background of Lie's method are finite transformations, these are local group with parameters. All the information about these finite transformations are contained in infinitesimal generators. Next observe that G_3IX

$$\begin{aligned}
 [X_1, X_2] &= X_3, [X_2, X_3] = X_1, \\
 [X_2, X_1] &= X_2,
 \end{aligned}$$

lives inside the projective group

$$\begin{aligned}
 X &= (a_1 + a_2x + a_3y + a_4xy + a_5x^2) \frac{\partial}{\partial x} \\
 &+ (a_6 + a_7x + a_8y + a_5xy + a_4y^2) \frac{\partial}{\partial y}
 \end{aligned}$$

as a subgroup. Before proving this assertion recall that G_3IX can be realized in 2 - space as

$$\begin{aligned}
 X_1 &= \frac{\partial}{\partial x}, \quad X_2 = \cot y \cos x \frac{\partial}{\partial x}, \\
 X_3 &= -\sin x \cot y \frac{\partial}{\partial x} + \cos x \frac{\partial}{\partial y}.
 \end{aligned}$$

The task could be accomplished if inside the projective group a subgroup that walks and talks like this representation could be found. The structural constants here when viewed as an abstract lie algebra is known as $so(3)$, (Gothic Font). $so(3)$ generates $SO(3)$. $so(3)$ can be realized as

$$\begin{aligned}
 X_1 &= y\partial_x - x\partial_y, \\
 X_2 &= \frac{1}{2}(1 + x^2 - y^2)\partial_x + xy\partial_y, \\
 X_3 &= xy\partial_x + \frac{1}{2}(1 - x^2 + y^2)\partial_y
 \end{aligned}$$

which loosely speaking shows its presence inside projective group. One can write the commutator table for projective group and look for a block of this three generators. This is a laborious job writing again and again in different basis. Standing on Stepanhi's shoulders [13] we can write

$$\begin{aligned}
 X_1 &= X_1\phi \frac{\partial}{\partial \phi} + X_1\psi \frac{\partial}{\partial \psi}, \\
 X_2 &= X_2\phi \frac{\partial}{\partial \phi} + X_2\psi \frac{\partial}{\partial \psi}, \\
 X_3 &= X_3\phi \frac{\partial}{\partial \phi} + X_3\psi \frac{\partial}{\partial \psi},
 \end{aligned}$$

as

$$\begin{aligned}
 X_1 &= y \frac{\partial}{\partial x} + (-x) \frac{\partial}{\partial y}, \\
 X_2 &= -xy \frac{\partial}{\partial x} + (-1 - y^2) \frac{\partial}{\partial y}, \\
 X_3 &= (1 + x^2) \frac{\partial}{\partial x} + xy \frac{\partial}{\partial y}.
 \end{aligned}$$

Coming from solution space into the xy plane won't change the commutators. The infinitesimal generators of the lie algebra corresponding is projective group

$$\begin{aligned}\hat{X}_1 &= x^2 \frac{\partial}{\partial x} + xy \frac{\partial}{\partial y}, & \hat{X}_2 &= xy \frac{\partial}{\partial x} + y^2 \frac{\partial}{\partial y}, \\ \hat{X}_3 &= x \frac{\partial}{\partial x}, & \hat{X}_4 &= y \frac{\partial}{\partial x}, & \hat{X}_5 &= \frac{\partial}{\partial x}, \\ \hat{X}_6 &= x \frac{\partial}{\partial y}, & \hat{X}_7 &= y \frac{\partial}{\partial y}, & \hat{X}_8 &= \frac{\partial}{\partial y}. \\ X_1 &= \hat{X}_4 - \hat{X}_6, \\ X_2 &= -\hat{X}_8 - \hat{X}_2, \\ X_3 &= \hat{X}_1 + \hat{X}_5.\end{aligned}$$

Hence $G3IX$ is a subgroup of the projective group. \triangleleft

Coming back again to the basic goal of solving differential equations. Two examples earlier showed that presence of G_2 is enough to solve a second order ordinary differential equation using line integral. A second order ordinary differential equation can admit up to eight symmetries. So for classification the only problem is any algebra that does not have G_2 inside. Only such algebra in the plane is group of rotations. Actually classification of differential equations was a project dear to Lie's heart. Although not for its own sake but for solutions of differential equations.

$y'' = 2(xy' - y)/x^2$ admits

$$\begin{aligned}X_1 &= y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y}, & X_2 &= \frac{-y}{x} \frac{\partial}{\partial x} + \left(x^2 - \frac{y^2}{x^2}\right) \frac{\partial}{\partial y}, \\ X_3 &= (1 + x^2) \frac{\partial}{\partial x} + \left(2xy + \frac{y}{x}\right) \frac{\partial}{\partial y}.\end{aligned}$$

It would be a different problem if inside this group there is no two dimensional subgroup with generators $Y_\alpha = a_\alpha^a X_i, \alpha = 1, 2$. To check this easiest way is to work out the commutator table and see if this is rotation group or not. Lets take a longer route. If one solution $U(x)$ of $\sum_{i=0}^n f_i(x)y^{(i)} = 0$ is known then $X = U(x) \frac{\partial}{\partial x}$ is a symmetry. $U(x) = x$ is already known

$$\begin{aligned}\text{check: } & y = x, \\ & y' = 1, \\ & y'' = 0, \\ & x^2 y'' - 2xy' + 2y = 0, \\ & x^2(0) - 2x(1) + 2(x) = 0, \\ & 0 = 0.\end{aligned}$$



Symmetry has suggested to introduce t, s , so that $Xs = 1, Xt = 0$.

$$\begin{aligned}x \frac{\partial t}{\partial y} &= 0, & x \frac{\partial s}{\partial y} &= 1 \\ \Rightarrow t &= x, & s &= \frac{y}{x}.\end{aligned}$$

Prolonging these coordinates

$$\begin{aligned}\frac{ds}{dt} &= \frac{s_x + s_y y'}{t_x + t_y y'}, \\ s' &= -\frac{y}{x^2} + \frac{y'}{x},\end{aligned}$$

$$\begin{aligned}
 s'' &= \frac{\left(\frac{\partial}{\partial x} + y' \frac{\partial}{\partial y} + y'' \frac{\partial}{\partial y'}\right) \left(-\frac{y}{x^2} + \frac{y'}{x}\right)}{\left(\frac{\partial}{\partial x} + y' \frac{\partial}{\partial y} + y'' \frac{\partial}{\partial y'}\right) x} \\
 &= \frac{2y}{x^3} - \frac{y'}{x^2} - \frac{y'}{x^2} + \frac{y''}{x} \\
 &= \frac{2y}{x^3} - \frac{2y'}{x^2} + \frac{2(xy' - y)}{x^3} \\
 &= \frac{2y}{x^3} - \frac{2y'}{x^2} + \frac{2y'}{x^2} - \frac{2y}{x^3} \\
 &= 0, \\
 s'' &= 0, \\
 s &= at + b, \\
 \frac{y}{x} &= ax + b, \\
 y &= ax^2 + bx.
 \end{aligned}$$


 Figure 1: st plane with random straight line solutions

 Figure 2: xy plane with translated parabolas

The transformation $(x, y) \rightarrow (t, s)$ has straightened the frame of differential equation. Symmetry unraveled that given differential equation is trivial $s'' = 0$. The relation

$$X_1 = \phi(x, y, y')X_2 + \psi(x, y, y')X_3 + v(x, y, y')A$$

could have directly given two first integrals if given symmetries were forming G_3IX .

$$\begin{aligned}
[X_1, X_2] &= \left\{ \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right) \left(-\frac{y}{x} \right) - \left(\left(-\frac{y}{x} \right) \frac{\partial}{\partial x} + \left(x^2 - \frac{y^2}{x^2} \right) \frac{\partial}{\partial y} \right) \right\} \\
&\quad \frac{\partial}{\partial x} + \left\{ \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right) \left(x^2 - \frac{y^2}{x^2} \right) - \left(-\frac{y}{x} \right) \frac{\partial}{\partial x} + \left(x^2 - \frac{y^2}{x^2} \right) \right. \\
&\quad \left. \frac{\partial}{\partial y} (-x) \right\} \frac{\partial}{\partial y} \\
&= \left(1 - x^2 + \frac{2y^2}{x^2} \right) \frac{\partial}{\partial x} + \left(2xy + 2\frac{y^3}{x^3} + \frac{3y}{x} \right) \frac{\partial}{\partial y} \\
&\neq X_3.
\end{aligned}$$

Changing the coordinates

$$\begin{aligned}
X_1(t) &= \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right) x = y = st, \\
X_1(t) &= st, \\
X_1(s) &= \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial x} \right) \left(\frac{y}{x} \right) = -\frac{y^2}{x^2} - 1 \\
&= -(1 + s^2) \\
X_1 &= -(1 + s^2) \frac{\partial}{\partial s} + st \frac{\partial}{\partial t} \\
X_2(s) &= \left(\left(-\frac{y}{x} \right) \frac{\partial}{\partial x} + \left(x^2 - \frac{y^2}{x^2} \right) \frac{\partial}{\partial y} \right) \frac{y}{x} \\
&= \frac{y^2}{x^3} + x - \frac{y^2}{x^3} = t \\
X_2(t) &= \frac{-y}{x} = -s, \\
X_2 &= t \frac{\partial}{\partial s} - s \frac{\partial}{\partial t}. \\
X_3(s) &= \left((1 + x^2) \frac{\partial}{\partial x} + \left(2xy + \frac{y}{x} \right) \frac{\partial}{\partial y} \right) \frac{y}{x} \\
&= -(1 + x^2) \frac{y}{x^2} + 2y + \frac{y}{x^2} \\
&= \frac{-y}{x^2} - y + 2y + \frac{y}{x^2} = y = st \\
X_3(t) &= (1 + x^2) = 1 + t^2, \\
X_3 &= st \frac{\partial}{\partial s} + (1 + t^2) \frac{\partial}{\partial t}.
\end{aligned}$$

Writing the realization of SO_3 in last question

$$\begin{aligned}
X_1 &= y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y}, \\
X_2 &= -xy \frac{\partial}{\partial x} + (-1 - y^2) \frac{\partial}{\partial y}, \\
X_3 &= (1 + x^2) \frac{\partial}{\partial x} + xy \frac{\partial}{\partial y}.
\end{aligned}$$

Comparing with this problem

$$X_1 = -(1 + s^2) \frac{\partial}{\partial s} + st \frac{\partial}{\partial t},$$

$$\begin{aligned} X_2 &= t \frac{\partial}{\partial s} - s \frac{\partial}{\partial t}, \\ X_3 &= st + (1 + t^2) \frac{\partial}{\partial t}. \end{aligned}$$

As the given algebra corresponds to the group of rotations, without integrating, the two first integrals could also be obtained alternatively. Rotation is certainly a symmetry. Being a member of projective group it is going to throw lines to lines (solutions in st -plane). The other two symmetries are hard to visualize but can be checked by $X^n \omega = \eta^n$. \triangleleft

If a chain of derived group terminates at the identity group then such a group is solvable. Easiest example of solvable groups are abelian groups. So G_2 is solvable. To show that let infinitesimal generators X_1 and X_2 be basis vectors for a two dimensional lie algebra. Suppose $[X_1, X_2] = kX_1 + lX_2 = Z$. If k and l are zero then there is nothing to show as L is abelian. If for c_1 and c_2 arbitrary constants; $c_1X_1 + c_2X_2 \in L$ then

$$\begin{aligned} [Z, c_1X_1 + c_2X_2] &= c_1 [Z, X_1] + c_2 [Z, X_2], \\ &= c_1 l [X_2, X_1] + c_2 k [X_1, X_2] \\ &= (c_2 k - c_1 l) Z \end{aligned}$$

hence Z is an ideal of L with dimension one. \triangleleft

To study a connection between differential equations and solvability observe that a solvable G_n is admitted by every linear differential equation of n^{th} order. It is not hard to see that generators occurring as symmetries of linear differential equations are

$$X_N = U_N(x) \frac{\partial}{\partial y}.$$

Here U_N are n linearly independent solutions of the corresponding homogeneous linear differential equations. The symmetries form an abelian group as structure constants are all zero; $[X_N, X_M] = 0$.

Note: Linear differential equations can have more than n symmetries. Take $y'' = 0$ for instance has translation along x axis and other symmetries as well. \triangleleft

Lie's theory works for higher order differential equations even better. To see, solve $y''' = \frac{3}{2} \frac{y''}{y'}$ using first strategy [13].

$$\text{Taking } X_1 = \frac{\partial}{\partial x}; X_2 = x \frac{\partial}{\partial x}.$$

$$[X_1, X_2] = \left[\frac{\partial}{\partial x}, x \frac{\partial}{\partial x} \right] = X_1.$$

$$t = y, \quad s = x.$$

$$X_1 = \frac{\partial}{\partial s},$$

$$X_2 = x \frac{\partial}{\partial x} (y) \frac{\partial}{\partial x} + x \frac{\partial}{\partial x} (x) \frac{\partial}{\partial s} = s \frac{\partial}{\partial s},$$

$$A = \frac{\partial}{\partial t} + s' \frac{\partial}{\partial s} + s'' \frac{\partial}{\partial s'} + (\dots) \frac{\partial}{\partial s''},$$

$$X_2^{(3)} = s \frac{\partial}{\partial s} + s' \frac{\partial}{\partial s'} + s'' \frac{\partial}{\partial s''} + s''' \frac{\partial}{\partial s'''},$$

$$y''' = \frac{3s''^2 - s' s'''}{s'^5}, \quad y'' = \frac{-s''}{s'^3}, \quad y' = \frac{1}{s'},$$

$$\frac{3s''^2 - s' s'''}{s'^5} = \frac{3}{2} \left(\frac{-s''}{s'^3} \right)^2 s',$$

$$s''' = \frac{3}{2} \frac{s''^2}{s'}. \tag{22}$$

$$Y_2 = s' \frac{\partial}{\partial s'} + s'' \frac{\partial}{\partial s''}$$

$$\begin{aligned}
Y_2 u(t, s') &= 1; \quad Y_2 v(t, s') = 0 \\
s' \frac{\partial}{\partial s'} + s'' \frac{\partial}{\partial s''} (u) &= 1 \\
s' \frac{\partial}{\partial s'} + s'' \frac{\partial}{\partial s''} (v) &= 0 \\
u &= \ln s', \quad v = t \\
e^u &= s' \\
s'' &= e^u u' \\
s''' &= e^u u'' + e^u v'^2
\end{aligned}$$

(22) is transformed to

$$\begin{aligned}
e^u u' + e^u u'^2 &= \frac{3}{2} \frac{(e^u u'^2)^2}{e^u} \\
u'' + u'^2 &= \frac{3}{2} u'^2 \\
u'' &= \frac{1}{2} u'^2 \\
\text{let } u' &= w \\
u'' &= w' \\
w' &= \frac{1}{2} w^2 \\
\frac{dw}{w^2} &= \frac{1}{2} dv \\
-\frac{1}{w} &= \frac{1}{2} v + c_1 \\
-2 &= vw + 2c_1 w \\
w &= \frac{-2}{2c_1 + v} \\
\frac{du}{dv} &= \frac{-2}{c'_1 + v} \\
u &= -2 \ln |c'_1 + v| + c_2 \\
\ln s' &= \ln |c'_1 + t|^{-2} c'_2 \\
s' &= \frac{c'_2}{(c'_1 + t)^2} \\
\frac{ds}{dt} &= \frac{c'_2}{(c'_1 + t)^2} \\
s &= \frac{-c'_2}{(c'_1 + t)} + c'_3 \\
x &= \frac{-c'_2}{(c'_1 + y)} + c'_3 \\
(c'_1 + y)x &= -c'_2 + c_3 c'_1 + c'_3 y \\
-c'_3 y + yx &= -c'_1 + c'_3 c'_1 - c'_1 x \\
y &= \phi_1 - \frac{\phi_2}{x - \phi_3}
\end{aligned}$$

So the given differential equation is the equation of rectangular hyperbolas that includes all the isotherms. Notice that two straight quadrature could have solved the given differential equation but goal was to implement first strategy. \triangleleft

As a last example use $X_1 = \frac{\partial}{\partial x}$, $X_2 = \frac{\partial}{\partial y}$, $X_3 = x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y}$, $X_4 = y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y}$ to solve $y''' (1 + y'^2) = (3y'^2 + a) y''$. If the group is not transitive then a linear relation between A and the generators X_N exists, and the coefficients of the X_N in the relation may lead to first integrals. One way is to take the matrix of operators, using Gaussian elimination and method of characteristics, find at least one differential invariant, this differential invariant will act as an orbit. But in the space of (x, y, y', y'') the five operators X_1, X_2, X_3, X_4 and A are always linearly dependent so

$$\begin{aligned}
X_1 &= \phi_1 X_2 + \phi_2 X_3 + \phi_3 X_4 + \nu A \\
\frac{\partial}{\partial x} &= \phi \frac{\partial}{\partial y} + \phi_2 \left(x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} - y'' \frac{\partial}{\partial y''} - 2y''' \frac{\partial}{\partial y'''} \right) + \\
\phi_3 \left(y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} - (1 + y'^2) \frac{\partial}{\partial y'} - 3y' y'' \frac{\partial}{\partial y''} - 3y''^2 - 4y' y''' \frac{\partial}{\partial y'''} \right) + \\
&\quad \nu \left(\frac{\partial}{\partial x} + y' \frac{\partial}{\partial y} + y'' \frac{\partial}{\partial y'} + \omega \frac{\partial}{\partial y''} \right) \\
\det &= \begin{vmatrix} x & y & 1 \\ 0 & -(1 + y'^2) & y'' \\ -y'' & -3y' y'' & \frac{(3y' + a)y''}{(1 + y'^2)} \end{vmatrix} \\
&= x \left(-(3y' + a) y'' + 3y' y'' \right) - y'' (y y'' + (1 + y'^2)) \\
a_{11} &= \begin{vmatrix} y & -x & y' \\ 0 & -(1 + y'^2) & y'' \\ y'' & -3y' y'' & \frac{(3y' + a)y''^2}{1 + y'^2} \end{vmatrix} \\
a_{11} &= y \left(-(3y' + a) y''^2 + 3y' y''^2 \right) - y'' \left(-x y'' + (1 + y'^2) y' \right) \\
\phi_1 &= \frac{y \left(-(3y' + a) y''^2 + 3y' y''^2 \right) - y'' \left(-x y'' + (1 + y'^2) y' \right)}{x \left(-(3y' + a) y'' + 3y' y'' \right) - y'' (y y'' + (1 + y'^2))} \\
a_{12} &= - \begin{vmatrix} 1 & -x & y' \\ 0 & -(1 + y'^2) & y'' \\ 0 & -3y' y'' & \frac{(3y' + a)y''^2}{1 + y'^2} \end{vmatrix} \\
\phi_2 &= \frac{- \left(-(3y' + a) y''^2 + 3y' y''^2 \right)}{x \left(-(3y' + a) y'' + 3y' y'' \right) - y'' (y y'' + (1 + y'^2))} \\
a_{13} &= \begin{vmatrix} 1 & y & y' \\ 0 & 0 & y'' \\ 0 & -y'' & \frac{(3y' + a)y''^2}{(1 + y'^2)} \end{vmatrix} \\
\phi_3 &= \frac{y''^2}{x \left(-(3y' + a) y''^2 + 3y' y''^2 - y'' (y y'' + (1 + y'^2)) \right)}
\end{aligned}$$

◁

This concludes our modest cookbook on how to solve odes using symmetry analysis. For pde see [1]

3 Linearizing FHND

Now that isolated fixed points have been found for CFHND. Stability analysis for solutions near fix points can be carried out easily. To perform the stability analysis one needs to derive the Jacobian for FHND first. Recall [8] that stability for an equilibrium solution for a first order ordinary differential equation $\frac{du}{dt} = f(u)$ can be studied doing the following procedure. Firstly perturbing u about a fix point u_* .

$$u(t) = u_* + \delta(u)(t),$$

$$\frac{d}{dt}(u_* + \delta u) = f(u_* + \delta u),$$

$$\frac{d}{dt}(u_* + \delta u) = \frac{d\delta u}{dt},$$

$$f(u_* + \delta u) \approx f(u_*) + f'(u_*) \delta u,$$

$$\frac{d\delta u}{dt} = f'(u_*),$$

$$\delta u = a e^{\sigma t},$$

$$\sigma = f'(u_*),$$

$$\delta u(t) = a e^{f'(u_*)t},$$

Now if $f'(u_*) < 0$ u_* is linearly stable

If $f'(u_*) > 0$ u_* is linearly unstable

Next given reaction diffusion equation $\partial_t u(t, x) = D \partial_{xx}^2 u + f(u)$, stability of an isolated fix point can be seen repeating a similar idea of perturbing substituting and solving.

$$u(t, x) = u_* + \delta u(t, x),$$

$$\partial_t (u_* + \delta u(t, x)) = \partial_{xx}^2 (u_* + \delta u) + f(u_* + \delta u),$$

$$\partial_t \delta u(t, x) = D \partial_{xx}^2 \delta u + f'(u_*) \delta u,$$

$$\begin{aligned} \delta u(t, x) &= a e^{\sigma t} e^{ikx}, \\ \sigma &= f'(u_k) - Dk^2. \end{aligned}$$

Now for a system of ordinary differential equations

$$\frac{du}{dt} = f_1(u, v); \quad \frac{dv}{dt} = f_2(u, v),$$

the analysis is not different.

$$u(t) = u_* + \delta u(t),$$

$$v(t) = v_* + \delta v(t);$$

$$\frac{d}{dt}[u_* + \delta u(t)] = \frac{d\delta u}{dt} = f_1(u_* + \delta u, v_* + \delta v),$$

$$\frac{d}{dt}[v_* + \delta v(t)] = \frac{d\delta v}{dt} = f_2(u_* + \delta u, v_* + \delta v),$$

$$\frac{d\delta u}{dt} = a_{11} \delta u + a_{12} \delta v,$$

$$\frac{d\delta v}{dt} = a_{21} \delta u + a_{22} \delta v,$$

$$a_{ij} = \frac{\partial f_i}{\partial u_j} |_{(u^*, v^*)},$$

$$\delta u(t) = \delta u_0 e^{\sigma t},$$

$$\delta v(t) = \delta v_0 e^{\sigma t};$$

$$\sigma \delta u_0 = a_{11} \delta u_0 + a_{12} \delta v_0,$$

$$\sigma \delta v_0 = a_{21} \delta u_0 + a_{22} \delta v_0,$$

$$A \delta \vec{u}_0 = \sigma \delta \vec{u}_0.$$

So finally let us gather these ideas to linearize FHND about an isolated fix point.

$$\partial_t u(t, x) = D \partial_{xx}^u u + f_1(u, v),$$

$$\partial_t v(t, x) = f_2(u, v);$$

$$u(t, x) = u_* + \delta u(t, x),$$

$$v(t, x) = v_* + \delta v(t, x);$$

$$\partial_t \delta u(t, x) = D \partial_{xx}^2 \delta u + a_{11} \delta u + a_{12} \delta v,$$

$$\partial_t \delta v(t, x) = a_{21} \delta u + a_{22} \delta v;$$

$$\delta u(t, x) = \delta u_0 e^{\sigma t} e^{ikx},$$

$$\delta v(t, x) = \delta v_0 e^{\sigma t} e^{ikx};$$

$$J = \begin{pmatrix} a_{11} - Dk^2 & a_{12} \\ a_{21} & a_{22} \end{pmatrix}.$$

4 Non-classical symmetry analysis

Qualitative analysis for differential equations mostly become meaningless if exact solution to the system is known. Lie symmetry analysis is one of the most powerful tool available to solve differential equations exactly and systematically. Especially for nonlinear differential equation where no general methods are available.

(1) has two obvious Lie symmetries. The spatio-temporal translation invariance. These two symmetries do not give any differential invariant so they are not enough to run quadrature. G. Bluman [2] extended the idea of Lie symmetries. He introduced this idea of non-classical symmetries. Most of the times it is difficult to tackle a system of differential equations. The system of two equations (1) were merged together and as a result a third order differential equation was obtained:

$$Du_{txx} - u_{tt} + \varepsilon \beta D u_{xx} - \varepsilon \beta u_t - \varepsilon u + u_t (-u^2 + 1) + \varepsilon \beta \left(u - \frac{u^3}{3} \right) = 0. \quad (23)$$

5 Infinite Exact Solutions of FitzHugh-Nagumo Model with Diffusion

When non-classical symmetry analysis (Appendix) for conditional symmetries [12] were carried out on (23), following non-classical symmetries were obtained.

$$Y_1 : \quad \xi_x = 1 \quad (24)$$

$$Y_2 : \quad \xi_t = 1, \xi_x = \sqrt{-D\beta\epsilon} \quad (25)$$

$$Y_3 : \quad \xi_t = 1, \xi_x = -\sqrt{-D\beta\epsilon} \quad (26)$$

$$Y_4 : \quad \xi_t = 1, \xi_x = c \quad (27)$$

$$Y_\infty : \quad \xi_t = 1, \eta_u = A(t, x)u + B(t, x) \quad (28)$$

Using Y_∞ following ansatz (29) together with the constraints (30-33) resulted.

$$u(x, t) = \frac{e^{A^2 F(x)A - B}}{A} \quad (29)$$

$$-\epsilon\beta F(x)^3 A^3 - 3F(x)^3 A^4 = 0 \quad (30)$$

$$3\epsilon\beta F(x)^2 B A^2 + 6F(x)^2 B A^3 = 0 \quad (31)$$

$$3\epsilon\beta D\left(\frac{d^2}{dx^2} F(x)\right)A^3 + 3D\left(\frac{d^2}{dx^2} F(x)\right)A^4 - 3\epsilon\beta F(x)A^4 \\ - 3A^5 F(x) + 3\epsilon\beta F(x)A^3 - 3\epsilon\beta F(x)B^2 A + 3F(x)A^4 \\ - 3\epsilon F(x)A^3 - 3F(x)B^2 A^2 = 0 \quad (32)$$

$$-3\epsilon\beta B A^2 + \epsilon\beta B^3 + 3\epsilon B A^2 = 0 \quad (33)$$

These differential and algebraic constraints produced following results.

$$F(x) = c_1 e^{\frac{\sqrt{A^3\beta\epsilon + A^4 - A^2\beta\epsilon + B^2\beta\epsilon - A^3 + A^2\epsilon + A B^2} x}{A\sqrt{D}\sqrt{\epsilon\beta + A}}} \\ + c_2 e^{-\frac{\sqrt{A^3\beta\epsilon + A^4 - A^2\beta\epsilon + B^2\beta\epsilon - A^3 + A^2\epsilon + A B^2} x}{A\sqrt{D}\sqrt{\epsilon\beta + A}}} \quad (34)$$

$$A = -\frac{\epsilon\beta}{3}, B = 0 \quad (35)$$

$$u(t, x) = e^{-\frac{\epsilon\beta t}{3}} \left(c_1 e^{-\frac{\sqrt{-2\epsilon^4\beta^4 - 6\epsilon^3\beta^3 + 9\epsilon^3\beta^2} x\sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}}} + c_2 e^{\frac{\sqrt{-2\epsilon^4\beta^4 - 6\epsilon^3\beta^3 + 9\epsilon^3\beta^2} x\sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}}} \right) \quad (36)$$

$$k = \frac{\sqrt{-2\epsilon^4\beta^4 - 6\epsilon^3\beta^3 + 9\epsilon^3\beta^2} \sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}} \quad (37)$$

$$u(t, x) = e^{-\frac{\epsilon\beta t}{3}} (c_1 e^{-kx} + c_2 e^{kx}) \quad (38)$$

$$v(t, x) = -\frac{1}{6\beta(e^{\frac{\epsilon\beta t}{3}})^3 (e^{kx})^3} [2(e^{kx})^6 (c_2)^3 \beta + 6(e^{kx})^4 c_1 (c_2)^2 \beta \\ - 9(e^{kx})^4 (e^{\frac{\epsilon\beta t}{3}})^2 c_2 + 6(e^{kx})^2 (c_1)^2 c_2 \beta - 9(e^{kx})^2 (e^{\frac{\epsilon\beta t}{3}})^2 c_1 + 2(c_1)^3 \beta] \quad (39)$$

6 A particular solution of FitzHugh-Nagumo

(36) and (39) to the best of our knowledge is the first ever exact solution to (1). Although the existence of traveling wave (36) was indicated (Fig.3) by numerous numerical and asymptotical works done earlier [7],[10]. The greatest gift symmetry analysis has presented to the age-long FHND is the algorithm from Y_∞ to generate infinite exact solutions for FHND.

$$u(t, x) = e^{-\frac{\epsilon\beta t}{3}} (c_1 e^{-kx} + c_2 e^{kx}) \quad (40)$$

$$k = \frac{\sqrt{-2\epsilon^4\beta^4 - 6\epsilon^3\beta^3 + 9\epsilon^3\beta^2} \sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}} \quad (41)$$




Figure 3: An exact solution (36) for FHND (1); A solution surface $u(t, x)$; $c_1 = 1, c_2 = 1, \epsilon = 0.3, \beta = 2, D = 1.03, c = 0$

7 Appendix

Following is partial calculation for non-classical symmetries of (1). Consider

$$\begin{aligned} u_t &= Du_{xx} - v + g(u) \\ v_t &= \epsilon[-\beta v + c + u] \end{aligned}$$

Writing differential consequence of the fast equation

$$u_{tt} = Du_{xxt} - v_t + g_t(u)$$

Substituting the value of v_t

$$u_{tt} = Du_{xxt} - \epsilon[-\beta v + c + u] + g_t(u)$$

Making v the subject in the first fast FHND

$$v = Du_{xx} - u_t + g(u)$$

Substituting this value in the differential consequence

$$u_{tt} - Du_{xxt} + \epsilon[-\beta(Du_{xx} - u_t + g(u)) + c + u] - g_t(u) = 0$$

FHND has now been transformed into the following third order pde

$$Du_{xxt} - u_{tt} + \epsilon\beta Du_{xx} + g_t(u) - \epsilon\beta u_t + \epsilon\beta g(u) - \epsilon c - \epsilon u = 0$$




Figure 4: An exact solution (39) for FHND (1); A solution surface $v(t, x)$; $c_1 = 1, c_2 = 1, \epsilon = 0.3, \beta = 2, D = 1.03, c = 0$

Consider next the infinitesimal generator for non-classical symmetry and its extension

$$\Gamma = \xi^0 \frac{\partial}{\partial t} + \xi^1 \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial u}$$

And its third extension

$$\Gamma^{(3)} = \Gamma + \rho_x \frac{\partial}{\partial u_x} + \rho_t \frac{\partial}{\partial u_t} + \sigma_{tt} \frac{\partial}{\partial u_{tt}} + \sigma_{tx} \frac{\partial}{\partial u_{tx}} + \sigma_{xx} \frac{\partial}{\partial u_{xx}} + \tau_{xxt} \frac{\partial}{\partial u_{xxt}} + \dots$$

$$D_t \tau_{xxt} - \sigma_{tt} + \epsilon \beta D \sigma_{xx} + \eta g_{tu} - \epsilon \beta \rho_t + \epsilon \beta \eta g_u - \eta \epsilon = 0$$

$$\eta, \rho_t, \sigma_{xx}, \sigma_{tt}, \tau_{xxt}$$

$$\rho_t = D_t(\eta) - u_t D_t(\xi^0) - u_x D_t(\xi^1)$$

$$D_t = \frac{\partial}{\partial t} + u_t \frac{\partial}{\partial u} + u_{tt} \frac{\partial}{\partial u_t} + u_{tx} \frac{\partial}{\partial u_x}$$

$$\rho_t = \eta_t + u_t \eta_u - u_t \xi_t^0 - u_t^2 \xi_t^0 - u_x \xi_t^1 - u_t u_x \xi_u^1$$

$$\sigma_{tt} = D_t(\rho_t) - u_{tt} D_t(\xi^0) - u_{tx} D_t(\xi^1)$$

$$\begin{aligned} \sigma_{tt} = & \eta_{tt} + u_t \eta_{ut} - u_t \xi_t - u_t \xi_t - 2u_t \xi_{tt} - 2u_t \xi_t - u_x \xi_{tt} + \eta u_{tt} \\ & - u_{tt} \xi_u - u_{tx} \xi_x - u_x \xi_{xt} + u_t \eta_{tt} - u_x \xi_t - u_t \xi_x - u_{tx} \xi_u + u_{tx} \xi - u_{tx} \xi - u_{tt} \xi \\ & - u_x^2 \xi_u - u_x^2 \xi_t - u_{tt} \xi_x - u_x \xi_t - 2u_t \xi_{tx} - u_{tx} \eta_x - u_x^2 \eta - u_t \xi - u_{tx} \eta_{tx} + \eta_t - u_t \xi_u \end{aligned}$$

$$\begin{aligned}
& -u_x^2 \eta_u - 2u_t \xi_u - u_t \xi_x - u_{tx} u_x \xi - 2u_{tx} u_x \xi_u - 2u_{tx} u_x \xi_x - u_{tx} \eta_{tx} - u_{tx} \xi_t - u_t u_x \xi_u \\
& -u_t u_x \xi_t - 2u_{tx} \eta_x - u_t \xi_{tx} - 2u_t \xi_t - u_x^2 \xi_u - 2u_{tx} \xi_x - u_x \eta_u - u_x \xi_t - u_{tx} \xi_u - u_{tx} \xi \\
& -u_{tx} \eta_u - u_x^2 \xi_t - u_x \xi_t - u_x \xi_x - u_t \xi - u_{tx} \eta_x - u_x \xi_t - u_x \xi_t - u_t \xi_{tt} - u_{tx} \xi_t - u_x \eta_u \\
& \quad -u_t \eta_x - u_{tx} \xi_t - u_x \xi_t - u_x^2 \xi - u_{tx} \xi_t - u_{tx} \xi_u
\end{aligned}$$

$$\sigma_{xx} = D_x(\rho_x) - u_{tx} D_x(\xi^0) - u_{xx} D_x(\xi^1)$$

$$\begin{aligned}
\rho_x &= D_x(\eta) - u_t D_x(\xi^0) - u_x D_x(\xi^1) \\
&= \eta_x + u_x \eta_u - u_t \xi_x^0 - u_t u_x \xi_u^0 - u_x \xi_x^1 - u_x^2 \xi_u^1
\end{aligned}$$

$$\begin{aligned}
\sigma_{xx} &= \eta_{xx} + u_x \eta_{ux} - u_t \xi_{xx}^0 - u_t u_x \xi_{ux}^0 \\
&\quad - u_x \xi_{xxx}^1 - u_x^2 \xi_{ux}^1 + u_x [\eta_u + u_x \eta_{ux} - u_t \xi_x^0 - u_x \xi_u^1] \\
&\quad - 2u_x \xi_{ux} - u_{tx} [\xi_x^0 + u_x \xi_u^0] - u_{xx} [\xi_x^1 + u_x \xi_u^1]
\end{aligned}$$

$$\begin{aligned}
\tau_{xxt} &= D_t(\sigma_{xx}) - u_{tx} D_t(\xi^0) - u_{xxt} D_t(\xi^1) \\
&= D_x(\sigma_{tx}) - u_{tx} D_x(\xi^0) - u_{xxt} D_x(\xi^1) \\
&= \eta_{xxt} + u_t \eta_{uxu} + u_x \eta_{uxu} + u_t u_x \eta_{uxu} + u_x u_x \eta_{uxu} \\
&\quad - u_t \xi_{xxt} - u_t u_x \xi_{uxxt} - u_t^2 \xi_{uxt} - u_t u_{xx} \xi_{uxu} - u_t u_x \xi_{uxt} \\
&\quad - u_t u_x \xi_{uxu} - u_t u_{xx} \xi_{uxu} - u_x u_t \xi_{uxu} - u_x u_t \xi_{uxu} \\
&\quad - u_{xxt} [\xi^0 + u_t \xi_u^0] - u_{xxt} [\xi^1 + u_x \xi_u^1] - u_{tx}^2 \xi_{xx}^1 - u_t^2 \xi_{uu} \\
&\quad - u_t u_x \xi_{uu} - u_t \xi_u^1 - u_t \xi_u^1 - 2u_t u_{xx} \xi_{uu} + u_x \eta_{xu} \\
&\quad + u_t u_x \eta_{ut} + u_{xx} \eta_{uu} + u_{tx} \eta_{ut} + u_x \eta_{uu} + u_x^2 \eta_{uu} \\
&\quad - u_{tx} u_x \xi_{ux} + u_{tx} \eta_u + \eta_{tx} \xi_u - u_{tx} \xi_{tx} + u_t \eta_{uu} \\
&\quad + \eta_{tx} \eta_u + \eta_{tx}^2 \xi_{tx} + u_x \eta_{xx} + u_t \eta_{tx} + u_t \eta_{ut} - u_{tt} \eta_u \\
&\quad - u_{tx} \eta_t + u_x^2 \eta_{uu} + u_t u_{tx} \eta_u + u_{tx} \eta_u + \eta_{tx} \eta_u - u_{tx} \xi_{tx} \\
&\quad + u_t \eta_{ut} - u_t u_x \xi_{uu} + u_{xx} \eta_{uu} + u_x^2 \eta_{ut} - u_x^3 \xi_{ut} + u_x \eta_{xx} \\
&\quad + \eta_{tx} \xi_{tt} + u_x^2 \eta_{uu} - 2u_t u_{xx} \xi_{tx} + u_x \eta_{xx} + u_t u_{xx} \eta_{ut} \\
&\quad + u_{xx} \eta_{ut} + \eta_{tx} \xi_u - 2u_t u_{tx} \xi_{uu} - u_{tx} u_x \xi_{tx} + u_x^2 \eta_{tx} \\
&\quad + u_{tx} \eta_u - u_{tx} \xi_{tx} - u_{tx} u_x \eta_{ut} + u_t u_{tx} \xi_{ut} + u_x^2 \eta_{uu} + u_t \eta_{tx} \\
&\quad - u_{tt} \eta_u - 2u_t u_x \xi_{uu} - u_x u_{xx} \eta_{ut} + \eta_{tx} \eta_{tx} + u_t^2 \eta_{ut} \\
&\quad + u_x \eta_{tx} + u_t u_{xx} \eta_{ut} + u_x^2 \eta_{tx} - u_x u_{xx} \eta_{ut} + u_{tx} \eta_u + u_x \eta_{xx} \\
&\quad - u_{tx} u_x \eta_{ut} + u_{tx} \eta_u + u_{tx} \eta_u - u_t u_{xx} \xi_{ut} + u_t \eta_{xx} + u_{tx} \eta_{ut} \\
&\quad + u_{tx} \eta_u - u_{tx} u_x \xi_{ut} - u_x u_{xx} \eta_{ut} + \eta_{tx} \xi_u - u_t u_x \xi_{ut} - u_x^2 \eta_{tx} \\
&\quad + u_t u_{tx} \eta_u + u_t^2 \eta_{tx} + u_x \eta_{xx} - u_t u_{xx} \eta_{ut} + u_x \eta_{xx} - u_t u_x \xi_{ut} \\
&\quad - u_{tx} u_x \eta_{ut} + \eta_{tx} \xi_u + u_t^2 \eta_{ut} + u_{tx} \eta_u + u_x^2 \eta_{tx} + u_x \eta_{xx} \\
&\quad - u_{tx} \xi_{tx} - u_x u_{xx} \eta_{ut} - u_{xx} \xi_{xt}^1 - u_t u_x \xi_{xt}^1 - u_{tx} \xi_{xt}^1 - u_x u_x \xi_{xt}^1 \\
&\quad - u_t u_x u_x \xi_{ut}^0 - u_x u_x \xi_t^1 - u_{tx} u_x \xi_u^1 - u_{txx} \xi_u^1 - u_{tx} \xi_t^1 - u_t u_{txx} \xi_u^1 \\
&\quad + D u_{xxt} + D u_x \nu_{xt} + D u_x \nu_{xt} + D u_x \nu_{xt} + D u_x \nu_{xt} + D u_{tx} \nu_{xt} \\
&\quad - D u_t \xi_{ut}^0 - D u_t \xi_{ut}^0 - D u_t \xi_{ut}^0 - D u_x u_{tx} \xi_{ut}^0 - D u_x \xi_{ut}^0 - D u_{tx} u_t \xi_u^1 \\
&\quad - D u_{tx} \xi_{ut}^1 - D u_x \xi_{ut}^1 - D u_{tx} u_x \xi_t^1 - D u_t \xi_u^1 - D u_x \xi_{xt}^1 - D u_t u_x \xi_{xt}^1 \\
&\quad - D u_t^2 \xi_u^1 - D u_t u_{xt} \xi_{ut} - 2D u_t u_{tx} \xi_u^1 + D u_x \nu_{xt} + D u_{xt} \nu_{xt} \\
&\quad + D u_{tx} u_{xt} + D u_t \xi_{xt} + D u_{xt} u_x + 2D u_{tx} u_{xt} \nu_{xt} - D u_x \xi_{ut}^1 \\
&\quad - D u_{xt} \xi_{ut}^1 - D u_{tx} \xi_u^1 - D u_t u_{tx} \xi_{ut}^0 - D u_{tx} u_x - D u_{tx} \xi_{ut} - D u_t u_{tx} \xi_{ut}^1 \\
&\quad - D u_t u_x \xi_{ut} - D u_{tx} u_t \xi_{ut} - 2D u_{tx} u_x \xi_t - D u_{tx} \xi_t - D u_t u_{tx} \xi_t
\end{aligned}$$

$$\begin{aligned}
& -Du_{tx}u_{tx}\xi_u^1 - Du_tu_{tx} - Du_tu_x\xi_{xt} - Du_{tx}\xi_u - Du_tu_{tx}\xi_{ut} \\
& -Du_{tx}u_t\xi_{ut}^1 - Du_{tx}u_x\xi_{ut} - Du_{tx}\xi_t^0 - Du_{tx}\xi_x - \eta_t - u_t\nu_t + u_t\xi_{tt}^1 \\
& + u_t\xi_{tt}^1 - Du_{tt}\xi_x^0 - Du_xu_{tx}\xi_{ut}^0 - Du_tu_{tx}\xi_{xt}^1 - Du_{tx}^2\xi_t^1 \\
& -Du_{tt}\xi_t^1 - Du_{tt}u_x\xi_n^1 + Du_{txx}u_t + Du_{tx}u_{tt}u_t \\
& + Du_{tx}\nu_t - Du_{tx}u_t\xi_t^1 - Du_{txx}\xi_t^1 - Du_{tt}u_x\xi_{xt}^1 - Du_{tx}u_x\xi_{xt}^1 \\
& -Du_{tx}\xi_{xt}^1 - 2Du_{tx}u_x\xi_{xt}^1 - Du_{txx}\xi_t^1 - 2Du_{tx}u_x\xi_t^1 - 2Du_{tt}u_x\xi_{xt}^1 \\
& -Du_{txx}\xi_{xt}^1 - Du_{tx}\xi_t^1 - Du_{tt}\xi_t^1 - Du_{txx}\xi_{xt}^1 \\
& -Du_{txx}\xi_t^1 - Du_{tx}u_x\xi_{xt}^1 - Du_{tt}u_x\xi_t^1 - Du_{tx}\xi_t^1 - Du_{txx}u_x\xi_t^1 \\
& -Du_{tx}\xi_{xt}^1 - Du_{txx}u_t - Du_{tx}\xi_t^1 - Du_{txx}\xi_t^1 - Du_{tx}\xi_{xt}^1 \\
& -Du_{tx}u_x\xi_{xt}^1 - Du_{txx}u_x\xi_t^1 - Du_{tx}\xi_{xt}^1 - Du_{txx}\xi_{xt}^1 \\
& -Du_{tx}\xi_t^1 - Du_{txx}\xi_t^1 - Du_{tx}\xi_{xt}^1 - Du_{txx}u_x\xi_{xt}^1 \\
& -Du_{tx}u_x\xi_{xt}^1 - Du_{tx}\xi_t^1 - Du_{txx}\xi_t^1 - Du_{tx}\xi_{xt}^1 - Du_{txx}\xi_t^1 \\
& -Du_{tx}\xi_{xt}^1 - Du_{txx}u_x\xi_{xt}^1 - Du_{tx}u_x\xi_{xt}^1 - Du_{tx}\xi_t^1 - Du_{txx}\xi_t^1 \\
& -Du_{tx}\xi_{xt}^1 - Du_{txx}\xi_t^1 - \eta_t - u_t\nu_t + u_t\xi_t^0 + u_t\xi_{tt}^1 \\
& + u_tu_x\xi_{ut}^1 - u_t\eta_u - u_{tt}\eta_u + u_x^2\xi_t + u_t^2\xi_u + u_tu_x\xi_u + u_x^2\xi_{uu} \\
& -\eta_uu_{tt} + u_{tt}\xi_t + u_{tt}u_x\xi_u + u_t\xi_t + u_tu_{xx}\xi_{tt} + \epsilon\beta D\eta_{ux} + \epsilon\beta\eta_u \\
& -\epsilon\beta Du_t\xi_uu_x - \epsilon\beta Du_t\xi_{ux} - \epsilon\beta Du_t^2\xi_tu_{xx} - \epsilon\beta Du_x\xi_t \\
& -\epsilon\beta Du_x^2\xi_{tt} - \epsilon\beta Du_{xx}u_x\xi_t - \epsilon\beta Du_t^2\xi_{tt}u_x - \epsilon\beta Du_t\xi_t \\
& -\epsilon\beta Du_x\xi_{tt} - \epsilon\beta Du_xu_{xx}\xi_u - \epsilon\beta Du_xu_{tt}\xi_t - \epsilon\beta Du_t\xi_x \\
& -\epsilon\beta Du_x\xi_{tt} - \epsilon\beta Du_xu_t\xi_u - \epsilon\beta Du_xu_{xx}\xi_t - \epsilon\beta Du_x\xi_u \\
& +\epsilon\beta Du_x\xi_u + \epsilon\beta Du_xu_t\xi_t + \epsilon\beta Du_x\xi_{tt} + \epsilon\beta Du_xu_x\xi_t \\
& +\epsilon\beta Du_t\xi_t + \epsilon\beta u_x\xi_u + \epsilon\beta Du_x\xi_u - \epsilon\eta = 0
\end{aligned}$$

$$\begin{aligned}
\text{Const} & : \quad D\eta_{xxt} - \eta_{tt} + \epsilon\beta D\eta_{xuu} + \xi_{uu} - u_{xx}\xi \\
& \quad -\epsilon\beta\xi_{ut} + \epsilon\beta\eta_{uu} - \epsilon\eta = 0 \\
u_t & : \quad D\eta_{uxu} - D\xi_{xuu} - \eta_{uuu} + \xi_{tt} - \eta_{ut} + \xi \\
& \quad -2\eta_{uu} + \xi_{ut} - u_x^2\eta + u_t^2\xi + \xi_t - \epsilon\beta D\xi_{xxu} \\
& \quad -\epsilon\beta\eta = 0 \\
u_x & : \quad D\eta_{xuu} - D\xi_{xuu} + D\eta_{xxu} + \xi_t + \xi_t u_x^2 \\
& \quad +\epsilon\beta\eta_{uu} - \epsilon\beta\xi_{xuu} + \epsilon\beta\xi_{ut} = 0 \\
u_t u_x & : \quad D\eta_{xuu} - D\xi_{xuu} - \eta_{xuu} \\
& \quad -\xi_{ut} + \epsilon\beta\eta_{xuu} - \epsilon\beta\xi_u = 0 \\
u_x^2 & : \quad D\eta_{uuu} - D\eta_{uux} + D\eta_{xxu} - D\eta_{uux} \\
& \quad -\xi_{ux} + \epsilon\beta\xi_{xuu} - \epsilon\beta\xi_{xuu} = 0 \\
u_t u_x^2 & : \quad -\xi_{ux} + \xi_{ux}u_t^2 + \epsilon\beta\eta_{uu} - D\xi_{xuu} \\
& \quad -\eta_{uuu} + \epsilon\beta\xi_{uu} - \epsilon\beta D\xi_u = 0 \\
u_{tt} & : \quad D\xi_{uux} - \eta_{uu} + \xi_t + \xi_t = 0 \\
u_t^2 u_x & : \quad \xi_{uu} - D\xi_{xuu} - D\xi_{xuuu} - D\xi_{uuxu} = 0 \\
u_{tt} u_x & : \quad -D\xi_{ux}^0 - D\xi_{ux} + \xi_{uu} = 0
\end{aligned}$$

$$\begin{aligned}
 u_{tx}u_t & : & -D\xi_{ux}^o - D\xi_{uxu} - D\xi_{uxu} &= 0 \\
 u_x^2 & : & D\eta_{uut} - 2D\xi_{xuu} - \epsilon\beta D\xi_{ux}^1 \\
 & & + \epsilon\beta D\eta_{uuu} - \epsilon\beta D\xi_{xu} &= 0 \\
 u_t u_x^2 & : & -D\xi_{ux}^1 + D\eta_{uu} - D\xi_{uut} \\
 & & - D\xi_{xuu} - \epsilon\beta D\xi_{uu}^o &= 0 \\
 u_{tx}u_{xx} & : & -2D\xi_{ux}^1 + 2D\eta_{uuu} - 2D\xi_{xuu} \\
 & & - D\xi_{ut} - \epsilon\beta D\xi_u - \epsilon\beta D\xi_u &= 0 \\
 u_x^2 & : & -D\xi_{uuu}^o &= 0 \\
 u_t u_x^2 & : & -D\xi_{uu}^o &= 0 \\
 u_x^3 & : & -D\xi_{uu}^1 - \epsilon\beta D\xi_{uux} &= 0 \\
 u_t u_x^3 & : & -D\xi_{uuu} &= 0 \\
 u_x^2 u_{tx} & : & -3D\xi_{uu}^1 &= 0 \\
 u_t u_x & : & -D\xi_x^o - D\xi_x^o &= 0 \\
 u_t u_{xx} u_{tx} & : & -2D\xi_{uu}^1 - D\xi_{xu}^o - D\xi_{uu}^o &= 0 \\
 u_t^2 u & : & -D\xi_u^o - D\xi_u^o &= 0 \\
 u_{tx} u_x & : & -D\xi_u^o - D\xi_u^o &= 0 \\
 u_{xx} & : & D\eta_{tt} - D\xi_{xt}^1 - D\xi_{xt}^1 + \epsilon\beta D\eta_u - \epsilon 3D\xi_x^1 - \epsilon\beta D\xi_u^1 &= 0 \\
 u_t u_{xxt} & : & -\epsilon\beta D\xi_u^o + D\eta_{uu} - D\xi_{ut}^o - D\xi_{xu}^1 - D\xi_{xu}^1 &= 0 \\
 u_t^2 u_{xx} & : & -D\xi_{uu}^o &= 0 \\
 u_{xx} & : & D\eta_u - D\xi_x^1 - D\xi_x^1 - D\xi_t^o &= 0 \\
 u_{tx} u_{xx} & : & -D\xi_u^o &= 0 \\
 u_{txx} u_t & : & -D\xi_u^o - D\xi_u^o &= 0 \\
 u_{xx} u_x u_x & : & -2D\xi_{ut}^1 - D\xi_{ut}^1 - 2\epsilon\beta D\xi_u^1 &= 0 \\
 u_t u_x u_{xx} u_{xx} & : & -2D\xi_{uu}^1 - D\xi_{uu}^1 - 0 &= 0 \\
 u_{xx} u_{xt} & : & -D\xi_u^1 &= 0 \\
 u_{tx} u_{xx} & : & -2D\xi_u^1 - D\xi_u^1 &= 0 \\
 u_{xx} u_{xt} & : & -D\xi_t^1 &= 0 \\
 u_{xxt} u_t & : & -D\xi_u^1 &= 0
 \end{aligned}$$

$$u_t u_{tt} \quad : \quad +2\xi_u^o + \xi_u^o = 0$$

$$u_t^3 \quad : \quad \xi_{uu}^o = 0$$

$$\xi^o = E^o(t, x, u)$$

$$= E^o(t, x)$$

$$= E^o(t)$$

$$\xi^1 = E^1(t, x, u)$$

$$= E^1(t, x)$$

$$= E^1(x)$$

$$D\eta_{xxt} - \eta_{tt} + \epsilon\beta D\eta_{xx} + \eta g_{tu} + \epsilon\beta\eta_t + \epsilon\beta\eta g_u - \epsilon\eta = 0$$

$$D\eta_{xxu} - \eta_{ut} + \xi_{tt}^o - \eta_{tu} - \epsilon\beta\eta_u + \epsilon\beta\xi_t^o = 0$$

$$2D\eta_{uxt} + 2\epsilon\beta D\eta_{ux} - \epsilon\beta D\xi_{xx}^1 = 0$$

$$D\eta_{ux} - D\xi_{xx}^1 + D\eta_{xu} = 0$$

$$-\eta_u + 2\xi_t^o = 0$$

$$\eta_{uu} = 0$$

[9]

> #First Ever Exact Sol of FHND

$$\begin{aligned} > FHND := \left[\left(\frac{\partial}{\partial t} u(t, x) \right) = D \cdot \left(\frac{\partial^2}{\partial x \partial x} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u^3(t, x)}{3}, \left(\frac{\partial}{\partial t} v(t, x) \right) \right. \\ & \left. = \epsilon \cdot (-\beta \cdot v(t, x) + u(t, x)) \right] \end{aligned}$$

$$FHND := \left[\frac{\partial}{\partial t} u(t, x) = D \left(\frac{\partial^2}{\partial x^2} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u(t, x)^3}{3}, \frac{\partial}{\partial t} v(t, x) = \epsilon (-\beta v(t, x) + u(t, x)) \right] \quad \text{(Check1)}$$

$$\begin{aligned} > Sol := \left\{ u(t, x) = e^{-\frac{\epsilon \beta t}{3}} \left(c_{-1} e^{-\frac{\sqrt{-2\epsilon^4 \beta^4 - 6\epsilon^3 \beta^3 + 9\epsilon^3 \beta^2} x \sqrt{6}}{6\epsilon \beta \sqrt{D} \sqrt{\epsilon \beta}}} \right. \right. \\ & \left. \left. + c_{-2} e^{\frac{\sqrt{-2\epsilon^4 \beta^4 - 6\epsilon^3 \beta^3 + 9\epsilon^3 \beta^2} x \sqrt{6}}{6\epsilon \beta \sqrt{D} \sqrt{\epsilon \beta}}} \right), v(t, x) = \right. \end{aligned}$$

$$\begin{aligned} & \left. - \frac{1}{6\beta \left(e^{\frac{\epsilon \beta t}{3}} \right)^3 \left(e^{\frac{\sqrt{-\epsilon^3 \beta^2 (2\beta^2 \epsilon + 6\beta - 9) x \sqrt{6}}{6\epsilon \beta \sqrt{D} \sqrt{\epsilon \beta}}} \right)^3 \left(e^{\frac{\sqrt{-\epsilon^3 \beta^2 (2\beta^2 \epsilon + 6\beta - 9) x \sqrt{6}}{6\epsilon \beta \sqrt{D} \sqrt{\epsilon \beta}}} \right)^6 \right.} \\ & \cdot (c_{-2})^3 \beta + 6 \left(e^{\frac{\sqrt{-\epsilon^3 \beta^2 (2\beta^2 \epsilon + 6\beta - 9) x \sqrt{6}}{6\epsilon \beta \sqrt{D} \sqrt{\epsilon \beta}}} \right)^4 c_{-1} \cdot (c_{-2})^2 \beta \\ & - 9 \left(e^{\frac{\sqrt{-\epsilon^3 \beta^2 (2\beta^2 \epsilon + 6\beta - 9) x \sqrt{6}}{6\epsilon \beta \sqrt{D} \sqrt{\epsilon \beta}}} \right)^4 \left(e^{\frac{\epsilon \beta t}{3}} \right)^2 c_{-2} + 6 \left(e^{\frac{\sqrt{-\epsilon^3 \beta^2 (2\beta^2 \epsilon + 6\beta - 9) x \sqrt{6}}{6\epsilon \beta \sqrt{D} \sqrt{\epsilon \beta}}} \right)^2 \\ & \left. \cdot (c_{-1})^2 c_{-2} \beta - 9 \left(e^{\frac{\sqrt{-\epsilon^3 \beta^2 (2\beta^2 \epsilon + 6\beta - 9) x \sqrt{6}}{6\epsilon \beta \sqrt{D} \sqrt{\epsilon \beta}}} \right)^2 \left(e^{\frac{\epsilon \beta t}{3}} \right)^2 c_{-1} + 2 \cdot (c_{-1})^3 \beta \right\} \end{aligned}$$

$$Sol := \left\{ u(t, x) = e^{-\frac{\epsilon \beta t}{3}} \left(c_{-1} e^{-\frac{\sqrt{-2\epsilon^4 \beta^4 - 6\epsilon^3 \beta^3 + 9\epsilon^3 \beta^2} x \sqrt{6}}{6\epsilon \beta \sqrt{D} \sqrt{\epsilon \beta}}} \right. \right. \quad \text{(Check2)}$$

$$\begin{aligned}
& + c_2 e^{\frac{\sqrt{-2\epsilon^4\beta^4 - 6\epsilon^3\beta^3 + 9\epsilon^3\beta^2} x\sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}}} \Big), v(t, x) = \\
& - \frac{1}{6\beta \left(e^{\frac{\epsilon\beta t}{3}} \right)^3 \left(e^{\frac{\sqrt{-\epsilon^3\beta^2(2\beta^2\epsilon + 6\beta - 9)} x\sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}}} \right)^3} \left(2 \left(e^{\frac{\sqrt{-\epsilon^3\beta^2(2\beta^2\epsilon + 6\beta - 9)} x\sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}}} \right)^6 \right. \\
& c_2^3 \beta + 6 \left(e^{\frac{\sqrt{-\epsilon^3\beta^2(2\beta^2\epsilon + 6\beta - 9)} x\sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}}} \right)^4 c_1 c_2^2 \beta \\
& - 9 \left(e^{\frac{\sqrt{-\epsilon^3\beta^2(2\beta^2\epsilon + 6\beta - 9)} x\sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}}} \right)^4 \left(e^{\frac{\epsilon\beta t}{3}} \right)^2 c_2 \\
& + 6 \left(e^{\frac{\sqrt{-\epsilon^3\beta^2(2\beta^2\epsilon + 6\beta - 9)} x\sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}}} \right)^2 c_1^2 c_2 \beta \\
& \left. - 9 \left(e^{\frac{\sqrt{-\epsilon^3\beta^2(2\beta^2\epsilon + 6\beta - 9)} x\sqrt{6}}{6\epsilon\beta\sqrt{D}\sqrt{\epsilon\beta}}} \right)^2 \left(e^{\frac{\epsilon\beta t}{3}} \right)^2 c_1 + 2 c_1^3 \beta \right) \Big\}
\end{aligned}$$

>

These answers can be tests by passing them to odetest as a list.

> pdetest(Sol, FHND)

[0, 0]

(Check3)

>

> #Steady State Verification

$$\begin{aligned} > FHND := & \left[\left(\frac{\partial}{\partial t} u(t, x) \right) = D \cdot \left(\frac{\partial^2}{\partial x \partial x} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u^3(t, x)}{3}, \left(\frac{\partial}{\partial t} v(t, x) \right) \right. \\ & \left. = \epsilon \cdot (-\beta \cdot v(t, x) + u(t, x)) \right] \end{aligned}$$

$$FHND := \left[\frac{\partial}{\partial t} u(t, x) = D \left(\frac{\partial^2}{\partial x^2} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u(t, x)^3}{3}, \frac{\partial}{\partial t} v(t, x) = \epsilon \left(-\beta v(t, x) + u(t, x) \right) \right] \quad \text{(Check1)}$$

$$> SS := \begin{cases} u(t, x) \end{cases}$$

$$\begin{aligned} & = -C2 \sqrt{6} \sqrt{\frac{\beta - 1}{\beta - C2^2 + 5\beta - 6}} \text{JacobiSN} \left(\frac{1}{6 D \beta} \left((6 - C1 D \beta \right. \right. \\ & \left. \left. + \sqrt{6} \sqrt{D \beta (-6 + 5\beta)} x \right) \sqrt{6} \sqrt{\frac{\beta - 1}{\beta - C2^2 + 5\beta - 6}} \right), \frac{-C2 \sqrt{5\beta^2 - 6\beta}}{-6 + 5\beta} \right), v(t, \end{aligned}$$

x)

$$\begin{aligned} & = \frac{1}{\beta} \left(-C2 \sqrt{6} \sqrt{\frac{\beta - 1}{\beta - C2^2 + 5\beta - 6}} \text{JacobiSN} \left(\frac{1}{6 D \beta} \left((6 - C1 D \beta \right. \right. \right. \\ & \left. \left. \left. + \sqrt{6} \sqrt{D \beta (-6 + 5\beta)} x \right) \sqrt{6} \sqrt{\frac{\beta - 1}{\beta - C2^2 + 5\beta - 6}} \right), \frac{-C2 \sqrt{5\beta^2 - 6\beta}}{-6 + 5\beta} \right) \right) \end{aligned}$$

$$SS := \begin{cases} u(t, x) \end{cases}$$

(Check2)

$$= -C2 \sqrt{6} \sqrt{\frac{\beta - 1}{\beta - C2^2 + 5\beta - 6}} \text{JacobiSN} \left(\frac{1}{6 D \beta} \left((6 - C1 D \beta \right. \right.$$

$$+ \sqrt{6} \sqrt{D \beta (-6 + 5 \beta)} x) \sqrt{6} \sqrt{\frac{\beta - 1}{\beta - C2^2 + 5 \beta - 6}}, \frac{-C2 \sqrt{5 \beta^2 - 6 \beta}}{-6 + 5 \beta} \Big), v(t,$$

x)

$$= \frac{1}{\beta} \left(-C2 \sqrt{6} \sqrt{\frac{\beta - 1}{\beta - C2^2 + 5 \beta - 6}} \text{JacobiSN} \left(\frac{1}{6 D \beta} \left((6 - C1 D \beta \right. \right. \right. \\ \left. \left. \left. + \sqrt{6} \sqrt{D \beta (-6 + 5 \beta)} x) \sqrt{6} \sqrt{\frac{\beta - 1}{\beta - C2^2 + 5 \beta - 6}}, \frac{-C2 \sqrt{5 \beta^2 - 6 \beta}}{-6 + 5 \beta} \right) \right) \right) \Bigg\}$$

>

These answers can be tests by passing them to odetest as a list.

> pdetest(SS, FHND)

[0, 0]

(Check3)

>

```

> #Steady State Verification
> FHND := [ ( (∂/∂t) u(t, x) ) = D · ( (∂²/∂x ∂x) u(t, x) ) - v(t, x) + u(t, x) - (u³(t, x)/3), (∂/∂t) v(t, x) )
           = ε · ( -β · v(t, x) + u(t, x) ) ]
FHND := [ (∂/∂t) u(t, x) = D ( (∂²/∂x²) u(t, x) ) - v(t, x) + u(t, x) - (u(t, x)³/3), (∂/∂t) v(t, x) = ε ( (Check1)
           -β v(t, x) + u(t, x) ) ]
> SS := { u(t, x) = (√3 √(β-1) / √β), v(t, x) = (√3 √(β-1) / β√β) }
           SS := { u(t, x) = (√3 √(β-1) / √β), v(t, x) = (√3 √(β-1) / β³/²) } (Check2)
>
These answers can be tests by passing them to odetest as a list.
> pdetest(SS, FHND)
           [0, 0] (Check3)
>

```

```

> #Steady State Verification
> FHND := [ ( (∂/∂t) u(t, x) ) = D · ( (∂²/∂x ∂x) u(t, x) ) - v(t, x) + u(t, x) - u³(t, x)/3, (∂/∂t) v(t, x) )
           = ε · ( -β · v(t, x) + u(t, x) ) ]
FHND := [ (∂/∂t) u(t, x) = D ( (∂²/∂x²) u(t, x) ) - v(t, x) + u(t, x) - u(t, x)³/3, (∂/∂t) v(t, x) = ε ( (Check1)
           -β v(t, x) + u(t, x) ) ]
> SS := { u(t, x) = -√3 √(β-1) / √β, v(t, x) = -√3 √(β-1) / (β√β) }
           SS := { u(t, x) = -√3 √(β-1) / √β, v(t, x) = -√3 √(β-1) / β³/² } (Check2)
>
These answers can be tests by passing them to odetest as a list.
> pdetest(SS, FHND)
           [0, 0] (Check3)
>

```

> #Steady State Verification

$$\begin{aligned} > FHND := & \left[\left(\frac{\partial}{\partial t} u(t, x) \right) = D \cdot \left(\frac{\partial^2}{\partial x \partial x} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u^3(t, x)}{3}, \left(\frac{\partial}{\partial t} v(t, x) \right) \right. \\ & \left. = \epsilon \cdot (-\beta \cdot v(t, x) + u(t, x)) \right] \end{aligned}$$

$$FHND := \left[\frac{\partial}{\partial t} u(t, x) = D \left(\frac{\partial^2}{\partial x^2} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u(t, x)^3}{3}, \frac{\partial}{\partial t} v(t, x) = \epsilon (-\beta v(t, x) + u(t, x)) \right] \quad \text{(Check1)}$$

$$\begin{aligned} > SS := & \left\{ \begin{aligned} u(t, x) = & - \frac{\tanh\left(\frac{1}{2} \sqrt{2} \sqrt{\frac{\beta-1}{D\beta}} (x+x0)\right) \sqrt{3} \sqrt{\beta-1}}{\sqrt{\beta}}, v(t, x) = \\ & - \frac{\tanh\left(\frac{1}{2} \sqrt{2} \sqrt{\frac{\beta-1}{D\beta}} (x+x0)\right) \sqrt{3} \sqrt{\beta-1}}{\beta \sqrt{\beta}} \end{aligned} \right\} \end{aligned}$$

$$\begin{aligned} SS := & \left\{ \begin{aligned} u(t, x) = & - \frac{\tanh\left(\frac{\sqrt{2} \sqrt{\frac{\beta-1}{D\beta}} (x+x0)}{2}\right) \sqrt{3} \sqrt{\beta-1}}{\sqrt{\beta}}, v(t, x) = \\ & - \frac{\tanh\left(\frac{\sqrt{2} \sqrt{\frac{\beta-1}{D\beta}} (x+x0)}{2}\right) \sqrt{3} \sqrt{\beta-1}}{\beta^{3/2}} \end{aligned} \right\} \quad \text{(Check2)} \end{aligned}$$

>

These answers can be tests by passing them to odetest as a list.

$$\begin{aligned} > pdetest(SS, FHND) & \quad \quad \quad [0, 0] \quad \quad \quad \text{(Check3)} \end{aligned}$$

>

> #Steady State Verification

$$\begin{aligned} > FHND := \left[\left(\frac{\partial}{\partial t} u(t, x) \right) = D \cdot \left(\frac{\partial^2}{\partial x \partial x} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u^3(t, x)}{3}, \left(\frac{\partial}{\partial t} v(t, x) \right) \right. \\ & \quad \left. = \epsilon \cdot (-\beta \cdot v(t, x) + u(t, x)) \right] \end{aligned}$$

$$FHND := \left[\frac{\partial}{\partial t} u(t, x) = D \left(\frac{\partial^2}{\partial x^2} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u(t, x)^3}{3}, \frac{\partial}{\partial t} v(t, x) = \epsilon \left(-\beta v(t, x) + u(t, x) \right) \right] \quad \text{(Check1)}$$

$$\begin{aligned} > SS := & \left\{ u(t, x) = \frac{\tanh\left(\frac{1}{2} \sqrt{2} \sqrt{\frac{\beta-1}{D\beta}} (x+x_0)\right) \sqrt{3} \sqrt{\beta-1}}{\sqrt{\beta}}, v(t, x) \right. \\ & \left. = \frac{\tanh\left(\frac{1}{2} \sqrt{2} \sqrt{\frac{\beta-1}{D\beta}} (x+x_0)\right) \sqrt{3} \sqrt{\beta-1}}{\beta \sqrt{\beta}} \right\} \end{aligned}$$

$$\begin{aligned} SS := & \left\{ u(t, x) = \frac{\tanh\left(\frac{\sqrt{2} \sqrt{\frac{\beta-1}{D\beta}} (x+x_0)}{2}\right) \sqrt{3} \sqrt{\beta-1}}{\sqrt{\beta}}, v(t, x) \right. \\ & \left. = \frac{\tanh\left(\frac{\sqrt{2} \sqrt{\frac{\beta-1}{D\beta}} (x+x_0)}{2}\right) \sqrt{3} \sqrt{\beta-1}}{\beta^{3/2}} \right\} \quad \text{(Check2)} \end{aligned}$$

> These answers can be tests by passing them to odetest as a list.

$$\begin{aligned} > pdetest(SS, FHND) \\ & \quad \quad \quad [0, 0] \quad \quad \quad \text{(Check3)} \end{aligned}$$

>

> #Eq. Pt. Verification

$$\begin{aligned} > FHND := \left[\left(\frac{\partial}{\partial t} u(t, x) \right) = D \cdot \left(\frac{\partial^2}{\partial x \partial x} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u^3(t, x)}{3}, \left(\frac{\partial}{\partial t} v(t, x) \right) \right. \\ & \quad \left. = \epsilon \cdot (-\beta \cdot v(t, x) + u(t, x)) \right] \end{aligned}$$

$$FHND := \left[\frac{\partial}{\partial t} u(t, x) = D \left(\frac{\partial^2}{\partial x^2} u(t, x) \right) - v(t, x) + u(t, x) - \frac{u(t, x)^3}{3}, \frac{\partial}{\partial t} v(t, x) = \epsilon (\textbf{Check1}) \right. \\ \left. -\beta v(t, x) + u(t, x) \right]$$

$$\begin{aligned} > FP := \left\{ \begin{aligned} u(t, x) &= \frac{\left(\sqrt{-\frac{(\beta-1)^3}{\beta} \beta^2} \right)^{2/3} + \beta^2 - \beta}{\left(\sqrt{-\frac{(\beta-1)^3}{\beta} \beta^2} \right)^{1/3} \beta}, v(t, x) = \\ &= \frac{\left(\sqrt{-\frac{(\beta-1)^3}{\beta} \beta^2} \right)^{2/3} + \beta^2 - \beta}{\left(\sqrt{-\frac{(\beta-1)^3}{\beta} \beta^2} \right)^{1/3} \beta} \end{aligned} \right\} \end{aligned}$$

$$\begin{aligned} FP := \left\{ \begin{aligned} u(t, x) &= \frac{\left(\sqrt{-\frac{(\beta-1)^3}{\beta} \beta^2} \right)^{2/3} + \beta^2 - \beta}{\left(\sqrt{-\frac{(\beta-1)^3}{\beta} \beta^2} \right)^{1/3} \beta}, v(t, x) \\ &= \frac{\left(\sqrt{-\frac{(\beta-1)^3}{\beta} \beta^2} \right)^{2/3} + \beta^2 - \beta}{\left(\sqrt{-\frac{(\beta-1)^3}{\beta} \beta^2} \right)^{1/3} \beta^2} \end{aligned} \right\} \end{aligned}$$

(Check2)

>

These answers can be tests by passing them to odetest as a list.

$$> pdetest(FP, FHND)$$

(Check3)



[0, 0]

(Check3)

Resource Availability Statement

The following resources are available on request from the first author: Hand written calculations for non-classical symmetry for FitzHugh-Nagumo with diffusion; Matlab code for FitzHugh-Nagumo with/without diffusion; Hand calculations of related system using symmetry analysis [4]; Conditional symmetry calculations for locally and non-locally related systems of FitzHugh-Nagumo with diffusion [5]; Maple code for exp and Riccati methods of traveling wave solutions; and Remaining steady state solutions of FitzHugh- Nagumo model.

Acknowledgements

The first author expresses his heartfelt gratitude to Prof. Dr. Asghar Qadir, who proselytized him into symmetries.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- [1] Bluman, G.W. and Cheviakov, A.F. and Anco, S.: Applications of Symmetry Methods to Partial Differential Equations, Springer New York, (2009)
- [2] Bluman George W. Construction of Solutions to Partial Differential Equations by the Use of Transformation Groups, Ph.D. Thesis, Caltech (1967)
- [3] Bluman George W., de la Rosa Rafael, Santos Bruzón María and Luz Gandarias María, Differential invariant method for seeking nonlocally related systems and nonlocal symmetries. I: General theory and examples, Proc. R. Soc., (2021)
- [4] George W. Bluman, Zhengzheng Yang, A symmetry-based method for constructing nonlocally related partial differential equation systems, J. Math. Phys. 1 September (2013)
- [5] Cherniha, R., Davydovych, V., Conditional symmetries and exact solutions of nonlinear reaction-diffusion systems with non-constant diffusivities, Communications in Nonlinear Science and Numerical Simulation, 17 (8), 3177-3188 (2012)
- [6] Nagumo J., Arimoto S., and Yoshizawa S. An active pulse transmission line simulating nerve axon, Proc. IRE. 50:2061–2070, (1962).
- [7] John Rinzel, Joseph B. Keller, Traveling Wave Solutions of a Nerve Conduction Equation, Biophysical Journal, Volume 13, Issue 12, (1973)
- [8] Roussel, Marc R. Nonlinear Dynamics: A Hands-On Introductory Survey. United States, Morgan & Claypool Publishers, (2019)
- [9] Tarcísio M. Rocha Filho, Annibal Figueiredo, [SADE] a Maple package for the symmetry analysis of differential equations, Computer Physics Communications, Volume 182, Issue 2, (2011)
- [10] Schiesser, W. E., Griffiths, G. Traveling Wave Analysis of Partial Differential Equations: Numerical and Analytical Methods with Matlab and Maple, Netherlands: Elsevier Science, (2010)
- [11] Shahid, Abbas M, Kwessi E. Nonstandard Nearly Exact Analysis of the FitzHugh–Nagumo Model. Symmetry. 2024; 16(5):585. <https://doi.org/10.3390/sym16050585>
- [12] Shih M, Momoniat E and Mahomed F M, Approximate conditional symmetries and approximate solutions of the perturbed FitzHugh-Nagumo equation, J. Math. Phys. 46, 1, 1-10, (2005)
- [13] Stephani, H. and MacCallum, M.A.H., Differential Equations: Their Solution Using Symmetries, Cambridge University Press, (1989)