

# A CONVERSE TO A THEOREM OF GAUSS ON GAUSS SUMS

JONATHAN W. BOBER AND LEO GOLDBAKHER

ABSTRACT. In this note we prove (under mild hypotheses) that  $f$  is a character of  $\mathbb{F}_p^\times$  if and only if the Fourier transform of  $f$  has magnitude 1 somewhere in  $\mathbb{F}_p^\times$ . This implies a converse to a theorem of Gauss on the magnitude of the Gauss sum, in addition to other consequences.

## 1. INTRODUCTION

Consider an arbitrary function  $f : \mathbb{F}_p^\times \rightarrow \mathbb{C}^\times$ . A famous result (essentially due to Gauss) is that the quantity

$$\tau(f) := \sum_{n \in \mathbb{F}_p^\times} f(n) e\left(\frac{n}{p}\right)$$

has magnitude  $\sqrt{p}$  whenever  $f$  is a nontrivial character mod  $p$  (here and throughout,  $e(\alpha) := e^{2\pi i\alpha}$ ). Does the converse hold? Our goal is to prove that it does, under some mild assumptions about  $f$ . In the special case that  $f$  produces a string of  $\pm 1$ 's, our approach gives a particularly clean result:

**Proposition 1.1.** *Suppose the image of  $f : \mathbb{F}_p^\times \rightarrow \mathbb{C}^\times$  is  $\{\pm 1\}$ , and that  $f(1) = 1$ . Then  $f$  is a character (namely, the Legendre symbol) if and only if  $|\tau(f)| = \sqrt{p}$ .*

Thus we can verify an algebraic condition (that  $f$  is a homomorphism) via an analytic one (that  $f$  correlates in a special way with exponentials).

There are two obvious paths to generalizing Proposition 1.1. The first is to weaken the hypotheses, but we cannot be too cavalier—for  $f$  to be a character it is necessary that  $f(1) = 1$  and that the image of  $f$  consists of roots of unity. The second natural generalization is to interpret the quantity  $\tau(f)$  in terms of the finite Fourier transform of  $f$ , and to attempt to replace our condition on  $\tau(f)$  by a condition on  $\widehat{f}$ . To be more precise, given any  $f : \mathbb{F}_p^\times \rightarrow \mathbb{C}^\times$  we can extend it to a function on  $\mathbb{F}_p$  by setting  $f(0) := 0$ , and then define its Fourier transform  $\widehat{f} : \mathbb{F}_p \rightarrow \mathbb{C}$  by

$$\widehat{f}(\xi) := \frac{1}{\sqrt{p}} \sum_{x \in \mathbb{F}_p} f(x) e\left(-\frac{x\xi}{p}\right).$$

In this language, the conclusion of Proposition 1.1 asserts that  $f$  is the Legendre symbol (mod  $p$ ) if and only if  $|\widehat{f}(-1)| = 1$ . Does this hold if we replace  $-1$  by some other input? Our main theorem, which generalizes Proposition 1.1 in both the ways we just described, answers this question in the affirmative:

**Theorem 1.2.** *Suppose the image of  $f : \mathbb{F}_p^\times \rightarrow \mathbb{C}^\times$  consists of  $n^{\text{th}}$  roots of unity, that  $f(1) = 1$ , and that  $p \nmid n$ . Then  $f$  is a nontrivial character if and only if  $\exists a \in \mathbb{F}_p^\times$  such that  $|\widehat{f}(a)| = 1$ .*

*Remark.* The divisibility hypothesis  $p \nmid n$  is necessary for two reasons. First, when  $p \mid n$  it's possible for a non-homomorphism to have Gauss sum of magnitude  $\sqrt{p}$ : taking  $p = 3$  and setting  $f(1) = 1$  and  $f(2) = e(\frac{5}{6})$  yields  $|\tau(f)| = \sqrt{3}$ . Second, if  $n$  is chosen to be minimal in Theorem 1.2, then  $p \mid n$  implies that  $f$  is not a character. To see this, observe that if  $f$  were a character then  $n$  would be the order of  $f$  as an element of the dual group  $\widehat{\mathbb{F}_p^\times}$ . This would imply  $n \mid p - 1$ , whence  $p \nmid n$ .

If  $f$  is a nontrivial character, a standard change of variables in  $\tau(f)$  implies  $|\widehat{f}(a)| = 1$  for every  $a \in \mathbb{F}_p^\times$ . This yields the following amusing result, which we don't know how to prove without using Theorem 1.2:

**Corollary 1.3.** *Suppose the image of  $f : \mathbb{F}_p^\times \rightarrow \mathbb{C}^\times$  consists of  $n^{\text{th}}$  roots of unity, and that  $p \nmid n$ . Then either  $|\widehat{f}(a)| = 1$  for all  $a \in \mathbb{F}_p^\times$ , or  $|\widehat{f}(a)| \neq 1$  for all  $a \in \mathbb{F}_p^\times$ .*

We conclude our introduction with a brief survey of some other characterizations of characters. Recall one of the standard definitions: a function  $f : \mathbb{Z} \rightarrow \mathbb{C}$  (or  $\mathbb{N} \rightarrow \mathbb{C}$ ) is a Dirichlet character (mod  $q$ ) if and only if

- $f$  is completely multiplicative,
- $f$  is periodic with period  $q$ , and
- the support of  $f$  is the set of integers coprime to  $q$ .

It turns out there are various ways to modify these hypotheses without including any other functions. One example of this is the following result due to Allouche and the second author [1], which built on previous work of Sárközy [12], Heppner-Maxsein [5], and Methfessel [11]:

**Proposition 1.4** (Allouche-Goldmakher). *A function  $f : \mathbb{Z} \rightarrow \mathbb{C}$  is a Dirichlet character if and only if  $f$  is completely multiplicative, eventually satisfies a linear recurrence, and has support strictly larger than  $\{\pm 1\}$ .*

There are also a number of related characterizations that use the notion of automaticity in place of linear recurrences; see [6] and [8] for a representative example.

A different genre of characterization, discovered recently by Konieczny [9], starts with the observation that for any Dirichlet character  $\chi$  and any integer  $a \geq 0$ , the function  $\chi(n)n^a$  agrees with the outputs of a *generalized polynomial*, i.e. any function that can be built out of polynomials, addition, multiplication, and the floor function.

**Theorem 1.5** (Konieczny). *Suppose  $f : \mathbb{N} \rightarrow \mathbb{C}$  is a completely multiplicative function that coincides with a generalized polynomial, and that  $f$  has support strictly larger than  $\{1\}$ . Then there exists a Dirichlet character  $\chi$  and an integer  $a \geq 0$  such that  $f(n) = \chi(n)n^a$ .*

A third type of characterization is in terms of the asymptotic behavior of the mean value. Orthogonality implies that for any Dirichlet character  $f$ ,

$$\sum_{n \leq x} f(n) = \alpha x + O(1) \tag{1.1}$$

for all large  $x$  (note that  $\alpha = 0$  if and only if  $f$  is nontrivial). Chudakov conjectured the converse should hold, and this was proved by Glazkov [4] in the case that  $\alpha \neq 0$  and by Klurman-Mangerel [7] in the case that  $\alpha = 0$ :

**Theorem 1.6** (Glazkov, Klurman-Mangerel). *Suppose  $f : \mathbb{N} \rightarrow \mathbb{C}$  is a completely multiplicative function whose image is finite and whose support contains all but finitely many primes. If (1.1) holds for some  $\alpha \in \mathbb{C}$ , then  $f$  must be a Dirichlet character.*

All three of the above results, as well as almost all other characterizations of characters we've seen, assume from the outset that  $f$  is multiplicative. Apart from our main theorem (Theorem 1.2), the only other characterization we're aware of that *deduces* multiplicativity from other hypotheses is the following result due to Kurlberg [10]:

**Theorem 1.7** (Kurlberg). *Suppose  $f : \mathbb{F}_p \rightarrow \mathbb{C}$  satisfies  $f(0) = 0$  and  $f(1) = 1$ , and that the image of  $\mathbb{F}_p^\times$  under  $f$  consists of roots of unity. Then  $f$  is a multiplicative character on  $\mathbb{F}_p$  if and only if*

$$\sum_{x \in \mathbb{F}_p} f(x) \overline{f(x+h)} = \begin{cases} -1 & \text{if } h \neq 0 \\ p-1 & \text{otherwise.} \end{cases}$$

In other words, the behavior of a character (mod  $p$ ) is completely determined by its autocorrelations. This result was motivated by a question of Harvey Cohn from the 1990s, which asked whether this happens more generally. It turns out that the natural analogue of Kurlberg's result in other finite fields fails to hold, as demonstrated by Choi and Siu; for any odd  $p$  and any  $k \geq 2$ , Theorem 3 of [3] produces a non-character  $f : \mathbb{F}_{p^k}^\times \rightarrow \{\pm 1\}$  that satisfies the analogous autocorrelation condition. Very recently, Benoist [2] used tools from symplectic geometry to demonstrate that the roots of unity assumption in Kurlberg's theorem is necessary.

## 2. PROOF

By abuse of notation, we call the quantity  $\tau(f)$  the *Gauss sum* of  $f$ . Our goal is to show that the value of the Gauss sum of a function carries a remarkable amount of information about the behavior of the function itself. As a warm up, observe that if  $g(x) = -1$  for all  $x \in \mathbb{F}_p^\times$ , then  $\tau(g) = 1$ ; a bit of playing around shows that this constant function is the *only* function  $\mathbb{F}_p^\times \rightarrow \{\pm 1\}$  that has Gauss sum 1. This phenomenon generalizes to functions whose Gauss sums live in any finite abelian extension of  $\mathbb{Q}$ :

**Lemma 2.1.** *Suppose  $g : \mathbb{F}_p^\times \rightarrow \mathbb{Q}(\zeta_n)$  such that  $p \nmid n$  and  $\tau(g) \in \mathbb{Q}(\zeta_n)$ . Then  $g(a) = -\tau(g)$  for all  $a \in \mathbb{F}_p^\times$ .*

*Proof.* Note that the polynomial

$$m(x) := -\tau(g) + g(1)x + g(2)x^2 + \cdots + g(p-1)x^{p-1}$$

has  $\zeta_p$  as a root. This means  $m(x)$  is a multiple of the minimal polynomial of  $\zeta_p$  over  $\mathbb{Q}(\zeta_n)$ , which (since  $p \nmid n$ ) is  $1 + x + \cdots + x^{p-1}$ . The claim instantly follows.  $\square$

Our proof of Theorem 1.2 proceeds in several stages of increasing generality. We start by proving a converse to Gauss' theorem on Gauss sums.

**Proposition 2.2.** *Suppose the image of  $f : \mathbb{F}_p^\times \rightarrow \mathbb{C}^\times$  consists of  $n^{\text{th}}$  roots of unity, that  $f(1) = 1$ , and that  $p \nmid n$ . Then  $f$  is a nontrivial character if and only if  $|\tau(f)| = \sqrt{p}$ .*

*Proof.* The forward direction is classical, so we henceforth assume  $|\tau(f)| = \sqrt{p}$  and prove that  $f$  is a character. Extend  $f$  to a function  $\mathbb{F}_p \rightarrow \mathbb{C}$  by setting  $f(0) := 0$ , and observe that

$$p = \left| \sum_{\ell \in \mathbb{F}_p} f(\ell) e\left(\frac{\ell}{p}\right) \right|^2 = \sum_{m, \ell \in \mathbb{F}_p} f(m) \overline{f(\ell)} e\left(\frac{m-\ell}{p}\right) = \sum_{k \in \mathbb{F}_p} e\left(\frac{k}{p}\right) \sum_{\ell \in \mathbb{F}_p} f(\ell+k) \overline{f(\ell)}.$$

The contribution from the  $k = 0$  term is  $\sum_{a \in \mathbb{F}_p} |f(a)|^2 = p - 1$ , whence

$$\sum_{k \in \mathbb{F}_p^\times} \underbrace{\left( \sum_{\ell \in \mathbb{F}_p} f(\ell+k) \overline{f(\ell)} \right)}_{g(k)} e\left(\frac{k}{p}\right) = 1.$$

Note that  $g(k) \in \mathbb{Q}(\zeta_n)$  and  $\tau(g) = 1$ , so Lemma 2.1 implies  $g(k) = -1$  for all  $k \in \mathbb{F}_p^\times$ . Since  $g(0) = p - 1$ , Kurlberg's theorem 1.7 implies that  $f$  must be a character.  $\square$

In practice, it's helpful to relax the hypotheses by removing the condition that  $f(1) = 1$ . The result becomes simpler to state if we abuse notation and refer to a *function*  $h : \mathbb{F}_p^\times \rightarrow \mathbb{C}^\times$  as an  $n^{\text{th}}$  root of unity if and only if everything in the image of  $h$  is an  $n^{\text{th}}$  root of unity.

**Corollary 2.3.** *Suppose  $f : \mathbb{F}_p^\times \rightarrow \mathbb{C}^\times$  is an  $n^{\text{th}}$  root of unity, and that  $p \nmid n$ . Then  $|\tau(f)| = \sqrt{p}$  if and only if  $f = \epsilon \chi$  for some constant  $\epsilon$  and some nontrivial character  $\chi$  that are both  $n^{\text{th}}$  roots of unity.*

*Proof.* The reverse direction immediately follows from Gauss' theorem, so we assume  $|\tau(f)| = \sqrt{p}$ . The function  $g := \overline{f(1)}f$  satisfies all the hypotheses of Proposition 2.2, so  $g$  must be a nontrivial character.  $\square$

Armed with these tools, we can now give a short proof of our main theorem.

*Proof of Theorem 1.2.* Recall our claim: given  $f : \mathbb{F}_p^\times \rightarrow \mathbb{C}^\times$  an  $n^{\text{th}}$  root of unity such that  $f(1) = 1$  and  $p \nmid n$ , we wish to prove that  $f$  is a nontrivial character if and only if  $\exists a \in \mathbb{F}_p^\times$  such that  $|\widehat{f}(a)| = 1$ . As before, the forward direction is classical (take  $a = -1$ ), so it suffices to prove the reverse direction. Pick  $a \in \mathbb{F}_p^\times$  such that  $|\widehat{f}(a)| = 1$ . A change of variables yields

$$\widehat{f}(a) = \frac{1}{\sqrt{p}} \sum_{m \in \mathbb{F}_p^\times} f(-\bar{a}m) e\left(\frac{m}{p}\right),$$

where  $\bar{a}$  denotes the multiplicative inverse of  $a$  in  $\mathbb{F}_p^\times$ . Corollary 2.3 implies  $f(-\bar{a}m) = \epsilon\chi(m)$  for some  $n^{\text{th}}$  roots of unity  $\epsilon$  and  $\chi$  (with the former a constant and the latter a nontrivial character). This in turn implies

$$f(\ell) = \epsilon\chi(-a)\chi(\ell) = \epsilon'\chi(\ell)$$

with  $\epsilon'$  an  $n^{\text{th}}$  root of unity. Since  $f(1) = 1 = \chi(1)$ , we deduce  $\epsilon' = 1$ , and the claim is proved.  $\square$

ACKNOWLEDGEMENTS. We're grateful to Andrew Granville, Oleksiy Klurman, and Sacha Mangerel for their comments and encouragement.

## REFERENCES

- [1] Jean-Paul Allouche, Leo Goldmakher, *Mock characters and the Kronecker symbol*, J. Number Theory **192** (2018), pp. 356–372.
- [2] Yves Benoist, *Fourier transform in cyclic groups*, preprint available at [arXiv:2406.11529](https://arxiv.org/abs/2406.11529)
- [3] Kwok-Kwong Choi, Man-Keung Siu, *Counter-examples to a Problem of Cohn on Classifying Characters*, Journal of Number Theory **84** (2000), pp. 40–48.
- [4] V. V. Glazkov, *Characters of multiplicative semigroup of natural numbers*, Studies in Number Theory (in Russian), No. 2, pp. 3–49, Izdat. Saratov. Univ., Saratov (1968).
- [5] E. Heppner, T. Maxsein, *Potenzreihen mit multiplikativen Koeffizienten*, Analysis **5** (1985), 87–95.
- [6] Oleksiy Klurman, Pär Kurlberg, *A note on multiplicative automatic sequences, II*, Bulletin of the London Mathematical Society Volume 52, Issue 1, pp. 185–188, 2020.
- [7] Oleksiy Klurman, Alexander P. Mangerel, *Rigidity theorems for multiplicative functions*, Mathematische Annalen (2018) **372** pp. 651–697.
- [8] Jakub Konieczny, *On multiplicative automatic sequences*, Bulletin of the London Mathematical Society, Volume 52, Issue 1, pp. 175–184, 2020.
- [9] Jakub Konieczny, *Multiplicative generalised polynomial sequences*, preprint available at [arXiv:2403.17590](https://arxiv.org/abs/2403.17590)
- [10] Pär Kurlberg, *On a character sum problem of Cohn*, J. Number Theory, **92** (2002), no. 1, pp. 174–181.
- [11] C. Methfessel, *Multiplicative and additive recurrent sequences*, Arch. Math. **63** (1994), 321–328.
- [12] A. Sárközy, *On multiplicative arithmetic functions satisfying a linear recursion*, Studia Sci. Math. Hungar. **13** (1978), 79–104.

HEILBRONN INSTITUTE FOR MATHEMATICAL RESEARCH, SCHOOL OF MATHEMATICS, UNIVERSITY OF BRISTOL, BRISTOL, UK  
Email address: [j.bober@bristol.ac.uk](mailto:j.bober@bristol.ac.uk)

DEPT OF MATHEMATICS AND STATISTICS, WILLIAMS COLLEGE, WILLIAMSTOWN, MA, USA 01267  
Email address: [Leo.Goldmakher@williams.edu](mailto:Leo.Goldmakher@williams.edu)