

ON THE NONLINEAR PERIODIC WESTERVELT EQUATION*

BENJAMIN RAINER[†] AND BARBARA KALTENBACHER[‡]

Abstract. This study focuses on numerically approximating solutions to the nonlinear periodic Westervelt equation within bounded domains in \mathbb{R}^d , where $d \in \{2, 3\}$, with Robin boundary conditions. We establish the existence and uniqueness of solutions in the linear and nonlinear case, emphasizing the importance of small excitations for existence and uniqueness in the nonlinear case. Adopting a multiharmonic formulation, we address the nonlinear propagation's harmonic frequencies. An iteration scheme is devised and its convergence under smallness conditions on the excitation is shown. We explore numerical solution to the resulting system of Helmholtz equations, with a conforming finite element method for its discretization. Using an implementation of the devised methodology, we showcase how acoustic waves propagate within nonlinear media. This study is supposed to enhance our understanding of ultrasound propagation dynamics, crucial for obtaining high-quality images from limited boundary measurements.

Key words. nonlinear periodic Westervelt equation, well-posedness, numerical approximation

AMS subject classifications. 35C09,35L05,35L70,65M12,65M60

1. Introduction. Ultrasound imaging plays a crucial role in medical diagnostics, relying on non-ionizing sound waves that pose no risk of cancer formation. In physics, ultrasound waves are characterized by a frequency above 20 kHz and typically ranging from 2 to 18 MHz in medical ultrasound imaging. However, the human body's nonlinear nature means its response to ultrasound is not linear [6], necessitating the understanding of nonlinear wave equations, crucial in various applications such as ultrasound diagnostics for tissue discrimination. In this study, we investigate the existence and uniqueness of the nonlinear periodic Westervelt equation, governing the practically relevant scenario of periodic (e.g., sinusoidal) excitations.

The Westervelt equation reads

$$(1.1) \quad \frac{1}{c^2} p_{tt} - \Delta p - \frac{b}{c^2} \Delta p_t = \frac{\beta_a}{\rho_0 c^4} (p^2)_{tt},$$

where $\beta_a = 1 + \frac{B}{2A}$ is the space dependent nonlinearity parameter. Multiplying (1.1) by $c^2 > 0$ and substituting $\eta(x) := \frac{\beta_a(x)}{\rho_0 c^2}$, equipping (1.1) with Robin boundary as well as periodic in time (with period T) conditions and adding a source term h , we obtain the following boundary value problem for the Westervelt equation on a bounded domain Ω with $C^{1,1}$ boundary $\partial\Omega$

$$(1.2) \quad \begin{cases} p_{tt}(t, x) - c^2 \Delta p(t, x) - b \Delta p_t(t, x) = \eta(x) (p(t, x)^2)_{tt} + h(t, x) & \text{in } (0, T) \times \Omega, \\ \beta p_t(t, x) + \gamma p(t, x) + \nabla p(t, x) \cdot \mathbf{n} = 0 & \text{on } (0, T) \times \partial\Omega, \\ p(0, x) = p(T, x), p_t(0, x) = p_t(T, x) & x \in \Omega, \end{cases}$$

where $\beta, \gamma \geq 0$ are the parameters for specifying absorption or impedance conditions

*Submitted to the editors 07/23/2024.

Funding: This work received funding from the ECSEL Joint Undertaking (JU) under grant agreement No. 101007350 and was partially supported by the Austrian Science Fund (FWF) [10.55776/P36318].

[†]Austrian Institute of Technology GmbH, Austria, Vienna (benjamin.rainer@ait.ac.at)
Department of Mathematics, University of Klagenfurt, Austria, Carinthia (benjamin.rainer@aau.at)

[‡]Department of Mathematics, University of Klagenfurt, Austria, Carinthia (barbara.kaltenbacher@aau.at).

on $\partial\Omega$, \mathbf{n} denotes the outer normal on $\partial\Omega$, and $b \geq 0$ is the diffusivity of sound in the media.

In our analysis we will assume $\eta \in L^\infty(\Omega)$, noting that (possibly nonsmoothly) varying B/A is highly relevant for imaging techniques that rely on the tissue specific values of this parameter. Indeed, there exists a vast corpus of literature, e.g., [5, 6, 10, 18, 29] that puts the value of the nonlinearity parameter in relation to different tissue types. Hence, the nonlinearity parameter $\eta(x)$ is image giving and its reconstruction in inhomogeneous media from (incomplete) measurements is of high interest.

We see that $h(t, x)$ has SI units $\text{kg}/(\text{m s}^4)$ and if we assume that the excitation/source is an ultrasound transducer $h(t, x)$ denotes the second time derivative of a periodic space dependent excitation g . That is, with $g(t, x) = \text{Re}\{\hat{h}(x)e^{i\omega t}\}$, $\omega = \frac{2\pi}{T}$, we have $h(t, x) = g(t, x)_{tt}$. Thus, possible solutions to (1.2) will consist of the fundamental frequency ω and, due to the nonlinearity, also multiples thereof – so-called higher harmonics. This will in fact be a key element in our reformulation of (1.2).

Taking a closer look at (1.2) and applying the identity $(p(t, x)^2)_{tt} = 2(p_t(t, x))^2 + p(t, x)p_{tt}(t, x)$ we obtain

$$(1.3) \quad (1 - 2\eta(x)p(t, x))p_{tt}(t, x) - c^2(x)\Delta p(t, x) - b\Delta p_t(t, x) = \eta(x)p_t(t, x)^2 + h(t, x).$$

This shows that the Westervelt equation degenerates if $1 - 2\eta(x)p(t, x) \leq 0$ for some $x \in \Omega$ and makes its analysis challenging.

Several studies, such as [14, 16, 19] have established local existence and uniqueness results for nonlinear Westervelt equations with different boundary conditions, as well as a priori estimates for finite element methods (FEM) [20]. Numerical methods, including time-stepping methods with fractional damping [4] and finite difference approaches [25] offer simulation based insights into the nonlinear propagation of ultrasound. Additionally, analytical solutions for specific boundary problems, like those presented in [28] contribute to our understanding of the spatial distribution of harmonic components.

Since excitations are often sinusoidal in ultrasound applications, an analysis and numerics in frequency domain are highly desirable to enhance efficiency and enable simulation times and precision guarantees that are compatible with the demands of imaging tasks. The difficulty here lies in the fact that the quadratic nonlinearity gives rise to autoconvolution terms in the Fourier transformed (with respect to time) equations and thus disables control of the potential degeneracy the way it is done in the time domain analysis, namely by bounding a pressure dependent coefficient (cf. $1 - 2\eta p$ in (1.3)).

Following up on the preliminary analysis in [13], we therefore

- provide a well-posedness analysis that allows to deal with the potential degeneracy in frequency domain, cf. Theorem 3.3;
- develop an iterative multi-level (with respect to frequency levels) scheme to solve (1.2) in frequency domain and prove its convergence, cf. Theorem 4.2;
- devise a finite element scheme for its implementation and provide experiments demonstrating its efficiency.

As compared to [13], we carry out our analysis relying on a perturbation of the strongly damped wave equation with unit inertial coefficient (rather than a space-time varying one), which leads to a fixed point scheme that is much more efficient to implement. Moreover, we consider a setting incorporating spatially (possibly nonsmoothly) varying sound speed, attenuation and nonlinearity coefficients, as highly relevant in imaging.

This paper is structured as follows: we introduce notation and relevant inequalities in [section 2](#), establish the main existence and uniqueness results in [section 3](#), present a multiharmonic expansion of the nonlinear periodic Westervelt equation including an iteration scheme and its convergence analysis for numerical computations in [section 4](#), provide experimental results in [section 5](#), and draw some conclusions in [section 6](#).

2. Notation and auxiliary results. We briefly set the notation and recall important inequalities used throughout this work. The (fractional) Sobolev spaces for an open and bounded C^1 domain Ω is denoted by $W^{s,p}(\Omega)$ with $1 \leq p \leq \infty$, $s \in \mathbb{R}_{\geq 0}$ and by real interpolation we have $W^{s,p}(\Omega) = [W^{k,p}(\Omega), W^{k+1,p}(\Omega)]_{\theta,p}$, where $k = \lfloor s \rfloor$, and $\theta = s - \lfloor s \rfloor$. The Hilbert space $W^{s,2}(\Omega)$ is denoted by $H^s(\Omega)$. We denote the norm on these spaces by $\|\cdot\|_{W^{s,p}(\Omega)}$. For the involved Bochner spaces we denote $\|\cdot\|_{L^p(0,T;L^q(\Omega))}$ as the norm on $L^p(0,T;L^q(\Omega))$. We will skip dependencies on time and space for the sake of readability when they are apparent from the context. For an open, bounded and connected C^1 domain $\Omega \subseteq \mathbb{R}^d$, $d \in \{2,3\}$ we make use of the following Sobolev embeddings and the resulting inequalities

$$(2.1) \quad \|u\|_{L^\infty} \leq C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)} \|u\|_{H^{13/8}(\Omega)},$$

$$(2.2) \quad \|u\|_{L^4(\Omega)} \leq C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)} \|u\|_{H^{9/8}(\Omega)},$$

$$(2.3) \quad \|u\|_{L^6(\Omega)} \leq C_{H^1(\Omega) \rightarrow L^6(\Omega)} \|u\|_{H^1(\Omega)}.$$

In the space-time domain we use the embedding inequalities

$$(2.4) \quad \|u\|_{L^\infty(0,T;X)} \leq C_{H^{3/4}(0,T) \rightarrow L^\infty(0,T)} \|u\|_{H^{3/4}(0,T;X)},$$

$$(2.5) \quad \|u\|_{L^4(0,T;X)} \leq C_{H^{1/4}(0,T) \rightarrow L^4(0,T)} \|u\|_{H^{1/4}(0,T;X)}.$$

Moreover, we apply the trace theorem for $\Omega \subset \mathbb{R}^d$, open, bounded, connected, $d \in \{2,3\}$ with a Lipschitz boundary $\partial\Omega$, which yields

$$(2.6) \quad \|u\|_{H^{1/2}(\partial\Omega)} \leq C_{\text{trace}} \|u\|_{H^1(\Omega)},$$

and the Poincaré-Friedrichs inequality which reads

$$(2.7) \quad \|u\|_{H^1(\Omega)}^2 \leq C_{\text{PF}}^2 \left(\|\nabla u\|_{L^2(\Omega)}^2 + \|u\|_{L^2(\partial\Omega)}^2 \right).$$

3. Existence and Uniqueness of solutions. In this section we investigate the existence and uniqueness of a solution to [\(1.1\)](#). We do so by first stating a general result in the linear case with variable coefficients and then extending this result towards the nonlinear case for a sufficiently small excitation. We start by investigating the linear case, thereby extending [\[13\]](#) to spatially variable coefficients c , b , and η , as relevant for imaging. Minimizing the smoothness assumptions on these coefficients to allow for jumps requires a dedicated weak and Galerkin formulation, cf. [\(6.1\)](#), [\(6.3\)](#) in the proof of [Theorem 3.1](#). Note that a strictly positive impedance parameter $\gamma > 0$ is needed to avoid constant solutions of the homogeneous problem, that is, to obtain uniqueness.

THEOREM 3.1. *[linear well-posedness]*

Let $T > 0$, $\Omega \subseteq \mathbb{R}^d$, $d \in \{2, 3\}$, open, bounded, connected, with $C^{1,1}$ boundary, $\beta, \gamma > 0$, $c, b, \frac{1}{c}, \frac{1}{b} \in L^\infty(\Omega)$, $c, b > 0$, $\alpha, \frac{1}{\alpha} \in L^\infty(0, T; L^\infty(\Omega))$, $\alpha_t \in L^\infty(0, T; L^\infty(\Omega))$, $\alpha > 0$, $\alpha(0) = \alpha(T)$, $\mu \in L^\infty(0, T; L^{2q/(q-1)}(\Omega))$, $\mu(0) = \mu(T)$ for some $q \in [1, \infty)$ and $f \in L^2(0, T; L^2(\Omega))$

Then there exists $r_0 > 0$ such that for coefficients satisfying the smallness condition

$$(3.1) \quad \|\alpha_t\|_{L^\infty(0, T; L^\infty(\Omega))} + \left\| \frac{\gamma}{\beta} - \frac{\mu}{\alpha} \right\|_{L^\infty(0, T; L^{2q/(q-1)}(\Omega))} < r_0$$

there exists a unique (weak) solution u of

$$(3.2) \quad \begin{cases} \alpha u_{tt} - c^2 \Delta u - b \Delta u_t + \mu u_t = f & \text{in } \Omega \times (0, T), \\ \beta u_t + \gamma u + \nabla u \cdot \mathbf{n} = 0 & \text{on } \partial\Omega \times (0, T), \\ u(0) = u(T), u_t(0) = u_t(T) & \text{in } \Omega, \end{cases}$$

with

$$u \in X = \{v \in H^2(0, T; L^2(\Omega)) \cap H^1(0, T; H^{3/2}(\Omega)) \cap L^2(0, T; H^2(\Omega)) : \|\nabla v \cdot \mathbf{n}\|_{H^1(0, T; L^2(\partial\Omega))} < \infty, v(0) = v(T), v_t(0) = v_t(T) \text{ a.e.}\}$$

and the solution u satisfies

$$\begin{aligned} \|u\|_X^2 &= \|u\|_{H^2(0, T; L^2(\Omega))}^2 + \|u\|_{H^1(0, T; H^{3/2}(\Omega))}^2 \\ &\quad + \|u\|_{L^2(0, T; H^2(\Omega))}^2 + \|\nabla v \cdot \mathbf{n}\|_{H^1(0, T; L^2(\partial\Omega))}^2 \\ &\leq C(\alpha, b, \gamma, c, \beta, T, \Omega)^2 \|f\|_{L^2(0, T; L^2(\Omega))}^2, \end{aligned}$$

where $C(C_\alpha, b, \gamma, c, \beta, T, \Omega) > 0$.

Proof. See the appendix. \square

Remark 3.2. By the Sobolev embedding Theorem we obtain that $u \in X$ is also in $C(0, T; H^{3/2}(\Omega)) \cap C^1(0, T; L^2(\Omega))$. We further want to highlight that this result does not need the periodicity of the right hand side f , but the periodicity of the coefficients α and μ must be ensured.

THEOREM 3.3. *[nonlinear well-posedness]*

Let $T > 0$, $\Omega \subseteq \mathbb{R}^d$, $d \in \{2, 3\}$, open, bounded, connected, with $C^{1,1}$ boundary, $\beta, \gamma > 0$, $c, b, \frac{1}{c}, \frac{1}{b} \in L^\infty(\Omega)$, $c, b > 0$ and $\eta \in L^\infty(\Omega)$. Then there exists $\delta > 0$ such that for all $h \in L^2(0, T; L^2(\Omega))$ with $\|h\|_{L^2(0, T; L^2(\Omega))} \leq \delta$ there exists a unique (weak) solution $u \in X$ of

$$(3.3) \quad \begin{cases} u_{tt} - c^2 \Delta u - b \Delta u_t = \eta(u^2)_{tt} + h & \text{in } (0, T) \times \Omega, \\ \beta u_t + \gamma u + \nabla u \cdot \mathbf{n} = 0 & \text{on } (0, T) \times \partial\Omega, \\ u(0) = u(T), u_t(0) = u_t(T) & \text{in } \Omega, \end{cases}$$

and the solution u satisfies

$$\|u\|_X \leq C(C_\alpha, b, \gamma, c, \beta, T, \Omega) \|h\|_{L^2(0, T; L^2(\Omega))},$$

for some constant $C(C_\alpha, b, \gamma, c, \beta, T, \Omega) > 0$ independent of h .

Proof. We define $\mathcal{F} : X \rightarrow X$ as the solution operator of

$$(3.4) \quad \begin{cases} u_{tt} - c^2 \Delta u - b \Delta u_t = \eta(v^2)_{tt} + h & \text{in } \Omega \times (0, T), \\ \beta u_t + \gamma u + \nabla u \cdot \mathbf{n} = 0 & \text{on } \partial\Omega \times (0, T), \\ u(0) = u(T), u_t(0) = u_t(T) & \text{in } \Omega, \end{cases}$$

$v \in X$ and show that \mathcal{F} , restricted to a suitable ball $B_r(0)$, $r > 0$, is a self-mapping and contraction. Since X is a Banach space we thus obtain uniqueness and existence of a solution u to (3.3) by the Banach fixed point theorem.

Now let $r > 0$ and $v \in B_r(0)$ be arbitrary. In order to apply Theorem 3.1 with $\alpha = 1$, $\mu = 0$, we have to verify that $f := \eta(v^2)_{tt} + h$ is in $L^2(0, T; L^2(\Omega))$. Note, that periodicity of the right hand side is not needed and in the linear case with setting $\alpha = 1$ we do not run into the problem of degeneracy. For the right hand side we obtain

$$\|\eta(v^2)_{tt} + h\|_{L^2(0, T; L^2(\Omega))} \leq 2\|\eta\|_{L^\infty(\Omega)} \|v_t^2 + v_{tt}v\|_{L^2(0, T; L^2(\Omega))} + \|h\|_{L^2(0, T; L^2(\Omega))}.$$

Exploiting the Sobolev embedding Theorem (see, [8] and [9]) and by (real) interpolation [1] we have

$$\begin{aligned} H^{3/4}(0, T; H^{13/8}(\Omega)) &= \\ H^{3/4}(0, T; [H^2(\Omega), H^{3/2}(\Omega)]_{3/4}) &= [L^2(0, T; H^2(\Omega)), H^1(0, T; H^{3/2}(\Omega))]_{3/4}, \end{aligned}$$

and

$$\begin{aligned} H^{1/4}(0, T; H^{9/8}(\Omega)) &= \\ H^{1/4}(0, T; [H^{3/2}(\Omega), L^2(\Omega)]_{1/4}) &= [L^2(0, T; H^{3/2}(\Omega)), H^1(0, T; L^2(\Omega))]_{1/4}, \end{aligned}$$

with this we obtain

$$(3.5) \quad \begin{aligned} \|v\|_{L^\infty(0, T; L^\infty(\Omega))} &\leq C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)} \|v\|_{L^\infty(0, T; H^{13/8}(\Omega))} \\ &\leq C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)} C_{H^{3/4}(0, T) \rightarrow L^\infty(0, T)} \|v\|_{H^{3/4}(0, T; H^{13/8}(\Omega))} \\ &\leq C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)} C_{H^{3/4}(0, T) \rightarrow L^\infty(0, T)} \\ &\quad \|v\|_{H^1(0, T; H^{3/2}(\Omega))}^{3/4} \|v\|_{L^2(0, T; H^2(\Omega))}^{1/4} \\ &\leq C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)} C_{H^{3/4}(0, T) \rightarrow L^\infty(0, T)} \|v\|_X \\ &\leq C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)} C_{H^{3/4}(0, T) \rightarrow L^\infty(0, T)} r, \end{aligned}$$

this yields that $v \in L^\infty(0, T; L^\infty(\Omega))$. We further show that v_t^2 is in $L^2(0, T; L^2(\Omega))$:

$$\begin{aligned} \|v_t^2\|_{L^2(0, T; L^2(\Omega))} &\leq \|v_t\|_{L^4(0, T; L^4(\Omega))}^2 \\ &\leq C_{H^{1/4}(0, T) \rightarrow L^4(0, T)}^2 C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 \|v_t\|_{H^{1/4}(0, T; H^{9/8}(\Omega))}^2 \\ &\leq C_{H^{1/4}(0, T) \rightarrow L^4(0, T)}^2 C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 \\ &\quad \left(\|v_t\|_{H^1(0, T; L^2(\Omega))}^{1/4} \|v_t\|_{L^2(0, T; H^{3/2}(\Omega))}^{3/4} \right)^2 \\ &\leq C_{H^{1/4}(0, T) \rightarrow L^4(0, T)}^2 C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 \|v\|_X^2 \\ &\leq C_{H^{1/4}(0, T) \rightarrow L^4(0, T)}^2 C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 r^2. \end{aligned}$$

Now we establish $f = \eta(v^2)_{tt} + h \in L^2(0, T; L^2(\Omega))$,

$$2\|\eta\|_{L^\infty(\Omega)}\|v_t^2 + v_{tt}v\|_{L^2(0, T; L^2(\Omega))} + \|h\|_{L^2(0, T; L^2(\Omega))} \leq \|h\|_{L^2(0, T; L^2(\Omega))} + 4\|\eta\|_{L^\infty(\Omega)} r^2 \left(C_{H^{1/4}(0, T) \rightarrow L^4(0, T)}^2 C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 + C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)} C_{H^{3/4}(0, T) \rightarrow L^\infty(0, T)} \right).$$

Thus, by Theorem 3.1 we obtain a constant $C > 0$ independent of f , η and v such that

$$\|\mathcal{F}(v)\|_X \leq C\|\eta(v^2)_{tt} + h\|_{L^2(0, T; L^2(\Omega))}.$$

In order to ensure that $\mathcal{F}(B_r(0)) \subseteq B_r(0)$, we impose the following conditions

$$(3.6) \quad C(4\|\eta\|_{L^\infty(\Omega)}(C_{H^{1/4}(0, T) \rightarrow L^4(0, T)}^2 C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 + C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)} C_{H^{3/4}(0, T) \rightarrow L^\infty(0, T)})r^2 + \delta) < r,$$

and

$$(3.7) \quad \|\eta\|_{L^\infty(\Omega)} C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)} C_{H^{3/4}(0, T) \rightarrow L^\infty(0, T)} r < \frac{1}{2}.$$

which can be achieved by choosing $\delta > 0$ and $r > 0$ sufficiently small. We are now ready to proceed by showing contractivity of \mathcal{F} . For $v_1, v_2 \in B_r(0)$ we set $w = u_1 - u_2 = \mathcal{F}(v_1) - \mathcal{F}(v_2)$, where both solutions exist and are unique according to Theorem 3.1. Then w solves

$$(3.8) \quad \begin{cases} w_{tt} - c^2 \Delta w - b \Delta w_t = \eta((v_1^2)_{tt} - (v_2^2)_{tt}) & \text{in } \Omega \times (0, T), \\ \beta w_t + \gamma w + \nabla w \cdot \mathbf{n} = 0 & \text{on } \partial\Omega \times (0, T), \\ w(0) = w(T), w_t(0) = w_t(T) & \text{in } \Omega. \end{cases}$$

Estimating $\eta((v_1^2)_{tt} - (v_2^2)_{tt})$ as before, allows us to apply Theorem 3.1 which yields

$$\|w\|_X \leq C\|\eta((v_1^2)_{tt} - (v_2^2)_{tt})\|_{L^2(0, T; L^2(\Omega))}.$$

We further have

$$(3.9) \quad \begin{aligned} (v_1^2)_{tt} - (v_2^2)_{tt} &= 2(v_{1_t}^2 + v_1 v_{1_{tt}} - v_{2_t}^2 - v_2 v_{2_{tt}}) \\ &= 2((v_{1_t} + v_{2_t})(v_{1_t} - v_{2_t}) + v_1 v_{1_{tt}} - v_2 v_{2_{tt}}) \\ &= 2((v_{1_t} + v_{2_t})(v_{1_t} - v_{2_t}) + v_1 v_{1_{tt}} - v_2 v_{2_{tt}} - v_1 v_{2_{tt}} + v_1 v_{2_{tt}}) \\ &= 2((v_{1_t} + v_{2_t})(v_{1_t} - v_{2_t}) + v_1(v_{1_{tt}} - v_{2_{tt}}) + v_{2_{tt}}(v_1 - v_2)). \end{aligned}$$

This yields the estimate

$$\begin{aligned} &\|(v_{1_t} + v_{2_t})(v_{1_t} - v_{2_t}) + v_1(v_{1_{tt}} - v_{2_{tt}}) + v_{2_{tt}}(v_1 - v_2)\|_{L^2(0, T; L^2(\Omega))} \\ &\leq \|v_{1_t} + v_{2_t}\|_{L^4(0, T; L^4(\Omega))} \|v_{1_t} - v_{2_t}\|_{L^4(0, T; L^4(\Omega))} \\ &\quad + \|v_1\|_{L^\infty(0, T; L^\infty(\Omega))} \|v_{1_{tt}} - v_{2_{tt}}\|_{L^2(0, T; L^2(\Omega))} \\ &\quad + \|v_1 - v_2\|_{L^\infty(0, T; L^\infty(\Omega))} \|v_{2_{tt}}\|_{L^2(0, T; L^2(\Omega))} \\ &= \|v_{1_t} + v_{2_t}\|_{L^4(0, T; L^4(\Omega))} \|(v_1 - v_2)_t\|_{L^4(0, T; L^4(\Omega))} \\ &\quad + \|v_1\|_{L^\infty(0, T; L^\infty(\Omega))} \|(v_1 - v_2)_{tt}\|_{L^2(0, T; L^2(\Omega))} \\ &\quad + \|v_1 - v_2\|_{L^\infty(0, T; L^\infty(\Omega))} \|v_{2_{tt}}\|_{L^2(0, T; L^2(\Omega))}. \end{aligned}$$

Applying the estimates from above we obtain

$$(3.10) \quad \|\mathcal{F}(v_1) - \mathcal{F}(v_2)\|_X \leq rC_0\|v_1 - v_2\|_X,$$

where C_0 reads

$$(3.11) \quad 4C\|\eta\|_{L^\infty(\Omega)}(C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)}C_{H^{3/4}(0,T) \rightarrow L^\infty(0,T)} \\ + C_{H^{1/4}(0,T) \rightarrow L^4(0,T)}^2C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2).$$

The imposed condition in equation (3.6) also ensures that $rC_0 < 1$ since by division with $r > 0$ we obtain

$$rC_0 = rC(4\|\eta\|_{L^\infty(\Omega)}(C_{H^{1/4}(0,T) \rightarrow L^4(0,T)}^2C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 \\ + C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)}C_{H^{3/4}(0,T) \rightarrow L^\infty(0,T)})) \\ \leq rC(4\|\eta\|_{L^\infty(\Omega)}(C_{H^{1/4}(0,T) \rightarrow L^4(0,T)}^2C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 \\ + C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)}C_{H^{3/4}(0,T) \rightarrow L^\infty(0,T)})) + C\frac{\delta}{r} < 1.$$

Thereby we obtain that \mathcal{F} is a contraction on $B_r(0)$ and, thus, has a unique fixed point $u \in B_r(0)$ with $\mathcal{F}(u) = u$ solving (3.3). We further note that (3.6) after rearranging yields

$$\delta < r \left(\frac{1}{C} - 4\|\eta\|_{L^\infty(\Omega)}(C_{H^{1/4}(0,T) \rightarrow L^4(0,T)}^2C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 \\ + C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)}C_{H^{3/4}(0,T) \rightarrow L^\infty(0,T)}r) \right).$$

Thus, $r > 0$ has to be sufficiently small such that

$$\frac{1}{C} > 4\|\eta\|_{L^\infty(\Omega)}(C_{H^{1/4}(0,T) \rightarrow L^4(0,T)}^2C_{H^{9/8}(\Omega) \rightarrow L^4(\Omega)}^2 \\ + C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)}C_{H^{3/4}(0,T) \rightarrow L^\infty(0,T)}r).$$

(3.7) ensures that we do not encounter degeneracy, since

$$\begin{aligned} & \sup\{c \in \mathbb{R} : 1 - 2\eta(x)u(t, x) \geq c, \text{ a.e. in } (0, T) \times \Omega\} \\ &= 1 - \inf\{c \in \mathbb{R} : 2\eta(x)u(t, x) \leq c, \text{ a.e. in } (0, T) \times \Omega\} \\ &\geq 1 - 2\|\eta u\|_{L^\infty(0,T;L^\infty(\Omega))} \\ &\geq 1 - 2\|\eta\|_{L^\infty(\Omega)}C_{H^{13/8}(\Omega) \rightarrow L^\infty(\Omega)}C_{H^{3/4}(0,T) \rightarrow L^\infty(0,T)}r > 0. \end{aligned}$$

This concludes the proof. \square

4. Multiharmonic expansion and iteration scheme. Thanks to periodicity, the solution to (1.2) can be represented in terms of the $L^2(0, T)$ orthogonal basis $(e^{im\omega t})_{m \in \mathbb{Z}}$ as $p(t, x) = \text{Re}\{\sum_{m=0}^{\infty} \hat{p}_k(x)e^{im\omega t}\}$, cf. [13]. Motivated by this, we use a multiharmonic ansatz $p^N(t, x) = \text{Re}\{\sum_{m=0}^N \hat{p}_k(x)e^{im\omega t}\}$, where $\hat{p} \in L^2(\Omega; \mathbb{C})$ and project (1.2) onto the subspace $X_N := \{\text{Re}\{\sum_{m=0}^N \alpha_m(x)e^{im\omega t}\} : \alpha_m \in H^2(\Omega; \mathbb{C})\}$ obtaining an approximation of the actual solution. We will further investigate under what assumptions this projection converges to the unique solution of (1.2) if it exists.

Another possible way to express our ansatz is given by the identity $\operatorname{Re}(z) = \frac{1}{2}(z + \bar{z})$ which yields

$$\operatorname{Re} \left\{ \sum_{m=0}^N \hat{p}(x) e^{\iota m \omega t} \right\} = \frac{1}{2} \left(\sum_{m=0}^N \hat{p}(x) e^{\iota m \omega t} + \sum_{m=0}^N \bar{\hat{p}}(x) e^{-\iota m \omega t} \right).$$

PROPOSITION 4.1. *Projecting (1.2) onto $X_N := \{\operatorname{Re}\{\sum_{m=0}^N \alpha_m(x) e^{\iota m \omega t}\} : \alpha_m \in H^2(\Omega; \mathbb{C})\}$ with periodic excitation $h(t, x) = g_{tt}$, $g(t, x) = \operatorname{Re}\{\sum_{k=1}^N \hat{h}_k(x) e^{\iota k \omega t}\}$, $\omega = \frac{2\pi}{T}$ yields the following coupled system of PDEs (Helmholtz equations) for $x \in \Omega$*

$$(4.1) \quad \begin{cases} -c^2 \Delta \hat{u}_0(x) = 0 & m = 0, \\ -\kappa^2 \hat{u}_1(x) - \Delta \hat{u}_1(x) = -\frac{\kappa^2 \eta(x)}{2} \left(\hat{u}_0(x) \hat{u}_1(x) + \sum_{j=1:2}^{2N-1} \bar{\hat{u}}_{\frac{j-m}{2}}(x) \hat{u}_{\frac{j+m}{2}}(x) \right) & m = 1, \\ -\kappa^2 m^2 \hat{u}_m(x) - \Delta \hat{u}_m(x) = -\frac{m^2 \kappa^2 \eta(x)}{2} \left(\sum_{j=1}^m \hat{u}_j(x) \hat{u}_{m-j}(x) + 2 \sum_{j=m:2}^{2N-m} \bar{\hat{u}}_{\frac{j-m}{2}}(x) \hat{u}_{\frac{j+m}{2}}(x) \right) & m > 1, \end{cases}$$

where $\kappa^2 = \frac{\omega^2}{c^2 + \iota m \omega b}$. This system is further equipped with the projected boundary conditions $\iota \beta m \omega \hat{u}_m + \gamma \hat{u}_m + \nabla \hat{u}_m \cdot \mathbf{n} = 0$ on $\partial\Omega$.

Proof. Let $u^N, v^N \in X_N$ then

$$(4.2) \quad \begin{aligned} u^N v^N &= \frac{1}{2} \left(\sum_{m=0}^N \hat{u}_m(x) e^{\iota m \omega t} + \sum_{m=0}^N \bar{\hat{u}}_m(x) e^{-\iota m \omega t} \right) \\ &\quad \frac{1}{2} \left(\sum_{m=0}^N \hat{v}_m(x) e^{\iota m \omega t} + \sum_{m=0}^N \bar{\hat{v}}_m(x) e^{-\iota m \omega t} \right) \\ &= \frac{1}{2} \operatorname{Re} \left\{ \sum_{m=0}^{2N} \sum_{j=\max\{m-N, 0\}}^{\min\{m, N\}} \hat{u}_j(x) \hat{v}_{m-j}(x) e^{\iota m \omega t} + \bar{\hat{u}}_j(x) \bar{\hat{v}}_{m-j}(x) e^{\iota(m-2j)\omega t} \right\}, \end{aligned}$$

where we used the formula for the finite Cauchy product. A thorough computation of the second sum yields

$$(4.3) \quad \begin{aligned} &\operatorname{Re} \left\{ \sum_{m=0}^{2N} \sum_{j=\max\{m-N, 0\}}^{\min\{m, N\}} \bar{\hat{u}}_j(x) \hat{v}_{m-j}(x) e^{\iota(m-2j)\omega t} \right\} \\ &= \operatorname{Re} \left\{ \sum_{j=0}^N \bar{\hat{u}}_j(x) \hat{v}_j(x) + \sum_{m=1}^N e^{\iota m \omega t} \sum_{k=m:2}^{2N-m} \left[\bar{\hat{u}}_{\frac{k-m}{2}}(x) \hat{v}_{\frac{k+m}{2}}(x) + \hat{u}_{\frac{k+m}{2}}(x) \bar{\hat{v}}_{\frac{k-m}{2}}(x) \right] \right\}. \end{aligned}$$

After projection, the first sum becomes (since we skip all harmonic frequencies above

the N -th harmonic)

$$\begin{aligned} \text{Proj}_{X_N} \left(\text{Re} \left\{ \sum_{m=0}^{2N} \sum_{j=\max\{m-N,0\}}^{\min\{m,N\}} \hat{u}_j(x) \hat{v}_{m-j}(x) e^{\iota m \omega t} \right\} \right) \\ = \text{Re} \left\{ \sum_{m=0}^N e^{\iota m \omega t} \sum_{j=0}^m \hat{u}_j(x) \hat{v}_{m-j}(x) \right\}. \end{aligned}$$

This finally yields

$$\begin{aligned} \text{Proj}_{X_N}(u^N v^N) &= \frac{1}{2} \text{Re} \left\{ \sum_{m=0}^N e^{\iota m \omega t} \sum_{j=0}^m \hat{u}_j(x) \hat{v}_{m-j}(x) \right\} \\ &+ \frac{1}{2} \text{Re} \left\{ \sum_{j=0}^N \hat{u}_j(x) \hat{v}_j(x) + \sum_{m=1}^N e^{\iota m \omega t} \sum_{k=m:2}^{2N-m} \left[\hat{u}_{\frac{k-m}{2}}(x) \hat{v}_{\frac{k+m}{2}}(x) + \hat{u}_{\frac{k+m}{2}}(x) \hat{v}_{\frac{k-m}{2}}(x) \right] \right\}. \end{aligned}$$

For $\text{Proj}_{X_N}(((p^N)^2)_{tt})$ we obtain the following expression

$$(4.4) \quad \text{Proj}_{X_N}(((p^N)^2)_{tt}) = -\frac{1}{2} \omega^2 \text{Re} \left\{ \sum_{m=1}^N m^2 e^{\iota m \omega t} \left(\sum_{k=1}^m \hat{p}_k(x) \hat{p}_{m-k}(x) + 2 \sum_{k=m:2}^{2N-m} \tilde{\hat{p}}_{\frac{k-m}{2}}(x) \hat{p}_{\frac{k+m}{2}}(x) \right) \right\}.$$

For the left hand side of (1.2) we obtain

$$\begin{aligned} \text{Proj}_{X_N}(p_{tt}^N - c^2 \Delta p^N - b \Delta p_t^N) &= \omega^2 \frac{1}{2} \sum_{k=0}^N -\hat{p}_k(x) k^2 e^{\iota k \omega t} - \bar{\hat{p}}_k(x) k^2 e^{-\iota k \omega t} \\ &- c^2 \frac{1}{2} \sum_{k=0}^N \Delta \hat{p}_k(x) e^{\iota k \omega t} + \Delta \bar{\hat{p}}_k(x) e^{-\iota k \omega t} \\ &- \omega b \frac{1}{2} \sum_{k=0}^N \Delta \hat{p}_k(x) k e^{\iota k \omega t} - \Delta \bar{\hat{p}}_k(x) k e^{-\iota k \omega t}. \end{aligned}$$

For the source term we further have $h(x, t) = g(x, t)_{tt}$ which yields $\text{Proj}_{X_N}(h(x, t)) = -\frac{\omega^2}{2} \sum_{k=1}^N k^2 \hat{h}_k e^{\iota k \omega t} + k^2 \bar{\hat{h}}_k e^{-\iota k \omega t}$. Considering the linear independence of $e^{\iota m \omega t}$ in $L^2(0, T)$ we obtain for each $0 \leq m \leq N$ that $\hat{p}_m(x)$ has to solve

$$\begin{aligned} -\omega^2 m^2 \hat{p}_m(x) - (c^2 + \iota m \omega b) \Delta \hat{p}_m(x) &= \\ -\frac{\eta(x) \omega^2 m^2}{2} \left(\sum_{k=1}^m \hat{p}_k(x) \hat{p}_{m-k}(x) + 2 \sum_{k=m:2}^{2N-m} \tilde{\hat{p}}_{\frac{k-m}{2}}(x) \hat{p}_{\frac{k+m}{2}}(x) \right) &- \omega^2 m^2 \hat{h}_m(x). \end{aligned}$$

Dividing by $(c^2 + \iota m \omega b)$ yields

$$\begin{aligned} -\kappa^2 m^2 \hat{p}_m(x) - \Delta \hat{p}_m(x) &= \\ -\frac{\eta(x) \kappa^2 m^2}{2} \left(\sum_{k=1}^m \hat{p}_k(x) \hat{p}_{m-k}(x) + 2 \sum_{k=m:2}^{2N-m} \tilde{\hat{p}}_{\frac{k-m}{2}}(x) \hat{p}_{\frac{k+m}{2}}(x) \right) &- \kappa^2 m^2 \hat{h}_m(x), \end{aligned}$$

$m = 0$ results in

$$(4.5) \quad -\Delta \hat{p}_0(x) = 0,$$

and the boundary conditions for each of the Helmholtz equations read

$$(4.6) \quad (\iota\beta m\omega + \gamma)\hat{p}_m(x) + \nabla \hat{p}_m(x) \cdot \mathbf{n} = 0,$$

where we define $\kappa^2 = \frac{\omega^2}{c^2 + \iota m \omega b}$. This concludes the proof. \square

THEOREM 4.2. *[convergence of projection scheme]*

Under the assumptions of Theorem 3.3 there exists $\delta > 0$ such that for all $h \in L^2(0, T, L^2(\Omega))$ with $\|h\|_{L^2(0, T; L^2(\Omega))} < \delta$ there exists a unique solution u to (3.3) and the solutions $\tilde{u}_N \in X_N$ of the iteration scheme

$$(4.7) \quad \text{Proj}_{X_N}((\tilde{u}_N)_{tt} - c^2 \Delta(\tilde{u}_N) - b \Delta(\tilde{u}_N)_t - \eta(\tilde{u}_{N-1}^2)_{tt} - h) = 0,$$

converge to the unique solution u .

Proof. We consider $L_{\alpha, \mu} : X \rightarrow X'$ and $\tilde{u}_N \in X_N$ as a Galerkin solution of

$$(4.8) \quad \langle L_{\alpha, \mu} \tilde{u}_N - f, \phi \rangle_{X', X} := \int_0^T (\alpha(\tilde{u}_N)_{tt} + \mu(\tilde{u}_N)_t, \phi)_{L^2(\Omega)} + (c^2 \nabla \tilde{u}_N + b \nabla(\tilde{u}_N)_t, \nabla \phi)_{L^2(\Omega)} + ((c^2 \beta + b\gamma)(\tilde{u}_N)_t + c^2 \gamma \tilde{u}_N + b\beta(\tilde{u}_N)_{tt}, \phi)_{L^2(\partial\Omega)} - (f, \phi)_{L^2(\Omega)} dt = 0,$$

for all $\phi \in X_N$ where we set $\alpha = 1$, $\mu = 0$, $f = \eta(\tilde{u}_{N-1}^2)_{tt} + h$. One readily checks that the solution to (4.8) exists and is unique using the respective test functions \tilde{u}_N , $(\tilde{u}_N)_t$, $(\tilde{u}_N)_{tt}$, $-\Delta \tilde{u}_N$, $-\Delta(\tilde{u}_N)_t$ which are indeed contained in X_N for α and f fulfilling the assumptions of Theorem 3.1. We further obtain the following coupled Helmholtz system for $\tilde{u}_N = \sum_{k=1}^N \hat{u}_k^N e^{\iota k \omega t}$,

$$(4.9) \quad \begin{cases} -c^2 \Delta \hat{u}_0^N(x) = 0 & m = 0, \\ -\kappa^2 \hat{u}_1^N(x) - \Delta \hat{u}_1^N(x) = \\ \quad -\kappa^2 \eta(x) \left(\hat{u}_0^{N-1}(x) \hat{u}_1^{N-1}(x) + \sum_{j=1:2}^{2N-3} \bar{u}_{\frac{j-1}{2}}^{N-1}(x) \hat{u}_{\frac{j+1}{2}}^{N-1}(x) \right) - \kappa^2 \hat{h}_1(x) & m = 1, \\ -\kappa^2 m^2 \hat{u}_m^N(x) - \Delta \hat{u}_m^N(x) = \\ \quad -\frac{\kappa^2 m^2 \eta(x)}{2} \left(\sum_{j=1}^m \hat{u}_j^{N-1}(x) \hat{u}_{m-j}^{N-1}(x) + 2 \sum_{j=m:2}^{2N-m-2} \bar{u}_{\frac{j-m}{2}}^{N-1}(x) \hat{u}_{\frac{j+m}{2}}^{N-1}(x) \right) \\ \quad - \kappa^2 m^2 \hat{h}_m(x) & m > 1, \end{cases}$$

with the boundary conditions $\iota\beta m\omega \hat{u}_m^N + \gamma \hat{u}_m^N + \nabla \hat{u}_m^N \cdot \mathbf{n} = 0$ on $\partial\Omega$. We obtain $\|\tilde{u}_N\|_X < r$ provided $\|\tilde{u}_{N-1}\|_X < r$, by (3.6) for $\|h\|_{L^2(0, T; L^2(\Omega))} < \delta$ and $r > 0$ chosen small enough according to Theorem 3.3. A direct investigation of the error $w := \tilde{u}_N - u$ is not possible because the difference need not necessarily be contained in X_N . Therefore, we circumvent this issue by investigating $\hat{w} := \tilde{u}_N - \text{Proj}_{X_N} u \in X_N$. For $\phi \in X_N$ we have

$$\begin{aligned} \langle L_{\alpha, \mu} \hat{w}, \phi \rangle_{X', X} &= \langle L_{\alpha, \mu}(\tilde{u}_N - u + u - \text{Proj}_{X_N} u), \phi \rangle_{X', X} \\ &= \langle L_{\alpha, \mu}(\tilde{u}_N - u) + L_{\alpha, \mu}(u - \text{Proj}_{X_N} u), \phi \rangle_{X', X} \\ &= (\eta(\tilde{u}_{N-1}^2)_{tt} - \eta(u^2)_{tt}, \phi)_{L^2(0, T; L^2(\Omega))} + \langle L_{\alpha, \mu}(u - \text{Proj}_{X_N} u), \phi \rangle_{X', X} \\ &= (2\eta[(\tilde{u}_{N-1})_t^2 - u_t^2] + \tilde{u}_{N-1}((\tilde{u}_{N-1})_{tt} - u_{tt}) \\ &\quad + u_{tt}(\tilde{u}_{N-1} - u), \phi)_{L^2(0, T; L^2(\Omega))} + \langle L_{\alpha, \mu}(u - \text{Proj}_{X_N} u), \phi \rangle_{X', X}, \end{aligned}$$

where we used the fact that $(L_{\alpha,\mu}u, \phi)_{L^2(0,T;L^2(\Omega))} = (\eta(u^2)_{tt} + h, \phi)_{L^2(0,T;L^2(\Omega))} \forall \phi \in X$ and $X_N \subset X$. Theorem 3.1 yields a $C > 0$ such that

$$\begin{aligned} \|\tilde{u}_N - \text{Proj}_{X_N} u\|_X &\leq C \|2\eta[(\tilde{u}_{N-1})_t^2 - u_t^2] + \tilde{u}_{N-1}((\tilde{u}_{N-1})_{tt} - u_{tt}) \\ &\quad + u_{tt}(\tilde{u}_{N-1} - u)\|_{L^2(0,T;L^2(\Omega))} + L_{\alpha,\mu}(u - \text{Proj}_{X_N} u) \|_{L^2(0,T;L^2(\Omega))}. \end{aligned}$$

We estimate similarly as in Theorem 3.3 and obtain

$$\begin{aligned} &C \|2\eta[(\tilde{u}_{N-1})_t^2 - u_t^2 + \tilde{u}_{N-1}((\tilde{u}_{N-1})_{tt} - u_{tt}) + u_{tt}(\tilde{u}_{N-1} - u)]\|_{L^2(0,T;L^2(\Omega))} \\ &\leq 2C \|\eta\|_{L^\infty(\Omega)} (\|(\tilde{u}_{N-1})_t - u_t\|_{L^4(0,T;L^4(\Omega))} \|(\tilde{u}_{N-1})_t + u_t\|_{L^4(0,T;L^4(\Omega))} \\ &\quad + \|\tilde{u}_{N-1}\|_{L^\infty(0,T;L^\infty(\Omega))} \|(\tilde{u}_{N-1} - u)_{tt}\|_{L^2(0,T;L^2(\Omega))} \\ &\quad + \|u_{tt}\|_{L^2(0,T;L^2(\Omega))} \|\tilde{u}_{N-1} - u\|_{L^\infty(0,T;L^\infty(\Omega))}) \\ &\leq rC_0 \|\tilde{u}_{N-1} - u\|_X, \end{aligned}$$

Since $\bigcup_{N \in \mathbb{N}} X_N$ is dense in X , it holds that

$$\|L_{\alpha,\mu}(u - \text{Proj}_{X_N} u)\|_{X'} \leq \|L_{\alpha,\mu}\|_{X,X'} \|u - \text{Proj}_{X_N} u\|_X \rightarrow 0, N \rightarrow \infty.$$

According to Theorem 3.3 we have $q(r) := rC_0 < 1$ and by the triangle inequality we obtain

$$\|\tilde{u}_N - u\|_X \leq q(r) \|\tilde{u}_{N-1} - u\|_X + (C+1) \|u - \text{Proj}_{X_N} u\|_X.$$

By induction, $\tilde{u}_0 = 0$ and setting $\tilde{C} = C+1$ we finally arrive at

$$\|\tilde{u}_N - u\|_X \leq q(r)^N \|u\|_X + \tilde{C} \sum_{i=1}^N q(r)^{N-i} \|u - \text{Proj}_{X_i} u\|_X.$$

Now, let $\varepsilon > 0$ be arbitrary, by the density of $\bigcup_{N \in \mathbb{N}} X_N$ in X there exists an $N(\varepsilon) \in \mathbb{N}$ such that for all $n > N(\varepsilon)$ we have $\|u - \text{Proj}_{X_n} u\|_X < \varepsilon$. We set $N > N(\varepsilon)$, large enough such that $q(r)^{N-N(\varepsilon)} < \varepsilon$ holds. This yields

$$\begin{aligned} \|\tilde{u}_N - u\|_X &\leq q(r)^N \|u\|_X + \tilde{C} \sum_{i=1}^N q(r)^{N-i} \|u - \text{Proj}_{X_i} u\|_X \\ &= q(r)^N \|u\|_X + \tilde{C} q(r)^{N-N(\varepsilon)} \sum_{i=1}^{N(\varepsilon)} q^{N(\varepsilon)-i} \|u - \text{Proj}_{X_i} u\|_X \\ &\quad + \tilde{C} \sum_{i=N(\varepsilon)+1}^N q(r)^{N-i} \|u - \text{Proj}_{X_i} u\|_X \\ &< \varepsilon \|u\|_X + \frac{\varepsilon \tilde{C} c_{\text{Proj}}}{1-q} + \frac{\varepsilon \tilde{C}}{1-q} \\ &= \varepsilon \left(\|u\|_X + \frac{\tilde{C}(1+c_{\text{Proj}})}{1-q} \right), \end{aligned}$$

where $c_{\text{Proj}} := \sup_{i \in \mathbb{N}} \|u - \text{Proj}_{X_i} u\|_X$ is finite because $\bigcup_{N \in \mathbb{N}} X_N$ is dense in X . This proves the convergence of \tilde{u}_N to u in X . \square

5. Numerical experiments. We revisit the result from Theorem 4.2 where we obtained (4.9) which simplifies to the following system of Helmholtz equations for $x \in \Omega$:

$$(5.1) \quad \begin{cases} -\kappa^2 \hat{u}_1^N(x) - \Delta \hat{u}_1^N(x) = -\kappa^2 \left(\eta(x) \sum_{j=1:2}^{2N-3} \bar{u}_{\frac{j-1}{2}}^{N-1}(x) \hat{u}_{\frac{j+1}{2}}^{N-1}(x) \hat{h}_1(x) \right) & m = 1, \\ -\kappa^2 m^2 \hat{u}_m^N(x) - \Delta \hat{u}_m^N(x) = -\kappa^2 m^2 \hat{h}_m(x) - \frac{\kappa^2 m^2 \eta(x)}{2} \\ \left(\sum_{j=1}^m \hat{u}_j^{N-1}(x) \hat{u}_{m-j}^{N-1}(x) + 2 \sum_{j=m:2}^{2(N-1)-m} \bar{u}_{\frac{j-m}{2}}^{N-1}(x) \hat{u}_{\frac{j+m}{2}}^{N-1}(x) \right) & m > 1, \end{cases}$$

with the following boundary conditions for $x \in \partial\Omega$:

$$(5.2) \quad i\beta m \omega \hat{u}_m^N(x) + \gamma \hat{u}_m^N(x) + \nabla \hat{u}_m^N(x) \cdot \mathbf{n} = 0,$$

where $c, \omega, b > 0$, $\kappa^2 = \frac{\omega^2}{c^2 + i m \omega b} \in \mathbb{C}$, $\Omega \subset \mathbb{R}^2$ with a $C^{1,1}$ boundary $\partial\Omega$ and in case of non constant $c(x)$, $b(x)$ we assume that $c, \frac{1}{c}, b, \frac{1}{b} \in L^\infty(\Omega)$ and obtain a space dependent wave number $\kappa(x)^2 = \frac{\omega^2}{c^2(x) + i \omega b(x)}$. The solution for $m = 0$ is zero. In Theorem 4.2 we have shown that this iteration scheme converges to the unique (weak) solution of (1.1) as $N \rightarrow \infty$. We notice that for each level $m \in \mathbb{N}$ the wave number increases. Thus, conforming Galerkin FEM methods applied to this system of Helmholtz equations will encounter the so called pollution effect [2, 3]. Moreover, in each iteration the solutions of the previous iteration are used. Therefore, the pollution effect affects not only solutions for higher frequencies but any of them and gets amplified throughout the iterations.

For our numerical computations we implement a conforming 2D FEM solver with linear Lagrange elements for the Helmholtz equation. The implementation of the FEM solver and the iteration scheme (5.1) is available at <https://shorturl.at/vBCN7> and as supplemental material enclosed to this paper. We select the domain of propagation to be $\Omega = B_{0.2}(0) \subset \mathbb{R}^2$ and we place a monopole source at $(0, 0)^T$ using a regularized Dirac function which reads

$$(5.3) \quad \hat{h}(x) = \tilde{\delta}(x) = \begin{cases} \frac{1}{4\zeta} \left(1 + \cos \left(\pi \frac{\|x-x_0\|_{l^2}}{2\zeta} \right) \right) & \|x-x_0\|_{l^2} \leq 2\zeta, \\ 0 & \text{else,} \end{cases}$$

where x_0 denotes the point source location. We fix the excitation frequency and consider the excitation to be periodic. Therefore, our constructed source is indeed in $L^2(0, T; L^2(\Omega))$. The parameters used in the simulations are chosen as typical for water, in particular $B/A = 5$ [22, 26], $\rho_0 = 1000 \text{ kg/m}^3$, $\omega = 2\pi \frac{1}{10^{-5}} \text{ rad/s}$, $b = 10^{-9} \text{ m}^2/\text{s}$, mesh parameter 0.0005 m , $c = 1480 \text{ m/s}$, $\gamma = 1$, $\zeta = 0.0005$, sound pressure 0.5 MPa , and number of iterations 15. We further set the impedance coefficient for each Helmholtz equation to $\beta = \frac{1}{c}$ to obtain absorbing boundary conditions mimicking the Sommerfeld radiation condition (if the source is placed in the center). Figure 1 shows the L^2 error of consecutive iterations for the second harmonic with respect to different mesh sizes. A large mesh parameter leads to a strong pollution effect in the higher harmonic solutions. The influence of the pollution effect increases with an increase in the number of iterations.

Figure 3 depicts the acoustic pressure along a line originating in $(0, 0)^T$. The high fundamental frequency leads to stronger attenuation and, therefore, the nonlinear effect is prevalent. Figure 2 shows the frequency components present at different locations in space. We notice that the three first harmonics are above -40 dB . In Figure 4, we further plot the acoustic pressure over time in a single point within Ω .




FIG. 1. L^2 error of the second level (harmonic) of consecutive iterations for different mesh sizes.




FIG. 2. Frequency components of $p(t, x)$ at $(0, -0.1)^T$, where we applied a Hann window [21] in time.

Comparing the nonlinear case to the linear one, the non-uniform phase velocity is clearly visible in the nonlinear case.

It is well known that if the wave vectors are not perpendicular to the tangent in boundary points the employed absorbing boundary conditions cause spurious reflections [24]. This is avoided in case of a source centered in a circular domain, where the wave hits the boundary in normal direction. This is not necessarily the case for other source positions relative to the boundary and needs to be taken into account accordingly. A self-adaptive boundary condition scheme for the nonlinear Westervelt equation in time domain has been introduced in [17]. The absorbing boundary conditions are adapted in *real-time* taking into account incidence wave vectors. Here, estimates of the solution computed in time step t_{n-1} are used to adapt the boundary conditions in the next time step t_n . This works remarkably well in space-time finite element methods. However, we are facing (5.1), a system of Helmholtz equations.




FIG. 3. The propagation of the wave along a line originating in $(0, 0)^T$ and ending in $(0, 0.2)^T$. The black dotted line denotes the linear case where $\beta_a = 0$.




FIG. 4. Evolution over time of the acoustic pressure at $(0, -0.1)^T$.

Thus adaptation over time is not an option. If there is only a single source or if ∇p is known, the gradient method can be employed [12]. In our iteration scheme, each iteration of (5.1) virtually adds new *sources* with different frequencies, though. A detailed investigation and the derivation of adaptive absorbing boundary conditions for (5.1) is beyond the scope of this paper.

Next, we turn to a scenario that resembles a watershed experiment (cf. Figure 5). Here, we are interested in boundary measurements of a domain filled with water. This type of experiment is of high interest for evaluating the accuracy of inverse methods [15] for the reconstruction of the nonlinearity parameter. We consider a circular air domain with a radius of one meter and a concentrically placed circular water tank with a radius of 0.35 meter including three phantoms. The phantoms differ in size and material. Their radii are 0.05, 0.04 and 0.06 meters, respectively. The sound speed is set accordingly; for air 344 m/s and for water we use 1450 m/s. We use the following B/A values which determine the nonlinearity parameter η : air (1), water (5), first phantom (9), second phantom (10), third phantom (12). A linear array with nine elements spaced by $\lambda/4$ in horizontal direction, is placed in the center of the domain. Each element of the linear array emits a 10 kHz sinusoidal ultrasound wave with a peak pressure of 40 kPa. We use (5.1) to compute the first six harmonic solutions which are then used to approximate $p(t, x)$. Figure 6 depicts the approximation of $p(0, x)$ with all three phantoms being present. The effect of the




FIG. 5. Container filled with water and three phantoms inside, placed in a circular domain filled with air.

nonlinearity parameter and the different propagation speeds is clearly visible. We are interested in boundary measurements of the domain filled with water for the cases of no phantoms, one phantom, two phantoms and three phantoms. To this end, $p_j(t, x)$, for $j \in \{0, 1, 2, 3\}$ denotes the solution with no phantoms, phantom one, phantom two, or phantom three present. Solutions with combinations of phantoms are denoted by the corresponding indices, i.e., $p_{1,2}(t, x)$ with phantoms one and two and $p_{1,2,3}(t, x)$ with all three phantoms present. Figure 7 displays the boundary values relative to the location of the source on the surface of the water tank. In the left column we show the difference between the solutions without phantoms and single phantoms present. The right column displays the solution without phantoms, and the difference to having two (the first and second) and three phantoms present. The first and second phantom have similar B/A values but differ in size and position. This is reflected in the first two figures in the left column of figure 7. The third phantom has a B/A value of 12 and lies closest to the source; this already causes the boundary values to differ by more than 1 kPa.




FIG. 6. Simulation result of $p_{1,2,3}(0, x)$, left x - y view of $p_{1,2,3}(0, x)$ and right a 3D plot of $p_{1,2,3}(0, x)$.

6. Conclusions. In conclusion, our study aims to provide analytic and numerical tools for ultrasound imaging, particularly focusing on the nonlinear periodic Westervelt equation. This equation is pivotal in understanding tissue interactions and dis-




FIG. 7. Boundary values of the container filled with water with different phantoms present. On the left side we show the difference between $p_0(0, x)$ and $p_j(0, x)$, $j \in \{1, 2, 3\}$, the right side of the figure shows the difference between $p_0(0, x)$ and the solutions including two, and three phantoms. The x -axis denotes the radians where the source is located at 0 radian.

crimination in ultrasound diagnostics. Through our investigation, we have shown that solutions to the nonlinear periodic Westervelt equation exist under certain smallness assumptions on the excitation. Additionally, we have introduced an iterative scheme that approximates solutions in frequency domain.

While in this work we considered an excitation in the interior of the domain, in practical application the ultrasound transducer array is located on a part of the boundary. Therefore, our future research will include investigations on the case where the excitation is present on $\Gamma \subset \partial\Omega$.

Also the development and use of enhanced absorbing boundary conditions [7, 11] for a nonlinear wave equation in frequency domain appears to be a relevant topic of research in our context.

Based on the results obtained here, we plan to provide precise and efficient reconstruction schemes for nonlinear ultrasound tomography, which amounts to reconstructing space dependent coefficients $c(x)$ and/or $\eta(x)$ in (1.1).

Appendix.

Proof of Theorem 3.1. First, we derive the weak form of (3.2). To this end, we multiply by $-\frac{1}{\alpha}\Delta\psi$ for some $\psi \in H^1(0, T; H^1(\Omega)) \cap L^2(0, T; H^2(\Omega))$, and integrate over space time, using integration by parts in the first term (which we have augmented in order to be able to use the boundary conditions)

$$\begin{aligned} & \int_0^T \int_{\Omega} (u_{tt} + \frac{\gamma}{\beta}u_t)(-\Delta\psi) dx dt \\ &= \int_0^T \left(\int_{\Omega} \nabla(u_{tt} + \frac{\gamma}{\beta}u_t) \cdot \nabla\psi dx - \int_{\partial\Omega} (u_{tt} + \frac{\gamma}{\beta}u_t)\nabla\psi \cdot \mathbf{n} dx \right) dt \\ &= \int_0^T \left(\int_{\Omega} (-\nabla u_t \cdot \nabla\psi_t + \frac{\gamma}{\beta}\nabla u_t \cdot \nabla\psi) dx + \int_{\partial\Omega} \frac{1}{\beta}(\nabla u_t \cdot \mathbf{n})(\nabla\psi \cdot \mathbf{n}) dS \right) dt \\ & \quad + \left[\int_{\Omega} \nabla u_t \cdot \nabla\psi dx \right]_0^T. \end{aligned}$$

Moreover, the compatibility condition

$$0 = \int_0^T (\beta u_t + \gamma u + \nabla u \cdot \mathbf{n}) dt = \int_0^T (\gamma u + \nabla u \cdot \mathbf{n}) dt$$

(the last identity following from $u(T) = u(0)$) that will be needed to conclude the boundary condition $\beta u_t + \gamma u + \nabla u \cdot \mathbf{n} = 0$ from its time differentiated version, is taken care of by introducing an auxiliary function h and imposing its periodicity. This yields the variational formulation

$$\begin{aligned} & u \in X, \quad u(0) = u(T), \quad u_t(0) = u_t(T), \quad h(0) = h(T) \\ & \text{and for all } \psi \in H^1(0, T; H^1(\Omega)) \cap L^2(0, T; H^2(\Omega)), \quad \zeta \in L^2(0, T; L^2(\partial\Omega)) : \\ (6.1) \quad & \int_0^T \left(\int_{\Omega} \left(-\nabla u_t \cdot \nabla\psi_t + \frac{\gamma}{\beta}\nabla u_t \cdot \nabla\psi + \frac{1}{\alpha}(c^2\Delta u + b\Delta u_t + (\frac{\gamma}{\beta}\alpha - \mu)u_t)\Delta\psi \right) dx \right. \\ & \quad \left. + \int_{\partial\Omega} \frac{1}{\beta}(\nabla u_t \cdot \mathbf{n})(\nabla\psi \cdot \mathbf{n}) dS + \int_{\partial\Omega} (h_t - u_t - [\gamma + \mathbf{n} \cdot \nabla]u)\zeta dS \right) dt \\ & \quad + \left[\int_{\Omega} \nabla u_t \cdot \nabla\psi dx \right]_0^T = - \int_0^T \int_{\Omega} \frac{1}{\alpha}f\Delta\psi dx dt. \end{aligned}$$

Indeed, reversing the above integration by parts step in (6.1) yields

$$\begin{aligned} & u \in X, \quad u(0) = u(T), \quad u_t(0) = u_t(T), \quad \int_0^T [\gamma + \mathbf{n} \cdot \nabla]u dt = 0 \\ & \text{and for all } \psi \in H^1(0, T; H^1(\Omega)) \cap L^2(0, T; H^2(\Omega)) : \\ (6.2) \quad & \int_0^T \left(\int_{\Omega} \frac{1}{\alpha}(\alpha u_{tt} - c^2\Delta u - b\Delta u_t + \mu u_t - f)(-\Delta\psi) dx \right. \\ & \quad \left. + \int_{\partial\Omega} \frac{1}{\beta}(\beta u_{tt} + \gamma u_t + \nabla u_t \cdot \mathbf{n})(\nabla\psi \cdot \mathbf{n}) dS \right) dt = 0 \end{aligned}$$

Second, we employ a Galerkin method. For this purpose, we utilize the fact that the eigenfunctions $(\phi_n)_{n \in \mathbb{N}}$ of the impedance Laplacian $-\Delta_{\gamma}$, that is,

$$-\Delta\phi_k = \lambda_k\phi_k \text{ in } \Omega, \quad [\gamma + \mathbf{n} \cdot \nabla]\phi_k = 0 \text{ on } \partial\Omega,$$

form an orthogonal basis of $H^1(\Omega)$, an orthonormal basis of $L^2(\Omega)$ and with $\tilde{V}_n := \text{lin}\{\phi_1, \dots, \phi_n\}$, $\bigcup_{n \in \mathbb{N}} \tilde{V}_n$ is dense in $H^1(\Omega)$ [8]. As an ansatz space for h we use the sequence $(\eta_k)_{k \in \mathbb{N}}$ defined by $\eta_k = \text{tr}_{\partial\Omega} \phi_k$ which is indeed linearly independent by construction and whose span is dense in $L^2(\Omega)$, due to the trace theorem; this choice guarantees that the boundary trace of $u_{nt} + [\gamma + \mathbf{n} \cdot \nabla]$ (which coincides with the boundary trace of u_{nt} by construction of the basis functions ϕ_k) can be used as a test function later on in the energy estimates. Thus altogether, we arrive at the Galerkin ansatz space $V_n := \text{lin}\{\phi_1, \dots, \phi_n\} \times \text{lin}\{\eta_1, \dots, \eta_n\}$ whose union $\bigcup_{n \in \mathbb{N}} V_n$ is dense in $H^1(\Omega) \times L^2(\partial\Omega)$. Now, plugging the ansatz $(u_n(\cdot, t), h_n(t)) := \sum_{k=1}^n (\mathbf{a}_k(t)\phi_k, \mathbf{c}_k(t)\eta_k) \in V_n$ into (6.1) and testing with $\phi_j, \eta_j, j \in \{1, \dots, n\}$ ¹ yields the system of ODEs

$$(6.3) \quad \begin{bmatrix} I_{n \times n} & 0_{n \times n} & 0_{n \times n} \\ 0_{n \times n} & \mathbf{H} & 0_{n \times n} \\ 0_{n \times n} & 0_{n \times n} & \mathbf{D} \end{bmatrix} \dot{\bar{z}}(t) = \begin{bmatrix} 0_{n \times n} & I_{n \times n} & 0_{n \times n} \\ -\mathbf{C} & -\mathbf{G} & 0_{n \times n} \\ 0_{n \times n} & \mathbf{D} & 0_{n \times n} \end{bmatrix} \bar{z}(t) + \begin{bmatrix} 0 \\ \mathbf{F} \\ 0 \end{bmatrix}$$

for $\bar{z}(t) = (\mathbf{a}_1(t), \dots, \mathbf{a}_n(t), \dot{\mathbf{a}}_1(t), \dots, \dot{\mathbf{a}}_n(t), \mathbf{c}_1(t), \dots, \mathbf{c}_n(t))^T$, with periodicity conditions $\bar{z}(0) = \bar{z}(T)$. Here

$$\begin{aligned} \mathbf{F}_j &= -\left(\frac{1}{\alpha} f, \Delta \phi_j\right)_{L^2(\Omega)} \\ \mathbf{H}_{i,j} &= (\nabla \phi_i, \nabla \phi_j)_{L^2(\Omega)} \\ \mathbf{C}_{i,j} &= \left(\frac{1}{\alpha} c^2 \Delta \phi_i, \Delta \phi_j\right)_{L^2(\Omega)}, \\ \mathbf{G}_{i,j} &= \frac{\gamma}{\beta} (\nabla \phi_i, \nabla \phi_j)_{L^2(\Omega)} + \left(\frac{1}{\alpha} (b \Delta \phi_i + (\frac{\gamma}{\beta} \alpha - \mu) \phi_i), \Delta \phi_j\right)_{L^2(\Omega)} \\ &\quad + \frac{1}{\beta} ((\nabla \phi_i \cdot \mathbf{n}), (\nabla \phi_j \cdot \mathbf{n}))_{L^2(\partial\Omega)} \\ \mathbf{D}_{i,j} &= (\eta_i, \eta_j)_{L^2(\partial\Omega)} \end{aligned}$$

Here, the matrices \mathbf{H}, \mathbf{D} are positive definite.

We can thus restate the problem as follows

$$(6.4) \quad \begin{aligned} \dot{\bar{z}}(t) &= \begin{bmatrix} I_{n \times n} & 0_{n \times n} & 0_{n \times n} \\ 0_{n \times n} & \mathbf{H} & 0_{n \times n} \\ 0_{n \times n} & 0_{n \times n} & \mathbf{D} \end{bmatrix}^{-1} \left(\begin{bmatrix} 0_{n \times n} & I_{n \times n} & 0_{n \times n} \\ -\mathbf{C} & -\mathbf{G} & 0_{n \times n} \\ 0_{n \times n} & \mathbf{D} & 0_{n \times n} \end{bmatrix} \bar{z}(t) + \begin{bmatrix} 0 \\ \mathbf{F} \\ 0 \end{bmatrix} \right) \\ &= \tilde{\mathbf{A}}(t) \bar{z}(t) + \tilde{\mathbf{F}}(t), \end{aligned}$$

where $\tilde{\mathbf{A}}(0) = \tilde{\mathbf{A}}(T)$. It is readily checked that the conditions for the Floquet-Lyapunov Theorem (see [23, p. 90]) are fulfilled, since the system with vanishing f only has the zero solution (see the energy estimates below). Therefore, we obtain the existence of a periodic solution in $C^2(0, T; V_n)$ to (6.4). Note that (6.3) is equivalent to the spatially discretized version of (6.1) after reverting the partial integration with respect to time in the first term and setting $\psi(s) = \chi_{[0,t]}(s)\phi_j(s)$.

Third, we derive energy estimates providing the boundedness of the sequence of Galerkin solutions. We do so by testing the spatially discretized version of (6.1) with

¹Formally, after reversing the time integration by parts in the highest order term, we set $\psi(x, s) = \delta_t(s)\phi_j(x)$, $\zeta(x, s) = \delta_t(s)\eta_j(x)$; more precisely we set $\psi(x, s) = \chi_{[0,t]}(s)\phi_j(x)$, $\zeta(x, s) = \chi_{[0,t]}(s)\eta_j(x)$ and differentiate with respect to t then.

$\psi = \frac{\gamma}{2\beta}u_n + u_{nt}$, $\zeta = h_{nt} - u_{nt} - [\gamma + \mathbf{n} \cdot \nabla]u_n$. Using the periodicity of \bar{z} (that is, of u_n and u_{nt}) yields the energy identity

$$\begin{aligned} \text{lhs} &:= \frac{\gamma}{2\beta} \|\nabla u_{nt}\|_{L^2(0,T;L^2(\Omega))}^2 + \|\sqrt{\frac{b}{\alpha}} \Delta u_{nt}\|_{L^2(0,T;L^2(\Omega))}^2 + \frac{\gamma}{2\beta} \|\frac{c}{\sqrt{\alpha}} \Delta u_n\|_{L^2(0,T;L^2(\Omega))}^2 \\ &\quad + \frac{1}{\beta} \|\nabla u_{nt} \cdot \mathbf{n}\|_{L^2(0,T;L^2(\partial\Omega))}^2 + \|h_{nt} - u_{nt} - [\gamma + \mathbf{n} \cdot \nabla]u_n\|_{L^2(0,T;L^2(\partial\Omega))}^2 \\ &= \int_0^T \int_{\Omega} \left((\frac{1}{2\alpha})_t (c^2 + \frac{\gamma}{2\beta}b) (\Delta u_n)^2 - \frac{1}{\alpha} (f + (\frac{\gamma}{\beta}\alpha - \mu)u_{nt}) (\Delta u_{nt} + \frac{\gamma}{2\beta} \Delta u_n) \right) dx dt \\ &=: \text{rhs}. \end{aligned}$$

Here the right hand side can be further estimated by means of the Holder's and Young's inequality

$$\begin{aligned} |\text{rhs}| &\leq \|(\frac{1}{2\alpha})_t (c^2 + \frac{\gamma}{2\beta}b)\|_{L^\infty(0,T;L^\infty(\Omega))} \|\Delta u_n\|_{L^2(0,T;L^2(\Omega))}^2 \\ &\quad + \frac{\varepsilon_1 + \varepsilon_2}{2} \|\Delta u_{nt} + \frac{\gamma}{2\beta} \Delta u_n\|_{L^2(0,T;L^2(\Omega))}^2 \\ &\quad + \frac{1}{2\varepsilon_1} \|\frac{\gamma}{\beta} - \frac{\mu}{\alpha}\|_{L^\infty(0,T;L^{2q/(q-1)}(\Omega))} \|u_{nt}\|_{L^2(0,T;L^{2q}(\Omega))}^2 + \frac{1}{2\varepsilon_2} \|f\|_{L^2(0,T;L^2(\Omega))}^2 \end{aligned}$$

Due to our ansatz space setting and elliptic regularity, skipping the h_{nt} term, the left hand side can be bounded from below by

$$\begin{aligned} \text{lhs} &\geq \frac{\gamma}{2\beta} \|\nabla u_{nt}\|_{L^2(0,T;L^2(\Omega))}^2 + \|\sqrt{\frac{b}{\alpha}} \Delta u_{nt}\|_{L^2(0,T;L^2(\Omega))}^2 + \frac{\gamma}{2\beta} \|\frac{c}{\sqrt{\alpha}} \Delta u_n\|_{L^2(0,T;L^2(\Omega))}^2 \\ &\quad + \frac{1}{\beta} \|\nabla u_{nt} \cdot \mathbf{n}\|_{L^2(0,T;L^2(\partial\Omega))}^2 + \frac{\gamma}{2\beta} \|u_{nt}\|_{L^2(0,T;L^2(\partial\Omega))}^2 \\ &\geq \frac{1}{C} \|u_{nt}\|_{L^2(0,T;H^{3/2}(\Omega))}^2 + \|\Delta u_n\|_{L^2(0,T;L^2(\Omega))}^2 \end{aligned}$$

for some $C > 0$. Together with continuity of the embedding $H^{3/2}(\Omega) \subseteq L^{2q}(\Omega)$, $q \geq 1$, and the imposed bounds on the coefficients, this yields an estimate of the form

$$(6.5) \quad \|u_{nt}\|_{L^2(0,T;H^{3/2}(\Omega))}^2 + \|\Delta u_n\|_{L^2(0,T;L^2(\Omega))}^2 \leq C \|f\|_{L^2(0,T;L^2(\Omega))}^2.$$

On the other hand, we clearly also have

$$\text{lhs} \geq \|h_{nt} - u_{nt} - [\gamma + \mathbf{n} \cdot \nabla]u_n\|_{L^2(0,T;L^2(\partial\Omega))}^2,$$

from which together with the above estimate on u_n we can extract an estimate of the form

$$(6.6) \quad \|h_{nt}\|_{L^2(0,T;L^2(\partial\Omega))}^2 \leq C \|f\|_{L^2(0,T;L^2(\Omega))}^2.$$

To also obtain an estimate on u_{ntt} , we test the spatially discretized version of (6.1) with $(\psi_n(t), \zeta_n(t)) = ((-\Delta_\gamma)^{-1}u_{ntt}(t), 0) \in V_n$, and use the identity

$$\begin{aligned} &\int_0^T \int_{\partial\Omega} \frac{1}{\beta} (\beta u_{ntt} + \gamma u_{nt} + \nabla u_{nt} \cdot \mathbf{n}) (\nabla \psi_n \cdot \mathbf{n}) dS dt \\ &= -\gamma \int_0^T \int_{\partial\Omega} u_{ntt} \psi_n dS dt = \int_0^T \int_{\partial\Omega} \psi_n \nabla u_{nt} \cdot \mathbf{n} dS dt \end{aligned}$$

to arrive at (cf. (6.2))

$$\begin{aligned} \|u_{n\,tt}\|_{L^2(0,T;L^2(\Omega))}^2 &= - \left(\frac{1}{\alpha} \left(c^2 \Delta u_n + b \Delta u_{n\,t} - \mu u_{n\,t} + f \right), u_{n\,tt} \right)_{L^2(0,T;L^2(\Omega))} \\ &\quad + \int_0^T \langle \nabla u_{n\,t} \cdot \mathbf{n}, \psi_n \rangle_{H^{3/2}(\partial\Omega)^*, H^{3/2}(\partial\Omega)} dt, \end{aligned}$$

where due to the trace theorem and elliptic regularity

$$\|\psi_n(t)\|_{H^{3/2}(\partial\Omega)} \leq C_{\text{trace}} \|\Delta_\gamma \psi_n(t)\|_{L^2(\Omega)} = C_{\text{trace}} \|u_{n\,tt}(t)\|_{L^2(\Omega)}.$$

Applying the Cauchy-Schwarz inequality as well as (6.5) yields

$$(6.7) \quad \begin{aligned} &\|u_{n\,tt}\|_{L^2(0,T;L^2(\Omega))}^2 + \|u_{n\,t}\|_{L^2(0,T;H^{3/2}(\Omega))}^2 + \|\Delta u_n\|_{L^2(0,T;L^2(\Omega))}^2 \\ &\leq C \|f\|_{L^2(0,T;L^2(\Omega))}^2. \end{aligned}$$

Fourth, we take weak limits to construct a solution. The space $\tilde{X} := \{v \in H^2(0, T; L^2(\Omega)) \cap H^1(0, T; H^{3/2}(\Omega)) : \Delta v \in L^2(0, T; L^2(\Omega))\}$ induced by the energy estimates so far is a Hilbert space, thus reflexive according to the Riesz-Frechet representation theorem (see [27, p. 228]). The estimates derived before state that $(u_n)_{n \in \mathbb{N}}$ is a bounded sequence in \tilde{X} and by the Eberlein-Šmulian Theorem (see [27, p. 107]), there exists a weakly convergent subsequence $(u_{n_k})_{k \in \mathbb{N}} \subseteq (u_n)_{n \in \mathbb{N}}$ such that $u_{n_k} \rightharpoonup u \in \tilde{X}$ for $k \rightarrow \infty$. Due to linearity of the problem it is readily checked that u indeed satisfies (6.1) and by combining the above estimates and using weakly lower semicontinuity of the norm, we obtain

$$(6.8) \quad \|u\|_{\tilde{X}} \leq C \|f\|_{L^2(0,T;L^2(\Omega))}.$$

In fact u solves (3.2) in an $L^2(0, T; L^2(\Omega))$ sense via (6.2).

Fifth, we derive higher spatial regularity. Our goal is to show that a solution to (3.2) is contained in X . Therefore, we want to apply elliptic regularity theory (see [8, p. 326]). To this end, besides using the L^2 estimate on Δu resulting from (6.8), we have to take into account the regularity of the boundary values. Using the boundary conditions and again estimate (6.8), we have that $[\gamma + \mathbf{n} \cdot \nabla]u(t) = -\beta u_t$ is the trace of an $H^1(\Omega)$ function and by the trace theorem (see [8, p. 272]) and the estimates above we further obtain

$$(6.9) \quad \|[\gamma + \mathbf{n} \cdot \nabla]u\|_{L^2(0,T,H^{1/2}(\partial\Omega))} = \|\beta u_t\|_{L^2(0,T,H^{1/2}(\partial\Omega))} \leq \tilde{C} \|f\|_{L^2(0,T;L^2(\Omega))}.$$

Applying elliptic regularity for the impedance Laplacian yields the estimate

$$(6.10) \quad \|u\|_{L^2(0,T;H^2(\Omega))} \leq \tilde{C} \|f\|_{L^2(0,T;L^2(\Omega))}.$$

This concludes the proof.

REFERENCES

- [1] H. AMANN, *Linear and quasilinear parabolic problems, Vol. I: Abstract linear theory*, vol. I, 01 1995, <https://doi.org/10.1007/978-3-0348-9221-6>.
- [2] I. M. BABUSKA AND S. A. SAUTER, *Is the pollution effect of the FEM avoidable for the Helmholtz equation considering high wave numbers?*, SIAM Review, 42 (2000), pp. 451–484, <http://www.jstor.org/stable/2653302>.

- [3] I. BABUŠKA, F. IHLENBURG, E. T. PAIK, AND S. A. SAUTER, *A Generalized Finite Element Method for solving the Helmholtz equation in two dimensions with minimal pollution*, *Computer Methods in Applied Mechanics and Engineering*, 128 (1995), pp. 325–359, [https://doi.org/10.1016/0045-7825\(95\)00890-X](https://doi.org/10.1016/0045-7825(95)00890-X).
- [4] K. BAKER, L. BANJAI, AND M. PTASHNYK, *Numerical analysis of a time-stepping method for the Westervelt equation with time-fractional damping*, *Mathematics of Computation*, 93 (2024), <https://doi.org/10.1090/mcom/3945>.
- [5] R. T. BEYER, *Parameter of nonlinearity in fluids*, *The Journal of the Acoustical Society of America*, 32 (2005), pp. 719–721, <https://doi.org/10.1121/1.1908195>.
- [6] F. A. DUCK, *Nonlinear acoustics in diagnostic ultrasound*, *Ultrasound in medicine & biology*, 28 (2002), pp. 1–18.
- [7] B. ENNGQUIST AND A. MAJDA, *Absorbing boundary conditions for numerical simulation of waves*, *Proceedings of the National Academy of Sciences*, 74 (1977), pp. 1765–1766.
- [8] L. C. EVANS, *Partial differential equations*, *Graduate Studies in Mathematics*, American Mathematical Society, 2010, <https://bookstore.ams.org/gsm-19-r/>.
- [9] X. FAN, J. SHEN, AND D. ZHAO, *Sobolev embedding theorems for spaces $W^{k,p(x)}(\Omega)$* , *Journal of Mathematical Analysis and Applications*, 262 (2001), pp. 749–760, <https://doi.org/10.1006/jmaa.2001.7618>.
- [10] W. S. GAN, *B/A nonlinear parameter acoustical imaging*, Springer Singapore, Singapore, 2021, pp. 37–48, https://doi.org/10.1007/978-981-16-7015-2_6.
- [11] D. GIVOLI, *High-order local non-reflecting boundary conditions: a review*, *Wave Motion*, 39 (2004), pp. 319–326, <https://doi.org/10.1016/j.wavemoti.2003.12.004>. New computational methods for wave propagation.
- [12] D. GORDON, R. GORDON, AND E. TURKEL, *Compact high order schemes with gradient-direction derivatives for absorbing boundary conditions*, *Journal of Computational Physics*, 297 (2015), pp. 295–315, <https://doi.org/10.1016/j.jcp.2015.05.027>.
- [13] B. KALTENBACHER, *Periodic solutions and multiharmonic expansions for the Westervelt equation*, *Evolution Equations and Control Theory*, 10 (2021), pp. 229–247, <https://doi.org/10.3934/eect.2020063>.
- [14] B. KALTENBACHER AND I. LASIECKA, *Global existence and exponential decay rates for the Westervelt equation*, *Discrete and Continuous Dynamical Systems, Series S*, 2 (2009), p. 503.
- [15] B. KALTENBACHER AND W. RUNDELL, *Nonlinearity parameter imaging in the frequency domain*, 2023, <https://arxiv.org/abs/2303.09796>.
- [16] K. MIZOHATA AND S. UKAI, *The global existence of small amplitude solutions to the nonlinear acoustic wave equation*, *Journal of Mathematics of Kyoto University*, 33 (1993), pp. 505–522.
- [17] M. MUHR, V. NIKOLIĆ, AND B. WOHLMUTH, *Self-adaptive absorbing boundary conditions for quasilinear acoustic wave propagation*, *Journal of Computational Physics*, 388 (2019), pp. 279–299, <https://doi.org/10.1016/j.jcp.2019.03.025>.
- [18] K. A. NAUGOL'NYKH, L. A. OSTROVSKY, E. A. ZABOLOTSKAYA, AND M. A. BREAZEALE, *Nonlinear acoustics*, *The Journal of the Acoustical Society of America*, 99 (1996), pp. 1815–1815, <https://doi.org/10.1121/1.415329>.
- [19] V. NIKOLIĆ, *Local existence results for the Westervelt equation with nonlinear damping and Neumann as well as absorbing boundary conditions*, *Journal of Mathematical Analysis and Applications*, 427 (2015), pp. 1131–1167, <https://doi.org/10.1016/j.jmaa.2015.02.076>.
- [20] V. NIKOLIĆ AND B. WOHLMUTH, *A priori error estimates for the finite element approximation of westervelt's quasi-linear acoustic wave equation*, *SIAM Journal on Numerical Analysis*, 57 (2019), pp. 1897–1918, <https://doi.org/10.1137/19M1240873>.
- [21] A. V. OPPENHEIM, *Discrete-time signal processing*, Pearson Education India, 1999.
- [22] C. PANTEA, C. F. OSTERHOUDT, AND D. N. SINHA, *Determination of acoustical nonlinear parameter β of water using the finite amplitude method*, *Ultrasonics*, 53 (2013), pp. 1012–1019, <https://doi.org/10.1016/j.ultras.2013.01.008>.
- [23] G. SCHMIDT, *Linear differential equations with periodic coefficients*, *Journal of Applied Mathematics and Mechanics*, 56 (1976), pp. 222–222, <https://doi.org/10.1002/zamm.19760560516>.
- [24] I. SHEVCHENKO AND B. WOHLMUTH, *Self-adapting absorbing boundary conditions for the wave equation*, *Wave Motion*, 49 (2012), pp. 461–473, <https://doi.org/10.1016/j.wavemoti.2011.12.007>.
- [25] M. SOLOVCHUK, T. W. SHEU, AND M. THIRIET, *Simulation of nonlinear Westervelt equation for the investigation of acoustic streaming and nonlinear propagation effects*, *The Journal of the Acoustical Society of America*, 134 (2013), pp. 3931–3942.
- [26] B. STURTEVANT, C. PANTEA, AND D. SINHA, *Determination of the Acoustic Nonlinearity*

- Parameter in Liquid Water up to 250 degrees C and 14 MPa*, 10 2012, pp. 285–288, <https://doi.org/10.1109/ULTSYM.2012.0070>.
- [27] D. WERNER, *Functional Analysis*, Springer, 2011, <http://www.math.fu-berlin.de>.
- [28] J. WÓJCIK, *Analytical solution of the nonlinear equations of acoustic in the form of gaussian beam*, *Ultrasonics*, 122 (2022), p. 106687, <https://doi.org/10.1016/j.ultras.2022.106687>.
- [29] D. ZHANG, X. FEN GONG, AND X. CHEN, *Experimental imaging of the acoustic nonlinearity parameter B/A for biological tissues via a parametric array*, *Ultrasound in Medicine & Biology*, 27 (2001), pp. 1359–1365, [https://doi.org/10.1016/S0301-5629\(01\)00432-X](https://doi.org/10.1016/S0301-5629(01)00432-X).