

# Extensions to the Navier-Stokes-Fourier Equations for Rarefied Transport: Variational Multiscale Moment Methods for the Boltzmann Equation

FA Baidoo<sup>1</sup>, IM Gamba<sup>1</sup>, TJR Hughes<sup>1</sup>, MRA Abdelmalik<sup>2</sup>

<sup>1</sup>*Oden Institute for Computational Sciences and Engineering, University of Texas at Austin*

<sup>2</sup>*Department of Mechanical Engineering, Eindhoven University of Technology*

## Abstract

We derive a fourth order entropy stable extension of the Navier-Stokes-Fourier equations into the transition regime of rarefied gases. We do this through a novel reformulation of the closure of conservation equations derived from the Boltzmann equation that subsumes existing methods such as the Chapman-Enskog expansion. We apply the linearized version of this extension to the stationary heat problem and the Poiseuille channel and compare our analytical solutions to asymptotic and numerical solutions of the linearized Boltzmann equation. In both model problems, our solutions compare remarkably well in the transition regime. For some macroscopic variables, this agreement even extends far beyond the transition regime.

## 1 Introduction

Gases in the transition regime have a mean free path that is large enough relative to macroscopic reference length scales to render continuum models like the Navier-Stokes-Fourier equations invalid while still being small enough that statistical methods such as Direct Simulation Monte Carlo are computationally expensive. This regime finds applications that span many orders of magnitude from the flows around objects in the thin atmosphere of near earth orbit to flows within microchannels of microelectromechanical systems.

Perhaps the most storied macroscopic model for gases in this regime is the Burnett equations. These third order moment equations are derived from the Chapman-Enskog expansion of the Boltzmann equation, where they arise as a correction to the Navier-Stokes-Fourier equations. Unfortunately, the Burnett equations prove to be a far from ideal extension into the transition regime. For example, Bobylev [1, 2] showed that the Burnett equations are unstable with respect to short wavelength perturbations while Comeaux et al. [3] showed that the Burnett equations violate the second law of thermodynamics at large enough Knudsen numbers. The Burnett equations have also been shown to produce unphysical stationary solutions [4].

Many modifications have been made to the Burnett equations to overcome these shortcomings. To name a few, Zhong [5] adds new terms to improve stability creating the augmented Burnett equations while Jin and Slemrod [6] propose relaxing the deviatoric pressure and heat flux with rate equations. Bobylev [2, 7] advocates for a transformation of the vector of hydrodynamic variables in order to obtain generalized Burnett equations

which do not suffer from the instabilities of the original. If we look beyond modifications to the Burnett equations and also consider alternatives to it, we can also talk about moment methods that build on the work of Grad [8]. For example, the regularized 13 (and 26) moment equations of Struchtrup and Torrilhon [9] are still an area of active theoretical [10–12] and numerical research [13, 14].

In this paper, we will use the framework of the Variational Multiscale (VMS) method to derive an alternative to the Burnett equations by modifying the process by which we obtain constitutive relations for the deviatoric stress and heat flux in the conservation equations obtained from the Boltzmann equation. The VMS method [15–17] was originally created as a framework for deriving stable finite element schemes for highly advective partial differential equations and, later, for closures for turbulence modeling [18]. If we think of the conservation equations (or moment equations more generally) as a weak form in microscopic velocity of the Boltzmann equation, then it is not too surprising that an idea from the finite element method can be brought to bear on the problem of finding closures. The methodology is conceptually simple. Assuming that the distribution is a perturbation of a Maxwellian, we separate the Boltzmann equation into a finite-dimensional coarse scale equation from which the macroscopic equations will arise and an infinite dimensional fine-scale equation which encodes the macroscopic variables we seek closures for. Within this framework, deriving a closure amounts to substituting an approximation of the fine-scale equation solution into the coarse-scale equation. The framework is general enough that it can be used to describe the Chapman-Enskog expansion whilst opening the door to considering new closures. For our particular alternative to the Burnett equation, we take advantage of the fact the coarse-scale equation also gives an equation for the macroscopic entropy of the system of conservation equations. We create an approximation of the fine-scale equation that ensures that the conservation equations derived are entropy stable. In our alternative to the Burnett equations, the momentum and energy equations are fourth order extensions to the Navier-Stokes-Fourier equations.

The paper is organized as follows. In section 2, we introduce the Boltzmann equation and the conservation equations that arise from it. We discuss the collision operator (with a special emphasis on the linearized collision operator) and the properties that make the analysis that follows possible. The material is presented in this section is well-known but is presented here in order for the work to be self-contained.<sup>1</sup> In section 3, we elaborate on the VMS moment method for deriving closures. We illustrate the process with the Chapman-Enskog expansion, showing that the Euler and Navier-Stokes-Fourier equations are entropy stable whilst the Burnett equations and higher are not. We then introduce the methodology to generate an entropy stable alternative to the Burnett equations. In section 4, we apply the linearized version of the extension to the stationary heat problem and the Poiseuille channel problem and compare the analytical solutions generated to asymptotic and numerical solutions to the Boltzmann equation found in [22, 23]. Supplementary calculations can be found in the Appendices at the end.

## 2 The Boltzmann Equation

Consider the distribution function  $F = F(t, \mathbf{x}, \mathbf{v})$  that at time  $t \in \mathbb{R}_{\geq 0}$  and position  $\mathbf{x} = (x_1, \dots, x_D) \in \mathbb{R}^D$  gives the density of particles of a monatomic fluid with velocity  $\mathbf{v} = (v_1, \dots, v_D) \in \mathbb{R}^D$ , where  $D$  is the number

---

<sup>1</sup>The reader can consult the works such as [19–21] for a more in depth and technical presentation of the material in section 2.

of dimensions. The evolution of  $F$  is governed by the Boltzmann equation

$$\text{St} \partial_t F(t, \mathbf{x}, \mathbf{v}) + v_i \partial_{x_i} F(t, \mathbf{x}, \mathbf{v}) = \frac{1}{\epsilon} \mathcal{C}(F)(t, \mathbf{x}, \mathbf{v}) \quad \text{where } (t, \mathbf{x}, \mathbf{v}) \in \mathbb{R}_+ \times \mathbb{R}^D \times \mathbb{R}^D, \quad (1)$$

where the collision operator  $\mathcal{C}(F)$  acts globally on the velocity  $\mathbf{v}$  dependence of  $F$  but locally in time  $t$  and position  $\mathbf{x}$ . The dimensionless number  $\text{St}$  is called the kinetic Strouhal number, defined as the ratio of a reference macroscopic length scale of the gas to the product of the *thermal speed of sound* of the gas (defined later) and a reference time scale. The dimensionless number  $\epsilon$  is the Knudsen number, defined as the ratio of the mean free path of the gas to a reference macroscopic length scale. In the above equation and throughout the text, we employ the convention that repeated indices imply summation up to dimension  $D$ .

## 2.1 Macroscopic Observables

Macroscopic fluid variables are obtained from  $F(t, \mathbf{x}, \mathbf{v})$  by taking its moments in the velocity variable:

$$\int_{\mathbb{R}^D} F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} = \rho \quad (2a)$$

$$\int_{\mathbb{R}^D} v_i F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} = \rho u_i \quad (2b)$$

$$\int_{\mathbb{R}^D} v_i v_j F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} = \rho u_i u_j + P_{ij} \quad (2c)$$

$$\int_{\mathbb{R}^D} v_i v_j v_k F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} = \rho u_i u_j u_k + u_i P_{jk} + u_j P_{ik} + u_k P_{ij} + Q_{ijk} \quad (2d)$$

where  $\rho = \rho(t, \mathbf{x})$ ,  $u_i = u_i(t, \mathbf{x})$  are the macroscopic mass and bulk velocity vector, respectively. In addition, the macroscopic stress tensor  $P_{ij} = P_{ij}(t, \mathbf{x})$  and the heat flux tensor  $Q_{ijk} = Q_{ijk}(t, \mathbf{x})$  are derived from the second and third moments of  $F$  with respect to random deviations of the particle velocities, that is

$$P_{ij}(t, \mathbf{x}) = \int_{\mathbb{R}^D} (v_i - u_i)(v_j - u_j) F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} \quad (3)$$

$$Q_{ijk}(t, \mathbf{x}) = \int_{\mathbb{R}^D} (v_i - u_i)(v_j - u_j)(v_k - u_k) F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} \quad (4)$$

The stress tensor is decomposed into an isotropic pressure  $p(t, \mathbf{x}) = \frac{1}{D} \text{Tr}(P_{ij})$  and the accompanying deviatoric stress tensor  $\sigma_{ij}(t, \mathbf{x}) = p \delta_{ij} - P_{ij}$ . With the internal energy per degree of freedom given by

$$\rho \theta(t, \mathbf{x}) = \frac{1}{D} \int_{\mathbb{R}^D} |\mathbf{v} - \mathbf{u}|^2 F d\mathbf{v} \quad (5)$$

we thus have that  $p = \rho \theta$ . For an ideal gas, this means that  $\theta(t, \mathbf{x})$  is the re-scaled temperature i.e.  $\theta = \frac{k_B T}{m}$ , where  $k_B = 1.38 \cdot 10^{-23} \text{JK}^{-1}$  is the Boltzmann constant and  $m$  is the mass of a gas particle. As such, given a reference temperature  $\theta_0$ , the thermal speed of sound of the gas is given by  $\sqrt{\gamma \theta_0}$  where  $\gamma$  is the adiabatic exponent. Furthermore, we may infer relations for the total energy and total energy flux by taking the traces of (2c) and (4), respectively:

$$\frac{1}{2} \int_{\mathbb{R}^D} |\mathbf{v}|^2 F d\mathbf{v} = \frac{1}{2} \rho |\mathbf{u}|^2 + \frac{D}{2} \rho \theta, \quad (6a)$$

$$\frac{1}{2} \int_{\mathbb{R}^D} |\mathbf{v}|^2 v_i F d\mathbf{v} = \frac{1}{2} \rho |\mathbf{u}|^2 u_i + \frac{(D+2)}{2} \rho \theta u_i - \sigma_{ij} u_j + q_i, \quad (6b)$$

where the vector  $q_i = Q_{ijj}/2$  denotes the heat flux vector.

For our purposes, it is also necessary to consider the situation where our distribution  $F$  is close to a stationary equilibrium state. In particular,

$$\rho(t, \mathbf{x}) = \rho_0 + \epsilon \tilde{\rho}(t, \mathbf{x})$$

$$\mathbf{u}(t, \mathbf{x}) = \epsilon \tilde{\mathbf{u}}(t, \mathbf{x})$$

$$\theta(t, \mathbf{x}) = \theta_0 + \epsilon \tilde{\theta}(t, \mathbf{x})$$

where  $\rho_0$  and  $\theta_0$  are a constant background density and temperature. In this situation, we consider the linearization of the macroscopic parameters (2) with respect to  $\epsilon$ :

$$\int_{\mathbb{R}^D} F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} = \rho_0 + \epsilon \tilde{\rho} \quad (7a)$$

$$\int_{\mathbb{R}^D} v_i F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} = \epsilon \rho_0 \tilde{u}_i \quad (7b)$$

$$\int_{\mathbb{R}^D} v_i v_j F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} = \rho_0 \theta_0 \delta_{ij} + \epsilon (\rho_0 \tilde{\theta} + \tilde{\rho} \theta_0) \delta_{ij} - \epsilon \tilde{\sigma}_{ij} \quad (7c)$$

$$\int_{\mathbb{R}^D} |\mathbf{v}|^2 v_i F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} = \epsilon (D+2) \rho_0 \theta_0 \tilde{u}_i + 2 \epsilon \tilde{q}_i \quad (7d)$$

where  $\epsilon \tilde{\sigma}_{ij}$  and  $\epsilon \tilde{q}_i$  represent the linearizations of  $\sigma_{ij}$  and  $q_i$  respectively.

## 2.2 The Collision Operator

The collision operator  $\mathcal{C}(F)$  is defined on a set of functions  $\mathcal{D}_v(\mathcal{C}) \subset L^1_2(\mathbb{R}_v^D) \cap L \log L(\mathbb{R}_v^D)$  where  $L^1_2(\mathbb{R}_v^D) := \{F : \int_{\mathbb{R}^D} (1 + |\mathbf{v}|^2) F(t, \mathbf{x}, \mathbf{v}) d\mathbf{v} < \infty\}$  and  $L \log L(\mathbb{R}_v^D) := \{F : F \log(F) d\mathbf{v} < \infty\}$ . Given any vector  $\mathbf{U} \in \mathbb{R}^D$  and orthonormal matrix  $\mathbf{O} \in \mathbb{R}^{D \times D}$ , translations and rotations in the  $\mathbf{v}$  variable are defined by  $(\mathcal{T}_U F)(\mathbf{v}) := F(\mathbf{v} - \mathbf{U})$  and  $(\mathcal{T}_O F)(\mathbf{v}) = F(\mathbf{O}^T \mathbf{v})$ . It will be assumed that for all  $F \in \mathcal{D}_v(\mathcal{C})$ , we also have that  $\mathcal{T}_U F$  and  $\mathcal{T}_O F$  are contained in  $\mathcal{D}_v(\mathcal{C})$ .

For our purposes, we will require that the collision operator satisfies certain properties we shall outline in the intervening subsections. These properties are well-known (see, for instance [24]) but are presented here to ensure coherence in the text.

**Property 1.** *We require that the collision operator commutes with translations and rotations, that is*

$$\mathcal{T}_U \mathcal{C}(F) = \mathcal{C}(\mathcal{T}_U F), \quad \text{and} \quad \mathcal{T}_O \mathcal{C}(F) = \mathcal{C}(\mathcal{T}_O F). \quad (8)$$

The invariances (8) are consistent with the Hamiltonian dynamics of the advective material derivative in the left hand side of the Boltzmann equation (1), meaning whenever  $F(t, \mathbf{x}, \mathbf{v})$  solves this equation, then so do  $F(t, \mathbf{x} - (\mathbf{v} - \mathbf{U})t, \mathbf{v} - \mathbf{U})$  and  $F(t, \mathbf{x} - \mathbf{O}^T \mathbf{v} t, \mathbf{O}^T \mathbf{v})$ .

### 2.2.1 Collision Invariance

A scalar-valued function  $\psi(\mathbf{v})$  is a collision invariant of  $\mathcal{C}$  if

$$\int_{\mathbb{R}^D} \psi \mathcal{C}(F) d\mathbf{v} = 0 \quad \forall F \in \mathcal{D}_v(\mathcal{C}), \quad (9)$$

The relation in (9) associates a scalar conservation law to (1) with each collision invariant:

$$\text{St } \partial_t \int_{\mathbb{R}^D} \psi F d\mathbf{v} + \partial_{x_i} \int_{\mathbb{R}^D} v_i \psi F d\mathbf{v} = 0 \quad (10)$$

**Property 2.** *The space of collision invariants of  $\mathcal{C}$  is  $\mathcal{I} := \text{span}\{1, v_1, \dots, v_D, |\mathbf{v}|^2\}$ . In other words,*

$$\int_{\mathbb{R}^D} \psi(\mathbf{v}) \mathcal{C}(F) d\mathbf{v} = 0 \quad \forall F \in \mathcal{D}_{\mathbf{v}}(\mathcal{C}) \Leftrightarrow \psi(\mathbf{v}) \in \mathcal{I}. \quad (11)$$

Property 2 implies that solutions of the Boltzmann equation (1) obey the conservation laws for compressible gas flow systems<sup>2</sup>

$$\text{St } \partial_t \rho + \partial_{x_i} (\rho u_i) = 0; \quad (12a)$$

$$\text{St } \partial_t (\rho u_j) + \partial_{x_i} (\rho u_i u_j + \rho \theta \delta_{ij} - \sigma_{ij}) = 0; \quad (12b)$$

$$\text{St } \partial_t \left( \frac{1}{2} \rho |\mathbf{u}|^2 + \frac{D}{2} \rho \theta \right) + \partial_{x_i} \left( \left( \frac{1}{2} \rho |\mathbf{u}|^2 u_i + \frac{D}{2} \rho \theta \right) u_i + \rho \theta u_i - \sigma_{ij} u_j + q_i \right) = 0; \quad (12c)$$

i.e. solutions of (1) conserve mass according to (12a), momentum according to (12b), and energy according to (12c). Note that the system of conservation laws (12) is not closed since there are  $(D^2 + 5D + 2)/2$  variables and only  $D + 2$  relations. The closure of (12) requires a constitutive modeling assumption that characterizes, both  $\sigma_{ij}$  and  $q_i$ , in terms of the proposed unknowns  $\rho$ ,  $u_i$  and  $\theta$ . Using equations (3), (4) and (6b), one finds that

$$\sigma_{ij} = - \int_{\mathbb{R}^D} \left( (v_i - u_i)(v_j - u_j) - \frac{1}{D} |\mathbf{v} - \mathbf{u}|^2 \delta_{ij} \right) F d\mathbf{v}, \quad (13a)$$

$$q_i = \frac{1}{2} \int_{\mathbb{R}^D} (|\mathbf{v} - \mathbf{u}|^2 (v_i - u_i) - (D + 2)\theta (v_i - u_i)) F d\mathbf{v}, \quad (13b)$$

Thus closures for (12) can be found by approximating  $F$  as a function that is solely parametrized by  $\rho$ ,  $u_i$  and  $\theta$  along with their derivatives.

The above dynamic holds true when we linearize about a constant equilibrium state. We obtain linearized conservation equations:

$$\text{St } \partial_t \tilde{\rho} + \rho_0 \partial_{x_i} \tilde{u}_i = 0; \quad (14a)$$

$$\text{St } \rho_0 \partial_t \tilde{u}_j + \partial_{x_i} (\tilde{\rho} \theta_0 \delta_{ij} + \rho_0 \tilde{\theta} \delta_{ij} - \tilde{\sigma}_{ij}) = 0; \quad (14b)$$

$$\frac{D}{2} \text{St } \rho_0 \partial_t \tilde{\theta} + \partial_{x_i} (\rho_0 \theta_0 \tilde{u}_i + \tilde{q}_i) = 0; \quad (14c)$$

where

$$\epsilon \tilde{\sigma}_{ij} = - \int_{\mathbb{R}^D} \left( v_i v_j - \frac{1}{D} |\mathbf{v}|^2 \delta_{ij} \right) F d\mathbf{v}, \quad (15a)$$

$$\epsilon \tilde{q}_i = \frac{1}{2} \int_{\mathbb{R}^D} (|\mathbf{v}|^2 v_i - (D + 2)\theta_0 v_i) F d\mathbf{v}, \quad (15b)$$

and closures for (14) can be found by approximating  $F$  as a function that is solely parametrized by  $\tilde{\rho}$ ,  $\tilde{u}_i$  and  $\tilde{\theta}$  along with their derivatives.

---

<sup>2</sup>The reader can recall that the classical adiabatic exponent  $\gamma = \gamma(D)$  depends on the space dimension  $D$ . For monoatomic gases this relation is  $\gamma = 1 + 2/D$ , recovering the familiar version of these conservation equations in terms of the adiabatic number.

### 2.2.2 $\mathcal{H}$ -theorem: Entropy Dissipation and Equilibria

**Property 3.**  $\mathcal{C}$  satisfies the local dissipation relation

$$\int_{\mathbb{R}^D} \ln(F) \mathcal{C}(F) d\mathbf{v} \leq 0 \quad \forall F \in \mathcal{D}(\mathcal{C}). \quad (16)$$

Relation (16) leads to a statement of Boltzmann's  $\mathcal{H}$ -theorem. This is accomplished by weighting the Boltzmann equation (1) with  $\ln(F)$  and integrating on the entire velocity space in order to obtain the local entropy-dissipation law:

$$\text{St } \partial_t \int_{\mathbb{R}^D} (F \ln(F) - F) d\mathbf{v} + \partial_{x_i} \int_{\mathbb{R}^D} v_i (F \ln(F) - F) d\mathbf{v} = \frac{1}{\epsilon} \int_{\mathbb{R}^D} \ln(F) \mathcal{C}(F) d\mathbf{v} \leq 0 \quad (17)$$

It is also important that when the Boltzmann equation is at equilibrium ( $\mathcal{C}(F) = 0$ ), the entropy equation (17) also be at equilibrium. As such

**Property 4.**

$$\int_{\mathbb{R}^D} \ln(F) \mathcal{C}(F) d\mathbf{v} = 0 \implies \mathcal{C}(F) = 0 \quad (18)$$

Using (11) we have that equality in (16) holds if and only if  $\ln(F) \in \mathcal{S}$  in accordance with (11). As a result,

$$\mathcal{C}(F) = 0 \iff \int_{\mathbb{R}^D} \ln(F) \mathcal{C}(F) d\mathbf{v} = 0 \iff \ln(F) \in \mathcal{S} \quad (19)$$

By virtue of (11), the second equivalence indicates that the form of such local equilibria is given by

$$F = e^{a+b_i v_i + c|\mathbf{v}|^2} \quad (20)$$

for some coefficients  $a(t, \mathbf{x})$ ,  $b_i(t, \mathbf{x})$  and  $c(t, \mathbf{x})$ . Furthermore, integrability implies that  $c < 0$ .

**Remark 2.1.** Without loss of generality we may reparameterize (20) as a Maxwellian distribution

$$\mathcal{M}_{\rho, \mathbf{u}, \theta}(\mathbf{v}) := \frac{\rho}{(2\pi\theta)^{D/2}} \exp\left(-\frac{|\mathbf{v} - \mathbf{u}|^2}{2\theta}\right) \quad (21)$$

for some  $(\rho, \mathbf{u}, \theta) \in \mathbb{R}_+ \times \mathbb{R}^D \times \mathbb{R}_+$ .

For the rest of the text, we shall use  $\mathcal{M}$  to refer to any arbitrary Maxwellian and only include the subscript when we want to emphasize the Maxwellian's parameterization by  $\rho$ ,  $\mathbf{u}$  and  $\theta$ . Furthermore, the mass, momentum and energy moments of a distribution  $F$ , given by Equations (2a), (2b) and (6a) respectively, can be used to derive what is called the self-consistent Maxwellian which we shall denote by  $\mu(F) := \mathcal{M}_{\rho_\mu, \mathbf{u}_\mu, \theta_\mu}$  where

$$\begin{pmatrix} \rho_\mu(F) \\ \rho_\mu(F) \mathbf{u}_\mu(F) \\ \rho_\mu(F) |\mathbf{u}_\mu(F)|^2 + D\rho_\mu(F)\theta_\mu(F) \end{pmatrix} := \begin{pmatrix} \int_{\mathbb{R}^D} F d\mathbf{v} \\ \int_{\mathbb{R}^D} \mathbf{v} F d\mathbf{v} \\ \int_{\mathbb{R}^D} |\mathbf{v}|^2 F d\mathbf{v} \end{pmatrix} = \begin{pmatrix} \int_{\mathbb{R}^D} \mu(F) d\mathbf{v} \\ \int_{\mathbb{R}^D} \mathbf{v} \mu(F) d\mathbf{v} \\ \int_{\mathbb{R}^D} |\mathbf{v}|^2 \mu(F) d\mathbf{v} \end{pmatrix}. \quad (22)$$

In other words,  $\mu(F)$  is the Maxwellian that has the same  $\rho(t, \mathbf{x})$ ,  $u_i(t, \mathbf{x})$  and  $\theta(t, \mathbf{x})$  as  $F$ . Because  $\rho(t, \mathbf{x})$ ,  $u_i(t, \mathbf{x})$  and  $\theta(t, \mathbf{x})$  completely determine a Maxwellian, we also have that  $F = \mu(F)$  if and only if  $F$  is a Maxwellian.

## 2.3 The Linearized Collision Operator

To derive macroscopic fluid equations from (1) near equilibrium, we assume that the distribution  $F$  is a perturbation in Knudsen number from some Maxwellian.

$$F(t, \mathbf{x}, \mathbf{v}) = \mathcal{M}(t, \mathbf{x}, \mathbf{v}) (1 + \epsilon f(t, \mathbf{x}, \mathbf{v})). \quad (23)$$

A corresponding Taylor series expansion of the collision operator about the Maxwellian  $\mathcal{M}$  gives

$$\mathcal{C}(\mathcal{M}(1 + \epsilon f)) = \epsilon \frac{d}{d\epsilon} [\mathcal{C}(\mathcal{M}(1 + \epsilon f))] \Big|_{\epsilon=0} + \frac{\epsilon^2}{2} \frac{d^2}{d\epsilon^2} [\mathcal{C}(\mathcal{M}(1 + \epsilon f))] \Big|_{\epsilon=0} + O(\epsilon^3). \quad (24)$$

Substituting (23) and (24) into the Boltzmann equation (1) gives

$$(\text{St} \partial_t + v_i \partial_{x_i})[\mathcal{M} + \epsilon \mathcal{M}f] = \mathcal{M} \mathcal{L}_{\mathcal{M}}[f] + \frac{\epsilon}{2} \frac{d^2}{d\epsilon^2} [\mathcal{C}(\mathcal{M}(1 + \epsilon f))] \Big|_{\epsilon=0} + O(\epsilon^2) \quad (25)$$

where the linearized collision operator

$$\mathcal{L}_{\mathcal{M}}[f] := \frac{1}{\mathcal{M}} \frac{d}{d\epsilon} [\mathcal{C}(\mathcal{M}(1 + \epsilon f))] \Big|_{\epsilon=0} \quad (26)$$

is of particular importance and thus requires further study.

**Remark 2.2.** For the bilinear collision operators  $\mathcal{C}(F) = \mathcal{Q}(F, F)$ , the linearized collision operator takes the form

$$\mathcal{L}_{\mathcal{M}}^{\mathcal{Q}}[f] = \frac{1}{\mathcal{M}} \left( \mathcal{Q}(\mathcal{M}f, \mathcal{M}) + \mathcal{Q}(\mathcal{M}, \mathcal{M}f) \right)$$

For example, the Hard Spheres operator will be

$$\mathcal{L}_{\mathcal{M}}^{\text{HS}}[f] = \frac{\epsilon}{2\pi\rho_0\ell\sqrt{2}} \int_{\mathbb{R}^3 \times \mathbb{S}^2} \mathcal{M}(\mathbf{v}_*) \left( f(\mathbf{v}') + f(\mathbf{v}'_*) - f(\mathbf{v}_*) - f(\mathbf{v}) \right) |\hat{\boldsymbol{\eta}} \cdot (\mathbf{v}_* - \mathbf{v})| d\hat{\boldsymbol{\eta}} d\mathbf{v}_*$$

where  $\ell$  is the mean free path of the gas<sup>3</sup>,  $\mathbf{v}' = \mathbf{v} + (\hat{\boldsymbol{\eta}} \cdot (\mathbf{v}_* - \mathbf{v}))\hat{\boldsymbol{\eta}}$  and  $\mathbf{v}'_* = \mathbf{v}_* - (\hat{\boldsymbol{\eta}} \cdot (\mathbf{v}_* - \mathbf{v}))\hat{\boldsymbol{\eta}}$ .

With the Bhatnagar-Gross-Krook (BGK) operator [22, 26],  $\mathcal{C}^{\text{BGK}}(F) = \frac{\mu(F) - F}{\tau(F)}$  where  $\tau(F)$  assumed to not depend on  $\mathbf{v}$ , the linearized collision operator will be

$$\mathcal{L}_{\mathcal{M}}^{\text{BGK}}[f] = -\frac{1}{\tau(\mathcal{M})} (\text{Id} - \Pi_{\mathcal{M}})[f] \quad (27)$$

where  $\text{Id}$  is the identity operator and given a Maxwellian  $\mathcal{M}$  with conservation moments  $(\varrho, \mathbf{w}, \vartheta)$  and for any function  $g \in \mathcal{L}^2(\mathcal{M}d\mathbf{v})$

$$\begin{aligned} \Pi_{\mathcal{M}}[g] := & \frac{1}{\varrho} \int_{\mathbb{R}^D} g \mathcal{M} d\mathbf{v} + \frac{\mathbf{v} - \mathbf{w}}{\varrho\vartheta} \cdot \int_{\mathbb{R}^D} (\mathbf{v} - \mathbf{w}) g \mathcal{M} d\mathbf{v} \\ & + \left( \frac{|\mathbf{v} - \mathbf{w}|^2}{2\vartheta} - \frac{D}{2} \right) \frac{2}{D\varrho} \int_{\mathbb{R}^D} \left( \frac{|\mathbf{v} - \mathbf{w}|^2}{2\vartheta} - \frac{D}{2} \right) g \mathcal{M} d\mathbf{v}, \quad (28) \end{aligned}$$

is an orthogonal projection from  $\mathcal{L}^2(\mathcal{M}d\mathbf{v})$  onto  $\mathcal{S}$ . A proof that the expression (28) is indeed an orthogonal projection can be found in Appendix B and the calculation to derive (27) can be found in Appendix C.

<sup>3</sup> $\ell = \frac{m}{d_m^2 \rho_0 \pi \sqrt{2}}$  where  $d_m$  is the diameter of the gas particle and  $m$  is the mass.

### 2.3.1 Properties of the Linearized Collision Operator

For our purposes, we require that the linearized collision operator satisfy a set of properties we will use extensively. Given an arbitrary Maxwellian  $\mathcal{M}_{\rho, \mathbf{u}, \theta}$ , we denote the space of functions that are square-integrable in  $\mathbf{v}$  when weighted by  $\mathcal{M}_{\rho, \mathbf{u}, \theta}$  by  $\mathcal{L}^2(\mathcal{M} d\mathbf{v})$ . We also denote by  $\mathcal{O}_{\mathcal{O}}$  the composition of operators  $\mathcal{T}_{\mathbf{u}} \mathcal{T}_{\mathcal{O}} \mathcal{T}_{\mathbf{u}}^{-1}$ . The linearized collision operators  $\mathcal{L}_{\mathcal{M}}$  we deal with are such that:

**Property 2.3.1.1.**  $\mathcal{L}_{\mathcal{M}} : \mathcal{L}^2(\mathcal{M} d\mathbf{v}) \rightarrow \mathcal{L}^2(\mathcal{M} d\mathbf{v})$  is a self-adjoint operator with respect to the weighted inner product of  $\mathcal{L}^2(\mathcal{M} d\mathbf{v})$ .

**Property 2.3.1.2.** The linearized collision operator commutes with  $\mathcal{O}_{\mathcal{O}}$  i.e.

$\mathcal{O}_{\mathcal{O}} \mathcal{L}_{\mathcal{M}}[f] = \mathcal{L}_{\mathcal{M}}[\mathcal{O}_{\mathcal{O}} f]$ . This property is inherited from the collision operator  $\mathcal{C}$  because if we apply (24) to both sides of (8) we get

$$\begin{aligned} \epsilon \mathcal{O}_{\mathcal{O}} \left[ \frac{d}{d\epsilon} \mathcal{C}(\mathcal{M}(1 + \epsilon f)) \Big|_{\epsilon=0} \right] + \frac{\epsilon^2}{2} \mathcal{O}_{\mathcal{O}} \left[ \frac{d^2}{d\epsilon^2} \mathcal{C}(\mathcal{M}(1 + \epsilon f)) \Big|_{\epsilon=0} \right] + O(\epsilon^3) \\ = \epsilon \frac{d}{d\epsilon} \mathcal{C} \left( \mathcal{O}_{\mathcal{O}} [\mathcal{M}(1 + \epsilon f)] \right) \Big|_{\epsilon=0} + \frac{\epsilon^2}{2} \frac{d^2}{d\epsilon^2} \mathcal{C}(\mathcal{O}_{\mathcal{O}} [\mathcal{M}(1 + \epsilon f)]) \Big|_{\epsilon=0} + O(\epsilon^3), \end{aligned} \quad (29)$$

Since  $\epsilon$  is arbitrary and  $\mathcal{O}_{\mathcal{O}} \mathcal{M} = \mathcal{M}$ , the property follows immediately. This property greatly simplifies the form in which the closures for the deviatoric stress and heat flux take.

**Property 2.3.1.3.**  $\mathcal{L}_{\mathcal{M}}$  is negative semi-definite on  $\mathcal{L}^2(\mathcal{M} d\mathbf{v})$ . This property is inherited from the dissipation property of the collision operator (Equation (16)) because

$$\int_{\mathbb{R}^D} f \mathcal{M} \mathcal{L}_{\mathcal{M}}[f] d\mathbf{v} = \frac{1}{2} \frac{d^2}{d\epsilon^2} \left[ \int_{\mathbb{R}^D} \ln(\mathcal{M} + \epsilon \mathcal{M} f) \mathcal{C}(\mathcal{M} + \epsilon \mathcal{M} f) d\mathbf{v} \Big|_{\epsilon=0} \right] \leq 0 \quad (30)$$

where the inequality in (30) follows from the fact that we attain a local maximum at  $\epsilon = 0$  whilst the equality in (30) can be obtained by a Taylor series expansion of the middle term

$$\frac{d^2}{d\epsilon^2} \left[ \int_{\mathbb{R}^D} \left( \overbrace{\ln(\mathcal{M})}^{\in \mathcal{I}} + \overbrace{\ln(1 + \epsilon f)}^{\epsilon f + O(\epsilon^2)} \right) \overbrace{\mathcal{C}(\mathcal{M} + \epsilon \mathcal{M} f)}^{\text{Equation (24)}} d\mathbf{v} \right] \Big|_{\epsilon=0} = \int_{\mathbb{R}^D} 2f \frac{d}{d\epsilon} \mathcal{C}(\mathcal{M}(1 + \epsilon f)) \Big|_{\epsilon=0} d\mathbf{v} \quad (31)$$

**Property 2.3.1.4.** The kernel of  $\mathcal{L}_{\mathcal{M}}$  is the space of collision invariants  $\mathcal{I}$ .

This motivates the decomposition  $\mathcal{L}^2(\mathcal{M} d\mathbf{v}) = \mathcal{I} \oplus \mathcal{I}^{\perp \mathcal{M}}$ , where  $\mathcal{I}^{\perp \mathcal{M}}$  is the orthogonal complement to the space of collision products with respect to the  $\mathcal{L}^2(\mathcal{M} d\mathbf{v})$  inner product.

**Property 2.3.1.5** (Fredholm Alternative). The linearized collision operator with domain restricted to  $\mathcal{I}^{\perp \mathcal{M}}$  is invertible. That is, the equation  $\mathcal{L}_{\mathcal{M}}[f] = g$  has a unique solution if and only if  $f, g \in \mathcal{I}^{\perp \mathcal{M}}$ . We denote the inverse by  $\mathcal{L}_{\mathcal{M}}^{-1}$  and note that it can be extended to the rest of  $\mathcal{L}^2(\mathcal{M} d\mathbf{v})$  if we assert  $\mathcal{I}$  as its kernel. Abusing notation, we will also denote the extension by  $\mathcal{L}_{\mathcal{M}}^{-1}$ . Thus for any  $g \in \mathcal{L}^2(\mathcal{M} d\mathbf{v})$ ,

$$\mathcal{L}_{\mathcal{M}} \mathcal{L}_{\mathcal{M}}^{-1}[g] = (\text{Id} - \Pi_{\mathcal{M}})[g]$$

where  $\Pi_{\mathcal{M}}$  is given by (28).

Of the properties outlined above, the Fredholm alternative property is the most restrictive since it does not apply to collision operators such as the soft-sphere collision operators. It is however known to hold for hard



potential and Maxwell molecule operators [8, 20, 27] as well as the BGK operator by virtue of it being a scaled orthogonal projection onto  $\mathcal{I}^{\perp\mathcal{M}}$ .

### 3 Variational Multiscale Moment Closures

The first step will be to apply the projection operator  $\Pi_{\mathcal{M}}$  to the equation

$$\frac{1}{\mathcal{M}} (\text{St } \partial_t + v_i \partial_{x_i}) [\mathcal{M} + \epsilon \mathcal{M} f] = \mathcal{L}_{\mathcal{M}}[f] \quad (32)$$

in order to separate it into a coarse-scale equation made up of the components of (32) from the space of collision invariants  $\mathcal{I}$  and a fine-scale equation made up of components of (32) from the orthogonal complement of  $\mathcal{I}^{\perp\mathcal{M}}$ :

$$\text{St } \partial_t \ln(\mathcal{M}) + \Pi_{\mathcal{M}} \left[ v_i \partial_{x_i} \ln(\mathcal{M}) + \frac{\epsilon}{\mathcal{M}} (\text{St } \partial_t + v_i \partial_{x_i}) [\mathcal{M} f] \right] = 0 \quad (33a)$$

$$(\text{Id} - \Pi_{\mathcal{M}}) \left[ v_i \partial_{x_i} \ln(\mathcal{M}) + \frac{\epsilon}{\mathcal{M}} (\text{St } \partial_t + v_i \partial_{x_i}) [\mathcal{M} f] \right] = \mathcal{L}_{\mathcal{M}}[f] \quad (33b)$$

When we test the first equation with an arbitrary element of  $\mathcal{I}$ , denoted by  $\bar{m}$ , in the  $\mathcal{L}(\mathcal{M} d\mathbf{v})$  inner product, we obtain conservation equations of the form:

$$\langle \bar{m}, \text{St } \partial_t \mathcal{M} \rangle + \langle \bar{m}, v_i \partial_{x_i} \mathcal{M} \rangle + \epsilon \langle \bar{m}, (\text{St } \partial_t + v_i \partial_{x_i}) [\mathcal{M} f] \rangle = 0 \quad (34)$$

Where we now use angular brackets to denote integration in  $\mathbf{v}$  over  $\mathbb{R}^D$

$$\langle h(\mathbf{v}), g(\mathbf{v}) \rangle := \int_{\mathbb{R}^D} h(\mathbf{v}) g(\mathbf{v}) d\mathbf{v}$$

The second equation can be re-written as

$$(\text{Id} - \Pi_{\mathcal{M}})[f] = \mathcal{L}_{\mathcal{M}}^{-1} (\text{Id} - \Pi_{\mathcal{M}}) \left[ v_i \partial_{x_i} \ln(\mathcal{M}) + \frac{\epsilon}{\mathcal{M}} (\text{St } \partial_t + v_i \partial_{x_i}) [\mathcal{M} f] \right] \quad (35)$$

giving an equation for the term  $(\text{Id} - \Pi_{\mathcal{M}})[f]$ . As will be elaborated upon in the coming sections, the basic idea for deriving variational multiscale moment closures is to substitute a suitable approximation of (35) into the conservation equation (34).

We work with two types of Maxwellian within the variational multiscale framework. The first is constant background  $M(\mathbf{v}) := \mathcal{M}_{\rho_0, \mathbf{0}, \theta_0}$ , which we will use to elaborate on the closure derivation process for the linearized Boltzmann equation. The second is the self-consistent Maxwellian  $\mu(F)$  to the distribution  $F$  which gives us closures for the conservation equations in their general form. For brevity, we shall denote this Maxwellian by  $\mu$  from now on. In both cases, the equations (33) simplify greatly.

**Remark 3.1.** Note that (32) leaves out the higher order terms like

$\left. \frac{d^2}{d\epsilon^2} [\mathcal{C}(\mathcal{M}(1 + \epsilon f))] \right|_{\epsilon=0}$ . This simplifies the analysis without fundamentally changing its form. In particular the coarse-scale equation (33a) remains unchanged whilst the fine-scale equation incorporates the higher-order

collision terms in a straightforward manner:

$$(Id - \Pi_{\mathcal{M}})[f] = \mathcal{L}_{\mathcal{M}}^{-1}(Id - \Pi_{\mathcal{M}}) \left[ v_i \partial_{x_i} \ln(\mathcal{M}) + \frac{\epsilon}{\mathcal{M}} (St \partial_t + v_i \partial_{x_i}) [\mathcal{M}f] - \frac{\epsilon}{2\mathcal{M}} \frac{d^2}{d\epsilon^2} [\mathcal{C}(\mathcal{M}(1 + \epsilon f))] \Big|_{\epsilon=0} + O(\epsilon^2) \right] \quad (36)$$

Thus a more complete closure procedure will involve adding terms that incorporate these higher order collision terms to whatever approximation of (35) we come up with.

### 3.1 Linear Theory: The Constant Background Maxwellian Formulation

With  $\mathcal{M} = M$ , we obtain the linearized Boltzmann equation from (32)

$$(St \partial_t + v_i \partial_{x_i}) [f] = \frac{1}{\epsilon} \mathcal{L}_M [f] \quad (37)$$

Noting that  $\langle \bar{m}, M \partial_t (Id - \Pi_M)[f] \rangle = 0$ , the conservation equations (34) take the form

$$\langle \bar{m}, M (St \partial_t + v_i \partial_{x_i}) \bar{f} \rangle + \langle \bar{m}, M v_i \partial_{x_i} \check{f} \rangle = 0 \quad (38)$$

where  $\bar{f} := \Pi_M[f]$  and  $\check{f} := (Id - \Pi_M)[f]$  and the fine-scale equation (35) simplifies to

$$\check{f} = \epsilon \mathcal{L}_M^{-1} (Id - \Pi_M) \left[ v_i \partial_{x_i} \bar{f} + (St \partial_t + v_i \partial_{x_i}) \check{f} \right] \quad (39)$$

Recalling the expressions (15a) and (15b) for the linearized deviatoric stress and heat flux and observing that  $(Id - \Pi_M)[v_i v_j] = v_i v_j - \frac{1}{D} |\mathbf{v}|^2 \delta_{ij}$  and  $(Id - \Pi_M)[|\mathbf{v}|^2 v_i] = |\mathbf{v}|^2 v_i - (D + 2)\theta_0 v_i$ , we see that the expressions for the deviatoric stress and heat flux are completely determined by the  $\langle v_i \bar{m}, M \partial_{x_i} \check{f} \rangle$  term in (38). This means that the process of closing the system of conservation equations derived from the Boltzmann equation amounts to a substitution of some approximation of (39) into (38). Our goal in this text is to ensure that the fine scale approximation is *entropy stable*.

With  $\bar{m} = \bar{f}$  in (38), we have that

$$\frac{St}{2} \partial_t \langle \bar{f}, M \bar{f} \rangle + \frac{1}{2} \partial_{x_i} \langle v_i \bar{f}, M \bar{f} \rangle + \partial_{x_i} \langle v_i \bar{f}, M \check{f} \rangle - \langle v_i \partial_{x_i} \bar{f}, M \check{f} \rangle = 0 \quad (40)$$

An entropy stable (or entropy dissipative) closure refers to a fine-scale approximation  $\check{f}$  such that the spatial integral of  $\langle v_i \partial_{x_i} \bar{f}, M \check{f} \rangle$  is non-positive. This is because under the assumption that any terms in divergence form can be ignored (for example in an infinite spatial domain), equation (40) gives

$$St \frac{d}{dt} \int_{\Omega} \frac{1}{2} \langle \bar{f}, M \bar{f} \rangle d\mathbf{x} = \int_{\Omega} \langle v_i \partial_{x_i} \bar{f}, M \check{f} \rangle d\mathbf{x} \leq 0 \quad (41)$$

The non-negative term  $\frac{1}{2} \langle \bar{f}, M \bar{f} \rangle$  defines the macroscopic entropy for the closed conservation equations derived from the approximation  $\check{f}$  and (41) says that this entropy is non-increasing in time.<sup>4</sup>

<sup>4</sup>To obtain the *non-decreasing* entropy more commonly used in physics, we would use  $-\frac{1}{2} \langle \bar{f}, M \bar{f} \rangle$  instead of  $\frac{1}{2} \langle \bar{f}, M \bar{f} \rangle$  as our macroscopic entropy.

An important motivation for (41) comes from testing the linearized Boltzmann equation (37) with  $f$  in the  $\mathcal{L}^2(M dv)$  inner-product and integrating in space (ignoring the divergence term) to get

$$\text{St} \frac{d}{dt} \int_{\Omega} \frac{1}{2} \langle f, Mf \rangle d\mathbf{x} = \frac{1}{\epsilon} \int_{\Omega} \langle f, M \mathcal{L}_M[f] \rangle d\mathbf{x} \leq 0 \quad (42)$$

The idea is to view the conservation equations as a weak form (in  $\mathbf{v}$ ) of the (linearized) Boltzmann equation. The process of closing these conservation equations then amounts to solving for a finite-dimensional Galerkin approximation  $\bar{f}$  of the unknown  $f$  with the entropy stability criterion in (41) serving to ensure that  $\bar{f}$  satisfies a "discrete" version of the entropy inequality (42) that  $f$  satisfies.

**Remark 3.2.** Using (7) and (28), we can show that

$$\bar{f} = \frac{\tilde{\rho}}{\rho_0} + \frac{\mathbf{v} \cdot \tilde{\mathbf{u}}}{\theta_0} + \left( \frac{|\mathbf{v}|^2}{2\theta_0} - \frac{D}{2} \right) \frac{\tilde{\theta}}{\theta_0}$$

As such, the macroscopic entropy is given by

$$\frac{1}{2} \langle \bar{f}, M \bar{f} \rangle = \frac{1}{2\rho_0} \tilde{\rho}^2 + \frac{\rho_0}{2\theta_0} |\tilde{\mathbf{u}}|^2 + \frac{D\rho_0}{4\theta_0^2} \tilde{\theta}^2$$

### 3.2 Self-consistent Maxwellian formulation

With  $\mathcal{M} = \mu$ , we have that  $f$  belongs in the orthogonal complement  $\mathcal{S}^{\perp\mu}$  because by the definition of  $\mu$ ,

$$\langle \bar{m}, \mu \rangle = \langle \bar{m}, F \rangle = \langle \bar{m}, \mu + \epsilon \mu f \rangle$$

and thus  $\langle \bar{m}, \mu f \rangle = 0$ . Furthermore, we can show that  $\langle \bar{m}, \partial_t[\mu f] \rangle = 0$ . Thus the conservation equation (34) takes the form

$$\langle \bar{m}, \text{St} \partial_t \mu \rangle + \langle \bar{m}, v_i \partial_{x_i} \mu \rangle + \epsilon \langle \bar{m}, v_i \partial_{x_i} [\mu f] \rangle = 0 \quad (43)$$

and the fine-scale equation takes the form

$$f = \mathcal{L}_{\mu}^{-1} (\text{Id} - \Pi_{\mu}) \left[ v_i \partial_{x_i} \ln(\mu) + \frac{\epsilon}{\mu} (\text{St} \partial_t + v_i \partial_{x_i}) [\mu f] \right] \quad (44)$$

As with the linear case, the closure for the deviatoric stress and heat flux is wholly determined by the approximation  $f$  through the fine-scale equation. If we use  $\bar{m} = \ln(\mu)$  in (43), the resulting equation can be written as

$$\text{St} \partial_t \langle (\mu \ln(\mu) - \mu), 1 \rangle + \partial_{x_i} \langle (\mu \ln(\mu) - \mu), v_i \rangle + \epsilon \partial_{x_i} \langle v_i \ln(\mu), \mu f \rangle - \epsilon \langle v_i \partial_{x_i} \ln(\mu), \mu f \rangle = 0 \quad (45)$$

Entropy stability in this context refers to when the approximation of  $f$  is such that the term  $\epsilon \langle v_i \partial_{x_i} \ln(\mu), \mu f \rangle$  is non-positive. This is because under circumstances in which we can do away with terms in divergence form when we integrate in space, we get

$$\text{St} \frac{d}{dt} \int_{\Omega} \langle (\mu \ln(\mu) - \mu), 1 \rangle d\mathbf{x} = \epsilon \int_{\Omega} \langle v_i \partial_{x_i} \ln(\mu), \mu f \rangle d\mathbf{x} \leq 0 \quad (46)$$

where  $\langle (\mu \ln(\mu) - \mu), 1 \rangle$  is now the macroscopic entropy for the resulting conservation equations. This inequality is the "discrete" analogue to the local entropy dissipation law (17) which when integrated in space under similar conditions gives

$$\text{St} \frac{d}{dt} \int_{\Omega} \langle (F \ln(F) - F), 1 \rangle d\mathbf{x} = \frac{1}{\epsilon} \int_{\Omega} \langle \ln(F), \mathcal{C}(F) \rangle d\mathbf{x} \leq 0$$

We can also draw a direct comparison with the linear case by making the substitution  $\bar{f} = \ln(\mu)$  and  $\check{f} = \epsilon f$  into (43) and (44) to get

$$\begin{aligned} \langle \bar{m}, \mu (\partial_t + v_i \partial_{x_i}) [\bar{f}] \rangle + \langle \bar{m}, \mu v_i \partial_{x_i} \check{f} \rangle + \langle \bar{m}, \mu \check{f} v_i \partial_{x_i} \bar{f} \rangle &= 0 \\ \check{f} &= \epsilon \mathcal{L}_{\mu}^{-1} (\text{Id} - \Pi_{\mu}) \left[ v_i \partial_{x_i} \bar{f} + (\text{St} \partial_t + v_i \partial_{x_i}) \check{f} + \check{f} v_i \partial_{x_i} \bar{f} \right] \end{aligned}$$

We see that the primary differences from the linear case are in the fact that the Maxwellian depends on the coarse-scale term (i.e.  $\mu = e^{\bar{f}}$ ) as well as the additional non-linear term  $\check{f} v_i \partial_{x_i} \bar{f}$  in both the coarse and fine scale equations. We would then think of the coarse-scale term  $\ln(\mu)$  as a finite-dimensional Galerkin approximation to  $\ln(F)$ . It is, however, much more convenient to work with  $\mu$  and  $f$  for the self-consistent formulation.

**Remark 3.3.** *In terms of the macroscopic variables, the "discrete" entropy is*

$$\langle (\mu \ln(\mu) - \mu), 1 \rangle = \rho \left( \ln \left( \frac{\rho}{(2\pi\theta)^{\frac{D}{2}}} \right) - \frac{D+2}{2} \right)$$

In the sections that follow, we shall use the framework developed here to describe the Chapman-Enskog expansion, showing how entropy stability exists for the Euler and Navier-Stokes-Fourier equations but not for the Burnett equations and beyond. We will then describe a way to induce entropy stability beyond the Navier-Stokes-Fourier equations.

### 3.3 Chapman-Enskog Closures

The first step to deriving the classical Chapman-Enskog closure from equations (44) and (39) is to assume that the orthogonal complement term can be written as a formal power series:

$$\check{f}(t, \mathbf{x}, \mathbf{v}) = \sum_{n=0}^{\infty} \epsilon^n \check{f}_n(t, \mathbf{x}, \mathbf{v}) \quad (47)$$

If we substitute the power series into (44) and (39) and arrange the resulting sequence of equations in orders of  $\epsilon$ , we get respectively

$$\check{f}_0 = 0; \quad \check{f}_1 = \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}]; \quad \check{f}_{n+1} = \mathcal{L}_M^{-1} (\text{St} \partial_t + v_i \partial_{x_i}) [\check{f}_n] \quad (48a)$$

$$f_0 = \mathcal{L}_{\mu}^{-1} [v_i \partial_{x_i} \ln(\mu)]; \quad f_{n+1} = \mathcal{L}_{\mu}^{-1} \left[ \frac{1}{\mu} (\text{St} \partial_t + v_i \partial_{x_i}) [\mu f_n] \right]; \quad (48b)$$

Truncating the power series at different orders of  $\epsilon$  leads to different closures for the conservation equations. For example the zeroth order truncation gives the Navier-Stokes-Fourier equations in the self-consistent Maxwellian formulation and the linearized Euler equations in the background Maxwellian formulation. For our purposes, it is more advantageous to obtain the Chapman-Enskog closure through a fixed point iteration on (44) and (39)

that generates the partial sums of the power series (47):

$$\check{f}_{(n+1)} = \epsilon \mathcal{L}_M^{-1} \left[ (\text{St } \partial_t + v_i \partial_{x_i}) [\check{f}_{(n)}] + v_i \partial_{x_i} \bar{f} \right]; \quad \check{f}_{(0)} = 0 \quad (49a)$$

$$f_{(n+1)} = \mathcal{L}_\mu^{-1} \left[ \frac{\epsilon}{\mu} (\text{St } \partial_t + v_i \partial_{x_i}) [\mu f_{(n)}] + v_i \partial_{x_i} \ln(\mu) \right]; \quad f_{(0)} = 0 \quad (49b)$$

with

$$\check{f}_{(n+1)} = \sum_{j=1}^{n+1} \epsilon^j \check{f}_j; \quad f_{(n+1)} = \sum_{j=0}^n \epsilon^j f_j$$

### 3.3.1 Euler Equations

The Euler equations arise from using  $f_{(0)} = \check{f}_{(0)} = 0$  as our closure for the fine-scale term. Thus the conservation equations (38) and (43) become

$$\langle \bar{m}, \text{St } \partial_t \mu \rangle + \langle \bar{m}, v_i \partial_{x_i} \mu \rangle = 0 \quad (50)$$

$$\langle \bar{m}, M \text{St } \partial_t \bar{f} \rangle + \langle \bar{m}, M v_i \partial_{x_i} \bar{f} \rangle = 0 \quad (51)$$

with  $\bar{m} = 1$ ,  $v_i$  and  $\frac{|v|^2}{2}$  macroscopic conservation of mass, momentum and energy explicitly take the form of (14) and (12) with  $\sigma_{ij} = \tilde{\sigma}_{ij} = 0$  and  $q_i = \tilde{q}_i = 0$ . The corresponding macroscopic entropy equations are given by

$$\frac{d}{dt} \int_{\Omega} \frac{1}{2} \langle \bar{f}, M \bar{f} \rangle = 0 \quad (52)$$

$$\frac{d}{dt} \int_{\Omega} \langle 1, (\mu \ln(\mu) - \mu) \rangle = 0 \quad (53)$$

which means that these equations are entropy stable.

### 3.3.2 Navier-Stokes-Fourier equations

The Navier-Stokes-Fourier equations arise from using the correction directly above that of the Euler equations:

$$\check{f}_{(1)} = \epsilon \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}] \quad \text{and} \quad f_{(1)} = \mathcal{L}_\mu^{-1} [v_i \partial_{x_i} \ln(\mu)]$$

This results in conservation equations (12) and (14) with deviatoric stress tensor and heat flux given by a Newtonian stress-tensor satisfying the Stokes hypothesis and Fourier's law of heat conduction respectively:

$$\begin{aligned} \tilde{\sigma}_{ij}^{(1)} &= \tilde{\omega} \left( \partial_{x_j} \tilde{u}_i + \partial_{x_i} \tilde{u}_j - \frac{2}{D} \partial_{x_k} \tilde{u}_k \delta_{ij} \right) & \tilde{q}_i^{(1)} &= -\tilde{\kappa} \partial_{x_i} \tilde{\theta} \\ \sigma_{ij}^{(1)} &= \omega \left( \partial_{x_j} u_i + \partial_{x_i} u_j - \frac{2}{D} \partial_{x_k} u_k \delta_{ij} \right) & q_i^{(1)} &= -\kappa \partial_{x_i} \theta \end{aligned}$$

where the viscosity and heat conductivity are given by

$$\tilde{\omega} = -\frac{\epsilon}{\theta_0} \langle A_{12}(\mathbf{v}), M \mathcal{L}_M^{-1} [A_{12}(\mathbf{v})] \rangle \quad (\geq 0)$$

$$\omega(t, \mathbf{x}) = -\frac{\epsilon}{\theta} \langle A_{12}(\mathbf{v} - \mathbf{u}), \mu \mathcal{L}_\mu^{-1} [A_{12}(\mathbf{v} - \mathbf{u})] \rangle \quad (\geq 0) \quad (54)$$

$$\tilde{\kappa} = -\frac{\epsilon}{\theta_0^2} \langle B_1(\mathbf{v}), M \mathcal{L}_M^{-1} [B_1(\mathbf{v})] \rangle \quad (\geq 0)$$

$$\kappa(t, \mathbf{x}) = -\frac{\epsilon}{\theta^2} \langle B_1(\mathbf{v} - \mathbf{u}), \mu \mathcal{L}_\mu^{-1} [B_1(\mathbf{v} - \mathbf{u})] \rangle \quad (\geq 0) \quad (55)$$

The details for this derivation and the definitions of the tensors  $A_{ij}$  and  $B_i$  are given in Appendices D and E.

The entropy relations read

$$\begin{aligned} \text{St} \frac{d}{dt} \int_{\Omega} \frac{1}{2} \langle \bar{f}, M \bar{f} \rangle d\mathbf{x} &= \epsilon \int_{\Omega} \langle v_i \partial_{x_i} \bar{f}, M \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}] \rangle d\mathbf{x} \leq 0 \\ \text{St} \frac{d}{dt} \int_{\Omega} \langle (\ln(\mu) - 1), \mu \rangle d\mathbf{x} &= \epsilon \int_{\Omega} \langle v_i \partial_{x_i} \ln(\mu), \mu \mathcal{L}_\mu^{-1} [v_i \partial_{x_i} \ln(\mu)] \rangle d\mathbf{x} \leq 0 \end{aligned}$$

where we use the negative-definiteness of the linearized collision operator to obtain the inequalities above. We also observe that entropy stability holds at all Knudsen number  $\epsilon$  regardless of whether or not the Navier-Stokes-Fourier equations remain a valid model of gas flow. We will see shortly that this is not the case for the Burnett equations.

### 3.3.3 Burnett equations (Part 1)

We have:

$$f_{(2B1)} = f_{(1)} + \epsilon \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} (\text{St} \partial_t + v_i \partial_{x_i}) [\mu \mathcal{L}_\mu^{-1} [v_i \partial_{x_i} \ln(\mu)]] \right] \quad (56)$$

$$\check{f}_{(2B1)} = \check{f}_{(1)} + \epsilon^2 \mathcal{L}_M^{-1} (\text{St} \partial_t + v_i \partial_{x_i}) \mathcal{L}_M^{-1} [v_i \partial_{x_i} \check{f}] \quad (57)$$

The stress and heat flux for the self-consistent Maxwellian formulation contains many terms so we will only write out the stress and heat flux for the background Maxwellian formulation:

$$\begin{aligned} \tilde{\sigma}_{ij}^{(2B1)} &= \tilde{\sigma}_{ij}^{(1)} - \Xi \text{St} \partial_t \left[ \partial_{x_j} \tilde{u}_i + \partial_{x_i} \tilde{u}_j - \frac{2}{D} \partial_{x_k} \tilde{u}_k \delta_{ij} \right] - \Psi \left( \partial_{x_i} \partial_{x_j} \tilde{\theta} - \frac{1}{D} \partial_{x_k} \partial_{x_k} \tilde{\theta} \delta_{ij} \right) \\ \tilde{q}_i^{(2B1)} &= \tilde{q}_i^{(1)} + \Upsilon \text{St} \partial_t \partial_{x_i} \tilde{\theta} + \frac{\theta_0 \Psi}{2} \left( \partial_{x_k} \partial_{x_k} \tilde{u}_i + \left( 1 - \frac{2}{D} \right) \partial_{x_i} \partial_{x_k} \tilde{u}_k \right) \end{aligned} \quad (58)$$

where

$$\Xi = \frac{\epsilon^2}{\theta_0} \langle \mathcal{L}_M^{-1} [A_{12}(\mathbf{v})], M \mathcal{L}_M^{-1} [A_{12}(\mathbf{v})] \rangle \quad (59)$$

$$\Psi = \frac{2\epsilon^2}{\theta_0^2} \langle \mathcal{L}_M^{-1} [A_{12}(\mathbf{v})], M D_{12}(\mathbf{v}) \rangle \quad (60)$$

$$\Upsilon = \frac{\epsilon^2}{\theta_0^2} \langle \mathcal{L}_M^{-1} [B_1(\mathbf{v})], M \mathcal{L}_M^{-1} [B_1(\mathbf{v})] \rangle \quad (61)$$

The derivation of these new terms and the definition of the tensor  $D_{mn}(\mathbf{v})$  are given in Appendices D and E.

To check for entropy stability, it suffices to analyse  $\langle v_i \partial_{x_i} \bar{f}, M \check{f}_{(2B1)} \rangle$  or, more specifically, the

$\epsilon^2 \langle v_i \partial_{x_i} \bar{f}, M \mathcal{L}_M^{-1} (\text{St } \partial_t + v_i \partial_{x_i}) \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}] \rangle$  term contained within it. We have that

$$\begin{aligned} & \langle v_i \partial_{x_i} \bar{f}, M \mathcal{L}_M^{-1} (\text{St } \partial_t + v_i \partial_{x_i}) \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}] \rangle \\ &= \text{St} \langle \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}], M \partial_t \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}] \rangle \\ &+ \langle \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}], M v_i \partial_{x_i} \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}] \rangle \\ &= \frac{\text{St}}{2} \partial_t \langle M, (\mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}])^2 \rangle + \frac{1}{2} \partial_{x_i} \langle v_i, M (\mathcal{L}_M^{-1} [v_j \partial_{x_j} \bar{f}])^2 \rangle \end{aligned}$$

We used the self-adjointness of the linearized operator in the first step and used the product rule for derivatives in the second step.

The time derivative term in the above calculation does not have a definite sign and as such for large Knudsen numbers (when the  $\epsilon^2$  terms dominate) a loss of entropy stability is possible. Considering that this would only be a problem at large enough Knudsen numbers, it might be argued that it only serves as a hard cap on the regime of validity of the Burnett equations [28]. We argue that this loss of entropy stability is an undesirable property for a macroscopic conservation equation that extends the Navier-Stokes-Fourier equations for two reasons. First of all, the Euler equations and the Navier-Stokes-Fourier equations remain entropy stable for all Knudsen numbers even beyond the regime of validity for these equations. This suggests a pattern we should strive to preserve when extending these equations. Secondly, even if we were to allow for the loss of entropy stability in derived extended hydrodynamic equations, the above calculation shows that the loss of entropy stability is problem dependent because the sign of the potentially troublesome term  $\frac{\text{St}}{2} \partial_t \langle M, (\mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}])^2 \rangle$  depends on the conservation variables we would be using the conservation equations to solve for since they are contained in  $\bar{f}$ . Thus it becomes difficult to make an a priori determination of when entropy stability is lost. This hampers the practicality of these equations for real world problems. The same issue is also present in the self-consistent Maxwellian formulation where the  $O(\epsilon)$  term  $\langle v_i \partial_{x_i} \ln(\mu), \mu \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} (\text{St } \partial_t + v_i \partial_{x_i}) [\mu \mathcal{L}_\mu^{-1} [v_i \partial_{x_i} \ln(\mu)]] \right] \rangle$  can be rewritten as

$$\frac{\text{St}}{2} \left\langle \frac{1}{\mu}, \partial_t (\mu \mathcal{L}_\mu^{-1} [v_i \partial_{x_i} \ln(\mu)])^2 \right\rangle + \frac{1}{2} \left\langle \frac{v_i}{\mu}, \partial_{x_i} (\mu \mathcal{L}_\mu^{-1} [v_j \partial_{x_j} \ln(\mu)])^2 \right\rangle$$

### 3.3.4 Burnett equations (Part 2)

The Burnett equations are usually derived in such a way as to get rid of the time derivatives in the closure for the deviatoric stress and heat flux. This is accomplished by using the Euler equations. For (58), this means that we make the following substitutions

$$\begin{aligned} \text{St } \partial_t \tilde{u}_j &= -\frac{1}{\rho_0} \partial_{x_j} [\rho_0 \tilde{\theta} + \tilde{\rho} \theta_0] \\ \text{St } \partial_t \tilde{\theta} &= -\frac{2}{D} \theta_0 \partial_{x_i} \tilde{u}_i \end{aligned}$$

in order to obtain stress and heat flux given by

$$\begin{aligned} \tilde{\sigma}_{ij}^{(2B^2)} &= \tilde{\sigma}_{ij}^{(1)} - (\Psi - 2\Xi) \left( \partial_{x_i} \partial_{x_j} \tilde{\theta} - \frac{1}{D} \partial_{x_k} \partial_{x_k} \tilde{\theta} \delta_{ij} \right) + \frac{2\Xi \theta_0}{\rho_0} \left( \partial_{x_i} \partial_{x_j} \tilde{\rho} - \frac{1}{D} \partial_{x_k} \partial_{x_k} \tilde{\rho} \delta_{ij} \right) \\ \tilde{q}_i^{(2B^2)} &= \tilde{q}_i^{(1)} + \theta_0 \left( \frac{\Psi}{2} \left( 1 - \frac{2}{D} \right) - \frac{2}{D} \Upsilon \right) \partial_{x_i} \partial_{x_k} \tilde{u}_k + \frac{\theta_0 \Psi}{2} \partial_{x_k} \partial_{x_k} \tilde{u}_i \end{aligned} \quad (62)$$

On the level of the coarse and fine-scale equations, this can be accomplished by observing that equation (38) can be written as

$$\text{St } \partial_t \bar{f} = -\Pi_M [v_i \partial_{x_i} \bar{f}] + O(\epsilon)$$

where  $O(\epsilon)$  represents any fine-scale corrections made. These correction terms will be at least an order of  $\epsilon$  higher than the other terms in  $\check{f}_{(2)}$ . As such to get the fine-scale term for the Burnett equations, we formally ignore these corrections when we make the substitution into (57) to obtain

$$\check{f}_{(2B2)} = \check{f}_{(1)} - \epsilon^2 \mathcal{L}_M^{-2} v_j \partial_{x_j} \Pi_M [v_i \partial_{x_i} \bar{f}] + \epsilon^2 \mathcal{L}_M^{-1} v_j \partial_{x_j} \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}] \quad (63)$$

The derivation of the above closure takes advantage of the fact that  $\mathcal{L}_M^{-1}$  commutes with the time derivative. This is not the case with the self-consistent formulation and as such we do not have a fine scale analogue to the substitution described in this section. That said, the fine-scale approximation (63) also does not provide unconditional entropy stability. We get

$$\begin{aligned} \left\langle v_i \partial_{x_i} \bar{f}, M \check{f}_{(2B2)} \right\rangle &= \underbrace{\left\langle v_i \partial_{x_i} \bar{f}, M \check{f}_{(1)} \right\rangle}_{\text{Stokes-Fourier (good)}} + \frac{\epsilon^2}{2} \underbrace{\partial_{x_i} \left\langle v_i, M (\mathcal{L}_M^{-1} [v_j \partial_{x_j} \bar{f}])^2 \right\rangle}_{\text{Divergence term (ok)}} \\ &\quad - \epsilon^2 \underbrace{\left\langle \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}], M \mathcal{L}_M^{-1} [v_j \partial_{x_j} \Pi_M [v_k \partial_{x_k} \bar{f}]] \right\rangle}_{\text{Indeterminate sign (not good)}} \end{aligned}$$

### 3.3.5 Super-Burnett equation

We note that the next level of closure of the Chapman-Enskog expansion in the background Maxwellian formulation is given by

$$\check{f}_{(3B)} = \check{f}_{(2B)} + \epsilon^3 \mathcal{L}_M^{-1} (\text{St } \partial_t + v_k \partial_{x_k}) \mathcal{L}_M^{-1} (\text{St } \partial_t + v_j \partial_{x_j}) \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}]$$

and the new terms that do not end up on the boundary in the entropy equation (40) will come from the term  $\langle v_l \partial_{x_l} \bar{f}, M \mathcal{L}_M^{-1} (\text{St } \partial_t + v_k \partial_{x_k}) \mathcal{L}_M^{-1} (\text{St } \partial_t + v_j \partial_{x_j}) \mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}] \rangle$ . Using self-adjointness and the product rule of derivatives this term can be rewritten to give

$$\begin{aligned} - \left\langle (\text{St } \partial_t + v_j \partial_{x_j}) \check{f}_{(1)}, M \mathcal{L}_M^{-1} \left[ (\text{St } \partial_t + v_j \partial_{x_j}) \check{f}_{(1)} \right] \right\rangle \\ + \text{St } \partial_t \left\langle \check{f}_{(1)}, M \mathcal{L}_M^{-1} (\text{St } \partial_t + v_j \partial_{x_j}) \check{f}_{(1)} \right\rangle \\ + \partial_{x_k} \left\langle v_k \check{f}_{(1)}, M \mathcal{L}_M^{-1} (\text{St } \partial_t + v_j \partial_{x_j}) \check{f}_{(1)} \right\rangle \end{aligned}$$

Due to the negative-definiteness of the linearized collision operator, we see that the first term above will be non-negative. As such it will have a detrimental effect on the entropy stability of the conservation equations that come from this closure. We are thus unable to deal with the problems of the Burnett equation by moving up to the closure above it.



### 3.4 An Entropy Stable Fine-Scale Closure

With the observations made of the entropy stability of the Burnett equation and Super-Burnett equations, we can now propose a fine-scale closure that does produce entropy stability. This closure is accomplished with two steps:

1. We assume that the Strouhal number  $St$  is at least of the same order as the Knudsen number  $\epsilon$ . This means that the term  $St \partial_t$  in the fine-scale equations (44) and (39) will be at least an order higher in Knudsen number than the other terms and thus when we come up with a new closure term beyond the Navier-Stokes closure, we formally ignore this term. Thus our starting point for our alternative to the Burnett equations will be

$$\begin{aligned} f &= \mathcal{L}_\mu^{-1} \left[ v_i \partial_{x_i} [\ln(\mu)] + \frac{\epsilon}{\mu} v_i \partial_{x_i} [\mu f] \right] + O(\epsilon^2 f) \\ \check{f} &= \epsilon \mathcal{L}_M^{-1} v_i \partial_{x_i} [\bar{f} + \check{f}] + O(\epsilon^2 \check{f}) \end{aligned}$$

2. The above equations can be trivially rewritten as:

$$\begin{aligned} f &= f + \epsilon \mathcal{L}_\mu^{-1} v_j \partial_{x_j} \left[ f - \mathcal{L}_\mu^{-1} \left[ v_i \partial_{x_i} [\ln(\mu)] + \frac{\epsilon}{\mu} v_i \partial_{x_i} [\mu f] \right] \right] \\ \check{f} &= \check{f} + \epsilon \mathcal{L}_M^{-1} v_j \partial_{x_j} \left[ \check{f} - \epsilon \mathcal{L}_M^{-1} v_i \partial_{x_i} [\bar{f} + \check{f}] \right] \end{aligned}$$

Initializing with  $f_{(1)}$  (resp.  $\check{f}_{(1)}$ ), we use the above form as the basis for a new recursive relation on the fine-scale:

$$f_{(k+1)} = f_{(k)} + \epsilon \mathcal{L}_\mu^{-1} v_j \partial_{x_j} \left[ f_{(k)} - \mathcal{L}_\mu^{-1} \left[ v_i \partial_{x_i} [\ln(\mu)] + \frac{\epsilon}{\mu} v_i \partial_{x_i} [\mu f_{(k)}] \right] \right] \quad (64)$$

$$\check{f}_{(k+1)} = \check{f}_{(k)} + \epsilon \mathcal{L}_M^{-1} v_j \partial_{x_j} \left[ \check{f}_{(k)} - \epsilon \mathcal{L}_M^{-1} v_i \partial_{x_i} [\bar{f} + \check{f}_{(k)}] \right] \quad (65)$$

In particular, we get an alternative closure to the Burnett equations of the form

$$f_{(2E)} = f_{(1)} - \epsilon^2 \mathcal{L}_\mu^{-1} v_k \partial_{x_k} \left[ \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} v_j \partial_{x_j} [\mu \mathcal{L}_\mu^{-1} v_i \partial_{x_i} \ln(\mu)] \right] \right] \quad (66)$$

$$\check{f}_{(2E)} = \check{f}_{(1)} - \epsilon^3 \mathcal{L}_M^{-1} v_k \partial_{x_k} \left[ \mathcal{L}_M^{-1} v_j \partial_{x_j} [\mathcal{L}_M^{-1} [v_i \partial_{x_i} \check{f}]] \right] \quad (67)$$

For the constant background formulation, this leads to a deviatoric stress and heat flux given by:

$$\tilde{\sigma}_{ij}^{(2E)} = \tilde{\sigma}_{ij}^{(1)} - \left( \tilde{\Phi}_1 \partial_{x_k} \partial_{x_k} \left[ \partial_{x_i} \tilde{u}_j + \partial_{x_j} \tilde{u}_i - \frac{2}{D} \partial_{x_l} \tilde{u}_l \delta_{ij} \right] + 2\tilde{\Phi}_2 \left( 1 - \frac{2}{D} \right) \left( \partial_{x_i} \partial_{x_j} - \frac{\delta_{ij}}{D} \partial_{x_l} \partial_{x_l} \right) [\partial_{x_k} \tilde{u}_k] \right) \quad (68)$$

$$\tilde{q}_i^{(2E)} = \tilde{q}_i^{(1)} + \tilde{\Lambda} \partial_{x_i} \partial_{x_k} \partial_{x_k} \tilde{\theta} \quad (69)$$

where

$$\tilde{\Phi}_1 = -\frac{\epsilon^3}{\theta_0} K_{121}^{121}; \quad \tilde{\Phi}_2 = -\frac{\epsilon^3}{\theta_0} \frac{K_{112}^{121} + K_{122}^{111}}{2\left(1 - \frac{2}{D}\right)}; \quad \tilde{\Lambda} = -\frac{\epsilon^3}{\theta_0^2} \langle D_{11}(\mathbf{v}), M \mathcal{L}_M^{-1} [D_{11}(\mathbf{v})] \rangle.$$

The process of deriving equations (68) and (69) and the definitions of the terms  $K_{ijk}^{pnm}$  and  $D_{11}(\mathbf{v})$  are elaborated upon in Appendices D and E. To prove entropy stability we need to show that the terms  $\langle v_i \partial_{x_i} \bar{f}, M \check{f}_{(2E)} \rangle$  and  $\langle v_i \partial_{x_i} \ln(\mu), \mu f_{(2E)} \rangle$  yield non-positive terms and divergence terms. We use the self-adjointness of  $\mathcal{L}_M^{-1}$  and the product rule of derivatives to show that

$$\begin{aligned} \langle v_i \partial_{x_i} \bar{f}, M (-\epsilon^3 \mathcal{L}_M^{-1} v_k \partial_{x_k} [\mathcal{L}_M^{-1} v_j \partial_{x_j} [\mathcal{L}_M^{-1} [v_i \partial_{x_i} \bar{f}]]]) \rangle &= -\epsilon \left\langle \check{f}_{(1)}, M v_k \partial_{x_k} [\mathcal{L}_M^{-1} v_j \partial_{x_j} [\check{f}_{(1)}]] \right\rangle \\ &= -\epsilon \partial_{x_k} \left\langle v_k \check{f}_{(1)}, M \mathcal{L}_M^{-1} v_j \partial_{x_j} [\check{f}_{(1)}] \right\rangle \\ &\quad + \underbrace{\epsilon \left\langle v_k \partial_{x_k} \check{f}_{(1)}, M \mathcal{L}_M^{-1} [v_j \partial_{x_j} \check{f}_{(1)}] \right\rangle}_{\leq 0} \end{aligned}$$

Likewise for the non-linear case, starting from the term

$$\left\langle v_i \partial_{x_i} \ln(\mu), \mu \left( -\epsilon^2 \mathcal{L}_\mu^{-1} v_k \partial_{x_k} \left[ \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} v_j \partial_{x_j} [\mu \mathcal{L}_\mu^{-1} v_i \partial_{x_i} \ln(\mu)] \right] \right] \right) \right\rangle$$

we have

$$\begin{aligned} -\epsilon^2 \left\langle f_{(1)}, \mu v_k \partial_{x_k} \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} v_j \partial_{x_j} [\mu f_{(1)}] \right] \right\rangle &= -\epsilon^2 \partial_{x_k} \left\langle v_k f_{(1)}, \mu \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} v_j \partial_{x_j} [\mu f_{(1)}] \right] \right\rangle \\ &\quad + \underbrace{\epsilon^2 \left\langle \frac{1}{\mu} v_k \partial_{x_k} [\mu f_{(1)}], \mu \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} v_j \partial_{x_j} [\mu f_{(1)}] \right] \right\rangle}_{\leq 0} \end{aligned}$$

The iteration (65) also leads to an entropy stable  $\check{f}_{(3E)}$ . Using  $\check{f}_{(1)} = \epsilon \mathcal{L}_M^{-1} v_k \partial_{x_k} \bar{f}$ , this approximation can be written

$$\begin{aligned} \check{f}_{(3E)} &= \check{f}_{(2E)} + \epsilon \mathcal{L}_M^{-1} v_k \partial_{x_k} \left[ \check{f}_{(2E)} - \epsilon \mathcal{L}_M^{-1} v_j \partial_{x_j} [\bar{f} + \check{f}_{(2E)}] \right] \\ &= \check{f}_{(2E)} + \epsilon \mathcal{L}_M^{-1} v_k \partial_{x_k} \left[ \check{f}_{(2E)} - \check{f}_{(1)} \right] - \epsilon^2 \mathcal{L}_M^{-1} v_k \partial_{x_k} \mathcal{L}_M^{-1} v_j \partial_{x_j} [\check{f}_{(2E)}] \\ &= \check{f}_{(2E)} - \epsilon^3 \mathcal{L}_M^{-1} v_k \partial_{x_k} \mathcal{L}_M^{-1} v_j \partial_{x_j} \mathcal{L}_M^{-1} v_i \partial_{x_i} \check{f}_{(1)} - \epsilon^2 \mathcal{L}_M^{-1} v_k \partial_{x_k} \mathcal{L}_M^{-1} v_j \partial_{x_j} [\check{f}_{(2E)}] \end{aligned}$$

Thus the term of interest for proving entropy stability takes the form

$$\begin{aligned} \langle v_m \partial_{x_m} \bar{f}, M \check{f}_{(3E)} \rangle &= \langle v_m \partial_{x_m} \bar{f}, M \check{f}_{(2E)} \rangle - \epsilon^2 \langle \check{f}_{(1)}, M v_k \partial_{x_k} \mathcal{L}_M^{-1} v_j \partial_{x_j} \mathcal{L}_M^{-1} v_i \partial_{x_i} \check{f}_{(1)} \rangle \\ &\quad - \epsilon \langle \check{f}_{(1)}, M v_k \partial_{x_k} \mathcal{L}_M^{-1} v_j \partial_{x_j} \check{f}_{(2E)} \rangle \end{aligned}$$

and we need only focus on just the last two terms from this expression. First

$$\begin{aligned} -\epsilon^2 \langle \check{f}_{(1)}, M v_k \partial_{x_k} \mathcal{L}_M^{-1} v_j \partial_{x_j} \mathcal{L}_M^{-1} v_i \partial_{x_i} \check{f}_{(1)} \rangle &= \\ -\epsilon^2 \partial_k \langle v_k \check{f}_{(1)}, M v_j \partial_{x_j} \mathcal{L}_M^{-1} v_i \partial_{x_i} \check{f}_{(1)} \rangle &+ \epsilon^2 \langle \mathcal{L}_M^{-1} v_k \partial_{x_k} \check{f}_{(1)}, M v_j \partial_{x_j} \mathcal{L}_M^{-1} v_i \partial_{x_i} \check{f}_{(1)} \rangle = \\ -\epsilon^2 \partial_k \langle v_k \check{f}_{(1)}, M v_j \partial_{x_j} \mathcal{L}_M^{-1} v_i \partial_{x_i} \check{f}_{(1)} \rangle &+ \frac{\epsilon^2}{2} \partial_{x_j} \left\langle v_j, M \left( \mathcal{L}_M^{-1} v_k \partial_{x_k} \check{f}_{(1)} \right)^2 \right\rangle \end{aligned}$$

With the other term, applying the product rule to  $\partial_{x_k}$  gives

$$-\epsilon \left\langle \check{f}_{(1)}, M v_k \partial_{x_k} \mathcal{L}_M^{-1} v_j \partial_{x_j} \check{f}_{(2^E)} \right\rangle = -\partial_{x_k} \langle \dots \rangle + \epsilon \left\langle \mathcal{L}_M^{-1} v_k \partial_{x_k} \check{f}_{(1)}, M v_j \partial_{x_j} \check{f}_{(2^E)} \right\rangle$$

Noting that  $\check{f}_{(2^E)} = \check{f}_{(1)} - \epsilon^2 \mathcal{L}_M^{-1} v_n \partial_{x_n} \mathcal{L}_M^{-1} v_l \partial_{x_l} \check{f}_{(1)}$ , the second term in the above can be written as

$$\epsilon \underbrace{\left\langle \mathcal{L}_M^{-1} v_k \partial_{x_k} \check{f}_{(1)}, M v_j \partial_{x_j} \check{f}_{(1)} \right\rangle}_{\leq 0} - \epsilon^3 \left\langle \mathcal{L}_M^{-1} v_k \partial_{x_k} \check{f}_{(1)}, M v_j \partial_{x_j} \mathcal{L}_M^{-1} v_n \partial_{x_n} \mathcal{L}_M^{-1} v_l \partial_{x_l} \check{f}_{(1)} \right\rangle$$

Applying the product rule to  $\partial_{x_j}$  in the second term above gives

$$-\epsilon^3 \partial_{x_j} \langle \dots \rangle + \epsilon^3 \underbrace{\left\langle v_j \partial_{x_j} \mathcal{L}_M^{-1} v_k \partial_{x_k} \check{f}_{(1)}, M \mathcal{L}_M^{-1} \left[ v_n \partial_{x_n} \mathcal{L}_M^{-1} v_l \partial_{x_l} \check{f}_{(1)} \right] \right\rangle}_{\leq 0}$$

At this time, we do not have a way to show whether the iteration (65) always leads to an entropy stable closure. Moreover, due to the dependence  $\mu$  has on  $\mathbf{x}$ , the steps followed above cannot be used to prove entropy stability for the fine-scale approximation  $f_{(3^E)}$  derived from (64). Compared to its linear counterpart, we will have terms like

$$\left\langle \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} v_k \partial_{x_k} [\mu f_{(1)}] \right], \mu v_j \partial_{x_j} \left[ \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} v_i \partial_{x_i} [\mu f_{(1)}] \right] \right] \right\rangle = \frac{\epsilon^2}{2} \left\langle v_j, \mu \partial_{x_j} \left( \mathcal{L}_\mu^{-1} \left[ \frac{1}{\mu} v_k \partial_{x_k} [\mu f_{(1)}] \right] \right)^2 \right\rangle$$

that cannot either be moved to the boundary or shown to be non-positive.

For the remainder of this text, we shall focus on the conservation equations that arise from the fine-scale approximation  $\check{f}_{(2^E)}$ .

### 3.5 Summary Of Fine-Scale Approximations and the Resulting Closures

We summarise the linearized closures derived in this section for  $D = 3$  in the table below

| <b>Chapman-Enskog Iteration</b>  |  |
|--|--|
| $\check{f}_{(0)} = 0; \quad \check{f}_{(n+1)} = \epsilon \mathcal{L}_M^{-1} \left[ \mathbf{v} \cdot \nabla \bar{f} + (\text{St} \partial_t + \mathbf{v} \cdot \nabla) [\check{f}_{(n)}] \right]$   |  |
| <i>Stokes-Fourier</i>  | $\tilde{\sigma}^{(1)} = \tilde{\omega} \left( \nabla \tilde{\mathbf{u}} + (\nabla \tilde{\mathbf{u}})^T - \frac{2}{3} \text{div} \tilde{\mathbf{u}} \text{Id} \right) =: 2 \tilde{\omega} \boldsymbol{\varepsilon}(\tilde{\mathbf{u}})$<br>$\tilde{\mathbf{q}}^{(1)} = -\tilde{\kappa} \nabla \tilde{\theta}$  |
| <i>Burnett I</i>   | $\tilde{\sigma}^{(2^{B1})} = \tilde{\sigma}^{(1)} - 2 \tilde{\Xi} \text{St} \partial_t \boldsymbol{\varepsilon}(\tilde{\mathbf{u}}) - \Psi \left( \nabla \nabla - \frac{\text{Id}}{3} \Delta \right) [\tilde{\theta}]$<br>$\tilde{\mathbf{q}}^{(2^{B1})} = \tilde{\mathbf{q}}^{(1)} + \Upsilon \text{St} \partial_t \nabla \tilde{\theta} + \frac{\theta_0 \Psi}{2} \left( \Delta \tilde{\mathbf{u}} - \frac{1}{3} \nabla \text{div} \tilde{\mathbf{u}} \right)$ |
| <i>Burnett II</i>  | $\tilde{\sigma}^{(2^{B2})} = \tilde{\sigma}^{(1)} - \left( \nabla \nabla - \frac{\text{Id}}{3} \Delta \right) \left[ (\Psi - 2\Xi) \tilde{\theta} - \frac{2\theta_0 \Xi}{\rho_0} \tilde{\rho} \right]$<br>$\tilde{\mathbf{q}}^{(2^{B2})} = \tilde{\mathbf{q}}^{(1)} + \frac{\theta_0 \Psi}{2} \Delta \tilde{\mathbf{u}} + \theta_0 \left( \frac{\Psi}{6} - \frac{2\Upsilon}{3} \right) \nabla \text{div} \tilde{\mathbf{u}}$                                     |
| <b>New Iteration</b>   |  |
| $\check{f}_{(1)} = \epsilon \mathcal{L}_M^{-1} \mathbf{v} \cdot \nabla \bar{f}; \quad \check{f}_{(k+1)} = \check{f}_{(k)} + \epsilon \mathcal{L}_M^{-1} \mathbf{v} \cdot \nabla \left[ \check{f}_{(k)} - \epsilon \mathcal{L}_M^{-1} \mathbf{v} \cdot \nabla \left[ \bar{f} + \check{f}_{(k)} \right] \right]$ |  |
| <i>Entropy Stable Extension</i>  | $\tilde{\sigma}^{(2^E)} = \tilde{\sigma}^{(1)} - 2 \left( \tilde{\Phi}_1 \Delta [\boldsymbol{\varepsilon}(\tilde{\mathbf{u}})] + \frac{1}{3} \tilde{\Phi}_2 \left( \nabla \nabla - \frac{\text{Id}}{3} \Delta \right) [\text{div} \tilde{\mathbf{u}}] \right)$<br>$\tilde{\mathbf{q}}^{(2^E)} = \tilde{\mathbf{q}}^{(1)} + \tilde{\Lambda} \nabla \Delta \tilde{\theta}$   |

The conservation equations for the linearized entropy stable extension will be

$$\begin{aligned}
0 &= \partial_t \tilde{\rho} + \rho_0 \operatorname{div} \tilde{\mathbf{u}} \\
0 &= \rho_0 \partial_t \tilde{\mathbf{u}} + \nabla [\rho_0 \tilde{\theta} + \tilde{\rho} \theta_0] - 2 \operatorname{div} \left[ \tilde{\omega} \varepsilon(\tilde{\mathbf{u}}) - \tilde{\Phi}_1 \Delta \varepsilon(\tilde{\mathbf{u}}) - \frac{\tilde{\Phi}_2}{3} \left( \nabla \nabla - \frac{\operatorname{Id}}{3} \Delta \right) [\operatorname{div} \tilde{\mathbf{u}}] \right] \\
0 &= \frac{3}{2} \rho_0 \partial_t \tilde{\theta} + \rho_0 \theta_0 \operatorname{div} \tilde{\mathbf{u}} + \operatorname{div} [-\tilde{\kappa} \nabla \tilde{\theta} + \tilde{\Lambda} \nabla \Delta \tilde{\theta}]
\end{aligned}$$

## 4 Model Problems

We now test the entropy stable extension on the parallel plates problem illustrated in Figure 1. The advantage of




Figure 1: The setup consists of two parallel plates, assumed to be infinite in two dimensions, that are a distance  $l$  apart. A constant force  $G$  parallel to the surface of the plate is applied to the gas within the channel.

this setup is that the macroscopic parameters of the system depend of just the coordinate perpendicular to the walls of the channel  $x_1$ . By imposing the condition  $\partial_{x_2} [\tilde{\rho} \theta_0 + \rho_0 \tilde{\theta}] = -G$  on the entire domain, the equations simplify to:

$$\begin{aligned}
\partial_{x_i} \tilde{u}_i &= 0 & -\tilde{\omega} \partial_{x_1}^2 \tilde{u}_2 + \tilde{\Phi}_1 \partial_{x_1}^4 \tilde{u}_2 &= G \\
\rho_0 \partial_{x_1} \tilde{\theta} + \theta_0 \partial_{x_1} \tilde{\rho} &= 0 & -\tilde{\kappa} \partial_{x_1}^2 \tilde{\theta} + \tilde{\Lambda} \partial_{x_1}^4 \tilde{\theta} &= 0
\end{aligned} \tag{70}$$

For our purposes, we only focus on the two equations on the right. The corresponding equations for the two versions of the Burnett equations discussed, as well as the Navier-Stokes-Fourier equations, are

$$-\tilde{\omega} \partial_{x_1}^2 \tilde{u}_2 = G; \quad -\tilde{\kappa} \partial_{x_1}^2 \tilde{\theta} = 0 \tag{71}$$

All variables and parameters are non-dimensionalized and the dependence on the Knudsen number extracted by introducing the following variables:

$$\begin{aligned}
x &= \frac{x_1}{l} & u &= \frac{\tilde{u}_2}{\sqrt{\theta_0}} & \theta &= \frac{\tilde{\theta}}{\theta_0} \\
B &= \epsilon \frac{Gl^2}{\tilde{\omega} \sqrt{\theta_0}} & k_u &= \epsilon l \sqrt{\frac{\tilde{\omega}}{\tilde{\Phi}_1}} & k_\theta &= \epsilon l \sqrt{\frac{\tilde{\kappa}}{\tilde{\Lambda}}}
\end{aligned}$$

Substituting these variables into (70) gives the following ODEs

$$-\frac{d^2u}{dx^2} + \frac{\epsilon^2}{k_u^2} \frac{d^4u}{dx^4} = \frac{B}{\epsilon} \quad (72)$$

$$-\frac{d^2\theta}{dx^2} + \frac{\epsilon^2}{k_\theta^2} \frac{d^4\theta}{dx^4} = 0 \quad (73)$$

We will use this setup to study two problems. The stationary heat problem will focus exclusively on (73) and the Poiseuille channel problem will focus exclusively on (72).

The constants in (70) take the form:

$$\begin{aligned} \tilde{\omega} &= \gamma_1 \epsilon \tau \rho_0 \theta_0 & \tilde{\kappa} &= \frac{5}{2} \gamma_2 \epsilon \tau \rho_0 \theta_0 \\ \tilde{\Phi}_1 &= \Gamma_1 \epsilon^3 \tau^3 \rho_0 \theta_0^2 & \tilde{\Lambda} &= \Gamma_2 \epsilon^3 \tau^3 \rho_0 \theta_0^2 \end{aligned}$$

where  $\gamma_1$ ,  $\gamma_2$ ,  $\Gamma_1$  and  $\Gamma_2$  are parameters that depend on the choice of collision operator and  $\tau = l \sqrt{\frac{\pi}{8\theta_0}}$  for what follows.<sup>5</sup> These can be calculated using the method described in Appendix F. We list their values for the Hard Spheres and BGK operators in Table 1.

From this point on unless otherwise indicated, we shall work exclusively with the Hard Spheres parameters when concrete solutions need to be computed.

| Calculated Fluid Parameters for BGK and Hard spheres |            |            |            |                |
|--|------------|------------|------------|----------------|
|  | $\gamma_1$ | $\gamma_2$ | $\Gamma_1$ | $\Gamma_2$     |
| <b>BGK</b>   | 1          | 1          | 2          | $\frac{28}{3}$ |
| <b>Hard Spheres</b>                                  | 1.270042   | 1.922284   | 2.838848   | 37.47015       |

Table 1

## 4.1 Boundary conditions

In general, the derivation of boundary conditions for a given macroscopic closure requires its own specialized analysis. For example, Aoki et al. [29] and Coron [30] employ an asymptotic boundary layer analysis on the Boltzmann equation to derive slip boundary conditions for the Navier-Stokes-Fourier equations while Gu and Emerson [14] use a Hermite polynomial expansion of the distribution function supplemented by correction factors obtained from DSMC modeling of planar Couette flow to derive a complete set of boundary conditions for the regularized 13 moment equations [9].

In order to obtain boundary conditions for the entropy stable extension, we first determine complementary pairs of essential and natural boundary conditions for the two main equations of (70) by deriving a suitable

<sup>5</sup>This value of  $\tau$  can be derived from equation (18b) and (21) of [29]. The mean collision frequency of a hard sphere gas at the equilibrium state described by the global Maxwellian  $M$  is  $\frac{1}{\epsilon\tau}$ .

weak form for them. Using test functions  $w, \vartheta \in H^2([0, l])$  and integrating by parts<sup>6</sup>, we get

$$\begin{aligned} 0 &= \tilde{\omega} \int_0^l \partial_{x_1} w \partial_{x_1} u_2 dx_1 + \tilde{\Phi}_1 \int_0^l \partial_{x_1}^2 w \partial_{x_1}^2 u_2 dx_1 + \left( -w n_1 \tilde{\sigma}_{12}^{(2E)} - \partial_{x_1} w n_1 \tilde{\Phi}_1 \partial_{x_1}^2 \tilde{u}_2 \right) \Big|_{x=0}^{x=l} \\ 0 &= \tilde{\kappa} \int_0^l \partial_{x_1} \vartheta \partial_{x_1} \tilde{\theta} dx_1 + \tilde{\Lambda} \int_0^l \partial_{x_1}^2 \vartheta \partial_{x_1}^2 \tilde{\theta} dx_1 + \left( \vartheta n_1 \tilde{q}_1^{(2E)} - \partial_{x_1} \vartheta n_1 \tilde{\Lambda} \partial_{x_1}^2 \tilde{\theta} \right) \Big|_{x=0}^{x=l} \end{aligned}$$

where  $n_1$  is the outward pointing normal at the boundary (so  $n_1(0) = -1$  and  $n_1(1) = 1$ ). We observe two pairs of complementary essential and natural boundary conditions inherent to the equations: the first pair indicates that we either specify the temperature  $\tilde{\theta}$  (resp.  $\tilde{u}_2$ ) at the boundary or specify the heat flux  $n_1 \tilde{q}_1^{(2E)}$  (resp.  $n_1 \tilde{\sigma}_{12}^{(2E)}$ ) at the boundary and the second pair indicates that we either specify  $n_1 \partial_{x_1} \tilde{\theta}$  (resp.  $n_1 \partial_{x_1} \tilde{u}_2$ ) or specify the second derivative term  $\tilde{\Lambda} \partial_{x_1}^2 \tilde{\theta}$  (resp.  $\tilde{\Phi}_1 \partial_{x_1}^2 \tilde{u}_2$ ).

This analysis only tells us where to impose boundary conditions but not what the imposed boundary conditions should be. As a starting point for this latter goal, we replace the linearized Boltzmann equation's distribution function with our closure in the kinetic boundary condition. For the diffuse reflection boundary condition, which will be our focus in this paper, this amounts to computing:

$$\int_{\{v_1 n_1 \geq 0\}} (\bar{f} + \check{f}_{(2E)}) M v_1 n_1 dv + \int_{\{v_1 n_1 \leq 0\}} \left( \frac{\rho_b}{\rho_0} + \frac{\tilde{\mathbf{u}}_b \cdot \mathbf{v}}{\theta_0} + \frac{\tilde{\varphi}_b}{2\theta_0^2} (|\mathbf{v}|^2 - D\theta_0) \right) M v_1 n_1 dv = 0 \quad (74)$$

$$\int_{\{v_1 n_1 \leq 0\}} \bar{m} (\bar{f} + \check{f}_{(2E)}) M v_1 n_1 dv - \int_{\{v_1 n_1 \leq 0\}} \bar{m} \left( \frac{\rho_b}{\rho_0} + \frac{\tilde{\mathbf{u}}_b \cdot \mathbf{v}}{\theta_0} + \frac{\tilde{\varphi}_b}{2\theta_0^2} (|\mathbf{v}|^2 - D\theta_0) \right) M v_1 n_1 dv = 0 \quad (75)$$

where  $\bar{m} \in \{1, v_2, |\mathbf{v}|^2\}$ ,  $\tilde{\varphi}_b$  and  $\tilde{\mathbf{u}}_b$  are the temperature and velocity of the boundary and  $\rho_b$  is the mass density at the boundary required to ensure that the zero mass flux condition (74) is satisfied. At  $x = 0$ , the computation of (74) and (75) yields<sup>7</sup>

$$\tilde{u}_1 = 0 \quad (76)$$

$$\frac{1}{\rho_0} (\rho(0) - \rho_b^L) + \frac{1}{2} (\theta(0) - \varphi^L) = 0 \quad (77)$$

$$\sqrt{\frac{2}{\pi}} (u(0) - u_b(0)) - \frac{\tilde{\omega}}{\epsilon l \rho_0 \sqrt{\theta_0}} \left( \epsilon \frac{du}{dx}(0) \right) + \frac{\tilde{\Phi}_1}{\epsilon^3 l^3 \rho_0 \sqrt{\theta_0}} \left( \epsilon^3 \frac{d^3 u}{dx^3}(0) \right) = 0 \quad (78)$$

$$\begin{aligned} \sqrt{\frac{2}{\pi}} \left( \frac{2(\rho(0) - \rho_b^L)}{\rho_0} + 3(\theta(0) - \varphi^L) \right) - \frac{\tilde{\kappa}}{\epsilon l \rho_0 \sqrt{\theta_0}} \left( \epsilon \frac{d\theta}{dx}(0) \right) \\ + \frac{\tilde{\Lambda}}{\epsilon^3 l^3 \rho_0 \sqrt{\theta_0}} \left( \epsilon^3 \frac{d^3 \theta}{dx^3}(0) \right) = 0 \end{aligned} \quad (79)$$

from which we obtain natural boundary conditions of the form

$$-\sigma_{12}^{(2E)}(x^b) n_1(x^b) := - \left( \epsilon \underline{\omega} \frac{du}{dx}(x^b) - \epsilon^3 \underline{\Phi} \frac{d^3 u}{dx^3}(x^b) \right) n_1(x^b) = \sqrt{\frac{2}{\pi}} \left( u(x^b) - u_b(x^b) \right) \quad (80)$$

$$q_1^{(2E)}(x^b) n_1(x^b) := \left( -\epsilon \underline{\kappa} \frac{d\theta}{dx}(x^b) + \epsilon^3 \underline{\Lambda} \frac{d^3 \theta}{dx^3}(x^b) \right) n_1(x^b) = 2\sqrt{\frac{2}{\pi}} \left( \theta(x^b) - \varphi_b \right) \quad (81)$$

where  $x^b \in \{0, 1\}$ ,  $\underline{\omega} = \frac{\tilde{\omega}}{\epsilon l \rho_0 \sqrt{\theta_0}}$ ,  $\underline{\kappa} = \frac{\tilde{\kappa}}{\epsilon l \rho_0 \sqrt{\theta_0}}$ ,  $\underline{\Phi} = \frac{\tilde{\Phi}_1}{\epsilon^3 l^3 \rho_0 \sqrt{\theta_0}}$  and  $\underline{\Lambda} = \frac{\tilde{\Lambda}}{\epsilon^3 l^3 \rho_0 \sqrt{\theta_0}}$ .

We would still need boundary conditions for  $\underline{\Phi} \frac{d^2 u}{dx^2}$  and  $\underline{\Lambda} \frac{d^2 \theta}{dx^2}$  in order to completely determine a solution. A prescription for each will be made in the model problems that follow.

<sup>6</sup>Once on the Stokes-Fourier terms and twice for the entropy stable extension terms.

<sup>7</sup>At  $x = 1$ , the signs on the derivative terms switch.

## 4.2 Stationary Heat Problem

The general solution to Equation (73) is

$$\theta(x) = c_1 \exp\left(\frac{k_\theta x}{\epsilon}\right) + c_2 \exp\left(-\frac{k_\theta x}{\epsilon}\right) + c_3 x + c_4 \quad (82)$$

and the heat flux across the channel will be

$$q_1^{(2^E)} := -\epsilon \underline{\kappa} \frac{d\theta}{dx} + \epsilon^3 \underline{\Lambda} \frac{d^3\theta}{dx^3} = -\epsilon \underline{\kappa} \left( \frac{d\theta}{dx} - \frac{\epsilon^2}{k_\theta^2} \frac{d^3\theta}{dx^3} \right) = -\epsilon \underline{\kappa} c_3 \quad (83)$$

In order solve the  $c_i$  uniquely, we use (81) in addition to the following boundary condition:

$$\epsilon^2 \underline{\Lambda} \frac{d^2\theta}{dx^2}(x^b) = \epsilon \underline{\zeta} \frac{d\theta}{dx}(x^b) n_1(x^b) + \underline{\chi} (\theta(x^b) - \varphi_b) \quad (84)$$

where  $\underline{\zeta}$  and  $\underline{\chi}$  are as yet undetermined constants. As a result,

$$\begin{aligned} c_1 &= -c_2 e^{-\frac{k_\theta}{\epsilon}} = -\frac{\epsilon}{2c_0} (4\underline{\zeta} - \underline{\chi} \underline{\kappa} \sqrt{2\pi}) (\varphi^R - \varphi^L) \\ c_3 &= \frac{2}{c_0} \left( k_\theta \underline{\zeta} (1 + e^{\frac{k_\theta}{\epsilon}}) + \underline{\kappa} (1 - e^{\frac{k_\theta}{\epsilon}}) \right) (\varphi^R - \varphi^L) \\ c_4 &= \frac{1}{2c_0} \left( \epsilon (\varphi^R + \varphi^L) \left( k_\theta \underline{\zeta} \underline{\kappa} \sqrt{2\pi} (1 + e^{\frac{k_\theta}{\epsilon}}) + (4\underline{\zeta} + \underline{\kappa} (\underline{\kappa} - \underline{\chi}) \sqrt{2\pi}) (1 - e^{\frac{k_\theta}{\epsilon}}) \right) \right. \\ &\quad \left. + 4\varphi^L \left( k_\theta \underline{\zeta} (1 + e^{\frac{k_\theta}{\epsilon}}) + \underline{\kappa} (1 - e^{\frac{k_\theta}{\epsilon}}) \right) \right) \end{aligned} \quad (85)$$

where  $c_0 = \underline{\zeta} k_\theta (2 + \epsilon \underline{\kappa} \sqrt{2\pi}) (1 + e^{\frac{k_\theta}{\epsilon}}) + (\epsilon (4\underline{\zeta} + \underline{\kappa} (\underline{\kappa} - \underline{\chi}) \sqrt{2\pi}) + 2\underline{\kappa}) (1 - e^{\frac{k_\theta}{\epsilon}})$ . We still need to determine good values for  $\underline{\zeta}$  and  $\underline{\chi}$ . To this end, we use the data points in Table 4.1 of [22] which were obtained by numerically solving the linearized Boltzmann equation with the hard spheres operator. With  $\varphi^R = 1$  and  $\varphi^L = 0$ , this heat flux data,  $q^{\text{Data}}$ , and our non-dimensional heat flux (83) are related by  $q^{\text{Data}} = -q_1^{(2^E)} \sqrt{2}$  while the Knudsen number  $k$  in [22] and our Knudsen number  $\epsilon$  are related by  $k = \frac{\sqrt{\pi}}{2} \epsilon$ . Leaving  $\underline{\zeta}$  and  $\underline{\chi}$  as free parameters, we use the *fitnlm* function in MATLAB to find the values that best fit (83) to the rescaled linearized Boltzmann solution data  $-\frac{1}{\sqrt{2}} q^{\text{Data}}$ . This process yields values of  $\underline{\zeta} = -0.96491$  and  $\underline{\chi} = -0.94298$  with  $R^2 = 1$  and root mean square error of 0.000671 between the data and our function.

In Figure 2, we plot Equation (83) with the obtained fitted values and  $-\frac{1}{\sqrt{2}} q^{\text{Data}}$ . We also plot the Navier-Stokes-Fourier heat flux solution with no temperature jump  $q_1^{(1)} = -\epsilon \underline{\kappa}$ , as well as a Navier-Stokes-Fourier heat flux solution with a temperature jump obtained by solving Equation (71) using the boundary condition (81) without the third order derivative term

$$q_1^{(1)} = -\frac{\epsilon \underline{\kappa}}{1 + \epsilon \underline{\kappa} \sqrt{\frac{\pi}{2}}}$$

We also include the solution due to the Grad-Hilbert expansion given in Equation (4.18c) of [22]. In our non-dimensionalization, this will be

$$q^{\text{GH}} = -\frac{\epsilon \underline{\kappa}}{1 + \epsilon d_1 \sqrt{\frac{\pi}{2}}},$$

where  $d_1 = 2.4001$  for hard spheres. Finally, we note that in the collisionless limit  $\epsilon \rightarrow \infty$ , the linearized Boltzmann equation can be solved exactly with heat flux calculated to be  $q_\infty^{\text{Exact}} = -\sqrt{\frac{2}{\pi}}$ . Due to the boundary




Figure 2: Heat flux as a function of the Knudsen number in the stationary heat problem. All the methods compared agree in the small  $\epsilon$  regime but the Navier-Stokes-Fourier solution without a jump condition (green dotted line) deviates first as  $\epsilon$  increases. The Navier-Stokes-Fourier solution with a jump condition (maroon dotted line) remains relatively close to the linearized Boltzmann solution and even converges to the correct collisionless limit heat flux. The Grad-Hilbert solution (orange dot-dashed line) does better than the Navier-Stokes-Fourier solution with a jump condition for  $\epsilon \leq 1$  but converges to the wrong value in the collisionless limit. Finally, the solution due to our entropy stable extension agrees with the linearized Boltzmann solution remarkably well over the entire range of Knudsen numbers observed.

condition (81), it turns out that the heat flux due to our entropy stable extension will always converge to  $q_\infty^{\text{Exact}}$  regardless of the value of  $\underline{\kappa}$ ,  $\underline{\Delta}$ ,  $\underline{\zeta}$  and  $\underline{\chi}$  used.<sup>8</sup>

In Figure 3, we plot the temperature distribution that results from using  $\underline{\zeta} = -0.96491$  and  $\underline{\chi} = -0.94298$  in Equation (85) at  $\epsilon = 0.125, 0.25, 1.0$  and  $100$ . For comparison, we also plot the temperature distribution due to the Grad-Hilbert extension (Equation (4.18b) of [22]<sup>9</sup>) and the Navier-Stokes-Fourier solution with jump conditions.

In the collisionless limit, the temperature distribution predicted by the linearized Boltzmann equation is  $\frac{\varphi^L + \varphi^R}{2} = \frac{1}{2}$  and we see that all three solutions plotted converge to this limit solution at large Knudsen numbers<sup>10</sup> approach this limit. Thus all solutions are again in agreement at very large Knudsen numbers.

### 4.3 Poiseuille Channel Flow and the Knudsen Minimum

The general solution to (72) is

$$u(x) = -\frac{Bx^2}{2\epsilon} + d_3x + d_4 + d_1 \exp\left(\frac{k_u x}{\epsilon}\right) + d_2 \exp\left(-\frac{k_u x}{\epsilon}\right) \quad (86)$$

As in the case of the stationary heat problem, we need to specify a boundary condition on  $\underline{\Phi} \frac{d^2 u}{dx^2}(x^b)$  in addition to the boundary condition (80), in order to fully determine the solution. However we cannot use a boundary

<sup>8</sup>In fact, the Navier-Stokes-Fourier solution with the jump condition derived from (81) also exhibits this behavior.

<sup>9</sup>With  $\varphi^R = 1$  and  $\varphi^L = 0$ , the non-dimensional temperature in [22]  $\frac{T_H - T_0}{T_1 - T_0}$  equals  $\theta$ .

<sup>10</sup>We again note that our solution converges to this limit regardless of the value of  $\underline{\kappa}$ ,  $\underline{\Delta}$ ,  $\underline{\zeta}$  and  $\underline{\chi}$  used.






Figure 3: Comparison of the non-dimensional temperature distribution obtained via the entropy stable extension (blue line), the Grad-Hilbert expansion (orange dashed) and the Navier-Stokes-Fourier equations with the jump boundary conditions described in the text (red dotted). The three solutions agree strongly in the interior of the channel for smaller Knudsen numbers with a discrepancy at the walls that extends into the domain as the Knudsen number increases ( $\epsilon = 0.125, 0.25$ , and  $1$ ). The solution due to the entropy stable extension hews more closely to the Grad-Hilbert temperature distribution than the Navier-Stokes-Fourier solution does. Because all three solutions converge to the correct collisionless limit temperature distribution, they again agree strongly for very large Knudsen numbers ( $\epsilon = 100$ ).

condition like (84) because Equation (72) shows that in the limit where  $\epsilon$  goes to zero  $\frac{d^2 u}{dx^2} = O(\epsilon^{-1})$  and this would be inconsistent with a boundary condition that takes a form similar to (84). With this in mind, we instead opt for the following boundary conditions

$$\left(-\epsilon \underline{\omega} \frac{du}{dx}(x^b) + \epsilon^3 \underline{\Phi} \frac{d^3 u}{dx^3}(x^b)\right) n_1(x^b) = u(x^b) \sqrt{\frac{2}{\pi}}; \quad \frac{d^2 u}{dx^2}(x^b) = \left(-1 + C(\epsilon)\right) \frac{B}{\epsilon}; \quad x^b \in \{0, 1\}$$

The second boundary condition can be thought of as a perturbation of the Navier-Stokes-Fourier solution's second derivative (i.e.  $-\frac{B}{\epsilon}$ ) by the function  $C(\epsilon)$ . The solution that results is

$$\frac{u(x)}{B} = \frac{(x-x^2)}{2\epsilon} + \frac{\underline{\omega}}{2} \sqrt{\frac{\pi}{2}} + \frac{\epsilon C(\epsilon)}{k_u^2} \left( \frac{\exp\left(\frac{k_u x}{\epsilon}\right) + \exp\left(\frac{k_u(1-x)}{\epsilon}\right)}{\exp\left(\frac{k_u}{\epsilon}\right) + 1} - 1 \right) \quad (87)$$

where the first two terms represent the Stokes-Fourier contribution to the solution whilst the term with  $C(\epsilon)$  is the contribution of our entropy stable extension. Notice that the velocity slip at the boundary is constant for all Knudsen numbers because the  $C(\epsilon)$  term vanishes at the boundaries. This means that the current set of boundary conditions produces a velocity slip at the boundary equal to that given by the Navier-Stokes-Fourier

solution. This shortcoming will be dealt with in the next section, but first we explore Equation (87) a bit further.

The perturbation  $C(\epsilon)$  determines how the extension to the Stokes-Fourier solution behaves within the domain and thus needs to be carefully chosen. To that end, we calculate the mass flux within the channel. It has been experimentally observed that the mass flux of the gas in the channel is observed to initially decrease with increasing Knudsen number until it reaches a particular Knudsen number, *the Knudsen minimum*, where the mass flux then increases monotonically [31]. This phenomenon cannot be replicated by the Navier-Stokes-Fourier equations which predict a monotonically decreasing mass flux. Equation (87) gives the mass flux in the channel as

$$\frac{Q(\epsilon)}{B} := \int_0^1 \frac{u(x)}{B} dx = \frac{1}{12\epsilon} + \frac{\omega}{2} \sqrt{\frac{\pi}{2}} + C(\epsilon) \underbrace{\frac{2\epsilon^2}{k_u^3} \left( \tanh\left(\frac{k_u}{2\epsilon}\right) - \frac{k_u}{2\epsilon} \right)}_{:= D(\epsilon)} \quad (88)$$

and we immediately see that the  $C(\epsilon)D(\epsilon)$  term will have to counteract the monotone decreasing behavior of the first two terms as  $\epsilon$  increases in order to produce a Knudsen minimum. Because  $D(\epsilon)$  is fixed, we can carefully choose  $C(\epsilon)$  to select for a particular Knudsen minimum and growth behavior in the collisionless limit.

To illustrate this point, we note that Cercignani and Daneri show in [32] that  $\frac{Q(\epsilon)}{B} \sim \frac{\gamma_1}{4} \ln(\epsilon)$  in the collisionless limit. We therefore choose a perturbation of the form  $C(\epsilon) = -\epsilon(h_1 \ln(1 + \epsilon) + h_2)$ , where  $h_1$  and  $h_2$  are constants. This will ensure that  $C(0) = 0$  and  $C(\epsilon)D(\epsilon) = O(\ln(\epsilon))$  in the collisionless limit. Roughly speaking,  $h_1$  tunes the logarithmic growth at large  $\epsilon$  whilst  $h_2$  tunes the Knudsen minimum in the transition regime. This ansatz is chosen for its simplicity; it is by no means unique nor provably the best possible choice. To determine good values for  $h_1$  and  $h_2$ , we use the data in Table V of the paper by Ohwada et al. [23], where this problem is solved numerically for the Hard Spheres linearized Boltzmann equation over a wide range of Knudsen numbers.

We first note that the relationship between the non-dimensional mass flux  $M_P$  in [23] and ours is  $\frac{Q(\epsilon)}{B} = -\frac{\gamma_1 \sqrt{\pi}}{2} M_P$  and that the conversion from our Knudsen parameter  $\epsilon$  to their Knudsen parameter  $k$  is given by  $k = \frac{\sqrt{\pi}}{2} \epsilon$ . Using the *fitnlm* function in MATLAB gives best fit parameters of  $h_1 = 2.1246$  and  $h_2 = 2.3066$  with  $R^2 = 0.978$  between Equation (88) and the data. In Figure 4, we compare the fitted solution  $\frac{2}{\gamma_1 \sqrt{\pi}} \frac{Q}{B}$  to the mass flux  $-M_P$  in Table V, the Navier-Stokes-Fourier equation solution (i.e.  $C(\epsilon) = 0$ ) and the regularized 13 moment (R13) equations solution calculated for Maxwell molecules by Struchtrup and Torrillon in [33]. A summary of the conversions used to compare these solutions is given in Table 2.

| Poiseuille Channel Conversions   |               |   |                                   |
|----------------------------------|---------------|---|-----------------------------------|
| <i>Linearized Boltzmann [23]</i> | <i>Boltz-</i> | <i>Entropy Stable Ex-</i>                   | <i>R13 moment equa-</i>           |
|                                  |               | <i>tension</i>                              | <i>tions [33]</i>                 |
| $k$                              |               | $\frac{\sqrt{\pi}}{2} \epsilon$             | $\frac{4\sqrt{2}}{5} \text{Kn}$   |
| $-u_P$                           |               | $\frac{2}{\gamma_1 \sqrt{\pi}} \frac{u}{B}$ | $\frac{4\gamma_1 \sqrt{2}}{5F} v$ |
| $-M_P$                           |               | $\frac{2}{\gamma_1 \sqrt{\pi}} \frac{Q}{B}$ | $\frac{4\gamma_1 \sqrt{2}}{5F} J$ |

Table 2: The conversion scheme for a direct comparison of our entropy stable extension solutions to that of Ohwada et al. [23] and Struchtrup and Torrillon [33]. Here,  $\gamma_1 = 1.270042$

Figure 4 shows that the mass flux due to our entropy stable extension is in reasonable agreement with the linearized Boltzmann solution for the range of Knudsen numbers studied. The R13 solution is just as good as




Figure 4: We use  $C(\epsilon) = -\epsilon (2.1246 \ln(1 + \epsilon) + 2.3066)$  in (88) and plot  $\frac{2}{B\gamma_1\sqrt{\pi}}Q\left(\frac{2}{\sqrt{\pi}}k\right)$  against  $k$  (blue line). Also plotted are the linearized Boltzmann data points due to Ohwada, Sone and Aoki [23] (red dots), the Navier-Stokes-Fourier solution ( $C(\epsilon) = 0$ ) (black dash line) and the R13 moments solution from Struchtrup and Torrilhon [33] (magenta dot-dash). We observe a reasonable agreement between our function and the data points from Ohwada et al., quantified by  $R^2 = 0.978$ . The Knudsen minimum for the extension occurs around attained at  $k = 0.48$ . The Stokes solution for the mass flux (black dashed) does not exhibit a Knudsen minimum and decreases monotonically whilst R13 solution is really good for smaller values of  $k$  but diverges from the Boltzmann solution as  $k$  increases.

our solution for smaller values of  $k$  but diverges wildly from linearized Boltzmann solution shortly after attaining a Knudsen minimum.

Next, we use the fitted parameters in the velocity equation (87) and compare the result to velocity data taken from Table I and II of [23] in Figures 5. Because our solution provides no correction to the velocity slip, the velocity given by (87) is a poor match to the Boltzmann solution. The best results are observed for low intermediate Knudsen numbers (eg.  $k = 0.4$  in Figure (5)) but the flow profile at smaller or bigger Knudsen regimes leaves something to be desired. For smaller Knudsen numbers, the inaccuracy could be remedied by a more judicious choice of  $C(\epsilon)$ . However at larger Knudsen numbers, no choice of  $C(\epsilon)$  could remedy the inaccuracy that results due to a lack of a correction to the velocity slip in Equation (87).

In the next section, we describe a set of boundary conditions that introduce a non-constant velocity slip for our solution, leading to a better solution overall.

#### 4.4 The Effects of an Added Slip Function

In order to introduce a non-constant velocity slip to our solution, we replace  $u_b(x^b) = 0$  with  $u_b(x^b) = B\mathfrak{s}(\epsilon)$  in the boundary condition (80), where  $\mathfrak{s}(\epsilon)$  is a slip function we shall describe later. To keep things as simple as possible, our boundary conditions will now be:

$$\left(-\epsilon\omega \frac{du}{dx}(x^b) + \epsilon^3\Phi \frac{d^3u}{dx^3}(x^b)\right) n_1(x^b) = \sqrt{\frac{2}{\pi}}\left(u(x^b) - B\mathfrak{s}(\epsilon)\right); \quad \epsilon^2 \frac{d^2u}{dx^2}(x^b) = -\alpha B\mathfrak{s}(\epsilon); \quad x^b \in \{0, 1\}$$




Figure 5: We use  $C(\epsilon) = -\epsilon(2.1246 \ln(1 + \epsilon) + 2.3066)$  in (87) to compare  $\frac{2}{\gamma_1 \sqrt{\pi}} u(x; \frac{2}{\sqrt{\pi}} k)$  (blue line) to the data points for the velocity profile provided in [23] (red dots), the reduced R13 moments solution in [34] (magenta dot-dash) and the Navier-Stokes-Fourier solution (black dash). We fix the range of the y-axis to  $[0.25, 1.6]$  in all the plots. Notice that the Navier-Stokes-Fourier solution and the entropy stable extension always agree at  $x = 1$ . The best results occur for low intermediate Knudsen numbers (represented by  $k = 0.4$ ). We omit the R13 solution at  $k = 2$  and  $k = 8$  because these are outside its range of validity.

where  $\alpha$  is a yet to be determined constant. Notice that requiring that  $\frac{d^2 u}{dx^2} = O(\epsilon^{-1})$  as  $\epsilon$  goes to zero implies that  $\mathfrak{s}(\epsilon) = O(\epsilon)$  in this limit. The velocity and mass flux that result from these boundary conditions are

$$\frac{u}{B} = \frac{x - x^2}{2\epsilon} + \frac{\omega}{2} \sqrt{\frac{\pi}{2}} + \mathfrak{s}(\epsilon) + \frac{\epsilon - \alpha \mathfrak{s}(\epsilon)}{k_u^2} \left( \frac{\exp\left(\frac{k_u x}{\epsilon}\right) + \exp\left(\frac{k_u(1-x)}{\epsilon}\right)}{\exp\left(\frac{k_u}{\epsilon}\right) + 1} - 1 \right) \quad (89)$$

$$\frac{Q}{B} = \frac{1}{12\epsilon} + \frac{\omega}{2} \sqrt{\frac{\pi}{2}} + \mathfrak{s}(\epsilon) + \frac{2\epsilon(\epsilon - \alpha \mathfrak{s}(\epsilon))}{k_u^3} \left( \tanh\left(\frac{k_u}{2\epsilon}\right) - \frac{k_u}{2\epsilon} \right) \quad (90)$$

We observe that  $\frac{Q}{B} \sim \mathfrak{s}(\epsilon)$  in the collisionless limit which motivates the ansatz  $\mathfrak{s}(\epsilon) = h_1 \ln(1 + h_2 \epsilon)$ . This ensures that  $\mathfrak{s}(\epsilon) = O(\epsilon)$  as  $\epsilon$  goes to zero and  $\mathfrak{s}(\epsilon) = O(\ln(\epsilon))$  for large  $\epsilon$ . There are now three parameters to determine, namely,  $h_1$ ,  $h_2$  and  $\alpha$ .

As before, we determine good values for  $\alpha$ ,  $h_1$  and  $h_2$  by comparing  $\frac{2}{B\gamma_1\sqrt{\pi}} Q\left(\frac{2}{\sqrt{\pi}} k\right)$  to the  $-M_p$  values in Table V of [23] using the *fitnlm* function in MATLAB. This process gives  $\alpha = 3.543$ ,  $h_1 = 0.18199$  and  $h_2 = 3.0858$  with  $R^2 = 0.998$ . In Figure 6, we compare our fitted solution to the data points of [23], the full R13 solution and the Navier-Stokes-Fourier solution.

We then use these parameters in the velocity equation (89) and compare the result to the velocity data in Table I and II of [23] in Figures 7 and 8. In general, the fitted velocity underestimates the peak velocity whilst




Figure 6: Plot of  $\frac{2}{B\gamma_1\sqrt{\pi}}Q\left(\frac{2}{\sqrt{\pi}}k\right)$  from (90) with  $\alpha = 3.543$ ,  $h_1 = 0.18199$ ,  $h_2 = 3.0858$ . Also plotted are the data points from Table V of [23], the R13 moment solution from [33] and the Navier-Stokes-Fourier solution. We observe an even stronger agreement to the data points than the solution given in Figure 4 quantified by a higher  $R^2$  of 0.998. The Knudsen minimum for the entropy stable extension with an added slip function, located at  $k = 0.76$ , is also in strong agreement with that of the Boltzmann solution from [23].

overestimating the velocity slip. However this solution is superior to Equation (87) in matching the linearized Boltzmann solution.

## 5 Conclusion

In this manuscript, we derived a set of fourth order PDEs that serve as an alternative to the Burnett equations through a reformulation of the process of deriving closures for the deviatoric stress and heat flux from the Boltzmann equation. This reformulation subsumes the Chapman-Enskog expansion whilst opening the door to other possibilities for closure. In particular, our closure is crafted so as to obtain entropy stability at all Knudsen numbers for both non-linear and linearized versions of the resulting conservation equations.

The rest of the paper focuses on the linearized Hard spheres version of the equations which we apply to the stationary heat problem and the Poiseuille channel. By deriving a symmetric weak form for the momentum and energy equations, we are able to deduce the left hand sides of the natural boundary conditions for these two problems. We obtain the full form of the first set of natural boundary conditions<sup>11</sup> by directly substituting our closure distribution into the linearized Boltzmann diffuse reflection boundary condition. For the second set of natural boundary conditions, we assume the right hand side takes a particular form with free parameters whose best values we determine by comparing to the Boltzmann equation.

For the stationary heat problem, we are able to obtain an almost perfect match to the linearized Boltzmann heat flux by fitting just two free parameters in the second set of natural boundary conditions. We then compute

<sup>11</sup>Those obtained when we integrate by parts the first time.




Figure 7: We use  $\alpha = 3.543$ ,  $h_1 = 0.18199$ ,  $h_2 = 3.0858$  in Equation (89) to compare  $\frac{2}{B\sqrt{\pi}}u\left(x; \frac{2\gamma_1}{\sqrt{\pi}}k\right)$  (blue line) with the Navier-Stokes-Fourier solution (black dash), the linearized Boltzmann solution in [23] (red dot) and the reduced R13 solution from [34] for  $k < 1$ . The range of the y-axis is fixed to  $[0.25, 1.6]$  for all plots. The added slip allows the solution of the entropy stable extension to better match the linearized Boltzmann solution (compare solution without added slip in Figure 5). In general, this solution underestimates the peak velocity and overestimates the slip at the boundary.

the temperature distribution within the channel with the obtained free parameters and find that it agrees remarkably well with the Grad-Hilbert temperature in the interior of the domain. The discrepancy between the two solutions at the boundary slowly increases as the Knudsen number is increased.

For the Poiseuille channel, we find that even though the fitted parameters lead to remarkable agreement of our mass flux to that of the linearized Boltzmann equation, the boundary conditions as set up do not provide any correction to the Navier-Stokes-Fourier velocity slip at the boundary. We found that introducing a slip function into the boundary conditions not only provided a correction to the velocity slip but also led to a mass flux that better matched the linearized Boltzmann mass flux.

Considering that these closures were derived as asymptotic corrections to the Navier-Stokes-Fourier equations into the early transition regime, it cannot be overemphasized how surprising it is that they produce accurate solutions far beyond the expected regime of validity. As the simple nature of the parallel plate test domain might help explain this, it is imperative to test these equations on more complicated domains analytically and numerically. Of particular interest for future research is the behavior of these equations in dynamic problems especially in situations where the assumption that  $St = O(\epsilon)$  is no longer true.

Deriving solutions for these equations will also require a far better understanding of the boundary conditions




Figure 8: We use  $\alpha = 3.543$ ,  $h_1 = 0.18199$ ,  $h_2 = 3.0858$  in Equation (89) to compare  $\frac{2}{B\sqrt{\pi}}u\left(x; \frac{2\gamma_1}{\sqrt{\pi}}k\right)$  (blue line) with the Navier-Stokes-Fourier solution (black dash) and the linearized Boltzmann solution (red dot) for  $k$  ten times larger than in Figure 7. Range of the y-axis is fixed to  $[0.25, 1.6]$  for all plots. The mismatch between the slip from the extension and that from the linearized Boltzmann equation is much more pronounced but the non-constant slip leads to a better overall flow profile than in the version without the slip function. The velocity slip mismatch could indicate that effects due to the Knudsen layer are not sufficiently resolved by the extension.

that need to be imposed. On this front, the model problems in this paper suggest the importance of picking extra boundary conditions that interpolate the continuum and collisionless limit behaviors of the system under consideration. What's more, we believe that the parameters in the boundary condition will depend on the surface derivative of the normal vector and, as a result, the local curvature of the boundary. A systematic derivation of the boundary conditions for these equations that correspond to kinetic analogues (eg. diffuse reflection) would be needed whatever the case may be.

We also plan to investigate the non-linear conservation equations we derived along lines similar to above outline for the linearized equations. These equations will present more complications compared to their linearized counterpart, particularly because the underlying Maxwellian now depends on space and time and thus interacts with those derivatives. However they potentially possess a much larger range of applicability compared to the linearized equations.

## References

- [1] A. V. Bobylev, The chapman-enskog and grad methods for solving the boltzmann equation, *Akademiia Nauk SSSR Doklady* 262 (1) (1982) 71–75.
- [2] A. V. Bobylev, Instabilities in the chapman-enskog expansion and hyperbolic burnett equations, *Journal of statistical physics* 124 (2) (2006) 371–399.
- [3] K. Comeaux, D. Chapman, R. MacCormack, An analysis of the burnett equations based on the second law of thermodynamics, in: *33rd Aerospace Sciences Meeting and Exhibit*, American Institute of Aeronautics and Astronautics, 1995, pp. 1 – 19. doi:10.2514/6.1995-415.
- [4] H. Struchtrup, Failures of the burnett and super-burnett equations in steady state processes, *Continuum Mechanics and Thermodynamics* 17 (2005) 43–50. doi:10.1007/s00161-004-0186-0.
- [5] X. Zhong, R. W. MacCormack, D. R. Chapman, Stabilization of the burnett equations and application to hypersonic flows, *AIAA Journal* 31 (1993) 1036–1043. doi:10.2514/3.11726.
- [6] S. Jin, M. Slemrod, Regularization of the burnett equations via relaxation, *Journal of Statistical Physics* 103 (2001) 1009–1033. doi:10.1023/A:1010365123288.
- [7] A. V. Bobylev, Boltzmann equation and hydrodynamics beyond navier-stokes, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 376 (4 2018). doi:10.1098/RSTA.2017.0227.
- [8] H. Grad, Asymptotic theory of the boltzmann equation, *Physics of Fluids* 6 (1963). doi:10.1063/1.1706716.
- [9] H. Struchtrup, M. Torrilhon, Regularization of grad’s 13 moment equations: Derivation and linear analysis, *Physics of Fluids* 15 (2003) 2668–2680. doi:10.1063/1.1597472.
- [10] H. Struchtrup, Derivation of 13 moment equations for rarefied gas flow to second order accuracy for arbitrary interaction potentials, *Multiscale Modeling & Simulation* 3 (2005) 221–243. doi:10.1137/040603115.
- [11] X.-J. Gu, D. R. Emerson, A high-order moment approach for capturing non-equilibrium phenomena in the transition regime, *Journal of Fluid Mechanics* 636 (2009) 177–216. doi:10.1017/S002211200900768X.
- [12] P. Kauf, M. Torrilhon, M. Junk, Scale-induced closure for approximations of kinetic equations, *Journal of Statistical Physics* 141 (2010) 848–888. doi:10.1007/s10955-010-0073-y.
- [13] A. Rana, M. Torrilhon, H. Struchtrup, A robust numerical method for the r13 equations of rarefied gas dynamics: Application to lid driven cavity, *Journal of Computational Physics* 236 (2013) 169 – 186.
- [14] X. Gu, D. Emerson, A computational strategy for the regularized 13 moment equations with enhanced wall-boundary conditions, *Journal of Computational Physics* 225 (7 2007). doi:10.1016/j.jcp.2006.11.032.
- [15] T. J. Hughes, Multiscale phenomena: Green’s functions, the dirichlet-to-neumann formulation, subgrid scale models, bubbles and the origins of stabilized methods, *Computer Methods in Applied Mechanics and Engineering* 127 (1995) 387–401. doi:10.1016/0045-7825(95)00844-9.



- [16] T. J. Hughes, G. R. Feijóo, L. Mazzei, J. B. Quincy, The variational multiscale method—a paradigm for computational mechanics, *Computer Methods in Applied Mechanics and Engineering* 166 (1998) 3–24. doi:10.1016/S0045-7825(98)00079-6.
- [17] T. J. Hughes, G. Sangalli, Variational multiscale analysis: the fine-scale green’s function, projection, optimization, localization, and stabilized methods, *SIAM Journal on Numerical Analysis* 45 (2) (2007) 539–557.
- [18] Y. Bazilevs, V. Calo, J. Cottrell, T. Hughes, A. Reali, G. Scovazzi, Variational multiscale residual-based turbulence modeling for large eddy simulation of incompressible flows, *Computer Methods in Applied Mechanics and Engineering* 197 (2007) 173–201. doi:10.1016/j.cma.2007.07.016.
- [19] C. Cercignani, R. Illner, M. Pulvirenti, *The Mathematical Theory of Dilute Gases*, Springer New York, New York, NY, 1994.
- [20] F. Golse, The boltzmann equation and its hydrodynamic limits, *Evolutionary equations* 2 (2005) 159–301.
- [21] L. Saint-Raymond, *Hydrodynamic Limits of the Boltzmann Equation*, no. v. 1971 in *Lecture Notes in Mathematics*, Springer, 2009.
- [22] Y. Sone, *Molecular Gas Dynamics*, Birkhäuser Boston, 2007.
- [23] T. Ohwada, Y. Sone, K. Aoki, Numerical analysis of the poiseuille and thermal transpiration flows between two parallel plates on the basis of the boltzmann equation for hard-sphere molecules, *Physics of Fluids A: Fluid Dynamics* 1 (1989) 2042–2049. doi:10.1063/1.857478.
- [24] C. Cercignani, *The Boltzmann Equation and Its Applications*, Springer New York, New York, NY, 1988.
- [25] C. Levermore, Moment closure hierarchies for kinetic theories, *Journal of Statistical Physics* 83 (1996) 1021–1065.
- [26] P. Bhatnagar, E. Gross, M. Krook, A model for collision processes in gases. I. Small amplitude processes in charged and neutral one-component systems., *Phys. Rev.* 94 (511—525) (1954).
- [27] L. Saint-Raymond, A mathematical pde perspective on the chapman–enskog expansion, *Bulletin of the American Mathematical Society* 51 (2) (2014) 247–275.
- [28] L. Garciacolin, R. Velasco, F. Uribe, Beyond the navier–stokes equations: Burnett hydrodynamics, *Physics Reports* 465 (2008) 149–189. doi:10.1016/j.physrep.2008.04.010.
- [29] K. Aoki, C. Baranger, M. Hattori, S. Kosuge, G. Martalò, J. Mathiaud, L. Mieussens, Slip boundary conditions for the compressible navier–stokes equations, *Journal of Statistical Physics* 169 (11 2017). doi:10.1007/s10955-017-1886-8.
- [30] F. Coron, Derivation of slip boundary conditions for the navier-stokes system from the boltzmann equation, *Journal of Statistical Physics* 54 (1989) 829–857. doi:10.1007/BF01019777.
- [31] M. Knudsen, Die gesetze der molekularströmung und der inneren reibungsströmung der gase durch röhren, *Annalen der Physik* 333 (1909) 75–130. doi:10.1002/andp.19093330106.

- [32] C. Cercignani, A. Daneri, Flow of a rarefied gas between two parallel plates, *Journal of Applied Physics* 34 (1963) 3509–3513. doi:10.1063/1.1729249.
- [33] H. Struchtrup, M. Torrilhon, H theorem, regularization, and boundary conditions for linearized 13 moment equations, *Physical Review Letters* 99 (2007) 014502. doi:10.1103/PhysRevLett.99.014502.
- [34] H. Struchtrup, M. Torrilhon, Higher-order effects in rarefied channel flows, *Physical Review E* 78 (2008) 046301. doi:10.1103/PhysRevE.78.046301.
- [35] K. Hashiguchi, *Mathematical fundamentals*, Elsevier, 2020, Ch. 1, pp. 1–59. doi:10.1016/B978-0-12-819428-7.00001-8.
- [36] E. D. Gust, L. E. Reichl, Molecular dynamics simulation of collision operator eigenvalues, *Physical Review E* 79 (2009) 031202. doi:10.1103/PhysRevE.79.031202.

## A Even Moments of the Maxwellian

We derive general expressions for integrals of the form

$$\int_{\mathbb{R}^D} c_{i_1} \dots c_{i_m} |\mathbf{c}|^k \mathcal{M}_{\rho, \mathbf{u}, \theta} d\mathbf{c} \quad (91)$$

where  $\mathbf{c} = \mathbf{v} - \mathbf{u}$  and  $m$  and  $k$  are positive integers such that  $m + k$  is even (one can show the integral evaluates to zero when  $m + k$  is odd).

First we note that generalized spherical coordinates can be used to show that

$$\int_{\mathbb{R}^D} |\mathbf{c}|^{2n} \mathcal{M}_{\rho, \mathbf{u}, \theta} d\mathbf{c} = \left( \prod_{k=0}^{n-1} (D + 2k) \right) \rho \theta^n \quad (92)$$

The key to finding a general form for (91) is to think of it as a tensor i.e.

$$\int_{\mathbb{R}^D} c_{i_1} \dots c_{i_m} |\mathbf{c}|^k \mathcal{M}_{\rho, \mathbf{u}, \theta} d\mathbf{c} := \mathbf{T}_{i_1 \dots i_m}^k \quad (93)$$

Because the Maxwellian and the Lebesgue measure  $d\mathbf{c}$  are invariant under rotational transformations, the tensor  $\mathbf{T}_{i_1 \dots i_m}^k$  will be invariant under a change of coordinates in the  $i_j$  subscripts. In other words for fixed  $k$ , the tensor  $\mathbf{T}_{i_1 \dots i_m}^k$  is an isotropic tensor. This is to our advantage because isotropic tensors take very particular forms [35]. For example all isotropic 2-tensors and 4-tensors take the form:

$$\begin{aligned} \mathbf{T}_{i_1 i_2}^k &= T \delta_{i_1 i_2} \\ \mathbf{T}_{i_1 i_2 i_3 i_4}^k &= T_a \delta_{i_1 i_2} \delta_{i_3 i_4} + T_b \delta_{i_1 i_3} \delta_{i_2 i_4} + T_c \delta_{i_1 i_4} \delta_{i_2 i_3} \end{aligned}$$

The scalars  $T, T_a, T_b, T_c$  can then be found by considering all possible complete contractions of the tensor <sup>12</sup>:

$$\begin{aligned} T &= \frac{\mathbf{T}_{ii}^k}{D} \\ T_a &= \frac{(D+1)\mathbf{T}_{iijj}^k - \mathbf{T}_{ijij}^k - \mathbf{T}_{ijji}^k}{D(D-1)(D+2)} \\ T_b &= \frac{(D+1)\mathbf{T}_{ijij}^k - \mathbf{T}_{iijj}^k - \mathbf{T}_{ijji}^k}{D(D-1)(D+2)} \\ T_c &= \frac{(D+1)\mathbf{T}_{ijji}^k - \mathbf{T}_{ijij}^k - \mathbf{T}_{iijj}^k}{D(D-1)(D+2)} \end{aligned}$$

The final observation is that these complete contractions (eg.  $\mathbf{T}_{ijij}^k$ ) will be integrals of the form shown in (92). We present below several integrals we will use in later sections of the appendix that were computed in the manner described.

$$\int_{\mathbb{R}^D} c_i c_j \mathcal{M} d\mathbf{c} = \frac{\delta_{ij}}{D} \int_{\mathbb{R}^D} |\mathbf{c}|^2 \mathcal{M} d\mathbf{w} = \rho\theta\delta_{ij} \quad (94)$$

$$\int_{\mathbb{R}^D} c_i c_j |\mathbf{c}|^2 \mathcal{M} d\mathbf{w} = (D+2)\rho\theta^2\delta_{ij} \quad (95)$$

$$\int_{\mathbb{R}^D} c_i c_j |\mathbf{c}|^4 \mathcal{M} d\mathbf{w} = (D+2)(D+4)\rho\theta^3\delta_{ij} \quad (96)$$

$$\int_{\mathbb{R}^D} c_i c_j c_k c_l \mathcal{M} d\mathbf{c} = \rho\theta^2 (\delta_{ij}\delta_{kl} + \delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}) \quad (97)$$

$$\int_{\mathbb{R}^D} c_i c_j c_k c_l |\mathbf{c}|^2 \mathcal{M} d\mathbf{c} = (D+4)\rho\theta^3 (\delta_{ij}\delta_{kl} + \delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}) \quad (98)$$

$$\int_{\mathbb{R}^D} c_i c_j c_k c_l c_m c_n \mathcal{M} d\mathbf{c} = \rho\theta^3 (\delta_{ij}\delta_{kl}\delta_{mn} + \text{fourteen other terms}) \quad (99)$$

## B Derivation of Projection Operator

In this section, we show that equation (28) (shown below as equation (100)) defines an orthogonal projection onto the space of collision invariants  $\mathcal{I}$  with respect to the  $\mathcal{L}^2(\mathcal{M}d\mathbf{v})$  inner product.

$$\begin{aligned} \Pi_{\mathcal{M}}[g] &= \frac{1}{\rho} \langle 1, \mathcal{M}g \rangle + \frac{(\mathbf{v} - \mathbf{u})}{\rho\theta} \cdot \langle (\mathbf{v} - \mathbf{u}), \mathcal{M}g \rangle \\ &\quad + \left( \frac{|\mathbf{v} - \mathbf{u}|^2}{2\theta} - \frac{D}{2} \right) \frac{2}{D\rho} \left\langle \left( \frac{|\mathbf{v} - \mathbf{u}|^2}{2\theta} - \frac{D}{2} \right), \mathcal{M}g \right\rangle \quad (100) \end{aligned}$$

The derivation amounts to switching from the basis  $\{1, v_1, \dots, v_D, |\mathbf{v}|^2/2\}$  to the  $\mathcal{L}^2(\mathcal{M}d\mathbf{v})$ -orthonormal basis

$$\left\{ \frac{1}{\sqrt{\rho}}, \frac{v_1 - u_1}{\sqrt{\rho\theta}}, \dots, \frac{v_D - u_D}{\sqrt{\rho\theta}}, \sqrt{\frac{2}{D\rho}} \left( \frac{|\mathbf{v} - \mathbf{u}|^2}{2\theta} - \frac{D}{2} \right) \right\}$$

and using this basis for our projection. To that end, we perform the orthogonalization via Gram-Schmidt.

The first basis element  $e_1 = \frac{1}{\sqrt{\rho}}$  follows immediately from the fact that  $\langle 1, \mathcal{M} \rangle = \rho$ . To deduce  $e_2$ , note that

$$q_2^1 = v_1 - \langle e_1, \mathcal{M}v_1 \rangle e_1 = v_1 - \frac{1}{\rho} \langle 1, \mathcal{M}v_1 \rangle = v_1 - u_1$$

<sup>12</sup>Actually  $T_a = T_b = T_c$  because equation (91) is symmetric in its subscript indices. We go through the trouble of writing out the explicit expressions for  $T_a, T_b$  and  $T_c$  in order to accommodate the instances in which an isotropic tensor is not symmetric.

has norm

$$\|q_2^1\|_{\mathcal{M}}^2 := \langle (v_1 - u_1), \mathcal{M}(v_1 - u_1) \rangle = \rho\theta$$

where we used equation (94) to obtain the final equality. Thus our second orthonormal basis term is  $e_2^1 = (v_1 - u_1)/\sqrt{\rho\theta}$ . More generally if we define  $e_2^i = (v_i - u_i)/\sqrt{\rho\theta}$ , we see that the set  $\{e_2^i\}_{i=1}^D$  is orthonormal because, by Equation (94),

$$\langle (v_i - u_i), \mathcal{M}(v_j - u_j) \rangle = \rho\theta\delta_{ij} = 0, \quad \text{for } i \neq j$$

Thus we are left with just  $|\mathbf{v}|^2$ . We first orthogonalize

$$q_3 = \frac{|\mathbf{v}|^2}{2} - \frac{1}{2} \langle e_1, \mathcal{M}|\mathbf{v}|^2 \rangle e_1 - \frac{1}{2} \langle e_2^i, \mathcal{M}|\mathbf{v}|^2 \rangle e_2^i = \frac{|\mathbf{v} - \mathbf{u}|^2 - D\theta}{2} \quad (101)$$

The norm of  $q_3$  gives us three terms to integrate

$$\begin{aligned} \left\langle \left( \frac{|\mathbf{v} - \mathbf{u}|^2 - D\theta}{2} \right)^2, \mathcal{M} \right\rangle &= \frac{1}{4} \langle |\mathbf{v} - \mathbf{u}|^4, \mathcal{M} \rangle - \frac{D\theta}{2} \langle |\mathbf{v} - \mathbf{u}|^2, \mathcal{M} \rangle + \frac{D^2\theta^2}{4} \langle 1, \mathcal{M} \rangle \\ &= \frac{D}{2} \rho\theta^2 \end{aligned} \quad (102)$$

where we have made use of the identity (92) in the above. Thus our final basis element can be written as

$$e_3 = \sqrt{\frac{2}{D\rho}} \left( \frac{|\mathbf{v} - \mathbf{u}|^2}{2\theta} - \frac{D}{2} \right)$$

We then perform an orthonormal projection onto  $\mathcal{S}$  with our orthonormal basis  $\{e_1, e_2^1, \dots, e_2^D, e_3\}$ , i.e.

$$\Pi_{\mathcal{M}}[g] = e_1 \langle e_1, \mathcal{M}g \rangle + e_2^i \langle e_2^i, \mathcal{M}g \rangle + e_3 \langle e_3, \mathcal{M}g \rangle$$

Notice that equation (100) isn't the only orthonormal projection onto the space of collision invariants as we could, for example, obtain other orthonormal bases by switching the order in which we carry out the Gram-Schmidt procedure. However the fact that (100) also appears when computing  $\mu'(\mathcal{M})[\mathcal{M}f]$  (see section C) suggests that this particular projection can be considered canonical.

## C BGK collision operator

In this section, we will detail how the BGK operator [26] gives rise to a linearized collision operator that satisfies the properties required for our formulation. The BGK operator is given by

$$\mathcal{C}^{\text{BGK}}(F) = \frac{\mu(F) - F}{\tau(F)} \quad (103)$$

where  $\tau(F) > 0$  denotes the rate of relaxation to equilibrium that may depend on moments of  $F$ . For example, [22] and [23] use  $\tau = A_c \rho(F)$  where  $A_c$  is a constant.

To show that  $\mathcal{C}^{\text{BGK}}$  complies with Galilean invariance, note that by a change of variables in (22) it can be shown that  $\rho_{\mu}(\mathcal{T}_{\mathbf{U}}F) = \rho_{\mu}(F) = \rho_{\mu}(\mathcal{T}_{\mathbf{O}}F)$  and  $\theta_{\mu}(\mathcal{T}_{\mathbf{U}}F) = \theta_{\mu}(F) = \theta_{\mu}(\mathcal{T}_{\mathbf{O}}F)$  whilst  $\mathbf{u}_{\mu}(\mathcal{T}_{\mathbf{U}}F) = \mathbf{u}_{\mu}(F) + \mathbf{U}$

and  $\mathbf{u}_\mu(\mathcal{T}_O F) = \mathcal{T}_O \mathbf{u}_\mu(F)$ . This means that  $\mu(\mathcal{T}_U F) = \mathcal{T}_U \mu(F)$  and  $\mathcal{T}_O \mu(F) = \mu(\mathcal{T}_O F)$ . Therefore, we establish that

$$\mathcal{C}^{\text{BGK}}(\mathcal{T}_U F) = \frac{\mu(\mathcal{T}_U F) - \mathcal{T}_U F}{\tau} = \mathcal{T}_U \mathcal{C}^{\text{BGK}}(F). \quad (104)$$

and

$$\mathcal{C}^{\text{BGK}}(\mathcal{T}_O F) = \frac{\mu(\mathcal{T}_O F) - \mathcal{T}_O F}{\tau} = \mathcal{T}_O \mathcal{C}^{\text{BGK}}(F), \quad (105)$$

provided that  $\tau$  is invariant under co-ordinate translations and rotations, which is a reasonable imposition on the relaxation rate. The collision invariance property 2 follows immediately from (22) by linearity.

To establish the dissipation relation (16) for  $\mathcal{C}^{\text{BGK}}$  we note that  $\ln(\mu(F)) \in \mathcal{S}$  and therefore by (22),

$$\int_{\mathbb{R}^D} \ln(\mu(F)) (F - \mu(F)) d\mathbf{v} = 0. \quad (106)$$

The dissipation inequality (16) for  $\mathcal{C}^{\text{BGK}}$  then follows because

$$\int_{\mathbb{R}^D} \ln(F) \mathcal{C}^{\text{BGK}}(F) d\mathbf{v} = -\frac{1}{\tau} \int_{\mathbb{R}^D} \ln\left(\frac{F}{\mu(F)}\right) (F - \mu(F)) d\mathbf{v} \leq 0. \quad (107)$$

because  $\ln\left(\frac{a}{b}\right)(a - b) \geq 0$  for  $a, b > 0$ . Moreover, because equality in (107) holds if and only if  $F = \mu(F)$ , the condition

$$\int_{\mathbb{R}^D} \ln(F) \mathcal{C}^{\text{BGK}}(F) d\mathbf{v} = 0$$

implies that  $F$  is a Maxwellian and as a result  $\ln(F) \in \mathcal{S}$ . Thus the equivalence in (18) is therefore also verified.

The linearization of the BGK collision operator about an arbitrary Maxwellian  $\mathcal{M}$  is given by

$$\left. \frac{d}{d\epsilon} \mathcal{C}^{\text{BGK}}(\mathcal{M} + \epsilon \mathcal{M}f) \right|_{\epsilon=0} = \frac{\left. \frac{d}{d\epsilon} \mu(\mathcal{M} + \epsilon \mathcal{M}f) \right|_{\epsilon=0} - \mathcal{M}f}{\tau(\mathcal{M})} \quad (108)$$

We note that the derivative of a Maxwellian  $\mathcal{M}_{\rho, \mathbf{u}, \theta}$  with respect to some arbitrary argument of  $(\rho, \mathbf{u}, \theta)$  is given by

$$\partial \mathcal{M}_{\rho, \mathbf{u}, \theta} = \mathcal{M}_{\rho, \mathbf{u}, \theta} \left( \frac{\partial \rho}{\rho} + \frac{(\mathbf{v} - \mathbf{u}) \cdot \partial \mathbf{u}}{\theta} + \left( \frac{|\mathbf{v} - \mathbf{u}|^2}{2\theta} - \frac{D}{2} \right) \frac{\partial \theta}{\theta} \right) \quad (109)$$

Thus in order to calculate  $\left. \frac{d}{d\epsilon} \mu(\mathcal{M} + \epsilon \mathcal{M}f) \right|_{\epsilon=0}$ , we need expressions for  $\left. \frac{d}{d\epsilon} \rho_\mu(\mathcal{M} + \epsilon \mathcal{M}f) \right|_{\epsilon=0}$ ,  $\left. \frac{d}{d\epsilon} \mathbf{u}_\mu(\mathcal{M} + \epsilon \mathcal{M}f) \right|_{\epsilon=0}$  and  $\left. \frac{d}{d\epsilon} \theta_\mu(\mathcal{M} + \epsilon \mathcal{M}f) \right|_{\epsilon=0}$ . Working from (2a), (2b) and (5), we have that:

$$\left. \frac{d}{d\epsilon} \rho_\mu(\mathcal{M} + \epsilon \mathcal{M}f) \right|_{\epsilon=0} = \langle f, \mathcal{M} \rangle \quad (110)$$

$$\left. \frac{d}{d\epsilon} \mathbf{u}_\mu(\mathcal{M} + \epsilon \mathcal{M}f) \right|_{\epsilon=0} = \frac{1}{\rho_\mu(\mathcal{M})} \langle (\mathbf{v} - \mathbf{u}_\mu(\mathcal{M})), \mathcal{M}f \rangle \quad (111)$$

$$\left. \frac{d}{d\epsilon} \theta_\mu(\mathcal{M} + \epsilon \mathcal{M}f) \right|_{\epsilon=0} = \frac{2}{\rho_\mu(\mathcal{M})} \left\langle \left( \frac{|\mathbf{v} - \mathbf{u}_\mu(\mathcal{M})|^2}{2D} - \frac{\theta_\mu(\mathcal{M})}{2} \right), \mathcal{M}f \right\rangle \quad (112)$$

Substituting the above into (109) gives (100). Thus we have proved that

$$\mathcal{L}_{\mathcal{M}}^{\text{BGK}}[f] = -\frac{1}{\tau(\mathcal{M})} (\text{Id} - \Pi_{\mathcal{M}})[f] \quad (113)$$

Compliance of  $\mathcal{L}_{\mathcal{M}}^{\text{BGK}}[f]$  with properties (2.3.1.1)–(2.3.1.5) follow directly from the fact that  $\Pi_{\mathcal{M}}$  is an orthogonal projection of  $\mathcal{L}^2(\mathcal{M}d\mathbf{v})$  onto  $\mathcal{S}$ .

## D Higher order tensors in the orthogonal complement

We define here a set of tensors that arise when computing the macroscopic fluid equations that arise from the various fine-scale closures in the text. For an arbitrary Maxwellian  $\mathcal{M}$  with density, bulk velocity and temperature given by  $\rho$ ,  $\mathbf{u}$  and  $\theta$  respectively, we have a corresponding canonical projection  $\Pi_{\mathcal{M}}[\cdot]$ , derived in the previous section and a (pseudo-inverse) linearized collision operator  $\mathcal{L}_{\mathcal{M}}^{-1}[\cdot]$ . With  $\mathbf{c} = \mathbf{v} - \mathbf{u}$ , we define the following tensors:

$$A_{ij}(\mathbf{c}) = (\text{Id} - \Pi_{\mathcal{M}})[c_i c_j] = c_i c_j - \frac{|\mathbf{c}|^2}{D} \delta_{ij} \quad (114)$$

$$B_i(\mathbf{c}) = (\text{Id} - \Pi_{\mathcal{M}}) \left[ \frac{|\mathbf{c}|^2 c_i}{2} \right] = \frac{1}{2} (|\mathbf{c}|^2 c_i - (D+2)\theta c_i) \quad (115)$$

$$C_{ijk}(\mathbf{c}) = (\text{Id} - \Pi_{\mathcal{M}}) [c_k \mathcal{L}_{\mathcal{M}}^{-1}[A_{ij}(\mathbf{c})]] \quad (116)$$

$$D_{ij}(\mathbf{c}) = (\text{Id} - \Pi_{\mathcal{M}}) [c_j \mathcal{L}_{\mathcal{M}}^{-1}[B_i(\mathbf{c})]] \quad (117)$$

$$E_{ijkl}(\mathbf{c}) = (\text{Id} - \Pi_{\mathcal{M}}) [c_l \mathcal{L}_{\mathcal{M}}^{-1}[C_{ijk}(\mathbf{c})]] \quad (118)$$

$$F_{ijk}(\mathbf{c}) = (\text{Id} - \Pi_{\mathcal{M}}) [c_k \mathcal{L}_{\mathcal{M}}^{-1}[D_{ij}(\mathbf{c})]] \quad (119)$$

The exact form of the tensors such as  $C_{ijk}$ ,  $D_{ij}$  depend on  $\mathcal{L}_{\mathcal{M}}^{-1}$ . For the BGK operator,  $\mathcal{L}_{\mathcal{M}}^{-1}[\cdot] = -\tau(\text{Id} - \Pi_{\mathcal{M}})[\cdot]$  these tensors can be computed explicitly with the tools developed in the previous sections.

$$C_{ijk}^{\text{BGK}}(\mathbf{c}) = -\tau \left( c_i c_j c_k - \frac{|\mathbf{c}|^2}{D} c_k \delta_{ij} - \theta \left( c_j \delta_{ik} + c_i \delta_{jk} - \frac{2}{D} c_k \delta_{ij} \right) \right) \quad (120)$$

$$D_{ij}^{\text{BGK}}(\mathbf{c}) = -\frac{\tau}{2} \left( |\mathbf{c}|^2 c_i c_j - (D+2)\theta c_i c_j - 2 \left( 1 + \frac{2}{D} \right) \left( \frac{|\mathbf{c}|^2}{2\theta} - \frac{D}{2} \right) \theta^2 \delta_{ij} \right) \quad (121)$$

$$F_{ijk}^{\text{BGK}}(\mathbf{c}) = \frac{\tau^2}{2} \left( |\mathbf{c}|^2 c_i c_j c_k - (D+2)\theta c_i c_j c_k - 2 \left( 1 + \frac{2}{D} \right) \left( \frac{|\mathbf{c}|^2}{2\theta} - \frac{D}{2} \right) \theta^2 c_k \delta_{ij} - 2\theta^2 \left( c_j \delta_{ik} + c_i \delta_{jk} - \frac{2}{D} c_k \delta_{ij} \right) \right) \quad (122)$$

## E Derivation of the higher order fluid mechanical parameters

Our goal is to show how to compute parameters such as  $\omega$ ,  $\kappa$ ,  $\Xi$ ,  $\Psi$  obtained from the various fluid equations we derived. We will only focus on the terms obtained from the constant background Maxwellian formulation. It is important to read through Appendix D before reading this section.

### E.1 Stokes-Fourier terms

The first step to computing  $\tilde{\omega}$  and  $\tilde{\kappa}$  is to substitute the Stokes-Fourier closure into the last term of (38), i.e. consider  $\epsilon \partial_k \langle \bar{m} v_k, M \mathcal{L}_M^{-1}[v_i \partial_{x_i} \bar{f}] \rangle$  where

$$\bar{f} = \frac{\bar{p}}{\rho_0} + \frac{\mathbf{v} \cdot \bar{\mathbf{u}}}{\theta_0} + \frac{\bar{\theta}}{\theta_0} \left( \frac{|\mathbf{v}|^2}{2\theta_0} - \frac{D}{2} \right).$$

Because the range of  $\mathcal{L}_M^{-1}$  is in the orthogonal complement of the collision invariants  $\mathcal{S}^{\perp M}$ , this term will be zero when  $\bar{m} = 1$ . This means there will be no "Stokes-Fourier correction" to the conservation of mass equation.<sup>13</sup> Furthermore, only the last two terms in  $\bar{f}$  will contribute to the correction term because  $\mathcal{L}_M^{-1}[v_i] = 0$ . We are thus left with the conservation of momentum closure ( $\bar{m} = v_l$ ) and conservation of energy closure ( $\bar{m} = \frac{|\mathbf{v}|^2}{2}$ ).

<sup>13</sup>As expected.

With  $\bar{m} = v_l$ , we have

$$\begin{aligned}
\epsilon \langle v_k v_l, M\mathcal{L}_M^{-1}[v_i \partial_i \bar{f}] \rangle &= \epsilon \frac{\partial_{x_i} \tilde{u}_j}{\theta_0} \langle (\text{Id} - \Pi_M)[v_k v_l], M\mathcal{L}_M^{-1}[v_i v_j] \rangle \\
&\quad + \epsilon \frac{\partial_{x_i} \tilde{\theta}}{2\theta_0^2} \langle v_k v_l, M\mathcal{L}_M^{-1}[v_i |\mathbf{v}|^2] \rangle \\
&= \epsilon \frac{\partial_{x_i} \tilde{u}_j}{\theta_0} \langle A_{kl}(\mathbf{v}), M\mathcal{L}_M^{-1}[A_{ij}(\mathbf{v})] \rangle + 0
\end{aligned} \tag{123}$$

In the above calculation,  $v_k v_l$  becomes  $A_{kl}(\mathbf{v})$  in the first term because the range of  $\mathcal{L}_M^{-1}$  is in the orthogonal complement of collision invariants. Because  $\mathcal{L}_M^{-1}$  commutes with rotational transformations  $\mathcal{O}_O$ , it preserves parity (i.e. maps even functions to even functions and odd functions to odd functions) and so the second term will be zero. Furthermore, the term  $\langle A_{kl}(\mathbf{v}), M\mathcal{L}_M^{-1}[A_{ij}(\mathbf{v})] \rangle$  will be a fourth rank isotropic tensor. Combined with the fact that  $A_{ij}(\mathbf{v}) = A_{ji}(\mathbf{v})$  and that  $A_{ii}(\mathbf{v}) = 0$ , we can conclude that

$$\langle A_{kl}(\mathbf{v}), M\mathcal{L}_M^{-1}[A_{ij}(\mathbf{v})] \rangle = \frac{\langle A_{mn}(\mathbf{v}), M\mathcal{L}_M^{-1}[A_{mn}(\mathbf{v})] \rangle}{(D-1)(D+2)} \left( \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk} - \frac{2}{D} \delta_{ij} \delta_{kl} \right) \tag{124}$$

Thus the right-hand side of (123) takes the form

$$\tilde{\sigma}_{kl}^{(1)} = -\frac{\epsilon}{\theta_0} \frac{\langle A_{mn}(\mathbf{v}), M\mathcal{L}_M^{-1}[A_{mn}(\mathbf{v})] \rangle}{(D-1)(D+2)} \left( \partial_{x_k} \tilde{u}_l + \partial_{x_l} \tilde{u}_k - \frac{2}{D} \delta_{kl} \partial_{x_i} \tilde{u}_i \right)$$

However from (124), we observe that<sup>14</sup>

$$\langle A_{12}(\mathbf{v}), M\mathcal{L}_M^{-1}[A_{12}(\mathbf{v})] \rangle = \frac{\langle A_{mn}(\mathbf{v}), M\mathcal{L}_M^{-1}[A_{mn}(\mathbf{v})] \rangle}{(D-1)(D+2)}$$

and thus the formula for viscosity can be written as

$$\tilde{\omega} = -\frac{\epsilon}{\theta_0} \langle A_{12}(\mathbf{v}), M\mathcal{L}_M^{-1}[A_{12}(\mathbf{v})] \rangle \tag{125}$$

With  $\bar{m} = \frac{|\mathbf{v}|^2}{2}$ , a similar calculation to the one in (123) gives

$$\begin{aligned}
\epsilon \left\langle \frac{v_k |\mathbf{v}|^2}{2}, M\mathcal{L}_M^{-1}[v_i \partial_{x_i} \bar{f}] \right\rangle &= \epsilon \frac{\partial_{x_i} \tilde{\theta}}{\theta_0^2} \langle B_k(\mathbf{v}), M\mathcal{L}_M^{-1}[B_i(\mathbf{v})] \rangle \\
&= \frac{\epsilon}{D\theta_0^2} \langle B_n(\mathbf{v}), M\mathcal{L}_M^{-1}[B_n(\mathbf{v})] \rangle \delta_{ik} \partial_{x_i} \tilde{\theta} \\
&= \frac{\epsilon}{D\theta_0^2} \langle B_n(\mathbf{v}), M\mathcal{L}_M^{-1}[B_n(\mathbf{v})] \rangle \partial_{x_k} \tilde{\theta}
\end{aligned}$$

By noting that  $\langle B_1(\mathbf{v}), M\mathcal{L}_M^{-1}[B_1(\mathbf{v})] \rangle = \frac{1}{D} \langle B_n(\mathbf{v}), M\mathcal{L}_M^{-1}[B_n(\mathbf{v})] \rangle$ , the formula for heat conductivity can be written as

$$\tilde{\kappa} = -\frac{\epsilon}{\theta_0^2} \langle B_1(\mathbf{v}), M\mathcal{L}_M^{-1}[B_1(\mathbf{v})] \rangle \tag{126}$$

**Remark E.1.** We define the fourth rank tensor  $\phi_{ijkl} := \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk} - \frac{2}{D} \delta_{ij} \delta_{kl}$ .

Notice that  $\phi_{ijkl} = \phi_{jikl} = \phi_{ijlk} = \phi_{klij}$  and  $\phi_{ijkk} = \phi_{jjkl} = 0$ .

<sup>14</sup>The formula also works with  $A_{13}$  or  $A_{23}$  in place of  $A_{12}$ .

## E.2 Burnett correction terms

The term of interest is

$$\begin{aligned} \epsilon^2 \langle \bar{m}, M v_l \partial_{x_l} \mathcal{L}_M^{-1} [(\text{St } \partial_t + v_k \partial_{x_k}) [\mathcal{L}_M^{-1} [v_j \partial_{x_j} \bar{f}]]] \rangle \\ = \epsilon^2 \partial_{x_l} \underbrace{\langle \mathcal{L}_M^{-1} [\bar{m} v_l], M [(\text{St } \partial_t + v_k \partial_{x_k}) \mathcal{L}_M^{-1} [v_j \partial_{x_j} \bar{f}]] \rangle}_{\text{Burnett } \sigma_{nl} \text{ and } q_l \text{ corrections}} \end{aligned} \quad (127)$$

where we have taken advantage of the self-adjointness of  $\mathcal{L}_M^{-1}$ . As before, equation (127) equals zero when  $\bar{m} = 1$ . Thus the correction only applies to the conservation of momentum and energy equations.

When  $\bar{m} = v_n$ , we are left with

$$\epsilon^2 \left( \frac{\text{St } \partial_t \partial_{x_j} \tilde{u}_i}{\theta_0} \langle \mathcal{L}_M^{-1} [A_{nl}(\mathbf{v})], M \mathcal{L}_M^{-1} [A_{ij}(\mathbf{v})] \rangle + \frac{\partial_{x_k} \partial_{x_j} \tilde{\theta}}{\theta_0^2} \langle \mathcal{L}_M^{-1} [A_{nl}(\mathbf{v})], M D_{jk}(\mathbf{v}) \rangle \right)$$

after we ignore the terms that give odd powered integrands (and thus integrate to zero). Because  $A_{nl}(\mathbf{v}) = A_{ln}(\mathbf{v})$  and  $A_{nn}(\mathbf{v}) = 0$ , we can simplify the fourth rank isotropic tensors above to

$$\begin{aligned} \left( \Xi \phi_{nl ij} \text{St } \partial_t \partial_j \tilde{u}_i + \frac{\Psi}{2} \phi_{nl jk} \partial_k \partial_j \tilde{\theta} \right) \\ = \Xi \text{St } \partial_t \left[ \partial_{x_i} \tilde{u}_n + \partial_{x_n} \tilde{u}_l - \frac{2}{D} \partial_{x_k} \tilde{u}_k \delta_{nl} \right] + \Psi \left( \partial_{x_l} \partial_{x_n} \tilde{\theta} - \frac{1}{D} \partial_{x_k} \partial_{x_k} \tilde{\theta} \delta_{nl} \right) \end{aligned}$$

where

$$\Xi = \frac{\epsilon^2}{\theta_0} \langle \mathcal{L}_M^{-1} [A_{12}(\mathbf{v})], M \mathcal{L}_M^{-1} [A_{12}(\mathbf{v})] \rangle \quad \Psi = \frac{\epsilon^2}{\theta_0} \langle \mathcal{L}_M^{-1} [A_{12}(\mathbf{v})], M D_{12}(\mathbf{v}) \rangle$$

With  $\bar{m} = \frac{|v|^2}{2}$  in (127), we get

$$\begin{aligned} \epsilon^2 \left( \frac{\text{St } \partial_t \partial_{x_j} \tilde{\theta}}{\theta_0^2} \langle \mathcal{L}_M^{-1} [B_l(\mathbf{v})], M \mathcal{L}_M^{-1} [B_j(\mathbf{v})] \rangle + \frac{\partial_{x_k} \partial_{x_j} \tilde{u}_i}{\theta_0} \langle D_{lk}(\mathbf{v}), M \mathcal{L}_M^{-1} [A_{ij}(\mathbf{v})] \rangle \right) \\ = \Upsilon \delta_{jl} \text{St } \partial_t \partial_{x_j} \tilde{\theta} + \frac{\theta_0 \Psi}{2} \phi_{lk ij} \partial_{x_k} \partial_{x_j} \tilde{u}_i \\ = \Upsilon \text{St } \partial_t \partial_{x_l} \tilde{\theta} + \frac{\theta_0 \Psi}{2} \left( \partial_{x_j} \partial_{x_j} \tilde{u}_l + \left( 1 - \frac{2}{D} \right) \partial_{x_l} \partial_{x_k} \tilde{u}_k \right) \end{aligned}$$

with

$$\Upsilon = \frac{\epsilon^2}{\theta_0^2} \langle \mathcal{L}_M^{-1} [B_1(\mathbf{v})], M \mathcal{L}_M^{-1} [B_1(\mathbf{v})] \rangle$$

For the BGK operator an explicit calculation gives

$$\Xi = \Psi = \epsilon^2 \tau^2 \rho_0 \theta_0 \quad \text{and} \quad \Upsilon = \frac{D+2}{2} \epsilon^2 \tau^2 \rho_0 \theta_0$$



### E.3 Entropy Stable Extension terms

The new fine scale term is

$$\begin{aligned}
& -\epsilon^3 \left\langle \bar{m}, M v_n \partial_{x_n} \left[ \mathcal{L}_M^{-1} v_m \partial_{x_m} \left[ \mathcal{L}_M^{-1} v_k \partial_{x_k} \left[ \mathcal{L}_M^{-1} v_j \partial_{x_j} \bar{f} \right] \right] \right] \right\rangle \\
& = -\epsilon^3 \partial_{x_n} \underbrace{\left\langle \mathcal{L}_M^{-1} [v_n \bar{m}], M v_m \partial_{x_m} \left[ \mathcal{L}_M^{-1} v_k \partial_{x_k} \left[ \mathcal{L}_M^{-1} [v_j \partial_{x_j} \bar{f}] \right] \right] \right\rangle}_{\text{correction to } \sigma_{np}^{(1)} \text{ and } q_n^{(1)}} \quad (128)
\end{aligned}$$

where we used the self-adjointness of  $\mathcal{L}_M^{-1}$  in the right-hand side of the above. The mass conservation equation is unchanged because when  $\bar{m} = 1$  as  $\mathcal{L}_M^{-1}[v_n] = 0$ .

When  $\bar{m} = \frac{|\mathbf{v}|^2}{2}$ , the only non-zero term that results can be written as

$$-\epsilon^3 \left( \frac{\partial_{x_m} \partial_{x_k} \partial_{x_j} \tilde{\theta}}{\theta_0^2} \left\langle \mathcal{L}_M^{-1} [D_{nm}(\mathbf{v})], M D_{jk}(\mathbf{v}) \right\rangle \right)$$

If we define the tensor  $S_{cd}^{ab} = \langle \mathcal{L}_M^{-1} [D_{ab}(\mathbf{v})], M D_{cd}(\mathbf{v}) \rangle$  and note that it is a rank four isotropic tensor, then this term can be written as

$$\begin{aligned}
& -\frac{\epsilon^3 \partial_{x_m} \partial_{x_k} \partial_{x_j} \tilde{\theta}}{\theta_0^2} \left( \frac{(D+1)S_{aa}^{bb} - S_{ab}^{ab} - S_{ab}^{ba}}{D(D-1)(D+2)} \delta_{nm} \delta_{jk} + \frac{(D+1)S_{ab}^{ab} - S_{aa}^{bb} - S_{ab}^{ba}}{D(D-1)(D+2)} \delta_{nj} \delta_{mk} \right. \\
& \quad \left. + \frac{(D+1)S_{ab}^{ba} - S_{ab}^{ab} - S_{aa}^{bb}}{D(D-1)(D+2)} \delta_{nk} \delta_{mj} \right) \\
& = -\frac{\epsilon^3 (S_{aa}^{bb} + S_{ab}^{ab} + S_{ab}^{ba})}{D(D+2)\theta_0^2} \partial_{x_n} \partial_{x_k} \partial_{x_j} \tilde{\theta}
\end{aligned}$$

Finally using the left hand side of the above expression, one can show that

$S_{11}^{11} = \frac{(S_{aa}^{bb} + S_{ab}^{ab} + S_{ab}^{ba})}{D(D+2)}$  which means that

$$\tilde{\Lambda} = -\frac{\epsilon^3}{\theta_0^2} S_{11}^{11} \quad (129)$$

From which we can surmise that  $\tilde{\Lambda}$  is non-negative because the negative definiteness of the linearized collision operator ensures that  $S_{11}^{11} \leq 0$ .

For the BGK operator, a direct computation gives

$$\tilde{\Lambda} = \frac{D^2 + 7D - 2}{D} \epsilon^3 \tau^3 \rho_0 \theta_0^2$$

With  $\bar{m} = v_p$ , the calculation is substantially more complicated. The non-zero term we get from the correction (128) can be written as

$$-\epsilon^3 \left( \frac{\partial_{x_m} \partial_{x_k} \partial_{x_j} \tilde{u}_i}{\theta_0} \left\langle \mathcal{L}_M^{-1} [C_{pnm}(\mathbf{v})], M C_{ijk}(\mathbf{v}) \right\rangle \right) \quad (130)$$

We observe that the rank six isotropic tensor  $K_{ijk}^{pnm} := \langle \mathcal{L}_M^{-1} [C_{pnm}(\mathbf{v})], M C_{ijk}(\mathbf{v}) \rangle$  is such that  $K_{ijk}^{pnm} = K_{ijk}^{npm} =$

$K_{jik}^{pnm}$  and  $K_{ijk}^{nmm} = 0 = K_{iik}^{pnm}$ . As such for  $D \geq 3$ , the tensor takes the form

$$K_{ijk}^{pnm} = a_1 \delta_{mk} \phi_{ijpn} + a_2 \left( \delta_{pm} \phi_{ijnk} + \delta_{nm} \phi_{ijpk} - \frac{2}{D} \delta_{pn} \phi_{ijmk} \right) + a_3 \left( \delta_{nk} \phi_{ijpm} + \delta_{pk} \phi_{ijmn} - \frac{2}{D} \delta_{pn} \phi_{ijmk} \right) \quad (131)$$

where  $a_1$ ,  $a_2$  and  $a_3$  satisfy the system of equations

$$\begin{aligned} \frac{K_{ijk}^{ijk}}{(D-1)(D+2)} &= D a_1 + 2a_2 + 2a_3 \\ \frac{DK_{ikj}^{ijk}}{(D-1)(D+2)} &= D a_1 + (D-2)a_2 + (D-1)(D+2)a_3 \\ \frac{DK_{ikk}^{inn}}{(D-1)(D+2)} &= D a_1 + (D-1)(D+2)a_2 + (D-2)a_3 \end{aligned}$$

Thus (130) takes the form

$$\begin{aligned} -\frac{\epsilon^3}{\theta_0} \left( (a_1 + a_2 + a_3) \partial_{x_k} \partial_{x_k} \left[ \partial_{x_n} \tilde{u}_p + \partial_{x_p} \tilde{u}_n - \frac{2}{D} \delta_{np} \operatorname{div} \tilde{\mathbf{u}} \right] \right. \\ \left. + 2(a_2 + a_3) \left( 1 - \frac{2}{D} \right) (\partial_{x_n} \partial_{x_p} - \frac{\delta_{np}}{D} \partial_{x_k} \partial_{x_k}) [\operatorname{div} \tilde{\mathbf{u}}] \right) \end{aligned}$$

Using (131), one can show that

$$a_1 + a_2 + a_3 = K_{121}^{121} \quad \text{and} \quad a_1 + a_2 = \frac{K_{112}^{121} + K_{122}^{111}}{2 \left( 1 - \frac{2}{D} \right)}$$

and we can thus define the parameters  $\tilde{\Phi}_1$  and  $\tilde{\Phi}_2$  as

$$\begin{aligned} \tilde{\Phi}_1 &= -\frac{\epsilon^3}{\theta_0} (a_1 + a_2 + a_3) = -\frac{\epsilon^3}{\theta_0} K_{121}^{121} \\ \tilde{\Phi}_2 &= -\frac{\epsilon^3}{\theta_0} (a_2 + a_3) = -\frac{\epsilon^3}{\theta_0} \frac{K_{112}^{121} + K_{111}^{122}}{2 \left( 1 - \frac{2}{D} \right)} \end{aligned}$$

Due to the negative definiteness of  $\mathcal{L}_M^{-1}$ , we surmise that  $\tilde{\Phi}_1$  is non-negative. The sign of  $\tilde{\Phi}_2$  is not immediately obvious from the above formula. However, we verified that  $\tilde{\Phi}_2$  is positive albeit smaller than  $\tilde{\Phi}_1$  for the BGK, Maxwell molecule and Hard Spheres collision operators.

For the BGK operator, a direct computation shows that

$$\tilde{\Phi}_1 = 2\tilde{\Phi}_2 = 2\epsilon^3 \tau^3 \rho_0 \theta_0^2$$

## F Hard Spheres calculations of fluid parameters

The calculation of the fluid parameters for the Hard sphere collision operator is based on the methodology outlined in [36]. With  $\mathbf{w} := \frac{\mathbf{v}}{\sqrt{2\theta_0}}$  and  $\hat{\mathbf{w}} := \frac{\mathbf{w}}{|\mathbf{w}|}$ , we define the Burnett functions

$$\chi_{nlm}(\mathbf{w}) = \sqrt{\frac{2(n!)}{(n+l+\frac{1}{2})!}} L_n^{l+\frac{1}{2}}(|\mathbf{w}|^2) |\mathbf{w}|^l Y_l^m(\hat{\mathbf{w}})$$

where  $L_n^\alpha$  is an associated Laguerre polynomial and  $Y_l^m$  is a spherical harmonic. This family of functions possesses the following orthogonality property:

$$\langle \chi_{nlm}^*, M \chi_{pqr} \rangle = \frac{\rho_0}{\sqrt{\pi^3}} \int_{\mathbb{R}^3} \chi_{nlm}^*(\mathbf{w}) e^{-|\mathbf{w}|^2} \chi_{pqr}(\mathbf{w}) d\mathbf{w} = \frac{\rho_0}{\sqrt{\pi^3}} \delta_{np} \delta_{lq} \delta_{mr}$$

where  $\chi_{nlm}^*$  is the complex conjugate of  $\chi_{nlm}$ . The decomposition of a function  $h \in \mathcal{L}^2(e^{-|\mathbf{w}|^2} d\mathbf{w})$  will be

$$h(\mathbf{w}) = \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \sum_{m=-l}^l h_{nlm} \chi_{nlm}(\mathbf{w}), \quad \text{where } h_{nlm} = \int_{\mathbb{R}^3} \chi_{nlm}^*(\mathbf{w}) e^{-|\mathbf{w}|^2} h(\mathbf{w}) d\mathbf{w}$$

If we orient the spherical coordinate system such that  $\hat{\mathbf{w}} = (\cos \theta, \sin \theta \sin \phi, \sin \theta \cos \phi)$ , then decomposition of the collision invariants will be

$$\begin{aligned} 1 &= \pi^{\frac{3}{4}} \chi_{000}(\mathbf{w}) & w_1 &= \frac{\pi^{\frac{3}{4}}}{\sqrt{2}} \chi_{010}(\mathbf{w}) \\ w_2 &= \frac{\mathbf{i}\pi^{\frac{3}{4}}}{2} (\chi_{011}(\mathbf{w}) + \chi_{01-1}(\mathbf{w})) & w_3 &= \frac{\pi^{\frac{3}{4}}}{2} (\chi_{01-1}(\mathbf{w}) - \chi_{011}(\mathbf{w})) \\ |\mathbf{w}|^2 &= \pi^{\frac{3}{4}} \left( \frac{3}{2} \chi_{000}(\mathbf{w}) - \sqrt{\frac{3}{2}} \chi_{100}(\mathbf{w}) \right) \end{aligned}$$

This means that the projection of a function onto the orthogonal complement of the collision invariants can be accomplished simply by removing all  $\chi_{000}$ ,  $\chi_{01m}$  and  $\chi_{100}$  components from the function's decomposition.

The action of the linearized collision operator and its inverse can be characterized as

$$\begin{aligned} \mathcal{L}_M[h] &= \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \sum_{m=-l}^l \left( \sum_{p=0}^{\infty} [\mathfrak{L}^l]_{np} h_{plm} \right) \chi_{nlm}(\mathbf{w}) \\ \mathcal{L}_M^{-1}[h] &= \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \sum_{m=-l}^l \left( \sum_{p=0}^{\infty} [\mathfrak{L}^l]_{np}^{-1} h_{plm} \right) \chi_{nlm}(\mathbf{w}) \end{aligned}$$

where the  $l$ -parametrized family of infinite symmetric matrices  $[\mathfrak{L}^l]$  are derived from the identity

$$\int_{\mathbb{R}^3} \chi_{nlm}^* e^{-|\mathbf{w}|^2} \mathcal{L}_M[\chi_{pqr}] d\mathbf{w} = [\mathfrak{L}^l]_{np} \delta_{lq} \delta_{mr}$$

and  $[\mathfrak{L}^l]^{-1}$  represents the corresponding (pseudo-)inverse.<sup>15</sup>

<sup>15</sup>Note that the indices for both  $[\mathfrak{L}^l]$  and  $[\mathfrak{L}^l]^{-1}$  start from  $n=0$ ,  $p=0$ .

For the hard spheres operator, the entries of  $[\mathcal{L}^l]$  are<sup>16</sup>

$$[\mathcal{L}^l]_{np} = \frac{1}{\tau\sqrt{2}} \sqrt{\frac{n!p!}{8\Gamma(n+l+\frac{3}{2})\Gamma(p+l+\frac{3}{2})}} \sum_{j=0}^{\min\{n,p\}} \sum_{k=0}^l \frac{l!\Gamma(-\frac{1}{2}+n+p+l-2j-k)}{(n-j)!(p-j)!(l-k)!2^{n+p+l-2j-k}} B_j^k$$

where  $B_j^k = \frac{(j+k+1)!}{j!k!} + \delta_{j0}\delta_{k0} - 2^{1-k} \binom{2j+k+1}{k}$ .

For our calculation, we found that it was enough to calculate the first 200 rows and columns of  $[\mathcal{L}^l]$  for  $l \in \{0, 1, 2, 3\}$ . We use these finite matrices to compute the corresponding pseudo-inverses  $[\mathcal{L}^l]^{-1}$ . For brevity, we will only describe the calculation of  $\tilde{\kappa}$  and  $\tilde{\Lambda}$ . The process for calculating  $\tilde{\omega}$ ,  $\tilde{\Phi}_1$  and  $\tilde{\Phi}_2$  follow similar steps.

We first note that

$$B_1(\mathbf{v}) = \frac{\theta_0^{\frac{3}{2}}}{2} (|\mathbf{w}|^2\sqrt{8} - 5\sqrt{2})w_1 = -\left(\pi^{\frac{3}{4}}\sqrt{\frac{5\theta_0^3}{2}}\right)\chi_{110}(\mathbf{w})$$

Thus,

$$\begin{aligned} \tilde{\kappa} &= -\frac{\epsilon}{\theta_0^2} \langle B_1(\mathbf{v}), M\mathcal{L}_M^{-1}[B_1(\mathbf{v})] \rangle \\ &= -\frac{5}{2}\epsilon\rho_0\theta_0 \int_{\mathbb{R}^3} \chi_{110}^*(\mathbf{w}) e^{-|\mathbf{w}|^2} \sum_{n=0}^{\infty} [\mathcal{L}^1]_{1n}^{-1} \chi_{n10}(\mathbf{w}) d\mathbf{w} \\ &= -\frac{5}{2}\epsilon\rho_0\theta_0 [\mathcal{L}^1]_{11}^{-1} = \frac{5}{2}(1.922284) \epsilon\tau\rho_0\theta_0 \end{aligned}$$

Next<sup>17</sup>

$$\begin{aligned} D_{11}(\mathbf{v}) &= v_1 \mathcal{L}_M^{-1}[B_1(\mathbf{v})] = -\pi^{\frac{3}{4}}\theta_0^2\sqrt{5} \sum_{n=0}^{\infty} [\mathcal{L}^1]_{1n}^{-1} (w_1 \chi_{n10}(\mathbf{w})) \\ &= -\pi^{\frac{3}{4}}\theta_0^2\sqrt{5} \sum_{n=0}^{\infty} [\mathcal{L}^1]_{1n}^{-1} \sum_{n_1=0}^{\infty} (\alpha_{nn_1} \chi_{n_100}(\mathbf{w}) + \beta_{nn_1} \chi_{n_120}(\mathbf{w})) \end{aligned}$$

where  $\alpha_{nn_1} = \sqrt{\frac{3+2n}{6}} \delta_{nn_1} - \sqrt{\frac{n+1}{3}} \delta_{n\{n_1-1\}}$  and  $\beta_{nn_1} = \sqrt{\frac{10+4n}{15}} \delta_{nn_1} - \sqrt{\frac{4n}{15}} \delta_{n\{n_1+1\}}$ . As a result

$$\begin{aligned} \mathcal{L}_M^{-1}[D_{11}(\mathbf{v})] &= -\pi^{\frac{3}{4}}\theta_0^2\sqrt{5} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \left( [\mathcal{L}^0]_{n_1n_2}^{-1} \left( \sum_{n=0}^{\infty} [\mathcal{L}^1]_{1n}^{-1} \alpha_{nn_2} \right) \chi_{n_100}(\mathbf{w}) \right. \\ &\quad \left. + [\mathcal{L}^2]_{n_1n_2}^{-1} \left( \sum_{n=0}^{\infty} [\mathcal{L}^1]_{1n}^{-1} \beta_{nn_2} \right) \chi_{n_120}(\mathbf{w}) \right) \end{aligned}$$

Thus

$$\begin{aligned} \tilde{\Lambda} &= -\frac{\epsilon^3}{\theta_0^2} \langle D_{11}(\mathbf{v}), M\mathcal{L}_M^{-1}[D_{11}(\mathbf{v})] \rangle \\ &= -5\epsilon^3\rho_0\theta_0^2 \left( \sum_{n_1=0}^{\infty} \left( \sum_{n=0}^{\infty} [\mathcal{L}^1]_{1n}^{-1} \alpha_{nn_1} \right) \sum_{n_1=0}^{\infty} [\mathcal{L}^0]_{n_1n_2}^{-1} \left( \sum_{n=0}^{\infty} [\mathcal{L}^1]_{1n}^{-1} \alpha_{nn_2} \right) \right. \\ &\quad \left. + \sum_{n_1=0}^{\infty} \left( \sum_{n=0}^{\infty} [\mathcal{L}^1]_{1n}^{-1} \beta_{nn_1} \right) \sum_{n_1=0}^{\infty} [\mathcal{L}^2]_{n_1n_2}^{-1} \left( \sum_{n=0}^{\infty} [\mathcal{L}^1]_{1n}^{-1} \beta_{nn_2} \right) \right) \\ &= 37.47015 \epsilon^3 \tau^3 \rho_0 \theta_0^2 \end{aligned}$$

<sup>16</sup>See [36] for a derivation

<sup>17</sup>Strictly speaking, we have to apply  $(\text{Id} - \Pi_M)$  in the definition of  $D_{11}$ . However, the inner product with  $\mathcal{L}_M^{-1}[D_{11}(\mathbf{v})]$  automatically accomplishes this projection.

Although we have presented the above calculation in terms of infinite sums, in practice, truncating each of the sums after about fifteen terms gives a result accurate to seven significant figures.

The above calculations can be repeated for any collision operator provided their corresponding collision matrices  $[\mathfrak{L}^l]$  are available. For example, by observing that the BGK operator has collision matrices of the form

$$\tau[\mathfrak{L}^l]_{np} = \frac{1}{\tau}[\mathfrak{L}^l]_{np}^{-1} = \begin{cases} 0, & l = 0 \text{ and } n \in \{0, 1\} \text{ or } p \in \{0, 1\} \\ 0, & l = 1 \text{ and } n = 0 \text{ or } p = 0 \\ -\delta_{np}, & \text{otherwise} \end{cases}$$

one can easily obtain the BGK values reported in this manuscript with the formulas derived above.