

Automorphisms of the DAHA of type \check{C}_1C_1 and their action on Askey–Wilson polynomials and functions. I.

The flip $(a, b, c, d) \mapsto (a, b, qd^{-1}, qc^{-1})$

Tom Koornwinder* and Marta Mazzocco†

Abstract

In this paper we consider the automorphisms of the double affine Hecke algebra (DAHA) of type \check{C}_1C_1 which have a relatively simple action on the generators and on the parameters, notably a symmetry t_4 which sends the AW parameters (a, b, c, d) to (a, b, qd^{-1}, qc^{-1}) . We study how these symmetries act on the basic representation and on the symmetric and non-symmetric Askey–Wilson (AW) polynomials and functions. Interestingly t_4 maps AW polynomials to functions. In the second part of the paper we look for a version of non-symmetric AW functions that behave well under these symmetries.

1 Introduction

This paper marks the beginning of a programme to consider and possibly classify the symmetries of the double affine Hecke algebra (DAHA) of type \check{C}_1C_1 and of the Zhedanov algebra, to see how the symmetries of these two algebras are related, and to study the actions of these symmetries on the basic representation and on the symmetric and non-symmetric Askey–Wilson (AW) polynomials and functions.

Some symmetries were earlier observed in the literature. Oblomkov [19, Proposition 2.4] gives an extended modular group $\bar{\Gamma}$ action on the DAHA, while Crampé et al. [2, §4] and the first author [12, §2.3] observe an action of $W(D_4)$ (the Weyl group of root system D_4) on the Zhedanov algebra. A natural question is therefore to understand what these actions do in the basic representation and whether they map AW polynomials to AW polynomials. Interestingly, it turns out that some of them actually map AW polynomials into AW functions, so that it is more natural to see whether they map AW functions to AW functions.

Our programme to systematically study the symmetries of the DAHA of type \check{C}_1C_1 is motivated by two main reasons. On one side, the action of a symmetry on the DAHA implies new relations between AW polynomials or functions at different parameters and/or different values of the variable. On the other side, since the \check{C}_1C_1 DAHA can be obtained as quantisation of the group algebra of the monodromy group associated to the sixth Painlevé equation (PVI) [17],

*thkmath@xs4a11.nl, Korteweg–de Vries Institute, University of Amsterdam

†m.mazzocco@bham.ac.uk, School of Mathematics, University of Birmingham

it is natural to ask whether the symmetry group G of PVI lifts to the \check{C}_1C_1 DAHA. Okamoto [20] proved that G is isomorphic to the affine Weyl group of the exceptional root system F_4 . The group G contains as subgroups the affine $W(D_4)$, its extension by the Klein four-group \mathcal{K}_4 and three more transformations which can be represented by the action of $\bar{\Gamma}$. So finding the action of affine $W(D_4)$ and $\bar{\Gamma}$ on the \check{C}_1C_1 DAHA would provide an important step to answer this question. This is what we do in the first part of this paper.

Then, in order to start the study of the actions of these symmetries on the basic representation and on the symmetric and non-symmetric Askey–Wilson (AW) polynomials and functions, we concentrate on the symmetry t_4 which sends the AW parameters (a, b, c, d) to (a, b, qd^{-1}, qc^{-1}) . We find that to study the action of t_4 on eigenfunctions of the AW operator, the second order q -difference operator of which the AW polynomials are eigenfunctions, we are naturally guided to AW functions. Indeed, on AW polynomials the permutations of (a, b, c, d) are acting, but not t_4 , while on AW functions the symmetries of (a, b, c, d) are acting which permute (a, b, c, qd^{-1}) .

Next we look for a version of non-symmetric AW functions as analytic continuations with respect to the degree of non-symmetric AW polynomials [10]. These are written as the sum of an ordinary (symmetric) AW function and an anti-symmetric term involving a symmetric AW function with shifted parameters. Expressions of one-variable non-symmetric special functions of this form are also well-known [13] for Bessel functions and for Jacobi polynomials. The analytic continuation from the non-symmetric AW polynomial case is not unique, but with some effort we arrive at a defining expression for non-symmetric AW functions which satisfies all the desired symmetries, and which moreover can be shown to be equal to the one-variable case of Stokman’s Cherednik kernel [25, Definition 5.16].

We should emphasize here that, when studying AW functions, we are mostly interested in eigenfunctions of the AW operator which act as a kernel in a Fourier type integral transform, as is the case with the AW functions considered in [9] and the non-symmetric AW functions coming from [25].

The contents of this paper are as follows. In Section 2 we introduce the DAHA of type \check{C}_1C_1 in two different but equivalent presentations. In Section 3 we introduce some DAHA automorphisms: the parameter inversions and the extended modular group. In Section 4 we define the basic DAHA representation and consider the action of automorphisms, notably of t_4 , on it. We also discuss the spherical subalgebra (which is isomorphic to the Zhedanov algebra at a special scalar value of the Casimir element), its polynomial representation, and its symmetries. In Section 5 we introduce AW polynomials and functions and give some of their properties. Next Section 6 deals with the action of DAHA automorphisms on AW polynomials and functions. Finally Section 7 culminates in an expression for non-symmetric AW functions together with their symmetries. It ends with showing that our functions are equal to the one-variable case of Stokman’s Cherednik kernel

As a follow-up to this paper we have two further papers in preparation, to which we will sometimes refer in this paper. The tentative titles are:

[P2] (with D. Dal Martello) II. Quantum middle convolution.

[P3] III. Further automorphisms.

Conventions For q -shifted factorials and q -hypergeometric series we use notation as in [4, §1.2]. In this paper $q \in (0, 1)$ will be fixed. Throughout we suppress the dependence on q as a separate parameter in the notation.

We will denote $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$. Omnipresent parameters a, b, c, d will be generic. In particular they are supposed to be complex, nonzero, and the parameters together with their inverses should be all distinct.

For $z \in \mathbb{C} \setminus (-\infty, 0]$ the square root \sqrt{z} , continuously depending on z , will be uniquely chosen such that $\sqrt{z} > 0$ if $z > 0$.

2 DAHA of type \check{C}_1C_1

We present the DAHA of type \check{C}_1C_1 as in our earlier papers [10, (3.1)–(3.4)], [16, (1.1)–(1.5)] and [14, (84)]. This is essentially the case $n = 1$ of Sahi’s presentation [23, §3].

Definition 2.1. The DAHA of type \check{C}_1C_1 , denoted by $\mathcal{H}(a, b, c, d)$, is the unital associative \mathbb{C} -algebra generated by T_1, T_0, Z with relations

$$(T_1 + ab)(T_1 + 1) = 0, \quad (2.1)$$

$$(T_0 + q^{-1}cd)(T_0 + 1) = 0, \quad (2.2)$$

$$(T_1Z + a)(T_1Z + b) = 0, \quad (2.3)$$

$$(qT_0Z^{-1} + c)(qT_0Z^{-1} + d) = 0. \quad (2.4)$$

Note that (2.1)–(2.3) imply that T_1, T_0, Z have two-sided inverses, respectively given by

$$\begin{aligned} T_1^{-1} &= -a^{-1}b^{-1}T_1 - (1 + a^{-1}b^{-1}), & T_0^{-1} &= -qc^{-1}d^{-1}T_0 - (1 + qc^{-1}d^{-1}), \\ Z^{-1} &= -a^{-1}b^{-1}T_1ZT_1 - (a^{-1} + b^{-1})T_1. \end{aligned} \quad (2.5)$$

Further important elements of $\mathcal{H}(a, b, c, d)$ are

$$Y := T_1T_0, \quad Y^{-1} = T_0^{-1}T_1^{-1}, \quad D := Y + q^{-1}abcdY^{-1}, \quad X := Z + Z^{-1}. \quad (2.6)$$

The following lemma is well-known. For completeness we sketch a proof.

Lemma 2.2. *The elements X and D commute with T_1 .*

Proof. It follows from (2.1) and (2.3) that

$$T_1Z - Z^{-1}T_1 = (ab + 1)Z^{-1} - (a + b) = ZT_1 - T_1Z^{-1}.$$

Hence T_1 commutes with $Z + Z^{-1} = X$. Also, it follows from (2.1) and (2.2) that

$$T_1^2T_0 - q^{-1}abcdT_0^{-1} = -(ab + 1)T_1T_0 + ab(q^{-1}cd + 1) = T_1T_0T_1 - q^{-1}abcdT_1T_0^{-1}T_1^{-1}.$$

Hence T_1 commutes with $T_1T_0 + q^{-1}abcdT_0^{-1}T_1^{-1} = D$. \square

Corollary 2.3. *If $f(Z)$ is a Laurent polynomial in Z such that $f(Z) = f(Z^{-1})$ then $f(Z)$ commutes with T_1 .*

Proof. By Lemma 2.2 T_1 commutes with $X^n = (Z + Z^{-1})^n$ ($n \in \mathbb{Z}_{\geq 0}$). Now it follows by complete induction that T_1 commutes with $Z^n + Z^{-n}$ ($n \in \mathbb{Z}_{> 0}$). \square

In order to compare relations (2.1)–(2.4) with an alternative presentation of the DAHA of type \check{C}_1C_1 below it is convenient to rewrite (2.1), (2.2) as the following equivalent relations.

$$(T_1^{-1}Z^{-1} + a^{-1})(T_1^{-1}Z^{-1} + b^{-1}) = 0, \quad (2.7)$$

$$(ZT_0^{-1} + qc^{-1})(ZT_0^{-1} + qd^{-1}) = 0. \quad (2.8)$$

In [18, Theorem 2.22], [19, Definition 2.1] the DAHA of type \check{C}_1C_1 is defined as an algebra $H(u_1, k_1, k_0, u_0)$ with generators $\check{V}_1, V_1, V_0, \check{V}_0$, and relations

$$(\check{V}_1 - u_1)(\check{V}_1 + u_1^{-1}) = 0, \quad (2.9)$$

$$(V_1 - k_1)(V_1 + k_1^{-1}) = 0, \quad (2.10)$$

$$(V_0 - k_0)(V_0 + k_0^{-1}) = 0, \quad (2.11)$$

$$(\check{V}_0 - u_0)(\check{V}_0 + u_0^{-1}) = 0, \quad (2.12)$$

$$\check{V}_1 V_1 V_0 \check{V}_0 = q^{-1/2}, \quad (2.13)$$

where $u_1, k_1, k_0, u_0 \in \mathbb{C}^*$. Here the Hecke relations (2.9)–(2.12) immediately yield explicit two-sided inverses of the four generators. Relation (2.13) remains unchanged under cyclic permutation of the factors.

The algebras $\mathcal{H}(a, b, c, d)$ and $H(u_1, k_1, k_0, u_0)$, with parameters related by

$$a = -u_1 k_1^{-1}, \quad b = u_1 k_1, \quad c = q^{1/2} u_0 k_0, \quad d = -q^{1/2} u_0 k_0^{-1}, \quad (2.14)$$

vice versa:

$$u_1 = i\sqrt{ab}, \quad k_1 = -i\sqrt{a^{-1}b}, \quad u_0 = -i\sqrt{q^{-1}cd}, \quad k_0 = i\sqrt{cd^{-1}}, \quad (2.15)$$

are isomorphic by an isomorphism $\phi = \phi_{a,b,c,d}$, explicitly given by

$$\begin{aligned} \phi(T_0) &= u_0 \check{V}_0 = -i\sqrt{q^{-1}cd} \check{V}_0, & \phi(T_1) &= u_1 \check{V}_1 = i\sqrt{ab} \check{V}_1, \\ \phi(Z) &= q^{1/2} V_0 \check{V}_0, & \phi(Z^{-1}) &= \check{V}_1 V_1, \end{aligned} \quad (2.16)$$

vice versa:

$$\begin{aligned} \phi^{-1}(\check{V}_0) &= i\sqrt{qc^{-1}d^{-1}} T_0 = u_0^{-1} T_0, & \phi^{-1}(\check{V}_1) &= -i\sqrt{a^{-1}b^{-1}} T_1 = u_1^{-1} T_1, \\ \phi^{-1}(V_0) &= -iq^{-1}\sqrt{cd} Z T_0^{-1} = q^{-1/2} u_0 Z T_0^{-1}, & \phi^{-1}(V_1) &= i\sqrt{ab} T_1^{-1} Z^{-1} = u_1 T_1^{-1} Z^{-1}. \end{aligned} \quad (2.17)$$

In (2.15) and (2.17) recall our convention about square roots, by which we also immediately see:

Lemma 2.4. *$a, b, c, d, ab, ab^{-1}, cd, cd^{-1}$ have real part > 0 iff $k_0, -k_1, -u_0, u_1$ have argument $> \pi/4$ and $< 3\pi/4$.*

Remark 2.5. By comparing the Hecke relations (2.1), (2.2), (2.7), (2.8) on the one hand and (2.9)–(2.12) on the other hand, we see that the particular choices made in (2.14)–(2.17) identify $\check{V}_1, V_1, V_0, \check{V}_0$ with the given constant multiples of $T_1, T_1^{-1}Z^{-1}, ZT_0^{-1}, T_0$, respectively, in this particular order, because the cyclic relation (2.13) must be equivalent to $ZZ^{-1} = 1$. The only freedom left is that we might have simultaneously performed cyclic permutation of $\check{V}_1, V_1, V_0, \check{V}_0$ and u_1, k_1, k_0, u_0 .

3 Some DAHA automorphisms

In this section we first define what we mean by a DAHA automorphism. Then we introduce the special automorphisms t_2, t_3, t_4, σ and τ which will be of special interest for their possible action on AW polynomials and functions because they leave T_1 invariant. In connection with σ we introduce dual parameters $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}$. The section ends with short discussions of the extended modular group and the spherical braid group acting on the DAHA. We postpone a more comprehensive treatment of DAHA automorphisms to a next paper [P3].

3.1 Definition of a DAHA automorphism

Definition 3.1. An *automorphism* t of the parametrized DAHA family $\{\mathcal{H}(a, b, c, d)\}$ is a bijective map $t: (a, b, c, d) \rightarrow (a', b', c', d')$ together with bijective DAHA isomorphisms $t = t_{a,b,c,d}: \mathcal{H}(a, b, c, d) \rightarrow \mathcal{H}(a', b', c', d')$. In other words, the relations (2.1)–(2.4) with (T_0, T_1, Z) replaced by $(t(T_0), t(T_1), t(Z))$ have to be equivalent to the same relations with (a, b, c, d) replaced by $t(a, b, c, d)$.

The definition of a DAHA automorphism for the parametrized family $\{H(u_1, k_1, k_0, u_0)\}$ is similar. We can relate a DAHA automorphism t of $\{\mathcal{H}(a, b, c, d)\}$ and a DAHA automorphism of $\{H(u_1, k_1, k_0, u_0)\}$ (by abuse of notation again called t) by the following commutative diagrams.

$$\begin{array}{ccc}
(a, b, c, d) & \xrightarrow{t} & (a', b', c', d') \\
\downarrow & & \downarrow \\
(u_1, k_1, k_0, u_0) & \xrightarrow{t} & (u'_1, k'_1, k'_0, u'_0)
\end{array}
\qquad
\begin{array}{ccc}
(T_0, T_1, Z) & \xrightarrow{t_{a,b,c,d}} & (T'_0, T'_1, Z') \\
\downarrow \phi_{a,b,c,d} & & \downarrow \phi_{a',b',c',d'} \\
(\check{V}_1, V_1, V_0, \check{V}_0) & \xrightarrow{t_{u_1, k_1, k_0, u_0}} & (\check{V}'_1, V'_1, V'_0, \check{V}'_0)
\end{array}$$

In the right vertical arrow in the right diagram we evaluate $\phi_{a',b',c',d'}(U') = \phi_{a',b',c',d'}(t_{a,b,c,d}(U))$ ($U = T_0, T_1$ or Z) by applying $\phi_{a',b',c',d'}$ as an algebra homomorphism to the algebraic expression in T_0, T_1, Z given by $t_{a,b,c,d}(TU)$. Here $\phi_{a',b',c',d'}$ only acts on the generators T_0, T_1, Z , but leaves scalars unchanged, even if they depend on a, b, c, d . See later in this section a few worked out examples, for instance for t_4 in Table 1.

If $s \circ t$ is a composition of bijections of (a, b, c, d) parameter space then we have correspondingly $(s \circ t)_{a,b,c,d} = s_{t(a,b,c,d)} \circ t_{a,b,c,d}: H(a, b, c, d) \rightarrow H(t(a, b, c, d)) \rightarrow H(s(t(a, b, c, d)))$.

An alternative definition of DAHA automorphism takes $\mathcal{H}(a, b, c, d)$ with (a, b, c, d) also being considered as generators and with relations added which just say that the generators a, b, c, d are in the center. Then we can define an automorphism t just of $\mathcal{H}(a, b, c, d)$ rather than of

the parametrized family $\{\mathcal{H}(a, b, c, d)\}$. This automorphism has to be such that it sends the generators a, b, c, d to expressions only depending on a, b, c, d and the generators T_0, T_1, Z to expressions possibly depending on all generators including a, b, c, d .

Definition 3.1 is similar to the one used by Oblomkov [19, Prop. 2.4], while the definition considering the parameters as central generators is used by Sahi [23, §4]. Clearly the two definitions give different outcomes for automorphisms in explicit situations. In the mentioned cases by Oblomkov and Sahi these differences are tempered because the automorphism acting on the parameters is involutive and the automorphism acting on the generators does not depend on the parameters. We keep to definition 3.1 because it is more suitable for action on the basic representation.

3.2 Parameter inversions

We call *parameter inversions* the transformations of $H(u_1, k_1, k_0, u_0)$ that preserve the generators $\check{V}_1, V_1, V_0, \check{V}_0$ and only act on one of the parameters u_1, k_1, k_0, u_0 by inversion and change of sign. We list them in Table 1, where we also give the action of these transformations on $\mathcal{H}(a, b, c, d)$.

	u_1	k_1	k_0	u_0	a	b	c	d	T_0	T_1	Z
t_1	$-\frac{1}{u_1}$	k_1	k_0	u_0	$\frac{1}{b}$	$\frac{1}{a}$	c	d	T_0	abT_1	Z
t_2	u_1	$-\frac{1}{k_1}$	k_0	u_0	b	a	c	d	T_0	T_1	Z
t_3	u_1	k_1	$-\frac{1}{k_0}$	u_0	a	b	d	c	T_0	T_1	Z
t_4	u_1	k_1	k_0	$-\frac{1}{u_0}$	a	b	$\frac{q}{d}$	$\frac{q}{c}$	$\frac{cd}{q}T_0$	T_1	Z

Table 1: Parameter inversions.

It is immediately seen that the actions on (a, b, c, d) as given in the table correspond with the parameter inversions in (u_1, k_1, k_0, u_0) . This correspondence also holds on the DAHA level. For instance, let us show that $t_4(T_0) = q^{-1}cdT_0$ corresponds with $t_4(\check{V}_0, V_0) = (\check{V}_0, V_0)$ (for brevity we write $\phi_{c,d}$ instead of $\phi_{a,b,c,d}$):

$$\begin{array}{ccc}
i\sqrt{qc^{-1}d^{-1}}T_0 & \xrightarrow{t_4} & i\sqrt{q^{-1}cd}T_0 \\
\phi_{c,d}^{-1} \uparrow & & \downarrow \phi_{q/d,q/c} \\
\check{V}_0 & \xrightarrow{t_4} & i\sqrt{q^{-1}cd}\phi_{q/d,q/c}(T_0) = (i\sqrt{q^{-1}cd})(-i\sqrt{q^{-1}cd})\check{V}_0 = \check{V}_0, \\
\\
-iq^{-1}\sqrt{cd}ZT_0^{-1} & \xrightarrow{t_4} & -i(cd)^{-1/2}ZT_0^{-1} \\
\phi_{c,d}^{-1} \uparrow & & \downarrow \phi_{q/d,q/c} \\
V_0 & \xrightarrow{t_4} & -i(cd)^{-1/2}\phi_{q/d,q/c}(Z)\phi_{q/d,q/c}(T_0)^{-1} = V_0.
\end{array}$$

Observe the following properties of the parameter inversions

- The automorphisms t_2 and t_3 even leave all generators T_0, T_1, Z invariant;

- t_4 leaves T_1 and Z invariant.

As a consequence, we will see that t_2, t_3, t_4 act on the spherical subalgebra as well. Interestingly though, while t_2 and t_3 act on symmetric and non-symmetric AW polynomials, we will show in this paper that t_2 and t_4 act on symmetric and non-symmetric AW functions. As mentioned above, we postpone a discussion of the action of t_1 to [P3] because it doesn't restrict to the spherical subalgebra.

Remark 3.2. Note that the parameter inversions in Table 1 exchange the positions of a and b or of c and d , but never mix the pair (a, b) with the pair (c, d) . One symmetry which does a further mixing of parameters is

$$t_0: (a, b, c, d) \mapsto \left(\frac{q}{d}, b, c, \frac{q}{a}\right).$$

However, the action of this transformation on the generators T_0, T_1, Z , or on $\check{V}_1, V_1, V_0, \check{V}_0$ is quite involved and will be the subject of [P2]. Note also that the two equivalent sets of constraints on the parameters in Lemma 2.4 are not preserved by t_0 .

Notice that

$$\hat{t}_0 := (t_2 t_4) t_0 (t_2 t_4)^{-1}: (a, b, c, d) \mapsto (a, c, b, d). \quad (3.1)$$

So, in their action on parameter space, t_0, t_2, t_3, t_4 generate a group isomorphic to the semi-direct product $S_4 \ltimes (\mathbb{Z}/2\mathbb{Z})^3$ (where $(\mathbb{Z}/2\mathbb{Z})^3$ is the group of an even number of sign changes of 1, 2, 3, 4). We recognize this group as the Weyl group $W(D_4)$, see [6, §12.1]. In [P2] the corresponding actions of t_0, t_2, t_3, t_4 on the DAHA will be studied. In Section 4.5 we will come back to $W(D_4)$ for its action on the spherical subalgebra.

If we take t_0, t_1, t_2, t_3 instead of t_0, t_2, t_3, t_4 as generators then another version of $W(D_4)$ acting on the parameter space is obtained. When we include both t_1 and t_4 , i.e., consider $\langle t_0, t_1, t_2, t_3, t_4 \rangle$, we obtain an action of the Weyl group of the affine root lattice $D_4^{(1)}$. This will also be discussed in [P2] and [P3].

3.3 Duality and the automorphism σ

Recall our convention about square roots. For $a, b, c, d \in \mathbb{C}^*$ assume that $q^{-1}abcd \in \mathbb{C} \setminus (-\infty, 0]$. Then *dual parameters* $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}$ are well-defined by

$$\tilde{a} = \sqrt{q^{-1}abcd}, \quad \tilde{a}\tilde{b} = ab, \quad \tilde{a}\tilde{c} = ac, \quad \tilde{a}\tilde{d} = ad. \quad (3.2)$$

Then we have also:

$$\begin{aligned} \tilde{b} &= \sqrt{qabc^{-1}d^{-1}}, & \tilde{c} &= \sqrt{qab^{-1}cd^{-1}}, & \tilde{d} &= \sqrt{qab^{-1}c^{-1}d}, \\ \tilde{b}\tilde{c} &= qad^{-1}, & \tilde{b}\tilde{d} &= qac^{-1}, & \tilde{c}\tilde{d} &= qab^{-1}, & \tilde{a}\tilde{b}^{-1} &= q^{-1}cd, & \tilde{a}\tilde{c}^{-1} &= q^{-1}bd, \\ \tilde{a}\tilde{d}^{-1} &= q^{-1}bc, & \tilde{b}\tilde{c}^{-1} &= bc^{-1}, & \tilde{b}\tilde{d}^{-1} &= bd^{-1}, & \tilde{c}\tilde{d}^{-1} &= cd^{-1}. \end{aligned} \quad (3.3)$$

Note that t_4 acting on a, b, c, d corresponds to t_2 acting on $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}$, while exchange of b, c corresponds to exchange of \tilde{b}, \tilde{c} . Also the following lemma is clear from (3.3).

Lemma 3.3. *If $a, ab, ab^{-1}, cd, cd^{-1}$ have real part > 0 then (i) b, c, d have real part > 0 and $q^{-1}abcd \in \mathbb{C} \setminus (-\infty, 0]$, (ii) the dual parameters are well-defined by (3.2), (iii) $\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}, \tilde{a}\tilde{b}, \tilde{a}\tilde{b}^{-1}, \tilde{c}\tilde{d}, \tilde{c}\tilde{d}^{-1}$ all have real part > 0 and $q^{-1}\tilde{a}\tilde{b}\tilde{c}\tilde{d} \in \mathbb{C} \setminus (-\infty, 0]$, (iv) taking duals of dual parameters once more, we recover a, b, c, d .*

If moreover $ac, ad, bc, bd, ac^{-1}, ad^{-1}, bc^{-1}, bd^{-1}$ have real part > 0 then $\tilde{a}\tilde{c}, \tilde{a}\tilde{d}, \tilde{b}\tilde{c}, \tilde{b}\tilde{d}, \tilde{a}\tilde{c}^{-1}, \tilde{a}\tilde{d}^{-1}, \tilde{b}\tilde{c}^{-1}, \tilde{b}\tilde{d}^{-1}$ have real part > 0 .

This is important because it allows us to uniquely define the square roots involved in the equations (2.14) and (2.15) also after the duality transformation (3.2), (3.3).

By (2.14) and (2.15) the duality map $(a, b, c, d) \rightarrow (\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d})$ translates to $(u_1, k_1, k_0, u_0) \rightarrow (u_1, -u_0^{-1}, k_0, -k_1^{-1})$. A corresponding DAHA automorphism σ which maps $H(u_1, k_1, k_0, u_0)$ to $H(u_1, -u_0^{-1}, k_0, -k_1^{-1})$ can be chosen in several ways. We follow the choice made in [18, Prop. 8.5(ii)], [25, Cor. 2.7] rather than the one in [19, (2.10)] and we define

$$\begin{aligned} \sigma: \quad H(u_1, k_1, k_0, u_0) &\rightarrow H(u_1, -u_0^{-1}, k_0, -k_1^{-1}) \\ (\check{V}_1, V_1, V_0, \check{V}_0) &\mapsto (\check{V}_1, \check{V}_1^{-1}\check{V}_0\check{V}_1, V_1V_0V_1^{-1}, V_1), \end{aligned} \quad (3.4)$$

or equivalently:

$$\begin{aligned} \sigma: \quad \mathcal{H}(a, b, c, d) &\rightarrow \mathcal{H}(\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}) \\ (T_0, T_1, Z) &\mapsto (\tilde{a}T_1^{-1}Z^{-1}, T_1, aT_1^{-1}T_0^{-1}). \end{aligned} \quad (3.5)$$

Let us show, for instance, that the action of σ on $\mathcal{H}(a, b, c, d)$ implies that $\sigma(\check{V}_0) = V_1$ and $\sigma(V_1) = \check{V}_1^{-1}\check{V}_0\check{V}_1$:

$$\begin{array}{ccc} i\sqrt{qc^{-1}d^{-1}}T_0 & \xrightarrow{\sigma} & i\sqrt{ab}T_1^{-1}Z \\ \phi_{a,b,c,d}^{-1} \uparrow & & \downarrow \phi_{\tilde{a},\tilde{b},\tilde{c},\tilde{d}} \\ \check{V}_0 & \xrightarrow{\sigma} & i\sqrt{ab}\phi_{\tilde{a},\tilde{b},\tilde{c},\tilde{d}}(T_1)^{-1}\phi_{\tilde{a},\tilde{b},\tilde{c},\tilde{d}}(Z) = V_1, \end{array}$$

$$\begin{array}{ccc} i\sqrt{ab}T_1^{-1}Z^{-1} & \xrightarrow{\sigma} & i\sqrt{a^{-1}b}T_1^{-1}T_0T_1 \\ \phi_{a,b,c,d}^{-1} \uparrow & & \downarrow \phi_{\tilde{a},\tilde{b},\tilde{c},\tilde{d}} \\ V_1 & \xrightarrow{\sigma} & i\sqrt{a^{-1}b}\phi_{\tilde{a},\tilde{b},\tilde{c},\tilde{d}}(T_1)^{-1}\phi_{\tilde{a},\tilde{b},\tilde{c},\tilde{d}}(T_0)\phi_{\tilde{a},\tilde{b},\tilde{c},\tilde{d}}(T_1) = \check{V}_1^{-1}\check{V}_0\check{V}_1. \end{array}$$

Iteration of (3.4) and (3.5) gives

$$\begin{aligned} \sigma^2: \quad H(u_1, k_1, k_0, u_0) &\rightarrow H(u_1, k_1, k_0, u_0) \\ (\check{V}_1, V_1, V_0, \check{V}_0) &\mapsto \check{V}_1^{-1}(\check{V}_1, V_1, V_0, \check{V}_0)\check{V}_1, \end{aligned} \quad (3.6)$$

or equivalently:

$$\begin{aligned} \sigma^2: \quad \mathcal{H}(a, b, c, d) &\rightarrow \mathcal{H}(a, b, c, d) \\ (T_0, T_1, Z) &\mapsto T_1^{-1}(T_0, T_1, Z)T_1. \end{aligned} \quad (3.7)$$

Remark 3.4. By (2.15) $u_1 u_0 = \sqrt{q^{-1}abcd} = \tilde{a}$, where \tilde{a} is the dual parameter given by (3.2). So $\phi(T_1 T_0) = \tilde{a} \check{V}_1 \check{V}_0$. So the element Y in [18, §2.15 and Theorem 2.22(2)] and the element Y_1 in case $n = 1$ of [25, (2.7)] can be identified with our $\tilde{a}^{-1} Y$. By (3.5) we have

$$\sigma(\tilde{a}^{-1} Y) = \tilde{a}^{-1} \sigma(T_1 T_0) = Z^{-1}, \quad (3.8)$$

compare with [25, Cor. 2.7].

3.4 The flip $(a, b, c, d) \mapsto (a, b, c, qd^{-1})$ and the automorphism τ

Let τ be the DAHA automorphism given on $H(u_1, k_1, k_0, u_0)$ by

$$\tau: (\check{V}_1, V_1, V_0, \check{V}_0) \mapsto (\check{V}_1, V_1, V_0 \check{V}_0 V_0^{-1}, V_0): H(u_1, k_1, k_0, u_0) \rightarrow H(u_1, k_1, u_0, k_0), \quad (3.9)$$

or equivalently on $\mathcal{H}(a, b, c, d)$ by

$$\tau: (T_0, T_1, Z) \mapsto (q^{-1} c Z T_0^{-1}, T_1, Z): \mathcal{H}(a, b, c, d) \rightarrow \mathcal{H}(a, b, c, qd^{-1}). \quad (3.10)$$

Then

$$\tau^{-1}: (T_0, T_1, Z) \mapsto (q^{-1} c T_0^{-1} Z, T_1, Z): \mathcal{H}(a, b, c, d) \rightarrow \mathcal{H}(a, b, c, qd^{-1}). \quad (3.11)$$

In combination with the data for t_3 and t_4 in Table 1 we see that

$$\begin{aligned} t_4 = \tau t_3 \tau^{-1}: T_0 &\mapsto q^{-1} c T_0^{-1} Z \mapsto q^{-1} c T_0^{-1} Z \mapsto q^{-1} c d T_0 \\ &: \mathcal{H}(a, b, c, d) \rightarrow \mathcal{H}(a, b, c, qd^{-1}) \rightarrow \mathcal{H}(a, b, qd^{-1}, c) \rightarrow \mathcal{H}(a, b, qd^{-1}, qc^{-1}). \end{aligned} \quad (3.12)$$

We also have

$$\tau^2: (T_0, T_1, Z) \mapsto (Z T_0 Z^{-1}, T_1, Z): \mathcal{H}(a, b, c, d) \rightarrow \mathcal{H}(a, b, c, d). \quad (3.13)$$

From (3.4), (3.9) or (3.5), (3.10) we obtain

$$\begin{aligned} (\sigma\tau)^3: H(u_1, k_1, k_0, u_0) &\rightarrow H(u_1, k_1, k_0, u_0) \\ (\check{V}_1, V_1, V_0, \check{V}_0) &\mapsto \check{V}_1^{-2} (\check{V}_1, V_1, V_0, \check{V}_0) \check{V}_1^2, \end{aligned} \quad (3.14)$$

or equivalently:

$$\begin{aligned} (\sigma\tau)^3: \mathcal{H}(a, b, c, d) &\rightarrow \mathcal{H}(a, b, c, d) \\ (T_0, T_1, Z) &\mapsto T_1^{-2} (T_0, T_1, Z) T_1^2. \end{aligned} \quad (3.15)$$

3.5 Extended modular group action on the DAHA

Clearly σ and τ in their action on u_1, k_1, k_0, u_0 parameter space generate the group S_3 of permutations of k_1, k_0, u_0 . Following Oblomkov [19, §2] we observe from (3.6) and (3.14) that σ and τ generate an action on the DAHA of a group which is up to global conjugation the modular group $\mathbb{P}SL_2(\mathbb{Z})$ presented as $\langle \sigma, \tau \mid \sigma^2 = (\sigma\tau)^3 = 1 \rangle$. If, again following Oblomkov, we also consider a further automorphism

$$\begin{aligned} \eta: (\check{V}_1, V_1, V_0, \check{V}_0) &\mapsto \left(-\check{V}_1^{-1}, -\check{V}_1 V_1^{-1} \check{V}_1^{-1}, -\check{V}_1 V_1 V_0^{-1} V_1^{-1} \check{V}_1^{-1}, -\check{V}_0^{-1} \right) \\ (u_1, k_1, k_0, u_0; q) &\mapsto (-u_1^{-1}, -k_1^{-1}, -k_0^{-1}, -u_0^{-1}; q^{-1}) \end{aligned} \quad (3.16)$$

or equivalently

$$\begin{aligned}\eta: (T_0, T_1, Z) &\mapsto (T_0^{-1}, T_1^{-1}, Z^{-1}) \\ (a, b, c, d; q) &\mapsto (a^{-1}, b^{-1}, c^{-1}, d^{-1}; q^{-1})\end{aligned}\tag{3.17}$$

(note that η does not leave q invariant) then we observe further relations $\eta^2 = (\sigma\eta)^2 = (\tau\eta)^2 = 1$ (exactly, not up to conjugation). So we are dealing with an action on the DAHA of a group which is up to global conjugation the *extended modular group* $\bar{\Gamma} := \mathbb{P}GL_2(\mathbb{Z})$ with presentation

$$\langle \sigma, \tau, \eta \mid \sigma^2 = (\sigma\tau)^3 = \eta^2 = (\sigma\eta)^2 = (\tau\eta)^2 = 1 \rangle.$$

Note that the action of $\bar{\Gamma}$ on parameter space is simply isomorphic with $S_3 \times \mathbb{Z}_2$.

Also note that, by (2.6) and (3.17),

$$\eta(Y) = T_1^{-1}Y^{-1}T_1, \quad \eta(D) = q(abcd)^{-1}D, \quad \eta(X) = X.\tag{3.18}$$

Remark 3.5. Notice that we have modified the definition of σ and η w.r.t. to Oblomkov's paper, let us denote the latter by $\hat{\sigma}$ and $\hat{\eta}$ respectively. The relation between the two is in terms the cyclic permutations

$$\begin{aligned}\gamma: (\check{V}_1, V_1, V_0, \check{V}_0) &:= (\check{V}_0, \check{V}_1, V_1, V_0) \\ (u_1, k_1, k_0, u_0) &\mapsto (u_0, u_1, k_1, k_0),\end{aligned}$$

and global conjugations:

$$\gamma\sigma = \hat{\sigma}\gamma, \quad \hat{\eta} = -V_1^{-1}\check{V}_1^{-1}\eta\check{V}_1V_1,$$

where the minus sign means that one has to take the minus in front of all resulting parameters and generators. The reason to define σ as in (3.4) is that it coincides with the duality operator. Moreover Oblomkov uses a generator $\phi = \sigma^{-1}\tau$ to present $\bar{\Gamma}$.

3.6 Spherical braid group

In [17], it was shown that the DAHA of type \check{C}_1C_1 appears naturally as quantisation of the group algebra of the monodromy group associated to the sixth Painlevé equation. In [3] it was shown that the analytic continuation of the solutions of the sixth Painlevé equation is described by a natural action of the spherical braid group $B_3 = \langle \beta_1^{\pm 1}, \beta_2^{\pm 1} \mid \beta_1\beta_2\beta_1 = \beta_2\beta_1\beta_2, (\beta_1\beta_2)^3 = 1 \rangle$. By quantisation, assigning $M_0 \mapsto \check{V}_0$, $M_t \mapsto V_0$, $M_1 \mapsto V_1$, this action can be brought to the DAHA as follows:

$$\begin{aligned}\beta_1: (\check{V}_1, V_1, V_0, \check{V}_0) &\mapsto (\check{V}_1, V_1, V_0\check{V}_0V_0^{-1}, V_0) \\ (u_1, k_1, k_0, u_0; q) &\mapsto (u_1, k_1, u_0, k_0; q)\end{aligned}$$

and

$$\begin{aligned}\beta_2: (\check{V}_1, V_1, V_0, \check{V}_0) &\mapsto (\check{V}_1, V_1V_0V_1^{-1}, V_1, \check{V}_0) \\ (u_1, k_1, k_0, u_0; q) &\mapsto (u_1, k_0, k_1, u_0; q).\end{aligned}$$

It is a straightforward computation to check the braid relation $\beta_1\beta_2\beta_1 = \beta_2\beta_1\beta_2$ and that, up to a global conjugation by \check{V}_1 , the Coxeter element $(\beta_1\beta_2)^3$ acts like the identity. The inversion η given by (3.16) was studied also in [15] as an extension of the braid group.

Note that in the formulae for β_1 and β_2 the element \check{V}_1 is always preserved, and η keeps its position (similarly both σ and τ preserve \hat{V}_1). This is because in fact the action of B_3 and $\bar{\Gamma}$ is natural on three elements.

We can express the modular group generators in terms of braids as follows

$$\sigma(\check{V}_1, V_1, V_0, \check{V}_0) := \beta_1 \beta_2 \beta_1(\check{V}_1, V_1, V_0, \check{V}_0),$$

$$\tau(\check{V}_1, V_1, V_0, \check{V}_0) := \beta_1(\check{V}_1, V_1, V_0, \check{V}_0).$$

Thanks to the fact that $\beta_1 = \tau$, the action of β_1 on $\mathcal{H}(a, b, c, d)$ is given by formula (3.10), while the action of β_2 can be easily derived using formulae (2.16) and (2.17):

$$\begin{aligned} \beta_2 : \quad (T_0, T_1, Z) &\mapsto (T_0, T_1, -q\sqrt{abc^{-1}d^{-1}}T_1^{-1}Z^{-1}T_0) \\ (a, b, c, d; q) &\mapsto (-\sqrt{q}\tilde{a}/c, -\sqrt{q}\tilde{a}/d, -\sqrt{q}\tilde{a}/a, -\sqrt{q}\tilde{a}/b; q). \end{aligned}$$

4 The basic DAHA representation and DAHA automorphisms acting on it

4.1 The basic DAHA representation

The *basic* (or polynomial) *representation* $\pi_{a,b,c,d}: U \mapsto \hat{U}$ [23], [18], [10, (3.10)–(3.12)], [14, §5.3] of $\mathcal{H}(a, b, c, d)$ on the space $\mathcal{A} := \mathbb{C}[z^{\pm 1}]$ of Laurent polynomials sends the generators Z, T_1, T_0 to operators $\hat{Z}, \hat{T}_1 = \hat{T}_1(a, b), \hat{T}_0 = \hat{T}_0(c, d)$ acting on \mathcal{A} as

$$(\hat{Z}f)(z) := z f(z), \tag{4.1}$$

$$\begin{aligned} (\hat{T}_1 f)(z) &:= \frac{(a+b)z - (1+ab)}{1-z^2} f(z) + \frac{(1-az)(1-bz)}{1-z^2} f(z^{-1}) \\ &= -abf(z) + z^{-1}(1-az)(1-bz) \frac{f(z) - f(z^{-1})}{z - z^{-1}}, \end{aligned} \tag{4.2}$$

$$\begin{aligned} (\hat{T}_0 f)(z) &:= \frac{(q^{-1}cd + 1)z^2 - (c+d)z}{q - z^2} f(z) - \frac{(c-z)(d-z)}{q - z^2} f(qz^{-1}) \\ &= -q^{-1}cdf(z) - z^{-1}(c-z)(d-z) \frac{f(z) - f(qz^{-1})}{z - qz^{-1}}. \end{aligned} \tag{4.3}$$

This representation is faithful. Clearly the expressions (4.1)–(4.3) are invariant under t_2 (exchange of a, b) and t_3 (exchange of c, d). We will discuss their behaviour under t_4 in the next subsection. The basic representation extends to the space \mathcal{O} of analytic functions on \mathbb{C}^* .

Further function spaces on which the basic representation is acting can be obtained as cyclic DAHA modules generated by a Gaussian or inverse Gaussian. In this context we define the *Gaussian* [25, Def. 2.9] by

$$G_e(z) := \frac{1}{(ez, ez^{-1}; q)_\infty}, \quad e, z \in \mathbb{C}^*, \quad z^{\pm 1} \notin eq^{\mathbb{Z}_{\geq 0}}. \tag{4.4}$$

From (4.2) we see that \hat{T}_1 commutes with $h(\hat{Z})$ for any function h if $h(z) = h(z^{-1})$ (compare also with Corollary 2.3). So \hat{T}_1 commutes with $G_e(\hat{Z})$.

Following [25, Prop. 2.11] we now compute the conjugation of \hat{T}_0 by suitable Gaussians.

Lemma 4.1. *We have*

$$G_d(\hat{Z})\hat{T}_0(c, d)G_d(\hat{Z})^{-1} = q^{-1}c\hat{Z}\hat{T}_0(c, qd^{-1})^{-1}, \quad (4.5)$$

$$G_{qd^{-1}}(\hat{Z})^{-1}\hat{T}_0(c, d)G_{qd^{-1}}(\hat{Z}) = q^{-1}c\hat{T}_0(c, qd^{-1})^{-1}\hat{Z}. \quad (4.6)$$

Proof. Note that

$$\frac{G_d(z)}{G_d(qz^{-1})} = \frac{1 - q^{-1}dz}{1 - dz^{-1}} \quad \text{and} \quad \frac{G_{qd^{-1}}(qz^{-1})}{G_{qd^{-1}}(z)} = \frac{q}{z^2} \frac{1 - q^{-1}dz}{1 - dz^{-1}}.$$

Then use (4.3) and (2.5). □

Since $G_d(\hat{Z})$ and $G_{qd^{-1}}(\hat{Z})$ commute with \hat{Z} and \hat{T}_1 , we conclude:

Corollary 4.2. *If the basic representation acts on a module \mathcal{M} consisting of functions on \mathbb{C}^* (like \mathcal{A} or \mathcal{O}) then it acts also on $\mathcal{M}G_d^{-1}$ and $\mathcal{M}G_{qd^{-1}}$.*

Remark 4.3. From (3.10), (3.11), (4.5) and (4.6) we see how the automorphisms τ and τ^{-1} act on the basic representation:

$$G_d(\hat{Z})\pi_{a,b,c,d}(U)G_d(\hat{Z})^{-1} = \pi_{a,b,c,qd^{-1}}(\tau(U)), \quad (4.7)$$

$$G_{qd^{-1}}(\hat{Z})^{-1}\pi_{a,b,c,d}(U)G_{qd^{-1}}(\hat{Z}) = \pi_{a,b,c,qd^{-1}}(\tau^{-1}(U)), \quad (4.8)$$

where $U \in \mathcal{H}(a, b, c, d)$ in both cases. Formula (4.7) is the case $n = 1$ of [25, Prop. 2.11].

4.2 Action of t_4 on the basic representation

Lemma 4.4. *Given an element $U \in \mathcal{H}(a, b, c, d)$, we have*

$$\frac{G_c(\hat{Z})}{G_{qd^{-1}}(\hat{Z})} \pi_{a,b,c,d}(U) \frac{G_{qd^{-1}}(\hat{Z})}{G_c(\hat{Z})} = \pi_{a,b,c,qd^{-1}}(t_4(U)). \quad (4.9)$$

In particular,

$$\frac{G_c(\hat{Z})}{G_{qd^{-1}}(\hat{Z})} \hat{T}_0(c, d) \frac{G_{qd^{-1}}(\hat{Z})}{G_c(\hat{Z})} = q^{-1}cd\hat{T}_0(qd^{-1}, qc^{-1}). \quad (4.10)$$

Hence also

$$\frac{G_c(\hat{Z})}{G_{qd^{-1}}(\hat{Z})} \hat{Y}(a, b, c, d) \frac{G_{qd^{-1}}(\hat{Z})}{G_c(\hat{Z})} = q^{-1}cd\hat{Y}(a, b, qd^{-1}, qc^{-1}), \quad (4.11)$$

$$\frac{G_c(\hat{Z})}{G_{qd^{-1}}(\hat{Z})} \hat{D}(a, b, c, d) \frac{G_{qd^{-1}}(\hat{Z})}{G_c(\hat{Z})} = q^{-1}cd\hat{D}(a, b, qd^{-1}, qc^{-1}). \quad (4.12)$$

Proof. For (4.9) apply (3.12) together with (4.7) and (4.8). Alternatively, (4.10) can be proved by direct computation using (4.3).

For (4.11) and (4.12) use (2.6). In particular, for (4.12) note that by (2.6) and (4.11) both sides of (4.12) are equal to $q^{-1}cd(\hat{Y}(a, b, qd^{-1}, qc^{-1}) + qabc^{-1}d^{-1}\hat{Y}(a, b, qd^{-1}, qc^{-1})^{-1})$. \square

As a corollary of (4.11) we obtain:

Proposition 4.5. *Let $\phi \in \mathcal{O}G_{qd^{-1}}$ be an eigenfunction of $\hat{Y}(a, b, c, d)$ with eigenvalue λ , then $\frac{G_c}{G_{qd^{-1}}}\phi \in \mathcal{O}G_c$ is an eigenfunction of $\hat{Y}(a, b, qd^{-1}, qc^{-1})$ with eigenvalue $q(cd)^{-1}\lambda$.*

If we have, for (a, b, c, d, λ) in a certain region of \mathbb{C}^5 , a family of eigenfunctions $\phi_\lambda^{(a, b, c, d)}$ of $\hat{Y}(a, b, c, d)$ with $\phi_\lambda^{(a, b, c, d)}$ analytic in (a, b, c, d, λ) for generic values of (a, b, c, d) then it is not yet guaranteed by Proposition 4.5 that

$$\frac{G_c}{G_{qd^{-1}}}\phi_\lambda^{(a, b, c, d)} = \text{canst.} \phi_{q(cd)^{-1}\lambda}^{(a, b, qd^{-1}, qc^{-1})}.$$

In (7.27) we will have an instance that such an equality holds.

4.3 Action of σ on the basic representation

Different from what we saw for τ and t_4 , the action of σ on the basic representation cannot simply be given by an elementary conjugation. Following Stokman [25, (3.1)] we see that having such an action of σ amounts to giving a corresponding Fourier transform, called *difference Fourier transform*.

Definition 4.6. Let V and W_σ be suitable function spaces on \mathbb{C}^* on which $\pi_{a, b, c, d}$ and $\pi_{\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}}$, respectively, act. A linear map $F: V \rightarrow W_\sigma$ is called a *difference Fourier transform* if

$$F(\pi_{a, b, c, d}(U)f) = \pi_{\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}}(\sigma(U))(F(f)) \quad (4.13)$$

for all $f \in V$ and all $U \in \mathcal{H}(a, b, c, d)$.

In Stokman's paper this is defined for general rank, but in [25, §8] it is worked out in detail in the rank one case. There F is an integral transform with integral kernel given by the Cherednik kernel (7.28) or its symmetric form (6.5).

4.4 The spherical subalgebra and the Zhedanov algebra

In this subsection let $\mathcal{M} = \mathcal{A}G$ or $\mathcal{O}G$ with $G = 1$ or $G_{qd^{-1}}$ or G_d^{-1} . We call $f \in \mathcal{M}$ *symmetric* if $f(z) = f(z^{-1})$ and we notate \mathcal{M}^+ for the subspace of symmetric functions in \mathcal{M} .

Lemma 4.7. *For any $g \in \mathcal{M}$*

$$(\hat{T}_1 g)(z) = -abg(z) \iff g \in \mathcal{M}^+ \quad (4.14)$$

$$(\hat{T}_1 g)(z) = -g(z) \iff g(z) = \frac{(1-az)(1-bz)}{z} h(z) \text{ with } h \in \mathcal{M}^+. \quad (4.15)$$

Proof. Observe that, for $g \in \mathcal{M}$,

$$((\hat{T}_1 + ab)g)(z) = \frac{(1-az)(1-bz)}{1-z^2} (g(z^{-1}) - g(z)).$$

This immediately implies (4.14). We can also write

$$((\hat{T}_1 + 1)g)(z) = \frac{(1-az)(1-bz)}{1-z^2} g(z^{-1}) - \frac{(a-z)(b-z)}{1-z^2} g(z). \quad (4.16)$$

If $g(z) = z^{-1}(1-az)(1-bz)h(z)$ with $h \in \mathcal{M}^+$ then the right-hand side of (4.16) is 0. Vice-versa, for $a \neq a^{-1}$ and $b \neq b^{-1}$, if the right-hand side of (4.16) is identically 0, then $g(a^{-1}) = 0$ and $g(b^{-1}) = 0$. Hence we can write $g(z) = z^{-1}(1-az)(1-bz)h(z)$ with $h \in \mathcal{M}$. Substituting this in

$$\frac{(1-az)(1-bz)}{1-z^2} g(z^{-1}) - \frac{(a-z)(b-z)}{1-z^2} g(z) \equiv 0$$

yields that $h(z) = h(z^{-1})$, which proves (4.15). \square

Corollary 4.8. *Given any $g \in \mathcal{M}$, there exist two elements $g_1, g_2 \in \mathcal{M}^+$ such that*

$$g(z) = g_1(z) - \frac{(1-az)(1-bz)}{z} g_2(z). \quad (4.17)$$

Proof. We define

$$g_1 = \frac{1}{1-ab} (\hat{T}_1 g + g), \quad (4.18)$$

$$g_2 = \frac{z}{(1-az)(1-bz)} \frac{1}{1-ab} (\hat{T}_1 g + abg), \quad (4.19)$$

which gives (4.17). To prove that g_1 and g_2 are symmetric, it is enough to show that $\hat{T}_1 g_1 = -abg_1$, and $\hat{T}_1 h_2 = -h_2$ with $h_2(z) = \frac{(1-az)(1-bz)}{z} g_2(z)$; both relations follow immediately from (2.1). \square

Let $e := (1-ab)^{-1}(T_1 + 1)$. By (2.1) we have $e^2 = e$.

Definition 4.9. The *spherical subalgebra* of $\mathcal{H}(a, b, c, d)$ is the image $\mathcal{SH}(a, b, c, d)$ of the projection operator $U \mapsto eUe$.

Proposition 4.10.

(a) $\mathcal{SH}(a, b, c, d)$ is generated as an algebra by its elements De and Xe .

(b) In the basic representation on \mathcal{M} the subspace \mathcal{M}^+ of symmetric functions is invariant under the action of $\mathcal{SH}(a, b, c, d)$. The corresponding representation is faithful.

(c) For $f \in \mathcal{M}^+$ we have

$$\pi_{a,b,c,d}(De)f = \hat{D}(a, b, c, d)f = L^{a,b,c,d}f, \quad (4.20)$$

where $L = L^{a,b,c,d}$ is the Askey–Wilson operator

$$(Lf)(z) = L_z f(z) := (1 + q^{-1}abcd)f(z) + \frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)}(f(qz) - f(z)) \\ + \frac{(a-z)(b-z)(c-z)(d-z)}{(1-z^2)(q-z^2)}(f(q^{-1}z) - f(z)), \quad f(z) = f(z^{-1}). \quad (4.21)$$

Also

$$(\pi_{a,b,c,d}(Xe)f)(z) = (\hat{X}f)(z) = (z + z^{-1})f(z).$$

(d) The algebra generated by the operators $L^{a,b,c,d}$ and \hat{X} acting on \mathcal{M}^+ is isomorphic with the quotient $AW(3, Q_0)$ of the Zhedanov algebra $AW(3)$ (as defined in [10, §2]) with the ideal generated by $Q - Q_0$ (see [10, (2,3), (2,8)]; here Q is the Casimir element). The generators K_0 and K_1 of $AW(3, Q_0)$ correspond under this isomorphism with $L^{a,b,c,d}$ and \hat{X} , respectively.

For the proof of parts (a), (b) and (c) see [11, §3]. For the proof of part (d) see [10, §2]. See also [29].

4.5 Action of $W(D_4)$ on the spherical subalgebra $\mathcal{SH}(a, b, c, d)$

In Remark 3.2, we observed that \hat{t}_0, t_2, t_3, t_4 , acting on (a, b, c, d) parameter space, generate $W(D_4)$. The action of these automorphisms on D and T_1 in the basic representation induces an action on \hat{D} . However, while t_2, t_3, t_4 preserve T_1 and therefore automatically restrict to the spherical subalgebra, \hat{t}_0 doesn't preserve T_1 and therefore its restriction to the spherical subalgebra is more complicated and it will be explored in [P2].

From the expression for L in (4.21) we see that L is invariant under permutations of a, b, c, d while acting on the space \mathcal{A}^+ of symmetric Laurent polynomials and that it is only invariant under permutations of a, b, c while acting on $\mathcal{O}^+G_{qd^{-1}}$. Then t_4 also induces an action on L . Indeed, by Lemma 4.4 and (4.20) we have

$$\frac{G_c(\hat{Z})}{G_{qd^{-1}}(\hat{Z})} L^{a,b,c,d} \frac{G_{qd^{-1}}(\hat{Z})}{G_c(\hat{Z})} = q^{-1}cd \hat{L}^{a,b,qd^{-1},qc^{-1}}. \quad (4.22)$$

Remark 4.11. On L considered as a formal operator the permutations of a, b, c, d together with t_4 generate a symmetry group which is isomorphic with $W(D_4)$, see [12, §2.3]. This symmetry group can also be identified with the $W(D_4)$ action discussed in [2, §4], as we will show in our upcoming paper [P2].

As a corollary to (4.22) we obtain:

Proposition 4.12. Let $\phi \in \mathcal{O}^+G_{qd^{-1}}$ be an eigenfunction of $L^{a,b,c,d}$ with eigenvalue λ , then $\frac{G_c}{G_{qd^{-1}}} \phi \in \mathcal{O}^+G_c$ is an eigenfunction of $L^{a,b,qd^{-1},qc^{-1}}$ with eigenvalue $q(cd)^{-1}\lambda$.

This Proposition is equivalent to Suslov's observation in [26, §6.3, item 2], already met just after (5.24).

From Proposition 4.5 it follows that if for (a, b, c, d, λ) in a certain region of \mathbb{C}^5 , one has a family of eigenfunctions $\phi_\lambda^{(a,b,c,d)}$ of $L^{a,b,c,d}$ with $\phi_\lambda^{(a,b,c,d)}$ analytic in (a, b, c, d, λ) for generic values of (a, b, c, d) then $\frac{G_c}{G_{qd^{-1}}}\phi_\lambda^{(a,b,c,d)}$ defines a family of eigenfunctions of $L^{(a,b,qd^{-1},qc^{-1})}$. However, this doesn't guarantee that we may have a relation of the following form

$$\frac{G_c}{G_{qd^{-1}}}\phi_\lambda^{(a,b,c,d)} = \text{const.} \phi_{q(cd)^{-1}\lambda}^{(a,b,qd^{-1},qc^{-1})}.$$

In (6.1) we will show an instance that such an equality holds.

5 Some special functions

5.1 Askey–Wilson polynomials

Denote and renormalize *Askey–Wilson polynomials* (or briefly *AW polynomials*) as

$$R_n(z) = R_n(z; a, b, c, d) := {}_4\phi_3\left(\begin{matrix} q^{-n}, q^{n-1}abcd, az, az^{-1} \\ ab, ac, ad \end{matrix}; q, q\right). \quad (5.1)$$

This is related to the usual notation and normalization [1, (7.5.2)] by

$$R_n(z; a, b, c, d) = \frac{a^n}{(ab, ac, ad; q)_n} p_n\left(\frac{1}{2}(z + z^{-1}); a, b, c, d | q\right). \quad (5.2)$$

By (5.2) and (5.1) p_n has the expansion

$$p_n\left(\frac{1}{2}(z + z^{-1}); a, b, c, d | q\right) = a^{-n} \sum_{k=0}^n (abq^k, acq^k, adq^k; q)_{n-k} \frac{(q^{-n}, q^{n-1}abcd, az, az^{-1}; q)_k}{(q; q)_k} q^k. \quad (5.3)$$

Although not completely visible from (5.3), p_n is symmetric in a, b, c, d [1, §7.5]. The nontrivial symmetry in a and b follows from Sears' transformation [1, (III.15)]. It can be written for $R_n(z)$ as

$$R_n(z; a, b, c, d) = \left(\frac{a}{b}\right)^n \frac{(bc, bd; q)_n}{(ac, ad; q)_n} R_n(z; b, a, c, d). \quad (5.4)$$

We will also need the (in z) *monic* AW polynomial

$$P_n(z) = P_n(z; a, b, c, d) := \frac{(ab, ac, ad; q)_n}{(q^{n-1}abcd; q)_n a^n} R_n(z; a, b, c, d) = \frac{p_n\left(\frac{1}{2}(z + z^{-1}); a, b, c, d | q\right)}{(q^{n-1}abcd; q)_n}, \quad (5.5)$$

which is also symmetric in a, b, c, d .

Under suitable constraints on the parameters the AW polynomials are orthogonal polynomials with explicit orthogonality relations [1]. They are eigenfunctions of the Askey–Wilson operator (AW operator) $L = L^{a,b,c,d}$ given by (4.21):

$$(LR_n)(z) = (q^{-n} + abcdq^{n-1})R_n(z). \quad (5.6)$$

It is worth remembering that the operator $L^{a,b,c,d}$ is the restriction of the operator $\hat{D}(a,b,c,d)$ (in the basic representation) to the symmetric functions in the representation space.

From (5.1) we see the *duality* for Askey–Wilson polynomials:

$$R_n(a^{-1}q^{-m}; a, b, c, d) = R_m(\tilde{a}^{-1}q^{-n}; \tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}), \quad m, n \in \mathbb{Z}_{\geq 0}. \quad (5.7)$$

5.2 Askey–Wilson functions

We will use the notation [4, (2.1.11)] for *very-well-poised series*, and we will further abbreviate this notation in the case of *very-well-poised-balanced series* by

$$\begin{aligned} {}_8W_7(a; b, c, d, e, f) &= {}_8W_7\left(a; b, c, d, e, f; q, \frac{q^2 a^2}{bcdef}\right) \\ &:= {}_8\phi_7\left(\begin{matrix} a, qa^{\frac{1}{2}}, -qa^{\frac{1}{2}}, b, c, d, e, f \\ a^{\frac{1}{2}}, -a^{\frac{1}{2}}, qa/b, qa/c, qa/d, qa/e, qa/f \end{matrix}; q, \frac{q^2 a^2}{bcdef}\right), \quad \left|\frac{q^2 a^2}{bcdef}\right| < 1. \end{aligned} \quad (5.8)$$

Note that this is symmetric in b, c, d, e, f .

Koelink and Stokman [9, (3.2)] define the *Askey–Wilson function* (or briefly *AW function*) ϕ_γ as follows.

$$\begin{aligned} \phi_\gamma(z) = \phi_\gamma(z; a, b, c, d) = \phi_\gamma(z; a, b, c, d; q) &:= \frac{(q\tilde{a}d^{-1}\gamma z, q\tilde{a}d^{-1}\gamma z^{-1}; q)_\infty}{(bc, q\gamma\tilde{a}ad^{-1}, \gamma\tilde{a}^{-1}bc, qd^{-1}z, qd^{-1}z^{-1}; q)_\infty} \\ &\times {}_8W_7(\tilde{a}ad^{-1}\gamma; az, az^{-1}, \tilde{a}\gamma, \tilde{b}\gamma, \tilde{c}\gamma) \quad |\gamma| > |q\tilde{d}^{-1}|. \end{aligned} \quad (5.9)$$

They use it as a kernel in a Fourier type integral transform, called the Askey–Wilson function transform, for which they give an inversion formula.

This function was earlier studied by Suslov [26, (6.18)], [27, (2.7)], [28, (2.7)] (notations and normalizations vary). Another representation of $\phi_\gamma(z)$ can be obtained from (5.9) by applying [4, (III.23)] (there replace $a; b, c, d, e, f$ by $\tilde{a}ad^{-1}\gamma; \tilde{b}\gamma, \tilde{c}\gamma, az, az^{-1}, \tilde{a}\gamma$):

$$\begin{aligned} \phi_\gamma(z; a, b, c, d) &= \frac{(qa\tilde{d}^{-1}\gamma^{-1}z^{-1}, qa\tilde{d}^{-1}\gamma z^{-1}; q)_\infty}{(\tilde{b}\tilde{c}, q\tilde{a}a\tilde{d}^{-1}z^{-1}, a^{-1}\tilde{b}\tilde{c}z^{-1}, q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1}; q)_\infty} \\ &\times {}_8W_7(\tilde{a}a\tilde{d}^{-1}z^{-1}; \tilde{a}\gamma, \tilde{a}\gamma^{-1}, az^{-1}, \tilde{b}z^{-1}, cz^{-1}), \quad |z| < |q^{-1}d|. \end{aligned} \quad (5.10)$$

The representation (5.10) of ϕ_γ earlier occurred in papers by Ismail & Rahman [7, (1.12)] and by Rahman [21, (1.32), (1.35)], [22, (1.19)] (again with varying notations and normalizations), see also Haine & Iliev [5, (2.4), (2.8a)]. Suslov [28, (2.7), (2.10)] gave both representations (5.9) and (5.10).

The combination of (5.9) and (5.10) states the *duality*

$$\phi_\gamma(z; a, b, c, d) = \phi_{z^{-1}}(\gamma^{-1}; \tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}). \quad (5.11)$$

By (5.9) and (5.10) $\phi_\gamma(z)$ is symmetric in z and γ :

$$\phi_\gamma(z) = \phi_\gamma(z^{-1}), \quad \phi_\gamma(z) = \phi_{\gamma^{-1}}(z).$$

Stokman [24, (3.1)] defines ϕ_γ with a normalization different from (5.9); we denote it here by ϕ_γ^S :

$$\phi_\gamma^S := \frac{(qabcd^{-1}; q)_\infty}{(qbd^{-1}, qcd^{-1}; q)_\infty} \phi_\gamma. \quad (5.12)$$

In [25, Dfn. 6.12, Thm. 6.20] Stokman uses yet another notation and normalization:

$$\mathfrak{E}^+(\gamma; z) = \mathfrak{E}^+(\gamma; z; a, b, c, d) := (bc, qad^{-1}, qa^{-1}d^{-1}; q)_\infty \phi_\gamma(z; a, b, c, d) \quad (5.13)$$

$$= \frac{(q\gamma\tilde{a}d^{-1}z, q\gamma\tilde{a}d^{-1}z^{-1}, qad^{-1}, qa^{-1}d^{-1}; q)_\infty}{(q\gamma\tilde{a}ad^{-1}, \gamma\tilde{a}^{-1}bc, qd^{-1}z, qd^{-1}z^{-1}; q)_\infty} {}_8W_7(\gamma\tilde{a}ad^{-1}; az, az^{-1}, \gamma\tilde{a}, \gamma ab\tilde{a}^{-1}, \gamma ac\tilde{a}^{-1}) \quad (5.14)$$

From (5.14) we observe the normalization

$$\mathfrak{E}^+(\gamma; a^{\pm 1}; a, b, c, d) = 1. \quad (5.15)$$

Again we have symmetry in z and γ ,

$$\mathfrak{E}^+(\gamma; z) = \mathfrak{E}^+(\gamma; z^{-1}), \quad \mathfrak{E}^+(\gamma; z) = \mathfrak{E}^+(\gamma^{-1}; z),$$

and duality

$$\mathfrak{E}^+(\gamma; z; a, b, c, d) = \mathfrak{E}^+(z, \gamma; \tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}). \quad (5.16)$$

Hence, by (5.15) and (5.16),

$$\mathfrak{E}^+(\tilde{a}^{\pm 1}; z; a, b, c, d) = 1. \quad (5.17)$$

Remark 5.1. If we substitute $\gamma = \tilde{a}$ in (5.14) then, by (5.17) and (5.8),

$$1 = \frac{(abcz, abc z^{-1}, qad^{-1}, qa^{-1}d^{-1}; q)_\infty}{(a^2bc, bc, qd^{-1}z, qd^{-1}z^{-1}; q)_\infty} {}_6W_5(q^{-1}a^2bc; az, az^{-1}, q^{-1}abcd; q, qa^{-1}d^{-1}). \quad (5.18)$$

Formula (5.18) is equivalent with the evaluation formula [4, (II.20)].

By [4, (III.36)] we can write

$$\begin{aligned} & \mathfrak{E}^+(\gamma; z; a, b, c, d) \\ &= {}_4\phi_3 \left(\begin{matrix} az, az^{-1}, \tilde{a}\gamma, \tilde{a}\gamma^{-1} \\ ab, ac, ad \end{matrix}; q, q \right) + \frac{(az, az^{-1}, \tilde{a}\gamma, \tilde{a}\gamma^{-1}, qbd^{-1}, qcd^{-1}, qa^{-1}d^{-1}; q)_\infty}{(qd^{-1}z, qd^{-1}z^{-1}, q\gamma\tilde{a}a^{-1}d^{-1}, q\gamma^{-1}\tilde{a}a^{-1}d^{-1}, ab, ac, q^{-1}ad; q)_\infty} \\ & \quad \times {}_4\phi_3 \left(\begin{matrix} qd^{-1}z, qd^{-1}z^{-1}, q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1} \\ qbd^{-1}, qcd^{-1}, q^2a^{-1}d^{-1} \end{matrix}; q, q \right) \end{aligned} \quad (5.19)$$

Hence, as observed in [9, §3], $\mathfrak{E}^+(\gamma; z; a, b, c, d)$ extends for generic parameters a, b, c, d to a meromorphic function in z and γ . More precisely,

Proposition 5.2.

$$(qd^{-1}z, qd^{-1}z^{-1}, q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1}; q)_\infty \mathfrak{E}^+(\gamma; z; a, b, c, d)$$

is an entire complex analytic function in $z + z^{-1}$ and $\gamma + \gamma^{-1}$. Equivalently, $\mathfrak{E}^+(\gamma; z; a, b, c, d)$ belongs to $\mathcal{O}^+G_{qd^{-1}}$ as a function of z and to $\mathcal{O}^+G_{q\tilde{d}^{-1}}$ as a function of γ .

Here $1/(qd^{-1}z, qd^{-1}z^{-1}; q)_\infty$ and $1/(q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1}; q)_\infty$ are *Gaussians* $G_\tau(z)$ as considered in [25, Definition 2.9].

For $\gamma = q^n \tilde{a}$, $n = 0, 1, 2, \dots$, we get from (5.19) an Askey–Wilson polynomial (5.1):

$$\mathfrak{E}^+(q^n \tilde{a}; z; a, b, c, d) = {}_4\phi_3 \left(\begin{matrix} q^{-n}, q^{n-1}abcd, az, az^{-1} \\ ab, ac, ad \end{matrix}; q, q \right) = R_n(z; a, b, c, d). \quad (5.20)$$

Hence the eigenvalue equation (5.6) for Askey–Wilson polynomials can be rewritten as

$$\tilde{a}^{-1}L_z \mathfrak{E}^+(q^n \tilde{a}; z) = (q^n \tilde{a} + (q^n \tilde{a})^{-1}) \mathfrak{E}^+(q^n \tilde{a}; z), \quad n = 0, 1, 2, \dots, \quad (5.21)$$

As stated in [9, §3], this extends to an eigenvalue equation for $\mathfrak{E}^+(\gamma; z)$:

$$\tilde{a}^{-1}L_z \mathfrak{E}^+(\gamma; z) = (\gamma + \gamma^{-1}) \mathfrak{E}^+(\gamma; z). \quad (5.22)$$

Here L is still given by (4.21). Note that the first term on the right in (4.21) is missing in [9, (2.4)], so that it occurs there in the eigenvalue [9, (2.5)].

Remark 5.3. The eigenvalue equation (5.22) was probably first given by Suslov [26, §6.3]. He derived it by observing that both terms on the right-hand side of (5.19) are also eigenfunctions of $\tilde{a}^{-1}L_z$ with eigenvalue $\gamma + \gamma^{-1}$. Indeed, put

$$\mathfrak{R}(\gamma; z; a, b, c, d) := {}_4\phi_3 \left(\begin{matrix} az, az^{-1}, \tilde{a}\gamma, \tilde{a}\gamma^{-1} \\ ab, ac, ad \end{matrix}; q, q \right), \quad (5.23)$$

so $\mathfrak{R}(q^\nu \tilde{a}; z; a, b, c, d) = R_\nu(z; a, b, c, d)$ with R_ν given by (5.1) with n replaced by ν . Now (5.19) can be rewritten as

$$\begin{aligned} \mathfrak{E}^+(\gamma; z; a, b, c, d) &= \mathfrak{R}(\gamma; z; a, b, c, d) \\ &+ \frac{(\tilde{a}\gamma, \tilde{a}\gamma^{-1}, qbd^{-1}, qcd^{-1}, qa^{-1}d^{-1}; q)_\infty}{(q\gamma\tilde{a}a^{-1}d^{-1}, q\gamma^{-1}\tilde{a}a^{-1}d^{-1}, ab, ac, q^{-1}ad; q)_\infty} \frac{(az, az^{-1}; q)_\infty}{(qd^{-1}z, qd^{-1}z^{-1}; q)_\infty} \mathfrak{R}(\gamma; z; qd^{-1}, b, c, qa^{-1}). \end{aligned} \quad (5.24)$$

The eigenvalue equation for $\mathfrak{R}(\gamma; z; a, b, c, d)$ is proved in the same way as in the polynomial case. For the second term on the right in (5.24) Suslov [26, §6.3, item 2] observes that, if $f(z; a, b, c, d)$ is an eigenfunction of $L_z^{a,b,c,d}$ with eigenvalue λ then so is $\frac{(az, az^{-1}; q)_\infty}{(qd^{-1}z, qd^{-1}z^{-1}; q)_\infty} f(z; qd^{-1}, b, c, qa^{-1})$. We will come back to this symmetry of L_z in Proposition 4.12.

6 Action of DAHA automorphisms on Askey–Wilson polynomials and functions

As observed in §4.5, the AW operator $L^{a,b,c,d}$ is invariant under permutations of a, b, c, d , while it satisfies the symmetry (4.22) under the action of $t_4: (a, b, c, d) \mapsto (a, b, qd^{-1}, qc^{-1})$. These symmetries generate a group isomorphic to $W(D_4)$. In this section we study the possible action of these symmetries on the eigenfunctions of L . First we consider this for the AW polynomials, but there no satisfactory action of t_4 can be found. Then we consider it for the AW functions, where t_4 acts nicely but t_3 (interchanging c and d) fails.

6.1 An attempted action of t_4 on Askey–Wilson polynomials

Let us try to apply Proposition 4.12 to Askey–Wilson polynomials. By (5.6) the eigenfunction $R_n(z; a, b, c, d)$ of $L_z^{a,b,c,d}$ has eigenvalue $q^{-n} + abcdq^{n-1}$ and the eigenfunction $R_m(z; a, b, qd^{-1}, qc^{-1})$ of $L_z^{a,b,qd^{-1},qc^{-1}}$ has eigenvalue $q^{-m} + abc^{-1}d^{-1}q^{m+1}$. If $cd = q^{m-n+1}$ then the second eigenvalue is equal to $qc^{-1}d^{-1}$ times the first eigenvalue. Conversely, if $q^{-m} + abc^{-1}d^{-1}q^{m+1} = qc^{-1}d^{-1}(q^{-n} + abcdq^{n-1})$ for all $m, n = 0, 1, 2, \dots$ with $n - m$ fixed then $cd = q^{m-n+1}$. This shows, in view of Proposition 4.12, that $R_n(z; a, b, c, d)$ and $\frac{G_c(z)}{G_{qd^{-1}}(z)}R_m(z; a, b, qd^{-1}, qc^{-1})$ are related if and only if $cd = q^{m-n+1}$. Without loss of generality we assume that $n \geq m$. It is also convenient to work with the normalization and expansion (5.3) because of the symmetry in a, b, c, d . Then we find

$$p_n\left(\frac{1}{2}(z + z^{-1}); a, b, c, d \mid q\right) = (-c)^{-(n-m)} q^{-\frac{1}{2}(n-m)(n-m-1)} (q^m ab; q)_{n-m} \\ \times (cz, cz^{-1}; q)_{n-m} p_m\left(\frac{1}{2}(z + z^{-1}); a, b, qd^{-1}, qc^{-1} \mid q\right), \quad cd = q^{m-n+1}. \quad (6.1)$$

Note that for $cd = q^{m-n+1}$ indeed $\frac{G_c(z)}{G_{qd^{-1}}(z)} = (cz, cz^{-1}; q)_{n-m}$.

Proof of (6.1). From (5.3), the symmetry in a, b, c, d , the condition $cd = q^{m-n+1}$ and the vanishing for $k < n - m$ of $(q^{m-n+k+1}; q)_{n-k}$ we see that

$$p_n\left(\frac{1}{2}(z + z^{-1}); c, a, b, d \mid q\right) = c^{-n} \sum_{k=n-m}^n (acq^k, bcq^k, q^{m-n+k+1}; q)_{n-k} \frac{(q^{-n}, q^m ab, cz, cz^{-1}; q)_k}{(q; q)_k} q^k \\ = c^{-n} \sum_{j=0}^m (acq^{n-m+j}, bcq^{n-m+j}, q^{j+1}; q)_{m-j} \frac{(q^{-n}, q^m ab, cz, cz^{-1}; q)_{n-m+j}}{(q; q)_{n-m+j}} q^{n-m+j} \\ = (-c)^{-(n-m)} q^{-\frac{1}{2}(n-m)(n-m-1)} (q^m ab; q)_{n-m} (cz, cz^{-1}; q)_{n-m} q^{-m(n-m)} c^{-m} \\ \times \sum_{j=0}^m (acq^{n-m+j}, bcq^{n-m+j}, q^{n-m+j+1}; q)_{m-j} \frac{(q^{-m}, q^n ab, q^{n-m} cz, q^{n-m} cz^{-1}; q)_j}{(q; q)_j} q^j \\ = (-c)^{-(n-m)} q^{-\frac{1}{2}(n-m)(n-m-1)} (q^m ab; q)_{n-m} (cz, cz^{-1}; q)_{n-m} (qd^{-1})^{-m} \\ \times \sum_{j=0}^m (ad^{-1}q^{j+1}, bd^{-1}q^{j+1}, c^{-1}d^{-1}q^{j+2}; q)_{m-j} \frac{(q^{-m}, q^{m+1}abc^{-1}d^{-1}, qd^{-1}z, qd^{-1}z^{-1}; q)_j}{(q; q)_j} q^j \\ = (-c)^{-(n-m)} q^{-\frac{1}{2}(n-m)(n-m-1)} (q^m ab; q)_{n-m} (cz, cz^{-1}; q)_{n-m} \\ \times p_m\left(\frac{1}{2}(z + z^{-1}); qd^{-1}, a, b, qc^{-1} \mid q\right). \quad \square$$

As we just saw, if we want to realize the action of t_4 on Askey–Wilson polynomials, then we have to assume that cd is an integer power of q . With a continuous spectrum of L , as for Askey–Wilson functions as eigenfunctions, the condition on cd can be dropped, as we will see in the next subsection.

6.2 Action of t_4 and t_2 on Askey–Wilson functions

We will work with the Askey–Wilson function $\mathfrak{E}^+(\gamma; z; a, b, c, d)$ as defined by (5.14). Note that this function is symmetric in b and c (by the symmetry in b, c in (5.19) or by the symmetry in b, c, d, e in (5.8)).

We now turn to the action of t_4 on the Askey–Wilson formula, as formulated in the following theorem. There formula (6.2) is immediately seen from (5.14), again by the symmetry in b, c, d, e in (5.8).

Theorem 6.1. *Let $\mathfrak{E}^+(\gamma; z; a, b, c, d)$ be the Askey–Wilson function defined in (5.14), which is an eigenfunction of $L^{a,b,c,d}$ with eigenvalue $(\gamma + \gamma^{-1})\sqrt{q^{-1}abcd}$. Then*

$$\mathfrak{E}^+(\gamma; z; a, b, c, d) = \frac{(qad^{-1}, qa^{-1}d^{-1}; q)_\infty}{(ac, a^{-1}c; q)_\infty} \frac{G_c(z)}{G_{qd^{-1}}(z)} \mathfrak{E}^+(\gamma; z; a, b, qd^{-1}, qc^{-1}). \quad (6.2)$$

The function $\mathfrak{E}^+(\gamma; z; a, b, qd^{-1}, qc^{-1})$ is an eigenfunction of $L_z^{a,b,qd^{-1},qc^{-1}}$ with eigenvalue $(\gamma + \gamma^{-1})\sqrt{qabc^{-1}d^{-1}}$.

Consider [4, (III.23)] with the following substitutions:

$$a \rightarrow \gamma \tilde{a} a d^{-1}, \quad b \rightarrow a z, \quad c \rightarrow a z^{-1}, \quad d \rightarrow \gamma \tilde{a}^{-1} a c, \quad e \rightarrow \gamma \tilde{a}, \quad f \rightarrow \gamma \tilde{a}^{-1} a b.$$

Then

$$\begin{aligned} & {}_8W_7(\gamma \tilde{a} a d^{-1}; a z, a z^{-1}, \gamma \tilde{a}, \gamma a b \tilde{a}^{-1}, \gamma a c \tilde{a}^{-1}) \\ &= \frac{(q \gamma \tilde{a} a d^{-1}, q \gamma^{-1} \tilde{a} b^{-1} d^{-1}, q b d^{-1}, b c; q)_\infty}{(q a d^{-1}, a c, q \gamma \tilde{a} b d^{-1}, q \gamma^{-1} \tilde{a} a^{-1} d^{-1}; q)_\infty} {}_8W_7(\gamma \tilde{a} b d^{-1}; b z, b z^{-1}, \gamma \tilde{a}, \gamma a b \tilde{a}^{-1}, \gamma b c \tilde{a}^{-1}). \end{aligned}$$

Hence, in combination with (5.14), we have the following action of t_2 :

$$\mathfrak{E}^+(\gamma; z; a, b, c, d) = \frac{(bc, qa^{-1}d^{-1}; q)_\infty}{(ac, qb^{-1}d^{-1}; q)_\infty} \frac{G_{\tilde{c}}(\gamma)}{G_{q\tilde{d}^{-1}}(\gamma)} \mathfrak{E}^+(\gamma; z; b, a, c, d). \quad (6.3)$$

For $\gamma = q^n \tilde{a}$ (6.3) combined with (5.20) simplifies to the well-known (5.4).

So altogether we have seen that the AW functions have symmetries which come from the subgroup of $W(D_4)$ consisting of permutations of a, b, c, qd^{-1} . In the AW function transform [9, §5] the weight function $W(z) = W_{a,b,c,d,t}(z)$ is symmetric in a, b, c , while, by (6.2)

$$\begin{aligned} \mathfrak{E}^+(\gamma; z; a, b, c, d) W_{a,b,c,d,t}(z) &= \frac{(qad^{-1}, qa^{-1}d^{-1}; q)_\infty}{(ac, a^{-1}c; q)_\infty} \\ &\quad \times \mathfrak{E}^+(\gamma; z; a, b, qd^{-1}, qc^{-1}) W_{a,b,qd^{-1},qc^{-1},q^{-1}cdt}(z). \end{aligned}$$

The weight function $\tilde{W}(\gamma)$ occurring in the inverse transform is symmetric in b, c, qd^{-1} , while an equality similarly as above can be given for the exchange of a and b .

Remark 6.2. [25, Proposition 6.18] is equivalent with (6.3). Indeed, note from Definition 2.9, p.414, and (4.7) in [25] that

$$G_\alpha(\gamma)^{-1} = (-q^{\frac{1}{2}}t_0u_0^{-1}\gamma, -q^{\frac{1}{2}}t_0u_0^{-1}\gamma^{-1}; q)_\infty = (d\gamma, d\gamma^{-1}; q)_\infty.$$

Then σ , which interchanges t_0 and u_1 , sends $d = -q^{\frac{1}{2}}t_0u_0^{-1}$ to $\tilde{d} = -q^{\frac{1}{2}}u_1u_0^{-1}$. Next τ , which interchanges in α_σ u_0 and u_1 sends $\tilde{d} = -q^{\frac{1}{2}}u_1u_0^{-1}$ to $q\gamma\tilde{a}a^{-1}d^{-1} = q\tilde{d}^{-1} = -q^{\frac{1}{2}}u_0u_1^{-1}$. Hence

$$G_{\alpha_{\sigma\tau}}(\gamma)^{-1} = (-q^{\frac{1}{2}}u_0u_1^{-1}\gamma, -q^{\frac{1}{2}}u_0u_1^{-1}\gamma^{-1}; q)_\infty = (q\gamma\tilde{a}a^{-1}d^{-1}, q\gamma^{-1}\tilde{a}a^{-1}d^{-1}; q)_\infty$$

Finally note from [25, (4.7) and Proposition 4.15a] that the difference multiplicity functions α and β are obtained from each other by interchanging a and b . This finishes the proof that [25, Proposition 6.18] and (6.3) are equivalent.

See also [24, Section 4], where the one-variable symmetric Askey–Wilson functions are considered and the involutions σ and τ and the Gaussian G_α are immediately defined in terms of a, b, c, d .

Remark 6.3. Use the duality (5.16) in the left-hand side of (6.2), and also in the right-hand side in the form

$$\begin{aligned} \mathfrak{E}^+(\gamma; z; a, b, qd^{-1}, qc^{-1}) &= \mathfrak{E}^+\left(z; \gamma; \sqrt{\frac{qab}{cd}}, ab\sqrt{\frac{cd}{qab}}, qad^{-1}\sqrt{\frac{cd}{qab}}, qac^{-1}\sqrt{\frac{cd}{qab}}\right) \\ &= \mathfrak{E}^+\left(z; \gamma; \frac{ab}{\sqrt{q^{-1}abcd}}, \sqrt{q^{-1}abcd}, \frac{ac}{\sqrt{q^{-1}abcd}}, \frac{ad}{\sqrt{q^{-1}abcd}}\right) = \mathfrak{E}^+(z, \gamma; \tilde{b}, \tilde{a}, \tilde{c}, \tilde{d}). \end{aligned} \quad (6.4)$$

Thus (6.3) can be obtained by combining (6.2) with (5.16) and (6.4).

Remark 6.4. From [24, (4.4), (4.7)] together with (5.12), (5.13) and (5.5) we obtain the following expansion formula.

$$\begin{aligned} \mathfrak{E}^+(\gamma; z; a, b, c, d) &= \frac{(bc, qad^{-1}, qbd^{-1}, qcd^{-1}, qa^{-1}d^{-1}; q)_\infty}{(qabcd^{-1}; q)_\infty (qd^{-1}z, qd^{-1}z^{-1}, q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1}; q)_\infty} \\ &\times \sum_{m=0}^{\infty} (-1)^m (ad)^{-m} q^{\frac{1}{2}m(m+1)} \frac{1 - q^{2m}abcd^{-1}}{1 - abcd^{-1}} \frac{(ab, ac, abcd^{-1}; q)_m}{(qbd^{-1}, qcd^{-1}, q; q)_m} \\ &\times R_m(z; a, b, c, qd^{-1}) R_m(\gamma; \tilde{a}, \tilde{b}, \tilde{c}, q\tilde{d}^{-1}) \end{aligned} \quad (6.5)$$

$$\begin{aligned} &= \frac{(bc, qad^{-1}, qbd^{-1}, qcd^{-1}, qa^{-1}d^{-1}; q)_\infty}{(qabcd^{-1}; q)_\infty (qd^{-1}z, qd^{-1}z^{-1}, q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1}; q)_\infty} \\ &\times \sum_{m=0}^{\infty} (-1)^m (abcd^{-1})^{\frac{1}{2}m} q^{\frac{1}{2}m^2} \frac{(q^m abcd^{-1}; q)_m (qabcd^{-1}; q)_{2m}}{(ab, ac, bc, qad^{-1}, qbd^{-1}, qcd^{-1}, q; q)_m} \\ &\times P_m(z; a, b, c, qd^{-1}) P_m(\gamma; \tilde{a}, \tilde{b}, \tilde{c}, q\tilde{d}^{-1}). \end{aligned} \quad (6.6)$$

The infinite sum in (6.5) and (6.6) converge absolutely for all $z, \gamma \in \mathbb{C} \setminus (0)$. Formula (6.6) shows that the infinite sum is symmetric in a, b, c, qd^{-1} because the monic Askey–Wilson polynomials P_m are symmetric in their parameters. The duality (5.16) is implied by (6.5).

7 Non-symmetric Askey–Wilson functions and their symmetries

We want to obtain explicit expressions for non-symmetric Askey–Wilson functions in terms of (symmetric) Askey–Wilson functions similar to the explicit expressions for non-symmetric Askey–Wilson polynomials in terms of Askey–Wilson polynomials as obtained in [10]. Let us emphasize that there are two pairs of equivalent formulae for $E_{\pm n}$ in [10]: the formulae of [10, Theorem 4.1], and the formulae [10, (4.9) and (4.10)]. When trying to mimic these formulae to construct Askey–Wilson functions, the results differ!

7.1 A first choice of a non-symmetric Askey–Wilson function

We will look for an extension of the non-symmetric Askey–Wilson polynomials as given by [10, (4.1)] such that the eigenvalue equation [10, (4.4)] also extends. In the proof of [10, (4.4)] a first step was the following q -difference formula [10, (4.8)] for Askey–Wilson polynomials.

$$\delta_{q,z} P_n(z; a, b, c, d) = (q^{\frac{1}{2}n} - q^{-\frac{1}{2}n})(z - z^{-1})P_{n-1}(z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d), \quad (7.1)$$

where $P_n(z)$ is the monic Askey–Wilson polynomial given by (5.5) and where

$$\delta_{q,z} f(z) := f(q^{\frac{1}{2}}z) - f(q^{-\frac{1}{2}}z).$$

We will extend (7.1) for Askey–Wilson functions as given by (5.24),

Lemma 7.1. *We have*

$$\delta_{q,z} \mathfrak{E}^+(\gamma; z; a, b, c, d) = \frac{q^{\frac{1}{2}}a(1 - \gamma\tilde{a})(1 - \gamma^{-1}\tilde{a})}{(1 - ab)(1 - ac)(1 - ad)} (z - z^{-1})\mathfrak{E}^+(\gamma; z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d). \quad (7.2)$$

Proof. Straightforward computations give

$$\delta_{q,z} (az, az^{-1}; q)_k = q^{-\frac{1}{2}}a(1 - q^k)(z - z^{-1})(q^{\frac{1}{2}}az, q^{\frac{1}{2}}az^{-1}; q)_{k-1}$$

and

$$\begin{aligned} \delta_{q,z} \frac{(az, az^{-1}; q)_{\infty}}{(q^{k+1}d^{-1}z, q^{k+1}d^{-1}z^{-1}; q)_{\infty}} \\ = q^{-\frac{1}{2}}a(1 - q^{k+1}a^{-1}d^{-1})(z - z^{-1}) \frac{(q^{\frac{1}{2}}az, q^{\frac{1}{2}}az^{-1}; q)_{\infty}}{(q^{k+\frac{1}{2}}d^{-1}z, q^{k+\frac{1}{2}}d^{-1}z^{-1}; q)_{\infty}}. \end{aligned}$$

Hence, by (5.23),

$$\delta_{q,z} \mathfrak{R}(\gamma; z; a, b, c, d) = \frac{q^{\frac{1}{2}}a(1 - \gamma\tilde{a})(1 - \gamma^{-1}\tilde{a})}{(1 - ab)(1 - ac)(1 - ad)} (z - z^{-1})\mathfrak{R}(\gamma; z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d) \quad (7.3)$$

and

$$\begin{aligned} \delta_{q,z} \frac{(az, az^{-1}; q)_\infty}{(qd^{-1}z, qd^{-1}z^{-1}; q)_\infty} \mathfrak{R}(\gamma; z; qd^{-1}, b, c, qa^{-1}) &= q^{-\frac{1}{2}}a(1 - qa^{-1}d^{-1})(z - z^{-1}) \\ &\times \frac{(q^{\frac{1}{2}}az, q^{\frac{1}{2}}az^{-1}; q)_\infty}{(q^{\frac{1}{2}}d^{-1}z, q^{\frac{1}{2}}d^{-1}z^{-1}; q)_\infty} \mathfrak{R}(\gamma; z; q^{\frac{1}{2}}d^{-1}, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}a^{-1}). \end{aligned}$$

We conclude that (7.3) remains true if there $\mathfrak{R}(\gamma; z; a, b, c, d)$ is replaced by the second term on the right in (5.24), and hence also if there $\mathfrak{R}(\gamma; z; a, b, c, d)$ is replaced by $\mathfrak{E}^+(\gamma; z; a, b, c, d)$. \square

Note that, for $\gamma = q^n \tilde{a}$, (7.2) reduces by (5.5) and (5.20) to (7.1).

Another important element in the proof of [10, Theorem 4.1] is the following lemma (only sketched there).

Lemma 7.2. *If $f \in \mathcal{O}_{qd^{-1}}^+$ then*

$$\begin{aligned} (\hat{Y}f)(z) &= \frac{(c-z)(d-z)(1+ab-(a+b)z)}{(1-z^2)(q-z^2)} (f(q^{-1}z) - f(z)) \\ &\quad + \frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)} (f(qz) - f(z)) + q^{-1}abcd f(z). \end{aligned} \quad (7.4)$$

Also, if $g(z) = z^{-1}(c-z)(d-z)h(q^{-\frac{1}{2}}z)$ with $h \in \mathcal{O}_{qd^{-1}}^+$ then

$$(Yg)(z) = \frac{(c-z)(d-z)(1+ab-(a+b)z)}{z(1-z^2)} h(q^{-\frac{1}{2}}z) - \frac{(1-az)(1-bz)(1-cz)(1-dz)}{z(1-z^2)} h(q^{\frac{1}{2}}z). \quad (7.5)$$

Proof. Recall from [10, (3.13)] that

$$\begin{aligned} (\hat{Y}f)(z) &= \frac{z(1+ab-(a+b)z)((c+d)q-(cd+q)z)}{q(1-z^2)(q-z^2)} f(z) \\ &\quad + \frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)} f(qz) \\ &\quad + \frac{(1-az)(1-bz)((c+d)qz-(cd+q))}{q(1-z^2)(1-qz^2)} f(z^{-1}) \\ &\quad + \frac{(c-z)(d-z)(1+ab-(a+b)z)}{(1-z^2)(q-z^2)} f(qz^{-1}). \end{aligned} \quad (7.6)$$

Then (7.4) and (7.5) follow by straightforward computations. \square

Proposition 7.3. *Define a non-symmetric Askey–Wilson function*

$$\begin{aligned} \mathfrak{F}(\gamma; z; a, b, c, d) &:= \mathfrak{E}^+(\gamma; z; a, b, c, d) - \frac{a(1-\tilde{a}\gamma)}{(1-ab)(1-ac)(1-ad)} \\ &\quad \times z^{-1}(c-z)(d-z)\mathfrak{E}^+(\gamma; q^{-\frac{1}{2}}z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d). \end{aligned} \quad (7.7)$$

It satisfies the eigenvalue equation

$$\tilde{a}^{-1}(\hat{Y} \mathfrak{F}(\gamma; \cdot; a, b, c, d))(z) = \gamma^{-1} \mathfrak{F}(\gamma; z; a, b, c, d). \quad (7.8)$$

Proof. Consider (7.4) and (7.5) with

$$f(z) = \mathfrak{E}^+(\gamma; z; a, b, c, d), \quad g(z) = z^{-1}(c-z)(d-z)h(q^{-\frac{1}{2}}z), \quad h(z) = \mathfrak{E}^+(\gamma; z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d).$$

Then

$$\begin{aligned} (\hat{Y}f)(z) - \tilde{a}\gamma^{-1}f(z) &= \frac{(c-z)(d-z)(1+ab-(a+b)z)}{(1-z^2)(q-z^2)} (f(q^{-1}z) - f(z)) \\ &\quad + \frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)} (f(qz) - f(z)) + (\tilde{a}^2 - \tilde{a}\gamma^{-1})f(z) \end{aligned}$$

and

$$\begin{aligned} (\hat{Y}g)(z) - \tilde{a}\gamma^{-1}g(z) &= \frac{(c-z)(d-z)}{q-z^2} \left(\frac{(1+ab-(a+b)z)}{z(1-z^2)} - \tilde{a}\gamma^{-1} \right) \frac{(1-ab)(1-ac)(1-ad)}{a(1-\gamma\tilde{a})(1-\gamma^{-1}\tilde{a})} (f(q^{-1}z) - f(z)) \\ &\quad + \frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)} \frac{(1-ab)(1-ac)(1-ad)}{a(1-\gamma\tilde{a})(1-\gamma^{-1}\tilde{a})} (f(qz) - f(z)), \end{aligned}$$

where we also used (7.2). Now take a linear combination of the last two equalities as in (7.7). Then we obtain

$$\begin{aligned} &(\hat{Y} \mathfrak{F}(\gamma; \cdot; a, b, c, d))(z) - \tilde{a}\gamma^{-1} \mathfrak{F}(\gamma; z; a, b, c, d) \\ &= \frac{1}{1-\tilde{a}^{-1}\gamma} \left(\frac{(1-az)(1-bz)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)} (f(qz) - f(z)) \right. \\ &\quad \left. + \frac{(a-z)(b-z)(c-z)(d-z)}{(1-z^2)(q-z^2)} (f(q^{-1}z) - f(z)) + (1-\tilde{a}\gamma)(1-\tilde{a}\gamma^{-1})f(z) \right), \end{aligned}$$

which vanishes because of (4.21) and (5.22). \square

By (5.20) and (5.5) formula (7.7) reduces for $\gamma = q^n \tilde{a}$ to

$$\begin{aligned} \mathfrak{F}(q^n \tilde{a}; z; a, b, c, d) &= R_n(z; a, b, c, d) \\ &\quad - \frac{a(1-q^{n-1}abcd)}{(1-ab)(1-ac)(1-ad)} z^{-1}(c-z)(d-z)R_{n-1}(q^{-\frac{1}{2}}z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d) \\ &= \frac{(q^{n-1}abcd; q)_n a^n}{(ab, ac, ad; q)_n} \left(P_n(z; a, b, c, d) - q^{\frac{1}{2}(n-1)} z^{-1}(c-z)(d-z)P_{n-1}(q^{-\frac{1}{2}}z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d) \right), \end{aligned}$$

which is a constant multiple of the non-symmetric Askey–Wilson polynomial E_{-n} as defined in [10, (4.1), (3.19)]. Also, for $\gamma = q^{-n}\tilde{a}^{-1}$, formula (7.7) reduces to

$$\begin{aligned} \mathfrak{F}(q^{-n}\tilde{a}^{-1}; z; a, b, c, d) &= R_n(z; a, b, c, d) \\ &- \frac{a(1-q^{-n})}{(1-ab)(1-ac)(1-ad)} z^{-1}(c-z)(d-z)R_{n-1}(q^{-\frac{1}{2}}z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d) = \frac{(q^{n-1}abcd; q)_n a^n}{(ab, ac, ad; q)_n} \\ &\times \left(P_n(z; a, b, c, d) - \frac{q^{\frac{1}{2}(n-1)}(1-q^{-n})}{1-q^{n-1}abcd} z^{-1}(c-z)(d-z)P_{n-1}(q^{-\frac{1}{2}}z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d) \right), \end{aligned}$$

which is a constant multiple of the non-symmetric Askey–Wilson polynomial E_n as defined in [10, (4.1), (3.21)]. For these two choices of γ the eigenvalue equation (7.8) also agrees with the eigenvalue equations [10, (4.4), (4.5)].

We would like to prove, similarly as (6.2), that

$$\mathfrak{F}(\gamma; z; a, b, c, d) = \frac{(qad^{-1}, qa^{-1}d^{-1}; q)_\infty}{(ac, a^{-1}c; q)_\infty} \frac{G_c(z)}{G_{qd^{-1}}(z)} \mathfrak{F}(\gamma; z; a, b, qd^{-1}, qc^{-1}).$$

Unfortunately, the second term in (7.7) gives problems in doing this. Also a proof of an extension of the duality (5.16) to (7.7) fails because of that second term.

7.2 A better choice of non-symmetric Askey–Wilson functions

We now consider a different non-symmetric Askey–Wilson function (7.12), which extends the non-symmetric Askey–Wilson polynomials $E_{\pm n}$ as given by [10, (4.9), (4.10)]. In order to prove that this function is an eigenfunction of Y in the basic representation we need two preparatory lemmas.

Lemma 7.4. *We have*

$$\begin{aligned} &z(1-q^{-\frac{1}{2}}az^{-1})(1-q^{-\frac{1}{2}}bz^{-1})\mathfrak{E}^+(\gamma; q^{-\frac{1}{2}}z; a, b, c, d) \\ &\quad - z^{-1}(1-q^{-\frac{1}{2}}az)(1-q^{-\frac{1}{2}}bz)\mathfrak{E}^+(\gamma; q^{\frac{1}{2}}z; a, b, c, d) \\ &= (1-q^{-1}ab)(z-z^{-1})\mathfrak{E}^+(\gamma; z; q^{-\frac{1}{2}}a, q^{-\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d). \end{aligned} \quad (7.9)$$

Proof. Straightforward computations give

$$\begin{aligned} &z(1-q^{-\frac{1}{2}}az^{-1})(1-q^{-\frac{1}{2}}bz^{-1})(q^{-\frac{1}{2}}az, q^{\frac{1}{2}}az^{-1}; q)_k - z^{-1}(1-q^{-\frac{1}{2}}az)(1-q^{-\frac{1}{2}}bz)(q^{\frac{1}{2}}az, q^{-\frac{1}{2}}az^{-1}; q)_k \\ &= (1-q^{k-1}ab)(z-z^{-1})(q^{-\frac{1}{2}}az, q^{-\frac{1}{2}}az^{-1}; q)_k \end{aligned}$$

and

$$\begin{aligned} &z(1-q^{-\frac{1}{2}}az^{-1})(1-q^{-\frac{1}{2}}bz^{-1}) \frac{(q^{-\frac{1}{2}}az, q^{\frac{1}{2}}az^{-1}; q)_\infty}{(q^{k+\frac{1}{2}}d^{-1}z, q^{k+\frac{3}{2}}d^{-1}z^{-1}; q)_\infty} - z^{-1}(1-q^{-\frac{1}{2}}az)(1-q^{-\frac{1}{2}}bz) \\ &\quad \times \frac{(q^{\frac{1}{2}}az, q^{-\frac{1}{2}}az^{-1}; q)_\infty}{(q^{k+\frac{3}{2}}d^{-1}z, q^{k+\frac{1}{2}}d^{-1}z^{-1}; q)_\infty} = (1-q^k bd^{-1})(z-z^{-1}) \frac{(q^{-\frac{1}{2}}az, q^{-\frac{1}{2}}az^{-1}; q)_\infty}{(q^{k+\frac{1}{2}}d^{-1}z, q^{k+\frac{1}{2}}d^{-1}z^{-1}; q)_\infty}. \end{aligned}$$

Hence, by (5.23),

$$\begin{aligned}
& z(1 - q^{-\frac{1}{2}}az^{-1})(1 - q^{-\frac{1}{2}}bz^{-1})\mathfrak{R}(\gamma; q^{-\frac{1}{2}}z; a, b, c, d) \\
& \quad - z^{-1}(1 - q^{-\frac{1}{2}}az)(1 - q^{-\frac{1}{2}}bz)\mathfrak{R}(\gamma; q^{\frac{1}{2}}z; a, b, c, d) \\
& \quad = (1 - q^{-1}ab)(z - z^{-1})\mathfrak{R}(\gamma; z; q^{-\frac{1}{2}}a, q^{-\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d) \quad (7.10)
\end{aligned}$$

and

$$\begin{aligned}
& z(1 - q^{-\frac{1}{2}}az^{-1})(1 - q^{-\frac{1}{2}}bz^{-1}) \frac{(q^{-\frac{1}{2}}az, q^{\frac{1}{2}}az^{-1}; q)_{\infty}}{(q^{\frac{1}{2}}d^{-1}z, q^{\frac{3}{2}}d^{-1}z^{-1}; q)_{\infty}} \mathfrak{R}(\gamma; q^{-\frac{1}{2}}z; qd^{-1}, b, c, qa^{-1}) \\
& \quad - z^{-1}(1 - q^{-\frac{1}{2}}az)(1 - q^{-\frac{1}{2}}bz) \frac{(q^{\frac{1}{2}}az, q^{-\frac{1}{2}}az^{-1}; q)_{\infty}}{(q^{\frac{3}{2}}d^{-1}z, q^{\frac{1}{2}}d^{-1}z^{-1}; q)_{\infty}} \mathfrak{R}(\gamma; q^{\frac{1}{2}}z; qd^{-1}, b, c, qa^{-1}) \\
& \quad = (1 - bd^{-1})(z - z^{-1})\mathfrak{R}(\gamma; z; q^{\frac{1}{2}}d^{-1}, q^{-\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{3}{2}}a^{-1}).
\end{aligned}$$

We conclude that (7.10) remains true if there $\mathfrak{R}(\gamma; z; a, b, c, d)$ is replaced by the second term on the right in (5.24), and hence also if there $\mathfrak{R}(\gamma; z; a, b, c, d)$ is replaced by $\mathfrak{E}^+(\gamma; z; a, b, c, d)$. \square

For $\gamma = q^n \tilde{a}$ formula (7.9) becomes a formula for Askey–Wilson polynomials which was earlier given in [8, (2.1B)].

Lemma 7.5. *If $k \in \mathcal{O}_{qd^{-1}}^+$ and $l(z) := z^{-1}(1 - az)(1 - bz)k(z)$ then*

$$\begin{aligned}
a^{-1}b^{-1}(\hat{Y}l)(z) &= \left(a + b + \frac{1}{q} \left(-\frac{1}{a} - \frac{1}{b} + c + d \right) \right. \\
& \quad \left. + abcd \left(z + \frac{1}{z} - \frac{1}{a} - \frac{1}{b} - \frac{1}{c} - \frac{1}{d} \right) + \left(\frac{1}{qab} + cd \left(1 - \frac{1}{q} \right) - 1 \right) z^{-1} \right) k(z) \\
& \quad + \frac{(qa - z)(qb - z)(c - z)(d - z)(1 + ab - (a + b)z)}{qabz(1 - z^2)(q - z^2)} (k(q^{-1}z) - k(z)) \\
& \quad + \frac{(1 - az)(1 - bz)(1 - cz)(1 - dz)(1 - qaz)(1 - qbz)}{qabz(1 - z^2)(1 - qz^2)} (k(qz) - k(z)). \quad (7.11)
\end{aligned}$$

Proof. This follows by straightforward computation from (7.6). \square

Theorem 7.6. *Define a non-symmetric Askey–Wilson function*

$$\begin{aligned}
\mathfrak{E}(\gamma; z; a, b, c, d) &:= \mathfrak{E}^+(\gamma; z; a, b, c, d) - \frac{\sqrt{qab^{-1}cd}}{(1 - ab)(1 - qab)(1 - ac)(1 - ad)} \\
& \quad \times \gamma^{-1}(1 - \tilde{a}\gamma)(1 - \tilde{b}\gamma) z^{-1}(1 - az)(1 - bz) \mathfrak{E}^+(\gamma; z; qa, qb, c, d). \quad (7.12)
\end{aligned}$$

It satisfies the eigenvalue equation

$$\tilde{a}^{-1}(\hat{Y} \mathfrak{E}(\gamma; \cdot; a, b, c, d))(z) = \gamma^{-1} \mathfrak{E}(\gamma; z; a, b, c, d). \quad (7.13)$$

Furthermore, there is the normalization

$$\mathfrak{E}(\gamma; a^{-1}; a, b, c, d) = 1. \quad (7.14)$$

Also,

$$(qd^{-1}z, qd^{-1}z^{-1}, q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1}; q)_\infty \mathfrak{E}(\gamma; z; a, b, c, d)$$

is a complex analytic function in z, γ for $z, \gamma \neq 0$.

Proof. Consider (7.7) and (7.12) with

$$\begin{aligned} f(z) &= \mathfrak{E}^+(\gamma; z; a, b, c, d), & g(z) &= z^{-1}(c-z)(d-z)h(q^{-\frac{1}{2}}z), \\ h(z) &= \mathfrak{E}^+(\gamma; z; q^{\frac{1}{2}}a, q^{\frac{1}{2}}b, q^{\frac{1}{2}}c, q^{\frac{1}{2}}d), \\ k(z) &:= \mathfrak{E}^+(\gamma; z; qa, qb, c, d), & l(z) &:= z^{-1}(1-az)(1-bz)k(z). \end{aligned}$$

On comparing (7.8) and (7.13) we see that it is sufficient to prove that

$$((\hat{Y} - \tilde{a}\gamma^{-1})g)(z) = \frac{d}{1-qab} \frac{1 - \gamma\sqrt{qabc^{-1}d^{-1}}}{\gamma\sqrt{q^{-1}abc^{-1}d}} ((\hat{Y} - \tilde{a}\gamma^{-1})l)(z). \quad (7.15)$$

On the left-hand side of (7.15) substitute (7.5) and next substitute in the resulting expression for $h(q^{\pm\frac{1}{2}}z)$ the following formulas obtained from (7.9):

$$\begin{aligned} h(q^{-\frac{1}{2}}z) &= \frac{(1-qa z^{-1})(1-qb z^{-1})}{(1-qab)(1-qz^{-2})} k(q^{-1}z) + \frac{(1-az)(1-bz)}{(1-qab)(1-q^{-1}z^2)} k(z), \\ h(q^{\frac{1}{2}}z) &= \frac{(1-az^{-1})(1-bz^{-1})}{(1-qab)(1-q^{-1}z^{-2})} k(z) + \frac{(1-qa z)(1-qb z)}{(1-qab)(1-qz^2)} k(qz). \end{aligned}$$

On the right-hand side of (7.15) substitute (7.11). From the resulting equality (5.24), which yet has to be proved, consider the left-hand side minus the right-hand side. We can show that this equals

$$\begin{aligned} &\frac{cd(1-az)(1-bz)}{q\tilde{a}} (qab-1)\gamma z \left(\frac{(1-qa z)(1-qb z)(1-cz)(1-dz)}{(1-z^2)(1-qz^2)} (k(qz) - k(z)) \right. \\ &\quad \left. + \frac{(qa-z)(qb-z)(c-z)(d-z)}{(1-z^2)(q-z^2)} (k(q^{-1}z) - k(z)) + (1-q\tilde{a}\gamma)(1-q\tilde{a}\gamma^{-1})k(z) \right), \end{aligned}$$

which vanishes because of (4.21) and (5.22).

Formula (7.14) follows from (7.12) and (5.15). The analyticity statement follows from (7.12) together with Proposition 5.2. \square

Following [9, (3.17), (4.3), (4.9), (4.10)] we put for $n = 1, 2, \dots$:

$$Q_n(z; a, b, c, d) := a^{-1}b^{-1}z^{-1}(1 - az)(1 - bz)P_{n-1}(z; qa, qb, c, d), \quad (7.16)$$

$$E_{-n}(z; a, b, c, d) := \frac{ab}{ab - 1} (P_n(z; a, b, c, d) - Q_n(z; a, b, c, d)), \quad (7.17)$$

$$E_n(z; a, b, c, d) := \frac{1}{(1 - ab)(1 - q^{2n-1}abcd)} \left((1 - q^n ab)(1 - q^{n-1}abcd)P_n(z; a, b, c, d) - ab(1 - q^n)(1 - q^{n-1}cd)Q_n(z; a, b, c, d) \right), \quad (7.18)$$

$$E_0(z; a, b, c, d) := 1. \quad (7.19)$$

Here $P_n(z; a, b, c, d)$ is a monic Askey–Wilson polynomial (5.5) and $E_n(z; a, b, c, d)$ ($n \in \mathbb{Z}$) is a *non-symmetric Askey–Wilson polynomial*.

By (5.20) and (5.5) formula (7.12) reduces for $\gamma = q^n \tilde{a}$ ($n = 1, 2, \dots$) to

$$\begin{aligned} \mathfrak{E}(q^n \tilde{a}; z; a, b, c, d) &= R_n(z; a, b, c, d) - \frac{q^{1-n}(1 - q^{n-1}abcd)(1 - q^n ab)}{b(1 - ab)(1 - qab)(1 - ac)(1 - ad)} \\ &\quad \times z^{-1}(1 - az)(1 - bz)R_{n-1}(z; qa, qb, c, d) \end{aligned} \quad (7.20)$$

$$\begin{aligned} &= \frac{(q^{n-1}abcd; q)_n a^n}{(ab, ac, ad; q)_n} (P_n(z; a, b, c, d) - a^{-1}b^{-1}z^{-1}(1 - az)(1 - bz)P_{n-1}(z; qa, qb, c, d)) \\ &= \frac{(q^{n-1}abcd; q)_n a^n}{(ab, ac, ad; q)_n} (1 - a^{-1}b^{-1})E_{-n}(z; a, b, c, d). \end{aligned} \quad (7.21)$$

Also, for $\gamma = q^{-n} \tilde{a}^{-1}$ ($n = 1, 2, \dots$), formula (7.12) reduces to

$$\begin{aligned} \mathfrak{E}(q^{-n} \tilde{a}^{-1}; z; a, b, c, d) &= R_n(z; a, b, c, d) - \frac{aq^{1-n}(1 - q^n)(1 - q^{n-1}cd)}{(1 - ab)(1 - qab)(1 - ac)(1 - ad)} \\ &\quad \times z^{-1}(1 - az)(1 - bz)R_{n-1}(z; qa, qb, c, d) \\ &= \frac{(q^{n-1}abcd; q)_n a^n}{(ab, ac, ad; q)_n} \left(P_n(z; a, b, c, d) - \frac{(1 - q^n)(1 - q^{n-1}cd)}{(1 - q^n ab)(1 - q^{n-1}abcd)} \right. \\ &\quad \left. \times z^{-1}(1 - az)(1 - bz)P_{n-1}(z; qa, qb, c, d) \right) \\ &= \frac{(q^n abcd; q)_n a^n}{(qab, ac, ad; q)_n} E_n(z; a, b, c, d). \end{aligned} \quad (7.22)$$

Finally,

$$\mathfrak{E}(\tilde{a}^{-1}; z; a, b, c, d) = 1 = E_0(z; a, b, c, d). \quad (7.24)$$

Theorem 7.7. *The non-symmetric Askey–Wilson function (7.12) satisfies duality*

$$\mathfrak{E}(\gamma; z; a, b, c, d) = \mathfrak{E}(z, \gamma; \tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}), \quad (7.25)$$

and symmetries

$$\mathfrak{E}(\gamma; z; a, b, c, d) = \frac{(bc, qa^{-1}d^{-1}; q)_\infty}{(ac, qb^{-1}d^{-1}; q)_\infty} \frac{G_{\tilde{c}}(\gamma)}{G_{q\tilde{d}^{-1}}(\gamma)} \mathfrak{E}(\gamma; z; b, a, c, d), \quad (7.26)$$

$$\mathfrak{E}(\gamma; z; a, b, c, d) = \frac{(qad^{-1}, qa^{-1}d^{-1}; q)_\infty}{(ac, a^{-1}c; q)_\infty} \frac{G_c(z)}{G_{qd^{-1}}(z)} \mathfrak{E}(\gamma; z; a, b, qd^{-1}, qc^{-1}). \quad (7.27)$$

Proof. Each of the three formulas follows by (7.12) from a corresponding formula ((5.16), (6.3) and (6.2), respectively) for the two terms there. The symmetries (6.3) and (6.2) also follow from each other by the duality (5.16) in a similar way as in Remark 6.3. \square

Open problem Is $\mathfrak{E}(\gamma; z; a, b, c, d)$ different from $\mathfrak{F}(\gamma; z; a, b, c, d)$? If yes, does $\mathfrak{F}(\gamma; z; a, b, c, d)$ have any symmetries?

7.3 Relation with Stokman's kernel formula

Theorem 7.8. *The non-symmetric Askey–Wilson function (7.12) can be expanded as*

$$\begin{aligned} \mathfrak{E}(\gamma; z; a, b, c, d) &= \frac{(bc, qad^{-1}, qbd^{-1}, qcd^{-1}, qa^{-1}d^{-1}; q)_{\infty}}{(qabcd^{-1}; q)_{\infty}(qd^{-1}z, qd^{-1}z^{-1}, q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1}; q)_{\infty}} \\ &\times \sum_{m=0}^{\infty} (-1)^m (ad)^{-m} q^{\frac{1}{2}m(m+1)} \frac{1 - q^{2m}abcd^{-1}}{1 - abcd^{-1}} \frac{(ab, ac, abcd^{-1}; q)_m}{(qbd^{-1}, qcd^{-1}, q; q)_m} \\ &\times \left(-\frac{ab(1 - q^m)(1 - q^mcd^{-1})}{(1 - ab)(1 - q^{2m}abcd^{-1})} \mathfrak{E}(q^m\sqrt{abcd^{-1}}; z; a, b, c, qd^{-1}) \mathfrak{E}(q^m\sqrt{abcd^{-1}}; \gamma; \tilde{a}, \tilde{b}, \tilde{c}, q\tilde{d}^{-1}) \right. \\ &\left. + \frac{(1 - q^mab)(1 - q^mabcd^{-1})}{(1 - ab)(1 - q^{2m}abcd^{-1})} \mathfrak{E}\left(\frac{1}{q^m\sqrt{abcd^{-1}}}; z; a, b, c, qd^{-1}\right) \mathfrak{E}\left(\frac{1}{q^m\sqrt{abcd^{-1}}}; \gamma; \tilde{a}, \tilde{b}, \tilde{c}, q\tilde{d}^{-1}\right) \right). \end{aligned} \quad (7.28)$$

Proof. On the one hand, by (7.12) and (6.5) the left-hand side of (7.28) equals

$$\begin{aligned} &\frac{(bc, qad^{-1}, qbd^{-1}, qcd^{-1}, qa^{-1}d^{-1}; q)_{\infty}}{(qabcd^{-1}; q)_{\infty}(qd^{-1}z, qd^{-1}z^{-1}, q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1}; q)_{\infty}} \\ &\times \sum_{m=0}^{\infty} (-1)^m (ad)^{-m} q^{\frac{1}{2}m(m+1)} \frac{1 - q^{2m}abcd^{-1}}{1 - abcd^{-1}} \frac{(ab, ac, abcd^{-1}; q)_m}{(qbd^{-1}, qcd^{-1}, q; q)_m} \\ &\times \left(R_m(z; a, b, c, qd^{-1}) R_m(\gamma; \tilde{a}, \tilde{b}, \tilde{c}, q\tilde{d}^{-1}) \right. \\ &\quad - \sqrt{q^{-1}ab^{-1}cd} \frac{q^{2-2m}(1 - q^m)(1 - q^mab)(1 - q^mcd^{-1})(1 - q^mabcd^{-1})}{(1 - ab)^2(1 - qab)^2(1 - ac)^2(1 - bc)(1 - qad^{-1})} \\ &\quad \left. \times z^{-1}(1 - az)(1 - bz) R_{m-1}(z; qa, qb, c, qd^{-1}) \gamma^{-1}(1 - \tilde{a}\gamma)(1 - \tilde{b}\gamma) R_{m-1}(\gamma; q\tilde{a}, q\tilde{b}, \tilde{c}, q\tilde{d}^{-1}) \right). \end{aligned}$$

On the other hand, the right-hand side of (7.28) can be written in this form by (7.20) and (7.22). \square

By Theorem 7.6 the non-symmetric Askey–Wilson function (7.12) is equal to the *Cherednik kernel* given by Stokman [25, Definition 5.16] (in case $n = 1$). Indeed, property i) in [25, Theorem 5.17] follows from the analyticity statement in Theorem 7.6. For Property ii) use [25, Proposition 3.2] and the eigenvalue equation (7.13). Thus the function (7.12) equals the Cherednik kernel up to a constant factor. By the normalization (7.14) this constant factor is

equal to 1. So the explicit expansion (7.28) should coincide with the case $n = 1$ of [25, (6.6)] for $\mathfrak{E}(\gamma, x)$. We can also check this.

First observe that the parameters t_1, u_1, t_0, u_0 as introduced in [25, §2.2, $n = 1$], can be expressed in terms of our parameters u_1, k_1, u_0, k_0 by

$$(t_1, u_1, t_0, u_0)_{\text{St}} = (u_1, -k_1^{-1}, u_0, k_0)_{\text{K-M}}$$

(compare (2.14) with [25, (4.7)]). So $(t_1, u_1, t_0, u_0)_{\text{St}}$ relates to (a, b, c, d) by

$$\begin{aligned} (a, b, c, d) &= (t_1 u_1, -t_1 u_1^{-1}, q^{\frac{1}{2}} t_0 u_0, -q^{\frac{1}{2}} t_0 u_0^{-1}), \\ (t_1, u_1, t_0, u_0) &= \left(i\sqrt{ab}, -i\sqrt{ab^{-1}}, -i\sqrt{q^{-1}cd}, i\sqrt{cd^{-1}} \right). \end{aligned} \quad (7.29)$$

By [25, p.414] the involution σ interchanges u_1 and t_0 and the involution τ interchanges u_0 and t_0 . Note that these involutions agree with our σ and τ as discussed in §3.3 and (3.10) (only regarding their action on parameter space).

By [25, §4.1] the spectrum S is the spectrum of $\tilde{a}^{-1}Y$ (Y in our notation) acting in the polynomial representation. Thus by (7.13) the spectral value $s = q^{-n}\tilde{a}^{-1}$ corresponds to $\mathfrak{E}(q^n \tilde{a}; z; a, b, c, d)$ ($n = 1, 2, \dots$) and $s = q^n \tilde{a}$ corresponds to $\mathfrak{E}(q^{-n} \tilde{a}^{-1}; z; a, b, c, d)$ ($n = 0, 1, 2, \dots$). Thus the summation in the right-hand side of [25, (6.6)] takes for $n = 1$ and $x = z$ the form

$$\begin{aligned} &\sum_{m=1}^{\infty} \mu_{\tau}(q^{-m}(abcd^{-1})^{-\frac{1}{2}}) \mathfrak{E}(q^m(abcd^{-1})^{\frac{1}{2}}; z; a, b, c, qd^{-1}) \mathfrak{E}(q^m(abcd^{-1})^{\frac{1}{2}}; \gamma; \tilde{a}, \tilde{b}, \tilde{c}, q\tilde{d}^{-1}) \\ &+ \sum_{m=0}^{\infty} \mu_{\tau}(q^m(abcd^{-1})^{\frac{1}{2}}) \mathfrak{E}(q^{-m}(abcd^{-1})^{-\frac{1}{2}}; z; a, b, c, qd^{-1}) \mathfrak{E}(q^{-m}(abcd^{-1})^{-\frac{1}{2}}; \gamma; \tilde{a}, \tilde{b}, \tilde{c}, q\tilde{d}^{-1}). \end{aligned}$$

By [25, Definition 2.9]

$$G(z) = G_d(z) = \frac{1}{(dz, dz^{-1}; q)_{\infty}}.$$

Hence

$$G_{\tau}(z)G_{\sigma\tau}(\gamma) = G_{qd^{-1}}(z)G_{q\tilde{d}^{-1}}(\gamma) = \frac{1}{(qd^{-1}z, qd^{-1}z^{-1}, q\tilde{d}^{-1}\gamma, q\tilde{d}^{-1}\gamma^{-1}; q)_{\infty}}.$$

By [25, (6.9)]

$$C_0 = \frac{(ad, bd, cd, bc, a^{-1}d; q)_{\infty}}{(abcd; q)_{\infty}}, \quad (C_0)_{\tau} = \frac{(qad^{-1}, qbd^{-1}, qcd^{-1}, bc, qa^{-1}d^{-1}; q)_{\infty}}{(qabcd^{-1}; q)_{\infty}}.$$

Hence the factor in front of the summation sign in (7.28) equals $(C_0)_{\tau}G_{\sigma\tau}(\gamma)G_{\tau}(z)$.

We now turn to the quotient

$$\frac{G_{\tau\sigma\tau}(s)N_{\tau\sigma}(s^{-1})}{G_{\tau\sigma\tau}(s_0)N_{\tau\sigma}(s_0^{\frac{1}{2}})}$$

(see [25, (6.7)]) which should give the coefficients after the summation sign in (7.28). By [25, (4.16)] this quotient is for $s = (q^m(abcd^{-1})^{\frac{1}{2}})^{\pm 1}$ equal to

$$\frac{G_{\tau\sigma\tau}(q^m(abcd^{-1})^{\frac{1}{2}})}{G_{\tau\sigma\tau}((abcd^{-1})^{\frac{1}{2}})} \frac{N_{\tau\sigma}^+(q^m(abcd^{-1})^{\frac{1}{2}})}{N_{\tau\sigma}^+((abcd^{-1})^{\frac{1}{2}})} \frac{\mathcal{C}_{\tau\sigma}((q^m(abcd^{-1})^{\frac{1}{2}})^{\mp 1})}{\mathcal{C}_{\tau\sigma}((abcd^{-1})^{-\frac{1}{2}})}$$

We have

$$G_{\tau\sigma\tau}(z) = G_{\sqrt{a^{-1}bcd}}(z) = \frac{1}{(\sqrt{a^{-1}bcd} z, \sqrt{a^{-1}bcd} z^{-1}; q)_{\infty}}.$$

Hence

$$\frac{G_{\tau\sigma\tau}(q^m(abcd^{-1})^{\frac{1}{2}})}{G_{\tau\sigma\tau}((abcd^{-1})^{\frac{1}{2}})} = \frac{(bc; q)_m}{(qad^{-1}; q)_m} (-ad^{-1})^m q^{\frac{1}{2}m(m+1)}$$

(see also this evaluation in the Proof of Theorem 4.2 in [24]). We also have

$$\frac{N_{\tau\sigma}^+(q^m(abcd^{-1})^{\frac{1}{2}})}{N_{\tau\sigma}^+((abcd^{-1})^{\frac{1}{2}})} = a^{-2m} \frac{1 - q^{2m}abcd^{-1}}{1 - abcd^{-1}} \frac{(ab, ac, qad^{-1}, abcd^{-1}; q)_m}{(bc, qbd^{-1}, qcd^{-1}, q; q)_m}$$

(see [25, (4.17)] or see this evaluation in the Proof of Theorem 4.2 in [24] which uses [24, (2.6)]).

Thus

$$\frac{G_{\tau\sigma\tau}(q^m(abcd^{-1})^{\frac{1}{2}})}{G_{\tau\sigma\tau}((abcd^{-1})^{\frac{1}{2}})} \frac{N_{\tau\sigma}^+(q^m(abcd^{-1})^{\frac{1}{2}})}{N_{\tau\sigma}^+((abcd^{-1})^{\frac{1}{2}})}$$

is equal to the coefficient after the summation sign and before the big brackets in (7.28). By [25, (2.3), (4.14)] we have

$$\mathcal{C}(x) = \frac{(1 - ax^{-1})(1 - b(x^{-1}))}{1 - x^{-2}}$$

(see also the expression for $\alpha(x)$ in [18, §§6.25]). Hence

$$\begin{aligned} \mathcal{C}_{\tau\sigma}((q^m(abcd^{-1})^{\frac{1}{2}})^{-1}) &= \frac{(1 - q^m ab)(1 - q^m abcd^{-1})}{(1 - q^{2m} abcd^{-1})}, \\ \mathcal{C}_{\tau\sigma}((abcd^{-1})^{-\frac{1}{2}}) &= 1 - ab, \\ \mathcal{C}_{\tau\sigma}(q^m(abcd^{-1})^{\frac{1}{2}}) &= -\frac{ab(1 - q^m)(1 - q^m cd^{-1})}{(1 - q^{2m} abcd^{-1})}. \end{aligned}$$

This gives the coefficients inside the big brackets in (7.28).

So we have indeed checked that the expansion (7.28) coincides with the case $n = 1$ of Stokman's kernel formula [25, (6.6)] for $\mathfrak{E}(\gamma, x)$.

Acknowledgments We thank Jasper Stokman for helping us to show that the non-symmetric Askey–Wilson function defined by (7.12) is equal to the function defined by the case $n = 1$ of his Cherednik kernel [25, Definition 5.16].

The work of Marta Mazzocco was funded by the Leverhulme Trust Research Project Grant RPG-2021-047.

References

- [1] R. Askey and J. A. Wilson, Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials, *Mem. Amer. Math. Soc.* **54**, No. 319 (1985).
- [2] N. Crampé , L. Frappat , J. Gaboriaud, L. Poulain d’Andecy, E. Ragoucy and L. Vinet, The Askey–Wilson algebra and its avatars, *J. Phys. A: Math. Theor.*, **54** (2021).
- [3] B. A. Dubrovin and M. Mazzocco, Monodromy of certain Painlevé-VI transcendents and reflection group, *Invent. Math.* **141** (2000), 55–147.
- [4] G. Gasper and M. Rahman, *Basic hypergeometric series*, Second edition, Cambridge University Press, Cambridge, 2004.
- [5] L. Haine and P. Iliev, Askey–Wilson type functions, with bound states, *Ramanujan J.* **11** (2006), 285–329.
- [6] J. E. Humphreys, *Introduction to Lie algebras and representation theory*, Springer-Verlag, Third printing, 1990.
- [7] M. E. H. Ismail and M. Rahman, The associated Askey–Wilson polynomials, *Trans. Amer. Math. Soc.* **328** (1991), 201–237.
- [8] E. G. Kalnins and W. Miller, Jr., Symmetry techniques for q -series: Askey–Wilson polynomials, *Rocky Mountain J. Math.* **19** (1989), 223–230.
- [9] E. Koelink and J. V. Stokman, The Askey–Wilson function transform, *Internat. Math. Res. Notices* (**2001**), No. 22, 1203–1227.
- [10] T. H. Koornwinder, The relationship between Zhedanov’s algebra $AW(3)$ and the double affine Hecke algebra in the rank one case, *SIGMA* **3** (2007), 063, 15 pages; [arXiv:math/0612730v4 \[math.QA\]](#).
- [11] T. H. Koornwinder, Zhedanov’s algebra $AW(3)$ and the double affine Hecke algebra in the rank one case. II. The spherical subalgebra, *SIGMA* **4** (2008), 052, 17 pages; [arXiv:0711.2320v3 \[math.QA\]](#).
- [12] T. H. Koornwinder, Charting the q -Askey scheme. II. The q -Zhedanov scheme, *Indag. Math. (N.S.)* **34** (2023) 317–337,1419–1420; [arXiv:2209.07995v3 \[math.CA\]](#).
- [13] T. H. Koornwinder and F. Bouzeffour, Non-symmetric Askey-Wilson polynomials as vector-valued polynomials *Appl. Anal.* **90** (2011), no. 3-4, 731–746.
- [14] T. H. Koornwinder and M. Mazzocco, Dualities in the q -Askey scheme and degenerate DAHA, *Studies Appl. Math.* **141** (2018), 424–473.
- [15] O. Lisovyy and Y. Tykhyy, Algebraic solutions of the sixth Painlevé equation, *J. Geom. Phys.* **85** (2014), 124–163.

- [16] M. Mazzocco, Confluences of the Painlevé equations, Cherednik algebras and q -Askey scheme, *Nonlinearity* **29** (2016), 2565–2608.
- [17] M. Mazzocco, Embedding of the rank 1 DAHA into $Mat(2, \mathbb{T}_q)$ and its automorphisms, in: *Representation theory, special functions and Painlevé equations, RIMS 2015*, Adv. Stud. Pure Math. **76**, pp. 449–468, Math. Soc. Japan, Tokyo, 2018.
- [18] M. Noumi and J. V. Stokman, Askey–Wilson polynomials: an affine Hecke algebraic approach, in: *Laredo Lectures on Orthogonal Polynomials and Special Functions*, Adv. Theory Spec. Funct. Orthogonal Polynomials, Nova Sci. Publ., Hauppauge, NY (2004) 111–144; [arXiv:math/0001033v1 \[math.QA\]](#).
- [19] A. Oblomkov, Double affine Hecke algebras of rank 1 and affine cubic surfaces, *Internat. Math. Res. Notices* (**2004**), No. 18, 877–912.
- [20] K. Okamoto, Studies on the Painlevé equations. I. Sixth Painlevé equation P_{VI} , *Ann. Mat. Pura Appl. (4)* **146** (1987), 337–381.
- [21] M. Rahman, Askey–Wilson functions of the first and second kinds: series and integral representations of $C_n^2(x; \beta | q) + D_n^2(x; \beta | q)$, *J. Math. Anal. Appl.* **164** (1992), 263–284.
- [22] M. Rahman, A q -extension of a product formula of Watson, *Quaest. Math.* **22** (1999), 27–42.
- [23] S. Sahi, non-symmetric Koornwinder polynomials and duality, *Ann. of Math. (2)* **150**, (1999), 267–282.
- [24] J. V. Stokman, An expansion formula for the Askey–Wilson function, *J. Approx. Theory* **114** (2002), 308–342.
- [25] J. V. Stokman, Difference Fourier transforms for nonreduced root systems *Selecta Math. (N.S.)* **9** (2003), 409–494.
- [26] S. K. Suslov, The theory of difference analogues of special functions of hypergeometric type, *Russian Math. Surveys* **44** (1989) no. 2, 227–278.
- [27] S. K. Suslov, Some orthogonal very well poised ${}_8\phi_7$ -functions. *J. Phys. A* **30** (1997), 5877–5885.
- [28] S. K. Suslov, Some orthogonal very-well-poised ${}_8\phi_7$ -functions that generalize Askey–Wilson polynomials, *Ramanujan J.* **5** (2001), 183–218.
- [29] P. Terwilliger, The universal Askey–Wilson algebra, *SIGMA* **7** (2011), 069, 24 pp.