

THE LOW DIMENSIONAL HOMOLOGY GROUPS OF THE ELEMENTARY GROUP OF RANK TWO

BEHROOZ MIRZAI AND ELVIS TORRES PÉREZ

ABSTRACT. In this article we study the first, the second and the third homology groups of the elementary group $E_2(A)$, where A is a commutative ring. In particular, we prove a refined Bloch-Wigner type exact sequence over a semilocal ring (with some mild restriction on its residue fields) such that $-1 \in (A^\times)^2$ or $|A^\times/(A^\times)^2| \leq 4$.

For a commutative ring A with unit, let $E_2(A)$ be the subgroup of the general linear group $GL_2(A)$ generated by elementary matrices. Clearly $E_2(A) \subseteq SL_2(A)$. We say that A is a $GE_2(A)$ -ring if $E_2(A) = SL_2(A)$.

In a series of papers [14], [15] and [16] we studied the low dimensional homology groups of the elementary group $E_2(A)$. In this paper, we further study these homology groups.

The first main result of this paper is over the first homology group of $E_2(A)$. We show that for any commutative ring A , we have the exact sequence

$$H_2(E_2(A), \mathbb{Z}) \rightarrow H_1(Y_\bullet(A^2))_{E_2(A)} \rightarrow A/M \rightarrow H_1(E_2(A), \mathbb{Z}) \rightarrow 0,$$

where $Y_\bullet(A^2)$ is a complex of $GE_2(A)$ -modules obtained from certain unimodular vectors in A^2 (see Theorem 2.1). Here M is the additive subgroup of A generated by $x(a^2 - 1)$ and $3(b + 1)(c + 1)$, where $x \in A$ and $a, b, c \in A^\times$. The right hand side map is induced by the map $A \rightarrow E_2(A)$, where sends x to the elementary matrix $E_{12}(x)$. The group A/M and its connection with the first homology of $E_2(A)$ first were studied by Cohn in [3] and [4].

From the complex $Y_\bullet(A^2)$ we obtain a spectral sequence $E_{p,q}^1 \Rightarrow H_{p+q}(E_2(A), \mathbb{Z})$ which will be our main tool in handling the low dimensional homology groups of $E_2(A)$. The above exact sequence is obtained from a careful analysing of this spectral sequence. This gives a different proof of the main result of [16, Theorem 3.1] over commutative rings.

We use further this spectral sequence to study the second homology group $H_2(E_2(A), \mathbb{Z})$, when A is a semilocal ring such that $2 \in A^\times$ and all its residue fields have more than four elements (see Theorem 3.4). More precisely, we give a set of cycles in $B_\bullet(E_2(A)) \otimes_{E_2(A)} \mathbb{Z}$, where their homology classes generate $H_2(E_2(A), \mathbb{Z})$. Here $B_\bullet(E_2(A)) \rightarrow \mathbb{Z}$ is the bar resolution of $E_2(A)$ over \mathbb{Z} [2, Chap. I, §5]. This result is used in the study of the third homology of $E_2(A)$.

The third main result of this paper is a refined Bloch-Wigner exact sequence over a semilocal domain. This exact sequence studies the third homology of $E_2(A)$. More precisely, let A be a semilocal ring such that for any maximal ideal $\mathfrak{m} \in \text{Specm}(A)$, the residue field A/\mathfrak{m} either is infinite or when $|A/\mathfrak{m}| = p^d$, then $(p - 1)d > 6$. If $-1 \in A^{\times 2}$ or

$|A^\times/(A^\times)^2| \leq 4$, then we obtain the refined Bloch-Wigner exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^\sim \rightarrow \frac{H_3(\mathbb{E}_2(A), \mathbb{Z})}{\mu_2(A) \otimes_{\mathbb{Z}} I^2(A)} \rightarrow \mathcal{RB}(A) \rightarrow 0.$$

Here $I(A)$ is the fundamental ideal of A and $\mathcal{RB}(A)$ is the refined Bloch group of A . Moreover, the map $\mu_2(A) \otimes_{\mathbb{Z}} I^2(A) \rightarrow H_3(\mathbb{E}_2(A), \mathbb{Z})$ is obtained from the product map

$$\mu_2(A) \otimes_{\mathbb{Z}} H_2(\mathbb{E}_2(A), \mathbb{Z}) \rightarrow H_3(\mathbb{E}_2(A), \mathbb{Z}),$$

which is induced by the map $\mu_2(A) \times \mathbb{E}_2(A) \rightarrow \mathbb{E}_2(A)$, $(-1, X) \mapsto -X$. This generalizes the main result of [15]. In [19, Question 0.1] we raised the question that over local rings is the product map

$$\mu_2(A) \otimes_{\mathbb{Z}} H_2(\mathbb{E}_2(A), \mathbb{Z}) \rightarrow H_3(\mathbb{E}_2(A), \mathbb{Z})$$

trivial, provided that $|A/\mathfrak{m}_A| \neq 2$?

Notations. In this paper all rings are commutative, except probably group rings, and have the unit element 1. For a commutative ring A , let $\mathcal{G}_A := A^\times/(A^\times)^2$ and set

$$\mathcal{W}_A := \{a \in A : a(1-a) \in A^\times\}.$$

The element of \mathcal{G}_A represented by $a \in A^\times$ is denoted by $\langle a \rangle$ and set $\langle\langle a \rangle\rangle := \langle a \rangle - 1 \in \mathbb{Z}[\mathcal{G}_A]$. For $n \in \mathbb{N}$, let $\mu_n(A)$ denotes the group of n -roots of unity in A . Moreover, let

$$\mu(A) := \bigcup_{n \in \mathbb{N}} \mu_n(A).$$

If $\mathcal{B} \rightarrow \mathcal{A}$ is a homomorphism of abelian groups, by \mathcal{A}/\mathcal{B} we mean $\mathrm{coker}(\mathcal{B} \rightarrow \mathcal{A})$.

1. THE COMPLEX OF UNIMODULAR VECTORS

Let A be a commutative ring. Let $\mathbb{E}_2(A)$ be the subgroup of $\mathrm{GL}_2(A)$ generated by the elementary matrices $E_{12}(a) := \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$ and $E_{21}(a) := \begin{pmatrix} 1 & 0 \\ a & 1 \end{pmatrix}$, $a \in A$. Let $\mathrm{GE}_2(A)$ be the subgroup of $\mathrm{GL}_2(A)$ generated by diagonal and elementary matrices. It is easy to see that $\mathbb{E}_2(A)$ is normal in $\mathrm{GE}_2(A)$ [3, Proposition 2.1] and the center of $\mathrm{GE}_2(A)$ is $A^\times I_2$. We say that A is a *GE₂-ring* if $\mathrm{GE}_2(A) = \mathrm{GL}_2(A)$. Semilocal rings and euclidean domains are examples of *GE₂-rings* [20, p. 245], [3, §2].

We call $\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \in A^2$ a *GE₂-unimodular vector* if there exists a vector $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \in A^2$ such that $(\mathbf{u}, \mathbf{v}) := \begin{pmatrix} u_1 & v_1 \\ u_2 & v_2 \end{pmatrix} \in \mathrm{GE}_2(A)$. For any non-negative integer n , let $Y_n(A^2)$ be the free abelian group generated by the set of all $(n+1)$ -tuples $(\langle \mathbf{v}_0 \rangle, \dots, \langle \mathbf{v}_n \rangle)$, where every vector $\mathbf{v}_i \in A^2$ is *GE₂-unimodular* and for any $i \neq j$, $(\mathbf{v}_i, \mathbf{v}_j) \in \mathrm{GE}_2(A)$.

We consider $Y_l(A^2)$ as a left $\mathrm{GE}_2(A)$ -module (resp. left $\mathbb{E}_2(A)$ -module) in a natural way. If necessary, we convert this action to a right action by the definition $m.g := g^{-1}m$. Let us define the l -th differential operator

$$\partial_l : Y_l(A^2) \rightarrow Y_{l-1}(A^2), \quad l \geq 1,$$

as an alternating sum of face operators which throws away the i -th component of generators. Hence we have the complex

$$Y_\bullet(A^2) : \cdots \longrightarrow Y_2(A^2) \xrightarrow{\partial_2} Y_1(A^2) \xrightarrow{\partial_1} Y_0(A^2) \rightarrow 0.$$

We say that the above complex is exact in dimension k if $H_k(Y_\bullet(A^2)) = 0$. Let $\partial_{-1} = \varepsilon : Y_0(A^2) \rightarrow \mathbb{Z}$ be defined by $\sum_i n_i \langle \mathbf{v}_{0,i} \rangle \mapsto \sum_i n_i$.

Proposition 1.1 (Hutchinson). *For any commutative ring A , $H_0(Y_\bullet(A^2)) \xrightarrow{\cong} \mathbb{Z}$. In other words, the complex $Y_1(A^2) \xrightarrow{\partial_1} Y_0(A^2) \xrightarrow{\varepsilon} \mathbb{Z} \rightarrow 0$ always is exact.*

Proof. See [11, Theorem 3.3] or [18, Proposition 2.1]. □

Let $Z_1(A^2) := \ker(\partial_1)$. From the complex $0 \rightarrow Z_1(A^2) \xrightarrow{\text{inc}} Y_1(A^2) \xrightarrow{\partial_1} Y_0(A^2) \rightarrow 0$, we obtain the double complex

$$0 \rightarrow B_\bullet(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Z_1(A^2) \rightarrow B_\bullet(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Y_1(A^2) \rightarrow B_\bullet(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Y_0(A^2) \rightarrow 0,$$

where $B_\bullet(\mathbb{E}_2(A)) \rightarrow \mathbb{Z}$ is the bar resolution of \mathbb{Z} over $\mathbb{E}_2(A)$. This gives us the first quadrant spectral sequence

$$E_{p,q}^1 = \begin{cases} H_q(\mathbb{E}_2(A), Y_p(A^2)) & p = 0, 1 \\ H_q(\mathbb{E}_2(A), Z_1(A^2)) & p = 2 \\ 0 & p > 2 \end{cases} \implies H_{p+q}(\mathbb{E}_2(A), \mathbb{Z})$$

(see [2, §5, Chap. VII] or [13, §1]).

The group $\mathbb{E}_2(A)$ acts transitively on the sets of generators of $Y_i(A^2)$ for $i = 0, 1$. Let

$$\infty := \langle \mathbf{e}_1 \rangle, \quad \mathbf{0} := \langle \mathbf{e}_2 \rangle, \quad \mathbf{a} := \langle \mathbf{e}_1 + a\mathbf{e}_2 \rangle, \quad a \in A^\times,$$

where $\mathbf{e}_1 := \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\mathbf{e}_2 := \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. We choose (∞) and $(\infty, \mathbf{0})$ as representatives of the orbit of the generators of $Y_0(A^2)$ and $Y_1(A^2)$, respectively. Then

$$Y_0(A^2) \simeq \text{Ind}_{\mathbb{B}(A)}^{\mathbb{E}_2(A)} \mathbb{Z}, \quad Y_1(A^2) \simeq \text{Ind}_{\mathbb{T}(A)}^{\mathbb{E}_2(A)} \mathbb{Z},$$

where

$$\mathbb{B}(A) := \text{Stab}_{\mathbb{E}_2(A)}(\infty) = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} : a \in A^\times, b \in A \right\},$$

$$\mathbb{T}(A) := \text{Stab}_{\mathbb{E}_2(A)}(\infty, \mathbf{0}) = \{D(a) : a \in A^\times\},$$

where $D(a) := \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$ (see [2, §3, Chap. I]). Note that $\mathbb{T}(A) \simeq A^\times$ and $\mathbb{B}(A) \simeq A \rtimes A^\times$.

In our calculations usually we identify $\mathbb{T}(A)$ with A^\times . By Shapiro's lemma we have

$$E_{0,q}^1 \simeq H_q(\mathbb{B}(A), \mathbb{Z}), \quad E_{1,q}^1 \simeq H_q(\mathbb{T}(A), \mathbb{Z}).$$

In particular, $E_{0,0}^1 \simeq \mathbb{Z} \simeq E_{1,0}^1$. Moreover, $d_{1,q}^1 = H_q(\sigma) - H_q(\text{inc})$, where $\sigma : T(A) \rightarrow B(A)$ is given by $\sigma(D(a)) = wD(a)w^{-1} = D(a^{-1})$ for $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. This easily implies that $d_{1,0}^1$ is trivial and

$$d_{1,1}^1 : T(A) \rightarrow B(A)^{\text{ab}} = H_1(B(A), \mathbb{Z}), \quad D(a) \mapsto D(a)^{-2}.$$

In fact, note that

$$B(A)^{\text{ab}} \simeq T(A) \oplus A_{A^\times} \simeq A^\times \oplus A_{A^\times},$$

where

$$A_{A^\times} = A / \langle a^2 - 1 : a \in A^\times \rangle.$$

Thus $d_{1,1}^1(D(a)) = (D(a)^{-2}, 0)$. This implies that

$$E_{0,1}^2 \simeq \mathcal{G}_A \oplus A_{A^\times}, \quad \ker(d_{1,1}^1) \simeq \mu_2(A).$$

Furthermore, it is not difficult to see that $d_{2,1}^1 : H_1(E_2(A), Z_1(A^2)) \rightarrow \mu_2(A)$ is surjective. In fact, for any $b \in \mu_2(A)$, we have

$$d_{2,1}^1([b] \otimes \partial_2(\infty, \mathbf{0}, \mathbf{a})) = b.$$

Therefore $E_{1,1}^2 = 0$.

The orbits of the action of $E_2(A)$ on $Y_2(A)$ are represented by

$$\langle a \rangle [] := (\infty, \mathbf{0}, \mathbf{a}), \quad \langle a \rangle \in \mathcal{G}_A.$$

Therefore $Y_2(A^2)_{E_2(A)} \simeq \mathbb{Z}[\mathcal{G}_A]$. Moreover, the map

$$\mathbb{Z}[\mathcal{G}_A] \simeq Y_2(A^2)_{E_2(A)} \rightarrow Y_1(A^2)_{E_2(A)} \simeq \mathbb{Z}$$

coincides with the augmentation map. Now from the composite

$$\mathbb{Z}[\mathcal{G}_A] \simeq Y_2(A^2)_{E_2(A)} \rightarrow Z_1(A^2)_{E_2(A)} \rightarrow Y_1(A^2)_{E_2(A)} \simeq \mathbb{Z}$$

it follows that $d_{2,0}^1 : H_0(E_2(A), Z_1(A^2)) \rightarrow \mathbb{Z}$ is surjective. Hence $E_{1,0}^2 = 0$. Let

$$\text{GW}(A) := H_0(E_2(A), Z_1(A^2)), \quad \epsilon := d_{2,0}^1 : \text{GW}(A) \rightarrow \mathbb{Z}.$$

We denote the kernel of ϵ by $I(A)$. Let

$$\overline{\text{GW}}(A) := \mathbb{Z}[\mathcal{G}_A] / \langle \langle a \rangle \langle 1 - a \rangle : a \in \mathcal{W}_A \rangle.$$

The augmentation map $\mathbb{Z}[\mathcal{G}_A] \rightarrow \mathbb{Z}$ induces the natural map $\bar{\epsilon} : \overline{\text{GW}}(A) \rightarrow \mathbb{Z}$. The kernel of this map is denoted by $\bar{I}(A)$. Note that

$$\bar{I}(A) := \mathcal{I}_A / \langle \langle a \rangle \langle 1 - a \rangle : a \in \mathcal{W}_A \rangle,$$

where \mathcal{I}_A is the augmentation ideal of $\mathbb{Z}[\mathcal{G}_A]$.

From the exact sequence $0 \rightarrow B_1(A^2) \rightarrow Z_1(A^2) \rightarrow H_1(Y_\bullet(A^2)) \rightarrow 0$, we obtain the exact sequence

$$H_0(E_2(A), B_1(A^2)) \rightarrow \text{GW}(A) \rightarrow H_1(Y_\bullet(A^2))_{E_2(A)} \rightarrow 0.$$

From the sequence $Y_3(A^2) \rightarrow Y_2(A^2) \rightarrow B_1(A^2)$ we obtain the sequence

$$H_0(E_2(A), Y_3(A^2)) \rightarrow H_0(E_2(A), Y_2(A^2)) \rightarrow H_0(E_2(A), B_1(A^2)).$$

We showed in above that $Y_2(A^2)_{E_2(A)} \simeq \mathbb{Z}[\mathcal{G}_A]$. The orbits of the action of $E_2(A)$ on $Y_3(A)$ are represented by

$$\langle a \rangle [x] := (\infty, \mathbf{0}, \mathbf{a}, \mathbf{ax}), \quad \langle a \rangle \in \mathcal{G}_A, x \in \mathcal{W}_A.$$

Thus $Y_3(A^2)_{E_2(A)}$ is the free $\mathbb{Z}[\mathcal{G}_A]$ -module generated by the symbols $[x]$, $x \in \mathcal{W}_A$. Moreover the map

$$Y_3(A^2)_{E_2(A)} \rightarrow Y_2(A^2)_{E_2(A)} \simeq \mathbb{Z}[\mathcal{G}_A]$$

is given by $[x] \mapsto \langle\langle x \rangle\rangle \langle\langle 1-x \rangle\rangle$. All these induce a surjective map

$$\overline{\text{GW}}(A) \twoheadrightarrow H_0(E_2(A), B_1(A^2)).$$

Thus we have the exact sequence

$$\overline{\text{GW}}(A) \rightarrow \text{GW}(A) \rightarrow H_1(Y_\bullet(A^2))_{E_2(A)} \rightarrow 0.$$

Finally from this we obtain the exact sequence

$$(1.1) \quad \bar{I}(A) \rightarrow I(A) \rightarrow H_1(Y_\bullet(A^2))_{E_2(A)} \rightarrow 0.$$

2. THE FIRST HOMOLOGY OF E_2

The following theorem first was proved in [16, Theorem 3.1]. Here we present a new proof of the theorem using the spectral sequence $E_{\bullet, \bullet}$ introduced in the previous section.

Theorem 2.1. *Let A be a commutative ring and M the additive subgroup of A generated by $x(a^2 - 1)$ and $3(b + 1)(c + 1)$, where $x \in A$ and $a, b, c \in A^\times$. Then there is an exact sequence of \mathcal{G}_A -modules*

$$H_2(E_2(A), \mathbb{Z}) \rightarrow H_1(Y_\bullet(A^2))_{E_2(A)} \rightarrow A/M \rightarrow H_1(E_2(A), \mathbb{Z}) \rightarrow 0.$$

Proof. The second page of the spectral sequence $E_{\bullet, \bullet}$ is as follows:

$$\begin{array}{ccccccc} H_2(B(A), \mathbb{Z}) & & * & & * & & 0 \\ & \swarrow & & & & & \\ & & d_{2,1}^2 & & & & \\ & & & & & & \\ \mathcal{G}_A \oplus A_{A^\times} & & 0 & & E_{2,1}^2 & & 0 \\ & \swarrow & & & & & \\ & & d_{2,0}^2 & & & & \\ & & & & & & \\ \mathbb{Z} & & 0 & & I(A) & & 0 \end{array}$$

Now by an easy analysis of the above spectral sequence, we obtain the exact sequence

$$H_2(B(A), \mathbb{Z}) \rightarrow H_2(E_2(A), \mathbb{Z}) \rightarrow I(A) \xrightarrow{d_{2,0}^2} \mathcal{G}_A \oplus A_{A^\times} \rightarrow H_1(E_2(A), \mathbb{Z}) \rightarrow 0.$$

Let study the composite map $\bar{I}(A) \rightarrow I(A) \xrightarrow{d_{2,0}^2} \mathcal{G}_A \oplus A_{A^\times}$. Take the element $\langle\langle a \rangle\rangle \in \bar{I}(A)$, $a \in A^\times$. This is represented by

$$X_a := [] \otimes ((\infty, \mathbf{0}, \mathbf{a}) - (\infty, \mathbf{0}, \mathbf{1})) \in B_0(E_2(A)) \otimes_{E_2(A)} Y_2(A^2).$$

By passing to $I(A)$, the image of $\langle\langle a \rangle\rangle$ is represented by

$$[] \otimes \partial_2((\infty, \mathbf{0}, \mathbf{a}) - (\infty, \mathbf{0}, \mathbf{1})) \in B_0(E_2(A)) \otimes_{E_2(A)} Z_1(A^2),$$

which we denote it again by X_a . Thus $d_{2,0}^2(\langle\langle a \rangle\rangle) = d_{2,0}^2(\overline{X_a})$. To calculate the above composition we look at the diagram

$$\begin{array}{ccc} B_1(\mathbb{E}_2(A)) \otimes Y_0(A^2) & \xleftarrow{\text{id} \otimes \partial_1} & B_1(\mathbb{E}_2(A)) \otimes Y_1(A^2) \\ & & \downarrow d_1 \otimes \text{id} \\ & & B_0(\mathbb{E}_2(A)) \otimes Y_1(A^2) \xleftarrow{\text{id} \otimes \text{inc}} B_0(\mathbb{E}_2(A)) \otimes Z_1(A^2). \end{array}$$

We have

$$\begin{aligned} (\text{id}_{B_0(\mathbb{E}_2(A))} \otimes \text{inc})(X_a) &= [] \otimes ((\mathbf{0}, \mathbf{a}) - (\infty, \mathbf{a}) - (\mathbf{0}, \mathbf{1}) + (\infty, \mathbf{1})) \\ &= [] \otimes (g_a(\infty, \mathbf{0}) - h_a(\infty, \mathbf{0}) - g_1(\infty, \mathbf{0}) + h_1(\infty, \mathbf{0})) \\ &= (g_a^{-1} - h_a^{-1} - g_1^{-1} + h_1^{-1})[] \otimes (\infty, \mathbf{0}), \end{aligned}$$

where $g_a = \begin{pmatrix} 0 & 1 \\ -1 & a \end{pmatrix}$, $h_a = \begin{pmatrix} 1 & a^{-1} \\ 0 & 1 \end{pmatrix}$. This element sits in the image of $d_1 \otimes \text{id}_{Y_1(A^2)}$. In fact

$$(d_1 \otimes \text{id}_{Y_1(A^2)})((g_a^{-1} - [h_a^{-1}] - [g_1^{-1}] + [h_1^{-1}]) \otimes (\infty, \mathbf{0})) = (g_a^{-1} - h_a^{-1} - g_1^{-1} + h_1^{-1})[] \otimes (\infty, \mathbf{0}).$$

Finally $(\text{id}_{B_1(\mathbb{E}_2(A))} \otimes \partial_1)((g_a^{-1} - [h_a^{-1}] - [g_1^{-1}] + [h_1^{-1}]) \otimes (\infty, \mathbf{0}))$ represents $d_{2,0}^2(\langle\langle a \rangle\rangle)$. This is the element

$$\left(w([g_a^{-1}] - [h_a^{-1}] - [g_1^{-1}] + [h_1^{-1}]) - ([g_a^{-1}] - [h_a^{-1}] - [g_1^{-1}] + [h_1^{-1}]) \right) \otimes (\infty) \in B_1(\mathbb{E}_2(A)) \otimes Y_0(A^2).$$

By adding the null element $(d_2 \otimes \text{id}_{Y_0(A^2)})((w[h_a^{-1}] - [w|g_a^{-1}] + [w|g_1^{-1}] - [w|h_1^{-1}]) \otimes (\infty))$, we see that $d_{2,0}^2(\langle\langle a \rangle\rangle)$ is represented by

$$([wg_a^{-1}] - [wh_a^{-1}] - [wg_1^{-1}] + [wh_1^{-1}] - [g_a^{-1}] + [h_a^{-1}] + [g_1^{-1}] - [h_1^{-1}]) \otimes (\infty).$$

Now by adding the null element $(d_2 \otimes \text{id}_{Y_0(A^2)})((D(a)^{-1}h_{a^{-1}}^{-1}|wh_a^{-1}] - [h_1^{-1}|wh_1^{-1}]) \otimes (\infty)$ (note that $D(z)^{-1}h_{z^{-1}}^{-1} \in B(A) = \text{Stab}_{\mathbb{E}_2(A)}(\infty)$ and $D(z)^{-1}h_{a^{-1}}^{-1}wh_a^{-1} = wg_a^{-1}$), we see that $d_{2,0}^2(\langle\langle a \rangle\rangle)$ is represented by

$$([D(a)^{-1}h_{a^{-1}}^{-1}] - [h_1^{-1}] - [g_a^{-1}] + [h_a^{-1}] + [g_1^{-1}] - [h_1^{-1}]) \otimes (\infty).$$

Once more, by adding the null element

$$(d_2 \otimes \text{id}_{X_0(A^2)})((h_1| - w] - [h_{a^{-1}}| - w]) \otimes (\infty) = ([g_a^{-1}] - [h_{a^{-1}}] - [g_1^{-1}] + [h_1]) \otimes (\infty),$$

we see that $d_{2,0}^2(\langle\langle a \rangle\rangle)$ is represented by

$$([D(a)^{-1}h_{a^{-1}}^{-1}] - [h_1^{-1}] + [h_a^{-1}] - [h_1^{-1}] - [h_{a^{-1}}] + [h_1]) \otimes (\infty).$$

Note that in $H_1(\mathbb{E}_2(A), Y_0(A^2))$, if $g, h \in B(A)$, then

$$[gh] \otimes (\infty) = ([g] + [h]) \otimes (\infty) \quad \text{and} \quad [h^{-1}] \otimes (\infty) = -[h] \otimes (\infty).$$

Thus $d_{2,0}^2(\langle\langle a \rangle\rangle)$ in $H_1(\mathbf{E}_2(A), Y_0(A^2))/\text{im}(d_{1,1}^1)$ is represented by

$$[D(a)^{-1}h_{a^{-1}}^{-2}h_a^{-1}h_1^3] \otimes (\infty) = \left[\begin{pmatrix} a^{-1} & 0 \\ 0 & a \end{pmatrix} \begin{pmatrix} 1 & -2a - a^{-1} + 3 \\ 0 & 1 \end{pmatrix} \right] \otimes (\infty).$$

Therefore in $\mathcal{G}_A \oplus A_{A^\times}$ we have

$$d_{2,0}^2(\langle\langle a \rangle\rangle) = (\langle a^{-1} \rangle, \overline{-2a - a^{-1} + 3}) = (\langle a \rangle, \overline{3(1 - a)}) = 3(\langle a \rangle, \overline{1 - a}).$$

Thus we get the exact sequence

$$H_2(\mathbf{E}_2(A), \mathbb{Z}) \rightarrow I(A)/\bar{I}(A) \rightarrow \frac{\mathcal{G}_A \oplus A_{A^\times}}{\langle\langle b \rangle, 3(b-1) \mid b \in A^\times \rangle} \rightarrow H_1(\mathbf{E}_2(A), \mathbb{Z}) \rightarrow 0.$$

The map

$$\frac{A}{M} \rightarrow \frac{\mathcal{G}_A \oplus A_{A^\times}}{\langle\langle b \rangle, 3(b-1) \mid b \in A^\times \rangle}, \quad \bar{a} \mapsto (\langle 1 \rangle, \bar{a}),$$

is an isomorphism with the inverse map

$$\frac{\mathcal{G}_A \oplus A_{A^\times}}{\langle\langle b \rangle, 3(b-1) \mid b \in A^\times \rangle} \rightarrow \frac{A}{M}, \quad (\langle b \rangle, \bar{a}) \mapsto \overline{a - 3b + 3}.$$

Now (1.1) completes the proof of the theorem. \square

The Steinberg group $\text{St}(2, A)$ is the group with generators $x_{12}(r)$ and $x_{21}(s)$, $r, s \in A$, subject to the Steinberg relations

- (α) $x_{ij}(r)x_{ij}(s) = x_{ij}(r+s)$ for any $r, s \in A$,
- (β) $w_{ij}(u)x_{ji}(r)w_{ij}(u)^{-1} = x_{ij}(-u^2r)$, for any $u \in A^\times$ and $r \in A$,
where $w_{ij}(u) := x_{ij}(u)x_{ji}(-u^{-1})x_{ij}(u)$.

The map $\text{St}(2, A) \rightarrow \mathbf{E}_2(A)$ given by $x_{ij}(r) \mapsto E_{ij}(r)$, is a well defined homomorphism of groups. The unstable K_2 -group of order 2 is the kernel of this map and is denoted by $K_2(2, A)$. For $u \in A^\times$, let $h_{ij}(u) := w_{ij}(u)w_{ij}(-1)$. An element of the form

$$\{v, u\} := \{v, u\}_{12} = h_{12}(uv)h_{12}^{-1}(u)h_{12}(v)^{-1}$$

is called a Steinberg symbol. This is an element of $K_2(2, A)$ and is a central element of $\text{St}(2, A)$ [6, §9]. Let $C(2, A)$ be the subgroup of $K_2(2, A)$ generated by the Steinberg symbols.

Proposition 2.2 (Hutchinson). *For any commutative ring A , we have*

$$H_1(Y_\bullet) \simeq \left(\frac{K_2(2, A)}{C(2, A)} \right)^{\text{ab}}.$$

Proof. See [11, Theorem 7.2]. \square

A ring A is called universal for GE_2 if $K_2(2, A)$ is generated by Steinberg symbols [6, p. 251]. Thus if A is universal for GE_2 , then

$$H_1(\mathbf{E}_2(A), \mathbb{Z}) \simeq A/M.$$

Example 2.3. A commutative semilocal ring A is universal for GE_2 if and only if none of the rings $\mathbb{Z}/2 \times \mathbb{Z}/2$ and $\mathbb{Z}/2 \times \mathbb{Z}/3$ is a direct factor of $A/J(A)$ [12, Theorem 2.14], where $J(A)$ is the Jacobson radical of A . Thus if A is semilocal with the above condition, then

$$H_1(\text{E}_2(A), \mathbb{Z}) \simeq A/M.$$

(i) If A is a local ring with maximal ideal \mathfrak{m}_A , then

$$H_1(\text{E}_2(A), \mathbb{Z}) \simeq \begin{cases} A/\mathfrak{m}_A^2 & \text{if } |A/\mathfrak{m}_A| = 2 \\ A/\mathfrak{m}_A & \text{if } |A/\mathfrak{m}_A| = 3 \\ 0 & \text{if } |A/\mathfrak{m}_A| \geq 4 \end{cases}$$

(see [16, Proposition 4.1]).

(ii) If A is semilocal such that all its residue fields have more than three elements, i.e. $|A/\mathfrak{m}| > 3$ for all $\mathfrak{m} \in \text{Specm}(A)$, then there is an element $a \in A^\times$ such that $a^2 - 1 \in A^\times$. Thus $A = M$ and hence

$$H_1(\text{E}_2(A), \mathbb{Z}) = 0.$$

(iii) If A is Artinian, then $A \simeq A_1 \times \cdots \times A_r$, where each A_i is Artinian local rings (see [1, Theorem 8.7]). Hence $\text{E}_2(A) \simeq \text{E}_2(A_1) \times \cdots \times \text{E}_2(A_r)$ and thus

$$H_1(\text{E}_2(A), \mathbb{Z}) \simeq H_1(\text{E}_2(A_1), \mathbb{Z}) \times \cdots \times H_1(\text{E}_2(A_r), \mathbb{Z}).$$

Now using (i) we can calculate $H_1(\text{E}_2(A), \mathbb{Z})$. For more applications of the above theorem see [16].

3. THE SECOND HOMOLOGY OF E_2

We have seen in the proof of Theorem 2.1 that the composite $\bar{I}(A) \rightarrow I(A) \xrightarrow{d_{2,0}^2} \mathcal{G}_A \oplus A_{A^\times}$, takes $\langle\langle a \rangle\rangle \in \bar{I}(A)$, $a \in A^\times$, to the element $(\langle a \rangle, \overline{3(1-a)})$.

Let either $2 \in A^\times$ or $A_{A^\times} = 0$. In both cases $\overline{3(1-a)} = 0$ in A_{A^\times} . So the above composite sends $\langle\langle a \rangle\rangle$ to $(\langle a \rangle, 0)$. Note that if $2 \in A^\times$, then we have a surjective map of abelian groups $A/3A \rightarrow A_{A^\times}$, since $3 = 2^2 - 1$.

Denote the kernel of $d_{2,0}^2 : I(A) \rightarrow \mathcal{G}_A \oplus A_{A^\times}$, by $I^2(A)$, i.e.

$$I^2(A) := \ker(d_{2,0}^2).$$

We know that $\bar{I}(A)/\bar{I}^2(A) \simeq \mathcal{G}_A$ [21, Theorem 6.1.11]. Consider the commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \bar{I}^2(A) & \longrightarrow & \bar{I}(A) & \longrightarrow & \mathcal{G}_A \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & I^2(A) & \longrightarrow & I(A) & \longrightarrow & \mathcal{G}_A \oplus A_{A^\times}, \end{array}$$

where the right vertical map is given by $\langle a \rangle \mapsto (\langle a \rangle, 0)$. By the Snake Lemma and (1.1) we obtain the exact sequence

$$(3.1) \quad \bar{I}^2(A) \rightarrow I^2(A) \rightarrow H_1(Y_\bullet(A^2))_{\text{E}_2(A)} \rightarrow A_{A^\times}.$$

Lemma 3.1. *If $A_{A^\times} = 0$, then we have the exact sequence*

$$\bar{I}^2(A) \rightarrow I^2(A) \rightarrow H_1(Y_\bullet(A^2))_{\mathbb{E}_2(A)} \rightarrow 0.$$

Proof. This follows immediately from the above exact sequence. \square

On the other hand, by an easy analysis of the main spectral sequence we have the exact sequence

$$(3.2) \quad H_2(B(A), \mathbb{Z}) \rightarrow H_2(\mathbb{E}_2(A), \mathbb{Z}) \rightarrow I^2(A) \rightarrow 0.$$

We denote the image of $\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \in \bar{I}^2(A)$ in $I^2(A)$ again by $\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \in I^2(A)$ to an element of $H_2(\mathbb{E}_2(A), \mathbb{Z})$ through the surjective map

$$H_2(\mathbb{E}_2(A), \mathbb{Z}) \rightarrow I^2(A).$$

Lemma 3.2. *Let A be a commutative ring with $2 \in A^\times$. If $a, b \in A^\times$, then under the surjective map*

$$H_2(\mathbb{E}_2(A), \mathbb{Z}) \rightarrow I^2(A),$$

the homology class of the cycles $([D(a)|D(b)] + R_{ab} - R_a - R_b + R_1) \otimes 1$ maps to $\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \in I^2(A)$, where for any $z \in A^\times$,

$$R_z := [w|g_z^{-1}] - [w|h_z^{-1}] - [h_z^{-1}D(z)^{-1}|wh_z^{-1}] + [h_{z^{-1}}|w^{-1}] - [h_z^{-1}|D(z)^{-1}] + [D(z)h_z|D(z)^{-1}] \\ - [D(z)h_z|h_z^{-1}] + 2[h_z^{-1}|h_z^{-1}] + [h_z^{-2}|h_z^{-2}] + [D(2)h_z^{-1}D(2)^{-1}|D(2)] - [D(2)|h_z^{-1}]$$

with $g_z = \begin{pmatrix} 0 & 1 \\ -1 & z \end{pmatrix}$ and $h_z = \begin{pmatrix} 1 & z^{-1} \\ 0 & 1 \end{pmatrix}$.

Proof. We know that $I^2(A) \subseteq I(A) \subseteq \text{GW}(A) = H_0(\mathbb{E}_2(A), Z_1(A^2))$. So we need to study the map

$$H_2(\mathbb{E}_2(A), \mathbb{Z}) \rightarrow H_0(\mathbb{E}_2(A), Z_1(A^2)).$$

From the exact sequence $0 \rightarrow Z_1(A^2) \rightarrow Y_1(A^2) \rightarrow Y_0(A^2) \rightarrow \mathbb{Z} \rightarrow 0$ we obtain two short exact sequences

$$0 \rightarrow Z_1(A^2) \rightarrow Y_1(A^2) \rightarrow Z_0(A^2) \rightarrow 0, \quad 0 \rightarrow Z_0(A^2) \rightarrow Y_0(A^2) \rightarrow \mathbb{Z} \rightarrow 0.$$

From these we obtain the connecting homomorphisms

$$\delta_1 : H_2(\mathbb{E}_2(A), \mathbb{Z}) \rightarrow H_1(\mathbb{E}_2(A), Z_0(A^2)), \\ \delta_2 : H_1(\mathbb{E}_2(A), Z_0(A^2)) \rightarrow H_0(\mathbb{E}_2(A), Z_1(A^2)).$$

Then

$$\delta_2 \circ \delta_1 : H_2(\mathbb{E}_2(A), \mathbb{Z}) \rightarrow H_0(\mathbb{E}_2(A), Z_1(A^2))$$

is the above mentioned map. For $z \in A^\times$, the element $\langle\langle z \rangle\rangle \in I(A)$ is represented by $[\] \otimes \partial_2(X_z - X_1)$, where $X_z := (\infty, \mathbf{0}, \mathbf{z})$. Consider the diagram

$$B_1(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Z_0(A^2) \xleftarrow{\text{id}_{B_1} \otimes \partial_1} B_1(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Y_1(A^2) \\ \downarrow d_1 \otimes \text{id}_{Y_1} \\ B_0(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Y_1(A^2) \xleftarrow{\text{id}_{B_0} \otimes \text{inc}} B_0(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Z_1(A^2).$$

Then

$$\begin{aligned}
(\text{id}_{B_0} \otimes \text{inc})\left([\] \otimes \partial_2(X_z - X_1)\right) &= [\] \otimes \left(\mathbf{0}, \mathbf{z}\right) - (\infty, \mathbf{z}) - (\mathbf{0}, \mathbf{1}) + (\infty, \mathbf{1}) \\
&= [\] \otimes \left((g_z - h_z - g_1 + h_1)(\infty, \mathbf{0})\right) \\
&= (d_1 \otimes \text{id}_{Y_1(A^2)})\left(\left([g_z^{-1}] - [h_z^{-1}] - [g_1^{-1}] + [h_1^{-1}]\right) \otimes (\infty, \mathbf{0})\right).
\end{aligned}$$

Thus the element

$$\begin{aligned}
U_z &:= (\text{id}_{B_1} \otimes \partial_1)\left(\left([g_z^{-1}] - [h_z^{-1}] - [g_1^{-1}] + [h_1^{-1}]\right) \otimes (\infty, \mathbf{0})\right) \\
&= \left([g_z^{-1}] - [h_z^{-1}] - [g_1^{-1}] + [h_1^{-1}]\right) \otimes \partial_1(\infty, \mathbf{0})
\end{aligned}$$

represents an element of $H_1(\mathbb{E}_2(A), Z_0(A^2))$, where maps to $\langle\langle z \rangle\rangle$ through δ_2 . Now consider the diagram

$$\begin{array}{ccc}
B_2(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} \mathbb{Z} & \xleftarrow{\text{id}_{B_2} \otimes \varepsilon} & B_2(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Y_0(A^2) \\
& & \downarrow d_2 \otimes \text{id}_{Y_0} \\
& & B_1(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Y_0(A^2) \xleftarrow{\text{id}_{B_1} \otimes \text{inc}} B_1(\mathbb{E}_2(A)) \otimes_{\mathbb{E}_2(A)} Z_0(A^2).
\end{array}$$

Then we have

$$\begin{aligned}
(\text{id}_{B_1} \otimes \text{inc})(U_z) &= \left([g_z^{-1}] - [h_z^{-1}] - [g_1^{-1}] + [h_1^{-1}]\right) \otimes \partial_1(\infty, \mathbf{0}) \\
&= (w - 1)\left([g_z^{-1}] - [h_z^{-1}] - [g_1^{-1}] + [h_1^{-1}]\right) \otimes (\infty) \\
&= -[D(z)] + (d_2 \otimes \text{id}_{Y_0(A^2)})\left(\left(R_z - R_1\right) \otimes (\infty)\right).
\end{aligned}$$

Note that in our calculation we used the formulas

$$w g_z^{-1} = D(z)^{-1} h_{z^{-1}}^{-1} w h_z^{-1}, \quad D(z) h_u D(z)^{-1} = h_{u/z^2},$$

with $u, z \in A^\times$. Now it is easy to see that

$$(d_2 \otimes \text{id}_{Y_0(A^2)})\left(\left([D(a)|D(b)] + R_{ab} - R_a - R_b + R_1\right) \otimes (\infty)\right) = (\text{id}_{B_1} \otimes \text{inc})(U_{ab} - U_a - U_b).$$

Finally, applying $\text{id}_{B_2} \otimes \varepsilon$ we have

$$(\text{id}_{B_2} \otimes \varepsilon)\left(\left([D(a)|D(b)] + R_{ab} - R_a - R_b + R_1\right) \otimes (\infty)\right) = \left([D(a)|D(b)] + R_{ab} - R_a - R_b + R_1\right) \otimes 1.$$

The element $\left([D(a)|D(b)] + R_{ab} - R_a - R_b + R_1\right) \otimes 1$ is a cycle and represents an element in $H_2(\mathbb{E}_2(A), \mathbb{Z})$ which maps to $\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \in I^2(A)$. This completes the proof of the lemma. \square

Remark 3.3. In [9, §7] Hutchinson has constructed a similar cycles as in above lemma over fields. Let F be a field with at least four elements. Let $a, b \in F^\times$ and $\lambda \in F^\times$ such that $\lambda^2 - 1 \in F^\times$. Then Hutchinson has constructed a cycle

$$F(a, b)_\lambda \in B_2(\text{SL}_2(F)) \otimes_{\text{SL}_2(F)} \mathbb{Z}$$

such that its homology class in $H_2(\mathrm{SL}_2(F), \mathbb{Z})$ is independent of the choice of λ and maps to $\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \in I^2(F)$ (see [9, Proposition 7.1, Lemma 7.5, Theorem 7.8]). (Note that the existence of $\lambda \in A^\times$ such that $\lambda^2 - 1 \in A^\times$ implies that $A_{A^\times} = 0$.)

Theorem 3.4. *Let A be a semilocal ring such that $2 \in A^\times$ and all its residue fields have more than four elements. Then the group $H_2(\mathbb{E}_2(A), \mathbb{Z})$ is generated by the homology class of the cycles $([D(a)|D(b)] + R_{ab} - R_a - R_b + R_1)$, $([D(a)|D(b)] - [D(b)|D(a)])$, $a, b \in A^\times$, and $([E_{12}(x)|E_{12}(y)] - [E_{12}(y)|E_{12}(x)])$, $x, y \in A$.*

Proof. Since A is universal for GE_2 (see Example 2.3), by Proposition 2.2, $H_1(Y_\bullet(A^2)) = 0$. Thus by Lemma 3.1, the map $\bar{I}^2(A) \rightarrow I^2(A)$ is surjective. This shows that $I^2(A)$ is generated by the elements of the form $\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle$, $a, b \in A^\times$.

Now consider the exact sequence

$$H_2(\mathrm{B}(A), \mathbb{Z}) \rightarrow H_2(\mathbb{E}_2(A), \mathbb{Z}) \xrightarrow{\theta} I^2(A) \rightarrow 0.$$

Let $X \in H_2(\mathbb{E}_2(A), \mathbb{Z})$ and let

$$\theta(X) = \sum \langle\langle a_i \rangle\rangle \langle\langle b_i \rangle\rangle.$$

If $F(a, b) := ([D(a)|D(b)] + R_{ab} - R_a - R_b + R_1)$, then $\theta(X - \sum \overline{F(a_i, b_i)}) = 0$. It follows from this that $X - \sum \overline{F(a_i, b_i)}$ sits in the image of $H_2(\mathrm{B}(A), \mathbb{Z})$.

Consider the extension

$$1 \rightarrow \mathrm{N}(A) \rightarrow \mathrm{B}(A) \rightarrow \mathrm{T}(A) \rightarrow 1,$$

where $\mathrm{N}(A) := \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} : a \in A^\times, b \in A \right\}$. Since any residue field of A has more than four elements, there is $\lambda \in A^\times$ such that $\lambda^2 - 1 \in A^\times$. This implies that $A_{A^\times} = 0$. It follows from this that for any $k \geq 0$, $H_k(A^\times, A) = 0$ [14, Corollary 3.2]. By studying the Lyndon/Hochschild-Serre spectral sequence of this extension we obtain the exact sequence

$$H_2(A, \mathbb{Z})_{A^\times} \rightarrow H_2(\mathrm{B}(A), \mathbb{Z}) \rightarrow H_2(\mathrm{T}(A), \mathbb{Z}) \rightarrow 0.$$

Note that we have the splitting map $\mathrm{inc}_* : H_2(\mathrm{T}(A), \mathbb{Z}) \rightarrow H_2(\mathrm{B}(A), \mathbb{Z})$. The image of $H_2(A, \mathbb{Z}) \simeq A \wedge A$ in $H_2(\mathrm{B}(A), \mathbb{Z})$ is generated by the homology classes of the cycles

$$G(x, y) := ([E_{12}(x)|E_{12}(y)] - [E_{12}(y)|E_{12}(x)]) \otimes 1, \quad x, y \in A.$$

Moreover, $H_2(\mathrm{T}(A), \mathbb{Z}) \simeq \mathrm{T}(A) \wedge \mathrm{T}(A)$ is generated by the homology classes of the cycles

$$H(a, b) := ([D(a)|D(b)] - [D(b)|D(a)]) \otimes 1.$$

Now the claim follows from the fact that $\ker(\theta)$ is generated by the images of the homology classes of the cycles $G(x, y)$ and $H(a, b)$. This completes the proof of the theorem. \square

4. THE MAP $H_n(T(A), \mathbb{Z}) \rightarrow H_n(B(A), \mathbb{Z})$

In the previous sections, sometime we assumed that $A_{A^\times} = 0$, i.e. the ring A is generated by the elements of the form $a^2 - 1$, where $a \in A^\times$. It follows from this that the inclusion $T(A) \rightarrow B(A)$ induces the isomorphism

$$H_1(T(A), \mathbb{Z}) \simeq H_1(B(A), \mathbb{Z}).$$

In the next section we will need such isomorphism for the second and the third homology groups. Hutchinson has proved the following result over local domains [10, Proposition 3.19].

Proposition 4.1. *Let A be a semilocal ring. For any maximal ideal $\mathfrak{m} \in \text{Specm}(A)$, let either A/\mathfrak{m} be infinite or if $|A/\mathfrak{m}| = p^d$, then $(p-1)d > 2(n+1)$ ($(p-1)d > 2n$ when A is a domain). Then $H_k(T(A), \mathbb{Z}) \simeq H_k(B(A), \mathbb{Z})$ for all $k \leq n$.*

Proof. If A is a domain, then the proof is very similar to the proof of a similar result over local domains presented in [10, §3]. Just there one should replace the maximal ideal of the local domain (denoted by \mathcal{M} in [10, §3]) with the Jacobson radical $J(A)$ of A .

Now let A be a general semilocal ring with the given condition. Then similar to the proof presented in [10, Proposition 3.19], we can show that $H_n(T(A), k) \simeq H_n(B(A), k)$, where k is a prime field and $(p-1)d > 2n$. Now the claim follows from [13, Lemma 2.3]. \square

5. THE THIRD HOMOLOGY OF E_2

Let $\overline{\mathcal{RP}}(A)$ be the quotient of the free $\mathbb{Z}[\mathcal{G}_A]$ -module generated by symbols $[a]$, $a \in \mathcal{W}_A$, by the $\mathbb{Z}[\mathcal{G}_A]$ -submodule generated by the elements

$$[a] - [b] + \langle a \rangle \left[\frac{b}{a} \right] - \langle a^{-1} - 1 \rangle \left[\frac{1 - a^{-1}}{1 - b^{-1}} \right] + \langle 1 - a \rangle \left[\frac{1 - a}{1 - b} \right],$$

where $a, b, a/b \in \mathcal{W}_A$. By a direct computation one can show that the map

$$\bar{\lambda}_1 : \overline{\mathcal{RP}}(A) \rightarrow \mathcal{I}_A^2 \subseteq \mathbb{Z}[\mathcal{G}_A], \quad [a] \mapsto \langle\langle a \rangle\rangle \langle\langle 1 - a \rangle\rangle$$

is a well-defined $\mathbb{Z}[\mathcal{G}_A]$ -homomorphism. Let

$$S_{\mathbb{Z}}^2(A^\times) := (A^\times \otimes_{\mathbb{Z}} A^\times) / \langle a \otimes b + b \otimes a : a, b \in A^\times \rangle.$$

If we consider $S_{\mathbb{Z}}^2(A^\times)$ as a trivial \mathcal{G}_A -module, then

$$\bar{\lambda}_2 : \overline{\mathcal{RP}}(A) \rightarrow S_{\mathbb{Z}}^2(A^\times), \quad [a] \mapsto a \otimes (1 - a),$$

is a homomorphism of $\mathbb{Z}[\mathcal{G}_A]$ -modules. Hutchinson defined the $\mathbb{Z}[\mathcal{G}_A]$ -modules $\overline{\mathcal{RP}}_1(A)$ and $\overline{\mathcal{RB}}(A)$ as follows

$$\begin{aligned} \overline{\mathcal{RP}}_1(A) &:= \ker(\bar{\lambda}_1 : \overline{\mathcal{RP}}(A) \rightarrow \mathcal{I}_A^2), \\ \overline{\mathcal{RB}}(A) &:= \ker(\bar{\lambda}_2|_{\overline{\mathcal{RP}}_1(A)} : \overline{\mathcal{RP}}_1(A) \rightarrow S_{\mathbb{Z}}^2(A^\times)) \end{aligned}$$

(see [7, page 28], [8, Subsection 2.3]).

Let $Z_i(A^2) = \ker(\partial_i : Y_i(A^2) \rightarrow Y_{i-1}(A^2))$. From the complex $Y_4(A^2) \rightarrow Y_3(A^2) \rightarrow Z_2(A^2) \rightarrow 0$ we obtain the complex of \mathcal{G}_A -modules

$$Y_4(A^2)_{E_2(A)} \rightarrow Y_3(A^2)_{E_2(A)} \rightarrow Z_2(A^2)_{E_2(A)} \rightarrow 0.$$

The orbits of the action of $E_2(A)$ on $Y_3(A)$ and $Y_4(A)$ are represented by

$$\langle a \rangle [x]' := (\infty, \mathbf{0}, \mathbf{a}, \mathbf{ax}), \quad \text{and} \quad \langle a \rangle [x, y]' := (\infty, \mathbf{0}, \mathbf{a}, \mathbf{ax}, \mathbf{ay}), \quad \langle a \rangle \in \mathcal{G}_A, x, y, x/y \in \mathcal{W}_A,$$

respectively. Thus $Y_3(A^2)_{E_2(A)}$ is the free $\mathbb{Z}[\mathcal{G}_A]$ -module generated by the symbols $[x]'$, $x \in \mathcal{W}_A$ and $Y_4(A^2)_{E_2(A)}$ is the free $\mathbb{Z}[\mathcal{G}_A]$ -module generated by the symbols $[x, y]'$, $x, y, x/y \in \mathcal{W}_A$. It is straightforward to check that

$$\overline{\partial}_4([x, y]') = [x]' - [y]' + \langle x \rangle \left[\frac{y}{x} \right]' - \langle x^{-1} - 1 \rangle \left[\frac{1 - x^{-1}}{1 - y^{-1}} \right]' + \langle 1 - x \rangle \left[\frac{1 - x}{1 - y} \right]'$$

Thus we obtain a natural map

$$\eta : \overline{\mathcal{RP}}(A) \rightarrow H_0(E_2(A), Z_2(A^2)).$$

If $Y_\bullet(A) \rightarrow \mathbb{Z}$ is exact in dimension < 4 , then the above map becomes an isomorphism. We define

$$\mathcal{RP}(A) := H_0(E_2(A), Z_2(A^2)) = H_0(PE_2(A), Z_2(A^2)).$$

If A is a GE_2 -ring, then this definition of $\mathcal{RP}(A)$ coincides with the definition of Coronado and Hutchinson given in [5, §3].

The inclusion $Z_2(A^2) \rightarrow Y_2(A^2)$ induces the natural map

$$\lambda_1 := \text{inc} : \mathcal{RP}(A) = Z_2(A^2)_{E_2(A)} \rightarrow Y_2(A^2)_{E_2(A)} \simeq \mathbb{Z}[\mathcal{G}_A].$$

It is straightforward to check that $\bar{\lambda}_1 = \lambda_1 \circ \eta$. We denote the kernel of λ_1 by $\mathcal{RP}_1(A)$ and call it the *refined scissors congruence group of A* . Note that we have a natural map

$$\overline{\mathcal{RP}}_1(A) \rightarrow \mathcal{RP}_1(A).$$

Let $Y_\bullet(A)$ be exact in dimension 1. From the short exact sequence

$$0 \rightarrow Z_2(A^2) \rightarrow Y_2(A^2) \xrightarrow{\partial_2} Z_1(A^2) \rightarrow 0,$$

we obtain the (long) exact sequence

$$H_1(E_2(A), Y_2(A^2)) \xrightarrow{\bar{\partial}_2} H_1(E_2(A), Z_1(A^2)) \rightarrow \mathcal{RP}(A) \xrightarrow{\lambda_1} \mathbb{Z}[\mathcal{G}_A].$$

Since

$$H_1(E_2(A), Y_2(A^2)) \simeq H_1\left(E_2(A), \bigoplus_{\langle a \rangle \in \mathcal{G}_A} \text{Ind}_{\mu_2(A)}^{E_2(A)} \mathbb{Z}\right) \simeq \mu_2(A) \otimes_{\mathbb{Z}} \mathbb{Z}[\mathcal{G}_A],$$

we have the exact sequence

$$\mu_2(A) \otimes_{\mathbb{Z}} \mathbb{Z}[\mathcal{G}_A] \xrightarrow{\bar{\partial}_2} E_{2,1}^1 \rightarrow \mathcal{RP}_1(A) \rightarrow 0.$$

From the commutative diagram

$$\begin{array}{ccc} Y_2(A^2) & \xrightarrow{\partial_2} & Z_1(A^2) \\ \downarrow \partial_2 & & \downarrow \text{inc} \\ Y_1(A^2) & \xlongequal{\quad} & Y_1(A^2) \end{array}$$

we obtain the commutative diagram

$$\begin{array}{ccccc} \mu_2(A) \otimes_{\mathbb{Z}} \mathbb{Z}[\mathcal{G}_A] & \xrightarrow{\bar{\partial}_2} & E_{2,1}^1 & \longrightarrow & \mathcal{R}\mathcal{P}_1(A) \longrightarrow 0 \\ \downarrow & & \downarrow d_{2,1}^1 & & \\ \mu_2(A) & \xlongequal{\quad} & \mu_2(A) & & \end{array}$$

It is not difficult to see that $(d_{2,1}^1 \circ \bar{\partial}_2)(b \otimes \langle a \rangle) = b$. Thus from the above commutative diagram we obtain the exact sequence

$$\mu_2(A) \otimes_{\mathbb{Z}} \mathcal{I}_A \rightarrow E_{2,1}^2 \rightarrow \mathcal{R}\mathcal{P}_1(A) \rightarrow 0.$$

Consider the differential $d_{2,1}^2 : E_{2,1}^2 \rightarrow H_2(\mathbb{B}(A), \mathbb{Z})$. From now on we assume that the surjective map $p : \mathbb{B}(A) \rightarrow \mathbb{T}(A)$ induces the isomorphism

$$H_2(\mathbb{B}(A), \mathbb{Z}) \xrightarrow{p^*} H_2(\mathbb{T}(A), \mathbb{Z}).$$

Then the composite

$$\mu_2(A) \otimes_{\mathbb{Z}} \mathcal{I}_A \rightarrow E_{2,1}^2 \xrightarrow{d_{2,1}^2} H_2(\mathbb{B}(A), \mathbb{Z}) \simeq H_2(\mathbb{T}(A), \mathbb{Z}) \simeq A^\times \wedge A^\times$$

is given by $b \otimes \langle a \rangle \mapsto b \wedge a$ (see [14, Lemma 4.1]). Thus we have the commutative diagram

$$\begin{array}{ccccccc} \mu_2(A) \otimes_{\mathbb{Z}} \mathcal{I}_A & \longrightarrow & E_{2,1}^2 & \longrightarrow & \mathcal{R}\mathcal{P}_1(A) & \longrightarrow & 0 \\ \downarrow & & \downarrow d_{2,1}^2 & & \downarrow & & \\ 0 & \longrightarrow & \mu_2(A) \wedge A^\times & \longrightarrow & A^\times \wedge A^\times & \longrightarrow & \frac{A^\times \wedge A^\times}{\mu_2(A) \wedge A^\times} \longrightarrow 0. \end{array}$$

Consider the injective map

$$\alpha : \frac{A^\times \wedge A^\times}{A^\times \wedge \mu_2(A)} \rightarrow S_{\mathbb{Z}}^2(A^\times), \quad a \wedge b \mapsto 2(a \otimes b).$$

One can show that the composite map

$$\overline{\mathcal{R}\mathcal{P}}_1(A) \rightarrow \mathcal{R}\mathcal{P}_1(A) \rightarrow \frac{A^\times \wedge A^\times}{\mu_2(A) \wedge A^\times} \rightarrow S_{\mathbb{Z}}^2(A^\times)$$

coincides with $\bar{\lambda}_2|_{\overline{\mathcal{R}\mathcal{P}}_1(A)} : \overline{\mathcal{R}\mathcal{P}}_1(A) \rightarrow S_{\mathbb{Z}}^2(A^\times)$. We denote the kernel of

$$\mathcal{R}\mathcal{P}_1(A) \rightarrow \frac{A^\times \wedge A^\times}{\mu_2(A) \wedge A^\times} \rightarrow S_{\mathbb{Z}}^2(A^\times)$$

with $\mathcal{RB}(A)$ and call it the *refined Bloch group of A* . Note that we have a natural map

$$\overline{\mathcal{RB}}(A) \rightarrow \mathcal{RB}(A).$$

Now from the above commutative diagram we obtain a surjective map $E_{2,1}^\infty \rightarrow \mathcal{RB}(A)$. By an easy analysis of the main spectral sequence we obtain the surjective map

$$H_3(\mathbf{E}_2(A), \mathbb{Z}) \rightarrow E_{2,1}^\infty.$$

Combining these two maps we obtain a surjective map

$$H_3(\mathbf{E}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A).$$

This map factors through $H_3(\mathbf{PE}_2(A), \mathbb{Z})$, where

$$\mathbf{PE}_2(A) := \mathbf{E}_2(A) / \{\pm I_2\}.$$

We have proved the following version of a refined Bloch-Wigner exact sequence in [17].

Theorem 5.1. *Let A be a semilocal ring such that there is a ring homomorphism $A \rightarrow F$, F a field, where $\mu(A) \rightarrow \mu(F)$ is injective. For any maximal ideal $\mathfrak{m} \in \text{Specm}(A)$ let either A/\mathfrak{m} be infinite or if $|A/\mathfrak{m}| = p^d$, then $(p-1)d > 8$ ($(p-1)d > 6$ when A is a domain). If either $-1 \in A^{\times 2}$ or $|\mathcal{G}_A| \leq 4$, then we have the exact sequence*

$$0 \rightarrow \text{Tor}_1^{\mathbb{Z}}(\tilde{\mu}(A), \tilde{\mu}(A)) \rightarrow H_3(\mathbf{PE}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0,$$

where $\tilde{\mu}(A) = \mu(A)/\mu_2(A)$.

Proof. See [17, Theorem 3.1, Theorem 3.2, Corollary 3.3] and use Proposition 4.1 and [17, Lemma 2.8]. \square

Let \mathcal{A} be a finite cyclic group. If $2 \mid |\mathcal{A}|$, let \mathcal{A}^\sim denote the unique non-trivial extension of \mathcal{A} by $\mathbb{Z}/2$. If $2 \nmid |\mathcal{A}|$, we define $\mathcal{A}^\sim := \mathcal{A}$. Thus if $n = |\mathcal{A}|$, then

$$\mathcal{A}^\sim \simeq \mathbb{Z} / \text{gcd}(n^2, 2n).$$

Let \mathcal{A} be an ind-cyclic group, i.e. \mathcal{A} is direct limit of its finite cyclic subgroups. Then by passing to the limit we can define \mathcal{A}^\sim . Note that if \mathcal{A} has no 2-torsion, then $\mathcal{A}^\sim = \mathcal{A}$. There is always a canonical injective homomorphism $\mathcal{A} \hookrightarrow \mathcal{A}^\sim$ whose composition with the projection $\mathcal{A}^\sim \rightarrow \mathcal{A}$ coincides with multiplication by 2.

The connection between the third homology groups of $\mathbf{E}_2(A)$ and $\mathbf{PE}_2(A)$ is given in [19, Proposition 5.1].

Theorem 5.2. *Let A be a semilocal ring such that there is a ring homomorphism $A \rightarrow F$, F a field, where $\mu_2(A) \simeq \mu_2(F)$. If all the residue fields of A have more than three elements, then we have the exact sequence*

$$0 \rightarrow \mu_2(A)^\sim \rightarrow \frac{H_3(\mathbf{E}_2(A), \mathbb{Z})}{\mu_2(A) \otimes_{\mathbb{Z}} H_2(\mathbf{E}_2(A), \mathbb{Z})} \rightarrow H_3(\mathbf{PE}_2(A), \mathbb{Z}) \rightarrow 0,$$

where the map $\mu_2(A) \otimes_{\mathbb{Z}} H_2(\mathbf{E}_2(A), \mathbb{Z}) \rightarrow H_3(\mathbf{E}_2(A), \mathbb{Z})$ is induced by the product map $\mu_2(A) \times \mathbf{E}_2(A) \rightarrow \mathbf{E}_2(A)$, $(-1, X) \mapsto -X$.

Proof. We may assume that $\text{char}(F) \neq 2$. Since for any maximal ideal $\mathfrak{m} \in \text{Specm}(A)$, $|A/\mathfrak{m}| \geq 4$, there is $a \in A^\times$ such that $a^2 - 1 \in A^\times$. It follows from this that $M = A$. Thus by Theorem 2.1 we have $H_1(E_2(A), \mathbb{Z}) = 0$. Now by [19, Theorem 3.1] we obtain the desired exact sequence. \square

The above two theorems are results of our attempt to find an answer to the following question of Coronado and Hutchinson [5, p.3].

Question 5.3. Is it true that for any local domain A one has the exact sequence

$$0 \rightarrow \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^\sim \rightarrow H_3(\text{SL}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0?$$

The main result of this section is as follows, which generalizes the main result of [15, Theorem 5.1, Corollary 5.2].

Theorem 5.4. *Let A be a semilocal ring such that there is a ring homomorphism $A \rightarrow F$, F a field, where $\mu(A) \rightarrow \mu(F)$ is injective. For any maximal ideal $\mathfrak{m} \in \text{Specm}(A)$ let either A/\mathfrak{m} be infinite or if $|A/\mathfrak{m}| = p^d$, then $(p-1)d > 8$ ($(p-1)d > 6$ when A is a domain). If either $-1 \in A^{\times 2}$ or $|\mathcal{G}_A| \leq 4$, then we have the exact sequence*

$$0 \rightarrow \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^\sim \rightarrow \frac{H_3(E_2(A), \mathbb{Z})}{\mu_2(A) \otimes_{\mathbb{Z}} I^2(A)} \rightarrow \mathcal{RB}(A) \rightarrow 0,$$

where $(-1) \otimes \langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \in \mu_2(A) \otimes_{\mathbb{Z}} I^2(A)$ maps in $H_3(E_2(A), \mathbb{Z})$ to the class of the cycle which is obtained from the shuffle product of $[-1] \in B_1(\mu_2(A))_{\mu_2(A)}$ with the cycle $[D(a)|D(b)] + R_{ab} - R_a - R_b + R_1 \in B_2(E_2(A))_{E_2(A)}$.

Proof. Consider the surjective maps

$$\frac{H_3(E_2(A), \mathbb{Z})}{\mu_2(A) \otimes_{\mathbb{Z}} H_2(E_2(A), \mathbb{Z})} \twoheadrightarrow H_3(\text{PE}_2(A), \mathbb{Z}) \twoheadrightarrow \mathcal{RB}(A).$$

Let \mathcal{K} be the kernel of this composite. Then we have the commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{K} & \longrightarrow & \frac{H_3(E_2(A), \mathbb{Z})}{\mu_2(A) \otimes_{\mathbb{Z}} H_2(E_2(A), \mathbb{Z})} & \longrightarrow & \mathcal{RB}(A) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \text{Tor}_1^{\mathbb{Z}}(\tilde{\mu}(A), \tilde{\mu}(A)) & \longrightarrow & H_3(\text{PE}_2(A), \mathbb{Z}) & \longrightarrow & \mathcal{RB}(A) \longrightarrow 0 \end{array}$$

(see Theorem 5.1). From this diagram and Theorem 5.2 we obtain the exact sequence

$$0 \rightarrow \mu_2(A)^\sim \rightarrow \mathcal{K} \rightarrow \text{Tor}_1^{\mathbb{Z}}(\tilde{\mu}(A), \tilde{\mu}(A)) \rightarrow 0.$$

This clearly implies that $\mathcal{K} \simeq \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^\sim$ and thus we have the exact sequence

$$0 \rightarrow \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^\sim \rightarrow \frac{H_3(E_2(A), \mathbb{Z})}{\mu_2(A) \otimes_{\mathbb{Z}} H_2(E_2(A), \mathbb{Z})} \rightarrow \mathcal{RB}(A) \rightarrow 0.$$

By Proposition 4.1, $H_2(B(A), \mathbb{Z}) \simeq H_2(T(A), \mathbb{Z})$. This shows that the homology class of the cycle $([E_{12}(x)|E_{12}(y)] - [E_{12}(y)|E_{12}(x)])$, $x, y \in A$, in $H_2(E_2(A), \mathbb{Z})$ is trivial (see

Theorem 3.4). Now consider the homology class of the cycle $([D(a)|D(b)] - [D(b)|D(a)])$, $a, b \in A^\times$. This is the image of $D(a) \wedge D(b)$ under the map

$$\mathrm{T}(A) \wedge \mathrm{T}(A) \simeq H_2(\mathrm{T}(A), \mathbb{Z}) \rightarrow H_2(\mathrm{E}_2(A), \mathbb{Z}).$$

Now from the commutative diagram

$$\begin{array}{ccccc} \mu_2(A) \otimes_{\mathbb{Z}} (\mathrm{T}(A) \wedge \mathrm{T}(A)) & \longrightarrow & \mu_2(A) \otimes_{\mathbb{Z}} H_2(\mathrm{T}(A), \mathbb{Z}) & \longrightarrow & \mu_2(A) \otimes_{\mathbb{Z}} H_2(\mathrm{E}_2(A), \mathbb{Z}) \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{T}(A) \wedge \mathrm{T}(A) \wedge \mathrm{T}(A) & \longrightarrow & H_3(\mathrm{T}(A), \mathbb{Z}) & \longrightarrow & H_3(\mathrm{E}_2(A), \mathbb{Z}) \end{array}$$

it follows that the right side vertical map is given by

$$(-1) \otimes (D(a) \wedge D(b)) \mapsto D(-1) \wedge D(a) \wedge D(b).$$

Note that here we denoted the image of $D(-1) \wedge D(a) \wedge D(b) \in \bigwedge_{\mathbb{Z}}^3 \mathrm{T}(A)$ in $H_3(\mathrm{E}_2(A), \mathbb{Z})$ again by $D(-1) \wedge D(a) \wedge D(b)$. If $i^2 = -1$, then

$$D(-1) \wedge D(a) \wedge D(b) = 2(D(i) \wedge D(a) \wedge D(b)) = 0.$$

Let $|\mathcal{G}_A| \leq 4$. If $-1 \in A^{\times 2}$, we proceed as in above. If $-1 \notin A^{\times 2}$, then $\mathcal{G}_A = \{1, -1, u, -u\}$ for some $u \in A^\times$. So we may take $a, b \in \{1, -1, u, -u\}$ and in any case

$$D(-1) \wedge D(a) \wedge D(b) = 0.$$

Thus by Theorem 3.4, we have

$$\frac{H_3(\mathrm{E}_2(A), \mathbb{Z})}{\mu_2(A) \otimes_{\mathbb{Z}} H_2(\mathrm{E}_2(A), \mathbb{Z})} = \frac{H_3(\mathrm{E}_2(A), \mathbb{Z})}{\mu_2(A) \otimes_{\mathbb{Z}} I^2(A)}.$$

This completes the proof of the theorem. \square

In order to have a positive answer to Question 5.3, we should have a positive answer to the following question, which was raised in [19].

Question 5.5. Let A be a local domain such that $|A/\mathfrak{m}_A| \neq 2$. Is the product map

$$\rho_* : \mu_2(A) \otimes_{\mathbb{Z}} H_2(\mathrm{E}_2(A), \mathbb{Z}) \rightarrow H_3(\mathrm{E}_2(A), \mathbb{Z})$$

trivial?

We do not know the answer of the above question even over a general infinite field. For more on this we refer the interested readers to [19, Example 4.4, Remark 5.2, Proposition 5.4].

Remark 5.6. Let A be a local domain such that $|A/\mathfrak{m}_A| \neq 2, 3, 4, 5, 7, 8, 9, 16, 27, 32, 64$. Then the results of this paper suggest that to give a positive answer to Question 5.3, one may show that

(i) we have the projective Bloch-Wigner exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\tilde{\mu}(A), \tilde{\mu}(A)) \rightarrow H_3(\mathrm{PE}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0,$$

(ii) the product map $\rho_* : \mu_2(A) \otimes_{\mathbb{Z}} H_2(\mathrm{E}_2(A), \mathbb{Z}) \rightarrow H_3(\mathrm{E}_2(A), \mathbb{Z})$ is trivial.

REFERENCES

- [1] Atiyah, M. F., MacDonald, I. G. Introduction To Commutative Algebra. Addison-Wesley series in mathematics, Avalon Publishing (1994) [8](#)
- [2] Brown, K. S. Cohomology of Groups. Graduate Texts in Mathematics, 87. Springer-Verlag, New York (1994) [1](#), [3](#)
- [3] Cohn, P. M. On the structure of the GL_2 of a ring. Inst. Hautes Études Sci. Publ. Math. **30** (1966), 5–33 [1](#), [2](#)
- [4] Cohn, P. M. A presentation of SL_2 for Euclidean imaginary quadratic number fields. Mathematika **15** no 2, (1968) 156–163 [1](#)
- [5] Coronado, R. C., Hutchinson, K. Bloch Groups of Rings. <https://arxiv.org/abs/2201.04996> [13](#), [16](#)
- [6] Dennis, R. K., Stein, M. R. The functor K_2 : a survey of computations and problems, in: Lecture Notes in Math. **342**, Springer (1973), 243–280. [7](#)
- [7] Hutchinson, K. A Bloch-Wigner complex for SL_2 . J. K -Theory **12** (2013), no. 1, 15–68 [12](#)
- [8] Hutchinson, K. A refined Bloch group and the third homology of SL_2 of a field. J. Pure Appl. Algebra **217** (2013), 2003–2035 [12](#)
- [9] Hutchinson, K. The second homology of SL_2 of S -integers. Journal of Number Theory **159** (2016), 223–272 [10](#), [11](#)
- [10] Hutchinson, K. The third homology of SL_2 of local rings. Journal of Homotopy and Related Structures **12**, 931–970 (2017) [12](#)
- [11] Hutchinson, K. GE_2 -rings and a graph of unimodular rows. J. Pure Appl. Algebra **226**, (2022), no. 10, 107074 [3](#), [7](#)
- [12] Menal, R. Remarks on the GL_2 of a ring. Journal of Algebra **61** (1979), 335–359 [8](#)
- [13] Mirzaii, B. A Bloch-Wigner exact sequence over local rings. Journal of Algebra **476**, 459–493 (2017) [3](#), [12](#)
- [14] Mirzaii, B., Torres Pérez, E. A refined scissors congruence group and the third homology of SL_2 . Journal of Pure and Applied Algebra **228** (2024), no. 6, 107615, 1–28 [1](#), [11](#), [14](#)
- [15] Mirzaii, B., Torres Pérez, E. A refined Bloch-Wigner exact sequence in characteristic 2. Journal of Algebra **657** (2024), no. 1, 141–158 [1](#), [2](#), [16](#)
- [16] Mirzaii, B., Torres Pérez, E. The abelianization of the elementary group of rank two. Available at <https://arxiv.org/abs/2401.06330> [1](#), [5](#), [8](#)
- [17] Mirzaii, B., Torres Pérez, E. The third homology of projective special linear group of rank 2. Available at <https://arxiv.org/abs/2401.04341>. [15](#)
- [18] Mirzaii, B., Torres Pérez, E. The low dimensional homology of projective linear group of rank two. Available at <https://arxiv.org/abs/2405.08950> [3](#)
- [19] Mirzaii, B., Torres Pérez, E. On the connections between the low dimensional homology groups of SL_2 and PSL_2 . Available at <https://arxiv.org/abs/2402.08074> [2](#), [15](#), [16](#), [17](#)
- [20] Silvester, J. R. On the GL_n of a semi-local ring. Algebraic K -Theory, Lecture Notes in Mathematics, Vol **966** (1982), 244–260 [2](#)
- [21] Weibel, C. A. An Introduction to Homological Algebra. Cambridge Studies in Advanced Mathematics, 38. Cambridge University Press, Cambridge, (1994). [8](#)

Instituto de Ciências Matemáticas e de Computação (ICMC), Universidade de São Paulo, São Carlos, Brasil
Email address: bmirzaii@icmc.usp.br
Email address: elvistorres@usp.br