

# THE AFFINE SUBSPACE CONCENTRATION INEQUALITY FOR CENTERED CONVEX BODIES

KATHARINA ELLER AND ANSGAR FREYER

ABSTRACT. An affine version of the linear subspace concentration inequality as proposed by Wu in [11] is established for centered convex bodies. This generalizes results from [11] and [8] on polytopes to convex bodies.

## 1. INTRODUCTION

Let  $\mathcal{K}^n$  denote the set of convex bodies in  $\mathbb{R}^n$ , i.e., the family of all convex and compact subsets  $K \subset \mathbb{R}^n$  with non-empty interior. The subfamily of convex bodies containing the origin in their interior is denoted by  $\mathcal{K}_{(o)}^n$  and the subset of origin-symmetric convex bodies, i.e., the sets  $K \in \mathcal{K}^n$  satisfying  $K = -K$ , is denoted by  $\mathcal{K}_c^n$ . A convex body  $K \in \mathcal{K}^n$  is called centered if its centroid  $c(K)$  is located at the origin, i.e.,

$$c(K) = \frac{1}{\text{vol}_n(K)} \int_K x \, d\mathcal{H}^n(x) = 0,$$

where  $\mathcal{H}^n$  denotes the  $n$ -dimensional Hausdorff measure. We denote by  $\text{vol}_n$  the  $n$ -dimensional volume functional, i.e., the restriction of  $\mathcal{H}^n$  to the set of  $n$ -dimensional convex bodies. The set of all centered convex bodies in  $\mathbb{R}^n$  is denoted by  $\mathcal{K}_c^n$ . For  $x, y \in \mathbb{R}^n$  let  $\langle x, y \rangle$  denote the standard inner product on  $\mathbb{R}^n$ , and  $|x| = \sqrt{\langle x, x \rangle}$  the Euclidean norm of  $x$ .

The study of geometric measures related to convex bodies is a corner stone of convex geometry. One of the central problems in the classical Brunn-Minkowski theory is the Minkowski-Christoffel problem asking for necessary and sufficient conditions characterizing the surface area measures of a convex body among the finite Borel measures on the sphere. For a definition of the surface area measures and an overview of the Minkowski-Christoffel problem we refer to [10, Chapter 8]. In addition to the surface area measures, another important geometric measure is the cone volume measure  $V_K$  of a convex body  $K \in \mathcal{K}_{(o)}^n$ . It is defined for a Borel set  $\omega \subseteq \mathbb{S}^{n-1}$  of the sphere by

$$V_K(\omega) = \frac{1}{n} \int_{\nu_K^{-1}(\omega)} \langle x, \nu_K(x) \rangle \, d\mathcal{H}^{n-1}(x),$$

where  $\nu_K$  is the Gauss map of  $K$ , i.e.,  $\nu_K$  is defined on the regular boundary points  $\text{bd}'(K)$  of  $K$  and assigns each such boundary point its unique outer unit normal vector. The interest in the cone volume measure also stems from the fact that it is a particular instance of an  $L_p$  surface area measure (see [10, Section 9.2]) that allows for a geometric interpretation. The geometry behind the cone volume measure is best explained for polytopes: Let  $P = \{x \in \mathbb{R}^n : \langle a_i, x \rangle \leq 1, 1 \leq i \leq m\}$  be a polytope in  $\mathcal{K}_{(o)}^n$ . We assume that the

inequalities  $\langle a_i, x \rangle \leq 1$  are irredundant, i.e., each of them defines a proper facet  $F_i$  of  $P$ . If we denote by  $C_i \subset P$  the convex hull of the facet  $F_i$  with the origin (a “cone” in  $P$ ), one can verify that

$$V_P(\omega) = \sum_{i: \frac{a_i}{|a_i|} \in \omega} \text{vol}(C_i)$$

holds for all  $\omega \subset \mathbb{S}^{n-1}$ . As in the case of the surface area measures, it is natural to ask for necessary and sufficient conditions on a Borel measure  $\mu$  to be the cone volume measure of a convex body  $K \in \mathcal{K}_{(o)}^n$ , which leads to the log-Minkowski problem. For *even* measures, i.e., measures  $\mu$  with  $\mu(-\omega) = \mu(\omega)$ , this question was settled by Böröczky, Lutwak, Yang and Zhang in [5]. They showed that an even non-zero finite Borel measure  $\mu$  on  $\mathbb{S}^{n-1}$  can be represented as  $\mu = V_K$  for some  $K \in \mathcal{K}_e^n$ , if and only if  $\mu$  satisfies the (linear) subspace concentration condition, i.e., we have

$$(1.1) \quad \mu(L \cap \mathbb{S}^{n-1}) \leq \frac{\dim(L)}{n} \mu(\mathbb{S}^{n-1}),$$

for all proper linear subspaces  $L \subset \mathbb{R}^n$ , and whenever equality holds in (1.1) for some  $L$  then there exists a complementary linear subspace  $L'$  such that  $\mu$  is concentrated on  $(L \cup L') \cap \mathbb{S}^{n-1}$ .

In the general case, the log-Minkowski problem is still open, although further results were obtained in recent years [1, 2, 3, 4, 7, 9, 12]. For an overview on the state of the art regarding the cone volume measure and related problems we refer to the survey article [6]. In particular, it was shown by Henk and Linke in [9] that the cone volume measures of centered polytopes satisfy the subspace concentration condition. This was generalized to arbitrary centered convex bodies Böröczky and Henk in [3]. We state the inequality here.

**Theorem 1.1** ([3]). *Let  $K \in \mathcal{K}_e^n$ , then*

$$(1.2) \quad V_K(L \cap \mathbb{S}^{n-1}) \leq \frac{\dim(L)}{n} V_K(\mathbb{S}^{n-1}),$$

*holds for all linear subspaces  $L \subset \mathbb{R}^n$ .*

Recently, Wu formulated an affine version of the subspace concentration condition for polytopes [11]: Again, let  $P = \{x \in \mathbb{R}^n : \langle a_i, x \rangle \leq 1, 1 \leq i \leq m\}$ , where each vector  $a_i$  defines a facet of  $P$ , and let  $C_i$ ,  $1 \leq i \leq m$ , be the corresponding cones in  $P$ . Then  $P$  is said to satisfy the affine subspace concentration condition, if

$$(1.3) \quad \sum_{i: a_i \in A} \text{vol}(C_i) \leq \frac{\dim(A) + 1}{n + 1} \text{vol}(P)$$

holds for all proper affine subspaces  $A \subset \mathbb{R}^n$  and, moreover, equality holds, if and only if there exists a complementary affine subspace  $A' \subset \mathbb{R}^n$  with  $\{a_1, \dots, a_m\} \subset A \cup A'$ . Here, the term “complementary” is meant in the sense of affine spaces, i.e.,  $A \cap A' = \emptyset$  and  $\text{aff}(A \cup A') = \mathbb{R}^n$ .

In [11] the affine subspace concentration condition is proven for polytopes that are in addition smooth and reflexive lattice polytopes. We shall not give the definition here, but refer to [11] instead. It seems that the extra condition

of  $P$  being a smooth reflexive lattice polytope might not be necessary. In [8] it is shown that the inequality (1.3) holds for all centered polytopes and all affine subspaces. However, the equality cases remain open in general.

Note that the normalization of the inequalities  $\langle a_i, x \rangle \leq 1$  determining  $P$  to have right hand side 1 is crucial in the affine subspace concentration condition. Renormalizing the vectors  $a_i$  would create new affine dependencies and thus a different type of affine subspace concentration condition. In our setting the normalization is natural in that the vectors  $a_i$  are the vertices of the polar polytope  $P^*$ . Since the cone volume measure  $V_P$  is supported on the vectors  $\frac{a_i}{|a_i|}$ , we can restate (1.3) in terms of the cone volume measure as follows:

$$V_P(\pi(\text{bd}(P^*) \cap A)) \leq \frac{\dim(A)}{n+1} V_P(\mathbb{S}^{n-1}),$$

where  $\pi: \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{S}^{n-1}$  denotes the central projection to the sphere. Presented this way, the inequality could be stated for any convex body in  $K \in \mathcal{K}_{(o)}^n$ , where  $K^* = \{a \in \mathbb{R}^n: \langle a, x \rangle \leq 1, \forall x \in K\}$  is the polar convex body. We show that it indeed extends to centered convex bodies.

**Theorem 1.2.** *Let  $K \in \mathcal{K}_{(o)}^n$  be centered and let  $A \subset \mathbb{R}^n$  be an affine subspace. Then it holds*

$$V_K(\pi(\text{bd}(K^*) \cap A)) \leq \frac{\dim(A)}{n+1} V_K(\mathbb{S}^{n-1}).$$

In the proof we make use of a lifting argument from [8] for polytopes that we adapt to convex bodies. Moreover, we provide characterizations of the equality case in Theorem 1.2 for two special cases in Section 4.

## 2. PRELIMINARIES AND NOTATION

For a comprehensive overview of the basic concepts and definitions in convex geometry we refer to [10]. We write  $\mathbb{S}^{n-1}$  for the  $(n-1)$ -dimensional Euclidean unit sphere, i.e.,  $\mathbb{S}^{n-1} = \{x \in \mathbb{R}^n : |x| = 1\}$ . For a convex body  $K$  let  $\text{bd}(K)$  denote its boundary points.

For a given convex body  $K \in \mathcal{K}^n$  the *support function*  $h_K: \mathbb{R}^n \rightarrow \mathbb{R}$  is defined by

$$h_K(u) = \max_{x \in K} \langle u, x \rangle.$$

The support function is continuous and positively homogeneous of degree 1. The hyperplane  $H_K(u) = \{x \in \mathbb{R}^n : \langle u, x \rangle = h_K(u)\}$  is a supporting hyperplane of  $K$  and for a boundary point  $v \in \text{bd} K \cap H_K(u)$ , the vector  $u$  will be called an *outer normal vector*. If in addition  $u \in \mathbb{S}^{n-1}$  then  $u$  is an *outer unit normal vector*. Let  $\text{bd}'(K) \subseteq \text{bd}(K)$  be the set of all boundary points having a unique outer unit normal vector, it is called the *regular boundary*. We remark that the set of boundary points not having a unique outer normal vector has measure zero, that is  $\mathcal{H}^{n-1}(\text{bd}(K) \setminus \text{bd}'(K)) = 0$ . The *Gauss map* (or *spherical image map*)  $\nu_K: \text{bd}'(K) \rightarrow \mathbb{S}^{n-1}$  maps a point  $x$  to its unique outer unit normal vector.

The *central projection*  $\pi_k: \mathbb{R}^k \setminus \{0\} \rightarrow \mathbb{S}^{k-1}$  is given by  $\pi_k(x) = |x|^{-1}x$ . We simply write  $\pi$  instead of  $\pi_k$  if the ambient dimension is clear from the context.

The cone volume can also be expressed as (cf. Section 9.2 in [10])

$$V_K(\omega) = \frac{1}{n} \int_{\omega} h_K(u) \, dS_K(u)$$

where  $S_K(\omega) = \mathcal{H}^{n-1}(\nu_K^{-1}(\omega))$  denotes the surface area measure. We will also use a description of the cone volume measure in terms of the volume of sets that we call *star pyramids*. A star pyramid  $[Q, v]$  with apex  $v$  and (not necessarily convex) base  $Q$  is the union of closed segments from  $Q$  to  $v$ , i.e.,  $[Q, v] = \bigcup_{x \in Q} [x, v]$ . If  $Q$  is convex, we have  $[Q, v] = \text{conv}(Q \cup \{v\})$ .

**Lemma 2.1** (Lemma 9.2.4 in [10]). *Let  $K \in \mathcal{K}_{(o)}^n$  and  $\omega \subset \mathbb{S}^{n-1}$  be a Borel set. Then,*

$$V_K(\omega) = \mathcal{H}^n([\nu_K^{-1}(\omega), 0]).$$

### 3. AFFINE SUBSPACE CONCENTRATION FOR CENTERED CONVEX BODIES

The volume formula for pyramids extends to star pyramids as introduced in Section 2.

**Lemma 3.1.** *Let  $Q \subset \mathbb{R}^{n-1}$ ,  $h \neq 0$ . Then it holds*

$$\mathcal{H}^n([Q \times \{h\}, 0]) = \frac{h}{n} \mathcal{H}^{n-1}(Q).$$

*Proof.* Note that it holds  $[Q \times \{h\}, 0] \cap \{x \in \mathbb{R}^n : x_n = t\} = \frac{t}{h}Q \times \{t\}$ . Then we obtain by Cavalieri's principle and the translation invariance of the Hausdorff measure

$$\begin{aligned} \mathcal{H}^n([Q \times \{h\}, 0]) &= \int_0^h \mathcal{H}^{n-1}([Q \times \{h\}, 0] \cap \{x \in \mathbb{R}^n : x_n = t\}) \, dt \\ &= \int_0^h \left(\frac{t}{h}\right)^{n-1} \mathcal{H}^{n-1}(Q) \, dt = \frac{h}{n} \mathcal{H}^{n-1}(Q). \quad \square \end{aligned}$$

For a set  $Q \subset \mathbb{R}^k$  we will denote  $\widehat{Q} := Q \times \{1\} \subset \mathbb{R}^{k+1}$  or  $(Q)^\wedge$  for longer arguments. For  $K \in \mathcal{K}_{(o)}^n$  and  $j \geq 1$  we define pyramids by

$$(3.1) \quad K^{(j)} = [(K^{(j-1)})^\wedge, -(n+j)e_{n+j}]$$

where  $K^{(0)} = K$ . The pyramids  $K^{(j)}$  are convex bodies in  $\mathbb{R}^{n+j}$ .

**Lemma 3.2.** *Let  $K \in \mathcal{K}_{(o)}^n$ . Then the following holds:*

- i) *If  $K$  is centered,  $K^{(1)}$  is centered.*
- ii)  $\text{vol}_{n+1}(K^{(1)}) = \frac{n+2}{n+1} \text{vol}_n(K)$ .
- iii) *If  $y \in \text{bd}(K^*)$  is an outer normal vector of  $x \in K$ , then  $(\frac{n+2}{n+1}y, -\frac{1}{n+1})^\top \in \text{bd}((K^{(1)})^*)$  is an outer normal vector of  $K^{(1)}$  at each point in  $[(x, 1)^\top, -(n+1)e_{n+1}]$ .*
- iv) *A regular boundary point of  $K^{(1)}$  is either in the relative interior of  $\widehat{K}$ , or of the form  $\mu(x, 1)^\top - (1-\mu)(n+1)e_{n+1}$  for some  $x \in \text{bd}'(K)$ ,  $\mu \in (0, 1)$ .*

*Proof.* The first statement is another application of Cavalieri's principle (see [8, Lemma 2.1] for details). The second statement follows directly from Lemma 3.1.

For the third statement let  $\bar{y} = (\frac{n+2}{n+1}y, -\frac{1}{n+1})^\top$  and  $\hat{z} = (z, 1)^\top$ , where  $z \in K$  as well as  $\mu \in [0, 1]$ . Then we have

$$\langle \bar{y}, \mu \hat{z} - (1 - \mu)(n + 1)e_{n+1} \rangle = \mu \frac{n+2}{n+1} \langle y, z \rangle - \mu \frac{1}{n+1} + (1 - \mu).$$

Since  $y \in \text{bd}(K)$  we see that

$$\langle \bar{y}, \mu \hat{z} - (1 - \mu)(n + 1)e_{n+1} \rangle \leq 1$$

with equality for  $z = x$ , which proves the claim. The last statement is a direct consequence of iii).  $\square$

To keep track of the part of the boundary corresponding to an affine subspace  $A \subset \mathbb{R}^n$  within the sequence of pyramids  $K^{(j)}$  we introduce the following notation for  $j \geq 1$

$$(3.2) \quad F_{K,A}^{(j)} = [(F_{K,A}^{(j-1)})^\wedge, -(n+j)e_{n+j}] \subset K^{(j)}$$

where

$$F_{K,A}^{(0)} = \nu_K^{-1}(\pi_n(\text{bd}(K^\star) \cap A)).$$

The corresponding ‘‘cones’’ are star pyramids defined by

$$C_{K,A}^{(j)} = [F_{K,A}^{(j)}, 0]$$

for  $j \geq 0$ .

**Lemma 3.3.** *Let  $K \in \mathcal{K}_{(o)}^n$  and  $A \subset \mathbb{R}^n$  be an affine subspace. Then for  $j \geq 1$ ,*

$$\mathcal{H}^{n+j}(C_{K,A}^{(j)}) = \mathcal{H}^n(C_{K,A}^{(0)}).$$

*Proof.* First we show  $\mathcal{H}^{n+1}(C_{K,A}^{(1)}) = \mathcal{H}^n(C_{K,A}^{(0)})$ . Set  $G_1 = [(C_{K,A}^{(0)})^\wedge, -(n+1)e_{n+1}]$  and  $G_2 = [(C_{K,A}^{(0)})^\wedge, 0]$ . We claim that

$$\mathcal{H}^{n+1}(C_{K,A}^{(1)}) = \mathcal{H}^{n+1}(G_1) - \mathcal{H}^{n+1}(G_2).$$

Because  $[(F_{K,A}^{(0)})^\wedge, 0]$  has  $\mathcal{H}^{n+1}$ -measure zero - as it is part of the boundary of the convex body  $[\widehat{K}, 0]$  - it suffices to show

$$C_{K,A}^{(1)} = (G_1 \setminus G_2) \cup [(F_{K,A}^{(0)})^\wedge, 0].$$

Let  $p \in C_{K,A}^{(1)}$ , then by construction there exists  $x \in (F_{K,A}^{(0)})^\wedge$  and  $\lambda, \beta \in [0, 1]$  with  $p = \beta(\lambda x + (1 - \lambda)(- (n + 1)e_{n+1}))$ . Let  $E = \text{lin}(\{x, -(n + 1)e_{n+1}\})$ , this is a two dimensional subspace with  $p \in E$ . In fact  $p$  is contained in the triangle  $\text{conv}(\{0, -(n + 1)e_{n+1}, x\})$ . So if  $p \notin [0, x]$  then it holds  $p \notin \text{conv}(\{0, e_{n+1}, x\})$ . We obtain

$$p \in \text{conv}(\{0, -(n + 1)e_{n+1}, x\}) \setminus \text{conv}(\{0, e_{n+1}, x\}) \subseteq G_1 \setminus G_2.$$

On the other hand let  $p \in G_1 \setminus G_2$ . Then there exists some  $x \in (F_{K,A}^{(0)})^\wedge$  such that  $p \in \text{conv}(\{-(n+1)e_{n+1}, e_{n+1}, x\})$  and also  $p \notin \text{conv}(\{0, e_{n+1}, x\})$ . In particular, there exist  $\lambda, \mu \in [0, 1]$  with

$$\begin{aligned} p &= \lambda x + (1 - \lambda)(\mu(-(n+1)e_{n+1}) + (1 - \mu) \cdot 0) \\ &= (\lambda + (1 - \lambda)\mu) \left( \frac{\lambda}{\lambda + (1 - \lambda)\mu} x + \frac{(1 - \lambda)\mu}{\lambda + (1 - \lambda)\mu} (-(n+1)e_{n+1}) \right). \end{aligned}$$

This implies  $p \in C_{K,A}^{(1)}$ . Furthermore by construction we have  $(F_{K,A}^{(0)})^\wedge \subseteq F_{K,A}^{(1)}$  and so  $[(F_{K,A}^{(0)})^\wedge, 0] \subseteq C_{K,A}^{(1)}$ . This yields the claim.

Using Lemma 3.2 we obtain

$$\begin{aligned} \mathcal{H}^{n+1}(C_{K,A}^{(1)}) &= \mathcal{H}^{n+1}(G_1) - \mathcal{H}^{n+1}(G_2) \\ &= \frac{n+2}{n+1} \mathcal{H}^n((C_{K,A}^{(0)})^\wedge) - \frac{1}{n+1} \mathcal{H}^n((C_{K,A}^{(0)})^\wedge) = \mathcal{H}^n(C_{K,A}^{(0)}). \end{aligned}$$

Analogously one can prove  $\mathcal{H}^{n+j}(C_{K,A}^{(j)}) = \mathcal{H}^{n+j-1}(C_{K,A}^{(j-1)})$  for  $j \geq 1$ . This yields the desired statement.  $\square$

*Proof of Theorem 1.1.* For any  $k \in \mathbb{Z}_{\geq 1}$ , we consider the embedding

$$\varphi_k : \mathbb{R}^k \rightarrow \mathbb{R}^{k+1}, \quad \varphi_k(x) = \begin{pmatrix} \frac{k+2}{k+1}x \\ \frac{1}{k+1} \end{pmatrix}$$

and we write  $\psi_{n+j} := \varphi_{n+j-1} \circ \dots \circ \varphi_n$ , where  $j \geq 1$ . For  $j = 0$  we define  $\psi_n$  to be the identity on  $\mathbb{R}^n$ . We seek to apply the linear subspace concentration inequality to the convex bodies  $K^{(j)}$  (introduced in (3.1)) and the linear spaces  $L^{(j)} := \text{lin}(\psi_{n+j}(A)) \subseteq \mathbb{R}^{n+j}$ . Note that we have  $\dim(L^{(j)}) = \dim(A) + 1$  for all  $j > 0$ .

Here we write  $F^{(j)} := F_{K,A}^{(j)}$  for the boundary part of the  $j$ -th pyramid associated to  $A$  (cf. (3.2)). By Lemmas 2.1 and 3.3 we have

$$(3.3) \quad \mathbb{V}_K(\pi_n(\text{bd}(K^*) \cap A)) = \mathcal{H}^n([F^{(0)}, 0]) = \mathcal{H}^{n+j}([F^{(j)}, 0]).$$

It follows from Lemma 3.2 iii) that

$$(3.4) \quad \psi_{n+j}(\text{bd } K^*) \subseteq \text{bd}((K^{(j)})^*)$$

holds for all  $j \geq 0$ . We claim that

$$(3.5) \quad F^{(j)} \cap \text{bd}'(K^{(j)}) \subseteq \nu_{K^{(j)}}^{-1}(\pi_{n+j} \circ \psi_{n+j}(\text{bd}(K^*) \cap A)).$$

For  $j = 0$ , this trivially follows from the definition of  $F^{(0)}$ . For  $j > 0$  we argue by induction. Let  $x \in F^{(j)} \cap \text{bd}'(K^{(j)})$ . Then  $x$  can be expressed as

$$x = \mu \hat{y} + (1 - \mu)(-(n+j)e_{n+j}),$$

where  $\mu \in (0, 1)$ ,  $y \in F^{(j-1)} \cap \text{bd}'(K^{(j-1)})$  and  $\hat{y} = (y, 1)^\top \in \mathbb{R}^{n+j}$  (cf. Lemma 3.2). By induction, there exists an outer normal vector of  $K^{(j-1)}$  at  $y$  of the form  $\psi_{n+j-1}(a)$ , where  $a \in \text{bd}(K^*) \cap A$ . By (3.4), we know that  $\psi_{n+j-1}(a) \in \text{bd}((K^{(j-1)})^*)$ . Hence, Lemma 3.2 iii) shows that  $\psi_{n+j}(a)$  is the unique (up to positive multiples) outer normal vector of  $K^{(j)}$  at  $x$  and (3.5) is proven.

It follows from (3.5) and the fact that  $L^{(j)}$  is a linear space that

$$F^{(j)} \cap \text{bd}'(K^{(j)}) \subseteq \nu_{K^{(j)}}^{-1}(L^{(j)} \cap \mathbb{S}^{n+j-1}).$$

Since the Hausdorff measure of the non-regular boundary points of  $K^{(j)}$  is zero, we obtain from (3.3) that

$$V_K(\pi_n(\text{bd}(K^*) \cap A)) \leq V_{K^{(j)}}(L^{(j)} \cap \mathbb{S}^{n+j-1}).$$

Since  $K^{(j)}$  is centered and  $L^{(j)}$  is  $(\dim(A) + 1)$ -dimensional, the linear subspace concentration inequality (cf. Theorem 1.1) yields

$$\begin{aligned} V_K(\pi_n(\text{bd}(K^*) \cap A)) &\leq \frac{\dim(A) + 1}{n + j} \text{vol}_{n+j}(K^{(j)}) \\ &= \frac{\dim(A) + 1}{n + j} \cdot \frac{n + j + 1}{n + 1} \text{vol}_{n+j}(K), \end{aligned}$$

where we used Lemma 3.2 to obtain the last line. Taking the limit  $j \rightarrow \infty$  now yields the desired inequality.  $\square$

#### 4. EQUALITY CASES

Here we extend the characterization of the equality case in Theorem 1.2 to the convex body setting. The corresponding results for centered polytopes are shown in [8].

**Theorem 4.1.** *Let  $K \in \mathcal{K}_c^n$ .*

- i) *If  $A = \{a\}$ , then equality holds in (1.2) if and only if  $K$  is a pyramid with base  $Q = H_K(a) \cap K$ .*
- ii) *If  $A$  is a supporting hyperplane of  $K^*$  spanned by boundary point of  $K^*$ , then equality holds in (1.2) if and only if  $K$  is a pyramid with apex  $v_A$ , where  $v_A$  is the unique point in  $K$  with  $\langle v, a \rangle = 1$  for all  $a \in A$ .*

Note that the point  $v_A$  from ii) indeed exists. Since  $A$  does not pass through the origin, it follows that  $A$  can be uniquely expressed as  $A = \{a \in \mathbb{R}^n : \langle a, v_A \rangle = 1\}$  for some  $v_A \in \mathbb{R}^n$ . Since  $A$  is supporting to  $K^*$  it follows that  $v_A \in K$ .

*Proof.* The proof of i) can be obtained along the same lines as in [8, Theorem 1.2 i)]; there, only the concavity properties of the section function w.r.t.  $a^\perp$  are used.

For ii), we adapt an argument from [8]: For a set  $\omega \subseteq \mathbb{S}^{n-1}$  we consider the function

$$f_\omega : K \rightarrow \mathbb{R}, \quad f_\omega(x) = V_{K-x}(\omega).$$

This is an affine function on  $K$ . Indeed,

$$\begin{aligned} (4.1) \quad V_{K-x}(\omega) &= \frac{1}{n} \int_\omega h_{K-x}(u) \, dS(K-x, u) \\ &= \frac{1}{n} \int_\omega h_K(u) - \langle x, u \rangle \, dS_K(u) \\ &= V_K(\omega) - \left\langle x, \int_\omega u \, dS_K(u) \right\rangle. \end{aligned}$$

In the following we consider the set  $\omega = \pi(\text{bd}(K^*) \cap A)$  and abbreviate  $f = f_\omega$ . Let  $X$  be a uniformly distributed random vector in  $K$ . For convenience, we assume that  $\text{vol}_n(K) = 1$  so that  $f(x) \in [0, 1]$  for all  $x \in K$ . Since  $f$  is affine and  $K$  is centered, we have  $V_K(\omega) = f(0) = \mathbb{E}[f(X)]$ . On the other hand, we have  $\mathbb{E}[f(X)] = 1 - \int_0^1 \mathbb{P}(f(X) < t) dt$ . For  $t \in [0, 1]$ , let  $K(t) = \{x \in K : f(x) \leq t\}$  be the corresponding sublevel set of  $f$ . Note that since  $f$  is affine,  $K(t)$  is the intersection of  $K$  with a halfspace whose normal vector  $\int_\omega u dS_K(u)$  is the same for all  $t$ .

Note that the vector  $v_A \in K$  satisfies (by definition)  $\langle v_A, u \rangle = h_K(u)$  for all  $u \in \omega$ . Thus, we obtain from (4.1) that  $v_A \in K(0)$ . Let  $m \in [0, 1]$  denote the maximum of  $f$ . Then for any  $t \in [0, 1]$  we have the inclusion,

$$K(t) \supseteq \frac{t}{m}K(m) + \frac{t-m}{m}K(0) \supseteq \frac{t}{m}K(m) + \frac{t-m}{m}v_A.$$

Thus,

$$(4.2) \quad \mathbb{P}(f(X) < t) = \text{vol}_n(K(t)) \geq \text{vol}_n\left(\frac{t}{m}K(m)\right) = \left(\frac{t}{m}\right)^n.$$

Since we have  $f(x) = 1$  for  $x \in [m, 1]$  we deduce

$$V_K(\omega) = 1 - \int_0^m \mathbb{P}(f(X) < t) dt - (1-m) \leq \frac{mn}{n+1} \leq \frac{n}{n+1}$$

This reproves the affine subspace concentration condition in this particular case. In order to have equality, it is necessary that  $m = 1$ . Moreover, for the first inequality in the above to be tight, we need equality in (4.2), i.e.,  $\text{vol}_n(K(t)) = t^n$ . Taking derivatives, it follows that  $\mathcal{H}^{n-1}(\{x \in K : f(x) = t\}) = nt^{n-1}$ . Since  $f$  is an affine function this implies that  $K$  is a pyramid with apex  $v_A$  and base  $\{x \in K : f(x) = m\}$ .  $\square$

#### ACKNOWLEDGEMENTS

Ansgar Freyer is partially supported by the Austrian Science Fund (FWF) Project P34446-N.

#### REFERENCES

- [1] K. Böröczky and P. Kalantzopoulos. Log-brunn-minkowski inequality under symmetry. *Transactions of the American Mathematical Society*, 375(08):5987–6013, 2022.
- [2] K. J. Böröczky, P. Hegedüs, and G. Zhu. On the discrete logarithmic Minkowski problem. *International Mathematics Research Notices. IMRN*, (6):1807–1838, 2016.
- [3] K. J. Böröczky and M. Henk. Cone-volume measure of general centered convex bodies. *Advances in Mathematics*, 286:703–721, 2016.
- [4] K. J. Böröczky and M. Henk. Cone-volume measure and stability. *Advances in Mathematics*, 306:24–50, 2017.
- [5] K. J. Böröczky, E. Lutwak, D. Yang, and G. Zhang. The logarithmic Minkowski problem. *Journal of the American Mathematical Society*, 26(3):831–852, 2013.
- [6] K. J. Böröczky. *The logarithmic Minkowski conjecture and the  $L_p$ -Minkowski problem*, pages 83–118. De Gruyter, Berlin, Boston, 2023.
- [7] S. Chen, Q.-R. Li, and G. Zhu. The logarithmic Minkowski problem for non-symmetric measures. *Transactions of the American Mathematical Society*, 371(4):2623–2641, 2019.
- [8] A. Freyer, M. Henk, and C. Kipp. Affine subspace concentration conditions for centered polytopes. *Mathematika*, 69(2):458–472, 2023.



- [9] M. Henk and E. Linke. Cone-volume measures of polytopes. *Advances in Mathematics*, 253:50–62, 2014.
- [10] R. Schneider. *Convex bodies: The Brunn–Minkowski theory*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2 edition, 2013.
- [11] K.-Y. Wu. Affine subspace concentration conditions. *Épjournal de Géométrie Algébrique*, 7, 2023.
- [12] G. Zhu. The logarithmic Minkowski problem for polytopes. *Advances in Mathematics*, 262:909–931, 2014.

INSTITUT FÜR MATHEMATIK, TECHNISCHE UNIVERSITÄT BERLIN, STRASSE DES 17.  
JUNI 136, 10623 BERLIN, DEUTSCHLAND

*Email address:* `eller@math.tu-berlin.de`

INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, TECHNISCHE UNIVERSITÄT  
WIEN, WIEDNER HAUPTSTRASSE 8-10/1046, 1040 WIEN, AUSTRIA

*Email address:* `ansgar.freyer@tuwien.ac.at`