

# Enhanced power enhancements for testing many moment equalities: Beyond the 2- and $\infty$ -norm

Anders Bredahl Kock

University of Oxford

St. Hilda's College

10 Manor Rd, Oxford OX1 3UQ

[anders.kock@economics.ox.ac.uk](mailto:anders.kock@economics.ox.ac.uk)

David Preinerstorfer

SEPS-SEW

University of St. Gallen

Varnbuelstrasse 14, 9000 St. Gallen

[david.preinerstorfer@unisg.ch](mailto:david.preinerstorfer@unisg.ch)

Preliminary version

July 2024

## Abstract

Tests based on the 2- and  $\infty$ -norm have received considerable attention in high-dimensional testing problems, as they are powerful against dense and sparse alternatives, respectively. The power enhancement principle of [Fan et al. \(2015\)](#) combines these two norms to construct tests that are powerful against both types of alternatives. Nevertheless, the 2- and  $\infty$ -norm are just *two* out of the whole spectrum of  $p$ -norms that one can base a test on. In the context of testing whether a candidate parameter satisfies a large number of moment equalities, we construct a test that harnesses the strength of all  $p$ -norms with exponents in  $[2, \infty]$  and is consistent against strictly more alternatives than any test based on a single  $p$ -norm. In particular, our test is consistent against more alternatives than tests based on the 2- and  $\infty$ -norm, which is what most implementations of the power enhancement principle target.

We illustrate the scope of our general results by using them to construct a test that simultaneously dominates the Anderson-Rubin test (based on  $p = 2$ ) and tests based on the  $\infty$ -norm in terms of consistency in the linear instrumental variable model with many (weak) instruments.

# 1 Introduction

In the era of big data, target parameters are often (partially) identified by the fact that they must satisfy a large number of moment equalities. Then, one frequently wishes to test whether a candidate (structural) parameter  $\beta_n^*$  satisfies  $\mathbb{E}h_n(\mathbf{X}, \beta_n^*) = \mathbf{0}_d$ , i.e., one wishes to test whether a candidate parameter  $\beta_n^*$  is an element of the identified set.<sup>1</sup> An important example that can be cast within this framework is testing restrictions on coefficients in regression models in the presence of endogenous regressors and many (weak) instruments, cf. Section 2.1.1. Testing whether a treatment has an effect on one of many outcome variables or one of several groups of individuals is another example of high practical relevance, cf. Section 2.1.2. More generally, true parameter(s) in  $M$ -,  $Z$ -, and GMM-estimation problems satisfy such moment restrictions, and hypotheses on the mean vector(s) in classic one- and two-sample testing problems can be accommodated too.

The number of moment restrictions,  $d$ , can be large for several reasons: First, if  $\beta_n^* \in \mathbb{R}^d$  with  $d$  large is defined by maximizing a population objective function, it often satisfies  $d$  first-order conditions of the form  $\mathbb{E}h_n(\mathbf{X}, \beta_n^*) = \mathbf{0}_d$ . Second, even if the dimension of the parameter vector itself is low, it may be (partially) identified by satisfying *many* moment restrictions. For example, when  $\beta_n^*$  satisfies a set of conditional moment equalities  $\mathbb{E}[h_n(\mathbf{X}, \beta_n^*) \mid \mathbf{W}] = \mathbf{0}$ , these are often transformed into unconditional ones of the form  $\mathbb{E}h_n(\mathbf{X}, \beta_n^*)g(\mathbf{W}) = \mathbf{0}_d$  for  $g$  in a family of functions  $\mathcal{G}_n$ . To avoid increasing the size of the identified set in the course of this transformation,  $\mathcal{G}_n$  may have to be large. This results in  $d$  being large. Similarly, as detailed in Section 2.1.1, in the linear IV model  $d$  equals the — nowadays often large — number of available instruments irrespectively of the dimension of  $\beta_n^*$ .

Given a sample  $\mathbf{X}_{1,n}, \dots, \mathbf{X}_{n,n}$  of random variables with the same marginal distribution as  $\mathbf{X}$ , denote the (scaled) empirical counterpart of the population moments by

$$\mathbf{H}_n(\beta_n^*) := \frac{1}{\sqrt{n}} \sum_{i=1}^n h_n(\mathbf{X}_{i,n}, \beta_n^*),$$

and let  $\hat{\Sigma}_n(\beta_n^*)$  be an estimator of the covariance matrix  $\Sigma_n(\beta_n^*)$  of  $\mathbf{H}_n(\beta_n^*)$ , which we assume to exist. It is then natural to base a test of  $\mathbb{E}h_n(\mathbf{X}, \beta_n^*) = \mathbf{0}_d$  on the distance of  $\hat{\Sigma}_n^{-1/2}(\beta_n^*)\mathbf{H}_n(\beta_n^*)$  from the origin. To measure this distance, one most commonly

---

<sup>1</sup>Here  $\beta_n^*$  may or may not be point-identified, cf., e.g., [Molinari \(2020\)](#) for many examples of how lack of point-identification may arise.

makes use of a  $p$ -norm, which then requires the choice of an exponent  $p$ . For example, in the context of testing with many instruments an appropriate choice of  $\hat{\Sigma}_n(\beta_n^*)$  and  $p = 2$  results in the Anderson-Rubin test (Anderson and Rubin (1949)) whereas  $p = \infty$  gives the sup-score test of Belloni et al. (2012). More generally, testing many moment inequalities based on  $p = \infty$  was studied in Chernozhukov et al. (2019) and  $p = 2$  was used for a fixed number of (in)equalities in Stock and Wright (2000) and Rosen (2008). Indeed, most tests used in practice are based on  $p = 2$  or  $p = \infty$  and it is well-understood that tests based on the former are relatively powerful against “dense” alternatives, that is they are good at detecting many small effects. On the other hand, tests based on the supremum-norm ( $p = \infty$ ) are relatively powerful against “sparse” alternatives, that is they are good at detecting few large effects, cf., e.g., Ingster and Suslina (2003). To construct a test that is simultaneously powerful against dense as well as many sparse alternatives Fan et al. (2015) combined tests based on the 2- and  $\infty$ -norm via their *power enhancement principle*. The idea of combining the 2- and  $\infty$ -norm based tests or a *finite* number of tests in order to construct a more powerful test has since gained considerable popularity, cf. Xu et al. (2016), Yang and Pan (2017), Tang et al. (2018), Kock and Preinerstorfer (2019), Liu et al. (2019), Jammalamadaka et al. (2020), He et al. (2021), Feng et al. (2022), Juodis and Reese (2022), Zhang et al. (2022), Fan et al. (2023), Yu et al. (2023), Ge et al. (2024), Li et al. (2024), Yu et al. (2024a), Yu et al. (2024b).

Despite the success of the power enhancement principle and related combination procedures, sparse and dense alternatives are merely two (conceptually useful) “endpoints” between which a continuum of semi-sparse alternatives exist. This continuum of structures is mirrored by a continuum of  $p$ -norms,  $p \in [2, \infty]$ , between the two extremes  $p = 2$  and  $p = \infty$  that most tests are based on. In particular, there exist semi-sparse alternatives against which tests based on the 2- and  $\infty$ -norm are inconsistent, but against which tests based on any  $p \in (2, \infty)$  are consistent. Thus, it is important to harness the power from all  $p$ -norms,  $p \in [2, \infty]$ : In the clean but restrictive testbed of the Gaussian sequence model Kock and Preinerstorfer (2023) characterized the consistency properties of all  $p$ -norm based tests and constructed a test  $\psi_n$  that is “dominant” in the following sense: If there exists *some*  $p$  for which the corresponding  $p$ -norm based test is consistent, then so is  $\psi_n$ . In particular, without inflating asymptotic size, this test is consistent against more alternatives than tests based on  $p = 2$  or  $p = \infty$ , which is what current applications of the power enhancement principle target to be consistent against.

In the present paper, we go beyond the power enhancement principle by leveraging the

findings of [Kock and Preinerstorfer \(2023\)](#) to *construct a test simultaneously dominating all  $p$ -norm based tests with  $p \in [2, \infty]$  for testing  $H_{0,n} : \mathbb{E}h_n(\mathbf{X}, \boldsymbol{\beta}_n^*) = \mathbf{0}_d$ .*<sup>2</sup> This requires rigorous high-dimensional Gaussian approximations to justify that results obtained in the stylized Gaussian sequence model carry over to more complex empirically relevant models. Indeed, a large and vibrant literature on high-dimensional Gaussian approximations over hyperrectangles was sparked by the work of [Chernozhukov et al. \(2013\)](#) and is nicely surveyed in [Chernozhukov et al. \(2023\)](#). Such approximations over hyperrectangles are suitable for studying tests based on  $p = \infty$ . However, since we study the full *family* of  $p$ -norm based tests we cannot rely on these approximations, because the rejection regions of these tests, albeit convex, are no hyperrectangles.

We now discuss our contributions in more detail. We first exhibit high-level conditions under which the characterization of the consistency properties of  $p$ -norm based tests from [Kock and Preinerstorfer \(2023\)](#) carries over to the setting considered in the present paper in [Theorem 3.1](#). We then show in [Theorem 3.2](#) that these high-level conditions are satisfied under

- (i) a precision guarantee for  $\hat{\boldsymbol{\Sigma}}_n(\boldsymbol{\beta}_n^*)$  as an estimator of  $\boldsymbol{\Sigma}_n(\boldsymbol{\beta}_n^*)$ , and
- (ii) a Gaussian approximation condition over convex sets for  $\boldsymbol{\Sigma}_n^{-1/2}(\boldsymbol{\beta}_n^*)\mathbf{H}_n(\boldsymbol{\beta}_n^*)$ .

In particular, although it is not our focus, dependent data can be accommodated. We also provide a catalogue of primitive sufficient conditions for (i) and (ii) in the i.i.d. case and discuss their implication on the allowed growth rate of  $d$ . To this end, because it is often not realistic to impose that the entries of  $h_n(\mathbf{X}_{i,n}, \boldsymbol{\beta}_n^*)$  have light tails, we focus on a setting wherein (essentially) only four moments exist. Here, to handle (i), we use the recent estimators  $\hat{\boldsymbol{\Sigma}}_n(\boldsymbol{\beta}_n^*)$  by [Abdalla and Zhivotovskiy \(2022\)](#) or [Oliveira and Rico \(2022\)](#), which remarkably provide sub-Gaussian precision guarantees without imposing any structure on  $\boldsymbol{\Sigma}_n(\boldsymbol{\beta}_n^*)$  and four moments only. Concerning (ii), [Fang and Koike \(2024\)](#) provides a suitable Gaussian approximation result. Under four moments, we need  $d = o(n^{2/5})$ , cf. [Corollary 3.3](#), and we discuss how this rate cannot be improved by much, even if one restricts attention to tests based on  $p = \infty$  only. Clearly,  $d$  larger than  $n$  is ruled out. Nevertheless, many empirically relevant settings involving a large  $d$  that is still

---

<sup>2</sup>The results in [Kock and Preinerstorfer \(2023\)](#) show that already in the Gaussian sequence model tests based on  $p$ -norms with  $p \leq 2$  are dominated (in terms of consistency) by the 2-norm based test. For simplicity, we therefore focus on  $p \in [2, \infty]$  in the present paper.

smaller than  $n$  are covered. Alternatively, one can verify (i) and (ii) under stronger moment assumptions or by imposing structure on  $\Sigma_n(\beta_n^*)$  to allow  $d$  to grow faster than  $n^{2/5}$ .

The high-level conditions (i) and (ii) allow us to establish a Gaussian approximation for  $\hat{\Sigma}_n^{-1/2}(\beta_n^*)\mathbf{H}_n(\beta_n^*)$  over convex sets when the “noncentrality term”

$$\boldsymbol{\theta}_n(\beta_n^*) = \sqrt{n}\Sigma_n^{-1/2}(\beta_n^*)\mathbb{E}h_n(\mathbf{X}, \beta_n^*)$$

is sufficiently close to  $\mathbf{0}_d$ . This enables us to deduce the power properties of tests based on the statistics  $\|\hat{\Sigma}_n^{-1/2}\mathbf{H}_n(\beta_n^*)\|_p$  for any  $p \in [2, \infty]$  from those of  $p$ -norm based tests in the Gaussian sequence model, which was analyzed in [Kock and Preinerstorfer \(2023\)](#). The case of “large”  $\boldsymbol{\theta}_n(\beta_n^*)$  is handled separately by techniques tailored to this regime. We find that a  $p$ -norm based test is consistent for  $p \in [2, \infty)$  if and only if

$$d^{-1/2} \sum_{j=1}^d [\theta_{j,n}^2(\beta_n^*) \vee |\theta_{j,n}(\beta_n^*)|^p] \rightarrow \infty,$$

and obtain a similar necessary and sufficient condition for consistency of the  $\infty$ -norm based test. Thus, consistency of a  $p$ -norm based test is *not* solely determined by the  $p$ -norm of  $\boldsymbol{\theta}_n(\beta_n^*)$ . Furthermore, the characterization reveals the following monotonicity: For  $2 \leq p < q < \infty$  tests based on the  $q$ -norm are consistent against more alternatives than tests based on the  $p$ -norm. This monotonicity does *not* extend to  $p = \infty$ . Although sufficient, yet not necessary, conditions are known for consistency of tests based on  $p = \infty$  (based on maximin rates of testing, cf. [Ingster and Suslina \(2003\)](#)), we are not aware of any full characterization providing necessary and sufficient conditions beyond the Gaussian setting considered in [Kock and Preinerstorfer \(2023\)](#). Thus, our characterization also sheds new light on tests based on  $p = \infty$ , which are frequently employed.

Based on the monotonicity property we then construct a test that harnesses the strength of all  $p$ -norms with exponents in  $[2, \infty]$  in the sense that it has asymptotic size  $\alpha \in (0, 1)$  and is consistent against a sequence of alternatives whenever some  $p$ -norm based test with  $p \in [2, \infty]$  and asymptotic size in  $(0, 1)$  is consistent against that sequence of alternatives. Our test thus simultaneously dominates all  $p$ -norm based tests. In particular, it goes beyond current implementations of the power enhancement principle, which typically target consistency against alternatives that either tests based on the 2- or  $\infty$ -norm are consistent against, i.e. only *two* norms are targeted.

Finally, we illustrate the scope of our results in the running example of the linear instrumental variable model with many (weak) instruments. Here our general construction yields a test that is consistent against more alternatives than i) the classic 2-norm based Anderson-Rubin test, ii) the  $\infty$ -norm based sup-score test and iii) power-enhancement based combinations of these.

## 2 The problem

We observe realizations of random variables  $\mathbf{X}_{1,n}, \dots, \mathbf{X}_{n,n}$ , each of which is defined on an underlying probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and takes its values in the measurable space  $(\mathcal{X}_n, \mathcal{A}_n)$ . We wish to test whether a candidate structural parameter  $\beta_n^* \in \mathbf{B}_n \neq \emptyset$  satisfies the *moment equalities*

$$\mathbb{E}h_{j,n}(\mathbf{X}_{1,n}, \beta_n^*) = 0, \quad j = 1, \dots, d(n), \quad (1)$$

for  $h_{j,n} : \mathcal{X}_n \times \mathbf{B}_n \rightarrow \mathbb{R}$  with  $h_{j,n}(\cdot, \beta_n^*)$  being  $\mathbb{P}$ -integrable and  $\mathbb{E}$  denoting the expectation with respect to  $\mathbb{P}$ . The  $\mathbf{X}_{i,n}$  are assumed to be identically distributed throughout, but are allowed to be dependent (although our detailed examples are for independent variables). *In this paper we are interested in a setting wherein  $d(n) \rightarrow \infty$  as the sample size  $n \rightarrow \infty$ .* That is, there are “many” moment equalities. To conserve on notation, we write  $d = d(n)$  whenever this causes no confusion. For our general results, no structure needs to be imposed on the parameter space  $\mathbf{B}_n$ . Of course, the complexity of constructing confidence sets for the parameters via test inversion depends on the size and structure of  $\mathbf{B}_n$ .

For  $h_n = (h_{1,n}, \dots, h_{d,n})'$  and  $\mathbf{0}_d$  denoting the  $d \times 1$  vector of zeros, the requirement in (1) is more conveniently expressed as

$$\mathbb{E}h_n(\mathbf{X}_{1,n}, \beta_n^*) = \mathbf{0}_d.$$

We thus consider the testing problem

$$H_{0,n} : \mathbb{E}h_n(\mathbf{X}_{1,n}, \beta_n^*) = \mathbf{0}_d \quad \text{against} \quad H_{1,n} : \mathbb{E}h_n(\mathbf{X}_{1,n}, \beta_n^*) \neq \mathbf{0}_d, \quad (2)$$

and are interested in asymptotic size and power properties of tests in situations in which  $d$  increases with  $n$ . Finally, we stress that since the tests to be constructed build on plugging in a candidate  $\beta_n^*$ , no assumptions need to be made on (the degree of) identification of

the parameter(s) satisfying (1). Thus, the tests are trivially robust to (weak) identification problems.

## 2.1 Examples

As pointed out already in Hansen and Singleton (1982), moment conditions as in (1) frequently arise in economic models as first-order conditions to an agent's optimization problem. Furthermore, as explained in the introduction, the true population parameters in  $M$ - and  $Z$ -estimation problems, such as (non-linear) least squares and maximum likelihood satisfy first-order conditions of the form (1). Our running example will be inference in the presence of many (weak) instruments, and since the 2-norm based test there is just the Anderson-Rubin test, we begin by recalling how this fits into our general framework. We also provide details for treatment effect testing examples and refer to the introduction for further examples falling within our general testing framework. For clarity, we assume i.i.d. sampling in the examples below.

### 2.1.1 Inference in the presence of many (weak) instruments

Consider the classic linear instrumental variable (IV) setting in which  $\mathbf{X}_{i,n} = (y_{i,n}, \mathbf{Y}'_{i,n}, \mathbf{z}'_{i,n})'$  for  $y_{i,n} \in \mathbb{R}$  an outcome of interest,  $\mathbf{Y}_{i,n} \in \mathbb{R}^k$  a vector of endogenous explanatory variables with  $k$  fixed and  $\mathbf{z}_{i,n} \in \mathbb{R}^d$  a vector of instruments. Thus,  $\mathcal{X}_n = \mathbb{R}^{1+k+d}$ ,  $\mathbf{B}_n = \mathbb{R}^k$  and  $h_n(\mathbf{X}_{1,n}, \boldsymbol{\beta}_n^*) = (y_{1,n} - \mathbf{Y}'_{1,n} \boldsymbol{\beta}_n^*) \mathbf{z}_{1,n}$ . In this context, we wish to test whether a candidate parameter vector  $\boldsymbol{\beta}_n^*$  lies in the identified set, i.e., whether

$$H_{0,n} : \mathbb{E} [(y_{1,n} - \mathbf{Y}'_{1,n} \boldsymbol{\beta}_n^*) \mathbf{z}_{1,n}] = \mathbf{0}_d \quad \text{against} \quad H_{1,n} : \mathbb{E} [(y_{1,n} - \mathbf{Y}'_{1,n} \boldsymbol{\beta}_n^*) \mathbf{z}_{1,n}] \neq \mathbf{0}_d.$$

Note that the number of moment equalities equals the number of instruments  $d$ . This can be large even for  $k$  fixed. Inference on  $\boldsymbol{\beta}_n^*$  in the presence of many (weak) instruments has received considerable recent attention as witnessed by, e.g., the works of Andrews and Stock (2007), Anatolyev and Gospodinov (2011), Belloni et al. (2012), Tchuente and Carrasco (2016), Kaffo and Wang (2017), Crudu et al. (2021), Mikusheva (2021), Matsushita and Otsu (2021), Mikusheva and Sun (2022), Matsushita and Otsu (2022), Dovì et al. (2022), Boot and Ligtenberg (2023). There are several important practical reasons for this interest. First, when identification is weak, researchers may seek to obtain more precise inference by using a large number of instruments in order to capture more of the exoge-

nous variation in the endogenous covariates. Second, approaches such as the granular IV approach of [Gabaix and Koijen \(2020\)](#), the saturation approach of [Blandhol et al. \(2022\)](#) or Mendelian Randomization ([Davey Smith and Ebrahim \(2003\)](#)), the latter using genetic variation as instruments, can lead to situations where the number of instruments is large compared to the sample size. The same is true for technical instruments such as transformations or interactions or empirical strategies such as “judge designs,” cf. [Mikusheva and Sun \(2024\)](#). However, as there is no guarantee that even a large number of instruments is jointly informative, it is important to develop tests that remain valid under no or weak identification.

Our general results allow us to construct a test that strictly dominates the Anderson-Rubin test, the sup-score test, and power enhancement combinations of these in terms of consistency properties when  $d \rightarrow \infty$ , cf. [Section 5](#).

The examples in the following two subsections revisit Examples 1 and 2 in [Belloni et al. \(2018\)](#) and both consider testing for treatment effects in a randomized controlled trial.

### 2.1.2 Randomized controlled trial with many outcomes

Consider a randomized controlled trial in which  $\mathbf{X}_{i,n} = (D_{i,n}, \mathbf{Y}'_{i,n}(0), \mathbf{Y}'_{i,n}(1))'$  for  $D_{i,n} \in \{0, 1\}$  indicating whether individual  $i$  is assigned to the treatment ( $D_{i,n} = 1$ ) or not ( $D_{i,n} = 0$ ). For each individual we observe a large number,  $d$ , of outcome variables and denote this by  $\mathbf{Y}_{i,n}(1) = (Y_{i1,n}(1), \dots, Y_{id,n}(1))' \in \mathbb{R}^d$  in case individual  $i$  is treated and  $\mathbf{Y}_{i,n}(0) = (Y_{i1,n}(0), \dots, Y_{id,n}(0))' \in \mathbb{R}^d$  if not. This framework of many outcome variables is relevant, because a researcher may, for example, be interested in whether a medication has an effect on any of a large number of health outcomes (blood pressure, weight, heart rate, ...) rather than only monitoring the effect on a single of these.<sup>3</sup>

Suppose that  $\mathbb{P}(D_{1,n} = 1) = \pi$  for  $\pi \in (0, 1)$  a known probability of being assigned to the treatment and that the treatments are assigned independently of the potential outcomes, that is  $D_{1,n} \perp\!\!\!\perp (\mathbf{Y}_{1,n}(0), \mathbf{Y}_{1,n}(1))$ . Furthermore,  $\mathcal{X}_n = \mathbb{R}^{1+2d}$ . The treatment does not have an effect on any of the outcomes being measured if  $\mathbb{E}\mathbf{Y}_{1,n}(1) = \mathbb{E}\mathbf{Y}_{1,n}(0)$ . More generally, for  $\boldsymbol{\beta}_n^* = (\beta_{n,1}^*, \dots, \beta_{n,d}^*)' \in \mathbb{R}^d$  a vector of coordinate-wise effects, we are interested in testing whether  $\mathbb{E}\mathbf{Y}_{1,n}(1) - \mathbb{E}\mathbf{Y}_{1,n}(0) = \boldsymbol{\beta}_n^*$ . Because  $D_{1,n} \perp\!\!\!\perp (\mathbf{Y}_{1,n}(0), \mathbf{Y}_{1,n}(1))$

---

<sup>3</sup>Although a treatment may be targeted to affect a single outcome of interest, it may nevertheless be important to scan for potential side effects on other outcome variables.



this is equivalent to

$$\mathbb{E} \left[ \frac{D_{1,n} \mathbf{Y}_{1,n}(1)}{\pi} - \frac{(1 - D_{1,n}) \mathbf{Y}_{1,n}(0)}{1 - \pi} - \boldsymbol{\beta}_n^* \right] = \mathbb{E} \left[ \frac{D_{1,n} \mathbf{Y}_{1,n}}{\pi} - \frac{(1 - D_{1,n}) \mathbf{Y}_{1,n}}{1 - \pi} - \boldsymbol{\beta}_n^* \right] = \mathbf{0}_d, \quad (3)$$

where  $\mathbf{Y}_{1,n} = D_{1,n} \mathbf{Y}_{1,n}(1) + (1 - D_{1,n}) \mathbf{Y}_{1,n}(0)$  is the observed outcome. This is a testing problem of the form in (2) with  $h_n(\mathbf{X}_{1,n}, \boldsymbol{\beta}_n^*) = \frac{D_{1,n} \mathbf{Y}_{1,n}}{\pi} - \frac{(1 - D_{1,n}) \mathbf{Y}_{1,n}}{1 - \pi} - \boldsymbol{\beta}_n^*$ .

### 2.1.3 Randomized controlled trials with many groups

Similar to the previous example of Section 2.1.2, consider a randomized controlled trial. However, let there be only one outcome variable for each individual. Suppose, however, that each subject falls into one of  $d$  groups and one wants to test whether a treatment has the hypothesized effect (e.g., no effect) in every one of these groups. Assuming for simplicity that there are  $n$  observations available in each group, one can thus again write the problem in the form of two  $d$ -dimensional outcome vectors  $\mathbf{Y}_{i,n}(0)$  and  $\mathbf{Y}_{i,n}(1)$  as in the previous example ( $\mathbf{Y}_{i,n}(0)$  is the vector of the stacked outcomes of the  $i$ -th subjects in each group under the no treatment condition, whereas  $\mathbf{Y}_{i,n}(1)$  is the vector of the stacked outcomes of the  $i$ -th subjects in each group under the treatment condition). The null hypotheses of the treatment having effect  $\beta_{n,j}^*$  in group  $j$  for  $j = 1, \dots, d$  then again amounts to (3).

As pointed out in, e.g., Belloni et al. (2018), the project STAR investigating the effect of class size on student learning falls within the framework of the present example as many background characteristics were collected on pupils and teachers. For example gender or race. For pupils, their month of birth was also recorded. Identifying each group with a particular combination of covariates, this results in a large number of groups.

## 3 Characterization of consistency properties of $p$ -norm based tests

Recall from the introduction that given  $\boldsymbol{\beta}_n^* \in \mathbf{B}_n$ , we write

$$\mathbf{H}_n(\boldsymbol{\beta}_n^*) := \frac{1}{\sqrt{n}} \sum_{i=1}^n h_n(\mathbf{X}_{i,n}, \boldsymbol{\beta}_n^*)$$

and let  $\hat{\Sigma}_n(\beta_n^*)$  be a positive semidefinite and symmetric estimator of the covariance matrix  $\Sigma_n(\beta_n^*)$  of  $\mathbf{H}_n(\beta_n^*)$ , which we assume to exist.

*Remark 3.1.* Already under independent sampling, the choice of  $\hat{\Sigma}_n(\beta_n^*)$  may depend on instance-specific additional structural information (e.g., bandedness, sparsity or factor structure), the incorporation of which can improve the estimator. We are particularly interested in a setting where no structural information on  $\Sigma_n(\beta_n^*)$  is available and leave the specific choice of estimator unspecified in the following. In Section 3.2 we highlight how the precision of the chosen  $\hat{\Sigma}_n(\beta_n^*)$  influences the allowed growth rate of  $d$  and discuss precision guarantees available in the literature.

A common way of measuring the empirical evidence against the null of  $\mathbb{E}h_n(\mathbf{X}_{1,n}, \beta_n^*) = \mathbf{0}_d$  is based on the distance of  $\hat{\Sigma}_n^{-1/2}(\beta_n^*)\mathbf{H}_n(\beta_n^*)$  to  $\mathbf{0}_d$  in the Euclidean norm, that is, one rejects  $H_{0,n}$  null whenever

$$S_{n,2}(\beta_n^*) := \sqrt{\mathbf{H}'_n(\beta_n^*)\hat{\Sigma}_n^{-1}(\beta_n^*)\mathbf{H}_n(\beta_n^*)} = \left\| \hat{\Sigma}_n^{-1/2}(\beta_n^*)\mathbf{H}_n(\beta_n^*) \right\|_2 \quad (4)$$

exceeds a critical value chosen to ensure that the resulting test has a desired (asymptotic) size  $\alpha \in (0, 1)$ , and where  $\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^d x_i^2}$  for  $\mathbf{x} \in \mathbb{R}^d$ .<sup>4</sup> For  $\mathbf{x} = (x_1, \dots, x_d)' \in \mathbb{R}^d$  and  $p \in [2, \infty]$ , define the  $p$ -norm

$$\|\mathbf{x}\|_p := \begin{cases} \left( \sum_{i=1}^d |x_i|^p \right)^{\frac{1}{p}} & \text{if } p < \infty, \\ \max_{i=1, \dots, d} |x_i| & \text{else} \end{cases}$$

and introduce, analogously to  $S_{n,2}(\beta_n^*)$  in (4) but based on the  $p$ -norm, the family of test statistics

$$S_{n,p}(\beta_n^*) := \left\| \hat{\Sigma}_n^{-1/2}(\beta_n^*)\mathbf{H}_n(\beta_n^*) \right\|_p, \quad p \in [2, \infty].$$

For sequences of critical values  $(\kappa_{n,p})_{n \in \mathbb{N}}$  that guarantee a desired asymptotic size, we shall

---

<sup>4</sup>In order to avoid taking a stance on  $\hat{\Sigma}_n(\beta_n^*)$  potentially not being invertible, we denote by  $\mathbf{A}^{-1}$  the Moore-Penrose pseudoinverse for any matrix  $\mathbf{A}$  and define  $\mathbf{A}^{-1/2} := (\mathbf{A}^{1/2})^{-1}$  in case  $\mathbf{A}$  is symmetric and positive semidefinite. Recall that the Moore-Penrose inverse is identical to the regular matrix inverse whenever the latter exists.

study consistency properties of the  $p$ -norm based tests

$$\mathbb{1}(S_{n,p}(\boldsymbol{\beta}_n^*) \geq \kappa_{n,p})$$

and coverage properties of the associated confidence sets

$$\{\boldsymbol{\beta} \in \mathbf{B}_n : S_{n,p}(\boldsymbol{\beta}) \leq \kappa_{n,p}\},$$

when  $d \rightarrow \infty$  as  $n \rightarrow \infty$ . Define

$$\mathbf{B}^* := \prod_{n=1}^{\infty} \mathbf{B}_n,$$

i.e., the set of possible *sequences* of parameters  $\boldsymbol{\beta}_n$  for  $n \in \mathbb{N}$ , along which power or coverage properties of tests for the sequence of testing problems in (2) can be studied asymptotically in the current setup. In particular, we write  $\boldsymbol{\beta}^* = (\boldsymbol{\beta}_1^*, \boldsymbol{\beta}_2^*, \dots)$ . Let

$$\mathbf{B}^{(0)} = \{\boldsymbol{\beta} = (\boldsymbol{\beta}_1, \boldsymbol{\beta}_2, \dots) \in \mathbf{B}^* : \mathbb{E}h_n(\mathbf{X}_{1,n}, \boldsymbol{\beta}_n) = \mathbf{0}_d \text{ for every } n \in \mathbb{N}\} \subseteq \mathbf{B}^*$$

be the set of sequences of parameters satisfying  $H_{0,n}$  in (2) for every  $n \in \mathbb{N}$ .

As we shall see, the scaled deviations from the null hypothesis

$$\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*) = \sqrt{n}\boldsymbol{\Sigma}_n^{-1/2}(\boldsymbol{\beta}_n^*)\mathbb{E}h_n(\mathbf{X}_{1,n}, \boldsymbol{\beta}_n^*) \in \mathbb{R}^d, \quad (5)$$

play the role of a noncentrality parameter in characterizing the consistency properties of tests based on  $S_{n,p}(\boldsymbol{\beta}_n^*)$ . We also let  $\sigma_p^2 := \mathbb{V}ar(|Z|^p)$  with  $Z \sim \mathbf{N}_1(0, 1)$  and define the functions

$$\lambda_p(x) = \mathbb{E}|Z + x|^p \quad \text{and} \quad g_p(x) = x^2 \vee |x|^p, \quad x \in \mathbb{R}, \quad p \in [2, \infty). \quad (6)$$

### 3.1 Power properties of $p$ -norm based tests: $p \in [2, \infty)$

We first consider the case of  $p \in [2, \infty)$  as the characterization of their power has a common structure. The case of  $p = \infty$  is then covered in Section 3.4.

Our most general characterization of the power properties of  $p$ -norm based tests is carried out under the following high-level conditions for which primitive conditions are provided in Section 3.2.

*Assumption 3.1* (Size).  $\beta^* \in \mathbf{B}^{(0)}$  and

$$\frac{S_{n,p}^p(\beta_n^*) - d\lambda_p(0)}{\sqrt{d}\sigma_p} \rightsquigarrow \mathbf{N}_1(0, 1). \quad (7)$$

Note that  $d\lambda_p(0)$  appearing in Assumption 3.1 is not the expected value of  $S_{n,p}^p(\beta_n^*)$ , which, in fact, may not even exist. Section 3.2 provides conditions under which Assumption 3.1 is satisfied for *all* powers  $p \in [2, \infty)$  even when the  $h_{j,n}(\mathbf{X}_{1,n}, \beta_n^*)$  have only four moments.

*Assumption 3.2* (Power). For  $\beta^* \in \mathbf{B}^*$  we summarize the following two properties:

1. If  $\frac{1}{\sqrt{d'}} \sum_{i=1}^{d'} [\lambda_p(\theta_{i,n'}(\beta_{n'}^*)) - \lambda_p(0)]$  is bounded for a subsequence  $n'$  of  $n$  and where  $d' = d(n')$ , then

$$\frac{S_{n',p}^p(\beta_{n'}^*) - \sum_{i=1}^{d'} \lambda_p(\theta_{i,n'}(\beta_{n'}^*))}{\sqrt{d'}\sigma_p} \rightsquigarrow \mathbf{N}_1(0, 1). \quad (8)$$

2. If  $\frac{1}{\sqrt{d}} \sum_{i=1}^d [\lambda_p(\theta_{i,n}(\beta_n^*)) - \lambda_p(0)] \rightarrow \infty$ , then

$$\frac{S_{n,p}^p(\beta_n^*) - \sum_{i=1}^d \lambda_p(\theta_{i,n}(\beta_n^*))}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}(\beta_n^*)) - \lambda_p(0)]} = o_{\mathbb{P}}(1). \quad (9)$$

The following theorem now characterizes asymptotic size and power properties of  $p$ -norm based tests for  $p \in [2, \infty)$ . We denote the cdf of a standard normal distribution by  $\Phi$ .

**Theorem 3.1.** *Let  $p \in [2, \infty)$  and  $\alpha \in (0, 1)$ .*

1. *Size control: Under Assumption 3.1, a sequence of real numbers  $(\kappa_{n,p})_{n \in \mathbb{N}}$  satisfies*

$$\mathbb{P}(S_{n,p}(\beta_n^*) \geq \kappa_{n,p}) \rightarrow \alpha, \quad (10)$$

*if and only if  $\kappa_{n,p} = \kappa_{n,p}(\alpha) = [(\Phi^{-1}(1 - \alpha) + o(1))d^{1/2}\sigma_p + d\lambda_p(0)]^{1/p}$ .*

2. *Local power: If Part 1 of Assumption 3.2 holds and if*

$$\frac{1}{\sqrt{d}} \sum_{i=1}^d [\lambda_p(\theta_{i,n}(\beta_n^*)) - \lambda_p(0)] \rightarrow c \in [0, \infty),$$

then, for  $(\kappa_{n,p})_{n \in \mathbb{N}}$  satisfying (10),

$$\mathbb{P}(S_{n,p}(\boldsymbol{\beta}_n^*) \geq \kappa_{n,p}) \rightarrow 1 - \Phi(\Phi^{-1}(1 - \alpha) - c/\sigma_p).$$

3. *Consistency:* Under Assumption 3.2 and for  $(\kappa_{n,p})_{n \in \mathbb{N}}$  satisfying (10), it holds that

$$\mathbb{P}(S_{n,p}(\boldsymbol{\beta}_n^*) \geq \kappa_{n,p}) \rightarrow 1 \iff \frac{\sum_{i=1}^d [\lambda_p(\theta_{i,n}(\boldsymbol{\beta}_n^*)) - \lambda_p(0)]}{\sqrt{d}} \rightarrow \infty \quad (11)$$

$$\iff \frac{\sum_{i=1}^d g_p(\theta_{i,n}(\boldsymbol{\beta}_n^*))}{\sqrt{d}} \rightarrow \infty. \quad (12)$$

Part 1 of Theorem 3.1 characterizes sequences of critical values that yield asymptotic size control. For any  $\alpha \in (0, 1)$ , a canonical choice is

$$\kappa_{n,p} = [\Phi^{-1}(1 - \alpha)d^{1/2}\sigma_p + d\lambda_p(0)]^{1/p}.$$

Part 2 provides results on local asymptotic power and Part 3 provides a complete characterization of the alternatives that a  $p$ -norm based test is consistent against. Apart from when  $p = 2$ , neither of these can be expressed in terms of only the  $p$ -norm of the (scaled) deviation from the null  $\|\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*)\|_p$ ; instead they depend on the asymptotic behaviour of

$$d^{-1/2} \sum_{i=1}^d [\lambda_p(\theta_{i,n}(\boldsymbol{\beta}_n^*)) - \lambda_p(0)].$$

Concerning consistency, the divergence of the latter is equivalent to divergence of

$$\frac{\sum_{i=1}^d g_p(\theta_{i,n}(\boldsymbol{\beta}_n^*))}{\sqrt{d}} = \frac{\sum_{i=1}^d [\theta_{i,n}^2(\boldsymbol{\beta}_n^*) \vee |\theta_{i,n}(\boldsymbol{\beta}_n^*)|^p]}{\sqrt{d}},$$

which is somewhat easier to interpret.

Concerning the consistency of a  $p$ -norm based test (with asymptotic size in  $(0, 1)$ ), (12) implies that a sufficient condition is that  $d^{-1/2}\|\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*)\|_2^2 \rightarrow \infty$  or  $d^{-1/2}\|\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*)\|_p^p \rightarrow \infty$ , cf. also the previous display. What is more, observing that

$$g_p(x) \leq g_q(x) \quad \text{for} \quad 2 \leq p < q < \infty \text{ and all } x \in \mathbb{R},$$

Part 3 of Theorem 3.1 shows that

$$\mathbb{P}(S_{n,p}(\boldsymbol{\beta}_n^*) \geq \kappa_{n,p}) \rightarrow 1 \quad \text{implies} \quad \mathbb{P}(S_{n,q}(\boldsymbol{\beta}_n^*) \geq \kappa_{n,q}) \rightarrow 1 \quad (13)$$

when  $(\kappa_{n,p})_{n \in \mathbb{N}}$  and  $(\kappa_{n,q})_{n \in \mathbb{N}}$  are chosen such that the corresponding tests have asymptotic sizes in  $(0, 1)$ . In words, *any violation of the moment conditions  $\mathbb{E}h_n(\mathbf{X}_{1,n}, \boldsymbol{\beta}_n^*) = \mathbf{0}_d$  that a test based on the  $p$ -norm is consistent against, a test based on the  $q$ -norm will also be consistent against*,  $2 \leq p < q < \infty$ . Thus, the  $q$ -norm based test always weakly dominates the  $p$ -norm based test in terms of consistency. This domination is *strict* if and only there exists a  $\boldsymbol{\beta}^* \in \mathbf{B}^*$  such that  $\liminf_{n \rightarrow \infty} \mathbb{P}(S_{n,p}(\boldsymbol{\beta}_n^*) \geq \kappa_{n,p}) < 1$  but  $\mathbb{P}(S_{n,q}(\boldsymbol{\beta}_n^*) \geq \kappa_{n,q}) \rightarrow 1$ , i.e., if and only if (cf. (12))

$$\liminf_{n \rightarrow \infty} \frac{\sum_{i=1}^d g_p(\theta_{i,n}(\boldsymbol{\beta}_n^*))}{\sqrt{d}} < \infty \quad \text{and} \quad \frac{\sum_{i=1}^d g_q(\theta_{i,n}(\boldsymbol{\beta}_n^*))}{\sqrt{d}} \rightarrow \infty. \quad (14)$$

Summarizing, given the conditions imposed in Theorem 3.1, the characterization and monotonicity properties of the consistency of  $p$ -norm based tests established in Kock and Preinerstorfer (2023) in the Gaussian sequence model continue to hold in the current setting.

By Theorem 3.4 in Kock and Preinerstorfer (2023) the  $\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*)$  that satisfy (14) are necessarily approximately sparse and unbalanced. Furthermore, if the limit inferior to the left in (14) is zero, then there exists a subsequence along which the  $p$ -norm based test has asymptotic power equal to its size, while the  $q$ -norm based test is consistent — a substantial difference in power.

### 3.2 Primitive conditions for Assumptions 3.1 and 3.2

We next provide conditions that will be shown to be sufficient for Assumptions 3.1 and 3.2 to be satisfied and hence for Theorem 3.1 to apply. To formulate the conditions, let  $\mathbf{Z}_d \sim \mathbf{N}_d(\mathbf{0}_d, \mathbf{I}_d)$ ,

$$\mathcal{C}_n = \{C \subseteq \mathbb{R}^{d(n)} : C \text{ is convex and Borel measurable}\},$$

and  $\|\mathbf{A}\|_2$  be the spectral norm of the matrix  $\mathbf{A}$ .

*Assumption 3.3.* Let  $\boldsymbol{\beta}^* \in \mathbf{B}^*$ , assume that  $d \rightarrow \infty$ , that the eigenvalues of  $\boldsymbol{\Sigma}_n(\boldsymbol{\beta}_n^*)$  are (uniformly) bounded away from zero and from above, and that

1.  $\|\hat{\boldsymbol{\Sigma}}_n(\boldsymbol{\beta}_n^*) - \boldsymbol{\Sigma}_n(\boldsymbol{\beta}_n^*)\|_2 = O_{\mathbb{P}}(a_n)$  with  $d^{3/4}a_n \rightarrow 0$ , and

$$2. \sup_{C \in \mathcal{C}_n} \left| \mathbb{P} \left( \hat{\Sigma}_n^{-1/2}(\beta_n^*) \mathbf{H}_n(\beta_n^*) \in C \right) - \mathbb{P} \left( \mathbf{Z}_d + \boldsymbol{\theta}_n(\beta_n^*) \in C \right) \right| \rightarrow 0.$$

Note that these conditions do not impose independence.

**Theorem 3.2.** *Suppose Assumption 3.3 is satisfied. Then, the following holds:*

- If  $\beta^* \in \mathbf{B}^{(0)}$ , then Assumption 3.1 holds for all  $p \in [2, \infty)$ .
- Assumption 3.2 holds for all  $p \in [2, \infty)$ .

*Remark 3.2.* The requirement  $d^{3/4} a_n \rightarrow 0$  in Part 1 of Assumption 3.3 can be relaxed to  $d^{3/4 \wedge (1-1/p)} a_n \rightarrow 0$ , which is milder for  $p \in [2, 4)$ , but we omit a formal statement.

Theorem 3.2 shows that for Theorem 3.1 to apply it suffices to upper bound the estimation error of the covariance matrix,  $\|\hat{\Sigma}_n(\beta_n^*) - \Sigma_n(\beta_n^*)\|_2$ , and a Gaussian approximation to hold.<sup>5</sup> As discussed below, for both of these requirements catalogues of sufficient conditions exist in the literature.

To prove Theorem 3.2 we show in Lemma C.3 that when  $d^{-1} \sum_{i=1}^d g_2(\theta_{i,n}(\beta_n^*))$  is bounded, a Gaussian approximation property holds even when  $\Sigma_n(\beta_n^*)$  is replaced by  $\hat{\Sigma}_n(\beta_n^*)$ :

$$\sup_{C \in \mathcal{C}_n} \left| \mathbb{P} \left( \hat{\Sigma}_n^{-1/2}(\beta_n^*) \mathbf{H}_n(\beta_n^*) \in C \right) - \mathbb{P} \left( \mathbf{Z}_d + \boldsymbol{\theta}_n(\beta_n^*) \in C \right) \right| \rightarrow 0. \quad (15)$$

This essentially allows us to deduce the desired properties of  $S_{n,p}(\beta_n^*) = \|\hat{\Sigma}_n^{-1/2}(\beta_n^*) \mathbf{H}_n(\beta_n^*)\|_p$  in Assumptions 3.1 and 3.2 from the ones of  $p$ -norms of the high-dimensional Gaussian mean shift  $\mathbf{Z}_d + \boldsymbol{\theta}_n(\beta_n^*)$  in the “limit experiment”, the latter of which was analyzed in Kock and Preinerstorfer (2023).<sup>6</sup> More broadly, (15) may be of independent interest beyond the scope of this paper as it allows one to deduce probabilistic properties of the scaled “central statistic”  $\hat{\Sigma}_n^{-1/2}(\beta_n^*) \mathbf{H}_n(\beta_n^*)$  from those of  $\mathbf{Z}_d + \boldsymbol{\theta}_n(\beta_n^*)$  even when  $d \rightarrow \infty$ . We next discuss Parts 1 and 2 of Assumption 3.3 in further detail.

Concerning 1 of Assumption 3.3, it is well-known that if the vectors  $h_n(\mathbf{X}_{i,n}, \beta_n^*)$  are i.i.d. and appropriately sub-Gaussian, then the empirical covariance matrix  $\hat{\Sigma}_{n,\text{emp}}(\beta_n^*)$  satisfies  $\|\hat{\Sigma}_{n,\text{emp}}(\beta_n^*) - \Sigma_n(\beta_n^*)\|_2 = O_{\mathbb{P}}(\sqrt{d/n})$ , cf., e.g., Vershynin (2012) or Koltchinskii

<sup>5</sup>We impose the Gaussian approximation to hold over convex sets as we study all  $p$ -norm based tests simultaneously. If one is only interested in a single  $p$ , it may be enough for the approximation to hold over  $p$ -norm balls with arbitrary centres and radii for this given  $p$  — potentially allowing for  $d$  to grow faster.

<sup>6</sup>This is done when  $d^{-1} \sum_{i=1}^d g_2(\theta_{i,n}(\beta_n^*)) = d^{-1} \|\boldsymbol{\theta}_n(\beta_n^*)\|_2^2$  is bounded. In regimes wherein this is not the case, arguments tailored to those settings are used to verify (9) of Assumption 3.2.

and Lounici (2017). If, in addition,  $\Sigma_n(\beta_n^*)$  possesses further structure, this can be utilised to construct estimators with even better performance guarantees. For example, Bickel and Levina (2008) showed that if  $\Sigma_n(\beta_n^*)$  is sparse,  $\hat{\Sigma}_{n,\text{emp}}(\beta_n^*)$  can be thresholded to yield an estimator satisfying  $\|\hat{\Sigma}_n(\beta_n^*) - \Sigma_n(\beta_n^*)\|_2 = O_{\mathbb{P}}(\sqrt{\log(d)/n})$ . The overviews in Fan et al. (2016) and Cai et al. (2016) list further estimators utilising structural properties of  $\Sigma_n(\beta_n^*)$  and their performance guarantees.

One is often not willing to assume that the vectors  $h_n(\mathbf{X}_{i,n}, \beta_n^*)$  are sub-Gaussian. In particular, much effort is currently being devoted to the construction of covariance matrix estimators with the sub-Gaussian performance guarantee  $\|\hat{\Sigma}_n(\beta_n^*) - \Sigma_n(\beta_n^*)\|_2 = O_{\mathbb{P}}(\sqrt{d/n})$  even when only *four* moments exist. Although the sample covariance matrix does not work well under such heavy tails,<sup>7</sup> the existence of estimators satisfying  $\|\hat{\Sigma}_n(\beta_n^*) - \Sigma_n(\beta_n^*)\|_2 = O_{\mathbb{P}}(\sqrt{d/n})$  — without imposing structure on  $\Sigma_n(\beta_n^*)$  — has been established in Abdalla and Zhivotovskiy (2022) and Oliveira and Rico (2022), cf. also Mendelson and Zhivotovskiy (2020) for bounds containing an extra factor  $\sqrt{\log(d)}$ . These results are proven under the assumption that the kurtosis of all one-dimensional marginals of  $h_n(\mathbf{X}_{1,n}, \beta_n^*)$  is bounded (in particular only four moments need to exist):

*Assumption 3.4.* Denote  $\mu_n(\beta_n^*) = \mathbb{E}h_n(\mathbf{X}_{1,n}, \beta_n^*)$  and let  $\langle \cdot, \cdot \rangle$  be the standard inner product in  $\mathbb{R}^d$ . There exists a real number  $L(\beta^*) \geq 1$ , such that for all  $t \in \mathbb{R}^d$  and every  $n \in \mathbb{N}$

$$\left(\mathbb{E}\langle [h_n(\mathbf{X}_{1,n}, \beta_n^*) - \mu_n(\beta_n^*)], t \rangle^4\right)^{\frac{1}{4}} \leq L(\beta^*) \left(\mathbb{E}\langle [h_n(\mathbf{X}_{1,n}, \beta_n^*) - \mu_n(\beta_n^*)], t \rangle^2\right)^{\frac{1}{2}}.$$

As pointed out in Mendelson and Zhivotovskiy (2020), Assumption 3.4 is satisfied if, e.g.,  $h_n(\mathbf{X}_{1,n}, \beta_n^*) - \mu_n(\beta_n^*)$  follows a multivariate  $t$ -distribution with  $\nu > 4$  degrees of freedom. This example is one of rather heavy tails as only moments strictly lower than  $\nu$  exist.

Concerning Part 2 of Assumption 3.3, note that by translation invariance of convex sets this is equivalent to

$$\sup_{C \in \mathcal{C}_n} \left| \mathbb{P} \left( n^{-1/2} \sum_{i=1}^n \Sigma_n^{-1/2}(\beta_n^*) [h_n(\mathbf{X}_{i,n}, \beta_n^*) - \mu_n(\beta_n^*)] \in C \right) - \mathbb{P}(\mathbf{Z}_d \in C) \right| \rightarrow 0;$$

that is (scaled) partial sums in  $\mathbb{R}^d$  with mean  $\mathbf{0}_d$  and identity covariance matrix  $\mathbf{I}_d$  should

---

<sup>7</sup>This is in analogy to the sub-optimality of the sample average under heavy tails in the one-dimensional mean estimation problem, cf. the overview in Lugosi and Mendelson (2019).



obey a Gaussian approximation. We show that by [Fang and Koike \(2024\)](#) this holds under i.i.d. sampling and Assumption 3.4 (or, slightly weaker, bounded fourth moments) if, up to logarithmic factors in  $n$ ,  $d/n^{2/5} \rightarrow 0$ .

Summarizing the above discussion results in the following statement.

**Corollary 3.3.** *Let  $\beta^* \in \mathbf{B}^*$ , let  $d \rightarrow \infty$ , and suppose that the eigenvalues of  $\Sigma_n(\beta_n^*)$  are (uniformly) bounded away from zero and from above. Furthermore, let  $\mathbf{X}_{1,n}, \dots, \mathbf{X}_{n,n}$  be i.i.d. for each  $n \in \mathbb{N}$ . Suppose  $\frac{d}{n^{2/5}}[\log(n)]^{2/5} \rightarrow 0$  and one uses the estimator  $\hat{\Sigma}_n(\beta_n^*)$  from [Abdalla and Zhivotovskiy \(2022\)](#) (with their  $\eta = 0$ ,  $\delta = 1/d$  and  $p = 4$ ) or [Oliveira and Rico \(2022\)](#) (with their  $\eta = 0$ ,  $\alpha = 1/d$  and  $p = 4$ ) applied to the auxiliary sample*

$$(h_n(\mathbf{X}_{2,n}, \beta_n^*) - h_n(\mathbf{X}_{1,n}, \beta_n^*)) / \sqrt{2}, \dots, ((h_n(\mathbf{X}_{2_{\lfloor n/2 \rfloor, n}}, \beta_n^*) - h_n(\mathbf{X}_{2_{\lfloor n/2 \rfloor - 1, n}}, \beta_n^*)) / \sqrt{2}.$$

*If Assumption 3.4 holds, then Assumption 3.3 is satisfied.*

Under the assumptions of Corollary 3.3, Theorem 3.1 provides a characterization of the consistency properties of  $p$ -norm based tests for all  $p \in [2, \infty)$ . In particular, even though the  $h_{j,n}(\cdot, \beta_n^*)$  only possess four moments, the characterization also holds for  $p$ -norm based tests with  $p > 4$ ; that is the consistency properties of  $p$ -norm based tests are characterized even when the  $p$ th moments of the entries of  $h_n(\mathbf{X}_{1,n}, \beta_n^*)$  need not exist.

*Remark 3.3.* Many recent tests in high-dimensional testing problems are based on the supremum-norm such that Gaussian approximations over the class of hyperrectangles (contained in  $\mathcal{C}_n$ ) suffice. This allows  $d$  to increase faster than  $n^{2/5}$ . However, even for this smaller class of sets, there exist distributions with bounded fourth moments such that the Gaussian approximation breaks down if  $d$  grows faster than  $n$ , cf. [Zhang and Wu \(2017\)](#) and [Kock and Preinerstorfer \(2024\)](#) for precise formulations. In particular, there exist distributions satisfying the null such that for all  $\alpha \in (0, 1)$  the rejection frequency of the supremum-norm based test using Gaussian size  $\alpha$  critical values test tends to *one*. Thus, not even supremum-norm based tests control size uniformly over distributions with bounded fourth moments beyond  $d = n$ . In Section 3.4 we characterize the consistency properties of supremum-norm based tests and in Section 6 we show how a sample split can be used to accommodate any growth rate of  $d$ .

*Remark 3.4.* As acknowledged by [Abdalla and Zhivotovskiy \(2022\)](#) and [Oliveira and Rico \(2022\)](#), their estimators of the population covariance matrix are targeting the best possible statistical performance guarantees at the expense of not being practical to implement.

“User-friendly” estimators, losing logarithmic factors in  $d$  in the performance guarantees, have been proposed in [Ke et al. \(2019\)](#).

### 3.3 The size of confidence sets based on $p$ -norms for $p \in [2, \infty)$

Theorem [3.1](#) also has implications for the coverage properties and size of confidence sets on the form

$$\text{CS}_{n,p,1-\alpha_p} := \{\boldsymbol{\beta} \in \mathbf{B}_n : S_{n,p}(\boldsymbol{\beta}) \leq \kappa_{n,p}(\alpha_p)\} \quad \text{for } p \in [2, \infty) \text{ and } \alpha_p \in (0, 1).$$

Under the assumptions of Part 1 of Theorem [3.1](#), these confidence sets have the desired (uniform) asymptotic coverage guarantee. That is, for any  $\boldsymbol{\beta}^* \in \mathbf{B}^{(0)}$ , it holds that

$$\lim_{n \rightarrow \infty} \mathbb{P}(\boldsymbol{\beta}_n^* \in \text{CS}_{n,p,1-\alpha_p}) = 1 - \alpha_p.$$

Furthermore, under the assumptions of Part 3 of the same theorem and  $2 \leq p < q < \infty$ , one has for any  $\boldsymbol{\beta}^*$  (not satisfying the moment conditions) that

$$\lim_{n \rightarrow \infty} \mathbb{P}(\boldsymbol{\beta}_n^* \in \text{CS}_{n,p,1-\alpha_p}) = 0 \quad \text{implies} \quad \lim_{n \rightarrow \infty} \mathbb{P}(\boldsymbol{\beta}_n^* \in \text{CS}_{n,q,1-\alpha_q}) = 0; \quad (16)$$

that is any  $\boldsymbol{\beta}^*$  that is excluded from  $\text{CS}_{n,p,1-\alpha_p}$  with probability tending to one will also be excluded from  $\text{CS}_{n,q,1-\alpha_q}$  with probability tending to one. Also, if  $\boldsymbol{\beta}^*$  satisfies the two conditions in [\(14\)](#), with the limit inferior being zero, then

$$\limsup_{n \rightarrow \infty} \mathbb{P}(\boldsymbol{\beta}_n^* \in \text{CS}_{n,p,1-\alpha_p}) \rightarrow 1 - \alpha_p \quad \text{yet} \quad \lim_{n \rightarrow \infty} \mathbb{P}(\boldsymbol{\beta}_n^* \in \text{CS}_{n,q,1-\alpha_q}) = 0. \quad (17)$$

In words,  $\boldsymbol{\beta}^*$  then does not satisfy the moment conditions, yet it is included in the confidence set  $\text{CS}_{n,p,1-\alpha_p}$  with the same asymptotic probability as those parameters that do, whereas  $\text{CS}_{n,q,1-\alpha_q}$  correctly excludes this  $\boldsymbol{\beta}^*$  asymptotically. Thus, [\(16\)](#) and [\(17\)](#) give a sense in which basing confidence sets on larger  $p \in [2, \infty)$  results in “smaller” confidence sets. However, using results in [Pinelis \(2010\)](#), it is not difficult to show that already in the high-dimensional Gaussian location model one can have that  $\mathbb{E}h_n(\mathbf{X}_{1,n}, \boldsymbol{\beta}_n^*) \neq \mathbf{0}_d$  and

$$0 < \lim_{n \rightarrow \infty} \mathbb{P}(\boldsymbol{\beta}_n^* \in \text{CS}_{n,p,1-\alpha_p}) < \lim_{n \rightarrow \infty} \mathbb{P}(\boldsymbol{\beta}_n^* \in \text{CS}_{n,q,1-\alpha_q}).$$

Thus, it is not generally the case that eventually  $\text{CS}_{n,q,1-\alpha_q} \subseteq \text{CS}_{n,p,1-\alpha_p}$ .

### 3.4 Tests based on the supremum-norm: $p = \infty$

Tests based on the supremum-norm have received considerable attention due to the work on high-dimensional Gaussian approximations over hyperrectangles

$$\mathcal{H}_n = \left\{ \prod_{j=1}^d [a_j, b_j] \cap \mathbb{R} : -\infty \leq a_j \leq b_j \leq \infty, j = 1, \dots, d \right\}.$$

The following theorem exactly characterizes which alternatives tests based on the supremum-norm are consistent against under the same assumptions as those of Theorem 3.2.

**Theorem 3.4.** *Let  $\alpha \in (0, 1)$  and suppose Assumption 3.3 is satisfied. Then, the following holds.*

a) *Size control: Suppose  $\beta^* \in \mathbf{B}^{(0)}$ . A sequence of real numbers  $(\kappa_{n,\infty})_{n \in \mathbb{N}}$  satisfies*

$$\mathbb{P}(S_{n,\infty}(\beta_n^*) \geq \kappa_{n,\infty}) \rightarrow \alpha \quad (18)$$

$$\text{if and only if } \kappa_{n,\infty} = \kappa_{n,\infty}(\alpha) = \sqrt{2 \log(d)} - \frac{\log \log(d) + \log(4\pi)}{2\sqrt{2 \log(d)}} - \frac{\log(-\log(1-\alpha)/2) + o(1)}{\sqrt{2 \log(d)}}.$$

b) *Consistency: For  $(\kappa_{n,\infty})_{n \in \mathbb{N}}$  as in (18)*

$$\mathbb{P}(S_{n,\infty}(\beta_n^*) \geq \kappa_{n,\infty}) \rightarrow 1 \iff \sum_{i=1}^d \frac{\overline{\Phi}(\mathbf{c}_d - |\theta_{i,d}(\beta_n^*)|)}{\Phi(\mathbf{c}_d - |\theta_{i,d}(\beta_n^*)|)} \rightarrow \infty,$$

$$\text{where } \mathbf{c}_d := \sqrt{2 \log(d)} - \frac{\log \log(d)}{2\sqrt{2 \log(d)}} \text{ for } d \geq 2 \text{ and } \overline{\Phi} = 1 - \Phi \text{ (and one may set } \mathbf{c}_1 := 0).$$

Well-known *sufficient*, yet not necessary, conditions for consistency of the supremum-norm based test, such as  $\|\theta_n(\beta_n^*)\|_\infty - \sqrt{2 \log(d)} \rightarrow \infty$ , are trivial special cases of Theorem 3.4. In particular, the  $\infty$ -norm based test is consistent against  $(\sqrt{3 \log(d)}, 0, \dots, 0)$  whereas by (12) of Theorem 3.1 no  $p$ -norm based test with  $p \in [2, \infty)$  is consistent.

Concerning dense alternatives of the form  $\theta_n(\beta_n^*) = (c_n, \dots, c_n)$ , it follows that the supremum-norm based test is consistent if and only if  $\sqrt{\log(d)}|c_n| \rightarrow \infty$ , cf. Appendix A.4 of Kock and Preinerstorfer (2023). Thus, the supremum-norm based test is *not* consistent if  $c_n = 1/\sqrt{\log(d)}$  whereas by (12) of Theorem 3.1 every  $p$ -norm based test with  $p \in [2, \infty)$  is consistent. Hence, i) the monotonicity in (13) does not extend to  $q = \infty$  and ii) no single  $p$ -norm based test dominates all others in terms of its consistency properties.

The exact characterization of consistency in Theorem 3.4 sheds new light on the frequently used supremum-norm based test for testing moment equalities. It is more informative than the currently available power analysis of the supremum-norm based test through a maximin-lens. The latter provides the sufficient condition for consistency  $\|\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*)\|_\infty - \sqrt{2\log(d)} \rightarrow \infty$ , but is silent about the alternatives not satisfying this condition that the supremum-norm based test is nevertheless consistent against. For example, if  $\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*) = (1, \dots, 1)'$ , then by the observation on dense alternatives in the previous paragraph the supremum-norm based test *is* consistent, yet  $\|\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*)\|_\infty - \sqrt{2\log(d)} \not\rightarrow \infty$ .

Concerning confidence sets, Theorem 3.4 shows that for a given  $\alpha \in (0, 1)$ ,  $\kappa_{n,\infty}(\alpha)$  as in (18), and  $\boldsymbol{\beta}^* \in \mathbf{B}^{(0)}$ , the sets  $\mathbf{CS}_{n,\infty,1-\alpha} := \{\boldsymbol{\beta} \in \mathbf{B}_n : S_{n,\infty}(\boldsymbol{\beta}) \leq \kappa_{n,\infty}(\alpha)\}$  satisfy  $\lim_{n \rightarrow \infty} \mathbb{P}(\boldsymbol{\beta}_n^* \in \mathbf{CS}_{n,\infty,1-\alpha}) = 1 - \alpha$ .

*Remark 3.5.* To prove Theorem 3.4, Lemma F.1 establishes the following variant of (15) which, crucially for our characterization of consistency, holds for  $\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*)$  further away from the origin than required for (15):

$$\sup_{H \in \mathcal{H}_n} \left| \mathbb{P}(\hat{\boldsymbol{\Sigma}}_n^{-1/2}(\boldsymbol{\beta}_n^*) \mathbf{H}_n(\boldsymbol{\beta}_n^*) \in H) - \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*) \in H) \right| \rightarrow 0.$$

## 4 A test dominating all $p$ -norm based test for $p \in [2, \infty]$

The discussion following Theorem 3.4 revealed that no single  $p$ -norm based test is “best” in terms of the amount of alternatives it is consistent against. This makes choosing a  $p$  to base a test on very difficult. Even if one *knows* that the deviation from the null  $\mathbb{E}h_n(\mathbf{X}_{1,n}, \boldsymbol{\beta}_n^*)$  is sparse, which *could* suggest using a test based on the supremum-norm, sparsity need not be inherited by the noncentrality parameter

$$\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*) = \sqrt{n} \boldsymbol{\Sigma}_n^{-1/2}(\boldsymbol{\beta}_n^*) \mathbb{E}h_n(\mathbf{X}_{1,n}, \boldsymbol{\beta}_n^*)$$

due to the presence of  $\boldsymbol{\Sigma}_n^{-1/2}(\boldsymbol{\beta}_n^*)$ . Therefore, one often does not know anything about the structure of  $\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*)$ , which is the actual quantity entering the characterization of the power of  $p$ -norm based tests in Theorems 3.1 and 3.4. This underscores the importance of having a test that is powerful irrespective of the unknown structure of  $\boldsymbol{\theta}_n(\boldsymbol{\beta}_n^*)$ .

One could hope that combining tests based on the two “endpoints”  $p = 2$  and  $p = \infty$ , e.g. by the power enhancement principle of Fan et al. (2015), of the interval  $[2, \infty]$  results in a test that is consistent whenever *some*  $p$ -norm based test for  $p \in [2, \infty]$  is

consistent. However, Corollary 4.2 in [Kock and Preinerstorfer \(2023\)](#) shows that already in the Gaussian sequence model this is not the case: There exist (semi-sparse) alternatives against which tests based on  $p = 2$  and  $p = \infty$  are inconsistent, but against which a test based on any single  $p \in (2, \infty)$  is consistent. To construct a test that is consistent whenever a test based on some  $p \in [2, \infty]$  is consistent, we use a construction similar in spirit to the one employed in the Gaussian sequence model in [Kock and Preinerstorfer \(2023\)](#). A crucial difference is that the present construction explicitly includes the supremum-norm based test rather than using a  $p_n$ -norm based test with sufficiently quickly increasing  $p_n$  to dominate the former.

#### 4.1 The dominant test

Let  $\alpha \in (0, 1)$  and  $\alpha_2, \alpha_I$  and  $\alpha_\infty$  be non-negative with  $\alpha_2 + \alpha_I + \alpha_\infty = \alpha$  as well as

$$\mathbb{P}(\|\mathbf{Z}_d\|_2 \geq \kappa_{n,2}) = \alpha_2 \quad \text{and} \quad \mathbb{P}(\|\mathbf{Z}_d\|_\infty \geq \kappa_{n,\infty}) = \alpha_\infty.$$

Furthermore, let  $p_n$  be a strictly increasing and unbounded sequence in  $(2, \infty)$  and let  $m_n$  be a non-decreasing and unbounded sequence in  $\mathbb{N}$ . Fix an array

$$\mathcal{A} = \left\{ \alpha_{n,p_j} \in (0, 1) : n \in \mathbb{N}, j = 1, \dots, m_n \right\}$$

such that

$$\sum_{j=1}^{m_n} \alpha_{n,p_j} = \alpha_I \text{ for every } n \in \mathbb{N} \quad \text{and} \quad \lim_{n \rightarrow \infty} \alpha_{n,p_j} > 0 \text{ for every } j \in \mathbb{N},$$

where the conditions implicitly impose the existence of the respective limits. For every  $n \in \mathbb{N}$  and every  $j = 1, \dots, m_n$ , choose  $\kappa_{n,p_j} > 0$  and  $c_n \in (0, 1]$  such that

$$\mathbb{P}\left(\|\mathbf{Z}_d\|_{p_j} \geq \kappa_{n,p_j}\right) = \alpha_{n,p_j} \quad \text{and} \quad \mathbb{P}\left(\max_{p \in \mathfrak{P}_n} \kappa_{n,p}^{-1} \|\mathbf{Z}_d\|_p \geq c_n\right) = \alpha, \quad (19)$$

where  $\mathfrak{P}_n = \{2, p_1, \dots, p_{m_n}, \infty\}$ . Define the test  $\psi_n(\boldsymbol{\beta}_n^*)$  as

$$\psi_n(\boldsymbol{\beta}_n^*) := \mathbf{1} \left\{ \max_{p \in \mathfrak{P}_n} \kappa_{n,p}^{-1} S_{n,p}(\boldsymbol{\beta}_n^*) \geq c_n \right\}. \quad (20)$$

Thus,  $\psi_n(\boldsymbol{\beta}_n^*)$  rejects  $H_{0,n} : \mathbb{E}h_n(\mathbf{X}_{1,n}, \boldsymbol{\beta}_n^*) = \mathbf{0}_d$  if for any  $p$ -norm based tests with  $p \in \mathfrak{P}_n$  it is the case that  $S_{n,p}(\boldsymbol{\beta}_n^*)$  exceeds  $c_n \kappa_{n,p}$ . The tests based on  $p = 2$  and  $p = \infty$  are included in the construction of  $\psi_n(\boldsymbol{\beta}_n^*)$  to make it powerful against dense and sparse alternatives, respectively. The tests based on  $p \in \{p_1, \dots, p_{m_n}\}$  are included to cover the (semi-sparse) alternatives that neither tests based on the 2- or supremum-norm are consistent against. The quantities  $\alpha_2$ ,  $\alpha_I$ , and  $\alpha_\infty$  control how much size is distributed to the components of  $\psi_n(\boldsymbol{\beta}_n^*)$  targeting dense, semi-sparse and sparse alternatives, respectively while the array  $\mathcal{A}$  distributes the size  $\alpha_I$  to the  $p$ -norm based tests with  $p \in \{p_1, \dots, p_{m_n}\}$ . The role of  $c_n$  is to ensure that  $\psi_n(\boldsymbol{\beta}_n^*)$  has asymptotic size  $\alpha$ , cf. Part 1 of Theorem 4.1 below. Choosing  $c_n = 1$  will generally result in  $\psi_n(\boldsymbol{\beta}_n^*)$  having asymptotic size not larger but potentially smaller than  $\alpha$ , i.e. a conservative asymptotic level  $\alpha$  test.

**Theorem 4.1.** *Let  $\alpha \in (0, 1)$  and suppose Assumption 3.3 is satisfied. Then, the following holds.*

1. *Size control: If  $\boldsymbol{\beta}^* \in \mathbf{B}^{(0)}$ , then  $\lim_{n \rightarrow \infty} \mathbb{E}\psi_n(\boldsymbol{\beta}_n^*) = \alpha$ .*
2. *Dominance: For any  $p \in [2, \infty]$ ,  $\bar{\alpha}_p \in (0, 1)$ , and  $(\bar{\kappa}_{n,p}(\bar{\alpha}_p))_{n \in \mathbb{N}}$  given by (10) in case  $p \in [2, \infty)$  and by (18) in case  $p = \infty$  it holds that*

$$\mathbb{P}(S_{n,p}(\boldsymbol{\beta}_n^*) \geq \bar{\kappa}_{n,p}(\bar{\alpha}_p)) \rightarrow 1 \quad \text{implies} \quad \mathbb{E}\psi_n(\boldsymbol{\beta}_n^*) \rightarrow 1.$$

Part 1 of Theorem 4.1 shows that the critical values  $\kappa_{n,p}$ ,  $p \in \mathfrak{P}_n$ , and the  $c_n$  that guarantee exact size  $\alpha \in (0, 1)$  under Gaussianity in (19) also guarantee that  $\psi_n(\boldsymbol{\beta}_n^*)$  has asymptotic size  $\alpha$ . This may not be obvious ex ante as the construction of  $\psi_n(\boldsymbol{\beta}_n^*)$  is based on combining  $|\mathfrak{P}_n| = (m_n + 2) \rightarrow \infty$  tests. However, having established (15), this becomes a trivial consequence of writing non-rejection of  $\psi_n(\boldsymbol{\beta}_n^*)$  as the event

$$\left\{ \hat{\boldsymbol{\Sigma}}_n^{-1/2}(\boldsymbol{\beta}_n^*) \mathbf{H}_n(\boldsymbol{\beta}_n^*) \in \bigcap_{p \in \mathfrak{P}_n} \mathbb{B}_p(c_n \kappa_{n,p}) \right\}$$

where for  $r \in (0, \infty)$  we set  $\mathbb{B}_p(r) = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_p < r\}$  and observe that  $\bigcap_{p \in \mathfrak{P}_n} \mathbb{B}_p(c_n \kappa_{n,p})$  is convex.

Part 2 shows that  $\psi_n(\boldsymbol{\beta}_n^*)$  is consistent against *every* deviation from the null hypothesis that *some*  $p$ -norm based test with  $p \in [2, \infty]$  is consistent against (irrespective of the asymptotic size of the latter, as long as it is a number in  $(0, 1)$ ). In particular,  $\psi_n(\boldsymbol{\beta}_n^*)$

is consistent whenever tests based on the 2- or  $\infty$ -norm with asymptotic size in  $(0, 1)$  are consistent. Recall that the (dense and sparse) alternatives that tests based on these norms are consistent against are the ones targeted by current applications of the power enhancement principle of [Fan et al. \(2015\)](#), cf. also [Kock and Preinerstorfer \(2019\)](#). In addition,  $\psi_n(\boldsymbol{\beta}_n^*)$  is consistent against further alternatives, as it is also consistent as soon as there exists a  $p \in (2, \infty)$  such that a test based on this  $p$  is consistent (a property that tests based on the power enhancement principle that combine 2- and  $\infty$ -norm based tests do not share, cf. [Kock and Preinerstorfer \(2023\)](#)).

Furthermore, as a consequence of [Theorems 3.1 and 3.4](#), it is not possible to choose a single best  $p$ -norm to base a test on: Our results show that no matter which *single*  $p \in [2, \infty]$  one chooses, there exists another  $p$ -norm based test that is consistent against alternatives that the test based on the chosen  $p$  is inconsistent against. [Theorem 4.1](#) suggests a way to overcome this impossibility of choosing a single “best” norm to base a tests on as  $\psi_n(\boldsymbol{\beta}_n^*)$  simultaneously dominates all  $p$ -norm based tests for  $p \in [2, \infty]$ .

[Theorem 4.1](#) is silent about the power properties of  $\psi_n(\boldsymbol{\beta}_n^*)$  when no  $p$ -norm based test is consistent. Thus, there could, in principle, exist alternatives against which a  $p$ -norm based test has substantially higher asymptotic power than  $\psi_n(\boldsymbol{\beta}_n^*)$  without the former converging to one. The following theorem shows that this possibility can be ruled out by allocating sufficient asymptotic size  $\alpha_p$  to a given  $p$ -norm in the construction of  $\psi_n(\boldsymbol{\beta}_n^*)$ .

**Theorem 4.2.** *Let  $\alpha \in (0, 1)$  and suppose [Assumption 3.3](#) is satisfied. In the context of the construction of  $\psi_n(\boldsymbol{\beta}_n^*)$  fix  $p \in \cup_{n \in \mathbb{N}} \mathfrak{P}_n$  and define  $\alpha_p := \lim_{n \rightarrow \infty} \alpha_{n,p} \in (0, 1)$  if  $p \in (2, \infty)$ .<sup>8</sup>*

1. *If  $p \in [2, \infty)$ , then for any sequence  $(\bar{\kappa}_{n,p}(\alpha))_{n \in \mathbb{N}}$  as in [\(10\)](#)*

$$\limsup_{n \rightarrow \infty} \left[ \mathbb{P} \left( S_{n,p}(\boldsymbol{\beta}_n^*) \geq \bar{\kappa}_{n,p}(\alpha) \right) - \mathbb{E} \psi_n(\boldsymbol{\beta}_n^*) \right] \leq \frac{\Phi^{-1}(1 - \alpha_p) - \Phi^{-1}(1 - \alpha)}{\sqrt{2\pi}}.$$

2. *If  $p = \infty$ , then for any sequence  $(\bar{\kappa}_{n,\infty}(\alpha))_{n \in \mathbb{N}}$  as in [\(18\)](#)*

$$\limsup_{n \rightarrow \infty} \left[ \mathbb{P} \left( S_{n,\infty}(\boldsymbol{\beta}_n^*) \geq \bar{\kappa}_{n,\infty}(\alpha) \right) - \mathbb{E} \psi_n(\boldsymbol{\beta}_n^*) \right] \leq f(\alpha) - f(\alpha_\infty),$$

where  $f : (0, 1) \rightarrow \mathbb{R}$  is defined via  $x \mapsto \log(-\log(1-x)/2)$ .

---

<sup>8</sup>Note that  $\alpha_2$  and  $\alpha_\infty$  have been defined already in the construction of  $\psi_n(\boldsymbol{\beta}_n^*)$ .

Theorem 4.2 shows that compared to the  $p$ -norm based test of asymptotic size  $\alpha$  and with exponent  $p$ , there are no alternatives for which much is lost in terms of asymptotic power by using a test  $\psi_n(\beta_n^*)$  with  $\alpha_p$  close to  $\alpha$ . On the other hand: i)  $\psi_n(\beta_n^*)$  is consistent against any alternative that some  $p$ -norm based test is consistent against and ii) alternatives can exist against which  $\psi_n(\beta_n^*)$  is consistent yet the  $p$ -norm based test has asymptotic power equal to its asymptotic size. Thus, there can be much to gain from using  $\psi_n(\beta_n^*)$ . Note, however, that the above construction of a test necessitates the choice of  $p$  to target. That is,  $\psi_n(\beta_n^*)$  with asymptotic size  $\alpha$  cannot have power arbitrarily close to, e.g., that of the 2-norm and  $\infty$ -norm based tests (each with asymptotic size  $\alpha$ ) simultaneously.

## 5 Many IVs and dominating the Anderson-Rubin test

Recall that in the context of the linear IV model in Section 2.1.1 one has

$$h_n(\mathbf{X}_{1,n}, \beta_n^*) = (y_{1,n} - \mathbf{Y}'_{1,n} \beta_n^*) \mathbf{z}_{1,n}.$$

The theory developed in the previous sections has the following consequences for testing whether a given  $\beta_n^*$  satisfies  $\mathbb{E}(y_{1,n} - \mathbf{Y}'_{1,n} \beta_n^*) \mathbf{z}_{1,n} = \mathbf{0}_d$ :

1. Choosing  $p = 2$  results in a classic weak identification robust Anderson-Rubin (AR) test. Thus, this AR test is dominated in terms of consistency by any  $p$ -norm based test with  $p \in (2, \infty)$ , cf. the relationship in (13). There is no ranking between tests based on  $p \in [2, \infty)$  and the sup-score type test corresponding to  $p = \infty$ ; see Belloni et al. (2012) for the definition of the original sup-score statistic.
2. For simplicity, let  $\mathbf{B}_n = \mathbb{R}$  and assume that  $\beta_n$  satisfies  $\mathbb{E}(y_{1,n} - \beta_n \mathbf{Y}_{1,n}) \mathbf{z}_{1,n} = \mathbf{0}_d$ . Even if only one instrument (say) is relevant in the sense of  $\mathbb{E} \mathbf{z}_{1,n} \mathbf{Y}_{1,n} = (a_n, 0, \dots, 0)'$  for some  $a_n \neq 0$  such that the moments

$$\mathbb{E}(y_{1,n} - \beta_n^* \mathbf{Y}_{1,n}) \mathbf{z}_{1,n} = \mathbb{E} \mathbf{z}_{1,n} \mathbf{Y}_{1,n} (\beta_n - \beta_n^*) = (a_n (\beta_n - \beta_n^*), 0, \dots, 0)'$$

only differ from zero in one entry, this sparsity need not be inherited by

$$\boldsymbol{\theta}_n(\beta_n^*) = \sqrt{n} [\boldsymbol{\Sigma}_n(\beta_n^*)]^{-1/2} (a_n (\beta_n - \beta_n^*), 0, \dots, 0)'$$

unless one is willing to impose structure on the covariance matrix  $\boldsymbol{\Sigma}_n(\beta_n^*)$  of  $(y_{1,n} -$



$\beta_n^* \mathbf{Y}_{1,n} \mathbf{z}_{1,n}$ . Thus, as it is  $\theta_n(\beta_n^*)$  that determines the power properties of  $p$ -norm based tests, one cannot advise on which  $p$  to use solely based on the number of instruments that one suspects to be relevant. Similarly, one should be cautious basing advice on the structure of  $\pi_n$  in the “first-stage”  $Y_{1,n} = \pi_n' \mathbf{z}_{1,n} + \nu_{1,n}$ : Since

$$\pi_n = [\mathbb{E}(\mathbf{z}_{1,n} \mathbf{z}_{1,n}')^{-1} \mathbb{E} \mathbf{z}_{1,n} \mathbf{Y}_{1,n}$$

(assuming that  $\mathbb{E}(\mathbf{z}_{1,n} \mathbf{z}_{1,n}')$  is invertible) one can write  $\mathbb{E} \mathbf{z}_{1,n} \mathbf{Y}_{1,n} = \mathbb{E}(\mathbf{z}_{1,n} \mathbf{z}_{1,n}') \pi_n$  and hence

$$\theta_n(\beta_n^*) = \sqrt{n} [\boldsymbol{\Sigma}_n(\beta_0)]^{-1/2} \mathbb{E}(\mathbf{z}_{1,n} \mathbf{z}_{1,n}') \pi_n \cdot (\beta_n - \beta_n^*),$$

cf. the first equality in the penultimate display. Again there is no link between the sparsity pattern of  $\pi_n$  and that of  $\theta_n(\beta_n^*)$ ; the latter determining the power properties of  $p$ -norm based tests.

3. In light of the previous point, it is difficult to give advice on which single  $p$ -norm to base a test on even if one is willing to impose assumptions on the number of relevant instruments. Thus, it is useful that the test  $\psi_n(\beta_n^*)$  in (20) is consistent whenever some  $p$ -norm based test (including the AR- or sup-score type tests) is consistent. Therefore,  $\psi_n(\beta_n^*)$  is consistent against strictly more alternatives than any test based on a single  $p$ . In this sense,  $\psi_n(\beta_n^*)$  does not rely on knowing whether  $\theta_n(\beta_n^*)$  is sparse or not.

## 6 Rapidly increasing dimension via sample splitting

The results presented so far generally restrict the growth rate of  $d$ . Suppose, however, that one wishes to test whether  $\beta_n^*$  satisfies  $D = D(n)$  moment conditions

$$\mathbb{E} h_{j,n}(\mathbf{X}_{1,n}, \beta_n^*) = 0, \quad j = 1, \dots, D, \quad (21)$$

with  $D$  *unrestricted*. Under independent sampling one can use a two-step procedure to test the validity of (21) by means of the tests developed in the previous sections: Based on a partition of the sample into two disjoint subsamples  $N_1 = N_1(n) \subset \{1, \dots, n\}$  and  $N_2 = N_2(n) \subset \{1, \dots, n\}$  of sizes  $n_1 = n_1(n)$  and  $n_2 = n_2(n)$ , respectively, one

first uses the subsample  $N_1$  to select a subset  $S = S(n_1, n_2, \{\mathbf{X}_{i,n}\}_{i \in N_1}) \subset \{1, \dots, D\}$  of moment functions. In a second step, using the subsample  $N_2$ , one tests whether  $\beta_n^*$  satisfies

$$\mathbb{E}h_{j,n}(\mathbf{X}_{1,n}, \beta_n^*) = 0, \quad j \in S$$

using only the data  $\{\mathbf{X}_{i,n}\}_{i \in N_2}$ . In Theorem J.1 in Section J in the appendix we show that if a predetermined fixed number of *selected* moments  $|S| = d(n_2)$  satisfies the growth conditions of the theory developed in the previous sections relative to the second step “sample size”  $n_2$ , then all asymptotic size guarantees established so far remain valid. The concrete moment selection mechanism  $S$  does not affect the asymptotic size of the subsequent tests as independent sampling guarantees that no selection bias is introduced. However, it is clear that the power of a test depends on  $S$  including moments that are violated. Some ways to select the subset of moments  $S$  to be tested in order to obtain powerful tests are discussed in Appendix J, where the size-control result informally sketched above is also formally established in Theorem J.1.

## A Notation

For brevity, we define  $\mathbf{Y}_{i,n} = h_n(\mathbf{X}_{i,n}, \boldsymbol{\beta}_n^*)$  and write  $\boldsymbol{\mu}_n = \mathbb{E}\mathbf{Y}_{i,n}$  as well as  $\boldsymbol{\Sigma}_n = \boldsymbol{\Sigma}_n(\boldsymbol{\beta}_n^*)$ . Write  $\boldsymbol{\theta}_n := \sqrt{n}\boldsymbol{\Sigma}_n^{-1/2}\boldsymbol{\mu}_n$  and let  $\rightsquigarrow$  denote convergence in distribution. We shall also use that by Lemma C.1 in [Kock and Preinerstorfer \(2023\)](#) for every  $p \in [2, \infty)$  there exists a constant  $c_p \in (0, \infty)$  such that

$$c_p^{-1}g_p(x) \leq \lambda_p(x) - \lambda_p(0) \leq c_p g_p(x) \quad \text{for all } x \in \mathbb{R}, \quad (\text{A.1})$$

where  $g_p$  and  $\lambda_p$  are defined in (6). Furthermore, for every  $p \in [1, \infty)$  and for a matrix  $\mathbf{A}$  we denote by  $\|\mathbf{A}\|_p$  the matrix norm induced by the vector  $p$ -norm  $\|\cdot\|_p$ .

## B Proof of Theorem 3.1

We start with Part 1 and note that  $\boldsymbol{\beta}^* \in \mathbf{B}^{(0)}$  in this case due to Assumption 3.1, which also implies  $\boldsymbol{\theta}_n = \mathbf{0}_d$ . Fix  $\alpha \in (0, 1)$ ,  $p \in [2, \infty)$  and a sequence of real numbers  $\kappa_n$ . First, observe that  $\left\|n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n}\right\|_p \geq \kappa_n$  is equivalent to

$$\begin{aligned} \frac{\left\|n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n}\right\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sqrt{d}\sigma_p} + \frac{\sum_{i=1}^d (\lambda_p(\theta_{i,n}) - \lambda_p(0))}{\sqrt{d}\sigma_p} \\ \geq \frac{\kappa_n^p - d\lambda_p(0)}{\sqrt{d}\sigma_p} =: \bar{\kappa}_n. \end{aligned} \quad (\text{B.1})$$

Under Assumption 3.1, the second term in the first line of (B.1) vanishes and the first converges in distribution to a standard normal distribution, so that Polya's theorem shows that

$$\mathbb{P}\left(\left\|n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n}\right\|_p \geq \kappa_n\right) \rightarrow \alpha$$

if and only if  $\bar{\kappa}_n \rightarrow \Phi^{-1}(1 - \alpha) =: z_{1-\alpha}$ , which establishes Part 1 of Theorem 3.1.

To establish Part 2 and “ $\implies$ ” in (11) of Part 3, let  $\boldsymbol{\beta}^* \in \mathbf{B}^*$  and consider (more

generally than needed for Part 2) a subsequence  $n'$  of  $n$  along which

$$\frac{1}{\sqrt{d'}} \sum_{i=1}^{d'} [\lambda_p(\theta_{i,n'}) - \lambda_p(0)] \rightarrow c \in \mathbb{R}. \quad (\text{B.2})$$

Then, by (8), (B.1), and  $\bar{\kappa}_n \rightarrow z_{1-\alpha}$ , along this subsequence

$$\lim_{n' \rightarrow \infty} \mathbb{P} \left( \left\| n'^{-1/2} \sum_{i=1}^{n'} \hat{\Sigma}_{n'}^{-1/2} \mathbf{Y}_{i,n'} \right\|_p \geq \kappa_{n'} \right) = 1 - \Phi(z_{1-\alpha} - c/\sigma_p) < 1.$$

Thus, Part 2 follows by the above with “ $n' = n$ ” while “ $\implies$ ” in (11) of Part 3 follows from observing that  $\frac{1}{\sqrt{d}} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)] \not\rightarrow \infty$  implies the existence of a subsequence along which (B.2) holds for some  $c \in \mathbb{R}$ .

We next establish “ $\Leftarrow$ ” of (11) of Part 3. Let  $\beta^* \in \mathbf{B}^*$ . The rejection event in (B.1) can be rewritten as

$$\frac{\left\| n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} + 1 - \bar{\kappa}_n \frac{\sqrt{d} \sigma_p}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} \geq 0.$$

Hence, from  $d^{-1/2} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)] \rightarrow \infty$  and (9) together with  $\bar{\kappa}_n \rightarrow \Phi^{-1}(1 - \alpha)$ , it follows that

$$\mathbb{P} \left( \left\| n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p \geq \kappa_n \right) \rightarrow 1.$$

Finally, (12) of Part 3 follows from (A.1).

## C Gaussian approximation over convex sets with estimated covariance matrix and $d \rightarrow \infty$

### C.1 Estimating $\Sigma_n^{-1/2}$

Recall that for any  $d \times d$  matrix  $\mathbf{A}$  we denote by  $\mathbf{A}^{-1}$  its Moore-Penrose generalized inverse and define  $\mathbf{A}^{-1/2} := (\mathbf{A}^{1/2})^{-1}$ , in case  $\mathbf{A}$  is symmetric and positive semidefinite, cf. Footnote 4. The following result is well known and immediately follows from, e.g., Equations (7.2.13) and (5.8.4) in Horn and Johnson (1985). We provide the argument for

completeness. Recall also that  $\hat{\Sigma}$  is positive semidefinite and symmetric by assumption.

**Lemma C.1.** *Let  $n'$  be a subsequence of  $n$ , assume that the eigenvalues of  $\Sigma_{n'}$  are (uniformly) bounded away from zero and from above, and that  $\|\hat{\Sigma}_{n'} - \Sigma_{n'}\|_2 = O_{\mathbb{P}}(a_{n'})$  for some  $a_{n'} \rightarrow 0$ . Then,  $\|\hat{\Sigma}_{n'}^{-1/2} - \Sigma_{n'}^{-1/2}\|_2 = O_{\mathbb{P}}(a_{n'})$ .*

*Proof.* Write  $n$  instead of  $n'$ . Since  $\|\hat{\Sigma}_n - \Sigma_n\|_2 \rightarrow 0$  in probability and the eigenvalues of  $\Sigma_n$  are uniformly bounded away from zero and infinity,  $\hat{\Sigma}_n$  is invertible with probability tending to one. Denote by  $G_n$  the event on which  $\hat{\Sigma}_n$ , and hence  $\hat{\Sigma}_n^{1/2}$ , is invertible. On  $G_n$  one has that  $\hat{\Sigma}_n^{-1/2}$  is the regular matrix inverse of  $\hat{\Sigma}_n^{1/2}$  and so

$$\|\hat{\Sigma}_n^{-1/2} - \Sigma_n^{-1/2}\|_2 = \|(\hat{\Sigma}_n^{1/2})^{-1} - (\Sigma_n^{1/2})^{-1}\|_2 \mathbf{1}_{G_n} + \|\hat{\Sigma}_n^{-1/2} - \Sigma_n^{-1/2}\|_2 \mathbf{1}_{G_n^c},$$

where, with a slight overload of notation, the matrix inverse is now a regular inverse on  $G_n$ . Equations (7.2.13) and (5.8.4) in [Horn and Johnson \(1985\)](#) guarantee that

$$\|\hat{\Sigma}_n^{-1/2} - \Sigma_n^{-1/2}\|_2 \leq O_{\mathbb{P}}(a_n) + \|\hat{\Sigma}_n^{-1/2} - \Sigma_n^{-1/2}\|_2 \mathbf{1}_{G_n^c},$$

and the conclusion follows from  $\mathbb{P}(G_n^c) \rightarrow 0$ .  $\square$

## C.2 An approximation result

**Lemma C.2.** *Let  $p \in [2, \infty)$ ,  $n'$  be a subsequence of  $n$  and  $d' \rightarrow \infty$ . Assume that the eigenvalues of  $\Sigma_{n'}$  are (uniformly) bounded away from zero and from above and that  $\|\hat{\Sigma}_{n'} - \Sigma_{n'}\|_2 = O_{\mathbb{P}}(a_{n'})$  for some  $a_{n'} \rightarrow 0$ , as well as*

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P} \left( \left\| n'^{-1/2} \sum_{i=1}^{n'} \Sigma_{n'}^{-1/2} \mathbf{Y}_{i,n'} \right\|_p \leq t \right) - \mathbb{P} (\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p \leq t) \right| \rightarrow 0. \quad (\text{C.1})$$

1. *If  $\frac{1}{d'} \sum_{i=1}^{d'} g_p(\theta_{i,n'})$  is bounded, then*

$$\left\| n'^{-1/2} \sum_{i=1}^{n'} \hat{\Sigma}_{n'}^{-1/2} \mathbf{Y}_{i,n'} - n'^{-1/2} \sum_{i=1}^{n'} \Sigma_{n'}^{-1/2} \mathbf{Y}_{i,n'} \right\|_p = O_{\mathbb{P}}(d'^{1/2} a_{n'}).$$

2. If  $\frac{1}{d'} \sum_{i=1}^{d'} g_p(\theta_{i,n'}) \rightarrow \infty$ , then

$$\left\| n'^{-1/2} \sum_{i=1}^{n'} \hat{\Sigma}_{n'}^{-1/2} \mathbf{Y}_{i,n'} - n'^{-1/2} \sum_{i=1}^{n'} \Sigma_{n'}^{-1/2} \mathbf{Y}_{i,n'} \right\|_p = O_{\mathbb{P}} \left( d'^{\frac{1}{2} - \frac{1}{p}} \left[ \sum_{i=1}^{d'} g_p(\theta_{i,n'}) \right]^{\frac{1}{p}} a_{n'} \right).$$

*Proof.* Let  $p \in [2, \infty)$  and write  $n$  instead of  $n'$ . Note that

$$\begin{aligned} & \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} - \frac{1}{\sqrt{n}} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p \\ & \leq \left\| (\hat{\Sigma}_n^{-1/2} - \Sigma_n^{-1/2}) \Sigma_n^{1/2} \right\|_p \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p \\ & \leq d^{\frac{1}{2} - \frac{1}{p}} \left\| \Sigma_n^{1/2} \right\|_2 \left\| \hat{\Sigma}_n^{-1/2} - \Sigma_n^{-1/2} \right\|_2 \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p \\ & = d^{\frac{1}{2} - \frac{1}{p}} O_{\mathbb{P}}(a_n) \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p \end{aligned} \tag{C.2}$$

where the second inequality follows from (1.2) of Theorem 1 in [Goldberg \(1987\)](#) and the equality follows from Lemma [C.1](#) together with the uniform boundedness assumption on the eigenvalues of  $\Sigma_n$ . To proceed, observe that by [\(A.1\)](#)

$$\sum_{i=1}^d \lambda_p(\theta_{i,n}) = \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)] + d\lambda_p(0) \leq c_p \sum_{i=1}^d g_p(\theta_{i,n}) + d\lambda_p(0). \tag{C.3}$$

Consider first the case where  $d^{-1/2} \sum_{i=1}^d g_p(\theta_{i,n})$  is bounded, which is only possible in Part 1: Then, Lemma C.3 in [Kock and Preinerstorfer \(2023\)](#) gives

$$\frac{\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sqrt{d \text{Var}|Z|^p}} \rightsquigarrow \mathbf{N}_1(0, 1),$$

which together with [\(C.1\)](#) yields [use “ $t = (x \cdot \sqrt{d \text{Var}|Z|^p} + \sum_{i=1}^d \lambda_p(\theta_{i,n}))^{1/p}$ ” and  $x \in \mathbb{R}$ ]

$$\frac{\|n^{-1/2} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n}\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sqrt{d \text{Var}|Z|^p}} \rightsquigarrow \mathbf{N}_1(0, 1).$$

Therefore, by (C.3), and exploiting boundedness of  $d^{-1/2} \sum_{i=1}^d g_p(\theta_{i,n})$ ,

$$\left\| n^{-1/2} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p = O_{\mathbb{P}} \left( \left[ \sum_{i=1}^d \lambda_p(\theta_{i,n}) \right]^{1/p} + \left[ d \text{Var}|Z|^p \right]^{1/2p} \right) = O_{\mathbb{P}}(d^{1/p}),$$

which together with (C.2) implies Part 1 when  $d^{-1/2} \sum_{i=1}^d g_p(\theta_{i,n})$  is bounded.

Consider next the case where  $d^{-1/2} \sum_{i=1}^d g_p(\theta_{i,n}) \rightarrow \infty$  (potentially along a subsequence  $n''$  of  $n'$ , which we again denote by  $n$  for simplicity of notation), a condition that is always satisfied in Part 2: By (C.23) in [Kock and Preinerstorfer \(2023\)](#),

$$\frac{\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d g_p(\theta_{i,n})} = o_{\mathbb{P}}(1),$$

which together with (C.1) yields

$$\frac{\|n^{-1/2} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n}\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d g_p(\theta_{i,n})} = o_{\mathbb{P}}(1).$$

Hence,

$$\left\| n^{-1/2} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p = O_{\mathbb{P}} \left( \left[ \sum_{i=1}^d \lambda_p(\theta_{i,n}) \right]^{1/p} + \left[ \sum_{i=1}^d g_p(\theta_{i,n}) \right]^{1/p} \right). \quad (\text{C.4})$$

Together with (C.3) this yields

$$\left\| n^{-1/2} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p = \begin{cases} O_{\mathbb{P}}(d^{1/p}) & \text{if } \frac{1}{d} \sum_{i=1}^d g_p(\theta_{i,n}) \text{ is bounded} \\ O_{\mathbb{P}}([\sum_{i=1}^d g_p(\theta_{i,n})]^{1/p}) & \text{if } \frac{1}{d} \sum_{i=1}^d g_p(\theta_{i,n}) \rightarrow \infty. \end{cases} \quad (\text{C.5})$$

Together with (C.2), the first case in (C.5) now also establishes Part 1 in the remaining case where  $d^{-1/2} \sum_{i=1}^d g_p(\theta_{i,n}) \rightarrow \infty$  (along subsequences). Furthermore, the second case in (C.5) establishes Part 2.  $\square$

### C.3 Gaussian approximation

For any non-empty  $A \subseteq \mathbb{R}^d$ ,  $\varepsilon > 0$ , and  $p \in [1, \infty]$ , denote

$$A^\varepsilon = \{\mathbf{x} \in \mathbb{R}^d : \inf_{\mathbf{y} \in A} \|\mathbf{x} - \mathbf{y}\|_2 \leq \varepsilon\} \quad \text{and} \quad \mathcal{B}_p(\mathbf{x}, \varepsilon) = \{\mathbf{y} \in \mathbb{R}^d : \|\mathbf{y} - \mathbf{x}\|_p \leq \varepsilon\}.$$

We abbreviate  $\mathcal{B}_p(\varepsilon) = \mathcal{B}_p(\mathbf{0}_d, \varepsilon)$ , and set

$$A^{-\varepsilon} = \{\mathbf{x} \in \mathbb{R}^d : \mathcal{B}_2(\mathbf{x}, \varepsilon) \subseteq A\}.$$

Recall that  $\mathcal{C}_n = \{C \subseteq \mathbb{R}^{d(n)} : C \text{ is convex and Borel measurable}\}$ .

**Lemma C.3.** *Let  $n'$  be a subsequence of  $n$  and  $d' \rightarrow \infty$ . Assume that the eigenvalues of  $\Sigma_{n'}$  are (uniformly) bounded away from zero and from above and that  $\|\hat{\Sigma}_{n'} - \Sigma_{n'}\|_2 = O_{\mathbb{P}}(a_{n'})$  with  $d'^{3/4} a_{n'} \rightarrow 0$  as well as*

$$\sup_{C \in \mathcal{C}_{n'}} \left| \mathbb{P} \left( n'^{-1/2} \sum_{i=1}^{n'} \Sigma_{n'}^{-1/2} \mathbf{Y}_{i,n'} \in C \right) - \mathbb{P}(\mathbf{Z}_{d'} + \boldsymbol{\theta}_{n'} \in C) \right| \rightarrow 0. \quad (\text{C.6})$$

Then, if  $\frac{1}{d'} \sum_{i=1}^{d'} g_2(\theta_{i,n'}) = \frac{1}{d'} \|\boldsymbol{\theta}_{n'}\|_2^2$  is bounded, it holds that

$$\sup_{C \in \mathcal{C}_{n'}} \left| \mathbb{P} \left( n'^{-1/2} \sum_{i=1}^{n'} \hat{\Sigma}_{n'}^{-1/2} \mathbf{Y}_{i,n'} \in C \right) - \mathbb{P}(\mathbf{Z}_{d'} + \boldsymbol{\theta}_{n'} \in C) \right| \rightarrow 0. \quad (\text{C.7})$$

*Proof.* The quantity inside the supremum in (C.7) vanishes for  $C = \emptyset$ . Thus, consider a sequence  $C_n$  with  $C_n \in \mathcal{C}_n$  and  $C_n \neq \emptyset$  for every  $n$ . Write  $n$  instead of  $n'$  and let

$$A_n := \left\| n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} - n^{-1/2} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \right\|_2$$

Since  $d^{-1} \sum_{i=1}^d g_2(\theta_{i,n})$  is bounded, Part 1 of Lemma C.2 implies that for every  $\delta \in (0, 1)$  there exists an  $M = M_\delta > 0$  such that for  $\varepsilon_n = M d^{1/2} a_n$  one has  $\mathbb{P}(A_n > \varepsilon_n) \leq \delta$  for all  $n$ . Therefore, we have

$$\left\{ n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \in C_n \right\} \subseteq \left\{ n^{-1/2} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \in C_n^{\varepsilon_n} \right\} \cup \{A_n > \varepsilon_n\}.$$



Denoting the supremum in (C.6) by  $r_n$  it thus follows from convexity of  $C_n^{\varepsilon_n}$  that

$$\mathbb{P}\left(n^{-1/2}\sum_{i=1}^n\hat{\Sigma}_n^{-1/2}\mathbf{Y}_{i,n}\in C_n\right)\leq\mathbb{P}\left(\mathbf{Z}_d+\boldsymbol{\theta}_n\in C_n^{\varepsilon_n}\right)+r_n+\delta.$$

Next, by  $C_n^{\varepsilon_n}-\boldsymbol{\theta}_n=(C_n-\boldsymbol{\theta}_n)^{\varepsilon_n}$ , convexity of  $C_n-\boldsymbol{\theta}_n$ , and Lemma 2.6 in Bentkus (2003) (cf. also Nazarov (2003)), it eventually holds that

$$\begin{aligned}\mathbb{P}\left(\mathbf{Z}_d+\boldsymbol{\theta}_n\in C_n^{\varepsilon_n}\right)&=\mathbb{P}\left(\mathbf{Z}_d\in C_n^{\varepsilon_n}-\boldsymbol{\theta}_n\right) \\ &=\mathbb{P}\left(\mathbf{Z}_d\in(C_n-\boldsymbol{\theta}_n)^{\varepsilon_n}\right) \\ &\leq\mathbb{P}\left(\mathbf{Z}_d\in C_n-\boldsymbol{\theta}_n\right)+d^{1/4}\varepsilon_n \\ &=\mathbb{P}\left(\mathbf{Z}_d+\boldsymbol{\theta}_n\in C_n\right)+d^{1/4}\varepsilon_n.\end{aligned}$$

Therefore,

$$\limsup_{n\rightarrow\infty}\left[\mathbb{P}\left(n^{-1/2}\sum_{i=1}^n\hat{\Sigma}_n^{-1/2}\mathbf{Y}_{i,n}\in C_n\right)-\mathbb{P}\left(\mathbf{Z}_d+\boldsymbol{\theta}_n\in C_n\right)\right]\leq\delta.$$

On the other hand, it also holds that

$$\left\{n^{-1/2}\sum_{i=1}^n\Sigma_n^{-1/2}\mathbf{Y}_{i,n}\in C_n^{-\varepsilon_n}\right\}\subseteq\left\{n^{-1/2}\sum_{i=1}^n\hat{\Sigma}_n^{-1/2}\mathbf{Y}_{i,n}\in C_n\right\}\cup\{A_n>\varepsilon_n\}.$$

Hence, by convexity of  $C_n^{-\varepsilon_n}$  and recalling that  $r_n$  denotes the supremum in (C.6) (with  $r_n=0$  for  $C_n^{-\varepsilon_n}=\emptyset$ )

$$\mathbb{P}\left(\mathbf{Z}_d+\boldsymbol{\theta}_n\in C_n^{-\varepsilon_n}\right)-r_n\leq\mathbb{P}\left(n^{-1/2}\sum_{i=1}^n\hat{\Sigma}_n^{-1/2}\mathbf{Y}_{i,n}\in C_n\right)+\delta$$

Next, by  $C_n^{-\varepsilon_n}-\boldsymbol{\theta}_n=(C_n-\boldsymbol{\theta}_n)^{-\varepsilon_n}$  (which remains true for  $C_n^{-\varepsilon_n}=\emptyset$ ), convexity of  $C_n-\boldsymbol{\theta}_n$

and Lemma 2.6 [Bentkus \(2003\)](#) it eventually holds that

$$\begin{aligned}
\mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in C_n^{-\varepsilon_n}) &= \mathbb{P}(\mathbf{Z}_d \in C_n^{-\varepsilon_n} - \boldsymbol{\theta}_n) \\
&= \mathbb{P}(\mathbf{Z}_d \in (C_n - \boldsymbol{\theta}_n)^{-\varepsilon_n}) \\
&\geq \mathbb{P}(\mathbf{Z}_d \in C_n - \boldsymbol{\theta}_n) - d^{1/4}\varepsilon_n \\
&= \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in C_n) - d^{1/4}\varepsilon_n.
\end{aligned}$$

Therefore,

$$\liminf_{n \rightarrow \infty} \left[ \mathbb{P} \left( n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \in C_n \right) - \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in C_n) \right] \geq -\delta.$$

Thus, [\(C.7\)](#) follows because the sequence  $C_n \neq \emptyset$  and  $\delta \in (0, 1)$  were arbitrary.  $\square$

## D Proof of Theorem [3.2](#)

It will be repeatedly used that by [\(A.1\)](#), for every  $p \in [2, \infty)$ , there exists a  $c_p \in (0, \infty)$  such that

$$c_p^{-1} \sum_{i=1}^d g_p(\theta_{i,n}) \leq \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)] \leq c_p \sum_{i=1}^d g_p(\theta_{i,n}). \tag{D.1}$$

Fix  $p \in [2, \infty)$ .

Assumption [3.1](#), in case we additionally have  $\boldsymbol{\beta}^* \in \mathbf{B}^{(0)}$ , and, more generally, [\(8\)](#) of Assumption [3.2](#) are verified by combining Lemma [C.3](#) and

$$\frac{\|\mathbf{Z}_{d'} + \boldsymbol{\theta}_{n'}\|_p^p - \sum_{i=1}^{d'} \lambda_p(\theta_{i,n'})}{\sqrt{d' \text{Var} |Z|^p}} \rightsquigarrow \mathbf{N}_1(0, 1),$$

for every subsequence  $n'$  of  $n$  along which  $(1/\sqrt{d}) \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]$  (and hence also  $d^{-1} \sum_{i=1}^d g_2(\theta_{i,n})$ ) is bounded — the previous display following from Lemma C.3 in [Kock and Preinerstorfer \(2023\)](#).

To verify [\(9\)](#) of Assumption [3.2](#), let  $d^{-1/2} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)] \rightarrow \infty$ , so that by [\(C.23\)](#) in [Kock and Preinerstorfer \(2023\)](#) together with [\(D.1\)](#)

$$\frac{\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} = o_{\mathbb{P}}(1). \tag{D.2}$$

We show that for any subsequence  $n'$  of  $n$  there exists a subsequence  $n''$  of  $n'$  along which (9) holds: Pick a subsequence  $n'$  of  $n$ .

**Case 1:** Suppose  $n'$  possesses a subsequence  $n''$ , say, along which  $d^{-1} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]$  (and hence also  $d^{-1} \sum_{i=1}^d g_2(\theta_{i,n})$ ) is bounded, then (9) follows along  $n''$  by combining Lemma C.3 with (D.2).

**Case 2:** Assume  $d'^{-1} \sum_{i=1}^{d'} [\lambda_p(\theta_{i,n'}) - \lambda_p(0)] \rightarrow \infty$ . In this case, set  $n'' = n'$  and write  $n$  instead of  $n'$ . We first observe that for any  $\delta \in (0, 1)$ ,

$$\left\{ \frac{\left\| n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} \leq \delta \right\} = \left\{ n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \in \mathcal{B}_p(r_{n,\delta}) \right\},$$

where  $r_{n,\delta} = \left( \delta \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)] + \sum_{i=1}^d \lambda_p(\theta_{i,n}) \right)^{1/p}$ . Letting

$$B_n = \left\| n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} - n^{-1/2} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p,$$

it follows from Lemma C.2 that for every  $\rho \in (0, 1)$  there exists an  $M = M_\rho$  such that for  $\varepsilon_n = M d^{\frac{1}{2} - \frac{1}{p}} a_n \left[ \sum_{i=1}^d g_p(\theta_{i,n}) \right]^{\frac{1}{p}}$  one has  $\mathbb{P}(B_n > \varepsilon_n) \leq \rho$ . Furthermore, note that since  $r_{n,\delta}^p \geq \delta c_p^{-1} \sum_{i=1}^d g_p(\theta_{i,n})$  it holds that

$$\frac{\varepsilon_n}{r_{n,\delta}} = O(d^{\frac{1}{2} - \frac{1}{p}} a_n) = o(1), \quad (\text{D.3})$$

such that, in particular,  $r_{n,\delta} - \varepsilon_n$  is eventually strictly positive. Next, observe that (eventually)

$$\left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \in \mathcal{B}_p(r_{n,\delta} - \varepsilon_n) \right\} \subseteq \left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \in \mathcal{B}_p(r_{n,\delta}) \right\} \cup \{B_n > \varepsilon_n\},$$

and note that by assumption

$$\liminf_{n \rightarrow \infty} \mathbb{P} \left( \frac{1}{\sqrt{n}} \sum_{i=1}^n \Sigma_n^{-1/2} \mathbf{Y}_{i,n} \in \mathcal{B}_p(r_{n,\delta} - \varepsilon_n) \right) = \liminf_{n \rightarrow \infty} \mathbb{P} (\mathbf{Z}_d + \boldsymbol{\theta}_n \in \mathcal{B}_p(r_{n,\delta} - \varepsilon_n)).$$

Furthermore, by Bernoulli's inequality (eventually)

$$(r_{n,\delta} - \varepsilon_n)^p = r_{n,\delta}^p (1 - \varepsilon_n/r_{n,\delta})^p \geq r_{n,\delta}^p (1 - p\varepsilon_n/r_{n,\delta})$$

so that the right-hand side of the penultimate display is no smaller than

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \mathbb{P} (\mathbf{Z}_d + \boldsymbol{\theta}_n \in \mathcal{B}_p(r_{n,\delta}(1 - p\varepsilon_n/r_{n,\delta})^{1/p})) \\ &= \liminf_{n \rightarrow \infty} \mathbb{P} \left( \frac{\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} < \delta - \frac{pr_{n,\delta}^p \varepsilon_n}{r_{n,\delta} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} \right). \end{aligned}$$

Because  $r_{n,\delta}^p / \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]$  is bounded (recall that  $d^{-1} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)] \rightarrow \infty$ ) and by (D.3)

$$\frac{pr_{n,\delta}^p \varepsilon_n}{r_{n,\delta} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} = O(\varepsilon_n/r_{n,\delta}) = o(1).$$

Hence, by (D.2) the penultimate display equals one and we conclude that

$$\liminf_{n \rightarrow \infty} \mathbb{P} \left( \frac{\left\| n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} \leq \delta \right) \geq 1 - \rho. \quad (\text{D.4})$$

Next, since  $r_{n,-\delta} = \left( -\delta \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)] + \sum_{i=1}^d \lambda_p(\theta_{i,n}) \right)^{1/p} > 0$  for  $\delta \in (0, 1)$ ,

$$\left\{ \frac{\left\| n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} \right\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} \leq -\delta \right\} = \left\{ n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} \in \mathcal{B}_p(r_{n,-\delta}) \right\},$$

which is contained in

$$\left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n \boldsymbol{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \in \mathcal{B}_p(r_{n,-\delta} + \varepsilon_n) \right\} \cup \{B_n > \varepsilon_n\}.$$

By assumption

$$\limsup_{n \rightarrow \infty} \mathbb{P} \left( \frac{1}{\sqrt{n}} \sum_{i=1}^n \boldsymbol{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \in \mathcal{B}_p(r_{n,-\delta} + \varepsilon_n) \right) = \limsup_{n \rightarrow \infty} \mathbb{P} (\mathbf{Z}_d + \boldsymbol{\theta}_n \in \mathcal{B}_p(r_{n,-\delta} + \varepsilon_n)). \quad (\text{D.5})$$

Furthermore, since  $r_{n,-\delta}^p \geq (1-\delta)c_p^{-1} \sum_{i=1}^d g_p(\theta_{i,n})$  it holds that

$$\frac{\varepsilon_n}{r_{n,-\delta}} = O(d^{\frac{1}{2}-\frac{1}{p}} a_n) = o(1). \quad (\text{D.6})$$

Thus, by the mean-value theorem, it eventually holds that

$$(r_{n,-\delta} + \varepsilon_n)^p = r_{n,-\delta}^p (1 + \varepsilon_n/r_{n,-\delta})^p \leq r_{n,-\delta}^p (1 + 2p\varepsilon_n/r_{n,-\delta}),$$

such that the right-hand side in (D.5) is no greater than

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in \mathcal{B}_p(r_{n,-\delta}(1 + 2p\varepsilon_n/r_{n,-\delta})^{1/p})) \\ &= \limsup_{n \rightarrow \infty} \mathbb{P}\left(\frac{\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} \leq -\delta + \frac{2pr_{n,-\delta}^p \varepsilon_n}{r_{n,-\delta} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]}\right). \end{aligned}$$

Because  $r_{n,-\delta}^p / \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]$  is bounded (recall that  $d^{-1} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)] \rightarrow \infty$ ) and by (D.6)

$$\frac{2pr_{n,-\delta}^p \varepsilon_n}{r_{n,-\delta} \sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} = O(\varepsilon_n/r_{n,-\delta}) = o(1).$$

Hence, the expression in the penultimate display equals zero by (D.2) and we conclude that

$$\limsup_{n \rightarrow \infty} \mathbb{P}\left(\frac{\|n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n}\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]} \leq -\delta\right) \leq \rho. \quad (\text{D.7})$$

The statement in (9) of Assumption 3.2 now follows, because Equations (D.4) and (D.7) imply that for every  $\delta \in (0, 1)$  and every  $\rho \in (0, 1)$  it holds that

$$\limsup_{n \rightarrow \infty} \mathbb{P}\left(\left|\frac{\|n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n}\|_p^p - \sum_{i=1}^d \lambda_p(\theta_{i,n})}{\sum_{i=1}^d [\lambda_p(\theta_{i,n}) - \lambda_p(0)]}\right| > \delta\right) \leq 2\rho.$$

## E Proof of Corollary 3.3

Concerning Condition 1 in Assumption 3.3, consider first the case where  $\hat{\boldsymbol{\Sigma}}_n(\boldsymbol{\beta}_n^*)$  is the estimator  $\hat{\boldsymbol{\Sigma}}_{\eta,\delta}$  from Theorem 1 of Abdalla and Zhivotovskiy (2022) for  $\eta = 0$ ,  $\delta = 1/d$ ,

and  $p = 4$  applied to the i.i.d. sample

$$(\mathbf{Y}_{2,n} - \mathbf{Y}_{1,n})/\sqrt{2}, \dots, (\mathbf{Y}_{2_{\lfloor n/2 \rfloor}, n} - \mathbf{Y}_{2_{\lfloor n/2 \rfloor - 1}, n})/\sqrt{2}$$

of size  $\lfloor n/2 \rfloor$  with covariance matrix  $\boldsymbol{\Sigma}_n$  and mean vector zero; cf. also Section 1.2 in [Mendelson and Zivotovskiy \(2020\)](#). We apply Theorem 1 of [Abdalla and Zivotovskiy \(2022\)](#) to show that  $\|\hat{\boldsymbol{\Sigma}}_{0,d-1} - \boldsymbol{\Sigma}_n\|_2 = O_{\mathbb{P}}(\sqrt{d/n})$ , which then verifies Condition 1 of Assumption [3.3](#) with  $a_n = \sqrt{d/n}$ , because  $d/n^{2/5} \rightarrow 0$  by assumption. To this end, we only need to show that the condition in Equation (1) of [Abdalla and Zivotovskiy \(2022\)](#) holds for  $p = 4$  (note that we only have to verify that condition for  $p = q = 4$  due to Lyapunov's inequality): By the Binomial theorem and Assumption [3.4](#), for every  $t \in \mathbb{R}^d$  and writing  $L = L(\boldsymbol{\beta}^*)$ ,

$$\mathbb{E}\langle \mathbf{Y}_{2,n} - \mathbf{Y}_{1,n}, t \rangle^4 = \mathbb{E}\langle (\mathbf{Y}_{2,n} - \boldsymbol{\mu}_n) - (\mathbf{Y}_{1,n} - \boldsymbol{\mu}_n), t \rangle^4 \leq (2L^4 + 6) (\mathbb{E}\langle \mathbf{Y}_{1,n} - \boldsymbol{\mu}_n, t \rangle^2)^2.$$

In addition,

$$\mathbb{E}\langle \mathbf{Y}_{1,n} - \boldsymbol{\mu}_n, t \rangle^2 = t' \boldsymbol{\Sigma}_n t = \frac{1}{2} t' \mathbb{E}(\mathbf{Y}_{2,n} - \mathbf{Y}_{1,n})(\mathbf{Y}_{2,n} - \mathbf{Y}_{1,n})' t = \mathbb{E}\langle (\mathbf{Y}_{2,n} - \mathbf{Y}_{1,n})/\sqrt{2}, t \rangle^2,$$

such that  $(\mathbf{Y}_{2,n} - \mathbf{Y}_{1,n})/\sqrt{2}$  satisfies the condition in their Equation (1) with  $\kappa(4) = [(2L^4 + 6)/4]^{1/4}$ . This also shows that Condition 1 of Assumption [3.3](#) holds if  $\hat{\boldsymbol{\Sigma}}_n(\boldsymbol{\beta}_n^*)$  is the estimator in Theorem 1.3 of [Oliveira and Rico \(2022\)](#) (with their  $\eta = 0$ ,  $\alpha = 1/d$  and  $p = 4$  and applied to the auxiliary sample as above).

Concerning Condition 2 of Assumption [3.3](#), we use Theorem 2.1 in [Fang and Koike \(2024\)](#) to establish that

$$\sup_{C \in \mathcal{C}_n} \left| \mathbb{P} \left( n^{-1/2} \sum_{i=1}^n \boldsymbol{\Sigma}_n^{-1/2} [\mathbf{Y}_{i,n} - \boldsymbol{\mu}_n] \in C \right) - \mathbb{P}(\mathbf{Z}_d \in C) \right| \rightarrow 0;$$

that is their  $\xi_i = n^{-1/2} \boldsymbol{\Sigma}_n^{-1/2} [\mathbf{Y}_{i,n} - \boldsymbol{\mu}_n]$  and their  $\boldsymbol{\Sigma} = \mathbf{I}_d$ . Thus, the Cauchy-Schwarz inequality yields that their

$$\mathbb{E}\|\boldsymbol{\Sigma}^{-1/2} \xi_1\|_2^4 = \mathbb{E}\|\xi_1\|_2^4 = \mathbb{E} \left( \sum_{j=1}^d \xi_{1,j}^2 \right)^2 \leq d \sum_{j=1}^d \mathbb{E} \xi_{1,j}^4.$$

In addition, letting  $e_j$  be the  $j$ th canonical basis vector of  $\mathbb{R}^d$ , it follows from Assumption 3.4 that for  $j = 1, \dots, d$  and  $L = L(\boldsymbol{\beta}^*)$  as above,

$$\mathbb{E}\xi_{1,j}^4 = n^{-2}\mathbb{E}\langle(\mathbf{Y}_{i,n} - \boldsymbol{\mu}_n), \boldsymbol{\Sigma}_n^{-1/2}e_j\rangle^4 \leq L^4n^{-2}\left(\mathbb{E}\langle\boldsymbol{\Sigma}_n^{-1/2}(\mathbf{Y}_{i,n} - \boldsymbol{\mu}_n), e_j\rangle^2\right)^2 = L^4n^{-2}.$$

Therefore,  $\sum_{i=1}^n \mathbb{E}\|\boldsymbol{\Sigma}_n^{-1/2}\xi_i\|_2^4 \leq \frac{L^4d^2}{n}$  and Theorem 2.1 in Fang and Koike (2024) together with  $\frac{d}{n^{2/5}}[\log(n)]^{2/5} \rightarrow 0$  yields that Condition 2 of Assumption 3.3 is satisfied, since

$$\sup_{C \in \mathcal{C}_n} \left| \mathbb{P}\left(n^{-1/2} \sum_{i=1}^n \boldsymbol{\Sigma}_n^{-1/2}[\mathbf{Y}_{i,n} - \boldsymbol{\mu}_n] \in C\right) - \mathbb{P}(\mathbf{Z}_d \in C)\right| = O\left(\frac{d^{5/4}}{n^{1/2}}\sqrt{\log(n)}\right) = o(1).$$

## F Gaussian approximation over hyperrectangles with estimated covariance matrix and $d \rightarrow \infty$

We begin by formulating a version of Lemma C.3 for hyperrectangles  $\mathcal{H}_n$ , cf. Section 3.4. Although a Gaussian approximation result over  $\mathcal{H}_n$  follows directly from Lemma C.3 and  $\mathcal{H}_n \subseteq \mathcal{C}_n$ , an approximation result applying to  $\boldsymbol{\theta}_n$  further from zero, necessary for our purposes, can be obtained for  $\mathcal{H}_n$ . The proof strategy is identical to that of Lemma C.3, but as other anti-concentration inequalities are employed we provide the proof for completeness. For any  $A \subseteq \mathbb{R}^d$  and  $\varepsilon > 0$ , let

$$A^{\varepsilon, \infty} = \{\mathbf{x} \in \mathbb{R}^d : \inf_{\mathbf{y} \in A} \|\mathbf{x} - \mathbf{y}\|_\infty \leq \varepsilon\}.$$

Furthermore,

$$A^{-\varepsilon, \infty} = \{\mathbf{x} \in \mathbb{R}^d : \mathcal{B}_\infty(\mathbf{x}, \varepsilon) \subseteq A\} \quad \text{and} \quad \mathcal{B}_\infty(\mathbf{x}, \varepsilon) = \{\mathbf{y} \in \mathbb{R}^d : \|\mathbf{y} - \mathbf{x}\|_\infty \leq \varepsilon\}.$$

**Lemma F.1.** *Let  $n'$  be a subsequence of  $n$  and  $d' \rightarrow \infty$ . Assume that the eigenvalues of  $\boldsymbol{\Sigma}_{n'}$  are (uniformly) bounded away from zero and from above and that  $\|\hat{\boldsymbol{\Sigma}}_{n'} - \boldsymbol{\Sigma}_{n'}\|_2 = O_{\mathbb{P}}(a_{n'})$  with  $d'^{3/4}a_{n'} \rightarrow 0$  as well as*

$$\sup_{C \in \mathcal{C}_{n'}} \left| \mathbb{P}\left(n'^{-1/2} \sum_{i=1}^{n'} \boldsymbol{\Sigma}_{n'}^{-1/2} \mathbf{Y}_{i,n'} \in C\right) - \mathbb{P}(\mathbf{Z}_{d'} + \boldsymbol{\theta}_{n'} \in C)\right| \rightarrow 0. \quad (\text{F.1})$$

Then, if  $\frac{1}{d^{3/2}/\log(d^d)} \sum_{i=1}^{d'} g_2(\theta_{i,n'}) = \frac{1}{d^{3/2}/\log(d^d)} \|\boldsymbol{\theta}_{n'}\|_2^2$  is bounded, it also holds that

$$\sup_{H \in \mathcal{H}_{n'}} \left| \mathbb{P} \left( n'^{-1/2} \sum_{i=1}^{n'} \hat{\boldsymbol{\Sigma}}_{n'}^{-1/2} \mathbf{Y}_{i,n'} \in H \right) - \mathbb{P}(\mathbf{Z}_{d'} + \boldsymbol{\theta}_{n'} \in H) \right| \rightarrow 0. \quad (\text{F.2})$$

*Proof.* Write  $n$  instead of  $n'$  and let

$$\begin{aligned} A_n &:= \left\| n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} - n^{-1/2} \sum_{i=1}^n \boldsymbol{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \right\|_{\infty} \\ &\leq \left\| n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} - n^{-1/2} \sum_{i=1}^n \boldsymbol{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \right\|_2. \end{aligned}$$

By Lemma C.2 (with  $p = 2$ ) it follows that  $A_n = O_{\mathbb{P}}([d^{1/2} \vee \|\boldsymbol{\theta}_n\|_2] a_n)$ , where we used that  $\sum_{i=1}^d g_2(\theta_{i,n}) = \|\boldsymbol{\theta}_n\|_2^2$ . Thus, for every  $\delta \in (0, 1)$  there exists an  $M = M_{\delta} > 0$  such that for  $\varepsilon_n = M[d^{1/2} \vee \|\boldsymbol{\theta}_n\|_2] a_n$  one has  $\mathbb{P}(A_n > \varepsilon_n) \leq \delta$  for all  $n \in \mathbb{N}$ . Therefore, for any sequence  $H_n$  with  $H_n \in \mathcal{H}_n$  for every  $n$  we have

$$\left\{ n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} \in H_n \right\} \subseteq \left\{ n^{-1/2} \sum_{i=1}^n \boldsymbol{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \in H_n^{\varepsilon_n, \infty} \right\} \cup \{A_n > \varepsilon_n\}.$$

Since  $H_n \in \mathcal{H}_n$ , one has  $H_n = \prod_{j=1}^d [a_j, b_j]$  for  $-\infty \leq a_j \leq b_j \leq \infty$ ,  $j = 1, \dots, d$ , and so  $H_n^{\varepsilon_n, \infty} = \prod_{j=1}^d [a_j - \varepsilon_n, b_j + \varepsilon_n] \in \mathcal{H}_n$ . Thus, denoting the supremum in (F.1) by  $r_n$ ,

$$\mathbb{P} \left( n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} \in H_n \right) \leq \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n^{\varepsilon_n, \infty}) + r_n + \delta.$$

Next, by  $H_n^{\varepsilon_n, \infty} - \boldsymbol{\theta}_n = \prod_{j=1}^d [a_j - \theta_{j,n} - \varepsilon_n, b_j - \theta_{j,n} + \varepsilon_n]$  it follows that

$$\begin{aligned} &\mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n^{\varepsilon_n, \infty}) \\ &= \mathbb{P} \left( \mathbf{Z}_d \in \prod_{j=1}^d [a_j - \theta_{j,n} - \varepsilon_n, b_j - \theta_{j,n} + \varepsilon_n] \right) \\ &= \mathbb{P} \left( (\mathbf{Z}'_d, -\mathbf{Z}'_d)' \in \prod_{j=1}^d (-\infty, b_j - \theta_{j,n} + \varepsilon_n] \times \prod_{j=1}^d (-\infty, -a_j + \theta_{j,n} + \varepsilon_n] \right). \end{aligned}$$



Lemma A.1 in Chernozhukov et al. (2017a) (which is attributed to Nazarov (2003), cf. also Chernozhukov et al. (2017b)) now yields that the far right-hand side of the previous display can (eventually) be bounded from above by

$$\begin{aligned} & \mathbb{P} \left( (\mathbf{Z}'_d, -\mathbf{Z}'_d)' \in \prod_{j=1}^d (-\infty, b_j - \theta_{j,n}] \times \prod_{j=1}^d (-\infty, -a_j + \theta_{j,n}] \right) + C \sqrt{\log(d)} \varepsilon_n \\ &= \mathbb{P} (\mathbf{Z}_d \in H_n - \boldsymbol{\theta}_n) + C \sqrt{\log(d)} \varepsilon_n \\ &= \mathbb{P} (\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n) + C \sqrt{\log(d)} \varepsilon_n. \end{aligned}$$

Therefore,

$$\limsup_{n \rightarrow \infty} \left[ \mathbb{P} \left( n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} \in H_n \right) - \mathbb{P} (\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n) \right] \leq \delta.$$

On the other hand, it also holds that

$$\left\{ n^{-1/2} \sum_{i=1}^n \boldsymbol{\Sigma}_n^{-1/2} \mathbf{Y}_{i,n} \in H_n^{-\varepsilon_n, \infty} \right\} \subseteq \left\{ n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} \in H_n \right\} \cup \{A_n > \varepsilon_n\}.$$

Since  $H_n^{-\varepsilon_n, \infty} = \prod_{j=1}^d [a_j + \varepsilon_n, b_j - \varepsilon_n] \in \mathcal{H}_n$ , similar arguments to the above, noting also that

$$|\mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n) - \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n^{-\varepsilon_n, \infty})| \leq \Phi(\varepsilon_n) - \Phi(-\varepsilon_n) \quad \text{if } H_n^{-\varepsilon_n, \infty} = \emptyset,$$

imply

$$\liminf_{n \rightarrow \infty} \left[ \mathbb{P} \left( n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n} \in H_n \right) - \mathbb{P} (\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n) \right] \geq -\delta.$$

Thus, (F.2) follows since the sequence  $H_n$  and  $\delta \in (0, 1)$  were arbitrary.  $\square$

*Remark F.1.* Under the assumptions of Lemma F.1 the supremum in (F.2) can be extended to also include all open rectangles of the form  $H_n^o = \prod_{j=1}^d (a_j, b_j)$  (where without loss of generality  $-\infty < a_j < b_j < \infty$ ) by the following squeezing argument: write  $H_n = \prod_{j=1}^d [a_j, b_j]$  and  $\mathbf{S}_n = n^{-1/2} \sum_{i=1}^n \hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{Y}_{i,n}$ . Using that  $\mathbf{Z}_d$  has no mass at  $H_n \setminus H_n^o$ , one

has (along any subsequence of  $n$ )

$$\mathbb{P}(\mathbf{S}_n \in H_n^o) - \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n^o) \leq \mathbb{P}(\mathbf{S}_n \in H_n) - \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n);$$

by Lemma F.1 the right-hand side tends to zero. Similarly, setting  $c_n = [\log(d)]^{-1} \wedge \min_{j=1, \dots, d} (b_j - a_j)/4$

$$\mathbb{P}(\mathbf{S}_n \in H_n^o) - \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n^o) \geq \mathbb{P}\left(\mathbf{S}_n \in \prod_{j=1}^d [a_j + c_n, b_j - c_n]\right) - \mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n).$$

Using Lemma A.1 in Chernozhukov et al. (2017a) as in the proof of Lemma F.1 it follows that there exists a constant  $C \in (0, \infty)$  such that

$$\mathbb{P}(\mathbf{Z}_d + \boldsymbol{\theta}_n \in H_n) \leq \mathbb{P}\left(\mathbf{Z}_d + \boldsymbol{\theta}_n \in \prod_{j=1}^d [a_j + c_n, b_j - c_n]\right) + C\sqrt{\log(d)}c_n,$$

such that Lemma F.1 can again be applied yielding the claimed uniform convergence as  $H_n^o$  was an arbitrary open rectangle. As any rectangle can be squeezed between its counterpart with all endpoints excluded and included, respectively, Lemma F.1 also applies to arbitrary rectangles.

## G Proof of Theorem 3.4

The size control in part a) follows by combining Lemma F.1 (using “ $n' = n$ ” and  $\boldsymbol{\theta}_n = \mathbf{0}_d$ ) and Lemma A.2 in Kock and Preinerstorfer (2023), cf. also Remark F.1 above (which is invoked tacitly in the remainder of the proof).

Regarding part b), suppose that  $b_n := \sum_{i=1}^d \frac{\bar{\Phi}(c_d - |\theta_{i,d}|)}{\Phi(c_d - |\theta_{i,d}|)} \rightarrow \infty$ . By Theorem 3.5 in Kock and Preinerstorfer (2023) one has  $\mathbb{P}(\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_\infty \geq \kappa_{n,\infty}) \rightarrow 1$ . Let  $n'$  be a subsequence of  $n$ . We show that it possesses a further subsequence along which  $\mathbb{P}(S_{n',\infty} \geq \kappa_{n',\infty}) \rightarrow 1$ . If  $\frac{1}{d^{3/2}/\log(d)} \|\boldsymbol{\theta}_n\|_2^2$  is bounded along  $n'$ , Lemma F.1 implies  $\mathbb{P}(S_{n',\infty} \geq \kappa_{n',\infty}) \rightarrow 1$ .

If, on the other hand,  $\frac{1}{d^{3/2}/\log(d)} \|\boldsymbol{\theta}_n\|_2^2$  is unbounded there exists a further subsequence  $n''$  along which  $\frac{1}{d^{3/2}/\log(d)} \|\boldsymbol{\theta}_n\|_2^2 \rightarrow \infty$ . Furthermore, note that  $\mathbb{P}(S_{n,\infty} \geq \kappa_{n,\infty})$

equals

$$\mathbb{P} \left( \left\| n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} (\mathbf{Y}_{i,n} - \boldsymbol{\mu}_n) + n^{1/2} \hat{\Sigma}_n^{-1/2} \boldsymbol{\mu}_n \right\|_{\infty} \geq \kappa_{n,\infty} \right). \quad (\text{G.1})$$

Because  $\mathbf{Y}_{i,n} - \boldsymbol{\mu}_n$  is centered and has the same covariance matrix as  $\mathbf{Y}_{i,n}$ , we can apply Lemma F.1 (with  $\boldsymbol{\theta}_n = \mathbf{0}_d$  there) in conjunction with part a) to conclude that (along the entire sequence  $n$ )

$$\left\| n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} (\mathbf{Y}_{i,n} - \boldsymbol{\mu}_n) \right\|_{\infty} = O_{\mathbb{P}}(\sqrt{\log(d)}). \quad (\text{G.2})$$

Next,

$$\|n^{1/2} \hat{\Sigma}_n^{-1/2} \boldsymbol{\mu}_n\|_{\infty} \geq \frac{\|n^{1/2} \hat{\Sigma}_n^{-1/2} \boldsymbol{\mu}_n\|_2}{\sqrt{d}} \geq \frac{\|n^{1/2} \boldsymbol{\Sigma}_n^{-1/2} \boldsymbol{\mu}_n\|_2 - \|n^{1/2} [\hat{\Sigma}_n^{-1/2} - \boldsymbol{\Sigma}_n^{-1/2}] \boldsymbol{\mu}_n\|_2}{\sqrt{d}}$$

and the numerator on the far right-hand side equals  $\|n^{1/2} \boldsymbol{\Sigma}_n^{-1/2} \boldsymbol{\mu}_n\|_2 = \|\boldsymbol{\theta}_n\|_2$  (which is eventually positive along  $n''$ ) times

$$1 - \frac{\|n^{1/2} [\hat{\Sigma}_n^{-1/2} - \boldsymbol{\Sigma}_n^{-1/2}] \boldsymbol{\mu}_n\|_2}{\|n^{1/2} \boldsymbol{\Sigma}_n^{-1/2} \boldsymbol{\mu}_n\|_2} \geq 1 - \frac{\|\hat{\Sigma}_n^{-1/2} - \boldsymbol{\Sigma}_n^{-1/2}\|_2 \|\boldsymbol{\mu}_n\|_2}{c \|\boldsymbol{\mu}_n\|_2} = 1 - O_{\mathbb{P}}(a_n),$$

for some real  $c > 0$ , due to the uniform bound on the eigenvalues of  $\boldsymbol{\Sigma}_n$ , the last equality following from Lemma C.1. The previous two displays thus yield that along  $n''$  there exists an  $M \in (0, \infty)$  such that

$$\|n^{1/2} \hat{\Sigma}_n^{-1/2} \boldsymbol{\mu}_n\|_{\infty} \geq \frac{\|\boldsymbol{\theta}_n\|_2}{\sqrt{d}} (1 - O_{\mathbb{P}}(a_n)) \geq M \frac{d^{1/4}}{\sqrt{\log(d)}} (1 - O_{\mathbb{P}}(a_n)) \quad (\text{G.3})$$

Finally, (G.1) and hence  $\mathbb{P}(S_{n,\infty}(\boldsymbol{\beta}_n^*) \geq \kappa_{n,\infty})$  is no smaller than

$$\mathbb{P} \left( \left\| n^{1/2} \hat{\Sigma}_n^{-1/2} \boldsymbol{\mu}_n \right\|_{\infty} - \left\| n^{-1/2} \sum_{i=1}^n \hat{\Sigma}_n^{-1/2} (\mathbf{Y}_{i,n} - \boldsymbol{\mu}_n) \right\|_{\infty} \geq \kappa_{n,\infty} \right) \rightarrow 1$$

along  $n''$ ; the convergence following from (G.2), (G.3),  $\kappa_{n,\infty} = O(\sqrt{\log(d)})$ , and  $a_n \rightarrow 0$ .

Suppose now that  $\mathbb{P}(S_{n,\infty}(\boldsymbol{\beta}_n^*) \geq \kappa_{n,\infty}) \rightarrow 1$  and choose a subsequence  $n'$ . We show

that  $n'$  possesses a further subsequence along which  $b_n \rightarrow \infty$ . If  $\frac{\log(d)}{d^{3/2}} \|\boldsymbol{\theta}_n\|_2^2$  is bounded along  $n'$ , then Lemma F.1 and an application of Theorem 3.5 in Kock and Preinerstorfer (2023) along  $n'$  imply that  $b_{n'} \rightarrow \infty$ . If, on the other hand,  $\frac{\log(d)}{d^{3/2}} \|\boldsymbol{\theta}_n\|_2^2$  is unbounded along  $n'$  then there is a further subsequence  $n''$  along which  $\frac{\log(d)}{d^{3/2}} \|\boldsymbol{\theta}_n\|_2^2 \rightarrow \infty$ . Since  $\|\boldsymbol{\theta}_n\|_\infty \geq d^{-1/2} \|\boldsymbol{\theta}_n\|_2$  and  $\mathfrak{c}_d = O(\sqrt{\log(d)})$ , this implies along  $n''$  that

$$b_n \geq \frac{\overline{\Phi}(\mathfrak{c}_d - \|\boldsymbol{\theta}_n\|_\infty)}{\Phi(\mathfrak{c}_d - \|\boldsymbol{\theta}_n\|_\infty)} \rightarrow \infty.$$

## H Proof of Theorem 4.1

For  $p \in [2, \infty]$  and  $r \in (0, \infty)$  define  $\mathbb{B}_p(r) = \{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\|_p < r\}$ . Concerning the asymptotic size control in Part 1, note that by convexity of  $\bigcap_{p \in \mathfrak{P}_n} \mathbb{B}_p(c_n \kappa_{n,p})$ , Lemma C.3 yields

$$\mathbb{E}\psi_n = 1 - \mathbb{P}\left(\hat{\boldsymbol{\Sigma}}_n^{-1/2} \mathbf{H}_n \in \bigcap_{p \in \mathfrak{P}_n} \mathbb{B}_p(c_n \kappa_{n,p})\right) = 1 - \mathbb{P}\left(\mathbf{Z}_d \in \bigcap_{p \in \mathfrak{P}_n} \mathbb{B}_p(c_n \kappa_{n,p})\right) + r_n;$$

for a sequence  $r_n \rightarrow 0$ . Finally, by (19),

$$1 - \mathbb{P}\left(\mathbf{Z}_d \in \bigcap_{p \in \mathfrak{P}_n} \mathbb{B}_p(c_n \kappa_{n,p})\right) = \mathbb{P}\left(\max_{p \in \mathfrak{P}_n} \kappa_{n,p}^{-1} \|\mathbf{Z}_d\|_p \geq c_n\right) = \alpha.$$

Concerning Part 2, first let  $p \in [2, \infty)$  and consider a test  $\mathbf{1}\{S_{n,p} \geq \bar{\kappa}_{n,p}\}$  as in the statement of the theorem (where we drop the dependence of  $S_{n,p}$  on  $\beta_n^*$  and of  $\bar{\kappa}_{n,p}$  on  $\bar{\alpha}_p$  for notational convenience). Observe that since  $m_n \rightarrow \infty$  and  $p_n$  is strictly increasing, there exists a  $J \in \mathbb{N}$  such that eventually  $p \leq p_J \leq p_{m_n} < \infty$  and

$$\psi_n \geq \mathbf{1}\{S_{n,p_J} \geq c_n \kappa_{n,p_J}\} \geq \mathbf{1}\{S_{n,p_J} \geq \kappa_{n,p_J}\}.$$

By Lemma C.3, the test  $\mathbf{1}\{S_{n,p_J} \geq \kappa_{n,p_J}\}$  has asymptotic null rejection probability equal to  $\lim_{n \rightarrow \infty} \alpha_{n,p_J} \in (0, \alpha_2)$ . Hence, by Theorem 3.1 (see also (13)), if  $\mathbb{P}(S_{n,p} \geq \bar{\kappa}_{n,p}) \rightarrow 1$  it also follows that  $\mathbb{P}(S_{n,p_J} \geq \kappa_{n,p_J}) \rightarrow 1$ , so that  $\mathbb{E}\psi_n \rightarrow 1$  follows from the previous display.

Next, let  $p = \infty$  and consider a test  $\mathbf{1}\{S_{n,\infty} \geq \bar{\kappa}_{n,\infty}\}$  as in the statement of the theorem (again dropping some dependencies for notational convenience). Clearly,

$$\psi_n \geq \mathbf{1}\{S_{n,\infty} \geq c_n \kappa_{n,\infty}\} \geq \mathbf{1}\{S_{n,\infty} \geq \kappa_{n,\infty}\}.$$

By Lemma F.1, the test  $\mathbb{1}\{S_{n,\infty} \geq \kappa_{n,\infty}\}$  has asymptotic null rejection frequency  $\alpha_\infty \in (0, 1)$ , cf. also Remark F.1. Hence, by Theorem 3.4, if  $\mathbb{P}(S_{n,\infty} \geq \bar{\kappa}_{n,\infty}) \rightarrow 1$  then also  $\mathbb{P}(S_{n,\infty} \geq \kappa_{n,\infty}) \rightarrow 1$  and  $\mathbb{E}\psi_n \rightarrow 1$  follows from the previous display.

## I Proof of Theorem 4.2

Also in this proof, we do not signify the dependence of several quantities on  $\beta^*$ . Let  $n$  be large enough to ensure that  $p \in \mathfrak{P}_n$ . The definition of  $\psi_n$  in (20) along with  $c_n \in (0, 1]$  implies that

$$\mathbb{P}(S_{n,p} \geq \bar{\kappa}_{n,p}(\alpha)) - \mathbb{E}\psi_n \leq \mathbb{P}(S_{n,p} \geq \bar{\kappa}_{n,p}(\alpha)) - \mathbb{P}(S_{n,p} \geq \kappa_{n,p}). \quad (\text{I.1})$$

By the definition of  $\kappa_{n,p}$  in the construction of  $\psi_n$  and Lemma C.3 it holds under the null that  $\lim_{n \rightarrow \infty} \mathbb{P}(S_{n,p} \geq \kappa_{n,p}) = \alpha_p \in (0, 1)$ . For clarity, we thus write  $\kappa_{n,p}(\alpha_p)$  in what follows. Next, denote by  $\mathfrak{s}$  the limit superior of the left-hand side in (I.1) and let  $n'$  be a subsequence satisfying

$$\lim_{n' \rightarrow \infty} \left[ \mathbb{P}(S_{n',p} \geq \bar{\kappa}_{n',p}(\alpha)) - \mathbb{E}\psi_{n'} \right] = \mathfrak{s}.$$

We shall also use that the right hand side of (I.1) equals

$$\begin{aligned} & \left[ \mathbb{P}(S_{n,p} \geq \bar{\kappa}_{n,p}(\alpha)) - \mathbb{P}(\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p \geq \bar{\kappa}_{n,p}(\alpha)) \right] \\ & - \left[ \mathbb{P}(S_{n,p} \geq \kappa_{n,p}(\alpha_p)) - \mathbb{P}(\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p \geq \kappa_{n,p}(\alpha_p)) \right] \\ & + \left[ \mathbb{P}(\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p \geq \bar{\kappa}_{n,p}(\alpha)) - \mathbb{P}(\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_p \geq \kappa_{n,p}(\alpha_p)) \right]. \end{aligned} \quad (\text{I.2})$$

Consider first the case of  $p \in \mathfrak{P}_n \cap [2, \infty)$ : If necessary, pass to a further subsequence  $n''$  along which  $d^{-1/2} \sum_{i=1}^d g_p(\theta_{i,n})$  converges to some  $b \in [0, \infty]$ . If  $b = \infty$ , applying Theorems 3.1 (Part 3) and 3.2 along this subsequence<sup>9</sup> yields

$$\mathfrak{s} \leq \lim_{n' \rightarrow \infty} \left[ \mathbb{P}(S_{n',p} \geq \bar{\kappa}_{n',p}(\alpha)) - \mathbb{P}(S_{n',p} \geq \kappa_{n',p}(\alpha_p)) \right] = 1 - 1 = 0.$$

If, on the other hand,  $b \in [0, \infty)$ , then  $d^{-1} \sum_{i=1}^d g_p(\theta_{i,n}) \rightarrow 0$ . Hence, also  $d^{-1} \sum_{i=1}^d g_2(\theta_{i,n}) \rightarrow 0$  and so (I.2) and Lemma C.3 together with the argument following (C.61) in the proof of

---

<sup>9</sup>Inspection of the proof of these theorems shows that they remain valid along subsequences satisfying the conditions of the theorem.

Theorem C.10 in [Kock and Preinerstorfer \(2023\)](#) yield the desired bound on  $\mathfrak{s}$ .

Next, consider the case where  $p = \infty$ : If necessary, pass to a further subsequence  $n''$  along which  $\sum_{i=1}^d \frac{\bar{\Phi}(c_d - |\theta_{i,d}|)}{\Phi(c_d - |\theta_{i,d}|)}$  converges to some  $c \in [0, \infty]$ . If  $c = \infty$ , applying Part 2 of Theorem 3.4 along this subsequence yields

$$\mathfrak{s} \leq \lim_{n' \rightarrow \infty} \left[ \mathbb{P}(S_{n',\infty} \geq \bar{\kappa}_{n',\infty}(\alpha)) - \mathbb{P}(S_{n',\infty} \geq \kappa_{n',\infty}(\alpha_\infty)) \right] = 1 - 1 = 0.$$

If, on the other hand  $c \in [0, \infty)$ , then  $\frac{1}{d^{3/2}/\log(d')} \sum_{i=1}^{d'} g_2(\theta_{i,n'}) = \frac{1}{d^{3/2}/\log(d')} \|\boldsymbol{\theta}_{n'}\|_2^2$  is bounded (cf. the argument at the end of the proof of Theorem 3.4). Therefore, by (I.2) and Lemma F.1 it suffices to study the limiting behaviour of

$$\begin{aligned} & \mathbb{P}(\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_\infty \leq \kappa_{n,\infty}(\alpha_\infty)) - \mathbb{P}(\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_\infty \leq \bar{\kappa}_{n,\infty}(\alpha)) \\ &= \mathbb{P}(\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_\infty \leq \bar{\kappa}_{n,\infty}(\alpha) + [\kappa_{n,\infty}(\alpha_\infty) - \bar{\kappa}_{n,\infty}(\alpha)]) - \mathbb{P}(\|\mathbf{Z}_d + \boldsymbol{\theta}_n\|_\infty \leq \bar{\kappa}_{n,\infty}(\alpha)), \end{aligned}$$

along  $n'$ . By (18)

$$\kappa_{n,\infty}(\alpha_\infty) - \bar{\kappa}_{n,\infty}(\alpha) = \frac{\log(-\log(1-\alpha)/2)}{\sqrt{2\log(d)}} - \frac{\log(-\log(1-\alpha_\infty)/2)}{\sqrt{2\log(d)}} + o\left(\frac{1}{\sqrt{\log(d)}}\right).$$

Therefore, by Theorem 1 in [Chernozhukov et al. \(2017b\)](#) and arguments similar to those in the proof of Lemma F.1 the limit superior of the right-hand side of the penultimate display along  $n'$  is no greater than  $\log(-\log(1-\alpha)/2) - \log(-\log(1-\alpha_\infty)/2)$ , which is the claimed upper bound.

## J Sample split

To state and prove the asymptotic size control of the proposed sample split, let  $I \subseteq \{1, \dots, D\}$  be non-empty. For any  $D \times 1$  vector  $\mathbf{a}$  and any  $D \times D$  matrix  $\mathbf{A}$ , let  $\mathbf{a}_I$  and  $\mathbf{A}_I$ , respectively, denote the sub-vector (sub-matrix) with rows (and columns) indexed by  $I$ . In this section  $h_n = (h_{1,n}, \dots, h_{D,n})'$  denotes the total number,  $D$ , of moment restriction

functions. In addition,

$$\mathbf{H}(\boldsymbol{\beta}_n^*, N) := \frac{1}{\sqrt{|N|}} \sum_{i \in N} h_n(\mathbf{X}_{i,n}, \boldsymbol{\beta}_n^*) \quad \text{where } N \subseteq \{1, \dots, n\}, N \neq \emptyset$$

$\hat{\boldsymbol{\Sigma}}(\boldsymbol{\beta}_n^*, N, I)$  is a symmetric, positive semi-definite estimator of the covariance matrix of  $\mathbf{H}_I(\boldsymbol{\beta}_0, N)$  using observations indexed by  $N$  only.

$$S_p(\boldsymbol{\beta}_n^*, N, I) := \left\| [\hat{\boldsymbol{\Sigma}}(\boldsymbol{\beta}_n^*, N, I)]^{-1/2} \mathbf{H}_I(\boldsymbol{\beta}_n^*, N) \right\|_p, \quad p \in [2, \infty]$$

$$\psi(\boldsymbol{\beta}_n^*, N, I) := \mathbb{1} \left\{ \max_{p \in \mathfrak{P}_{|N|}} \kappa_{|N|,p}^{-1} S_p(\boldsymbol{\beta}_n^*, N, I) \geq c_{|N|} \right\},$$

with all other quantities as in the construction of  $\psi_n(\boldsymbol{\beta}_n^*)$  in Section 4, but with the  $|N|$  replacing  $n$ . Thus, the test (statistics) based on the observations in  $N_2$  and the selected moments  $S$  (depending on observations in  $N_1$  only) are  $S_p(\boldsymbol{\beta}_n^*, N_2, S)$ ,  $p \in [2, \infty]$ , and  $\psi(\boldsymbol{\beta}_n^*, N_2, S)$ . The allowed growth rate of  $d = d(n_2)$  is now dependent on size  $n_2$  of the second fold. Finally, let  $J_{n_2} \subseteq J_{n_2}^* := \{I \subseteq \{1, \dots, D\} : |I| = d\}$  be non-empty.

For concreteness, we work under the appropriate “sample split version” of the assumptions of Theorem 3.2. These essentially impose the assumptions of that theorem to be valid under the null only, but uniformly over all possible sets of selected moments (of size  $d$ ) replacing the sample size  $n$  by the sample size  $n_2$  in the second step (recall that  $d$  is now the number *selected* moment restrictions out of a total of  $D$  moment restrictions). Under independent sampling, appropriate versions of all other asymptotic size control results that we have stated can also be adjusted to incorporate a sample split. Recall that  $d = d(n_2)$ .

**Theorem J.1.** *Let  $\boldsymbol{\beta}^* \in \mathbf{B}^{(0)}$ ,  $n_2 \rightarrow \infty$ ,  $d \rightarrow \infty$  and  $\boldsymbol{\Sigma}_{n,I}(\boldsymbol{\beta}_n^*)$  be invertible for all  $I \in J_{n_2}$  with  $\max_{I \in J_{n_2}} \|\boldsymbol{\Sigma}_{n,I}(\boldsymbol{\beta}_n^*)\|_2$  as well as  $\max_{I \in J_{n_2}} \|\boldsymbol{\Sigma}_{n,I}^{-1}(\boldsymbol{\beta}_n^*)\|_2$  (uniformly) bounded from above. Furthermore,  $\mathbf{X}_{1,n}, \dots, \mathbf{X}_{n,n}$  is i.i.d. for each  $n \in \mathbb{N}$  and, as  $n_2 \rightarrow \infty$ ,*

1.  $\max_{I \in J_{n_2}} \|\hat{\boldsymbol{\Sigma}}(\boldsymbol{\beta}_n^*, N_2, I) - \boldsymbol{\Sigma}_{n,I}(\boldsymbol{\beta}_n^*)\|_2 = O_{\mathbb{P}}(a_{n_2})$  with  $d^{3/4} a_{n_2} \rightarrow 0$  and
2.  $\sup_{C \in \mathcal{C}_{n_2}} \max_{I \in J_{n_2}} |\mathbb{P}([\boldsymbol{\Sigma}(\boldsymbol{\beta}_n^*, N_2, I)]^{-1/2} \mathbf{H}_I(\boldsymbol{\beta}_n^*, N_2) \in C) - \mathbb{P}(\mathbf{Z}_d \in C)| \rightarrow 0$ .

In addition, let  $\alpha \in (0, 1)$  and  $S = S(n_1, n_2, \{\mathbf{X}_{i,n}\}_{i \in N_1}) \in J_{n_2}$  be measurable. Then:

1. For  $p \in [2, \infty]$  a sequence of real numbers  $(\kappa_{n_2,p})_{n_2 \in \mathbb{N}}$  satisfies

$$\mathbb{P}(S_p(\boldsymbol{\beta}_n^*, N_2, S) \geq \kappa_{n_2,p}) \rightarrow \alpha,$$

if and only if  $\kappa_{n_2,p}$  is as in (10) if  $p \in [2, \infty)$  or as in (18) if  $p = \infty$ .

2.  $\mathbb{E}\psi(\beta_n^*, N_2, S) \rightarrow \alpha$ .

Theorem J.1 shows that *irrespective* of how the set of moments  $S$  to be tested in the second step is selected — as long as this selection depends (measurably) on observations in  $N_1$  only — all tests studied have the desired asymptotic size if  $d$  satisfies the growth conditions of our previous theorems relative to the second step sample size  $n_2$ . Thus, one can use *any* state-of-the-art procedure to select the moments to be tested (some examples are given in Section J.1 below). We also reiterate that no conditions need to be imposed on the total number of available moment conditions,  $D$ .

*Proof of Theorem J.1.* Fix  $\alpha \in (0, 1)$  and let  $\varphi_{n_2} : \mathcal{X}_n^{n_2} \times J_{n_2} \rightarrow [0, 1]$  (measurable) be a test. Then, by  $N_1 \cap N_2 = \emptyset$  and independent sampling,

$$\begin{aligned} \mathbb{E}\varphi(\{\mathbf{X}_{i,n}\}_{i \in N_2}, S) &= \sum_{I \in J_{n_2}} \mathbb{E}[\varphi_{n_2}(\{\mathbf{X}_{i,n}\}_{i \in N_2}, I) \mathbf{1}(S = I)] \\ &= \sum_{I \in J_{n_2}} \mathbb{E}\varphi_{n_2}(\{\mathbf{X}_{i,n}\}_{i \in N_2}, I) \mathbb{P}(S = I). \end{aligned}$$

Thus, to show that  $\lim_{n \rightarrow \infty} \mathbb{E}\varphi_{n_2}(\{\mathbf{X}_{i,n}\}_{i \in N_2}, S) = \alpha$ , it suffices to establish (as  $n_2 \rightarrow \infty$ )

$$\max_{I \in J_{n_2}} |\mathbb{E}\varphi_{n_2}(\{\mathbf{X}_{i,n}\}_{i \in N_2}, I) - \alpha| \rightarrow 0. \quad (\text{J.1})$$

We now argue that (J.1) holds for  $\varphi_{n_2}$  a  $p$ -norm based tests with  $p \in [2, \infty)$ . Thus, let

$$\varphi_{n_2}(\{\mathbf{X}_{i,n}\}_{i \in N_2}, I) = \mathbf{1}(S_p(\beta_n^*, N_2, I) \geq \kappa_{n_2,p}).$$

Then (J.1) follows from Part 1 of Theorem 3.1 since the sufficient conditions of Theorem 3.2 are satisfied on the second subsample  $N_2$  uniformly in  $J_{n_2}$  with “ $\{1, \dots, n\} = N_2$ ” and sample size “ $n = n_2 = |N_2|$ ”. Also, “ $d = d(n_2)$ ”. The cases of  $p = \infty$  and  $\psi(\beta_n^*, N_2, S)$  are handled similarly via Theorems 3.4 and 4.1, respectively.  $\square$



## J.1 Choosing the set of moment restrictions in order to maximize power

- *Maximizing the test statistic in the first step:* If one intends to use  $S_p(\beta_n^*, N_2, S)$  in the second step, then one may choose

$$S \in \arg \max_{I \in J_{n_2}} S_p(\beta_n^*, N_1, I),$$

with  $d$  satisfying the growth conditions of Theorem J.1. Since choosing  $S$  in this fashion may be computationally burdensome for very large  $D$ , one can alternatively add moments one-by-one in a set-up approach (one could also use a step-down procedure) until arriving at  $d$  moments, in each step adding the moment that maximizes the value of the test statistic given the previously selected moments. A similar procedure can be used for  $\psi(\beta_n^*, N_2, S)$ . In this case, one chooses

$$S \in \arg \max_{I \in J_{n_2}} \left[ \max_{p \in \mathfrak{P}_{n_2}} \kappa_{n_1, p}^{-1} S_p(\beta_n^*, N_1, I) \right].$$

- *The  $d$  largest scaled moments:* Let  $S$  be the indices of the  $d$  largest elements of

$$\left\{ \left| \left[ \hat{\Sigma}(\beta_n^*, N_1, \{j\}) \right]^{-1/2} \mathbf{H}_{\{j\}}(\beta_n^*, N_1) \right|, j = 1, \dots, D \right\}, \quad N_1 \neq \emptyset.$$

This amounts to testing those  $S$  individual moments that most strongly indicate a violation of (21) (taking into account the scale of these, but ignoring their correlation structure). It is computationally cheaper than the method in the previous point.

- One can use *any* state-of-the-art procedure to select the moments to be tested. For example, in the context of hypothesis testing in the presence of many instruments, cf. Sections 2.1.1 and 5, one can use the Lasso in the first step with a choice of penalty parameter ensuring that the number of selected instruments  $d$  satisfies the growth conditions imposed in the previous sections.

## References

ABDALLA, P. AND N. ZHIVOTOVSKIY (2022): “Covariance estimation: Optimal dimension-free guarantees for adversarial corruption and heavy tails,” *Journal of the*

*European Mathematical Society (forthcoming).*

- ANATOLYEV, S. AND N. GOSPODINOV (2011): “Specification testing in models with many instruments,” *Econometric Theory*, 27, 427–441.
- ANDERSON, T. (1955): “The integral of a symmetric unimodal function over a symmetric convex set and some probability inequalities,” *Proceedings of the American Mathematical Society*, 6, 170–176.
- ANDERSON, T. W. AND H. RUBIN (1949): “Estimation of the parameters of a single equation in a complete system of stochastic equations,” *The Annals of Mathematical Statistics*, 20, 46–63.
- ANDREWS, D. W. AND J. H. STOCK (2007): “Testing with many weak instruments,” *Journal of Econometrics*, 138, 24–46.
- BELLONI, A., D. CHEN, V. CHERNOZHUKOV, AND C. HANSEN (2012): “Sparse models and methods for optimal instruments with an application to eminent domain,” *Econometrica*, 80, 2369–2429.
- BELLONI, A., V. CHERNOZHUKOV, D. CHETVERIKOV, C. HANSEN, AND K. KATO (2018): “High-dimensional econometrics and regularized GMM,” *arXiv preprint arXiv:1806.01888*.
- BENTKUS, V. (2003): “On the dependence of the Berry–Esseen bound on dimension,” *Journal of Statistical Planning and Inference*, 113, 385–402.
- BICKEL, P. AND E. LEVINA (2008): “Covariance regularization by thresholding,” *The Annals of Statistics*, 36, 2577–2604.
- BLANDHOL, C., J. BONNEY, M. MOGSTAD, AND A. TORGOVITSKY (2022): “When is tsls actually late?” Tech. rep., National Bureau of Economic Research.
- BOOT, T. AND J. W. LIGTENBERG (2023): “Identification-and many instrument-robust inference via invariant moment conditions,” *arXiv preprint arXiv:2303.07822*.
- CAI, T., Z. REN, AND H. ZHOU (2016): “Estimating structured high-dimensional covariance and precision matrices: Optimal rates and adaptive estimation,” *Electronic Journal of Statistics*, 10, 1–59.

- CHERNOZHUKOV, V., D. CHETVERIKOV, AND K. KATO (2013): “Gaussian approximations and multiplier bootstrap for maxima of sums of high-dimensional random vectors,” *The Annals of Statistics*, 41, 2786–2819.
- (2017a): “Central limit theorems and bootstrap in high dimensions,” *The Annals of Probability*, 45, 2309–2352.
- (2017b): “Detailed proof of Nazarov’s inequality,” *arXiv preprint arXiv:1711.10696*.
- (2019): “Inference on causal and structural parameters using many moment inequalities,” *The Review of Economic Studies*, 86, 1867–1900.
- CHERNOZHUKOV, V., D. CHETVERIKOV, K. KATO, AND Y. KOIKE (2023): “High-dimensional data bootstrap,” *Annual Review of Statistics and Its Application*, 10, 427–449.
- CRUDU, F., G. MELLACE, AND Z. SÁNDOR (2021): “Inference in instrumental variable models with heteroskedasticity and many instruments,” *Econometric Theory*, 37, 281–310.
- DAVEY SMITH, G. AND S. EBRAHIM (2003): ““Mendelian randomization”: can genetic epidemiology contribute to understanding environmental determinants of disease?” *International journal of epidemiology*, 32, 1–22.
- DOVÌ, M.-S., A. B. KOCK, AND S. MAVROEIDIS (2022): “A Ridge-Regularised Jackknifed Anderson-Rubin Test,” *arXiv preprint arXiv:2209.03259*.
- FAN, J., Y. LIAO, AND H. LIU (2016): “An overview of the estimation of large covariance and precision matrices,” *The Econometrics Journal*, 19, C1–C32.
- FAN, J., Y. LIAO, AND J. YAO (2015): “Power enhancement in high-dimensional cross-sectional tests,” *Econometrica*, 83, 1497–1541.
- FAN, Q., Z. GUO, AND M. ZIWEI (2023): “A Heteroskedasticity-Robust Overidentifying Restriction Test with High-Dimensional Covariates,” *arXiv preprint arXiv:2205.00171*.
- FANG, X. AND Y. KOIKE (2024): “Large-dimensional central limit theorem with fourth-moment error bounds on convex sets and balls,” *Annals of Applied Probability*, 34, 2065–2106.

- FENG, L., T. JIANG, B. LIU, AND W. XIONG (2022): “Max-sum tests for cross-sectional independence of high-dimensional panel data,” *Annals of Statistics*, 50, 1124 – 1143.
- GABAIX, X. AND R. S. KOIJEN (2020): “Granular instrumental variables,” Tech. rep., National Bureau of Economic Research.
- GE, S., S. LI, AND O. LINTON (2024): “Dynamic peer groups of arbitrage characteristics,” *Journal of Business & Economic Statistics*, 42, 367–390.
- GOLDBERG, M. (1987): “Equivalence constants for  $\ell_p$  norms of matrices,” *Linear and Multilinear Algebra*, 21, 173–179.
- HANSEN, L. P. AND K. SINGLETON (1982): “Generalized instrumental variables estimation of nonlinear rational expectations models,” *Econometrica*, 1269–1286.
- HE, Y., G. XU, C. WU, AND W. PAN (2021): “Asymptotically independent U-statistics in high-dimensional testing,” *The Annals of Statistics*, 49, 154–181.
- HORN, R. A. AND C. R. JOHNSON (1985): *Matrix analysis*, Cambridge University Press.
- INGSTER, Y. AND I. SUSLINA (2003): *Nonparametric goodness-of-fit testing under Gaussian models*, vol. 169, Springer Science & Business Media.
- JAMMALAMADAKA, S. R., S. MEINTANIS, AND T. VERDEBOUT (2020): “On Sobolev tests of uniformity on the circle with an extension to the sphere,” *Bernoulli*, 26, 2226–2252.
- JUODIS, A. AND S. REESE (2022): “The incidental parameters problem in testing for remaining cross-section correlation,” *Journal of Business & Economic Statistics*, 40, 1191–1203.
- KAFFO, M. AND W. WANG (2017): “On bootstrap validity for specification testing with many weak instruments,” *Economics Letters*, 157, 107–111.
- KE, Y., S. MINSKER, Z. REN, Q. SUN, AND W.-X. ZHOU (2019): “User-friendly covariance estimation for heavy-tailed distributions,” *Statistical Science*, 34, 454–471.
- KOCK, A. B. AND D. PREINERSTORFER (2019): “Power in High-Dimensional Testing Problems,” *Econometrica*, 87, 1055–1069.

- (2023): “Consistency of p-norm based tests in high dimensions: Characterization, monotonicity, domination,” *Bernoulli*, 29, 2544–2573.
- (2024): “A remark on moment-dependent phase transitions in high-dimensional Gaussian approximations,” *Statistics & Probability Letters*.
- KOLTCHINSKII, V. AND K. LOUNICI (2017): “Concentration inequalities and moment bounds for sample covariance operators,” *Bernoulli*, 110–133.
- LI, D., L. XUE, H. YANG, AND X. YU (2024): “Power-Enhanced Two-Sample Mean Tests for High-Dimensional Compositional Data with Application to Microbiome Data Analysis,” *arXiv preprint arXiv:2405.02551*.
- LIU, Y., W. SUN, A. P. REINER, C. KOOPERBERG, AND Q. HE (2019): “Statistical inference of genetic pathway analysis in high dimensions,” *Biometrika*, 106, 651–664.
- LUGOSI, G. AND S. MENDELSON (2019): “Mean estimation and regression under heavy-tailed distributions: A survey,” *Foundations of Computational Mathematics*, 19, 1145–1190.
- MATSUSHITA, Y. AND T. OTSU (2021): “Second-order refinements for t-ratios with many instruments,” *Journal of Econometrics*, 232, 346–366.
- (2022): “A jackknife Lagrange multiplier test with many weak instruments,” *Econometric Theory*.
- MENDELSON, S. AND N. ZHIVOTOVSKIY (2020): “Robust covariance estimation under  $L_4 - L_2$  norm equivalence,” *The Annals of Statistics*, 48, 1648–1664.
- MIKUSHEVA, A. (2021): “Many Weak Instruments in Time Series Econometrics,” .
- MIKUSHEVA, A. AND L. SUN (2022): “Inference with many weak instruments,” *The Review of Economic Studies*, 89, 2663–2686.
- (2024): “Weak Identification with Many Instruments,” *Econometrics Journal*.
- MOLINARI, F. (2020): “Microeconometrics with partial identification,” *Handbook of econometrics*, 7, 355–486.

- NAZAROV, F. (2003): “On the maximal perimeter of a convex set in  $\mathbb{R}^n$  with respect to a gaussian measure,” in *Geometric Aspects of Functional Analysis*, Springer, 169–187.
- OLIVEIRA, R. I. AND Z. F. RICO (2022): “Improved covariance estimation: optimal robustness and sub-Gaussian guarantees under heavy tails,” *arXiv preprint arXiv:2209.13485*.
- PINELIS, I. (2010): “Asymptotic efficiency of p-mean tests for means in high dimensions,” *arXiv preprint arXiv:1006.0505*.
- ROSEN, A. M. (2008): “Confidence sets for partially identified parameters that satisfy a finite number of moment inequalities,” *Journal of Econometrics*, 146, 107–117.
- STOCK, J. H. AND J. H. WRIGHT (2000): “GMM with weak identification,” *Econometrica*, 68, 1055–1096.
- TANG, Y., H. J. WANG, AND E. BARUT (2018): “Testing for the presence of significant covariates through conditional marginal regression,” *Biometrika*, 105, 57–71.
- TCHUENTE, G. AND M. CARRASCO (2016): “Regularization Based Anderson Rubin Tests for Many Instruments,” .
- VERSHYNIN, R. (2012): “How close is the sample covariance matrix to the actual covariance matrix?” *Journal of Theoretical Probability*, 25, 655–686.
- XU, G., L. LIN, P. WEI, AND W. PAN (2016): “An adaptive two-sample test for high-dimensional means,” *Biometrika*, 103, 609–624.
- YANG, Q. AND G. PAN (2017): “Weighted statistic in detecting faint and sparse alternatives for high-dimensional covariance matrices,” *Journal of the American Statistical Association*, 112, 188–200.
- YU, X., D. LI, AND L. XUE (2024a): “Fisher’s combined probability test for high-dimensional covariance matrices,” *Journal of the American Statistical Association*, 119, 511–524.
- YU, X., D. LI, L. XUE, AND R. LI (2023): “Power-enhanced simultaneous test of high-dimensional mean vectors and covariance matrices with application to gene-set testing,” *Journal of the American Statistical Association*, 118, 2548–2561.

YU, X., J. YAO, AND L. XUE (2024b): “Power enhancement for testing multi-factor asset pricing models via Fisher’s method,” *Journal of Econometrics*, 239.

ZHANG, D. AND W. B. WU (2017): “Gaussian approximation for high dimensional time series,” *Annals of Statistics*, 45, 1895–1919.

ZHANG, Y., R. WANG, AND X. SHAO (2022): “Adaptive inference for change points in high-dimensional data,” *Journal of the American Statistical Association*, 117, 1751–1752.