

# Fast convergence of the Expectation Maximization algorithm under a logarithmic Sobolev inequality

Rocco Caprio and Adam M. Johansen  
Department of Statistics, University of Warwick.

Email: {[rocco.caprio](mailto:rocco.caprio@warwick.ac.uk), [a.m.johansen](mailto:a.m.johansen@warwick.ac.uk)}@warwick.ac.uk.

July 26, 2024

## Abstract

By utilizing recently developed tools for constructing gradient flows on Wasserstein spaces, we extend an analysis technique commonly employed to understand alternating minimization algorithms on Euclidean space to the Expectation Maximization (EM) algorithm via its representation as coordinate-wise minimization on the product of a Euclidean space and a space of probability distributions due to Neal and Hinton (1998). In so doing we obtain finite sample error bounds and exponential convergence of the EM algorithm under a natural generalisation of a log-Sobolev inequality. We further demonstrate that the analysis technique is sufficiently flexible to allow also the analysis of several variants of the EM algorithm.

**Keywords:** EM algorithm, first-order EM algorithm, maximum likelihood estimator, empirical Bayes, latent variable models, non-asymptotic bounds, functional inequalities, log-Sobolev inequality, Wasserstein gradient.

## 1 Introduction

The Expectation Maximization (EM) algorithm has been a central part of the statistician’s toolbox since being formalised by [22] as an effective general computational solution to the marginal maximum likelihood problem. At that time the algorithm had been proposed previously in numerous special contexts, including that of empirical Bayes [27]. Empirical Bayes methods have received considerable attention in the modern machine learning literature, where they are widely used to specify hyper-parameters in high-dimensional models.

In recent years there has been a great deal of interest within the Bayesian statistics and machine learning communities in the construction of gradient flows, especially Wasserstein gradient flows, which underlie Langevin Monte Carlo algorithms. Some recent work has focussed on the intersection of empirical Bayes type methods and gradient flow-based algorithms. Our aim is to demonstrate here that some of the tools, particularly those emerging from optimal transport and Wasserstein geometry, which have been developed in the context of these modern computational methods provide a natural approach to the analysis of the EM algorithm itself—and many of its approximations. Such analysis is quite direct, requires limited further technical work and yields state-of-the-art conclusions under conditions which are, if anything, weaker than those ordinarily employed in the quantitative analysis of EM algorithms.

In this paper we utilize the connection between EM and a coordinate-wise minimization algorithm applied to the free energy functional identified by [43] to provide non-asymptotic error bounds for EM algorithms under an extended form of the log-Sobolev inequality. To do this, we extend an argument commonly used to understand Euclidean coordinate descent algorithms by comparison with gradient descent via the descent lemma [9, 8, 10], together with recently developed results for using and understanding gradients on the product of Euclidean and Wasserstein spaces [13].

## 1.1 The Expectation Maximization Algorithm

Consider the problem of fitting a probabilistic model, with Lebesgue density  $p_\theta(x, y)$ , featuring latent variables,  $x \in \mathbb{R}^{d_x}$ , to data,  $y$ , which have been observed. Within the maximum likelihood framework one seeks model parameters,  $\theta \in \mathbb{R}^{d_\theta}$ , that maximize the (marginal) likelihood — the probability density,  $p_\theta(y)$ , of observing the obtained data upon integrating out the latent variables: that is,  $\theta_\star$  belonging to

$$\mathcal{O}_\star := \arg \max_{\theta \in \mathbb{R}^{d_\theta}} p_\theta(y) = \arg \max_{\theta \in \mathbb{R}^{d_\theta}} \int p_\theta(x, y) dx. \quad (1)$$

One can also characterize the conditional distribution of the latent variables given the observed data under the model specified by parameter vector  $\theta_\star$  as:

$$p_{\theta_\star}(x|y) := \frac{p_{\theta_\star}(x, y)}{p_{\theta_\star}(y)},$$

and due to the Bayesian flavour of this computation and the connection with empirical Bayes [49] we will term this the *posterior* distribution of  $x$  throughout. For most models of practical interest, the integral in (1) is intractable, we have no closed-form expressions for  $p_\theta(y)$  or its derivatives, and we are unable to directly optimize  $p_\theta(y)$ . The most popular algorithm to solve this problem is the Expectation Maximization (EM) algorithm.

**Notation.** As the data  $y$  can be treated as fixed, in the description of the algorithm below and in the remainder we shorten our notation and write  $\rho_\theta(x)$  for  $p_\theta(x, y)$ ,  $Z_\theta$  for its normalizing constant  $p_\theta(y)$ ,  $\pi_\theta(x) = \rho_\theta(x)/Z_\theta$  for  $p_\theta(x|y)$ , and additionally  $\ell(\theta; x)$  for the (complete) log-likelihood  $\log(\rho_\theta(x))$ . When dealing with measures absolutely continuous with respect to the Lebesgue measure we use the same symbol to refer to both the measure and its Lebesgue density. A summary of this and other relevant notation can be found in Appendix A.

---

### Algorithm 1 EM Algorithm

---

- 1: **Inputs:** Initial values  $(\theta_0, q_0)$ .
- 2: **for**  $k \geq 0$  **do**
- 3: (M-step) Update the parameter estimate:

$$\theta_{k+1} = \arg \max_{\theta} \int \ell(\theta; x) q_k(dx). \quad (2)$$

- 4: (E-step) Update the posterior estimate:

$$q_{k+1} = \pi_{\theta_{k+1}}. \quad (3)$$

- 5: **end for**
-

In Algorithm 1, we perform the M-step first, hence the initial value  $\theta_0$  is not used. In some cases, it might be more convenient to perform the E-step first by specifying an initial guess  $\theta_0$  and then setting  $q_0 := \pi_{\theta_0}$ . Our results apply to both cases.

**Related literature.** The importance of EM in statistics, machine learning and science more broadly, means that its convergence properties have been extensively studied. Beyond the characterization provided by [22], early analysis was provided by [12, 58]. These and other works [30, 40, 39] focus on asymptotic convergence rates. However, relatively little is known about the non-asymptotic performance. [33, 34] leverage interesting connections between the EM algorithm, generalized surrogate optimization methods and mirror descent, respectively, to study this problem. In particular, [33] shows global sub-exponential rates for the parameter estimates under the hypothesis of strong concavity of the *surrogate function*

$$\theta \mapsto Q(\theta|\theta') = \int \ell(\theta; x) \pi_{\theta'}(dx). \quad (4)$$

for all  $\theta' \in \mathbb{R}^{d_\theta}$ . [34] obtains global sub-exponential rates for the posterior estimates when  $(\rho_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  forms a minimal exponential family, which implies that the surrogate is concave. [7] studies the related but different problem of deriving convergence rates to the true population parameter, and not the MLE, in the case of an infinite sample (‘population EM’) and of a finite sample (‘sample EM’), by also considering an assumption of strong concavity of the surrogate and provided one starts the algorithm in some neighbourhood of the optimum. Other lines of work have focused on more specific models or situations, such as Gaussian mixtures [59, 29, 56], the case of misspecified models [24], or stochastic EM methods [32].

## 1.2 A differential analysis of the EM Algorithm

In this work, we take a somewhat different approach to earlier literature and we connect EM’s exponential convergence with the concepts of gradients in the space of probability distributions, appropriate log-Sobolev inequalities, and other results and techniques in optimal transport and sampling. Let  $\mathcal{P}(\mathbb{R}^{d_x})$  denote the space of the Borel probability measures on the Euclidean space  $\mathbb{R}^{d_x}$ . We start from the following key observation, due to [43], that connects the EM algorithm to an alternating minimization procedure on  $\mathcal{M} := \mathbb{R}^{d_\theta} \times \mathcal{P}(\mathbb{R}^{d_x})$ .

**Proposition 1.** *The steps of the EM iteration, (2) and (3), are, respectively, equivalent to*

$$\theta_{k+1} \in \arg \min_{\theta \in \mathbb{R}^{d_\theta}} F(\theta, q_k), \quad (2')$$

$$q_{k+1} \in \arg \min_{q \in \mathcal{P}(\mathbb{R}^{d_x})} F(\theta_{k+1}, q), \quad (3')$$

where  $F : \mathbb{R}^{d_\theta} \times \mathcal{P}(\mathbb{R}^{d_x}) \rightarrow \mathbb{R}$  is the *free energy* functional:

$$F(\theta, q) := \begin{cases} \int \log \left( \frac{q(x)}{\rho_\theta(x)} \right) q(dx) & \text{if } q \ll \rho_\theta(dx) \\ +\infty & \text{otherwise} \end{cases} \quad \forall (\theta, q) \in \mathcal{M}. \quad (5)$$

For reasons that will become clear later, we focus our attention on  $\mathcal{M}_2 := \mathbb{R}^{d_\theta} \times \mathcal{P}_2(\mathbb{R}^{d_x})$ , where  $\mathcal{P}_2(\mathbb{R}^{d_x})$  is the restriction of  $\mathcal{P}(\mathbb{R}^{d_x})$  to elements which are absolutely continuous with

respect to Lebesgue measure and admit finite second moments. Since EM is a minimization procedure on  $F$ , it is key to the analysis to understand  $F$ 's variations along the EM iterates. Proposition 1 shows that the free energy can only decrease along the EM iterations:

$$F(\theta_0, q_0) \geq F(\theta_1, q_0) \geq F(\theta_1, q_1) \geq \dots \geq F(\theta_k, q_k) \geq F(\theta_{k+1}, q_k) \geq F(\theta_{k+1}, q_{k+1}),$$

and to characterize EM's exponential convergence, we would like to quantify this decrease. The alternating minimization representation of EM suggests we might be able to adapt the analysis of alternating minimization algorithms on Euclidean space [9, 8, 10]. When minimizing a smooth function  $f : \mathbb{R}^{d_x} \rightarrow \mathbb{R}$  satisfying a gradient growth condition

$$2\lambda[f(x) - f_\star] \leq \|\nabla_x f(x)\|^2 \quad \forall x \in \mathbb{R}^{d_x}, \quad \text{where } \lambda > 0 \quad \text{and} \quad f_\star := \inf f$$

known as Polyak–Łojasiewicz inequality (PLI), this analysis can be conducted by comparing the alternating minimization updates with appropriate gradient steps. Since the free energy is a function on  $\mathcal{M}_2$ , it is not immediately clear how to translate this approach to our setting. However, recent advances in optimal transport provide natural solutions to this problem.

In particular, we show that under a smoothness assumption, we can lower bound EM's free energy decrease abstractly in terms of the norm of  $\text{grad}_{\mathcal{M}_2} F$ ,  $F$ 's gradient in the geometry induced by the product of the Euclidean and Wasserstein metrics on  $\mathcal{M}_2$ . Because the posterior updates is on the space of probability measures, the notion of a gradient step with which to compare the EM step relates to the concepts of Wasserstein gradient flows. Having lower-bounded the decrease in free energy in terms of  $\text{grad}_{\mathcal{M}_2} F$  via smoothness, we assume that  $\text{grad}_{\mathcal{M}_2} F$ 's norm grows at least quadratically away from  $F$ 's minimizers along EM iterations, a natural analogue of the PLI on  $\mathcal{M}_2$ . More precisely:

$$2\lambda[F(\theta, q) - F_\star] \leq \|\text{grad}_{\mathcal{M}_2} F(\theta, q)\|_{\mathcal{M}_2}^2 \tag{6}$$

for all relevant EM iterates  $(\theta, q)$ , where  $\lambda > 0$  is a positive constant and where

$$F_\star := \inf_{(\theta, q) \in \mathcal{M}_2} F(\theta, q) = \inf_{\theta \in \mathbb{R}^{d_\theta}} F(\theta, \pi_\theta) = -\log \left( \sup_{\theta \in \mathbb{R}^{d_\theta}} Z_\theta \right) =: -\log Z_\star.$$

Inequality (6) is a generalization of the log-Sobolev and Polyak–Łojasiewicz inequalities (LSI and PLI, respectively) of optimal transport and optimization, and was studied in [13] in the context of a gradient flow on  $\mathcal{M}_2$ , where it appears naturally. Whether the log-Sobolev type inequality (6) holds depends solely on the model. In particular, it is always verified if the model is strongly log-concave.

Once the convergence of the free energy of EM iterates to  $F_\star$  via (6) has been established, we investigate fast convergence of the EM iterates themselves  $(\theta_k, q_k)$  to  $\mathcal{M}_\star := \arg \min F(\theta, q) = \{(\theta_\star, \pi_{\theta_\star}) : \theta_\star \in \mathcal{O}_\star\}$ , i.e. to a (local) maximum of the marginal likelihood and its corresponding posterior. The decomposition

$$F(\theta_k, q_k) - F_\star = \text{KL}(q_k || \pi_{\theta_k}) + [\log(Z_\star) - \log(Z_{\theta_k})] \tag{7}$$

where KL denotes the Kullback–Leibler divergence, suggests that the convergence in free energy should also give information on convergence of both the EM iterates, since  $\text{KL}(q_k || \pi_{\theta_k})$  gives information on the convergence of the posterior updates  $q_k$  in terms of KL, while  $\log(Z_\star) - \log(Z_{\theta_k})$  tracks the convergence of the parameter updates  $\theta_k$ . An natural extension of the Otto–Villani Theorem [46] enables this intuition: inequality (6) actually implies an extension of the Talagrand inequality [52]:

$$2[F(\theta, q) - F_\star] \geq \lambda d((\theta, q), \mathcal{M}_\star)^2, \tag{8}$$

where  $\mathbf{d}$  is the product metric

$$\mathbf{d}((\theta, q), (\theta', q')) := \sqrt{\mathbf{d}_E(\theta, \theta')^2 + \mathbf{d}_{W_2}(q, q')^2}, \quad (9)$$

with  $\mathbf{d}_E$  and  $\mathbf{d}_{W_2}$  denoting the Euclidean and Wasserstein-2 distances, respectively. As a result, we are able to derive non-asymptotic convergence error bounds in  $\mathbf{d}$ -distance for both the posterior and parameter estimates of EM by assuming solely a smoothness condition and that the model satisfies the log-Sobolev type inequality (6).

### 1.3 Paper structure

This paper is organized as follows. In Section 2 we study the inequality (6), primarily using results from [13]. In particular, we derive an expression for  $\text{grad}_{\mathcal{M}_2} F$ , study the relationship between (6) and (8), and derive some sufficient conditions on the model to verify these. In Section 3 we derive non-asymptotic rates of convergence for the EM algorithm under condition (6) and a smoothness assumption using the approach we outlined. We further analyze three alternatives to the EM that can be used when either the E- or M- step (2)–(3) is intractable (or both are intractable) and compare their convergence properties illustrating the potential of the method to study the many variants of the EM algorithm.

## 2 Differential inequalities for EM

As outlined above, the EM algorithm can be viewed as a minimization procedure applied to the free energy  $F$ . We aim to understand how to think about free energy dissipation along the EM iterations, and in particular how to think about its gradient, and hence to understand (6). In Section 2.1, we derive an expression for  $\text{grad}_{\mathcal{M}_2} F(\theta, q)$  and its norm. Our aim there is solely motivating the use of a certain functional (given in (12) below) for that later, and in this sense it can be skipped on a first reading. In Section 2.2 we study inequality (6), characterizing it as a generalization of the well-known log-Sobolev inequality. Using results recently derived in [13], we connect it with the generalization of the Talagrand inequality (8). In Section 2.3 we study sufficient conditions on the model to verify (6), and show how we can leverage advances from the functional inequalities literature to investigate the performance of models with different completions.

### 2.1 Differentiating the free energy in $\mathcal{M}_2$

In this section we show how we can derive the gradient of the free energy in  $\mathcal{M}_2$ 's geometry, which is key for our analysis, and which we need to make sense of the inequality (6). To do so, we leverage the formal interpretation of  $\mathcal{P}_2(\mathbb{R}^{d_x})$  endowed with the Wasserstein-2 distance  $\mathbf{d}_{W_2}$  as a Riemannian-manifold, an approach pioneered in [45]. These computations were carried out in [35] to make sense of gradient descent for  $F$ , which we discuss later. In this section we skim over technical details; this approach is not completely rigorous but more technical arguments following the approach in [2] yield the same result. None of our results depend upon this informal reasoning.

If we think of  $\mathcal{M}_2$  as a Riemannian manifold, in order to define gradients on  $\mathcal{M}_2$ , we need a sensible notion of tangent space and inner product. Because of the product structure of  $\mathcal{M}_2 = \mathbb{R}^{d_\theta} \times \mathcal{P}_2(\mathbb{R}^{d_x})$ , our tangent space will be the product of the tangent spaces of each component. For the  $\mathbb{R}^{d_\theta}$  component, we identify the tangent space at any point  $\theta$  as

$\mathbb{R}^{d_\theta}$  itself, so that  $\mathcal{T}_\theta \mathbb{R}^{d_\theta} = \mathbb{R}^{d_\theta}$ . For  $\mathcal{P}_2(\mathbb{R}^{d_x})$  we consider the geometry induced by the Wasserstein-2 distance  $\mathbf{d}_{W_2}$ . With this choice, we think at the tangent space at any point  $q \in \mathcal{P}_2(\mathbb{R}^{d_x})$  as the space of Lebesgue-integrable functions with zero integral

$$\mathcal{T}_q \mathcal{P}_2(\mathbb{R}^{d_x}) := \left\{ h : \int h(x) dx = 0 \right\},$$

which we equip with the following inner product.

**Definition 1** (Wasserstein-2 inner product). *Given two elements  $h_1, h_2 \in \mathcal{T}_q \mathcal{P}_2(\mathbb{R}^{d_x})$ , we define their Wasserstein-2 inner product at  $q \in \mathcal{P}_2(\mathbb{R}^{d_x})$  as*

$$\langle h_1, h_2 \rangle_{W_2} := \int \langle \nabla_x \psi_1(x), \nabla_x \psi_2(x) \rangle q(dx), \quad \text{where } \psi_i \text{ solves } \nabla_x \cdot (q \nabla_x \psi_i) = -h_i \quad i = 1, 2,$$

and we denote with  $\|\cdot\|_{W_2}$  the induced norm.

One motivation for this choice of tangent space and inner product comes from comparing the Benamou–Brenier formula [55, p.159],

$$\mathbf{d}_{W_2}(q, p) = \inf_{q_t} \left\{ \int_0^1 \inf_{v_t} \left\{ \int \|v_t\|^2 q_t(dx) dt : \nabla_x \cdot (q_t v_t) = -\partial_t q_t \right\} : q_0 = q, q_1 = p \right\}, \quad (10)$$

with the formula for the distance between two points in a Riemannian manifold  $(M, \mathbf{d}_M)$

$$\mathbf{d}_M(p, q) = \inf_{q_t} \left\{ \int_0^1 \|\partial_t q_t\|_M dt : q_0 = q, q_1 = p \right\},$$

upon noting that the optimal ‘velocity’ field  $v_t$  realizing the infimum in (10) is achieved by a gradient of a function  $\nabla_x \psi$ , and that for  $\nabla_x \cdot (q \nabla_x \psi) = -h$  to be solvable it is required that  $h$  has zero Lebesgue integral. See [26, 55] for more details.

For a point  $(\theta, q) \in \mathcal{M}_2$ , we set  $\mathcal{T}_{(\theta, q)} \mathcal{M}_2 := \mathcal{T}_\theta \mathbb{R}^{d_\theta} \times \mathcal{T}_q \mathcal{P}_2(\mathbb{R}^{d_x})$  and we endow  $\mathcal{M}_2$  with the following inner product which arises naturally from the product-space structure.

**Definition 2** ( $\mathcal{M}_2$  inner product). *Given two elements  $(a_1, h_1), (a_2, h_2) \in \mathcal{T}_{(\theta, q)} \mathcal{M}_2$ , we define their  $\mathcal{M}_2$  inner product as*

$$\langle (a_1, h_1), (a_2, h_2) \rangle_{\mathcal{M}_2} := \langle a_1, a_2 \rangle + \langle h_1, h_2 \rangle_{W_2}$$

and we denote with  $\|\cdot\|_{\mathcal{M}_2}$  the induced norm.

We can now define the gradient on  $\mathcal{M}_2$  in analogy with Riemannian geometry (see, e.g., Section 3 in [11]).

**Definition 3** (Gradients in  $\mathcal{M}_2$ ). *For a functional  $F$  on  $\mathcal{M}_2$ , its  $\mathcal{M}_2$ -gradient at  $(\theta, q) \in \mathcal{M}_2$  is the unique function  $\text{grad}_{\mathcal{M}_2} F(\theta, q) = (\text{grad}_{\mathbb{R}^{d_\theta}} F(\theta, q), \text{grad}_{\mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta, q))$  such that*

$$\frac{d}{dt} \Big|_{t=0} F(\theta_t, q_t) = \left\langle \text{grad}_{\mathcal{M}_2} F(\theta, q), \left( \frac{d}{dt} \Big|_{t=0} \theta_t, \frac{d}{dt} \Big|_{t=0} q_t \right) \right\rangle_{\mathcal{M}_2}$$

for any smooth curve  $t \mapsto (\theta_t, q_t)$  such that  $(\theta_0, q_0) = (\theta, q)$ , provided that it exists.

The first variation will prove useful in calculating gradients, as the following lemma shows.

**Definition 4** (First variation). *For a functional  $F$  on  $\mathcal{M}_2$ , its first variation at  $(\theta, q) \in \mathcal{M}_2$  is the unique (up to an additive constant) function  $\delta_{\mathcal{M}_2}F(\theta, q) = (\delta_{\mathbb{R}^{d_\theta}}F(\theta, q), \delta_{\mathcal{P}_2(\mathbb{R}^{d_x})}F(\theta, q))$  such that*

$$\frac{d}{dt}\Big|_{t=0} F(\theta_t, q_t) = \left\langle \delta_{\mathcal{M}_2}F(\theta, q), \left( \frac{d}{dt}\Big|_{t=0} \theta_t, \frac{d}{dt}\Big|_{t=0} q_t \right) \right\rangle_{\mathcal{L}^2(dx)}$$

for any smooth curve  $t \mapsto (\theta_t, q_t)$  such that  $(\theta_0, q_0) = (\theta, q)$ , provided that it exists.

**Lemma 2.** *If it exists,  $\text{grad}_{\mathcal{M}_2}F(\theta, q)$  satisfies*

$$\text{grad}_{\mathcal{M}_2}F(\theta, q) = \left( \delta_{\mathbb{R}^{d_\theta}}F(\theta, q), -\nabla_x \cdot (q \nabla_x \delta_{\mathcal{P}_2(\mathbb{R}^{d_x})}F(\theta, q)) \right).$$

*Proof.* See Appendix B.1. □

Using Definition 4 we readily compute  $\delta_{\mathcal{M}_2}F(\theta, q) = (-\int \nabla_\theta \ell(\theta; x)q(dx), \log(q/\rho_\theta))$ , and using the above lemma we identify the following gradient of the free energy:

$$\text{grad}_{\mathcal{M}_2}F(\theta, q) = \left( -\int \nabla_\theta \ell(\theta; x)q(dx), -\nabla_x \cdot (q(\nabla_x \log(q) - \nabla_x \log(\rho_\theta))) \right). \quad (11)$$

## 2.2 Log-Sobolev and Talagrand Inequalities on $\mathcal{M}_2$

Having a notion of gradient on  $\mathcal{M}_2$ , we are ready to define a very important quantity—the squared norm of the gradient of the free energy in  $\mathcal{M}_2$ :

$$I(\theta, q) := \|\text{grad}_{\mathcal{M}_2}F(\theta, q)\|_{\mathcal{M}_2}^2 = \left\| \int \nabla_\theta \ell(\theta; x)q(dx) \right\|^2 + \int \left\| \nabla_x \log \left( \frac{q(x)}{\rho_\theta(x)} \right) \right\|^2 q(dx). \quad (12)$$

We refer to this quantity as the *extended Fisher information functional*, because if the parameter space is the trivial space  $\{\theta\}$ , this quantity reduces to the relative Fisher information functional (e.g. see equation (8) in [46]):

$$I(q|\pi_\theta) := \int \left\| \nabla_x \log \left( \frac{q(x)}{\pi_\theta(x)} \right) \right\|^2 q(dx). \quad (13)$$

In light of this, upon considering the restriction  $\mathcal{P}_2^1(\mathbb{R}^{d_x})$  of  $\mathcal{P}_2(\mathbb{R}^{d_x})$  to probability measures having at least a.e. differentiable densities to ensure (12) is well defined, we can interpret the following as a gradient growth condition on the free energy in  $\mathcal{M}_2$ 's geometry.

**Definition 5** (Extended log-Sobolev inequality; xLSI). *The measures  $(\rho_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  satisfy the extended log-Sobolev inequality with constant  $\lambda > 0$  if (6) holds for all  $(\theta, q) \in \mathcal{M}_2 := \mathbb{R}^{d_\theta} \times \mathcal{P}_2^1(\mathbb{R}^{d_x})$ .*

This inequality generalizes both the well-known log-Sobolev inequality (LSI) in optimal transport [55, Definition 21.1] and the Polyak–Łojasiewicz inequality (PLI) in optimization [48, 38], which arise when  $\pi_\theta$  is independent of  $\theta$  and when we consider  $q = \pi_\theta$ , respectively. While the LSI is a statement about a single probability distribution, and PLI about a single function, the xLSI is a statement about a parameterized family of finite measures,

our (complete data) model. The xLSI does not imply that the marginal likelihood has a unique maximizer, and it allows for multiple stationary points. However, it implies that any stationary point is a global maximizer.

We will use the xLSI in conjunction with a smoothness assumption to conclude convergence of the free energy along EM iterates, but we would like to infer convergence of the iterates themselves in some appropriate distance. We consider the product metric on  $\mathcal{M}_2$  given by (9).

**Definition 6** (Extension of the Talagrand inequality; xT<sub>2</sub>I). *The measures  $(\rho_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  satisfy an extension of the Talagrand inequality with constant  $\lambda > 0$  if (8) holds for all  $(\theta, q) \in \mathcal{M}_2$ .*

We can show that xT<sub>2</sub>I generalizes both the Talagrand inequality (‘T<sub>2</sub>I’) [46, 52] and the quadratic growth (QG) condition [4] by considering again the cases when the parameter or the distribution spaces are trivial as above. To investigate the link between xT<sub>2</sub>I and the xLSI we impose some regularity conditions on the model.

**Assumption 1.** (i) For all  $x$  in  $\mathbb{R}^{d_x}$ ,  $\theta \mapsto \pi_\theta(x)$  is differentiable; and  $\theta \mapsto Z_\theta$  is differentiable; (ii) for all  $\theta$  in  $\mathbb{R}^{d_\theta}$ ,  $\pi_\theta$  is twice continuously differentiable; (iii) for all  $\theta$  in  $\mathbb{R}^{d_\theta}$  and  $x$  in  $\mathbb{R}^{d_x}$ ,  $\rho_\theta(x) > 0$ ; (iv)  $\pi_\theta$  has finite second moments for all  $\theta \in \mathbb{R}^{d_\theta}$ , and these are continuous in  $\theta$ ; (v)  $\mathcal{O}_*$  is compact.

**Assumption 2.** *The log-likelihood  $\ell$  is differentiable and its gradient  $\nabla \ell = (\nabla_\theta \ell, \nabla_x \ell)$  satisfies*

$$\|\nabla \ell(\theta; x) - \nabla \ell(\theta'; x')\| \leq L_\theta \|\theta - \theta'\| + L_x \|x - x'\|$$

for some constants  $L_\theta$  and  $L_x$ . In particular,  $\nabla \ell$  is  $L := \max(L_\theta, L_x)$ -Lipschitz.

Assumption 1 imposes only very mild regularity conditions on the model and it is typically satisfied by models used in practice. Conditions (iv,v) are only required for the results in Lemma 11 and Theorem 12 to ensure that  $\pi_{\theta_*}$ ’s second moments are uniformly bounded across  $\theta_* \in \mathcal{O}_*$ . If  $\mathcal{O}_*$  consists of finitely many points (e.g. if Assumption 3 below holds), in (iv) the continuity of second moments is not even needed. Assumption 2 is a classical, but more stringent, smoothness condition. We have the following generalization of the Otto–Villani Theorem [46]:

**Theorem 3** (Theorem 2 in [13]; xLSI  $\Rightarrow$  xT<sub>2</sub>I). *If Assumptions 1–2 hold, and the measures  $(\rho_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  satisfy the xLSI with constant  $\lambda$ , then they also satisfy the xT<sub>2</sub>I with the same constant.*

For our analysis, this Theorem tells us that it is enough to establish convergence of free energy of EM iterates under the xLSI, as then we can just use the xT<sub>2</sub>I to translate that into the convergence of the iterates themselves in d-distance.

### 2.3 Bakry–Émery and the perturbation and contraction principles

In this section we study some sufficient conditions on the model to verify the xLSI. A generalization of the Bakry and Émery criterion [5] shows that a convenient sufficient (but not necessary) condition for the xLSI is strong log-concavity.



**Assumption 3** (Strong log-concavity). *There exists a  $\lambda > 0$  such that*

$$\ell((1-t)\theta + t\theta'; (1-t)x + tx') \geq (1-t)\ell(\theta; x) + t\ell(\theta'; x') + \frac{\lambda t(1-t)}{2} \|(\theta, x) - (\theta', x')\|^2,$$

for all  $(\theta, x), (\theta', x')$  in  $\mathbb{R}^{d_\theta} \times \mathbb{R}^{d_x}$  and  $0 \leq t \leq 1$ .

**Theorem 4** (Theorem 3 in [13]; Strong log-concavity  $\Rightarrow$  xLSI). *Any model satisfying Assumptions 1(iii) and 3 satisfies the xLSI.*

We now show a simple example of a model class in which this Bakry–Émery argument allows us to verify that the xLSI holds.

**Example 1.** For  $m \in \mathbb{N}$ ,  $C_i \in \mathbb{R}^{d_x/m \times d_x/m}$ ,  $D \in \mathbb{R}^{d_\theta \times d_x/m}$ , consider the hierarchical model

$$Y_i = C_i X_i + U_i \quad X_i = D\theta + V_i \quad \text{for } i = 1, \dots, m,$$

where  $U_i$  and  $V_i$  are i.i.d. symmetric around 0 random vectors with distribution densities  $p_u$  and  $p_v$ , respectively. We think of this model as describing an underlying process of interest  $X$ , that depends on some parameter  $\theta$ , and which is only observed with noise via  $Y$ . The model is given by

$$\rho_\theta(x) := \prod_{i=1}^m p_u(y_i - C_i x_i) p_v(x_i - D\theta)$$

and within the Empirical Bayes setting, we would like to perform inference on the states using  $\pi_{\theta_\star}$ , with  $\theta_\star$  being the MLE, via the EM algorithm. In this setting, whether the xLSI holds depends solely on the tail behaviour of both the noise distributions  $p_u$  and  $p_v$ . It is common to assume that  $p_v$  is a Normal density. In this case, any strongly log-concave density  $p_v$  returns a model satisfying the xLSI by Theorem 4, and any gradient-Lipschitz density  $p_v$  returns a model satisfying Assumption 2. The Normal hierarchical model, which assumes that  $p_u$  and  $p_v$  are both Normal, also satisfies both requirements.

### 2.3.1 Operations preserving the xLSI and models with different completions

The literature on functional inequalities contains many results which show preservation of functional inequalities under various operations. For instance, the LSI is known to be preserved under contractive mappings [6], bounded perturbations [31], mixtures [17], convolutions with Gaussians and some class of smooth perturbations [15] and many more. These results extend considerably the settings where one can verify the LSI, showing that it goes far beyond strong log-concavity. These sorts of considerations also motivated the use of functional inequalities for the analysis of Langevin Monte Carlo [54, 18]. Here our goal is to illustrate that similar results hold for the xLSI, by generalizing the aforementioned contractive mapping and the bounded perturbation results.

For the LSI, these results allow us to conclude that if a probability distribution satisfies the LSI, then another distribution not too different from the original, as obtained as a result of these operations, still does. To understand what the xLSI analogues say, recall that for any given marginal likelihood,  $Z_\theta$ , there are many possible choices of complete likelihood,  $\rho_\theta = \pi_\theta Z_\theta$ , depending on the completion,  $\pi_\theta$ , the choice which is known to dramatically impact the performance and convergence properties of the associated EM algorithm [22, 41]. The following results then say that if a given model satisfies the xLSI, a model using another completion, that is not too different from the original, still does. As we will connect the fast convergence of EM with the xLSI, this in principle also gives an estimate on the performance we can expect from models obtained with different completions.

**Proposition 5** (Perturbation principle). *Let Assumption 1(i) hold, and suppose that the measures  $(\rho_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  satisfy the xLSI with constant  $\lambda > 0$ . Consider the measures  $(\tilde{\rho}_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  defined by  $\tilde{\rho}_\theta := \tilde{\pi}_\theta Z_\theta$ , where  $\tilde{\pi}_\theta$  is a bounded perturbation of  $\pi_\theta$  in the sense that  $c^{-1} \leq d\pi_\theta/d\tilde{\pi}_\theta \leq c$  for some  $c > 1$  independent of  $\theta$ , and that*

$$b := \sup \frac{\|[\nabla_\theta \rho_\theta - \nabla_\theta \tilde{\rho}_\theta]/\tilde{\rho}_\theta\|^2}{[\log(Z_\star) - \log(Z_\theta)]} < \infty. \quad (14)$$

*Then the measures  $(\tilde{\rho}_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  also satisfy the xLSI with constant  $(\lambda - c^2 \cdot b)/2c^2$ .*

*Proof.* See Appendix B.2. □

Notice that in the result above, the marginal likelihood  $Z_\theta$  of the perturbed model is the same as that of the unperturbed one, so we are comparing models with different completions defined on a common space. In the degenerate case in which  $\pi_\theta$  is independent of  $\theta$ , we can take  $b = 0$  and this result reduces, up to a factor of 2, to the Holley–Stroock perturbation lemma [31] (also see [6, Lemma 5.1.6]) which asserts that if a probability measure  $\pi_\theta$  satisfies the LSI with constant  $\lambda$ , then its bounded perturbation  $\tilde{\pi}_\theta$  also does, but with constant  $\lambda/c^2$ . In this case, Theorem 4 and Proposition 5 immediately imply that, under Assumption 1(i,iii), if the log-likelihood  $\ell$  is strongly concave only in the ‘tails’ i.e. over  $A^c \times \mathbb{R}^{d_\theta}$ , where  $A$  is some compact set in  $\mathbb{R}^{d_x}$ , then the xLSI will still hold. When  $\pi_\theta$  depends on  $\theta$ , (14) says that the gradients of the completions need to be identical in the stationary points of the marginal likelihood (at the MLE, where  $\log(Z_\theta) = \log(Z_\star)$ ), and that outside there it should not be too large relative to the tail behaviour of the marginal likelihood. On the other hand, the marginal log-likelihood is typically intractable so verifying (14) is complicated in real settings. Using this result, we speculate it should be possible to show that the xLSI can hold in situations where the surrogate (4) is not even concave.

As is the case for the LSI, the xLSI is preserved under the action of Lipschitz maps. For a probability measure  $\mu \in \mathcal{P}(\mathbb{R}^{d_x})$  and a measurable map  $T$  with domain  $\mathbb{R}^{d_x}$ , let  $T_\# \mu = \mu \circ T^{-1}$  denote the pushforward of  $\mu$  by  $T$ .

**Proposition 6** (Contraction principle). *Let Assumption 1(i) hold, and suppose that the measures  $(\rho_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  satisfy the xLSI with constant  $\lambda > 0$ . If  $\tilde{\pi}_\theta$  can be written as  $\tilde{\pi}_\theta = T_\# \pi_\theta$ , for some  $L_T$ -Lipschitz diffeomorphism  $T$ , then the measures  $(\tilde{\rho}_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  defined by  $\tilde{\rho}_\theta := \tilde{\pi}_\theta Z_\theta$  also satisfy the xLSI with constant  $\lambda/\max(1, L_T^2)$ .*

*Proof.* See Appendix B.3. □

Rather than deriving results in the most general form our aim here was to illustrate the flavour of this sort of results. We believe that such arguments have the potential to further characterize EM’s fast convergence; developing a theory similar to that other standard functional inequalities and Langevin Monte Carlo.

## 3 Non-asymptotic analysis of EM and related algorithms

### 3.1 EM Algorithm

The goal is to establish exponential convergence of the EM iterates under the xLSI. To do so, we first establish the exponential convergence of the free energy to its minimizer and

then transfer the result, via the extension of the Talagrand inequality and Theorem 3, to the EM iterates themselves. Proposition 1 shows that the free energy can only decrease along the EM iterations, and in order to prove EM's exponential convergence we need to quantify the decrease. Since we think of the extended Fisher information functional  $I$  (12) as the squared norm of the free energy gradient, we might expect that under a smoothness assumption we can lower bound the magnitude of this decrease in terms of  $I$ :

**Lemma 7.** *If Assumption 2 holds, for all  $k \in \mathbb{N}$ ,*

$$F(\theta_k, q_k) - F(\theta_{k+1}, q_k) \geq \frac{1}{2L_\theta} I(\theta_k, q_k) \quad (15)$$

*Proof.* Consider  $\vartheta_k := \theta_k + (1/L_\theta) \int \nabla_\theta \ell(\theta_k; x) q_k(\mathrm{d}x)$ . By the minimality of  $\theta_{k+1}$ ,

$$F(\theta_k, q_k) - F(\theta_{k+1}, q_k) \geq F(\theta_k, q_k) - F(\vartheta_k, q_k), \quad (16)$$

and because  $\theta \mapsto \nabla_\theta \ell(\theta; x)$  is  $L_\theta$ -Lipschitz for all  $x \in \mathbb{R}^{d_x}$ ,

$$\ell(x; \theta_k) - \ell(x; \vartheta_k) + \langle \nabla_\theta \ell(\theta_k; x), \vartheta_k - \theta_k \rangle \leq \frac{L_\theta}{2} \|\vartheta_k - \theta_k\|^2,$$

so that

$$\begin{aligned} F(\theta_k, q_k) - F(\vartheta_k, q_k) &= \int (\ell(\vartheta_k; x) - \ell(\theta_k; x)) q_k(\mathrm{d}x) \\ &\geq \int \left( \frac{1}{L_\theta} \left\langle \nabla_\theta \ell(\theta_k; x), \int \nabla_\theta \ell(\theta_k; \tilde{x}) q_k(\mathrm{d}\tilde{x}) \right\rangle - \frac{1}{2L_\theta} \left\| \int \nabla_\theta \ell(\theta_k; \tilde{x}) q_k(\mathrm{d}\tilde{x}) \right\|^2 \right) q_k(\mathrm{d}x) \\ &= \frac{1}{2L_\theta} \left\| \int \nabla_\theta \ell(\theta_k; x) q_k(\mathrm{d}x) \right\|^2, \end{aligned} \quad (17)$$

(an inequality known as the *descent lemma* in optimization). Now, (15) follows by combining this estimate with (16) while also noting that we have

$$q_k \in \arg \min_{q \in \mathcal{P}(\mathbb{R}^{d_x})} F(\theta_k, q) = \pi_{\theta_k} \Rightarrow I(\theta_k, q_k) = \left\| \int \nabla_\theta \ell(\theta_k; x) q_k(\mathrm{d}x) \right\|^2. \quad (18)$$

□

Now we just need to know that, along EM iterations,  $F$ 's gradient grows at least quadratically away from the set of minimizers, but that is precisely what the xLSI says.

**Proposition 8.** *Let Assumption 2 hold, and assume that the measures  $(\rho_\theta(\mathrm{d}x))_{\theta \in \mathbb{R}^{d_\theta}}$  satisfy the xLSI with constant  $\lambda > 0$ . Then, for all  $k \in \mathbb{N}$ ,*

$$F(\theta_k, q_k) - F_\star \leq (1 - \lambda/L_\theta)^k [F(\theta_0, q_0) - F_\star] \leq e^{-k\lambda/L_\theta} [F(\theta_0, q_0) - F_\star].$$

*Proof.* By Lemma 7 and then the xLSI,

$$F(\theta_k, q_k) - F(\theta_{k+1}, q_{k+1}) \geq F(\theta_k, q_k) - F(\theta_{k+1}, q_k) \geq \frac{1}{2L_\theta} I(\theta_k, q_k) \geq \frac{\lambda}{L_\theta} [F(\theta_k, q_k) - F_\star]$$

for all  $k \in \mathbb{N}$ , and the claim follows upon rearranging this inequality as

$$F(\theta_{k+1}, q_{k+1}) - F_\star \leq (1 - \lambda/L_\theta) [F(\theta_k, q_k) - F_\star]$$

and iterating. □

We can use the extension of Talagrand’s inequality provided by Theorem 3 to conclude convergence in  $\mathbf{d}$ -distance of the EM iterates.

**Corollary 9.** *Under Assumption 1 and the conditions of Proposition 8,*

$$\lambda \mathbf{d}((\theta_k, q_k), \mathcal{M}_\star)^2 \leq 2(1 - \lambda/L_\theta)^k [F(\theta_0, q_0) - F_\star] \leq 2e^{-k\lambda/L_\theta} [F(\theta_0, q_0) - F_\star].$$

Having demonstrated that differential arguments provide an efficient way to obtain a convergence bound for both the parameter and the posterior estimates in EM to the MLE, before commenting on the significance and the interpretation of this result, we first explore how this bound can be sharpened.

To prove the above results, we considered only the decrease in free energy due the parameter updates. We can improve the bound by also considering the decreases due posterior updates. To do so, we would like to follow the same principles as the proofs above, by comparing EM’s updates to appropriate gradient steps. We need a notion of a gradient step for  $q \mapsto F(\theta_{k+1}, q) \propto \text{KL}(q|\pi_{\theta_{k+1}})$ . As is well-known in some areas of optimal transport and sampling, in the Wasserstein-2 geometry this coincides with a Langevin step, in the sense that following this gradient direction coincides with evolving the probability density according to the Langevin Fokker–Planck equation. Actually, from (11) we are already in a good position to prove (at least formally) this fact: when the parameter space is the trivial space  $\{\theta_{k+1}\}$  from (11) we have

$$\text{grad}_{\mathcal{P}_2(\mathbb{R}^{d_x})} \text{KL}(q|\pi_{\theta_{k+1}}) = \nabla_x \cdot \left( q(\nabla_x \log(q) - \nabla_x \log(\pi_{\theta_{k+1}})) \right)$$

which implies that the curve  $t \mapsto q_t$  defined by

$$\partial_t q_t = \text{grad}_{\mathcal{P}_2(\mathbb{R}^{d_x})} \text{KL}(q_t|\pi_{\theta_{k+1}}) = \nabla_x \cdot \left( q_t(\nabla_x \log(q_t) - \nabla_x \log(\pi_{\theta_{k+1}})) \right)$$

is the ‘Wasserstein’ gradient flow of  $q \mapsto \text{KL}(q|\pi_{\theta_{k+1}})$  in  $\mathcal{P}_2(\mathbb{R}^{d_x})$ —the steepest descent curve on the space of probability distributions that connects an initial distribution to the invariant  $\pi_{\theta_{k+1}}$ . This differential equation is the Fokker–Planck equation of the overdamped Langevin diffusion

$$dX_t = \nabla_x \log(\pi_{\theta_{k+1}}(X_t)) dt + \sqrt{2} dW_t = \nabla_x \ell(\theta_{k+1}; X_t) dt + \sqrt{2} dW_t$$

where  $W_t$  denotes a Brownian Motion. Hence, as an analogue of (16), we wish to bound

$$F(\theta_{k+1}, q_k) - F(\theta_{k+1}, q_{k+1}) \geq F(\theta_{k+1}, q_k) - F(\theta_{k+1}, \text{Law}(X_k + h\nabla_x \ell(\theta_{k+1}; X_k) + \sqrt{2h}\xi_k)) \quad (19)$$

from below, where  $\text{Law}(X_k) = q_k$ ,  $h > 0$  is to be chosen based upon  $L_x$  and  $\xi_k$  is a standard normal random variable. We would like to obtain a lower bound on the right hand side quantity that depends on  $L_x$  and the extended Fisher information functional, giving something that looks like (15). The only difficulty is that there is not an immediate candidate for an analogue of the descent lemma (inequality (17)) on  $\mathcal{P}_2(\mathbb{R}^{d_x})$ . However, we have the following:

**Lemma 10** (Descent lemma on  $\mathcal{P}_2(\mathbb{R}^{d_x})$ ). *Let Assumption 2 hold. Let  $(p_t)$  be an interpolation in  $\mathcal{P}_2(\mathbb{R}^{d_x})$  between  $q_k$  at  $t = kh$  and  $\text{Law}(X_k + h\nabla_x \ell(\theta_{k+1}; X_k) + \sqrt{2h}\xi_k)$ , where  $\text{Law}(X_k) = q_k$ , at  $t = (k+1)h$ , defined by the law of*

$$Z_{t-} = X_k \quad dZ_t = \nabla_x \ell(\theta_{k+1}; Z_{t-}) dt + \sqrt{2} dW_t,$$

where  $t \in [kh, (k+1)h]$  and  $t_- = kh$ . If  $h \leq 1/4L_x$ ,

$$\partial_t \text{KL}(p_t | \pi_{\theta_{k+1}}) \leq -\frac{1}{2}I(p_t | \pi_{\theta_{k+1}}) + 6L_x^2 d_x(t - t_-) \quad (20)$$

This result has been used to study Langevin Monte Carlo and was established in [54] and refined in [18, Section 4.2]. In Section 3.4 we present a generalization of this result, whose proof also illustrates how the above lemma can be established. The term  $6L_x^2 d_x(t - t_-)$  is a bias term which has no analogue in the Euclidean case, and it is essentially due to the fact that the naïve discretization of the Langevin diffusion does not preserve the stationary distribution of the continuous time process.

We need one last ingredient: in the proof of Proposition 8 we used the fact that  $q_k$  minimizes  $F(\theta_{k+1}, \cdot)$  to obtain (18). Now, we can use the fact that  $\theta_{k+1}$  minimizes  $F(\cdot, q_k)$ , to obtain

$$\theta_{k+1} \in \arg \min_{\theta \in \mathbb{R}^{d_\theta}} F(\theta, q_k) \Rightarrow \int \nabla_\theta \ell(\theta_{k+1}, x) q_k(dx) = 0 \Rightarrow I(\theta_{k+1}, q_k) = I(q_k | \pi_{\theta_{k+1}}). \quad (21)$$

However, since we will work with the interpolation as in Lemma 16 we also need to know that  $\int \nabla_\theta \ell(\theta_{k+1}, x) p_t(dx)$  is small and establishing that is the focus of the next lemma.

**Lemma 11.** *Let Assumption 2 hold and let  $(p_t)$  be as in Lemma 10. For  $h \leq 1/4L_x$ ,*

$$\left\| \int \nabla_\theta \ell(\theta_{k+1}, x) p_t(dx) \right\|^2 \leq L_x(t - t_-)[C + 4d_x L_x], \quad t \in [kh, (k+1)h]$$

with  $C := L^2 \sup_k \int \|x - x_\dagger\|^2 q_k(dx) + \|\theta_{k+1} - \theta_\dagger\|^2$ , and where  $(\theta_\dagger, x_\dagger)$  is a stationary point of  $\ell$ . In particular, when  $q_k$  is given by the EM update,  $q_k = \arg \min_{q \in \mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta_{k+1}, q)$ , we have the bound

$$C \leq L^2 \sup_{\theta_\star \in \mathcal{O}_\star} \left\{ \int \|x - x_\dagger\|^2 \pi_{\theta_\star}(dx) + \|\theta_\star - \theta_\dagger\|^2 \right\} + \frac{2L^2}{\lambda} [F(\theta_0, q_0) - F_\star]. \quad (22)$$

*Proof.* See Appendix C.1. □

(22) shows that  $C$  is finite under Assumption 1. It is common to shift coordinates so that  $(\theta_\dagger, x_\dagger) = (0, 0)$ . We now improve on Proposition 8.

**Theorem 12.** *Let Assumptions 1–2 hold, and assume that the measures  $(\rho_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  satisfy the xLSI with constant  $\lambda > 0$ . Then, for all  $k \in \mathbb{N}$ ,*

$$F(\theta_k, q_k) - F_\star \leq \inf_{h \leq 1/4L_x} \left\{ e^{-k\lambda(h+1/L_\theta)} [F(\theta_0, q_0) - F_\star] + \frac{h^2 B}{1 - e^{-\lambda(h+1/L_\theta)}} \right\} \quad (23)$$

where  $B := [8L_x^2 d_x + CL_x/2]$  and  $C$  is as per Lemma 11.

*Proof.* Because  $\partial_t F(\theta_{k+1}, p_t) = \partial_t \text{KL}(p_t | \pi_{\theta_{k+1}})$ , we can write, by equation (20), the xLSI and Lemma 11, that for all  $h \leq 1/4L_x$ ,  $t \in [kh, (k+1)h]$ , with  $t_- = kh$ :

$$\begin{aligned} \partial_t F(\theta_{k+1}, p_t) &\leq -\frac{1}{2}I(\theta_{k+1}, p_t) + \frac{1}{2} \left\| \int \nabla_\theta \ell(\theta_{k+1}, x) p_t(dx) \right\|^2 + 6L_x^2 d_x(t - t_-) \\ &\leq -\lambda[F(\theta_{k+1}, p_t) - F_\star] + (t - t_-)B. \end{aligned} \quad (24)$$

Hence,

$$\partial_t \left[ e^{t\lambda} [F(\theta_{k+1}, p_t) - F_\star] \right] = e^{t\lambda} \left[ \lambda [F(\theta_{k+1}, p_t) - F_\star] + \partial_t F(\theta_{k+1}, p_t) \right] \leq e^{t\lambda} hB.$$

Integrating between  $t = kh$  and  $t = (k+1)h$  we obtain

$$e^{(k+1)h\lambda} [F(\theta_{k+1}, p_{(k+1)h}) - F_\star] - e^{kh\lambda} [F(\theta_{k+1}, p_{kh}) - F_\star] \leq e^{(k+1)h\lambda} h^2 B,$$

and thus:

$$F(\theta_{k+1}, p_{(k+1)h}) - F_\star \leq e^{-h\lambda} [F(\theta_{k+1}, q_k) - F_\star] + h^2 B. \quad (25)$$

Combining the left hand side with the observation in (19) and the right hand side with the bound in the proof of Proposition 8,

$$F(\theta_{k+1}, q_{k+1}) - F_\star \leq e^{-\lambda(h+1/L_\theta)} [F(\theta_k, q_k) - F_\star] + h^2 B.$$

Iterating this inequality we finally prove

$$F(\theta_k, q_k) - F_\star \leq e^{-k\lambda(h+1/L_\theta)} [F(\theta_0, q_0) - F_\star] + h^2 B \sum_{j=0}^{k-1} e^{-j\lambda(h+1/L_\theta)}$$

and we conclude by bounding the geometric sum above with its limit, and taking the infimum across  $h \leq 1/4L_x$ .  $\square$

As before, convergence in  $\mathbf{d}$ -distance of the both the sequence of parameters and the posterior now follows by the extension of the Talagrand inequality via Theorem 3.

**Corollary 13.** *Under the same conditions as Theorem 12,*

$$\lambda \mathbf{d}((\theta_k, q_k), \mathcal{M}_\star)^2 \leq 2 \inf_{h \leq 1/4L_x} \left\{ e^{-k\lambda(h+1/L_\theta)} [F(\theta_0, q_0) - F_\star] + \frac{h^2 B}{1 - e^{-\lambda(h+1/L_\theta)}} \right\}.$$

Since  $\mathcal{M}_\star = \{(\theta_\star, \pi_{\theta_\star}) : \theta_\star \in \mathcal{O}_\star\}$ , we are bounding the distance of the EM iterates to a (local) maxima of the marginal likelihood and the corresponding posterior, which can be identified as the projection of  $(\theta_k, q_k)$  onto the optimal set  $\mathcal{M}_\star$ . By definition, the log-Sobolev constant  $\lambda$ , which dictates the convergence rate of EM, has the interpretation of being bounded by the maximum ratio between information and the free energy produced along the EM iterates, as given by the functionals  $I$  and  $F$ .

By the results in Section 2.3, we know, for instance, that the same bound holds if we replace the xLSI assumption with  $\ell$ 's strong concavity (Assumption 3), or any bounded perturbation, in the sense of Proposition 5, of such model. When Assumption 3 holds,  $\ell$  has a unique maximizer, as does the marginal likelihood  $Z_\theta$ , and  $\mathcal{M}_\star$  is a singleton. At least without additional assumptions, we are required to start the algorithm with an initial distribution  $q_0$  having a density, otherwise the term  $F(\theta_0, q_0)$  would be infinite. In practice, this is never an issue, and we can avoid this problem altogether by performing an additional E-step as part of the initialization procedure, so given  $\theta_0$  we set  $q_0 = \pi_{\theta_0}$  which is, in any case, the optimal choice of  $q_0$  for a given  $\theta_0$  in terms of the resulting error bound, since  $F(\theta_0, \pi_{\theta_0}) = \inf_{q_0 \in \mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta_0, q_0)$ .

The bound above is reminiscent of the convergence bounds in Langevin Monte Carlo (LMC) when  $h$  is its discretization step size, which we need to be small enough, and that

appears both in a bias term and in the convergence rate, showing that the LMC algorithm is biased [23]. However, the rightmost term in (23) is *not* a bias term in the sense that the EM algorithm converges to the correct limit—we can always just take  $h = 0$  and eliminate this term, while retaining a non-zero convergence rate. Of course, small choices of  $h$  reduce the rightmost term, but they also make the rate of convergence smaller. When  $k$  is small, larger choices of  $h$  provide a much sharper bound than  $h = 0$ . In principle one could obtain still sharper bounds by considering different values of  $h$  for each step of the algorithm but we do not pursue that.

We now show that we can use similar techniques to analyze alternatives to the EM algorithm that we consider when either, or both, the E-M steps (2)–(3) are intractable.

### 3.2 First-order EM

Often, the M-step (2) is intractable. In these cases, rather than solving (2) exactly, one often performs instead a gradient step to find a  $\theta$  with a smaller, but not optimal, value of  $F(\theta, q_k) \propto -\int \ell(\theta; x)q_k(dx)$ . This results in the first-order EM (or gradient EM) algorithm (see, e.g. [7]).

---

#### Algorithm 2 First-order EM Algorithm

---

1: **Inputs:** Step size  $h$ , initial parameters  $(\theta_0, q_0)$ .

2: **for**  $k \geq 0$  **do**

3:   Update the parameter estimate

$$\theta_{k+1} = \theta_k + h \int \nabla_{\theta} \ell(\theta_k; x) q_k(dx).$$

4:   Update the posterior estimate

$$q_{k+1} = \pi_{\theta_{k+1}}.$$

5: **end for**

---

We can easily use the tools developed above to study first-order EM. Indeed, provided  $h \leq 1/L_{\theta}$ , we can verbatim adapt the proofs of Lemma 7, Proposition 8 and Corollary 9 with  $h$  in place of  $1/L_{\theta}$ , yielding:

**Theorem 14.** *Let Assumption 1–2 hold, and assume that the measures  $(\rho_{\theta}(dx))_{\theta \in \mathbb{R}^{d_{\theta}}}$  satisfy the  $x$ LSI with constant  $\lambda > 0$ . Then, if  $h \leq 1/L_{\theta}$ , for all  $k \in \mathbb{N}$ ,*

$$\lambda d((\theta_k, q_k), \mathcal{M}_{\star})^2 \leq 2e^{-k\lambda h} [F(\theta_0, q_0) - F_{\star}]$$

Because the parameter update here is not a minimization step anymore, we cannot leverage the posterior updates to improve the convergence bound in the same way as for the basic EM algorithm. Therefore, we can ensure first-order EM convergence with a small enough choice of the step size, but the error bounds for first-order EM algorithm will in general decrease more slowly than those for EM, which of course is consistent with the way in which we would expect the actual error to behave. Using similar techniques, it should be straightforward to analyze the case in which  $h$  is specified adaptively.

### 3.3 Langevin EM

In a complementary setting, the E-step (3) might be intractable. Hypothetically, in these cases, we could consider following the same strategy as the first-order EM algorithm, al-

though it is perhaps less obvious how to implement such a scheme: rather than performing (3), which is minimizing the free energy for a fixed  $\theta_{k+1}$  (Proposition 1), one can take a gradient step in the space of probability measures to find a new  $q$  which reduces  $F(\theta_{k+1}, q) \propto \text{KL}(q || \pi_{\theta_{k+1}})$ . For a step size  $h > 0$ , as we argued in Section 3.1, in the Wasserstein-2 geometry this step consists in  $q_k \rightarrow \text{Law}(X_{k+1})$ , where

$$X_{k+1} = X_k + h\nabla_x \ell(\theta_{k+1}; X_k) + \sqrt{2h}\xi_k, \quad \text{Law}(X_k) = q_k, \quad (26)$$

and where  $\xi_k$  is a standard normal random variable. We term the resulting algorithm, which appears in Appendix D of [35] as the gradient flow approximated by “marginal particle gradient descent”, Langevin EM. Although this algorithm may appear somewhat contrived, we view it as an idealisation of a form of Monte Carlo EM in which  $q_k$  is approximated with a particle system and the particles within the approximation are updated by an application of an unadjusted Langevin kernel of step-size  $h$  and invariant distribution  $\pi_{\theta_k}$  at each step. The use of (unadjusted) Langevin steps with EM type algorithms has been explored in a number of contexts, e.g. [28], and we anticipate that the analysis of Langevin EM can also be adapted to study these implementations.

---

**Algorithm 3** Langevin EM Algorithm

---

- 1: **Inputs:** Step size  $h$ , initial parameters  $(\theta_0, q_0 =: \text{Law}(X_0))$ .
- 2: **for**  $k \geq 0$  **do**
- 3:   Update the parameter estimate

$$\theta_{k+1} = \arg \max_{\theta} \int \ell(\theta_k; x) q_k(\mathrm{d}x)$$

- 4:   Update the posterior estimate, with  $\xi_k$  denoting a  $\mathcal{N}(0, I_{d_x})$  r.v.,

$$q_{k+1} = \text{Law}(X_{k+1}), \quad \text{where}$$

$$X_{k+1} = X_k + h\nabla_x \ell(\theta_{k+1}; X_k) + \sqrt{2h}\xi_k$$

- 5: **end for**
- 

We can study Langevin EM almost immediately with our results: we follow verbatim the proof of Theorem 12 until (25). Then, rather than combining the bound with the one in Proposition 8, which we cannot use as the posterior update is not a minimization step here, we use the looser bound  $F(\theta_{k+1}, q_k) \leq F(\theta_k, q_k)$ , iterate the resulting inequality and then combine it with Theorem 3 as usual.

**Theorem 15.** *Let Assumptions 1–2 hold, and assume that the measures  $(\rho_{\theta}(\mathrm{d}x))_{\theta \in \mathbb{R}^{d_{\theta}}}$  satisfy the xLSI with constant  $\lambda > 0$ . Then, if  $h \leq 1/4L_x$ , for all  $k \in \mathbb{N}$ ,*

$$\lambda d((\theta_k, q_k), \mathcal{M}_{\star})^2 \leq 2e^{-k\lambda h} [F(\theta_0, q_0) - F_{\star}] + \frac{2h^2 B}{1 - e^{-\lambda(h+1/L_{\theta})}}.$$

$B$  is as defined as in Theorem 12. In contrast to the EM case, we have not shown that  $B$  is always finite here. However, doing so is not typically onerous and is, for example, immediately true if the log-likelihood is strongly concave in the tails uniformly on  $\theta$ , as can be shown by adapting e.g. the arguments of Corollary 2.3 in [16].

We notice that, as opposed to EM or first-order EM, Langevin EM is actually biased: there is no way to cancel the rightmost term without getting a null rate of convergence. However, we can control the bias by choosing a small step size  $h$ . This is because the discretization of the Langevin diffusion (26) introduces a bias, and it is consistent with known results for Langevin Monte Carlo [23].



### 3.4 (Alternating) Gradient Descent

When both E- and M- steps are intractable, we can take a gradient step in both directions at each iteration; essentially replacing the usual alternating minimization optimization of the free energy with an alternating gradient descent method, as one might in Euclidean optimization problems for which both the coordinate-wise minimization problems are not tractable. This results in the Alternating Gradient Descent Algorithm 4 below, a mean-field limit version of the SOUL algorithm [21] when the batch sizes are set to one, and when the parameter updates can be carried out explicitly.

---

#### Algorithm 4 Alternating Gradient Descent Algorithm

---

1: **Inputs:** Step size  $h$ , initial parameters  $(\theta_0, q_0 =: \text{Law}(X_0))$ .

2: **for**  $k \geq 0$  **do**

3: Update the parameter estimate

$$\theta_{k+1} = \theta_k + h \int \nabla_{\theta} \ell(\theta_k; x) q_k(dx), \quad (27)$$

4: Update the posterior estimate: with  $\xi_k$  denoting a  $\mathcal{N}(0, I_{d_x})$  r.v.,

$$\begin{aligned} q_{k+1} &= \text{Law}(X_{k+1}), \quad \text{where} \\ X_{k+1} &= X_k + h \nabla_x \ell(\theta_{k+1}; X_k) + \sqrt{2h} \xi_k \end{aligned} \quad (28)$$

5: **end for**

---

To study Algorithm 4, we need a generalization of the descent lemma on the whole  $\mathcal{M}_2$

**Lemma 16** (Descent lemma on  $\mathcal{M}_2$ ). *Let Assumptions 1–2 hold. Let  $(\vartheta_t, p_t)$  be an interpolation in  $\mathcal{M}_2$  between  $(\theta_k, q_k)$  at  $t = kh$  and  $(\theta_k + h \int \nabla_{\theta} \ell(\theta_k; x) q_k(dx), \text{Law}(X_k + h \nabla_x \ell(\theta_{k+1}; X_k) + \sqrt{2h} \xi_k))$ , where  $\text{Law}(X_k) = q_k$ , at  $t = (k+1)h$ . If  $h \leq 1/4L$ , for  $t \in [kh, (k+1)h]$ ,*

$$\partial_t F(\theta_t, p_t) \leq -\frac{1}{2} I(\theta_t, p_t) + 6L^2 d_x (t - t_-), \quad t_- := kh.$$

*Proof.* See Appendix C.2. □

The interpolating  $(\vartheta_t, p_t)$  is specified in the proof. When the parameter space is trivial and equal to  $\{\theta_{k+1}\}$ , this result reduces in to Lemma 10.

**Theorem 17.** *Let Assumptions 1–2 hold, and assume that the measures  $(\rho_{\theta}(dx))_{\theta \in \mathbb{R}^{d_{\theta}}}$  satisfy the xLSI with constant  $\lambda > 0$ . Then, if  $h \leq 1/4L$ , for all  $k \in \mathbb{N}$ ,*

$$\lambda d((\theta_k, q_k), \mathcal{M}_{\star})^2 \leq 2e^{-k\lambda h} [F(\theta_0, q_0) - F_{\star}] + \frac{12L^2 d_x h^2}{1 - e^{-\lambda h}}$$

*Proof.* See Appendix C.3. □

It is also possible to update both parameters and distributions simultaneously (rather than in the alternating fashion described here), in which case we would obtain (a mean-field limit version of) the algorithm recently proposed in [35], studied and extended in [36, 13]. Such algorithm in fact corresponds to the (non alternating) gradient descent method on the free energy  $F$  in  $\mathcal{M}_2$ 's geometry.

A comparison with EM, first-order EM and Langevin EM results shows that the vanilla EM algorithm is the fastest of the four. While comparing error *bounds* cannot provide

definitive statements, this is consistent with the way one would expect the errors themselves to behave in general—because EM is solving two entire minimization problems at each iterations, whereas first-order EM and Langevin EM are solving one and taking a gradient step for the other and the alternative gradient descent approach is actually only taking a gradient step in both directions. Like Langevin EM, the alternating gradient descent approach is expectedly biased, and this is in line with previous analyses of similar algorithms [13, 1, 25, 44]. This hierarchy between alternating minimization and gradient descent type algorithms is consistent with the corresponding results and behaviour in Euclidean space; on this see [9] for a comparative convergence analysis and [51] for a qualitative comparison.

These results do not necessarily imply that implementable variants of EM algorithm outperforms the implementable form of any of these alternatives in practice, although it will certainly be the case when the minimization steps can be implemented exactly. Algorithms 1–4 are idealized algorithms, and for many models of interest, particularly in large-scale modern applications, the iterations therein are still intractable. In practice, one often resorts to Monte Carlo approximations. As noted above, for EM and first-order EM, it is common to substitute the E-steps with  $N$  Monte Carlo samples from  $\pi_{\theta_{k+1}}$  (the Monte Carlo EM algorithm [57]) whereas one would make Algorithms 3 and 4 implementable by approximating  $\text{Law}(X_{k+1})$  with the empirical distribution of  $N$  particles following the above Langevin dynamics. An analysis of these Monte Carlo versions of the algorithms under the xLSI and using the techniques we introduced here would potentially be interesting but introduces a number of further technical complications, and is left for future work.

## 4 Discussion

This paper established non-asymptotic error bounds and convergence rates for the EM algorithm—and some of its variants—under a log-Sobolev type inequality. Starting from the observation of [43] that EM corresponds to an alternating procedure on the product of Euclidean and the space of probability distributions, this approach can be considered a generalization of standard arguments used to study alternating minimization on Euclidean space via some concepts in optimal transport and sampling.

The log-Sobolev constant, which dictates the convergence rate in EM (Corollary 13), has the natural interpretation of being bounded by the maximum ratio between information and free energy produced along the EM iterates (as measured by the functionals  $I$  and  $F$ ). The convergence bounds seem to be state-of-the-art in terms of characterizing the exponential regime of EM, at least when the latent space is continuous (but see comments below). A complete comparison with the results of [33, 34] is hard to carry out due the very different approaches and settings: while they do not assume any smoothness, they characterize the *sub*-exponential convergence of EM parameter estimates [33] or posterior estimates [34] assuming (or implying) at least concavity of the surrogate.

The connections established herein mean there is potential to leverage the literatures of optimal transport, functional inequalities and Langevin Monte Carlo to better understand the EM algorithm and its many variants theoretically. Below we mention some potential connections and extensions, some of which form part of current or future research.

In Section 2.3 we presented a couple of generalizations of standard results (the Bakry–Émery criterion, the Holley–Stroock and the contraction principles) to verify the xLSI and compare the performance of models with different completions, and we expect that many other results of this flavour can be adapted from the LSI literature, further characteriz-

ing the exponential convergence regime of EM. Moreover, we expect that it is possible to consider ‘weak’ or ‘modified’ versions of the xLSI, as done in [53, 14, 50, 3, 42] for other standard functional inequalities, and similarly to consider gradient Hölder continuity assumptions in place of the assumed Lipschitz condition, to characterize the sub-exponential regime of convergence for a wide class of models.

The fact that the EM iterations are actually agnostic to any type of underlying metric on  $\mathcal{M}$  means that we could have considered different ones, and thus different induced log-Sobolev type inequalities; for instance, by considering the gradient of the free energy  $\text{grad}_{\mathcal{M}_2} F$  induced by the product of Euclidean and Stein’s geometry on  $\mathcal{P}_2(\mathbb{R}^{d_x})$ . These might translate into different practical conditions on the underlying model (as analogues of the results in Section 2.3) and could better characterize EM’s convergence in some settings, and it would be interesting to consider these.

Lastly, we mention that a limitation of our approach is that we focus on continuous Euclidean state spaces. While these constitute an important part of the models EM is applied to, in particular within the Empirical Bayes framework, many other typical applications, such as Gaussian Mixture Models, involve discrete latent spaces. We believe that it is possible to obtain analogues of our results in the discrete setting by leveraging advances on the study of Wasserstein gradients on discrete spaces [20]. Similarly, within a continuous state spaces setting, it should be possible to consider appropriate Riemannian manifolds rather than  $\mathbb{R}^{d_x}$  by adapting the relevant results in the literature [46].

## Acknowledgements

The authors thank Juan Kuntz for very helpful comments on an earlier version of this manuscript. RC was funded by the UK Engineering and Physical Sciences Research Council (EPSRC) via studentship 2585619 as part of grant number EP/W523793/1. AMJ acknowledges further EPSRC support under grant numbers EP/R034710/1 and EP/Y014650/1.

## References

- [1] Ö. Deniz Akyıldız, Francesca R. Crucinio, Mark Girolami, Tim Johnston, and Sotirios Sabanis. “Interacting particle Langevin algorithm for maximum marginal likelihood estimation”. In: *arXiv:2303.13429* (2023).
- [2] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. *Gradient Flows: In Metric Spaces and in the Space of Probability Measures*. Springer Science & Business Media, 2005.
- [3] Christophe Andrieu, Anthony Lee, Samuel Power, and Andi Q Wang. “Comparison of Markov chains via weak Poincaré inequalities with application to pseudo-marginal MCMC”. In: *The Annals of Statistics* 50.6 (2022), pp. 3592–3618.
- [4] Mihai Anitescu. “Degenerate nonlinear programming with a quadratic growth condition”. In: *SIAM Journal on Optimization* 10 (2000), pp. 1116–1135.
- [5] Dominique Bakry and Michel Émery. “Diffusions hypercontractives”. In: *Séminaire de Probabilités XIX 1983/84*. Lecture Notes in Mathematics Springer, 1985.
- [6] Dominique Bakry, Ivan Gentil, and Michel Ledoux. *Analysis and Geometry of Markov Diffusion Operators*. Springer, 2014.

- [7] Sivaraman Balakrishnan, Martin J. Wainwright, and Bin Yu. “Statistical guarantees for the EM algorithm: From population to sample-based analysis”. In: *The Annals of Statistics* 40 (2017), pp. 77–120.
- [8] Amir Beck. “On the convergence of alternating minimization for convex programming with applications to iteratively reweighted least squares and decomposition schemes”. In: *SIAM Journal on Optimization* 25.1 (2015), pp. 185–209.
- [9] Amir Beck and Luba Tretuashvili. “On the convergence of block coordinate descent type methods”. In: *SIAM Journal on Optimization* 23.4 (2013), pp. 2037–2060.
- [10] Jakub Wiktor Both. “On the rate of convergence of alternating minimization for non-smooth non-strongly convex optimization in Banach spaces”. In: *Optimization Letters* 16.2 (2022), pp. 729–743.
- [11] Nicolas Boumal. *An Introduction to Optimization on Smooth Manifolds*. Cambridge University Press, 2023.
- [12] Russell A Boyles. “On the convergence of the EM algorithm”. In: *Journal of the Royal Statistical Society Series B: Statistical Methodology* 45.1 (1983), pp. 47–50.
- [13] Rocco Caprio, Juan Kuntz, Samuel Power, and Adam M Johansen. “Error bounds for particle gradient descent, and extensions of the log-Sobolev and Talagrand inequalities”. In: *arXiv:2403.02004* (2024).
- [14] Patrick Cattiaux, Ivan Gentil, and Arnaud Guillin. “Weak logarithmic Sobolev inequalities and entropic convergence”. In: *Probability Theory and Related Fields* 139 (2007), pp. 563–603.
- [15] Patrick Cattiaux and Arnaud Guillin. “Functional inequalities for perturbed measures with applications to log-concave measures and to some Bayesian problems”. In: *Bernoulli* 28.4 (2022), pp. 2294–2321.
- [16] Patrick Cattiaux, Arnaud Guillin, and Florent Malrieu. “Probabilistic approach for granular media equations in the non-uniformly convex case”. In: *Probability Theory and Related Fields* 140 (2008), pp. 19–40.
- [17] Hong-Bin Chen, Sinho Chewi, and Jonathan Niles-Weed. “Dimension-free log-Sobolev inequalities for mixture distributions”. In: *Journal of Functional Analysis* 281.11 (2021), p. 109236.
- [18] Sinho Chewi. “Log-concave Sampling”. Book draft. 2024. URL: <https://chewisinho.github.io>.
- [19] Sinho Chewi, Murat A Erdogdu, Mufan Bill Li, Ruoqi Shen, and Matthew Zhang. “Analysis of Langevin Monte Carlo from Poincaré to Log-Sobolev”. In: *Proceedings of Thirty Fifth Conference on Learning Theory*. PMLR. 2022.
- [20] Shui-Nee Chow, Wuchen Li, and Haomin Zhou. “Entropy dissipation of Fokker-Planck equations on graphs”. In: *Discrete & Continuous Dynamical Systems* 38 (2018).
- [21] Valentin De Bortoli, Alain Durmus, Marcelo Pereyra, and Ana F Vidal. “Efficient stochastic optimisation by unadjusted Langevin Monte Carlo: Application to maximum marginal likelihood and empirical Bayesian estimation”. In: *Statistics and Computing* 31 (2021), pp. 1–18.
- [22] Arthur P. Dempster, Nan M. Laird, and Donald B. Rubin. “Maximum likelihood from incomplete data via the EM Algorithm”. In: *Journal of the Royal Statistical Society, Series B* 39 (1977), pp. 2–38.

- [23] Alain Durmus and Éric Moulines. “High-dimensional Bayesian inference via the unadjusted Langevin algorithm”. In: *Bernoulli* 25.4A (2019), pp. 2854–2882.
- [24] Raaz Dwivedi, Nhat Ho, Koulik Khamaru, Martin J Wainwright, Michael I Jordan, and Bin Yu. “Singularity, misspecification and the convergence rate of EM”. In: *The Annals of Statistics* 48.6 (2020), pp. 3161–3182.
- [25] Paula C. Encinar, Ö. Deniz Akyıldız, and Francesca R. Crucinio. “Proximal Interacting Particle Langevin Algorithms”. In: *arXiv:2406.14292* (2024).
- [26] Alessio Figalli and Federico Glaudo. *An Invitation to Optimal Transport, Wasserstein Distances, and Gradient Flows*. EMS Press, 2021.
- [27] Irving J Good. “On the estimation of small frequencies in contingency tables”. In: *Journal of the Royal Statistical Society: Series B (Methodological)* 18.1 (1956), pp. 113–124.
- [28] Samuel Gruffaz, Kyurae Kim, Alain Durmus, and Jacob Gardner. “Stochastic Approximation with Biased MCMC for Expectation Maximization”. In: *Proceedings of The 27th International Conference on Artificial Intelligence and Statistics*. Vol. 238. PMLR. 2024, pp. 2332–2340.
- [29] Botao Hao, Will Wei Sun, Yufeng Liu, and Guang Cheng. “Simultaneous clustering and estimation of heterogeneous graphical models”. In: *Journal of Machine Learning Research* 18.217 (2018), pp. 1–58.
- [30] Alfred O Hero and Jeffrey A Fessler. “Convergence in norm for alternating expectation-maximization (EM) type algorithms”. In: *Statistica Sinica* 5 (1995), pp. 41–54.
- [31] Richard Holley and Daniel W Stroock. “Logarithmic Sobolev inequalities and stochastic Ising models”. In: *Journal of Statistical Physics* 46 (1987), pp. 1159–1194.
- [32] Belhal Karimi, Hoi-To Wai, Eric Moulines, and Marc Lavielle. “On the Global Convergence of (Fast) Incremental Expectation Maximization Methods”. In: *Advances in Neural Information Processing Systems* 32 (2019).
- [33] Raunak Kumar and Mark Schmidt. “Convergence Rate of Expectation-Maximization”. In: *10th NeurIPS Workshop on Optimization for Machine Learning*. 2017.
- [34] Frederik Kunstner, Raunak Kumar, and Mark Schmidt. “Homeomorphic-invariance of EM: Non-Asymptotic Convergence in KL divergence for Exponential Families via Mirror Descent”. In: *Proceedings of The 24th International Conference on Artificial Intelligence and Statistics*. PMLR. 2021, pp. 3295–3303.
- [35] Juan Kuntz, Jen Ning Lim, and Adam M. Johansen. “Particle algorithms for maximum likelihood training of latent variable models”. In: *Proceedings of The 26th International Conference on Artificial Intelligence and Statistics*. Vol. 206. 2023, pp. 5134–5180.
- [36] Jen Ning Lim, Juan Kuntz, Samuel Power, and Adam M. Johansen. “Momentum particle maximum likelihood”. In: *Proceedings of 41st International Conference on Machine Learning (ICML)*. In press. PMLR. 2024.
- [37] Michel Loève. “Probability Concepts”. In: *Probability Theory I*. Springer New York, 1977, pp. 151–176.
- [38] Stanislaw Łojasiewicz. “Une propriété topologique des sous-ensembles analytiques réels”. In: *Les équations aux dérivées partielles* 117 (1963), pp. 87–89.

- [39] Geoffrey J. McLachlan and Thriyambakam Krishnan. *The EM Algorithm and Extensions*. John Wiley & Sons, 2007.
- [40] Xiao-Li Meng and Donald B. Rubin. “On the global and componentwise rates of convergence of the EM algorithm”. In: *Linear Algebra and its Applications* 199 (1994), pp. 413–425.
- [41] Xiao-Li Meng and David Van Dyk. “The EM algorithm—an old folk-song sung to a fast new tune”. In: *Journal of the Royal Statistical Society Series B: Statistical Methodology* 59.3 (1997), pp. 511–567.
- [42] Alireza Mousavi-Hosseini, Tyler Farghly, Ye He, Krishnakumar Balasubramanian, and Murat A Erdogdu. “Towards a complete analysis of Langevin Monte Carlo: Beyond Poincaré inequality”. In: *Proceedings of Thirty Sixth Conference on Learning Theory*. 2023, pp. 1–35.
- [43] Radford M. Neal and Geoffrey E. Hinton. “A View of the EM Algorithm that Justifies Incremental, Sparse, and other Variants”. In: *Learning in Graphical Models*. Springer Netherlands, 1998, pp. 355–368.
- [44] Paul Felix Valsecchi Oliva and Ö Deniz Akyildiz. “Kinetic Interacting Particle Langevin Monte Carlo”. In: *arXiv:2407.05790* (2024).
- [45] Felix Otto. “The Geometry of Dissipative Evolution Equations: the Porous Medium Equation”. In: *Communications in Partial Differential Equations* 26 (2001), pp. 101–174.
- [46] Felix Otto and Cédric Villani. “Generalization of an Inequality by Talagrand and Links with the Logarithmic Sobolev Inequality”. In: *Journal of Functional Analysis* 173 (2000), pp. 361–400.
- [47] Grigorios A. Pavliotis. *Diffusion Processes, the Fokker-Planck and Langevin Equations*. 1st. Texts in Applied Mathematics. New York: Springer, 2014.
- [48] Boris T. Polyak. “Gradient methods for the minimisation of functionals (in Russian)”. In: *Zhurnal Vychislitel’noi Matematiki i Matematicheskoi Fiziki* 3 (1963), pp. 643–653.
- [49] Herbert Robbins. “An empirical Bayes approach to statistics”. In: *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability*. Vol. 3.1. 1956, pp. 157–164.
- [50] Michael Röckner and Feng-Yu Wang. “Weak Poincaré inequalities and L2-convergence rates of Markov semigroups”. In: *Journal of Functional Analysis* 185.2 (2001), pp. 564–603.
- [51] James C. Spall. “Cyclic Seesaw Process for Optimization and Identification”. In: *Journal of Optimization Theory and Applications* 154 (2012), pp. 187–208.
- [52] Michel Talagrand. “Transportation cost for Gaussian and other product measures”. In: *Geometric & Functional Analysis* 6 (1996), pp. 587–600.
- [53] Guiseppe Toscani and Cédric Villani. “On the trend to equilibrium for some dissipative systems with slowly increasing a priori bounds”. In: *Journal of Statistical Physics* 98 (2000), pp. 1279–1309.
- [54] Santosh Vempala and Andre Wibisono. “Rapid Convergence of the Unadjusted Langevin Algorithm: Isoperimetry Suffices”. In: *Advances in Neural Information Processing Systems*. Vol. 32. 2019, pp. 8094–8106.

- [55] Cédric Villani. *Optimal Transport: Old and New*. Springer Science & Business Media, 2009.
- [56] Zhaoran Wang, Quanquan Gu, Yang Ning, and Han Liu. “High dimensional expectation-maximization algorithm: Statistical optimization and asymptotic normality”. In: *Advances in Neural Information Processing Systems* 28 (2015).
- [57] Greg C.G. Wei and Martin A. Tanner. “A Monte Carlo implementation of the EM algorithm and the poor man’s data augmentation algorithms”. In: *Journal of the American Statistical Association* 85.411 (1990), pp. 699–704.
- [58] Jeff CF Wu. “On the convergence properties of the EM algorithm”. In: *The Annals of Statistics* 11 (1983), pp. 95–103.
- [59] Ji Xu, Daniel J Hsu, and Arian Maleki. “Global analysis of expectation maximization for mixtures of two Gaussians”. In: *Advances in Neural Information Processing Systems* 29 (2016).

## A Notation

We collect together the most widely used notation here for convenience:

$\rho_\theta(x)$	Joint distribution $p_\theta(x, y)$ of latent variables $x$ and data $y$
$\ell(\theta; x)$	Logarithm of $\rho_\theta(x)$
$Z_\theta$	The marginal likelihood, $Z_\theta := \int \rho_\theta(x) dx$
$Z_\star$	Value of $Z_\theta$ at a stationary point
$\pi_\theta(x)$	Posterior of $x$ given $y$ , $\pi_\theta(x) = p_\theta(x y) = \rho_\theta(x)/Z_\theta$
$F(\theta, q)$	Free energy functional (5)
$I(\theta, q)$	Extended Fisher information functional (12)
$I(q  \pi_\theta)$	Relative Fisher information functional (13)
$\mathcal{O}_\star$	Local maxima of the marginal likelihood
$\mathcal{P}_2(\mathbb{R}^{d_x})$	Space of probability measures with densities and finite second moments
$\mathcal{M}_2$	Product spaces, $\mathcal{M}_2 := \mathbb{R}^{d_\theta} \times \mathcal{P}_2(\mathbb{R}^{d_x})$
$\mathcal{M}_\star$	$F$ ’s optimal set in $\mathcal{M}$ , $\mathcal{M}_\star = \arg \min F = \{(\theta_\star, \pi_{\theta_\star}) : \theta_\star \in \mathcal{O}_\star\}$
$\langle \cdot, \cdot \rangle$ and $\ \cdot\ $	Euclidean inner product and norm
$\langle \cdot, \cdot \rangle_{\mathcal{L}_2(dx)}$	$\mathcal{L}_2(\mathbb{R}^{d_x}, dx)$ ’s inner product
$\langle \cdot, \cdot \rangle_{W_2}$ , $\langle \cdot, \cdot \rangle_{\mathcal{M}_2}$	Wasserstein-2 and $\mathcal{M}_2$ inner products (Definitions 1 and 2)
$d_E, d_{W_2}$ and $d$	Euclidean distance, Wasserstein-2 distance, and their product (9)
$\nabla$ and $\nabla \cdot$	Gradient and divergence operators
$\Delta$	Laplacian operator, $\Delta := \nabla \cdot \nabla$

## B Proofs for Section 2

### B.1 Proof of Lemma 2

By comparing Definitions 3 and 4 we observe

$$\begin{aligned}
 \frac{d}{dt} \Big|_{t=0} F(\theta_t, q_t) &= \left\langle \delta_{\mathbb{R}^{d_\theta}} F(\theta, q), \frac{d}{dt} \Big|_{t=0} \theta_t \right\rangle + \left\langle \delta_{\mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta, q), \frac{d}{dt} \Big|_{t=0} q_t \right\rangle_{\mathcal{L}^2(dx)} \\
 &= \left\langle \text{grad}_{\mathbb{R}^{d_\theta}} F(\theta, q), \frac{d}{dt} \Big|_{t=0} \theta_t \right\rangle + \left\langle \text{grad}_{\mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta, q), \frac{d}{dt} \Big|_{t=0} q_t \right\rangle_{W_2}.
 \end{aligned}$$

Let  $\psi$  denote the solution of  $\nabla_x \cdot (q \nabla_x \psi) = -(dq_t/dt)|_{t=0}$ . The first terms on the RHS clearly coincide. We then have, upon equating the second terms on the RHS of the above display and inserting  $\psi$  so defined:

$$\begin{aligned} \left\langle \text{grad}_{\mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta, q), \frac{d}{dt} \Big|_{t=0} q_t \right\rangle_{W_2} &= - \int \delta_{\mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta, q)(x) \nabla_x \cdot (q(x) \nabla_x \psi(x)) \, dx \\ &= \int \left\langle \nabla_x \delta_{\mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta, q)(x), \nabla_x \psi(x) \right\rangle q(dx) \end{aligned}$$

where we integrated by parts. We conclude by noting that by Definition 1 we know that  $\text{grad}_{\mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta, q)$  has to satisfy  $-\nabla_x \cdot (q \nabla_x \delta_{\mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta, q)) = \text{grad}_{\mathcal{P}_2(\mathbb{R}^{d_x})} F(\theta, q)$ .  $\square$

## B.2 Proof of Proposition 5

Let  $\phi(r) := r \log(r)$  and for a non-negative function  $f$  on  $\mathbb{R}^{d_x} \times \mathbb{R}^{d_\theta}$  define the functionals

$$\begin{aligned} E_\theta(f) &:= \int \phi(f) \, d\pi_\theta - \phi \left( \int f \, d\pi_\theta \right), \\ J_\theta(f) &:= \int \frac{|\nabla_x f|^2}{f} \, d\pi_\theta, \end{aligned}$$

with  $\tilde{E}_\theta(f)$  and  $\tilde{J}_\theta(f)$  denoting the corresponding functionals when the model is  $\tilde{\rho}_\theta(dx) := \tilde{\pi}_\theta(dx) Z_\theta$ . Importantly, notice that if  $f := dq/d\pi_\theta$ ,  $E_\theta(f) - \log(Z_\theta) = F(\theta, q)$  and  $J_\theta(f) + \|\int \nabla_x \ell(\theta; x) q(dx)\|^2 = I(\theta, q)$ . We begin as in the proof of [6, Proposition 5.1.6] and use the following variational formula.

**Lemma 18.** *Letting  $\phi'(r)$  denote the derivative of  $\phi$  at  $r$ , we have,*

$$E_\theta(f) = \inf_{r>0} \int [\phi(f) - \phi(r) - \phi'(r)(f - r)] \, d\pi_\theta$$

*Proof.* Let  $X \sim \pi_\theta$ . Since  $\phi$  is convex, for any  $r > 0$ ,

$$\phi(\mathbb{E}[f(X)]) - \phi(r) - \phi'(r)(\mathbb{E}[f(X)] - r) \geq 0$$

so that, adding  $\mathbb{E}[\phi(f(X))] - \phi(\mathbb{E}[f(X)])$  both sides,

$$\mathbb{E}[\phi(f(X)) - \phi(r) - \phi'(r)(f(X) - r)] \geq \mathbb{E}[\phi(f(X))] - \phi(\mathbb{E}[f(X)]) = E_\theta(f)$$

with equality if and only if  $r = \mathbb{E}[f(X)]$ .  $\square$

By  $\phi$ 's convexity  $\phi(f) - \phi(r) - \phi'(r)(f - r) \geq 0$  for all  $r > 0$ , so using the formula above we can readily write

$$\tilde{E}_\theta(f) \leq c \inf_{r>0} \int [\phi(f) - \phi(r) - \phi'(r)(f - r)] \, d\pi_\theta = c E_\theta(f).$$

Next, we take  $f := dq/d\pi_\theta$ , which gives

$$\tilde{E}_\theta(f) + c(\log(Z_\star) - \log(Z_\theta)) \leq c[E_\theta(f) + \log(Z_\star) - \log(Z_\theta)] = c[F(\theta, q) - F_\star]$$



and because  $(\rho_\theta(dx))_{\theta \in \mathbb{R}^{d_\theta}}$  satisfy the xLSI

$$2\lambda[\tilde{E}_\theta(f) + c(\log(Z_\star) - \log(Z_\theta))] \leq cI(\theta, q) \leq c^2 \int \frac{|\nabla_x f|^2}{f} d\tilde{\pi}_\theta + c \left\| \int \nabla_\theta \ell(\theta; x) q(dx) \right\|^2.$$

Define now

$$\tilde{q}(dx) := \frac{1}{\tilde{c}} \frac{d\tilde{\pi}_\theta}{d\pi_\theta}(x) q(dx), \quad \tilde{c} := \int \frac{d\tilde{\pi}_\theta}{d\pi_\theta}(x) q(dx), \quad \tilde{f}(x) := \frac{f}{\tilde{c}}(x) = \frac{d\tilde{q}}{d\tilde{\pi}_\theta}(x),$$

noting that  $\tilde{q}$  is a probability measure in  $\mathcal{P}_2^1(\mathbb{R}^{d_x})$  under our assumptions. Since the functional  $E$  is homogeneous, i.e.  $\tilde{E}_\theta(af) = a\tilde{E}_\theta(f)$  for all  $a > 0$ , we may write

$$\begin{aligned} 2\lambda[\tilde{F}(\theta, p) - \tilde{F}_\star] &= 2\lambda[\tilde{E}_\theta(\tilde{f}) - \log(Z_\theta) + \log(Z_\star)] \\ &\leq 2\lambda[\tilde{E}_\theta(\tilde{f}) + \frac{c}{\tilde{c}}(\log(Z_\star) - \log(Z_\theta))] \\ &\leq \frac{c^2}{\tilde{c}} \int \frac{|\nabla_x f|^2}{f} d\tilde{\pi}_\theta + \frac{c}{\tilde{c}} \left\| \int \nabla_\theta \ell(\theta; x) q(dx) \right\|^2 \\ &\leq c^2 \left[ \tilde{J}_\theta(\tilde{f}) + \left\| \int \nabla_\theta \ell(\theta; x) \frac{d\rho_\theta}{d\tilde{\rho}_\theta}(x) \tilde{q}(dx) \right\|^2 \right] \\ &= c^2 \left[ \tilde{J}_\theta(\tilde{f}) + \left\| \int \frac{\nabla_\theta \rho_\theta(x)}{\tilde{\rho}_\theta(x)} \tilde{q}(dx) \right\|^2 \right] \end{aligned} \quad (29)$$

where in the first and penultimate inequalities we used  $\tilde{c} \leq c$  and  $\log(Z_\star) - \log(Z_\theta) \geq 0$ . By the inequality  $(a + b)^2 \leq 2a^2 + 2b^2$  ([37, p. 157]), Jensen's inequality and the assumed growth condition on  $\|\nabla_\theta \tilde{\rho}_\theta(x) - \nabla_\theta \rho_\theta(x)\|$ ,

$$\begin{aligned} \left\| \int \frac{\nabla_\theta \rho_\theta(x)}{\tilde{\rho}_\theta(x)} \tilde{q}(dx) \right\|^2 &\leq 2 \left\| \int \nabla_\theta \tilde{\ell}(\theta; x) \tilde{q}(dx) \right\|^2 + 2 \int \left\| \frac{\nabla_\theta \tilde{\rho}_\theta(x) - \nabla_\theta \rho_\theta(x)}{\tilde{\rho}_\theta(x)} \right\|^2 \tilde{q}(dx) \\ &\leq 2 \left\| \int \nabla_\theta \tilde{\ell}(\theta; x) \tilde{q}(dx) \right\|^2 + 2b[\log(Z_\star) - \log(Z_\theta)] \\ &\leq 2 \left\| \int \nabla_\theta \tilde{\ell}(\theta; x) \tilde{q}(dx) \right\|^2 + 2b[\tilde{F}(\theta, \tilde{q}) - \tilde{F}_\star] \end{aligned}$$

where in the last inequality we used  $\text{KL}(\tilde{q}|\tilde{\pi}_\theta) \geq 0$ . Combining this estimate with (29),

$$2(\lambda - c^2 \cdot b) [\tilde{F}(\theta, \tilde{q}) - \tilde{F}_\star] \leq 2c^2 \left[ \tilde{J}_\theta(\tilde{f}) + \left\| \int \nabla_\theta \tilde{\ell}(\theta; x) \tilde{q}(dx) \right\|^2 \right] = 2c^2 \tilde{I}(\theta, \tilde{q})$$

□

### B.3 Proof of Proposition 6

Denote with  $\tilde{F}$  and  $\tilde{I}$  the free energy and extended Fisher information for the model  $\tilde{\rho}_\theta$ . Consider an arbitrary  $(\theta, \tilde{q}) \in \mathcal{M}_2^1$  and set  $q := T_{\#}\tilde{q}$ . By assumption,  $\tilde{\pi}_\theta(T(x)) \det(\nabla_x T(x)) = \pi_\theta(x)$  and similarly for  $q$ 's density, hence, by a change of variables  $x \mapsto T(x)$ ,

$$\begin{aligned} \tilde{F}(\theta, \tilde{q}) - \log(Z_\theta) &= \int \log \left( \frac{\tilde{q}(x)}{\tilde{\pi}_\theta(x)} \right) \tilde{q}(x) dx = \int \log \left( \frac{\tilde{q}(T(x))}{\tilde{\pi}_\theta(T(x))} \right) \tilde{q}(T(x)) \det(\nabla_x T(x)) dx \\ &= \int \log \left( \frac{q(x)}{\pi_\theta(x)} \right) q(x) dx = F(\theta, q) - \log(Z_\theta) \end{aligned}$$

Thus, since  $q$  belongs to  $\mathcal{P}_2^1(\mathbb{R}^{d_x})$  by the Lipschitz property of  $T$ ,  $2\lambda[\tilde{F}(\theta, \tilde{q}) - F_\star] = 2\lambda[F(\theta, q) - F_\star] \leq I(\theta, q)$  by the xLSI. On the other hand, if we let  $\|\cdot\|_\infty^2$  denote the uniform norm,

$$\begin{aligned}
I(\theta, q) &= \int \left\| \frac{\nabla_x q(x)}{q(x)} - \frac{\nabla_x \pi_\theta(x)}{\pi_\theta(x)} \right\|^2 q(\mathrm{d}x) + \left\| \int \left( \frac{\nabla_\theta \pi_\theta(x)}{\pi_\theta(x)} + \frac{\nabla_\theta Z_\theta}{Z_\theta} \right) q(\mathrm{d}x) \right\|^2 \\
&= \int \left\| \frac{\nabla_x(\tilde{q}(T(x)))}{\tilde{q}(T(x))} - \frac{\nabla_x(\tilde{\pi}_\theta(T(x)))}{\tilde{\pi}_\theta(T(x))} \right\|^2 q(\mathrm{d}x) + \left\| \int \left( \frac{\nabla_\theta(\tilde{\pi}_\theta(T(x)))}{\tilde{\pi}_\theta(T(x))} + \frac{\nabla_\theta Z_\theta}{Z_\theta} \right) q(\mathrm{d}x) \right\|^2 \\
&\leq \|\nabla_x T\|_\infty^2 \int \left\| \frac{\nabla_{T(x)} \tilde{q}(T(x))}{\tilde{q}(T(x))} - \frac{\nabla_{T(x)} \tilde{\pi}_\theta(T(x))}{\tilde{\pi}_\theta(T(x))} \right\|^2 q(\mathrm{d}x) + \left\| \int \left( \frac{\nabla_\theta \tilde{\pi}_\theta(T(x))}{\tilde{\pi}_\theta(T(x))} + \frac{\nabla_\theta Z_\theta}{Z_\theta} \right) q(\mathrm{d}x) \right\|^2 \\
&\leq \|\nabla_x T\|_\infty^2 \tilde{I}(\tilde{q}|\tilde{\pi}_\theta) + \left\| \int \nabla_\theta \log(\tilde{\rho}(\theta, x)) \tilde{q}(x) \mathrm{d}x \right\|^2 \leq \max(1, \|\nabla_x T\|_\infty^2) \tilde{I}(\theta, \tilde{q})
\end{aligned}$$

where in the second line we substituted the expression for the density of  $\tilde{q}(T(x))$  and  $\tilde{\pi}_\theta(T(x))$  (noting that the terms with the determinant cancel), in the third we used the chain rule and the fact that  $T$  does not depend on  $\theta$ , and in the fourth we used the formula for  $q(x)$  and a change of variables  $T(x) \mapsto x$ . We conclude by noting that  $L_T$  provides a bound on  $\|\nabla_x T\|_\infty$ .  $\square$

## C Proofs for Section 3

### C.1 Proof of Lemma 11

Because of  $\theta_{k+1}$ 's minimality (21), by Jensen's inequality and Assumption 2,

$$\left\| \int \nabla_\theta \ell(\theta_{k+1}, x) p_t(\mathrm{d}x) \right\|^2 = \left\| \mathbb{E} [\nabla_\theta \ell(\theta_{k+1}, Z_t) - \nabla_\theta \ell(\theta_{k+1}, X_k)] \right\|^2 \leq L_x^2 \mathbb{E} [\|Z_t - X_k\|^2]$$

for any coupling of  $(Z_t, X_k)$ . Let  $(\theta_\dagger, x_\dagger)$  denote a stationary point of  $\ell$ , so that  $\nabla \ell(\theta_\dagger, x_\dagger) = 0$ . Inserting the expression for  $Z_t$ , since  $Z_{kh} = X_k$  a.s., using  $(a+b)^2 \leq a^2 + b^2$ , the fact  $h \leq 1/4L_x \Rightarrow 2L_x(t-t_-) \leq 1/2$  and again Assumption 2,

$$\begin{aligned}
\left\| \int \nabla_\theta \ell(\theta_{k+1}, x) p_t(\mathrm{d}x) \right\|^2 &\leq L_x^2 \mathbb{E} \left[ \mathbb{E} \left[ \left\| (t-t_-) \nabla_x \ell(\theta_{k+1}, X_k) + \sqrt{2}(W_t - W_{t_-}) \right\|^2 \middle| X_k \right] \right] \\
&\leq L_x(t-t_-) \left( 2L_x(t-t_-) \mathbb{E} [\|\nabla_x \ell(\theta_{k+1}, X_k)\|^2] + 4d_x L_x \right) \\
&\leq L_x(t-t_-) \left( \frac{1}{2} \mathbb{E} [\|\nabla_x \ell(\theta_{k+1}, X_k)\|^2] + 4d_x L_x \right) \\
&\leq L_x(t-t_-) \left( \frac{L^2}{2} \mathbb{E} [\|X_k - x_\dagger\|^2 + \|\theta_{k+1} - \theta_\dagger\|^2] + 4d_x L_x \right).
\end{aligned}$$

When  $q_k$  is given by an EM update, we now derive an uniform bound for  $\mathbb{E}[\|X_k - x_\dagger\|^2 + \|\theta_{k+1} - \theta_\dagger\|^2]$ . Consider the point  $(\theta_\star, \pi_{\theta_\star}) \in \mathcal{M}_\star$  corresponding to the projection of  $(\theta_{k+1}, q_k)$  onto  $\mathcal{M}_\star$ , let  $X_\star \sim \pi_{\theta_\star}$  and consider an optimal coupling between  $X_\star$  and  $X_k \sim q_k$ .

By the definition of  $d_{W_2}$ , Theorem 3, together with the fact that the free energy can only decrease along EM iterations,

$$\lambda \mathbb{E} \left[ \|X_k - X_\star\|^2 + \|\theta_{k+1} - \theta_\star\|^2 \right] \leq 2[F(\theta_{k+1}, q_k) - F_\star] \leq 2[F(\theta_0, q_0) - F_\star].$$

We conclude upon combining the two previously obtained inequalities via the bounds  $\|X_k - x_\dagger\|^2 \leq 2\|X_\star - x_\dagger\|^2 + 2\|X_k - X_\star\|^2$ , and the analogous bound for  $\|\theta_{k+1} - \theta_\dagger\|^2$ .  $\square$

## C.2 Proof of Lemma 16

The following proof is a suitable generalization of the one of [54] and in particular of the refinement in [18, Section 4.2] under the hypothesis of non-trivial parameter space, and the proof is therefore somewhat similar. Consider the continuous time process given for all  $t \in [kh, (k+1)h)$  and any  $k \in \mathbb{N}_0$  by

$$\begin{aligned} \vartheta_{kh} &= \theta_k & d\vartheta_t &= \int \nabla_{\theta} \ell(\vartheta_{t_-}; x) p_{t_-}(dx) dt, \\ Z_{kh} &= X_k & dZ_t &= \nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) dt + \sqrt{2} dW_t, \end{aligned} \tag{30}$$

where we set  $t_- := kh, p_t := \text{Law}(Z_t)$  and where  $W_t$  is a Brownian Motion. Notice that this is an interpolation of the Algorithm 4 iterates. To prove Lemma 16 we need three auxiliary results.

**Lemma 19.** *If Assumptions 1–2 hold,  $\{(\vartheta_t, p_t) : t_- \leq t < t_- + h\}$  is a solution of the PDE*

$$\begin{aligned} \partial_t \vartheta_t &= \int \nabla_{\theta} \ell(\vartheta_t; x) p_t(dx) + \mathbb{E}[\nabla_{\theta} \ell(\vartheta_{t_-}; Z_{t_-}) - \nabla_{\theta} \ell(\vartheta_t; Z_t)] \\ \partial_t p_t &= \nabla_x \cdot \left[ p_t \nabla_x \log \left( \frac{p_t}{\rho_{\vartheta_t}} \right) + p_t \mathbb{E}[\nabla_x \ell(\vartheta_t; \cdot) - \nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) | Z_t = \cdot] \right]. \end{aligned} \tag{31}$$

*Proof.* Conditionally on  $(\vartheta_{t_-}, Z_{t_-}) = (\vartheta, z)$ ,  $\{(\vartheta_t, Z_t) : t_- \leq t < t_- + h\}$  has a time independent drift, hence  $p_{t|t_-}(\cdot|z)$ , the conditional density of  $Z_t$  given  $Z_{t_-} = z$ , is the density of a Normal with mean  $z + (t - t_-)\nabla_x \ell(\vartheta; z)$  and variance  $2(t - t_-)$ . Hence, we can directly check that  $p_{t|t_-}$  solves the following PDE

$$\begin{aligned} \partial_t \vartheta_t &= \int \nabla_{\theta} \ell(\vartheta_{t_-}; x) p_{t_-}(dx), \\ \partial_t p_{t|t_-} &= \nabla_x \cdot \left[ p_{t|t_-} \nabla_x \log \left( \frac{p_{t|t_-}}{\rho_{\vartheta_{t_-}}} \right) \right] = \Delta_x p_{t|t_-} - \nabla_x \cdot (p_{t|t_-} \nabla_x \ell(\vartheta_{t_-}; Z_{t_-})). \end{aligned}$$

One can also derive the above immediately by using formulæ for the Fokker–Planck equation of Itô diffusions, see [47, Chapters 2 and 4]. We notice that all the derivatives of  $x \mapsto p_{t|t_-}(x|z)$  exist and are continuous in all arguments by Assumption 2, that  $z \mapsto p_{t|t_-}(x|z)$  is at least continuous for all  $x \in \mathbb{R}^{d_x}$ , and that  $t \mapsto p_{t|t_-}(x|z)$  is continuously differentiable for all  $x, z \in \mathbb{R}^{d_x}$ . By Bayes' rule, if we take expectation w.r.t.  $p_{t_-}$ ,

$$\begin{aligned} \mathbb{E} \left[ p_{t|t_-}(x|Z_{t_-}) \nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) \right] &= \int p_{t|t_-}(x|z) \nabla_x \ell(\vartheta_{t_-}; z) p_{t_-}(dz) \\ &= p_t(x) \int p_{t_-|t}(dz|x) \nabla_x \ell(\vartheta_{t_-}; z) = p_t(x) \mathbb{E} \left[ \nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) | Z_t = x \right] \end{aligned}$$

Hence, taking expectation in the Fokker–Planck equation, the regularity properties of  $p_{t|t_-}$  allow us to use the tower rule and compute

$$\partial_t \vartheta_t = \int \nabla_{\theta} \ell(\vartheta_{t_-}; x) p_{t_-}(dx), \quad \partial_t p_t = \Delta_x p_t - \nabla_x \cdot (p_t \mathbb{E}[\nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) | Z_t = \cdot]).$$

Adding and subtracting  $\mathbb{E}[\nabla_{\theta} \ell(\vartheta_t; Z_t)]$  in the first expression and  $\nabla_x \cdot (p_t \mathbb{E}[\nabla_x \ell(\vartheta_t; \cdot) | Z_t = \cdot]) = \nabla_x \cdot (p_t \nabla_x \log(\rho_{\vartheta_t}))$  in the second yields the claim.  $\square$

**Lemma 20.** *If Assumptions 1–2 hold, for all  $t \in [t_-, t_- + h)$ ,*

$$\partial_t F(\vartheta_t, p_t) \leq -\frac{3}{4} I(\vartheta_t, p_t) + \mathbb{E} \left[ \left\| \nabla \ell(\vartheta_t; Z_t) - \nabla \ell(\vartheta_{t_-}; Z_{t_-}) \right\|^2 \right] \quad (32)$$

*Proof.* By the regularity properties of  $p_{t|t_-}$  illustrated in the proof of the previous Lemma,  $p_t(x) = \mathbb{E}[p_{t|t_-}(x, Z_{t_-})]$  is smooth in  $x$  and at least continuously differentiable in  $t$ . This allows us to compute

$$\begin{aligned} \partial_t F(\vartheta_t, p_t) &= \int \left[ \log \left( \frac{p_t(x)}{\rho_{\vartheta_t}(x)} \right) + 1 \right] \nabla_x \cdot \left[ p_t(x) \nabla_x \log \left( \frac{p_t(x)}{\rho_{\vartheta_t}(x)} \right) \right] dx - \left\| \int \nabla_{\theta} \ell(\vartheta_t; x) p_t(dx) \right\|^2 \\ &\quad + \int \left[ \log \left( \frac{p_t(x)}{\rho_{\vartheta_t}(x)} \right) + 1 \right] \nabla_x \cdot [p_t(x) \mathbb{E}[\nabla_x \ell(\vartheta_t, x) - \nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) | Z_t = x]] dx \\ &\quad + \mathbb{E}[\nabla_{\theta} \ell(\vartheta_t; Z_t) - \nabla_{\theta} \ell(\vartheta_{t_-}; Z_{t_-})] \mathbb{E}[\nabla_{\theta} \ell(\vartheta_t; Z_t)] \\ &= -I(\vartheta_t, p_t) + \int \nabla_x \log \left( \frac{p_t(x)}{\rho_{\vartheta_t}(x)} \right) \mathbb{E}[\nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) - \nabla_x \ell(\vartheta_t; x) | Z_t = x] p_t(dx) \\ &\quad + \mathbb{E}[\nabla_{\theta} \ell(\vartheta_t; Z_t) - \nabla_{\theta} \ell(\vartheta_{t_-}; Z_{t_-})] \mathbb{E}[\nabla_{\theta} \ell(\vartheta_t; Z_t)] \end{aligned} \quad (33)$$

where we used the regularity of  $\{(\vartheta_t, p_t) : t_- \leq t < t_- + h\}$  to differentiate inside the integral and the chain rule in the first equality, and we integrated by parts for the second. Now, by the Young’s inequality  $ab \leq (a^2/4) + b^2$  and then Jensen’s inequality, we have

$$\begin{aligned} &\int \nabla_x \log \left( \frac{p_t(x)}{\rho_{\vartheta_t}(x)} \right) \mathbb{E}[\nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) - \nabla_x \ell(\vartheta_t; x) | Z_t = x] p_t(dx) \\ &\leq \frac{1}{4} I(p_t | \pi_{\vartheta_t}) + \int \left\| \mathbb{E}[\nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) - \nabla_x \ell(\vartheta_t; x) | Z_t = x] \right\|^2 p_t(dx) \\ &\leq \frac{1}{4} I(p_t | \pi_{\vartheta_t}) + \mathbb{E} \left[ \left\| \nabla_x \ell(\vartheta_t; Z_t) - \nabla_x \ell(\vartheta_{t_-}; Z_{t_-}) \right\|^2 \right] \end{aligned}$$

and similarly

$$\begin{aligned} &\mathbb{E}[\nabla_{\theta} \ell(\vartheta_t; Z_t) - \nabla_{\theta} \ell(\vartheta_{t_-}; Z_{t_-})] \mathbb{E}[\nabla_{\theta} \ell(\vartheta_t; Z_t)] \\ &\leq \frac{1}{4} \left\| \int \nabla_{\theta} \ell(\vartheta_t; x) p_t(dx) \right\|^2 + \mathbb{E} \left[ \left\| \nabla_{\theta} \ell(\vartheta_t; Z_t) - \nabla_{\theta} \ell(\vartheta_{t_-}; Z_{t_-}) \right\|^2 \right] \end{aligned}$$

now by combining these last two estimates with (33) we prove the desired result.  $\square$

Our goal now is to obtain a bound on the rightmost term in (32) in terms of  $I$ .

**Lemma 21.** *If Assumption 2 holds, for any  $\theta \in \mathbb{R}^{d_{\theta}}$  and  $q \in \mathcal{P}_2(\mathbb{R}^{d_x})$  we have*

$$\left\| \mathbb{E}[\nabla_{\theta} \ell(\theta; X)] \right\|^2 + \mathbb{E} \left[ \left\| \nabla_x \ell(\theta; X) \right\|^2 \right] \leq I(\theta, q) + 2Ld_x, \quad \text{where } \text{Law}(X) = q.$$

*Proof.* This inequality follows almost immediately from [18, Lemma 4.2.5] or [19, Lemma 16] and the definition of  $I(\theta, q)$ . We reproduce those proofs here in our notation for convenience. Consider the (overdamped) Langevin diffusion with stationary distribution  $\pi_\theta \propto e^{\ell(\theta; \cdot)}$ . Its generator satisfies  $\mathcal{L}\ell(\theta, \cdot) = \Delta_x \ell(\theta; \cdot) + \|\nabla_x \ell(\theta; \cdot)\|^2$ . We can estimate directly

$$\begin{aligned} \mathbb{E} \left[ \|\nabla_x \ell(\theta; X)\|^2 \right] &= \mathbb{E} \left[ -\Delta_x \ell(\theta; X) + \mathcal{L}\ell(\theta; X) \right] \\ &\leq Ld_x + \int \mathcal{L}\ell(\theta; x) \frac{dq}{d\pi_\theta}(x) \pi_\theta(dx) \\ &= Ld_x + \int \|\nabla_x \ell(\theta; x)\|^2 \frac{dq}{d\pi_\theta}(x) \pi_\theta(x) + \Delta_x \ell(\theta; x) \frac{dq}{d\pi_\theta}(x) \pi_\theta(x) dx \\ &= Ld_x + \int \|\nabla_x \ell(\theta; x)\|^2 \frac{dq}{d\pi_\theta}(x) \pi_\theta(x) - \left\langle \nabla_x \ell(\theta; x), \nabla_x \left( \frac{dq}{d\pi_\theta}(x) \pi_\theta(x) \right) \right\rangle dx \end{aligned}$$

where in the second line we used  $-\Delta_x \ell(\theta, \cdot) \leq Ld_x$  by Assumption 2, in the fourth integration by parts. Now, with the product rule and the fact  $\nabla_x \ell(\theta; x) \pi_\theta(x) = \nabla_x \pi_\theta(x)$ ,

$$\begin{aligned} \mathbb{E} \left[ \|\nabla_x \ell(\theta; X)\|^2 \right] &= Ld_x - \int \left\langle \nabla_x \ell(\theta; x), \nabla_x \frac{dq}{d\pi_\theta}(x) \right\rangle \pi_\theta(dx) \\ &= Ld_x - 2 \int \left\langle \sqrt{\frac{dq}{d\pi_\theta}}(x) \nabla_x \ell(\theta; x), \nabla_x \sqrt{\frac{dq}{d\pi_\theta}}(x) \right\rangle \pi_\theta(dx) \\ &\leq Ld_x + \frac{1}{2} \mathbb{E} \left[ \|\nabla_x \ell(\theta; X)\|^2 \right] + 2 \int \left\| \nabla_x \sqrt{\frac{dq}{d\pi_\theta}}(x) \right\|^2 \pi_\theta(dx) \\ &= Ld_x + \frac{1}{2} \mathbb{E} \left[ \|\nabla_x \ell(\theta; X)\|^2 \right] + \frac{1}{2} I(q|\pi_\theta), \end{aligned}$$

where we used the chain rule, and the Young's inequality  $ab \leq (a^2/4) + b^2$  again. Now rearranging, adding  $\|\mathbb{E} [\nabla_\theta \ell(\theta; x)]\|^2$  to both sides and using the definition of  $I(\theta, q)$  proves the bound.  $\square$

Having established these intermediate results we can return to the now straightforward proof of the lemma of interest.

*Proof of Lemma 16.* By Assumption 2, the inequality  $(a + b)^2 \leq 2a^2 + 2b^2$ , and Jensen's,

$$\begin{aligned} \mathbb{E} \left[ \|\nabla \ell(\vartheta_t, Z_t) - \nabla \ell(\vartheta_{t-}, Z_{t-})\|^2 \right] &\leq L^2 \mathbb{E} \left[ \|\vartheta_t - \vartheta_{t-}\|^2 \right] \\ &= L^2 (t - t_-)^2 \left\{ \mathbb{E} \left[ \|\nabla_\theta \ell(\vartheta_{t-}, Z_{t-})\|^2 \right] + \mathbb{E} \left[ \|\nabla_x \ell(\vartheta_{t-}, Z_{t-})\|^2 \right] \right\} + 2L^2 \mathbb{E} \left[ \|W_t - W_{t-}\|^2 \right] \\ &\leq 2L^2 (t - t_-)^2 \left\{ \mathbb{E} \left[ \|\nabla_\theta \ell(\vartheta_t, Z_t)\|^2 \right] + \mathbb{E} \left[ \|\nabla_x \ell(\vartheta_t, Z_t)\|^2 \right] + \mathbb{E} \left[ \|\nabla \ell(\vartheta_t, Z_t) - \nabla \ell(\vartheta_{t-}, Z_{t-})\|^2 \right] \right\} \\ &\quad + 2L^2 \mathbb{E} \left[ \|W_t - W_{t-}\|^2 \right] \end{aligned}$$

Hence, if  $h \leq 1/2L \Rightarrow 1/2 \leq (1 - 2L^2(t - t_-)^2)$  we can then rearrange into

$$\begin{aligned} \frac{1}{2} \mathbb{E} \left[ \|\nabla \ell(\vartheta_t, Z_t) - \nabla \ell(\vartheta_{t-}, Z_{t-})\|^2 \right] &\leq (1 - 2L^2(t - t_-)^2) \mathbb{E} \left[ \|\nabla \ell(\vartheta_t, Z_t) - \nabla \ell(\vartheta_{t-}, Z_{t-})\|^2 \right] \\ &\leq 2L^2 (t - t_-)^2 \left\{ \mathbb{E} \left[ \|\nabla_\theta \ell(\vartheta_t, Z_t)\|^2 \right] + \mathbb{E} \left[ \|\nabla_x \ell(\vartheta_t, Z_t)\|^2 \right] \right\} + 2L^2 d_x (t - t_-) \end{aligned}$$

if further  $h \leq 1/4L \Rightarrow 4L^2(t - t_-)^2 \leq 1/4$ , we can use Lemma 21 to estimate

$$\begin{aligned} \mathbb{E} \left[ \left\| \nabla \ell(\vartheta_t, Z_t) - \nabla \ell(\vartheta_{t_-}, Z_{t_-}) \right\|^2 \right] &\leq \frac{1}{4} I(\vartheta_t, p_t) + 8L^3 d_x (t - t_-)^2 + 4L^2 d_x (t - t_-) \\ &\leq \frac{1}{4} I(\vartheta_t, p_t) + 6L^2 d_x (t - t_-) \end{aligned}$$

where we used  $8L(t - t_-) \leq 2$ . We conclude by combining this inequality with Lemma 20.  $\square$

### C.3 Proof of Theorem 17

By the descent Lemma 16 and the xLSI we have

$$\partial_t [F(\vartheta_t, p_t) - F_\star] \leq -\frac{1}{2} I(\vartheta_t, p_t) + 6L^2 d_x (t - t_-) \leq -\lambda [F(\vartheta_t, p_t) - F_\star] + 6L^2 d_x (t - t_-).$$

Now we proceed similarly to the proof of Theorem 12, and write

$$\partial_t \left[ e^{t\lambda} [F(\vartheta_t, p_t) - F_\star] \right] \leq 6L^2 d_x h e^{t\lambda}.$$

so that integrating from  $t = kh$  to  $t = (k+1)h$  gives

$$[F(\vartheta_{(k+1)h}, p_{(k+1)h}) - F_\star] \leq e^{-h\lambda} [F(\vartheta_{kh}, p_{kh}) - F_\star] + 6L^2 d_x h^2.$$

We conclude by iterating the above inequality, and by combining the result with the extension of Talagrand inequality via Theorem 3.  $\square$