# Identification and multiply robust estimation of causal effects via instrumental variables from an auxiliary heterogeneous population

Wei Li and Jiapeng Liu

Center for Applied Statistics and School of Statistics, Renmin University of China

and

Peng Ding

Department of Statistics, University of California, Berkeley

and

Zhi Geng

School of Mathematics and Statistics, Beijing Technology and Business University

#### Abstract

Evaluating causal effects in a primary population of interest with unmeasured confounders is challenging. Although instrumental variables (IVs) are widely used to address unmeasured confounding, they may not always be available in the primary population. Fortunately, IVs might have been used in previous observational studies on similar causal problems, and these auxiliary studies can be useful to infer causal effects in the primary population, even if they represent different populations. However, existing methods often assume homogeneity or equality of conditional average treatment effects between the primary and auxiliary populations, which may be limited in practice. This paper aims to remove the homogeneity requirement and establish a novel identifiability result allowing for different conditional average treatment effects across populations. We also construct a multiply robust estimator that remains consistent despite partial misspecifications of the observed data model and achieves local efficiency if all nuisance models are correct. The proposed approach is illustrated through simulation studies. We finally apply our approach by leveraging data from lower income individuals with cigarette price as a valid IV to evaluate the causal effect of smoking on physical functional status in higher income group where strong IVs are not available.

*Keywords:* Causal inference; Instrumental variable; Multiple robustness; Semiparametric inference; Unmeasured confounding

# 1 Introduction

It is challenging to evaluate causal effect of a treatment on an outcome in observational studies due to the presence of unmeasured confounding. A commonly employed strategy to address unmeasured confounding is the use of instrumetal variables (IVs). However, IVs may be unavailable in the target or primary population of interest. In such cases, previous auxiliary studies with complete instrument information can be leveraged to evaluate causal effects in the primary population. For example, suppose we are interested in the effect of education on earnings in a specific industry within a developing country where no suitable IVs are available. Similar studies in developed countries have used changes in compulsory schooling laws as IVs to study the same causal problem (Oreopoulos, 2006). By appropriately using these studies from the developed countries, we can still infer the causal effect of education on earnings in the developing country without IVs. Even in a primary population where an IV is available, its strength can vary across different subgroups. For some subgroups, the IV may be strong, but for others it can be very weak. When estimating causal effects in a subgroup with a weak IV, results may be unreliable if we rely on that weak IV. To address this, we can leverage information from subgroups with strong IVs to aid in estimating causal effects in the subgroup with the weak IV. For instance, in our motivating example, we aim to study the causal effect of smoking on physical functioning among higher income individuals. Leigh and Schembri (2004) used the price of cigarettes as an IV. However, in the higher income group, the relationship between cigarette price and smoking behavior is weak, making it a weak IV, whereas in the lower income group, the cigarette price is a strong IV. Therefore, we can use the lower income group with strong IV information to help identify and estimate the causal effect of smoking on physical functioning in the higher income group.

Integrating data from other sources has become increasingly popular for achieving valid causal inference in the primary population of interest (Yang and Ding, 2020; Shi et al., 2023; Degtiar and Rose, 2023; Li et al., 2024; Colnet et al., 2024). A strand of literature focuses on generalizing or transporting findings from auxiliary randomized trials to the primary population based on various versions of the selection exchangeability assumption (Stuart et al., 2011; Buchanan et al., 2018; Dahabreh et al., 2020; Yang et al., 2023). This assumption typically requires that the conditional average treatment effects given observed covariates are equal across both the primary and auxiliary populations. In the absence of randomized trials, some studies use an auxiliary dataset with supplemental confounding information to mitigate estimation bias in the primary population (McCandless et al., 2012; Yang and Ding, 2020; Luo et al., 2023). In particular, Yang and Ding (2020) proposed a general framework to estimate causal effects in the setting where the big primary data have unmeasured confounders, but the smaller external auxiliary data provide supplementary information on these confounders. However, such auxiliary datasets with full confounder information may not always be available in practice. Another line of research leverages auxiliary data with IVs to establish identification and estimation of causal effects when the primary sample lacks important variables for subsequent analysis (Ridder and Moffitt, 2007; Gamazon et al., 2015; Zhao et al., 2019). Specifically, in the classical twosample IV framework, the primary sample contains information on the outcome, IV, and covariates, but lacks information on the treatment; whereas the auxiliary sample provides information on the treatment, IV, covariates, but lacks information on the outcome. This methodology has been widely used in econometrics and social sciences, and mendelian randomization studies with genetic factors as instruments (Graham et al., 2016; Shu and Tan, 2020; Zhao et al., 2019). Recently, Sun and Miao (2022) developed semiparametric efficient estimation of average treatment effects in the two-sample IV framework. However, this framework requires measurements of instruments in both primary data and auxiliary data, which is fundamentally different from our setting where only auxiliary sample contains IVs. A more relevant paper by Shuai et al. (2024) considered similar settings where only the auxiliary sample has IVs and the primary sample includes the treatment and outcome, but their identifiability relies on a structural equation model with homogeneous treatment effects across the two samples. Despite rapid progress in the data fusion literature, most studies impose strong homogeneity assumptions or require the transportability of the conditional average treatment effects between the primary and auxiliary populations.

In this paper, we aim to remove the homogeneity requirement and propose a novel approach for identifying and estimating causal effects in the primary population with unmeasured confounders using IVs from an auxiliary population. Specifically, our study includes a primary sample from the population of interest without IVs and an auxiliary sample from possibly heterogeneous populations with a valid IV. Under a structural equation model, we show that the average treatment effect in the primary population can be uniquely identified from the observed data distribution by combining information from the auxiliary sample. Different from existing methods, our identifiability result allows for different conditional average treatment effects between the primary and auxiliary populations. Building on the identification, we derive the efficient influence function and propose a locally semiparametric efficient estimator that is multiply robust in the sense that it is consistent even if the observed data model is partially misspecified. Our proposed estimator remains its consistency and asymptotic normality when leveraging highly data-adaptive machine learning techniques for nuisance models. We finally develop a flexible sensitivity analysis approach to assess the impact of departures from identification assumptions on inference about the causal effect in the primary population.

The rest of this paper is organized as follows. Section 2 introduces notation, assumptions, and identification results. Section 3 presents the semiparametric efficient multiply robust estimator and discusses the use of flexible machine learning for nuisance models. Section 4 compares the multiply robust estimator with other semiparametric estimators. Section 5 provides a sensitivity analysis approach for assessing the impact of a departure from identifying assumptions. Section 6 uses simulation studies to evaluate the finite sample performance of the proposed estimators. Section 7 applies our approach to estimate the causal effect of smoking on physical functional status in the higher income individuals. We conclude with a brief discussion in Section 8 and relegate proofs to the supplementary material.

### 2 Identification

Suppose there are *n* individuals merged from two different sources. Let *R* denote the source indicator, where R = 0 if a subject is observed in the primary sample from the target population, and R = 1 if observed in the auxiliary sample from potentially heterogeneous populations. In the primary sample, we observe a binary treatment *X* with 0 and 1 for the labels of control and active treatments, respectively. Let *Y* denote an outcome of interest and *V* denote a vector of pre-treatment covariates. We assume that *U* contains all unmeasured common causes of *X* and *Y*. We use the Neyman-Rubin potential outcomes framework to define causal effects. Let  $Y_x$  denote the potential outcome if an individual, possibly contrary to the fact, were assigned to treatment *x*. We impose the consistency assumption that the observed outcome is  $Y = XY_1 + (1 - X)Y_0$ . We are interested in the average treatment effect of the primary population:

$$\tau = E(Y_1 - Y_0 \mid R = 0).$$

In the auxiliary sample, besides the treatment X, the outcome Y, and observed covariates V, a binary instrumental variable Z is also observed. We use  $f(\cdot)$  to denote the probability density or mass function of a random variable (vector). The notation  $A \perp B \mid C$  denotes the conditional independence between A and B given C for generic variables A, B, C. We make the following assumptions to identify causal effects.

Assumption 1 (Latent ignorability).  $Y_x \perp X \mid V, U, R$ .

Assumption 2 (IV for auxiliary population). (i)  $Z \perp U \mid V, R = 1$ ; (ii)  $Z \perp Y \mid X, V, U, R = 1$ ; (iii)  $Z \not\perp X \mid V, R = 1$ .

Assumption 1 means that the treatment X is latent ignorable given observed covariates V and unmeasured variables U in both primary and auxiliary populations (Frangakis and Rubin, 1999). Assumption 2 characterizes standard conditions for a valid instrumental variable Z in the auxiliary population (Angrist et al., 1996). Assumption 2(i) requires the instrument Z to be independent of unmeasured confounders U given V; Assumption 2(ii) formalizes no direct effect of the instrument Z on the outcome Y; Assumption 2(iii) ensures that the instrument Z is correlated with the treatment X even after conditioning on V. Although these assumptions are generally untestable without further conditions, potential IVs may be obtained through some natural or quasi-experiment in observational studies (Baiocchi et al., 2014; Wang and Tchetgen Tchetgen, 2018).

Assumptions 1 and 2 are commonly made in the causal inference literature. However, they are not sufficient to ensure identification of the causal parameter  $\tau$  in the primary population, because no suitable IVs are available in the primary population and the auxiliary population with IVs may be heterogeneous from the primary population. To address this issue, we further impose a semiparametric structural equation model to link the relationships between the primary and auxiliary populations, as outlined in the following assumption. Let  $\varepsilon(V, U) = E(X \mid V, U, R = 1) - E(X \mid V, U, R = 0)$  denote the difference of the treatment propensity scores given all confounders (V, U) between the two populations.

**Assumption 3** (Structural equation model). Suppose that  $U \perp R \mid V$ , and we consider the following structural equation model:

$$E(Y \mid X, V, U, R) = \zeta_0(V, U) + \zeta(V)R + \beta(V, R)X, \tag{1}$$

where  $\operatorname{Cov}\{\zeta_0(V,U), \varepsilon(V,U) \mid V, R\} = 0$ , and  $\{\zeta_0(\cdot), \zeta(\cdot), \beta(\cdot)\}$  are unknown scalar functions.

The outcome model in (1) allows  $\beta(V, R)$  to depend on R, indicating that the effect of X on Y conditional on the observed covariates V may vary between the primary and auxiliary populations. The first part of Assumption 3, i.e.,  $U \perp R \mid V$ , is generally weaker than the selection exchangeability assumption, i.e.,  $Y_x \perp R \mid V$ , which has been widely used in the literature for generalizing study's findings to the primary population. For instance, Degtiar and Rose (2023), Shi et al. (2023), and Colnet et al. (2024) all incorporate similar assumptions. Model (1) requires that there are no additive U-R and U-X interactions. We show in Section S2 of the supplementary material that model (1) may be relaxed so that  $\{\zeta(\cdot), \beta(\cdot)\}$  vary with U even after controlling for V under certain conditions, which will become saturated models for binary variables (R, X). A sufficient condition for the zerocovariance equation in Assumption 3 is  $\varepsilon(V, U) = \varepsilon(V)$ , which implies no additive U-R interaction in predicting the treatment after conditioning on observed covariates. Similar no-interaction conditions have been widely used in the recent literature for the identification of average treatment effect with IVs (Wang and Tchetgen Tchetgen, 2018; Sun and Miao, 2022) or the GENIUS approach with invalid IVs (Tchetgen Tchetgen et al., 2021; Sun et al., 2022). A design implication of Assumption 3 is that it requires us to collect as many causes of the treatment and outcome as possible such that no residual effect modification involving U remains within strata of the observed covariates V.

Below we simply let  $\beta_r(V) = \beta(V, r)$  for r = 0, 1, and we will use these two ways of notation interchangeably depending on the situation. Under Assumptions 1 and 3,

$$\beta_r(v) = E(Y_1 - Y_0 \mid V = v, R = r),$$

which encodes the conditional average treatment effect within levels of V among the population R = r, and hence  $\tau = E\{\beta_0(V) \mid R = 0\}$ . Note that  $\beta_1(V)$  can be identified by the instrumental variable Z, but generally  $\beta_0(V)$  is not equal to  $\beta_1(V)$ . In fact, under Assumptions 2 and 3, we show in Section S1.1 of the supplementary material that the identification equation of  $\beta_1(V)$  is given by

$$\beta_1(V) = \frac{E(Y \mid Z = 1, V, R = 1) - E(Y \mid Z = 0, V, R = 1)}{E(X \mid Z = 1, V, R = 1) - E(X \mid Z = 0, V, R = 1)} \triangleq \frac{\delta^Y(V)}{\delta^X(V)}.$$
(2)

This nonparametric representation is the well-known conditional Wald estimand, which has been derived under similar conditions in the IV literature (Wang and Tchetgen Tchetgen, 2018; Sun and Miao, 2022). For identification of  $\tau$ , a common strategy is to impose the selection exchangeability or some wearker mean exchangeability assumptions. These assumptions require that the primary and auxiliary populations are homogeneous conditional on the observed covariates V, and thus  $\beta_0(V) = \beta_1(V)$  is identifiable. However, these exchangeability assumptions could be violated if some shifted treatment effect modifiers are not captured in the concatenated data. This is a plausible scenario, particularly considering that the data are often collected independently and may represent heterogeneous populations.

In this paper, we do not assume  $\beta_0(V) = \beta_1(V)$  in the underlying model and we establish identification of  $\tau$  under Assumptions 1-3. Let  $\omega(V) = f(R = 1 | V)$ ,  $\mu(V, R) = E(X | V, R)$ ,  $\eta = X - \mu(V, R)$ , and  $\sigma^2(V, R) = \mu(V, R)\{1 - \mu(V, R)\}$  denote the sampling mechanism, treatment propensity score, treatment regression residual, and conditional variance of the treatment, respectively. To simplify notation, we use  $\mu_r(v)$  and  $\sigma_r^2(v)$  to represent  $\mu(v, r)$  and  $\sigma^2(v, r)$ , respectively. Since no restrictions are placed on the observed data distribution in Assumptions 1–3, the models we consider are essentially nonparametric, and we use  $\mathcal{M}$  to denote the corresponding nonparametric model. Letting q = f(R = 0), we then have the following theorem.

**Theorem 1.** Suppose that Assumptions 1–3 hold. Then  $\beta_0(V)$  is identified in  $\mathcal{M}$  by

$$\beta_0(V) = E \left\{ \frac{\beta_1(V)\sigma_1^2(V)}{\sigma_0^2(V)} - \frac{(2R-1)\eta Y}{f(R \mid V)\sigma_0^2(V)} \mid V \right\},\tag{3}$$

where  $\beta_1(V)$  is identified by (2). In addition, we have the following equation for  $\tau$ :

$$\tau = E\left[\frac{(1-R)\beta_1(V)\sigma_1^2(V)}{q\sigma_0^2(V)} - \left\{\frac{R}{f(R \mid V)} - 1\right\}\frac{\eta Y}{q\sigma_0^2(V)}\right].$$

The nonparametric identification formula about  $\beta_0(V)$  in (3) is new in the data fusion literature. Existing methods typically assume that  $\beta_0(V) = \beta_1(V)$ , but these two functions may not generally be equal, as shown in Theorem 1. The identifiability result in this theorem has some connections with the GENIUS approach (Tchetgen Tchetgen et al., 2021; Sun et al., 2022), which is used to identify average treatment effects in the context of a single data set with invalid instruments that violate the exclusion restriction assumption. Given  $U \perp R \mid V$  as stated in Assumption 3, it is possible to interprete R as an invalid IV that may potentially violate the relevance or exclusion restriction assumption. However, it should be noted that the framework considered here is fundamentally different from the GENIUS approach. The GENIUS approach precludes possible interactions between the invalid instrument (i.e., R) and the treatment in the outcome model. Additionally, it requires that the conditional variance of the treatment,  $\sigma^2(V, R)$ , must depend on the invalid instrument R. In contrast, our identifiability result does not require these conditions. Specifically, model (1) allows for a potential interaction between R and X, or equivalently permits  $\beta_r(V)$  to vary with r. Moreover, the conditional variance  $\sigma^2(V, R)$  in our setting does not necessarily depend on R, meaning that we allow  $\sigma_1^2(V) = \sigma_0^2(V)$ . The main reason we can still achieve identifiability results without these conditions is that we leverage data fusion settings where an auxiliary data set has provided relevant information about  $\beta_1(V)$ , which leaves us with the only task of identifying  $\beta_0(V)$  under the outcome model in (1).

Theorem 1 provides only one identification formula for  $\tau$  and there are several alternative identification formulas based on different components of the observed data likelihood, which are ommitted due to space contraints. It is important to note that the IV assumption 2 is specifically imposed to identify the conditional average treatment effect  $\beta_1(V)$  in the auxiliary sample, which can be replaced by any other assumptions as long as  $\beta_1(V)$  is guaranteed to be identifiable. For example, in cases where U is fully observable in the auxiliary sample, as considered in Yang and Ding (2020), then under Assumption 1, we can identify  $\beta_1(V)$  as follows:

$$\beta_1(V) = E\{E(Y \mid X = 1, V, U, R = 1) - E(Y \mid X = 0, V, U, R = 1) \mid V, R = 1\}.$$

In this scenario, the identification expression of  $\beta_0(V)$  in Theorem 1 is unchanged except that the expression of  $\beta_1(V)$  therein is replaced by the above equation.

#### 3 Multiply robust estimation

We have considered identification under a non-parametric model for the observed data distribution. The nuisance parameters in Theorem 1 can, in principle, be estimated nonparametrically using methods such as kernel smoothing or series estimation; however, as will typically be so in practice, the observed set of covariates V may be of moderate or high dimension, and nonparametric estimators may exhibit poor performance due to the curse of dimensionality. In this section, we turn our attention to a dicussion of semiparametric efficiency theory for estimation of the parameter  $\tau$ . Any regular and asymptotically linear estimator has an influence function. Accordingly, the influence function with the lowest variance is referred to as the efficient influence function (EIF), which is a key object in general semiparametric estimation theory (Bickel et al., 1993). The EIF is often useful in constructing locally efficient and multiply robust estimators, which allow for some components of the observed data distribution to be inconsistently estimated while preserving consistency of the estimators.

To derive the EIF, we let  $\pi(V) = f(Z = 1 | V, R = 1)$ ,  $\mu_0^X(V) = E(X | Z = 0, V, R = 1)$  and  $\mu_0^Y(V) = E(Y | Z = 0, V, R = 1)$  denote the conditional probability of Z = 1 given V, the conditional expectations of X and Y given (Z = 0, V) in the auxiliary population, respectively. We further define

$$\phi(V,R) = E\{Y - \beta(V,R)X \mid V,R\} \text{ and } \rho(V) = E[\eta\{Y - \beta(V,R)X\} \mid V,R = 0].$$
(4)

Similar to the simplifications in the previous section, we may use  $\phi_r(v)$  to represent  $\phi(v, r)$  for r = 0, 1. Here,  $\phi_r(v)$  is the conditional expectation of the residual  $Y - \beta(V, R)X$  given V = v in the population R = r, and  $\rho(V)$  is shown in Section S1.1 of the supplementary material to be the conditional unmeasured confounding bias given V.

Let  $\Delta = (\pi, \mu_0^X, \mu_0^Y, \delta^X, \beta_1, \omega, \mu_0, \beta_0, \phi_0, \rho)$  denote the set of all nuisance functions. Using the nuisances in  $\Delta$ , we can express the functions  $\mu_1(v)$  and  $\phi_1(v)$  as follows:

$$\mu_1(v) = \delta^X(v)\pi(v) + \mu_0^X(v), \quad \text{and} \quad \phi_1(v) = \mu_0^Y(v) - \beta_1(v)\mu_0^X(v).$$
(5)

Letting  $\gamma(v) = \{1 - \omega(v)\}/\omega(v)$ , we have the following theorem characterizing EIF for  $\tau$  under the nonparametric model of the observed data distribution, which serves as the foundation for our multiply robust estimator.

**Theorem 2.** Suppose that Assumptions 1–3 hold. The EIF for estimating  $\tau$  in  $\mathcal{M}$  is given by the following equation:

$$\begin{split} \varphi_{\text{eff}}(O;\Delta,q,\tau) = & \frac{1-R}{q} \{\beta_0(V) - \tau\} - \left\{ \frac{R}{f(R \mid V)} - 1 \right\} \frac{\eta\{Y - \beta(V,R)X - \phi(V,R)\} - \rho(V)}{q\sigma_0^2(V)} \\ & + \frac{\gamma(V)\sigma_1^2(V)}{q\sigma_0^2(V)} \varphi_1(O;\Delta), \end{split}$$

where

$$\varphi_1(O;\Delta) = \frac{R(2Z-1)}{f(Z \mid V, R=1)} \frac{1}{\delta^X(V)} \{ Y - \mu_0^Y(V) - X\beta_1(V) + \mu_0^X(V)\beta_1(V) \}.$$

Therefore, the semiparametric efficiency bound for estimating  $\tau$  in  $\mathcal{M}$  is  $E\{\varphi_{\text{eff}}(O;\Delta,q,\tau)^2\}$ .

The function  $\varphi_1(O; \Delta)$  in Theorem 2 involves only the auxiliary sample and plays an important role in characterizing EIF for the average treatment effect using IVs. For example, if we are interested in the average treatment effect in the auxiliary population, i.e.,  $\tau_{\text{aux}} = E(Y_1 - Y_0 \mid R = 1)$ , it can be identified by  $\tau_{\text{aux}} = E\{\beta_1(V) \mid R = 1\}$  under our model assumptions, where  $\beta_1(V)$  is identified by (2). The EIF for  $\tau_{\text{aux}}$  is expressed as follows:

$$\frac{R}{1-q}\left\{\beta_1(V) - \tau_{\text{aux}}\right\} + \frac{1}{1-q}\varphi_1(O;\Delta).$$

This EIF expression also involves the function  $\varphi_1(O; \Delta)$ , and it closly relates to the one derived by Wang and Tchetgen Tchetgen (2018) for the average treatment effect in the context of a single data set with IVs. The expression of the EIF for the average treatment effect  $\tau$  in the primary population, as presented in Theorem 2, is more complicated. Notably, we do not assume  $\beta_0(V) = \beta_1(V)$  when establishing the identifiability of  $\tau$ , and hence, the EIF for estimating  $\tau$  includes additional treatment and outcome information from the primary sample, as evidenced by the second term in  $\varphi_{\text{eff}}(O; \Delta, q, \tau)$ .

Let  $\mathbb{P}_n$  denote the empirical average, for example,  $\mathbb{P}_n\{h(V)\} = n^{-1}\sum_{i=1}^n h(V_i)$  for any  $h(\cdot)$ . Theorem 2 implies that we can construct an estimator of  $\tau$  by solving  $\mathbb{P}_n\{\varphi_{\text{eff}}(O; \hat{\Delta}, \hat{q}, \tau)\} = 0$  when nuisance functions in  $\Delta$  have been estimated and  $\hat{q} = \sum_{i=1}^n (1 - R_i)/n$ . The strategy of constructing estimators from estimating equations based on influence functions has been widely used in many causal inference literature (Wang and Tchetgen Tchetgen, 2018; Shi et al., 2020; Sun et al., 2022; Sun and Miao, 2022; Jiang et al., 2022). We will consider two different ways for modeling the nuisance functions in  $\Delta$ : one relies on standard parametric models, and the other uses modern machine learning techniques. We show that an estimator of  $\tau$  based on the EIF is multiply robust, in the sense that only certain combinations of the working models need to be correctly specified for consistent estimation. To construct the multiply robust estimator, we define the following two sets of nuisance models related to the auxiliary sample:  $\Delta_{\text{aux},1} = (\pi, \mu_0^X, \delta^X)$ , and  $\Delta_{\text{aux},2} = (\mu_0^X, \mu_0^Y, \beta_1)$ ; and the following three sets of nuisance models related to the primary sample:  $\Delta_{\text{pri},1} = (\omega, \mu_0)$ ,  $\Delta_{\text{pri},2} = (\omega, \beta_0, \phi_0)$ , and  $\Delta_{\text{pri},3} = (\mu_0, \beta_0, \rho)$ . Then consider the following semiparametric models that impose certain restrictions on the observed data distribution:

- $\mathcal{M}_1$ : models for  $\Delta_{\text{aux},1}$  and  $\Delta_{\text{pri},1}$  are correct;
- $\mathcal{M}_2$ : models for  $\Delta_{aux,2}$  and  $\Delta_{pri,1}$  are correct;
- $\mathcal{M}_3$ : models for  $\Delta_{aux,1}$  and  $\Delta_{pri,2}$  are correct;
- $\mathcal{M}_4$ : models for  $\Delta_{\text{aux},2}$  and  $\Delta_{\text{pri},2}$  are correct;
- $\mathcal{M}_5$ : models for  $\Delta_{aux,1}$  and  $\Delta_{pri,3}$  are correct;
- $\mathcal{M}_6$ : models for  $\Delta_{\text{aux},2}$  and  $\Delta_{\text{pri},3}$  are correct.

Let  $\mathcal{M}_{\text{union}} = \bigcup_{k=1}^{6} \mathcal{M}_k$  be defined as the semiparametric model that holds when any of  $\{\mathcal{M}_1, \ldots, \mathcal{M}_6\}$  is true. Below we separately consider parametric and machine learning methods for estimating the nuisance models in  $\Delta$ .

We first consider parametric working models  $\Delta(\Theta)$  with  $\Theta = (\theta_1^{\mathrm{T}}, \ldots, \theta_{10}^{\mathrm{T}})^{\mathrm{T}}$ , where  $\theta_j$  is a vector of finite-dimensional parameters associated with the *j*th nuisance model in  $\Delta$  for  $j = 1, \ldots, 10$ ; that is,  $\Delta(\Theta) = \{\pi(v; \theta_1), \mu_0^X(v; \theta_2), \mu_0^Y(v; \theta_3), \delta^X(v; \theta_4), \beta_1(v; \theta_5), \omega(v; \theta_6), \mu_0(v; \theta_7), \ldots, 10\}$ 

 $\beta_0(v;\theta_8), \phi_0(v;\theta_9), \rho(v;\theta_{10})$ }. Below we develop a three-step procedure to estimate the nuisance parameters. In the first step, we use maximum likelihood estimation to obtain estimators  $\hat{\theta}_1, \hat{\theta}_2, \hat{\theta}_3, \hat{\theta}_6, \hat{\theta}_7$  of the parameters in the nuisance models  $\pi, \mu_0^X, \mu_0^Y, \omega, \mu_0$ , respectively, because these nuisances are either models for binary variables or simply have conditional mean forms.

In the second step, we solve the following estimating equations to obtain the estimators  $\hat{\theta}_4$  and  $\hat{\theta}_5$  of the parameters associated with models  $\delta^X$  and  $\beta_1$ :

$$\mathbb{P}_{n}\Big[D_{4}(V)R\big\{X - \delta^{X}(V;\theta_{4})Z - \mu_{0}^{X}(V;\hat{\theta}_{2})\big\}\Big] = 0,$$
(6)

$$\mathbb{P}_n\Big[D_5(V)R\big\{Y - \beta_1(V;\theta_5)X - \mu_0^Y(V;\hat{\theta}_3) + \mu_0^X(V;\hat{\theta}_2)\beta_1(V;\theta_5)\big\}\Big] = 0,\tag{7}$$

where  $D_j(V)$  is a user-specified vector function of the same dimension as  $\theta_j$  for j = 4, 5. One can easily verify that the estimator  $\hat{\theta}_4$  is consistent if the models in  $\Delta_{\text{aux},1}$  are correct and  $\hat{\theta}_5$  is consistent if the models in  $\Delta_{\text{aux},2}$  are correct by noting that

$$E\{X - \delta^X(V)Z - \mu_0^X(V) \mid V, R = 1\} = 0, E\{Y - \beta_1(V)X - \mu_0^Y(V) + \mu_0^X(V)\beta_1(V) \mid V, R = 1\} = 0.$$

Based on the estimation procedures in the first and second steps, we can see that the estimators of the parameters associated with the nuisance modes in  $\Delta_{\text{aux},1}$ ,  $\Delta_{\text{aux},2}$ , or  $\Delta_{\text{pri},1}$  are consistent, provided that the corresponding models in each of these three sets are correctly specified.

We use  $\hat{\pi}(v)$  to simply denote  $\pi(v; \hat{\theta}_1)$ , and other estimated nuisance models are denoted similarly. After obtaining the estimators of the first seven nuisance functions in  $\Delta$ , we then plug them into (5) to obtain the estimates  $(\hat{\mu}_1, \hat{\phi}_1)$  of  $(\mu_1, \phi_1)$ . Similarly, we can obtain the estimates  $(\hat{\eta}, \hat{\sigma}_1^2, \hat{\gamma})$  by plugging the estimated nuisance functions into the corresponding equations of  $(\eta, \sigma_1^2, \gamma)$  introduced earlier; that is,

$$\hat{\eta} = X - R\hat{\mu}_1(V) - (1 - R)\hat{\mu}_0(V), \quad \hat{\sigma}_1^2(V) = \hat{\mu}_1(V)\{1 - \hat{\mu}_1(V)\}, \quad \hat{\gamma}(V) = \frac{1 - \hat{\omega}(V)}{\hat{\omega}(V)}.$$

Consequently, we can also obtain a plug-in estimate of  $\varphi_1(O; \Delta)$  defined in Theorem 2 and denote it by  $\hat{\varphi}_1$ .

In the third step, we aim to estimate the remaining nuisance models  $(\beta_0, \phi_0, \rho)$ . We first define two functions based on estimates of previous nuisance functions:

$$\hat{\beta}(\theta_8) = R\hat{\beta}_1(V) + (1-R)\beta_0(V;\theta_8), \qquad \hat{\phi}(\theta_9) = R\hat{\phi}_1(V) + (1-R)\phi_0(V;\theta_9).$$

Then we solve the following equations in (8)–(10) to obtain  $\hat{\theta}_8$ ,  $\hat{\theta}_9$ , and  $\hat{\theta}_{10}$  of the parameters associated with the three nuisance functions  $\beta_0$ ,  $\phi_0$ , and  $\rho$ :

$$\mathbb{P}_{n}\left[D_{8}(V)\left\{\left(\frac{R}{\hat{f}(R\mid V)}-1\right)\left(\hat{\eta}(Y-\hat{\beta}(\theta_{8})X-\hat{\phi}(\theta_{9}))-\rho(V;\theta_{10})\right)-\hat{\gamma}(V)\hat{\sigma}_{1}^{2}(V)\hat{\varphi}_{1}\right\}\right]=0,$$
(8)

$$\mathbb{P}_n\Big[D_9(V)(1-R)\big\{Y - \beta_0(V;\theta_8)X - \phi_0(V;\theta_9)\big\}\Big] = 0,$$
(9)

$$\mathbb{P}_n\Big[D_{10}(V)(1-R)\big\{(X-\hat{\mu}_0(V))(Y-\beta_0(V;\theta_8)X)-\rho(V;\theta_{10})\big\}\Big]=0,$$
(10)

where  $\hat{f}(R \mid V) = R\hat{\omega}(V) + (1-R)\{1-\hat{\omega}(V)\}$ , and  $D_j(V)$  is a user-specified vector function of the same dimension as  $\theta_j$  for j = 8, 9, 10. The estimating equations in (8) are similar to those in Sun et al. (2022) where they develop a GENIUS-approach based multiply robust efficient estimator of the average treatment effect in the context of a single data set with invalid IVs. The following lemma shows consistency of the estimators  $\hat{\theta}_8$ ,  $\hat{\theta}_9$ , and  $\hat{\theta}_{10}$  under certain conditions.

**Lemma 1.** The estimators  $\hat{\theta}_8$  and  $\hat{\theta}_9$  are consistent if  $\mathcal{M}_3 \cup \mathcal{M}_4$  holds, and the estimators  $\hat{\theta}_8$  and  $\hat{\theta}_{10}$  are consistent if  $\mathcal{M}_5 \cup \mathcal{M}_6$  holds.

In summary, the three-step procedures described above guarantee the consistency of the estimators for the parameters associated with the models in each  $\mathcal{M}_j$  if  $\mathcal{M}_j$  holds for  $j = 1, \ldots, 6$ . Let  $\hat{\Theta} = (\hat{\theta}_1^{\mathrm{T}}, \ldots, \hat{\theta}_{10}^{\mathrm{T}})^{\mathrm{T}}$  and  $\hat{\Delta} = \Delta(\hat{\Theta})$ . A multiply robust estimator  $\hat{\tau}_{\mathrm{mr}}$  is then obtained by solving the following estimating equation based on the EIF:

$$\mathbb{P}_n\left\{\varphi_{\text{eff}}(O;\hat{\Delta},\hat{q},\tau)\right\} = 0.$$
(11)

**Theorem 3.** Suppose that Assumptions 1–3 hold. Then  $\hat{\tau}_{mr}$  is a consistent and asymptotically normal estimator of  $\tau$  in  $\mathcal{M}_{union} = \bigcup_{k=1}^{6} \mathcal{M}_k$  under standard regularity conditions. Furthermore, if all the nuisance models in  $\Delta$  are correct, then  $\hat{\tau}_{mr}$  attains the semiparametric efficiency bound in  $\mathcal{M}$ .

The estimator  $\hat{\tau}_{mr}$  offers six genuine opportunities to estimate  $\tau$  consistently and also has valid inference provided that one of the six strategies holds. The asymptotic variance for  $\hat{\tau}_{mr}$  can be constructed using the sandwich estimator from the standard M-estimation theory (Newey and McFadden, 1994). However, due to computational complexity considerations, an alternative option in practice is nonparametric bootstrap procedures.

An important advantage of multiply robust estimators constructed using the EIF often yields second-order bias terms, which require slow convergence rates for the nuisance parameters involved, thereby enabling the use of flexible regression techniques in estimating these quantities (Chernozhukov et al., 2018; Kennedy, 2022). Below we propose such an estimation procedure using machine learning methods. With a bit abuse of notation, we still use  $\hat{\Delta}$  to denote the estimate of  $\Delta$  obtained by machine learning methods. Firstly, we obtain estimators  $(\hat{\pi}, \hat{\mu}_0^X, \hat{\mu}_0^Y, \hat{\omega}, \hat{\mu}_0)$  by machine learning of the conditional mean functions  $(\pi, \mu_0^X, \mu_0^Y, \omega, \mu_0)$ . Secondly, given the estimators  $(\hat{\pi}, \hat{\mu}_0^X, \hat{\mu}_0^Y)$ , we obtain  $\hat{\delta}^X$  and  $\hat{\beta}_1$  sequentially by machine learning methods based on the following conditional mean relationships:

$$\delta^X(V) = E\left\{\frac{X - \mu_0^X(V)}{\pi(V)} \mid V, R = 1\right\}, \quad \beta_1(V) = E\left\{\frac{Y - \mu_0^Y(V)}{\pi(V)\delta^X(V)} \mid V, R = 1\right\}.$$

Thirdly, given the estimators  $(\hat{\pi}, \hat{\mu}_0^X, \hat{\omega}, \hat{\mu}_0, \hat{\delta}^X, \hat{\beta}_1)$ , we obtain  $\hat{\beta}_0$ ,  $\hat{\phi}_0$ , and  $\hat{\rho}$  sequentially by machine learning methods based on the conditional mean relationships in (3) and (4). Finally, the machine learning based multiply robust estimator  $\hat{\tau}_{\rm mr}$  is again obtained by solving (11). Let  $||h||_2 = \{\int h(o)^2 dP(o)\}^{1/2}$  denote the  $L_2(P)$  norm of any real-valued function  $h(\cdot)$ . We next derive the asymptotic properties of the machine learning based estimator  $\hat{\tau}_{\rm mr}$  in the following theorem.

**Theorem 4** (Multiple machine learning estimation). Suppose that Assumptions 1-3 hold. We further assume the following conditions hold:

- (i)  $\hat{\Delta}(v) \to \Delta(v)$  in probability for all v in the support of V;
- (ii)  $\hat{\Delta}(v)$  and  $\Delta(v)$  are in a Donsker class;
- (*iii*)  $c < \{\hat{\pi}(v), \pi(v)\} < 1 c, \{\hat{\mu}_0(v), \mu_0(v), \hat{\omega}(v), \omega(v), |\hat{\delta}^X(v)|, |\delta^X(v)|\} > c, and$  $\{|\hat{\beta}_1(v)|, |\beta_1(v)|\} < C \text{ for some } c \in (0, 1), C > 0, and all v in the support of V;$
- (iv)  $n^{1/2}$ -convergence of second-order terms, i.e.,

$$(\|\hat{\beta}_1 - \beta_1\| + \|\hat{\mu}_0^Y - \mu_0^Y\| + \|\hat{\mu}_0^X - \mu_0^X\|) (\|\hat{\pi} - \pi\| + \|\hat{\delta}^X - \delta^X\| + \|\hat{\mu}_0^X - \mu_0^X\|) + \|\hat{\mu}_0 - \mu_0\| (\|\hat{\beta}_0 - \beta_0\| + \|\hat{\phi}_0 - \phi_0\|) + \|\hat{\omega} - \omega\| \|\hat{\rho} - \rho\| = o_p (n^{-1/2}).$$

Then  $\hat{\tau}_{mr}$  is asymptotically normal, has the influence function  $\varphi_{eff}(O; \Delta, q, \tau)$ , and achieves the semiparametric efficiency bound.

Conditions (i)-(iv) are similar to those for double machine learning estimation of average treatment effects (Kennedy et al., 2017; Chernozhukov et al., 2018; Kennedy, 2022). These conditions are all common in standard non-parametric regression problems (Van der Vaart, 2000). Among them, condition (ii) involves the complexity of the nuisance estimators  $\hat{\Delta}$  and their limits, and is a usual minimal regularity condition for problems involving nuisance functions. The cross-fitting techniques can be used to relax this condition (Chernozhukov et al., 2018; Kennedy, 2022). Condition (iv) requires that all the nuisance parameters are estimated with mean-squared error rates diminishing faster than  $n^{-1/4}$ . Such rates are achievable for many highly data-adaptive machine learning methods, including lasso, random forests or ensembles of these methods.

#### 4 Comparison with other semiparametric estimators

Under particular specifications of the nuisance models,  $\hat{\gamma}_{mr}$  reduces to other different semiparametric estimators. We introduce six such estimators, dependent on specific model specifications in  $\mathcal{M}_1, \ldots, \mathcal{M}_6$ , respectively. These estimators are based on parametric working models  $\Delta(\Theta)$  as introduced in Section 3.

The estimator  $\hat{\tau}_1$  under  $\mathcal{M}_1$  is defined as follows:

$$\hat{\tau}_1 = \mathbb{P}_n \left[ \frac{R(2Z-1)}{\hat{f}(Z \mid V, R=1)} \frac{\hat{\gamma}(V)\hat{\sigma}_1^2(V)}{\hat{q}\hat{\sigma}_0^2(V)} \frac{Y}{\hat{\delta}^X(V)} - \left\{ \frac{R}{\hat{f}(R \mid V)} - 1 \right\} \frac{\hat{\eta}Y}{\hat{q}\hat{\sigma}_0^2(V)} \right],$$

where  $\hat{f}(Z \mid V, R = 1) = Z\hat{\pi}(V) + (1 - Z)\{1 - \hat{\pi}(V)\}$ . The estimator  $\hat{\tau}_1$  depends only on the nuisance models in  $\mathcal{M}_1$  and is consistent and asymptotically normal if the models in  $\mathcal{M}_1$  are correctly specified. As shown in Section 3, the estimation of the nuisance models in  $\mathcal{M}_1$  can be obtained by maximizing likelihood functions along with solving estimating equations in (6), which does not involve outcome data. This estimator  $\hat{\tau}_1$  separates the design stage from the analysis stage in the sense that models in  $\mathcal{M}_1$  are specified before seeing any outcome data, and it helps to prevent selecting models that favour 'publishable' results (Rubin, 2007). However,  $\hat{\tau}_1$  fails to be multiply robust or locally efficient.

We next introduce the estimator  $\hat{\tau}_2$  under  $\mathcal{M}_2$  as below:

$$\hat{\tau}_2 = \mathbb{P}_n \bigg[ -\bigg\{ \frac{R}{\hat{f}(R \mid V)} - 1 \bigg\} \frac{\{X - (1 - R)\hat{\mu}_0(V)\}\{Y - R\hat{\beta}_1(V)X - R\hat{\phi}_1(V)\}}{\hat{q}\hat{\sigma}_0^2(V)} \bigg].$$

The estimator  $\hat{\tau}_2$  depends only on the nuisance models in  $\mathcal{M}_2$ , and the estimation of the nuisance models in  $\mathcal{M}_2$  can be obtained by maximizing likelihood functions along with solving estimating equations in (7).

In the sets of model specifications  $\mathcal{M}_3, \ldots, \mathcal{M}_6, \beta_0(V)$  is always required to be correct. Consequently, the estimators of  $\tau$  introduced below have the same expression across these settings, except that the parameter  $\theta_8$  involved in  $\beta_0(V; \theta_8)$  is estimated by solving different estimating equations. Specifically, the estimator  $\hat{\tau}_j$  under  $\mathcal{M}_j$  is given as follows:

$$\hat{\tau}_j = \mathbb{P}_n \left\{ \frac{1-R}{\hat{q}} \beta_0(V; \hat{\theta}_{8, \mathcal{M}_j}) \right\}, \quad (j = 3, \dots, 6),$$

where  $\hat{\theta}_{8,\mathcal{M}_j}$  is obtained by solving corresponding estimating equations that are shown in Section S1.6 of the supplementary material. We would like to highlight that the estimator  $\hat{\tau}_j$  relies soly

on the nuisance models in  $\mathcal{M}_j$  for  $j = 3, \ldots, 6$ . The estimation of the nuisance models in  $\mathcal{M}_j$  can be obtained by maximizing likelihood functions along with solving several estimating equations, as detailed in Section S1.6 of the supplementary material.

**Theorem 5.** Suppose that Assumptions 1–3 hold. Then  $\hat{\tau}_j$  is consistent and asymptotically normal in  $\mathcal{M}_j$  for  $j = 1, \ldots, 6$  under standard regularity conditions.

### 5 Flexible sensitivity analysis

Assumption 3 is crucial for the identification and estimation developed in previous sections. In this section, we develop a flexible machine learning based sensitivity analysis approach to assess the impact of partial violations of this assumption. For r = 0, 1, we define  $\varepsilon_r(V, U) = E(X | V, U, R = r)$  and  $\zeta_r(V, U) = E(Y | V, U, R = r, X = 0)$ , which denote the treatment propensity score given all confounders (V, U) and the baseline confounding effect on the outcome among the population R = r, respectively. We consider the following relaxed version of model (1) as imposed in Assumption 3:

$$E(Y \mid X, V, U, R) = \zeta_0(V, U) + \zeta(V, U)R + \beta(V, R)X,$$
(1')

where  $\{\zeta_0(\cdot), \zeta(\cdot), \beta(\cdot)\}\$  are unknown scalar functions. Compared with model (1) in Assumption 3, model (1') precludes only the existence of the U-X interaction rather than both U-R and U-X interactions. We next introduce the following sensitivity parameter:

$$\xi(V) = \operatorname{Cov}\{\varepsilon_0(V, U), \zeta_0(V, U) \mid V, R = 0\} - \operatorname{Cov}\{\varepsilon_1(V, U), \zeta_1(V, U) \mid V, R = 1\},\$$

which measures the difference of unmeasured confounding biases in the primary and auxiliary population. Under Assumption 3, the sensitivity parameter  $\xi(V) = 0$ , and hence, a large absolute value of  $\xi(V)$  indicates a strong deviation from this assumption.

**Proposition 1.** Suppose that Assumptions 1–2 and model (1') hold. Given the sensitivity parameter  $\xi(V), \beta_0(V)$  is identified by

$$\beta_0(V) = E \left\{ \frac{\beta_1(V)\sigma_1^2(V) - \xi(V)}{\sigma_0^2(V)} - \frac{(2R - 1)\eta Y}{f(R \mid V)\sigma_0^2(V)} \mid V \right\},\tag{12}$$

where  $\beta_1(V)$  is defined in (2). In addition, the parameter  $\tau$  is identified by

$$\tau = E\left[\frac{(1-R)\{\beta_1(V)\sigma_1^2(V) - \xi(V)\}}{q\sigma_0^2(V)} - \left\{\frac{R}{f(R \mid V)} - 1\right\}\frac{\eta Y}{q\sigma_0^2(V)}\right].$$

Proposition 1 shows that given a value of the sensitivity parameter  $\xi(V)$ ,  $\tau$  is identifiable under Assumptions 1–2 and model (1'). Similar to Theorem 2, we can also derive the EIF under the nonparametric model of observed data distribution, as shown in Proposition 2. **Proposition 2.** Suppose that the conditions in Proposition 1 hold. Given the sensitivity parameter  $\xi(V)$ , the EIF for estimating  $\tau$  under  $\mathcal{M}$  is given by

$$\begin{split} \varphi_{\text{eff},\xi}(O) = & \frac{1-R}{q} \{ \beta_0(V) - \tau \} - \left\{ \frac{R}{f(R \mid V)} - 1 \right\} \frac{\eta \{ Y - \beta(V,R)X - \phi(V,R) \} - \rho(V) + R\xi(V)}{q\sigma_0^2(V)} \\ & + \frac{\gamma(V)\sigma_1^2(V)}{q\sigma_0^2(V)} \varphi_1(O;\Delta), \end{split}$$

where  $\varphi_1(\cdot)$  is defined in Theorem 2.

Given the sensitivity parameter  $\xi(V)$ , we then propose a flexible estimation procedure using machine learning methods. The estimators  $(\hat{\pi}, \hat{\mu}_0^X, \hat{\mu}_0^Y, \hat{\omega}, \hat{\mu}_0, \hat{\delta}^X, \hat{\beta}_1)$  are obtained in the same way as those by machine learning methods in Section 3. Based on the expressions in (5), we obtain the estimators  $(\hat{\mu}_1, \hat{\phi}_1)$ . Similarly, we obtain  $(\hat{\eta}, \hat{\sigma}_1^2, \hat{\gamma})$  by plugging the nuisance model estimators into their corresponding equations, and we also obtain the estimator  $\hat{\varphi}_1$  of  $\varphi_1(O; \Delta)$  based on the expression defined in Theorem 2. Finally, we obtain  $\hat{\beta}_{0,\xi}$  by machine learning methods based on the conditional mean relationship in (12). The estimators  $\hat{\phi}_{0,\xi}$  and  $\hat{\rho}_{\xi}$  are obtained in the same way as those in Section 3. We further let

$$\hat{\beta}_{\xi}(V,R) = R\hat{\beta}_1 + (1-R)\hat{\beta}_{0,\xi}, \qquad \hat{\phi}_{\xi}(V,R) = R\hat{\phi}_1 + (1-R)\hat{\phi}_{0,\xi}.$$

Based on the above nuisance estimators and the sensitivity parameter  $\xi(V)$ , we define the following machine learning based multiply robust estimator for  $\tau$ :

$$\begin{split} \hat{\tau}_{\mathrm{mr},\xi} = & \mathbb{P}_n \bigg[ \frac{1-R}{q} \hat{\beta}_{0,\xi}(V) - \Big\{ \frac{R}{\hat{f}(R \mid V)} - 1 \Big\} \frac{\hat{\eta} \big\{ Y - \hat{\beta}_{\xi}(V,R) X - \hat{\phi}_{\xi}(V,R) \big\} - \hat{\rho}_{\xi}(V) + R\xi(V)}{q \hat{\sigma}_0^2(V)} \\ & \quad + \frac{\hat{\gamma}(V) \hat{\sigma}_1^2(V)}{q \hat{\sigma}_0^2(V)} \hat{\varphi}_1 \bigg]. \end{split}$$

**Proposition 3.** Suppose that the conditions in Proposition 1 and conditions (i)–(iv) in Theorem 4 hold. Then given the sensitivity parameter  $\xi(V)$ ,  $\hat{\tau}_{mr,\xi}$  is consistent, asymptotically normal, and has the influence function  $\varphi_{\text{eff},\xi}$  defined in Proposition 2.

#### 6 Simulation studies

In this section, we conduct simulation studies to evaluate the finite-sample performance of the proposed estimators. In our simulation setting, baseline covariates  $V = (V_1, \ldots, V_4)^{\mathrm{T}}$  are generated from independent standard normal distribution N(0,1). We consider the functional form  $V_k^* = \{1 + \exp(1 - 2V_k)\}^{-1}$  for  $k = 1, \ldots, 4$ . The unmeasured confounder U is generated from a uniform distribution  $\mathrm{Unif}\{-\rho(V^*), \rho(V^*)\}$ , where  $\rho^2(V^*) = 0.75 + 1.5V_1^* + 1.5V_2^* - 0.3V_3^* - 0.3V_4^*$ . Then the sampling indicator R, instrumental variable Z in the auxiliary sample, treatment X, and

outcome Y are separately generated according to the following mechanism:

$$R \mid V \sim \text{Bernoulli} \{ p = (1 + \exp(-0.8 - 3V_1^* + 2V_2^* + V_3^* + V_4^*))^{-1} \},$$
  

$$Z \mid V, R = 1 \sim \text{Bernoulli} \{ p = (1 + \exp(0.5 + V_1^* - 1.5V_2^* - 2V_3^* + V_4^*))^{-1} \},$$
  

$$X \mid Z, V, U, R \sim \text{Bernoulli} \{ p = R\alpha_1(Z) + (1 - R)\alpha_0(V) + 0.1U \},$$
  

$$Y \mid X, V, U, R \sim N\{ (2 - R + 2V_1^* - V_2^* + V_3^* - 0.5V_4^*)(1 + X) + U, 0.5^2 \},$$

where  $\alpha_1(Z) = \{1 + \exp(-1.3 + 2.4Z)\}^{-1}$  and  $\alpha_0(V) = \{1 + \exp(-V_1^* + V_2^* - 0.8V_3^* + 0.6V_4^*)\}^{-1}$ . Under the above data generating mechanism, the correct models for the nuisance models  $\{\pi(V), \mu_0^X(V), \omega(V), \mu_0(V)\}$ in  $\Delta$  are given by a logistic regression with V as linear predictors, and the correct models for the remaining models  $\{\mu_0^Y(V), \delta^X(V), \beta_1(V), \beta_1$ 

 $\beta_0(V), \phi_0(V), \rho(V)$  in  $\Delta$  are given by a linear regression with V as predictors. We are interested in estimating the average treatment effect among the primary population  $\tau$ , whose true value is 2.22.

Because the selection exchangeability assumption does not hold in our simulation setting, existing estimators based on this assumption generally have bias for estimating  $\tau$ . We apply the proposed multiply robust estimator  $\hat{\tau}_{mr}$  to estimate  $\tau$ , and use the estimators  $\hat{\tau}_1, \hat{\tau}_2, \hat{\tau}_3, \hat{\tau}_4, \hat{\tau}_5, \hat{\tau}_6$  introduced in Section 4 for comparison. We evaluate the performances of these estimators in situations where some models may be misspecified. In these cases, the quadratic functional form  $V_k^{**} = (V_k - 0.5)^2$  is used in place of  $V_k^*$  when constructing the nuisance models in  $\Delta$  for  $k = 1, \ldots, 4$ . Specifically, we present results from the following eight scenarios:

- $\mathcal{S}_0$ , all the models are correct;
- $\mathcal{S}_i$ , only the models in  $\mathcal{M}_i$  are correct for  $j = 1, \ldots, 6$ ;
- $\mathcal{S}_7$ , none of the models is correct.

Table 1 summarizes the simulation results for all estimators under the eight scenarios for n = 2000, 10000, where the bias, standard deviation, and 95% coverage probability of these estimators averaged across 200 replications are reported. As expected, the proposed estimator  $\hat{\tau}_{mr}$  exhibits small bias and has coverage probability close to the nominal level in  $\{S_j\}_{j=0}^6$ , confirming its multiple robustness property. In contrast, the other six estimators  $\hat{\tau}_1, \hat{\tau}_2, \hat{\tau}_3, \hat{\tau}_4, \hat{\tau}_5, \hat{\tau}_6$  generally show substantial bias in scenarios where their respective models are misspecified. For example, the estimator  $\hat{\tau}_1$  performs well in scenarios  $\{S_j\}_{j=0,1}$ , while exhibiting notable bias in scenarios  $\{S_j\}_{j=2,3,4,6,7}$ . We also observe that  $\hat{\tau}_1$  has comparable performance in some scenarios where models are misspecified, e.g.,  $S_5$ , to those in scenarios where models are correct. Similar phenomenon may appear in specific scenarios for other estimators, which is subject to the simulation setting. In general, these six estimators can maintain their consistency only when their corresponding models are correct.

We also implement the proposed multiply robust estimator by using machine learning methods for nuisance model estimation. Specifically, we consider estimators  $\hat{\tau}_{mr,lasso}$  and  $\hat{\tau}_{mr,sl}$ , whereby the nuisance models are estimated by lasso and super learner, respectively. Figure 1 shows the corresponding

Model		$\hat{ au}_1$		$\hat{ au}_2$		$\hat{ au}_3$		$\hat{ au}_4$		$\hat{\tau}_5$		$\hat{\tau}_6$		$\hat{\tau}_{\mathrm{mr}}$	
	2000	10000	2000	10000	2000	10000	2000	10000	2000	10000	2000	10000	2000	10000	
							В	ias							
$\mathcal{S}_0$	0	1	-2	0	0	1	-2	0	-1	1	-2	0	1	0	
$\mathcal{S}_1$	0	1	216	214	37	37	239	236	-11	-8	8	8	-2	0	
$\mathcal{S}_2$	-95	-104	-2	0	-52	-61	35	37	8	8	-11	-9	2	0	
$\mathcal{S}_3$	37	37	238	236	0	1	215	212	7	6	-58	-63	-2	0	
$\mathcal{S}_4$	-52	-62	35	37	-94	-102	-2	0	47	43	6	7	2	0	
$S_5$	-1	-2	-2	-3	12	12	60	58	-1	1	240	238	-4	1	
$\mathcal{S}_6$	10	11	-3	-2	24	27	11	12	-132	-141	-2	0	-5	-1	
$\mathcal{S}_7$	50	50	49	49	56	50	50	50	44	41	43	42	49	50	
							S	D							
$\mathcal{S}_0$	27	11	18	7	27	11	17	7	23	9	16	7	18	8	
$\mathcal{S}_1$	27	11	31	14	27	11	27	12	20	11	22	11	34	13	
$\mathcal{S}_2$	83	13	18	7	84	13	18	8	117	11	15	9	40	8	
$\mathcal{S}_3$	27	11	27	12	27	11	29	13	23	17	48	42	32	12	
$\mathcal{S}_4$	87	13	18	8	82	13	17	7	45	20	22	17	44	8	
$\mathcal{S}_5$	23	10	26	10	21	8	25	10	23	9	28	13	32	12	
$\mathcal{S}_6$	38	20	13	5	34	8	13	5	78	16	16	7	34	12	
$\mathcal{S}_7$	40	16	27	9	81	9	23	9	24	12	22	12	24	11	
							C	P							
$\mathcal{S}_0$	94	96	95	95	94	95	96	95	98	95	97	96	95	94	
$\mathcal{S}_1$	94	96	1	0	71	6	1	0	94	76	94	66	95	95	
$\mathcal{S}_2$	20	0	95	95	63	0	59	1	95	70	82	62	98	94	
$\mathcal{S}_3$	72	1	1	0	94	95	1	0	80	65	69	63	94	94	
$\mathcal{S}_4$	64	0	59	1	21	0	96	95	84	40	76	61	98	95	
$S_5$	93	94	95	95	89	64	26	1	98	95	1	0	94	94	
$\mathcal{S}_6$	98	83	95	93	87	6	87	34	28	0	97	96	98	97	
$S_7$	62	3	39	2	57	2	37	1	62	20	49	17	58	1	
0.6 U U U U U U U U U U U U U U U U U U U							Coverage	.00							
.70	ی میں میں میں میں میں میں میں میں میں می						Sample Size								

Table 1: Bias, standard deviation (SD), and 95% coverage probability (CP) of the semiparametric estimators under different sample sizes n = 2000, 10000. All results have been multiplied by  $10^2$ .

Figure 1: Root mean squared error (RMSE) and 95% coverage probability of two multiply robust machine learning estimators under different sample sizes. The red circles indicate the multiply robust lasso estimator  $\tau_{\rm mr,lasso}$ , and the green triangles indicate the multiply robust super learner estimator  $\tau_{\rm mr,sl}$ .

results about root mean squared error (RMSE) and 95% coverage probability of these two multiply robust machine learning estimators based on 200 repeated experiments with sample size from 2000 to 20000. We observe that as sample size increases, the RMSEs of both estimators become smaller, and the 95% coverage probabilities are more close to the nominal level. These results corroborate our previous theoretical findings and demonstrate the advantages of the proposed multiply robust estimators.

# 7 Application

In this section, we demonstrate an application of the proposed method by estimating the effects of smoking on physical functional status based on data from the Household Survey portion of Community Tracking Study (CTS) for the years 1996-1997, which are accessible through application at https: //www.icpsr.umich.edu/web/HMCA/studies/2524/datadocumentation. The CTS is desgined to provide health care system change and its effects on people in the United States (Kemper et al., 1996). Smoking is a risk factor of many diseases including heart disease, stroke, diabetes, and most cancers. These diseases affect physical functioning, that is, physical disability. To mitigate potential unmeasured confounding baises, Leigh and Schembri (2004) used cigarette price as an IV to assess the impact of smoking on physical functional status. We found that cigarette price is a strong IV only for the lower income group but is a weak IV for the higher income group. The higher income group with strong IV to evaluate the effect of smoking on physical functional status for the lower income group.

Let the treatment variable X = 1 if the subject smokes every day or on some days, and X = 0 if the subject does not smoke at all. The outcome Y of interest is physical functional status, measured by the Physical Component Summary of the SF-12 (Ware et al., 1995). The baseline covariates include sex, age, mental functional status, education, health condition and race. However, there may still exist unobserved variables confounding the relationship between smoking and physical functional status, such as genetic factors, psychological stress, etc. Let the instrumental variable Z = 1 if the cigarette price is greater than the median and Z = 0 otherwise. Since our purpose is to estimate the causal effect in the higher income subpopulation, we use the subset of data containing higher income subjects as the primary sample. The primary sample consists of 4198 subjects, with 1123 having X = 1 and 3075 having X = 0. The auxiliary sample includes 18603 lower income subjects, with 7834 having X = 1 and 10769 having X = 0.

While the IV (i.e., cigarette price) is available in the higher income subpopulation, it shows weak correlation with the treatment. This is because the decision of whether or not higher income individuals smoke may not be significantly influenced by cigarette price. In fact, after conducting a logistic regression of the treatment on the IV in the primary data and adjusting for observed covariates, we found that the coefficient of the IV is not statistically significant. When we apply the classical IV approach to estimate the effect of smoking on physical functional status using only the primary data, the point estimate and standard error are -0.37 and 6.75, respectively. The corresponding confidence interval (-13.6, 12.86) is too wide. These results are possibly unreliable due to the weak IV issue

Estimators	Point estimate	Standard error	95% Confidence interval
$\hat{ au}_1$	-2.04	0.60	(-3.22, -0.386)
$\hat{ au}_2$	-0.15	1.48	(-3.05, 2.75)
$\hat{ au}_3$	-2.13	0.53	(-3.17, -1.09)
$\hat{ au}_4$	-0.31	0.84	(-1.96, 1.34)
$\hat{ au}_5$	-0.27	0.11	(-0.49, -0.05)
$\hat{ au}_6$	-0.39	0.93	(-2.21, 1.43)
$\hat{ au}_{ m mr}$	-0.25	0.17	(-0.58, 0.08)
$\hat{\tau}_{\mathrm{mr,lasso}}$	-0.22	0.11	(-0.44, -0.01)
$\hat{ au}_{ m mr,sl}$	-0.24	0.11	(-0.46, -0.02)

Table 2: Estimates of the causal effect of smoking on the physical functional status in the higher income subpopulation.

in the primary data. Fortunately, the IV is strongly correlated with the treatment in the auxiliary data, and thus we can leverage this information and use the proposed semiparametric estimators  $\hat{\tau}_1, \hat{\tau}_2, \hat{\tau}_3, \hat{\tau}_4, \hat{\tau}_5, \hat{\tau}_6, \hat{\tau}_{mr}$  and the multiply robust machine learning estimators  $\hat{\tau}_{mr,lasso}, \hat{\tau}_{mr,sl}$  that were employed in the simulation studies to analyze the data. Table 2 presents the point estimates, standard errors, and 95% confidence intervals for these estimators. The multiply robust machine learning estimators  $\hat{\tau}_{mr,lasso}, \hat{\tau}_{mr,sl}$  are close to each other. Different from the result obtained by the classical IV approach, both of the multiply robust machine learning estimators that leverage the strong IV from the auxiliary lower income group are negative and statistically significant, suggesting that smoking significantly decreases physical functioning for higher income subjects. The multiply robust estimator  $\hat{\tau}_m$  based on parametric models performs similarly, but with a slightly larger standard error. For the remaining estimators,  $\hat{\tau}_1, \hat{\tau}_3$  yield substantially smaller point estimates than the multiply robust estimators, whereas  $\hat{\tau}_2, \hat{\tau}_4, \hat{\tau}_6$  have much larger standard errors, despite their point estimates being closer to the multiply robust estimators. This is likely due to the misspecification of one or several models utilized in these estimators.

#### 8 Discussion

Instrumental variables are widely used to identify causal effects in the presence of unmeasured confounding. However, the stringent constraints for a valid instrumental variable may render it unavailable in the primary population of interest. This paper proposes an approach to identify and estimate the average causal effect in the primary population via instrumental variables from an auxiliary dataset. Unlike previous data fusion methods, our approach does not rely on homogeneous assumptions or transportability of the conditional average treatment effect across the two populations. We have developed a semiparametric efficient multiply robust estimator and investigated its asymptotic properties, even when employing highly data-adaptive machine learning techniques for nuisance model estimation. Additionally, we also present a flexible sensitivity analysis approach for potential violations of crucial identifying assumptions.

This paper can be extended or improved in several directions. Firstly, our study focuses on the canonical case of binary instrument and treatment variables. A promising avenue for future work is to broaden this framework to encompass various types of these variables. Secondly, the proposed framework has the potential for extension to incorporate multiple auxiliary datasets with instrumental variables, and the efficiency of the proposed estimators could be further improved. Thirdly, it is also of interest to investigate the use of negative controls, a concept gaining popularity in recent years (Miao et al., 2018; Tchetgen Tchetgen et al., 2024), to identify and estimate causal effects under data fusion. We intend to explore these and other related issues in future research.

# References

- Angrist, J., Imbens, G., and Rubin, D. B. (1996). Identification of causal effects using instrumental variables. Journal of the American Statistical Association, 91:444–455.
- Baiocchi, M., Cheng, J., and Small, D. S. (2014). Instrumental variable methods for causal inference. Statistics in Medicine, 33(13):2297–2340.
- Bickel, P. J., Klaassen, C. A., Ritov, Y., and Wellner, J. A. (1993). Efficient and Adaptive Estimation for Semiparametric Models. Johns Hopkins University Press Baltimore.
- Buchanan, A. L., Hudgens, M. G., Cole, S. R., Mollan, K. R., Sax, P. E., Daar, E. S., Adimora, A. A., Eron, J. J., and Mugavero, M. J. (2018). Generalizing evidence from randomized trials using inverse probability of sampling weights. *Journal of the Royal Statistical Society Series A: Statistics* in Society, 181(4):1193–1209.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1):C1–C68.
- Colnet, B., Mayer, I., Chen, G., Dieng, A., Li, R., Varoquaux, G., Vert, J.-P., Josse, J., and Yang, S. (2024). Causal inference methods for combining randomized trials and observational studies: a review. *Statistical Science*, 39(1):165–191.
- Dahabreh, I. J., Robertson, S. E., Steingrimsson, J. A., Stuart, E. A., and Hernan, M. A. (2020). Extending inferences from a randomized trial to a new target population. *Statistics in Medicine*, 39(14):1999–2014.
- Degtiar, I. and Rose, S. (2023). A review of generalizability and transportability. Annual Review of Statistics and Its Application, 10:501–524.

- Frangakis, C. E. and Rubin, D. B. (1999). Addressing complications of intention-to-treat analysis in the combined presence of all-or-none treatment-noncompliance and subsequent missing outcomes. *Biometrika*, 86(2):365–379.
- Gamazon, E. R., Wheeler, H. E., Shah, K. P., Mozaffari, S. V., Aquino-Michaels, K., Carroll, R. J., Eyler, A. E., Denny, J. C., Consortium, G., Nicolae, D. L., et al. (2015). A gene-based association method for mapping traits using reference transcriptome data. *Nature Genetics*, 47(9):1091–1098.
- Graham, B. S., Pinto, C. C. d. X., and Egel, D. (2016). Efficient estimation of data combination models by the method of auxiliary-to-study tilting (ast). *Journal of Business & Economic Statistics*, 34(2):288–301.
- Jiang, Z., Yang, S., and Ding, P. (2022). Multiply robust estimation of causal effects under principal ignorability. Journal of the Royal Statistical Society Series B: Statistical Methodology, 84(4):1423– 1445.
- Kemper, P., Blumenthal, D., Corrigan, J. M., Cunningham, P. J., Felt, S. M., Grossman, J. M., Kohn, L. T., Metcalf, C. E., St. Peter, R. F., Strouse, R. C., et al. (1996). The design of the community tracking study: a longitudinal study of health system change and its effects on people. *Inquiry*, 33:195–206.
- Kennedy, E. H. (2022). Semiparametric doubly robust targeted double machine learning: a review. arXiv preprint arXiv:2203.06469.
- Kennedy, E. H., Ma, Z., McHugh, M. D., and Small, D. S. (2017). Non-parametric methods for doubly robust estimation of continuous treatment effects. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 79(4):1229–1245.
- Leigh, J. P. and Schembri, M. (2004). Instrumental variables technique: cigarette price provided better estimate of effects of smoking on sf-12. *Journal of Clinical Epidemiology*, 57(3):284–293.
- Li, W., Luo, S., and Xu, W. (2024). Calibrated regression estimation using empirical likelihood under data fusion. *Computational Statistics & Data Analysis*, 190:107871.
- Luo, S., Zhang, Y., and Li, W. (2023). Multiply robust estimation of causal effects using linked data. arXiv preprint arXiv:2309.08199.
- McCandless, L. C., Richardson, S., and Best, N. (2012). Adjustment for Missing Confounders Using External Validation Data and Propensity Scores. *Journal of the American Statistical Association*, 107(497):40–51.
- Miao, W., Geng, Z., and Tchetgen Tchetgen, E. J. (2018). Identifying causal effects with proxy variables of an unmeasured confounder. *Biometrika*, 105(4):987–993.

- Newey, W. K. and McFadden, D. (1994). Large sample estimation and hypothesis testing. *Handbook* of *Econometrics*, 4:2111–2245.
- Oreopoulos, P. (2006). Estimating average and local average treatment effects of education when compulsory schooling laws really matter. *American Economic Review*, 96(1):152–175.
- Ridder, G. and Moffitt, R. (2007). The econometrics of data combination. *Handbook of Econometrics*, 6:5469–5547.
- Rubin, D. B. (2007). The design versus the analysis of observational studies for causal effects: parallels with the design of randomized trials. *Statistics in Medicine*, 26(1):20–36.
- Shi, X., Miao, W., Nelson, J. C., and Tchetgen Tchetgen, E. J. (2020). Multiply robust causal inference with double-negative control adjustment for categorical unmeasured confounding. *Journal* of the Royal Statistical Society Series B: Statistical Methodology, 82(2):521–540.
- Shi, X., Pan, Z., and Miao, W. (2023). Data integration in causal inference. Wiley Interdisciplinary Reviews: Computational Statistics, 15(1):e1581.
- Shu, H. and Tan, Z. (2020). Improved methods for moment restriction models with data combination and an application to two-sample instrumental variable estimation. *Canadian Journal of Statistics*, 48(2):259–284.
- Shuai, K., Luo, S., Li, W., and He, Y. (2024). Identifying causal effects using instrumental variables from the auxiliary population. *Statistica Sinica, to appear.*
- Stuart, E. A., Cole, S. R., Bradshaw, C. P., and Leaf, P. J. (2011). The use of propensity scores to assess the generalizability of results from randomized trials. *Journal of the Royal Statistical Society Series A: Statistics in Society*, 174(2):369–386.
- Sun, B., Cui, Y., and Tchetgen, E. T. (2022). Selective machine learning of the average treatment effect with an invalid instrumental variable. *The Journal of Machine Learning Research*, 23(204):1– 40.
- Sun, B. and Miao, W. (2022). On semiparametric instrumental variable estimation of average treatment effects through data fusion. *Statistica Sinica*, 32:569–590.
- Tchetgen Tchetgen, E., Sun, B., and Walter, S. (2021). The genius approach to robust mendelian randomization inference. *Statistical Science*, 36(3):443–464.
- Tchetgen Tchetgen, E. J., Ying, A., Cui, Y., Shi, X., and Miao, W. (2024). An introduction to proximal causal learning. *Statistical Science*, 39(3):375–390.
- Van der Vaart, A. W. (2000). Asymptotic Statistics. Cambridge University Press.

- Wang, L. and Tchetgen Tchetgen, E. (2018). Bounded, efficient and multiply robust estimation of average treatment effects using instrumental variables. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 80(3):531–550.
- Ware, J. E., Keller, S. D., and Kosinski, M. (1995). SF-12: How to score the SF-12 physical and mental health summary scales. Health Institute, New England Medical Center.
- Yang, S. and Ding, P. (2020). Combining Multiple Observational Data Sources to Estimate Causal Effects. Journal of the American Statistical Association, 115(531):1540–1554.
- Yang, S., Gao, C., Zeng, D., and Wang, X. (2023). Elastic integrative analysis of randomised trial and real-world data for treatment heterogeneity estimation. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 85(3):575–596.
- Zhao, Q., Wang, J., Spiller, W., Bowden, J., and Small, D. S. (2019). Two-sample instrumental variable analyses using heterogeneous samples. *Statistical Science*, 34(2):317–333.