

# AUTOMORPHISM GROUP OF THE HOLOMORPH OF A CYCLIC GROUP

KAZUKI SATO

ABSTRACT. We show that the holomorph of a cyclic group of order  $n$  is isomorphic to its own automorphism group when  $n$  is twice of a power of an odd prime.

## 1. INTRODUCTION

Let  $C_n$  be a cyclic group of order  $n$ . In this paper we prove the following.

**Theorem 1.1** (Theorem 3.8). *Let  $p$  be an odd prime and  $e > 0$  a positive integer. Assume  $n = 2p^e$ . Let  $G = \text{Hol}(C_n)$  be the holomorph of  $C_n$ . Then there exists an isomorphism  $G \cong \text{Aut}(G)$ .*

It is known that the automorphism group of the dihedral group  $D_{2n}$  of order  $2n$  is isomorphic to the holomorph  $\text{Hol}(C_n)$  of  $C_n$  [4]. It is also known that the holomorph  $\text{Hol}(C_n)$  of  $C_n$  is a complete group when  $n$  is odd [2]. In particular,  $\text{Hol}(C_n)$  is isomorphic to  $\text{Aut}(\text{Hol}(C_n))$  when  $n$  is odd. But when  $n$  is even, the center  $Z(\text{Hol}(C_n))$  is nontrivial (see Lemma 2.2), so  $\text{Hol}(C_n)$  is not complete. Despite that fact, Theorem 1.1 asserts that two groups  $\text{Hol}(C_n)$  and  $\text{Aut}(\text{Hol}(C_n))$  are isomorphic as abstract groups when  $n = 2p^e$ . We note that the case  $e = 1$ , that is,  $n = 2p$  in Theorem 1.1 is known to be true [3, Theorem 2.10].

In section 2, we recall preliminary results from elementary group theory and number theory. We show our main theorem in section 3.

## 2. PRELIMINARIES

**2.1. Semi-direct product.** We will review briefly the notion of a semi-direct product. Given two groups  $A$  and  $B$ , and a homomorphism  $\theta : B \rightarrow \text{Aut}(A)$ , the semi-direct product gives a group structure to the set  $G = A \times B$  by

$$(a_1, b_1) \cdot (a_2, b_2) = (a_1\theta(b_1)(a_2), b_1b_2).$$

We will denote the semi-direct product of  $A$  by  $B$  with  $\theta : B \rightarrow \text{Aut}(A)$  by  $A \rtimes_{\theta} B$ , where  $\theta$  may be omitted if it is clear from the context. We identify an element  $a \in A$  with  $(a, 1)$  and  $b \in B$  with  $(1, b)$ . With these identifications we can see that all the elements of  $A \rtimes_{\theta} B$  can be written uniquely in the form  $ab$  with  $a \in A$  and  $b \in B$ . Multiplication of elements from  $A$  and  $B$  is determined by the relation  $bab^{-1} = \theta(b)(a)$ . From these identifications we can write  $A \rtimes_{\theta} B = AB$  with  $A \triangleleft A \rtimes_{\theta} B$  and  $A \cap B = 1$ . In particular  $(A \rtimes_{\theta} B)/A \cong B$ . If  $B = \text{Aut}(A)$  and  $\theta$  is the identity map, then we call  $A \rtimes \text{Aut}(A)$  the holomorph of  $A$  and write it as  $\text{Hol}(A)$ .

---

*Date:* July 29, 2024.

*2010 Mathematics Subject Classification.* 20D45.

We use the following theorem in section 3.

**Theorem 2.1** ([1, Theorem 3.35]). *Let  $G$  and  $G_0$  be groups, and let  $N \triangleleft G$  and  $N_0 \triangleleft G_0$ , where  $G/N$  and  $G_0/N_0$  are cyclic of finite order  $m$ . Suppose that  $y : N \rightarrow N_0$  is an isomorphism, and let  $gN$  and  $g_0N_0$  be generating cosets of  $G/N$  and  $G_0/N_0$  respectively. Assume that*

$$y(g^m) = (g_0)^m \quad \text{and} \quad y(gxg^{-1}) = g_0y(x)g_0^{-1}$$

for all elements  $x \in N$ . Then there is a unique isomorphism  $\alpha : G \rightarrow G_0$  that extends the given isomorphism  $y : N \rightarrow N_0$  and that satisfies  $\alpha(g) = g_0$ .

**2.2. Dihedral Groups.** Let  $n$  be a positive integer. In this paper we let  $C_n$  denote the cyclic group of order  $n$  and  $D_{2n}$  the dihedral group of order  $2n$ . As mentioned in the Introduction, it is known that the automorphism group  $\text{Aut}(D_{2n})$  of  $D_{2n}$  is isomorphic to the holomorph  $\text{Hol}(C_n)$  of  $C_n$ .

**Lemma 2.2.** *Let  $n$  be an even integer. Let  $G = \text{Hol}(C_n)$  be the holomorph of  $C_n$ . Then the center  $Z(G)$  of  $G$  is equal to  $\{1, z\}$ , where  $z \in C_n$  is the unique involution.*

*Proof.* Since  $y(z) \in C_n$  is also an involution for any  $y \in \text{Aut}(C_n)$ , we have  $yz y^{-1} = y(z) = z$ . Hence  $\{1, z\} \subset Z(G)$ . We have to show the reverse inclusion.

Let  $xy \in Z(G)$ , where  $x \in C_n, y \in \text{Aut}(C_n)$ . Then we have  $(xy)x' = x'(xy)$  for any  $x' \in C_n$ . Note that  $yx' = y(x')y$ , so  $xy(x')y = x(yx') = (xy)x' = x'(xy) = xx'y$  for any  $x' \in C_n$ . This implies that  $y(x') = x'$  for any  $x' \in C_n$ , hence  $y = 1 \in \text{Aut}(C_n)$ . Let  $\iota \in \text{Aut}(C_n)$  be the automorphism given by  $\iota(x) = x^{-1}$ . Since  $x \in Z(G)$ , we have  $x = \iota x \iota^{-1} = \iota(x) = x^{-1}$ . Therefore  $x^2 = 1$ , that is,  $x = 1$  or  $x = z$  is the involution. This shows that  $Z(G) \subset \{1, z\}$ .  $\square$

If  $G$  is an arbitrary group, then  $G/Z(G)$  is naturally isomorphic to the group of inner automorphisms  $\text{Inn}(G)$ . If  $Z(G) = 1$ , therefore,  $G \cong \text{Inn}(G)$ , and so we can identify  $G$  with  $\text{Inn}(G)$  via the natural isomorphism, and this embeds  $G$  as a subgroup of the automorphism group  $\text{Aut}(G)$ . We say that  $G$  is complete if  $Z(G) = 1$  and  $\text{Aut}(G) = \text{Inn}(G)$ . It is known that  $\text{Aut}(D_{2n}) \cong \text{Hol}(C_n)$  is complete when  $n$  is odd. But when  $n$  is even,  $\text{Aut}(D_{2n})$  can never be complete.

### 2.3. Elementary number theory.

**Lemma 2.3.** *Let  $k$  be an integer and  $p$  a prime. Assume that  $k \equiv 1 \pmod{p}$ . Then  $k^{p^{e-1}} \equiv 1 \pmod{p^e}$  for any  $e \geq 1$ .*

*Proof.* By induction on  $e$ . Assume that  $k^{p^{e-1}} \equiv 1 \pmod{p^e}$  for some  $e \geq 1$ . Then  $k^{p^{e-1}} = 1 + rp^e$  for some integer  $r$ . Hence we have

$$k^{p^e} = (1 + rp^e)^p = 1 + \sum_{i=1}^{p-1} \binom{p}{i} r^i p^{ei} + r^p p^{ep}.$$

For each  $1 \leq i \leq p-1$ , since  $\binom{p}{i}$  is divisible by  $p$ , we see that  $\binom{p}{i} p^{ei} \equiv 0 \pmod{p^{e+1}}$ . Clearly  $r^p p^{ep} \equiv 0 \pmod{p^{e+1}}$ . This shows that  $k^{p^e} \equiv 1 \pmod{p^{e+1}}$ .  $\square$

**Lemma 2.4.** *Let  $p$  be an odd prime and  $e > 0$ . Let  $k$  be an integer coprime to  $n = 2p^e$ . Assume that the order of  $k \in (\mathbb{Z}/n\mathbb{Z})^*$  is  $\phi(n) = p^{e-1}(p-1)$ , where  $\phi$  is Euler's totient function. Then*

- (i)  $k-1$  is not divisible by  $p$ ,

- (ii) the greatest common divisor of  $n$  and  $k - 1$  is  $(n, k - 1) = 2$ ,
- (iii)  $1 + k + k^2 + \dots + k^{\phi(n)-1} \equiv 0 \pmod{n}$ .

*Proof.* (i) Assume  $k \equiv 1 \pmod{p}$ . Then  $k^{p^{e-1}} \equiv 1 \pmod{p^e}$  from Lemma 2.3. Since  $k$  is odd, we also have  $k^{p^{e-1}} \equiv 1 \pmod{2}$ . Hence  $k^{p^{e-1}} \equiv 1 \pmod{n}$ , so the order of  $k \in (\mathbb{Z}/n\mathbb{Z})^*$  divides  $p^{e-1} < p^{e-1}(p-1) = \phi(n)$ , contradiction.

(ii) Since  $n$  and  $k - 1$  are even,  $2|(n, k - 1)$ . On the other hand,  $(n, k - 1)$  is not divisible by  $p$  by (i). This shows (ii).

(iii) See [3, Theorem 2.5].  $\square$

### 3. MAIN THEOREM

In this section, we fix the notation as follows. Let  $e > 0$  be an integer and  $p$  an odd prime. Put  $n := 2p^e$ . Then the automorphism group  $\text{Aut}(C_n)$  is a cyclic group of order  $\phi(n) = p^{e-1}(p-1)$ . We fix generators  $x \in C_n$  and  $y \in \text{Aut}(C_n)$ . Assume that  $y(x) = x^k$ , where  $k$  is an integer coprime to  $n$  with  $1 < k < n$ . In this situation, there exists an isomorphism  $\text{Aut}(C_n) \cong (\mathbb{Z}/n\mathbb{Z})^*$  that maps  $y$  to  $k$ . We note that the order of  $k \in (\mathbb{Z}/n\mathbb{Z})^*$  is  $\phi(n)$ . Let  $G := \text{Hol}(C_n) = C_n \rtimes \text{Aut}(C_n)$  be the holomorph of  $C_n$ . Then  $G$  is generated by  $x$  and  $y$ , and the elements of  $G$  can be written in the form  $x^a y^b$ . We see that  $x^a y^b = x^c y^d$  if and only if  $a \equiv c \pmod{n}$  and  $b \equiv d \pmod{\phi(n)}$ . We have the following relations in  $G$ ;

$$x^n = 1, \quad y^{\phi(n)} = 1, \quad yxy^{-1} = y(x) = x^k.$$

**Lemma 3.1.**  $y^b x^a y^{-b} = x^{ak^b}$  holds for any integer  $a$  and nonnegative integer  $b$ .

*Proof.* By induction on  $b$ . The case  $b = 0$  is trivial. Assume that  $b > 0$  and that  $y^{b-1} x^a y^{-(b-1)} = x^{ak^{b-1}}$  for any  $a$ . Then we have  $y^b x^a y^{-b} = y^{b-1} y x^a y^{-1} y^{-(b-1)} = y^{b-1} x^{ak} y^{-(b-1)} = x^{ak^b}$ . Here, for the second equality, we used the relation in  $G$  and for the third equality, we used the inductive hypothesis.  $\square$

**Lemma 3.2.**  $(x^a y^b)^m = x^{a(1+k^b+\dots+k^{b(m-1)})} y^{bm}$  holds for any integer  $a$ , nonnegative integer  $b$  and positive integer  $m$ .

*Proof.* By induction on  $m$ . The case  $m = 1$  is trivial. Assume that  $m > 1$ . Then

$$\begin{aligned} (x^a y^b)^m &= (x^a y^b)^{m-1} x^a y^b \\ &= x^{a(1+k^b+\dots+k^{b(m-2)})} y^{b(m-1)} x^a y^b \\ &= x^{a(1+k^b+\dots+k^{b(m-2)})} y^{b(m-1)} x^a y^{-b(m-1)} y^{bm} \\ &= x^{a(1+k^b+\dots+k^{b(m-2)})} x^{ak^{b(m-1)}} y^{bm} \\ &= x^{a(1+k^b+\dots+k^{b(m-1)})} y^{bm}. \end{aligned}$$

Here, for the second equality, we used the inductive hypothesis and for the fourth equality, we used Lemma 3.1.  $\square$

**Lemma 3.3.** Let  $\alpha \in \text{Aut}(G)$ . Then  $\alpha(x) = x^a$  for some integer  $a$  coprime to  $n$ . Hence there exists an integer  $j$  such that  $\alpha(x) = y^j(x)$ .

*Proof.* The last statement is clear from the first one. Let  $\alpha \in \text{Aut}(G)$ . Assume  $\alpha(x) = x^a y^b$ ,  $\alpha(y) = x^c y^d$  for some integers  $a, b, c, d$ , with  $0 \leq b < \phi(n)$ . First, we will show that  $b = 0$ . We must have  $\alpha(y)\alpha(x)\alpha(y^{-1}) = \alpha(x^k)$ . It follows that

$$(x^c y^d)(x^a y^b)(x^c y^d)^{-1} = (x^a y^b)^k,$$

which considered modulo  $\langle x \rangle$  implies that  $y^b = y^{bk}$ , and thus consequently that  $y^{bk-b} = 1$ . So  $bk \equiv b \pmod{\phi(n)}$ . Assume  $0 < b < \phi(n)$ . We note that  $k^b - 1 \neq 0$ , since  $1 < k < n$ . By Lemma 3.2, we see that

$$(x^a y^b)^{k-1} = x^{a(1+k^b+\dots+k^{b(k-2)})} y^{b(k-1)} = x^{a(k^{b(k-1)}-1)/(k^b-1)}.$$

Since the order of  $x^a y^b = \alpha(x)$  is  $n$ , the order of  $(x^a y^b)^{k-1}$  is  $n/(n, k-1) = n/2$  from Lemma 2.4, (ii). On the other hand, the order of  $x^{a(k^{b(k-1)}-1)/(k^b-1)}$  is  $n/\left(n, a \frac{k^{b(k-1)}-1}{k^b-1}\right)$ . Hence

$$(1) \quad \left(n, a \frac{k^{b(k-1)}-1}{k^b-1}\right) = 2.$$

Recall that  $b(k-1) \equiv 0 \pmod{\phi(n)}$ . Hence

$$(2) \quad k^{b(k-1)} \equiv 1 \pmod{n}.$$

(1) and (2) imply that  $k^b - 1 \equiv 0 \pmod{p^e}$ . Since  $k$  is odd, we also have  $k^b - 1 \equiv 0 \pmod{2}$ . Therefore we can conclude that  $k^b - 1 \equiv 0 \pmod{n}$ . This contradicts the assumptions that  $k \in (\mathbb{Z}/n\mathbb{Z})^*$  has order  $\phi(n)$  and that  $0 < b < \phi(n)$ . Hence  $b = 0$  and  $\alpha(x) = x^a$ . Since  $\alpha(x) = x^a$  has order  $n$ ,  $a$  is coprime to  $n$  and this completes the proof.  $\square$

**Lemma 3.4.** *Let  $\alpha \in \text{Aut}(G)$ . Then  $\alpha(y) = x^c y$  for some integer  $c$ .*

*Proof.* Let  $\alpha \in \text{Aut}(G)$ . By Lemma 3.3, we can assume that  $\alpha(x) = x^a$ ,  $\alpha(y) = x^c y^d$  for some integers  $a, c, d$  with  $a$  coprime to  $n$  and  $0 \leq d < \phi(n)$ . As above we must have

$$(x^c y^d)(x^a)(x^c y^d)^{-1} = (x^a)^k.$$

This implies  $x^{ak^d} = x^{ak}$ , so  $ak^d \equiv ak \pmod{n}$ . Since  $ak$  is coprime to  $n$ , we have  $k^{d-1} \equiv 1 \pmod{n}$ . Hence  $d-1 \equiv 0 \pmod{\phi(n)}$ . Therefore  $y^d = y$ , and we have  $\alpha(y) = x^c y^d = x^c y$ .  $\square$

We define a map  $\psi : \text{Aut}(G) \rightarrow G$  as follows. Let  $\alpha \in \text{Aut}(G)$ . By Lemmas 3.3 and 3.4, there exist integers  $j$  and  $c$  such that  $\alpha(x) = y^j(x)$ ,  $\alpha(y) = x^c y$ . We note that  $y^j$  and  $x^c$  are uniquely determined from  $\alpha$ . We define

$$\psi(\alpha) = x^c y^j.$$

**Proposition 3.5.** *The map  $\psi : \text{Aut}(G) \rightarrow G$  is a group homomorphism.*

*Proof.* Let  $\alpha, \beta \in \text{Aut}(G)$ . Assume  $\psi(\alpha) = x^c y^j$ ,  $\psi(\beta) = x^d y^m$ . Then we have  $(\alpha\beta)(x) = \alpha(\beta(x)) = \alpha(y^m(x)) = y^j(y^m(x)) = y^{j+m}(x)$  and  $(\alpha\beta)(y) = \alpha(\beta(y)) = \alpha(x^d y) = \alpha(x)^d \alpha(y) = y^j(x^d) x^c y$ . Hence  $\psi(\alpha\beta) = y^j(x^d) x^c y^{j+m}$ . On the other hand, we have  $\psi(\alpha)\psi(\beta) = x^c y^j x^d y^m = x^c y^j(x^d) y^{j+m} = y^j(x^d) x^c y^{j+m}$ . Therefore  $\psi(\alpha)\psi(\beta) = \psi(\alpha\beta)$ .  $\square$

**Proposition 3.6.** *The map  $\psi : \text{Aut}(G) \rightarrow G$  is injective.*

*Proof.* Let  $\alpha \in \text{Aut}(G)$ . Assume  $\psi(\alpha) = x^c y^j = 1$ . Then  $x^c = 1$  and  $y^j = 1$ , so we have  $\alpha(x) = y^j(x) = x$  and  $\alpha(y) = x^c y = y$ . Since  $G$  is generated by  $x$  and  $y$ , we see that  $\alpha$  is the identity map on  $G$ .  $\square$

**Proposition 3.7.** *The map  $\psi : \text{Aut}(G) \rightarrow G$  is surjective.*

*Proof.* Let  $z \in G$ . Then  $z = x^c y^j$  for some  $0 \leq c < n$  and  $0 \leq j < \phi(n)$ . We note that  $yC_n = (x^c y)C_n$  is a generating coset of  $G/C_n$ . We have  $(x^c y)^{\phi(n)} = x^{c(1+k+\dots+k^{\phi(n)-1})} y^{\phi(n)} = 1$  by Lemmas 3.2 and 2.4, (iii). Furthermore,  $y^j (yxy^{-1}) = y^j (y(x)) = y^{j+1}(x) = x^c y (y^j(x)) x^{-c} = (x^c y) y^j(x) (x^c y)^{-1}$ . Taking  $i$ -th power of both sides, we have  $y^j (yx^i y^{-1}) = (x^c y) y^j(x^i) (x^c y)^{-1}$  for all integers  $i$ . By Theorem 2.1, there exists a unique isomorphism  $\alpha : G \rightarrow G$  that extends the isomorphism  $y^j : \langle x \rangle \rightarrow \langle x \rangle$  so that  $\alpha(y) = x^c y$ . Then  $\psi(\alpha) = x^c y^j = z$  by definition of  $\psi$ .  $\square$

**Theorem 3.8.** *The map  $\psi : \text{Aut}(G) \rightarrow G$  is an isomorphism.*

*Proof.* This follows from Propositions 3.5, 3.6 and 3.7.  $\square$

#### REFERENCES

- [1] I. M. Isaacs, *Finite group theory*, Graduate Studies in Mathematics, vol. 92, American Mathematical Society, Providence, RI, 2008.
- [2] G. A. Miller, *On the holomorph of a cyclic group*, Trans. Amer. Math. Soc. **4** (1903), no. 2, 153–160.
- [3] S. Sajikumar, S. Vinod, and G. S. Biju, *Automorphisms of automorphism group of dihedral groups*, Creat. Math. Inform. **32** (2023), no. 2, 229–235.
- [4] G. L. Walls, *Automorphism groups*, Amer. Math. Monthly **93** (1986), no. 6, 459–462.

NATIONAL INSTITUTE OF TECHNOLOGY, ICHINOSEKI COLLEGE, JAPAN  
*Email address:* kazuki-s@ichinoseki.ac.jp