

The Redei-Berge function and related combinatorial Hopf algebras

Stefan Mitrović

Faculty of Mathematics
University of Belgrade
Serbia

stefan.mitrovic@matf.bg.ac.rs

Tanja Stojadinović

Faculty of Mathematics
University of Belgrade
Serbia

tanja.stojadinovic@matf.bg.ac.rs

July 29, 2024

Abstract

Stanley and Grinberg introduced the symmetric function associated to digraphs, called the Redei-Berge symmetric function. In [8] is shown that this symmetric function arises from a suitable structure of combinatorial Hopf algebra on digraphs. In this paper, we introduce two new combinatorial Hopf algebras of posets and permutations and define corresponding Redei-Berge functions for them. By using both theories, of symmetric functions and of combinatorial Hopf algebras, we prove many properties of the Redei-Berge function. These include some forms of deletion-contraction property, which make it similar to the chromatic symmetric function. We also find some invariants of digraphs that are detected by the Redei-Berge function.

Keywords: digraph, Redei-Berge symmetric function, combinatorial Hopf algebra

MSC2020: 05C20, 05E05, 16T30

1 Introduction

Redei's and Berge's theorems are two beautiful graph-theoretical results that deserve to be better known than they are. Berge's theorem says that if X is a simple digraph and \overline{X} is its complement, then the number of Hamiltonian paths of \overline{X} is congruent to the same number for X modulo 2. Redei's theorem says that if X is a tournament, then the number of Hamiltonian paths of X is odd.

Recent paper of Grinberg and Stanley [7] has revealed these results as a consequence of symmetric function theory, by constructing a symmetric function U_X (the Redei-Berge symmetric function) whose p -basis expansion easily yields both theorems. The function U_X first appeared in the 1990s, in Chow's paper [5],

and later in Wiseman's [14] in 2007, in connection with the path-cycle symmetric function of a digraph. The function U_X is in [8] identified as the image of the isomorphism class of digraph X under a certain canonical Hopf algebra morphism, defined through the Aguiar-Bergeron-Sottile universal construction applied to a combinatorial Hopf algebra of digraphs. This reconceptualization is used to prove some properties of U_X , including the antipode formula. The authors in [8] also reprove Berge's theorem by combining this antipode formula with a deletion-contraction-like recurrence for the principal evaluation of U_X . This principal specialization, $u_X(m)$, is called the Redei-Berge polynomial.

In this paper, we further investigate properties of the Redei-Berge function of digraphs. We introduce two combinatorial Hopf subalgebras of the combinatorial Hopf algebra of digraphs, which enables us to define the Redei-Berge function of permutations and posets. In the sequel, we investigate which invariants of digraphs and posets can be obtained from the Redei-Berge function and prove some deletion properties of this function. Finally, we construct numerous new bases for the algebra of symmetric functions whose generators are Redei-Berge functions.

2 Preliminaries

The theory of combinatorial Hopf algebras is founded in [1]. We review some basic notions and facts. A *partition* $\{V_1, \dots, V_k\} \vdash V$ of length k of a finite set V is a set of disjoint nonempty subsets with $V_1 \cup \dots \cup V_k = V$. A *composition* $(V_1, \dots, V_k) \models V$ is an ordered partition.

A composition $\alpha \models n$, where $n \in \mathbb{N}$, is a sequence $\alpha = (a_1, \dots, a_k)$ of positive integers with $a_1 + \dots + a_k = n$. The type of a composition $(V_1, \dots, V_k) \models V$ is the composition type $(V_1, \dots, V_k) = (|V_1|, \dots, |V_k|) \models |V|$. A partition $\lambda \vdash n$ is a composition $(a_1, \dots, a_k) \models n$ such that $a_1 \geq a_2 \geq \dots \geq a_k$. There is a bijection between the set of compositions of n , denoted by $\text{Comp}(n)$ and the power set $2^{[n-1]}$ given by $(a_1, \dots, a_k) \mapsto \{a_1, a_1 + a_2, \dots, a_1 + \dots + a_{k-1}\}$. We denote the inverse of this bijection by $I \mapsto \text{comp}(I)$.

A *combinatorial Hopf algebra*, CHA for short, (\mathcal{H}, ζ) over a field \mathbf{k} is a graded, connected Hopf algebra $\mathcal{H} = \bigoplus_{n \geq 0} \mathcal{H}_n$ over \mathbf{k} together with a multiplicative functional $\zeta : \mathcal{H} \rightarrow \mathbf{k}$ called the *character*, [1].

The terminal object in the category of CHAs is the CHA of quasisymmetric functions $(QSym, \zeta_Q)$. For basics of quasisymmetric functions see [13] and for the Hopf algebra structure on $QSym$ see [1]. A composition $\alpha = (a_1, \dots, a_k) \models n$ defines the monomial quasisymmetric function

$$M_\alpha = \sum_{i_1 < \dots < i_k} x_{i_1}^{a_1} \dots x_{i_k}^{a_k}.$$

Alternatively, we write $M_I = M_{\text{comp}(I)}$, $I \subset [n-1]$. Another basis consists of fundamental quasisymmetric functions

$$F_I = \sum_{\substack{1 \leq i_1 \leq i_2 \leq \dots \leq i_n \\ i_j < i_{j+1} \text{ for each } j \in I}} x_{i_1} x_{i_2} \cdots x_{i_n}, \quad I \subset [n-1], \quad (1)$$

which are expressed in the monomial basis by

$$F_I = \sum_{I \subset J} M_J, \quad I \subset [n-1]. \quad (2)$$

The unique canonical morphism $\Psi : (\mathcal{H}, \zeta) \rightarrow (QSym, \zeta_Q)$ is given on homogeneous elements with

$$\Psi(h) = \sum_{\alpha \models n} \zeta_\alpha(h) M_\alpha, \quad h \in \mathcal{H}_n, \quad (3)$$

where ζ_α is the convolution product

$$\zeta_\alpha = \zeta_{a_1} \cdots \zeta_{a_k} : \mathcal{H} \xrightarrow{\Delta^{(k-1)}} \mathcal{H}^{\otimes k} \xrightarrow{proj} \mathcal{H}_{a_1} \otimes \cdots \otimes \mathcal{H}_{a_k} \xrightarrow{\zeta^{\otimes k}} \mathbf{k}. \quad (4)$$

The algebra of symmetric functions Sym is a subalgebra of $QSym$ and it is the terminal object in the category of cocommutative combinatorial Hopf algebras. This algebra consists of quasisymmetric functions that are invariant under the action of permutations on the set of variables. As a vector space, it has many natural bases, see [13]. We will be particularly interested in the *power sum basis*. The i th *power sum symmetric function* is defined as

$$p_0 = 1, \\ p_i = \sum_{j=1}^{\infty} x_j^i,$$

for $i \geq 1$. For partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$, we define

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_k}.$$

A *digraph* X is a pair $X = (V, E)$, where V is a finite set and E is a collection $E \subseteq V \times V$. Elements $u \in V$ are vertices and elements $(u, v) \in E$ are directed edges of the digraph X .

A *V-listing* is a list of all vertices with no repetitions, i.e. a bijective map $\pi : [n] \rightarrow V$. We write Σ_V for the set of all V -listings. For a V -listing $\pi = (\pi_1, \dots, \pi_n) \in \Sigma_V$, define the *X-descent set* as

$$XDes(\pi) = \{1 \leq i \leq n-1 \mid (\pi_i, \pi_{i+1}) \in E\}.$$

If V is the set $[n] = \{1, \dots, n\}$ and $E = \{(i, j) \mid 1 \leq j < i \leq n\}$ then X -descent sets are standard descent sets of permutations $\pi \in \Sigma_{[n]}$.

If $X = (V, E)$ is a digraph, its complementary digraph is the digraph $\overline{X} = (V, (V \times V) \setminus E)$ and its opposite digraph is $X^{op} = (V, E')$, where $E' = \{(v, u) \mid (u, v) \in E\}$.

Grinberg and Stanley associated to a digraph X a generating function for X -descent sets expanded in the basis of fundamental quasisymmetric functions

$$U_X = \sum_{\pi \in \Sigma_V} F_{X\text{Des}(\pi)} \quad (5)$$

and named it the Redei-Berge symmetric function, see [7], [12].

Definition 2.1. Let $X = (V, E)$ be a digraph. Then, we define

$$\Sigma_V(X) = \{\pi \in \Sigma_V \mid \text{each non-trivial cycle of } \pi \text{ is a cycle of } X\},$$

$$\Sigma_V(X, \overline{X}) = \{\pi \in \Sigma_V \mid \text{each cycle of } \pi \text{ is a cycle of } X, \text{ or a cycle of } \overline{X}\}.$$

The following theorem shows that U_X is, indeed, a symmetric function for any digraph X . For a permutation π , let $\text{type}(\pi)$ denote the partition whose entries are the lengths of the cycles of π .

Theorem 2.2. [7] *If $X = (V, E)$ is a digraph, then*

$$U_X = \sum_{\pi \in \Sigma_V(X, \overline{X})} (-1)^{\varphi(\pi)} p_{\text{type}(\pi)},$$

with $\varphi(\pi) = \sum_{\gamma} (\ell(\gamma) - 1)$, where the summation runs over all cycles γ of π that are cycles in X and $\ell(\gamma)$ denotes the length of the cycle γ .

We say that two digraphs $X = (V, E)$ and $Y = (V', E')$ are isomorphic if there is a bijection of vertices $f : V \rightarrow V'$ such that $(u, v) \in E$ if and only if $(f(u), f(v)) \in E'$. Denote by $[X]$ the isomorphism class of a digraph X .

Let $\mathcal{D} = \bigoplus_{n \geq 0} \mathcal{D}_n$ be the graded vector space over the field of rational numbers \mathbb{Q} , which is linearly spanned by the set of all isomorphism classes of digraphs, where the grading is given by the number of vertices. In [8] the space \mathcal{D} is endowed with the structure of combinatorial Hopf algebra.

For digraphs $X = (V, E)$ and $Y = (V', E')$ we define the product $X \cdot Y$ as the digraph on the disjoint union $V \sqcup V'$ with the set of directed edges

$$E \cup E' \cup \{(u, v) \mid u \in V, v \in V'\}.$$

The product of digraphs is obviously an associative, but not a commutative operation. The linear extension of the product on digraphs determines the multiplication $\mu : \mathcal{D} \otimes \mathcal{D} \rightarrow \mathcal{D}$, which turns the space \mathcal{D} into a noncommutative algebra. The restriction of a digraph $X = (V, E)$ on a subset $S \subset V$ is the digraph $X|_S = (S, E|_S)$, where $E|_S = \{(u, v) \in E \mid u, v \in S\}$. The restrictions of digraphs may be used to define a comultiplication $\Delta : \mathcal{D} \rightarrow \mathcal{D} \otimes \mathcal{D}$ by

$$\Delta([X]) = \sum_{(V_1, V_2) \models V} [X|_{V_1}] \otimes [X|_{V_2}],$$

where V is the vertex set of a digraph X . Evidently, this is a coassociative and a cocommutative operation. It is easy to check that Δ is an algebra morphism.

The unit element is given by the class $[\emptyset]$ of the digraph on the empty set of vertices, while the counit is given by $\epsilon([\emptyset]) = 1$ and $\epsilon([X]) = 0$ otherwise.

Let $\zeta : \mathcal{D} \rightarrow \mathbb{Q}$ be the multiplicative linear functional given with

$$\zeta([X]) = \#\{\pi \in \Sigma_V \mid X\text{Des}(\pi) = \emptyset\}. \quad (6)$$

Note that $\zeta([X])$ actually counts Hamiltonian paths of the complementary digraph \overline{X} (which is an invariant of isomorphism classes of digraphs).

The fact that the space \mathcal{D} , together with operations defined in this way forms a CHA, sheds a new light on the Redei-Berge function.

Theorem 2.3. [8] *Let $\Psi : \mathcal{D} \rightarrow QSym$ be the universal morphism from the combinatorial Hopf algebra of digraphs to quasisymmetric functions. Then, for a digraph X ,*

$$\Psi([X]) = U_X.$$

Theorem 2.4. [8] *For any digraph X ,*

$$U_X = U_{X^{op}}.$$

3 The Redei-Berge function of permutations and posets

In this section, we introduce two combinatorial Hopf algebras compatible with the combinatorial Hopf algebra structure on digraphs. As a consequence, we show how the definition of the Redei-Berge function can be naturally extended to permutations and posets.

3.1 The combinatorial Hopf algebra of permutations \mathcal{S}

Let \mathcal{S}_n denote the \mathbf{k} -span of all permutations of $[n]$. An element $\sigma \in \mathcal{S}_n$ is given by a list $(\sigma(1), \sigma(2), \dots, \sigma(n))$. By \emptyset , we denote the unique permutation of the empty set \emptyset . With the operations defined as follows, the vector space $\mathcal{S} = \bigoplus_{n \geq 0} \mathcal{S}_n$ has a structure of a graded connected Hopf algebra, as it is shown in [2].

For the unit, we take the map $u : 1 \mapsto \emptyset$. The graded multiplication is given by

$$\pi \otimes \sigma \mapsto \pi\sigma \text{ for } \pi \in \mathcal{S}_m, \sigma \in \mathcal{S}_n,$$

where $\pi\sigma$ denotes the permutation $(\pi(1), \dots, \pi(m), \sigma(1) + m, \dots, \sigma(n) + m)$, and linearly extended on $\mathcal{S}_m \otimes \mathcal{S}_n \rightarrow \mathcal{S}_{m+n}$.

The graded comultiplication on \mathcal{S} is given with

$$\sigma \mapsto \sum_{\substack{(S,T) \models [m+n] \\ |S|=m, |T|=n}} \text{st}(\sigma|_S) \otimes \text{st}(\sigma|_T), \text{ for } \sigma \in \mathcal{S}_{m+n},$$

where st denotes the standardization map, and linearly extended on \mathcal{S}_{m+n} . We define the counit to be $\epsilon(\emptyset) = 1$.

Equipped with these operations, \mathcal{S} becomes a graded connected bialgebra, and hence a graded connected Hopf algebra [2]. It is cocommutative, but not commutative. Nevertheless, we will not follow [2] for the definition of character. We want to define a CHA structure accordant with the structure of CHA on digraphs.

If σ is a permutation on $[n]$, we say that a permutation $\pi \in S_n$ is σ -reversing if $\pi(i) > \pi(i+1)$, or $\sigma(\pi(i)) > \sigma(\pi(i+1))$ for every $i \in [n-1]$. Finally, by taking $\zeta(\sigma)$ to be the number of σ -reversing permutations, we obtain a multiplicative functional on \mathcal{S} .

3.2 The combinatorial Hopf algebra of posets \mathcal{P}

Let \mathcal{P}_n denote the \mathbf{k} -span of all partial orders on the set $[n] = \{1, 2, \dots, n\}$. The relation that induces poset P will be denoted as $<_P$. For the graded vector space $\mathcal{P} = \bigoplus_{n \geq 0} \mathcal{P}_n$, it is possible to define the structure of combinatorial Hopf algebra in many ways, see [1]. We define the operations on \mathcal{P} as in [2].

The unit is the map $u : \mathbf{k} \rightarrow \mathcal{P}$ given by $1 \mapsto \emptyset$, where we denoted by \emptyset the unique poset on the empty set. The graded multiplication $\mathcal{P}_m \otimes \mathcal{P}_n \rightarrow \mathcal{P}_{m+n}$ is the linear extension of the operation given by

$$P \otimes Q \mapsto PQ \text{ for } P \in \mathcal{P}_m, Q \in \mathcal{P}_n,$$

where PQ denotes the poset on $[m+n]$ with $<_{PQ}$ defined as follows: $<_{PQ}$ coincides with $<_P$ on $[m]$, $m+x <_{PQ} m+y$ iff $x <_Q y$ and $i <_{PQ} j$ for every $i \in [m]$ and every $j \in \{m+1, m+2, \dots, m+n\}$.

The graded comultiplication on \mathcal{P} is given by

$$P \mapsto \sum_{\substack{(S,T) \models [m+n] \\ |S|=m, |T|=n}} \text{st}(P|_S) \otimes \text{st}(P|_T),$$

where $P|_X$ denotes the restriction of relation $<_P$ on X , while st denotes the standardization of poset, and linearly extended on \mathcal{P}_{m+n} . We define the counit to be $\epsilon(\emptyset) = 1$.

With operations defined this way, \mathcal{P} becomes a graded connected Hopf algebra that is cocommutative, but not commutative, see [2]. For the same reason as for the CHA of permutations defined above, for the character $\zeta(P)$, we will take the number of linear extensions of P .

3.3 The connection between CHAs \mathcal{S} , \mathcal{P} and \mathcal{D}

There is a natural connection between the three combinatorial Hopf algebras from the previous text.

Theorem 3.1. *CHA \mathcal{S} is a combinatorial Hopf subalgebra of CHA \mathcal{P} .*

Proof. Define

$$f_n : \mathcal{S}_n \rightarrow \mathcal{P}_n, \text{ with } \sigma \mapsto P_\sigma,$$

where P_σ is a poset on $[n]$ given by $i <_{P_\sigma} j$ if $i < j$ and $\sigma(i) < \sigma(j)$. By extending linearly this map, we obtain an injective morphism of graded Hopf algebras, $f : \mathcal{S} \rightarrow \mathcal{P}$, see [2]. Hence, we only need to check that the respective characters are compatible.

However, σ -reversing permutations are exactly the linear extensions of P_σ^* , the dual poset of P_σ . Since a poset P and its dual P^* have the same number of linear extensions, we see that the characters agree. \square

On the other hand, we cannot say that \mathcal{P} is a combinatorial Hopf subalgebra of \mathcal{D} . This is due to the fact that for the generators of \mathcal{P} we have taken all posets on the set $[n]$, while for the generators of \mathcal{D} we have taken only all isomorphism classes of digraphs on the set $[n]$. Yet, if we define \mathcal{D} to be generated by all digraphs on the set $[n]$, which is only a technical difference, the following theorem would yield an embedding of \mathcal{P} into \mathcal{D} .

Theorem 3.2. *There is a natural morphism of combinatorial Hopf algebras*

$$g : \mathcal{P} \rightarrow \mathcal{D}.$$

Proof. Define

$$g_n : \mathcal{P}_n \rightarrow \mathcal{D}_n, \text{ with } P \mapsto [D_P],$$

where $[D_P]$ is an isomorphism class of a digraph D_P on the vertex set $[n]$ defined by the following condition: (i, j) is an edge in D_P if $i <_P j$. It is straightforward to check that these maps respect multiplication, comultiplication, unit, counit and grading. Notice that $\overline{D_P}$, the complementary digraph of D_P , is actually $g_n(P^*)$ with addition that it doesn't contain loops and that two incomparable elements of P (equivalently, of P^*) are connected by a double edge. Hence, Hamiltonian paths of $\overline{D_P}$ are exactly the linear extensions of P^* . Since P and P^* have the same number of linear extensions, the characters are compatible. \square

These two morphisms allow us to naturally define the Redei-Berge function for permutations and posets as the Redei-Berge function of their respective digraphs.

Definition 3.3. Let σ be a permutation. The Redei-Berge function U_σ of σ is

$$U_\sigma = U_{g(f(\sigma))}.$$

Let P be a poset. The Redei-Berge function U_P of P is

$$U_P = U_{g(P)}.$$

Since f and g are homomorphisms of CHAs, we see immediately that, according to Theorem 2.3, these functions are exactly the images of σ and P by the universal homomorphism of CHAs \mathcal{S} and \mathcal{P} respectively to $QSym$.

Theorem 3.4. *If $\Psi : \mathcal{S} \rightarrow QSym$ is the universal morphism of combinatorial Hopf algebras, then for a permutation σ ,*

$$\Psi(\sigma) = U_\sigma.$$

If $\Psi : \mathcal{P} \rightarrow QSym$ is the universal morphism of combinatorial Hopf algebras, then for a poset P ,

$$\Psi(P) = U_P.$$

In fact, we could define our morphism g differently. Namely, another natural way to get a connection between posets and digraphs is to assign a digraph to a poset by taking (u, v) as an edge if $v <_P u$. However, this would not change the Redei-Berge function of the poset, since $U_X = U_{X^{op}}$ for any digraph X by Theorem 2.4. Similarly, since $g(P)$ and $g(P^*)$ are two opposite digraphs, we have the following.

Lemma 3.5. *If P is a poset and P^* its dual poset, then*

$$U_P = U_{P^*}.$$

Consequently, the number of minimal and the number of maximal elements of some poset P cannot be uncovered from U_P .

Notice that permutations and posets always induce digraphs without non-trivial directed cycles (acyclic digraphs). Also, the collection of acyclic digraphs is closed under the multiplication and comultiplication in \mathcal{D} and, hence, forms a subalgebra of \mathcal{D} .

Theorem 3.6. *If D is an acyclic digraph, then U_D is p -positive. Consequently, for any permutation σ and any poset P , U_σ and U_P are p -positive.*

Proof. This follows immediately from Theorem 2.2, since if D has no non-trivial cycles, $\varphi(\pi) = 0$ for any listing π . \square

Example 3.7. 1) Let L_n denote the natural linear order on the set $[n]$. It is obvious that $L_n = L_{n-1} \cdot L_1$, hence $U_{L_n} = \Psi(L_n^n) = \Psi(L_1)^n = U_{L_1}^n = (F_\emptyset)^n = p_{(1, \dots, 1)} = h_n$. Notice that we could conclude the same by using Theorem 2.2. Since D_{L_n} has no non-trivial cycles and $\overline{D_{L_n}}$ has no cycles, it follows that $\Sigma_V(D_{L_n}, \overline{D_{L_n}}) = \{\text{Id}\}$. In Id , all cycles have length 1, hence we obtain $U_{L_n} = p_{(1, \dots, 1)}$.

2) Let D_n denote the discrete order on the set $[n]$. Then, D_{D_n} has no edges. Directly from the Definition 5, we get that $U_{D_n} = n!F_\emptyset$. On the other hand, since D_{D_n} has only trivial cycles and $\overline{D_{D_n}}$ has all possible nontrivial cycles, $\Sigma_V(D_{D_n}, \overline{D_{D_n}}) = S_n$, and by Theorem 2.2, we have that $U_{D_n} = \sum_{\pi \in S_n} p_{\text{type}(\pi)}$.

4 Some further properties of the Redei-Berge function

Motivated by some properties of the chromatic symmetric function, in the sequel we will further investigate the Redei-Berge symmetric function.

4.1 Decomposition techniques for digraphs

The Redei-Berge polynomial of a digraph, as an analogue of the chromatic polynomial of a graph, is defined in [8].

Definition 4.1. The *Redei-Berge polynomial* of a digraph X is the principal specialization of its Redei-Berge function:

$$u_X(m) = \text{ps}^1(U_X)(m) = U_X(\underbrace{1, \dots, 1}_{m \text{ ones}}, 0, 0, \dots).$$

The Redei-Berge polynomial, just like the chromatic polynomial, satisfies the deletion-contraction property, see [8]. The same property does not hold for the Redei-Berge function since the contraction of an edge gives a digraph with less vertices than the original digraph. However, there is a way to overcome this difficulty and express the Redei-Berge function of a digraph X in terms of the same function of some more simple digraphs. Similar decompositions for the chromatic symmetric function can be found in [6] and [10].

Theorem 4.2. (*k-deletion*): If e_1, e_2, \dots, e_k is a list of edges that form a directed k -cycle in a digraph $X = (V, E)$, then

$$U_X = \sum_{\substack{S \subseteq \{e_1, e_2, \dots, e_k\} \\ S \neq \emptyset}} (-1)^{|S|-1} U_{X \setminus S}. \quad (7)$$

Proof. If π is a V -listing, then there exists an edge e_l of the cycle that is not "detected" by $X\text{Des}(\pi)$. The subsets of $\{e_1, e_2, \dots, e_k\}$ that do not contain e_l are in bijection with the subsets of $\{e_1, e_2, \dots, e_k\}$ that contain e_l by $S \mapsto S \cup \{e_l\}$. Notice that S and $S \cup \{e_l\}$ induce the same descent set assigned to π . However, in

$$\sum_{\substack{S \subseteq \{e_1, e_2, \dots, e_k\} \\ S \neq \emptyset}} (-1)^{|S|-1} U_{X \setminus S},$$

they have opposite signs, hence, they cancel each other. Therefore, the only term that remains is the one for $S = \{e_l\}$. It is paired with the one for \emptyset , which is exactly the term that appears on the left side of equation (7). \square

Corollary 4.3. (*triple-deletion*): If e_1, e_2, e_3 is a list of edges that form a triangle in a digraph $X = (V, E)$, then

$$U_X = U_{X \setminus \{e_1\}} + U_{X \setminus \{e_2\}} + U_{X \setminus \{e_3\}} - U_{X \setminus \{e_1, e_2\}} - U_{X \setminus \{e_2, e_3\}} - U_{X \setminus \{e_3, e_1\}} + U_{X \setminus \{e_1, e_2, e_3\}}.$$

The key part of the previous proof was the fact that if a digraph contains a cycle, then for every listing of its vertices, there exists an edge not detected by the descent set of that listing. This condition, however, holds for every digraph different from a disjoint union of paths.

Theorem 4.4. *If $X = (V, E)$ is a digraph that is not a disjoint union of paths, then*

$$U_X = \sum_{S \subseteq E, S \neq \emptyset} (-1)^{|S|-1} U_{X \setminus S}. \quad (8)$$

The previous property can be a very powerful tool for calculating the Redei-Berge function of some particular digraph. There is no an analogous feature of the chromatic function. On the other hand, the chromatic function is multiplicative on the disjoint union of two graphs, since it is an image of the universal morphism of CHA of graphs where multiplication is given by disjoint union of graphs. The same property does not hold for the Redei-Berge function.

4.2 Properties preserved by the Redei-Berge function

Motivated by Stanley's question from [15], to what extent chromatic symmetric function distinguishes non-isomorphic graphs, many researchers have studied the properties of graphs that can be determined from their chromatic function, [9], [10] etc. In this subsection, we will try to answer some similar questions concerning the Redei-Berge function.

We do not expect that the Redei-Berge function differs non-isomorphic digraphs (equivalently, that $\Psi : \mathcal{D} \rightarrow Sym$ is injective). We already noticed that $U_X = U_{X^{op}}$ and X and X^{op} are not necessarily isomorphic. Even for $n = 2$, there is an n -vertex digraph X that is not isomorphic to its opposite digraph : $X = (\{1, 2\}, \{(1, 1), (1, 2)\})$. Also, adding new loops does not affect the Redei-Berge function. However, we can say the following.

Theorem 4.5. *Two digraphs having the same Redei-Berge function have the same number of vertices and the same number of edges that are not loops.*

Proof. The first part is obvious since for digraph $X = (V, E)$, U_X is homogeneous of degree $|V|$. By definition, $U_X = \sum_{\pi \in \Sigma_V} F_{X^{Des}(\pi)} = \sum_{S \subseteq [n-1]} c_S F_S$. If (i, j) is an edge of X , there are $(n-1)!$ permutations of $[n]$ in which i and j are consecutive. On the other hand, in the expansion of U_X , F_S "detects" $|S|$ distinct edges. Hence, $(n-1)!|E| = \sum_{S \subseteq [n-1]} c_S |S|$, and therefore

$$|E| = \frac{\sum_{S \subseteq [n-1]} c_S |S|}{(n-1)!}.$$

□

Corollary 4.6. *Two posets having the same Redei-Berge function have the same number of elements and the same number of comparable pairs.*

In fact, the previous two results can be obtained from the following for $k = 1$ and $k = 2$.

Theorem 4.7. *If two digraphs with n vertices have the same Redei-Berge function, they also have the same number of paths of length k for any $k \in [n]$. Consequently, their longest paths have the same length.*

Proof. If v_1, v_2, \dots, v_k is a path of length k in $X = (V, E)$ with $|V| = n$, then the vertices v_1, v_2, \dots, v_k appear in this order in $(n - k + 1)!$ listings of V . By definition, $U_X = \sum_{\pi \in \Sigma_V} F_{X \text{Des}(\pi)} = \sum_{S \subseteq [n-1]} c_S F_S$. In this expansion, F_S "detects" λ_S^k paths of length k , where

$$\lambda_S^k = \#\{i \in [n-1] \mid i, i+1, \dots, i+k-1 \in S\}.$$

Notice that λ_S^k depends only on S and k and not on X . If the number of paths of length k in X is denoted by P_X^k , it follows that

$$(n - k + 1)! P_X^k = \sum_{S \subseteq [n-1]} c_S \lambda_S^k.$$

Therefore,

$$P_X^k = \frac{\sum_{S \subseteq [n-1]} c_S \lambda_S^k}{(n - k + 1)!}.$$

□

Corollary 4.8. *If two posets have the same Redei-Berge function, they have the same number of chains of length k for any $k \in [n]$. Accordingly, their longest chains have the same length.*

From Corollary 4.6, we see that the number of incomparable pairs in some poset P can be uncovered from U_P . This also follows directly from Theorem 2.2, since

$$[p_{(2,1,\dots,1)}]U_P = \#\{(k, l) \mid k < l \text{ and } k \text{ and } l \text{ are incomparable in } P\},$$

due to the fact that D_P does not contain cycles of length 2.

Now we will focus our attention on some poset P that is a disjoint union of at most 4 incomparable chains L_x, L_y, L_z, L_w , where $x \geq y \geq z \geq w \geq 0$. From what we have said so far, we see that the following parameters can be revealed from U_P : the length of the longest chain $a = x$, the number of elements $b = x + y + z + w$, the number of comparable pairs $c = \binom{x}{2} + \binom{y}{2} + \binom{z}{2} + \binom{w}{2}$ and the number of incomparable pairs $d = xy + xz + xw + yz + yw + zw$. However, since $2c = b^2 - 2d - b$, these four equations are dependent, and do not necessarily induce a unique solution. Nevertheless, if we set $w = 0$, we get the following.

Theorem 4.9. *If two posets P and Q that are both disjoint unions of at most three incomparable chains have the same Redei-Berge function, then they are isomorphic.*

The previous theorem can be further generalized if we exploit Corollary 4.8 in a more detailed manner.

Theorem 4.10. *If two posets P and Q that are both disjoint unions of incomparable chains have the same Redei-Berge function, then they are isomorphic.*

Proof. Let $(1^{r_1}, 2^{r_2}, \dots, k^{r_k})$ and $(1^{s_1}, 2^{s_2}, \dots, l^{s_l})$ be the lists of lengths of chains of P and Q respectively. According to Corollary 4.8, the longest paths of P and Q have the same length, hence $k = l$ and, furthermore, $r_k = s_k$.

If we now suppose that we have already proved that $r_i = s_i$ for every $i \in \{m+1, m+2, \dots, k\}$, we are able to deduce that the same thing holds for $i = m$ as well. Namely, we know from Corollary 4.8 that the numbers of chains of length m in P and Q are the same. However, since there are $\binom{n}{m}$ subchains of length m in any chain of length n for $n \geq m$, this implies that

$$r_m + \binom{m+1}{m} r_{m+1} + \dots + \binom{k}{m} r_k = s_m + \binom{m+1}{m} s_{m+1} + \dots + \binom{k}{m} s_k,$$

which gives $r_m = s_m$, as desired. Hence, $r_i = s_i$ for every i , and therefore, P and Q are isomorphic. \square

In the same manner, we can obtain the following.

Theorem 4.11. *If two digraphs that are both disjoint unions of paths have the same Redei-Berge function, then they are isomorphic.*

The importance of disjoint unions of paths comes from the following. According to the Theorem 4.4, the Redei-Berge function of any digraph that is not a disjoint union of paths can be expressed as a linear combination of the Redei-Berge functions of appropriate subdigraphs. The same expansion can be applied again to any digraph appearing on the right side of the Equation 8 that is not a disjoint union of paths. If we continue with this procedure, we will be able to express the Redei-Berge function of the original digraph as a linear combination of the Redei-Berge functions of its spanning subdigraphs that are disjoint unions of paths.

Finally, there is one more thing we can say about a special class of digraphs.

Definition 4.12. A *tournament* is a loopless digraph $X = (V, E)$ such that whenever u and v are two distinct vertices in V , exactly one of (u, v) and (v, u) is an edge of X .

Theorem 4.13. *If two tournaments have the same Redei-Berge function, then they have the same number of k -cycles for any odd $k \geq 3$.*

Proof. If $X = (V, E)$ is a tournament and (u_1, u_2, \dots, u_k) is a k -cycle in X , then (u_k, \dots, u_2, u_1) is a k -cycle in \overline{X} and vice versa. According to Theorem 2.2, the number of k -cycles in X is exactly $\frac{[p_{(k,1,\dots,1)}]U_X}{2}$. \square

The previous theorem can not be expanded to the case of cycles of an even length since the terms appearing in Theorem 2.2 would cancel each other out.

4.3 Some new bases of Sym

In this section, motivated by the result from [4], we provide abundance of new algebraic bases for the algebra of symmetric functions, whose generators are Redei-Berge functions.

Definition 4.14. A digraph $X = (V, E)$ is irreducible if there do not exist non-trivial digraphs X_1 and X_2 such that $X = X_1 \cdot X_2$.

Notice that digraphs that are reducible do not have Hamiltonian cycles. The same holds for their complements since if $X = X_1 \cdot X_2$, then $\overline{X} = \overline{X_2} \cdot \overline{X_1}$.

Theorem 4.15. Let $(X_n)_{n \in \mathbb{N}_0}$ be a list of digraphs such that:

- 1) X_n has n vertices
- 2) at least one of X_n and $\overline{X_n}$ has a Hamiltonian cycle if n is odd
- 3) the numbers of Hamiltonian cycles in X_n and in $\overline{X_n}$ are different if n is even.

Then $(U_{X_n})_{n \in \mathbb{N}_0}$ is an algebraic basis of Sym .

Proof. Let $h(X)$ denote the number of Hamiltonian cycles in digraph X . According to Theorem 2.2, if n is odd, $[p_n]U_{X_n} = h(X_n) + h(\overline{X_n})$. Similarly, if n is even, $[p_n]U_{X_n} = h(\overline{X_n}) - h(X_n)$. If the conditions 2) and 3) are fulfilled, these numbers are non-zero.

For $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$, define $X_\lambda = X_{\lambda_1} \cdots X_{\lambda_k}$. The statement of this theorem is equivalent with the condition that the family $\{U_{X_\lambda}\}$ is a linear basis of Sym , since

$$U_{X_\lambda} = U_{X_{\lambda_1}} \cdots U_{X_{\lambda_k}}.$$

From Theorem 2.2 it follows that in the expansion of U_{X_λ} in the p -basis, there can only appear p_μ for $\mu \leq \lambda$. Here, $\mu \leq \lambda$ denotes the usual dominance order on partitions, given by $\mu_1 + \cdots + \mu_j \leq \lambda_1 + \cdots + \lambda_j$ for any j .

Therefore, $(U_{X_\lambda})_{\lambda \vdash n}$ induces a triangular transition matrix to the power sum basis, with non-zero entries on the main diagonal. Hence, $(U_{X_\lambda})_{\lambda \vdash n}$ is a linear basis of Sym as well. □

As we have already noticed, reducible digraphs can not appear in this list. Nevertheless, the conditions given in the previous theorem are not too strong. For example, we could take a discrete digraph on n vertices as X_n .

Theorem 4.16. There are infinitely many lists $(X_n)_{n \in \mathbb{N}_0}$ of loopless digraphs such that $(U_{X_n})_{n \in \mathbb{N}_0}$ is an algebraic basis of Sym .

Proof. For $n \in \mathbb{N}$, where $n \neq 0$ and $n \neq 1$, we are going to construct $n - 1$ non-isomorphic digraphs with n vertices such that each of these digraphs satisfies the conditions of the previous theorem. Moreover, their Redei-Berge functions will also be different.

For $k \in \{0, 1, \dots, n - 2\}$, let $X_{n,k}$ denote the digraph with the vertex set $[n]$ and with edges $(1, 1 + i)$ for $1 \leq i \leq k$. This digraph, obviously, does not contain a Hamiltonian cycle, but its complement $\overline{X_{n,k}}$ has at least one such cycle - $(n, n - 1, \dots, 2, 1, n)$. Hence, $X_{n,k}$ satisfies the conditions of the previous theorem for any $k \in \{0, 1, \dots, n - 2\}$. Moreover, since $X_{n,k}$ has exactly k edges, by Theorem 4.5, $X_{n,k}$ and $X_{n,l}$ for $k \neq l$ have different Redei-Berge functions. □

5 Further avenues

The Redei-Berge function is a symmetric function associated to digraphs, introduced in the 1990s. Unlike its peer, the chromatic symmetric function, it did not attract attention of many researchers until recently. In this paper, we tried to derive some of its properties that, to a certain extent, resemble properties of the chromatic symmetric function.

A natural avenue to pursue is to give an expansion of the Redei-Berge function in other bases of the space of symmetric functions and to examine its positivity in these bases. For now, it is not quite clear how this could be achieved. One may use formulas given in Theorems 2.2 and 4.4 to obtain such expansions. Though, the path of finding combinatorial interpretations of coefficients in other bases might be completely independent of these formulas. This would enable us to explicitly calculate the Redei-Berge function and the corresponding polynomial of a larger class of digraphs. We could also try to get an insight into the relationship between the standard bases of Sym and the new bases described in the previous section. Finally, we can seek some further properties of digraphs that are preserved by the Redei-Berge function.

As we have already mentioned, permutations and posets necessarily induce acyclic digraphs. This fact could imply some additional nice properties of their Redei-Berge function. In [15], Stanley gives combinatorial connection between the coefficients of the chromatic function of a graph in its expansion in elementary basis and the number of acyclic orientations of that graph, with a certain number of *sinks*. In our terminology, the sinks represent the maximal elements of the poset, and their counterparts represent its minimal elements. However, we have seen in Lemma 3.5 that the number m_P of minimal and the number M_P of maximal elements of poset P cannot be revealed from U_P . Nevertheless, we were not able to find a counterexample dealing with the value $m_P + M_P$.

Conjecture 5.1. If two posets P and Q have the same Redei-Berge function, then $m_P + M_P = m_Q + M_Q$.

References

- [1] M. Aguiar, N. Bergeron, F. Sottile, Combinatorial Hopf algebras and generalized Dehn-Sommerville relations, *Compositio Mathematica* 142 (2006) 1-30.
- [2] J. C. Aval, N. Bergeron, J. Machacek, New invariants for permutations, orders and graphs, *Advances in Applied Mathematics* 121, (2020), 102080
- [3] C. Berge, *Graphs and Hypergraphs*, North-Holland Mathematical Library 6, 2nd edition, North-Holland 1976.
- [4] S. Cho, S. van Willigenburg, Chromatic bases for symmetric functions, *Electron. J. Combin.* 23, 6pp. (2016)

- [5] T. Y. Chow, The Path-Cycle Symmetric Function of a Digraph, *Adv. Math.* 118 (1996), 71-98.
- [6] S. Dahlberg and S. van Willigenburg, Lollipop and lariat symmetric functions, *SIAM J. Discrete Math.* 32, 1029-1039 (2018).
- [7] D. Grinberg, R. Stanley, The Redei-Berge symmetric function of a directed graph, [arXiv:2307.05569v1](https://arxiv.org/abs/2307.05569v1)
- [8] V. Grujić, T. Stojadinović, The Redei-Berge Hopf algebra of digraphs, [arXiv:2402.07606v2](https://arxiv.org/abs/2402.07606v2), accepted in *Periodica Mathematica Hungarica*
- [9] J. Martin, M. Morin and J. Wagner, On distinguishing trees by their chromatic symmetric functions, *J. Combin. Theory Ser. A* 115, 237-253 (2008).
- [10] R. Orellana and G. Scott, Graphs with equal chromatic symmetric function, *Discrete Math.* 320, 1-14 (2014).
- [11] L. Redei, Ein kombinatorischer Satz, *Acta Litteraria Szeged* 7 (1934), 39-43.
- [12] R. Stanley, The X-Descent set of a permutation, *Algorithmic and Enumerative Combinatorics*, Vienna, 2022, available at: <https://math.mit.edu/~rstan/transparencies/vienna.pdf>
- [13] R. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge Univ. Press, Cambridge, 1999.
- [14] G. Wiseman, Enumeration of paths and cycles and e -coefficients of incomparability graphs, [arXiv:0709.0430v1](https://arxiv.org/abs/0709.0430v1)
- [15] R. Stanley, A symmetric function generalization of the chromatic polynomial of a graph, *Adv. Math.* 111, 166-194 (1995).

6 Declarations and Statements

The authors declare that no funds, grants, or other support were received during the preparation of this manuscript. The authors have no relevant financial or non-financial interests to disclose. All authors contributed to the study conception and design. All authors read and approved the final manuscript.