

# EXISTENCE OF MULTISOLITON SOLUTIONS OF THE GRAVITATIONAL HARTREE EQUATION IN THREE DIMENSIONS

YUTONG WU

ABSTRACT. We consider the three-dimensional gravitational Hartree equation. We prove the existence of multisoliton solutions with gravitational interaction of hyperbolic, parabolic and hyperbolic-parabolic types, generalizing and improving the result by Krieger, Martel and Raphaël [8] which dealt with two soliton solutions.

## 1. INTRODUCTION

**1.1. Background.** In 1927, soon after the Schrödinger equation was proposed, Douglas Hartree derived the Hartree equation, which provided a way to study many body quantum systems. It has then attracted the interest of both physicists and mathematicians.

In this paper, we consider the gravitational Hartree equation in 3D

$$(1.1) \quad iu_t + \Delta u - \phi_{|u|^2} u = 0,$$

where  $u : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{C}$  and

$$\phi_{|u|^2} = \Delta^{-1}(|u|^2) = -\frac{1}{4\pi|x|} * |u|^2.$$

We begin with some properties of the equation.

The equation possesses a large family of symmetries. Namely, if  $u$  solves (1.1), then for any  $(t_0, \alpha_0, \beta_0, \lambda_0, \gamma_0) \in \mathbb{R} \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}_+ \times (\mathbb{R}/2\pi\mathbb{Z})$ ,

$$(1.2) \quad v(t, x) = \lambda_0^2 u(\lambda_0^2 t + t_0, \lambda_0 x - \alpha_0 - \beta_0 t) e^{i(\frac{1}{2}\beta_0 \cdot x - \frac{1}{4}|\beta_0|^2 t + \gamma_0)}$$

also solves (1.1). In view of Noether's theorem, we expect the equation to have some conservation laws. The following quantities are conserved by the equation:

$$\begin{aligned} \text{Mass:} & \quad \int |u(t, x)|^2 dx, \\ \text{Momentum:} & \quad \text{Im} \int \nabla u(t, x) \overline{u(t, x)} dx, \\ \text{Hamiltonian:} & \quad \int |\nabla u(t, x)|^2 dx - \frac{1}{2} \int |\nabla \phi_{|u|^2}(t, x)|^2 dx. \end{aligned}$$

In other words, if  $u$  solves (1.1), then these quantities are independent of  $t$ .

Consider  $u_\lambda(x) = \lambda^2 u(\lambda x)$  for  $\lambda > 0$  as a special case of (1.2). Then  $\frac{d}{d\lambda} \|u_\lambda\|_{L^2} \geq 0$ , which means (1.1) is (mass) subcritical. By Theorem 6.1.1 in [3], we know the Cauchy problem of (1.1) is wellposed in  $H^1$ . To be more precise, for any  $u_0 \in H^1(\mathbb{R}^3)$ , there exists a unique  $u \in C(\mathbb{R}; H^1(\mathbb{R}^3))$  satisfying (1.1) and  $u(0, x) = u_0(x)$ . Moreover, this  $u$  depends on  $u_0$  continuously.

There is a special type of solutions called solitary waves. A solitary wave is a solution to (1.1) of the form  $u(t, x) = e^{it} W(x)$ . We deduce that  $W$  satisfies  $\Delta W - \phi_{|W|^2} W = W$ . From [10] we know there exists a unique radial and positive solution  $Q$  of

$$(1.3) \quad \Delta Q - \phi_{Q^2} Q = Q,$$

called the ground state. One of the properties of  $Q$  is the exponential decay:

$$(1.4) \quad Q(x) \leq Ce^{-c|x|}, \quad \forall x \in \mathbb{R}^3.$$

It is proved in [11] that this  $Q$  can also be characterized as the radial minimizer of the Hamiltonian subject to a given  $L^2$  norm. More precisely, it minimizes

$$\int |\nabla u|^2 - \frac{1}{2} \int |\nabla \phi_{|u|^2}|^2,$$

among all  $u \in H^1$  with the same  $L^2$  norm as  $Q$ . Using (1.2) we can construct a family of ground state solitary waves. The ground state solitary wave is orbitally stable as proved in [4].

The next object of interest is the multisolitary wave, which can be roughly understood as the sum of several solitary waves.

Multisolitary waves are important because they reflect the long time dynamics of the solution. For nonlinear dispersive PDEs such as the generalized Korteweg-de Vries equation (gKdV), the nonlinear Schrödinger equation (NLS), and the nonlinear wave equation (NLW), it was conjectured that a generic solution evolves asymptotically as the sum of a radiation term (which behaves like the solution to the linear equation) and multisolitary waves. There are substantial materials on the subject of soliton resolution. We list [20], [21], [2], [15], [14], [5] and [7] for reference.

However, the long time dynamics for the Hartree equation is largely unknown. One of the difficulties is the so-called long range effect of (1.1). For some other equations such as (NLS), the sum of two ground state solitary waves moving away at a constant speed solves the equation up to a term that decays exponentially in time. But for the Hartree equation, since

$$\phi_{Q^2}(x) \sim \frac{1}{|x|} \quad \text{as } x \rightarrow \infty,$$

the sum of two solitary waves does not satisfy the equation in a satisfactory sense. We will give a quantitative estimate of such errors in Lemma 2.3.

**1.2. The main result.** As a starting point of the study of long time dynamics, Krieger, Martel and Raphaël [8] studied the existence of 2-soliton solutions of (1.1). This pioneer paper revealed that one should expect a gravitational 2-body interaction within the two solitons.

Based on their method, we generalize their result to  $N$ -soliton solutions, which, as one would expect, involve an  $N$ -body interaction. We will also improve their result in the parabolic dynamic. Our result seems a satisfactory counterpart of [15] and [14], which dealt with the existence of multisolitary waves of (NLS) and (gKdV), respectively.

While  $N$  is the accustomed letter for the  $N$ -body problem, for the sake of other notations in this paper, we shall use  $m$  hereinafter.

To begin with, we recall the equation of the  $m$ -body problem.

**Definition (m-body problem).**

Let  $m \geq 2$ . The  $m$ -body problem is an ODE system

$$(1.5) \quad \dot{\alpha}_j(t) = 2\beta_j(t), \quad \dot{\beta}_j(t) = - \sum_{k \neq j} \frac{\|Q\|_{L^2}^2}{4\pi\lambda_k} \cdot \frac{\alpha_j(t) - \alpha_k(t)}{|\alpha_j(t) - \alpha_k(t)|^3}, \quad \forall 1 \leq j \leq m,$$

where  $\alpha_j, \beta_j \in C^1(\mathbb{R}, \mathbb{R}^3)$  and  $\lambda_j \in \mathbb{R}_+$  for  $1 \leq j \leq m$ .

In this paper, we will consider the following three types of solutions.

- **hyperbolic:** for all  $j \neq k$ , we have  $\lim_{t \rightarrow +\infty} \frac{|\alpha_j(t) - \alpha_k(t)|}{t} \in (0, +\infty)$ .
- **parabolic:** for all  $j \neq k$ , we have  $\lim_{t \rightarrow +\infty} \frac{|\alpha_j(t) - \alpha_k(t)|}{t^{2/3}} \in (0, +\infty)$ .

- *hyperbolic-parabolic: both cases appear, and they are the only cases.*

In striking contrast to the 2-body problem where solutions are classified, the  $m$ -body problem for  $m \geq 3$  is known to be unsolvable. Even for 3-body, chaotic dynamics may occur [18]. Despite of the difficulty, many papers [6], [12], [1], [13], [19] investigated the existence of the above types of solutions. We summarize the main results in [13] and [19] as follows.

Consider the sets of configuration centered at the origin

$$\mathcal{X} = \left\{ (x_1, \dots, x_m) \in \mathbb{R}^{3m} \mid \sum_{j=1}^m \lambda_j^{-1} x_j = 0 \right\},$$

$$\mathcal{Y} = \left\{ (x_1, \dots, x_m) \in \mathcal{X} \mid x_j \neq x_k, \forall j \neq k \right\} \quad \text{and} \quad \Delta = \mathcal{X} \setminus \mathcal{Y}.$$

We say  $(x_1, \dots, x_m) \in \mathcal{Y}$  is **minimal** if

$$\sum_{j < k} \frac{1}{\lambda_j \lambda_k |y_j - y_k|} \geq \sum_{j < k} \frac{1}{\lambda_j \lambda_k |x_j - x_k|}$$

for any  $(y_1, \dots, y_m) \in \mathcal{Y}$  such that

$$\sum_{j=1}^m \lambda_j^{-1} |y_j|^2 = \sum_{j=1}^m \lambda_j^{-1} |x_j|^2.$$

**Theorem** (Thm 1.1, [13]; Thm 1.6, 1.7, 1.8, [19]). *Given  $\lambda_j \in \mathbb{R}_+$  for  $1 \leq j \leq m$ .*

- (1) *There exists a hyperbolic solution to (1.5) of the form*

$$(1.6) \quad \alpha_j(t) = a_j t + o(t) \quad \text{as } t \rightarrow +\infty$$

*for any  $(a_1, \dots, a_m) \in \mathcal{Y}$  and initial configuration in  $\mathcal{X}$ .*

- (2) *There exists a parabolic solution to (1.5) of the form*

$$(1.7) \quad \alpha_j(t) = c b_j t^{\frac{2}{3}} + o(t^{\frac{1}{3}+}) \quad \text{as } t \rightarrow +\infty$$

*for any minimal  $(b_1, \dots, b_m) \in \mathcal{Y}$  and initial configuration in  $\mathcal{X}$ , where  $c > 0$  is determined by  $b_1, \dots, b_m$ .*

- (3) *There exists a hyperbolic-parabolic solution to (1.5) of the form*

$$(1.8) \quad \alpha_j(t) = a_j t + c_j b_j t^{\frac{2}{3}} + o(t^{\frac{1}{3}+}) \quad \text{as } t \rightarrow +\infty$$

*for any  $(a_1, \dots, a_m) \in \Delta$ , minimal  $(b_1, \dots, b_m) \in \mathcal{Y}$  and initial configuration in  $\mathcal{X}$ , where  $c_j > 0$  is determined by  $a_1, \dots, a_m, b_1, \dots, b_m$  and  $c_j = c_k$  whenever  $a_j = a_k$ .*

For hyperbolic-parabolic solutions, the relation on  $\{1, 2, \dots, m\}$  given by  $a_j = a_k$  is an equivalence relation. Let  $M$  denote the set of equivalent classes. For  $J \in M$ , let  $\alpha_J$  be any of  $\{\alpha_j \mid j \in J\}$  and  $\beta_J$  be any of  $\{\beta_j \mid j \in J\}$ . Then we have

$$\begin{aligned} |\alpha_j(t)| &\lesssim t, & |\alpha_j(t) - \alpha_J(t)| &\lesssim t^{\frac{2}{3}}, & \forall j \in J, \\ |\beta_j(t)| &\lesssim 1, & |\beta_j(t) - \beta_J(t)| &\lesssim t^{-\frac{1}{3}}, & \forall j \in J. \end{aligned}$$

The results in this paper assert the existence of multisoliton solutions to (1.1) reproducing the above three non-trapped dynamics. For the last two cases, an assumption on the masses of the bodies is needed.

Before stating our results, we introduce some notations. When there are  $\alpha_j, \beta_j, \lambda_j$  and  $\gamma_j$ , we will denote

$$(1.9) \quad \begin{aligned} \alpha &= (\alpha_1, \dots, \alpha_m), & \beta &= (\beta_1, \dots, \beta_m), \\ \lambda &= (\lambda_1, \dots, \lambda_m), & \gamma &= (\gamma_1, \dots, \gamma_m), \\ P &= (\alpha, \beta, \lambda), & g_j &= (\alpha_j, \beta_j, \lambda_j, \gamma_j), & g &= (P, \gamma), \\ \alpha_{jk} &= \alpha_j - \alpha_k, & \beta_{jk} &= \beta_j - \beta_k, & a &= \min_{j \neq k} |\alpha_{jk}|. \end{aligned}$$

We use similar notation when  $\alpha_j, \beta_j, \lambda_j, \gamma_j$  depend on  $t$  or there are superscripts.

For  $u : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{C}$ , we define  $g_j u : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{C}$  by

$$(1.10) \quad g_j u(t, x) = \frac{1}{\lambda_j^2} u\left(t, \frac{x - \alpha_j}{\lambda_j}\right) e^{-i\gamma_j + i\beta_j \cdot x}.$$

The notation is valid when some components of  $g_j$  depend on  $t$  or  $u : \mathbb{R}^3 \rightarrow \mathbb{C}$ .

Then we are ready to state the main results of this paper.

**Theorem 1.** *Let  $P^\infty(t)$  be a hyperbolic solution to the  $m$ -body problem (1.5) of the form (1.6). Then there exists a solution  $u$  to (1.1) and  $\gamma^\infty(t)$  that is  $C^1$  in  $t$  such that*

$$\lim_{t \rightarrow +\infty} \left\| u(t, \cdot) - \sum_{j=1}^m g_j^\infty Q(t, \cdot) \right\|_{H^1} = 0.$$

**Theorem 2.** *Let  $P^\infty(t)$  be a parabolic solution to the  $m$ -body problem (1.5) of the form (1.7). Assume  $\exists \lambda^\infty > 0$  such that  $\lambda_j^\infty = \lambda^\infty, \forall j$ . Then there exists a solution  $u$  to (1.1) and  $\gamma^\infty(t)$  that is  $C^1$  in  $t$  such that*

$$\lim_{t \rightarrow +\infty} \left\| u(t, \cdot) - \sum_{j=1}^m g_j^\infty Q(t, \cdot) \right\|_{H^1} = 0.$$

**Theorem 3.** *Let  $P^\infty(t)$  be a hyperbolic-parabolic solution to the  $m$ -body problem (1.5) of the form (1.8). Assume  $\exists \lambda_j^\infty > 0, J \in M$  such that  $\lambda_j^\infty = \lambda_j^\infty, \forall j \in J$ . Then there exists a solution  $u$  to (1.1) and  $\gamma^\infty(t)$  that is  $C^1$  in  $t$  such that*

$$\lim_{t \rightarrow +\infty} \left\| u(t, \cdot) - \sum_{j=1}^m g_j^\infty Q(t, \cdot) \right\|_{H^1} = 0.$$

**Remark.**

1. Theorem 2 improves the result in [8] as we take  $\alpha_j$  (in their statement) to be identical to  $\alpha_j^\infty$ , which trivially answers their Comment 2. This improvement is possible because of a cancellation of errors displayed in the proof of Proposition 5.1 (the construction of  $\tilde{P}$ ).

2. The assumption that  $(b_1, \dots, b_m)$  is minimal is not directly used when proving Theorem 2 and 3. But this assumption is needed in [19] to guarantee the existence of solutions of the  $m$ -body problem.

3. The elliptic case, the uniqueness of  $u$ , and the stability of multisolitary waves are some of the open problems after this work. They are actually open even for 2-solitons of Hartree. On the other hand, some of these questions were solved for other dispersive PDEs. For instance, [14] studied the uniqueness for  $(gKdV)$ , and [16], [17] studied the stability for  $(gKdV)$ ,  $(NLS)$ , respectively.

As we largely used the method in [8], the structure of the paper will also be similar, except we will first deal with the hyperbolic case, and then the other two cases.

Compared to [8], the difficulty of the  $m$ -soliton problem lies mainly in the parabolic and hyperbolic-parabolic cases. We need to study an approximate system of the  $m$ -body problem, which is essentially harder than the 2-body problem. For this purpose, we have to perform delicate computation of the constants involved.

The article is organized as follows. In Section 2, we construct approximate multisolitary solutions of (1.1) up to the  $N$ -th order for any  $N \geq 1$  to overcome the long range effect. We then focus on the hyperbolic problem. We reduce the problem to a uniform estimate and furthermore a modulation estimate in Section 3. Then the modulation estimate is proved in Section 4, finishing the proof of Theorem 1. Finally, in section 5, we modify some arguments to prove Theorem 2 and 3 and improve the conclusion in [8]. The only place where we need the assumption on the masses is pointed out at the end.

## 2. APPROXIMATE SOLUTIONS

As preparation, we do some basic calculation. Assume  $u = g_j v$  and the components of  $g_j$  may depend on  $t$ . More precisely, we assume

$$u(t, x) = \frac{1}{\lambda_j(t)^2} v \left( t, \frac{x - \alpha_j(t)}{\lambda_j(t)} \right) e^{-i\gamma_j(t) + i\beta_j(t) \cdot x}.$$

Then we have

$$(2.1) \quad \begin{aligned} u_t &= \frac{1}{\lambda_j^4} \left( \lambda_j^2 v_t - \lambda_j \dot{\alpha}_j \cdot \nabla v - \dot{\lambda}_j (x - \alpha_j) \cdot \nabla v - 2\lambda_j \dot{\lambda}_j v \right. \\ &\quad \left. - i\lambda_j^2 \dot{\gamma}_j v + i\lambda_j^2 \dot{\beta}_j \cdot x v \right) e^{-i\gamma_j + i\beta_j \cdot x}, \\ \nabla u &= \frac{1}{\lambda_j^3} (\nabla v + i\lambda_j \beta_j v) e^{-i\gamma_j + i\beta_j \cdot x}, \\ \Delta u &= \frac{1}{\lambda_j^4} (\Delta v + 2i\lambda_j \beta_j \cdot \nabla v - \lambda_j^2 |\beta_j|^2 v) e^{-i\gamma_j + i\beta_j \cdot x}, \\ \phi_{|u|^2}(x) &= \frac{1}{\lambda_j^2} \phi_{|v|^2} \left( \frac{x - \alpha_j}{\lambda_j} \right). \end{aligned}$$

Therefore, if we let

$$u(t, x) := \sum_{j=1}^m u_j(t, x) := \sum_{j=1}^m g_j v_j(t, x),$$

and set  $y_j = \frac{x - \alpha_j(t)}{\lambda_j(t)}$ ,  $\Lambda v_j = 2v_j + y_j \cdot \nabla v_j$ , then

$$iu_t + \Delta u - \phi_{|u|^2} u = \sum_{j=1}^m \frac{1}{\lambda_j^4} E_j(t, y_j) e^{-i\gamma_j + i\beta_j \cdot x} - \sum_{k \neq j} \phi_{\text{Re}(u_k \bar{u}_j)} u,$$

where

$$\begin{aligned} E_j(t, y_j) &= i\lambda_j^2 \partial_t v_j + \Delta v_j - v_j - i\lambda_j \dot{\lambda}_j \Lambda v_j - \lambda_j^3 \dot{\beta}_j \cdot y_j v_j \\ &\quad - i\lambda_j (\dot{\alpha}_j - 2\beta_j) \cdot \nabla v_j + \lambda_j^2 \left( \dot{\gamma}_j + \frac{1}{\lambda_j^2} - |\beta_j|^2 - \dot{\beta}_j \cdot \alpha_j \right) v_j \\ &\quad - \left[ \phi_{|v_j|^2} + \sum_{k \neq j} \left( \frac{\lambda_j}{\lambda_k} \right)^2 \phi_{|v_k|^2} \left( t, \frac{\lambda_j}{\lambda_k} y_j + \frac{\alpha_{jk}}{\lambda_k} \right) \right] v_j. \end{aligned}$$

To be clear, the space variable of the right hand side is  $y_j$  unless explicitly written out.

**2.1. Definition of approximate solutions.** We need to approximate the last term of  $E_j$ . Since

$$\left(\frac{\lambda_j}{\lambda_k}\right)^2 \phi_{|v_k|^2} \left(t, \frac{\lambda_j}{\lambda_k} y_j + \frac{\alpha_{jk}}{\lambda_k}\right) = -\frac{\lambda_j^2}{4\pi\lambda_k} \int_{\mathbb{R}^3} \frac{|v_k(t, \xi)|^2}{|\lambda_j y_j + \alpha_{jk} - \lambda_k \xi|} d\xi,$$

we consider the Taylor expansion

$$\frac{1}{|\alpha - \zeta|} = \sum_{n=1}^N F_n(\alpha, \zeta) + O\left(\frac{|\zeta|^{N+1}}{|\alpha|^{N+1}}\right) \quad \text{as } \zeta \rightarrow 0,$$

where  $F_n(\alpha, \zeta)$  is homogeneous of degree  $-n$  in  $\alpha$  and of degree  $n-1$  in  $\zeta$ . We define the approximation to be

$$\begin{aligned} \phi_{|v_k|^2}^{(N)}(t, y_j) &:= \sum_{n=1}^N \psi_{|v_k|^2}^{(n)}(t, y_j) \\ &:= \sum_{n=1}^N -\frac{\lambda_j^2}{4\pi\lambda_k} \int_{\mathbb{R}^3} |v_k(t, \xi)|^2 F_n(\alpha_{jk}, \lambda_k \xi - \lambda_j y_j) d\xi. \end{aligned}$$

Explicit formulae for the first few terms are as follows:

$$\begin{aligned} \psi_{|v_k|^2}^{(1)}(t, y_j) &= -\frac{\lambda_j^2}{4\pi\lambda_k|\alpha_{jk}|} \int |v_k(t, \xi)|^2 d\xi, \\ \psi_{|v_k|^2}^{(2)}(t, y_j) &= -\frac{\lambda_j^2}{4\pi\lambda_k|\alpha_{jk}|^3} \int \left( \lambda_k(\alpha_{jk} \cdot \xi) |v_k(t, \xi)|^2 - \lambda_j(y_j \cdot \alpha_{jk}) |v_k(t, \xi)|^2 \right) d\xi. \end{aligned}$$

We will need to take  $v_k = Q$ . In this case we denote  $\psi_{|v_k|^2}^{(n)}$  by  $\psi_{Q^2, k}^{(n)}$ . Namely,

$$\psi_{Q^2, k}^{(n)}(y_j) = -\frac{\lambda_j^2}{4\pi\lambda_k} \int_{\mathbb{R}^3} Q^2(\xi) F_n(\alpha_{jk}, \lambda_k \xi - \lambda_j y_j) d\xi.$$

We shall let  $v_j$  vary in  $N$ , and we also assume  $v_j$  depends on time only through the parameters  $t \mapsto P(t)$ , which means  $v_j(t, y_j) = V_j^{(N)}(P(t), y_j)$  for some  $V_j^{(N)}$ .

Define

$$(2.2) \quad R_g^{(N)}(t, x) := \sum_{j=1}^m R_{j, g}^{(N)}(t, x) := \sum_{j=1}^m g_j V_j^{(N)}(P(t), x).$$

Let us omit the subscript  $g$  of  $R^{(N)}$  for now. We have

$$\begin{aligned} (2.3) \quad & i\partial_t R^{(N)} + \Delta R^{(N)} - \phi_{|R^{(N)}|^2} R^{(N)} \\ &= \sum_{j=1}^m \frac{1}{\lambda_j^4} E_j^{(N)}(t, y_j) e^{-i\gamma_j + i\beta_j \cdot x} - \sum_{k \neq j} \phi_{\operatorname{Re}(R_k^{(N)} \overline{R_j^{(N)}})} R^{(N)} \\ &+ \sum_{j=1}^m \frac{1}{\lambda_j^4} \sum_{k \neq j} \left[ \phi_{|V_k^{(N)}|^2} - \left(\frac{\lambda_j}{\lambda_k}\right)^2 \phi_{|V_k^{(N)}|^2} \left(P(t), \frac{\lambda_j}{\lambda_k} y_j + \frac{\alpha_{jk}}{\lambda_k}\right) \right] V_j^{(N)} e^{-i\gamma_j + i\beta_j \cdot x}, \end{aligned}$$

where

$$\begin{aligned} E_j^{(N)} &= i\lambda_j^2 \partial_t V_j^{(N)} + \Delta V_j^{(N)} - V_j^{(N)} - i\lambda_j \dot{\lambda}_j \Lambda V_j^{(N)} - \lambda_j^3 \dot{\beta}_j \cdot y_j V_j^{(N)} \\ &\quad - i\lambda_j (\dot{\alpha}_j - 2\beta_j) \nabla V_j^{(N)} + \lambda_j^2 \left( \dot{\gamma}_j + \frac{1}{\lambda_j^2} - |\beta_j|^2 - \dot{\beta}_j \cdot \alpha_j \right) V_j^{(N)} \\ &\quad - \left( \phi_{|V_j^{(N)}|^2} + \sum_{k \neq j} \phi_{|V_k^{(N)}|^2} \right) V_j^{(N)}. \end{aligned}$$

For functions  $M_j^{(N)}(P)$  and  $B_j^{(N)}(P)$  of the parameters, we can decompose

$$E_j^{(N)} = \tilde{E}_j^{(N)} + S_j^{(N)},$$

where

$$\begin{aligned} \tilde{E}_j^{(N)}(t, y_j) &= \Delta V_j^{(N)} - V_j^{(N)} - i\lambda_j M_j^{(N)} \Lambda V_j^{(N)} - \lambda_j^3 B_j^{(N)} \cdot y_j V_j^{(N)} \\ (2.4) \quad &\quad + i\lambda_j^2 \sum_{k=1}^m \left( \frac{\partial V_j^{(N)}}{\partial \alpha_k} \cdot 2\beta_k + \frac{\partial V_j^{(N)}}{\partial \beta_k} \cdot B_k^{(N)} + \frac{\partial V_j^{(N)}}{\partial \lambda_k} M_k^{(N)} \right) \\ &\quad - \left( \phi_{|V_j^{(N)}|^2} + \sum_{k \neq j} \phi_{|V_k^{(N)}|^2} \right) V_j^{(N)} \end{aligned}$$

and

$$\begin{aligned} S_j^{(N)}(t, x) &= -i\lambda_j (\dot{\alpha}_j - 2\beta_j) \nabla V_j^{(N)} - \lambda_j^3 (\dot{\beta}_j - B_j^{(N)}) \cdot y_j V_j^{(N)} \\ (2.5) \quad &\quad - i\lambda_j (\dot{\lambda}_j - M_j^{(N)}) \Lambda V_j^{(N)} + \lambda_j^2 \left( \dot{\gamma}_j + \frac{1}{\lambda_j^2} - |\beta_j|^2 - \dot{\beta}_j \cdot \alpha_j \right) V_j^{(N)} \\ &\quad + i\lambda_j^2 \sum_{k=1}^m \left[ \frac{\partial V_j^{(N)}}{\partial \alpha_k} \cdot (\dot{\alpha}_k - 2\beta_k) + \frac{\partial V_j^{(N)}}{\partial \beta_k} \cdot (\dot{\beta}_j - B_j^{(N)}) \right. \\ &\quad \left. + \frac{\partial V_j^{(N)}}{\partial \lambda_k} (\dot{\lambda}_j - M_j^{(N)}) \right]. \end{aligned}$$

Note that  $\tilde{E}_j^{(N)}$  is set to be a function of  $y_j$  instead of  $x$ . This is to align with a later statement. It does not matter whether  $S_j^{(N)}$  is a function of  $y_j$  or  $x$ , but we will let it be a function of  $x$  for preciseness.

Next we show that we can choose  $V_j^{(N)}$ ,  $M_j^{(N)}$  and  $B_j^{(N)}$  so that  $\tilde{E}_j^{(N)}$  is small. This smallness will be a result of homogeneity, so we give the following definition.

**Definition (Admissible functions).**

Recalling (1.9), let  $\Omega$  denote the space of non-collision positions:

$$\Omega := \left\{ P = (\alpha, \beta, \lambda) \in \mathbb{R}^{3m} \times \mathbb{R}^{3m} \times \mathbb{R}_+^m \mid \alpha_j \neq \alpha_k, \forall j \neq k \right\}.$$

(1) Let  $n \in \mathbb{N}$ . Define  $S_n$  to be the set of functions  $\sigma : \Omega \rightarrow \mathbb{C}$  that is homogeneous in  $\alpha$  of degree  $-n$  and is a finite sum of

$$c \prod_{j \neq k} (\alpha_j - \alpha_k)^{p_{jk}} |\alpha_j - \alpha_k|^{-q_{jk}} \prod_{j=1}^m \beta_j^{k_j} \lambda_j^{l_j},$$

where  $c \in \mathbb{C}$ ,  $p_{jk} \in \mathbb{N}^3$ ,  $q_{jk} \in \mathbb{N}$ ,  $k_j \in \mathbb{N}^3$ ,  $l_j \in \mathbb{Z}$  and  $|p_{jk}| \leq q_{jk}$ .

(2) We say a function  $u : \Omega \times \mathbb{R}^3 \rightarrow \mathbb{C}$  is **admissible** if  $u$  is a finite sum of

$$\sigma(\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_m, \lambda_1, \dots, \lambda_m) \tau(x),$$

where  $\sigma \in S_n$  for some  $n \in \mathbb{N}$  and  $\tau \in C^\infty$  satisfies

$$|\nabla^k \tau(x)| \leq e^{-c_k |x|}, \quad \forall k \geq 0, x \in \mathbb{R}^3.$$

If  $n$  is the same for all addends, then we say  $u$  is admissible of degree  $n$ . Otherwise, taking  $n$  as the minimal one among all addends, we say  $u$  is admissible of degree  $\geq n$ .

Here are some properties of admissible functions.

**Lemma 2.1.** *Let  $n, m \in \mathbb{N}$  and  $u, v$  be admissible of degree  $n, m$ , respectively. Then*

- (1)  $\forall j$ ,  $\frac{\partial u}{\partial \alpha_j}$  is admissible of degree  $1 + n$ , and  $\frac{\partial u}{\partial \beta_j}, \frac{\partial u}{\partial \lambda_j}$  are admissible of degree  $n$ .
- (2)  $uv$  is admissible of degree  $n + m$ ;
- (3)  $\phi_u v$  is admissible of degree  $n + m$ ;
- (4)  $\forall k \geq 1$ ,  $\psi_u^{(k)} v$  is admissible of degree  $k + n + m$ ;
- (5)  $\forall N \geq 1$ ,  $\phi_u^{(N)} v$  is admissible of degree  $\geq 1 + n + m$ ;
- (6)  $\exists c > 0$  such that for any compact set  $K \subset \mathbb{R}^{3m} \times \mathbb{R}_+^m$ ,  $\exists C_K > 0$  such that

$$(2.6) \quad |u(\alpha, \beta, \gamma, x)| \leq \frac{C_K}{a^n} e^{-c|x|}, \quad \forall (\beta, \gamma) \in K,$$

where  $a = \min_{j \neq k} |\alpha_j - \alpha_k|$  as in (1.9).

The proof of these properties is direct so we shall omit it. The point of considering admissible functions is that according to (6), they decay rapidly when  $n$  is large.

Consider the linearized operators  $L_+, L_-$  around  $Q$  defined by

$$L_+ f := -\Delta f + f + \phi_Q^2 f + 2\phi_Q f Q, \quad L_- f := -\Delta f + f + \phi_Q^2 f.$$

By Theorem 4 in [9],  $\{\partial_1 Q, \partial_2 Q, \partial_3 Q\}$  spans  $\ker L_+$ , and  $\{Q\}$  spans  $\ker L_-$ . Moreover, Lemma 2.4 in [8] asserts that when restricted to admissible functions,  $\ker(L_\pm)^\perp$  is exactly the range of  $L_\pm$ . A precise statement is as follows.

**Lemma 2.2.** *Let  $n \in \mathbb{N}$  and  $f$  be real-valued and admissible of degree  $n$ .*

- (1) If  $(f, \nabla Q) = 0$ , then  $L_+ u = f$  has a real-valued solution  $u$  admissible of degree  $n$ .
- (2) If  $(f, Q) = 0$ , then  $L_- u = f$  has a real-valued solution  $u$  admissible of degree  $n$ .

Furthermore, if  $f$  is radial, then  $u$  can be chosen to be radial.

The following proposition constructs the approximate solutions.

**Proposition 2.1.** *For  $n \geq 1$  and  $1 \leq j \leq m$ , there exist real-valued  $m_j^{(n)}, b_j^{(n)} \in S_n$  and  $T_j^{(n)}$  that is admissible of degree  $n$  such that: for any  $N \geq 1$ , if setting*

$$V_j^{(N)}(t, y_j) = Q(y_j) + \sum_{n=1}^N T_j^{(n)}(P(t), y_j),$$

$$M_j^{(N)}(P) = \sum_{n=1}^N m_j^{(n)}(P) \quad \text{and} \quad B_j^{(N)}(P) = \sum_{n=1}^N b_j^{(n)}(P),$$

then  $\tilde{E}_j^{(N)}$  defined by (2.4) is admissible of degree  $\geq N + 1$ .



*Proof.* We construct the functions by induction in  $N$ .

For  $N = 1$ , we take  $m_j^{(1)} = b_j^{(1)} = 0$ . Suppose  $T_j^{(1)}$  is admissible of degree 1 and real-valued. By Lemma 2.1 and (1.3), we have

$$\begin{aligned} \tilde{E}_j^{(1)} &= \Delta V_j^{(1)} - V_j^{(1)} + i\lambda_j^2 \sum_{k=1}^m \frac{\partial V_j^{(1)}}{\partial \alpha_k} \cdot 2\beta_k - \left( \phi_{|V_j^{(1)}|^2} + \sum_{k \neq j} \psi_{|V_k^{(1)}|^2}^{(1)} \right) V_j^{(1)} \\ &= \Delta T_j^{(1)} - T_j^{(1)} - \phi_{Q^2} T_j^{(1)} - 2\phi_{QT_j^{(1)}} Q - \sum_{k \neq j} \psi_{Q^2, k}^{(1)} Q + error \\ &= -L_+ T_j^{(1)} + \sum_{k \neq j} \frac{\lambda_j^2 \|Q\|_{L^2}^2}{4\pi \lambda_k |\alpha_{jk}|} Q + error, \end{aligned}$$

where *error* is admissible of degree  $\geq 2$ . Since  $L_+(\Lambda Q) = -2Q$ , we may take

$$(2.7) \quad T_j^{(1)} = - \sum_{k \neq j} \frac{\lambda_j^2 \|Q\|_{L^2}^2}{8\pi \lambda_k |\alpha_{jk}|} \Lambda Q$$

to cancel the first two terms. This proves the conclusion when  $N = 1$ .

Next, we construct  $m_j^{(N+1)}, b_j^{(N+1)}$  and  $T_j^{(N+1)}$  from the first  $N$  terms. We have

$$\begin{aligned} \tilde{E}_j^{(N+1)} - \tilde{E}_j^{(N)} &= \Delta T_j^{(N+1)} - T_j^{(N+1)} - i\lambda_j m_j^{(N+1)} \Lambda Q - \lambda_j^3 b_j^{(N+1)} \cdot y_j Q \\ &\quad - \phi_{Q^2} T_j^{(N+1)} - 2\phi_{\operatorname{Re}(QT_j^{(N+1)})} Q - \sum_{k \neq j} \psi_{Q^2, k}^{(N+1)} Q + error \\ &= - \left( L_+ X_j^{(N+1)} + \lambda_j^3 b_j^{(N+1)} \cdot y_j Q + \sum_{k \neq j} \psi_{Q^2, k}^{(N+1)} Q \right) \\ &\quad - i \left( L_- Y_j^{(N+1)} + \lambda_j m_j^{(N+1)} \Lambda Q \right) + error, \end{aligned}$$

where

$$X_j^{(N+1)} = \operatorname{Re} T_j^{(N+1)}, \quad Y_j^{(N+1)} = \operatorname{Im} T_j^{(N+1)}$$

and

$$\begin{aligned}
error &= -i\lambda_j m_j^{(N+1)} \Lambda \left( V_j^{(N+1)} - Q \right) - i\lambda_j M_j^{(N)} \Lambda T_j^{(N+1)} \\
&\quad - \lambda_j^3 b_j^{(N+1)} \cdot y_j \left( V_j^{(N+1)} - Q \right) - \lambda_j^3 B_j^{(N)} \cdot y_j T_j^{(N+1)} \\
&\quad - \phi_{|V_j^{(N)}|^2 - Q^2} T_j^{(N+1)} - 2\phi_{\operatorname{Re}\left(V_j^{(N)} \overline{T_j^{(N+1)}}\right)} \left( V_j^{(N+1)} - Q \right) \\
&\quad - 2\phi_{\operatorname{Re}\left(\left(V_j^{(N)} - Q\right) \overline{T_j^{(N+1)}}\right)} Q - \phi_{|T_j^{(N+1)}|^2} V_j^{(N+1)} \\
&\quad - \sum_{k \neq j} \left( \psi_{|V_k^{(N)}|^2 - Q^2} V_j^{(N+1)} + \psi_{Q^2, k}^{(N+1)} \left( V_j^{(N+1)} - Q \right) + \phi_{|V_k^{(N)}|^2} T_j^{(N+1)} \right. \\
&\quad \quad \quad \left. + 2\phi_{\operatorname{Re}\left(V_k^{(N)} \overline{T_k^{(N+1)}}\right)} V_j^{(N+1)} + \phi_{|T_k^{(N+1)}|^2} V_j^{(N+1)} \right) \\
&\quad + i\lambda_j^2 \sum_{k=1}^m \left( \frac{\partial T_j^{(N+1)}}{\partial \alpha_k} \cdot 2\beta_k + \frac{\partial T_j^{(N+1)}}{\partial \beta_k} \cdot B_k^{(N)} + \frac{\partial V_j^{(N+1)}}{\partial \beta_k} \cdot b_k^{(N+1)} \right. \\
&\quad \quad \quad \left. + \frac{\partial T_j^{(N+1)}}{\partial \lambda_k} \cdot M_k^{(N)} + \frac{\partial V_j^{(N+1)}}{\partial \lambda_k} \cdot m_k^{(N+1)} \right).
\end{aligned}$$

Assume  $m_j^{(N+1)}, b_j^{(N+1)} \in S_{N+1}$  and  $T_j^{(N+1)}$  is admissible of degree  $N+1$ . Using Lemma 2.1, we see  $error$  is admissible of degree  $\geq N+2$ . Thus it suffices to require

$$\begin{cases} L_+ X_j^{(N+1)} = -\lambda_j^3 b_j^{(N+1)} \cdot y_j Q - \sum_{k \neq j} \psi_{Q^2, k}^{(N+1)} Q + \operatorname{Re} \hat{E}_j^{(N)}, \\ L_- Y_j^{(N+1)} = -\lambda_j m_j^{(N+1)} \Lambda Q + \operatorname{Im} \hat{E}_j^{(N)}, \end{cases}$$

where  $\hat{E}_j^{(N)}$  is the sum of terms in  $\tilde{E}_j^{(N)}$  that are admissible of degree  $N+1$ .

Recall that we let the right hand sides be functions of  $y_j$ , so they are admissible of degree  $N+1$ . By Lemma 2.2, it suffices to require

$$\begin{cases} \left( \lambda_j^3 b_j^{(N+1)} \cdot y_j Q + \sum_{k \neq j} \psi_{Q^2, k}^{(N+1)} Q - \operatorname{Re} \hat{E}_j^{(N)}, \nabla Q \right) = 0, \\ \left( \lambda_j m_j^{(N+1)} \Lambda Q - \operatorname{Im} \hat{E}_j^{(N)}, Q \right) = 0. \end{cases}$$

Such  $b_j^{(N+1)}$  and  $m_j^{(N+1)}$  exist because  $(y_j Q, \nabla Q) \neq 0$  and  $(\Lambda Q, Q) \neq 0$ .  $\square$

**2.2. Accuracy of approximate solutions.** We verify the accuracy of  $R^{(N)}$  as an approximate solution where  $V_j^{(N)}$  is determined in Proposition 2.1. We start with some estimates following from the definition of admissible functions.

Let  $\tilde{\Omega} = \Omega \times (\mathbb{R}/2\pi\mathbb{Z})^m$  denote the space of modulation parameters and  $g \in \tilde{\Omega}$ . If  $K$  is a compact set in  $\mathbb{R}^{3m} \times \mathbb{R}_+^m$  and  $(\beta, \gamma) \in K$ , then by (1.4) and (2.6), we have

$$(2.8) \quad |V_j^{(N)}| \leq C e^{-c|y_j|} + C_N a^{-1} e^{-c_N |y_j|},$$

which also yields

$$(2.9) \quad |R_{j,g}^{(N)}| \leq C e^{-c|x-\alpha_j|} + C_N a^{-1} e^{-c_N|x-\alpha_j|}.$$

By definition,  $R_g^{(N)}$  is  $C^1$  in  $g$ . Thus, if  $g' \in \Omega$  and  $(\beta', \gamma') \in K$ , then

$$(2.10) \quad \left\| R_g^{(N)} - R_{g'}^{(N)} \right\|_{H^1} \leq C_N \|g - g'\|.$$

Finally, since  $m_j^{(1)} = b_j^{(1)} = 0$ , we have

$$(2.11) \quad |M_j^{(N)}| \leq C a^{-2} + C_N a^{-3}, \quad |B_j^{(N)}| \leq C a^{-2} + C_N a^{-3}.$$

Here, the capital constants depend on  $K$ , while the little ones do not.

The next lemma consists of two localization properties. The first item shows that the cross term about  $R_j$  in (2.3) does not matter. The second item will be used later.

**Lemma 2.3.** *Let  $p \neq q \in \mathbb{R}^3$  and  $u, v$  be functions such that*

$$|u(x)| \leq e^{-|x-p|}, \quad |v(x)| \leq e^{-|x-q|}, \quad \forall x \in \mathbb{R}^3.$$

*Then there exist absolute constants  $C, c > 0$  such that:*

- (1)  $\|\phi_{uv}\|_{L^\infty} \leq C e^{-c|p-q|}$ ;
- (2) *If  $f \in L^2$ , then  $\|\phi_{fu}fv\|_{L^1} \leq C \max\left\{\frac{e^{-\frac{1}{2}|p-q|}}{|p-q|^{\frac{1}{2}}}, \frac{1}{|p-q|}\right\} \|f\|_{L^2}^2$ .*

*Proof.*

(1) Using the Hardy-Littlewood-Sobolev inequality, we get  $\|\phi_{uv}\|_{L^\infty} \leq C \|uv\|_{L^{3/2}}$ . Note that either  $|x-p| \geq \frac{1}{2}|p-q|$  or  $|x-q| \geq \frac{1}{2}|p-q|$ . In the first case, we use  $|u(x)| \leq e^{-c|p-q|}$  and  $\|v\|_{L^{3/2}} \leq C$  to conclude, and the second case is similar.

(2) We have

$$\|\phi_{fu}fv\|_{L^1} \leq C \iint \frac{|f(x)||f(y)|}{|x-y|} e^{-|x-p|-|y-q|} dx dy$$

The integral on the region  $|x-y| \geq \frac{1}{2}|p-q|$  is easily bounded by  $\frac{C}{|p-q|} \|f\|_{L^2}^2$ .

If  $|x-y| < \frac{1}{2}|p-q|$ , then  $|x-p| + |y-q| \geq \frac{1}{2}|p-q| + \frac{1}{4}|x-p|$ , so by Cauchy-Schwarz,

$$\begin{aligned} & \iint_{|x-y| < \frac{1}{2}|p-q|} \frac{|f(x)||f(y)|}{|x-y|} e^{-|x-p|-|y-q|} dx dy \\ & \leq e^{-\frac{1}{2}|p-q|} \int |f(x)| e^{-\frac{1}{4}|x-p|} \left( \int_{|y-x| \leq \frac{1}{2}|p-q|} \frac{|f(y)|}{|y-x|} dy \right) dx \\ & \leq C e^{-\frac{1}{2}|p-q|} |p-q|^{-\frac{1}{2}} \|f\|_{L^2} \int |f(x)| e^{-\frac{1}{4}|x-p|} dx \leq C \frac{e^{-\frac{1}{2}|p-q|}}{|p-q|^{\frac{1}{2}}} \|f\|_{L^2}^2. \end{aligned}$$

We then obtain the conclusion. □

The following is the main result in this subsection. It estimates the extent to which  $R_g^{(N)}$ , defined by (2.2), satisfies the Hartree equation (1.1).

**Proposition 2.2.** *Let  $c_0, C_0 > 0$  and suppose  $g \in C^1(\mathbb{R}_+, \tilde{\Omega})$  satisfies*

$$(2.12) \quad a \geq c_0, \quad |\beta| \leq C_0, \quad c_0 \leq \lambda_j \leq C_0.$$

Let  $V_j^{(N)}$ ,  $M_j^{(N)}$  and  $B_j^{(N)}$  be as in Proposition 2.1,  $R_g^{(N)}$  be defined by (2.2), and

$$(2.13) \quad \Psi^{(N)} = i\partial_t R_g^{(N)} + \Delta R_g^{(N)} - \phi_{|R_g^{(N)}|^2} R_g^{(N)} - \sum_{j=1}^m \frac{1}{\lambda_j^4} S_j^{(N)} e^{-i\gamma_j + i\beta_j \cdot x}.$$

Then there exist  $c, C > 0$  depending on  $c_0, C_0$  and  $N$  such that

$$(2.14) \quad |\Psi^{(N)}(t, x)| \leq \frac{C}{a^{N+1}(t)} \max_j e^{-c|x - \alpha_j(t)|}.$$

*Proof.* For simplicity, we omit the superscript  $N$  and the subscript  $g$ .

By (2.3), we have

$$\begin{aligned} \Psi &= \sum_{j=1}^m \frac{1}{\lambda_j^4} \tilde{E}_j(t, y_j) e^{-i\gamma_j + i\beta_j \cdot x} + \sum_{j \neq k} \phi_{\operatorname{Re}(R_j \bar{R}_k)} R \\ &+ \sum_{j=1}^m \frac{1}{\lambda_j^4} \sum_{k \neq j} \left[ \phi_{|V_k|^2} - \left( \frac{\lambda_j}{\lambda_k} \right)^2 \phi_{|V_k|^2} \left( P(t), \frac{\lambda_j}{\lambda_k} y_j + \frac{\alpha_{jk}}{\lambda_k} \right) \right] V_j e^{-i\gamma_j + i\beta_j \cdot x}. \end{aligned}$$

The first term is controlled using Proposition 2.1 and (2.6). The second term is controlled using Lemma 2.3 and (2.9). For the last term, we claim that

$$(2.15) \quad \left| \phi_{|V_k|^2} - \left( \frac{\lambda_j}{\lambda_k} \right)^2 \phi_{|V_k|^2} \left( \frac{\lambda_j}{\lambda_k} y_j + \frac{\alpha_{jk}}{\lambda_k} \right) \right| \leq \begin{cases} C(1 + |y_j|)^{N+1}, & \lambda_j |y_j| \geq \frac{|\alpha_{jk}|}{3}, \\ \frac{C(1 + |y_j|)^{N+1}}{|\alpha_{jk}|^{N+1}}, & \lambda_j |y_j| \leq \frac{|\alpha_{jk}|}{3}. \end{cases}$$

Since  $F_n$  is homogeneous of degree  $n$  in  $\zeta$ , using (2.8) and the Hardy-Littlewood-Sobolev inequality, we see the left hand side of (2.15) is always bounded by  $C(1 + |y_j|)^{N+1}$ , so we focus on the case when  $\lambda_j |y_j| \leq \frac{|\alpha_{jk}|}{3}$ . We write

$$\begin{aligned} LHS &= \frac{\lambda_j^2}{4\pi\lambda_k} \int |V_k(\xi)|^2 \left| \frac{1}{\alpha_{jk} - \lambda_k \xi + \lambda_j y_j} - \sum_{n=1}^N F_n(\alpha_{jk}, \lambda_k \xi - \lambda_j y_j) \right| d\xi \\ &= \frac{\lambda_j^2}{4\pi\lambda_k} \left( \int_{\lambda_k |\xi| \geq \frac{|\alpha_{jk}|}{3}} + \int_{\lambda_k |\xi| \leq \frac{|\alpha_{jk}|}{3}} \right) \triangleq \frac{\lambda_j^2}{4\pi\lambda_k} (I_1 + I_2). \end{aligned}$$

By Hardy-Littlewood-Sobolev, the definition of  $F_n$  and (2.8), we have

$$\begin{aligned} I_1 &\leq C \left\| V_k(\xi) \chi_{\{\lambda_k |\xi| \geq \frac{|\alpha_{jk}|}{3}\}} \right\|_{L^{8/3}}^2 + C \left\| |V_k(\xi)|^2 (1 + |\xi|)^N \chi_{\{\lambda_k |\xi| \geq \frac{|\alpha_{jk}|}{3}\}} \right\|_{L^1} \\ &\leq C e^{-c|\alpha_{jk}|} \leq \frac{C(1 + |y_j|)^{N+1}}{|\alpha_{jk}|^{N+1}}. \end{aligned}$$

By the Taylor formula,

$$\left| \frac{1}{|\alpha - \zeta|} - \sum_{n=1}^N F_n(\alpha, \zeta) \right| \leq C \frac{|\zeta|^{N+1}}{|\alpha|^{N+1}}, \quad \text{if } |\zeta| \leq \frac{|\alpha|}{3},$$

so by the assumption that  $\lambda_j |y_j| \leq \frac{|\alpha_{jk}|}{3}$  and (2.8), we have

$$I_2 \leq \frac{C(1 + |y_j|)^{N+1}}{|\alpha_{jk}|^{N+1}} \int |V_k(\xi)|^2 (1 + |\xi|)^{N+1} d\xi \leq \frac{C(1 + |y_j|)^{N+1}}{|\alpha_{jk}|^{N+1}}.$$

Thus (2.15) holds. Note that this yields

$$(2.16) \quad \left| \phi_{|V_k|^2}^{(N)} - \left( \frac{\lambda_j}{\lambda_k} \right)^2 \phi_{|V_k|^2} \left( \frac{\lambda_j}{\lambda_k} y_j + \frac{\alpha_{jk}}{\lambda_k} \right) \right| \leq \frac{C(1 + |y_j|)^{2N+2}}{a^{N+1}}.$$

Then by (2.8), we can control the first term.  $\square$

From the estimate (2.14), if  $N$  is large, then the error  $\Psi^{(N)}$  will decay rapidly in time. This helps us overcome the long range effect of (1.1). This is also why we need to construct the approximate solution. From now on, the strategy becomes also similar to that of [15].

### 3. REDUCTION OF THE PROBLEM

Now we will focus on the hyperbolic case. But we may still state the result in a more general way, for instance writing  $a(t)$  instead of  $t$ , so that it is easier to apply it to the other two cases.

In this section, we perform two steps of reduction of the hyperbolic problem.

**3.1. Uniform estimates.** Due to (2.13), we want  $S_j^{(N)}$  to vanish. Thus we need the following ODE result.

**Proposition 3.1.** *Let  $P^\infty$  be a hyperbolic solution to (1.5) of the form (1.6), and  $B_j^{(N)}, M_j^{(N)}$  be as determined in Proposition 2.1. Then there exist  $T_0 = T_0(N) > 0$  and  $P^{(N)} \in C^1([T_0, +\infty), \Omega)$  such that*

$$(3.1) \quad \begin{cases} \dot{\alpha}_j^{(N)}(t) = 2\beta_j^{(N)}(t), \\ \dot{\beta}_j^{(N)}(t) = B_j^{(N)}(P^{(N)}(t)), \\ \dot{\lambda}_j^{(N)}(t) = M_j^{(N)}(P^{(N)}(t)), \end{cases} \quad \forall t \geq T_0$$

and

$$(3.2) \quad \left\| P^{(N)}(t) - P^\infty(t) \right\| \leq t^{-1/2}, \quad \forall t \geq T_0.$$

We need the exact expression of  $b_j^{(2)}$ . Since  $T_j^{(1)}$  is real-valued, we have

$$\operatorname{Re} \hat{E}_j^{(1)} = -2\phi_{QT_j^{(1)}} T_j^{(1)} - \phi_{|T_j^{(1)}|^2} Q - \sum_{k \neq j} \left( 2\psi_{QT_k^{(1)}}^{(1)} Q + \psi_{Q^2, k}^{(1)} T_k^{(1)} \right).$$

Since  $T_j^{(1)}$  is even, by the explicit formula of  $\psi^{(1)}$ ,  $\operatorname{Re} \hat{E}_j^{(1)}$  is also even, and thus orthogonal to  $\nabla Q$ . We then obtain by the proof of Proposition 2.1 that

$$\left( \lambda_j^3 b_j^{(2)} \cdot y_j Q + \sum_{k \neq j} \psi_{Q^2, k}^{(2)} Q, \nabla Q \right) = 0.$$

By the explicit formula of  $\psi^{(2)}$  and using  $Q$  is even, we deduce

$$(3.3) \quad b_j^{(2)}(P) = - \sum_{k \neq j} \frac{\|Q\|_{L^2}^2 \alpha_{jk}}{4\pi \lambda_k |\alpha_{jk}|^3}.$$

Note that this is exactly the gravitational force acting on the  $j$ -th body. This explains the reason why we expect the  $m$ -body interaction and our choice of the coefficients. Also, (3.1) can be viewed as a perturbation of the  $m$ -body equation (1.5).

*Proof of Proposition 3.1.*

Let  $\epsilon = \frac{1}{10}$  and  $T_0 > 0$ . Define the norm  $\|\cdot\|_1$  of  $P \in C([T_0, +\infty), \Omega)$  by

$$\|P\|_1 := \sum_{j=1}^m \sup_{t \geq T_0} \left( t^{1-3\epsilon} |\alpha_j(t)| + t^{2-2\epsilon} |\beta_j(t)| + t^{1-\epsilon} |\lambda_j(t)| \right).$$

Let  $X = \left\{ P \in C([T_0, +\infty), \Omega) \mid \|P - P^\infty\|_1 \leq 1 \right\}$  and define  $\Gamma P(t)$  by

$$\begin{aligned} \Gamma \alpha_j(t) &= \alpha_j^\infty(t) + \int_t^\infty 2(\beta_j^\infty(\tau) - \beta_j(\tau)) d\tau, \\ \Gamma \beta_j(t) &= \lim_{t \rightarrow \infty} \beta_j^\infty(t) - \int_t^\infty B_j^{(N)}(P(\tau)) d\tau, \\ \Gamma \lambda_j(t) &= \lambda_j^\infty - \int_t^\infty M_j^{(N)}(P(\tau)) d\tau. \end{aligned}$$

Because of the decay of  $a$  in  $t$ , we know  $\lim_{t \rightarrow \infty} \beta_j^\infty(t)$  does exist.

We claim that: if  $T_0(N)$  is large enough, then  $\Gamma$  maps  $X$  into  $X$ , and for  $P, P' \in X$ , we have  $\|\Gamma P - \Gamma P'\|_1 \leq \frac{1}{2} \|P - P'\|_1$ .

Assume  $P \in X$ . Since  $P^\infty$  is hyperbolic, we have  $a(t) \gtrsim t$ . First,

$$|\Gamma \alpha_j(t) - \alpha_j^\infty(t)| \leq 2 \int_t^\infty |\beta_j(\tau) - \beta_j^\infty(\tau)| d\tau \leq 2 \int_t^\infty \tau^{2\epsilon-2} d\tau \leq C t^{2\epsilon-1}.$$

Using  $b_j^{(1)} = 0$  and  $b_j^{(2)}(P^\infty(t)) = \dot{\beta}_j^\infty(t)$ , which comes from (3.3), we have

$$|\Gamma \beta_j(t) - \beta_j^\infty(t)| \leq \int_t^\infty |b_j^{(2)}(P(\tau)) - b_j^{(2)}(P^\infty(\tau))| d\tau + \sum_{n=3}^N \int_t^\infty |b_j^{(n)}(P(\tau))| d\tau.$$

By the fundamental theorem of calculus, we have

$$|b_j^{(2)}(P(\tau)) - b_j^{(2)}(P^\infty(\tau))| \leq C \left( \frac{|\alpha - \alpha^\infty|}{a^3} + \frac{|\beta - \beta^\infty|}{a^2} + \frac{|\lambda - \lambda^\infty|}{a^2} \right) \leq C \tau^{\epsilon-3},$$

and thus, using  $b_j^{(n)} \in S_n$ , we get

$$|\Gamma \beta_j(t) - \beta_j^\infty(t)| \leq C \int_t^\infty \tau^{\epsilon-3} d\tau + C_N \sum_{n=3}^N \int_t^\infty \tau^{-n} d\tau \leq C_N t^{\epsilon-2}.$$

Using  $m_j^{(1)} = 0$  and  $m_j^{(n)} \in S_n$ , we have

$$|\Gamma \lambda_j(t) - \lambda_j^\infty(t)| \leq \sum_{n=2}^N \int_t^\infty |m_j^{(n)}(P(\tau))| d\tau \leq C_N \sum_{n=2}^N \int_t^\infty \tau^{-n} d\tau \leq C_N t^{-1}.$$

Collecting the above estimates, we get  $\|\Gamma P\|_1 \leq C_N T_0^{-\epsilon}$ .

Thus for  $T_0(N)$  large enough, we have  $\Gamma : X \rightarrow X$ . The contraction property can be checked in the same way. By the contraction mapping theorem,  $\Gamma$  has a unique fixed point in  $X$ . Taking this fixed point as  $P^{(N)}$ , then the requirements are satisfied.  $\square$

From Proposition 2.2 and 3.1, we know  $R_{g^{(N)}}^{(N)}$  is almost a solution of (1.1). We then reduce Theorem 1 to the following uniform estimate with a bootstrap assumption.

**Proposition 3.2.** *Let  $P^{(N)}$  be defined as in Proposition 3.1 and  $\gamma_j^{(N)}(t)$  be such that*

$$\gamma_j^{(N)}(0) = 0, \quad \dot{\gamma}_j^{(N)}(t) = -\frac{1}{\lambda_j^{(N)}(t)^2} + |\beta_j^{(N)}(t)|^2 + \dot{\beta}_j^{(N)}(t) \cdot \alpha_j^{(N)}(t).$$

Let  $T_n \rightarrow +\infty$  and  $u_n$  be the solution to

$$(3.4) \quad \begin{cases} i\partial_t u_n + \Delta u_n - \phi_{|u_n|^2} u_n = 0, \\ u_n(T_n, \cdot) = R_{g^{(N)}}^{(N)}(T_n, \cdot). \end{cases}$$

Then  $\exists T_0 = T_0(N)$  such that for  $N$  large and  $T_* \in [T_0, T_n]$ , if

$$(3.5) \quad \left\| u_n(t) - R_{g^{(N)}}^{(N)}(t) \right\|_{H^1} \leq 2t^{-\frac{N}{9}}, \quad \forall n \geq 1, \quad \forall t \in [T_*, T_n],$$

then

$$\left\| u_n(t) - R_{g^{(N)}}^{(N)}(t) \right\|_{H^1} \leq t^{-\frac{N}{9}}, \quad \forall n \geq 1, \quad \forall t \in [T_*, T_n].$$

*Proof of Theorem 1 by Proposition 3.2.*

Fix a large  $N$  such that the conclusion holds. By the standard bootstrap argument, we know (3.5) actually holds with  $T_* = T_0$ . Using (2.9), we know  $\exists C > 0$  such that

$$(3.6) \quad \|u_n(t)\|_{H^1} \leq C, \quad \forall n \geq 1, \quad \forall t \in [T_0, T_n].$$

Also, for any  $\delta > 0$ , there exist  $r = r(\delta) > 0$  and  $t_0 = t_0(\delta) > T_0$  such that

$$\int_{|x|>r} |u_n(t_0, x)|^2 dx < \delta, \quad \forall n \geq 1.$$

We claim that there exists  $r' = r'(\delta) > 0$  such that

$$(3.7) \quad \int_{|x|>r'} |u_n(T_0, x)|^2 dx < 2\delta, \quad \forall n \geq 1.$$

To prove (3.7), let  $\Phi \in C^\infty(\mathbb{R})$  be a cutoff such that

$$0 \leq \Phi \leq 1, \quad 0 \leq \Phi' \leq 2, \quad \Phi(x) = \begin{cases} 0, & x \leq 0, \\ 1, & x \geq 1. \end{cases}$$

Let  $L > 0$  and define  $z(t) = \int |u_n(t, x)|^2 \Phi\left(\frac{|x|-r}{L}\right) dx$ . Then  $z(t_0) \leq \delta$ . Since

$$z'(t) = -2\text{Im} \int \Delta u_n \overline{u_n} \Phi\left(\frac{|x|-r}{L}\right) dx = \frac{2}{L} \text{Im} \int \nabla u_n \overline{u_n} \cdot \frac{x}{|x|} \Phi'\left(\frac{|x|-r}{L}\right) dx,$$

we have  $|z'(t)| \leq \frac{4}{L} \|u_n\|_{H^1}^2$ . Integrating in  $t$  and using (3.6), we get

$$\int_{|x|>L+r} |u_n(T_0, x)|^2 dx \leq z(T_0) \leq \frac{4C^2(t_0 - T_0)}{L} + \delta.$$

We deduce (3.7) by taking  $L = L(\delta)$  large enough and  $r' = L + r$ .

Now, (3.6) and (3.7) imply the existence of a subsequence  $u_{n_k}(T_0)$  of  $u_n(T_0)$  that converges in  $L^2$  to some  $U_0$  and  $U_0 \in H^1$ . Let  $U$  be the solution to

$$\begin{cases} i\partial_t U + \Delta U - \phi_{|U|^2} U = 0, \\ U(T_0) = U_0. \end{cases}$$

By the well-posedness of (1.1), we have  $u_{n_k}(t) \rightarrow U(t)$  in  $L^2$  for any  $t \geq T_0$ . Thanks to (3.6), by passing to subsequence, we may assume  $u_{n_k}(t) \rightharpoonup U(t)$  in  $H^1$ . Using (3.5) and Fatou's lemma, we deduce

$$\left\| U(t) - R_{g^{(N)}}^{(N)}(t) \right\|_{H^1} \leq 2t^{-\frac{N}{9}}, \quad \forall t \geq T_0.$$

Let  $\gamma^\infty(t)$  be such that

$$\gamma_j^\infty(0) = 0, \quad \dot{\gamma}_j^\infty(t) = -\frac{1}{(\lambda_j^\infty)^2} + |\beta_j^\infty(t)|^2 + \dot{\beta}_j^\infty(t) \cdot \alpha_j^\infty(t).$$

Then by (2.10), (3.2) and (2.6), we obtain the conclusion of Theorem 1.  $\square$

**3.2. Modulation estimates.** We want to find a family of modulation parameters  $\alpha, \beta, \lambda$  and  $\gamma$  such that  $R_g^{(N)}$  is an orthogonal projection of  $u_n$ . More precisely, we prove the following lemma.

**Lemma 3.1.** *Let  $N, n \geq 1$ . Then there exist  $T_0 = T_0(N) > 0$  and a unique modulation parameter  $g \in C^1([T_0, +\infty), \tilde{\Omega})$  such that: if*

$$\varepsilon(t, x) = u_n(t, x) - R_g^{(N)}(t, x),$$

then for  $t \geq T_0$  and  $1 \leq j \leq m$ , we have

$$\begin{aligned} (3.8) \quad & \operatorname{Re}\left(\varepsilon(t), g_j V_j^{(N)}\right) = \operatorname{Re}\left(\varepsilon(t), g_j(y_j V_j^{(N)})\right) \\ & = \operatorname{Im}\left(\varepsilon(t), g_j(\Lambda V_j^{(N)})\right) = \operatorname{Im}\left(\varepsilon(t), g_j(\nabla V_j^{(N)})\right) = 0. \end{aligned}$$

In particular, we have

$$(3.9) \quad g(T_n) = g^{(N)}(T_n), \quad \varepsilon(T_n) = 0.$$

To prove the above result, we first work on a time-independent version.

**Lemma 3.2.** *Let  $N \geq 1$  and  $K \subset \mathbb{R}_+^m$  be compact. Then there exist  $\delta, A > 0$  such that: if  $g^0 \in \tilde{\Omega}$  and  $u \in H^1$  satisfy  $a^0 > A$ ,  $\lambda^0 \in K$  and  $\|u - R_{g^0}^{(N)}\|_{H^1} < \delta$ , then there exists a unique parameter  $g \in \tilde{\Omega}$  that  $C^1$ -depends on  $u$  and*

$$\begin{aligned} & \operatorname{Re}\left(u - R_g^{(N)}, g_j V_j^{(N)}\right) = \operatorname{Re}\left(u - R_g^{(N)}, g_j(y_j V_j^{(N)})\right) \\ & = \operatorname{Im}\left(u - R_g^{(N)}, g_j(\Lambda V_j^{(N)})\right) = \operatorname{Im}\left(u - R_g^{(N)}, g_j(\nabla V_j^{(N)})\right) = 0. \end{aligned}$$

*Proof.* Let  $p = (g, u)$  and  $\varepsilon(p) = u - R_g^{(N)}$ . Set  $u_0 = R_{g^0}^{(N)}$  and  $p_0 = (g_0, u_0)$ . Define

$$\begin{aligned} \rho_j^1(p) &= \operatorname{Re}\left(\varepsilon(p), g_j V_j^{(N)}\right), & \rho_j^2(p) &= \operatorname{Re}\left(\varepsilon(p), g_j(y_j V_j^{(N)})\right), \\ \rho_j^3(p) &= \operatorname{Im}\left(\varepsilon(p), g_j(\Lambda V_j^{(N)})\right), & \rho_j^4(p) &= \operatorname{Im}\left(\varepsilon(p), g_j(\nabla V_j^{(N)})\right). \end{aligned}$$

Then  $\varepsilon(p_0) = 0$  and  $\rho_j^\nu(p_0) = 0$  for  $\nu = 1, 2, 3, 4$ .



We would like to compute  $\frac{\partial \rho_j'}{\partial g}(p_0)$ . Since  $\varepsilon(p_0) = 0$ , the partial derivative of  $\rho_j'$  evaluated at  $p_0$  only falls on  $\varepsilon$ . We compute

$$\begin{aligned}\frac{\partial \varepsilon}{\partial \alpha_j} &= -\frac{1}{\lambda_j} g_j (\nabla V_j^{(N)}) + g_j \left( \frac{\partial V_j^{(N)}}{\partial \alpha_j} \right), \\ \frac{\partial \varepsilon}{\partial \beta_j} &= i \alpha_j g_j V_j^{(N)} + i \lambda_j g_j (y_j V_j^{(N)}) + g_j \left( \frac{\partial V_j^{(N)}}{\partial \beta_j} \right), \\ \frac{\partial \varepsilon}{\partial \lambda_j} &= g_j (\Lambda V_j^{(N)}) + g_j \left( \frac{\partial V_j^{(N)}}{\partial \lambda_j} \right), \\ \frac{\partial \varepsilon}{\partial \gamma_j} &= -i g_j V_j^{(N)}.\end{aligned}$$

Using (2.6), (2.8) and that  $Q$  is real and even, we can represent  $\frac{\partial \rho_j'}{\partial g}(p_0)$  by

$$\begin{array}{c|cccc} & 1 & 2 & 3 & 4 \\ \hline \alpha & 0 & 1 & 0 & 0 \\ \beta & 0 & 0 & * & 1 \\ \lambda & 1 & 0 & 0 & 0 \\ \gamma & 0 & 0 & 1 & 0\end{array}$$

where 0, for instance the  $(\alpha, 1)$  entry, represents

$$\frac{\partial \rho_j^1}{\partial \alpha_k}(p_0) = o(1), \quad \forall j, k,$$

while 1, for instance the  $(\alpha, 2)$  entry, represents

$$\frac{\partial \rho_j^2}{\partial \alpha_k}(p_0) = \begin{cases} o(1), & j \neq k, \\ c_j + o(1), & j = k. \end{cases}$$

Here  $c_j$  is invertible and independent of  $a$ , and  $o(1)$  means goes to 0 as  $a \rightarrow +\infty$ .

Therefore, for  $A$  large enough,  $\frac{\partial \rho_j'}{\partial g}(p_0)$  is an invertible matrix. Then we can conclude by the implicit function theorem. The last comment is that  $g \in \tilde{\Omega}$  because  $g$  is closed to  $g^{(N)}$ , which means we have  $a > \frac{A}{2}$  when  $\delta$  is small.  $\square$

*Proof of Lemma 3.1.*

By (3.5), for  $T_0(N)$  large enough and  $\delta, A$  determined in Lemma 3.2, if  $t \geq T_0$ , then  $a^{(N)}(t) > A$  and  $\|u_n(t) - R_{g^{(N)}}^{(N)}(t)\|_{H^1} < \delta$ . By Lemma 3.2, there exists a unique  $g(t) \in \tilde{\Omega}$  such that (3.8) holds. Moreover,  $g \in C^1$  because  $g$  is  $C^1$  in  $u_n$  and  $u_n$  is  $C^1$  in  $t$ .  $\square$

It follows from the implicit function theorem that  $g$  is closed to  $g^{(N)}$ . But to prove Proposition 3.2, we need a quantitative estimate of  $g - g^{(N)}$  and  $\varepsilon$ .

**Proposition 3.3.** For  $N$  and  $T_0 = T_0(N)$  large enough,  $\forall n \geq 1$ ,  $T_* \in [T_0, T_n]$ , if

$$(3.10) \quad \begin{cases} \|\varepsilon(t)\|_{H^1} \leq t^{-\frac{N}{4}}, \\ \sum_{j=1}^m \left| \lambda_j(t) - \lambda_j^{(N)}(t) \right| + \left| \beta_j(t) - \beta_j^{(N)}(t) \right| \leq t^{-1-\frac{N}{8}}, \\ \sum_{j=1}^m \left| \gamma_j(t) - \gamma_j^{(N)}(t) \right| + \left| \alpha_j(t) - \alpha_j^{(N)}(t) \right| \leq t^{-\frac{N}{8}}, \end{cases} \quad \forall t \in [T_*, T_n],$$

then

$$(3.11) \quad \begin{cases} \|\varepsilon(t)\|_{H^1} \leq \frac{1}{2}t^{-\frac{N}{4}}, \\ \sum_{j=1}^m \left| \lambda_j(t) - \lambda_j^{(N)}(t) \right| + \left| \beta_j(t) - \beta_j^{(N)}(t) \right| \leq \frac{1}{2}t^{-1-\frac{N}{8}}, \\ \sum_{j=1}^m \left| \gamma_j(t) - \gamma_j^{(N)}(t) \right| + \left| \alpha_j(t) - \alpha_j^{(N)}(t) \right| \leq \frac{1}{2}t^{-\frac{N}{8}}, \end{cases} \quad \forall t \in [T_*, T_n].$$

*Proof of Proposition 3.2 by Proposition 3.3.*

As the left hand sides are continuous in  $t$ , by a bootstrap argument, we know (3.10) actually holds for any  $t \in [T_0, T_n]$ . Then by (2.10), we have

$$\begin{aligned} \|u_n(t) - R_{g^{(N)}}^{(N)}\|_{H^1} &\leq \|u_n(t) - R_g^{(N)}\|_{H^1} + \|R_g^{(N)} - R_{g^{(N)}}^{(N)}\|_{H^1} \\ &\leq \|\varepsilon(t)\|_{H^1} + C_N \|g - g^{(N)}\| \leq t^{-\frac{N}{4}} + C_N t^{-\frac{N}{8}} \leq t^{-\frac{N}{9}} \end{aligned}$$

when  $T_0(N)$  is large enough. This finishes the proof of Proposition 3.2.  $\square$

So far, we have reduced Theorem 1 to Proposition 3.3, a modulation estimate with bootstrap assumptions.

#### 4. ESTIMATES OF THE MODULATION

To simplify notations, we write  $R_j = R_{j,g}^{(N)}$ ,  $R = R_g^{(N)}$ ,  $M_j = M_j^{(N)}$ ,  $B_j = B_j^{(N)}$ ,  $V_j = V_j^{(N)}$ ,  $S_j = S_j^{(N)}$  and  $\Psi = \Psi^{(N)}$ . But we will make clear whether a constant depends on  $N$ .

By (3.4) and (2.13), we have

$$(4.1) \quad i\partial_t \varepsilon + \Delta \varepsilon - 2\phi_{\text{Re}(\varepsilon \bar{R})} R - \phi_{|R|^2} \varepsilon = \mathcal{N}(\varepsilon) - \Psi - \sum_{j=1}^m \frac{1}{\lambda_j^4} S_j(t, x) e^{-i\gamma_j + i\beta_j \cdot x},$$

where

$$\mathcal{N}(\varepsilon) = 2\phi_{\text{Re}(\varepsilon \bar{R})} \varepsilon + \phi_{|\varepsilon|^2} R + \phi_{|\varepsilon|^2} \varepsilon.$$

By the Sobolev inequality and the Hardy-Littlewood-Sobolev inequality, we have

$$(4.2) \quad \|\mathcal{N}(\varepsilon)\|_{L^2} \leq C_N \|\varepsilon\|_{H^1}^2.$$

By (1.6), (3.2), (3.10) and (2.11), we have the following asymptotic properties:

$$(4.3) \quad a(t) \sim t, \quad |\alpha_j| \lesssim t, \quad |\beta_j| \lesssim 1, \quad |\dot{\beta}_j| \lesssim \frac{1}{t^2}, \quad \lambda_j \sim 1, \quad |\dot{\lambda}_j| \lesssim \frac{1}{t^2}.$$

In particular, (2.12) is satisfied.

**4.1. Control of the parameters.** In this subsection, we aim at proving the second and the third line of (3.11).

Define the modulation error

$$\begin{aligned} Mod(t) = \sum_{j=1}^m & \left( \left| \dot{\alpha}_j(t) - 2\beta_j(t) \right| + \left| \dot{\beta}_j(t) - B_j(P(t)) \right| + \left| \dot{\lambda}_j(t) - M_j(P(t)) \right| \right. \\ & \left. + \left| \dot{\gamma}_j(t) + \frac{1}{\lambda_j^2(t)} - |\beta_j(t)|^2 - \dot{\beta}_j(t) \cdot \alpha_j(t) \right| \right). \end{aligned}$$

First notice that  $S_j$  is controlled by  $Mod$ :

$$(4.4) \quad |S_j(t, x)| \leq C_N Mod(t) e^{-c_N |x - \alpha_j(t)|}.$$

Let  $\theta(t, x)$  be a function such that

$$(4.5) \quad |\theta(t, x)| \leq C e^{-c|x|} + C_N a^{-1} e^{-c_N |x|}, \quad \forall t > 0, x \in \mathbb{R}^3,$$

which corresponds to (2.8), (2.9), and let  $\theta_j = g_j \theta$ .

Using (4.1) and integration by parts, we can compute

$$\begin{aligned} \frac{d}{dt} \text{Im} \int \varepsilon \bar{\theta}_j &= \text{Re} \int \varepsilon \left( \overline{i \partial_t \theta_j + \Delta \theta_j - 2 \phi_{\text{Re}(\theta_j \bar{R})} R - \phi_{|R|^2} \theta_j} \right) \\ &+ \text{Re} \int (\Psi - \mathcal{N}(\varepsilon)) \bar{\theta}_j + \sum_{j=1}^m \text{Re} \int \frac{1}{\lambda_j^4} S_j(t, x) e^{-i\gamma_j + i\beta_j \cdot x} \bar{\theta}_j. \end{aligned}$$

By (2.1), (2.11) and (4.5), we have

$$\begin{aligned} i \partial_t \theta_j + \Delta \theta_j &= \frac{1}{\lambda_j^4} (i \lambda_j^2 \partial_t \theta + \Delta \theta - \theta) e^{-i\gamma_j + i\beta_j \cdot x} \\ &+ O \left( \frac{1}{a^2} + \frac{C_N}{a^3} + Mod \right) \left( e^{-c|x - \alpha_j|} + \frac{C_N}{a} e^{-c_N |x - \alpha_j|} \right). \end{aligned}$$

By Lemma 2.3, (2.9) and (4.5), we have

$$\begin{aligned} \phi_{\text{Re}(\theta_j \bar{R})} R &= \frac{1}{\lambda_j^4} \phi_{\text{Re}(\theta \bar{V}_j)} V_j e^{-i\gamma_j + i\beta_j \cdot x} + \phi_{\text{Re}(\theta_j \bar{R}_j)} \sum_{k \neq j} R_k \\ &+ O_N \left( e^{-c_N a} \max_k e^{-c_N |x - \alpha_k|} \right). \end{aligned}$$

By Lemma 2.3, (2.9), (4.5), (2.16) and the explicit formula of  $\psi^{(2)}$ , we have

$$\begin{aligned} \phi_{|R|^2} \theta_j &= \frac{1}{\lambda_j^4} \left( \phi_{|V_j|^2} + \sum_{k \neq j} \psi_{|V_k|^2}^{(1)} \right) \theta e^{-i\gamma_j + i\beta_j \cdot x} + O \left( \frac{1}{a^2} e^{-c|x - \alpha_j|} \right) \\ &+ O_N \left( (e^{-c_N a} + a^{-3}) e^{-c_N |x - \alpha_j|} \right). \end{aligned}$$

We collect the terms of degree 1 in  $\theta$

$$L_j \theta := -\Delta \theta + \theta + 2 \phi_{\text{Re}(\theta \bar{V}_j)} V_j + \left( \phi_{|V_j|^2} + \sum_{k \neq j} \psi_{|V_k|^2}^{(1)} \right) \theta.$$

With (4.3), if we take  $T_0(N)$  large enough, then we have

$$\begin{aligned} & i\partial_t\theta_j + \Delta\theta_j - 2\phi_{\text{Re}(\theta_j\bar{R}_j)}R - \phi_{|R|^2}\theta_j \\ &= \frac{1}{\lambda_j^4} (i\lambda_j^2\partial_t\theta - L_j\theta) e^{-i\gamma_j+i\beta_j\cdot x} - 2\phi_{\text{Re}(\theta_j\bar{R}_j)} \sum_{k\neq j} R_k \\ & \quad + O\left(\frac{1}{a^2} + \text{Mod}\right) e^{-c|x-\alpha_j|} + O_N\left(\frac{1}{a^3} + \frac{\text{Mod}}{a}\right) e^{-c_N \min_j |x-\alpha_j|}. \end{aligned}$$

Inserting this into the previous formula, using (2.14), (4.5), (4.2), and again taking  $T_0(N)$  large enough, we get

$$\begin{aligned} \frac{d}{dt}\text{Im} \int \varepsilon\bar{\theta}_j &= \text{Re} \int \frac{\varepsilon}{\lambda_j^4} \overline{(i\lambda_j^2\partial_t\theta - L_j\theta) e^{-i\gamma_j+i\beta_j\cdot x}} - 2\text{Re} \int \varepsilon\phi_{\text{Re}(\theta_j\bar{R}_j)} \sum_{k\neq j} \bar{R}_k \\ & \quad + \frac{1}{\lambda_j^6} \text{Re} \int S_j\bar{\theta} + O\left(\frac{\|\varepsilon\|_{H^1}}{a^2} + \text{Mod}\|\varepsilon\|_{H^1}\right) + O_N\left(\frac{1}{a^{N+1}} + \|\varepsilon\|_{H^1}^2\right). \end{aligned}$$

We will take  $\theta$  to be  $iV_j$ ,  $\nabla V_j$ ,  $\Lambda V_j$  and  $iy_jV_j$ . By (3.8), the left hand side always vanishes. By (2.8), (4.3) and (2.11), we always have

$$(4.6) \quad \partial_t\theta = O\left(\frac{1}{a^2} + \frac{C_N}{a^3} + \frac{\text{Mod}}{a}\right) \left(e^{-c|x-\alpha_j|} + \frac{C_N}{a}e^{-c_N|x-\alpha_j|}\right).$$

By the proof of Proposition 2.1, we know

$$W_j := -\Delta V_j + V_j + \left(\phi_{|V_j|^2} + \sum_{k\neq j} \psi_{|V_k|^2}^{(1)}\right)V_j$$

is admissible of degree  $\geq 2$ . Direct computation yields

$$\begin{aligned} L_j(iV_j) &= iW_j, \quad L_j(\Lambda V_j) = (\Lambda + 2)W_j - 2\left(1 + \sum_{k\neq j} \psi_{|V_k|^2}^{(1)}\right)V_j, \\ L_j(\nabla V_j) &= \nabla W_j, \quad L_j(iy_jV_j) = iy_jW_j - 2i\nabla V_j. \end{aligned}$$

By (2.6), (3.8) and that  $\psi^{(1)}$  is constant, we always have

$$L_j\theta = f + O\left(\frac{1}{a^2} + \frac{C_N}{a^3}\right) \left(e^{-c|x-\alpha_j|} + \frac{C_N}{a}e^{-c_N|x-\alpha_j|}\right),$$

where  $f$  is a function such that  $\text{Re} \int \varepsilon(\overline{g_j}f) = 0$ . Thus

$$(4.7) \quad \text{Re} \int \frac{\varepsilon}{\lambda_j^4} \overline{L_j\theta} e^{-i\gamma_j+i\beta_j\cdot x} = O\left(\frac{\|\varepsilon\|_{H^1}}{a^2} + \frac{C_N\|\varepsilon\|_{H^1}}{a^3}\right).$$

By (2.16) and the explicit formula of  $\psi^{(2)}$ , we have

$$\phi_{\text{Re}(\theta_j\bar{R}_j)} = \psi_{\text{Re}(\theta_j\bar{R}_j)}^{(1)} + O\left(\frac{1}{a^2} + \frac{C_N}{a^3}\right) (1 + |x - \alpha_j|)^2.$$

Since  $\psi^{(1)}$  is constant, by (3.8), we always have

$$(4.8) \quad 2\text{Re} \int \varepsilon\phi_{\text{Re}(\theta_j\bar{R}_j)} \sum_{k\neq j} \bar{R}_k = O\left(\frac{\|\varepsilon\|_{H^1}}{a^2} + \frac{C_N\|\varepsilon\|_{H^1}}{a^3}\right).$$

Finally, using (2.5), (2.8) and that  $Q$  is even, for  $\theta$  taken as the four functions,

$$(4.9) \quad \sum_{\theta} \sum_{j=1}^m \left| \operatorname{Re} \int S_j \bar{\theta} \right| \geq c \operatorname{Mod} - \frac{C_N \operatorname{Mod}}{a}.$$

Therefore, gathering (4.6), (4.7), (4.8) and (4.9), we obtain

$$\operatorname{Mod}(t) \leq \frac{C \|\varepsilon\|_{H^1}}{a^2} + \frac{C_N \|\varepsilon\|_{H^1}}{a^3} + \frac{C_N \operatorname{Mod}(t)}{a} + \frac{C_N}{a^{N+1}} + C_N \|\varepsilon\|_{H^1}^2.$$

Taking  $T_0(N)$  large enough to absorb some  $O_N$  terms, we get

$$(4.10) \quad \operatorname{Mod}(t) \leq \frac{C \|\varepsilon\|_{H^1}}{a^2} + \frac{C_N}{a^{N+1}} + C_N \|\varepsilon\|_{H^1}^2.$$

Using (3.10) and (4.3), we can get the decay of  $\operatorname{Mod}$ :

$$(4.11) \quad \operatorname{Mod}(t) \leq t^{-\frac{N}{4}}, \quad \forall t \in [T_*, T_n].$$

We are now going to deduce the second and third lines of (3.11). By the fundamental theorem of calculus, we have

$$\begin{aligned} & |M_j(P) - M_j(P^{(N)})| + |B_j(P) - B_j(P^{(N)})| \\ & \leq C \left( \frac{|\alpha - \alpha^{(N)}|}{a^3} + \frac{|\beta - \beta^{(N)}|}{a^2} + \frac{|\lambda - \lambda^{(N)}|}{a^2} \right) \\ & \quad + C_N \left( \frac{|\alpha - \alpha^{(N)}|}{a^4} + \frac{|\beta - \beta^{(N)}|}{a^3} + \frac{|\lambda - \lambda^{(N)}|}{a^3} \right). \end{aligned}$$

Using (3.1), (3.10), (4.3) and (4.11), if  $T_0(N)$  is large enough, then we get

$$\begin{aligned} & |\dot{\lambda}_j - \dot{\lambda}_j^{(N)}| + |\dot{\beta}_j - \dot{\beta}_j^{(N)}| \\ & \leq \operatorname{Mod} + |M_j(P) - M_j(P^{(N)})| + |B_j(P) - B_j(P^{(N)})| \leq C t^{-2-\frac{N}{8}}. \end{aligned}$$

Integrating in  $t$  and using (3.9), we deduce

$$(4.12) \quad |\lambda_j - \lambda_j^{(N)}| + |\beta_j - \beta_j^{(N)}| \leq \frac{C}{N} t^{-1-\frac{N}{8}} \leq \frac{1}{2} t^{-1-\frac{N}{8}}$$

when  $N$  is large enough. This is the second line of (3.11).

By (4.11) and (4.3), we also have

$$|\dot{\alpha}_j - \dot{\alpha}_j^{(N)}| \leq 2|\beta_j - \beta_j^{(N)}| + t^{-\frac{N}{4}}$$

and

$$|\dot{\gamma}_j - \dot{\gamma}_j^{(N)}| \leq C \left( |\lambda_j - \lambda_j^{(N)}| + |\beta_j - \beta_j^{(N)}| + t|\dot{\beta}_j - \dot{\beta}_j^{(N)}| + |\dot{\alpha}_j - \dot{\alpha}_j^{(N)}| \right) + t^{-\frac{N}{4}}.$$

We then deduce the third line of (3.11) for large  $N$  by (3.9) and (4.12).

**4.2. Control of the error.** In this subsection, we aim at proving the first line of (3.11). We start with the construction of cutoff functions.

**Lemma 4.1.** *There exist  $c, C > 0$  and  $\varphi_j \in C^{1,\infty}(\mathbb{R}_+ \times \mathbb{R}^3)$  for  $1 \leq j \leq m$  such that*

$$(4.13) \quad \begin{aligned} 0 \leq \varphi_j(t, x) \leq 1, \quad \sum_{j=1}^m \varphi_j(t, x) &\equiv 1, \\ |\partial_t \varphi_j| + |\nabla \varphi_j| &\leq \frac{C}{a}, \quad |\partial_t \sqrt{\varphi_j}| + |\nabla \sqrt{\varphi_j}| \leq \frac{C}{a}, \\ \varphi_j(t, x) &= \begin{cases} 1, & |x - \alpha_j(t)| \leq ca(t), \\ 0, & |x - \alpha_k(t)| \leq ca(t), \quad k \neq j. \end{cases} \end{aligned}$$

*Proof.* There exist  $c_0, C_0 > 0$  such that  $c_0 t < a(t) < C_0 t$ . Take  $c \in (0, \frac{1}{2})$  and  $r, R > 0$  such that  $cC_0 < r < R < \frac{1}{2}c_0$ . Let  $\Phi \in C^\infty(\mathbb{R}^3)$  be such that

$$0 \leq \Phi \leq 1, \quad \text{supp } \Phi \subset B_R, \quad \Phi = 1 \text{ in } B_r, \quad |\nabla \Phi| \leq C\sqrt{1 - \Phi}.$$

Here, the last property can be satisfied by taking  $\Phi_0$  satisfying the other properties and setting  $\Phi = 1 - (1 - \Phi_0)^2$ . Then we define

$$\varphi_j(t, x) = \Phi^2\left(\frac{x - \alpha_j(t)}{t}\right), \quad j = 1, 2, \dots, m-1,$$

and

$$\varphi_m(t, x) = 1 - \sum_{j=1}^{m-1} \varphi_j(t, x).$$

We claim that (4.13) holds. Only the estimates on  $\varphi_m$  need to be checked.

We have  $\varphi_m \in [0, 1]$  because  $\text{supp } \varphi_j$  are pairwise disjoint. The derivatives of  $\sqrt{\varphi_m}$  are bounded because of the last property of  $\Phi$ .  $\square$

Combining the properties of the cutoff functions and (2.6), we have

$$(4.14) \quad |\varphi_j R - R_j| \leq C_N e^{-ca(t)}, \quad \forall t > 0, \quad x \in \mathbb{R}^3.$$

This means  $\varphi_j$  localizes the multisoliton solutions.

Consider the sum of truncated conserved quantities of the Hartree equation

$$\begin{aligned} \mathcal{E}(u) &= \int |\nabla u|^2 - \frac{1}{2} \int |\nabla \phi_{|u|^2}|^2 \\ &\quad + \sum_{j=1}^m \left[ \left( \frac{1}{\lambda_j^2} + |\beta_j|^2 \right) \int \varphi_j |u|^2 - 2\beta_j \int \varphi_j \text{Im}(\nabla u \bar{u}) \right]. \end{aligned}$$

By the decomposition  $u = R + \varepsilon$ , we can expand  $\mathcal{E}$  in terms of  $\varepsilon$ . Then the second or higher order terms is  $\mathcal{G}(\varepsilon) = \mathcal{G}_1 + \mathcal{G}_2 + \mathcal{G}_3$ , where

$$\begin{aligned} \mathcal{G}_1 &= \int |\nabla \varepsilon|^2 + \int \phi_{|R|^2} |\varepsilon|^2 - 2 \int |\nabla \phi_{\text{Re}(\varepsilon \bar{R})}|^2 + 2 \int \phi_{\text{Re}(\varepsilon \bar{R})} |\varepsilon|^2 - \frac{1}{2} \int |\nabla \phi_{|\varepsilon|^2}|^2, \\ \mathcal{G}_2 &= \sum_{j=1}^m \left( \frac{1}{\lambda_j^2} + |\beta_j|^2 \right) \int \varphi_j |\varepsilon|^2, \quad \mathcal{G}_3 = -2 \sum_{j=1}^m \beta_j \int \varphi_j \text{Im}(\nabla \varepsilon \bar{\varepsilon}). \end{aligned}$$

We point out that the first order term of  $\varepsilon$  would vanish, by using the orthogonality condition (3.8), if  $R$  solved (1.3). Unfortunately, as  $R$  is an approximate solution, it does not solve (1.3).

However, for other equations such as (NLS) and (gKdV) where this is true, the property can be used to derive the estimate of  $\varepsilon$  (see [14] and [15]).

Therefore, we will not use the functional  $\mathcal{E}$  in our proof. Instead, we will prove the following two propositions. The first one states the positiveness of  $\mathcal{G}$ , which follows because of orthogonality (3.8). The second follows by a direct calculation and gives an estimate on the upper bound of  $\mathcal{G}$ .

**Proposition 4.1.** *Let  $N \geq 2$ . For  $T_0 = T_0(N)$  large enough, there exists  $c_0 > 0$  such that*

$$\mathcal{G}(\varepsilon(t)) \geq c_0 \|\varepsilon\|_{H^1}^2, \quad \forall t \in [T_*, T_n].$$

**Proposition 4.2.** *Let  $N \geq 2$ . For  $T_0 = T_0(N)$  large enough, if (3.10) holds, then there exists  $C > 0$  such that*

$$\left| \frac{d}{dt} \mathcal{G}(\varepsilon(t)) \right| \leq C t^{-1-\frac{N}{2}}, \quad \forall t \in [T_*, T_n].$$

If taking these two results for granted, then by (3.9), we have

$$c_0 \|\varepsilon\|_{H^1}^2 \leq |\mathcal{G}(\varepsilon(t))| \leq \int_t^{T_n} C \tau^{-1-\frac{N}{2}} d\tau \leq \frac{C}{N} t^{-\frac{N}{2}}.$$

Taking  $N$  large enough, then we obtain the first line of (3.11).

The rest of this subsection is for the proof of the two propositions. Once they are proved, we will have completed the proof of Theorem 1.

*Proof of Proposition 4.1.*

The main ingredient of the proof is the following coercivity result on the linearized operators  $L_-$  and  $L_+$ .

**Lemma 4.2.** *There exist  $\delta, c > 0$  such that: if  $v \in H^1$  is real-valued, then*

$$\begin{aligned} |(v, Q)| + |(v, xQ)| < \delta \|v\|_{H^1} &\implies (L_+ v, v) \geq c \|v\|_{H^1}^2, \\ |(v, \Lambda Q)| + |(v, \nabla Q)| < \delta \|v\|_{H^1} &\implies (L_- v, v) \geq c \|v\|_{H^1}^2. \end{aligned}$$

*Proof.* All functions in this proof are assumed to be real-valued.

It suffices to prove that for some  $c > 0$ , if  $v \in H^1$ , then

$$(4.15) \quad \begin{aligned} (v, Q) = (v, xQ) = 0 &\implies (L_+ v, v) \geq c \|v\|_{H^1}^2, \\ (v, \Lambda Q) = (v, \nabla Q) = 0 &\implies (L_- v, v) \geq c \|v\|_{H^1}^2. \end{aligned}$$

Let us only prove the sufficiency for the estimate on  $L_+$ . For  $v \in H^1$ , we have

$$(L_+ v, v) = \int |\nabla v|^2 + \int |v|^2 + \int \phi_{Q^2} v^2 - 2 \int |\nabla \phi_{Qv}|^2.$$

If  $u = v - w$ , then by the Sobolev inequality, we have

$$(L_+ v, v) - (L_+ u, u) \geq -C \|u\|_{H^1} \|w\|_{H^1} - C \|w\|_{H^1}^2.$$

We take

$$w = \frac{(v, Q)}{\|Q\|_{L^2}^2} Q + \frac{(v, xQ)}{\|xQ\|_{L^2}^2} \cdot xQ.$$

Then  $\|w\|_{H^1} \leq C\delta \|v\|_{H^1}$ , and  $u$  is orthogonal to both  $Q$  and  $xQ$ , so we have  $(L_+ u, u) \geq c \|u\|_{H^1}^2$ . We thus deduce  $(L_+ v, v) \geq \frac{c}{2} \|v\|_{H^1}^2$  by Cauchy-Schwarz and taking  $\delta$  small.

We then turn to prove (4.15). We only prove the first line, as the proof of the second line is similar and easier.

Recall that  $Q$  is a minimizer of

$$\int |\nabla u|^2 - \frac{1}{2} \int |\nabla \phi_{|u|^2}|^2, \quad \text{where } u \in H^1 \text{ and } \|u\|_{L^2} = \|Q\|_{L^2}.$$

Thus,  $Q$  is a minimizer in  $H^1 \setminus \{0\}$  of

$$\mathcal{J}(u) := \frac{\|Q\|_{L^2}^2}{\|u\|_{L^2}^2} \int |\nabla u|^2 - \frac{\|Q\|_{L^2}^4}{2\|u\|_{L^2}^4} \int |\nabla \phi_{|u|^2}|^2.$$

Assume  $f \in H^1$  and  $(f, Q) = 0$ . By direct computation, we have

$$\begin{aligned} \left. \frac{1}{2} \frac{d^2}{d\epsilon^2} \right|_{\epsilon=0} \mathcal{J}(Q + \epsilon f) &= \int |\nabla f|^2 - \frac{\|\nabla Q\|_{L^2}^2}{\|Q\|_{L^2}^2} \int f^2 \\ &\quad - 2 \int |\nabla \phi_{Qf}|^2 - \int \nabla \phi_{Q^2} \cdot \nabla \phi_{f^2} + \frac{\|\nabla \phi_{Q^2}\|_{L^2}^2}{\|Q\|_{L^2}^2} \int f^2. \end{aligned}$$

Using (1.3) and integration by parts, we get

$$\|\nabla Q\|_{L^2}^2 - \|\nabla \phi_{Q^2}\|_{L^2}^2 + \|Q\|_{L^2}^2 = 0.$$

Thus we obtain

$$\left. \frac{1}{2} \frac{d^2}{d\epsilon^2} \right|_{\epsilon=0} \mathcal{J}(Q + \epsilon f) = (L_+ f, f).$$

By the minimality of  $\mathcal{J}(Q)$ , we deduce that

$$f \in H^1, (f, Q) = 0 \implies (L_+ f, f) \geq 0.$$

Therefore, if the conclusion fails to be true, then there exist  $f_n \in H^1$  such that  $(f_n, Q) = (f_n, xQ) = 0$ ,  $\|f_n\|_{H^1} = 1$  and  $(L_+ f_n, f_n) \rightarrow 0$ . By passing to subsequence, we may assume  $f_n \rightharpoonup f_0$  in  $H^1$ . We have  $(f_0, Q) = (f_0, xQ) = 0$ , and by the Rellich-Kondrachov theorem and the decay of  $Q$ , we have

$$\int \phi_{Q^2} f_n^2 \rightarrow \int \phi_{Q^2} f_0^2 \quad \text{and} \quad \int |\nabla \phi_{Qf_n}|^2 \rightarrow \int |\nabla \phi_{Qf_0}|^2.$$

On the other hand,  $\|f_0\|_{H^1} \leq \liminf \|f_n\|_{H^1}$ . We deduce that  $(L_+ f_0, f_0) \leq 0$ .

By the non-negativity of  $L^+$  on  $Q^\perp$ , we know that  $f_0$  is a nonzero minimizer of  $(L_+ u, u)$ , where  $u \in H^1$  and  $(u, Q) = 0$ . By computing the first variation, we get

$$L_+ f_0 = aQ \quad \text{for some } a \in \mathbb{R}.$$

Since  $L_+(\Lambda Q) = -2Q$  and  $\ker L_+$  is spanned by  $\nabla Q$  (Theorem 4 in [9]), we know  $f_0$  is a linear combination of  $\Lambda Q$  and  $\nabla Q$ . But using  $(f_0, Q) = (f_0, \nabla Q) = 0$ , we must have  $f_0 = 0$ , which is a contradiction.  $\square$

Let  $\varepsilon_j = \varepsilon \sqrt{\varphi_j}$  and  $\tilde{\varepsilon}_j = g_j^{-1} \varepsilon_j$ , or more precisely, define

$$\tilde{\varepsilon}_j(t, y_j) = \lambda_j^2(t) \varepsilon_j(t, \lambda_j(t) y_j + \alpha_j(t)) e^{i\gamma_j(t) - i\beta_j(t) \cdot (\lambda_j(t) y_j + \alpha_j(t))}.$$

By (3.8), (2.6) and (4.3), for  $t$  large enough, we can apply Lemma 4.2 to  $\text{Re}(\tilde{\varepsilon}_j)$  and  $\text{Im}(\tilde{\varepsilon}_j)$ . Thus for  $T_0(N)$  large enough, we have

$$\left( L_+ \text{Re}(\tilde{\varepsilon}_j), \text{Re}(\tilde{\varepsilon}_j) \right) + \left( L_- \text{Im}(\tilde{\varepsilon}_j), \text{Im}(\tilde{\varepsilon}_j) \right) \geq c \|\tilde{\varepsilon}_j\|_{H^1}^2, \quad \forall t \geq T_0.$$

In the following, we always assume  $T_0$  is large enough so that the above holds.



Set  $Q_j = g_j Q$ . By computation similar to (2.1) and (2.6), we have

$$\begin{aligned}
& \left( L_+ \operatorname{Re}(\tilde{\varepsilon}_j), \operatorname{Re}(\tilde{\varepsilon}_j) \right) + \left( L_- \operatorname{Im}(\tilde{\varepsilon}_j), \operatorname{Im}(\tilde{\varepsilon}_j) \right) \\
&= \int |\nabla \tilde{\varepsilon}_j|^2 + \int |\tilde{\varepsilon}_j|^2 + \int \phi_{Q^2} |\tilde{\varepsilon}_j|^2 + 2 \int \phi_{\operatorname{Re}(Q\tilde{\varepsilon}_j)} \operatorname{Re}(Q\tilde{\varepsilon}_j) \\
&= \lambda_j^3 \left( \int |\nabla \varepsilon_j|^2 - 2\beta_j \int \operatorname{Im}(\nabla \varepsilon_j \bar{\varepsilon}_j) + |\beta_j|^2 \int |\varepsilon_j|^2 \right) + \lambda_j \int |\varepsilon_j|^2 \\
&\quad + \lambda_j^3 \int \phi_{Q_j^2} |\varepsilon_j|^2 + 2\lambda_j^3 \int \phi_{\operatorname{Re}(\varepsilon_j \bar{Q}_j)} \operatorname{Re}(\varepsilon_j \bar{Q}_j) \\
&= \lambda_j^3 \left( \int |\nabla \varepsilon_j|^2 + \int \phi_{|R_j|^2} |\varepsilon_j|^2 - 2 \int |\nabla \phi_{\operatorname{Re}(\varepsilon_j \bar{R}_j)}|^2 \right. \\
&\quad \left. + \left( \frac{1}{\lambda_j^2} + |\beta_j|^2 \right) \int |\varepsilon_j|^2 - 2\beta_j \int \operatorname{Im}(\nabla \varepsilon_j \bar{\varepsilon}_j) \right) + O_N \left( \frac{\|\varepsilon_j\|_{H^1}^2}{a} \right)
\end{aligned}$$

We then deduce that

$$(4.16) \quad \mathcal{H}_j(\varepsilon_j) \geq c \|\varepsilon_j\|_{H^1} - \frac{C_N}{a} \|\varepsilon_j\|_{H^1}^2 \geq c \int \varphi_j (|\nabla \varepsilon|^2 + |\varepsilon|^2) - \frac{C_N}{a} \|\varepsilon\|_{H^1}^2,$$

where

$$\begin{aligned}
\mathcal{H}_j(\varepsilon_j) &= \int |\nabla \varepsilon_j|^2 + \int \phi_{|R_j|^2} |\varepsilon_j|^2 - 2 \int |\nabla \phi_{\operatorname{Re}(\varepsilon_j \bar{R}_j)}|^2 \\
&\quad + \left( \frac{1}{\lambda_j^2} + |\beta_j|^2 \right) \int |\varepsilon_j|^2 - 2\beta_j \int \operatorname{Im}(\nabla \varepsilon_j \bar{\varepsilon}_j).
\end{aligned}$$

Next, we consider the truncated functional

$$\begin{aligned}
\mathcal{H}_{j,\varphi}(\varepsilon) &= \int \varphi_j |\nabla \varepsilon|^2 + \int \phi_{|R_j|^2} |\varepsilon|^2 - 2 \int |\nabla \phi_{\operatorname{Re}(\varepsilon \bar{R}_j)}|^2 \\
&\quad + \left( \frac{1}{\lambda_j^2} + |\beta_j|^2 \right) \int \varphi_j |\varepsilon|^2 - 2\beta_j \int \varphi_j \operatorname{Im}(\nabla \varepsilon \bar{\varepsilon}).
\end{aligned}$$

By (4.13), (4.14) and (4.16), we have

$$\begin{aligned}
(4.17) \quad \mathcal{H}_{j,\varphi}(\varepsilon) &= \int |\nabla \varepsilon_j|^2 + \int \phi_{|R_j|^2} |\varepsilon|^2 - 2 \int |\nabla \phi_{\operatorname{Re}(\varepsilon \bar{R}_j)}|^2 \\
&\quad + \left( \frac{1}{\lambda_j^2} + |\beta_j|^2 \right) \int |\varepsilon_j|^2 - 2\beta_j \int \operatorname{Im}(\nabla \varepsilon_j \bar{\varepsilon}_j) + O \left( \frac{\|\varepsilon\|_{H^1}^2}{a} \right) \\
&= \mathcal{H}_j(\varepsilon_j) + \int (1 - \sqrt{\varphi_j}) \phi_{|R_j|^2} |\varepsilon|^2 - 2 \int |\nabla \phi_{\operatorname{Re}((1-\sqrt{\varphi_j})\varepsilon \bar{R}_j)}|^2 \\
&\quad + 4 \int \phi_{\operatorname{Re}(\varepsilon_j \bar{R}_j)} \operatorname{Re}((1 - \sqrt{\varphi_j})\varepsilon \bar{R}_j) + O \left( \frac{\|\varepsilon\|_{H^1}^2}{a} \right) \\
&\geq c \int \varphi_j (|\nabla \varepsilon|^2 + |\varepsilon|^2) - \frac{C_N}{a} \|\varepsilon\|_{H^1}^2.
\end{aligned}$$

Finally, using the sum-to-1 property of  $\varphi_j$ , we write

$$\begin{aligned} \mathcal{G}(\varepsilon) &= \sum_{j=1}^m \mathcal{H}_{j,\varphi}(\varepsilon) + 2 \int \phi_{\text{Re}(\varepsilon\bar{R})} |\varepsilon|^2 - \frac{1}{2} \int |\nabla \phi_{|\varepsilon|^2}|^2 \\ &\quad + \sum_{j \neq k} \int \phi_{\text{Re}(R_k \bar{R}_j)} |\varepsilon|^2 - 2 \sum_{j \neq k} \int \nabla \phi_{\text{Re}(\varepsilon \bar{R}_j)} \cdot \nabla \phi_{\text{Re}(\varepsilon \bar{R}_k)}. \end{aligned}$$

The first term is controlled by (4.17). The other terms in the first line are  $O(t^{-N/4} \|\varepsilon\|_{H^1}^2)$  because of (3.10). Using Lemma 2.3, we know the two terms in the second line are  $O_N(e^{-ca} \|\varepsilon\|_{H^1}^2)$  and  $O_N\left(\frac{\|\varepsilon\|_{H^1}^2}{a}\right)$ , respectively. We thus have

$$\mathcal{G}(\varepsilon) \geq c \|\varepsilon\|_{H^1}^2 - \frac{C_N}{a} \|\varepsilon\|_{H^1}^2.$$

Thanks to (4.3), we conclude by taking  $T_0(N)$  large enough.  $\square$

*Proof of Proposition 4.2.* We deal with  $\mathcal{G}_1$ ,  $\mathcal{G}_2$  and  $\mathcal{G}_3$  separately.

(1) Using integration by parts, we have

$$\begin{aligned} \frac{d\mathcal{G}_1}{dt} &= -2\text{Im} \int i\partial_t \bar{\varepsilon} \left( \Delta \varepsilon - \phi_{|R|^2} \varepsilon - 2\phi_{\text{Re}(\varepsilon \bar{R})} R - \mathcal{N}(\varepsilon) \right) \\ &\quad + 4\text{Re} \int \phi_{\text{Re}(\varepsilon \bar{R})} \varepsilon \partial_t \bar{R} + 2 \int \phi_{\text{Re}(\partial_t R \bar{R})} |\varepsilon|^2 + 2\text{Re} \int \phi_{|\varepsilon|^2} \varepsilon \partial_t \bar{R}. \end{aligned}$$

By (4.1), (2.14), (4.4) and (4.2), the first line is  $O_N\left(\left(\frac{1}{a^{N+1}} + \text{Mod}\right) \|\varepsilon\|_{H^1}\right)$ . For the second line, using (2.1), (4.3) and (2.11), we have

$$\begin{aligned} \partial_t R_j &= \frac{1}{\lambda_j^2} \left( -\dot{\alpha}_j \cdot \nabla V_j - i(\dot{\gamma}_j - \dot{\beta}_j \cdot \alpha_j) V_j \right) e^{-i\gamma_j + i\beta \cdot x} + O_N\left(\frac{1}{a^2}\right) e^{-c_N|x-\alpha_j|} \\ &= -2\beta_j \cdot \nabla R_j + i\left(\frac{1}{\lambda_j^2} + |\beta_j|^2\right) R_j + O_N\left(\frac{1}{a^2} + \text{Mod}\right) e^{-c_N|x-\alpha_j|}. \end{aligned}$$

Combining these with (4.10), (2.9) and Lemma 2.3, we get

$$\begin{aligned} \frac{d\mathcal{G}_1}{dt} &= \sum_{j=1}^m \left( 4\left(\frac{1}{\lambda_j^2} + |\beta_j|^2\right) \int \phi_{\text{Re}(\varepsilon \bar{R}_j)} \text{Im}(\varepsilon \bar{R}_j) - 8 \int \phi_{\text{Re}(\varepsilon \bar{R})} \text{Re}(\varepsilon \beta_j \cdot \nabla \bar{R}_j) \right. \\ (4.18) \quad &\quad \left. - 4 \int \phi_{\text{Re}(\beta_j \cdot \nabla R_j \bar{R}_j)} |\varepsilon|^2 \right) + O_N\left(\frac{\|\varepsilon\|_{H^1}}{a^{N+1}} + \frac{\|\varepsilon\|_{H^1}^2}{a^2} + \|\varepsilon\|_{H^1}^3\right). \end{aligned}$$

(2) Using (4.3) and (4.13), we have

$$\frac{d\mathcal{G}_2}{dt} = \sum_{j=1}^m 2\left(\frac{1}{\lambda_j^2} + |\beta_j|^2\right) \int \varphi_j \text{Im}(i\partial_t \varepsilon \bar{\varepsilon}) + O\left(\frac{\|\varepsilon\|_{H^1}^2}{t}\right).$$

Then by (4.1), (2.14), (4.4) and (4.2), we have

$$\begin{aligned} \frac{d\mathcal{G}_2}{dt} &= \sum_{j=1}^m 2\left(\frac{1}{\lambda_j^2} + |\beta_j|^2\right) \int \varphi_j \text{Im}(2\phi_{\text{Re}(\varepsilon \bar{R})} R \bar{\varepsilon}) + O\left(\frac{\|\varepsilon\|_{H^1}^2}{t}\right) \\ &\quad + O_N\left(\frac{\|\varepsilon\|_{H^1}}{a^{N+1}} + \text{Mod} \|\varepsilon\|_{H^1} + \|\varepsilon\|_{H^1}^3\right). \end{aligned}$$

Finally using (4.10) and (4.14), we get

$$(4.19) \quad \begin{aligned} \frac{d\mathcal{G}_2}{dt} &= \sum_{j=1}^m -4 \left( \frac{1}{\lambda_j^2} + |\beta_j|^2 \right) \int \phi_{\operatorname{Re}(\varepsilon \bar{R}_j)} \operatorname{Im}(\varepsilon \bar{R}_j) + O \left( \frac{\|\varepsilon\|_{H^1}^2}{t} \right) \\ &\quad + O_N \left( \frac{\|\varepsilon\|_{H^1}}{a^{N+1}} + \frac{\|\varepsilon\|_{H^1}^2}{a^2} + \|\varepsilon\|_{H^1}^3 \right). \end{aligned}$$

(3) Similarly, we can compute

$$(4.20) \quad \begin{aligned} \frac{d\mathcal{G}_3}{dt} &= \sum_{j=1}^m -4\beta_j \int \varphi_j \operatorname{Re}(i\partial_t \varepsilon \nabla \bar{\varepsilon}) + O \left( \frac{\|\varepsilon\|_{H^1}^2}{t} \right) \\ &= \sum_{j=1}^m -4\beta_j \int \varphi_j \operatorname{Re} \left( 2\phi_{\operatorname{Re}(\varepsilon \bar{R}_j)} R \nabla \bar{\varepsilon} + \phi_{|R|^2} \varepsilon \nabla \bar{\varepsilon} \right) + O \left( \frac{\|\varepsilon\|_{H^1}^2}{t} \right) \\ &\quad + O_N \left( \frac{\|\varepsilon\|_{H^1}}{a^{N+1}} + \operatorname{Mod} \|\varepsilon\|_{H^1} + \|\varepsilon\|_{H^1}^3 \right) \\ &= \sum_{j=1}^m \left( 8 \int \phi_{\operatorname{Re}(\varepsilon \bar{R}_j)} \operatorname{Re}(\varepsilon \beta_j \varphi_j \cdot \nabla \bar{R}_j) + 4 \int \phi_{\operatorname{Re}(\beta_j \varphi_j \cdot \nabla R \bar{R}_j)} |\varepsilon|^2 \right) \\ &\quad + O \left( \frac{\|\varepsilon\|_{H^1}^2}{t} \right) + O_N \left( \frac{\|\varepsilon\|_{H^1}}{a^{N+1}} + \operatorname{Mod} \|\varepsilon\|_{H^1} + \|\varepsilon\|_{H^1}^3 \right) \\ &= \sum_{j=1}^m \left( 8 \int \phi_{\operatorname{Re}(\varepsilon \bar{R}_j)} \operatorname{Re}(\varepsilon \beta_j \cdot \nabla \bar{R}_j) + 4 \int \phi_{\operatorname{Re}(\beta_j \cdot \nabla R_j \bar{R}_j)} |\varepsilon|^2 \right) \\ &\quad + O \left( \frac{\|\varepsilon\|_{H^1}^2}{t} \right) + O_N \left( \frac{\|\varepsilon\|_{H^1}}{a^{N+1}} + \frac{\|\varepsilon\|_{H^1}^2}{a^2} + \|\varepsilon\|_{H^1}^3 \right). \end{aligned}$$

By combining (4.18), (4.19), (4.20) together, using (4.3), (3.10) and taking  $T_0(N)$  large enough, we get the desired result.  $\square$

Now we have finished the proof of Theorem 1.

## 5. THE PARABOLIC AND THE HYPERBOLIC-PARABOLIC CASE

One of the difficulties of dealing with these two cases is to establish Proposition 3.1. Due to the lower rates of expansion, we need more delicate computation.

**5.1. The approximate trajectory.** The goal of this subsection is to prove the alternative of Proposition 3.1.

**Proposition 5.1.** *Let  $P^\infty$  be a parabolic or hyperbolic-parabolic solution to (1.5) of the form (1.7) or (1.8), and  $B_j^{(N)}, M_j^{(N)}$  be as in Proposition 2.1. Then for  $N \geq 3$ ,  $\exists T_0 = T_0(N) > 0$  and  $P^{(N)} \in C^1([T_0, +\infty), \Omega)$  satisfying (3.1) and for any  $t \geq T_0$ ,*

$$(5.1) \quad |\alpha_j^{(N)}(t) - \alpha_j^\infty(t)| \leq t^{-\frac{1}{4}}, \quad |\beta_j^{(N)}(t) - \beta_j^\infty(t)| + |\lambda_j^{(N)}(t) - \lambda_j^\infty| \leq t^{-\frac{1}{2}}.$$

**Remark.** *This proposition is stronger than the one in [8] because: (1) we do not need to assume  $\lambda_j^\infty$  are identical in the parabolic case, or  $\lambda_j^\infty$  are identical for  $j \in J$  in the hyperbolic-parabolic case; (2) we know  $\alpha_j^{(N)}(t) - \alpha_j^\infty(t) \rightarrow 0$  as  $t \rightarrow +\infty$ .*

While before we have only used the formula of  $b_j^{(2)}$ , here we also need the explicit expression of  $m_j^{(2)}$  and  $b_j^{(3)}$ . Since  $T_j^{(1)}$  is real-valued, by (2.4), we have

$$\operatorname{Im} \hat{E}_j^{(1)} = \lambda_j^2 \sum_{k=1}^m \frac{\partial T_j^{(1)}}{\partial \alpha_k} \cdot 2\beta_k.$$

By the formula of  $T_j^{(1)}$  (2.7) and the requirement of  $m_j^{(2)}$ , we deduce

$$(5.2) \quad m_j^{(2)}(P) = \sum_{k \neq j} \frac{\lambda_j^3 \|Q\|_{L^2}^2 \alpha_{jk} \cdot \beta_{jk}}{4\pi \lambda_k |\alpha_{jk}|^3}.$$

Combining this and calculation in the proof of Proposition 2.1, we have

$$\begin{cases} L_+ \operatorname{Re} T_j^{(2)} = -2\phi_{QT_j^{(1)}} T_j^{(1)} - \phi_{|T_j^{(1)}|^2} Q - \sum_{k \neq j} \left( 2\psi_{QT_k^{(1)}}^{(1)} Q + \psi_{Q^2, k}^{(1)} T_k^{(1)} \right), \\ L_- \operatorname{Im} T_j^{(2)} = 0. \end{cases}$$

Thus we may take  $T_j^{(2)}$  as a real-valued radial function. Then we compute

$$\begin{aligned} \operatorname{Re} \hat{E}_j^{(2)} &= -\lambda_j^3 b_j^{(2)} \cdot y_j T_j^{(1)} - 2\phi_{T_j^{(1)} T_j^{(2)}} Q - 2\phi_{QT_j^{(2)}} T_j^{(1)} - 2\phi_{QT_j^{(1)}} T_j^{(2)} \\ &\quad - \phi_{|T_j^{(1)}|^2} T_j^{(1)} - \sum_{k \neq j} \left( \psi_{Q^2, k}^{(2)} T_j^{(1)} + 2\psi_{QT_k^{(1)}}^{(2)} Q + 2\psi_{QT_k^{(2)}}^{(1)} Q \right. \\ &\quad \left. + \psi_{|T_j^{(1)}|^2}^{(1)} Q + 2\psi_{QT_k^{(1)}}^{(1)} T_j^{(1)} + \psi_{Q^2, k}^{(1)} T_j^{(2)} \right). \end{aligned}$$

Recall that

$$\left( \lambda_j^3 b_j^{(3)} \cdot y_j Q + \sum_{k \neq j} \psi_{Q^2, k}^{(3)} Q - \operatorname{Re} \hat{E}_j^{(2)}, \nabla Q \right) = 0.$$

Since  $Q, T_j^{(1)}, T_j^{(2)}$  are all even, the terms with  $\phi, \psi^{(1)}, \psi^{(3)}$  are orthogonal to  $\nabla Q$ . Removing those terms and using (3.3), we obtain

$$\left( \lambda_j^3 b_j^{(3)} \cdot y_j Q + 2 \sum_{k \neq j} \psi_{QT_k^{(1)}}^{(2)} Q, \nabla Q \right) = 0.$$

Then by (2.7) and  $(\Lambda Q, Q) = \frac{1}{2} \|Q\|_{L^2}^2$ , a result of integration by parts, we get

$$(5.3) \quad b_j^{(3)}(P) = - \sum_{k \neq j} \frac{(Q, T_k^{(1)}) \alpha_{jk}}{2\pi \lambda_k |\alpha_{jk}|^3} = \frac{\|Q\|_{L^2}^4}{32\pi} \sum_{k \neq j} \left( \sum_{l \neq k} \frac{1}{\lambda_l |\alpha_{kl}|} \right) \frac{\lambda_k \alpha_{jk}}{|\alpha_{jk}|^3}.$$

Now we are ready to prove the proposition.

*Proof of Proposition 5.1.*

Let  $\epsilon = \frac{1}{100}$  and  $T_0 > 0$ . Define the norm  $\|\cdot\|_2$  of  $P \in C([T_0, +\infty), \Omega)$  by

$$\|P\|_2 := \sum_{j=1}^m \sup_{t \geq T_0} \left( t^{\frac{1}{3}-3\epsilon} |\alpha_j(t)| + t^{\frac{4}{3}-2\epsilon} |\beta_j(t)| + t^{1-\epsilon} |\lambda_j(t)| \right),$$

and let  $Y = \left\{ P \in C([T_0, +\infty), \Omega) \mid \|P - \tilde{P}\|_2 \leq 1 \right\}$ . Then it suffices to find a solution of (3.1) in  $Y$ . We will assume  $P \in Y$  hereinafter.

Take  $\tilde{P}(t) = (\alpha^\infty(t), \beta^\infty(t), \tilde{\lambda}(t))$ , where for each  $j$ ,

$$\tilde{\lambda}_j(t) = \lambda_j^\infty - \int_t^\infty \sum_{k \neq j} \frac{(\lambda_j^\infty)^3 \|Q\|_{L^2}^2 \alpha_{jk}^\infty(\tau) \cdot \beta_{jk}^\infty(\tau)}{4\pi \lambda_k^\infty |\alpha_{jk}^\infty(\tau)|^3} d\tau.$$

Using  $\dot{\alpha}_j^\infty = 2\beta_j^\infty$ , we may simplify the expression as

$$\tilde{\lambda}_j(t) = \lambda_j^\infty - \sum_{k \neq j} \frac{(\lambda_j^\infty)^3 \|Q\|_{L^2}^2}{8\pi \lambda_k^\infty |\alpha_{jk}^\infty(t)|}.$$

Recalling the expression (3.3), (5.2) and (5.3), if we set

$$\begin{aligned} \tilde{b}_j^{(2)}(P) &= b_j^{(2)}(P) + \sum_{k \neq j} \frac{\|Q\|_{L^2}^2 \alpha_{jk}}{4\pi \lambda_k^\infty |\alpha_{jk}|^3} - \sum_{k \neq j} \frac{\|Q\|_{L^2}^2 (\lambda_k - \lambda_k^\infty) \alpha_{jk}}{4\pi (\lambda_k^\infty)^2 |\alpha_{jk}|^3}, \\ \tilde{b}_j^{(3)}(P) &= b_j^{(3)}(P) - \frac{\|Q\|_{L^2}^4}{32\pi} \sum_{k \neq j} \left( \sum_{l \neq k} \frac{1}{\lambda_l^\infty |\alpha_{kl}|} \right) \frac{\lambda_k^\infty \alpha_{jk}}{|\alpha_{jk}|^3}, \\ \tilde{m}_j^{(2)}(P) &= m_j^{(2)}(P) - \sum_{k \neq j} \frac{(\lambda_j^\infty)^3 \|Q\|_{L^2}^2 \alpha_{jk} \cdot \beta_{jk}}{4\pi \lambda_k^\infty |\alpha_{jk}|^3}, \end{aligned}$$

then using  $\tilde{a}(t) \sim t^{\frac{2}{3}}$  and  $\|\tilde{P} - P^\infty\| \leq Ct^{-\frac{2}{3}}$ , we get

$$(5.4) \quad \left| \tilde{b}_j^{(2)}(\tilde{P}(t)) \right| + \left| \tilde{b}_j^{(3)}(\tilde{P}(t)) \right| \leq Ct^{-\frac{8}{3}}, \quad \left| \tilde{m}_j^{(2)}(\tilde{P}(t)) \right| \leq Ct^{-2}.$$

Moreover, direct computation yields

$$\dot{\tilde{\beta}}_j = b_j^{(2)}(\tilde{P}) - \tilde{b}_j^{(2)}(\tilde{P}) + b_j^{(3)}(\tilde{P}) - \tilde{b}_j^{(3)}(\tilde{P}), \quad \dot{\tilde{\lambda}}_j = m_j^{(2)}(\tilde{P}) - \tilde{m}_j^{(2)}(\tilde{P}).$$

This means the error caused by  $b_j^{(2)}$  and  $m_j^{(2)}$  is cancelled with the error of  $b_j^{(3)}$ . Our  $\tilde{P}$  serves as  $P^{(app)}$  in [8], but we have an explicit expression of  $\tilde{P}$  and most importantly, we know  $\tilde{\alpha} = \alpha^\infty$  and  $\tilde{\beta} = \beta^\infty$ .

Write  $P^{(N)} = P$  for simplicity. Then we can rewrite (3.1) as

$$\left\{ \begin{aligned} \dot{\alpha}_j(t) - \dot{\tilde{\alpha}}_j(t) &= 2\beta_j(t) - 2\tilde{\beta}_j(t), \\ \dot{\beta}_j(t) - \dot{\tilde{\beta}}_j(t) &= \left[ b_j^{(2)}(P(t)) - b_j^{(2)}(\tilde{P}(t)) \right] + \left[ b_j^{(3)}(P(t)) - b_j^{(3)}(\tilde{P}(t)) \right] \\ &\quad + \tilde{b}_j^{(2)}(\tilde{P}(t)) + \tilde{b}_j^{(3)}(\tilde{P}(t)) + \sum_{n=4}^N b_j^{(n)}(P(t)), \\ \dot{\lambda}_j(t) - \dot{\tilde{\lambda}}_j(t) &= \left[ m_j^{(2)}(P(t)) - m_j^{(2)}(\tilde{P}(t)) \right] + \tilde{m}_j^{(2)}(\tilde{P}(t)) + \sum_{n=3}^N m_j^{(n)}(P(t)). \end{aligned} \right.$$

By (5.4), estimates of  $b_j^{(n)}$ ,  $m_j^{(n)}$  and  $a(t) \sim t^{\frac{2}{3}}$ , we have

$$(5.5) \quad \left\{ \begin{aligned} \dot{\alpha}_j(t) - \dot{\tilde{\alpha}}_j(t) &= 2\beta_j(t) - 2\tilde{\beta}_j(t), \\ \dot{\beta}_j(t) - \dot{\tilde{\beta}}_j(t) &= b_j^{(2)}(P(t)) - b_j^{(2)}(\tilde{P}(t)) + O(t^{-\frac{8}{3}}), \\ \dot{\lambda}_j(t) - \dot{\tilde{\lambda}}_j(t) &= O(t^{-2}). \end{aligned} \right.$$

In this proof,  $O(t^{-\kappa})$  represents a continuous function of  $\tilde{P}$  and  $P$ , whose  $C^1$  norm in  $P$  is bounded by  $Ct^{-\kappa}$  when evaluated at  $(\tilde{P}(t), P(t))$ .

We still need to estimate  $b_j^{(2)}(P(t)) - b_j^{(2)}(\tilde{P}(t))$ . We have

$$b_j^{(2)}(P) - b_j^{(2)}(\tilde{P}) = \frac{\|Q\|_{L^2}^2}{4\pi} \sum_{k \neq j} \left[ \frac{1}{\tilde{\lambda}_k} \left( \frac{\tilde{\alpha}_{jk}}{|\tilde{\alpha}_{jk}|^3} - \frac{\alpha_{jk}}{|\alpha_{jk}|^3} \right) + \frac{(\lambda_k - \tilde{\lambda}_k)\alpha_{jk}}{\lambda_k \tilde{\lambda}_k |\alpha_{jk}|^3} \right].$$

By the Taylor formula,

$$\frac{\alpha_{jk}}{|\alpha_{jk}|^3} - \frac{\tilde{\alpha}_{jk}}{|\tilde{\alpha}_{jk}|^3} = \frac{\alpha_{jk} - \tilde{\alpha}_{jk}}{|\tilde{\alpha}_{jk}|^3} - \frac{3\tilde{\alpha}_{jk} \cdot (\alpha_{jk} - \tilde{\alpha}_{jk})}{|\tilde{\alpha}_{jk}|^5} \tilde{\alpha}_{jk} + O(t^{-\frac{8}{3}+6\epsilon}).$$

Using (1.7) and (1.8), there exists a matrix  $A_{jk} \in \mathbb{R}^{3 \times 3}$  such that

$$\frac{\alpha_{jk}}{|\alpha_{jk}|^3} - \frac{\tilde{\alpha}_{jk}}{|\tilde{\alpha}_{jk}|^3} = \frac{A_{jk}}{t^2} (\alpha_{jk} - \tilde{\alpha}_{jk}) + O(t^{-\frac{7}{3}+\frac{\epsilon}{2}}),$$

so there exists  $A_j \in \mathbb{R}^{3 \times 3m}$  such that

$$b_j^{(2)}(P) - b_j^{(2)}(\tilde{P}) = \frac{A_j(\alpha - \tilde{\alpha})^T}{t^2} + O(t^{-\frac{7}{3}+\epsilon}).$$

Set  $A = (A_1^T, \dots, A_m^T)^T$ . Then we can further rewrite (5.5) as

$$(5.6) \quad \begin{cases} \dot{\alpha}(t) - \dot{\tilde{\alpha}}(t) = 2\beta(t) - 2\tilde{\beta}(t), \\ \dot{\beta}(t) - \dot{\tilde{\beta}}(t) = \frac{A(\alpha - \tilde{\alpha})^T}{t^2} + O(t^{-\frac{7}{3}+\epsilon}), \\ \dot{\lambda}(t) - \dot{\tilde{\lambda}}(t) = O(t^{-2}). \end{cases}$$

The following lemma deals with an ODE of the above structure.

**Lemma 5.1.** *Let  $0 < \delta < \kappa$ ,  $n \in \mathbb{N}$ ,  $A \in \mathbb{R}^{n \times n}$  and  $F \in C(\mathbb{R}_+ \times \mathbb{R}^n; \mathbb{R}^n)$ . Assume*

$$(5.7) \quad \sup_{|x| \leq 1} \left( |F(t, x)| + |\nabla_x F(t, x)| \right) \leq t^{-2-\kappa}, \quad \forall t > 0.$$

*Then there exists  $T > 0$  and  $x \in C^2([T, +\infty), \mathbb{R}^n)$  such that*

$$\ddot{x}(t) = \frac{Ax(t)^T}{t^2} + F(t, x(t)) \quad \text{and} \quad |x(t)| \leq t^{-\delta}, \quad \forall t \geq T.$$

*Proof.* We may work instead on  $\mathbb{C}$  by setting  $F(t, z) = F(t, \operatorname{Re}(z))$  for  $z \in \mathbb{C}^n$  and allowing  $x(t)$  to take value in  $\mathbb{C}^n$ . If the complex counterpart is proved, then the lemma follows by taking the real part.

For  $T > 0$  and  $x \in C([T, +\infty), \mathbb{C}^n)$ , define the norm  $\|x\|_3 := \sup_{t \geq T} t^\delta |x(t)|$  and let  $B =$

$\{x \in C([T, +\infty), \mathbb{C}^n) \mid \|x\|_3 \leq 1\}$ . We want to find a solution in  $B$ .

First we consider the case when  $A$  is diagonalizable over  $\mathbb{C}$ . We may take  $n$  linear independent eigenvectors  $v_1, \dots, v_n \in \mathbb{C}^n$  of  $A$ , with eigenvalues  $c_1, \dots, c_n$ , respectively. Let  $a_j, b_j$  be the two roots of  $\lambda^2 - \lambda = c_j$ . Write  $F(t, x) = \sum_{j=1}^n f_j(t, x)v_j$ . Then  $f_j$  also satisfies (5.7). For  $x \in B$ , we define  $\Gamma x$  by

$$(\Gamma x)(t) = \sum_{j=1}^n G_{a_j, b_j} f_j(t, x)v_j,$$

where

$$G_{a,b}f(t, x) = \begin{cases} \frac{G_a f(t, x) - G_b f(t, x)}{a - b}, & a \neq b, \\ \frac{d}{da} G_a f(t, x), & a = b, \end{cases}$$

and

$$G_a f(t, x) = \begin{cases} t^a \int_1^t \tau^{1-a} f(\tau, x(\tau)) d\tau, & \operatorname{Re}(a) \leq -\kappa, \\ -t^a \int_t^\infty \tau^{1-a} f(\tau, x(\tau)) d\tau, & \operatorname{Re}(a) > -\kappa, \end{cases}$$

for any  $a, b \in \mathbb{C}$ , function  $f(\cdot, \cdot)$  satisfying (5.7) and  $x \in B$ .

Note that, if  $f$  satisfies (5.7), then  $G_a f(t, x)$  is well-defined and satisfies

$$|G_a f(t, x)| \leq Ct^{-\kappa}, \quad |G_a f(t, x) - G_a f(t, y)| \leq Ct^{-\kappa-\delta} \|x - y\|_3, \quad \forall x, y \in B.$$

Same results hold for  $\frac{d}{da} G_a f$  with the right hand sides multiplied by  $\log t$ .

Therefore,  $\Gamma$  is a well-define map on  $B$ . Moreover, if  $T$  is large enough, then  $\Gamma$  maps  $B$  to itself and is a contraction. By direct computation, we have

$$\frac{d^2}{dt^2}(\Gamma x) = \frac{A}{t^2}(\Gamma x) + F(t, x),$$

thus the unique fixed point of  $\Gamma$  in  $B$ , guaranteed by the contraction mapping theorem, is the desired  $x \in C([T, +\infty), \mathbb{C}^n)$ .

For the general case, for any  $c_0 > 0$ , there exists  $\tilde{A} \in \mathbb{R}^{n \times n}$  such that  $\tilde{A}$  is diagonalizable over  $\mathbb{C}$ ,  $\|A - \tilde{A}\| \leq c_0$ , and  $-\frac{1}{4}$  is not an eigenvalue of  $\tilde{A}$ . Consider

$$\ddot{x}(t) = \frac{\tilde{A}x(t)^T}{t^2} + \tilde{F}(t, x(t)), \quad \text{where } \tilde{F}(t, x) = F(t, x) + \frac{(A - \tilde{A})x^T}{t^2}.$$

Instead of (5.7), we have

$$\sup_{x \in B} \left( |\tilde{F}(t, x(t))| + |\nabla_x \tilde{F}(t, x(t))| \right) \leq c_0 t^{-2-\delta} + t^{-2-\kappa}, \quad \forall t > 0.$$

We repeat the construction of  $\Gamma$  with  $\tilde{A}$  and  $\tilde{F}$ . Note that there would not appear  $G_{a,b}f$  with  $a = b$  because  $-\frac{1}{4}$  is not an eigenvalue of  $\tilde{A}$ . Thus we will get

$$\|\Gamma x\|_3 \leq Cc_0 \quad \text{and} \quad \|\Gamma x - \Gamma y\|_3 \leq Cc_0 T^{-\delta} \|x - y\|_3, \quad \forall x, y \in B$$

for some  $C > 0$ . We can still conclude upon taking  $c_0$  small enough.  $\square$

Back to the proposition, we easily obtain  $\lambda - \tilde{\lambda} = O(t^{-1})$  by (5.6). Also the leading term of  $O(t^{-\frac{7}{3}+\epsilon})$  does not depend on  $\beta$  because it comes from  $b_j^{(2)}(P) - b_j^{(2)}(\tilde{P})$ . Thus we can obtain the conclusion by a slight modification of the lemma.  $\square$

**5.2. Review of the hyperbolic case.** Now, let us go over the proof of the hyperbolic case and see what has to be changed in the other two cases.

Everything in Section 2 works here, because it does not depend on the dynamics. Proposition 3.1 is replaced by Proposition 5.1. The rest of Section 3 will work because we have only used  $a^{(N)}(t) \rightarrow \infty$  as  $t \rightarrow \infty$ . Therefore, it suffices to prove Proposition 3.3 in the other two settings.

In Section 4, the asymptotic properties (4.3) need to be changed. In the parabolic setting, by (1.7) and (5.1), we have

$$(5.8) \quad a(t) \sim t^{\frac{2}{3}}, \quad |\alpha_j| \lesssim t^{\frac{2}{3}}, \quad |\beta_j| \lesssim t^{-\frac{1}{3}}, \quad |\dot{\beta}_j| \lesssim t^{-\frac{4}{3}}, \quad \lambda_j \sim 1, \quad |\dot{\lambda}_j| \lesssim t^{-\frac{4}{3}}.$$

In the hyperbolic-parabolic setting, for  $J \in M$ , we let  $\alpha_J(t)$ ,  $\beta_J(t)$ ,  $\lambda_J(t)$  be any of  $\alpha_j(t)$ ,  $\beta_j(t)$ ,  $\lambda_j(t)$ ,  $j \in J$ , respectively. Then by (1.8) and (5.1), we have

$$(5.9) \quad \begin{aligned} a(t) &\sim t^{\frac{2}{3}}, \quad |\alpha_J| \lesssim t, \quad |\beta_J| \lesssim 1, \quad |\dot{\beta}_J| \lesssim t^{-\frac{4}{3}}, \quad \lambda_J \sim 1, \quad |\dot{\lambda}_J| \lesssim t^{-\frac{4}{3}}, \\ &|\alpha_j - \alpha_J| \lesssim t^{\frac{2}{3}}, \quad |\beta_j - \beta_J| \lesssim t^{-\frac{1}{3}}, \quad |\lambda_j - \lambda_J| \lesssim t^{-\frac{1}{3}}, \quad \forall j \in J. \end{aligned}$$

With these one can check that all the estimates in Section 4.1, in particular (4.10) and (4.11), hold. This is mainly because we did not use the sharp bounds in (4.3).

However, we need some modification in Section 4.2. Lemma 4.1 is valid for the parabolic case. For the hyperbolic-parabolic case, we prove the following:

**Lemma 5.2.** *There exist  $c, C > 0$  and  $\varphi_j \in C^{1,\infty}(\mathbb{R}_+ \times \mathbb{R}^3)$  for  $1 \leq j \leq m$  such that (4.13) holds and, moreover, for any  $J \in M$ ,*

$$(5.10) \quad |\partial_t \varphi_J| + |\nabla \varphi_J| \leq Ct^{-1}, \quad \text{where } \varphi_J = \sum_{j \in J} \varphi_j.$$

*Proof.* For  $J \in M$ , let  $\alpha_J$  be any of  $\alpha_j$ ,  $j \in J$ . Applying Lemma 4.1 to  $\{\alpha_J | J \in M\}$ , we can find  $\varphi_J \in C^{1,\infty}(\mathbb{R}_+ \times \mathbb{R}^3)$  such that

$$\begin{aligned} 0 \leq \varphi_J(t, x) \leq 1, \quad \sum_{J \in M} \varphi_J(t, x) &\equiv 1, \\ |\partial_t \varphi_J| + |\nabla \varphi_J| \leq Ct^{-1}, \quad |\partial_t \sqrt{\varphi_J}| + |\nabla \sqrt{\varphi_J}| &\leq Ct^{-1}, \\ \varphi_J(t, x) &= \begin{cases} 1, & |x - \alpha_J(t)| \leq ct, \\ 0, & |x - \alpha_K(t)| \leq ct, \quad K \neq J. \end{cases} \end{aligned}$$

Then applying Lemma 4.1 to  $\{\alpha_j | j \in J\}$  for each  $J \in M$ , we find  $\psi_j \in C^{1,\infty}(\mathbb{R}_+ \times \mathbb{R}^3)$  such that

$$\begin{aligned} 0 \leq \psi_j(t, x) \leq 1, \quad \sum_{j \in J} \psi_j(t, x) &\equiv 1, \\ |\partial_t \psi_j| + |\nabla \psi_j| \leq Ct^{-\frac{2}{3}}, \quad |\partial_t \sqrt{\psi_j}| + |\nabla \sqrt{\psi_j}| &\leq Ct^{-\frac{2}{3}}, \\ \psi_j(t, x) &= \begin{cases} 1, & |x - \alpha_j(t)| \leq ct^{\frac{2}{3}}, \\ 0, & |x - \alpha_k(t)| \leq ct^{\frac{2}{3}}, \quad k \neq j. \end{cases} \end{aligned}$$

Finally, take  $\varphi_j = \varphi_J \psi_j$ , where  $J$  contains  $j$ . Then all the conditions are satisfied.  $\square$

It still suffices to prove Proposition 4.1 and 4.2.

We can prove Proposition 4.1 exactly as before. For Proposition 4.2, we need to check (4.18), (4.19) and (4.20). The proof of (4.18) need not to be changed. The difficulty of the other two estimates is that  $|\partial_t \varphi_j| + |\nabla \varphi_j|$  does not have an  $O(t^{-1})$  decay. By checking the previous computation, we need to show

$$(5.11) \quad \sum_{j=1}^m \left( \frac{1}{\lambda_j^2} + |\beta_j|^2 \right) \int \left( \partial_t \varphi_j |\varepsilon|^2 + 2 \nabla \varphi_j \operatorname{Im}(\nabla \varepsilon \bar{\varepsilon}) \right) = O \left( \frac{\|\varepsilon\|_{H^1}^2}{t} \right)$$

and

$$(5.12) \quad \begin{aligned} \sum_{j=1}^m \beta_j \int \left( \nabla \varphi_j \left( 2 |\nabla \varepsilon|^2 + 2 \phi_{\operatorname{Re}(\varepsilon \bar{R})} \operatorname{Re}(\varepsilon \bar{R}) + \phi_{|R|^2} |\varepsilon|^2 \right) \right. \\ \left. + \partial_t \varphi_j \operatorname{Im}(\nabla \varepsilon \bar{\varepsilon}) \right) = O \left( \frac{\|\varepsilon\|_{H^1}^2}{t} \right). \end{aligned}$$



In view of (5.8) and (5.9), we may understand the parabolic case as a special case of the hyperbolic-parabolic case, so we shall focus on the hyperbolic-parabolic case.

Our argument is easier than that in [8]. In fact, it is not clear whether the argument there can be applied here. Using (4.13) and (5.9), we have

$$|\beta_j - \beta_J| \cdot \left( |\partial_t \varphi_j| + |\nabla \varphi_j| \right) \leq \frac{C}{t}.$$

Combining this and (5.10), we derive (5.12). For (5.11), similarly, if we replace  $\lambda_j$  by  $\lambda_J$  and  $\beta_j$  by  $\beta_J$ , then the difference is at most  $O(t^{-1})$ . Finally the terms with  $\lambda_J$  or  $\beta_J$  are controlled using (5.10). We remark that this is the only place we need the assumption on the masses.

We have thus completed the proof of Theorem 2 and 3.

#### ACKNOWLEDGEMENT

The author would like to thank Professor Wilhelm Schlag for useful discussions.

#### REFERENCES

1. V. Barutello, S. Terracini, and G. Verzini, *Entire minimal parabolic trajectories: the planar anisotropic Kepler problem*, Arch. Ration. Mech. Anal. **207** (2013), no. 2, 583–609. MR 3005324
2. V. S. Buslaev and G. S. Perel'man, *On the stability of solitary waves for nonlinear Schrödinger equations*, Nonlinear evolution equations, Amer. Math. Soc. Transl. Ser. 2, vol. 164, Amer. Math. Soc., Providence, RI, 1995, pp. 75–98. MR 1334139
3. T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics, vol. 10, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003. MR 2002047
4. T. Cazenave and P.-L. Lions, *Orbital stability of standing waves for some nonlinear Schrödinger equations*, Comm. Math. Phys. **85** (1982), no. 4, 549–561. MR 677997
5. T. Duyckaerts, C. Kenig, and F. Merle, *Soliton resolution for the radial critical wave equation in all odd space dimensions*, Acta Math. **230** (2023), no. 1, 1–92. MR 4567713
6. D. L. Ferrario and S. Terracini, *On the existence of collisionless equivariant minimizers for the classical  $n$ -body problem*, Invent. Math. **155** (2004), no. 2, 305–362. MR 2031430
7. J. Jendrej and A. Lawrie, *Soliton resolution for the energy-critical nonlinear wave equation in the radial case*, Ann. PDE **9** (2023), no. 2, Paper No. 18, 117. MR 4650926
8. J. Krieger, Y. Martel, and P. Raphaël, *Two-soliton solutions to the three-dimensional gravitational Hartree equation*, Comm. Pure Appl. Math. **62** (2009), no. 11, 1501–1550. MR 2560043
9. E. Lenzmann, *Uniqueness of ground states for pseudorelativistic Hartree equations*, Anal. PDE **2** (2009), no. 1, 1–27. MR 2561169
10. E. H. Lieb, *Existence and uniqueness of the minimizing solution of Choquard's nonlinear equation*, Studies in Appl. Math. **57** (1976/77), no. 2, 93–105. MR 471785
11. P.-L. Lions, *Applications de la méthode de concentration-compacité à l'existence de fonctions extrémales*, C. R. Acad. Sci. Paris Sér. I Math. **296** (1983), no. 15, 645–648. MR 705681
12. J. Liu, D. Yan, and Y. Zhou, *Existence of hyperbolic motions to a class of Hamiltonians and generalized  $N$ -body system via a geometric approach*, Arch. Ration. Mech. Anal. **247** (2023), no. 4, Paper No. 64, 54. MR 4600219
13. E. Maderna and A. Venturelli, *Viscosity solutions and hyperbolic motions: a new PDE method for the  $N$ -body problem*, Ann. of Math. (2) **192** (2020), no. 2, 499–550. MR 4151083
14. Y. Martel, *Asymptotic  $N$ -soliton-like solutions of the subcritical and critical generalized Korteweg-de Vries equations*, Amer. J. Math. **127** (2005), no. 5, 1103–1140. MR 2170139
15. Y. Martel and F. Merle, *Multi solitary waves for nonlinear Schrödinger equations*, Ann. Inst. H. Poincaré C Anal. Non Linéaire **23** (2006), no. 6, 849–864. MR 2271697
16. Y. Martel, F. Merle, and T.-P. Tsai, *Stability and asymptotic stability in the energy space of the sum of  $N$  solitons for subcritical  $gKdV$  equations*, Comm. Math. Phys. **231** (2002), no. 2, 347–373. MR 1946336
17. ———, *Stability in  $H^1$  of the sum of  $K$  solitary waves for some nonlinear Schrödinger equations*, Duke Math. J. **133** (2006), no. 3, 405–466. MR 2228459

18. H. Poincaré, *The three-body problem and the equations of dynamics*, Astrophysics and Space Science Library, vol. 443, Springer, Cham, 2017, Poincaré's foundational work on dynamical systems theory, Translated from the 1890 French original and with a preface by Bruce D. Popp. MR 3642673
19. D. Polimeni and S. Terracini, *On the existence of minimal expansive solutions to the n-body problem*, arXiv: 2310.06360 (2023).
20. T. Tao, *A global compact attractor for high-dimensional defocusing non-linear Schrödinger equations with potential*, Dyn. Partial Differ. Equ. **5** (2008), no. 2, 101–116. MR 2435463
21. ———, *Why are solitons stable?*, Bull. Amer. Math. Soc. (N.S.) **46** (2009), no. 1, 1–33. MR 2457070

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06511

*Email address:* yutong.wu.yw894@yale.edu