# Exact expressions for the maximal probability that all k-wise independent bits are 1

D. Berend<sup>\*†</sup> P. A. Ernst<sup>‡</sup> A. Kontorovich<sup>§</sup> R. Kumar<sup>¶</sup>

### Abstract

Let M(n,k,p) denote the maximum probability of the event  $X_1 = X_2 = \cdots = X_n = 1$  under a k-wise independent distribution whose marginals are Bernoulli random variables with mean p. A long-standing question is to calculate M(n,k,p) for all values of n, k, p. This question has been partially addressed by several authors, primarily with the goal of answering asymptotic questions. The present paper focuses on obtaining exact expressions for this probability. To this end, we provide closed-form formulas of M(n,k,p) for p near 0 as well as p near 1.

# 1 Introduction

Let  $X_1, X_2, \ldots, X_n$  be a sequence of *n* random bits, modelled as *n* Bernoulli random variables with mean *p* for some  $p \in [0, 1]$ . To specify the joint distribution of the random variables, one needs to specify the  $2^n$  probabilities

$$\mathbf{P}(X_1 = \varepsilon_1, X_2 = \varepsilon_2, \dots, X_n = \varepsilon_n), \qquad (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n) \in \{0, 1\}^n.$$

The simplest setting for calculating the joint distribution occurs when the *n* variables are independent. Often, however, there is some form of weak independence between the *n* bits. It then becomes natural to inquire what this weak dependence implies about the distribution of various functions of the  $X_i$ -s. This question was studied by Benjamini et al. [4] with respect to several natural functions. For simplicity, Benjamini et al. [4] assumed that all  $X_i$ -s have the same marginals, namely  $\mathbf{P}(X_i = 1) = p, 1 \leq i \leq n$ , for some  $p \in [0, 1]$ . One of the functions they studied is the product  $\prod_{i=1}^{n} X_i$ . A vector

<sup>\*</sup>Departments of Mathematics and Computer Science, Ben-Gurion University, Beer Sheva 84105, Israel. E-mail: berend@cs.bgu.ac.il

<sup>&</sup>lt;sup>†</sup>Research supported in part by the Milken Families Foundation Chair in Mathematics.

<sup>&</sup>lt;sup>‡</sup>Department of Mathematics, Imperial College London, United Kingdom. E-mail: p.ernst@imperial.ac.uk

<sup>&</sup>lt;sup>§</sup>Department of Computer Science, Ben-Gurion University, Beer Sheva 84105, Israel. E-mail: karyeh@cs.bgu.ac.il

<sup>&</sup>lt;sup>¶</sup>Department of Mathematics, Ben-Gurion University, Beer Sheva 84105, Israel. E-mail: kumarr@post.bgu.ac.il

<sup>2020</sup> Mathematics Subject Classification: Primary 60E15; Secondary 62E99, 90C05, 90C08.

Key words and phrases. Bonferroni inequalities, linear programming problem, lexicographic order.

 $(X_1, \ldots, X_n)$  of random variables is k-wise independent if each subset of k of the random variables is independent. Denote by A(n, k, p) the space of all probability measures on  $\{0, 1\}^n$  that make the  $X_i$ -s k-wise independent with Bernoulli mean p marginals. In other words,  $\mathbf{P} \in A(n, k, p)$  if

$$\mathbf{P}(X_{i_1} = \dots = X_{i_s} = 1) = p^s, \qquad s \leq k, \ 1 \leq i_1 < \dots < i_s \leq n.$$

The authors of Benjamini et al. [4] were especially interested in the maximum M(n, k, p) of the functional  $\mathbf{P}(X_1 = X_2 = \cdots = X_n = 1)$  over the space A(n, k, p). Roughly speaking, they asked how large k should be in order to ensure that this maximum is not much larger than the value of  $p^n$  for the i.i.d. case. This study was continued in [19], where the function M(n, k, p) was specifically studied.

The main focus of [4] and [19] was to obtain both explicit bounds and asymptotic results for M(n, k, p). For example, it was shown in [4, Theorem 23] that for even k

$$M(n,k,p) \leq \frac{p^n}{\mathbb{P}(\operatorname{Bin}(n,1-p) \leq k/2)},$$

where  $\mathbb{P}$  is the fully independent distribution on *n* bits. Lower bounds on M(n, k, p) were given in [19, Theorem 1.1].

The main interest of the present paper is in obtaining exact results for M(n, k, p). Benjamini et al. [4] note that, for k = 2 and k = 3, closed-form expressions for M(n, k, p)over the entire interval [0, 1] can be obtained by invoking the closed-form results in [7] (see Propositions A and B below). An additional closed-form result, established by Benjamini et al. [4, Proposition 35], is that

$$M(n,k,p) = p^k, \qquad 0 \le p \le \frac{1}{n-1}.$$
(1)

We shall generalize this result in Theorem 2.2 of the present paper. To the best of our knowledge, the result in (1) and the exact results for M(n, 2, p) and M(n, 3, p) constitute the only exact results in the literature for M(n, k, p).

The primary purpose of the present paper is to help close this gap by obtaining further exact expressions for M(n, k, p). This paper's key contributions are the closed-form formulas of M(n, k, p) for p near 0 (Theorems 2.2, 2.5) and closed-form formulas for p near 1 (Theorems 2.3, 2.7). We also obtain tight upper bounds on M(n, k, p) (see Theorem 2.1). The second purpose of this paper is to provide observations (based on numerical studies) about M(n, k, p) which we conjecture will hold in general. Our hope is that these observations will help future researchers with the very difficult task of providing exact formulas for M(n, k, p) over the whole interval [0, 1].

A key motivation for the study of k-wise independent distributions arises from the extensive literature in computer science regarding derandomization of algorithms. The objective of derandomization is to analyze how a given algorithm behaves on k-wise independent data in comparison to how it would behave on completely independent data. The use of k-wise independent distributions for derandomization of algorithms was pioneered by [3,9,15,18]. We refer the reader to [2,5,8,13,14,17,23] for a more extensive overview of this literature.

The remainder of the paper is organized as follows. In Section 2 we state our main results. In Section 3 we recall preliminaries needed for the main proofs. In Section 4

we provide these proofs. In Section 5, we conduct numerical studies and present our key observations as eight open conjectures.

# 2 The Main Results

Let  $1 \leq k \leq n$  be fixed integers and  $p \in [0,1]$ . Let A(n,k,p) be the set of all laws **P** making the collection  $\{X_i : 1 \leq i \leq n\}$  into k-wise random bits with mean p. Let M(n,k,p) be the maximum value of  $\mathbf{P}(X_1 = \cdots = X_n = 1)$  over all laws  $\mathbf{P} \in A(n,k,p)$ . A law  $\mathbf{P} \in A(n,k,p)$  is an *optimizing measure* for (n,k,p) if it attains the maximum M(n,k,p).

Clearly, M(n, k, 0) = 0 and M(n, 1, p) = p. Also, M(n, k, 1) = 1 and  $M(n, n, p) = p^n$ . Thus, we will implicitly assume that  $2 \leq k \leq n-1$  and  $p \in (0, 1)$ . We first notice that the set A(n, k, p) is convex since, if  $\mathbf{P}_1$  and  $\mathbf{P}_2$  both make  $\{X_i : 1 \leq i \leq n\}$  into k-wise random bits with mean p, then so does  $\alpha \mathbf{P}_1 + (1 - \alpha)\mathbf{P}_2$ . As A(n, k, p) is compact, there always exists an optimizing measure for (n, k, p). The following lemma shows that we can restrict the search for an optimizing measure to a small subset of A(n, k, p). For self-containedness, we also provide a proof.

**Lemma A.** [19, p.507] There exists an optimizing measure for (n, k, p) that makes  $\{X_i : 1 \leq i \leq n\}$  exchangeable.

Recall that a sequence  $C_1, C_2, \ldots, C_n$  of events is *exchangeable* if, for every  $1 \le t \le n$ , the probabilities  $\mathbf{P}(C_{i_1} \cap C_{i_2} \cap \cdots \cap C_{i_t})$  are the same for every  $1 \le i_1 < i_2 < \cdots < i_t \le n$ .

*Proof.* Let **P** be an optimizing measure for (n, k, p). Under the special case where all k-wise marginals have the same law, k-wise independence is closed under symmetrization. It follows that there is a measure **P'** that makes  $\{X_i : 1 \le i \le n\}$  exchangeable and gives  $\sum_{i=1}^{n} X_i$  the same law as does **P**. Thus  $\mathbf{P}'(\sum_{i=1}^{n} X_i = n) = \mathbf{P}(\sum_{i=1}^{n} X_i = n) = M(n, k, p)$ , so that **P'** is an optimizing measure for (n, k, p).

By Lemma A, the problem is the same if we restrict to exchangeable measures. Let  $A_E(n, k, p)$  be the subset of A(n, k, p), consisting all exchangeable measures. An exchangeable measure **P** is uniquely determined by the quantities

$$v_i = \mathbf{P}\left(\sum_{j=1}^i X_j = i, \sum_{j=i+1}^n X_j = 0\right), \qquad 0 \le i \le n.$$

$$(2)$$

By symmetry, for any set of *i* bits,  $v_i$  is the probability of the event that these bits are all 1 and the rest are all 0. There are  $\binom{n}{i}$  such subsets, and hence  $\sum_{i=0}^{n} \binom{n}{i} v_i = 1$ .

By contrast,

$$\mathbf{P}\left(\sum_{j=1}^{i} X_j = i\right) = \sum_{j=i}^{n} \binom{n-i}{j-i} v_j = p^i, \qquad 0 \le i \le k,$$
(3)

but this relation is no longer (necessarily) true for i > k. Thus, the set of all k-wise independent distributions, whose marginals are all defined by  $\mathbf{P}(X_i = 1) = p$ , obeys the k + 1 equality constraints in (3), in addition to the non-negativity constraints. A subset  $I \subseteq \{0, 1, \ldots, n\}$  is a *weak support* of an exchangeable measure **P** if  $v_i = 0$  for all  $i \in \{0, 1, \ldots, n\} - I$ . The set I is the *support* of **P** if  $v_i > 0$  for each  $i \in I$  and  $v_i = 0$  for all  $i \in \{0, 1, \ldots, n\} - I$ . The following two propositions provide explicit formulas for M(n, 2, p) and M(n, 3, p). These formulas were obtained in [7] (see also [4] and [22]). For completeness, we also present the corresponding optimizing measures as in [22].

**Proposition A.** [22, Theorem 4.1] Let  $n \ge 3$  and  $p \in \left[\frac{j-1}{n-1}, \frac{j}{n-1}\right]$  for some  $1 \le j \le n-1$ . Then,

$$M(n,2,p) = \frac{1}{\binom{n-j+1}{2}} \left( \binom{n}{2} p^2 - n(j-1)p + \binom{j}{2} \right).$$
(4)

Moreover, the optimizing distribution is supported on  $\{j - 1, j, n\}$  and given by

$$v_{i} = \begin{cases} \frac{1}{\binom{n-1}{j-1}} \cdot (1-p) \cdot (j+p-np), & i = j-1, \\ \frac{1}{\binom{n-1}{j}} \cdot (1-p) \cdot (np-p-j+1), & i = j, \\ \frac{1}{\binom{n-j+1}{2}} \cdot \left(\binom{n}{2}p^{2} - n(j-1)p + \binom{j}{2}\right), & i = n. \end{cases}$$
(5)

For self-containedness, we provide a proof of the proposition in Section 4. Once we know the value of M(n, k, p) for a certain even k (for all n), we will also know its value when k is replaced by k + 1. In fact (see [19, p.504])

$$M(n, k+1, p) = pM(n-1, k, p), \qquad k \equiv 0 \pmod{2}, \ n \ge k-2, \ p \in [0, 1].$$
(6)

In particular, Proposition A readily yields a formula for M(n, 3, p). In Proposition B we also provide the optimizing measure.

**Proposition B.** Let  $n \ge 4$  and  $p \in \left[\frac{j-1}{n-2}, \frac{j}{n-2}\right]$  for some  $1 \le j \le n-1$ . Then,

$$M(n,3,p) = p \cdot \frac{1}{\binom{n-j}{2}} \left( \binom{n-1}{2} \cdot p^2 - (n-1)(j-1)p + \binom{j}{2} \right).$$
(7)

Moreover, the optimizing distribution is supported on  $\{0, j, j+1, n\}$  and given by

$$w_{i} = \begin{cases} \frac{1}{\binom{j+1}{2}} \cdot (1-p) \left(\binom{n-1}{2} p^{2} - j(n-1)p + \binom{j+1}{2}\right) \right), & i = 0, \\ p \cdot \frac{1}{\binom{n-2}{j-1}} \cdot (1-p)(j-(n-2)p), & i = j, \\ p \cdot \frac{1}{\binom{n-2}{j-1}} \cdot (1-p)(1-j+(n-2)p), & i = j+1, \\ p \cdot \frac{1}{\binom{n-j}{2}} \left(\binom{n-1}{2} \cdot p^{2} - (n-1)(j-1)p + \binom{j}{2}\right), & i = n. \end{cases}$$

$$(8)$$

Since the proof of the proposition is similar to that of Proposition A, it is omitted.

It will follow from the discussion in Subsection 3.2 that M(n, k, p) is a piecewise polynomial function of degree k in p for every fixed n and k. In Proposition A and B, these functions are presented explicitly for k = 2, 3. We will refer to the (maximal) intervals on which M(n, k, p) is polynomial, starting at p = 0 and continuing towards p = 1, as the *first interval* of M(n, k, p), second interval, and so forth. Notably, for k = 2and k = 3, the intervals have constant lengths of  $\frac{1}{n-1}$  and  $\frac{1}{n-2}$ , respectively. We will see that the intervals are of varying lengths for  $k \ge 4$ .

For integers  $m \ge l \ge 0$ , denote

$$B(m,l,x) = 1 - \binom{m}{l}x + \binom{m}{2}x^2 - \dots + (-1)^l \binom{m}{l}x^l, \qquad x \in \mathbf{R}$$
(9)

where B(m, l, x) = 0 for negative integers l. We refer to these polynomials as *Bonferroni* polynomials for reasons that will be explained in Subsection 3.1.

The following theorem provides upper bounds on M(n, k, p) in terms of Bonferroni polynomials.

### Theorem 2.1.

1. For every k:

$$M(n,k,p) \leq \frac{1}{\binom{n-k+m-1}{m}} \cdot p^{k-m} \cdot B(n-k+m,m,p), \qquad 0 \leq m \leq k, \ m \equiv 0 \ (\text{mod } 2).$$

2. For even k:

$$M(n,k,p) \leqslant B(n,k,1-p).$$

As mentioned in (1), it was proven in Benjamini et al. [4, Proposition 35] that, if  $p \leq \frac{1}{n-1}$ , then  $M(n, k, p) = p^k$ . We generalize this result in Theorem 2.2 below.

**Theorem 2.2.**  $M(n,k,p) = p^k$  if and only if  $p \leq \frac{1}{n-k+1}$ . Moreover, the optimizing distribution for such p is given by

$$v_{i} = \begin{cases} (1-p) \cdot p^{i} \cdot B(n-i-1,k-i-1,p), & 0 \leq i \leq k-1, \\ 0, & k \leq i \leq n-1, \\ p^{k}, & i = n. \end{cases}$$
(10)

For general k, Theorem 2.2 states the value of M(n, k, p) on the first interval (i.e., near 0). We will also find the value of M(n, k, p) on the second interval (Theorem 2.5) and on the last two intervals (Theorem 2.3 and 2.7).

**Theorem 2.3.** Let  $2 \le k \le n-1$  be positive integers with k even. Then

$$M(n,k,p) = B(n,k,1-p), \qquad 1 - \frac{1}{n-k+1} \le p \le 1.$$
(11)

Moreover, the optimizing distribution is given by

$$v_i = \begin{cases} 0, & 0 \le i \le n - k - 1, \\ (1 - p)^{n - i} \cdot B(i, i - (n - k), 1 - p), & n - k \le i \le n. \end{cases}$$
(12)

From Theorem 2.2 and 2.3, we obtain

### Corollary 2.4.

$$M(n, n-1, p) = \begin{cases} p^{n-1}, & p \le 1/2, \\ B(n, n-1, 1-p), & p > 1/2. \end{cases}$$

For  $p \leq 1/2$ , the optimal distribution is given by (10). For p > 1/2, it is given by (12).

We now proceed to state Theorem 2.5.

### Theorem 2.5.

1. Let  $2 \leq k \leq n-2$ , and

$$p_1 = \frac{1}{n-k+1}.$$

Then, for some  $p_2 > p_1$ ,

$$M(n,k,p) = \frac{1}{\binom{n-k+1}{2}} \cdot p^{k-2} \cdot B(n-k+2,2,p), \qquad p \in [p_1,p_2].$$
(13)

Moreover, the optimizing distribution is given by

$$v_{i} = \begin{cases} p^{i}B(n-i,k-i,p) + (-1)^{k-i-1}\frac{\binom{n-i-1}{k-i}}{\binom{n-k+1}{2}}p^{k-2}B(n-k+2,2,p), & i \in I - \{n\}, \\ 0, & i \notin I, \\ \frac{1}{\binom{n-k+1}{2}} \cdot p^{k-2} \cdot B(n-k+2,2,p), & i = n, \end{cases}$$
(14)

where  $I = \{0, 1, \dots, k - 3, k - 1, k, n\}.$ 

- 2. The largest  $p_2$  for which (13) and (14) hold is
  - $\frac{2}{n-1} \text{for } k = 2;$ •  $\frac{2}{n-2} - \text{for } k = 3;$
  - the zero of the cubic polynomial

$$1 - \binom{n-k+3}{1}p + \frac{5}{6}\binom{n-k+3}{2}p^2 - \frac{1}{2}\binom{n-k+3}{3}p^3$$
(15)

in the interval  $\left[\frac{1}{n-k+1}, \frac{2}{n-k+1}\right] - \text{for } k \ge 4.$ 

We mention that, using Lemma 3.1.1 below, we may rewrite the first case in (14) alternatively in the form

$$v_i = (1-p) \left( p^i B(n-i-1,k-i-1,p) + (-1)^{k-i-1} \frac{\binom{n-i-1}{k-i}}{\binom{n-k+1}{2}} p^{k-2} B(n-k+1,1,p) \right),$$
  
for  $i \in I - \{n\}.$ 

**Remark 2.6.** In view of Theorems 2.2 and 2.5, the upper bound in Theorem 2.1.1 is tight for m = 0, 2, and by Theorem 2.3 the same holds for Theorem 2.1.2. M(n, k, p) is given by the bound with m = 0 in Theorem 2.1.1 on the first interval, by the bound with m = 2 on the second interval, and by the bound in Theorem 2.1.2 on the last interval. In all of the examples we have checked, M(n, k, p) coincided with the bound for m = 4on the third interval. However, M(n, k, p) only sometimes coincided with the bound for m = 6 on the fourth interval. For example, for n = 10, k = 6, in the first four intervals, M(n, k, p) is given by the bounds as in Theorem 2.1.1 for m = 0, 2, 4, 6, respectively, but for n = 11, only the first three are as in Theorem 2.1.1, but not the 4-th, as follows:

$$M(11, 6, p) = \begin{cases} p^{6}, & 0 (16)$$

The first three equalities in (16) are the same as the bound in Theorem 2.1.1, namely

$$\frac{1}{\binom{n-k+m-1}{m}} \cdot p^{k-m} B(n-k+m,m,p),$$

for m = 0, 2, 4, respectively, while the fourth equality is different from that bound for m = 6, which is

$$\frac{1}{\binom{n-k+m-1}{m}} \cdot p^{k-m} B(n-k+m,m,p) = \frac{1}{120} (1-11p+55p^2-165p^3+330p^4-462p^5+462p^6).$$

The following theorem deals with the second to last interval.

### Theorem 2.7.

1. For k even let  $2 \leq k \leq n-2$  be positive integers and

$$\overline{p}_1 = 1 - \frac{1}{n-k+1}$$

Then, for some  $\overline{p}_2 < \overline{p}_1$ ,

$$M(n,k,p) = B(n,k,1-p) + \frac{\binom{n}{k+1}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1} B(n-k+1,1,1-p), \qquad p \in [\overline{p}_2,\overline{p}_1].$$
(17)

Moreover, the optimizing distribution is given by

$$v_{n-i} = \begin{cases} (1-p)^{i}B(n-i,k-i,1-p) - \\ (-1)^{k-i-1}\frac{\binom{n-i}{k+1-i}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1}B(n-k+1,1,1-p), & i \in (n-I), \\ 0 & i \notin n-I, \end{cases}$$
(18)

where  $I = \{n - k - 1, n - k, n - k + 2, \dots, n - 1, n\}.$ 

- 2. The smallest  $\overline{p}_2$  for which (13) and (18) hold is
  - $\frac{2}{n-1} for \ k = 2;$
  - the zero of the cubic polynomial

$$1 - \binom{n-k+3}{1}(1-p) + \frac{5}{6}\binom{n-k+3}{2}(1-p)^2 - \frac{1}{2}\binom{n-k+3}{3}(1-p)^3 \quad (19)$$
  
in the interval  $\left[1 - \frac{2}{n-k+1}, 1 - \frac{1}{n-k+1}\right] - for \ k \ge 4.$ 

We conclude this section with two remarks.

### Remark 2.8.

- (1) Unlike the polynomials describing M(n, k, p) on the first, second, and last intervals, which form upper bounds on M(n, k, p) on the entire interval [0, 1], the polynomial on the right-hand side of (17) is not an upper bound for M(n, k, p). In all of the examples we have checked, the polynomial is negative on some sub-interval of [0, 1].
- (2) It follows from (15) and (19) that the smallest  $\overline{p}_2$  in Theorem 2.7 and the largest  $p_2$  in Theorem 2.5 are symmetric with respect to the point p = 1/2, namely  $\overline{p}_2 = 1 p_2$ .

**Remark 2.9.** Theorems 2.3, 2.7, and the numerical data suggest that, for even k between 2 and n-2, there exists a point  $\overline{p}_3 < \overline{p}_2$  such that

$$M(n,k,p) = B(n,k,1-p) + \frac{\binom{n}{k+1}}{\binom{n-k+3}{4}} (1-p)^{k-3} B(n-k+3,3,1-p), \qquad p \in [\overline{p}_3,\overline{p}_2],$$

where the endpoints  $\overline{p}_2, \overline{p}_3$  are symmetric, respectively, to the endpoints  $p_2, p_3$  of the third interval with respect to the point p = 1/2.

This concludes the section of the main results, all of which will be proven in Section 4.

# 3 Preliminaries

In this section, we recall the necessary preliminaries for the proofs in Section 4.

### 3.1 Bonferroni Inequalities

Let  $A_1, A_2, \ldots, A_n$  be arbitrary events in a probability space  $(\Omega, \mathcal{B}, \mathbf{P})$ . Let N be the (random) number of events  $A_i$ -s that occur. Denote

$$S_t = \sum_{1 \le i_1 < i_2 < \dots < i_t \le n} \mathbf{P}(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_t}), \qquad t = 1, 2, \dots, n,$$
(20)

where  $S_0 = 1$ . Bonferroni's inequalities, first proved by Bonferroni [6] (see also [24]), state that  $2m+1 \qquad 2m$ 

$$\sum_{i=0}^{2m+1} (-1)^i S_i \leqslant \mathbf{P}(N=0) \leqslant \sum_{i=0}^{2m} (-1)^i S_i, \qquad m \ge 0.$$
(21)

Inequalities of the form

$$\sum_{i=0}^{n} c_i S_i \leqslant \mathbf{P}(N=r) \leqslant \sum_{i=0}^{n} d_i S_i,$$
(22)

where  $c_i = c_i(r, n)$  and  $d_i = d_i(r, n)$  are constants (unrelated to the specific events  $A_j$ ,  $1 \leq j \leq n$ ), possibly zero, are *Bonferroni-type inequalities* (see [10]). Such inequalities have a long history (cf. [6, 10, 12, 16, 20]). An extension of (21) is given by *Jordan's inequalities*, which are as follows (cf. [16, p.8])

$$\sum_{i=0}^{2m+1} (-1)^i \binom{i+r}{r} S_{i+r} \leq \mathbf{P}(N=r) \leq \sum_{i=0}^{2m} (-1)^i \binom{i+r}{r} S_{i+r}, \qquad 0 \leq r \leq n, \ m \geq 0.$$

It follows from [10, Theorem 2] that (22) holds for arbitrary events  $A_1, A_2, \ldots, A_n$  if and only it holds for exchangeable events. In other words, the inequality in (22) holds for arbitrary  $A_1, A_2, \ldots, A_n$  if and only if

$$\sum_{i=0}^{n} c_i \binom{n}{k} P_k \leqslant \mathbf{P}(N'=r) \leqslant \sum_{i=0}^{n} d_i \binom{n}{k} P_k$$

for arbitrary exchangeable events  $C_1, C_2, \ldots, C_n$ , where N' is the number of events  $C_i$  that occur and  $P_k = \mathbf{P}(C_{i_i} \cap C_{i_2} \cap \cdots \cap C_{i_k})$ . Furthermore, the inequality in (22) holds for arbitrary events  $A_1, A_2, \ldots, A_n$  if and only if it holds when the  $A_j$ 's are independent and  $P(A_j) = p$  for each  $1 \leq j \leq n$  (see [10, Theorem 3]).

Let  $X_1, X_2, \ldots, X_n$  be **P**-distributed random bits, where  $\mathbf{P} \in A_E(n, k, p)$ , namely  $\mathbf{P}(X_i = 1) = p$  and the  $X_i$ -s are k-wise independent. Let  $A_i = \{X_i = 1\}, 1 \leq i \leq n$ . By (20),

$$S_t = \binom{n}{t} p^t, \qquad t = 1, 2, 3, \dots, k.$$
(23)

Employing (21) and (23), we obtain

$$\mathbf{P}(N=0) \begin{cases} \leq B(n,k,p) = 1 - \binom{n}{1}p + \dots + \binom{n}{k}p^k, & k \equiv 0 \pmod{2}, \\ \geq B(n,k,p) = 1 - \binom{n}{1}p + \dots - \binom{n}{k}p^k, & k \equiv 1 \pmod{2}, \end{cases}$$
(24)

which explains the term "Bonferroni polynomials" introduced in Section 2. We refer the reader to [12] for further details on Bonferroni inequalities.

Several useful properties of Bonferroni polynomials are listed in the following lemma. So as to ease the flow of presentation, the proof is postponed to Section 4.

**Lemma 3.1.** For any positive integers  $0 \le k \le n$ ,

1. 
$$B(n,k,p) = B(n-1,k,p) - pB(n-1,k-1,p)$$
$$= (1-p)B(n-1,k-1,p) + (-1)^k \binom{n-1}{k} p^k.$$
2. 
$$\frac{d}{dp}B(n,k,p) = -nB(n-1,k-1,p).$$

3.

$$B(n,k,p) = B(k,k,p) - \sum_{i=0}^{k-1} \binom{n-k}{k-i} p^{k-i} B(n-k+i,i,p).$$
(25)

Another essential inequality concerning the Bonferroni polynomials, crucial for us in the sequel, is presented in the following proposition. So as to ease the flow of presentation, its proof is postponed to Section 4.

**Proposition 3.2.** Let  $0 \leq l \leq m$ . Then

$$B(m,l,p) \ge 0, \qquad 0 \le p \le \frac{1}{m-l+1}.$$
(26)

Moreover, the inequality in (26) is strict except for the following cases:

- 1. l = 1 and p = 1/m.
- 2. *m* is odd, l = m 1, and p = 1/2.
- 3. l = m and p = 1.

# **3.2** M(n,k,p) as the Solution of a Linear Programming Problem

Let  $X_1, X_2, \ldots, X_n$  be **P**-distributed random bits, where  $\mathbf{P} \in A_E(n, k, p)$ . Let  $A_i = \{X_i = 1\}, 1 \leq i \leq n$ . Let N be the number of events  $A_i$  which occur. By (2) and (3), one may recast the problem of calculating M(n, k, p) for given n, k, p as a primal linear programming problem, as follows:

 $\begin{array}{ll} \text{maximize} & v_n \\ \text{subject to} \end{array}$ 

$$\sum_{j=0}^{n} {n-i \choose j-i} v_j = p^i, \qquad i = 0, 1, 2, \dots, k,$$

$$v_j \ge 0, \qquad j = 0, 1, 2, \dots, n.$$
(27)

Letting

$$w_i = \binom{n}{i} v_i, \qquad i = 0, 1, 2, \dots, n,$$
$$S_i = \binom{n}{i} p^i, \qquad i = 0, 1, 2, \dots, k,$$

and

we may rewrite 
$$(27)$$
 in the form:

 $\begin{array}{ll} \text{maximize} & w_n \\ \text{subject to} \end{array}$ 

$$\sum_{j=0}^{n} {j \choose i} w_{j} = S_{i}, \qquad i = 0, 1, 2, \dots, k,$$

$$w_{j} \ge 0, \qquad j = 0, 1, 2, \dots, n.$$
(28)

We now pause to introduce some notation which will enable us to rewrite (28) in a more compact form.

$$a_{i} = \left(1, i, {\binom{i}{2}}, \dots, {\binom{i}{k}}\right)^{T} \in \mathbf{R}^{k+1}, \qquad i = 0, 1, 2, \dots, n,$$

$$A = [a_{0}, a_{1}, \dots, a_{n}] \in M_{(k+1) \times (n+1)},$$

$$w = (w_{0}, w_{1}, \dots, w_{n})^{T},$$

$$c = (0, 0, \dots, 0, 1)^{T} \in \mathbf{R}^{n+1},$$

$$d = (S_{0}, S_{1}, \dots, S_{k})^{T} \in \mathbf{R}^{n+1},$$
(29)

where T denotes matrix transpose. With this notation, (28) becomes

maximize 
$$c^T \cdot w$$
  
subject to (30)

The dual of (30) is

$$\begin{array}{ll}\text{minimize} & d^T \cdot y\\ \text{subject to} & (31) \end{array}$$

 $A^T \cdot y \ \geqslant \ c.$ 

We now mention some definitions and recall some key results of Boros and Prékopa [7], and Prékopa [20,21], who considered a more general problem than (30). With slight abuse of notation, let B be a basis consisting of k + 1 columns of A. Then B is *dual feasible* if

$$c_B^T B^{-1} a_j \ge c_j, \qquad j \in \{0, 1, \dots, n\} - I,$$
(32)

where I is the set of subscripts of the basis vectors and  $c_B$  is the vector of basic components of c. The basis B is *dual nondegenerate* if the inequality in (32) is strict for every j.

Let  $I = \{i_0, i_1, \dots, i_k\}, 0 \le i_0 < i_1 < \dots < i_k \le n$ . Denote

$$B(I) = \lfloor a_{i_0}, a_{i_1}, \dots, a_{i_k} \rfloor.$$

It follows from [7, Theorem 7.1] that

$$B(I)^{-1} = [b_{s,t}]_{s \in I}^{t=0,1,\dots,k},$$

where

$$b_{s,t} = \frac{(-1)^t \sum_{\alpha=0}^t (-1)^\alpha {t \choose \alpha} \prod_{u \in I - \{s\}} (u - \alpha)}{\prod_{u \in I - \{s\}} (u - s)}, \qquad s \in I, \ 0 \le t \le k.$$
(33)

**Theorem 3.3.** [7, Theorem 7.2] A basis B(I), corresponding to problem (30), is dual feasible if and only if I satisfies the following conditions:

- (1)  $n \in I$ .
- (2) If k is odd then  $0 \in I$ .
- (3) The elements of  $I \{n\}$  in the case of even k, and of  $I \{0, n\}$  in the case of odd k, come in pairs of consecutive numbers. Namely,

$$I = \begin{cases} \{i_1, i_1 + 1, i_2, i_2 + 1, \dots, i_{k/2}, i_{k/2} + 1, n\}, & k \equiv 0 \pmod{2}, \\ \{0, i_1, i_1 + 1, i_2, i_2 + 1, \dots, i_{(k-1)/2}, i_{(k-1)/2} + 1, n\}, & k \equiv 1 \pmod{2}, \end{cases}$$
(34)

where  $i_{j+1} \ge i_j + 2$  for each j.

**Theorem 3.4.** [7, Theorem 7.3], [12, Theorem IV.1.] Let  $I \subseteq \{0, 1, 2, ..., n\}$  be a set of size k + 1 of the type in (34). Then

$$M(n,k,p) \leqslant \sum_{i=0}^{k} b_{n,i} S_i, \tag{35}$$

where the  $b_{n,i}$  are as in (33). Moreover, If  $B(I)^{-1}d \ge 0$ , the bound in (35) is tight, and therefore

$$M(n,k,p) = \sum_{i=0}^{k} b_{n,i} S_i.$$

**Theorem 3.5.** [21, Theorem 6] Let B(I) be a dual feasible basis corresponding to (30). For  $i \in I$ , if all the bases  $B(I_{i,j})$ , where

$$I_{i,j} = (I - \{i\}) \cup \{j\}, \qquad j \in \{0, 1, \dots, n\} - I,$$

are not dual feasible basis, then we will have  $w_i \ge 0$ .

All dual feasible bases for (30) are dual nondegenerate [21, p.232]. Moreover, the optimal distribution is unique [21, Theorem 5]. We refer the reader to [12, Chapter IV] for more details on dual feasibility of the basis B(I) and related results.

# 4 Proofs

We first recall for later reference several well-known identities involving binomial coefficients (see [11, Lemma 1.4.2] and [12, p.14]):

(1)

$$\sum_{i=0}^{t} (-1)^{i} \binom{s}{i} = (-1)^{t} \binom{s-1}{t}, \qquad s \ge 1, \ t \ge 0,$$
(36)

(2)

$$\sum_{i=0}^{s} (-1)^{i} \binom{s}{i} = 0, \qquad s \ge 0, \tag{37}$$

(3)

$$\binom{n}{i}\binom{n-i}{j-i} = \binom{j}{i}\binom{n}{j}, \qquad 0 \le i \le j \le n.$$
(38)

# 4.1 Proof of Theorem 2.1

1. For  $p \in [0, 1]$ , we have

$$v_{0} + \binom{n}{1}v_{1} + \binom{n}{2}v_{2} + \dots + \binom{n}{k}v_{k} + \dots + \binom{n}{n}v_{n} = 1,$$

$$v_{1} + \binom{n-1}{1}v_{2} + \dots + \binom{n-1}{k-1}v_{k} + \dots + \binom{n-1}{n-1}v_{n} = p, \qquad (39)$$

$$\vdots$$

$$v_{k} + \dots + \binom{n-k}{n-k}v_{n} = p^{k}.$$

We proceed to multiply the last (m + 1) equalities in (39) by, respectively,

$$\binom{n-k+m}{0}, -\binom{n-k+m}{1}, \binom{n-k+m}{2}, -\binom{n-k+m}{3}, \dots, \binom{n-k+m}{m}.$$

We then add up the resulting equalities, which gives an equality of the form

$$c_{k-m}v_{k-m} + c_{k-m+1}v_{k-m+1} + \dots + c_kv_k + \dots + c_nv_n = \sum_{i=0}^m (-1)^i \binom{n-k+m}{i} \cdot p^{k-m+i}$$
$$= p^{k-m} \cdot B(n-k+m,m,p).$$
(40)

This equality will yield an upper bound on  $v_n$  if we find  $c_n$  and show that all other coefficients  $c_l$  on the left-hand side of (40) are non-negative. Using (36), (37), and (38), we clearly have:

•

$$c_{k-m} = 1. \tag{41}$$

• The following m coefficients vanish because

$$c_{k-m+j} = \sum_{i=0}^{j} (-1)^{i} \binom{n-k+m}{i} \binom{n-(k-m+i)}{j-i}$$
$$= \sum_{i=0}^{j} (-1)^{i} \binom{j}{i} \binom{n-k+m}{j} \qquad j = 1, 2, \dots, m.$$
(42)
$$= \binom{n-k+m}{j} \cdot \sum_{i=0}^{j} (-1)^{i} \binom{j}{i} = 0,$$

• The final n - k coefficients are strictly positive:

$$c_{k+j} = \sum_{i=0}^{m} (-1)^{i} {\binom{n-k+m}{i}} {\binom{n-(k-m+i)}{k+j-(k-m+i)}} = \sum_{i=0}^{m} (-1)^{i} {\binom{n-k+m}{i}} {\binom{n-k+m-i}{j+m-i}} = \sum_{i=0}^{m} (-1)^{i} {\binom{j+m}{i}} {\binom{n-k+m}{j+m}} = {\binom{n-k+m}{j+m}} \cdot \sum_{i=0}^{m} (-1)^{i} {\binom{j+m}{i}} = {\binom{n-k+m}{j+m}} \cdot {\binom{j+m-1}{m}} > 0.$$
(43)

In particular,  $c_n = \binom{n-k+m-1}{m}$ , and therefore

$$v_n \leqslant \frac{p^{k-m}}{\binom{n-k+m-1}{m}} \cdot B(n-k+m,m,p).$$

Hence,

$$M(n,k,p) \leq \frac{p^{k-m}}{\binom{n-k+m-1}{m}} \cdot B(n-k+m,m,p).$$

2. Recall that

$$M(n,k,p) = \max_{\mathbf{P} \in A_E(n,k,p)} \mathbf{P}(X_1 = \dots = X_n = 1).$$
(44)

Let  $A_i = \{X_i = 1\}, 1 \le i \le n$ . It follows from (20) and (21) that

$$\mathbf{P}(X_{1} = \dots = X_{n} = 1) \leq \sum_{t=0}^{k} (-1)^{t} \sum_{1 \leq i_{1} < i_{2} < \dots < i_{t} \leq n} \mathbf{P}(A_{i_{1}}^{C} \cap A_{i_{2}}^{C} \cap \dots \cap A_{i_{t}}^{C})$$

$$= \sum_{t=0}^{k} (-1)^{t} \binom{n}{t} (1-p)^{t} = B(n,k,1-p).$$
(45)

By (44) and (45),

 $M(n,k,p) \leqslant B(n,k,1-p).$ 

This concludes the proof.

# 4.2 Proof of Lemma 3.1

1. We have

$$B(n,k,p) = \sum_{i=0}^{k} (-1)^{i} {n \choose i} p^{i} = \sum_{i=0}^{k} (-1)^{i} \left( {n-1 \choose i} + {n-1 \choose i-1} \right) p^{i}$$
  

$$= \sum_{i=0}^{k} (-1)^{i} {n-1 \choose i} p^{i} + \sum_{i=0}^{k} (-1)^{i} {n-1 \choose i-1} p^{i}$$
  

$$= \sum_{i=0}^{k} (-1)^{i} {n-1 \choose i} p^{i} + \sum_{i=0}^{k-1} (-1)^{i+1} {n-1 \choose i} p^{i+1}$$
  

$$= B(n-1,k,p) - pB(n-1,k-1,p).$$
(46)

Simplifying the right-hand side of (46) further, we obtain

$$B(n,k,p) = B(n-1,k-1,p) + (-1)^k \binom{n-1}{k} p^k - pB(n-1,k-1,p)$$
  
=  $(1-p)B(n-1,k-1,p) + (-1)^k \binom{n-1}{k} p^k.$  (47)

2. We have

$$\frac{d}{dp}B(n,k,p) = \frac{d}{dp}\left(\sum_{i=0}^{k}(-1)^{i}\binom{n}{i}p^{i}\right) = \sum_{i=0}^{k}(-1)^{i}\binom{n}{i}ip^{i-1} = \sum_{i=1}^{k}(-1)^{i}\binom{n-1}{i-1}p^{i-1}$$
$$= -n\sum_{i=0}^{k-1}(-1)^{i}\binom{n-1}{i}p^{i} = -nB(n-1,k-1,p).$$

3. We use induction on k. For k = 0, both sides of (25) are 1. Assume that (25) holds with k-1 instead of k. Note that both sides of (25) are polynomials of degree at most k in p and they coincide at p = 0. Hence, it suffices to prove that the derivatives with respect to p of both sides of (25) coincide, namely that

$$\frac{d}{dp}B(n,k,p) = \frac{d}{dp}\left((1-p)^k - \sum_{i=0}^{k-1} \binom{n-k}{k-i} p^{k-i}B(n-k+i,i,p)\right).$$
(48)

By the preceding part, the right-hand side of (48) is

$$\begin{split} &-k(1-p)^{k-1} - \sum_{i=0}^{k-1} \binom{n-k}{k-i} (k-i)^{p^{k-1-i}} B(n-k+i,i,p) \\ &+ \sum_{i=0}^{k-1} \binom{n-k}{k-i} (n-k+i) p^{k-i} B(n-k+i-1,i-1,p) \\ &= -k(1-p)^{k-1} - \sum_{i=0}^{k-2} \binom{n-k}{k-i} (k-i) p^{k-1-i} B(n-k+i,i,p) - (n-k) B(n-1,k-1,p) \\ &+ \sum_{i=0}^{k-2} \binom{n-k}{k-(i+1)} (n-k+i+1) p^{k-(i+1)} B(n-k+i,i,p) \\ &+ \binom{n-k}{k} (n-k) p^k B(n-k-1,-1,p) \\ &= -\sum_{i=0}^{k-2} \left( \binom{n-k}{k-i} (k-i) - \binom{n-k}{k-1-i} (n-k+i+1) \right) p^{k-1-i} B(n-k+i,i,p) \\ &- k(1-p)^{k-1} - (n-k) B(n-1,k-1,p) + 0 \\ &= -k(1-p)^{k-1} - \sum_{i=0}^{k-2} (-k) \binom{n-k}{k-1-i} p^{k-1-i} B(n-k+i,i,p) \\ &- (n-k) B(n-1,k-1,p) \\ &= -k \left( (1-p)^{k-1} - \sum_{i=0}^{k-2} \binom{n-k}{k-1-i} \right) p^{k-1-i} B(n-k+i,i,p) \\ &- (n-k) B(n-1,k-1,p) \\ \end{split}$$

By the induction hypothesis, the last expression reduces to

$$-kB(n-1, k-1, p) - (n-k)B(n-1, k-1, p) = -nB(n-1, k-1, p).$$

In view of part 2 of the lemma, this proves (48).

# 4.3 Proof of Proposition 3.2

It follows from [11, Lemma 1.3.2] (see also [10, p.562]) that, for even l,

$$\binom{m}{1}p - \binom{m}{2}p^2 + \binom{m}{3}p^3 - \dots - \binom{m}{l}p^l \le 1 - (1-p)^m.$$
(49)

Thus, for even l, we have  $B(m, l, p) \ge (1 - p)^m$  for all  $0 \le p \le 1$ . It remains to prove the proposition for odd l. We first deal with a few simple cases.

1. If l = 1, then B(m, l, p) = 1 - mp, which is trivially non-negative for  $p \leq \frac{1}{m-1+1} = \frac{1}{m}$ , and vanishes only at  $p = \frac{1}{m}$ .

- 2. If l = m-1, then  $B(m, l, p) = (1-p)^m p^m$ . Since  $p \leq \frac{1}{2}$ , this is certainly nonnegative, vanishing only at p = 1/2.
- 3. If l = m, then  $B(m, l, p) = (1 p)^m$ , which is trivially non-negative, vanishing only at p = 1.

Now, let l be odd,  $3 \leq l \leq m - 2$ . Let

$$p_1 = \frac{1}{m-k+1}.$$

It follows from Lemma 3.1.2 (since l-1 is even) that  $\frac{d}{dp}B(m, l, p)$  is negative, and hence it suffices to check that the required inequality is satisfied and is strict at the right endpoint of the interval, namely that

$$1 - \binom{m}{1}p_1 + \binom{m}{2}p_1^2 - \dots - \binom{m}{l}p_1^l > 0.$$

Since l - 1 is even, we have by [11, Lemma 1.3.2]

$$\binom{m}{1}p_1 - \binom{m}{2}p_1^2 + \binom{m}{3}p_1^3 - \dots - \binom{m}{l-1}p_1^{l-1} \le 1 - (1-p_1)^m,$$

which yields

$$1 - \binom{m}{1}p_1 + \binom{m}{2}p_1^2 - \binom{m}{3}p_1^3 + \dots + \binom{m}{l-1}p_1^{l-1} - \binom{m}{l}p_1^l \ge (1-p_1)^m - \binom{m}{l}p_1^l.$$
(50)

Thus, it suffices to show that for odd l

$$\binom{m}{l}p_1^l < (1-p_1)^m,$$

which is equivalent to

$$\frac{m \cdot (m-1) \cdots (m-l+1)}{l! (m-l)^l} < \left(\frac{m-l}{m-l+1}\right)^{m-l}.$$
(51)

The right-hand side of (51) is at least 1/e, so that it suffices to show that

$$\frac{m-l+1}{1\cdot(m-l)} \cdot \frac{m-l+2}{2\cdot(m-l)} \cdot \frac{m-l+3}{3\cdot(m-l)} \cdot \dots \cdot \frac{m}{l\cdot(m-l)} = \prod_{j=1}^{l} \left(\frac{1}{j} + \frac{1}{m-l}\right) < \frac{1}{e}.$$
 (52)

We show this by distinguishing between several cases:

1.  $l \ge 7$ : We have

$$\prod_{j=1}^{l} \left( \frac{1}{j} + \frac{1}{m-l} \right) \leqslant \prod_{j=1}^{7} \left( \frac{1}{j} + \frac{1}{2} \right) = \frac{9}{32} < \frac{1}{e}.$$

- 2. l = 3, 5 and  $m \ge 10$ : For every fixed l, as m increases the product in (52) consists of the same number of factors, but they become smaller and smaller. Therefore, if (52) holds for some (m, l), then it must hold for  $(m + 1, l), (m + 2, l), \ldots$  We verified (52) numerically for m = 10, and l = 3 and l = 5; hence it must be true for  $m \ge 10$ .
- 3. l = 3, 5 and m < 10: It is easily verified that (26) holds (with a strict inequality) via straightforward calculation.

# 4.4 Proof of Theorem 2.2

Suppose first that  $M(n, k, p) \equiv v_n = p^k$ . By (3), all the  $v_i, k \leq i \leq n-1$ , must vanish. We need to show that

$$v_i = (1-p) \cdot p^i B(n-i-1,k-i-1,p), \qquad 0 \le i \le k-1.$$

Using the generalized inclusion-exclusion principle (see [1, Corollary 5.2]) and the fact that  $\mathbf{P}$  is an exchangeable measure, we have

$$v_i = \sum_{j=i}^n (-1)^{j-i} \binom{n-i}{j-i} \mathbf{P}(X_1 = X_2 = \dots = X_j = 1), \qquad i = 0, 1, \dots, n.$$
(53)

Since the  $X_i$ -s are k-wise independent,

$$\mathbf{P}(X_1 = X_2 = \dots = X_j = 1) = p^j, \qquad j = 0, 1, \dots, k.$$
(54)

For  $j \ge k$ ,

$$= p^{k} = \mathbf{P}(X_{1} = \dots = X_{k} = 1) \ge \mathbf{P}(X_{1} = \dots = X_{j} = 1) \ge \mathbf{P}(X_{1} = \dots = X_{n} = 1) = p^{k}$$

so that

$$\mathbf{P}(X_1 = X_2 = \dots = X_j = 1) = p^k, \qquad j = k+1, k+2, \dots, n.$$
(55)

By (53), (54), and (55), we have for  $0 \le i < k$ 

$$\begin{aligned} v_{i} &= p^{i} - (n-i)p^{i+1} + \binom{n-i}{2}p^{i+2} - \binom{n-i}{3}p^{i+3} + \cdots \\ &+ (-1)^{k-i-1}\binom{n-i}{k-i-1}p^{k-1} + (-1)^{k-i}\binom{n-i}{k-i}p^{k} + (-1)^{k-i+1}\binom{n-i}{k-i+1}p^{k} + \cdots \\ &+ (-1)^{n-i}\binom{n-i}{n-i}p^{k} \\ &= p^{i} - (n-i)p^{i+1} + \binom{n-i}{2}p^{i+2} - \binom{n-i}{3}p^{i+3} + \cdots \\ &+ (-1)^{k-1-i}\binom{n-i}{k-i-1}p^{k-1} + (-1)^{k-i}\binom{n-i-1}{k-i-1}p^{k}, \end{aligned}$$
(56)

where in the last step we have relied on the identity

$$\binom{m}{j} - \binom{m}{j+1} + \binom{m}{j+2} - \dots + (-1)^{m-j} \binom{m}{m} = \binom{m-1}{j-1}.$$

By (9), (56), and Lemma 3.1.1,

$$v_{i} = p^{i}B(n-i, k-i-1, p) + (-1)^{k-i} \binom{n-i-1}{k-i-1} p^{k}$$
  
=  $p^{i}(B(n-i-1, k-i-1, p) - pB(n-i-1, k-i-2, p))$   
 $- p^{i+1}(B(n-i-1, k-i-1, p) - B(n-i-1, k-i-2, p))$   
=  $p^{i}(1-p)B(n-i-1, k-i-1, p).$  (57)

In particular,

$$0 \leq v_{k-2} = p^{k-2}(1-p)B(n-k+1,1,p) = p^{k-2}(1-p)(1-(n-k+1)p),$$

which implies that the condition  $p \leq \frac{1}{n-k+1}$  is necessary. It follows from Proposition 3.2 that the solution in (10) is a feasible solution for  $p \leq \frac{1}{n-k+1}$ . Using Theorem 2.1.1, we conclude that the solution is optimal. Hence the condition  $p \leq \frac{1}{n-k+1}$  is sufficient.

#### 4.5Proof of Theorem 2.3

By the generalized inclusion-exclusion principle (see [1, Corollary 5.2]) and the fact that **P** is exchangeable,

$$\sum_{j=i}^{n} \binom{n-k}{j-i} v_j = p^i (1-p)^{k-i}, \qquad 0 \le i \le k, \qquad (58a)$$

$$v_i \ge 0, \qquad \qquad 0 \le i \le n. \tag{58b}$$

Since k is even, it follows from Theorem 2.1.2 that, for every feasible solution,

$$\mathbf{P}(X_1 = X_2 = \dots = X_n = 1) \leqslant B(n, k, 1 - p).$$
(59)

Now moving in the opposite direction, we have to find a feasible solution of the system of constraints (58a)-(58b) such that  $v_n = B(n, k, 1-p)$ . We choose

$$v_i = \begin{cases} 0, & 0 \le i \le n - k - 1, \\ (1 - p)^{n - i} \cdot B(i, i - (n - k), 1 - p), & n - k \le i \le n. \end{cases}$$
(60)

By Proposition 3.2, the  $v_i$ -s in (60) are non-negative for  $p \ge 1 - \frac{1}{n-k+1}$ . We rewrite (58a) in the form

$$\sum_{j=0}^{k} \binom{n-k}{n-k-i+j} v_{n-k+j} = p^{i}(1-p)^{k-i}, \qquad 0 \le i \le k$$

Equivalently,

$$\sum_{j=0}^{i} \binom{n-k}{i-j} v_{n-k+j} = p^{i} (1-p)^{k-i}, \qquad 0 \le i \le k.$$
(61)

By Lemma 3.1.3 and (60), we have that for  $0 \le i \le k$ 

$$\sum_{j=0}^{i} \binom{n-k}{i-j} v_{n-k+j} = \sum_{j=0}^{i} \binom{n-k}{i-j} (1-p)^{k-j} B(n-k+j,j,1-p)$$
$$= (1-p)^{k-i} \sum_{j=0}^{i} \binom{n-k}{i-j} (1-p)^{i-j} B(n-k+j,j,1-p)$$
$$= (1-p)^{k-i} p^{i}.$$

Hence, the  $v_i$ -s in (60) form the optimal distribution. This completes the proof.

# 4.6 Proof of Proposition A

Let  $I = \{j - 1, j, n\}$  for some  $1 \leq j \leq n - 1$ . The inverse  $B(I)^{-1} = [b_{s,t}]_{s \in I}^{t=0,1,2}$  is easily calculated as

$$B(I)^{-1} = \begin{bmatrix} \frac{nj}{n-j+1} & -\frac{n+j-1}{n-j+1} & \frac{2}{n-j+1} \\ -\frac{n(j-1)}{n-j} & \frac{n+j-2}{n-j} & -\frac{2}{n-j} \end{bmatrix}.$$
 (62)

$$\frac{j(j-1)}{(n-j+1)(n-j)} - \frac{2(j-1)}{(n-j+1)(n-j)} - \frac{2}{(n-j+1)(n-j)}$$

It follows from the discussion in Subsection 3.2 that

$$v_{i} = \frac{1}{\binom{n}{i}} \sum_{l=0}^{2} b_{i,l} \binom{n}{l} p^{l}, \qquad i = j - 1, j, n.$$
(63)

By (62) and (63),

$$v_{j-1} = \frac{1}{\binom{n}{j-1}} \cdot \frac{n}{n-j+1} \cdot (1-p) \cdot (j+p-np)$$

$$= \frac{1}{\binom{n-1}{j-1}} \cdot (1-p) \cdot (j+p-np),$$

$$v_{j} = \frac{1}{\binom{n}{j}} \cdot \frac{n}{n-j} \cdot (1-p) \cdot (np-p-j+1)$$

$$= \frac{1}{\binom{n-1}{j}} \cdot (1-p) \cdot (np-p-j+1),$$
(65)

and

$$v_n = \frac{1}{(n-j)(n-j+1)} \cdot (n(n-1)p^2 - 2n(j-1)p + j(j-1))$$
  
=  $\frac{1}{\binom{n-j+1}{2}} \left( \binom{n}{2}p^2 - n(j-1)p + \binom{j}{2} \right).$  (66)

By (64)-(66), considered as functions of p on the interval  $\left[\frac{j-1}{n-1}, \frac{j}{n-1}\right]$ :

- $v_{j-1}$  is decreasing and vanishes at the right endpoint;
- $v_j$  vanishes at the left endpoint and is increasing;
- $v_n$  attains its minimum, which is positive, at the left endpoint.

It then follows from Theorem 3.4 that

$$M(n,2,p) = \frac{1}{\binom{n-j+1}{2}} \left( \binom{n}{2} p^2 - n(j-1)p + \binom{j}{2} \right), \qquad p \in \left[ \frac{j-1}{n-1}, \frac{j}{n-1} \right],$$

and the distribution is given by (64)-(66).

# 4.7 Proof of Theorem 2.5

1. By Theorem 2.2, the optimal distribution for  $0 \leq p \leq p_1$  is given by

$$v_{i} = \begin{cases} (1-p) \cdot p^{i} \cdot B(n-i-1,k-i-1,p), & 0 \leq i \leq k-1, \\ 0, & k \leq i \leq n-1, \\ p^{k}, & i = n. \end{cases}$$
(67)

In particular, at the point  $p = p_1$ ,

$$v_{k-2} = (1-p_1)p_1^{k-2}B(n-k+1,1,p_1) = (1-p_1)p_1^{k-2}\left(1-(n-k+1)\cdot\frac{1}{n-k+1}\right) = 0,$$

while from Proposition 3.2 it follows that

$$v_i > 0, \qquad i \in \{0, 1, \dots, k - 3, k - 1, n\}.$$

The optimal distribution changes continuously with p, and therefore

$$v_i > 0, \qquad i \in \{0, 1, \dots, k - 3, k - 1, n\}, \ p \in [p_1, p_2],$$
(68)

for some  $p_2 > p_1$ . Let  $I = \{0, 1, 2, ..., k - 3, k - 1, k, n\}$ . It follows from the discussion in Subsection 3.2 and (33) that

$$v_i = \frac{1}{\binom{n}{i}} \cdot \sum_{l=0}^k b_{i,l} \binom{n}{l} p^l, \qquad i \in I,$$
(69)

where

$$b_{s,t} = \frac{(-1)^t \sum_{\alpha=0}^t (-1)^\alpha {t \choose \alpha} \prod_{u \in I - \{s\}} (u - \alpha)}{\prod_{u \in I - \{s\}} (u - s)}, \qquad s \in I, \ 0 \le t \le k.$$
(70)

By (70),

$$b_{0,j} = \begin{cases} (-1)^{j}, & 0 \leq j \leq k-3, \\ (-1)^{k-2} + (-1)^{k-1}C_{0}, & j = k-2, \\ (-1)^{k-1} - (-1)^{k-1} \binom{k-1}{k-2}C_{0}, & j = k-1, \\ (-1)^{k} + (-1)^{k-1} \binom{k}{k-2}C_{0}, & j = k, \end{cases}$$
(71)

where

$$C_0 = \frac{(k-2)! \cdot 2 \cdot (n-k+2)}{k! \cdot n} = \frac{\binom{n-1}{k}}{\binom{n-k+1}{2} \cdot \binom{n}{k-2}}.$$
(72)

In view of (69), (71), and (72), we obtain

$$v_{0} = \sum_{j=0}^{k} b_{0,j} {\binom{n}{j}} p^{j}$$

$$= \sum_{j=0}^{k} (-1)^{j} {\binom{n}{j}} p^{j} + (-1)^{k-1} \left( {\binom{n}{k-2}} p^{k-2} - {\binom{k-1}{k-2}} {\binom{n}{k-1}} p^{k-1} + {\binom{k}{k-2}} {\binom{n}{k}} p^{k-2} \right) C_{0}$$

$$= B(n,k,p) + (-1)^{k-1} {\binom{n}{k-2}} \cdot p^{k-2} B(n-k+2,2,p) C_{0}$$

$$= B(n,k,p) + (-1)^{k-1} \frac{{\binom{n-1}{k}}}{{\binom{n-k+1}{2}}} \cdot p^{k-2} B(n-k+2,2,p).$$
(73)

By (70), for  $1 \leq i \leq k - 3$ ,

$$b_{i,j} = \begin{cases} (-1)^{j} (-1)^{i} {j \choose i}, & i \leq j \leq k-3, \\ (-1)^{k-2} (-1)^{i} {k-2 \choose i} + (-1)^{k-i-1} C_{i}, & j = k-2, \\ (-1)^{k-1} (-1)^{i} {k-1 \choose i} - (-1)^{k-i-1} {k-1 \choose k-2} C_{i}, & j = k-1, \\ (-1)^{k} (-1)^{i} {k \choose i} + (-1)^{k-i-1} {k \choose k-2} C_{i}, & j = k, \end{cases}$$

$$(74)$$

where

$$C_{i} = \frac{(k-2)! \cdot 2 \cdot (n-k+2)}{i! \cdot (k-i)! \cdot (n-i)} = \frac{\binom{n}{i} \cdot \binom{n-i-1}{k-i}}{\binom{n-k+1}{2} \cdot \binom{n}{k-2}}. \qquad 1 \le i \le k-3.$$
(75)

(Here and below, we hint implicitly that all unspecified coefficients vanish. For example, in (74) we have  $b_{i,j} = 0$  for j < i.) By (69), (74), and (75), for  $1 \le i \le k - 3$ ,

$$\binom{n}{i} v_{i} = \sum_{j=0}^{k} b_{i,j} \binom{n}{j} p^{j}$$

$$= \sum_{j=i}^{k} (-1)^{j} (-1)^{i} \binom{j}{i} \binom{n}{j} p^{j}$$

$$+ (-1)^{k-i-1} p^{k-2} \left( \binom{n}{k-2} - \binom{k-1}{k-2} \binom{n}{k-1} p + \binom{k}{k-2} \binom{n}{k} p^{2} \right) C_{i}$$

$$= \binom{n}{i} p^{i} B(n-i,k-i,p) + (-1)^{k-i-1} p^{k-2} \binom{n}{k-2} B(n-k+2,2,p) C_{i},$$

and therefore

$$v_i = p^i B(n-i, k-i, p) + (-1)^{k-i-1} p^{k-2} \frac{\binom{n-i-1}{k-i}}{\binom{n-k+1}{2}} B(n-k+2, 2, p).$$
(76)

By (70),

$$b_{k-1,j} = \begin{cases} C_{k-1}, & j = k-2, \\ -\binom{k-1}{k-2}C_{k-1} + 1, & j = k-1, \\ \binom{k}{k-2}C_{k-1} - \binom{k}{k-1}, & j = k, \end{cases}$$
(77)

where

$$C_{k-1} = \frac{(k-2)! \cdot 2 \cdot (n-k+2)}{(k-1)! \cdot (n-k+1)} = \frac{\binom{n}{k-1}}{\binom{n}{k-2}} \cdot \frac{\binom{n-k}{1}}{\binom{n-k+1}{2}}.$$
(78)

By (69), (77), and (78), we have

$$\binom{n}{k-1} v_{k-1} = b_{k-1,k-2} \binom{n}{k-2} p^{k-2} + b_{k-1,k-1} \binom{n}{k-1} p^{k-1} + b_{k-1,k} \binom{n}{k} p^k$$

$$= C_{k-1} \cdot p^{k-2} \binom{n}{k-2} - \binom{k-1}{k-2} \binom{n}{k-1} C_{k-1} \cdot p^{k-1} + \binom{n}{k-1} p^{k-1}$$

$$+ \binom{k}{k-2} \binom{n}{k} C_{k-1} \cdot p^{k-1} - \binom{k}{k-1} \binom{n}{k} p^k$$

$$= \binom{n}{k-1} p^{k-1} B(n-k+1,1,p) + \binom{n}{k-2} p^{k-2} B(n-k+2,2,p) C_{k-1}$$

$$= \binom{n}{k-1} p^{k-1} B(n-k+1,1,p) + \frac{\binom{n}{k-1} \cdot \binom{n-k}{1}}{\binom{n-k+1}{2}} p^{k-2} B(n-k+2,2,p),$$

and therefore

$$v_{k-1} = p^{k-1}B(n-k+1,1,p) + \frac{\binom{n-k}{1}}{\binom{n-k+1}{2}}p^{k-2}B(n-k+2,2,p).$$
(79)

By (70),

$$b_{k,j} = \begin{cases} -C_k, & j = k - 2, \\ \binom{k-1}{k-2}C_k, & j = k - 1, \\ -\binom{k}{k-2}C_k + 1, & j = k, \end{cases}$$
(80)

where

$$C_k = \frac{(k-2)! \cdot 2 \cdot (n-k+2)}{k! \cdot (n-k)} = \frac{\binom{n}{k}}{\binom{n}{k-2}} \cdot \frac{1}{\binom{n-k+1}{2}}.$$
(81)

In view of (69), (80), and (81), we obtain

$$\binom{n}{k} v_{k} = b_{k,k-2} p^{k-2} \binom{n}{k-2} + b_{k,k-1} p^{k-1} \binom{n}{k-1} + b_{k,k} p^{k} \binom{n}{k}$$

$$= -C_{k} p^{k-2} \binom{n}{k-2} + \binom{k-1}{k-2} C_{k} p^{k-1} \binom{n}{k-1} - \binom{k}{k-2} C_{k} p^{k} \binom{n}{k} + p^{k} \binom{n}{k}$$

$$= \binom{n}{k} p^{k} - p^{k-2} \binom{n}{k-2} B(n-k+2,2,p) C_{k}$$

$$= \binom{n}{k} p^{k} - p^{k-2} \cdot \frac{\binom{n}{k}}{\binom{n-k+1}{2}} B(n-k+2,2,p),$$

and therefore

$$v_k = p^k - p^{k-2} \cdot \frac{1}{\binom{n-k+1}{2}} B(n-k+2,2,p).$$
(82)

Using Lemma 3.1.1, we further simplify (82) as

$$v_{k} = p^{k} - p^{k-2} \frac{1-p}{\binom{n-k+1}{2}} B(n-k+1,1.p) - p^{k}$$
  
=  $-p^{k-2} \frac{1-p}{\binom{n-k+1}{2}} B(n-k+1,1,p)$   
=  $\frac{1-p}{\binom{n-k+1}{2}} \cdot (-1 + (n-k+1)p).$  (83)

By (83), we observe that  $v_k = 0$  at the point  $p_1$  while  $v_k > 0$  for  $p > p_1$ . Using (69), we obtain

$$v_n = \sum_{l=0}^k b_{n,l} \binom{n}{l} p^l.$$
(84)

By (70),

$$b_{n,j} = \begin{cases} C_n, & j = k - 2, \\ -\binom{k-1}{k-2}C_n, & j = k - 1, \\ \binom{k}{k-2}C_n, & j = k, \end{cases}$$
(85)

where

$$C_n = \frac{1}{\binom{n}{k-2}} \cdot \frac{1}{\binom{n-k+1}{2}}.$$
(86)

In view of (84), (85), (86), we conclude that

$$v_{n} = b_{n,k-2} \binom{n}{k-2} p^{k-2} + b_{n,k-1} \binom{n}{k-1} p^{k-1} + b_{n,k} \binom{n}{k} p^{k}$$

$$= \binom{n}{k-2} - \binom{k-1}{k-2} \binom{n}{k-2} + \binom{k}{k-2} \binom{n}{k} C_{n}$$

$$= p^{k-2} \binom{n}{k-2} B(n-k+2,2,p) C_{n}$$

$$= \frac{1}{\binom{n-k+1}{2}} \cdot p^{k-2} \cdot B(n-k+2,2,p).$$
(87)

Using Theorem 2.1.1, (68), (83), and (87), we finally arrive at

$$M(n,k,p) = \frac{1}{\binom{n-k+1}{2}} \cdot p^{k-2} \cdot B(n-k+2,2,p), \qquad p \in [p_1,p_2].$$

2. For k = 2, 3, the claim follows readily from Propositions A-B. Thus, let  $k \ge 4$ . Using Lemma 3.1.1 and (79), we obtain

$$v_{k-1} = p^{k-1}B(n-k+1,1,p) + \frac{\binom{n-k}{1}}{\binom{n-k+1}{2}} \cdot p^{k-2}B(n-k+2,2,p)$$

$$= p^{k-1}\left((1-p)B(n-k,0,p) - \binom{n-k}{1}p\right)$$

$$+ \frac{2}{n-k+1} \cdot p^{k-2}\left((1-p)B(n-k+1,1,p) + \binom{n-k+1}{2}p^2\right) \qquad (88)$$

$$= p^{k-1}(1-p) + \frac{2}{n-k+1} \cdot p^{k-2}(1-p)B(n-k+1,1,p)$$

$$= \frac{1-p}{n-k+1} \cdot p^{k-2} \cdot (2-p(n-k+1)) = \frac{1-p}{n-k+1} \cdot p^{k-2} \cdot f_{k-1}(p),$$

where  $f_{k-1}(p) = 2 - (n - k + 1)p$ . Using Lemma 3.1.1 and (76), we obtain

$$v_{k-3} = p^{k-3}B(n-k+3,3,p) + \frac{\binom{n-k+2}{3}}{\binom{n-k+1}{2}} \cdot p^{k-2}B(n-k+2,2,p)$$

$$= p^{k-3}\left((1-p)B(n-k+2,2,p) - \binom{n-k+2}{3}p^3\right)$$

$$+ \frac{\binom{n-k+2}{3}}{\binom{n-k+1}{2}} \cdot p^{k-2}\left((1-p)B(n-k+1,1,p) + \binom{n-k+1}{2}p^2\right)$$

$$= p^{k-3}(1-p)\left(B(n-k+2,2,p) + \frac{n-k+2}{3} \cdot p \cdot B(n-k+1,1,p)\right)$$

$$= (1-p) \cdot p^{k-3} \cdot \left(1 - \frac{2}{3}(n-k+2)p + \frac{1}{6}(n-k+1)(n-k+2)p^2\right),$$

$$= (1-p) \cdot p^{k-3} \cdot f_{k-3}(p),$$
(89)

where

$$f_{k-3}(p) = 1 - \frac{2}{3} \binom{n-k+2}{1} p + \frac{1}{3} \binom{n-k+2}{2} p^2.$$
(90)

Using Lemma 3.1.1 and (76), we obtain

$$v_{k-4} = p^{k-4}B(n-k+4,4,p) - \frac{\binom{n-k+3}{4}}{\binom{n-k+1}{2}} \cdot p^{k-2}B(n-k+2,2,p)$$

$$= p^{k-4}\left((1-p)B(n-k+3,3,p) + \binom{n-k+3}{4}p^4\right)$$

$$-\frac{\binom{n-k+3}{4}}{\binom{n-k+1}{2}} \cdot p^{k-2}\left((1-p)B(n-k+1,1,p) + \binom{n-k+1}{2}p^2\right)$$

$$= p^{k-4}(1-p)\left(B(n-k+3,3,p) - \frac{\binom{n-k+3}{4}}{\binom{n-k+1}{2}} \cdot p^2 \cdot B(n-k+1,1,p)\right)$$

$$= (1-p) \cdot p^{k-4} \cdot f_{k-4}(p),$$
(91)

where

$$f_{k-4}(p) = 1 - \binom{n-k+3}{1}p + \frac{5}{6}\binom{n-k+3}{2}p^2 - \frac{1}{2}\binom{n-k+3}{3}p^3.$$
(92)

By (88)-(92), considered as functions of p on the interval  $\left[\frac{1}{n-k+1}, \frac{2}{n-k+1}\right]$ :

- $v_{k-1}$  is a decreasing function and vanishes at the right endpoint;
- $v_{k-3}$  attains its minimum, which is positive, at the right endpoint;
- $v_{k-4}$  is strictly positive at the left endpoint and strictly negative at the right endpoint. The derivative of  $f_{k-4}$  is strictly negative in the interval. Hence,  $v_{k-4}$  vanishes exactly once in the interval, at the point  $p_2$ ;
- $v_i$  is non-negative throughout the sub-interval  $[p_1, p_2]$  for i = 0, 1, 2, ..., k 5 by Theorem 3.5.

Using Theorem 3.4, we conclude the proof.

### 4.8 Proof of Theorem 2.7

By Theorem 2.3, the optimal distribution for  $[\bar{p}_1, 1]$  is given by

$$v_i = \begin{cases} 0, & 0 \le i \le n - k - 1, \\ (1 - p)^{n - i} \cdot B(i, i - (n - k), 1 - p), & n - k \le i \le n. \end{cases}$$
(93)

In particular, at the point  $p = \overline{p}_1$ 

$$v_{n-k+1} = (1 - \overline{p}_1)^{k-1} B(n-k+1, 1, 1 - \overline{p}_1) = (1 - \overline{p}_1)^{k-1} (1 - (n-k+1)(1 - \overline{p}_1)) = 0.$$

The optimal distribution changes continuously with p, and therefore,

 $v_i > 0,$   $i \in \{n - k, n - k + 2, \dots, n - 1, n\}, p \in [\overline{p}_2, \overline{p}_1],$ 

for some  $\overline{p}_2 < \overline{p}_1.$  Denote

$$\begin{aligned}
v'_{i} &= \mathbf{P}(X_{1} = X_{2} = \dots = X_{i} = 0, X_{i+1} = \dots = X_{n} = 1), & i = 0, 1, 2, \dots, n, \\
w'_{i} &= \binom{n}{i} v'_{i}, & i = 0, 1, 2, \dots, n, \\
S'_{0} &= 1, \\
S'_{i} &= \binom{n}{i} (1 - p)^{i}, & i = 1, 2, \dots, k, \\
I' &= \{0, 1, 2, \dots, k - 2, k, k + 1\}.
\end{aligned}$$
(94)

(Note that  $v'_i = v_{n-i}$ , we are using this notation to make the calculation simple and easily relate the calculation below to the proof of Theorem 2.5) It follows from the discussion in Subsection 3.2 and (33) that

$$v'_{i} = \frac{1}{\binom{n}{i}} \cdot \sum_{l=0}^{k} b_{i,l} \binom{n}{l} (1-p)^{l}, \qquad i \in I',$$
(95)

where

$$b_{s,t} = \frac{(-1)^t \sum_{\alpha=0}^t (-1)^\alpha {t \choose \alpha} \prod_{u \in I' - \{s\}} (u - \alpha)}{\prod_{u \in I' - \{s\}} (u - s)}, \qquad s \in I', \ 0 \leqslant t \leqslant k.$$
(96)

By (96),

$$b_{0,j} = \begin{cases} (-1)^j, & 0 \le j \le k-2, \\ (-1)^{k-1} - (-1)^{k-1} D_0, & j = k-1, \\ (-1)^k - (-1)^k \binom{k}{k-1} D_0, & j = k, \end{cases}$$
(97)

where

$$D_0 = \frac{1}{\binom{k+1}{2}}.$$
(98)

In view of (95), (97), and (98), we obtain

$$\begin{aligned} v_0' &= \sum_{j=0}^k b_{0,j} \binom{n}{j} (1-p)^j \\ &= \sum_{j=0}^k (-1)^j \binom{n}{j} (1-p)^j \\ &- (-1)^{k-1} \left( \binom{n}{k-1} (1-p)^{k-1} - \binom{k}{k-1} \binom{n}{k} (1-p)^k \right) D_0 \\ &= B(n,k,1-p) - (-1)^{k-1} \binom{n}{k-1} \cdot (1-p)^{k-1} B(n-k+1,1,1-p) D_0 \\ &= B(n,k,1-p) - (-1)^{k-1} \frac{\binom{n}{k-1}}{\binom{k+1}{2}} \cdot (1-p)^{k-1} B(n-k+1,1,1-p) \\ &= B(n,k,1-p) + \frac{\binom{n}{k+1}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1} B(n-k+1,1,1-p). \end{aligned}$$
(99)

By (99), and using the fact that  $v'_0 = v_n$ , we have

$$v_n = B(n,k,1-p) + \frac{\binom{n}{k+1}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1} B(n-k+1,1,1-p).$$
(100)

By (96), for  $1 \leq i \leq k-2$ ,

$$b_{i,j} = \begin{cases} (-1)^{j} (-1)^{i} {j \choose i}, & i \leq j \leq k-2, \\ (-1)^{k-1} (-1)^{i} {k-1 \choose i} - (-1)^{k-i-1} D_{i}, & j = k-1, \\ (-1)^{k} (-1)^{i} {k \choose i} + (-1)^{k-i-1} {k \choose k-1} D_{i}, & j = k, \end{cases}$$
(101)

where

$$D_{i} = \frac{\binom{n-i}{k+1-i} \cdot \binom{n}{i}}{\binom{n}{k-1} \cdot \binom{n-k+1}{2}}. \qquad 1 \le i \le k-2.$$
(102)

It follows from (95), (101), and (102) that, for  $1 \leq i \leq k-2$ ,

$$\binom{n}{i}v'_{i} = \sum_{j=0}^{k} b_{i,j}\binom{n}{j}(1-p)^{j}$$

$$= \sum_{j=i}^{k} (-1)^{j}(-1)^{i}\binom{j}{i}\binom{n}{j}(1-p)^{j}$$

$$- (-1)^{k-i-1}(1-p)^{k-1}\left(\binom{n}{k-1} - \binom{k}{k-1}\binom{n}{k}(1-p)\right)D_{i}$$

$$= \binom{n}{i}(1-p)^{i}B(n-i,k-i,1-p)$$

$$- (-1)^{k-i-1}(1-p)^{k-1}\binom{n}{k-1}B(n-k+1,1,1-p)D_{i},$$

and therefore

$$v_{n-i} = v'_i = (1-p)^i B(n-i, k-i, 1-p) - (-1)^{k-i-1} \frac{\binom{n-i}{k+1-i}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1} B(n-k+1, 1, 1-p).$$
(103)

By (96),

$$b_{k,j} = \begin{cases} D_k, & j = k - 1, \\ -\binom{k}{k - 1} D_k + 1, & j = k, \end{cases}$$
(104)

where

$$D_k = \frac{\binom{k-1}{1}}{\binom{k}{2}}.$$
 (105)

According to (95), (104), and (105),

$$\binom{n}{k}v'_{k} = \binom{n}{k}(1-p)^{k} + (1-p)^{k-1}\left(\binom{n}{k-1} - \binom{k}{k-1}\binom{n}{k}(1-p)\right)D_{k}$$

$$= \binom{n}{k}(1-p)^{k} + \binom{n}{k-1}(1-p)^{k-1}B(n-k+1,1,1-p)D_{k}$$

$$= \binom{n}{k}(1-p)^{k} + \frac{\binom{n}{k-1}\cdot\binom{k-1}{1}}{\binom{k}{2}}\cdot(1-p)^{k-1}B(n-k+1,1,1-p),$$

and therefore

$$v_{n-k} = v'_k = (1-p)^k + \frac{\binom{n-k}{1}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1} B(n-k+1,1,1-p).$$
(106)

By (96),

$$b_{k+1,j} = \begin{cases} -D_{k+1}, & j = k - 1, \\ \binom{k}{k-1} D_{k+1}, & j = k, \end{cases}$$
(107)

where

$$D_{k+1} = \frac{1}{\binom{k+1}{2}}.$$
(108)

Finally (95), (107), and (108) imply

$$\binom{n}{k+1}v'_{k+1} = -(1-p)^{k-1}\left(\binom{n}{k-1} - \binom{k}{k-1}\binom{n}{k}(1-p)\right)D_{k+1}$$
$$= -\binom{n}{k-1}(1-p)^k B(n-k+1,1,1-p)D_{k+1}$$
$$= -\frac{\binom{n}{k-1}}{\binom{k+1}{2}}(1-p)^k B(n-k+1,1,1-p),$$

and thus

$$v_{n-k-1} = v'_{k+1} = -\frac{1}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1} B(n-k+1,1,1-p).$$
(109)

By Proposition 3.2 and (109),

$$v_{n-k-1} \ge 0, \qquad p \le \overline{p}_1. \tag{110}$$

Using Theorem 3.4, (100), and (110), we arrive at the desired result

$$M(n,k.p) = B(n,k,1-p) + \frac{\binom{n}{k+1}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1}B(n-k+1,1,1-p), \qquad p \in [\overline{p}_2,\overline{p}_1].$$

2. For k = 2, the claim follows readily from Proposition A. Thus, let  $k \ge 4$ . Using (106), we have

$$v_{n-k} = (1-p)^k + \frac{\binom{n-k}{1}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1} B(n-k+1,1,1-p)$$
  
=  $(1-p)^{k-1} \left(\frac{2}{n-k+1} - (1-p)\right)$   
=  $(1-p)^{k-1} g_{n-k}(p),$  (111)

where

$$g_{n-k}(p) = \frac{2}{n-k+1} - (1-p).$$

By (103),

$$v_{n-k+2} = (1-p)^{k-2}B(n-k+2,2,1-p) + \frac{\binom{n-k+2}{3}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1}B(n-k+1,1,1-p)$$
  
=  $(1-p)^{k-2}\left(B(n-k+2,2,1-p) + \frac{n-k+2}{3} \cdot (1-p)B(n-k+1,1,1-p)\right)$   
=  $(1-p)^{k-2}\left(1 - \frac{2}{3}\binom{n-k+2}{1}(1-p) + \frac{1}{3}\binom{n-k+2}{2}(1-p)^2\right)$   
=  $(1-p)^{k-2}g_{n-k+2}(p),$  (112)

where

$$g_{n-k+2}(p) = 1 - \frac{2}{3} \binom{n-k+2}{1} (1-p) + \frac{1}{3} \binom{n-k+2}{2} (1-p)^2.$$
(113)

Employing (103), we obtain

$$v_{n-k+3} = (1-p)^{k-3}B(n-k+3,3,1-p) - \frac{\binom{n-k+3}{4}}{\binom{n-k+1}{2}} \cdot (1-p)^{k-1}B(n-k+1,1,1-p)$$
  
=  $(1-p)^{k-3}\left(B(n-k+3,3,1-p) - \frac{\binom{n-k+3}{4}}{\binom{n-k+1}{2}} \cdot (1-p)^2B(n-k+1,1,1-p)\right)$   
=  $(1-p)^{k-3} \cdot g_{n-k+3}(p),$  (114)

where

$$g_{n-k+3}(p) = 1 - \binom{n-k+3}{1}(1-p) + \frac{5}{6}\binom{n-k+3}{2}(1-p)^2 - \frac{1}{2}\binom{n-k+3}{3}(1-p)^3.$$
(115)

By (110)-(115), considered as function of p on the interval  $\left[1 - \frac{2}{n-k+1}, 1 - \frac{1}{n-k+1}\right]$ :

- $v_{n-k-1} \ge 0$  throughout the interval;
- $v_{n-k}$  vanishes at the left endpoint of the interval and is increasing throughout;
- $v_{n-k+2}$  attains it minimum, which is positive, at the left endpoint;
- $v_{n-k+3}$  is strictly positive at the right endpoint and strictly negative at the left endpoint. It vanishes only at the point  $\overline{p}_2$  of this interval;
- $v_{n-k+i}$  is non-negative throughout the sub-interval  $[\overline{p}_2, \overline{p}_1]$  for  $i = 4, 5, \ldots, k$  by Theorem 3.5.

Using Theorem 3.4, we conclude the proof.

# 5 Numerical Data

In this final section, we conduct numerical studies and present our key observations from these studies as eight open conjectures.

In the sequel, the data we will consider is for the following values of k and n:

- (1) k = 4 and  $6 \le n \le 500$ .
- (2) k = 6 and  $8 \le n \le 200$ .
- (3) k = 8 and  $10 \le n \le 100$ .
- (4) k = 10 and  $12 \le n \le 50$ .

All eight conjectures hold for this observed data.

Let  $2 \le k \le n-1$  with even k. Recall that a basis B is dual feasible if it consists of pairs of consecutive columns of A, say  $a_1, a_1 + 1, \ldots, a_{k/2}, a_{k/2} + 1$ , and in addition column n. By abuse of language, we will refer to the set of column numbers

$$I = \{a_1, a_1 + 1, a_2, a_2 + 1, \dots, a_{k/2}, a_{k/2} + 1, n\} \subset \{0, 1, 2, \dots, n\},\$$

as the basis. In all collected data, for every dual feasible basis I, the set

$$L(I) = \{ p \in [0, 1] : v_i \ge 0, i \in I \}$$

is a closed interval. We will thus refer to L(I) as the *interval corresponding to I*. For the vast majority of bases I, the interval L(I) is empty or degenerate (see Proposition 5.8 below). We will be interested in *realizable bases* – those for which L(I) is of positive length.

### 5.1 Ordering the Realizable Bases

Recall that the lexicographic order  $\leq$  on  $\mathbf{Z}^m$  is defined as follows: for any two points  $\mathbf{x} = (x_1, \ldots, x_m)$  and  $\mathbf{y} = (y_1, \ldots, y_m)$  in  $\mathbf{Z}^m$ , we have  $\mathbf{x} \leq \mathbf{y}$  if either

(1) for some  $1 \leq i \leq n$  we have  $x_l = y_l$  for all  $1 \leq l \leq i - 1$  and  $x_i < y_i$ , or

(2) 
$$\mathbf{x} = \mathbf{y}$$
.

Denote by  $\mathcal{R}$  the set of all realizable bases. The lexicographic order on  $\mathbb{Z}^m$  induces an order relation on the set  $\mathcal{R}$ , also denoted by  $\leq$ . We simply order the elements of each basis in increasing order and view the basis as a tuple in  $\mathbb{Z}^{k+1}$ .

Let  $\mathcal{L}$  be the set of all closed sub-intervals of positive length of [0,1]. We define an additional order relation, denoted by  $\leq$ , on  $\mathcal{L}$ , as follows: given two intervals  $J_1 = [a_1, b_1], J_2 = [a_2, b_2] \in \mathcal{L}$ , we have  $J_1 \ll J_2$  if  $J_1$  is to the left of  $J_2$ , namely  $b_1 \leqslant a_2$ , or  $J_1 = J_2$ . Note that, unlike  $\leq$ , the order  $\leq$  is not total.

We now fix n and k and write  $\mathcal{R} = \{I_1, I_2, \dots, I_N\}$ , ordered lexicographically:

$$I_1 \prec I_2 \prec \cdots \prec I_N.$$

Example 5.1.

(1) For k = 2 and n = 10, we have

$$\mathcal{R} = \{\{0, 1, 10\}, \{1, 2, 10\}, \{2, 3, 10\}, \{3, 4, 10\}, \{4, 5, 10\}, \{5, 6, 10\}, \{6, 7, 10\}, \{7, 8, 10\}, \{8, 9, 10\}\}.$$

(2) For k = 4 and n = 6, we have

$$\mathcal{R} = \{\{0, 1, 2, 3, 6\}, \{0, 1, 3, 4, 6\}, \{1, 2, 3, 4, 6\}, \{1, 2, 4, 5, 6\}, \{2, 3, 4, 5, 6\}\}.$$

**Conjecture 5.2.** The mapping  $L : \mathcal{R} \to \mathcal{L}$ , taking each realizable basis to the corresponding interval, is order-preserving, namely

$$I, I' \in \mathcal{R}, I \prec I' \implies L(I) \preccurlyeq L(I').$$

In other words, writing  $I_i = [c_i, d_i]$  for  $1 \le i \le N$ , we have

$$0 = c_0 < d_0 = c_1 < d_1 = c_2 < \dots = c_N < d_N = 1.$$

**Example 5.3.** For k = 4 and n = 10, there are 28 dual feasible bases, 12 of which are realizable. Table 1 lists all  $I_i$ -s, as well as the intervals  $L(I_i)$  and their lengths  $|L(I_i)|$ . (Here, all numerical values are rounded to two or three significant digits in their decimal expansion.) The graph of the function M(10, 4, p) is plotted in Figure 1. The intervals  $L(I_i)$  are depicted alternately in red and cyan to emphasize the piecewise polynomiality of the function.

i	$I_i$	$L(I_i)$	$ L(I_i) $
1	$\{0,1,2,3,10\}$	[0,1/7]	1/7
2	$\{0,1,3,4,10\}$	[1/7, 0.23]	0.090
3	$\{1,2,3,4,10\}$	[0.23, 0.26]	0.022
4	$\{1,2,4,5,10\}$	[0.26, 0.37]	0.117
5	$\{2,3,4,5,10\}$	$\left[0, 37, 0.38\right]$	0.004
6	$\{2,3,5,6,10\}$	[0.38, 1/2]	0.123
7	$\{3,4,6,7,10\}$	$\left[1/2, 0.62\right]$	0.123
8	$\{4,5,6,7,10\}$	[0.62, 0.63]	0.004
9	$\{4,5,7,8,10\}$	[0.63, 0.74]	0.117
10	$\{5,6,7,8,10\}$	[0.74, 0.77]	0.022
11	$\{5,6,8,9,10\}$	$\left[0.77,6/7\right]$	0.090
12	$\{6,7,8,9,10\}$	[6/7,1]	1/7

Table 1: Realizable bases, corresponding intervals, and their lengths for k = 4 and n = 10.




Figure 1: A graphical depiction of M(10, 4, p).

# 5.2 Realizable Bases

Let  $I_i$  be an arbitrary fixed realizable basis. As p varies from the left endpoint of  $L(I_i) = [c_i, d_i]$  to its right endpoint, the functions  $v_j = v_j(p), j \in I_i$  are non-negative. As we move

beyond the right endpoint  $d_i$ , one of these functions becomes negative – the solution of equation (27) involving only  $v_j$ -s with  $j \in I_i$  stops being feasible, and the control passes to the next basis  $I_{i+1}$ , with corresponding interval  $L(I_{i+1}) = [c_{i+1}, d_{i+1}]$ , where  $c_{i+1} = d_i$ .

**Conjecture 5.4.** For at most  $d_i$  breakpoints, the transition from  $I_i$  to  $I_{i+1}$  involves a minimal change  $-I_{i+1}$  is obtained from  $I_i$  by disposing of some  $a_j \in I_i$  and adjoining  $a_j + 2$  in its stead. Namely,  $I_i$  and  $I_{i+1}$  are of the form

$$I_{i} = \{a_{1}, a_{1} + 1, \dots, a_{j-1}, a_{j-1} + 1, \underline{a_{j}, a_{j} + 1}, \dots, n\},$$

$$I_{i+1} = \{a_{1}, a_{1} + 1, \dots, a_{j-1}, a_{j-1} + 1, \overline{a_{j} + 1}, a_{j} + 2, \dots, n\},$$
(116)

for some  $1 \leq j \leq k/2$ .

We note that a minimal change may look somewhat different from (116). For example,  $I_i$  could have contained some block of the form l, l + 1, l + 2, l + 3 (in addition to other elements), and l would be replaced by l + 4. However, this is never the case in the collected data.

Sometimes, though, the basis undergoes more than one change at a point; several elements  $i_1, i_2, \ldots, i_r$  leave the basis simultaneously, and r other elements join the basis for some  $r \ge 2$ . Points  $d_i$ , where such a transition occurs, are *exceptional* (see Section 5.3).

**Conjecture 5.5.** In the transition from  $I_i$  to  $I_{i+1}$ , if several elements  $i_1, i_2, \ldots, i_r$  leave the basis simultaneously, they are replaced by  $i_1 + 2, i_2 + 2, \ldots, i_r + 2$ .

### Example 5.6.

- (1) For k = 4 and n = 10, the bases  $\{2, 3, 5, 6, 10\}$  and  $\{3, 4, 6, 7, 10\}$  are consecutive realizable bases. We have two simultaneous changes: 2, 5 leave, and 4, 7 join at the exceptional point 1/2.
- (2) For k = 6 and n = 23, the bases  $\{6, 7, 10, 11, 14, 15, 23\}$  and  $\{7, 8, 11, 12, 15, 16, 23\}$  are consecutive realizable bases. We have three simultaneous changes: 6, 10, 14 leave, and 8, 12, 16 join at the exceptional point 1/2.

**Remark 5.7.** In view of Conjecture 5.5, at exceptional points the basis undergoes several simultaneous changes as in (116). Conjecture 5.2 does not exclude the possibility that a basis such as  $\{10, 11, 20, 21, n\}$ , for example, changes to  $\{11, 12, 19, 20, n\}$  or to  $\{11, 12, 22, 23\}$ . However, we did not encounter any such transitions.

If both Conjectures 5.4 and 5.5 hold, then Proposition 5.8 below holds.

**Proposition 5.8.** Assume that both Conjectures 5.4 and 5.5 hold. Then the following two statements hold:

(1) For every n and even  $k \leq n-2$ ,

$$|\mathcal{R}| \leqslant \frac{k(n-k)}{2} + 1. \tag{117}$$

(2) If the transitions from  $I_i$  to  $I_{i+1}$ , for all  $1 \le i \le N-1$ , involve minimal changes (in other words, there are no exceptional points), then

$$|\mathcal{R}| = \frac{k(n-k)}{2} + 1.$$

Proof. Denote

$$S(I) = \sum_{j=1}^{k/2} a_j, \qquad I = \{a_1, a_1 + 1, \dots, a_{k/2}, a_{k/2} + 1, n\}.$$

From Conjectures 5.4 and 5.5, it follows that for every two consecutive realizable bases  $I_i, I_{i+1}$ , the difference  $S(I_{i+1}) - S(I_i)$  is twice the number of elements of  $I_i$  replaced by others in  $I_{i+1}$ . In particular,

$$S(I_{i+1}) - S(I_i) \ge 2, \qquad 1 \le i \le N - 1.$$

Hence

$$S(I_N) - S(I_1) = \sum_{i=1}^{N-1} \left( S(I_{i+1}) - S(I_i) \right) \ge 2(N-1).$$
(118)

By Thereoms 2.2 and 2.3,

$$I_1 = \{0, 1, \dots, k-1, n\}, \qquad I_N = \{n-k, n-k+1, \dots, n-1, n\}.$$

We then have

$$S(I_1) = 0 + 2 + \dots + (k - 2) = \frac{k}{2}(k - 2),$$
(119)

and

$$S(I_N) = (n-k) + (n-k+2) + \dots + (n-2)$$
  
=  $nk - (2+4+\dots+k) = nk - k\left(\frac{k}{2}+1\right).$  (120)

By (118), (119), and (120), we obtain

$$N \leq 1 + \frac{1}{2}(S(I_N) - S(I_1)) = 1 + \frac{k(n-k)}{2},$$

which is equivalent to (117). Now, in the case when there is no exceptional point, we have  $S(I_{i+1}) - S(I_i) = 2$  for all  $1 \le i \le N - 1$ . We thus have equality in (118), and the second claimed statement follows.

## 5.3 Exceptional Points

A pair (n, k) is an *exceptional pair* if there exists at least one exceptional point for (n, k). Numerical data suggest that, for a fixed k, exceptional pairs are quite rare (see Table 2).

k	Range of $n$	Number of exceptional pairs $(n, k)$	
	[6,100]	19	
	[101,200]	15	
4	[201, 300]	13	
	[301, 400]	13	
	[401, 500]	11	
6	[8,100]	6	
0	[101,200]	3	
8	[10,100]	0	
10	[12,50]	0	

Table 2: Statistics for exceptional pairs.

Conjecture 5.9. All the exceptional points are rational.

Table 3 lists all exceptional points for n in the range [7, 100] and k = 4. For k = 6 in the range [8, 200]

23, 28, 35, 67, 76, 87, 135, 148, 163,

are the only *n*-s for which (n, 6) is exceptional. For each of these, p = 1/2 is the only exceptional point.

### 5.4 Symmetry of Bases and of Intervals

Theorems 2.2 and 2.3 show that  $L(I_1)$  and  $L(I_N)$  are mirror images of each other with respect to the point p = 1/2. The same holds by Theorems 2.5 and 2.7 for  $L(I_2)$  and  $L(I_{N-1})$ . Moreover, the bases themselves enjoy a similar symmetry property. Namely, ignoring the number n in the bases (as it belongs to all of them),  $I_1$  and  $I_N$  are mirror images with respect to the number  $\frac{n-1}{2}$ , and so are  $I_2$  and  $I_{N-1}$ . These symmetries hold in general for the collected data.

### Conjecture 5.10.

(1) If

$$I_i = \{a_1, a_1 + 1, a_2, a_2 + 1, \dots, a_{k/2}, a_{k/2} + 1, n\},\$$

where  $1 \leq i \leq N$ , then

$$I_{N+1-i} = \{n - a_{k/2} - 2, n - a_{k/2} - 1, \dots, n - a_1 - 1, n - a_1, n\}.$$

n	Exceptional points	n	Exceptional point
10	1/2	52	2/5, 3/5
11	1/3, 2/3	56	1/3, 2/3
17	1/2	65	1/6, 1/2, 5/6
27	2/5, 3/5	66	1/4, 3/4
29	1/6, 1/3, 2/3, 5/6	77	1/5, 3/10, 7/10, 4/5
34	1/4, 3/4	82	1/2
37	1/2	83	4/9, 5/9
50	1/2	92	1/3, 2/3
51	1/7, 3/7, 4/7, 6/7	100	1/7, 2/7, 5/7, 6/7

Table 3: Exceptional points for  $6 \le n \le 100$  for k = 4.

(2) Denoting 
$$L(I_i) = [c_i, d_i]$$
 for each *i*, we have  $c_{N+1-i} = 1 - d_i$  and  $d_{N+1-i} = 1 - c_i$ .

This implies

**Conjecture 5.11.** If d is an exceptional point for some (n, k), then so is 1 - d.

### 5.5 The Spread of the Bases

Let  $I = \{a_1, a_1 + 1, a_2, a_2 + 1, \dots, a_{k/2}, a_{k/2} + 1, n\}$  be a dual feasible basis. Since n belongs to all such bases, we ignore it and define the *spread* of I by

$$Sp(I) = |a_{k/2} + 1 - a_1|.$$

By Theorems 2.2 and 2.3,  $I_1$  and  $I_N$  are the densest realizable bases, i.e.,

$$Sp(I_1) = Sp(I_N) = k - 1 = \min\{Sp(I_i), : i = 1, 2, \dots, N\}.$$
(121)

There may be several realizable bases other than  $I_1$  and  $I_N$  with spread k-1. For example, for n = 10 and k = 4, we have  $I_3 = \{1, 2, 3, 4, 10\}$ ,  $I_5 = \{2, 3, 4, 5, 10\}$ ,  $I_8 = \{4, 5, 6, 7, 10\}$ , and  $I_{10} = \{5, 6, 7, 8, 10\}$ , all of which are of minimal spread (see Table 1). It follows from Conjecture 5.10 that

$$Sp(I_i) = Sp(I_{N+1-i}), \quad i = 1, 2, \dots, N.$$

The spread of a random dual feasible basis is usually not much less than n. However, our numerical studies suggest that all realizable bases are relatively quite concentrated; namely, the spread is much smaller than n. In fact, for fixed n and k, the maximum spread Msp(n, k) is defined by

$$Msp(n,k) = max{Sp(I_i) : i = 1, 2, ..., N}.$$

**Conjecture 5.12.** For each fixed k, Msp(n, k) is a monotonic non-decreasing function of n.

Tables 4.a, 4.b, 4.c, 4.d list Msp(n, 4), Msp(n, 6), Msp(n, 8), and Msp(n, 10), respectively, for the ranges of n in our collected data.

### Conjecture 5.13.

$$Msp(n,4) = \left\lfloor \sqrt{n-2} \right\rfloor + 2, \qquad n \ge 6.$$
(122)

Naturally, as k grows, the function Msp(n, k), considered as a function of n, grows faster. However, the collected data does not give such a simple formula as in (122). Nor is it enough to guess whether, for a fixed  $k \ge 6$ , the function grows as  $\Theta(\sqrt{n})$  or faster.

Msp(n,4)	Range of $n$	Msp(n,4)	Range of $n$	Msp(n,4)	Range of $n$
4	[6, 10]	11	[83,101]	18	[258, 290]
5	[11, 17]	12	[102,122]	19	[291,325]
6	[18,26]	13	[123,145]	20	[326,362]
7	[27,37]	14	[146,170]	21	[363,401]
8	[38, 50]	15	[171, 197]	22	[402, 442]
9	[51, 65]	16	[198,226]	23	[443,485]
10	[66,82]	17	[227,257]	24	[486,500]

Table 4.a: The spread for  $6 \le n \le 500$  and k = 4.

Msp(n,6)	Range of $n$	Msp(n, 6)	Range of $n$	Msp(n, 6)	Range of $n$
6	[8,9]	13	[42,49]	20	[110,121]
7	[10,13]	14	[50, 57]	21	[122,135]
8	[14,17]	15	[58,67]	22	[136,148]
9	[18,23]	16	[68,76]	23	[149,163]
10	[24,28]	17	[77,87]	24	[164,177]
11	[29, 35]	18	[88,97]	25	[178,193]
12	[36,41]	19	[98,109]	26	[194,200]

Table 4.b: The spread for  $8 \le n \le 200$  and k = 6.

Msp(n,8)	Range of $n$	Msp(n, 8)	Range of $n$	Msp(n, 8)	Range of $n$
8	[10,11]	14	[29,32]	20	[62,67]
9	[12,13]	15	[33,37]	21	[68,75]
10	[14,17]	16	[38,43]	22	[76,82]
11	[18,20]	17	[44,48]	23	[83,90]
12	[21,23]	18	[49,54]	24	[91,98]
13	[24,28]	19	[55,61]	25	[99,100]

Table 4.c: The spread for  $10 \le n \le 100$  and k = 8.

Msp(n, 10)	Range of $n$	Msp(n, 10)	Range of $n$	Msp(n, 10)	Range of $n$
10	[12, 12]	14	[21,23]	18	[35, 37]
11	[13, 14]	15	[24,26]	19	[38, 42]
12	[15,17]	16	[27,30]	20	[43,46]
13	[18,20]	17	[31,34]	21	[47,50]

Table 4.d: The spread for  $12 \le n \le 50$  and k = 10.

The numerical studies suggest that

$$|L(I_1)| = |L(I_N)| > \max\{|L(I_i)| : i = 2, 3, \dots, N-1\}.$$
(123)

The equality in (121) and the inequality in (123) raise the following question.

**Question 5.14.** For fixed n and k, are the sequences  $(\text{Sp}(I_i))_{i=1}^N$  and  $(|L(I_i)|)_{i=1}^N$  negatively correlated?

In Figure 2, we have plotted the correlation coefficients as a function of n for k = 4, 6, 8, 10. Note that the correlation coefficient is usually negative for k = 4 and is negative throughout for k = 6, 8, 10. However, it seems to converge to 0 as  $n \to \infty$ .




Figure 2: Correlation between the spread of realizable bases and the lengths of the corresponding intervals.

As we have previously noted, we hope that the above conjectures will help future researchers with the very difficult task of deriving further exact formulas for M(n, k, p) over the whole interval [0, 1].

Acknowledgements: We wish to thank J. Dudek, R. Pemantle, and A. Yadin for helpful discussions:

# References

- M. Aigner, A Course in Enumeration, Graduate Texts in Mathematics 238, Springer-Verlag Berlin, 2007.
- [2] N. Alon, A. Andoni, T. Kaufman, K. Matulef, R. Rubinfeld, and N. Xie, Testing kwise and almost k-wise independence, STOC-Proceedings of the 39th Annual ACM Symposium on Theory of Computing (2007), 496–505.
- [3] N. Alon, L. Babai, and A. Itai, A fast and simple randomized parallel algorithm for the maximal independent set problem, *J. Algorithms* **7(4)** (1986), 567–583.
- [4] I. Benjamini, O. Gurel-Gurevich, and R. Peled, On k-wise independent distributions and boolean functions, arXiv preprint arXiv:1201.3261, 2012.
- [5] A. Z. Broder, M. Charikar, and M. Mitzenmacher, A derandomization using min-wise independent permutations, *International Workshop on Randomization and Approxi*mation Techniques in Computer Science (1998), 15–24.
- [6] C. E. Bonferroni, Teoria statistica delle classi e calcolo delle probabilità, Pubblicazioni del R Istituto Superiore di Scienze Economiche e Commericiali di Firenze 8 (1936), 3–62.
- [7] E. Boros and A. Prékopa, Closed form two-sided bounds for probabilities that at least r and exactly r out of n events occur, *Math. Oper. Res.* **14(2)** (1989), 317–342.
- [8] K. Censor-Hillel, M. Parter, and G. Schwartzman, Derandomizing local distributed algorithms under bandwidth restrictions, *Distributed Computing* 33(3) (2020), 349– 366.
- [9] B. Chor, O. Goldreich, J. Hastad, J. Friedman, S. Rudich, and R. Smolensky, The bit extraction problem or t-resilient functions, Proc. 26th IEEE Symposium on Foundations of Computer Science (1985), 396–407.
- [10] J. Galambos, Methods for proving Bonferroni type inequalities, J. London Math. Soc. 2(4) (1975), 561–564.
- [11] J. Galambos, The Asymptotic Theory of Extreme Order Statistics, 2nd ed. Robert E. Krieger Publishing Co.Inc., Malabar, Florida, 1987.
- [12] J. Galambos and I. Simonelli, Bonferroni-type Inequalities with Applications, Springer, New York, 1996.
- [13] E. Kaplan, M. Naor, and O. Reingold, Derandomized constructions of k-wise (almost) independent permutations, Algorithmica 55(1) (2009), 113–133.
- [14] H. Karloff and Y. Mansour, On construction of k-wise independent random variables, Proceedings of the 26th Annual ACM Symposium on Theory of Computing (1994), 564–573.

- [15] R. Karp and A. Wigderson, A fast parallel algorithm for the maximum independent set problem, J. ACM 32(4) (1985), 762-773.
- [16] M.-Y. Lee, Bonferroni-type Inequalities, Ph.D. Thesis, Temple University, Philadelphia, PA, 1991.
- [17] M. Luby and A. Wigderson, Pairwise independence and derandomization, Found. Trends. Theor. Comput. Sci. 1(4) (2006), 237–301.
- [18] M. Luby, A simple parallel algorithm for the maximal independent set problem, *Proc* of the 17th Annual ACM Symposium on Theory of Computing (1985), 1-10.
- [19] R. Peled, A. Yadin, and A. Yehudayoff, The maximal probability that k-wise independent bits are all 1, Random Structures & Algorithms 38(4) (2011), 502–525.
- [20] A. Prékopa, Boole-Bonferroni inequalities and linear programming, Oper. Res. 36(1) (1988), 145–162.
- [21] A. Prékopa, Sharp bounds on probabilities using linear programming, Oper. Res. 38
   (2) (1990), 227–239.
- [22] A. K. Ramachandra and K. Natarajan, Tight probability bounds with pairwise independence, SIAM J. Discrete Math. 37(2) (2023), 516–555.
- [23] J. P. Schmidt, A. Siegel, and A. Srinivasan, Chernoff-Hoeffding bounds for applications with limited independence, SIAM J. Discrete Math. 8(2) (1995), 223–250.
- [24] T. H. Spencer, Generalized Bonferroni inequalities, J. Appl. Probab. 31(2) (1994), 409–417.
- [25] L. Takaćs, On the method of inclusion and exclusion, J. Amer. Statist. Assoc. 62(317) (1967), 102–113.