Richard G. Everitt^{1,2}

¹Department of Statistics, University of Warwick

²The Zeeman Institute for Systems Biology & Infectious Disease Epidemiology Research,

University of Warwick

¹Email: richard.everitt@warwick.ac.uk

29th July 2024

Abstract

Approximate Bayesian computation (ABC) is the most popular approach to inferring parameters in the case where the data model is specified in the form of a simulator. It is not possible to directly implement standard Monte Carlo methods for inference in such a model, due to the likelihood not being available to evaluate pointwise. The main idea of ABC is to perform inference on an alternative model with an approximate likelihood (the ABC) likelihood), estimated at each iteration from points simulated from the data model. The central challenge of ABC is then to trade-off bias (introduced by approximating the model) with the variance introduced by estimating the ABC likelihood. Stabilising the variance of the ABC likelihood requires a computational cost that is exponential in the dimension of the data, thus the most common approach to reducing variance is to perform inference conditional on summary statistics. In this paper we introduce a new approach to estimating the ABC likelihood: using iterative ensemble Kalman inversion (IEnKI) [\(Iglesias, 2016;](#page-18-0) [Iglesias](#page-18-1) et al., [2018\)](#page-18-1). We first introduce new estimators of the marginal likelihood in the case of a Gaussian data model using the IEnKI output, then show how this may be used in ABC. Performance is illustrated on the Lotka-Volterra model, where we observe substantial improvements over standard ABC and other commonly-used approaches.

1 Introduction

1.1 Intractable likelihoods and simulation-based inference

This paper considers the problem of Bayesian inference for parameters θ of a data generating process $P(\cdot | \theta)$ used as a model for data y_{obs} , using a prior p on θ . We use π to denote the posterior distribution of θ given y_{obs} , with $\tilde{\pi}$ denoting the unnormalised posterior given by $\tilde{\pi}(\theta | y_{obs}) = p(\theta)P(y_{obs} | \theta)$. We consider the case where the likelihood $P(y_{\text{obs}} | \theta)$ is intractable in that it cannot be evaluated pointwise at θ . There are several situations in which this can occur:

- 1. in the case of "tall" data [\(Bardenet](#page-18-2) *et al.*, [2017\)](#page-18-2), where the P consists of a product of a very large number of terms;
- 2. where π is "doubly intractable" [\(Murray](#page-19-0) *et al.*, [2006\)](#page-19-0), referring to P having an intractable normalising constant;
- 3. where latent variables x are involved in the specification of P as follows

$$
P(y | \theta) = \int_{x} g(y | x, \theta) f(x | \theta) dx;
$$
 (1)

4. where P is specified by a piece of software that can be simulated - it is possible to simulate $y \sim P(\cdot | \theta)$ - but it is impractical to write out P in mathematical form.

These cases overlap: for example usually in situation 4, it would in principle be possible to write the model as specified in equation [\(1\)](#page-0-0), were the details of the software implementation available. Inference in situation 4 has become known as *simulation-based inference* [\(Cranmer](#page-18-3) *et al.*, [2020\)](#page-18-3), a term given to collectively describe methods that approximate the posterior when P can be simulated, but not evaluated pointwise at θ .

The fundamental SBI technique is approximate Bayesian computation (ABC; [Tavaré](#page-19-1) et al. [\(1997\)](#page-19-1)). ABC approximates the likelihood using simulations from P , based on the simple principle that "good" parameters are more likely to generate simulations y that are close to the observed data y_{obs} . For each θ , M simulations $\{y^j\}_{j=1}^M$ are taken from $P(\cdot | \theta)$, summary statistics $s^j = S(y^j)$ of each y^j are calculated, then a kernel K_{ε} $(s_{\text{obs}} | s^j)$ (where $s_{\text{obs}} = S(y_{\text{obs}})$ is calculated for each s^j . The likelihood is then approximated using

$$
\frac{1}{M} \sum_{j=1}^{M} K_{\varepsilon} \left(s_{\text{obs}} \mid s^{j} \right), \tag{2}
$$

this being a Monte Carlo approximation of the ABC likelihood

$$
\int_{s} K_{\varepsilon} (s_{\text{obs}} \mid s) P_{s} (s \mid \theta) ds, \tag{3}
$$

where P_s is the distribution resulting from the transformation $s = S(y)$ of $y \sim P(\cdot | \theta)$. The kernel K_{ε} takes values in $[0,\infty)$ and is chosen to take higher values the closer s^j is to s_{obs} , with the "tolerance" $\varepsilon > 0$ controlling the "width" of the kernel. Popular choices for K_ε in include:

• Uniform

$$
K_{\varepsilon} (s_{\text{obs}} \mid s) = \begin{cases} 1 & \text{if } d(s_{\text{obs}}, s) < \varepsilon \\ 0 & \text{otherwise} \end{cases}
$$

where d is some distance metric. A common choice is the weighted Euclidean distance

$$
d(s_{\text{obs}}, s) = \left(\sum_{i=1}^{d_s} \left(\frac{s_{\text{obs},i} - s_i}{\sigma_i}\right)^2\right)^{1/2},
$$

where d_s is the dimension of s and each σ_i is chosen such that each of the terms in the sum have a similar scale, so that all summary statistics have a comparable influence.

• Gaussian

$$
K_{\varepsilon} (s_{\text{obs}} \mid s) = (2\pi\varepsilon)^{-d_y/2} \prod_{i=1}^{d_s} \frac{1}{\sigma_i^2} \exp\left(-\frac{1}{2\varepsilon^2} \sum_{i=1}^{d_s} \left(\frac{s_{\text{obs},i} - s_i}{\sigma_i}\right)^2\right),
$$

i.e. the multivariate Gaussian distribution with mean s and covariance $\varepsilon^2 \Sigma_s$ where Σ_s is diagonal with ith diagonal element σ_i^2 , similarly to scale the summary statistics.

The reduction of the data y to the summary s means that the ABC posterior (the posterior using the ABC likelihood) is a posterior conditional on s_{obs} rather than y_{obs} which represents a loss of informaton about θ unless s is sufficient. As $\varepsilon \to 0$, the ABC posterior converges to the true posterior conditional on s_{obs} . Using $\varepsilon > 0$, which is almost always required in practice, introduces a further approximation. Most other approaches to SBI also rely on approximations of some kind (see, e.g. [Cranmer](#page-18-3) et al. [\(2020\)](#page-18-3)).

The estimated ABC likelihood in equation [\(2\)](#page-1-0) is usually used within a Monte Carlo method that explores the space of θ, for example within a Markov chain Monte Carlo (MCMC) algorithm. The resultant algorithm [\(Marjoram](#page-19-2) *et al.*, [2003\)](#page-19-2) is known as ABC-MCMC, which can be seen to be an instance of a pseudo-marginal [\(Beaumont, 2003;](#page-18-4) [Andrieu and Roberts, 2009\)](#page-16-0) algorithm. It is well-established (e.g. [Andrieu and Vihola](#page-16-1) [\(2014\)](#page-16-1)) that for a pseudo-marginal algorithm to be efficienct, the estimated marginal likelihood needs to have a relatively small variance.

The past 25 years have seen the development of a range of alternative ABC and other SBI methods. A core challenge is that of performing SBI in "high" dimensions: most approaches rely on d_s being small. In particular, the variance of the likelihood estimator in equation [\(2\)](#page-1-0) requires M to be exponential in d_s to stablise its variance. This also limits the use of ABC to low-dimensional parameter spaces, since a dimension d_{θ} of θ larger than d_s results in identifiability problems in the posterior.

1.2 Alternatives and extensions to ABC

For the intractable likelihood in situation 3, SBI is often not the most effective choice of method. Whilst the integral P may be intractable, it is usually possible to evaluate the functions f and g in equation [\(1\)](#page-0-0), so we may use Monte Carlo approaches whose target is the joint posterior $\pi(\theta, x \mid y_{\text{obs}})$, then use the θ -points to approximate $\pi(\theta \mid y_{\text{obs}})$. One such approach is particle MCMC [\(Andrieu](#page-16-2) et al., [2010\)](#page-16-2), which uses sequential Monte Carlo (SMC) methods to explore x-space conditional on θ . In situation 3, such an approach might be seen to be favourable to SBI, since it has the does not suffer the approximations introduced in SBI. Indeed, some recent work [\(Prangle](#page-19-3) *et al.*, [2018;](#page-19-3) [Kerama](#page-18-5) [et al.](#page-18-5), [2022;](#page-18-5) [Zhang, 2022\)](#page-19-4) has focussed on using a reparameterisation to cast models that might usually be tackled using ABC in the form in equation [\(1\)](#page-0-0). Specifically, the idea is to rewrite the simulator model P as a deterministic transformation $G(\theta, u)$ of the parameter θ and random variables u whose prior $\phi(u | \theta)$ is a normalised tractable distribution. The ABC likelihood then changes from equation [\(3\)](#page-1-1) to

$$
\int_{u} K_{\varepsilon} \left(s_{\text{obs}} \mid S(G(\theta, u)) \right) \phi(u \mid \theta) du. \tag{4}
$$

We refer to this situation as *rare event* (RE) ABC after the interpretation of the estimation of equation [\(4\)](#page-2-0) as rare event estimation. Although there are situations where using such a reparameterisation may be difficult (for example when given a simulator in the form of legacy code), the significant advantage over standard ABC is that, now that the intractable P does not appear in the expression, we are free to use alternative Monte Carlo methods to sample from the target distribution. For example, particle MCMC is used in [Prangle](#page-19-3) et al. [\(2018\)](#page-19-3), referred to as RE-ABC-MCMC, using SMC to explore u-space conditional on θ . The SMC yields a lower variance estimator than equation [\(2\)](#page-1-0), hence usually yields a substantially more efficient MCMC chain compared to standard ABC-MCMC.

1.3 Proposed approach

Despite the advances described in the preceding section, there remain many situations where ABC might be a more efficient approach than, say, particle MCMC, when in situation 3. When the dimension d_x of x-space is high, or x-space has a complex structure, using Monte Carlo methods to explore this space may be inefficient: we may need a very large computational time to obtain low variance Monte Carlo estimates. Examples include: the situation for which ABC was originally designed in population genetics, where x is a tree describing the ancestry of a number of individuals [\(Tavaré](#page-19-1) et al., [1997\)](#page-19-1); or time-series data with a large number of noisy observations, where x is a latent time series (e.g. [Andrieu](#page-16-2) *et al.* [\(2010\)](#page-16-2)). The same issue is present in rare event ABC when d_u , the dimension of u, is high. ABC largely avoids this issue by using a likelihood that involves integration over space that is typically lower dimension: over dimension d_s (equation [\(3\)](#page-1-1)) compared to d_x (equation [\(1\)](#page-0-0)). Therefore, despite the bias present in an ABC posterior, the lower variance may result in smaller errors given a fixed computational effort.

In this paper we propose a new approach to estimating the ABC likelihood in equation [\(3\)](#page-1-1) using iterative ensemble Kalman inversion (IEnKI) [\(Iglesias, 2016;](#page-18-0) [Iglesias](#page-18-1) *et al.*, [2018\)](#page-18-1), and use this with an MCMC algorithm. IEnKI is a Monte Carlo approach that has some similarities with SMC, so in this sense our new approach has the flavour of RE-ABC-MCMC. However, it has the crucial difference that rather than using the form of the ABC likelihood in equation [\(4\)](#page-2-0) which integrates over a space of dimension d_u , it uses the original form in equation [\(3\)](#page-1-1) which integrates over a space of dimension d_s . We show empirically that this estimator can have substantially lower variance than the standard estimated ABC likelihood in equation [\(2\)](#page-1-0), and that the variance can be relatively low when d_s is much larger than is possible for standard ABC. We show the new approach compares favourably with a number of alternative approaches on the Lotka-Volterra model, including standard ABC, synthetic likelihood and particle MCMC.

The structure of the paper is as follows. In section [2](#page-2-1) we recap IEnKI and introduce new marginal likelihood estimators using the output of this algorithm. In section [3](#page-6-0) we show how to use the new marginal likelihood estimators to estimate the ABC likelihood, highlighting some properties of the new IEnKI that facilitate its use for ABC. We illustrate the performance of the new methods in section [4,](#page-9-0) before a discussion in section [5.](#page-13-0)

2 Iterative ensemble Kalman inversion

In this section we describe IEnKI, then outline how a marginal likelihood estimator may be devised from the output of this algorithm.

2.1 Algorithm

2.1.1 SMC samplers

The structure of IEnKI follows that of SMC samplers [\(Del Moral](#page-18-6) et al., [2006\)](#page-18-6): a method designed to produce a Monte Carlo approximation to a sequence of targets. We begin by introducing notation. Let π_t (for $t = 0 : T$) be a sequence of target distributions, with $\tilde{\pi}_t$ denoting an unnormalised form of each distribution and Z_t its normalising constant. SMC samplers are commonly used for approximating Bayesian posterior distributions, so a popular sequence of targets is

$$
\tilde{\pi}_t(x) = p(x)l^{\alpha_t}(x),\tag{5}
$$

where p is a prior distribution and l a likelihood, and $0 = \alpha_0 < ... < \alpha_T = 1$. This sequence has several appealing properties: the first distribution is equal to the prior and is usually easy to simulate; the final distribution is equal to the posterior; and the sequence of α_t can be chosen so as to create a path of distributions between the prior and posterior such that the approximation to the target π_{t-1} may be used (via the SMC algorithm) to constuct an approximation to π_t .

The output of an SMC sampler is a population of weighted particles $\{(x_t^j, w_t^j)\}_{j=1}^M$ at each iteration that is used to approximate π_t and estimate the normalising constant Z_t . Here we present the most commonly-used case, where at iteration t each particle is moved using K_t , an MCMC kernel with invariant distribution π_t At iteration t the unnormalised weight of each particle is calculated using $\tilde{w}_t^j = l^{\alpha_t - \alpha_{t-1}}(x_{t-1}^j)$, then the unnormalised weights $\{\tilde{w}_t^j\}_{j=1}^M$ are normalised to give $\{w_t^j\}_{j=1}^M$ and the new particle population is simulated from $\sum_{j=1}^M w_t^j K_t(\cdot \mid x_{t-1}^j)$. The performance of estimators from SMC samplers is critically dependent on choices made in the design of the algorithm, namely:

- 1. The choice of the the sequence (α_t) . If α_t is too far from α_{t-1} the algorithm will become *degenerate*, where one particle has most of the weight, resulting in high-variance estimators. If α_t is too close to α_{t-1} , the computational cost of the algorithm will be high. It is common to automatically construct the sequence (α_t) adaptively as the algorithm is running, where α_t is chosen at each iteration such that degeneracy is avoided [\(Del Moral](#page-18-7) et al., [2012\)](#page-18-7).
- 2. The choice of MCMC kernel K_t . The kernel needs to move efficiently around the space, with the critical point that K_t does indeed need to change with t, since a kernel that is efficient in exploring target π_1 is usually not efficient at exploring π_T . Adaptive schemes are commonly used, for example choosing the scale of a random walk proposal in a Metropolis-Hastings move at iteration t based on a scale estimated from the current particle population.

2.1.2 Iterative ensemble Kalman inversion

Whilst adaptive schemes can reduce the burden on the user of designing the SMC sampler, in some applications the implemention of an effective algorithm can still be non-trivial and the computational cost of the method prohibitively high. IEnKI was designed with the motivation of avoiding these issues.

The starting point in our description of the algorithm is in using the same sequence of targets as in equation (5) , but with the assumption that the prior p is Gaussian and that the data model q is also Gaussian, i.e.

$$
p(x) = (2\pi)^{-d_x/2} \det(\Sigma_0)^{-1/2} \exp\left(-\frac{1}{2} (x - \mu_0)^T \Sigma_0^{-1} (x - \mu_0)\right)
$$

$$
g(y_{\text{obs}} | x, \Sigma_y) = (2\pi)^{-d_y/2} \det(\Sigma_y)^{-1/2} \exp\left(-\frac{1}{2} (y_{\text{obs}} - H(x))^T \Sigma_y^{-1} (y_{\text{obs}} - H(x))\right),
$$

where $H: \mathbb{R}^{d_x} \to \mathbb{R}^{d_y}$. Let $l(x) := g(y_{obs} | x, \Sigma_y)$ be the likelihood of x and let $\tilde{\pi}_t(x) = p(x)l^{\alpha_t}(x)$ be the sequence of unnormalised targets used as in SMC, each with normalising constant Z_t . This choice results in the recursive relationship $\tilde{\pi}_t(x) = \tilde{\pi}_{t-1}(x) l^{\alpha_t-\alpha_{t-1}}(x)$. Simplifying notation by taking $\gamma_t := (\alpha_t - \alpha_{t-1})^{-1}$ we have

$$
l^{\alpha_t - \alpha_{t-1}}(x) = (2\pi)^{-\frac{dy}{2\gamma_t}} \det\left(\Sigma_y\right)^{-\frac{1}{2\gamma_t}} \exp\left(-\frac{1}{2\gamma_t} \left(y_{\text{obs}} - H(x)\right)^T \Sigma_y^{-1} \left(y_{\text{obs}} - H(x)\right)\right).
$$

Note that $l^{\gamma_t}(x)$ is proportional to the Gaussian distribution with covariance $\gamma_t \Sigma_y$. Temporarily, for the purposes of exposition, we assume that H is a linear function and switch the notation so that $H \in \mathbb{R}^{d_x \times d_y}$. Since p is Gaussian, this observation makes it clear that every member of sequence of normalised target distributions π_t is also Gaussian, and a conjugate update may be used to find $\pi_t(x)$ from $\pi_{t-1}(x)$. Let μ_t^x and C_t^{xx} respectively be the mean and covariance of π_t . We then have, following the Kalman filter update equations:

$$
S_t = HC_{t-1}^{xx}H^T + \gamma_t \Sigma_y
$$

$$
K_t = C_{t-1}^{xx}H^T S_t^{-1}
$$

$$
\mu_t^x = \mu_{t-1}^x + K_t \left(y_{\text{obs}} - H \mu_{t-1}^x \right)
$$

$$
C_t^{xx} = \left(I_{d_x} - K_t H \right) C_{t-1}^{xx}.
$$

When the Gaussian assumptions hold, there is clearly no need to use such a recursive algorithm to calculate π_t : the same result would be achieved through a single conjugate update from $p(x)$ to $p(x)l(x)$. The interest in this formulation arises when the sequence of targets is non-Gaussian. In place of the exact update, ensemble Kalman methods use a Monte Carlo approximation of the update: a population of ensemble members $\{x_t^j\}_{j=1}^M$ is used to represent the target π_t , and at each iteration these points are updated such that their sample mean and variance estimate μ_t^x and C_t^{xx} . Whilst convergence to the exact updates is guaranteed for the Gaussian case [\(Mandel](#page-18-8) *et al.*, [2011\)](#page-18-8), ensemble Kalman approaches are frequently used outside of this setting, as an approximate approach where convergence is not guaranteed [\(Katzfuss](#page-18-9) et al., [2016;](#page-18-9) [Roth](#page-19-5) et al., [2017\)](#page-19-5). IEnKI is the name given to the method that applies the ensemble Kalman idea to the sequence of targets described in this section.

We describe this approach precisely in algorithm [1,](#page-4-0) where we revert back to the non-linear/Gaussian case, taking $H: \mathbb{R}^{d_x} \to \mathbb{R}^{d_y}$. To simplify notation we use the notation $h_t^{(j)} := H(x_t^{(j)})$, using μ^h and C^{hh} to denote respectively the mean and covariance of h, and C^{xh} to denote the cross-covariance of x and h.

2.2 Marginal likelihood estimators

In this section we describe three estimators for the normalising constant of $\tilde{\pi}_T$:

$$
Z_T = \int_x p(x)l(x)dx.
$$
 (6)

2.2.1 Direct estimator

The direct estimator is derived from considering the ratio of normalising constants in the recursive algorithm in the linear-Gaussian case. We begin by noting that the incremental likelihood is given by $l^{\alpha_t-\alpha_{t-1}}(x)$ $c_t g(y_{obs} | x, \gamma_t \Sigma_u)$, where

$$
c_t = \gamma_t^{d_y/2} (2\pi)^{(1-\gamma_t)d_y/2} \det(\Sigma_y)^{(1-\gamma_t)/2},
$$

given by the ratio of the normalising constants of the Gaussian to the power $1/\gamma_t$ and the Gaussian with covariance $\gamma_t \Sigma_y$. Then

$$
\frac{Z_t}{Z_{t-1}} = \frac{\int_x p(x)l^{\alpha_t}(x)dx}{Z_{t-1}}
$$

\n
$$
= \int_x \frac{p(x)l^{\alpha_{t-1}}(x)}{Z_{t-1}}l^{\alpha_t-\alpha_{t-1}}(x)dx
$$

\n
$$
= c_t \int_x \pi_{t-1}(x)g(y_{\text{obs}} | x, \gamma_t \Sigma_y) dx
$$

\n
$$
= c_t \mathcal{N}(y_{\text{obs}} | \mu_{t-1}^h, C_{t-1}^{hh} + \gamma_t \Sigma_y),
$$

using in the last step that the integral is the marginal distribution of a multivariate Gaussian evaluated at y_{obs} . The direct estimator then defined by using

$$
\hat{Z}_T^d = \prod_{t=1}^T \frac{\widehat{Z_t}}{Z_{t-1}}\tag{7}
$$

with

$$
\frac{\widehat{Z_t}}{Z_{t-1}} = c_t \mathcal{N} \left(y_{\text{obs}} \mid \widehat{\mu}_{t-1}^h, \widehat{C}_{t-1}^{hh} + \gamma_t \Sigma_y \right). \tag{8}
$$

In the linear/Gaussian case, convergence as $M \to \infty$ of ensemble Kalman techniques [\(Mandel](#page-18-8) *et al.*, [2011\)](#page-18-8) yields convergence of $\frac{Z_t}{Z_{t-1}}$ and hence \hat{Z}_T . Note that to compute this estimate we need only run the loop in algorithm [1](#page-4-0) up to iteration $t = T - 1$.

2.2.2 Unbiased estimator

The direct estimator is consistent, but not unbiased, since $\mathcal{N}\left(y_{\text{obs}} \mid \hat{\mu}_{t-1}^h, \hat{C}_{t-1}^{hh} + \gamma_t \Sigma_y\right)$ is not an unbiased estimator of $\mathcal{N}\left(y_{\text{obs}} \mid \hat{\mu}_{t-1}^h, \hat{C}_{t-1}^{hh} + \gamma_t \Sigma_y\right)$. We can follow [Price](#page-19-6) *et al.* [\(2017\)](#page-19-6); [Drovandi](#page-18-10) *et al.* [\(2019\)](#page-18-10) in devising an unbiased version by instead using the estimator of [Ghurye and Olkin](#page-18-11) [\(1969\)](#page-18-11), which, for $M > d_y + 3$ takes the form

$$
\overline{\mathcal{N}}(y; \hat{\mu}, \hat{\Sigma}) = (2\pi)^{-d_y/2} \frac{\rho(d_y, M-2)}{\rho(d, M-1)(1 - 1/N)^{d_y/2}} |(M-1)\hat{\Sigma}|^{-(M-d_y-2)/2}
$$

$$
\psi \left((M-1)\hat{\Sigma} - (y-\hat{\mu})(y-\hat{\mu})^{\top}/(1 - 1/M) \right)^{(M-d_y-3)/2},
$$

where

$$
\rho(k, v) = \frac{2^{-kv/2} \pi^{-k(k-1)/4}}{\prod_{i=1}^{k} \Gamma(\frac{1}{2}(v-i+1))}
$$

and for a square matrix A, $\psi(A) = |A|$ if A is positive definite and $\psi(A) = 0$ otherwise. The *unbiased* estimator is then given by

$$
\hat{Z}_T^u = \prod_{t=1}^T \frac{\overline{Z_t}}{\overline{Z_{t-1}}} \qquad \overline{\frac{Z_t}{Z_{t-1}}} = \prod_{t=1}^T c_t \overline{\mathcal{N}} \left(y_{\text{obs}} \mid \hat{\mu}_{t-1}^h, \hat{C}_{t-1}^{hh} + \gamma_t \Sigma_y \right),\tag{9}
$$

where the unbiasedness of this estimator holds only in the linear/Gaussian case. We found empirically that, analogous to the finding in [Price](#page-19-6) *et al.* [\(2017\)](#page-19-6), this estimator has similar properties to the direct estimator thus we do not consider it further in this paper.

2.2.3 Path sampling estimator

Zhou [et al.](#page-19-7) [\(2015\)](#page-19-7) show how a path sampling estimator may be constructed from SMC output; here we show that this is also possible for IEnKI. [Gelman and Meng](#page-18-12) [\(1998\)](#page-18-12); [Friel and Pettitt](#page-18-13) [\(2008\)](#page-18-13) show that, for the path, over $\alpha \in [0,1]$, of distributions proportional to $p(x)l^{\alpha}(x)$, under mild regularity conditions

$$
\log(Z) = \int_{\alpha} \mathbb{E}_{\alpha} \left[\log l(.) \right] d\alpha,
$$

where \mathbb{E}_{α} denotes the expectation under the distribution proportional to $p(x)l^{\alpha}(x)$, and $Z = \int_x p(x)l(x)dx$. The sequence of distributions defined in equation [\(5\)](#page-3-0) is a discretised form of this path. Zhou *[et al.](#page-19-7)* [\(2015\)](#page-19-7) propose to use the points generated from an SMC algorithm on this sequence of targets in to approximate $log(Z)$ using a trapezoidal scheme to estimate the integral. Here we propose the analogous idea for IEnKI output:

$$
\widehat{\log(Z_T)}^p = \sum_{t=1}^T \frac{1}{2\gamma_t} \left(U_t + U_{t-1} \right),\tag{10}
$$

where

$$
U_t = \frac{1}{M} \sum_{j=1}^{M} \log l \left(x_t^j \right)
$$

=
$$
\frac{1}{M} \sum_{j=1}^{M} \log \left[\mathcal{N} \left(y_{\text{obs}} \mid h_t^j, \Sigma_y \right) \right].
$$

2.3 Alternative shifters

The "shift" step in algorithm [1](#page-4-0) involves simulating variables \tilde{y}_t^j in order that the measurement noise is correctly taken into account in the updates of the ensemble. Ensemble Kalman filters are typically used in settings where the size of the ensemble is small, thus the variance introduced by these simulations can lead to large errors in estimates from the filter. As a remedy, deterministic approaches were introduced: Monte Carlo implementations of square-root Kalman filters [\(Tippett](#page-19-8) et al., [2003\)](#page-19-8) that ensure that the ensemble has consistent first and second moments. In this paper we use both the *square root* and *adjustment* shifters. Both take the form

$$
x_t^j = x_{t-1}^j + \hat{K}_t \left(y_{\text{obs}} - \hat{\mu}_{t-1}^h \right) + b_t,
$$

where b_t is chosen such that the covariance of the sample estimates the desired posterior covariance [\(Whitaker and](#page-19-9) [Hamill, 2002;](#page-19-9) [Livings](#page-18-14) et al., [2008\)](#page-18-14). The square root and adjustment filters produce produce different ensembles that have the same covariance, thus exhibit different performance when applied to non-linear/non-Gaussian models. We will refer to IEnKI with each shifter as sIEnKI (stochastic), rIEnKI (square root) and aIEnKI (adjustment). The rIEnK and aIEnKI algorithms are given the Appendix.

3 IEnKI-ABC

In this section we investigate the use of iterative ensemble Kalman inversion in ABC (IEnKI-ABC), specifically as an alternative estimator of the ABC likelihood in equation [3.](#page-1-1) In section [3.1](#page-6-1) we introduce the new approach as an alternative to the SMC-based method introduced for rare-event ABC in [Prangle](#page-19-3) et al. [\(2018\)](#page-19-3), then in section [3.2](#page-8-0) we describe an approach to automatically chose the sequence of targets used in the method.

3.1 Method

3.1.1 From rare-event ABC to IEnKI-ABC

To estimate the ABC likelihood for parameter θ with tolerance ε , the rare-event ABC approaches of [Prangle](#page-19-3) *et al.* (2018) and [Kerama](#page-18-5) *et al.* (2022) make use of the parameterisation in equation (4) . [Prangle](#page-19-3) *et al.* (2018) introduces the idea of using an SMC algorithm on the sequence of targets $\tilde{\pi}_t(u \mid \theta) = K_{\varepsilon_t}(s_{\text{obs}} | S(G(\theta, u))) \phi(u \mid \theta)$ for $t = 0$: T, to estimate the ABC likelihood at θ , which is equivalent to the normalising constant of $\tilde{\pi}_T$. Our new

approach uses the same sequence of distributions under a reparameterisation, taking the unnormalised conditional (on θ) posterior at iteration t to be

$$
\tilde{\pi}_t(s \mid \theta) = f_s(s \mid \theta) K_{\varepsilon_t} (s_{\text{obs}} \mid s), \qquad (11)
$$

where f_s is the distribution of the transformed variable $S(x)$ with $x \sim f(\cdot | \theta)$. IEnKI-ABC then uses the estimators in section [2](#page-2-1) to estimate the integral

$$
\int_{s} f_s(s \mid \theta) K_{\varepsilon} (s_{\text{obs}} \mid s) ds,
$$
\n(12)

through the use of a sequence of distributions $\tilde{\pi}_t(s | \theta) = f_s(s | \theta) K_{\varepsilon_t}(s_{obs} | s)$ for $\infty = \varepsilon_0 > ... > \varepsilon_T = \varepsilon$. It is straightforward to see that, for $\varepsilon > 0$, this corresponds to a sequence of temperatures in IEnKI of $\alpha_t = (\varepsilon/\varepsilon_t)^2$, so that $\gamma_t = \varepsilon^{-2} \left(\varepsilon_t^{-2} - \varepsilon_{t-1}^{-2} \right)^{-1}$. As an example of the use of these methods in this context, we show IEnKI with a stochastic shifter in algorithm [2,](#page-7-0) the output of which may be used for any of the estimators in section [2.2.](#page-4-1)

Algorithm 2: Stochastic iterative ensemble Kalman inversion for ABC, for a given θ and sequence (ε_t) .

1 Simulate *M* points,
$$
\{x_0^j\}_{j=1}^M \sim f(\cdot | \theta)
$$
;
\n2 For $j = 1 : M$, let $s_0^j = S(x_0^j)$;
\n3 $\hat{\mu}_0^s = \frac{1}{M} \sum_{j=1}^M s_0^j$;
\n4 $\hat{C}_0^{ss} = \frac{1}{M-1} \sum_{j=1}^M \left(s_0^j - \hat{\mu}_0^s\right) \left(s_0^j - \hat{\mu}_0^s\right)^T$;
\n5 for $t = 1 : T$ do
\n6 $\hat{K}_t = \hat{C}_{t-1}^{ss} \left(\hat{C}_{t-1}^{ss} + \varepsilon^{-2} \left(\varepsilon_t^{-2} - \varepsilon_{t-1}^{-2}\right)^{-1} \Sigma_s\right)^{-1}$;
\n7 $\text{for } n = 1 : N_a$ do
\n8 $\left| \begin{array}{l} \text{Simulate } \tilde{s}_t^j \sim \mathcal{N}\left(s_{t-1}^j, \varepsilon^{-2} \left(\varepsilon_t^{-2} - \varepsilon_{t-1}^{-2}\right)^{-1} \Sigma_s\right) \\ \text{Similars } \tilde{s}_t^j \sim \mathcal{N}\left(s_{t-1}^j, \varepsilon^{-2} \left(\varepsilon_t^{-2} - \varepsilon_{t-1}^{-2}\right)^{-1} \Sigma_s\right); \end{array} \right.$
\n9 $\left| \begin{array}{l} \text{Shift } s_t^j = s_{t-1}^{(j)} + \hat{K}_t \left(s_{\text{obs}} - \tilde{s}_t^j\right); \\ \text{end} \right.$
\n10 $\hat{\mu}_t^s = \frac{1}{M} \sum_{j=1}^M s_t^j;$
\n12 $\hat{C}_t^{ss} = \frac{1}{M-1} \sum_{j=1}^M \left(s_t^j - \hat{\mu}_t^s\right) \left(s_t^j - \hat{\mu}_t^s\right)^T;$
\n13 end

3.1.2 Remarks

This approach has the following desirable properties:

- In contrast to rare-event ABC, IEnKI-ABC estimates the likelihood through exploring the space of s, rather than x, which is considerably easier given that we typically expect $d_s \ll d_x$.
- The use of the shifting operations of IEnKI means that no proposals for exploring s-space must be designed.
- We observe that the use of IEnKI within ABC is more straightforward than the general case presented in section [2,](#page-2-1) since the "state" (x) and "measurement" (y) spaces are the same: no H function is required to map x to y. Non-linearity in this function is one of the potential sources of error in estimators based on IEnKI, but this issue is avoided in the ABC context.
- Given simulations from f , the approach is very easy to implement. The main tuning parameter is the sequence (ε_t) , the choice of which we describe in section [3.2.](#page-8-0)
- Given that models commonly encountered in ABC are expensive to simulate, the computational cost of the approach will often be dominated by their simulation. If τ is the cost of simulating from f, the complexity of: standard ABC is $O(M\tau)$; stochastic IEnKI-ABC is $O(M\tau + Md_s^2 + d_s^3)$; square root IEnKI-ABC is $O(M\tau + Md_s^2)$; and adjustment IEnKI-ABC is $O(M\tau + M^2d_s + M^3)$ [\(Whitaker and Hamill, 2002\)](#page-19-9). The scaling of adjustment IEnKI-ABC with M means that it is only of practical use in cases where $d_s < M$, which are not considered in this paper.

3.1.3 Comparison to alternative methods

IEnKI has previously been used in the context of SBI by [Duffield and Singh](#page-18-15) [\(2022\)](#page-18-15), where IEnKI was used on the parameter space θ , rather than in the ABC likelihood estimator. The main difference between our new approach and this previous work is that [Duffield and Singh](#page-18-15) [\(2022\)](#page-18-15) provides a Monte Carlo approximation of the true posterior only when the joint distribution of θ and s is Gaussian, whereas IEnKI-ABC relies only on the conditional distribution P_s of s θ being Gaussian.

Synthetic likelihood (SL) is an alternative to ABC introduced by [Wood](#page-19-10) [\(2010\)](#page-19-10) and further developed by [Price](#page-19-6) [et al.](#page-19-6) [\(2017\)](#page-19-6). This approach also assumes P_s to be Gaussian, taking its mean and variance to be the sample mean $\hat{\mu}^s$ and covariance \hat{C}^{ss} from simulations from P_s , and taking the likelihood to be $\mathcal{N}\left(s_{\text{obs}} \mid \hat{\mu}^s, \hat{C}^{ss}\right)$ (or using the unbiased estimator in section [2.2.2,](#page-5-0) proposed in [Price](#page-19-6) et al. [\(2017\)](#page-19-6)).

SL is a limiting case of IEnKI-ABC, where the sequence of distributions contains only π_0 and π_1 , with $\varepsilon_0 = \infty$ and $\varepsilon_1 = 0$. Here K_{ε_1} is a Dirac delta and the integral in equation [12](#page-7-1) collapses to $f_s(s_{obs} | \theta)$. SL corresponds to implementing lines 1-4 of algorithm [2](#page-7-0) and using the result to calculate the SL estimator. As we will see in section [4.1,](#page-9-1) SL gives almost identical results to IEnKI-ABC when P_s is Gaussian. IEnKI-ABC is likely to improve upon the performance of SL when P_s is non-Gaussian by using a sequence of distributions as described in the paper. We observe such an improvement in section [4.2.](#page-10-0)

3.2 Sequences of targets

3.2.1 An adaptive procedure

The choice of the temperatures (α_t) signficantly affects the performance of the approach. [Iglesias](#page-18-1) *et al.* [\(2018\)](#page-18-1) introduces the idea of adapting the sequence of temperatures as the algorithm is running, following the corresponding idea in SMC introduced by [Del Moral](#page-18-7) et al. [\(2012\)](#page-18-7). At the beginning of each iteration of the algorithm, a new temperature α_t is found by using a bisection algorithm that aims to ensure that the distance between successive targets is not too large. In SMC, the *effective sample size* (ESS), calculated from particle weights, is used as an approximation to the chi-square distance between successive targets. [Iglesias](#page-18-1) *et al.* [\(2018\)](#page-18-1) proposes to use the same approach, where weights assigned to ensemble members (to be used only in this adaptive procedure) are defined to be precisely those that would be assigned to particles in an SMC sampler with MCMC moves. Ensemble member j at iteration t is assigned (unnormalised) weight

$$
\tilde{w}_t^j = \exp\left(-\frac{\varepsilon_t^{-2} - \varepsilon_{t-1}^{-2}}{2} \left(s_{\text{obs}} - s_{t-1}^j\right)^T \Sigma_s^{-1} \left(s_{\text{obs}} - s_{t-1}^j\right)\right).
$$

Each weight is a function of the choice of ε_t , and is calculated at each ε_t considered in the bisection procedure. The bisection aims to find ε_t such that the ESS:

$$
ESS = \frac{\left(\frac{1}{M}\sum_{j=1}^{M}\tilde{w}_t^j\right)^2}{\frac{1}{M}\sum_{j=1}^{M}\left(\tilde{w}_t^j\right)^2}
$$

is a proportion β of the sample size M.

 \mathcal{L}

As in SMC this adaptive procedure introduces a small bias into estimates produced from the resultant IEnKI: the properties of the analogous scheme for SMC are investigated empirically in [Prangle](#page-19-3) *et al.* [\(2018\)](#page-19-3). The approach has the appeal that it reduces the tuning of the sequence of temperatures to the choice of a single parameter β. However, the length of the sequence chosen for parameters with a small likelihood can be large (resulting in additional computation), so we instead consider an alternative approach, tailored to the particular sequence of targets in equation [\(11\)](#page-7-2).

3.2.2 Choice of tempering scheme

Consider an infinite sequence of (unnormalised) distributions $\tilde{\pi}_t(s | \theta) = P_s(s | \theta) K_{\varepsilon_t}(s_{\text{obs}} | s)$ where $\varepsilon_t \to 0$, recalling that $K_{\varepsilon_t}(s_{\text{obs}} | s)$ is Gaussian with mean s and covariance $\varepsilon_t^2 \Sigma_s$. For a fixed θ , this sequence has similar properties to a sequence of Bayesian posterior distributions with an increasing number n of data points, with $P_s(\cdot | \theta)$ playing the role of the prior and K_{ε_t} the role of a Gaussian likelihood with mean s_{obs} , covariance Σ_s with $n = \varepsilon_t^{-2}$ observations. As long as $P_s(\cdot \mid \theta)$ is absolutely continuous in a neighbourhood of s_{obs} with continuous positive density at s_{obs} , direct application of the Bernstein-von Mises theorem [\(van der Vaart](#page-19-11) [\(1998\)](#page-19-11); Chapter 10.2) yields that as $t \to \infty$ (i.e. $\varepsilon_t \to 0$ equating to $n \to \infty$), $\tilde{\pi}_t(s | \theta) \to \mathcal{MVN}(s_{obs}, \varepsilon_t^2 \Sigma_s)$.

In this section we make use of this property, together with the recent work of [Chopin](#page-18-16) et al. [\(2024\)](#page-18-16) which suggests a principled approach to choosing a tempering sequence (α_t) , and hence a sequence (ε_t) , through connection to mirror descent. We follow this approach, using the assumption that the sequence of targets is Gaussian. We suppose that π_0 and π_T are both multivariate normal distributions with common mean s_{obs} and covariance Σ_0 and $\varepsilon^2 \Sigma_s$ respectively. Through the argument above, the normality assumption for π_T is likely to be realistic when ε is small. Recall that Σ_s is a diagonal matrix chosen to scale the summary statistics. We assume that Σ_0 is some multiple κ^2 of Σ_s : this is not unreasonable, since in practice a common choice of Σ_s is to use the diagonal entries of an estimate $\hat{\Sigma}_0$, obtained through simulation from f, of Σ_0 (in this case we would have $\kappa^2 = 1$). In the empirical studies below, we take κ to be the mean of $\hat{\sigma}_{i,0}/\sigma_{i,s}$, where $\sigma_{i,s}$ is the square root of the *i*th element on the diagonal of Σ_s and $\hat{\sigma}_{i,0}$ is the sample standard deviation of the *i*th statistic over the M simulations from P_s . The argument of [Chopin](#page-18-16) [et al.](#page-18-16) [\(2024\)](#page-18-16) (details in the Appendix) results in continuous tempering schedule of

$$
\alpha(t) = \exp\left(2\log\left(\frac{\kappa}{\varepsilon}\right)t + \log\left(\frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}\right)\right) - \frac{\varepsilon^2}{\kappa^2 - \varepsilon^2} \tag{13}
$$

for $t \in [0,1]$. We propose using a discretised version (α_t) of this schedule, taking $\alpha_t = \alpha(t/T)$ for $t = 0$: T.

3.2.3 Target skipping

The result that $\tilde{\pi}_t(s | \theta) \to \mathcal{MVN}(s_{obs}, \varepsilon_t^2 \Sigma_s)$ has a further implication: when using a finite sequence of targets $t = 0$: T in IEnKI, when ε_T is chosen to be small, we might expect that for some $t < T$ the ensemble representing a from $\pi_t(s | \theta)$ will be close to a sample from a Gaussian distribution. Recall that the motivation for using the iterative EnKI algorithm arose purely from the acknowledgement that we do not expect the sequence of targets to be Gaussian. As soon as we detect that a target $\pi_t(s | \theta)$ is close to Gaussian, we may also assume that targets for $t' > t$ are also close to Gaussian, and we may use a single EnK update step to move our ensemble from $\pi_t(s | \theta)$ to $\pi_T(s \mid \theta)$. We envisage that this approach will lead to computational savings without sacrifiing much accuracy.

Based on this principle, we propose the following approach. At each iteration of algorithm [2,](#page-7-0) directly after line 5, we perform a hypothesis test where the null hypothesis is that the ensemble is drawn from a multivariate Gaussian distribution. If the null hypothesis is accepted at iteration t , in the remainder of the IEnKI we "skip" targets $t + 1$: T – 1 and perform the update (lines 6-10) moving directly to target π_T . In this paper we used the Henze-Zirkler test [\(Henze and Zirkler, 1990\)](#page-18-17) which has been illustrated to be powerful in a range of scenarios (e.g. [\(Farrell](#page-18-18) et al., [2007;](#page-18-18) [Ebner and Henze, 2020\)](#page-18-19).

4 Applications

4.1 Gaussian marginal likelihood

In this section we investigate the empirical properties of the new marginal likelihood estimators introduced in this paper. We study a toy example where the true marginal likelihood is available, and where the Gaussian assumptions underpinning IEnKI are all satisfied. We take $P_s(s | \theta) = \mathcal{N}(s | \theta, 1)$, $K_{\varepsilon}(s_{\text{obs}} | s) = \mathcal{N}(s_{\text{obs}} | s, \varepsilon^2)$, so that the exact ABC likelihood is available via

$$
\int_{s} K_{\varepsilon} (s_{\text{obs}} | s) P_{s} (s | \theta) ds = \mathcal{N} (s_{\text{obs}} | \theta, 1 + \varepsilon^{2})
$$

.

In this section we use $s_{obs} = 0$ and $\theta = 0$ and examine the properties of ABC, SL and IEnKI-ABC estimates of the ABC likelihood. To accurately compare SL with the other two approaches we set the data generating process used in this approach to be $\mathcal{N}(\cdot \mid 0, 1 + \varepsilon^2)$. We compared the algorithms for tolerances $\varepsilon = 0.1, 0.01, 0.001, 0.0001$, $M = 10, 50, 100, 200$ points simulated from P_s . and, for IEnKI-ABC, $T = 5, 10, 20$ targets (plus $T = 50, 100, 200$ for the path sampling estimator). The stochastic and square root shifting approaches were compared for IEnKI-ABC, using the direct (equation [\(7\)](#page-5-1)) and path sampling (equation [\(10\)](#page-6-2)) estimators. Since the model is linear/Gaussian, the square root and adjustment approaches produce almost identical results, thus the adjustment approach is omitted. All experiments were performed using the [ilike](https://github.com/richardgeveritt/ilike) R package, with the model written in C_{++} . Each algorithm was run 100 times, and the bias, standard deviation, root mean squared error (RMSE) of IEnKI marginal likelihood estimators was estimated from these 100 runs. In this section we examine the properties of the direct estimator from IEnKI-ABC as ε changes, and as T changes.

Figure 1: Properties of IEnKI normalising constant estimators on the Gaussian example.

Figure [1a](#page-10-1) shows how the RMSE of the different approaches scales with ε , where $M = 200$ and $T = 5$. We observe a dramatic increase in RMSE for ABC as ε gets closer to 0; the other schemes have the desirable property that the RMSE is similar for all values of ε . The rIEnKI-ABC and SL results are identical up to rounding errors, whereas we observe additional error, due to the additional variance introduced by the shifter in sIEnKI-ABC. Figure [1b,](#page-10-2) shows the log RMSE of the direct estimator for $M = 200$ and $\varepsilon = 0.01$, clearly showing the effect on the direct estimator from sIEnKI-ABC of increasing the number of targets T . We observe an increased variance as T increases for sIEnKI-ABC, whereas rIEnKI-ABC and aIEnKI-ABC are not affected by changes in T. In this example, where a conjugate Gaussian update holds, the best choice for T (giving lower variance and computational cost) is 1. The non-Gaussian example in section [4.2](#page-10-0) will show a case where it is useful to choose $T > 1$.

Further results and discussion are found in the Appendix, including an illustration that the error of path sampling IEnKI estimators increases as ε decreases, reducing the appeal of this approach (compared to the direct estimator) for use in ABC.

4.2 Lotka-Volterra model

In this section we apply IEnKI-ABC to the well-studied stochastic Lotka-Volterra (LV) predator-prey model. We use the Markov jump process version of the model with three parameters, as previously studied in [Wilkinson](#page-19-12) [\(2013\)](#page-19-12); [Papamakarios and Murray](#page-19-13) [\(2016\)](#page-19-13). Let x_1 be the number of predators and x_2 the number of prey, evolving with the following reactions:

- A prey may be born, with rate $\theta_1 x_2$, increasing x_2 by one.
- The predator-prey interaction in which x_1 increases by one and x_2 decreases by one, with rate $\theta_2 x_1 x_2$.
- A predator may die, with rate θ_3x_1 , decreasing x_1 by one.

ABC is a popular approach to inference in such models, even though it can be significantly less efficient than other approaches, such as particle MCMC. In this section we take the opportunity to compare several different approaches, alongside IEnKI-ABC. Let $x_{t,1}$ and $x_{t,2}$ be, respectively, the number of predator and prey at time t and $x_t = (x_{t,1}, x_{t,2})^T$. We simulate the LV model using the Gillespie algorithm via C++ code adapted from the smfsb R package [\(Wilkinson, 2024\)](#page-19-14), using the LVperfect data in the smfsb package to be our observed data y_{obs} , consisting of measurements of $x_{t,1}$ and $x_{t,2}$ at each time t (consisting of the the initial states at time 0, plus measurements at 15 further times). We use the same model for every inference algorithm we test, with the model having slightly different interpretations for the different algorithms used to estimate the marginal likelihood of θ :

• For ABC and IEnKI-ABC we take P to be the LV model so that $y = x$ (i.e. simulations from the LV model are treated as our observed data), take $s = y$ (i.e. use the full data rather than summary statistics), and choose K_{ε} to be Gaussian with covariance $\varepsilon^2 \Sigma_s$, where Σ_s is taken to be the identity.

- For SL, we take $y_{t,1} \sim \mathcal{N}(\cdot \mid x_{t,1}, \varepsilon^2), y_{t,2} \sim \mathcal{N}(\cdot \mid x_{t,2}, \varepsilon^2)$ to be noisy measurements of $x_{t,1}, x_{t,2}$, which are simulated from the LV model for $t = 0$: 15. We again take $s = y$.
- We also use a bootstrap particle filter (PF) [\(Gordon](#page-18-20) *et al.*, [1993\)](#page-18-20) and ensemble Kalman filters (EnKF) [\(Evensen, 1994\)](#page-18-21). These filters also take $y_{t,1} \sim \mathcal{N}(\cdot \mid x_{t,1}, \varepsilon^2), y_{t,2} \sim \mathcal{N}(\cdot \mid x_{t,2}, \varepsilon^2)$ to be noisy measurements of $x_{t,1}, x_{t,2}$ simulated from the LV model, and perform filtering on x given y. We use EnKFs with stochastic, square root and adjustment shifters - previously only the stochastic shifter has been investigated in the context of marginal likelihood estimation [\(Drovandi](#page-18-10) et al., [2019\)](#page-18-10).

4.2.1 Marginal likelihood estimation

In this section we investigate the empirical properties of the marginal likelihood estimators mentioned above, in a similar manner to section [4.1.](#page-9-1) This section provides a clearer picture of the relative efficiency of IEnKI-ABC in a more realistic situation, using a model that does not satisfy the Gaussian assumptions outlined in section [2.1.2.](#page-3-1) All experiments were performed using the [ilike](https://github.com/richardgeveritt/ilike) R package, with the model written in C_{++} . In this section we study the standard deviation (SD) of the likelihood estimator, since this is the main determining factor in the efficiency of an MCMC algorithm using the likelihood estimator, and we found the bias is difficult to estimate accurately.

We estimated the likelihood at the point at which the LVperfect data was generated, $\theta = (1, 0.005, 0.6)^T$. All methods used $M = 100$ points: for the PF and EnKF this corresponds to the number of particles and ensemble members respectively. We used EnKF with stochastic (sEnKF), square root (rEnKF) and adjustment (aEnKF) shifters. We also compared three IEnKI-ABC estimators, all using the stochastic shifter and with $T = 100$: the direct estimator (labelled "sIEnKI-ABC"); the path estimator ("sIEnKI-ABCpath") and the direct estimator where the target skipping approach is used with significance $\alpha = 0.1$ ("sIEnKI-ABCskip"). Results for the deterministic shifters in IEnKI-ABC are not presented here: poorly conditioned covariance matricies (due to diverging LV simulations) rendered these methods inaccurate. The cost of most of the methods was dominated by the cost of simulating from the LV model $M = 100$ times, taking approximately 0.1s. The cost of aEnKF was marginally higher at 0.13s, and the use of $T = 100$ targets increased the cost of the IEnKI methods, with sIEnKI-ABC taking 0.25s and sIEnKI-ABCpath 0.34s, except when using sIEnKI-ABCskip whose cost was again 0.1s. [This video](https://youtu.be/nERovsRTDzI) illustrates a run of sIEnKI-ABC for in this situation.

In figure [2a](#page-12-0) we compare the log SD of the likelihood estimators from each of the methods mentioned in the previous section, as a function of the log of ε . ABC and PF exhibit apparently exponentially increasing SD with the decrease in ε , with SL also following this trend when ε is small. The EnKF exhibits the best performance for all ε and the SD only slightly increases as ε decreases. The deterministic shifters offer slight improvements over the stochastic shifter. The SD of the sIEnKI-ABC approaches is also not dramatically affected by the decrease in ε . The target skipping approach results in a smaller SD due to omitting IEnKI steps that would not dramatically change the likelihood estimates but would result in additional noise via the stochastic shifter.

Figure [2b](#page-12-1) examines the effect on sIEnKI-ABC and sIEnKI-ABCpath of changing the number of targets T (for $\varepsilon = 0.1$ and $M = 100$). For the path estimator we observe that a relatively large number of targets are required in order for the numerical integration approach to achieve a small error. For the direct approach there is again an increase in the error as T increases due to the stochastic shifter, as observed in section [4.1.](#page-9-1) However, taking $T > 1$ results in significant benefits compared to the $T = 1$ case, represented in figure [2a](#page-12-0) by the SL results which are very poor due to f_s being far from Gaussian.

4.2.2 Use in MCMC

We ran Metropolis-Hastings (MH) algorithms using each of the likelihood estimators in the previous section, taking $M = 100$ for all algorithms. Each algorithm was run for 10^6 iterations and was initialised at $\theta = (1, 0.005, 0.6)^T$. For the IEnKI-ABC approaches, we used only the target skipping approach, taking $T = 100$ and $\alpha = 0.01$ which results in a comparable computational cost for all algorithms used in this section. A multivariate Gaussian proposal was used, with a covariance determined through pilot runs. The estimated posterior distributions for $\varepsilon = 10$ and $\varepsilon = 0.1$ are shown in figure [3](#page-12-2) and the multiESS (from the mcmcse R package) for each run in table [1.](#page-12-3) For several algorithms there are only a few acceptances in the whole run: for these the multiESS cannot be estimated reliably and so we report it simply as < 50 .

ABC-MCMC and SL-MCMC perform very poorly for both choices of ε . Particle marginal MH (using the PF estimate in MH) performs well for $\varepsilon = 10$, but poorly for $\varepsilon = 0.1$. Ensemble MCMC (using the EnKF estimate in MH) performs very well for both choices of ε . For $\varepsilon = 0.1$ IEnKI-ABC-MCMC significantly outperforms all approaches except ensemble MCMC, although some bias is evident in the IEnKI-ABC-MCMC posteriors.

(a) log SD as a function of the log tolerance $\varepsilon.$ (b) log SD of estimator as a function of the number of targets T.

Figure 2: Properties of alternative likelihood estimators on the Lotka-Volterra example.

(a) Estimated marginal posterior densities for the Lotka-Volterra model (b) Estimated marginal posterior densities for the Lotka-Volterra model when $\varepsilon = 10$. when $\varepsilon = 0.1$.

Figure 3: Estimated marginal posterior densities for the Lotka-Volterra model when $\varepsilon = 10$ (left) and $\varepsilon = 0.1$ (right).

| ε | ABC | | РF | EnKF | sEnKLABC |
|----|-----------|---------|-----------|-------|----------|
| 10 | $< \! 50$ | ${<}50$ | 27307 | 56665 | 6085 |
| | $< \! 50$ | ${<}50$ | $< \! 50$ | 47181 | 2408 |

Table 1: The multiESS for each MCMC algorithm for the Lotka-Volterra model.

5 Discussion

This paper introduces new, straightforward to implement, methods for estimating the ABC likelihood based on IEnKI that we show empirically to have substantially lower variance than the standard approach, in particular we see that the variance grows much more slowly as $\varepsilon \to 0$. The direct IEnKI-ABC estimator is also related to the SL approach. For cases where the summary statistic distribution f_s is close to Gaussian, the performance of IEnKI-ABC is likely to be similar to that of SL, however we see empirically that IEnKI-ABC outperforms SL for the LV model when f_s is non-Gaussian. IEnKI-ABC uses sequence of target distributions analogous to the sequence used in SMC approaches. We introduce an approach to choosing this sequence automatically, leaving the only tuning parameter to be the number of targets, and also introduce a method for automatically skipping to the end of the target sequence to save computation and reduce variance. IEnKI-ABC is applicable in all cases where ABC may be applied, thus is more generally applicable that some alternative approaches: for example the EnKF methods used in section [4.2.](#page-10-0)

Since IEnKI is being used outside of the linear-Gaussian setting, estimators based on its output will have a bias that is difficult to quantify: a feature avoided when using rare-event ABC. In particular, we might expect a large bias in situations where members of the sequence $\tilde{\pi}_t(s | \theta)$ are not unimodal. However, we expect in many applications that IEnKI-ABC posteriors are likely to have lower bias than those from SL, which are more influenced by the Gaussian assumption, and ABC, due to the necessity in using a large tolerance.

A IEnKI with deterministic shifters

IEnKI algorithms using square root and adjustment filters are given in algorithms [3](#page-13-1) and [4](#page-14-0) respectively.

Algorithm 3: Square root iterative ensemble Kalman inversion.

1 Simulate *M* points,
$$
\left\{x_0^j\right\}_{j=1}^M \sim p
$$
;
\n2 For $j = 1 : M$, let $h_0^j = H(x_0^j)$;
\n3 $\hat{\mu}_0^x = \frac{1}{M} \sum_{j=1}^M x_0^j$;
\n4 $\hat{\mu}_0^h = \frac{1}{M} \sum_{j=1}^M h_0^j$;
\n5 $\hat{C}_0^{xh} = \frac{1}{M-1} \sum_{j=1}^M \left(x_0^j - \hat{\mu}_0^x\right) \left(h_0^j - \hat{\mu}_0^h\right)^T$;
\n6 $\hat{C}_0^{hh} = \frac{1}{M-1} \sum_{j=1}^M \left(h_0^j - \hat{\mu}_0^h\right) \left(h_0^j - \hat{\mu}_0^h\right)^T$;
\n7 for $t = 1 : T$ do
\n8 $\begin{array}{r} \hat{K}_t = \hat{C}_{t-1}^{xh} \left(\hat{C}_{t-1}^{hh} + \gamma_t \Sigma_y\right)^{-1}$;
\n9 $\hat{S}_t = \hat{C}_{t-1}^{hh} + \gamma_t \Sigma_y$;
\n10 $\hat{K}_t = \hat{C}_{t-1}^{xh} \left(\hat{S}_t^{1/2} + \sqrt{\gamma_t} \Sigma_y^{1/2}\right)^{-1}$;
\n11 $\begin{array}{r} \text{for } n = 1 : N_a \text{ do} \\ \text{Shift } x_t^j = x_{t-1}^j + \hat{K}_t \left(y_{\text{obs}} - \hat{\mu}_{t-1}^h\right) - \bar{K}_t \left(h_{t-1}^{(j)} - \hat{\mu}_{t-1}^h\right); \\ \text{Left } h_t^j = H(x_t^j); \end{array}$

Each approach has a different computational complexity: $O(M d_x d_y + M d_y^2 + d_y^3 + d_x d_y^2)$ for sIEnKI, $O(M d_y +$ Md_xd_y) for rIEnKI and $O(M^2d_y + M^3 + M^2d_x)$ for aIEnKI [\(Whitaker and Hamill, 2002\)](#page-19-9).

Algorithm 4: Adjustment iterative ensemble Kalman inversion.

1 Simulate *M* points, $\left\{x_0^j\right\}^M$ $_{j=1} \sim p;$ 2 For $j = 1 : M$, let $h_0^j = H(x_0^j);$ **3** $\hat{\mu}_0^x = \frac{1}{M} \sum_{j=1}^M x_0^j;$ 4 $\hat{\mu}_0^h = \frac{1}{M} \sum_{j=1}^M h_0^j;$ 5 $\hat{Z}_0^x = \frac{1}{\sqrt{M}}$ $\frac{1}{M-1}\sum_{j=1}^M\left(x_0^j-\hat{\mu}_0^x\right);$ 6 $\hat{Z}_0^h = \frac{1}{\sqrt{M}}$ $\frac{1}{\overline{M-1}}\sum_{j=1}^M\Big(h_0^j-\hat{\mu}_0^h\Big);$ 7 $P_0 W_0^{1/2} V_0^T = {\rm SVD}\left(\hat{Z}_0^x\right);$ 8 for $t = 1 : T$ do $\mathbf{9} \quad \Big\vert \quad U_t^T D_t U_t = \text{SVD}\left(V_{t-1}^T \left(I_{d_x} + (\hat{Z}_{t-1}^h)^T (\gamma_t \Sigma_y)^{-1} \hat{Z}_{t-1}^h \right) V_{t-1} \right);$ $\texttt{10} \quad \hat{K}_t = \hat{Z}_{t-1}^x(\hat{Z}_{t-1}^h)^T \left(\hat{Z}_{t-1}^h(\hat{Z}_{t-1}^h)^T + \gamma_t \Sigma_y \right)^{-1};$ 11 $A_t = P_{t-1} W_{t-1}^{1/2} U_t D_t^{1/2} W_{t-1}^{-1/2} P_{t-1}^T;$ 12 for $n = 1 : N_a$ do 13 Shift $x_t^j = \hat{\mu}_{t-1}^x + \hat{K}_t (y_{\text{obs}} - \hat{\mu}_{t-1}^h) + A_t (x_{t-1}^{(j)} - \hat{\mu}_{t-1}^x);$ 14 | Let $h_t^j = H(x_t^j);$ ¹⁵ end 16 $\hat{\mu}_t^x = \frac{1}{M} \sum_{j=1}^M x_t^j;$ 17 $\hat{\mu}_t^h = \frac{1}{M}\sum_{j=1}^M h_t^j;$ 18 $\hat{Z}_t^x = \frac{1}{\sqrt{M}}$ $\frac{1}{M-1}\sum_{j=1}^M\left(x_t^j-\hat{\mu}_t^x\right);$ 19 $\hat{Z}_t^h = \frac{1}{\sqrt{M}}$ $\frac{1}{\overline{M-1}}\sum_{j=1}^M \left(h^j_t - \hat{\mu}^h_t\right);$ 20 $P_t W_t^{1/2} V_t^T = \text{SVD}(\hat{Z}_t^x);$ ²¹ end

B Sequence of targets for IEnKI-ABC

When d_s is large, [Chopin](#page-18-16) *et al.* [\(2024\)](#page-18-16) suggests deriving a tempering sequence by solving the ODE

$$
\dot{\alpha} = cI(\alpha)^{-1/2},\tag{14}
$$

where $I(\alpha)$ is the Fisher information of α under the tempered target. We follow this approach here, using the assumption that the sequence of targets is Gaussian. We suppose that π_0 and π_T are both multivariate normal distributions with common mean s_{obs} and covariance Σ_0 and $\varepsilon^2 \Sigma_s$ respectively. Through the argument at the beginning of section [3.2.2,](#page-8-1) the normality assumption for π_T is likely to be realistic when ε is small. Recall that Σ_s is a diagonal matrix chosen to scale the summary statistics. To simplify the following derivation we assume that Σ_0 is some multiple κ^2 of Σ_s : this is not unreasonable, since in practice a common choice of Σ_s is to use the diagonal entries of an estimate $\hat{\Sigma}_0$, obtained through simulation from f, of Σ_0 (in this case we would have $\kappa^2 = 1$). Following [Chopin](#page-18-16) et al. [\(2024\)](#page-18-16) we obtain that $\pi_t(s | \theta)$ is a multivariate normal distribution $\mathcal{N}(s | s_{\text{obs}}, \frac{\varepsilon^2 \kappa^2}{\varepsilon^2 + (\kappa^2 - \varepsilon^2)})$ $\frac{\varepsilon^2 \kappa^2}{\varepsilon^2 + (\kappa^2 - \varepsilon^2) \alpha_t} \Sigma_s$), and hence

$$
I(\alpha) = -\mathbb{E}\left[\frac{\partial^2 \log \pi_{\ell}(s|\theta)}{\partial \alpha^2}\right]
$$

\n
$$
= -\mathbb{E}\left[\frac{\partial^2}{\partial \alpha^2}\left(-\frac{d_s}{2}\log(2\pi) - \frac{1}{2}\log\det\left(\frac{\varepsilon^2 \kappa^2}{\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha}\Sigma_s\right) - \frac{\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha}{2\varepsilon^2 \kappa^2} (s - s_{\text{obs}})^T \Sigma_s^{-1} (s - s_{\text{obs}})\right)\right]
$$

\n
$$
= \frac{1}{2}\mathbb{E}\left[\frac{\partial}{\partial \alpha}\left(\frac{\det\left(\frac{\varepsilon^2 \kappa^2}{\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha}\Sigma_s\right) \operatorname{tr}\left(\left(\frac{\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha}{\varepsilon^2 \kappa^2}\Sigma_s\right) - \frac{\varepsilon^2 \kappa^2 (\varepsilon^2 - \kappa^2)}{(\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha)^2}\Sigma_s\right)\right)}{\det\left(\frac{\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha}{\varepsilon^2 \kappa^2}\Sigma_s\right)}\right)\right]
$$

\n
$$
+ \frac{1}{2}\mathbb{E}\left[\frac{\partial}{\partial \alpha}\left((s - s_{\text{obs}})^T\left(\frac{\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha}{2\varepsilon^2 \kappa^2}\Sigma_s^{-1} + \left(\frac{\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha}{2\varepsilon^2 \kappa^2}\Sigma_s^{-1}\right)^T\right)\right)\right]
$$

\n
$$
= \frac{1}{2}\mathbb{E}\left[\frac{\partial}{\partial \alpha}\left(\frac{(\varepsilon^2 - \kappa^2)d_s}{\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha}\right)\right]
$$

\n
$$
+ \mathbb{E}\left[\frac{\partial}{\partial \alpha}\left((s - s_{\text{obs}})^T\left(\frac{\varepsilon^2 + (\kappa^2 - \varepsilon^2)\alpha}{2\varepsilon^2 \kappa^
$$

The specific instance of equation [14](#page-15-0) is then

$$
\dot{\alpha} = c \left(\frac{\varepsilon^2}{(\kappa^2 - \varepsilon^2) \sqrt{d_s}} + \frac{\alpha}{\sqrt{d_s}} \right),
$$

$$
\alpha(t) = \exp \left(\frac{ct + k}{\sqrt{d_s}} \right) - \frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}.
$$

whose solution is

We use this solution to determine a continuous tempting schedule that has
$$
\alpha(0) = 0
$$
 and $\alpha(1) = 1$. We obtain

$$
\frac{\varepsilon^2}{\kappa^2 - \varepsilon^2} = \exp\left(\frac{k}{\sqrt{d_s}}\right)
$$

$$
\frac{k}{\sqrt{d_s}} = \log\left(\frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}\right)
$$

$$
k = \sqrt{d_s} \log\left(\frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}\right)
$$

giving

$$
\alpha(t) = \exp\left(\frac{ct + \sqrt{d_s} \log\left(\frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}\right)}{\sqrt{d_s}}\right) - \frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}
$$

,

then

$$
\exp\left(\frac{c+\sqrt{d_s}\log\left(\frac{\varepsilon^2}{\kappa^2-\varepsilon^2}\right)}{\sqrt{d_s}}\right) = 1 + \frac{\varepsilon^2}{\kappa^2-\varepsilon^2}
$$
\n
$$
c+\sqrt{d_s}\log\left(\frac{\varepsilon^2}{\kappa^2-\varepsilon^2}\right) = \sqrt{d_s}\log\left(1+\frac{\varepsilon^2}{\kappa^2-\varepsilon^2}\right)
$$
\n
$$
c = \sqrt{d_s}\log\left(1+\frac{\varepsilon^2}{\kappa^2-\varepsilon^2}\right) - \sqrt{d_s}\log\left(\frac{\varepsilon^2}{\kappa^2-\varepsilon^2}\right)
$$
\n
$$
c = \sqrt{d_s}\left(\log\left(\frac{\kappa^2}{\kappa^2-\varepsilon^2}\right)-\log\left(\frac{\varepsilon^2}{\kappa^2-\varepsilon^2}\right)\right)
$$
\n
$$
c = \sqrt{d_s}\log\left(\frac{\kappa^2}{\kappa^2-\varepsilon^2}\frac{\kappa^2-\varepsilon^2}{\varepsilon^2}\right)
$$
\n
$$
c = \sqrt{d_s}\log\left(\frac{\kappa^2}{\varepsilon^2-\varepsilon^2}\frac{\kappa^2-\varepsilon^2}{\varepsilon^2}\right)
$$
\n
$$
c = 2\sqrt{d_s}\log\left(\frac{\kappa}{\varepsilon}\right),
$$

which gives us

$$
\alpha(t) = \exp\left(\frac{2\sqrt{d_s}\log\left(\frac{\kappa}{\varepsilon}\right)t + \sqrt{d_s}\log\left(\frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}\right)}{\sqrt{d_s}}\right) - \frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}
$$

$$
= \exp\left(2\log\left(\frac{\kappa}{\varepsilon}\right)t + \log\left(\frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}\right)\right) - \frac{\varepsilon^2}{\kappa^2 - \varepsilon^2}.
$$

C Gaussian marginal likelihood: additional results

In this section we provide an additional study of the new methods on the model in section [4.1.](#page-9-1)

Figure [4a](#page-17-0) shows the log RMSE of the direct estimator for $\varepsilon = 0.01$ as a function of M for ABC, SL and IEnKI-ABC (where $T = 5$) for the two different shifters. The bias is negligible compared to the variance for this estimator, and we observe the usual Monte Carlo inverse scaling of the variance with M . ABC is outperformed by the other approaches, and SL has almost identical performance to IEnKI-ABC with deterministic shifters (the square root shifter and SL results overlap so only one line can be seen). The estimator from sIEnKI-ABC has a similar bias to rIEnKI-ABC and aIEnKI-ABC, but an increased variance due to the noise added in each update step, as also illustrated in the main text.

For the path sampling estimator we observe the need for an increasing number of targets to stabilise the RMSE as ε goes to zero, reflecting the increased distance between the targets at the start and end of the sequence. Figure [4b](#page-17-1) shows how the log estimated RMSE changes with log ε (for $T = 200$ targets), highlighting the poor scaling of this approach as ε goes to 0, in contrast with the direct estimator in figure [1a.](#page-10-1) Figure [4c](#page-17-2) shows how the log RMSE of the path sampling estimator scales with T for $\varepsilon = 0.01$, illustrating the appealing property (compared with the direct estimator in [1b\)](#page-10-2) that an increased T reduces the error, even though this error is significantly larger than the direct estimator. Figure [4d](#page-17-3) shows the estimated log MSE of the path sampling estimator times the computational time required to produce the estimate, highlighting that the additional noise introduced by the stochastic shifter severely affects the strategy of increasing T to reduce the error.

References

- Andrieu, C., Doucet, A., and Holenstein, R. (2010). Particle Markov chain Monte Carlo methods. Journal of the Royal Statistical Society: Series B 72 (3), 269–342.
- Andrieu, C. and Roberts, G. O. (2009). The pseudo-marginal approach for efficient Monte Carlo computations. The Annals of Statistics $37(2)$, 697–725.
- Andrieu, C. and Vihola, M. (2014). Establishing some order amongst exact approximations of MCMCs. $arXiv$, 1–20.

(a) log RMSE as a function of the number of draws M from P .

(b) log RMSE of the path sampling estimator as a function of the log tolerance $\varepsilon.$

(c) log RMSE of the path sampling estimator as a function of the number of targets T. (d) MSE times computational time for the path sampling estimator as a function of T.

Figure 4: Properties of IEnKI normalising constant estimators on the Gaussian example.

- Bardenet, R., Doucet, A., and Holmes, C. (2017). On Markov chain Monte Carlo methods for tall data. Journal of Machine Learning Research, 18, 1–43.
- Beaumont, M. A. (2003). Estimation of population growth or decline in genetically monitored populations. Genetics $164(3)$, 1139–1160.
- Chopin, N., Crucinio, F. R., and Korba, A. (2024, June). A connection between Tempering and Entropic Mirror Descent.
- Cranmer, K., Brehmer, J., and Louppe, G. (2020, December). The frontier of simulation-based inference. Proceedings of the National Academy of Sciences 117 (48), 30055–30062.
- Del Moral, P., Doucet, A., and Jasra, A. (2006). Sequential Monte Carlo samplers. Journal of the Royal Statistical Society: Series B 68(3), 411-436.
- Del Moral, P., Doucet, A., and Jasra, A. (2012). An adaptive sequential Monte Carlo method for approximate Bayesian computation. Statistics and Computing 22 (5), 1009–1020.
- Drovandi, C., Everitt, R., Golightly, A., and Prangle, D. (2019). Ensemble MCMC: Accelerating pseudo-marginal MCMC for state space models using the ensemble Kalman filter.
- Duffield, S. and Singh, S. S. (2022, August). Ensemble Kalman inversion for general likelihoods. Statistics \mathscr{C} Probability Letters, 187, 109523.
- Ebner, B. and Henze, N. (2020, April). Tests for multivariate normality – a critical review with emphasis on weighted \$L^2\$-statistics.
- Evensen, G. (1994). Sequential data assimilation with a nonlinear quasi-geostrophic model using Monte Carlo methods to forecast error statistics. Journal of Geophysical Research: Oceans 99(C5), 10143-10162.
- Farrell, P. J., Salibian-Barrera, M., and Naczk, K. (2007, December). On tests for multivariate normality and associated simulation studies. Journal of Statistical Computation and Simulation 77 (12), 1065–1080.
- Friel, N. and Pettitt, a. N. (2008, July). Marginal likelihood estimation via power posteriors. Journal of the Royal Statistical Society: Series B (Statistical Methodology) 70 (3), 589–607.
- Gelman, A. and Meng, X.-L. (1998, May). Simulating normalizing constants: From importance sampling to bridge sampling to path sampling. *Statistical Science 13*(2), 163–185.
- Ghurye, S. G. and Olkin, I. (1969, August). Unbiased Estimation of Some Multivariate Probability Densities and Related Functions. The Annals of Mathematical Statistics $40(4)$, 1261–1271.
- Gordon, N. J., Salmond, D. J., and Smith, A. F. M. (1993). Novel approach to nonlinear/non-Gaussian Bayesian state estimation. In Radar and Signal Processing, IEE Proceedings F, Volume 140, pp. 107–113. IET.
- Henze, N. and Zirkler, B. (1990, January). A class of invariant consistent tests for multivariate normality. Communications in Statistics - Theory and Methods 19(10), 3595–3617.
- Iglesias, M., Park, M., and Tretyakov, M. V. (2018, July). Bayesian inversion in resin transfer molding. Inverse Problems 34 (10), 105002.
- Iglesias, M. A. (2016, January). A regularizing iterative ensemble Kalman method for PDE-constrained inverse problems. Inverse Problems 32 (2), 025002.
- Katzfuss, M., Stroud, J. R., and Wikle, C. K. (2016, October). Understanding the Ensemble Kalman Filter. The American Statistician $70(4)$, 350–357.
- Kerama, I., Thorne, T., and Everitt, R. G. (2022). Rare event ABC-SMC^2.
- Livings, D. M., Dance, S. L., and Nichols, N. K. (2008, June). Unbiased ensemble square root filters. *Physica D:* Nonlinear Phenomena 237 (8), 1021–1028.
- Mandel, J., Cobb, L., and Beezley, J. D. (2011, December). On the convergence of the ensemble Kalman filter. Applications of Mathematics 56 (6), 533–541.
- Marjoram, P., Molitor, J., Plagnol, V., and Tavare, S. (2003, December). Markov chain Monte Carlo without likelihoods. Proceedings of the National Academy of Sciences of the United States of America 100(26), 15324– 15328.
- Murray, I., Ghahramani, Z., and MacKay, D. J. C. (2006). MCMC for doubly-intractable distributions. In UAI, pp. 359–366.
- Papamakarios, G. and Murray, I. (2016). Fast Epsilon-Free Inference of Simulation Models with Bayesian Conditional Density Estimation. arXiv.
- Prangle, D., Everitt, R. G., and Kypraios, T. (2018). A rare event approach to high dimensional approximate Bayesian computation. Statistics and Computing 28 (4), 819–834.
- Price, L. F., Drovandi, C. C., Lee, A., and Nott, D. J. (2017). Bayesian Synthetic Likelihood. Journal of Computational and Graphical Statistics.
- Roth, M., Hendeby, G., Fritsche, C., and Gustafsson, F. (2017, August). The Ensemble Kalman filter: A signal processing perspective. EURASIP Journal on Advances in Signal Processing 2017 (1), 56.
- Tavaré, S., Balding, D. J., Griffiths, R. C., and Donnelly, P. (1997, February). Inferring Coalescence Times From DNA Sequence Data. Genetics 145 (2), 505–518.
- Tippett, M. K., Anderson, J. L., Bishop, C. H., Hamill, T. M., and Whitaker, J. S. (2003, July). Ensemble Square Root Filters.
- van der Vaart, A. W. (1998). Asymptotic Statistics. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge: Cambridge University Press.
- Whitaker, J. S. and Hamill, T. M. (2002, July). Ensemble Data Assimilation without Perturbed Observations.
- Wilkinson, D. J. (2013). Summary Stats for ABC. https://darrenjw.wordpress.com/ 2013/09/01/summary-statsfor-abc/.
- Wilkinson, D. J. (2024). Smfsb: Stochastic Modelling for Systems Biology.
- Wood, S. N. (2010, August). Statistical inference for noisy nonlinear ecological dynamic systems. Nature 466 (August), 1102–1104.
- Zhang, C. (2022). On the Improvements and Innovations of Monte Carlo Methods. Ph. D. thesis, University of Bristol.
- Zhou, Y., Johansen, A. M., and Aston, J. A. D. (2015). Towards automatic model comparison: An adaptive sequential Monte Carlo approach. Journal of Computational and Graphical Statistics 25 (3), 701–726.