# An Accelerated Multi-level Monte Carlo Approach for Average Reward Reinforcement Learning with General Policy Parametrization

Swetha Ganesh<sup>1,2</sup> and Vaneet Aggarwal<sup>2</sup>

<sup>1</sup>Indian Institute of Science <sup>2</sup>Purdue University

#### Abstract

In our study, we delve into average-reward reinforcement learning with general policy parametrization. Within this domain, current guarantees either fall short with suboptimal guarantees or demand prior knowledge of mixing time. To address these issues, we introduce Randomized Accelerated Natural Actor Critic, a method that integrates Multi-level Monte-Carlo and Natural Actor Critic. Our approach is the first to achieve global convergence rate of  $\tilde{O}(1/\sqrt{T})$  without requiring knowledge of mixing time, significantly surpassing the state-of-the-art bound of  $\tilde{O}(1/T^{1/4})$ .

### 1 Introduction

In reinforcement learning (RL), the temporal dependence of data violates the independent and identically distributed (i.i.d.) assumption common in machine learning, making the theoretical analysis of RL methods challenging. In average-reward RL, stationary behavior under induced policies is fundamentally important to the performance guarantees. Specifically, understanding the effect of mixing time, which measures how long a Markov chain takes to reach stationarity, is critical for developing and analyzing average-reward RL methods. Policy Gradient and Actor Critic based approaches have wide applications due to their scalability and efficiency in handling large and complex reinforcement learning problems. The key goal of this paper is to study efficient algorithms for average reward RL with general parametrized policies.

RL problems are typically analyzed through three primary setups: episodic, infinite horizon discounted reward, and infinite horizon average reward. The infinite horizon average reward setup is particularly significant for real-world applications, including robotics [Gonzalez et al., 2023], transportation [Al-Abbasi et al., 2019], communication networks [Agarwal et al., 2022], and healthcare [Tamboli et al., 2024]. In this setup, the key challenge is that we have a Markovian trajectory, which restricts obtaining independent samples for gradient estimation. One key approach that had been used in the past works for model-free algorithms [Wei et al., 2020, Bai et al., 2024] is based on the observation that if the trajectory is divided into sub-trajectories that are order of mixing time apart, the sub-trajectories become near independent. However, such an approach requires the knowledge of mixing time. We note that model-based tabular Markov Decision Process (MDP) approaches [Jaksch et al., 2010, Agrawal and Jia, 2017, Agarwal and Aggarwal, 2023] do not have that issue, since transition probabilities can be learned from Markovian trajectories, enabling Martingale-based analyses (e.g., employing Azuma-Hoeffding's inequality), while even the algorithms for linear MDP [Wei et al., 2020] still use either the span of the optimal value function or the mixing time. The values of these are not known in practice for a general MDP. Recently, the authors of [Patel et al., 2024] showed that the algorithms that wait for orders of mixing time to have independent sub-trajectories require extremely large time-horizons to even have the sub-linear regret guarantees (even for mixing and hitting times of 10, the time-horizon needed for the guarantees is  $6.6 \times 10^9$ ). In order to alleviate the issue, [Patel et al., 2024] proposed an algorithm, called Multi-level Actor Critic (MAC), where the estimates are obtained directly from the Markovian trajectory rather than obtaining independent samples from subtrajectories, using the Multi-level Monte-Carlo (MLMC) gradient estimator. However, this approach is

Algorithm	Mixing Time	Global	Model-Free?	Policy
	Known?	Convergence		Parametrization
UCRL2 [Jaksch et al., 2010]	-	$\tilde{\mathcal{O}}(1/\sqrt{T})$	No	Tabular
PSRL [Agrawal and Jia, 2017]	-	$\tilde{\mathcal{O}}(1/\sqrt{T})$	No	Tabular
FOPO [Wei et al., 2021]	Yes	$\tilde{\mathcal{O}}(1/\sqrt{T})$	No	Linear
OLSVI.FH [Wei et al., 2021]	Yes	$\tilde{\mathcal{O}}(1/T^{1/4})$	No	Linear
MDP-EXP2 [Wei et al., 2021]	Yes	$\tilde{\mathcal{O}}(1/\sqrt{T})$	No	Linear
MDP-OOMD [Wei et al., 2020]	Yes	$\tilde{\mathcal{O}}(1/\sqrt{T})$	Yes	Tabular
PPGAE [Bai et al., 2024]	Yes	$\tilde{\mathcal{O}}(1/T^{1/4})$	Yes	General
MAC [Patel et al., 2024]	No	$\tilde{\mathcal{O}}(1/T^{1/4})$	Yes	General
PHAPG [Ganesh et al., 2024]	Yes	$\mathcal{O}(1/\sqrt{T})$	Yes	General
RANAC (Algorithm 1)	No	$\tilde{\mathcal{O}}(1/\sqrt{T})$	Yes	General

Table 1: This table summarizes the key related results on global convergence guarantees for average reward reinforcement learning.

shown to achieve a global convergence rate of  $\tilde{\mathcal{O}}(1/T^{1/4})$ , which is not order optimal. This raises the following question, that is addressed in this paper.

Is it possible to achieve a global convergence rate of  $\tilde{O}\left(\frac{1}{\sqrt{T}}\right)$  for a model-free algorithm without the algorithm requiring knowledge of mixing time, in the average-reward setup with general policy parametrization?

In this paper, we answer this question in positive. In order to do that, we propose a novel algorithm, that makes use of the MLMC method for reducing the bias of the Markovian estimators. The proposed algorithm does not use the mixing time, and achieves global convergence which is order-optimal in T. The key comparison of this work to the related works in the literature is summarized in Table 1.

**Novelty and Main Contributions:** The key technical novelty in this paper is to handle the estimation of the policy gradients from the Markovian trajectories for a model-free RL with parametrized policies, and using them efficiently in the Natural Actor Critic based algorithm. We will describe the key differences as compared to three key areas of work in the following.

- Discounted Reward RL: We note that while optimal sample complexity has been studied for the discounted reward setups [Mondal and Aggarwal, 2024] for policy-gradient based approach assuming the availability of a simulator and for actor-critic approaches under Markovian sampling [Xu et al., 2020b, Gaur et al., 2024], we do not need to wait till the mixing time to get independence. The discount factor helps with obtaining decayed impact of the sub-trajectories. Thus, the analysis with average reward setup uses significantly different approaches than those for discounted reward setups. Additionally, we note that our result implies a global sample complexity of order  $\tilde{O}(1/\epsilon^2)$ , which improves over the best known existing bound  $\tilde{O}(1/\epsilon^3)$ , for actor-critic algorithms in both the discounted and average-reward setup [Wang et al., 2024].
- Average Reward RL with Policy Gradients: The key works for average reward setup with parametrized policy are [Bai et al., 2024, Ganesh et al., 2024]. The key in such an approach is that the policy gradient estimation is through the use of sub-trajectories that are far enough so that they are nearly independent. However, such an approach require knowledge of mixing time. In contrast, we directly learn gradient estimation from the Markovian trajectories.
- Average Reward RL with Markovian Gradient Estimator: In order to avoid creating independent subtrajectories, we need to obtain the gradient estimates from Markovian data. Since for a given policy, the estimation approach becomes that of a Markov chain, the authors of [Patel et al., 2024] used the Markovian data algorithm in [Dorfman and Levy, 2022] to obtain the estimate of the policy gradient. In this approach, the gradient is estimated using a multi-level Monte Carlo (MLMC) technique [Giles,

2015] and adaptive learning rate of AdaGrad [Duchi et al., 2011]. We note that employing AdaGrad with MLMC alone fails to ensure global convergence rate of  $\tilde{\mathcal{O}}(1/\sqrt{T})$ , as seen in [Patel et al., 2024], where the convergence rate guarantee is  $\tilde{\mathcal{O}}(1/T^{1/4})$ . This requires multiple novel techniques in the approach as will be described next.

We propose a Randomized Accelerated Natural Actor Critic Algorithm, which employs MLMC to efficiently compute both the critic and the natural policy gradient direction. The term "randomized" refers to the MLMC method's use of a randomized batch length at each iteration for efficient bias reduction. For calculating the NPG direction, we employ a specific variant of Nesterov's momentum, as detailed in [Beznosikov et al., 2023], but modified to eliminate dependence on mixing time.

For the critic update, we demonstrate that TD learning combined with MLMC for bias reduction achieves a convergence rate of  $\tilde{O}(1/T)$ , significantly improving upon the existing guarantees of  $\tilde{O}(1/\sqrt{T})$  provided by previous works [Dorfman and Levy, 2022, Suttle et al., 2023, Patel et al., 2024] using MLMC. Notably, despite utilizing MLMC with Adagrad-based acceleration, these prior analyses obtained suboptimal convergence rates. This is because, while Adagrad can provide optimal-order convergence for non-strongly convex problems, it does not exploit the strong convexity of the TD problem. Our findings indicate that the proposed variant without Adagrad can achieve order-optimal bounds.

Additionally, the Natural Actor Critic framework utilizes a Natural Policy Gradient (NPG) update, in contrast to the vanilla Policy Gradient update studied in [Patel et al., 2024]. It is known that the NPG direction can be computed by solving a stochastic strongly convex optimization problem [Mondal and Aggarwal, 2024]. However, existing algorithms for stochastic strongly convex optimization with Markovian noise either assume knowledge of mixing time [Beznosikov et al., 2023], or require bounded domains/gradients, which necessitate projection operations [Doan et al., 2020, Even, 2023]. In this paper, we introduce an approach that mitigates these issues and utilizes a version of Nesterov's momentum to accelerate the stochastic strongly convex optimization problem, achieving a convergence rate of  $\tilde{O}(1/T)$ .

We summarize the key contributions in this work as follows:

- This paper provides a novel model-free algorithm for average reward reinforcement learning with parametrized policies, which does not use the knowledge of mixing time. The algorithm uses a Natural Actor Critic based approach that utilizes MLMC for estimation from Markovian data and momentum-based acceleration.
- The proposed algorithm is shown to achieve order-optimal global convergence rate of  $\tilde{\mathcal{O}}(1/\sqrt{T})$ , significantly surpassing the state-of-the-art bound of  $\tilde{\mathcal{O}}(1/T^{1/4})$  for such algorithms.
- In order to achieve the result, we propose TD learning with MLMC which achieves convergence rate of  $\tilde{O}(1/T)$ , without requiring knowledge of the mixing time.

### 2 Setup

In this paper, we explore an infinite horizon reinforcement learning problem with an average reward criterion, modeled by a Markov Decision Process (MDP) represented as a tuple  $\mathcal{M} = (S, \mathcal{A}, r, P, \rho)$ . Here, S denotes the state space,  $\mathcal{A}$  is the action space with a size of  $A, r : S \times \mathcal{A} \to [0, 1]$  represents the reward function,  $P : S \times \mathcal{A} \to \Delta^{|S|}$  is the state transition function, where  $\Delta^{|S|}$  denotes the probability simplex with dimension |S|, and  $\rho : S \to [0, 1]$  signifies the initial distribution of states. A policy  $\pi : S \to \Delta^{|\mathcal{A}|}$  determines the distribution of the action to be taken given the current state. It gives rise to a transition function  $P^{\pi} : S \to \Delta^{|S|}$  defined as  $P^{\pi}(s, s') = \sum_{a \in \mathcal{A}} P(s'|s, a)\pi(a|s)$ , for all  $s, s' \in S$ . It can be seen that for any given policy  $\pi$ , the sequence of states produced by the MDP forms a Markov chain. We will be assuming the following throughout the paper:

**Assumption 1.** The MDP  $\mathcal{M}$  is ergodic. That is, the Markov chain induced under every policy  $\pi$ ,  $\{s_t\}_{t\geq 0}$ , is irreducible and aperiodic.

Before proceeding further, we point out that we consider a parameterized class of policies  $\Pi$ , which consists of all policies  $\pi_{\theta}$  such that  $\theta \in \Theta$ , where  $\Theta \subset \mathbb{R}^d$ . It is well-established that if  $\mathcal{M}$  is ergodic, then for all  $\theta \in \Theta$ , there exists a unique stationary distribution denoted as  $d^{\pi_{\theta}} \in \Delta^{|\mathcal{S}|}$ , such that it satisfies  $P^{\pi_{\theta}} d^{\pi_{\theta}} = d^{\pi_{\theta}}$ . With this notation in place, we define the mixing time of an MDP.

**Definition 1.** The mixing time of an MDP  $\mathcal{M}$  with respect to a policy parameter  $\theta$  is defined as,

$$\tau_{\min}^{\theta} \coloneqq \min\left\{t \ge 1 \left| \| (P^{\pi_{\theta}})^t(s, \cdot) - d^{\pi_{\theta}} \| \le \frac{1}{4}, \forall s \in \mathcal{S} \right\}.$$
(1)

We also define  $\tau_{\min} := \sup_{\theta \in \Theta} \tau_{\min}^{\theta}$  as the the overall mixing time. In this paper,  $\tau_{\min}$  is finite due to ergodicity.

The mixing time of an MDP measures how quickly the MDP approaches its stationary distribution when the same policy is executed repeatedly. In the average reward setting, we seek to find a policy  $\pi_{\theta}$  such that the long-term average reward is given by  $J(\pi_{\theta}) := \lim_{T\to\infty} \mathbb{E}\left[\frac{1}{T}\sum_{t=0}^{T} r(s_t, a_t)\right]$ is maximized. In this work, we use an actor-critic approach to optimize *J*. Before proceeding further, we introduce a few important terms. The actionvalue (*Q*) function is defined as

$$Q^{\pi_{\theta}}(s,a) = \mathbb{E}\left[\sum_{t=0}^{\infty} \left[r(s_t, a_t) - J(\pi_{\theta})\right]\right], \quad (2)$$

such that  $s_0 = s, a_0 = a$ , and action  $a \sim \pi_{\theta}$ . We can then further write the state value function as

$$V^{\pi_{\theta}}(s) = \mathbb{E}_{a \sim \pi_{\theta}}(\cdot|s) [Q^{\pi_{\theta}}(s,a)].$$
(3)

Using the Bellman's Equation, we can write, from (2) and (3), the value of a state *s*, in terms of another as [Puterman, 2014]

$$V^{\pi_{\theta}}(s) = \mathbb{E}[r(s,a) - J(\pi_{\theta}) + V^{\pi_{\theta}}(s')], \qquad (4)$$

where the expectation is over  $a \sim \pi_{\theta}(\cdot|s), s' \sim \mathbb{P}(\cdot|a, s)$ . We also define the advantage term as follows,  $A^{\pi_{\theta}}(s, a) \triangleq Q^{\pi_{\theta}}(s, a) - V^{\pi_{\theta}}(s)$ . With the above notations in place, we can now state the well-known policy gradient theorem established by [Sutton et al., 1999]:  $\nabla_{\theta} J(\theta) = \mathbb{E}_{s \sim d^{\pi_{\theta}}, a \sim \pi_{\theta}(\cdot|s)} \left[ A^{\pi_{\theta}}(s, a) \nabla_{\theta} \log \pi_{\theta}(a|s) \right]$ .

PG algorithms maximize the average reward by updating  $\theta$  using gradient ascent, by updating  $\theta$  along  $\nabla_{\theta} J(\theta)$ . Natural Policy Gradient (NPG) methods update  $\theta$  along  $\omega_{\theta}^{*}$  instead, where Algorithm 1 Randomized Accelerated Natural Actor Critic (RANAC)

1: **Input:** Initial parameters  $\theta_0$ ,  $\omega_0 = \omega_0^f$ , and  $\xi_0 = \xi_0^f$ , policy update stepsize  $\alpha$ , parameters for NPG update  $(p_{\omega}, q_{\omega}, \beta_{\omega}, \gamma_{\omega})$ , parameters for critic update  $(p_{\xi}, q_{\xi}, \beta_{\xi}, \gamma_{\xi})$ , initial state  $s_0 \sim \rho(\cdot)$ , outer loop size K, inner loop size H, batch size B2: for  $k = 0, 1, \dots, K$  do 3:  $T_k = 0;$ > Average reward and critic estimation 4: for  $h = 1, 2, \dots, H$  do 5: Sample level length  $P_h^k \sim \text{Geom}(1/2)$ 6: for  $i = 1, ..., 2^{P_h^k} B$  do 7: Take action  $a_k^i \sim \pi_{\theta_k}(\cdot|s_k^i)$ Collect next state  $s_k^{i+1} \sim P(\cdot|s_k^i, a_k^i)$ 8: 9: Receive reward  $r_k^i = r(s_k^i, a_k^i)$ 10: 11: end for  $T_k + = 2^{P_h^k} B$ 12: Update  $\xi_h$  using (16) and (17) 13: end for 14: > Natural Policy Gradient (NPG) estimation 15: 16: for  $h = 1, 2, \dots, H$  do Sample level length  $Q_h^k \sim \text{Geom}(1/2)$ 17: for  $i = 1, ..., 2^{Q_h^k} B$  do 18:  $\begin{array}{l} \text{Take action } a_k^i \sim \pi_{\theta_k}(\cdot|s_k^i) \\ \text{Collect next state } s_k^{i+1} \sim P(\cdot|s_k^i,a_k^i) \end{array}$ 19: 20: Receive reward  $r_k^i = r(s_k^i, a_k^i)$ 21: end for 22:  $T_k + = 2^{Q_h^k} B$ 23: Update  $\omega_h$  using (14) and (15) 24: 25: end for ▷ Policy update 26: 27:  $v_k = \omega_H$  $\theta_{k+1} = \theta_k - \alpha v_k$ 28: 29: end for

$$\omega_{\theta}^* = F(\theta)^{-1} \nabla_{\theta} J(\theta),$$

and  $F(\theta)$  is the Fisher information matrix as defined as:

$$F(\theta) = \mathbb{E}_{s \sim d^{\pi_{\theta}}} \mathbb{E}_{a \sim \pi_{\theta}(\cdot|s)} \left[ \nabla_{\theta} \log \pi_{\theta}(a|s) (\nabla_{\theta} \log \pi_{\theta}(a|s))^{\top} \right].$$
(5)

### **3** Proposed Algorithm

Algorithm 2 Randomized Accelerated Gradient Descent [Beznosikov et al., 2023]

1: **Parameters:** stepsize  $\gamma > 0$ , momentums  $m, p, q, \beta$ , number of iterations T, MLMC limit  $T_{\text{max}}$  and batch size B2: **Initialization:** Set  $x_0^f = x_0, T_0 = 0$ 3: **for**  $t = 0, 1, 2, \dots, T - 1$  **do** 4:  $x_t^g = mx_t^f + (1 - m)x_t$ 5: Sample  $J_t \sim \text{Geom}(1/2)$ 6:  $g_t = g_t^0 + \begin{cases} 2^{J_t} (g_t^{J_t} - g_t^{J_t-1}), & \text{if } 2^{J_t} \leq T_{\text{max}} \\ 0, & \text{otherwise} \end{cases}$  with  $g_t^j = 2^{-j}B^{-1}\sum_{i=1}^{2^j B} \nabla F(x_t^g, Z_{T_t+i})$ 7:  $x_{t+1}^f = x_t^g - p\gamma g_t$ 8:  $x_{t+1} = qx_{t+1}^f + (p - q)x_t^f + (1 - p)(1 - \beta)x_t + (1 - p)\beta x_t^g$ 9:  $T_{t+1} = T_t + 2^{J_t}B$ 10: **end for** 

We propose Randomized Accelerated Natural Actor Critic (Algorithm 1). The algorithm is comprised of the following key components:

**Natural Policy Gradient (NPG):** For a fixed policy parameter  $\theta \in \Theta$ , we compute the NPG direction by viewing it as a stochastic optimization problem with Markovian noise. The NPG direction is the solution to the following strongly convex optimization problem:

$$\min f(\omega) \coloneqq (1/2) \cdot \omega^{\top} F(\theta) \omega - \nabla_{\theta} J(\theta) \omega.$$
(6)

Note that  $\nabla_{\omega} f(\omega) = F(\theta)\omega - \nabla_{\theta} J(\theta)$ . Consider  $\theta_k \in \Theta$  and a trajectory induced by policy  $\pi_{\theta_k}$ ,  $\mathcal{T}^i_k = (s^0_k, a^0_k, s^1_k, a^1_k \cdots, s^i_k, a^i_k)$ . Let  $z^j_k = (s^j_k, a^j_k, s^{j+1}_k)$  and  $d_Z$  denote the distribution determined by  $d_Z(z^j_k) = d^{\pi_{\theta_k}}(s^j_k)\pi(a^j_k|s^j_k)P(s^{j+1}_k|s^j_k, a^j_k)$ . Using these notations, we can write the estimate of as  $\nabla_{\omega} f(\omega)$  as

$$u(\omega, \theta_k; z_k^j) \coloneqq F(\theta_k; z_k^j) \omega - h(\theta_k; z_k^j), \tag{7}$$

where  $F(\theta_k; z_k^j) = \nabla_{\theta} \log \pi_{\theta_k}(a_k^j | s_k^j) \nabla_{\theta} \log \pi_{\theta}(a_k^j | s_k^j)^{\top}$ , and  $h(\theta_k; z_k^j)$  is the vanilla policy gradient estimator, which we provide in the next section in (11).

**Average-reward and critic estimation:** We consider the critic setup where the critic function is the inner product between a given feature map  $\phi(s) : S \to \mathbb{R}^m$  and a weight vector  $\nu \in \mathbb{R}^m$ . We denote the critic estimation for  $V^{\pi_{\theta}}(s)$  as  $V_{\nu}(s) \coloneqq \langle \phi(s), \nu \rangle$  and assume that  $\|\phi(s)\| \leq 1$  for all  $s \in S$ . The critic aims to minimize the error below

$$\min_{\nu \in \Omega} \sum_{s \in S} d^{\pi_{\theta}}(s) (V^{\pi_{\theta}}(s) - V_{\nu}(s))^{2}.$$
(8)

Consider  $\theta_k \in \Theta$  and a trajectory induced by policy  $\pi_{\theta_k}$ ,  $\mathcal{T}_k^i = (s_k^0, a_k^0, s_k^1, a_k^1, \cdots, s_k^i, a_k^i)$ . Let  $z_k^j = (s_k^j, a_k^j, s_k^{j+1})$ . The stochastic gradient estimate of (8) is typically approximated using the Temporal Difference (TD) method as

$$\left(r(s_k^j, a_k^j) - J(\pi_{\theta_k}) + \langle \phi(s_k^{j+1}), \nu_k \rangle - \langle \phi(s_k^j), \nu_k \rangle \right) \phi(s_k^j).$$

$$\tag{9}$$

Note that due to the use of bootstrapping, the above estimator is not a "true" gradient estimator [Maei, 2011], which requires a different analysis than regular stochastic optimization methods. Since the critic update in (9) relies on  $J(\pi_{\theta_t})$ , which we do not have access to, we can substitute with a recursive estimate for the average reward as  $\eta_{k+1} = \eta_k - \gamma_k(\eta_k - r(s_k^j, a_k^j))$ . The critic and average reward estimates are typically updated as:

$$\eta_{k+1} = \eta_k - c_\beta \beta_k (\eta_k - r(s_k^j, a_k^j)) \\ \nu_{k+1} = \nu_k - \beta_k (\phi(s_k^j)(\phi(s_k^{j+1}) - \phi(s_k^j))^\top \nu_k + (r(s_k^j, a_k^j) - \eta_k) \phi(s_k^j)),$$
(10)

where  $c_{\beta} > 0$  is some constant. With these estimates, we can estimate the vanilla policy gradient direction as

$$h(\theta_k; z_k^j) = (r(s_k^j, a_k^j) - \eta_k + \langle \phi(s_k^{j+1}) - \phi(s_k^j), \nu_k \rangle) \cdot \nabla_\theta \log \pi_{\theta_k}(a_k^j | s_k^j).$$

$$\tag{11}$$

We can combine the average-reward and critic estimates as a single vector  $\xi_k = [\eta_k, \nu_k^{\top}]^{\top}$  as in [Zhang et al., 2021b]. Then, the updates in (10) can be compactly expressed as

$$\xi_{k+1} = \xi_k - \beta_k v(\theta_k, \xi_k; z_k^{\mathcal{I}}), \tag{12}$$

where

We now state some assumptions we will be making regarding the critic updates.

**Assumption 2.** For a given feature mapping  $\phi$ , we define the worst-case approximation error to be

$$\epsilon_{\rm app} = \sup_{\theta} \mathbb{E}_{s \sim d^{\pi_{\theta}}} \left[ \phi(s)^T \nu^*(\theta) - V^{\pi_{\theta}}(s) \right]^2, \tag{13}$$

which we assume to be finite.

**Assumption 3.** There exist  $\lambda > 0$  such that, for all  $\theta$ , the matrix  $\mathbb{E}[\phi(s_k^j)(\phi(s_k^j) - \phi(s_k^{j+1}))^{\top}] - \lambda I$  is positive definite, i.e. for all  $x, x^{\top} \mathbb{E}[\phi(s_k^j)(\phi(s_k^j) - \phi(s_k^{j+1}))^{\top}] x \ge \lambda \cdot ||x||^2$ .

Both Assumptions 2 and 3 are frequently employed in the analysis of actor-critic methods [Bhandari et al., 2018, Zou et al., 2019, Qiu et al., 2021, Xu et al., 2020a, Suttle et al., 2023]. Assumption 2 intuitively relates to the quality of the feature mapping, where  $\epsilon_{app}$  measures this quality: well-designed features lead to a small or even zero  $\epsilon_{app}$ , whereas poorly designed features result in a higher worst-case error. Assumption 3, on the other hand, is essential for guaranteeing the solvability of the MSPBE minimization problem and the uniqueness of its solutions.

Let  $A_{\theta} = \mathbb{E}_{d_Z} A(\theta; z)$  and  $b_{\theta} = \mathbb{E}_{d_Z} b(\theta; z)$ . By setting large enough  $c_{\beta}$ , Assumption 3 implies that  $A_{\theta} - (\lambda/2)I$  is also positive definite (see Lemma 6). This also implies that  $A_{\theta}$  is invertible. For a fixed policy parameter  $\theta$ , let  $\xi^*$  denote  $A_{\theta}^{-1}b_{\theta}$ .

**Multi-level Monte Carlo**: Let  $\mathcal{T}_k^j := (s_k^0, a_k^0, s_k^1, a_k^1 \cdots, s_k^j, a_k^j)$  denote a trajectory generated by policy  $\pi_{\theta_k}$  with length j and  $z_k^i = (s_k^i, a_k^i, s_k^{i+1})$ . Sample  $Q_h^k \sim \text{Geom}(1/2)$ . Then the MLMC estimator of the gradient in the NPG subproblem at iteration

*h* can be calculated as:

$$\tilde{u}_{h} = u_{h}^{0} + \begin{cases} 2^{Q_{h}^{k}} (u_{h}^{Q_{h}^{k}} - u_{h}^{Q_{h}^{k}-1}), & \text{if } 2^{Q_{h}^{k}} \le T_{\max} \\ 0, & \text{otherwise} \end{cases}$$
(14)

with  $u_h^j = \frac{1}{2^j} \sum_{k=1}^{2^j} u(\omega_h^g, \theta_k; z_h^k)$  and where  $T_{\max} \ge 2$ . Using the MLMC estimator, the NPG direction  $\omega_h$  is updated using momentum-based acceleration as:

$$\omega_{h}^{g} = m_{\omega}\omega_{h}^{f} + (1 - m_{\omega})\omega_{h}$$

$$\omega_{h+1}^{f} = \omega_{h}^{g} - p_{\omega}\gamma_{\omega}\tilde{u}_{h}$$

$$\omega_{h+1} = q_{\omega}\omega_{h+1}^{f} + (p_{\omega} - q_{\omega})\omega_{h}^{f} + (1 - p_{\omega})(1 - \beta_{\omega})\omega_{h} + (1 - p_{\omega})\beta_{\omega}\omega_{h}^{g}$$
(15)

Similarly, for the critic updates, sample  $P_h^k \sim \text{Geom}(1/2)$ . Then the MLMC estimator of the critic update can be calculated as:

$$\tilde{v}_{h} = v_{h}^{0} + \begin{cases} 2^{Q_{h}^{k}} (v_{h}^{P_{h}^{k}} - v_{h}^{P_{h}^{k}-1}), & \text{if } 2^{P_{h}^{k}} \le T_{\max} \\ 0, & \text{otherwise} \end{cases}$$
(16)

with  $v_h^j = \frac{1}{2^j} \sum_{k=1}^{2^j} v(\xi_h, \theta_k; z_h^k)$  and where  $T_{\max} \ge 2$ . Using the MLMC estimator, the critic is updated as:

$$\xi_{h+1} = \xi_h - \beta_h \tilde{v}_h. \tag{17}$$

The advantage of the MLMC estimator is that it achieves the same bias as averaging  $T_{\text{max}}$  gradients, but with only O(1) samples. Additionally, since drawing from a geometric distribution does not require knowledge of the mixing time, we can eliminate the need for the oracle knowledge assumption used in previous works like [Bai et al., 2024, Ganesh et al., 2024].

However, the MLMC estimator can result in increased variance of the estimator. There are a line of works using AdaGrad to reduce the impact of the increased variance and accelerate convergence. However, this approach does not provide improved convergence rate of O(1/T) when the objective is strongly-convex. Thus, we use a version of Nesterov's acceleration considered with the above mentioned MLMC estimator. This version of Accelerated MLMC was studied in [Beznosikov et al., 2023], however, their choice of algorithm parameters required knowledge of mixing time. Moreover, our gradient estimates also suffer from bias due to the average-reward and critic estimation error, whereas the result in [Beznosikov et al., 2023] requires unbiased estimates. Since the critic update is not a pure gradient descent approach due to the use of bootstrapping, the result obtained for the strongly convex problem cannot be directly used. As a result, we use MLMC (without momentum) with TD learning.

#### 4 Main results

We first state some assumptions that we will be using before proceeding to the main results.

**Assumption 4.** For any  $\theta \in \mathbb{R}^d$ , the transferred compatible function approximation error,  $L_{\nu^\star}(\omega^{\sharp}_{\star};\theta)$ , satisfies

$$L_{d^{\pi^{\star}}}(\omega_{\star}^{\theta};\theta) \coloneqq \mathbb{E}_{s \sim d^{\pi^{\star}}} \mathbb{E}_{a \sim \pi^{\star}(\cdot|s)} \left[ \left( A^{\pi_{\theta}}(s,a) - (\omega_{\star}^{\theta})^{\top} \nabla_{\theta} \log \pi_{\theta}(a|s) \right)^{2} \right] \le \epsilon_{\text{bias}}, \tag{18}$$

where  $\omega_{+}^{\theta}$  is the exact NPG direction at  $\theta$ .

**Assumption 5.** For all  $\theta$ ,  $\theta_1$ ,  $\theta_2 \in \Theta$  and  $(s, a) \in S \times A$ , the following statements hold:

(a) 
$$\|\nabla_{\theta} \log \pi_{\theta}(a|s)\| \le G_1$$
 (b)  $\|\nabla_{\theta} \log \pi_{\theta}(a|s) - \nabla_{\theta} \log \pi_{\theta_2}(a|s)\| \le G_2 \|\theta_1 - \theta_2\|.$  (19)

**Assumption 6** (Fisher non-degenerate policy). There exists a constant  $\mu > 0$  such that  $F(\theta) - \mu I_d$  is positive semidefinite where  $I_d$  denotes an identity matrix.

Comments on Assumptions 4-6: We would like to highlight that all these assumptions are commonly found in PG literature [Liu et al., 2020, Agarwal et al., 2021, Papini et al., 2018, Xu et al., 2019, Fatkhullin et al., 2023]. We elaborate more on these assumptions below.

 $\epsilon_{\text{bias}}$  captures the parametrization capacity of  $\pi_{\theta}$ . For  $\pi_{\theta}$  using the softmax parametrization, we have  $\epsilon_{\text{bias}} = 0$  [Agarwal et al., 2021]. When  $\pi_{\theta}$  is a restricted parametrization, which may not contain all stochastic policies, we have  $\epsilon_{\text{bias}} > 0$ . It is known that  $\epsilon_{\text{bias}}$  is very small when rich neural parametrizations are used [Wang et al., 2019]. Assumption 5 requires that the score function is bounded and Lipschitz continuous. This assumption is widely used in the analysis of PG based methods [Liu et al., 2020, Agarwal et al., 2021, Papini et al., 2018, Xu et al., 2019, Fatkhullin et al., 2023]. Assumption 6 requires that the eigenvalues of the Fisher information matrix can be bounded from below and is commonly used in obtaining global complexity bounds for PG based methods [Liu et al., 2020, Zhang et al., 2021a, Bai et al., 2022, Fatkhullin et al., 2023]. Assumptions 5-6 were shown to hold for various examples recently including Gaussian policies with linearly parameterized means and certain neural parametrizations [Liu et al., 2020, Fatkhullin et al., 2023].

**Theorem 1.** Consider Algorithm 1 with  $K = \Theta(\sqrt{T})$ ,  $H = \Theta(\sqrt{T}/\log(T))$ ,  $B = \Theta(\log(T))$  and  $\alpha = \frac{\mu^2}{4G_1^2 L}$ . Let Assumptions 1-6 hold and assume J is L-smooth and  $\|\nabla J(\theta)\|$ ,  $h(\theta; z) \leq C$  for all  $\theta \in \Theta$  and  $z \in S \times A \times S$ . Then there exists a choice of parameters for the critic and NPG estimation such that for sufficiently large T, the following statement holds:

$$\frac{1}{K}\sum_{k=0}^{K-1} (J^* - \mathbb{E}[J(\theta_k)]) \le \mathcal{O}\left(\sqrt{\epsilon_{\mathrm{app}}} + \sqrt{\epsilon_{\mathrm{bias}}} + \frac{\tau_{\mathrm{mix}}^3(\log T)^2 + \tau_{\mathrm{mix}}^4(\log T)}{\sqrt{T}}\right).$$
(20)

**Remark 1.** A detailed version of this result can be found in Appendix D. We point out that the above bound is only  $\tilde{\mathcal{O}}(1/\sqrt{T})$ , in contrast to existing bounds of order  $\tilde{\mathcal{O}}(1/T^{1/4})$  in the average reward general policy setting without knowledge of mixing time [Bai et al., 2024, Patel et al., 2024]. Moreover, our bounds do not depend on the size of the action space and hitting time unlike [Bai et al., 2024, Ganesh et al., 2024]. Though [Patel et al., 2024] provides bounds with only  $\mathcal{O}(\sqrt{\tau_{mix}})$  dependence, these bounds depend on the projection radius of the critic updates,  $R_{\omega}$ , which can be large and scale with  $\tau_{mix}$  [Wei et al., 2020]. In contrast, our algorithm does not use such projection operators and do not scale with  $R_{\omega}$ .

### 5 **Proof Outline**

We structure our analysis into three parts: policy update, NPG estimation, and critic analysis.

### 5.1 Policy update analysis

Lemma 1. Consider any policy update rule of form

$$\theta_{k+1} = \theta_k + \alpha \omega_k. \tag{21}$$

If Assumptions 4 and 5 hold, then the following inequality is satisfied

$$J^{*} - \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E}[J(\theta_{k})] \leq \sqrt{\epsilon_{\text{bias}}} + \frac{G_{1}}{K} \sum_{k=0}^{K-1} \mathbb{E} \left\| \left( \mathbb{E} \left[ \omega_{k} | \theta_{k} \right] - \omega_{k}^{*} \right) \right\| \\ + \frac{\alpha G_{2}}{2K} \sum_{k=0}^{K-1} \mathbb{E} \left\| \omega_{k} \right\|^{2} + \frac{1}{\alpha K} \mathbb{E}_{s \sim d^{\pi^{*}}} \left[ KL(\pi^{*}(\cdot|s) \| \pi_{\theta_{0}}(\cdot|s)) \right],$$

$$(22)$$

where  $KL(\cdot \| \cdot)$  is the Kullback-Leibler divergence,  $\omega_k^*$  is the NPG direction  $F(\theta_k)^{-1} \nabla J(\theta_k)$ ,  $\pi^*$  is the optimal policy and  $J^*$  is the optimal value of the function  $J(\cdot)$ .

The last term above is of order  $\mathcal{O}(1/K)$  since  $\mathbb{E}_{s \sim d^{\pi^*}}[KL(\pi^*(\cdot|s) \| \pi_{\theta_0}(\cdot|s))]$  is constant. The term  $\mathbb{E} \| \omega_k \|^2$  can be further decomposed as

$$\frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\omega_k\|^2 \leq \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\omega_k - \omega_k^*\|^2 + \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\omega_k^*\|^2$$

$$\stackrel{(a)}{\leq} \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\omega_k - \omega_k^*\|^2 + \frac{\mu^{-2}}{K} \sum_{k=0}^{K-1} \mathbb{E} \|\nabla_\theta J(\theta_k)\|^2$$
(23)

where (*a*) follows from Assumption 6 and the definition that  $\omega_k^* = F(\theta_k)^{-1} \nabla_{\theta} J(\theta_k)$ . Thus, we can obtain a global convergence bound by bounding the terms  $\mathbb{E} \|\omega_k - \omega_k^*\|^2$ ,  $\mathbb{E} \|(\mathbb{E} [\omega_k | \theta_k] - \omega_k^*)\|$  and  $\mathbb{E} \|\nabla_{\theta} J(\theta_k)\|^2$ . The first two terms are the variance and bias of the NPG estimator,  $\omega_k$ , and the third term indicates the local convergence rate. Further,  $\mathbb{E} \|\nabla_{\theta} J(\theta_k)\|^2$  can be expressed in terms of  $\mathbb{E} \|\omega_k - \omega_k^*\|^2$ , and thus, in the next subsections we briefly describe how we obtain these bounds.

#### 5.2 NPG estimation analysis

In this section, we focus on the NPG estimation loop for a fixed  $\theta_k \in \Theta$ . We drop the subscript k for ease of exposition. We look at how the bounds are derived for  $\mathbb{E} \|\omega_H - \omega^*\|^2$  and  $\mathbb{E} \|(\mathbb{E} [\omega_H | \theta] - \omega^*)\|$ . Recall that Algorithm 1 computes  $\omega_H$  using Accelerated MLMC for H iterations.

Before we provide results for the NPG analysis, we first provide a result for Algorithm 2 for strongly convex optimization with Markovian noise that does not require knowledge of mixing time or bounded gradient/domain/projection. Consider a time-homogeneous, ergodic Markov chain  $(Z_i)_{i\geq 0}$  with with a unique invariant distribution  $d_Z$  and Markov kernel Q. Further, let Q be uniformly geometrically ergodic with mixing time  $\tau_{\text{mix}}$  as in Assumption 1. Let F denote the strongly convex function we wish to optimize and  $\nabla F(x, Z_i)$  denote the stochastic gradient estimate. Further, let F be L-smooth and  $\mu$ -strongly convex with  $x^* = \arg \min_{x \in \mathbb{R}^d} F(x)$ . An approach to optimize for F under this setup was provided in [Beznosikov et al., 2023] and provided in Algorithm 2. However, in [Beznosikov et al., 2023], the parameter choices for the algorithm requires knowledge of mixing time and does not consider biased estimates. Below, we provide a result which does not require mixing time for parameter choices and accounts for biased estimates.

**Theorem 2.** Consider Algorithm 2 with  $\|\mathbb{E}_{d_Z}[\nabla F(x,Z)] - \nabla F(x)\|^2 \leq \Delta$  and  $\|\nabla F(x,Z) - \nabla F(x)\|^2 \leq \sigma^2 + \delta^2 \|\nabla F(x)\|^2$ . Then for  $\gamma, \beta, m, q, p, T_{\max}, B$  satisfying

$$\begin{split} p &\simeq \sqrt{\frac{L}{\mu}} \cdot \frac{\log T}{T}, \quad \beta \simeq \sqrt{p^2 \mu \gamma}, \quad q \simeq \sqrt{\frac{1}{\mu \gamma}}, \quad \gamma = \frac{3}{4L} \\ m &\simeq \frac{1 - p \eta^{-1}}{1 - \beta p q^{-1}}, \quad T_{\max} \simeq \sqrt{T}, \quad B = \lceil \log T_{\max} \rceil, \end{split}$$

*it holds for*  $T \geq \sqrt{\frac{L}{\mu}} \cdot \tau_{\min}$  *that* 

$$\mathbb{E}\left[\|x_T - x^*\|^2 + \frac{6}{\mu}(F(x_T^f) - F(x^*))\right] \le \mathcal{O}\left(\frac{1}{T^2}\left[\|x_0 - x^*\|^2 + \frac{6}{\mu}(F(x_0) - F(x^*))\right] + \frac{\log T}{\mu T}\left(\sigma^2 \tau_{\min}^2\right) + \Delta\right).$$
(24)

Now, for the NPG analysis, we have  $f(\omega)$  is  $L_{\omega}$ -smooth and  $\mu$ -strongly convex where  $L_{\omega}$  and  $\mu$  are the largest and smallest eigenvalues of  $F(\theta)$ , respectively. From Assumption 6, it follows that  $\mu > 0$ . We also show that  $\|\mathbb{E}_{d_Z}[u(\omega, \theta; z)] - \nabla_{\omega} f(\omega)\|^2 \leq \mathcal{O}(\|\xi_H - \xi^*\|^2 + G_2^2 \epsilon_{app})$  and  $\|u(\omega, \theta; z) - \nabla_{\omega} f(\omega)\|^2 \leq \sigma_{\omega}^2 + \delta_{\omega}^2 \|\nabla_{\omega} f(\omega)\|^2$ , for some  $\delta_{\omega}$  and  $\sigma_{\omega}$  (see Appendix B for details). As a result, Theorem 2 can be invoked to provide the following bounds.

**Theorem 3.** Consider Algorithm 1 with  $\gamma_{\omega}$ ,  $\beta_{\omega}$ ,  $m_{\omega}$ ,  $q_{\omega}$ ,  $p_{\omega}$ ,  $T_{\max}$ , B satisfying

$$p_{\omega} \simeq \sqrt{\frac{L}{\mu}} \cdot \frac{\log H}{H}, \quad \beta_{\omega} \simeq \sqrt{p_{\omega}^2 \mu \gamma}, \quad q_{\omega} \simeq \sqrt{\frac{1}{\mu \gamma_{\omega}}}, \quad \gamma_{\omega} = \frac{3}{4L}, \quad m_{\omega} \simeq \frac{1 - p_{\omega} q_{\omega}^{-1}}{1 - \beta_{\omega} p_{\omega} q_{\omega}^{-1}}, \quad B = \lceil \log T_{\max} \rceil.$$

When all assumptions in Theorem 1 hold, then it holds for sufficiently large H and  $T_{max}$  that

$$\mathbb{E}\left[\|\omega_{H} - \omega^{*}\|^{2}\right] \leq \mathcal{O}\left(\frac{1}{H^{2}}\left[\|\omega_{0} - \omega^{*}\|^{2}\right] + \frac{\log H}{\mu^{2}H}\left(\sigma_{\omega}^{2}\tau_{\mathrm{mix}}^{2}\right) + G_{1}^{2}\mathbb{E}\left\|\xi_{H} - \xi^{*}\right\|^{2} + G_{1}^{2}\epsilon_{\mathrm{app}}\right)$$
(25)

and

$$\mathbb{E}\left[\|\mathbb{E}[\omega_{H}|\theta] - \omega^{*}\|^{2}\right] \leq \mathcal{O}\left(\frac{1}{H^{2}}[\|\omega_{0} - \omega^{*}\|^{2}] + \frac{G_{1}^{2}\tau_{\text{mix}}^{4}}{T_{\text{max}}^{2}} + G_{1}^{2}\|\mathbb{E}[\xi_{H}|\theta] - \xi^{*}\|^{2} + G_{1}^{2}\epsilon_{\text{app}}\right).$$
(26)

#### 5.3 Critic analysis

As discussed in Section 3, the average-reward and critic update can be viewed as a linear function. However, the matrix A is asymmetric and this update cannot be viewed as a gradient of a quadratic function. Moreover, its eigenvalues may not be real. However, it is still satisfies useful properties, such as  $(1/2)\xi^{\top}A\xi > \lambda/2 \cdot ||\xi||^2$ , for all  $\xi$ . We can still utilize this positive-definiteness property to obtain our results. Using the fact that the critic update estimates satisfy  $||A(\theta; z)|| \le L_A$  and  $||b(\theta; z)|| \le L_b$ , for all  $\theta$  and z (see Section C), we obtain the following bounds: **Theorem 4.** Consider Algorithm 1 and let Assumptions 1-6 hold. Then for  $\gamma_{\xi}$ ,  $\beta_{\xi}$ ,  $m_{\xi}$ ,  $q_{\xi}$ ,  $p_{\xi}$ ,  $T_{\max}$ , B satisfying

$$p_{\xi} \simeq \sqrt{\frac{L_A}{\lambda}} \cdot \frac{\log H}{H}, \quad \beta_{\xi} \simeq \sqrt{p_{\xi}^2 \gamma_{\xi} \lambda}, \quad q_{\xi} \simeq \sqrt{\frac{1}{\lambda \gamma_{\xi}}}, \quad \gamma_{\xi} = \frac{3}{4L_A}, \quad m_{\xi} \simeq \frac{1 - p_{\xi} q_{\xi}^{-1}}{1 - \beta_{\xi} p_{\xi} q_{\xi}^{-1}}, \quad B = \lceil \log T_{\max} \rceil$$

*it holds for sufficiently large* H *and*  $T_{max}$  *that* 

$$\mathbb{E}\left[\|\xi_{H} - \xi^{*}\|^{2}\right] \leq \mathcal{O}\left(\frac{1}{H^{2}} \mathbb{E}\left\|\xi_{0} - \xi^{*}\right\|^{2} + \frac{\log H}{H}(\lambda^{-3}L_{A}^{2} + L_{b}^{2}\lambda^{-1})(\tau_{\min}\log T_{\max} + \tau_{\min}^{2}) + \frac{L_{b}^{2}\tau_{\max}^{2}}{\lambda^{2}T_{\max}^{2}}\right)$$
(27)

and

$$\mathbb{E}\left[\|\mathbb{E}[\xi_{H}|\theta] - \xi^{*}\|^{2}\right] \leq \mathcal{O}\left(\frac{1}{H^{2}}\|\mathbb{E}[\xi_{0}|\theta] - \xi^{*}\|^{2} + \frac{L_{A}^{4}\tau_{\min}^{2}}{\lambda^{4}T_{\max}^{2}} + \frac{L_{b}^{2}\tau_{\min}^{2}}{\lambda^{2}T_{\max}^{2}}\right).$$
(28)

**Remark 2.** Note that the critic error  $\mathbb{E} \left[ \|\xi_T - \xi^*\|^2 \right]$  is  $\tilde{O}(1/T)$ , which is significantly improves upon existing guarantees of  $\tilde{O}(1/\sqrt{T})$  for TD with MLMC [Dorfman and Levy, 2022, Suttle et al., 2023, Patel et al., 2024]. Though these works use TD learning with MLMC and AdaGrad, this approach does not exploit the strong convexity of the underlying problem and results in suboptimal convergence rates.

### 6 Conclusions

In this work, we introduce the Randomized Accelerated Natural Actor Critic (Algorithm 1), which achieves a global convergence rate of order  $\tilde{O}(1/\sqrt{T})$  without requiring prior knowledge of the mixing time. Additionally, the algorithm employs an MLMC TD method for the critic update, ensuring convergence of order  $\tilde{O}(1/T)$ . This paper also presents the first result for stochastic strongly-convex optimization with Markovian noise that does not require knowledge of the mixing time, avoids projections, and does not assume a bounded domain or gradient, yet achieves a convergence rate of  $\tilde{O}(1/T)$ . Extension of the proposed approach to general critic functions is an important future direction, while we note that the corresponding results for the discounted reward setup leads to an increased sample complexity [Gaur et al., 2024].

#### References

- Alekh Agarwal, Sham M Kakade, Jason D Lee, and Gaurav Mahajan. On the theory of policy gradient methods: Optimality, approximation, and distribution shift. *The Journal of Machine Learning Research*, 22 (1):4431–4506, 2021.
- Mridul Agarwal and Vaneet Aggarwal. Reinforcement learning for joint optimization of multiple rewards. *Journal of Machine Learning Research*, 24(49):1–41, 2023.
- Mridul Agarwal, Qinbo Bai, and Vaneet Aggarwal. Concave utility reinforcement learning with zeroconstraint violations. *Transactions on Machine Learning Research*, 2022.
- Shipra Agrawal and Randy Jia. Optimistic posterior sampling for reinforcement learning: worst-case regret bounds. *Advances in Neural Information Processing Systems*, 30, 2017.
- Abubakr O Al-Abbasi, Arnob Ghosh, and Vaneet Aggarwal. Deeppool: Distributed model-free algorithm for ride-sharing using deep reinforcement learning. *IEEE Transactions on Intelligent Transportation Systems*, 20(12):4714–4727, 2019.
- Qinbo Bai, Amrit Singh Bedi, Mridul Agarwal, Alec Koppel, and Vaneet Aggarwal. Achieving zero constraint violation for constrained reinforcement learning via primal-dual approach. *Proceedings of the AAAI Conference on Artificial Intelligence*, 36:3682–3689, Jun. 2022. doi: 10.1609/aaai.v36i4.20281.
- Qinbo Bai, Washim Uddin Mondal, and Vaneet Aggarwal. Regret analysis of policy gradient algorithm for infinite horizon average reward markov decision processes. In *AAAI Conference on Artificial Intelligence*, 2024.

- Aleksandr Beznosikov, Sergey Samsonov, Marina Sheshukova, Alexander Gasnikov, Alexey Naumov, and Eric Moulines. First order methods with markovian noise: from acceleration to variational inequalities. In *Thirty-seventh Conference on Neural Information Processing Systems*, 2023.
- Jalaj Bhandari, Daniel Russo, and Raghav Singal. A finite time analysis of temporal difference learning with linear function approximation. *CoRR*, abs/1806.02450, 2018. URL http://arxiv.org/abs/1806.02450.
- Thinh T. Doan, Lam M. Nguyen, Nhan H. Pham, and Justin Romberg. Convergence rates of accelerated markov gradient descent with applications in reinforcement learning, 2020.
- Ron Dorfman and Kfir Yehuda Levy. Adapting to mixing time in stochastic optimization with Markovian data. In *Proceedings of the 39th International Conference on Machine Learning*, Jul 2022.
- Randal Douc, Eric Moulines, Pierre Priouret, and Philippe Soulier. *Markov chains*. Operation research and financial engineering. Springer, 2018. URL https://hal.science/hal-02022651.
- John Duchi, Elad Hazan, and Yoram Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of machine learning research*, 12(7), 2011.
- Mathieu Even. Stochastic gradient descent under markovian sampling schemes, 2023.
- Ilyas Fatkhullin, Anas Barakat, Anastasia Kireeva, and Niao He. Stochastic policy gradient methods: Improved sample complexity for fisher-non-degenerate policies. In *International Conference on Machine Learning*, pages 9827–9869. PMLR, 2023.
- Swetha Ganesh, Washim Uddin Mondal, and Vaneet Aggarwal. Variance-reduced policy gradient approaches for infinite horizon average reward markov decision processes, 2024. URL https://arxiv.org/abs/2404.02108.
- Mudit Gaur, Vaneet Aggarwal, Amrit Singh Bedi, and Di Wang. Closing the gap: Achieving global convergence (last iterate) of actor-critic under markovian sampling with neural network parametrization. In *International Conference on Machine Learning*, 2024.
- Michael B Giles. Multilevel monte carlo methods. Acta numerica, 24:259–328, 2015.
- Glebys Gonzalez, Mythra Balakuntala, Mridul Agarwal, Tomas Low, Bruce Knoth, Andrew W. Kirkpatrick, Jessica McKee, Gregory Hager, Vaneet Aggarwal, Yexiang Xue, Richard Voyles, and Juan Wachs. Asap: A semi-autonomous precise system for telesurgery during communication delays. *IEEE Transactions on Medical Robotics and Bionics*, 5(1):66–78, 2023. doi: 10.1109/TMRB.2023.3239674.
- Thomas Jaksch, Ronald Ortner, and Peter Auer. Near-optimal regret bounds for reinforcement learning. *The Journal of Machine Learning Research*, 11:1563–1600, 2010.
- Yanli Liu, Kaiqing Zhang, Tamer Basar, and Wotao Yin. An improved analysis of (variance-reduced) policy gradient and natural policy gradient methods. *Advances in Neural Information Processing Systems*, 33:7624– 7636, 2020.
- Hamid Reza Maei. *Gradient temporal-difference learning algorithms*. PhD thesis, University of Alberta, CAN, 2011. AAINR89455.
- Washim U Mondal and Vaneet Aggarwal. Improved sample complexity analysis of natural policy gradient algorithm with general parameterization for infinite horizon discounted reward markov decision processes. In *International Conference on Artificial Intelligence and Statistics*, pages 3097–3105. PMLR, 2024.
- Matteo Papini, Damiano Binaghi, Giuseppe Canonaco, Matteo Pirotta, and Marcello Restelli. Stochastic variance-reduced policy gradient. In *International conference on machine learning*, pages 4026–4035, 2018.
- Bhrij Patel, Wesley A Suttle, Alec Koppel, Vaneet Aggarwal, Brian M Sadler, Amrit Singh Bedi, and Dinesh Manocha. Global optimality without mixing time oracles in average-reward rl via multi-level actor-critic. In *International Conference on Machine Learning*, 2024.

- Martin L Puterman. *Markov decision processes: discrete stochastic dynamic programming*. John Wiley & Sons, 2014.
- Shuang Qiu, Zhuoran Yang, Jieping Ye, and Zhaoran Wang. On finite-time convergence of actor-critic algorithm. *IEEE Journal on Selected Areas in Information Theory*, 2(2):652–664, 2021.
- Wesley A Suttle, Amrit Bedi, Bhrij Patel, Brian M Sadler, Alec Koppel, and Dinesh Manocha. Beyond exponentially fast mixing in average-reward reinforcement learning via multi-level monte carlo actorcritic. In *International Conference on Machine Learning*, pages 33240–33267, 2023.
- Richard S Sutton, David McAllester, Satinder Singh, and Yishay Mansour. Policy gradient methods for reinforcement learning with function approximation. *Advances in neural information processing systems*, 12, 1999.
- Dipesh Tamboli, Jiayu Chen, Kiran Pranesh Jotheeswaran, Denny Yu, and Vaneet Aggarwal. Reinforced sequential decision-making for sepsis treatment: The posnegdm framework with mortality classifier and transformer. *IEEE Journal of Biomedical and Health Informatics*, 2024.
- Lingxiao Wang, Qi Cai, Zhuoran Yang, and Zhaoran Wang. Neural policy gradient methods: Global optimality and rates of convergence. In *International Conference on Learning Representations*, 2019.
- Yudan Wang, Yue Wang, Yi Zhou, and Shaofeng Zou. Non-asymptotic analysis for single-loop (natural) actor-critic with compatible function approximation. In *International Conference on Machine Learning*, 2024.
- Chen-Yu Wei, Mehdi Jafarnia Jahromi, Haipeng Luo, Hiteshi Sharma, and Rahul Jain. Model-free reinforcement learning in infinite-horizon average-reward markov decision processes. In *International conference on machine learning*, pages 10170–10180. PMLR, 2020.
- Chen-Yu Wei, Mehdi Jafarnia Jahromi, Haipeng Luo, and Rahul Jain. Learning infinite-horizon averagereward mdps with linear function approximation. In *International Conference on Artificial Intelligence and Statistics*, pages 3007–3015. PMLR, 2021.
- Pan Xu, Felicia Gao, and Quanquan Gu. Sample efficient policy gradient methods with recursive variance reduction. In *International Conference on Learning Representations*, 2019.
- Pan Xu, Felicia Gao, and Quanquan Gu. An improved convergence analysis of stochastic variance-reduced policy gradient. In *Uncertainty in Artificial Intelligence*, pages 541–551, 2020a.
- Tengyu Xu, Zhe Wang, and Yingbin Liang. Improving sample complexity bounds for (natural) actor-critic algorithms. *Advances in Neural Information Processing Systems*, 33:4358–4369, 2020b.
- Junyu Zhang, Chengzhuo Ni, Zheng Yu, Csaba Szepesvari, and Mengdi Wang. On the convergence and sample efficiency of variance-reduced policy gradient method. In A. Beygelzimer, Y. Dauphin, P. Liang, and J. Wortman Vaughan, editors, *Advances in Neural Information Processing Systems*, 2021a.
- Sheng Zhang, Zhe Zhang, and Siva Theja Maguluri. Finite sample analysis of average-reward td learning and q-learning. In *Advances in Neural Information Processing Systems*, volume 34, pages 1230–1242, 2021b.
- Shaofeng Zou, Tengyu Xu, and Yingbin Liang. Finite-sample analysis for sarsa with linear function approximation. *Advances in neural information processing systems*, 32, 2019.

### A Proof of Theorem 2

Consider  $x_0, \dots, x_{T-1}$  of Algorithm 2 and define the filtration  $\mathcal{F}_t = \sigma(x_j, j \leq t)$ . Let  $\mathbb{E}_t[\cdot]$  denote the expectation  $\mathbb{E}[\cdot|\mathcal{F}_t]$ . We start the proof with some guarantees on the MLMC estimator  $g_t$ . Similar guarantees exist in literature with MLMC estimation [Dorfman and Levy, 2022, Suttle et al., 2023, Beznosikov et al., 2023]. The difference between the result below and [Dorfman and Levy, 2022, Beznosikov et al., 2023] is that we consider a biased gradient estimate, in order for the results to be applicable to NPG subproblem with imperfect critic and average reward estimates. [Suttle et al., 2023] provides a bound with biased estimates as well, however these results are sharper, with the term  $\|\nabla F(x_t) - \mathbb{E}_t[g_t]\|^2$  bounded by  $\mathcal{O}(T_{\text{max}}^{-2})$  instead of  $\mathcal{O}(T_{\text{max}}^{-1})$ .

**Lemma 2.** Consider a time-homogeneous, ergodic Markov chain  $(Z_i)_{i\geq 0}$  with with a unique invariant distribution  $d_Z$  and Markov kernel Q. Further, let Q be uniformly geometrically ergodic with mixing time  $\tau_{mix}$ . Assume  $\|\mathbb{E}_{d_Z}[\nabla F(x,Z)] - \nabla F(x)\|^2 \leq \Delta^2$  and  $\|\nabla F(x,Z_i) - \nabla F(x)\|^2 \leq \sigma^2 + \delta^2 \|\nabla F(x)\|^2$  for all  $i \geq 0$ . Denote  $\mathbb{E}_t$  as the expectation conditioned on  $J_t$ . Then the MLMC estimator  $g_t$ 

$$g_t = g_t^0 + \begin{cases} 2^{J_t} \left( g_t^{J_t} - g_t^{J_t - 1} \right), & \text{if } 2^{J_t} \le T_{\max} \\ 0, & \text{otherwise} \end{cases} \quad \text{with} \quad g_t^j = 2^{-j} B^{-1} \sum_{i=1}^{2^j B} \nabla F(x_t, Z_{T_t + i})$$

satisfies the following:

- (a)  $\mathbb{E}_t[g_t] = \mathbb{E}_t[g_t^{\lfloor \log T_{\max} \rfloor}]$ (b)  $\mathbb{E}_t[\|\nabla F(x_t) - g_t\|^2] \le \mathcal{O}\left(\left(\tau_{\min} \log T_{\max} + \tau_{\min}^2\right)(\sigma^2 + \delta^2 \|\nabla F(x_t)\|^2)\right)$
- (c)  $\|\nabla F(x_t) \mathbb{E}_t[g_t]\|^2 \le \mathcal{O}\left(\tau_{\min}^2 T_{\max}^{-2}(\sigma^2 + \delta^2 \|\nabla F(x_t)\|^2) + \Delta\right).$

*Proof of Lemma 2.* Before proceeding to the proof, we state a useful lemma below:

**Lemma 3** (Lemma 1, [Beznosikov et al., 2023]). Let Assumption 1 hold. Then, for any  $n \ge 1$  and  $x \in \mathbb{R}^d$ , it holds that

$$\mathbb{E}_{d_Z}[\left\|n^{-1}\sum_{i=1}^n \nabla F(x, Z_i) - \nabla F(x)\right\|^2] \le \frac{8\tau_{\min}}{n} \left(\sigma^2 + \delta^2 \|\nabla F(x)\|^2\right).$$
<sup>(29)</sup>

where  $\|\nabla F(x, Z_i) - \nabla F(x)\| \le \sigma^2 + \delta^2 \|\nabla F(x)\|^2$  and  $d_Z$  is the stationary distribution of Z. Moreover, for any initial distribution d

$$\mathbb{E}_{d}[\left\|n^{-1}\sum_{i=1}^{n}\nabla F(x, Z_{i}) - \nabla F(x)\right\|^{2}] \leq \frac{C_{1}\tau_{\min}}{n} \left(\sigma^{2} + \delta^{2}\|\nabla F(x)\|^{2}\right),$$
(30)

where  $C_1 = 16(1 + \frac{1}{\ln^2 4})$ .

We first show that  $\mathbb{E}_t[g_t] = \mathbb{E}_t[g_t^{\lfloor \log_2 T_{\max} \rfloor}]$ . The proof is standard, where we simply compute the expectation with respect to  $J_t$ :

$$\mathbb{E}_t[g_t] = \mathbb{E}_t\left[\mathbb{E}_{J_t}[g_t]\right] = \mathbb{E}_t[g_t^0] + \sum_{i=1}^{\lfloor \log_2 T_{\max} \rfloor} \Pr\{J_t = i\} \cdot 2^i \mathbb{E}_t[g_t^i - g_t^{i-1}]$$
$$= \mathbb{E}_t[g_t^0] + \sum_{i=1}^{\lfloor \log_2 T_{\max} \rfloor} \mathbb{E}_t[g_t^i - g_t^{i-1}] = \mathbb{E}_t[g_t^{\lfloor \log_2 T_{\max} \rfloor}].$$

For the proof of (b), notice that

$$\begin{split} & \mathbb{E}_{t}[\|\nabla F(x_{t}) - g_{t}\|^{2}] \leq 2 \mathbb{E}_{t}[\|\nabla F(x_{t}) - g_{t}^{0}\|^{2}] + 2 \mathbb{E}_{t}[\|g_{t} - g_{t}^{0}\|^{2}] \\ &= 2 \mathbb{E}_{t}[\|\nabla F(x_{t}) - g_{t}^{0}\|^{2}] + 2 \sum_{i=1}^{\lfloor \log_{2} T_{\max} \rfloor} \Pr\{J_{t} = i\} \cdot 4^{i} \mathbb{E}_{t}[\|g_{t}^{i} - g_{t}^{i-1}\|^{2}] \\ &= 2 \mathbb{E}_{t}[\|\nabla F(x_{t}) - g_{t}^{0}\|^{2}] + 2 \sum_{i=1}^{\lfloor \log_{2} T_{\max} \rfloor} 2^{i} \mathbb{E}_{t}[\|g_{t}^{i} - g_{t}^{i-1}\|^{2}] \\ &\leq 2 \mathbb{E}_{t}[\|\nabla F(x_{t}) - g_{t}^{0}\|^{2}] + 4 \sum_{i=1}^{\lfloor \log_{2} T_{\max} \rfloor} 2^{i} \left(\mathbb{E}_{t}[\|\nabla F(x_{t}) - g_{t}^{i-1}\|^{2} + \mathbb{E}_{t}[\|g_{t}^{i} - \nabla F(x_{t})\|^{2}]\right) \,. \end{split}$$

To bound  $\mathbb{E}_t[\|\nabla F(x_t) - g_t\|^2]$ ,  $\mathbb{E}_t[\|\nabla F(x_t) - g_t^{i-1}\|^2$ ,  $\mathbb{E}_t[\|g_t^i - \nabla F(x_t)\|^2]$ , we apply Lemma 3 and get  $\mathbb{E}_t[\|\nabla F(x_t) - g_t\|^2]$ 

$$\mathbb{E}_{t}[\|\nabla F(x_{t}) - g_{t}\|^{2}]$$

$$\leq 2\sigma^{2} + 4\sum_{i=1}^{\lfloor \log_{2} T_{\max} \rfloor} 2^{i} \left( \frac{C_{1}\tau_{\min}}{2^{i}B} (\sigma^{2} + \delta^{2} \|\nabla F(x_{t})\|^{2}) + \frac{C_{1}\tau_{\min}^{2}}{2^{2i}B^{2}} (\sigma^{2} + \delta^{2} \|\nabla F(x_{t})\|^{2}) \right)$$

$$\leq \frac{4C_{1}(\sigma^{2} + \delta^{2} \|\nabla F(x_{t})\|^{2})\tau_{\min}\log_{2} T_{\max}}{B} + \frac{(4C_{1} + 2)(\sigma^{2} + \delta^{2} \|\nabla F(x_{t})\|^{2})\tau_{\min}^{2}}{B^{2}}.$$

For part (c), we have

$$\begin{aligned} \|\nabla F(x_t) - \mathbb{E}_t[g_t]\|^2 \\ &= \|\nabla F(x_t) - \mathbb{E}_t[g_t^{\lfloor \log_2 T_{\max} \rfloor}]\|^2 \\ &\leq 2\|\nabla F(x_t) - \mathbb{E}_{d_Z}[g_t^{\lfloor \log_2 T_{\max} \rfloor}]\|^2 + 2\|\mathbb{E}_{d_Z}[g_t^{\lfloor \log_2 T_{\max} \rfloor}] - \mathbb{E}_t[g_t^{\lfloor \log_2 T_{\max} \rfloor}]\|^2 \\ &\leq 2\Delta^2 + \underbrace{2\|\mathbb{E}_{d_Z}[g_t^{\lfloor \log_2 T_{\max} \rfloor}] - \mathbb{E}_t[g_t^{\lfloor \log_2 T_{\max} \rfloor}]\|^2}_{(A)}. \end{aligned}$$

$$(31)$$

(*A*) can be bounded using the maximal exact coupling argument in [Douc et al., 2018, Lemma 19.3.6 and Theorem 19.3.9] as:

$$\|\mathbb{E}_{d_{Z}}[g_{t}^{\lfloor \log_{2} T_{\max} \rfloor}] - \mathbb{E}_{t}[g_{t}^{\lfloor \log_{2} T_{\max} \rfloor}]\|^{2} \leq \mathcal{O}\left(\frac{\tau_{\max}^{2}}{T_{\max}^{2}} \cdot \|\nabla F(x_{t}, Z) - \nabla F(x_{t})\|^{2}\right)$$
$$\leq \mathcal{O}\left(\frac{\tau_{\max}^{2}}{T_{\max}^{2}} \cdot (\delta^{2} \|\nabla F(x_{t})\|^{2} + \sigma^{2})\right).$$
(32)

This concludes the proof of Lemma 2.

Towards proving the main claim, we state the following helpful lemmas that we will be using: Lemma 4 (Lemma 5, [Beznosikov et al., 2023]). *Consider Algorithm 2. Then for any*  $u \in \mathbb{R}^d$ , we get

$$\mathbb{E}_{t}[F(x_{t+1}^{f})] \leq F(u) - \langle \nabla F(x_{t}^{g}), u - x_{t}^{g} \rangle - \frac{\mu}{2} \|u - x_{t}^{g}\|^{2} - \frac{\gamma}{2} \|\nabla F(x_{t}^{g})\|^{2} + \frac{\gamma}{2} \|\mathbb{E}_{t}[g_{t}] - \nabla F(x_{t}^{g})\|^{2} + \frac{L\gamma^{2}}{2} \mathbb{E}_{t}[\|g_{t}\|^{2}].$$

**Lemma 5** (Lemma 6, [Beznosikov et al., 2023]). Consider Algorithm 2. Then for the iterates with  $m = (pq^{-1} - 1)/(\beta pq^{-1} - 1)$ , m > 0,  $q \ge 1$ , p > 0, it holds that

$$\mathbb{E}_{t}[\|x_{t+1} - x^{*}\|^{2}] \leq (1 + \alpha p \gamma q)(1 - \beta)\|x_{t} - x^{*}\|^{2} + (1 + \alpha p \gamma q)\beta\|x_{t}^{g} - x^{*}\|^{2} \\
+ (1 + \alpha p \gamma q)(\beta^{2} - \beta)\|x_{t} - x_{t}^{g}\|^{2} + p^{2}q^{2}\gamma^{2}\mathbb{E}_{t}[\|g_{t}\|^{2}] \\
- 2q^{2}\gamma\langle\nabla F(x_{t}^{g}), x_{t}^{g} + (pq^{-1} - 1)x_{t}^{f} - q^{-1}px^{*}\rangle \\
+ \frac{pq\gamma}{\alpha}\|\mathbb{E}_{t}[g_{t}] - \nabla F(x_{t}^{g})\|^{2},$$
(33)

where  $\alpha > 0$  is any positive constant.

The arguments for proving this claim are similar to ones used in [Beznosikov et al., 2023]. The key difference lies in tracking the effect of the parameter changes and the additional bias term. We must ensure that these changes do not accumulate and severely impact the bounds. Now using Lemma 4 with  $u = x^*$  and  $u = x_t^f$ , we obtain the following two inequalities

$$\begin{split} \mathbb{E}_t[F(x_{t+1}^f)] &\leq F(x^*) - \langle \nabla F(x_t^g), x^* - x_t^g \rangle - \frac{\mu}{2} \|x^* - x_t^g\|^2 - \frac{p\gamma}{2} \|\nabla F(x_t^g)\|^2 \\ &+ \frac{p\gamma}{2} \|\mathbb{E}_t[g_t] - \nabla F(x_t^g)\|^2 + \frac{Lp^2\gamma^2}{2} \mathbb{E}_t[\|g_t\|^2], \end{split}$$

$$\begin{split} \mathbb{E}_t[F(x_{t+1}^f)] \leq & F(x_t^f) - \langle \nabla F(x_t^g), x_t^f - x_t^g \rangle - \frac{\mu}{2} \|x_t^f - x_t^g\|^2 - \frac{p\gamma}{2} \|\nabla F(x_t^g)\|^2 \\ &+ \frac{p\gamma}{2} \|\mathbb{E}_t[g_t] - \nabla F(x_t^g)\|^2 + \frac{Lp^2\gamma^2}{2} \mathbb{E}_t[\|g_t\|^2]. \end{split}$$

We multiply the first inequality with  $2p\gamma q$ , the second with  $2\gamma q(q-p)$  and add them with (33) to obtain

$$\begin{split} \mathbb{E}_{t}[\|x_{t+1} - x^{*}\|^{2} + 2\gamma q^{2}F(x_{t+1}^{f})] \\ &\leq (1 + \alpha p\gamma q)(1 - \beta)\|x_{t} - x^{*}\|^{2} + (1 + \alpha p\gamma q)\beta\|x_{t}^{g} - x^{*}\|^{2} \\ &+ (1 + \alpha p\gamma q)(\beta^{2} - \beta)\|x_{t} - x_{t}^{g}\|^{2} - 2q^{2}\gamma \langle \nabla F(x_{t}^{g}), x_{t}^{g} + (pq^{-1} - 1)x_{t}^{f} - q^{-1}px^{*} \rangle \\ &+ p^{2}q^{2}\gamma^{2}\mathbb{E}_{t}[\|g_{t}\|^{2}] + \frac{pq\gamma}{\alpha}\|\mathbb{E}_{t}[g_{t}] - \nabla F(x_{t}^{g})\|^{2} \\ &+ 2p\gamma q \Big(F(x^{*}) - \langle \nabla F(x_{t}^{g}), x^{*} - x_{t}^{g} \rangle - \frac{\mu}{2}\|x^{*} - x_{t}^{g}\|^{2} - \frac{p\gamma}{2}\|\nabla F(x_{t}^{g})\|^{2} \\ &+ \frac{p\gamma}{2}\|\mathbb{E}_{t}[g_{t}] - \nabla F(x_{t}^{g})\|^{2} + \frac{Lp^{2}\gamma^{2}}{2}\mathbb{E}_{t}[\|g_{t}\|^{2}]\Big) \\ &+ 2\gamma q(q - p) \Big(F(x_{t}^{f}) - \langle \nabla F(x_{t}^{g}), x_{t}^{f} - x_{t}^{g} \rangle - \frac{\mu}{2}\|x_{t}^{f} - x_{t}^{g}\|^{2} - \frac{p\gamma}{2}\|\nabla F(x_{t}^{g})\|^{2} \\ &+ \frac{p\gamma}{2}\|\mathbb{E}_{t}[g_{t}] - \nabla F(x_{t}^{g})\|^{2} + \frac{Lp^{2}\gamma^{2}}{2}\mathbb{E}_{t}[\|g_{t}\|^{2}]\Big) \\ &= (1 + \alpha p\gamma q)(1 - \beta)\|x_{t} - x^{*}\|^{2} + 2\gamma q(q - p)F(x_{t}^{f}) + 2p\gamma qF(x^{*}) \\ &+ ((1 + \alpha p\gamma q)\beta - p\gamma q\mu)\|x_{t}^{g} - x^{*}\|^{2} \\ &+ (1 + \alpha p\gamma q)(\beta^{2} - \beta)\|x_{t} - x_{t}^{g}\|^{2} - p\gamma^{2}q^{2}\|\nabla F(x_{t}^{g})\|^{2} \\ &+ ((1 + \alpha p\gamma q)(1 - \beta)\|x_{t} - x^{*}\|^{2} + 2\gamma q(q - p)F(x_{t}^{f}) + 2p\gamma qF(x^{*}) \\ &+ ((1 + \alpha p\gamma q)(\beta^{2} - \beta)\|x_{t} - x_{t}^{g}\|^{2} - p\gamma^{2}q^{2}\|\nabla F(x_{t}^{g})\|^{2} \\ &+ (1 + \alpha p\gamma q)(\beta^{2} - \beta)\|x_{t} - x_{t}^{g}\|^{2} - p\gamma^{2}q^{2}\|\nabla F(x_{t}^{g})\|^{2} \\ &+ (1 + \alpha p\gamma q)(\beta^{2} - \beta)\|x_{t} - x_{t}^{g}\|^{2} - p\gamma^{2}q^{2}\|\nabla F(x_{t}^{g})\|^{2} \\ &+ (1 + \alpha p\gamma q)(\beta^{2} - \beta)\|x_{t} - x_{t}^{g}\|^{2} - p\gamma^{2}q^{2}\|\nabla F(x_{t}^{g})\|^{2} \\ &+ pq\gamma\left(\frac{1}{\alpha} + \gamma q\right)\|\mathbb{E}_{t}[g_{t}] - \nabla F(x_{t}^{g})\|^{2} + 2p^{2}q^{2}\gamma^{2}(1 + \gamma L)\mathbb{E}_{t}[\|g_{t} - \nabla F(x_{t}^{g})\|^{2}]. \end{split}$$

where (a) follows from the fact that  $||a + b||^2 \leq 2||a||^2 + 2||b||^2$ . Since  $\gamma \leq \frac{3}{4L}$ , the choice of  $\alpha = \frac{\beta}{2pq\gamma}$ ,  $\beta = \sqrt{4p^2 \mu \gamma/3}$ , and  $p \mu \gamma q = 3\beta/2$  gives

$$\beta = \sqrt{4p^2 \mu \gamma/3} \le \sqrt{p^2 \mu/L} \le 1,$$
  

$$(1 + \alpha pq\gamma)(1 - \beta) = \left(1 + \frac{\beta}{2}\right)(1 - \beta) \le \left(1 - \frac{\beta}{2}\right),$$
  

$$((1 + \alpha pq\gamma)\beta - p\mu\gamma q) = \left(\beta + \frac{\beta^2}{2} - p\mu\gamma q\right) \le \left(\frac{3\beta}{2} - p\mu\gamma q\right) \le 0,$$

and, therefore,

$$\begin{split} \mathbb{E}_t \Big[ \|x_{t+1} - x^*\|^2 + 2\gamma q^2 F(x_{t+1}^f) \Big] &\leq (1 - \beta/2) \|x_t - x^*\|^2 + 2\gamma q \left(q - p\right) F(x_t^f) + 2p\gamma q F(x^*) \\ &+ pq^2 \gamma^2 \left(1 + 2p/\beta\right) \|\mathbb{E}_t[g_t] - \nabla F(x_t^g)\|^2 \\ &+ 2p^2 q^2 \gamma^2 \left(1 + \gamma L\right) \mathbb{E}_t[\|g_t - \nabla F(x_t^g)\|^2] \\ &- p\gamma^2 q^2 (1 - 2p(1 + \gamma L)) \|\nabla F(x_t^g)\|^2. \end{split}$$

Adding  $-2\gamma q^2 F(x^*)$  on both sides yields

$$\mathbb{E}_t \Big[ \|x_{t+1} - x^*\|^2 + 2\gamma q^2 (F(x_{t+1}^f) - F(x^*)) \Big] \le (1 - \beta/2) \|x_t - x^*\|^2 + (1 - p/q) \cdot 2\gamma q^2 (F(x_t^f) - F(x^*)) \\ + pq^2 \gamma^2 (1 + 2p/\beta) \|\mathbb{E}_t[g_t] - \nabla F(x_t^g)\|^2 \\ + 2p^2 q^2 \gamma^2 (1 + \gamma L) \mathbb{E}_t[\|g_t - \nabla F(x_t^g)\|^2] \\ - p\gamma^2 q^2 (1 - 2p(1 + \gamma L)) \|\nabla F(x_t^g)\|^2.$$

Using Lemma 2 along with the fact that  $\|\mathbb{E}_{d_Z}[\nabla F(x,Z)] - \nabla F(x)\|^2 \le \Delta_1^2 + \Delta_2^2 \|\nabla F(x)\|^2$  and  $\|\nabla F(x,Z) - \nabla F(x)\|^2 \le \sigma^2 + \delta^2 \|\nabla F(x)\|^2$ , we obtain

$$\begin{split} \mathbb{E}_t \Big[ \|x_{t+1} - x^*\|^2 + 2\gamma q^2 (F(x_{t+1}^f) - F(x^*)) \Big] \\ &\leq (1 - \beta/2) \|x_t - x^*\|^2 + (1 - p/q) \cdot 2\gamma q^2 (F(x_t^f) - F(x^*)) \\ &+ pq^2 \gamma^2 (1 + 2p/\beta) \cdot (C_2 \tau_{\min}^2 T_{\max}^{-2} B^{-2} (\sigma^2 + \delta^2 \|\nabla F(x_t^g)\|^2) + \Delta_1^2 + \Delta_2^2 \|\nabla F(x_t^g)\|^2) \\ &+ 2p^2 q^2 \gamma^2 (1 + \gamma L) \cdot \left( (4C_1 \tau_{\min} B^{-1} \log T_{\max} + (4C_1 + 2)\tau_{\min}^2 B^{-2}) (\sigma^2 + \delta^2 \|\nabla F(x_t^g)\|^2) \right) \\ &- p\gamma^2 q^2 (1 - 2p(1 + \gamma L)) \|\nabla F(x_t^g)\|^2 \\ &\leq (1 - \beta/2) \|x_t - x^*\|^2 + (1 - p/q) \cdot 2\gamma q^2 (F(x_t^f) - F(x^*)) \\ &+ pq^2 \gamma^2 (1 + 2p/\beta) \cdot (C_2 \tau_{\min}^2 T_{\max}^{-2} B^{-2} (\sigma^2 + \delta'^2 \|\nabla F(x_t^g)\|^2) + \Delta_1^2) \\ &+ 2p^2 q^2 \gamma^2 (1 + \gamma L) \cdot \left( (4C_1 \tau_{\min} B^{-1} \log T_{\max} + (4C_1 + 2)\tau_{\min}^2 B^{-2}) (\sigma^2 + \delta'^2 \|\nabla F(x_t^g)\|^2) \right) \\ &- p\gamma^2 q^2 (1 - 2p(1 + \gamma L)) \|\nabla F(x_t^g)\|^2, \end{split}$$

where  $\delta'^2 \triangleq \delta^2 + \Delta_2^2 \cdot (T_{\max}^2 B^2) / (C_2 \tau_{\min}^2)$ . With  $T_{\max} \ge \sqrt{C_2 p^{-1} (1 + 2p/\beta)}$ , we have

$$\begin{split} \mathbb{E}_{t} \Big[ \|x_{t+1} - x^{*}\|^{2} + 2\gamma q^{2} (F(x_{t+1}^{f}) - F(x^{*})) \Big] \\ &\leq (1 - \beta/2) \|x_{t} - x^{*}\|^{2} + (1 - p/q) \cdot 2\gamma q^{2} (F(x_{t}^{f}) - F(x^{*})) \\ &+ p^{2} q^{2} \gamma^{2} \tau_{\text{mix}}^{2} B^{-2} (\sigma^{2} + \delta'^{2} \|\nabla F(x_{t}^{g})\|^{2}) \\ &+ 2p^{2} q^{2} \gamma^{2} (1 + \gamma L) \cdot (4C_{1} \tau_{\text{mix}} B^{-1} \log T_{\text{max}} + (4C_{1} + 2) \tau_{\text{mix}}^{2} B^{-2}) (\sigma^{2} + \delta'^{2} \|\nabla F(x_{t}^{g})\|^{2}) \\ &- p \gamma^{2} q^{2} (1 - 2p(1 + \gamma L)) \|\nabla F(x_{t}^{g})\|^{2} + p q^{2} \gamma^{2} (1 + 2p/\beta) \Delta_{1}^{2} \\ &\leq (1 - \beta/2) \|x_{t} - x^{*}\|^{2} + (1 - p/q) \cdot 2\gamma q^{2} (F(x_{t}^{f}) - F(x^{*})) \\ &+ 8p^{2} q^{2} \gamma^{2} (1 + \gamma L) \cdot (C_{1} \tau_{\text{mix}} B^{-1} \log T_{\text{max}} + (C_{1} + 1) \tau_{\text{mix}}^{2} B^{-2}) \sigma^{2} \\ &- p \gamma^{2} q^{2} \left[ 1 - 2p(1 + \gamma L) \left( 1 + 4 \left[ C_{1} \tau_{\text{mix}} B^{-1} \log T_{\text{max}} + (C_{1} + 1) \tau_{\text{mix}}^{2} B^{-2} \right] \delta'^{2} \right) \right] \|\nabla F(x_{t}^{g})\|^{2} \\ &+ p q^{2} \gamma^{2} (1 + 2p/\beta) \Delta_{1}^{2}. \end{split}$$

Since  $p = \sqrt{\frac{L}{\mu}} \cdot \frac{\log T}{T}$ ,  $B = \lceil \log T_{\max} \rceil$  and  $T_{\max} \ge 2$ , we obtain for large enough T

$$p \leq \left[1 + 2\left(1 + \gamma L\right) \left(1 + 4\left[C_{1}\tau_{\min}B^{-1}\log T_{\max} + (C_{1} + 1)\tau_{\min}^{2}B^{-2}\right]\delta^{\prime 2}\right)\right]^{-1},$$

and then,

$$\begin{split} \mathbb{E}_{t} \left[ \|x_{t+1} - x^{*}\|^{2} + 2\gamma q^{2} (F(x_{t+1}^{f}) - F(x^{*})) \right] &\leq (1 - \beta/2) \|x_{t} - x^{*}\|^{2} + (1 - p/q) \cdot 2\gamma q^{2} (F(x_{t}^{f}) - F(x^{*})) \\ &+ 8p^{2}q^{2}\gamma^{2} (1 + \gamma L) \cdot \left(C_{1}\tau_{\min}B^{-1}\log T_{\max} + (C_{1} + 1)\tau_{\min}^{2}B^{-2}\right)\sigma^{2} \\ &+ pq^{2}\gamma^{2} (1 + 2p/\beta) \Delta_{1}^{2}. \\ &\leq \max\left\{ (1 - \beta/2), (1 - p/q) \right\} \left[ \|x_{t} - x^{*}\|^{2} + 2\gamma q^{2} (F(x_{t}^{f}) - F(x^{*})) \right] \\ &+ 8p^{2}q^{2}\gamma^{2} (1 + \gamma L) \cdot \left(C_{1}\tau_{\min}B^{-1}\log T_{\max} + (C_{1} + 1)\tau_{\min}^{2}B^{-2}\right)\sigma^{2} \\ &+ pq^{2}\gamma^{2} (1 + 2p/\beta) \Delta_{1}^{2}. \end{split}$$

Using that  $pq\gamma = 3\beta/(2\mu)$ ,  $\beta/2 = p/q$ ,  $B = \lceil \log T_{\max} \rceil$  and  $\gamma \leq L^{-1}$ , we have

$$\mathbb{E}_{t}\left[\|x_{t+1} - x^{*}\|^{2} + 2\gamma q^{2}(F(x_{t+1}^{f}) - F(x^{*}))\right] \leq (1 - \beta/2) \left[\|x_{t} - x^{*}\|^{2} + 2\gamma q^{2}(F(x_{t}^{f}) - F(x^{*}))\right] + 36\beta^{2}\mu^{-2} \left(C_{1}\tau_{\min} + (C_{1} + 1)\tau_{\min}^{2}\right)\sigma^{2} + 2\beta q\gamma\Delta_{1}^{2}.$$
(34)

Unrolling the recursion and replacing  $\beta$  with  $\sqrt{4p^2\mu\gamma/3}$  yields

$$\mathbb{E}\left[\|x_{T} - x^{*}\|^{2} + 2\gamma q^{2} (F(x_{T}^{f}) - F(x^{*}))\right] \\
\leq \left(1 - \sqrt{\frac{p^{2}\mu\gamma}{3}}\right)^{T} \left[\|x^{0} - x^{*}\|^{2} + 2\gamma q^{2} (F(x_{f}^{0}) - F(x^{*}))\right] + 72\beta\mu^{-2} \left(C_{1}\tau_{\min} + (C_{1} + 1)\tau_{\min}^{2}\right)\sigma^{2} \\
\leq \exp\left(-\sqrt{\frac{p^{2}\mu\gamma T^{2}}{3}}\right) \left[\|x^{0} - x^{*}\|^{2} + 2\gamma q^{2} (F(x_{f}^{0}) - F(x^{*}))\right] + \frac{144p\sqrt{\gamma}}{\sqrt{3}\mu^{3/2}} \left(C_{1}\sigma^{2}\tau_{\min} + (C_{1} + 1)\sigma^{2}\tau_{\min}^{2}\right) + \frac{8}{\sqrt{3}}\Delta_{1}^{2}$$

Substituting  $p = \sqrt{\frac{L}{\mu}} \cdot \frac{\log T}{T}$ ,  $\gamma = \frac{3}{4L}$  and  $q = \sqrt{\frac{3}{\mu\gamma}}$  concludes the proof.

### **B** Proof of Theorem 3

In order to prove Theorem 3, we need to show that  $\|\mathbb{E}_{d_Z}[u(\omega,\theta;z)] - \nabla_{\omega}f(\omega)\|^2 \leq \mathcal{O}(\|\xi_H - \xi^*\|^2 + G_2^2\epsilon_{app})$ and  $\|u(\omega,\theta;z) - \nabla_{\omega}f(\omega)\|^2 \leq \sigma_{\omega}^2 + \delta_{\omega}^2 \|\nabla_{\omega}f(\omega)\|^2$ , for some  $\delta_{\omega}$  and  $\sigma_{\omega}$ . First, recall that  $u(\omega,\theta,z^j) := E(\theta,z^j)\omega = h(\theta,z^j)$ . It follows that

First, recall that 
$$u(\omega, \theta_k; z_k^J) \coloneqq F(\theta_k; z_k^J) \omega - h(\theta_k; z_k^J)$$
. It follows that

$$\begin{split} \|u(\omega,\theta;z) - \nabla_{\omega}f(\omega)\|^{2} &\leq \|(F(\theta_{k};z_{k}^{j}) - F(\theta_{k}))\omega - (h(\theta_{k};z_{k}^{j}) - \nabla J(\theta_{k}))\|^{2} \\ &\leq 2\|F(\theta_{k};z_{k}^{j}) - F(\theta_{k})\|^{2}\|\omega\|^{2} + 2\|h(\theta_{k};z_{k}^{j}) - \nabla J(\theta_{k})\|^{2} \\ &\leq 2(2\|F(\theta_{k};z_{k}^{j})\|^{2} + 2\|F(\theta_{k})\|^{2})\|\omega\|^{2} + 2\|h(\theta_{k};z_{k}^{j}) - \nabla J(\theta_{k})\|^{2} \\ &\stackrel{(a)}{\leq} 8G_{1}^{2}\|\omega\|^{2} + 4(\|h(\theta_{k};z_{k}^{j})\|^{2} + \|\nabla J(\theta_{k})\|^{2}) \\ &\stackrel{(b)}{\leq} \frac{16G_{1}^{2}}{\mu^{2}}\|\nabla_{\omega}f(\omega)\|^{2} + 4C^{2}(1 + 4G_{1}^{2}\mu^{-2}), \end{split}$$

where (a) follows from the fact that  $||F(\theta_k; z_k^j)||$ ,  $||F(\theta)|| \leq G_1$  since  $F(\theta_k; z_k^j) = \nabla \log \pi_{\theta_k}(a_k^j | s_k^j) \log \pi_{\theta_k}(a_k^j | s_k^j)^\top$ ,  $F(\theta_k) = \mathbb{E}_{d_Z}[F(\theta_k; z_k^j)]$  and  $||\nabla \log \pi_{\theta_k}(a_k^j | s_k^j)|| \leq G_1$  (from Assumption 5). Whereas (b) follows from the fact that  $\omega = F(\theta)^{-1}(\nabla_{\omega}f(\omega) + \nabla J(\theta_k))$  and that eigenvalues of  $F(\theta)^{-1}$  are upper bounded by  $1/\mu$  (from Assumption 6). The statement now follows by setting  $\sigma_{\omega} = 4C^2(1 + 4G_1^2\mu^{-2})$  and  $\delta_{\omega} = \frac{16G_1^2}{\mu^2}$ .

We overload notation from the general strongly convex optimization with Markovian noise to re-define the filtration  $\{\mathcal{F}_h\}_{h\geq 0}$  as  $\mathcal{F}_0 = \sigma(\theta)$  and  $\mathcal{F}_h = \sigma(\theta, \omega_1, \cdots, \omega_h)$  for  $h \geq 1$ . Additionally, let  $\mathbb{E}_h[\cdot] = \mathbb{E}[\cdot|\mathcal{F}_h]$ . Towards bounding  $\|\mathbb{E}_{d_Z}[u(\omega, \theta; z)] - \nabla_{\omega}f(\omega)\|^2$ , we introduce a few terms to facilitate exposition. Before proceeding, we note that  $\mathbb{E}_{d_Z}[u(\omega_h, \theta_k; z_k^j) - \nabla f_\omega(\omega_h)] = \mathbb{E}_{d_Z}[h(\theta_k; z_k^j)] - \nabla J(\theta_k)$  since we have  $\mathbb{E}_{d_Z}[F(\theta_k; z_k^j)] = F(\theta_k)$ . Hence, we focus on bounding  $\|\mathbb{E}_{d_Z}[h(\theta_k; z_k^j)] - \nabla J(\theta_k)\|$ .

Let

$$h(\theta_k; z_k^j) = (r(s_k^j, a_k^j) - \eta_k + \langle \phi(s_k^{j+1}) - \phi(s_k^j), \nu_k \rangle) \cdot \nabla_\theta \log \pi_{\theta_k}(a_k^j | s_k^j)$$

$$(35)$$

$$\bar{h}(\theta_k; z_k^j) = (r(s_k^j, a_k^j) - \eta_k^* + \langle \phi(s_k^{j+1}) - \phi(s_k^j), \nu_k^* \rangle) \cdot \nabla_\theta \log \pi_{\theta_k}(a_k^j | s_k^j)$$
(36)

$$h^*(\theta_k; z_k^j) = (r(s_k^j, a_k^j) - \eta_k^* + V(s_k^{j+1}) - V(s_k^j)) \cdot \nabla_\theta \log \pi_{\theta_k}(a_k^j | s_k^j)$$
(37)

where  $\eta_k$  and  $\nu_k$  denote the average reward and critic parameter estimate obtained at iteration k, respectively. Additionally, let  $\eta_k^* = J(\theta_k)$  and  $\omega_k^*$  represent the limiting point of TD(0) applied to evaluating the policy  $\pi_{\theta_k}$ . It's important to note that this notation differs from the algorithm, where we utilize  $\eta_H$  and  $\nu_H$  at every iteration k, as each of these estimates are acquired after H iterations of the accelerated MLMC TD

algorithm. However, to maintain clarity regarding the dependence of k, we opt for the subscript k instead of H for this proof. Notice that

$$h(\theta_k; z_k^j) - \nabla J(\theta_k) = \left(\underbrace{h(\theta_k; z_k^j) - \bar{h}(\theta_k; z_k^j)}_{(a)}\right) + \left(\underbrace{\bar{h}(\theta_k; z_k^j) - h^*(\theta_k; z_k^j)}_{(b)}\right) + \left(\underbrace{h^*(\theta_k; z_k^j) - \nabla J(\theta_k)}_{(c)}\right), \quad (38)$$

where (a) and (b) can be written as

$$h(\theta_k; z_k^j) - \bar{h}(\theta_k; z_k^j) = \left( (\eta_k^* - \eta_k) + \langle \phi(s_k^{j+1}) - \phi(s_k^j), \nu_k - \nu_k^* \rangle \right) \nabla \log \pi_{\theta_k}(a_k^j | s_k^j)$$
$$\bar{h}(\theta_k; z_k^j) - h^*(\theta_k; z_k^j) = \left[ \left( \langle \phi(s_k^{j+1}), \nu_k^* \rangle - V_{\theta_k}(s_k^{j+1}) \right) - \left( \langle \phi(s_k^j), \nu_k^* \rangle - V_{\theta_k}(s_k^j) \right) \right] \nabla \log \pi_{\theta_k}(a_k^j | s_k^j)$$

and (c) is such that  $\mathbb{E}_{d_Z}\left[h^*(\theta_k; z_k^j) - \nabla J(\theta_k)\right] = 0.$ We then obtain the following

$$\mathbb{E}_{d_{Z}}[h(\theta_{k};z_{k}^{j}) - \nabla J(\theta_{k})] = \mathbb{E}_{d_{Z}}\left[\left((\eta_{k}^{*} - \eta_{k}) + \langle \phi(s_{k}^{j+1}) - \phi(s_{k}^{j}), \nu_{k} - \nu_{k}^{*} \rangle\right) \nabla \log \pi_{\theta_{k}}(a_{k}^{j}|s_{k}^{j})\right] \\ + \mathbb{E}_{d_{Z}}\left[\left[\left(\langle \phi(s_{k}^{j+1}), \nu_{k}^{*} \rangle - V_{\theta_{k}}(s_{k}^{j+1})\right) - \left(\langle \phi(s_{k}^{j}), \nu_{k}^{*} \rangle - V_{\theta_{k}}(s_{k}^{j})\right)\right] \nabla \log \pi_{\theta_{k}}(a_{k}^{j}|s_{k}^{j})\right] + 0$$

Let  $\bar{C} \coloneqq \sup_{s,s'} \|\phi(s) - \phi(s')\|$ . Then,

$$\begin{split} \| \mathbb{E}_{d_{Z}} [h(\theta_{k}; z_{k}^{j}) - \nabla J(\theta_{k})] \| \\ &= \mathbb{E}_{d_{Z}} \| h(\theta_{k}; z_{k}^{j}) - \nabla J(\theta_{k}) \| \\ &\leq \mathbb{E}_{d_{Z}} \left[ \left( \| \eta_{k}^{*} - \eta_{k} \| + \bar{C} \| \nu_{k} - \nu_{k}^{*} \| \right) \| \nabla \log \pi_{\theta_{k}}(a_{k}^{j} | s_{k}^{j}) \| \right] + \mathbb{E}_{d_{Z}} \left[ \epsilon_{\mathrm{app}} \| \nabla \log \pi_{\theta_{k}}(a_{k}^{j} | s_{k}^{j}) \| \right] \\ &\leq G_{1} \| \eta_{k}^{*} - \eta_{k} \| + \bar{C}G_{1} \| \nu_{k} - \nu_{k}^{*} \| + G_{1} \sqrt{\epsilon_{\mathrm{app}}} \\ &\leq G_{1} \bar{C} \| \xi_{k}^{*} - \xi_{k} \| + G_{1} \sqrt{\epsilon_{\mathrm{app}}}. \end{split}$$

Since all conditions of Theorem 2 are satisfied, we obtain

$$\mathbb{E}\left[\|\omega_{H} - \omega^{*}\|^{2} + \frac{6}{\mu}(f_{\omega}(\omega_{H}^{f}) - f(\omega^{*}))\right] \leq \mathcal{O}\left(\frac{1}{H^{2}}\left[\|\omega_{0} - \omega^{*}\|^{2} + \frac{6}{\mu}(f(\omega_{0}) - f(\omega^{*}))\right] + \frac{\log H}{\mu H}\left(\sigma_{\omega}^{2}\tau_{\min}^{2}\right) + \Delta^{2}\right).$$
(39)

where  $\Delta = G_1 \overline{C} \mathbb{E} \|\xi_k^* - \xi_k\| + G_1 \sqrt{\epsilon_{\text{app}}}$  and  $\sigma_{\omega} = 4C^2(1 + 4G_1^2\mu^{-2})$ . Now we focus on the second part of Theorem 3. Notice that  $(\mathbb{E}[\omega_h|\theta])_{h\geq 1}$  can be recursively computed as:

$$\begin{split} & \mathbb{E}[\omega_{h}^{g}|\theta] = m_{\omega} \mathbb{E}[\omega_{h}^{f}|\theta] + (1 - m_{\omega}) \mathbb{E}[\omega_{h}|\theta] \\ & \mathbb{E}[\omega_{h+1}^{f}|\theta] = \mathbb{E}[\omega_{h}^{g}|\theta] - p_{\omega}\gamma_{\omega} \mathbb{E}[\tilde{u}_{h}|\theta] \\ & \mathbb{E}[\omega_{h+1}|\theta] = q_{\omega} \mathbb{E}[\omega_{h+1}^{f}|\theta] + (p_{\omega} - q_{\omega}) \mathbb{E}[\omega_{h}^{f}|\theta] + (1 - p_{\omega})(1 - \beta_{\omega}) \mathbb{E}[\omega_{h}|\theta] + (1 - p_{\omega})\beta_{\omega} \mathbb{E}[\omega_{h}^{g}|\theta] \end{split}$$

Thus, the above updates correspond to the case where the NPG loop uses  $\mathbb{E}[\tilde{u}_h|\theta]$  instead of  $\tilde{u}_h$ . As a result, we can again invoke Theorem 2 to obtain a bound for  $\|\mathbb{E}[\omega_H|\theta] - \omega^*\|$ , provided we obtain an upper bound for  $\|\mathbb{E}[\tilde{u}_h|\theta] - \nabla_{\omega} f(\mathbb{E}[\omega_h|\theta])\|$ , which will serve as a bound for both the bias and the variance for the update rule. Note that:

$$\mathbb{E}[u_h|\theta] = \mathbb{E}[\tilde{F}(\theta)\omega_h - \tilde{h}|\theta] = \mathbb{E}[\mathbb{E}[\tilde{F}(\theta)\omega_h - \tilde{h}|\omega_h, \theta]|\theta] = \mathbb{E}[\tilde{F}(\theta)|\theta] \mathbb{E}[\omega_h|\theta] - \mathbb{E}[\tilde{h}(\theta)|\theta],$$

It follows that:

$$\begin{aligned} \| \mathbb{E}[u_{h}|\theta] - \nabla_{\omega} f(\mathbb{E}[\omega_{h}|\theta]) \| &= \| \mathbb{E}[\tilde{F}(\theta)|\theta] - F(\theta)] \mathbb{E}[\omega_{h}|\theta] - (\mathbb{E}[\tilde{h}(\theta)|\theta] - \nabla J(\theta)) \| \\ &\leq \| \mathbb{E}[\tilde{F}(\theta)|\theta] - F(\theta)] \| \| \mathbb{E}[\omega_{h}|\theta] \| + \| \mathbb{E}[\tilde{h}(\theta)|\theta] - \nabla J(\theta) \| \\ &\leq \mu^{-1} \| \mathbb{E}[\tilde{F}(\theta)|\theta] - F(\theta)] \| \| \nabla f_{\omega}(\mathbb{E}[\omega_{h}|\theta]) \| + \| \mathbb{E}[\tilde{h}(\theta)|\theta] - \nabla J(\theta) \| \end{aligned}$$

Recall that  $||F(\theta) - F(\theta)|| \le ||F(\theta, z)|| + ||F(\theta)|| \le 2G_1$  and  $||\mathbb{E}_{d_Z}[F(\theta, z)] - F(\theta)|| = 0$ . It then follows from Lemma 2(c):

$$\|\mathbb{E}[\tilde{F}(\theta)|\theta] - F(\theta)\| = \|\mathbb{E}_h[\tilde{F}(\theta)] - F(\theta)\| \le \mathcal{O}\left(\frac{G_1\tau_{\min}}{T_{\max}}\right).$$
(40)

 $\text{Separately,} \| \, \mathbb{E}[\tilde{h}(\theta)|\theta] - \nabla J(\theta) \| = \| \, \mathbb{E}[\mathbb{E}_{h}[\tilde{h}(\theta)]|\theta] - \nabla J(\theta) \|.$ 

$$\begin{split} & \mathbb{E}_{h}[\bar{h}(\theta)] - \nabla J(\theta) \\ &= \mathbb{E}_{h} \left[ \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} (h(\theta; z_{i}) - \nabla J(\theta)) \right] \\ &= \mathbb{E}_{h} \left[ \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} (h(\theta; z_{i}) - \bar{h}(\theta; z_{i})) + (\bar{h}(\theta; z_{i}) - h^{*}(\theta; z_{i})) + (h^{*}(\theta; z_{i}) - \nabla J(\theta)) \right] \\ &= \mathbb{E}_{h} \left[ \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} (h(\theta; z_{i}) - \bar{h}(\theta; z_{i})) \right] + \mathbb{E}_{h} \left[ \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} (\bar{h}(\theta; z_{i}) - h^{*}(\theta; z_{i})) \right] \\ &+ \mathbb{E}_{h} \left[ \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} (h^{*}(\theta; z_{i}) - \nabla J(\theta)) \right]. \end{split}$$
(41)

Note that  $\mathbb{E}_{d_z}[h^*(\theta; z_i)] = \nabla J(\theta)$  and  $||h^*(\theta; z_i)|| \leq \mathcal{O}(G_1 \tau_{\min})$  since  $V^{\pi}(s) \leq \mathcal{O}(\tau_{\min})$  for all states *s* and policies  $\pi$  [Wei et al., 2020]. Thus, using Lemma 2:

$$\left\| \mathbb{E}\left[ \mathbb{E}_{h}\left[ \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} (h^{*}(\theta; z_{i}) - \nabla J(\theta)) \right] \middle| \theta \right] \right\| \leq \mathcal{O}\left( \frac{G_{1} \tau_{\min}^{2}}{T_{\max}} \right).$$
(42)

Recall that  $\|\bar{h}(\theta;z_i) - h^*(\theta;z_i)\| \le G_1 \sqrt{\epsilon_{\mathrm{app}}}$ , which implies

$$\left\| \mathbb{E}\left[ \mathbb{E}_{h}\left[ \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} (\bar{h}(\theta; z_{i}) - h^{*}(\theta; z_{i})) \right] \right| \theta \right] \right\| \leq \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} \|\bar{h}(\theta; z_{i}) - h^{*}(\theta; z_{i})\| \leq G_{1} \sqrt{\epsilon_{\exp}}$$

Also,

$$\mathbb{E}\left[\mathbb{E}_{h}\left[\frac{1}{T_{\max}}\sum_{i=1}^{T_{\max}}(\bar{h}(\theta;z_{i})-h(\theta;z_{i}))\right]\right|\theta\right] \\
=\mathbb{E}\left[\mathbb{E}\left[\frac{1}{T_{\max}}\sum_{i=1}^{T_{\max}}(\bar{h}(\theta;z_{i})-h^{*}(\theta;z_{i}))\right|\hat{\mathcal{F}}_{h},\theta\right]\right|\theta\right] \\
=\mathbb{E}\left[\mathbb{E}\left[\frac{1}{T_{\max}}\sum_{i=1}^{T_{\max}}(\eta_{k}-\eta_{k}^{*}+\langle\phi(s_{i+1})-\phi(s_{i}),\nu_{k}-\nu_{k}^{*}\rangle)\nabla\log\pi_{\theta}(a_{i}|s_{i})\right|\hat{\mathcal{F}}_{h},\theta\right]\right|\theta\right] \\
=\mathbb{E}\left[\frac{1}{T_{\max}}\sum_{i=1}^{T_{\max}}(\mathbb{E}[\eta_{k}|\theta]-\eta_{k}^{*}+\langle\phi(s_{i+1})-\phi(s_{i}),\mathbb{E}[\nu_{k}|\theta]-\nu_{k}^{*}\rangle)\nabla\log\pi_{\theta}(a_{i}|s_{i})\right|\theta\right].$$
(43)

It follows that

$$\left\| \mathbb{E} \left[ \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} \left( \bar{h}(\theta; z_i) - h(\theta; z_i) \right) \middle| \theta \right] \right\|$$

$$\leq \frac{1}{T_{\max}} \sum_{i=1}^{T_{\max}} \left\| \left( \mathbb{E}[\eta_k | \theta] - \eta_k^* + \langle \phi(s_{i+1}) - \phi(s_i), \mathbb{E}[\nu_k | \theta] - \nu_k^* \rangle \right) \nabla \log \pi_{\theta}(a_i | s_i) \right\|$$

$$\leq G_1 \left\| \mathbb{E}[\eta_k | \theta] - \eta_k^* \right\| + G_1 \left\| \mathbb{E}[\nu_k | \theta] - \nu_k^* \right\|.$$
(44)

Combining the above bounds, we obtain:

$$\begin{aligned} \|\mathbb{E}[u_{h}|\theta] - \nabla_{\omega}f(\mathbb{E}[\omega_{h}|\theta])\| \\ &\leq \mathcal{O}\left(\frac{G_{1}\tau_{\min}}{\mu T_{\max}}\|\nabla_{\omega}f_{\omega}(\mathbb{E}[u_{h}|\theta])\| + \frac{G_{1}\tau_{\min}^{2}}{T_{\max}} + G_{1}\|\mathbb{E}[\eta_{k}|\theta] - \eta_{k}^{*}\| + G_{1}\|\mathbb{E}[\nu_{k}|\theta] - \nu_{k}^{*}\| + G_{1}\sqrt{\epsilon_{\mathrm{app}}}\right). \end{aligned}$$

Set  $\Delta_1 = \sigma = \frac{G_1 \tau_{\text{mix}}^2}{T_{\text{max}}} + G_1 \|\mathbb{E}[\eta_k|\theta] - \eta_k^*\| + G_1 \|\mathbb{E}[\nu_k|\theta] - \nu_k^*\| + G_1 \sqrt{\epsilon_{\text{app}}} \text{ and } \Delta_2 = \delta = \frac{G_1 \tau_{\text{mix}}}{\mu T_{\text{max}}} \text{ in Theorem 2 to obtain:}$ 

$$\begin{split} & \mathbb{E}\left[\|\mathbb{E}[\omega_{H}|\theta] - \omega^{*}\|^{2}\right] \\ & \leq \mathcal{O}\left(\exp\left(-\sqrt{\frac{p^{2}\mu\gamma H^{2}}{3}}\right)\left[\|\omega_{0} - \omega^{*}\|^{2}\right] + \frac{p\sqrt{\gamma}}{\mu^{3/2}}\left(\sigma^{2}\tau_{\mathrm{mix}} + \sigma^{2}\tau_{\mathrm{mix}}^{2}\right) + \Delta_{1}^{2}\right) \\ & \leq \mathcal{O}\left(\frac{1}{H^{2}}[\|\omega_{0} - \omega^{*}\|^{2}] + \frac{G_{1}^{2}\tau_{\mathrm{mix}}^{4}}{T_{\mathrm{max}}^{2}} + G_{1}^{2}\|\mathbb{E}[\eta_{k}|\theta] - \eta_{k}^{*}\|^{2} + G_{1}^{2}\|\mathbb{E}[\nu_{k}|\theta] - \nu_{k}^{*}\|^{2} + G_{1}^{2}\epsilon_{\mathrm{app}} \right. \\ & + \frac{(\log H)\tau_{\mathrm{mix}}^{2}}{\mu H}\left(\frac{G_{1}^{2}\tau_{\mathrm{mix}}^{4}}{T_{\mathrm{max}}^{2}} + G_{1}^{2}\|\mathbb{E}[\eta_{k}|\theta] - \eta_{k}^{*}\|^{2} + G_{1}^{2}\|\mathbb{E}[\nu_{k}|\theta] - \nu_{k}^{*}\|^{2} + G_{1}^{2}\epsilon_{\mathrm{app}}\right)\right). \end{split}$$

The result follows by setting  $T_{max} = H$  and neglecting lower order terms.

### C Proof of Theorem 4

Unless specified otherwise, we drop the subscript  $\theta$  for ease of exposition. We overload notation again to re-define the filtration  $\{\mathcal{F}_h\}_{h\geq 0}$  as  $\mathcal{F}_0 = \sigma(\theta)$  and  $\mathcal{F}_h = \sigma(\theta, \xi_1, \dots, \xi_h)$  for  $h \geq 1$ .  $\mathbb{E}_h[\cdot]$  is defined as before as  $\mathbb{E}_h[\cdot] = \mathbb{E}[\cdot|\mathcal{F}_h]$ .

**Lemma 6.** Let  $A = \mathbb{E}_{d_Z}[A(\theta; z)]$ . By setting large enough  $c_\beta$ , Assumption 3 implies that A is positive definite, i.e.,  $\xi^{\top} A \xi \geq \lambda/2 \cdot \|\xi\|^2$ , for all  $\xi$ .

*Proof of Lemma 6.* The proof of this lemma follows similarly as in proof of Lemma 2 in [Zhang et al., 2021b]. We begin by noting that *A* can be written as

$$A = \begin{pmatrix} c_{\beta} & 0\\ -\Phi^{\top} D e & \Phi^{\top} D (P - I) \Phi \end{pmatrix},$$
(45)

where *e* denotes the vector of all ones, *D* denotes the diagonal matrix with  $D_{ii} = d^{\pi}(s_i)$ , and  $\Phi$  is the  $|S| \times d$  whose  $i^{th}$  column is  $\phi(s_i)$ . Observe that

$$\min_{\|\xi\|^{2}=1} \xi^{\top} A\xi = \min_{(\eta^{2}+\|\nu\|^{2})=1} c_{\beta} \eta^{2} - \eta \nu^{\top} (\Phi^{\top} De) + \nu^{\top} \Phi^{\top} D(I-P) \Phi \nu$$

$$\stackrel{(a)}{\geq} \min_{(\eta^{2}+\|\nu\|^{2})=1} c_{\beta} \eta^{2} - \eta \nu^{\top} (\Phi^{\top} De) + \lambda \|\nu\|^{2}$$

$$\stackrel{(b)}{\geq} \min_{(\eta^{2}+\|\nu\|^{2})=1} c_{\beta} \eta^{2} - |\eta| \|\nu\| + \lambda \|\nu\|^{2},$$
(46)

where (a) follows from Assumption 3 and (b) follows from the fact that

$$|\nu^{\top} \Phi^{\top} De| = |(\Phi\nu)^{\top} d^{\pi_{\theta}}| \le \|\Phi\nu\|_{\infty} \|d^{\pi_{\theta}}\|_{1} \le \|\Phi\nu\|_{\infty} \le \max_{i \in \mathcal{S}} \|\phi(i)\|_{2} \|\nu\|_{2} \le \|\nu\|_{2}.$$
(47)

It follows that

$$\min_{\|\xi\|^2 = 1} \xi^{\top} A\xi \ge \min_{\|\eta\| \le 1} c_{\beta} \eta^2 - |\eta| \sqrt{1 - \eta^2} + \lambda (1 - \eta^2) 
= \min_{u \in [0,1]} c_{\beta} u - \sqrt{u(1 - u)} + \lambda (1 - u) 
= \lambda + \min_{u \in [0,1]} (c_{\beta} - \lambda) u - \sqrt{u(1 - u)} 
\ge \lambda/2,$$
(48)

for  $c_{\beta} \geq \lambda + \sqrt{\frac{1}{\lambda^2} - 1}$ .

Let  $\tilde{A}_h$  and  $\tilde{b}_h$  denote the MLMC estimators, i.e.

$$\tilde{A}_{h} = A_{h}^{0} + \begin{cases} 2^{Q_{h}^{k}} (A_{h}^{P_{h}^{k}} - A_{h}^{P_{h}^{k} - 1}), & \text{if } 2^{P_{h}^{k}} \le T_{\max} \\ 0, & \text{otherwise} \end{cases}$$
(49)

with  $A_h^j = \frac{1}{2^j} \sum_{k=1}^{2^j} A(\theta_k; z_h^k)$  and

$$\tilde{b}_{h} = b_{h}^{0} + \begin{cases} 2^{Q_{h}^{k}} (b_{h}^{P_{h}^{k}} - b_{h}^{P_{h}^{k}-1}), & \text{if } 2^{P_{h}^{k}} \le T_{\max} \\ 0, & \text{otherwise} \end{cases}$$
(50)

with  $b_h^j = \frac{1}{2^j} \sum_{k=1}^{2^j} b(\theta_k; z_h^k)$ . For brevity, we henceforth denote the samples obtained at iteration h of the critic loop,  $A(\theta; z)$  and  $b(\theta; z)$ , as  $A_h$  and  $b_h$ , respectively. Notice that

$$\|b_h\| \le |c_\beta r(s,a)| + \| - r(s,a)\phi(s)\| \stackrel{(a)}{\le} c_\beta + 1 \triangleq L_b,$$
(51)

where (*a*) follows since we assume the rewards obtained r(s, a) and the critic features  $\phi(s)$  are bounded by 1 for all  $s \in S$  and  $a \in A$ . Additionally,

$$\|A_{h}\| \leq |c_{\beta}| + \| - \phi(s)\| + \|\phi(s)(\phi(s') - \phi(s))^{\top}\| \\ \leq c_{\beta} + \|\phi(s)\| + \|\phi(s)\|(\|\phi(s')\| + \|\phi(s)\|) \\ \stackrel{(a)}{\leq} c_{\beta} + 3 \triangleq L_{A},$$
(52)

where (*a*) holds since  $\phi(s) \leq 1$  for all  $s \in S$ . Using these inequalities, we can bound the bias and variance of  $\tilde{A}_h$  and  $\tilde{b}_h$  as follows:

**Lemma 7.** = Consider Algorithm 1. For  $||A_h|| \le L_A$  and  $||b_h|| \le L_b$ , the following statements hold:

$$(a) \|\mathbb{E}_{h}[\tilde{A}_{h}] - A\| \leq \mathcal{O}\left(\frac{L_{A}\tau_{\min}}{T_{\max}}\right)$$

$$(b) \|\mathbb{E}_{h}[\tilde{b}_{h}] - b\| \leq \mathcal{O}\left(\frac{L_{b}\tau_{\min}}{T_{\max}}\right)$$

$$(c) \mathbb{E}_{h} \|\tilde{A}_{h} - A\|^{2} \leq \mathcal{O}\left(L_{A}^{2}\left(\tau_{\min}\log T_{\max} + \tau_{\min}^{2}\right)\right)$$

$$(d) \mathbb{E}_{h} \|\tilde{b}_{h} - b\| \leq \mathcal{O}\left(L_{b}^{2}\left(\tau_{\min}\log T_{\max} + \tau_{\min}^{2}\right)\right)$$

$$(e) \|\mathbb{E}_{h}[\tilde{v}_{h}] - (A\xi_{h} - b)\| \leq \mathcal{O}\left(\frac{(L_{A}\|\xi_{h} - \xi^{*}\| + L_{b})\tau_{\max}}{T_{\max}}\right).$$

Lemma 7(a) and (b) follow from Lemma 2(c), while Lemma 7(c) and (d) follow from Lemma 2(b). Meanwhile, Lemma 7(e) follows from noting that:

$$\mathbb{E}_{h}[\tilde{v}_{h}] = \mathbb{E}_{h}[\tilde{A}_{h}\xi_{h} - \tilde{b}_{h}] = \mathbb{E}_{h}[\tilde{A}_{h}]\xi_{h} - \mathbb{E}_{h}[\tilde{b}_{h}] = \mathbb{E}_{h}[\tilde{A}_{h}]\xi_{h} - \mathbb{E}_{h}[\tilde{b}_{h}]$$
(53)

and applying Lemma 7(a) and (b). Now, for the proof of Theorem 4, observe that

$$\begin{split} \|\xi_{h+1} - \xi^*\|^2 &= \|\xi_h - \beta \tilde{v}_h - \xi^*\|^2 \\ &= \|\xi_h - \xi^*\|^2 + 2\beta\langle\xi_h - \xi^*, \tilde{v}_h\rangle + \beta^2 \|\tilde{v}_h\|^2 \\ &\stackrel{(a)}{=} \|\xi_h - \xi^*\|^2 + 2\beta\langle\xi_h - \xi^*, A(\xi_h - \xi^*)\rangle + 2\beta\langle\xi_h - \xi^*, \tilde{v}_h - A(\xi_h - \xi^*)\rangle \\ &+ \beta^2 \|\tilde{v}_h\|^2 \\ &\leq \|\xi_h - \xi^*\|^2 - \beta\lambda\|\xi_h - \xi^*\|^2 + 2\beta\langle\xi_h - \xi^*, \tilde{v}_h - A(\xi_h - \xi^*)\rangle \\ &+ 2\beta^2 \|\tilde{v}_h - A(\xi_h - \xi^*)\|^2 + 2\beta^2 \|A(\xi_h - \xi^*)\|^2 \\ &\leq \|\xi_h - \xi^*\|^2 - \beta\lambda\|\xi_h - \xi^*\|^2 + 2\beta\langle\xi_h - \xi^*, \tilde{v}_h - A(\xi_h - \xi^*)\rangle \\ &+ 2\beta^2 \|\tilde{v}_h - A(\xi_h - \xi^*)\|^2 + 2L_A^2\beta^2 \|\xi_h - \xi^*\|^2, \end{split}$$

where (a) follows since  $A\xi_h - b = A\xi_h - A\xi^* = A(\xi_h - \xi^*)$ . Taking expectation conditioned on  $\mathcal{F}_h$  on both sides

$$\mathbb{E}_{h}[\|\xi_{h+1} - \xi^{*}\|^{2}] \leq (1 - \beta\lambda + 2L_{A}^{2}\beta^{2})\|\xi_{h} - \xi^{*}\|^{2} + 2\beta\langle\xi_{h} - \xi^{*}, \mathbb{E}_{h}[\tilde{v}_{h} - A(\xi_{h} - \xi^{*})]\rangle + 2\beta^{2}\mathbb{E}_{h}\|\tilde{v}_{h} - A(\xi_{h} - \xi^{*})\|^{2}.$$
(54)

Observe that

$$\begin{split} \|\tilde{v}_h - A(\xi_h - \xi^*)\|^2 &= \|(\tilde{A}_h - A)(\xi_h - \xi^*) + (\tilde{A}_h - A)(\xi^*) + (\tilde{b}_h - b)\|^2 \\ &\leq 3\|\tilde{A}_h - A\|^2\|\xi_h - \xi^*\|^2 + 3\|\tilde{A}_h - A\|^2\|\xi^*\|^2 + 3\|\tilde{b}_h - b\|^2 \\ &\leq 3\|\tilde{A}_h - A\|^2\|\xi_h - \xi^*\|^2 + 3\lambda^{-2}\|\tilde{A}_h - A\|^2 + 3\|\tilde{b}_h - b\|^2. \end{split}$$

Taking expectation and applying Lemma 7 yields

$$\mathbb{E}_{h} \|v_{h} - A(\xi_{h} - \xi^{*})\|^{2} \\
\leq 3 \mathbb{E}_{h} \|\tilde{A}_{h} - A\|^{2} \|\xi_{h} - \xi^{*}\|^{2} + 3\lambda^{-2} \mathbb{E}_{h} \|\tilde{A}_{h} - A\|^{2} + 3 \mathbb{E}_{h} \|\tilde{b}_{h} - b\|^{2} \\
\leq (6L_{A}^{2} \|\xi_{h} - \xi^{*}\|^{2} + 6\lambda^{-2}L_{A}^{2} + 6L_{b}^{2})(\tau_{\min} \log T_{\max} + \tau_{\min}^{2}).$$
(55)

Separately,

$$\langle \xi_{h} - \xi^{*}, \mathbb{E}_{h} [\tilde{v}_{h} - A(\xi_{h} - \xi^{*})] \rangle \leq \frac{\lambda}{4} \|\xi_{h} - \xi^{*}\|^{2} + \frac{1}{\lambda} \|\mathbb{E}_{h} [\tilde{v}_{h} - A(\xi_{h} - \xi^{*})]\|^{2}$$
  
 
$$\leq \frac{\lambda}{4} \|\xi_{h} - \xi^{*}\|^{2} + \frac{(L_{A}^{2} \|\xi_{h} - \xi^{*}\|^{2} + L_{b}^{2})\tau_{\text{mix}}^{2}}{\lambda^{2} T_{\text{max}}}$$
 (56)

Substituting the above bounds in (54), we obtain

$$\begin{split} & \mathbb{E}_{h}[\|\xi_{h+1} - \xi^{*}\|^{2}] \\ & \leq \left(1 - \frac{\beta\lambda}{2} + 6L_{A}^{2}\beta^{2}(\tau_{\min}\log T_{\max} + \tau_{\min}^{2}) + \frac{\beta L_{A}^{2}\tau_{\min}^{2}}{\lambda T_{\max}^{2}}\right)\|\xi_{h} - \xi^{*}\|^{2} + \frac{\beta L_{b}^{2}\tau_{\min}^{2}}{\lambda T_{\max}^{2}} \\ & + (6\beta^{2}\lambda^{-2}L_{A}^{2} + 6\beta^{2}L_{b}^{2})(\tau_{\min}\log T_{\max} + \tau_{\min}^{2}) \end{split}$$

For  $T_{\max}^2 \ge \tau_{\min}^2(\lambda\beta)^{-1}$  and  $\beta \le \lambda (28L_A^2(\tau_{\min}\log T_{\max} + \tau_{\min}^2))^{-1}$ , we have  $\frac{\beta\lambda}{2} + 8L_A^2\beta^2 \le 1 - \frac{\beta\lambda}{4}$ .

$$\mathbb{E}_{h}[\|\xi_{h+1} - \xi^{*}\|^{2}]$$

$$\leq \left(1 - \frac{\beta\lambda}{4}\right) \|\xi_{h} - \xi^{*}\|^{2} + \frac{\beta L_{b}^{2} \tau_{\min}^{2}}{\lambda T_{\max}^{2}} + (6\beta^{2}\lambda^{-2}L_{A}^{2} + 6\beta^{2}L_{b}^{2})(\tau_{\min}\log T_{\max} + \tau_{\min}^{2}).$$

Taking expectation on both sides and unrolling the recursion yields

$$\begin{split} & \mathbb{E}[\|\xi_{H} - \xi^{*}\|^{2}] \\ & \leq \left(1 - \frac{\beta\lambda}{4}\right)^{H} \mathbb{E}\left\|\xi_{0} - \xi^{*}\right\|^{2} + \sum_{h=0}^{H-1} \left(1 - \frac{\beta\lambda}{4}\right)^{H-h} \left((6\beta^{2}\lambda^{-2}L_{A}^{2} + 6\beta^{2}L_{b}^{2})(\tau_{\min}\log T_{\max} + \tau_{\min}^{2}) + \frac{\beta L_{b}^{2}\tau_{\min}^{2}}{\lambda T_{\max}^{2}}\right) \\ & \leq e^{-\frac{H\beta\lambda}{4}} \mathbb{E}\left\|\xi_{0} - \xi^{*}\right\|^{2} + \frac{4}{\beta\lambda} \left((6\beta^{2}\lambda^{-2}L_{A}^{2} + 6\beta^{2}L_{b}^{2})(\tau_{\min}\log T_{\max} + \tau_{\min}^{2}) + \frac{\beta L_{b}^{2}\tau_{\min}^{2}}{\lambda T_{\max}^{2}}\right) \\ & = e^{-\frac{H\beta\lambda}{4}} \mathbb{E}\left\|\xi_{0} - \xi^{*}\right\|^{2} + \left(24\beta(\lambda^{-3}L_{A}^{2} + 24L_{b}^{2}\lambda^{-1})(\tau_{\min}\log T_{\max} + \tau_{\min}^{2}) + \frac{L_{b}^{2}\tau_{\max}^{2}}{\lambda^{2}T_{\max}^{2}}\right). \end{split}$$

Set  $\beta = \frac{8 \log H}{\lambda H}$  to get

$$\mathbb{E}[\|\xi_H - \xi^*\|^2] \le \frac{1}{H^2} \mathbb{E} \|\xi_0 - \xi^*\|^2 + \mathcal{O}\left(\frac{\log H}{H}(\lambda^{-3}L_A^2 + L_b^2\lambda^{-1})(\tau_{\min}\log T_{\max} + \tau_{\min}^2) + \frac{L_b^2\tau_{\max}^2}{\lambda^2 T_{\max}^2}\right).$$

Towards proving Theorem 4(b), note that

$$\mathbb{E}[\tilde{v}_h|\theta] = \mathbb{E}[\mathbb{E}_h[\tilde{v}_h]|\theta] = \mathbb{E}[\mathbb{E}_h[\tilde{A}_h]\xi_h - \mathbb{E}[\tilde{b}_h|\theta] = \mathbb{E}_h[\tilde{A}_h]\mathbb{E}[\xi_h|\theta] - \mathbb{E}[\tilde{b}_h] \triangleq \bar{A}\mathbb{E}[\xi_h|\theta] - \bar{b}.$$
(57)

With the above,  $\{\mathbb{E}[\xi_h|\theta]\}_{0 \le h \le H}$  can be recursively expressed with the initialization  $\mathbb{E}[\xi_0|\theta] = \xi_0$  as follows:

$$\mathbb{E}[\xi_{h+1}|\theta] = \mathbb{E}[\xi_h|\theta] - \beta \mathbb{E}[\tilde{v}_h] = \mathbb{E}[\xi_h|\theta] - \beta(\bar{A} \mathbb{E}[\xi_h|\theta] - \bar{b}).$$
(58)

Using this update rule, we can bound  $\|\mathbb{E}[\xi_{h+1}|\theta] - \xi^*\|^2$  as:

$$\| \mathbb{E}[\xi_{h+1}|\theta] - \xi^* \|^2 = \| \mathbb{E}[\xi_h|\theta] - \beta \mathbb{E}[\tilde{v}_h|\theta] - \xi^* \|^2$$
  

$$= \| \mathbb{E}[\xi_h|\theta] - \xi^* \|^2 + 2\beta \langle \mathbb{E}[\xi_h|\theta] - \xi^*, \mathbb{E}[\tilde{v}_h|\theta] \rangle + \beta^2 \| \mathbb{E}[\tilde{v}_h|\theta] \|^2$$
  

$$\leq \| \mathbb{E}[\xi_h|\theta] - \xi^* \|^2 - \beta \lambda \| \mathbb{E}[\xi_h|\theta] - \xi^* \|^2 + 2\beta \langle \mathbb{E}[\xi_h|\theta] - \xi^*, \mathbb{E}[\tilde{v}_h - A(\xi_h - \xi^*)|\theta] \rangle$$
  

$$+ 2\beta^2 \| \mathbb{E}[\tilde{v}_h - A(\xi_h - \xi^*)|\theta] \|^2 + 2L_A^2\beta^2 \| \mathbb{E}[\xi_h|\theta] - \xi^* \|^2$$
  

$$= (1 - \lambda\beta + 2L_A\beta^2) \| \mathbb{E}[\xi_h|\theta] - \xi^* \|^2 + 2\beta \langle \mathbb{E}[\xi_h|\theta] - \xi^*, \mathbb{E}[\tilde{v}_h - A(\xi_h - \xi^*)|\theta] \rangle$$
  

$$+ 2\beta^2 \| \mathbb{E}[\tilde{v}_h - A(\xi_h - \xi^*)|\theta] \|^2.$$
(59)

Using Lemma 7, we obtain

$$\|\mathbb{E}[\tilde{v}_{h} - A(\xi_{h} - \xi^{*})|\theta]\|^{2} = \|(\bar{A} - A)(\mathbb{E}[\xi_{h}|\theta] - \xi^{*}) + (\bar{A} - A)(\xi^{*}) + (\bar{b} - b)\|^{2}$$

$$\leq 3\|\bar{A} - A\|^{2}\|\mathbb{E}[\xi_{h}|\theta] - \xi^{*}\|^{2} + 3\lambda^{-2}L_{A}^{2}\|\bar{A} - A\|^{2} + 3\|\bar{b} - b\|^{2}$$

$$\leq \frac{3L_{A}^{2}\tau_{\text{mix}}^{2}}{T_{\text{max}}^{2}}\|\mathbb{E}[\xi_{h}|\theta] - \xi^{*}\|^{2} + \frac{3L_{A}^{4}\tau_{\text{mix}}^{2}}{\lambda^{2}T_{\text{max}}^{2}} + \frac{3L_{b}^{2}\tau_{\text{mix}}^{2}}{T_{\text{max}}^{2}}.$$
(60)

Also,

$$\langle \mathbb{E}[\xi_{h}|\theta] - \xi^{*}, \mathbb{E}[\tilde{v}_{h} - A(\xi_{h} - \xi^{*})|\theta] \rangle \leq \frac{\lambda}{4} \| \mathbb{E}[\xi_{h}|\theta] - \xi^{*}\|^{2} + \frac{1}{\lambda} \| \mathbb{E}[\tilde{v}_{h} - A(\xi_{h} - \xi^{*})|\theta] \|^{2}$$

$$\leq \frac{\lambda}{4} \| \mathbb{E}[\xi_{h}|\theta] - \xi^{*}\|^{2} + \frac{1}{\lambda} \| \mathbb{E}[\tilde{v}_{h} - A(\xi_{h} - \xi^{*})|\theta] \|^{2}$$

$$\leq \frac{\lambda}{4} \| \mathbb{E}[\xi_{h}|\theta] - \xi^{*}\|^{2} + \frac{3L_{A}^{2}\tau_{\min}^{2}}{\lambda T_{\max}^{2}} \| \mathbb{E}[\xi_{h}|\theta] - \xi^{*}\|^{2}$$

$$+ \frac{3L_{A}^{4}\tau_{\min}^{2}}{\lambda^{3}T_{\max}^{2}} + \frac{3L_{b}^{2}\tau_{\max}^{2}}{\lambda T_{\max}^{2}}$$

$$(61)$$

Substituting the above bounds in (59) yields

$$\begin{split} \| \mathbb{E}[\xi_{h+1}|\theta] - \xi^* \|^2 \\ &\leq \left(1 - \frac{\beta\lambda}{2} + \frac{6\beta L_A^2 \tau_{\min}^2}{\lambda T_{\max}^2} + \frac{6\beta^2 L_A^2 \tau_{\min}^2}{T_{\max}^2}\right) \| \mathbb{E}[\xi_h|\theta] - \xi^* \|^2 + \frac{6\beta(1+\beta\lambda)L_A^4 \tau_{\min}^2}{\lambda^3 T_{\max}^2} + \frac{6\beta(1+\beta\lambda)L_b^2 \tau_{\min}^2}{\lambda T_{\max}^2} \\ &\leq \left(1 - \frac{\beta\lambda}{2} + \frac{12\beta L_A^2 \tau_{\min}^2}{\lambda T_{\max}^2}\right) \| \mathbb{E}[\xi_h|\theta] - \xi^* \|^2 + \frac{6\beta(1+\beta\lambda)L_A^4 \tau_{\min}^2}{\lambda^3 T_{\max}^2} + \frac{6\beta(1+\beta\lambda)L_b^2 \tau_{\min}^2}{\lambda T_{\max}^2}. \end{split}$$

For  $T_{\max}^2 \geq \frac{12L_A^2 \tau_{\max}^2}{\beta \lambda^2}$ , we have  $\left(1 - \frac{\beta \lambda}{2} + \frac{12\beta L_A^2 \tau_{\max}^2}{\lambda T_{\max}^2}\right) \leq \left(1 - \frac{\beta \lambda}{4}\right)$ . With this choice, the above bound becomes

$$\|\mathbb{E}[\xi_{h+1}|\theta] - \xi^*\|^2 \le \left(1 - \frac{\beta\lambda}{4}\right) \|\mathbb{E}[\xi_h|\theta] - \xi^*\|^2 + \frac{6\beta(1+\beta\lambda)L_A^4\tau_{\min}^2}{\lambda^3 T_{\max}^2} + \frac{6\beta(1+\beta\lambda)L_b^2\tau_{\min}^2}{\lambda T_{\max}^2}.$$

Unrolling the recursion

$$\begin{split} &\| \mathbb{E}[\xi_{H}|\theta] - \xi^{*} \|^{2} \\ &\leq \left(1 - \frac{\beta\lambda}{4}\right)^{H} \| \mathbb{E}[\xi_{0}|\theta] - \xi^{*} \|^{2} + \sum_{h=0}^{H-1} \left(1 - \frac{\beta\lambda}{4}\right)^{H-h} \left(\frac{6\beta(1 + \beta\lambda)L_{A}^{4}\tau_{\min}^{2}}{\lambda^{3}T_{\max}^{2}} + \frac{6\beta(1 + \beta\lambda)L_{b}^{2}\tau_{\min}^{2}}{\lambda T_{\max}^{2}}\right) \\ &\leq e^{-\frac{H\beta\lambda}{4}} \| \mathbb{E}[\xi_{0}|\theta] - \xi^{*} \|^{2} + \frac{4}{\beta\lambda} \left(\frac{6\beta(1 + \beta\lambda)L_{A}^{4}\tau_{\min}^{2}}{\lambda^{3}T_{\max}^{2}} + \frac{6\beta(1 + \beta\lambda)L_{b}^{2}\tau_{\min}^{2}}{\lambda T_{\max}^{2}}\right) \\ &= e^{-\frac{H\beta\lambda}{4}} \| \mathbb{E}[\xi_{0}|\theta] - \xi^{*} \|^{2} + \left(\frac{24(1 + \beta\lambda)L_{A}^{4}\tau_{\max}^{2}}{\lambda^{4}T_{\max}^{2}} + \frac{24(1 + \beta\lambda)L_{b}^{2}\tau_{\max}^{2}}{\lambda^{2}T_{\max}^{2}}\right). \end{split}$$

Set  $\beta = \frac{8 \log H}{\lambda H}$  to get

$$\|\mathbb{E}[\xi_{H}|\theta] - \xi^{*}\|^{2} \leq \frac{1}{H^{2}} \|\mathbb{E}[\xi_{0}|\theta] - \xi^{*}\|^{2} + \mathcal{O}\left(\frac{L_{A}^{4}\tau_{\min}^{2}}{\lambda^{4}T_{\max}^{2}} + \frac{L_{b}^{2}\tau_{\min}^{2}}{\lambda^{2}T_{\max}^{2}}\right).$$

## D Proof of Theorem 1

Recall that the global convergence of any update of form  $\theta_{k+1} = \theta_k + \alpha \omega_k$  can be bounded as

$$J^{*} - \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E}[J(\theta_{k})] \leq \sqrt{\epsilon_{\text{bias}}} + \frac{G_{1}}{K} \sum_{k=0}^{K-1} \mathbb{E} \| (\mathbb{E}[\omega_{k}|\theta_{k}] - \omega_{k}^{*}) \| + \frac{\alpha G_{2}}{K} \sum_{k=0}^{K-1} \mathbb{E} \| \omega_{k} - \omega_{k}^{*} \|^{2} + \frac{\alpha \mu^{-2}}{K} \sum_{k=0}^{K-1} \mathbb{E} \| \nabla_{\theta} J(\theta_{k}) \|^{2} + \frac{1}{\alpha K} \mathbb{E}_{s \sim d^{\pi^{*}}} [KL(\pi^{*}(\cdot|s) \| \pi_{\theta_{0}}(\cdot|s))].$$
(62)

We note that our algorithm updates  $\theta$  at each iteration k using  $\omega_H$  and  $\xi_H$  obtained after H iterations of the NPG and critic estimation inner loops. Therefore, we use  $\omega_H$  and  $\xi_H$  instead of  $\omega_k$  and  $\xi_k$ . We begin by

deriving a bound for  $\frac{1}{K}\sum_{k=0}^{K-1} \|\nabla_{\theta} J(\theta_k)\|^2$ . Given that the function *J* is *L*-smooth, we obtain:

$$J(\theta_{k+1}) \geq J(\theta_{k}) + \langle \nabla_{\theta}J(\theta_{k}), \theta_{k+1} - \theta_{k} \rangle - \frac{L}{2} \|\theta_{k+1} - \theta_{k}\|^{2}$$

$$= J(\theta_{k}) + \alpha \langle \nabla_{\theta}J(\theta_{k}), \omega_{k} \rangle - \frac{\alpha^{2}L}{2} \|\omega_{k}\|^{2}$$

$$= J(\theta_{k}) + \alpha \langle \nabla_{\theta}J(\theta_{k}), \omega_{k}^{*} \rangle + \alpha \langle \nabla_{\theta}J(\theta_{k}), \omega_{k} - \omega_{k}^{*} \rangle - \frac{\alpha^{2}L}{2} \|\omega_{k} - \omega_{k}^{*} + \omega_{k}^{*}\|^{2}$$

$$\stackrel{(a)}{\geq} J(\theta_{k}) + \alpha \langle \nabla_{\theta}J(\theta_{k}), F(\theta_{k})^{-1}\nabla_{\theta}J(\theta_{k}) \rangle + \alpha \langle \nabla_{\theta}J(\theta_{k}), \omega_{k} - \omega_{k}^{*} \rangle - \alpha^{2}L \|\omega_{k} - \omega_{k}^{*}\|^{2} - \alpha^{2}L \|\omega_{k}^{*}\|^{2}$$

$$\stackrel{(b)}{\geq} J(\theta_{k}) + \frac{\alpha}{G_{1}^{2}} \|\nabla_{\theta}J(\theta_{k})\|^{2} + \alpha \langle \nabla_{\theta}J(\theta_{k}), \omega_{k} - \omega_{k}^{*} \rangle - \alpha^{2}L \|\omega_{k} - \omega_{k}^{*}\|^{2} - \alpha^{2}L \|\omega_{k}^{*}\|^{2}$$

$$= J(\theta_{k}) + \frac{\alpha}{2G_{1}^{2}} \|\nabla_{\theta}J(\theta_{k})\|^{2} + \frac{\alpha}{2G_{1}^{2}} [\|\nabla_{\theta}J(\theta_{k})\|^{2} + 2G_{1}^{2} \langle \nabla_{\theta}J(\theta_{k}), \omega_{k} - \omega_{k}^{*} \rangle + G_{1}^{4} \|\omega_{k} - \omega_{k}^{*}\|^{2}]$$

$$= J(\theta_{k}) + \frac{\alpha}{2G_{1}^{2}} \|\nabla_{\theta}J(\theta_{k})\|^{2} + \frac{\alpha}{2G_{1}^{2}} \|\nabla_{\theta}J(\theta_{k}) + G_{1}^{2}(\omega_{k} - \omega_{k}^{*})\|^{2} - \left(\frac{\alpha G_{1}^{2}}{2} + \alpha^{2}L\right) \|\omega_{k} - \omega_{k}^{*}\|^{2}$$

$$= J(\theta_{k}) + \frac{\alpha}{2G_{1}^{2}} \|\nabla_{\theta}J(\theta_{k})\|^{2} - \left(\frac{\alpha G_{1}^{2}}{2G_{1}^{2}} + \alpha^{2}L\right) \|\omega_{k} - \omega_{k}^{*}\|^{2} - \alpha^{2}L \|F(\theta_{k})^{-1}\nabla_{\theta}J(\theta_{k})\|^{2}$$

$$\geq J(\theta_{k}) + \left(\frac{\alpha}{2G_{1}^{2}} - \frac{\alpha^{2}L}{\mu^{2}}\right) \|\nabla_{\theta}J(\theta_{k})\|^{2} - \left(\frac{\alpha G_{1}^{2}}{2} + \alpha^{2}L\right) \|\omega_{k} - \omega_{k}^{*}\|^{2}$$

where (*a*) utilizes the Cauchy-Schwarz inequality and the definition that  $\omega_k^* = F(\theta_k)^{-1} \nabla_{\theta} J(\theta_k)$ . Inequalities (*b*), and (*c*) follow from Assumption 5(a) and 6 respectively. We take the above inequality, sum over  $k = 0, \dots, K - 1$ , rearrange the terms and substitute  $\alpha = \frac{\mu^2}{4G_1^2L}$ , to obtain:

$$\frac{\mu^2}{16G_1^4 L} \left( \frac{1}{K} \sum_{k=0}^{K-1} \|\nabla_\theta J(\theta_k)\|^2 \right) \le \frac{J(\theta_K) - J(\theta_0)}{K} + \left( \frac{\mu^2}{8L} + \frac{\mu^4}{16G_1^4 L} \right) \left( \frac{1}{K} \sum_{k=0}^{K-1} \|\omega_k - \omega_k^*\|^2 \right)$$

$$\stackrel{(a)}{\le} \frac{2}{K} + \left( \frac{\mu^2}{8L} + \frac{\mu^4}{16G_1^4 L} \right) \left( \frac{1}{K} \sum_{k=0}^{K-1} \|\omega_k - \omega_k^*\|^2 \right)$$
(64)

where (a) uses the fact that  $J(\cdot)$  is absolutely bounded above by 1. Using (64), we obtain

$$\frac{\mu^{-2}}{K} \left( \sum_{k=0}^{K-1} \|\nabla_{\theta} J(\theta_k)\|^2 \right) \le \frac{32LG_1^4}{\mu^4 K} + \left( \frac{2G_1^4}{\mu^2} + 1 \right) \left( \frac{1}{K} \sum_{k=0}^{K-1} \|\omega_k - \omega_k^*\|^2 \right)$$
(65)

Now all that is left is to bound  $\mathbb{E}\left[\|\omega_H - \omega^*\|^2\right]$  and  $\|\mathbb{E}[\omega_H|\theta] - \omega^*\|$ . From Theorem 3 and 4, we have

$$\mathbb{E}\left[\|\omega_{H} - \omega^{*}\|^{2}\right] \leq \mathcal{O}\left(\frac{1}{H^{2}}\left[\|\omega_{0} - \omega^{*}\|^{2}\right] + \frac{\log H}{\mu^{2}H}\left(\sigma_{\omega}^{2}\tau_{\min}^{2}\right) + \mathbb{E}\left\|\xi_{H} - \xi^{*}\right\|^{2} + G_{1}^{2}\epsilon_{\mathrm{app}}\right)$$
(66)

and

$$\mathbb{E}[\|\xi_H - \xi^*\|^2] \le \frac{1}{H^2} \|\xi_0 - \xi^*\|^2 + \mathcal{O}\left(\frac{\log H}{H}(\lambda^{-3}L_A^2 + L_b^2\lambda^{-1})(\tau_{\min}\log T_{\max} + \tau_{\min}^2) + \frac{L_b^2\tau_{\min}^2}{\lambda^2 T_{\max}^2}\right).$$

This gives us

$$\mathbb{E}\left[\|\omega_{H} - \omega^{*}\|^{2}\right] \leq \mathcal{O}\left(\frac{1}{H^{2}}\left[\|\omega_{0} - \omega^{*}\|^{2} + \|\xi_{0} - \xi^{*}\|^{2}\right] + \frac{L_{b}^{2}\tau_{\min}^{2}}{\lambda^{2}T_{\max}^{2}} + G_{1}^{2}\epsilon_{\mathrm{app}} + \frac{\log H}{H}(\sigma_{\omega}^{2}\tau_{\min}^{2}\mu^{-2} + \lambda^{-3}L_{A}^{2} + L_{b}^{2}\lambda^{-1})(\tau_{\min}\log T_{\max} + \tau_{\min}^{2})\right).$$
(67)

Summing the above inequality from k = 0 to K - 1 and multiplying by  $\frac{\alpha G_2}{K}$  with  $\alpha = \frac{\mu^2}{4G_1^2 L}$  then yields

$$\frac{\alpha G_2}{K} \sum_{k=0}^{K-1} \mathbb{E} \left[ \|\omega_H - \omega^*\|^2 \right] \le \mathcal{O} \left( \frac{\mu^2}{G_1^2 L H^2} \left[ \|\omega_0 - \omega^*\|^2 + \|\xi_0 - \xi^*\|^2 \right] + \frac{G_2 \mu^2 L_b^2 \tau_{\min}^2}{G_1^2 L \lambda^2 T_{\max}^2} + \mu^2 L^{-1} G_2 \epsilon_{\text{app}} + \frac{\mu^2 G_2 \log H}{L G_1^2 H} (\sigma_\omega^2 \tau_{\min}^2 \mu^{-2} + \lambda^{-3} L_A^2 + L_b^2 \lambda^{-1}) (\tau_{\min} \log T_{\max} + \tau_{\min}^2) \right).$$

$$(68)$$

Again from Theorems 3 and 4, we have

$$\mathbb{E}\left[\|\mathbb{E}[\omega_{H}|\theta] - \omega^{*}\|^{2}\right] \leq \mathcal{O}\left(\frac{1}{H^{2}}[\|\omega_{0} - \omega^{*}\|^{2}] + \frac{G_{1}^{2}\tau_{\min}^{4}}{T_{\max}^{2}} + G_{1}^{2}\|\mathbb{E}[\xi_{H}|\theta] - \xi^{*}\|^{2} + G_{1}^{2}\epsilon_{\exp}\right)$$

and

$$\|\mathbb{E}[\xi_{H}|\theta] - \xi^{*}\|^{2} \leq \frac{1}{H^{2}} \|\xi_{0} - \xi^{*}\|^{2} + \mathcal{O}\left(\frac{L_{A}^{4}\tau_{\min}^{2}}{\lambda^{4}T_{\max}^{2}} + \frac{L_{b}^{2}\tau_{\min}^{2}}{\lambda^{2}T_{\max}^{2}}\right).$$

This gives us

$$\mathbb{E}\left[\|\mathbb{E}[\omega_{H}|\theta] - \omega^{*}\|^{2}\right] \leq \mathcal{O}\left(\frac{1}{H^{2}}\left[\|\omega_{0} - \omega^{*}\|^{2} + \|\xi_{0} - \xi^{*}\|^{2}\right] + \frac{L_{A}^{4}\tau_{\min}^{2}}{\lambda^{4}T_{\max}^{2}} + \frac{L_{b}^{2}\tau_{\min}^{2}}{\lambda^{2}T_{\max}^{2}} + \frac{G_{1}^{2}\tau_{\min}^{4}}{T_{\max}^{2}} + G_{1}^{2}\epsilon_{\mathrm{app}}\right).$$

Taking square root on both sides yields

$$\mathbb{E}\left[\left\|\mathbb{E}[\omega_{H}|\theta] - \omega^{*}\right\|\right] \leq \mathcal{O}\left(\frac{1}{H}\left[\left\|\omega_{0} - \omega^{*}\right\| + \left\|\xi_{0} - \xi^{*}\right\|\right] + \frac{L_{A}^{2}\tau_{\min}}{\lambda^{2}T_{\max}} + \frac{L_{b}\tau_{\min}}{\lambda T_{\max}} + \frac{G_{1}\tau_{\min}^{2}}{T_{\max}} + G_{1}\sqrt{\epsilon_{\mathrm{app}}}\right)$$

Consequently, we can bound  $\frac{G_1}{K} \sum_{k=0}^{K-1} \mathbb{E} \left\| \left( \mathbb{E} \left[ \omega_k | \theta_k \right] - \omega_k^* \right) \right\|$  as

$$\frac{G_1}{K} \sum_{k=0}^{K-1} \mathbb{E} \left\| \left( \mathbb{E} \left[ \omega_k | \theta_k \right] - \omega_k^* \right) \right\| \\
\leq \mathcal{O} \left( \frac{G_1}{H} \left[ \left\| \omega_0 - \omega^* \right\| + \left\| \xi_0 - \xi^* \right\| \right] + \frac{G_1 L_A^2 \tau_{\text{mix}}}{\lambda^2 T_{\text{max}}} + \frac{G_1 L_b \tau_{\text{mix}}}{\lambda T_{\text{max}}} + \frac{G_1^2 \tau_{\text{mix}}^2}{T_{\text{max}}} + G_1^2 \sqrt{\epsilon_{\text{app}}} \right)$$

Substituting these bounds in Lemma 1 and setting  $K = \Theta(\sqrt{T})$ ,  $H = \Theta(\sqrt{T}/\log(T))$ ,  $B = \Theta(\log(T))$ ,  $T_{\max} = \Theta(\sqrt{T})$  and  $\alpha = \frac{\mu^2}{4G_1^2 L}$ , we obtain the following bound. We highlight that the  $G_1^2$  factor in the function approximation error term is a standard component in actor-critic results with a linear critic [Suttle et al., 2023, Patel et al., 2024]. However, this factor is often not explicitly mentioned in previous works, whereas we have included it here for completeness.

$$J^{*} - \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E}[J(\theta_{k})] \leq \mathcal{O}\left(\sqrt{\epsilon_{\text{bias}}} + G_{1}^{2}\sqrt{\epsilon_{\text{app}}} + \frac{G_{1}}{\sqrt{T}} \left[ \|\omega_{0} - \omega^{*}\| + \|\xi_{0} - \xi^{*}\| \right] + \frac{G_{1}L_{A}^{2}\tau_{\text{mix}}}{\lambda^{2}\sqrt{T}} + \frac{G_{1}^{2}\tau_{\text{mix}}^{2}}{\sqrt{T}} + \frac{\mu^{2}}{G_{1}^{2}LT} \left[ \|\omega_{0} - \omega^{*}\|^{2} + \|\xi_{0} - \xi^{*}\|^{2} \right] + \frac{G_{2}\mu^{2}L_{b}^{2}\tau_{\text{mix}}^{2}}{G_{1}^{2}L\lambda^{2}T} + \mu^{2}L^{-1}G_{2}\epsilon_{\text{app}} + \frac{G_{1}L_{b}\tau_{\text{mix}}}{\lambda\sqrt{T}} + \frac{\mu^{2}G_{2}\log T}{LG_{1}^{2}\sqrt{T}} (\sigma_{\omega}^{2}\tau_{\text{mix}}^{2}\mu^{-2} + \lambda^{-3}L_{A}^{2} + L_{b}^{2}\lambda^{-1})(\tau_{\text{mix}}\log T + \tau_{\text{mix}}^{2}) \right)$$

$$(69)$$

### E Proof of Lemma 1

The proof of the lemma is straightforward and follows a similar structure to the one in [Mondal and Aggarwal, 2024]. The key distinction is that the bound derived in [Mondal and Aggarwal, 2024] applies to the discounted setting, whereas our derivation pertains to the average-reward case. We begin by stating a useful lemma:

**Lemma 8** (Lemma 4, [Bai et al., 2024]). *The difference in the performance for any policies*  $\pi_{\theta}$  *and*  $\pi_{\theta'}$  *is bounded as follows* 

$$J(\theta) - J(\theta') = \mathbb{E}_{s \sim d^{\pi_{\theta}}} \mathbb{E}_{a \sim \pi_{\theta}}(\cdot|s) \left[ A^{\pi_{\theta'}}(s,a) \right].$$
(70)

Continuing with the proof, we have:

$$\begin{split} &\mathbb{E}_{s\sim d^{\pi^*}} [KL(\pi^*(\cdot|s) \| \pi_{\theta_k}(\cdot|s)) - KL(\pi^*(\cdot|s) \| \pi_{\theta_{k+1}}(\cdot|s))] \\ &= \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} \left[ \log \frac{\pi_{\theta_{k+1}(a|s)}}{\pi_{\theta_k}(a|s)} \right] \\ \stackrel{(a)}{\cong} \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} [\nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot (\theta_{k+1} - \theta_k)] - \frac{G_2}{2} \| \theta_{k+1} - \theta_k \|^2 \\ &= \alpha \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} [\nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot \omega_k] - \frac{G_2 \alpha^2}{2} \| \omega_k \|^2 \\ &= \alpha \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} [\nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot \omega_k^*] + \alpha \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} [\nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot (\omega_k - \omega_k^*)] \\ &- \frac{G_2 \alpha^2}{2} \| \omega_k \|^2 \\ &= \alpha [J^* - J(\theta_k)] + \alpha \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} [\nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot \omega_k^*] - \alpha [J^* - J(\theta_k)] \\ &+ \alpha \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} [\nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot (\omega_k - \omega_k^*)] - \frac{G_2 \alpha^2}{2} \| \omega_k \|^2 \\ \stackrel{(b)}{\bigoplus} \alpha [J^* - J(\theta_k)] + \alpha \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} \left[ \nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot \omega_k^* - A^{\pi_{\theta_k}}(s, a) \right] \\ &+ \alpha \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} [\nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot (\omega_k - \omega_k^*)] - \frac{G_2 \alpha^2}{2} \| \omega_k \|^2 \\ \stackrel{(c)}{\cong} \alpha [J^* - J(\theta_k)] - \alpha \sqrt{\mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} \left[ \left( \nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot \omega_k^* - A^{\pi_{\theta_k}}(s, a) \right)^2 \right]} \\ &+ \alpha \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} [\nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot (\omega_k - \omega_k^*)] - \frac{G_2 \alpha^2}{2} \| \omega_k \|^2 \\ \stackrel{(c)}{\cong} \alpha [J^* - J(\theta_k)] - \alpha \sqrt{\mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} \left[ \left( \nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot \omega_k^* - A^{\pi_{\theta_k}}(s, a) \right)^2 \right]} \\ &+ \alpha \mathbb{E}_{s\sim d^{\pi^*}} \mathbb{E}_{a\sim\pi^*(\cdot|s)} [\nabla_{\theta} \log \pi_{\theta_k}(a|s) \cdot (\omega_k - \omega_k^*)] - \frac{G_2 \alpha^2}{2} \| \omega_k \|^2 . \end{aligned}$$

Here, (a) and (b) stem from Assumption 5(b) and Lemma 8, respectively. Inequality (c) arises from the convexity of the function  $f(x) = x^2$ . Lastly, (d) is a consequence of Assumption 4. By taking expectations on both sides, we derive:

$$\mathbb{E}\left[\mathbb{E}_{s\sim d^{\pi^{*}}}\left[KL(\pi^{*}(\cdot|s)\|\pi_{\theta_{k}}(\cdot|s)) - KL(\pi^{*}(\cdot|s)\|\pi_{\theta_{k+1}}(\cdot|s))\right]\right] \\
\geq \alpha[J^{*} - \mathbb{E}\left[J(\theta_{k})\right]] - \alpha\sqrt{\epsilon_{\text{bias}}} \\
+ \alpha \mathbb{E}\left[\mathbb{E}_{s\sim d^{\pi^{*}}}\mathbb{E}_{a\sim\pi^{*}}(\cdot|s)\left[\nabla_{\theta}\log\pi_{\theta_{k}}(a|s) \cdot \left(\mathbb{E}[\omega_{k}|\theta_{k}] - \omega_{k}^{*}\right)\right]\right] - \frac{G_{2}\alpha^{2}}{2}\mathbb{E}\left[\|\omega_{k}\|^{2}\right] \\
\geq \alpha[J^{*} - \mathbb{E}\left[J(\theta_{k})\right]\right] - \alpha\sqrt{\epsilon_{\text{bias}}} \\
- \alpha \mathbb{E}\left[\mathbb{E}_{s\sim d^{\pi^{*}}}\mathbb{E}_{a\sim\pi^{*}}(\cdot|s)\left[\|\nabla_{\theta}\log\pi_{\theta_{k}}(a|s)\|\|\mathbb{E}[\omega_{k}|\theta_{k}] - \omega_{k}^{*}\|\right]\right] - \frac{G_{2}\alpha^{2}}{2}\mathbb{E}\left[\|\omega_{k}\|^{2}\right] \\
\overset{(a)}{\geq}\alpha[J^{*} - \mathbb{E}\left[J(\theta_{k})\right]\right] - \alpha\sqrt{\epsilon_{\text{bias}}} - \alpha G_{1}\mathbb{E}\left\|(\mathbb{E}[\omega_{k}|\theta_{k}] - \omega_{k}^{*})\| - \frac{G_{2}\alpha^{2}}{2}\mathbb{E}\left[\|\omega_{k}\|^{2}\right]$$
(71)

where (a) follows from Assumption 5(a). Rearranging the terms, we get,

$$J^* - \mathbb{E}[J(\theta_k)] \leq \sqrt{\epsilon_{\text{bias}}} + G_1 \mathbb{E} \left\| \left( \mathbb{E}[\omega_k | \theta_k] - \omega_k^* \right) \right\| + \frac{G_2 \alpha}{2} \mathbb{E} \left\| \omega_k \right\|^2 + \frac{1}{\alpha} \mathbb{E} \left[ \mathbb{E}_{s \sim d^{\pi^*}} \left[ KL(\pi^*(\cdot | s) \| \pi_{\theta_k}(\cdot | s)) - KL(\pi^*(\cdot | s) \| \pi_{\theta_{k+1}}(\cdot | s)) \right] \right]$$
(72)

Adding the above inequality from k = 0 to K - 1, using the non-negativity of KL divergence and dividing the resulting expression by K, we obtain the final result.