

# BIRATIONAL GEOMETRY OF FANO VARIETIES OF LINES ON CUBIC FOURFOLDS CONTAINING PAIRS OF CUBIC SCROLLS

COREY BROOKE, SARAH FREI, LISA MARQUAND, AND XUQIANG QIN

ABSTRACT. We characterize the birational geometry of some hyperkähler fourfolds of Picard rank 3 obtained as the Fano varieties of lines on cubic fourfolds containing pairs of cubic scrolls. In each of the two cases considered, we provide a census of the birational models, relating each model to familiar geometric constructions. We also provide structural results about the birational automorphism groups, giving generators in both cases and a full set of relations in one case. Finally, as a byproduct of our analysis, we obtain non-isomorphic cubic fourfolds whose Fano varieties of lines are birationally equivalent.

## 1. INTRODUCTION

Cubic fourfolds are a central object in algebraic geometry, studied with respect to rationality questions and for their connections to hyperkähler manifolds. In a very general cubic fourfold, any algebraic surface is homologous to a complete intersection, but a countably infinite collection of divisors  $\mathcal{C}_d$  in the moduli space of cubic fourfolds parametrize cubics containing extra algebraic surfaces [Has00]. Cubics contained in these divisors are often more interesting from the point of view of rationality questions: it is conjectured that the rational cubic fourfolds are contained in the union of certain divisors  $\mathcal{C}_d$ . The Fano variety of lines  $F$  on a cubic fourfold  $X$  is a hyperkähler manifold by [BD85], and if  $X \in \mathcal{C}_d$  then  $F$  exhibits richer birational geometry, which can be studied via the Global Torelli theorem (due to Huybrechts, Markman and Verbitsky). Here, we focus on the Fano varieties of lines on cubic fourfolds belonging to the family  $\mathcal{C}_{12}$ , whose general member contains a cubic scroll.

The birational geometry of the Fano variety of lines  $F$  of a very general cubic  $X \in \mathcal{C}_{12}$  was first studied in [HT10], where they exhibited a birational transformation of infinite order. Building on their results, we prove the following:

**Theorem 1.1.** *Let  $F$  be the Fano variety of lines of a very general member  $X \in \mathcal{C}_{12}$ . Then  $F$  has three isomorphism classes of birational hyperkähler models, represented by  $F$  itself and two non-isomorphic Mukai flops, both of which are isomorphic to double EPW sextics. Moreover,  $\text{Bir}(F)$  is generated by the covering involutions on these two double EPW sextics.*

In addition to amending Theorem 7.4 of [HT10], the novelty of this result is the connection between  $F$  and double EPW sextics. In [DIM15] (see also [KP18, Proposition 5.6]), the authors show that a general  $X \in \mathcal{C}_{12}$  is birational to a Gushel-Mukai fourfold  $Z$  containing a plane. The fourfold  $Z$  in turn determines a double EPW sextic, a hyperkähler manifold introduced by O’Grady ([O’G05, O’G06, O’G08b]). In [IM11], the authors show these manifolds can be constructed from considering conics on the Gushel-Mukai fourfold. Combining both the birational map between

$X$  and  $Z$  and this geometric description in terms of conics on  $Z$ , we construct birational maps between  $F$  and two non-isomorphic double EPW sextics explicitly.

The divisor  $\mathcal{C}_{12}$  contains many geometrically interesting family of cubic fourfolds. For example, there is a ten dimensional family of cubics admitting an involution fixing a line pointwise, denoted by  $\mathcal{M}_{\phi_2}$  in the notation of [Mar23]. For  $X \in \mathcal{M}_{\phi_2}$ , the middle algebraic cohomology of  $X$  is spanned by classes represented by cubic scrolls, along with the square of the hyperplane class. Any pair of cubic scrolls spanning different hyperplane sections of  $X$  is either **syzygetic** or **non-syzygetic**, meaning they intersect in three or one points, respectively (Definition 2.9). Further, such a cubic fourfold is conjecturally irrational [Mar23, Theorem 1.2], and one could hope this is reflected in the birational geometry of the associated hyperkähler manifolds.

Motivated by this, we study the birational geometry of the Fano variety  $F$  of lines on a very general cubic fourfold  $X$  containing either a syzygetic or a non-syzygetic pair of cubic scrolls. In both cases,  $F$  has Picard rank three, and we describe the movable and ample cones of  $F$  by identifying the wall divisors following techniques from [Mon15]. We first consider a cubic fourfold  $X$  with a syzygetic pair of cubic scrolls — in this case, the movable cone is bounded by infinitely many walls corresponding to prime exceptional divisors (see Section 2.4). We prove the following result:

**Theorem 1.2.** *Let  $F$  be the Fano variety of lines on a very general cubic fourfold  $X$  containing a syzygetic pair of cubic scrolls. Then  $F$  has **five** isomorphism classes of birational hyperkähler models, represented by the following:*

- $F$  itself, and
- four non-isomorphic Mukai flops of  $F$ , each isomorphic to a double EPW sextic.

In the non-syzygetic case, the movable cone coincides with the positive cone, but surprisingly the birational geometry of  $F(X)$  is more complicated. This is partially explained by the fact that cubics containing a non-syzygetic pair of cubic scrolls automatically contain a third family of cubic scrolls (see Lemma 2.11).

**Theorem 1.3.** *Let  $F$  be the Fano variety of lines on a very general cubic fourfold  $X$  containing a non-syzygetic pair of cubic scrolls. Then  $F$  has **eight** isomorphism classes of birational hyperkähler models, represented by the following:*

- $F$  itself,
- six non-isomorphic Mukai flops of  $F$ , each isomorphic to a double EPW sextic, and
- a Mukai flop of  $F$  along a pair of disjoint planes in  $F$ , isomorphic to the Fano variety of lines on another cubic fourfold containing a non-syzygetic pair of cubic scrolls.

As a consequence, we obtain examples of the following phenomenon:

**Corollary 1.4.** *There exist non-isomorphic cubic fourfolds with birationally equivalent Fano varieties of lines.*

These results are of interest for two main reasons. First, they provide examples of hyperkähler fourfolds of Picard rank three whose birational geometry is explicitly understood; to our knowledge, this has only been done in the significantly easier

case of Picard rank two. Second, our techniques also allow us to deduce information about the birational automorphism group of  $F$ , even though understanding the structure of  $\text{Bir}(F)$  is in general more difficult than enumerating the birational models. Specifically, we obtain the following structural result for  $\text{Bir}(F)$ :

**Theorem 1.5.** *Let  $F$  be the Fano variety of lines on  $X$ , a very general cubic fourfold containing a syzygetic or non-syzygetic pair of cubic scrolls. Then:*

- (1) *The infinite order group  $\text{Bir}(F)$  is generated by the covering involutions of the double EPW sextics obtained as Mukai flops of  $F$ .*
- (2) *In particular, in the syzygetic case we have:*

$$\text{Bir}(F) \cong \langle a, b, c, d \mid a^2 = b^2 = c^2 = d^2 = 1 \rangle.$$

**Outline.** In Section 2, we recall the relevant definitions and results about cubic threefolds and fourfolds containing cubic scrolls, the Fano variety of lines on such a cubic fourfold, and the birational geometry of hyperkähler manifolds of K3<sup>[2]</sup>-type. We also introduce Gushel-Mukai fourfolds and double EPW sextics. In Section 3, we study the birational geometry of the Fano variety of lines  $F$  on a very general cubic fourfold  $X$  containing a cubic scroll, proving Theorem 1.1. In Sections 4 and 5, we prove Theorems 1.2 and 1.3 (Theorems 4.1 and 5.1), respectively, as well as Theorem 1.5 (Theorems 4.2 and 5.2), by carrying out an analysis of the chambers of the movable cone of  $F$  when  $X$  contains a syzygetic and non-syzygetic, respectively, pair of cubic scrolls. Explicit examples in Appendix A illustrate generic behavior of such cubic fourfolds.

**Acknowledgments.** We thank Nicolas Addington, Brendan Hassett, Alex Perry, Jack Petok, and Yuri Tschinkel for valuable conversations. In particular, Nicolas Addington suggested the possible connection to double EPW sextics explored in Section 3.3, and Brendan Hassett suggested techniques that streamlined our arguments enumerating the birational models of  $F$ . The computations in Appendix A were done in Magma [BCP97].

This material is based upon work supported by the NSF Grant DMS-1929284 while the authors were in residence at the Institute for Computational and Experimental Research in Mathematics in Providence, RI, during the Hyperkähler Manifolds and Special Cubic Fourfolds Collaborate@ICERM Program. The authors were also partially supported by the Hausdorff Research Institute for Mathematics, while in residence for the Junior Trimester Program on Algebraic geometry: derived categories, Hodge theory, and Chow groups, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2047/1 – 390685813. S.F. was supported in part by an AMS-Simons travel grant.

## 2. PRELIMINARIES

We consider complex cubic threefolds and fourfolds that contain rational normal cubic scrolls, hereafter referred to as *cubic scrolls*.

**Definition 2.1.** A *cubic scroll*  $T \subset \mathbb{P}^4$  is a non-degenerate surface of degree 3. Equivalently,  $T$  is isomorphic to the blow up of  $\mathbb{P}^2$  in a point embedded into  $\mathbb{P}^4$ .

In Section 2.1 we recall generalities on nodal cubic threefolds  $Y \subset \mathbb{P}^4$  that contain cubic scrolls. In particular, we recall the description of the components of the Fano

variety of lines  $F(Y)$  for such a threefold due to [HT10]. In Section 2.2, we focus on smooth cubic fourfolds  $X \subset \mathbb{P}^5$  containing cubic scrolls, defining the notion of syzygetic and non-syzygetic pairs of scrolls (Definition 2.9). In Section 2.3, we describe the Abel-Jacobi map relating the cohomology of  $X$  to its Fano variety of lines. In Section 2.4, we describe the structure of the movable and ample cone of a hyperkähler manifold of K3<sup>[2]</sup>-type (Theorem 2.17). Finally, in Section 2.5 we recall the relevant results on Gushel-Mukai fourfolds and associated double EPW sextics, emphasizing a geometric point of view.

**2.1. Cubic threefolds containing cubic scrolls.** Let  $Y$  be a cubic threefold containing a cubic scroll  $T$ , as studied in detail in [HT10]. A generic such cubic threefold has six nodes in general position. Conversely, any cubic threefold with six nodes in general position contains a cubic scroll. We summarize some of their geometric properties in this section.

By [HT10, Proposition 4.7],  $Y$  contains another cubic scroll  $T^\vee$  residual to  $T$  in a quadric, and two nets of cubic scrolls homologous to  $T$  and  $T^\vee$  respectively. Each node of  $Y$  is contained in all of the cubic scrolls in  $Y$ , and the complete linear system  $|\mathcal{I}_{T/Y}(2)|$  induces a rational map  $q: Y \dashrightarrow \mathbb{P}^2$ , resolved by passing to a small resolution  $Y^+$  of  $Y$  as in the diagram below.

$$(2.2) \quad \begin{array}{ccc} & Y^+ & \\ f \swarrow & & \searrow p \\ Y & \xrightarrow{\quad q \quad} & \mathbb{P}^2 \end{array}$$

As noted in [CTZ24, Remark 7.1], the map  $p$  is a  $\mathbb{P}^1$ -bundle, and for a general line  $\ell \subset \mathbb{P}^2$ ,  $f_*p^*(\ell)$  is a cubic scroll homologous to  $T^\vee$ .

*Remark 2.3.* If  $T' \subset Y$  is a smooth cubic scroll homologous to  $T^\vee$ , there are two types of lines in  $T'$ : an exceptional line  $E$  with  $E^2 = -1$  under the intersection form and a pencil of lines  $L$  with  $L^2 = 0$ . The first type is a section of the  $\mathbb{P}^1$ -bundle  $q|_{T'}: T' \rightarrow \mathbb{P}^1$  whereas the second is a fiber.

Similarly, the linear system  $|\mathcal{I}_{T^\vee/Y}(2)|$  induces a map  $q^\vee: Y \dashrightarrow \mathbb{P}^2$  resolved by a small resolution  $Y^-$  which differs from  $Y^+$  by six Atiyah flops [CTZ24, Section 7].

The Fano variety of lines on  $Y$  has three components:

**Proposition 2.4.** [HT10, Proposition 4.1] *The Fano variety of lines  $F(Y)$  on  $Y$  decomposes as  $P \cup S' \cup P^\vee$  where  $P, P^\vee \simeq \mathbb{P}^2$  and  $S'$  is a singular surface whose normalization is a smooth cubic surface.*

*Remark 2.5.* We summarize another account of the components of  $F(Y)$ , as outlined in [Dol16]. Given a node  $y \in Y$ , a union of twisted cubic curves  $C_y$  and  $C_y^\vee$  parametrizes the lines on  $Y$  passing through  $y$  [HT10, Lemma 4.4]. Hence the surface in  $Y$  swept out by the lines through  $y$  decomposes as a union of cones  $A_y \cup A_y^\vee$  over twisted cubics; by [HT10, Proposition 4.7], the surfaces  $A_y$  and  $A_y^\vee$  are homologous to  $T$  and  $T^\vee$ , respectively. One obtains a birational map

$$\mathrm{Hilb}^2(C_y \cup C_y^\vee) \dashrightarrow F(Y)$$

by taking the residual line to the two lines determined by  $\xi \in \mathrm{Hilb}^2(C_y \cup C_y^\vee)$ . Considering instead bisecant lines, which intersect  $C_y \cup C_y^\vee$  with multiplicity at

least 2, Dolgachev upgrades this to an isomorphism

$$\text{Bis}(C_y \cup C_y^\vee) \xrightarrow{\sim} F(Y)$$

in [Dol16, Section 4]. The planes  $P, P^\vee \subset F(Y)$  can be identified with  $\text{Bis}(C_y)$  and  $\text{Bis}(C_y^\vee)$ , i.e. the set of lines bisecant to  $C_y$  and  $C_y^\vee$ , respectively. The surface  $S'$  consists of lines meeting both  $C_y$  and  $C_y^\vee$ .

**2.2. Special cubic fourfolds in  $\mathcal{C}_{12}$ .** For a smooth cubic fourfold  $X$ , we define the lattice of algebraic cycles

$$A(X) = H^4(X, \mathbb{Z}) \cap H^{2,2}(X, \mathbb{C}),$$

and letting  $\eta_X$  be the square of the hyperplane class, we also define

$$H^4(X, \mathbb{Z})_{\text{prim}} = \langle \eta_X \rangle^\perp \subset H^4(X, \mathbb{Z})$$

and

$$A(X)_{\text{prim}} = H^4(X, \mathbb{Z})_{\text{prim}} \cap A(X).$$

Each of the above is a lattice under the intersection pairing. For a very general cubic fourfold,  $A(X)_{\text{prim}} = 0$ , but countably many divisors  $\mathcal{C}_d$  in the moduli space of cubic fourfolds parametrize those with  $A(X)_{\text{prim}} \neq 0$ . Whenever  $\eta_X \in K \subset A(X)$  where  $K$  is a primitive sublattice of rank two, and the discriminant of the intersection form on  $K$  is  $d$ , we say  $X \in \mathcal{C}_d$ , and  $K$  is called a labeling of  $X$ .

We focus in particular on smooth cubic fourfolds  $X \subset \mathbb{P}^5$  that contain a cubic scroll  $T$ , or equivalently contain a hyperplane section  $Y = X \cap H$  with six nodes in general position [HT10, Prop. 6.1]. A cubic scroll  $T \subset X$  determines a labelling  $K_{12} \hookrightarrow A(X)$  (see [Has00, Section 4.13]), so  $X \in \mathcal{C}_{12}$ . Indeed, the intersection pairing on  $\langle \eta_X, T \rangle \subset A(X)$  is given by:

	$\eta_X$	$T$
$\eta_X$	3	3
$T$	3	7

As a consequence of Section 2.1, we immediately see the following:

**Lemma 2.6.** [Has96, Lemma 2.11] *Let  $X$  be a general cubic fourfold containing a cubic scroll  $T$ . Then:*

- (1) *there is a net of scrolls homologous to  $T$ , all contained in  $H$ ,*
- (2) *there exists a residual cubic scroll  $T^\vee \subset X$  spanning the same hyperplane as  $T$  such that  $[T] + [T^\vee] = 2\eta_X \in A(X)$ .*

In much of what follows, we take  $X$  to contain two cubic scrolls  $T_1$  and  $T_2$  spanning different hyperplanes  $H_1$  and  $H_2$ . For a very general such  $X$ , the classes  $\eta_X$ ,  $[T_1]$ , and  $[T_2]$  span  $A(X)$ , and the intersection form is given by

	$\eta_X$	$T_1$	$T_2$
$\eta_X$	3	3	3
$T_1$	3	7	$\tau$
$T_2$	3	$\tau$	7

where  $\tau \in \{1, 2, 3, 4, 5\}$  [Mar23, Lemma 4.11]. Note that if  $[T_1] \cdot [T_2] = \tau$ , then  $[T_1^\vee] \cdot [T_2] = 6 - \tau$  since  $[T_1] + [T_1^\vee] = 2\eta_X$ ; hence we can take  $\tau \in \{1, 2, 3\}$ .

*Remark 2.7.* Note that there is a small error in [Mar23, Lemma 4.11], which originally ruled out  $\tau = 2, 4$ : such an intersection gives a vector  $v = 2\eta_X - [T_1] - [T_2]$  with  $v^2 = 6$ , but  $\text{div}(v) = 1$  in the full primitive cohomology  $H^4(X, \mathbb{Z})$ .

**Lemma 2.8.** *Let  $X \subset \mathbb{P}^5$  be a general cubic fourfold containing two cubic scrolls  $T_1, T_2$  spanning distinct hyperplanes  $H_1, H_2 \subset \mathbb{P}^5$ . Then the cubic surface  $\Sigma := H_1 \cap H_2 \cap X$  is smooth.*

*Proof.* We provide examples of such cubics in Appendix A when  $\tau = 1, 3$  and since smoothness is an open condition, the result follows in those cases. A similar example proves the claim for  $\tau = 2$  but is not included because we do not treat such cubics in this paper.  $\square$

Thus, one can interpret the intersection of  $T_1$  and  $T_2$  in terms of the intersection of two twisted cubics on  $\Sigma = H_1 \cap H_2 \cap X$ , (see [Mar23, Appendix B]). This observation motivates the following definition:

**Definition 2.9.** We say that  $T_1$  and  $T_2$  are *syzygetic scrolls* provided  $[T_1] \cdot [T_2] = 3$ , and *azygetic scrolls* provided  $[T_1] \cdot [T_2] = 2$  or  $4$ . If  $[T_1] \cdot [T_2] = 1$  or  $5$  we say the pair is *non-syzygetic*.

Indeed, the associated twisted cubics form a syzygetic duad in the first case (see [Dol12, Lemma 9.1.5]), and an azygetic pair in the second.

In the non-syzygetic case, there are exactly two other algebraic cycles with the same numerics as a cubic scroll, namely:

$$(2.10) \quad \begin{aligned} [T_3] &= 3\eta_X - [T_1] - [T_2], \\ [T_3^\vee] &= [T_1] + [T_2] - \eta_X. \end{aligned}$$

**Lemma 2.11.** *The classes  $[T_3]$  and  $[T_3^\vee]$  are represented by cubic scrolls in  $X$ .*

*Proof.* Let  $\mathcal{X} \rightarrow B$  be a local universal family of marked cubic fourfolds, where we identify each lattice  $A(X_b)_{\text{prim}}$  with a sublattice of  $H^4(X_0, \mathbb{Z})_{\text{prim}}$ , with  $X_0 = X$ . We let  $B' \subset B$  denote the Hodge locus of the class  $[T_3]$ ; i.e. the locus parametrizing fibers  $\mathcal{X}_b$  where  $[T_3]$  remains algebraic, so  $[T_3] \in A(X_b)$ . If  $b \in B'$  is very general, then the cubic  $X_b$  has  $A(X_b) \cong \langle \eta_{X_b}, [T_3] \rangle$  and  $X_b$  contains a cubic scroll  $T'$ , necessarily with class  $[T_3]$  [Has96, Section 4.1.2]. Then by specialization, in  $X$  the class  $[T_3]$  is also represented by an effective cycle of degree 3 in  $\mathbb{P}^5$ . Since  $X$  does not contain a plane, this is necessarily irreducible, and since  $[T_3]^2 = 7$  the only option is for  $[T_3]$  to be represented by a cubic scroll. Since  $[T_3^\vee] = 2\eta_X - [T_3]$ , we know  $[T_3^\vee]$  is also represented by a cubic scroll.  $\square$

We will also make use of the following fact:

**Proposition 2.12.** *For  $X$  a general cubic fourfold in  $\mathcal{C}_{12}$  or a general cubic fourfold containing a pair of cubic scrolls,  $\text{Aut}(X) = 0$ .*

*Proof.* The locus of cubic fourfolds with an order  $p$  automorphism has dimension at most dimension 14 (see [GAL11, Theorem 3.8]). It follows that a general  $X$  is not contained in any of these loci.  $\square$

*Remark 2.13.* Our main motivation for studying the cubics containing syzygetic and non-syzygetic pairs of scrolls stems from the desire to investigate the rationality of cubics  $X$  with an involution fixing a line pointwise. The twelve dimensional moduli space of such cubics  $\mathcal{M}_{\phi_2}$  is contained in  $\mathcal{C}_{12}$ : however, a general  $X \in \mathcal{M}_{\phi_2}$  contains 120 classes represented by cubic scrolls, and  $\text{rank} A(X)_{\text{prim}} = 8$  [Mar23, Theorem 4.5]. Thus the results discussed here can be viewed as a step towards understanding cubics contained in this family, to be treated in future work.

**2.3. The Fano variety of lines on a cubic fourfold.** Let  $X$  be a smooth cubic fourfold and  $F$  its Fano variety of lines, a hyperkähler fourfold of  $\mathrm{K3}^{[2]}$ -type [BD85]. Note that if  $Y = H \cap X$  is a hyperplane section of  $X$ , then  $F(Y)$  is a Lagrangian subvariety of  $F$ .

The Abel-Jacobi map

$$\alpha: H^4(X, \mathbb{Z}) \rightarrow H^2(F, \mathbb{Z})$$

restricts to an isomorphism  $H^4(X, \mathbb{Z})_{\mathrm{prim}} \rightarrow H^2(F, \mathbb{Z})_{\mathrm{prim}}$  compatible with the Hodge filtrations and (up to sign) with the quadratic forms on each lattice; more specifically, we have

$$q(\alpha(x), \alpha(y)) = -x \cdot y$$

where  $q$  is the Beauville-Bogomolov-Fujiki form (BBF form) of  $F$ . We write  $v^2 := q(v, v)$  for  $v \in H^2(F, \mathbb{Z})$ . Letting  $g$  be the Plücker polarization on  $F$ , we have that  $\alpha(\eta_X) = g$  and, moreover,  $g^2 = 6$  [BD85, Proposition 6].

If  $A(X) = \langle \eta_X, T \rangle$  where  $T$  is a cubic scroll, then setting  $\lambda = \alpha(T - \eta_X)$ , the BBF form on  $\mathrm{NS}(F)$  is given by

$$J_{12} = \begin{array}{c|cc} & g & \lambda \\ \hline g & 6 & 0 \\ \lambda & 0 & -4 \end{array}$$

and the discriminant group of  $\mathrm{NS}(F)$  is

$$D_{\mathrm{NS}(F)} \cong \mathbb{Z}/6 \oplus \mathbb{Z}/4,$$

with factors generated by  $[\frac{g}{6}]$  and  $[\frac{\lambda}{4}] \in D_{\mathrm{NS}(F)}$ .

Next, suppose  $A(X) = \langle \eta_X, T_1, T_2 \rangle$  where  $T_1$  and  $T_2$  are cubic scrolls spanning different hyperplanes. Let  $\lambda_i = \alpha(T_i - \eta_X)$ . If  $T_1$  and  $T_2$  are a syzygetic pair, then the BBF form on  $\mathrm{NS}(F)$  is

$$J_{\mathrm{syz}} = \begin{array}{c|ccc} & g & \lambda_1 & \lambda_2 \\ \hline g & 6 & 0 & 0 \\ \lambda_1 & 0 & -4 & 0 \\ \lambda_2 & 0 & 0 & -4 \end{array}$$

and the discriminant group

$$D_{\mathrm{NS}(F)} \cong \mathbb{Z}/6 \oplus (\mathbb{Z}/4)^2$$

has factors generated by  $[\frac{g}{6}]$ ,  $[\frac{\lambda_1}{4}]$ , and  $[\frac{\lambda_2}{4}]$ . On the other hand, if  $T_1$  and  $T_2$  are a non-syzygetic pair labeled so that  $[T_1] \cdot [T_2] = 1$ , the BBF form on  $\mathrm{NS}(F)$  is

$$J_{\mathrm{nonsyz}} = \begin{array}{c|ccc} & g & \lambda_1 & \lambda_2 \\ \hline g & 6 & 0 & 0 \\ \lambda_1 & 0 & -4 & 2 \\ \lambda_2 & 0 & 2 & -4 \end{array}$$

and the discriminant group

$$D_{\mathrm{NS}(F)} \cong \mathbb{Z}/6 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/6$$

has factors generated by  $[\frac{g}{6}]$ ,  $[\frac{\lambda_1}{2}]$ , and  $[\frac{\lambda_1}{3} + \frac{\lambda_2}{6}]$ .

**2.4. Birational geometry of hyperkähler fourfolds.** In order to study the birational geometry of  $F$ , we use the Global Torelli theorem for hyperkähler manifolds of  $K3^{[2]}$ -type due to Huybrechts, Markman and Verbitsky. In particular, we use Markman’s Hodge-theoretic version. We denote by  $\text{Mon}_{\text{Hdg}}^2(F)$  the subgroup of monodromy operators in  $\text{Mon}^2(F)$  that preserve the Hodge structure. Recall that for a manifold  $F$  of  $K3^{[2]}$ -type,  $\text{Mon}^2(F)$  is equal to the subgroup of elements of  $O^+(H^2(F, \mathbb{Z}))$  that act by  $\pm \text{Id}$  on the discriminant group [Mar11, Lemma 9.2].

**Theorem 2.14.** [Mar11, Theorem 1.3] *Let  $F$  be a projective hyperkähler manifold. Let  $g \in \text{Mon}_{\text{Hdg}}^2(F)$ . Then there exists  $f \in \text{Bir}(F)$  such that  $f^* = g$  if and only if  $g^* \text{Mov}(F) = \text{Mov}(F)$ . Further,  $f \in \text{Aut}(F)$  if and only if  $g^* \text{Amp}(F) = \text{Amp}(F)$ .*

*Remark 2.15.* In particular, it follows that a birational map  $f: F \dashrightarrow F'$  between hyperkähler manifolds is regular (hence an isomorphism) if and only if  $f^* \omega$  is ample for an ample class  $\omega$ .

The structure of the nef and movable cones of hyperkähler manifolds are well known: for  $K3^{[m]}$ -type, they are described in [Mar11, §9] and [BHT15, §1] using the extended Mukai-lattice. Here, we focus on the simpler case of manifolds of  $K3^{[2]}$ -type.

Let  $F$  be a projective hyperkähler manifold of  $K3^{[2]}$ -type. Let  $\overline{\text{Pos}}(F)$  be the component of the cone  $\{x \in \text{NS}(F) \otimes \mathbb{R} \mid x^2 \geq 0\}$  that contains an ample class. We denote by  $\text{Mov}(F) \subset \overline{\text{Pos}}(F)$  the (closed) convex cone generated by classes of line bundles on  $F$  whose base locus has codimension at least 2. By [HT09, Theorem 7],

$$\text{Mov}(F) = \overline{\bigcup_{f: F \dashrightarrow F'} f^* \text{Nef}(F')},$$

where  $f: F \dashrightarrow F'$  is a birational map with  $F'$  a hyperkähler manifold.

*Remark 2.16.* For hyperkähler manifolds with  $b_2 > 5$ , there are finitely many birational models (see [MY15], [AV17]). In particular, this is the case for hyperkähler manifolds of  $K3^{[2]}$ -type.

We define the following set of divisors:

$$\begin{aligned} \mathcal{W}_{\text{pex}} &:= \{\rho \in \text{NS}(F) \mid \rho^2 = -2\} \\ \mathcal{W}_{\text{flip}} &:= \{\rho \in \text{NS}(F) \mid \rho^2 = -10, \text{div}(\rho) = 2\}. \end{aligned}$$

A divisor  $\rho \in \mathcal{W}_{\text{pex}} \cup \mathcal{W}_{\text{flip}}$  is called a wall divisor [Mon15, Definition 1.2, Proposition 2.12], and  $\rho \in \mathcal{W}_{\text{pex}}$  is moreover referred to as a stably prime exceptional divisor. The following structure theorem for  $\text{Mov}(F)$  and  $\text{Amp}(F)$  in the case of hyperkähler manifolds of  $K3^{[2]}$ -type combines the results of Markman [Mar11, Lemma 6.22, Proposition 6.10, 9.12, Theorem 9.17] and Bayer, Hassett and Tschinkel [BHT15, Theorem 1].

**Theorem 2.17.** [Deb20, Theorem 3.16] *Let  $F$  be a hyperkähler manifold of  $K3^{[2]}$ -type.*

- (1) *The interior of  $\text{Mov}(F)$  is the connected component of*

$$\overline{\text{Pos}}(F) \setminus \bigcup_{\rho \in \mathcal{W}_{\text{pex}}} \rho^\perp$$

*that contains the class of an ample divisor.*

(2) The ample cone  $\text{Amp}(F)$  is the connected component of

$$\overline{\text{Pos}(F)} \setminus \bigcup_{\rho \in \mathcal{W}_{\text{pex}} \cup \mathcal{W}_{\text{flop}}} \rho^\perp$$

that contains the class of an ample divisor.

Note that each connected component of

$$\text{Int}(\text{Mov}(F)) \setminus \bigcup_{\rho \in \mathcal{W}_{\text{flop}}} \rho^\perp$$

corresponds to  $f^*(\text{Amp}(F'))$  for a birational map  $f: F \dashrightarrow F'$  with  $F'$  a hyperkähler manifold.

To obtain this chamber decomposition of  $\overline{\text{Pos}(F)}$  and of  $\text{Mov}(F)$ , Markman uses a more explicit description of the group  $\text{Mon}_{\text{Hdg}}^2(F)$  as follows. Let  $W_{\text{Exc}} \subset \text{Mon}_{\text{Hdg}}^2(F)$  be the subgroup generated by reflections  $R_\rho$  for all  $\rho \in \mathcal{W}_{\text{pex}}$ , and let  $W_{\text{Bir}} \subset \text{Mon}_{\text{Hdg}}^2(F)$  be the subgroup generated by monodromy operators induced from birational transformations of  $F$ . Let  $\pi: \text{Mon}_{\text{Hdg}}^2(F) \rightarrow \text{O}(\text{NS}(F))$  be the restriction homomorphism. We record some useful facts:

**Proposition 2.18.** *Let  $F$  be a hyperkähler manifold of  $K3^{[2]}$ -type. Then:*

- (1)  $\text{Mon}_{\text{Hdg}}^2(F) = W_{\text{Exc}} \rtimes W_{\text{Bir}}$ ;
- (2)  $\text{Mov}(F)$  is a fundamental domain for the action of  $W_{\text{Exc}}$  on  $\overline{\text{Pos}(F)}$ ;
- (3) The kernel of  $\pi$  is a subgroup of  $W_{\text{Aut}}$ , the set of monodromy operators induced by automorphisms of  $F$ .

*Proof.* The first statement is [Mar11, Theorem 6.18]. The second statement is [Mar11, Lemma 6.22]. For the third, [Mar11, Lemma 6.23] asserts that the kernel of  $\pi$  is a subgroup of  $W_{\text{Bir}}$ . By [Fuj81], a birational map acting trivially on  $\text{NS}(F)$  is regular.  $\square$

*Remark 2.19.* If  $F$  is the Fano variety of lines on a cubic fourfold  $X$  with  $\text{Aut}(X) = 0$ , then  $\text{Bir}(F) \rightarrow \text{O}(\text{NS}(F))$  is an embedding. Indeed,  $\text{Aut}(F, g) = \text{Aut}(X) = 0$ , so no automorphisms of  $F$  act trivially on  $\text{NS}(F)$  (see [Cha12, Proposition 4]). In particular, by Proposition 2.12, this always applies for the cubic fourfolds we study.

**2.5. GM fourfolds and double EPW sextics.** Later, we will exhibit birational models of the Fano variety  $F$  of lines on a general cubic fourfold  $X \in \mathcal{C}_{12}$  as double Eisenbud-Popescu-Walter (EPW) sextics constructed from smooth Gushel-Mukai (GM) fourfolds. In this section we recall the relevant background and terminology.

Let  $V_5$  be a complex vector space of dimension 5.

**Definition 2.20.** An (ordinary) GM fourfold  $Z$  is a smooth transverse intersection of the form

$$Z := \text{Gr}(2, V_5) \cap \mathbb{P}^8 \cap Q \subset \mathbb{P}(\bigwedge^2 V_5),$$

where  $\text{Gr}(2, V_5)$  is given the Plücker embedding,  $\mathbb{P}^8$  is a linear subspace, and  $Q$  is a quadric.

Such a fourfold is a Fano variety with Picard number 1, index 2 and degree 10 [Muk89]. In [IM11], the authors relate a GM fourfold  $Z$  to an EPW sextic  $W$  in the following way. Let  $I := |\mathcal{O}_Z(2)|$  be the linear system of quadrics in  $\mathbb{P}^8$  containing  $Z$ , and let  $\text{Disc}(Z) \subset I$  be the irreducible component of the discriminant hypersurface that parametrizes singular quadrics that are not restrictions of the Plücker quadrics.

**Theorem 2.21** ([IM11, Proposition 2.4], [DK18, Proposition 3.18]). *Let  $Z$  be an ordinary GM fourfold. The component  $\text{Disc}(Z) \subset I$  of the discriminant locus is an EPW sextic.*

Taking an appropriate double cover of  $W := \text{Disc}(Z)$ , one obtains a double EPW sextic associated to  $Z$  coinciding with the one constructed and studied by O’Grady in [O’G05, O’G06, O’G08b]. We instead review the construction of the double EPW sextic dual to this double cover. Let  $W^\vee \subset I^\vee$  be the hypersurface dual to  $\text{Disc}(Z)$ ; when  $Z$  contains no indecomposable vectors,  $W^\vee$  is again an EPW sextic [O’G08a, Corollary 3.6] (see also [DK18, Proposition B.3]). We make use of the following geometric interpretation of  $W^\vee$ , explained in [IM11] (see the construction of the map  $\alpha$  preceding Proposition 4.9).

**Lemma 2.22.** *The EPW sextic  $W^\vee$  satisfies the following properties:*

- (1) *Let  $w \in W^\vee$  be a general point. Then  $w$  corresponds to a unique singular quadric threefold  $Q_w \subset \mathbb{P}(\wedge^2 V_4) \subset \mathbb{P}^8$  for some  $V_4 \subset V_5$ , and vice versa.*
- (2) *Let  $C \subset Z$  be a general conic not contained in any plane in  $Z$ . Then there is a unique point  $w \in W^\vee$  such that  $Q_w$  contains the plane spanned by  $C$ .*

*Proof.* A point  $w \in W^\vee$  determines a hyperplane  $H \subset I$ ; the base locus of this system of quadrics is a quadric threefold  $Q_w$ . Since  $H$  is tangent to  $W \subset I$ ,  $Q_w$  is singular. Moreover, for general  $w$ , [IM11, Proposition 4.7] implies there is a unique hyperplane  $V_4 \subset V_5$  such that  $Q_w \subset \mathbb{P}(\wedge^2 V_4)$ .

Conversely, suppose  $Q' \subset \mathbb{P}(\wedge^2 V_4)$  is a singular quadric threefold. Then the complete linear system of quadrics containing  $Q'$  determines a hypersurface  $H \subset I$  which is tangent to  $W$  since  $Q'$  is singular. Hence we obtain a point  $w \in W^\vee$ , proving (1).

For (2), let  $C \subset Z$  be a conic such that the plane  $\langle C \rangle$  is not contained in  $Z$ . A general such conic  $C$  is not a  $\rho$ -conic (see the discussion at the beginning of [IM11, Section 4.4]), so there is a unique  $V_4 \subset V_5$  such that  $\mathbb{P}(\wedge^2 V_4)$  contains  $C$ . Let  $P_{V_4}$  denote the intersection of the Plücker quadric  $G(2, V_4)$  with  $\mathbb{P}^8 \subset \mathbb{P}^9$ , and let  $Q_{V_4} := Q \cap \mathbb{P}(\wedge^2 V_4) \subset \mathbb{P}^8$ . This gives a pencil  $\langle P_{V_4}, Q_{V_4} \rangle$  of quadric threefolds in  $\mathbb{P}(\wedge^2 V_4)$ , uniquely determined by  $Z$  and  $C$ . Since  $\langle C \rangle \not\subset Z$ , there is a unique quadric  $Q_C$  in the pencil containing this plane. The quadric  $Q_C$  is thus a singular quadric threefold, and by (1) there exists a unique  $w \in W^\vee$  such that  $Q_C = Q_w$ .  $\square$

*Remark 2.23.* In light of the above, a general point on the the double EPW sextic  $\widetilde{W}^\vee$  associated to  $W^\vee$  can be regarded as a ruling on one of the quadric threefolds  $Q_w$ . The covering involution on  $\widetilde{W}^\vee$  exchanges the rulings of each  $Q_w$ .

### 3. CUBIC FOURFOLDS CONTAINING ONE CUBIC SCROLL

Let  $X$  be a smooth cubic fourfold containing a cubic scroll  $T$ , and let  $F$  be its Fano variety of lines. We assume  $X$  is *very general*, by which we mean  $\text{rank}(A(X)) = 2$ ,  $\text{Aut}(X) = 0$ , and  $\text{End}_{\text{Hdg}}(T(F)) = \{\pm 1\}$ ; the third assumption can be made in light of [BFvGS24, Section 14]. In [HT10], the authors compute the ample and movable cone of  $F$ , and they exhibit a birational automorphism of  $F$  of infinite order. They also describe hyperkähler fourfolds birational to  $F$  as follows. The scroll  $T$  spans a hyperplane  $H$ , and by Proposition 2.4, the Fano variety of lines on  $Y := X \cap H$  is a Lagrangian subvariety of  $F$  decomposing as  $F(Y) = P \cup S' \cup P^\vee$ ,

where  $P$  and  $P^\vee$  are planes. Let  $F_1$  and  $F_1^\vee$  be the Mukai flops of  $F$  along  $P$  and  $P^\vee$ , respectively.

We expand on their study using the more recent techniques outlined in Section 2.4 for studying the birational geometry of hyperkähler manifolds, proving Theorem 1.1 as follows. In Section 3.1, we study isometries of  $\text{NS}(F)$ , and in particular those induced by birational automorphisms of  $F$ . In Section 3.2, we provide a correction to Hassett and Tschinkel's count on the number of non-isomorphic birational hyperkähler models, showing that  $F$  has exactly three birational models up to isomorphism, represented by  $F$ ,  $F_1$ , and  $F_1^\vee$ . Finally, in Section 3.3, we prove that  $F_1$  (and similarly  $F_1^\vee$ ) is in fact isomorphic to a double EPW sextic associated to the pair  $(X, T^\vee)$  (similarly, the pair  $(X, T)$ ).

**3.1. The Lattice  $\text{NS}(F)$ .** Recall from Section 2.3 that  $\text{NS}(F)$  is isomorphic to the lattice  $J_{12}$ . The discriminant group is  $\mathbb{Z}/6 \oplus \mathbb{Z}/4$ , with factors generated by  $[\frac{g}{6}]$  and  $[\frac{\lambda}{4}]$ . The BBF form represents  $-10$  but not  $-2$ , so  $F$  contains no prime exceptional divisors, and  $\text{Mov}(F) = \overline{\text{Pos}(F)}$ .

As noted in [HT10, Section 7], the isometry group of  $\text{NS}(F)$  is the product  $\{\pm 1\} \times \Gamma$ , where  $\Gamma$  is the infinite dihedral group

$$\Gamma = \langle R_1, R_2 \mid R_1^2 = R_2^2 = 1 \rangle;$$

explicitly, the generators are

$$R_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \text{and} \quad R_2 = \begin{pmatrix} 5 & -4 \\ 6 & -5 \end{pmatrix}.$$

We determine which isometries are induced by birational automorphisms of  $F$ :

**Lemma 3.1.** *An isometry  $\varphi \in \text{O}(\text{NS}(F))$  is induced by a birational automorphism of  $F$  if and only if  $\varphi$  preserves the positive cone and acts on the subgroup  $H$  of  $D_{\text{NS}(F)}$  generated by  $[\frac{g}{3}]$  and  $[\frac{\lambda}{4}]$  by  $\pm \text{Id}$ .*

*Proof.* Recall that  $\text{Mov}(F) = \overline{\text{Pos}(F)}$ , so if  $\varphi \in \text{O}(\text{NS}(F))$  satisfies  $\varphi = g^*$  for some  $g \in \text{Bir}(F)$ , then Theorem 2.14 implies that  $\varphi$  preserves the positive cone. Moreover,  $\varphi$  restricts to a Hodge isometry of  $T(F)$ , and since  $F$  is very general, the only Hodge isometries of  $T(F)$  are  $\pm \text{Id}_{T(F)}$ . Such an isometry necessarily acts on the discriminant group  $H' := D_{T(F)}$  by  $\pm \text{Id}$ . The overlattice  $H^2(F, \mathbb{Z}) \supset \text{NS}(F) \oplus T(F)$  corresponds to an index two subgroup  $H' \subset D_{\text{NS}(F)}$ , via Nikulin's theory of overlattices [Nik79, Propositions 1.4.1, 1.4.2]. In particular, it follows that  $(H, q_{\text{NS}(F)}) \cong (H', -q_{T(F)})$ , and hence  $\varphi$  acts on  $H$  by  $\pm \text{Id}$ .

Conversely, if  $\varphi$  preserves  $\overline{\text{Pos}(F)} = \text{Mov}(F)$  and acts on  $H$  by  $\pm \text{Id}$ , then by Theorem 2.14 it suffices to show that  $\varphi \in \text{Im}(\pi: \text{Mon}_{Hdg}^2(F) \rightarrow \text{O}(\text{NS}(F)))$ . The action of  $\varphi$  on  $H$  along with Nikulin's theory of overlattices implies that  $\varphi$  extends to an isometry  $\tilde{\varphi}$  of  $H^2(F, \mathbb{Z})$ , acting as  $\pm \text{Id}_{T(F)}$ . Since by construction  $\tilde{\varphi}$  preserves  $T(F)$ , it preserves the Hodge structure, hence comes from  $\text{Mon}_{Hdg}^2(F)$ .  $\square$

**Lemma 3.2.** *The subgroup of  $\text{O}(\text{NS}(F))$  consisting of isometries induced by birational automorphisms of  $F$  is generated by the reflections  $R_2$  and  $R_1 R_2 R_1$ . Moreover,  $\text{Bir}(F) \cong \langle a, b \mid a^2 = b^2 = 1 \rangle$ .*

*Proof.* The induced actions of  $R_1$  and  $R_2$  on the subgroup  $\langle [\frac{g}{3}] \rangle \oplus \langle [\frac{\lambda}{4}] \rangle \subset D_{\text{NS}(F)}$  are given, respectively, by the matrices

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \text{ and } \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}.$$

By Lemma 3.1, it is easy to see that an isometry of  $\text{NS}(F)$  is induced by a birational automorphism of  $F$  if and only if, written as a word in  $R_1$  and  $R_2$ , the generator  $R_1$  appears an even number of times. Since the induced actions of  $R_1$  and  $R_2$  on  $D_{\text{NS}(F)}$  are involutions that commute, this proves the first claim.

The second claim follows from the complete description of generators and relations on  $\Gamma \subset \text{O}(\text{NS}(F))$  and the fact that  $\text{Bir}(F)$  embeds in  $\text{O}(\text{NS}(F))$  by Remark 2.19.  $\square$

*Remark 3.3.* The isometries of  $\text{NS}(F)$  given by  $R_1 R_2 R_1$  and  $R_2$  are induced by birational involutions  $\iota$  and  $\iota^\vee$ , becoming regular on  $F_1$  and  $F_1^\vee$ , respectively (cf. [HT10, proof of Theorem 7.3] for computing  $\iota^*$  and  $(\iota^\vee)^*$ ). We review the geometric descriptions of these involutions, given in [HT10, Theorem 6.2]. Recall that  $T \subset X \subset \mathbb{P}^5$  spans a hyperplane  $H \subset \mathbb{P}^5$ , and  $Y := H \cap X$  has  $F(Y) = P \cup S' \cup P^\vee$  (see Proposition 2.4). Given a line  $[m] \in F \setminus F(Y)$ , the point  $p = H \cap m$  lies on unique lines  $\ell$  and  $\ell^\vee$  in the families  $P$  and  $P^\vee$  of  $F(Y)$ , respectively [HT10, Corollary 4.2]. Let  $\Pi_m = \text{span}\langle m, \ell \rangle$  and  $\Pi_m^\vee = \text{span}\langle m, \ell^\vee \rangle$ . Then the decompositions

$$\Pi_m \cap X = m \cup \ell \cup \iota^\vee(m)$$

and

$$\Pi_m^\vee \cap X = m \cup \ell^\vee \cup \iota(m)$$

define  $\iota$  and  $\iota^\vee$  away from  $F(Y)$ . In fact,  $\iota$  extends over  $P^\vee \setminus (P \cup S')$ , and  $\iota^\vee$  extends over  $P \setminus (P^\vee \cup S')$ .

**3.2. Census of birational models of  $F$ .** The movable cone of  $F$  contains infinitely many chambers, corresponding to the nef cones of birational models of  $F$ , as outlined in [HT10, Theorem 7.4] and Section 2.4. To enumerate the chambers, we follow [HT10, Section 7] by enumerating the wall divisors, i.e.  $v \in \text{NS}(F)$  with  $v^2 = -10$ . Let  $\rho_1 = g - 2\lambda$ ,  $\rho_2 = 3g - 4\lambda$ ,  $\rho_i^\vee = R_1(\rho_i)$ , and for all integers  $n$ ,

$$\rho_{2n+i} = (R_1 R_2)^n \rho_i$$

where we interpret  $\rho_i = \rho_{-i}^\vee$  for  $i < 0$ . Using standard propagation techniques, one sees that all the classes  $v$  with  $v^2 = -10$  and  $q(v, g) \geq 0$  are of the form  $\rho_i$  or  $\rho_i^\vee$  for some  $i$ .

Let  $\alpha_i$  be the class spanning  $\rho_i^\perp$  and pairing positively with  $g$ ; similarly, let  $\alpha_i^\vee$  span  $(\rho_i^\vee)^\perp$  and pair positively with  $g$ . Then the walls of  $\text{Mov}(F)$  are spanned by the  $\alpha_i$  and  $\alpha_i^\vee$ . In [HT10, Proposition 7.2], it is shown that  $\text{Nef}(F) = \text{Cone}(\alpha_1, \alpha_1^\vee)$ , and by [HT10, Proposition 7.3], we have  $\text{Nef}(F_1) = \text{Cone}(\alpha_1, \alpha_2)$  and  $\text{Nef}(F_1^\vee) = \text{Cone}(\alpha_1^\vee, \alpha_2^\vee)$ . Moreover, for all  $n$ ,  $\text{Cone}(\alpha_n, \alpha_{n+1})$  and  $\text{Cone}(\alpha_n^\vee, \alpha_{n+1}^\vee)$  are chambers of  $\text{Mov}(F)$  representing nef cones of birational models  $F_n$  and  $F_n^\vee$ .

Essentially by construction,

$$R_1 R_2 \cdot \text{Nef}(F_n) = \text{Nef}(F_{n+2}),$$

so the authors of [HT10] conclude in Theorem 7.4 that  $F_n \simeq F_{n+2}$  for all  $n$  (again interpreting  $F_n = F_{-n}^\vee$  when  $n < 0$ ). In particular, this would mean that  $F$  has at most two birational models up to isomorphism, represented by  $F$  and  $F_1$ . However,

by Lemma 3.2, we see that  $R_1R_2$  is **not** induced by a birational automorphism of  $F$  so by Theorem 2.14 need not send the nef cone of one model to the nef cone of some isomorphic model. Instead, we provide the following correction to [HT10, Theorem 7.4]:

**Proposition 3.4.** *Up to isomorphism,  $F$  admits exactly three birational hyperkähler models, represented by  $F$ ,  $F_1$ , and  $F_1^\vee$ .*

*Proof.* First, note  $(R_1R_2)^2 = \iota^* \circ (\iota^\vee)^*$  by Remark 3.3, so from

$$(R_1R_2)^2 \cdot \text{Nef}(F_n) = \text{Nef}(F_{n+4}),$$

we deduce  $F_n \simeq F_{n+4}$  for all  $n$ . Moreover, it is straightforward to calculate

$$\iota^* \text{Nef}(F) = \text{Nef}(F_2) \quad \text{and} \quad (\iota^\vee)^* \text{Nef}(F) = \text{Nef}(F_2^\vee),$$

so  $F_2 \simeq F \simeq F_2^\vee$ , by Remark 2.15. It follows that  $F$  has at most three birational hyperkähler models up to isomorphism, represented by  $F$ ,  $F_1$ , and  $F_1^\vee$ . We now show that these three are non-isomorphic.

To distinguish  $F$  from  $F_1$  and  $F_1^\vee$ , note that  $F_1$  and  $F_1^\vee$  both admit nontrivial involutions whereas  $F$  does not: indeed, the only nontrivial automorphism of  $\text{Nef}(F)$  preserving the positive cone is  $R_1$  which, by Lemma 3.2, is not induced by a birational automorphism of  $F$ . To distinguish  $F_1$  from  $F_1^\vee$ , note that the only two isometries of  $\text{NS}(F)$  sending  $\text{Nef}(F_1)$  to  $\text{Nef}(F_1^\vee)$  are  $R_1$  and  $R_2R_1$ . Again, by Lemma 3.2, neither of these lattice automorphisms are induced by a birational automorphism of  $F$ .  $\square$

**3.3. Connection with Gushel–Mukai fourfolds.** Here, we show that  $F_1$  and  $F_1^\vee$  are isomorphic to double EPW sextics. This relates each birational model of  $F$  to a familiar family of hyperkähler fourfolds and explains the birational involutions  $\iota$  and  $\iota^\vee$  on  $F$ —they are induced by the covering involutions associated to the double EPW sextics.

Recalling the construction in [DIM15, Section 7.2] and [KP18, Proposition 5.6], the complete linear system of quadrics containing the scroll  $T$  induces a rational map  $q: X \dashrightarrow \mathbb{P}^8$  which is birational onto its image, a GM fourfold  $Z_T \subset \text{Gr}(2, V_5)$  containing a plane  $\Pi$ . Projection from  $\Pi$  induces a rational inverse  $f: Z_T \dashrightarrow X$ . We obtain the following diagram, where  $Y^+ \rightarrow Y$  is the small resolution from (2.2), and  $E$  is the exceptional divisor.

$$\begin{array}{ccccc}
 Y^+ & \xrightarrow{\quad \sim \quad} & & & E \\
 \downarrow & \searrow & & \swarrow & \downarrow \\
 Y & & \text{Bl}_T X & \xrightarrow{\quad \sim \quad} & \text{Bl}_\Pi Z_T & & \Pi \\
 \downarrow & \swarrow & & \searrow & & & \downarrow \\
 X & & & & & & Z_T \\
 & & & & \xrightarrow{\quad q \quad} & & 
 \end{array}$$

Note that the isomorphism  $\text{Bl}_T X \simeq \text{Bl}_\Pi Z_T$  identifies  $Y^+$  with  $E$ , so the image of  $Y^+$  in  $Z_T$  is  $\Pi$ .

Let  $W$  be the EPW sextic associated to  $Z_T$  and  $W^\vee$  its dual, as defined in Section 2.5. While  $Z_T$  depends on the choice of scroll  $T$  in its homology class,  $W$  does not [DIM15, Proposition 7.2]. Let  $\widetilde{W}^\vee$  be the double cover of  $W^\vee$ , also introduced in Section 2.5, equipped with its covering involution  $\tau$ .

**Lemma 3.5.** *Let  $Q_w$  be a singular quadric threefold associated via Lemma 2.22 to a point  $w \in W^\vee$ . Then either  $\Pi \subset Q_w$  or  $\Pi$  and  $Q_w$  meet in a point. For general  $w$ , the quadric  $Q_w$  is unique, and  $\Pi$  and  $Q_w$  meet in one point which is not the cone point of  $Q_w$ .*

*Proof.* By [DIM15, Section 7.2],  $\Pi = \mathbb{P}(\wedge^2 V_3)$  for some three-dimensional space  $V_3 \subset V_5$ . For general  $Q_w$ , there is a unique hyperplane  $V_4 \subset V_5$  such that  $Q_w = \mathbb{P}(\wedge^2 V_4) \cap Z_T$  by Lemma 2.22(1). Note  $\Pi \cap Q_w = \mathbb{P}(\wedge^2(V_3 \cap V_4))$  which is either a point or all of  $\Pi$  depending on whether  $V_3 \subset V_4$ . Only a two-dimensional family of hypersurfaces  $V_4$  contain  $V_3$ . Moreover, by the proof of [IM11, Lemma 2.3], each point in  $\mathbb{P}^8$  is the cone point of  $Q_w$  for at most one  $w \in W^\vee$ ; in particular, the cone point of  $Q_w$  belongs to  $\Pi$  for at most a two-dimensional locus in  $W^\vee$ .  $\square$

**Proposition 3.6.**  *$F$  and  $\widetilde{W}^\vee$  are birational.*

*Proof.* We define a rational map  $\beta: F \dashrightarrow \widetilde{W}^\vee$  as follows. Recall that  $H \subset \mathbb{P}^5$  is the hyperplane spanned by  $T$ . A general line  $m \subset X$  meets  $H$  in a point  $p$ , and  $C_m = q(m)$  is a conic in  $Z_T$  meeting  $\Pi$  in a point. In particular,  $C_m$  is not contained in any plane of  $Z$ , thus by Lemma 2.22(2) there exists a unique singular quadric  $Q_m \subset \mathbb{P}(\wedge^2 V_4)$  for some  $V_4 \subset \mathbb{P}^5$ . As  $C_m$  spans a plane contained in  $Q_m$ , it determines a ruling of  $Q_m$  and hence a point  $\beta(m) \in \widetilde{W}^\vee$  by Remark 2.23.

To show  $\beta$  is a birational equivalence, we describe its inverse. A general point  $w \in \widetilde{W}^\vee$  specifies a ruling on a singular quadric threefold  $Q_w$ . Lemma 3.5 implies there is a unique plane  $P_w$  in that ruling of  $Q_w$  containing the point  $Q_w \cap \Pi$ . The plane  $P_w$  meets  $Z_T$  in a conic  $C_w$  intersecting  $\Pi$ , so projection from  $\Pi$  yields a line in  $X$ , which we take as  $\beta^{-1}(w)$ . One can check that this construction is inverse to the construction of  $\beta$ .  $\square$

**Proposition 3.7.** *The birational map  $\beta$  induces an isomorphism  $F_1^\vee \simeq \widetilde{W}^\vee$ .*

*Proof.* It suffices to show that  $\iota^\vee$  becomes regular on  $\widetilde{W}^\vee$ , and in particular, we show  $\beta \circ \iota^\vee \circ \beta^{-1}$  and the covering involution  $\tau$  agree on an open set.

Let  $w \in W^\vee$  be a point defining a singular quadric threefold  $Q_w$ . By Lemma 3.5, for general  $Q_w$  we have  $Q_w \cap \Pi = \{x\}$  where  $x$  is not the cone point of  $Q_w$ . Hence there is a unique plane in each ruling of  $Q$  containing  $x$ ; these planes intersect  $Z$  in conics  $C$  and  $\tau(C)$ . The conics  $C$  and  $\tau(C)$  intersect in two points—once at  $x$  and again along the line joining  $x$  to the cone point of  $Q_w$ . In general,  $x$  and the second point  $y$  are distinct, so  $y \notin \Pi$ .

Projecting from  $\Pi$ , we obtain lines  $m = q^{-1}(C)$  and  $m' = q^{-1}(\tau(C))$  intersecting at  $q^{-1}(y) \notin H$ . Moreover, by Remark 2.3 the fiber of  $Y^+ \rightarrow \Pi$  over  $x$  is a line whose image  $\ell$  in  $Y$  is contained in a cubic scroll  $T'$  homologous to  $T^\vee$ , and  $\ell^2 = 0$  on  $T'$ . Since any two of  $m$ ,  $m'$ , and  $\ell$  intersect, but not all at the same point, the three lines are coplanar. By [HT10, Proposition 4.7],  $[\ell] \in P$ , so, recalling the definition of  $\iota^\vee$  from Remark 3.3, we have  $\iota^\vee(m) = m'$ . In other words,  $\iota^\vee(\beta^{-1}([C])) = \beta^{-1}(\tau([C]))$ , proving the claim.  $\square$

Starting instead with  $\text{Bl}_{T^\vee} X$ , one obtains an isomorphism between  $F_1$  and a double EPW sextic whose covering involution induces  $\iota$ . By Lemma 3.2 and Remark 3.3, the covering involutions on the two double EPW sextics generate  $\text{Bir}(F)$ . This observation, together with Propositions 3.4 and Proposition 3.7, completes the proof of Theorem 1.1.

*Remark 3.8.* It would be interesting to know whether  $Z_T$  and  $Z_{T^\vee}$  are dual GM fourfolds, or equivalently if the double EPW sextics associated as above to  $(X, T)$  and  $(X, T^\vee)$  are dual (see [DK18, Definition 3.26]). By [KP23, Theorem 1.6], dual GM fourfolds are Fourier-Mukai partners. Kuznetsov and Perry in [KP18, Theorem 5.8] show directly that  $X$  and  $Z_T$  have derived equivalent Kuznetsov components, which implies the same is true of  $Z_T$  and  $Z_{T^\vee}$ .

#### 4. A SYZYGETIC PAIR OF CUBIC SCROLLS

Let  $X \subset \mathbb{P}^5$  be a smooth cubic fourfold containing a syzygetic pair of cubic scrolls  $T_1$  and  $T_2$ . We also assume  $X$  is *very general*, meaning  $\text{rank}(A(X)) = 3$ ,  $\text{Aut}(X) = 0$ , and  $\text{End}_{\text{Hdg}}(T(F)) = \{\pm 1\}$ . In this section, we completely describe the birational geometry of the Fano variety  $F$  of lines on  $X$ . Specifically, we prove the following:

**Theorem 4.1.** *Let  $F$  be the Fano variety of lines on a very general cubic fourfold  $X$  containing a syzygetic pair of cubic scrolls. Then  $F$  has **five** isomorphism classes of birational hyperkähler models, represented by the following:*

- $F$  itself, and
- four non-isomorphic Mukai flops of  $F$ , each isomorphic to a double EPW sextic.

Furthermore, we determine the birational automorphism group of  $F$ :

**Theorem 4.2.** *Let  $F$  be as above. Then*

$$\text{Bir}(F) \cong \langle a, b, c, d \mid a^2 = b^2 = c^2 = d^2 = 1 \rangle.$$

*Moreover, the four generators can be identified with the covering involutions on the double EPW sextics obtained as Mukai flops of  $F$ .*

The outline is as follows: in Section 4.1, we study the arrangement of Lagrangian planes in  $F$ , showing that any two planes explained by the presence of  $T_1$  and  $T_2$  intersect. We then identify which isometries of  $\text{NS}(F)$  are induced by birational automorphisms of  $F$  in Section 4.2, including identifying how the involutions from Remark 3.3 act. After enumerating the (infinitely many) walls of the movable cone of  $F$  in Section 4.3, we enumerate the birational models and identify the birational automorphism group in Section 4.4.

**4.1. Lagrangian planes in  $F$ .** Let  $H_i$  be the hyperplane spanned by  $T_i$ , and  $Y_i = X \cap H_i$ . Recall from Section 2.1 that

$$F(Y_i) = P_i \cup S'_i \cup P_i^\vee$$

where  $P_i$  and  $P_i^\vee$  are Lagrangian planes. The intersection  $F(Y_1) \cap F(Y_2)$  parametrizes lines on the cubic surface  $\Sigma = X \cap H_1 \cap H_2$ . For a general cubic fourfold  $X$  satisfying the hypotheses above,  $\Sigma$  is smooth by Lemma 2.8. The following lemma implies the intersection  $F(Y_1) \cap F(Y_2)$  is transverse in general.

**Lemma 4.3.** *Let  $X$  be a smooth cubic fourfold containing a line  $L$ , let  $F$  be the Fano variety of lines on  $X$ , and let  $H_1$  and  $H_2$  be distinct hyperplanes containing  $L$ . Suppose further that  $L$  passes through no singular point of  $Y_i = X \cap H_i$ . Then the surfaces  $F(Y_1), F(Y_2) \subset F$  intersect transversely at  $[L]$  if and only if  $\Sigma = Y_1 \cap Y_2$  is smooth along  $L$ .*

*Proof.* Since  $Y_i$  is smooth along  $L$ , we have  $\dim(T_{[L]}F(Y_i)) = 2$ . From the exact sequence

$$0 \rightarrow T_{[L]}F(\Sigma) \rightarrow T_{[L]}F(Y_1) + T_{[L]}F(Y_2) \rightarrow T_{[L]}F(X),$$

it is therefore clear that  $F(Y_1)$  intersects  $F(Y_2)$  transversely at  $[L]$  if and only if  $\dim(T_{[L]}F(\Sigma)) = 0$ , which is the case precisely when  $\Sigma$  is smooth along  $L$ .  $\square$

We also compare intersections on  $X$  to intersections on  $\Sigma$ :

**Lemma 4.4.** *Let  $X$  be as above, suppose  $\Sigma$  is smooth, and let  $y_i \in Y_i$  be a node. Further, let  $A_i \cup A_i^\vee$  be the surface swept out by lines on  $X$  through  $y_i$ , with  $[A_i] = [T_i]$  and  $[A_i^\vee] = [T_i^\vee]$  as in Remark 2.5, and let  $\gamma_i = A_i \cap \Sigma$  and  $\gamma_i^\vee = A_i^\vee \cap \Sigma$ . Then  $\gamma_i$  and  $\gamma_i^\vee$  are twisted cubic curves, and  $[T_1] \cdot [T_2] = [\gamma_1] \cdot [\gamma_2]$  where the first intersection pairing happens on  $X$  and the second on  $\Sigma$ .*

*Proof.* Since  $\Sigma$  is smooth, none of the nodes of  $Y_1$  lie on  $Y_2$ . Hence the cubic curve  $\gamma_1 = A_1 \cap H_2$  does not pass through the cone point of  $A_1$ , so it is a smooth twisted cubic. Similarly,  $\gamma_2$  is a smooth twisted cubic. To verify the equality of the intersection numbers, note that  $A_1 \cap A_2 = (A_1 \cap H_2) \cap (A_2 \cap H_1) = \gamma_1 \cap \gamma_2$  scheme-theoretically, and  $[A_i] = [T_i]$ .  $\square$

**Proposition 4.5.** *Let  $X$  be a cubic fourfold as above with  $[T_1] \cdot [T_2] = 3$ . Then*

- $\deg(P_1 \cap P_2) = \deg(P_1 \cap P_2^\vee) = \deg(P_1^\vee \cap P_2) = \deg(P_1^\vee \cap P_2^\vee) = 1$ ,
- $\deg(P_1 \cap S_2') = \deg(P_1^\vee \cap S_2') = \deg(S_1' \cap P_2) = \deg(S_1' \cap P_2^\vee) = 4$ ,
- $\deg(S_1' \cap S_2') = 7$ ,

*and these intersections are all transverse when  $\Sigma = X \cap H_1 \cap H_2$  is smooth.*

*Proof.* It suffices to calculate the intersection degrees for any cubic fourfold satisfying the hypotheses, so by Lemma 2.8 we may assume that  $\Sigma$  is smooth. In particular,  $\Sigma$  being smooth requires that  $H_1$  contains none of the nodes of  $Y_2$ , and  $H_2$  contains none of the nodes of  $Y_1$ . Then, as mentioned in Lemma 4.3,  $F(Y_1)$  and  $F(Y_2)$  intersect transversely. Hence we need only count points in the intersections set-theoretically.

As in Lemma 4.4, let  $\gamma_i = A_i \cap \Sigma$  and  $\gamma_i^\vee = A_i^\vee \cap \Sigma$ . The strategy of the proof is as follows: we determine the incidence relations between each line  $\ell \subset \Sigma$  and each of the curves  $\gamma_i, \gamma_i^\vee$  for  $i = 1, 2$ . By Remark 2.5,  $\ell$  is bisecant to  $\gamma_i$  or  $\gamma_i^\vee$  if and only if  $[\ell] \in P_i$  or  $P_i^\vee$ , respectively;  $\ell$  is secant to both  $\gamma_i$  and  $\gamma_i^\vee$  if and only if  $[\ell] \in S_i'$ .

As noted in [Dol16], the linear system  $|\gamma_1|$  defines a morphism  $f: \Sigma \rightarrow \mathbb{P}^2$  blowing down a sixer  $E_1, \dots, E_6$ . Writing  $E_0 = f^*\mathcal{O}(1)$ , and using the facts that  $\gamma_1 \cup \gamma_1^\vee \subset Q$  for some quadric, and  $Q \cap \Sigma \sim -2K_\Sigma$ , we have

$$\gamma_1 \sim E_0 \quad \text{and} \quad \gamma_1^\vee \sim 5E_0 - 2 \sum_{i=1}^6 E_i.$$

Now, write  $\gamma_2 \sim \sum_{i=0}^6 a_i E_i$ . By Lemma 4.4,  $a_0 = [\gamma_1] \cdot [\gamma_2] = 3$ . Similarly,

$$3 = [\gamma_1^\vee] \cdot [\gamma_2] = 15 + 2 \sum_{i=1}^6 a_i.$$

Since  $\gamma_2$  is a twisted cubic, we also have  $1 = \gamma_2^2 = 9 - a_1^2 - \dots - a_6^2$ , so, possibly after relabeling  $E_1, \dots, E_6$ , we obtain

$$\gamma_2 \sim 3E_0 - 2E_1 - E_2 - E_3 - E_4 - E_5.$$

From here, the claim follows readily using the incidence relations for lines on  $\Sigma$ .  $\square$

In particular, any two of the four Lagrangian planes in  $F$  studied here intersect. This clarifies what we prove later: after flopping any one of the planes in  $F$ , one can no longer flop any of the other three.

**4.2. The lattice  $\text{NS}(F)$ .** Letting  $\lambda_i = \alpha(T_i - \eta_X)$ , recall from Section 2.3 that  $\text{NS}(F)$  with the BBF form is isomorphic to the lattice  $J_{xyz}$  and has discriminant group  $D_{\text{NS}(F)} \cong \mathbb{Z}/6 \oplus (\mathbb{Z}/4)^2$  with factors generated by  $[\frac{g}{6}]$ ,  $[\frac{\lambda_1}{4}]$ , and  $[\frac{\lambda_2}{4}]$ . The BBF form represents both  $-2$  and  $-10$ .

The isometry group of  $\text{NS}(F)$  with basis  $\{g, \lambda_1, \lambda_2\}$  is generated by  $\pm 1$  and the four reflections below. We calculated the isometry group  $\text{O}(\text{NS}(F))$  via the algorithm outlined in [Mer14] and implemented using the Magma package `AutHyp.m`.

$$R_1 = \begin{pmatrix} 5 & 4 & 0 \\ -6 & -5 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad R_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$R_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad R_4 = \begin{pmatrix} 7 & -4 & 4 \\ 6 & -3 & 4 \\ -6 & 4 & -3 \end{pmatrix}.$$

The following lemma helps distinguish which isometries of the lattice  $\text{NS}(F)$  are induced by birational automorphisms of  $F$ .

**Lemma 4.6.** *Any isometry  $\varphi \in \text{O}(\text{NS}(F))$  is induced by a birational automorphism of  $F$  if and only if  $\varphi$  preserves  $\text{Mov}(F)$  and acts on the index two subgroup*

$$H := \left\langle \left[ \frac{g}{3} \right] \right\rangle \oplus \left\langle \left[ \frac{\lambda_1}{4} \right] \right\rangle \oplus \left\langle \left[ \frac{\lambda_2}{4} \right] \right\rangle$$

of  $D_{\text{NS}(F)}$  by  $\pm \text{Id}$ .

*Proof.* This is similar to Lemma 3.1, the only difference being that preserving  $\text{Pos}(F)$  is no longer equivalent to preserving  $\text{Mov}(F)$ .  $\square$

Let  $F_i$  and  $F_i^\vee$  denote the flops of  $F$  along  $P_i$  and  $P_i^\vee$ , respectively. As mentioned in Remark 3.3, each of these models admits a regular involution:  $\iota_i$  and  $\iota_i^\vee$ , respectively. These involutions can also be regarded as birational involutions on  $F$ .

**Lemma 4.7.** *The birational involutions  $\iota_i$  and  $\iota_i^\vee$  for  $i = 1, 2$  act on  $\text{NS}(F)$  by  $\iota_1^* = R_1$ ,  $(\iota_1^\vee)^* = R_2 R_1 R_2$ ,  $\iota_2^* = R_3 R_1 R_3$ , and  $(\iota_2^\vee)^* = R_3 R_2 R_1 R_2 R_3$ .*

*Proof.* We give the proof for  $\iota_1^*$ . By [HT10, Proposition 6.5],  $\iota_1^*$  fixes the class  $g - \lambda_1$ . Using this fact, along with the properties that  $\iota_1^*$  preserves the BBF form and  $(\iota_1^*)^2 = \text{Id}$ , a direct computation verifies that the only involutions of  $\text{NS}(F)$  fixing  $g - \lambda_1$  are given by the matrices

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \pm 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 5 & 4 & 0 \\ -6 & -5 & 0 \\ 0 & 0 & \pm 1 \end{pmatrix}.$$

The first two candidates are impossible: they preserve  $g$  and therefore preserve  $\text{Nef}(F)$  whereas  $\iota_1$  is not regular on  $F$ . To distinguish between the two remaining

candidates, we inspect actions on the discriminant group of  $\text{NS}(F)$ . The matrix

$$\begin{pmatrix} 5 & 4 & 0 \\ -6 & -5 & 0 \\ 0 & 0 & \pm 1 \end{pmatrix}$$

acts on  $\langle [\frac{g}{3}] \rangle \times \langle [\frac{\lambda_1}{4}] \rangle \times \langle [\frac{\lambda_2}{4}] \rangle$  by

$$\begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & \pm 1 \end{pmatrix}.$$

By Lemma 4.6, the sign in the last entry of this matrix must be negative, yielding the desired result. The other three actions are calculated similarly.  $\square$

**4.3. Wall and chamber structure of  $\text{Mov}(F)$ .** In order to study the birational geometry of  $F$  and, in particular, to enumerate the birational hyperkähler models of  $F$  up to isomorphism, we give a description of the wall and chamber decomposition of the movable cone of  $F$ .

As before, we fix the orthogonal basis  $\{g, \lambda_1, \lambda_2\}$  on  $\text{NS}(F)$ , and we denote a class

$$v = ag + b\lambda_1 + c\lambda_2 \in \text{NS}(F) \otimes \mathbb{R}$$

by the vector  $v = (a, b, c)$ .

**Lemma 4.8.** *The nef cone of  $F$  is bounded by the four walls  $(1, \pm 2, 0)^\perp$  and  $(1, 0, \pm 2)^\perp$ . The only other walls in  $\text{Mov}(F)$  intersecting  $\text{Nef}(F)$  are the walls  $(1, 1, 1)^\perp$ ,  $(1, 1, -1)^\perp$ ,  $(1, -1, 1)^\perp$ , and  $(1, -1, -1)^\perp$ , each of which intersects  $\text{Nef}(F)$  in codimension two.*

*Proof.* By [HT10, Proposition 7.2], we know that each of the walls  $(1, \pm 2, 0)^\perp$  and  $(1, 0, \pm 2)^\perp$  induces a small contraction of  $F$ , hence lies on the boundary of the nef cone. We prove that the only walls intersecting the chamber enclosed by the above walls are  $(1, 1, 1)^\perp$ ,  $(1, 1, -1)^\perp$ ,  $(1, -1, 1)^\perp$ , and  $(1, -1, -1)^\perp$ .

To do this, consider the cross-section of  $\text{Mov}(F)$  by the plane  $x = 4$ : the cross-section of  $\text{Cone}((1, \pm 2, 0)^\perp, (1, 0, \pm 2)^\perp)$  is the square bounded by the four lines  $y = \pm 3$  and  $z = \pm 3$ . In this slice, any wall of  $\text{Nef}(F)$  is a line cutting through this square so a point on the line has distance at most  $\sqrt{18}$  from the origin.

First, suppose  $v = (a, b, c)$  satisfies  $v^2 = -2$ . Then any point on the line

$$(v^\perp) \cap \{x = 4\} = \{6a - by - cz = 0\}$$

has distance

$$\frac{6|a|}{\sqrt{b^2 + c^2}} = \frac{6\sqrt{2}|a|}{\sqrt{3a^2 + 1}}$$

from the origin. This distance is no more than  $\sqrt{18}$  if and only if  $|a| \leq 1$ . The only four such classes with  $|a| \leq 1$  are  $(1, 1, 1)^\perp$ ,  $(1, 1, -1)^\perp$ ,  $(1, -1, 1)^\perp$ , and  $(1, -1, -1)^\perp$ , which intersect the boundary of  $\text{Cone}((1, \pm 2, 0)^\perp, (1, 0, \pm 2)^\perp)$ .

A similar calculation shows that no other classes with  $v^2 = -10$  intersect  $\text{Cone}((1, \pm 2, 0)^\perp, (1, 0, \pm 2)^\perp)$ .  $\square$

**Lemma 4.9.** *The nef cones of the four flops of  $F$  are*

$$\begin{aligned} \text{Nef}(F_1) &= \text{Cone}((1, -2, 0)^\perp, (1, -1, 1)^\perp, (1, -1, -1)^\perp, (3, -4, 0)^\perp), \\ \text{Nef}(F_1^\vee) &= \text{Cone}((1, 2, 0)^\perp, (1, 1, 1)^\perp, (1, 1, -1)^\perp, (3, 4, 0)^\perp), \\ \text{Nef}(F_2) &= \text{Cone}((1, 0, -2)^\perp, (1, 1, -1)^\perp, (1, -1, -1)^\perp, (3, 0, -4)^\perp), \\ \text{Nef}(F_2^\vee) &= \text{Cone}((1, 0, 2)^\perp, (1, 1, 1)^\perp, (1, -1, 1)^\perp, (3, 0, 4)^\perp). \end{aligned}$$

*In particular, the chamber  $\text{Nef}(F_i)$  shares a face with two other chambers of  $\text{Mov}(F)$ :  $\text{Nef}(F)$  and  $\iota_i^*\text{Nef}(F)$ . Similarly,  $\text{Nef}(F_i^\vee)$  shares a face with the two chambers  $\text{Nef}(F)$  and  $(\iota_i^\vee)^*\text{Nef}(F)$ .*

*Proof.* First, we give the proof for  $F_1$ . Since  $F_1$  is the flop of  $F$  along  $P_1$ , we know  $(1, -2, 0)^\perp$  lies on the boundary of  $\text{Nef}(F_1)$ . Moreover,  $\iota_1$  is regular on  $F_1$  so acts on  $\text{Nef}(F_1)$  by an involution; using Lemma 4.7, we calculate  $\iota_1^*(1, -2, 0) = (-3, 4, 0)$ , so  $(3, -4, 0)^\perp$  must also lie on the boundary of  $\text{Nef}(F)$ . Since the prime exceptional classes cut the movable cone out of the positive cone, we have

$$\text{Nef}(F_1) \subset \text{Cone}((1, -2, 0)^\perp, (1, -1, 1)^\perp, (1, -1, -1)^\perp, (3, -4, 0)^\perp).$$

Using Lemma 4.8, we see that any other wall of  $\text{Nef}(F_1)$  intersecting the interior of the cone above would also cut into  $\text{Nef}(F)$  or  $\iota_1^*\text{Nef}(F)$ , a contradiction. Hence

$$\text{Nef}(F_1) = \text{Cone}((1, -2, 0)^\perp, (1, -1, 1)^\perp, (1, -1, -1)^\perp, (3, -4, 0)^\perp).$$

The three other cone descriptions are proved similarly.  $\square$

Figure 1 illustrates the movable cone with labels on the chambers we have described so far. Figure 2 gives more detail further out from the central chamber.

We finish the section by demonstrating how to propagate all of the (infinitely many) walls of the movable cone. This is not necessary for enumerating the birational models of  $F$  up to isomorphism, but we will obtain a description of the birational automorphism group of  $F$  as a byproduct.

Let  $\Gamma \subset \text{O}(\text{NS}(F))$  be the subgroup generated by  $\iota_1^*$ ,  $(\iota_1^\vee)^*$ ,  $\iota_2^*$ , and  $(\iota_2^\vee)^*$ . Since these lattice automorphisms are induced by birational automorphisms of  $F$ , they preserve  $\text{Mov}(F)$  and act on the sets  $\mathcal{W}_{\text{flop}}$  and  $\mathcal{W}_{\text{pex}}$  of wall divisors (see Section 2.4 for the definitions). In particular, they also act on the sets

$$\Delta_{\text{flop}} := \{\rho \in \mathcal{W}_{\text{flop}} \mid \rho^\perp \cap \text{Mov}(F) \neq \emptyset\}$$

and

$$\Delta_{\text{pex}} := \{\rho \in \mathcal{W}_{\text{pex}} \mid \rho^\perp \cap \text{Mov}(F) \neq \emptyset\},$$

which define walls between chambers in  $\text{Mov}(F)$  and walls bounding  $\text{Mov}(F)$ , respectively.

**Lemma 4.10.**  $\Gamma$  acts freely on  $\Delta_{\text{flop}}$  with four orbits, represented by the classes  $(1, \pm 2, 0)$  and  $(1, 0, \pm 2)$ .

*Proof.* Suppose  $v = (a, b, c) \in \mathcal{W}_{\text{flop}}$  is a class such that  $v^\perp \cap \text{Mov}(F) \neq \emptyset$ , i.e.  $v^\perp$  is a wall divisor. If  $a = 1$ , then  $v$  is one of the four vectors listed above, thus we assume  $a > 1$ . Note that  $v^\perp$  intersects one of the walls  $(1, 1, 1)^\perp$ ,  $(1, 1, -1)^\perp$ ,  $(1, -1, -1)^\perp$ , and  $(1, -1, 1)^\perp$ , from which we deduce  $|a| < |b|$  or  $|a| < |c|$ . Since




FIGURE 1. A cross-section of the chambers of the movable cone of  $F$ , bounded by the positive cone. Prime exceptional walls are drawn in red, and the remaining wall divisors are drawn in blue.




FIGURE 2. A detail of the figure above illustrating the chambers adjacent to  $F_1^V$ . Chambers are labeled by the isomorphism type of the model they represent. The involution  $\iota_1^V$  exchanges the two chambers adjacent to  $\text{Nef}(F_1^V)$ .

also  $3a^2 - 2b^2 - 2c^2 = -5$ , we have either  $|c| < |a| < |b|$  or  $|b| < |a| < |c|$ . We act by  $\Gamma$  and find:

$$\begin{aligned} \iota_1^* v &= (5a + 4b, -6a - 5b, -c) & \iota_2^* v &= (5a + 4c, -b, -6a - 5c) \\ (\iota_1^\vee)^* v &= (5a - 4b, 6a - 5b, -c) & (\iota_2^\vee)^* v &= (5a - 4c, -b, 6a - 5c). \end{aligned}$$

The four classes above all define walls of the movable cone of  $F$ . We claim that exactly one of the classes above has a first coordinate with smaller magnitude than  $|a|$ . Explicitly, if  $|c| < |a| < |b|$ , then:

- The inequality  $|c| < |a|$  implies  $|5a \pm 4c| > |a|$ .
- From  $|a| > 1$ ,  $|a| > |c|$ , and  $3a^2 - 2b^2 - 2c^2 = -5$ , we obtain the inequality  $|3a| > |2b|$ .
- If  $\frac{a}{b} > 0$ , then the inequalities  $a < b$  and  $|3a| > |2b|$  yield  $|5a - 4b| < |a|$  and  $|5a + 4b| > |a|$ .
- Conversely, if  $\frac{a}{b} < 0$ , then the inequalities  $a < b$  and  $|3a| > |2b|$  yield  $|5a + 4b| < |a|$  and  $|5a - 4b| > |a|$ .

The argument for the case  $|b| < |a| < |c|$  is similar. From this, we observe that the action by  $\Gamma$  must be free and that there is a unique element of  $\Gamma$  taking  $v$  to a  $(-10)$ -class whose first coordinate is 1. Hence there are exactly four orbits of the action of  $\Gamma$  on  $\Delta_{\text{flop}}$ .  $\square$

**Lemma 4.11.**  $\Gamma$  acts freely on  $\Delta_{\text{pex}}$  with four orbits, each containing one of the classes  $(1, \pm 1, \pm 1)$  and  $(1, \pm 1, \mp 1)$

*Proof.* The argument is essentially identical to the proof of Lemma 4.10.  $\square$

*Remark 4.12.* In the proof of Lemma 4.10, we did not just prove that there is a unique element of  $\Gamma$  taking  $v = (a, b, c) \in \Delta_{\text{flop}}$  to a class with first coordinate 1: we proved there is a unique reduced word in the generators  $\iota_i^*$  and  $(\iota_i^\vee)^*$  (subject to  $(\iota_i^*)^2 = ((\iota_i^\vee)^*)^2 = 1$ ). From this, we deduce the relations on  $\Gamma$ .

**Corollary 4.13.** *There is an isomorphism*

$$\langle a_1, b_1, a_2, b_2 \mid a_1^2 = a_2^2 = b_1^2 = b_2^2 = 1 \rangle \xrightarrow{\sim} \Gamma$$

given by  $a_i \mapsto \iota_i^*$  and  $b_i \mapsto (\iota_i^\vee)^*$ .

We will return to the group  $\Gamma$  at the end of this section, proving that it is isomorphic to  $\text{Bir}(F)$  in Proposition 4.17.

**4.4. Birational geometry of  $F$ .** With our understanding of the geometry of  $\text{Mov}(F)$ , we conclude by enumerating the birational hyperkähler models of  $F$  up to isomorphism, completing the proof of Theorem 4.1. We also describe the birational automorphism group of  $F$  completely.

**Lemma 4.14.** *The only birational automorphism  $\varphi$  of  $F$  such that  $\varphi^* \text{Nef}(F) = \text{Nef}(F)$  is the identity.*

*Proof.* We know the walls of  $\text{Nef}(F)$  from Lemma 4.8, and the isometries permuting these walls and preserving the positive cone belong to the dihedral group  $D_8 \cong \langle R_2, R_3 \rangle$ . Inspecting actions on the discriminant group, Lemma 4.6 verifies that the only one of these isometries that can be induced by a birational automorphism of  $F$  is the identity. By Proposition 2.18(3), such a birational automorphism is thus an automorphism of  $F$ , and it is one that preserves the polarization  $g$ . By Remark 2.19, it must be the identity.  $\square$

**Proposition 4.15.** *The five birational models  $F$ ,  $F_i$ , and  $F_i^\vee$  for  $i = 1, 2$  are pairwise non-isomorphic.*

*Proof.* By Lemma 4.14, we verified that no isometry of  $\text{NS}(F)$  induced by a birational automorphism of  $F$  fixes  $\text{Nef}(F)$ ; from this, we see  $F$  has no nontrivial regular automorphisms. This distinguishes  $F$  from the four models  $F_i$  and  $F_i^\vee$ .

To see that  $F_1$  is non-isomorphic to the other four flops of  $F$ , first suppose  $\varphi \in \text{O}(\text{NS}(F))$  sends  $\text{Nef}(F_1)$  to  $\text{Nef}(F_2)$ . Then either  $\varphi$  or  $\iota_2^* \circ \varphi$  acts on  $\text{Nef}(F)$  by a nontrivial automorphism. Similarly, if  $\varphi$  sends  $\text{Nef}(F_1)$  to  $\text{Nef}(F_i^\vee)$ , then either  $\varphi$  or  $(\iota_i^\vee)^* \circ \varphi$  acts on  $\text{Nef}(F)$  by a nontrivial automorphism. Again using Lemma 4.14, we conclude that  $\varphi$  cannot be induced by a birational automorphism of  $F$ . In particular, no isomorphism exists between  $F_1$  and any of the other three flops of  $F$ .

A symmetric argument shows that any two flops of  $F$  are non-isomorphic.  $\square$

**Proposition 4.16.** *Up to isomorphism,  $F$  has exactly five birational hyperkähler models.*

*Proof.* By Proposition 4.15,  $F$  has at least five birational hyperkähler models, each of which can be obtained from  $F$  by flopping a plane in  $F$ . We will prove there are no more.

Any birational hyperkähler model of  $F$  can be obtained via a finite sequence of Mukai flops, shown in [WW03, Theorem 1.2], building on [BHL03, Theorem 1.1]. Starting from  $F$ , the Mukai flops are  $F_i$  and  $F_i^\vee$  for  $i = 1, 2$ . Using Lemmas 4.7, 4.8, and 4.9, we see by Remark 2.15 that the two Mukai flops of  $F_i$  (respectively  $F_i^\vee$ ) are both isomorphic to  $F$ . Thus any sequence of two Mukai flops starting from  $F$  yields a model isomorphic to  $F$ . Inductively, we see that any birational model of  $F$  can be obtained via a single Mukai flop.  $\square$

Together with the content of Section 3.3, identifying the flops of  $F$  with pairs of dual double EPW sextics, Propositions 4.15 and 4.16 complete the proof of Theorem 4.1.

As mentioned previously, Proposition 4.5 explains geometrically why, after flopping any one of the four planes in  $F$ , the other three planes cannot also be flopped. We see this in Figure 1, where the nef cones of  $F_i, F_i^\vee$  have two flopping walls and two prime exceptional walls.

We conclude by characterizing the birational automorphism group of  $F$ .

**Proposition 4.17.** *The involutions  $\iota_i^*$  and  $(\iota_i^\vee)^*$  for  $i = 1, 2$  generate the birational automorphism group of  $F$ , i.e.  $\Gamma \cong \text{Bir}(F)$ .*

*Proof.* Let  $\varphi \in \text{Bir}(F)$ , and let  $v^\perp$  be one of the walls of  $\varphi^* \text{Nef}(F)$ . By Lemma 4.10, there is some  $\psi \in \Gamma$  such that  $\psi(v)$  has first coordinate 1. We will argue that  $\varphi^* = \psi^{-1} \in \Gamma$ .

There are five chambers of  $\text{Nef}(F)$  having a wall of the form  $w^\perp$  where  $w \in \mathcal{W}_{\text{flop}}$  has first coordinate 1: they are  $\text{Nef}(F)$ ,  $\text{Nef}(F_i)$ , and  $\text{Nef}(F_i^\vee)$  for  $i = 1, 2$ , so  $\psi \circ \varphi^* \text{Nef}(F)$  is one of these five. Since  $\psi \circ \varphi^*$  is induced by a birational automorphism of  $F$ , if  $\psi \circ \varphi^* \text{Nef}(F) = \text{Nef}(F')$ , then  $F \simeq F'$ . Using Proposition 4.15, we conclude  $\psi \circ \varphi^* \text{Nef}(F) = \text{Nef}(F)$ , and Lemma 4.14 forces  $\psi \circ \varphi^* = 1$ .

It follows that  $\Gamma$  is the image of  $\text{Bir}(F) \rightarrow \text{O}(\text{NS}(F))$ . This map is an embedding by Remark 2.19, completing the proof.  $\square$

Along with Corollary 4.13, Proposition 4.17 proves Theorem 4.2.

5. A NON-SYZYGETIC PAIR OF CUBIC SCROLLS

Let  $X \subset \mathbb{P}^5$  be a smooth cubic fourfold containing a non-syzygetic pair of cubic scrolls. We also assume  $X$  is *very general*, meaning  $\text{rank}(A(X)) = 3$ ,  $\text{Aut}(X) = 0$ , and  $\text{End}_{\text{Hdg}}(T(F)) = \{\pm 1\}$ . As with the syzygetic case, we describe the birational geometry of the Fano variety  $F$  of lines on  $X$ , proving the following main result:

**Theorem 5.1.** *Let  $F$  be the Fano variety of lines on a very general cubic fourfold  $X$  containing a non-syzygetic pair of cubic scrolls. Then  $F$  has **eight** isomorphism classes of birational hyperkähler models, represented by the following:*

- $F$  itself,
- six non-isomorphic Mukai flops of  $F$ , each isomorphic to a double EPW sextic, and
- a Mukai flop of  $F$  along a pair of disjoint planes in  $F$ , isomorphic to the Fano variety of lines on another cubic fourfold containing a non-syzygetic pair of cubic scrolls.

As before, we also obtain generators for  $\text{Bir}(F)$ :

**Theorem 5.2.** *Let  $F$  be as above. There is a surjection*

$$\langle a, b, c, d, e, f \mid a^2 = b^2 = c^2 = d^2 = e^2 = f^2 = 1 \rangle \twoheadrightarrow \text{Bir}(F)$$

*identifying each generator with a covering involution on one of the double EPW sextics obtained as a Mukai flop of  $F$ .*

In contrast to Theorem 4.2, this surjection has nontrivial kernel; for further discussion, see Remark 5.17.

The outline of this section is similar to that of Section 4: in Section 5.1, we study the arrangement of Lagrangian planes in  $F$ , showing that some pairs of planes intersect and others do not. We outline in Section 5.2 properties of the lattice  $\text{NS}(F)$  and describe the actions on this lattice by the various involutions on the flops of  $F$ . This allows us to describe the structure of the movable cone of  $F$  in Section 5.3, which further enables a census of the birational hyperkähler models of  $F$  and a description of  $\text{Bir}(F)$  in Section 5.4.

**5.1. Lagrangian planes in  $F$ .** For a general cubic  $X$  containing a non-syzygetic pair of cubic scrolls  $T_1$  and  $T_2$ , we saw in Lemma 2.11 that there is a third cubic scroll  $T_3 \subset X$  and  $[T_i] \cdot [T_j] = 1$  for all  $i, j$ . Let  $H_i$  be the hyperplane spanned by  $T_i$ , and  $Y_i = X \cap H_i$ . As in Section 2.1, we have

$$F(Y_i) = P_i \cup S'_i \cup P_i^\vee \subset F,$$

where  $P_i$  and  $P_i^\vee$  are Lagrangian planes. For  $i \neq j$ , let  $\Sigma_{ij} = X \cap H_i \cap H_j$ ; for general  $X$ , all three cubic surfaces  $\Sigma_{ij}$  are smooth, by Lemma 2.8. The lines on  $\Sigma_{ij}$  are parametrized by  $F(Y_i) \cap F(Y_j)$ .

**Proposition 5.3.** *Let  $X$  be a cubic fourfold containing a non-syzygetic pair of cubic scrolls  $T_i$  and  $T_j$ , with  $[T_i] \cdot [T_j] = 1$ . Then*

- $\deg(P_i \cap P_j) = \deg(P_i^\vee \cap P_j^\vee) = 6$ ;
- $\deg(S'_i \cap S'_j) = 15$ ;
- all other pairwise intersections between a component of  $F(Y_i)$  and a component of  $F(Y_j)$  are empty.

Moreover, these intersections are all transverse when  $\Sigma = X \cap H_i \cap H_j$  is smooth.

*Proof.* As in the proof of Proposition 4.5, we use Lemmas 4.3 and 4.4 to reduce the problem to studying incidence relations between the twisted cubics  $\gamma_i, \gamma_j, \gamma_i^\vee, \gamma_j^\vee$ , and the lines on  $\Sigma_{ij}$ . After finding that

$$\gamma_i \sim \gamma_j \sim E_0 \quad \text{and} \quad \gamma_i^\vee \sim \gamma_j^\vee \sim 5E_0 - 2 \sum_{k=1}^6 E_k$$

on  $\Sigma_{ij}$ , the result follows from the incidence relations for lines on  $\Sigma_{ij}$ .  $\square$

In particular, there are six pairs of disjoint planes among the  $P_i$  and  $P_i^\vee$ , namely,  $P_i \cup P_j^\vee$  for  $i \neq j$ . This means that in addition to the six flops of  $F$  along a plane  $P_i$  or  $P_i^\vee$ , there are six models which can be obtained by flopping a pair of disjoint planes. Perhaps surprisingly, we will show in Lemma 5.12 that all six models of the last type are isomorphic to one another.

**5.2. The lattice  $\text{NS}(F)$ .** Again, let  $\lambda_i = \alpha(T_i - \eta_X)$ . Recall from Section 2.3 that  $\text{NS}(F)$  with the BBF form is isomorphic to the lattice  $J_{\text{non-syz}}$  and has discriminant group  $D_{\text{NS}(F)} \cong \mathbb{Z}/6 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/6$  with factors generated by  $[\frac{g}{6}]$ ,  $[\frac{\lambda_1}{2}]$ , and  $[\frac{\lambda_1}{3} + \frac{\lambda_2}{6}]$ . It is easy to check that this quadratic form represents  $-10$  but not  $-2$ .

The isometry group of  $\text{NS}(F)$  with basis  $\{g, \lambda_1, \lambda_2\}$  is generated by  $\pm 1$  and the four transformations below. We calculated  $\text{Aut}(\text{NS}(F))$  via [Mer14] and the Magma package `Authyp.m`.

$$R_1 = \begin{pmatrix} 5 & 4 & -2 \\ -6 & -5 & 2 \\ 0 & 0 & -1 \end{pmatrix}, \quad R_2 = \begin{pmatrix} 3 & 2 & -2 \\ -2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix},$$

$$R_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad R_4 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 1 & 0 \end{pmatrix}.$$

Note that  $R_1, R_2$ , and  $R_3$  are reflections whereas  $R_4$  has order 6. The transformations  $R_3$  and  $R_4$  generate the dihedral group  $D_{12}$ .

**Lemma 5.4.** *An isometry  $\varphi \in \text{O}(\text{NS}(F))$  is induced by a birational automorphism of  $F$  if and only if  $\varphi$  preserves  $\overline{\text{Pos}}(F)$  and acts on the index two subgroup*

$$\left\langle \left[ \frac{g}{3} \right] \right\rangle \oplus \left\langle \left[ \frac{\lambda_1}{2} \right] \right\rangle \oplus \left\langle \left[ \frac{\lambda_1}{3} + \frac{\lambda_2}{6} \right] \right\rangle$$

of  $D_{\text{NS}(F)}$  by  $\pm \text{Id}$ .

*Proof.* The proof is exactly the same as in Lemma 3.1.  $\square$

Let  $F_i$  and  $F_i^\vee$  denote the flops of  $F$  along  $P_i$  and  $P_i^\vee$ , respectively, for  $i = 1, 2, 3$ . Recall from Remark 3.3 that each of these models admits a regular involution  $\iota_i$  or  $\iota_i^\vee$ . These involutions can also be regarded as birational involutions on  $F$ .

**Lemma 5.5.** *The birational involutions  $\iota_i$  and  $\iota_i^\vee$  act on  $\text{NS}(F)$  by  $\iota_1^* = R_1, (\iota_1^\vee)^* = R_4^3 R_1 R_4^3, \iota_2^* = R_4^2 R_1 R_4^4, (\iota_2^\vee)^* = R_4^5 R_1 R_4, \iota_3^* = R_4^4 R_1 R_4^2, \text{ and } (\iota_3^\vee)^* = R_4 R_1 R_4^5$ .*

*Proof.* The proof is similar to that of Lemma 4.7, and we omit most details. By [HT10, Proposition 6.5] and (2.10),

$$\begin{aligned} \iota_1^*(1, -1, 0) &= (1, -1, 0) & (\iota_1^\vee)^*(1, 1, 0) &= (1, 1, 0) \\ \iota_2^*(1, 0, -1) &= (1, 0, -1) & (\iota_2^\vee)^*(1, 0, 1) &= (1, 0, 1) \\ \iota_3^*(1, 1, 1) &= (1, 1, 1) & (\iota_3^\vee)^*(1, -1, -1) &= (1, -1, -1). \end{aligned}$$

The only matrices satisfying these conditions, squaring to the identity, preserving the quadratic form, and acting by  $\pm \text{Id}$  on the index two subgroup of the discriminant group from Lemma 5.4 are the ones proposed.  $\square$

**5.3. Wall and chamber structure of  $\text{Mov}(F)$ .** As in the syzygetic case, we enumerate birational models of  $F$  by studying the geometry of the movable cone. Unlike before, there are no prime exceptional divisors, so  $\text{Mov}(F) = \overline{\text{Pos}(F)}$ .

**Lemma 5.6.** *The nef cone of  $F$  is bounded by the six walls  $v^\perp$  where*

$$v \in \{(1, 2, 0), (1, 2, 2), (1, 0, 2), (1, -2, 0), (1, -2, -2), (1, 0, -2)\}.$$

*These are the six classes with first coordinate 1 that square to  $-10$ .*

*Proof.* Similar to the proof of Lemma 4.8, we know each of these walls induces a small contraction of  $F$  so lies on the boundary of  $\text{Nef}(F)$ . Moreover, there are six pairs of disjoint Lagrangian planes in  $F$  by Proposition 5.3; therefore, the six edges of the cone

$$\text{Cone}((1, 2, 0)^\perp, (1, 2, 2)^\perp, (1, 0, 2)^\perp, (1, -2, 0)^\perp, (1, -2, -2)^\perp, (1, 0, -2)^\perp)$$

also lie on the boundary of  $\text{Nef}(F)$ . It follows that no other walls cut into  $\text{Nef}(F)$ .  $\square$

**Corollary 5.7.** *The six planes in  $F$  coming from components of  $F(Y_i)$  where  $Y_i = X \cap H_i$  for  $i = 1, 2, 3$  are the only Lagrangian planes in  $F$ .*

Using the involutions on the six flops of  $F$ , we can also describe the nef cones of the flops:

**Lemma 5.8.** *The nef cones of flops of  $F$  are*

$$\begin{aligned} \text{Nef}(F_1) &= \text{Cone}((1, -2, 0)^\perp, (1, 0, 2)^\perp, (3, -4, 0)^\perp, (1, -2, -2)^\perp), \\ \text{Nef}(F_1^\vee) &= \text{Cone}((1, 2, 0)^\perp, (1, 0, -2)^\perp, (3, 4, 0)^\perp, (1, 2, 2)^\perp), \\ \text{Nef}(F_2) &= \text{Cone}((1, 0, -2)^\perp, (1, 2, 0)^\perp, (3, 0, -4)^\perp, (1, -2, -2)^\perp), \\ \text{Nef}(F_2^\vee) &= \text{Cone}((1, 0, 2)^\perp, (1, -2, 0)^\perp, (3, 0, 4)^\perp, (1, 2, 2)^\perp), \\ \text{Nef}(F_3) &= \text{Cone}((1, 2, 2)^\perp, (1, 0, 2)^\perp, (3, 4, 4)^\perp, (1, 2, 0)^\perp), \\ \text{Nef}(F_3^\vee) &= \text{Cone}((1, -2, -2)^\perp, (1, -2, 0)^\perp, (3, -4, -4)^\perp, (1, 0, -2)^\perp). \end{aligned}$$

*Proof.* We give the argument for  $\text{Nef}(F_1)$ , the other calculations being similar. Since  $F_1$  is the flop of  $F$  along  $P_1$ , we know  $(1, -2, 0)^\perp$  lies on the boundary of  $\text{Nef}(F_1)$ . The involution  $\iota_1$  is regular on  $F_1$ , so  $\iota_1^*(1, -2, 0)^\perp = (3, -4, 0)^\perp$  also lies on the boundary of  $\text{Nef}(F_1)$  by Theorem 2.14. Moreover, the walls  $(1, 0, 2)^\perp$  and  $(1, -2, -2)^\perp$  meet each of  $(1, -2, 0)^\perp$  and  $(3, -4, 0)^\perp$ , and  $\iota_1^*(1, 0, 2)^\perp = (1, -2, -2)^\perp$ , so

$$\text{Nef}(F_1) \subset \text{Cone}((1, -2, 0)^\perp, (1, 0, 2)^\perp, (3, -4, 0)^\perp, (1, -2, -2)^\perp).$$

Since  $P_1 \cap P_i^\vee = \emptyset$  for  $i = 2, 3$  by Proposition 5.3, one can flop  $F_1$  along the strict transform of  $P_i^\vee$ , so the lines

$$(1, -2, 0)^\perp \cap (1, 0, 2)^\perp = \text{span}(2, -1, 1)$$

and

$$(1, -2, 0)^\perp \cap (1, -2, -2)^\perp = \text{span}(2, -2, -1)$$

lie on the boundary of  $\text{Nef}(F_1)$ . Their images, under  $\iota_1^*$ ,

$$\text{span}(4, -5, -1) = (1, -2, -2)^\perp \cap (3, -4, 0)^\perp$$

and

$$\text{span}(4, -4, 1) = (1, 0, 2)^\perp \cap (3, -4, 0)^\perp,$$

must also lie on the boundary of  $\text{Nef}(F_1)$ . It follows that

$$\text{Nef}(F_1) = \text{Cone}((1, -2, 0)^\perp, (1, 0, 2)^\perp, (3, -4, 0)^\perp, (1, -2, -2)^\perp),$$

as claimed.  $\square$

For  $1 \leq i, j \leq 3$ , let  $F_{ij}$  be the flop of  $F$  along  $P_i$  and  $P_j^\vee$ .

**Lemma 5.9.** *The nef cones of the  $F_{ij}$  are*

$$\begin{aligned} \text{Nef}(F_{12}) &= R_2 \cdot \text{Nef}(F), & \text{Nef}(F_{21}) &= \iota_2^*(\iota_3^\vee)^* \iota_1^* R_2 \cdot \text{Nef}(F), \\ \text{Nef}(F_{13}) &= \iota_1^* R_2 \cdot \text{Nef}(F), & \text{Nef}(F_{31}) &= \iota_3^*(\iota_2^\vee)^* R_2 \cdot \text{Nef}(F), \\ \text{Nef}(F_{23}) &= (\iota_3^\vee)^* \iota_1^* R_2 \cdot \text{Nef}(F), & \text{Nef}(F_{32}) &= (\iota_2^\vee)^* R_2 \cdot \text{Nef}(F). \end{aligned}$$

*Proof.* Since  $R_2$  preserves the BBF form, it acts by an automorphism on  $\text{Mov}(F)$ , sending chambers to chambers. It is straightforward to verify that the only four chambers of  $\text{Mov}(F)$  containing  $(2, -1, 1)$  are  $\text{Nef}(F)$ ,  $\text{Nef}(F_1)$ ,  $\text{Nef}(F_2^\vee)$ , and  $R_2 \cdot \text{Nef}(F)$ . The edge  $\text{span}(2, -1, 1)$  of  $\text{Nef}(F)$  corresponds to simultaneously flopping  $P_1$  and  $P_2^\vee$ , since

$$(1, -2, 0)^\perp \cap (1, 0, 2)^\perp = \text{span}(2, -1, 1).$$

Hence  $\text{Nef}(F_{12})$  contains  $(2, -1, 1)$ , from which we deduce  $\text{Nef}(F_{12}) = R_2 \cdot \text{Nef}(F)$ . The other five verifications are similar.  $\square$

**Corollary 5.10.** *The model  $F_{12}$  contains exactly six Lagrangian planes, six pairs of which are disjoint.*

*Proof.* By Lemmas 5.9 and 5.6, the nef cone of  $F_{12}$  is a cone over a hexagon, the same as  $\text{Nef}(F)$ . The six faces of this chamber correspond to Lagrangian planes in  $F_{12}$ ; the six edges correspond to pairs of disjoint Lagrangian planes in  $F_{12}$ .  $\square$

We illustrate the movable cone as described so far in Figure 3. In the next section, we will need to know the chambers bordering  $\text{Nef}(F_{12})$ , enumerated by the following lemma and illustrated analogously for  $\text{Nef}(F_{32})$  in Figure 4.

**Lemma 5.11.** *Let  $\{i, j, k\} = \{1, 2, 3\}$ . Then the chambers sharing a face with  $\text{Nef}(F_{ij})$  are  $\text{Nef}(F_i)$ ,  $\text{Nef}(F_j^\vee)$ ,  $\iota_i^* \text{Nef}(F_k^\vee)$ ,  $(\iota_j^\vee)^* \text{Nef}(F_k)$ ,  $(\iota_j^\vee)^* \iota_k^* \text{Nef}(F_i^\vee)$ , and  $\iota_i^*(\iota_k^\vee)^* \text{Nef}(F_j)$ .*

*Proof.* This is verified by direct calculation: using Lemmas 5.9 and 5.8, one can identify the walls of  $\text{Nef}(F_{ij})$  and  $\text{Nef}(F_i)$  for all  $i, j, k$ . Using Lemma 5.5, one can check that each of the chambers above shares a face with  $\text{Nef}(F_{ij})$ . By Corollary 5.10, these six faces cover the entire boundary of  $\text{Nef}(F_{ij})$ .  $\square$




FIGURE 3. A slice of the chambers of the movable cone of  $F$ , bounded by the positive cone.

5.4. **Birational geometry of  $F$ .** We now classify birational hyperkähler models of  $F$  up to isomorphism, proving Theorem 5.1.

**Lemma 5.12.** *The six birational models  $F_{ij}$  for  $1 \leq i, j \leq 3$  are all isomorphic.*

*Proof.* Since, by Theorem 2.14,  $\iota_i^*$  and  $(\iota_i^V)^*$  act on  $\text{Mov}(F)$  by exchanging chambers corresponding to the same isomorphism type, this follows from Lemma 5.9.  $\square$

**Proposition 5.13.** *The eight birational models  $F$ ,  $F_i$ ,  $F_i^V$ , and  $F_{12}$  are pairwise non-isomorphic.*

*Proof.* The idea of the proof is to show that no isometry of  $\text{NS}(F)$  taking a chamber of one of the models above to a chamber of another satisfies the conditions of Lemma 5.4. We begin with  $F_1$ . Since the chamber  $\text{Nef}(F_1)$  has four walls, whereas  $\text{Nef}(F)$  and  $\text{Nef}(F_{12})$  each have six walls, we know  $F_1 \not\cong F$  and  $F_1 \not\cong F_{12}$ .

Let  $\varphi$  be an isometry sending  $\text{Nef}(F_1)$  to  $\text{Nef}(F_i)$  for  $i \neq 1$  or  $\text{Nef}(F_i^V)$  for  $i = 1, 2, 3$ . The four edges of  $\text{Nef}(F_1)$ , corresponding to simultaneous flops of disjoint planes, are rays spanned by the vectors

$$v_1 = (2, -1, 1), v_2 = (2, -2, -1), v_3 = (4, -4, 1), \text{ and } v_4 = (4, -5, -1);$$

let  $w_i = \varphi(v_i)$ , so the vectors  $w_i$  span the edges of the chamber  $\varphi(\text{Nef}(F_1))$ . Then

$$\varphi(0, 1, 2) = \varphi(v_1 - v_2) = w_1 - w_2.$$




FIGURE 4. A detail of the figure above illustrating the chambers adjacent to  $F_3$  and  $F_{32}$ . Chambers are labeled by the isomorphism type of the model they represent.

For  $\varphi$  to be induced by a birational automorphism of  $F$ , we must have  $\varphi([\lambda_1/2]) = [\lambda_1/2]$  in the discriminant group, by Lemma 5.4. Hence  $\varphi(0, 1, 2) = (a, b, c)$  where  $a$  and  $c$  are even and  $b$  is odd. By direct inspection, the only chamber other than  $\text{Nef}(F_1)$  having walls  $w_1, w_2$  such that  $w_1 - w_2 = (a, b, c)$  with  $a$  and  $c$  even and  $b$  odd is  $\text{Nef}(F_1^\vee)$ . Hence  $F_1$  is not isomorphic to  $F_i$  or  $F_i^\vee$  for  $i = 2, 3$ .

We now verify  $F_1 \not\cong F_1^\vee$ . There are eight possibilities for  $w_1$  and  $w_2$ :

- |  |  |
|--|--|
| (a) $w_1 = (2, 2, 1), w_2 = (2, 1, -1)$  | (e) $w_1 = (4, 4, -1), w_2 = (4, 5, 1)$    |
| (b) $w_1 = (2, 1, -1), w_2 = (2, 2, 1)$  | (f) $w_1 = (4, 5, 1), w_2 = (4, 4, -1)$    |
| (c) $w_1 = (2, 1, -1), w_2 = (4, 4, -1)$ | (g) $w_1 = (4, 5, 1), w_2 = (2, 2, 1)$     |
| (d) $w_1 = (2, 2, 1), w_2 = (4, 5, 1)$   | (h) $w_1 = (4, 4, -1), w_2 = (2, 1, -1)$ . |

Since  $\iota_1^\vee$  is regular on  $F_1^\vee$ , and  $(\iota_1^\vee)^*(2, 2, 1) = (4, 4, -1)$  and  $(\iota_1^\vee)^*(2, 1, -1) = (4, 5, 1)$ , composing  $\varphi$  with  $(\iota_1^\vee)^*$  reduces cases (e), (f), (g), and (h) to (a), (b), (c), and (d), respectively. In case (a), we find  $\varphi = R_4R_3$ ; in (b), we find  $\varphi = R_4^3$ ; in case (c), we find  $\varphi = R_4^3R_2$ ; in case (d), we find  $\varphi = R_4R_3R_2$ . We again check the action of  $\varphi$  on the discriminant group, concluding by Lemma 5.4 that  $\varphi$  cannot be induced by a birational automorphism of  $F$ .

By symmetry, all six models  $F_i$  and  $F_i^\vee$  are pairwise non-isomorphic. It remains to check that  $F \not\cong F_{12}$ . Suppose for the sake of contradiction that  $\varphi$  is an isometry of  $\text{NS}(F)$  induced by an isomorphism  $F_{12} \xrightarrow{\sim} F$ , so  $\varphi(\text{Nef}(F)) = \text{Nef}(F_{12})$ . Then  $\varphi$  maps the six chambers  $F_i, F_i^\vee$  for  $i = 1, 2, 3$  to the six chambers sharing faces with  $\text{Nef}(F_{12})$ . In light of Lemma 5.11 and the above, we deduce  $\varphi(\text{Nef}(F_1)) = \text{Nef}(F_1)$  and  $\varphi(\text{Nef}(F_2^\vee)) = \text{Nef}(F_2^\vee)$ . Direct computation verifies that the only isometry with  $\varphi(\text{Nef}(F)) = F_{12}$  satisfying these conditions is  $R_2$ , but by Lemma 5.4,  $R_2$  is not induced by a birational automorphism of  $F$ , yielding a contradiction.  $\square$

**Proposition 5.14.** *Up to isomorphism,  $F$  has eight birational hyperkähler models.*

*Proof.* By Proposition 5.13,  $F$  has at least eight birational hyperkähler models, each of which can be obtained from  $F$  by flopping a single plane or a pair of disjoint planes. We will prove there are no more.

As in Proposition 4.15, any birational hyperkähler model of  $F$  can be obtained via a finite sequence of Mukai flops. Starting from  $F$ , the Mukai flops are  $F_i$  and  $F_i^\vee$  for  $i = 1, 2, 3$ . Using Lemmas 5.8 and 5.9, the four Mukai flops of  $F_i$  (respectively  $F_i^\vee$ ) are isomorphic to  $F$ ,  $F_{ij}$ , and  $F_{ik}$  (respectively  $F$ ,  $F_{ji}$ , and  $F_{ki}$ ). Applying Lemma 5.12, we see that a sequence of two Mukai flops starting at  $F$  yields a model isomorphic either to  $F$  or to  $F_{12}$ . By Lemma 5.11, a third flop yields a model isomorphic to  $F_i$  or  $F_i^\vee$  for some  $i = 1, 2, 3$ . Inductively, we conclude that an odd number of Mukai flops starting from  $F$  yields a model isomorphic to  $F_i$  or  $F_i^\vee$ , and an even number of Mukai flops yields a model isomorphic to  $F$  or  $F_{12}$ ; in particular, any birational hyperkähler model of  $F$  is isomorphic to one of these.  $\square$

The content of Section 3.3 identifies each flop of  $F$  with a double EPW sextic. To complete the proof of Theorem 5.1, we identify  $F_{12}$  with the Fano variety of lines on another smooth cubic fourfold:

**Proposition 5.15.** *There is a unique smooth cubic fourfold  $X'$  containing a non-syzygetic pair of cubic scrolls whose Fano variety of lines is isomorphic to  $F_{12}$ . Moreover,  $X \not\cong X'$ .*

*Proof.* Lemma 5.9 states  $\text{Nef}(F_{12}) = R_2 \cdot \text{Nef}(F)$ , so  $F_{12}$  contains an ample class of square 6 and divisibility 2, namely  $g' = R_2 \cdot g$ . Hence the pair  $(F_{12}, g')$  defines a point in the moduli space  $\mathcal{M}_6^{(2)}$  of hyperkähler fourfolds of K3<sup>[2]</sup>-type with a polarization of square 6 and divisibility 2. Let  $\mathcal{M}_{cub}$  be the moduli space of marked cubic fourfolds, so there is a commutative diagram of period maps

$$\begin{array}{ccc} \mathcal{M}_{cub} & \xrightarrow{F} & \mathcal{M}_6^{(2)} \\ & \searrow p' & \swarrow p \\ & & \mathcal{P}_6^{(2)} \end{array}$$

By [Laz09] and [Loo09], the complement of the image of  $p'$  is the union of the Heegner divisors  $\mathcal{D}_{6,2}^{(2)}$  and  $\mathcal{D}_{6,6}^{(2)}$ ; for more discussion, see [Deb20, Appendix B]. Hence to show that  $F_{12}$  is the Fano variety of lines on a smooth cubic fourfold, it suffices to check that there are no classes  $v \in g'^\perp \cap \text{NS}(F_{12})$  with  $\text{div}(v) = 2$  and  $v^2 \in \{-2, -6\}$ . Indeed, if such a vector existed, then  $R_2 \cdot v$  would have the same numerics and lie in  $g^\perp \cap \text{NS}(F)$ ; recalling the Abeli-Jacobi map described in Section 2.3, this would force  $X$  to be of discriminant 2 or 6. But  $X$  is smooth, so  $X \notin \mathcal{C}_2 \cup \mathcal{C}_6$ , and no such class  $v$  exists.

We have shown  $F_{12}$  is the Fano variety of a smooth cubic fourfold  $X'$ . Moreover,  $X'$  is unique since  $p'$  is injective [Cha12, Proposition 6]. The Abel-Jacobi map allows us to deduce the intersection form on  $A(X')$ , proving that  $X'$  contains a non-syzygetic pair of cubic scrolls. Finally, the fact from Proposition 5.13 that  $F \not\cong F_{12}$  forces  $X \not\cong X'$ .  $\square$

We end by providing some information about the structure of  $\text{Bir}(F)$ . Let

$$\Gamma = \langle \iota_i^*, (\iota_i^\vee)^* \mid i = 1, 2, 3 \rangle \subset \text{O}(\text{NS}(F)).$$

Unlike in the syzygetic case, we are not able to make use of the action of  $\Gamma$  on  $\Delta_{\text{Hop}}$  in order to deduce generators and relations for  $\text{Bir}(F)$ ; largely, this is because in the syzygetic case, each wall of  $\text{Mov}(F)$  was adjacent to finitely many chambers whereas in the non-syzygetic case, each wall borders infinitely many chambers. On the other hand, the fact that the movable cone coincides with the positive cone somewhat streamlines the argument in the following lemma, analogous to Lemma 4.10.

**Lemma 5.16.** *The group  $\Gamma$  acts on the set  $\{v \in \text{NS}(F) \mid v^2 = 6\}$  with at most seven orbits, represented by the classes  $(1, 0, 0)$ ,  $(3, \pm 2, \pm 4)$ ,  $(3, \pm 4, \pm 2)$ , and  $(3, \pm 2, \mp 2)$ .*

*Proof.* As in Lemma 4.10, one starts with an arbitrary class  $v = (a, b, c)$  such that  $v^2 = 6$  and  $a > 3$  and finds that at least one of the involutions  $(\iota_i)^*$  or  $(\iota_i^\vee)^*$  for  $i = 1, 2, 3$  reduces the magnitude of the first coordinate. Iterating this process, one obtains one of the seven classes  $v \in \text{NS}(F)$  with  $v^2 = 6$  and first coordinate no larger than 3.  $\square$

*Remark 5.17.* Unlike in the syzygetic case, there is not a unique sequence of  $(\iota_i)^*$  and  $(\iota_i^\vee)^*$  taking an arbitrary class  $v \in \text{NS}(F)$  with  $v^2 = 6$  to one with first coordinate at most 3: indeed, for  $\{i, j, k\} = \{1, 2, 3\}$ , we note  $\iota_i \circ \iota_j^\vee \circ \iota_k = \iota_k^\vee \circ \iota_j \circ \iota_i^\vee$ . In particular, the action of Lemma 5.16 does not afford a characterization of the relations among the generators of  $\Gamma$ . Nevertheless, we obtain generators for  $\text{Bir}(F)$ .

**Proposition 5.18.** *The birational involutions  $\iota_i^*$  and  $(\iota_i^\vee)^*$  for  $i = 1, 2, 3$  generate the birational automorphism group of  $F$ , i.e.  $\Gamma \cong \text{Bir}(F)$ .*

*Proof.* Let  $\varphi \in \text{Bir}(F)$ , and let  $\varphi^* \text{Nef}(F) = \text{Nef}(F')$ . Then  $\varphi^*(g) = v$  for some class  $v$  with  $q(v) = 6$ . By Lemma 5.16, there is some  $f \in \Gamma$  such that  $f(v)$  has first coordinate at most 3. Moreover, since the generators of  $\Gamma$  are induced by birational automorphisms, we have  $f = \psi^*$  for some  $\psi \in \text{Bir}(F)$ . Now,

$$(\varphi \circ \psi)^*(g) \in \{(1, 0, 0), (3, \pm 2, \pm 4), (3, \pm 4, \pm 2), (3, \pm 2, \mp 2)\},$$

but on the other hand,  $(\varphi \circ \psi)^* \text{Nef}(F) = \text{Nef}(F'')$  where  $F'' \simeq F$ . Using Lemma 5.9, the classes  $(3, \pm 2, \pm 4)$ ,  $(3, \pm 4, \pm 2)$ , and  $(3, \pm 2, \mp 2)$  belong to the nef cones of  $F_{ij}$  for  $1 \leq i \neq j \leq 3$ , so by Proposition 5.13 we obtain  $(\varphi \circ \psi)^*(g) = g$ . The subgroup of  $\text{O}(\text{NS}(F))$  of isometries fixing  $g$  is the dihedral group generated by  $R_3$  and  $R_4$ , and using Lemma 5.4, the only one of these isometries induced by a birational automorphism of  $F$  is the identity. By Remark 2.19, the map  $\text{Bir}(F) \rightarrow \text{O}(\text{NS}(F))$  is an embedding, so  $\varphi \circ \psi = \text{Id}$ , and  $\varphi = \psi^{-1} \in \Gamma$ .  $\square$

This completes the proof of Theorem 5.2.

## APPENDIX A. EXAMPLES

Here, we provide explicit examples of cubic fourfolds containing pairs of cubic scrolls in order to justify earlier assertions about generic behavior (cf. Lemma 2.8). The computational claims in the proof below can be verified with Magma code provided on the arXiv as an ancillary file.

We work over the field  $\mathbb{F}_{29}$ , but the choices made in producing our examples amount to picking a point in a tower of projective bundles, as can be seen in the Magma code. Hence our examples lift to characteristic zero.

**A.1. An explicit cubic fourfold with a syzygetic pair of cubic scrolls.** Let  $X$  be the smooth cubic fourfold with defining equation

$$\begin{aligned} f = & 17x_0x_1x_2 + 19x_1^2x_2 + 9x_0x_2^2 + 10x_1x_2^2 + 18x_2^3 + 12x_0^2x_3 + 10x_0x_1x_3 + 8x_0x_2x_3 \\ & + 4x_1x_2x_3 + 27x_2^2x_3 + 2x_0x_3^2 + 3x_2x_3^2 + 20x_0^2x_4 + 11x_0x_1x_4 \\ & + 23x_1^2x_4 + 11x_0x_2x_4 + 24x_1x_2x_4 + 14x_2^2x_4 + 7x_0x_3x_4 + 26x_1x_3x_4 \\ & + 19x_2x_3x_4 + 15x_0x_4^2 + 10x_1x_4^2 + 7x_0^2x_5 + 16x_0x_1x_5 + 18x_1^2x_5 \\ & + 22x_0x_3x_5 + 8x_1x_3x_5 + 23x_3^2x_5 + 18x_0x_4x_5 + 5x_1x_4x_5 + 7x_3x_4x_5 \\ & + 22x_4^2x_5 + 21x_1x_5^2 + 5x_3x_5^2 + 28x_4x_5^2 + 2x_5^3. \end{aligned}$$

The hyperplanes  $H_1$  and  $H_2$  defined by  $x_5 = 0$  and  $x_2 = 0$ , respectively, intersect  $X$  in six-nodal cubic threefolds  $Y_1$  and  $Y_2$ . The cubic scroll  $T_1 \subset H_1$ , defined by the vanishing of the minors of the matrix

$$M_1 = \begin{pmatrix} x_0 & x_1 & x_2 \\ x_2 & x_3 & x_4 \end{pmatrix},$$

is contained in  $Y_1$ . Similarly, the cubic scroll  $T_2 \subset H_2$ , defined by the vanishing of the minors of the matrix

$$M_2 = \begin{pmatrix} l_0 & l_1 & l_2 \\ l_2 & l_3 & l_4 \end{pmatrix},$$

where

$$\begin{aligned} l_0 &:= 17x_0 + 12x_1 + 12x_2 + 17x_3 + 7x_4 + 6x_5, \\ l_1 &:= 17x_0 + 4x_1 + 17x_2 + 25x_3 + 18x_4 + 13x_5, \\ l_2 &:= x_5, \\ l_3 &:= 10x_0 + 13x_1 + 12x_2 + 15x_3 + 14x_4 + 17x_5, \\ l_4 &:= 16x_0 + 13x_1 + 9x_2 + 10x_3 + 19x_4 + 7x_5, \end{aligned}$$

is contained in  $Y_2$ . The pair of cubic scrolls is syzygetic;  $T_1$  and  $T_2$  intersect transversely in three points, namely  $(1 : 1 : 0 : 0 : 0 : 0)$ ,  $(0 : 0 : 0 : 1 : 1 : 0)$ , and  $(0 : 1 : 0 : 1 : 0 : 0)$ . The cubic surface  $\Sigma = Y_1 \cap Y_2$  is smooth, as desired.

**A.2. An explicit cubic fourfold with a non-syzygetic pair of cubic scrolls.** Let  $X$  be the smooth cubic fourfold with defining equation

$$\begin{aligned} f = & 20x_0x_1x_2 + 4x_1^2x_2 + 15x_0x_2^2 + 17x_1x_2^2 + 6x_2^3 + 9x_0^2x_3 + 25x_0x_1x_3 + 27x_0x_2x_3 \\ & + 15x_1x_2x_3 + 19x_2^2x_3 + 5x_0x_3^2 + 19x_2x_3^2 + 14x_0^2x_4 + 14x_0x_1x_4 \\ & + 9x_1^2x_4 + 23x_0x_2x_4 + 25x_1x_2x_4 + 21x_2^2x_4 + 14x_0x_3x_4 + 10x_1x_3x_4 \\ & + 18x_2x_3x_4 + 8x_0x_4^2 + 11x_1x_4^2 + 10x_0^2x_5 + 4x_0x_1x_5 + 24x_1^2x_5 \\ & + 22x_0x_3x_5 + 16x_1x_3x_5 + 17x_3^2x_5 + 5x_0x_4x_5 + 18x_1x_4x_5 + 25x_3x_4x_5 \\ & + 27x_4^2x_5 + 2x_0x_5^2 + 28x_1x_5^2 + 21x_3x_5^2 + 28x_4x_5^2 + 13x_5^3. \end{aligned}$$

We consider the following hyperplanes:

$$\begin{aligned} H_1 &:= \{x_5 = 0\}, \\ H_2 &:= \{x_2 = 0\}, \\ H_3 &:= \{x_0 + 24x_1 + x_2 + x_3 + 20x_4 + 9x_5 = 0\}. \end{aligned}$$

One sees that  $H_1, H_2$  and  $H_3$  intersect  $X$  in six-nodal cubic threefolds  $Y_1, Y_2$ , and  $Y_3$  respectively. The cubic scroll  $T_1 \subset H_1$  defined by the vanishing of the minors of the matrix

$$M_1 = \begin{pmatrix} x_0 & x_1 & x_2 \\ x_2 & x_3 & x_4 \end{pmatrix},$$

is contained in  $Y_1$ . The cubic scroll  $T_2 \subset H_2$ , defined by the vanishing of the minors of the matrix

$$M_2 = \begin{pmatrix} l_0 & l_1 & l_2 \\ l_2 & l_3 & l_4 \end{pmatrix},$$

where

$$\begin{aligned} l_0 &:= 26x_0 + 2x_1 + 8x_2 + 4x_3 + 5x_4 + 24x_5, \\ l_1 &:= 13x_0 + 11x_1 + 22x_2 + 18x_3 + 6x_4 + 15x_5, \\ l_2 &:= 12x_0 + 19x_1 + 15x_2 + 16x_3 + 17x_4 + 14x_5, \\ l_3 &:= 18x_0 + 3x_1 + 18x_2 + 26x_3 + 18x_4 + 10x_5, \\ l_4 &:= 28x_0 + 14x_1 + 5x_2 + 21x_3 + x_4 + 3x_5. \end{aligned}$$

is contained in  $Y_2$ . The cubic scroll  $T_3 \subset H_3$ , cut out by the quadrics

$$\begin{aligned} Q_{31} &:= x_1^2 + 17x_1x_3 + 27x_2x_3 + 9x_3^2 + 27x_1x_4 + 23x_2x_4 + 11x_3x_4 + 14x_4^2 + 24x_1x_5 \\ &\quad + 13x_2x_5 + 10x_3x_5 + 2x_4x_5 + 8x_5^2, \\ Q_{32} &:= x_1x_2 + 25x_1x_3 + 20x_2x_3 + 5x_3^2 + 5x_1x_4 + 6x_2x_4 + 23x_3x_4 + 5x_4^2 + 20x_1x_5 \\ &\quad + 2x_2x_5 + 24x_3x_5 + 3x_4x_5 + 8x_5^2, \\ Q_{33} &:= x_2^2 + 28x_1x_3 + 5x_2x_3 + 25x_3^2 + 20x_1x_4 + 14x_2x_4 + 15x_3x_4 + x_4^2 + 27x_1x_5 \\ &\quad + 15x_2x_5 + 10x_3x_5 + 2x_4x_5 + x_5^2, \end{aligned}$$

is contained in  $Y_3$ . The pairs  $(T_1, T_2)$ ,  $(T_1, T_3)$ , and  $(T_2, T_3)$  all form non-syzygetic pairs. Indeed, any two intersect transversely in one point; explicitly,

$$\begin{aligned} T_1 \cap T_2 &= \{(0 : 1 : 0 : 1 : 0 : 0)\}, \\ T_1 \cap T_3 &= \{(22 : 19 : 15 : 9 : 1 : 0)\}, \\ T_2 \cap T_3 &= \{(15 : 9 : 0 : 15 : 9 : 1)\}. \end{aligned}$$

To verify that  $[T_3] = 3\eta_X - [T_1] - [T_2]$ , it suffices to check  $[T_3] \in \langle \eta_X, [T_1], [T_2] \rangle$ . If not, then each of the non-syzygetic pairs gives rise to another  $\mathbb{F}_{29}$ -rational hyperplane slicing  $X$  in a cubic threefold singular along at least a length 6 zero-dimensional subscheme. In that case,  $X$  has at least six such hyperplane sections. Direct computation verifies that  $X$  has only four six-nodal hyperplane sections, cut out by  $H_1, H_2, H_3$ , and

$$x_0 + 16x_1 + 8x_2 + 11x_3 + 26x_4 + 13x_5 = 0.$$

So,  $T_3$  represents the desired class in cohomology.

Now, let  $\Sigma_{ij} = Y_i \cap Y_j$ . All three cubic surfaces  $\Sigma_{ij}$  are smooth, so the same is true for a general cubic fourfold containing a non-syzygetic pair of cubic scrolls.

## REFERENCES

- [AV17] Ekaterina Amerik and Misha Verbitsky, *Morrison-Kawamata cone conjecture for hyperkähler manifolds*, Ann. Sci. Éc. Norm. Supér. (4) **50** (2017), no. 4, 973–993. MR3679618
- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265. Computational algebra and number theory (London, 1993). MR1484478
- [BD85] Arnaud Beauville and Ron Donagi, *La variété des droites d'une hypersurface cubique de dimension 4*, C. R. Acad. Sci. Paris Sér. I Math. **301** (1985), no. 14, 703–706. MR818549
- [BFvGS24] Eva Bayer-Fluckiger, Bert van Geemen, and Matthias Schütt, *K3 surfaces with real or complex multiplication* (2024), available at [2401.04072](#).
- [BHL03] Dan Burns, Yi Hu, and Tie Luo, *HyperKähler manifolds and birational transformations in dimension 4*, Vector bundles and representation theory (Columbia, MO, 2002), 2003, pp. 141–149. MR1987745
- [BHT15] Arend Bayer, Brendan Hassett, and Yuri Tschinkel, *Mori cones of holomorphic symplectic varieties of K3 type*, Ann. Sci. Éc. Norm. Supér. (4) **48** (2015), no. 4, 941–950. MR3377069
- [Cha12] François Charles, *A remark on the Torelli theorem for cubic fourfolds* (2012), available at [1209.4509](#).
- [CTZ24] Ivan Cheltsov, Yuri Tschinkel, and Zhijia Zhang, *Equivariant geometry of singular cubic threefolds* (2024), available at [2401.10974](#).
- [Deb20] Olivier Debarre, *Hyperkähler manifolds* (2020), available at [1810.02087](#).
- [DIM15] O. Debarre, A. Iliev, and L. Manivel, *Special prime Fano fourfolds of degree 10 and index 2*, Recent advances in algebraic geometry, 2015, pp. 123–155. MR3380447
- [DK18] Olivier Debarre and Alexander Kuznetsov, *Gushel-Mukai varieties: classification and birationalities*, Algebr. Geom. **5** (2018), no. 1, 15–76. MR3734109
- [Dol12] Igor V. Dolgachev, *Classical algebraic geometry*, Cambridge University Press, Cambridge, 2012. A modern view. MR2964027
- [Dol16] Igor Dolgachev, *Corrado Segre and nodal cubic threefolds*, From classical to modern algebraic geometry, 2016, pp. 429–450. MR3776662
- [Fuj81] Akira Fujiki, *A theorem on bimeromorphic maps of Kähler manifolds and its applications*, Publ. Res. Inst. Math. Sci. **17** (1981), no. 2, 735–754. MR642659
- [GAL11] Víctor González-Aguilera and Alvaro Liendo, *Automorphisms of prime order of smooth cubic  $n$ -folds*, Arch. Math. (Basel) **97** (2011), no. 1, 25–37. MR2820585
- [Has00] Brendan Hassett, *Special cubic fourfolds*, Compositio Math. **120** (2000), no. 1, 1–23. MR1738215
- [Has96] Brendan Edward Hassett, *Special cubic hypersurfaces of dimension four*, ProQuest LLC, Ann Arbor, MI, 1996. Thesis (Ph.D.)—Harvard University. MR2694332
- [HT09] Brendan Hassett and Yuri Tschinkel, *Moving and ample cones of holomorphic symplectic fourfolds*, Geom. Funct. Anal. **19** (2009), no. 4, 1065–1080. MR2570315
- [HT10] Brendan Hassett and Yuri Tschinkel, *Flops on holomorphic symplectic fourfolds and determinantal cubic hypersurfaces*, J. Inst. Math. Jussieu **9** (2010), no. 1, 125–153. MR2576800
- [IM11] Atanas Iliev and Laurent Manivel, *Fano manifolds of degree ten and EPW sextics*, Ann. Sci. Éc. Norm. Supér. (4) **44** (2011), no. 3, 393–426. MR2839455
- [KP18] Alexander Kuznetsov and Alexander Perry, *Derived categories of Gushel-Mukai varieties*, Compos. Math. **154** (2018), no. 7, 1362–1406. MR3826460
- [KP23] Alexander Kuznetsov and Alexander Perry, *Categorical cones and quadratic homological projective duality*, Ann. Sci. Éc. Norm. Supér. (4) **56** (2023), no. 1, 1–57. MR4563868
- [Laz09] Radu Laza, *The moduli space of cubic fourfolds*, J. Algebraic Geom. **18** (2009), no. 3, 511–545. MR2496456
- [Loo09] Eduard Looijenga, *The period map for cubic fourfolds*, Invent. Math. **177** (2009), no. 1, 213–233. MR2507640

- [Mar11] Eyal Markman, *A survey of Torelli and monodromy results for holomorphic-symplectic varieties*, Complex and differential geometry, 2011, pp. 257–322. MR2964480
- [Mar23] Lisa Marquand, *Cubic fourfolds with an involution*, Trans. Amer. Math. Soc. **376** (2023), no. 2, 1373–1406. MR4531678
- [Mer14] Michael H. Mertens, *Automorphism groups of hyperbolic lattices*, J. Algebra **408** (2014), 147–165. MR3197177
- [Mon15] Giovanni Mongardi, *A note on the Kähler and Mori cones of hyperkähler manifolds*, Asian J. Math. **19** (2015), no. 4, 583–591. MR3423735
- [Muk89] Shigeru Mukai, *Biregular classification of Fano 3-folds and Fano manifolds of coindex 3*, Proc. Nat. Acad. Sci. U.S.A. **86** (1989), no. 9, 3000–3002. MR995400
- [MY15] Eyal Markman and Kota Yoshioka, *A proof of the Kawamata-Morrison cone conjecture for holomorphic symplectic varieties of  $K3^{[n]}$  or generalized Kummer deformation type*, Int. Math. Res. Not. IMRN **24** (2015), 13563–13574. MR3436156
- [Nik79] V. V. Nikulin, *Integer symmetric bilinear forms and some of their geometric applications*, Izv. Akad. Nauk SSSR Ser. Mat. **43** (1979), no. 1, 111–177, 238. MR525944
- [O’G05] K. G. O’Grady, *Involutions and linear systems on holomorphic symplectic manifolds*, Geom. Funct. Anal. **15** (2005), no. 6, 1223–1274. MR2221247
- [O’G06] Kieran G. O’Grady, *Irreducible symplectic 4-folds and Eisenbud-Popescu-Walter sextics*, Duke Math. J. **134** (2006), no. 1, 99–137. MR2239344
- [O’G08a] Kieran G. O’Grady, *Dual double EPW-sextics and their periods*, Pure Appl. Math. Q. **4** (2008), no. 2, 427–468. MR2400882
- [O’G08b] Kieran G. O’Grady, *Irreducible symplectic 4-folds numerically equivalent to  $(K3)^{[2]}$* , Commun. Contemp. Math. **10** (2008), no. 4, 553–608. MR2444848
- [WW03] Jan Wierzbka and Jarosław A. Wiśniewski, *Small contractions of symplectic 4-folds*, Duke Math. J. **120** (2003), no. 1, 65–95. MR2010734

DEPARTMENT OF MATHEMATICS AND STATISTICS, CARLETON COLLEGE, NORTHFIELD, MN 55057  
*Email address:* [cbrooke@carleton.edu](mailto:cbrooke@carleton.edu)

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, KEMENY HALL, HANOVER, NH 03755  
*Email address:* [sarah.frei@dartmouth.edu](mailto:sarah.frei@dartmouth.edu)

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012, USA  
*Email address:* [lisa.marquand@nyu.edu](mailto:lisa.marquand@nyu.edu)

DEPARTMENT OF MATHEMATICS, 329 PHILLIPS HALL, CHAPEL HILL, NC 27599  
*Email address:* [russellqin@gmail.com](mailto:russellqin@gmail.com)