

CENTROIDS AND EQUILIBRIUM POINTS OF CONVEX BODIES

ZSOLT LÁNGI AND PÉTER L. VÁRKONYI

ABSTRACT. The aim of this note is to survey the results in some geometric problems related to the centroids and the static equilibrium points of convex bodies. In particular, we collect results related to Grünbaum’s inequality and the Busemann-Petty centroid inequality, describe classifications of convex bodies based on equilibrium points, and investigate the location and structure of equilibrium points, their number with respect to a general reference point as well as the static equilibrium properties of convex polyhedra.

1. INTRODUCTION

The investigation of the centroids and static equilibrium points of solids has been in the focus of research since ancient Greece. These concepts play an important role in many different disciplines from physics to engineering to geology, and also appear in many problems in pure mathematics. The aim of this paper is to collect the results in the geometric literature related to some of these problems.

Apart from a few remarks, the objects of our paper are convex bodies in the d -dimensional Euclidean space \mathbb{R}^d , defined as compact, convex sets with nonempty interior. We denote the standard inner product of this space by $\langle \cdot, \cdot \rangle$, and the Euclidean norm of a vector $p \in \mathbb{R}^d$ by $|p| = \sqrt{\langle p, p \rangle}$. The *closed segment* with endpoints $p, q \in \mathbb{R}^d$ is denoted as $[p, q]$. An important convex body in \mathbb{R}^d is the closed unit ball centered at the origin o , which we denote by \mathbf{B}^d . The boundary of this ball is denoted by \mathbb{S}^{d-1} . We use the standard notation $\text{int}(X)$, $\text{bd}(X)$, $\text{conv}(X)$, $\text{vol}_d(X)$ for the *interior*, *boundary*, *convex hull* and d -dimensional *volume* of a set $X \in \mathbb{R}^d$. Two-dimensional volume is also called *area*, denoted by $\text{area}(\cdot)$, and 3-dimensional volume is denoted by $\text{vol}(\cdot)$. For any $s > 0$, we set $\kappa_s = \frac{\pi^{s/2}}{\Gamma(\frac{s}{2}+1)}$, where $\Gamma(x) = \int_0^\infty t^{x-1}e^{-t} dt$ is the usual Gamma function. Then, for any nonnegative integer d , $\kappa_d = \text{vol}_d(\mathbf{B}^d)$.

As usual, the orthogonal complement of a linear subspace L of \mathbb{R}^d is denoted by L^\perp , and the orthogonal projection of a set X onto L is denoted by $X|L$. For brevity, if L is a line generated by a nonzero vector u , we set $u^\perp = L^\perp$, $K|u = K|L$, and $K|u^\perp = K|L^\perp$. Finally, for any $u \in \mathbb{S}^{d-1}$ and $\alpha \in \mathbb{R}$, we denote the hyperplane

2020 *Mathematics Subject Classification.* 52A20, 52A38.

Key words and phrases. centroid, equilibrium point, Grünbaum’s inequality, Busemann-Petty centroid inequality, Gömböc, monostable polyhedra.

The authors thank Tímea Szabó and Gábor Domokos for their permissions to use their images of bodies in various equilibrium classes (Fig. 1). Zsolt Lángi is partially supported by the ERC Advanced Grant “ERMID”, and the NKFIH grant K147544. P. L. Várkonyi is supported by the NKFIH grant 143175.

$\{x \in \mathbb{R}^d : \langle x, u \rangle = \alpha\}$ by $H(u, \alpha)$. In this notation u is a *normal vector* of $H(u, \alpha)$ and $|\alpha|$ is its distance from o .

Let K be a convex body with $o \in \text{int}(K)$. The *radial function* $\rho_K : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$ of K is defined as

$$(1.1) \quad \rho_K(u) = \max\{\alpha \in \mathbb{R} : \alpha u \in K\},$$

and its *support function* $h_K : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$ as

$$(1.2) \quad h_K(u) = \max\{\langle x, u \rangle : x \in K\}.$$

The latter function is often defined for any $u \in \mathbb{R}^d$ using the formula in (1.2). To distinguish these two functions, we use the notation $\bar{h}_K : \mathbb{R}^d \rightarrow \mathbb{R}$ for the one extended to \mathbb{R}^d . We note that according to our definition, the function \bar{h}_K is a positively homogeneous, convex, nonnegative function which is zero only at o . It is also known [100] that any function satisfying these properties is the support function of some convex body K with $o \in \text{int}(K)$.

A set $S \subseteq \mathbb{R}^d$ is called *star-shaped* with respect to p , if for any $q \in S$, we have $[p, q] \subseteq S$. In particular, a convex set K is star-shaped with respect to any $p \in K$. We define the radial function $\rho_S : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$ of any compact set, star-shaped with respect to o , by the formula in (1.1). Such a set is called a *star body* if its radial function is strictly positive and continuous.

In the remaining part of the paper we present the results related to our basic concepts. In particular, in Section 2 we investigate the centroids of convex bodies, and discuss results related to two important problems concerning them: Grünbaum's centroid inequality in Subsection 2.1, and the Busemann-Petty centroid inequality in Subsection 2.2. In Section 3 we study various problems related to the static equilibrium points of convex bodies. Within this, in Subsection 3.1 we present a classification system of convex bodies based on the number of their equilibria. In Subsection 3.2 we refine this system by distinguishing convex bodies based on the topological properties of the gradient flow of their radial functions. In Subsection 3.3 we investigate how difficult it is to move a convex body from one class to another one. In Subsections 3.4, 3.5 and 3.6 we deal with the locations of equilibrium points on a convex body, equilibria with respect to a general reference point, and equilibria of convex polyhedra. Finally, in Section 4, we collect some applications and problems related to the presented topics.

2. THE CENTROID OF A CONVEX BODY

The main concept of this section is the following.

Definition 2.1. Let X be a compact set in \mathbb{R}^d with nonempty interior. Then the *centroid* of X is the point

$$c(X) = \frac{1}{\text{vol}_d(X)} \int_X x \, dx.$$

In the above definition, if $c(X) = o$, we say that X is *centered*. A concept related to the centroid of a set is its first moment with respect to a hyperplane.

Definition 2.2. Let X be a compact set in \mathbb{R}^d with nonempty interior, and let $u \in \mathbb{S}^{d-1}$, $\alpha \in \mathbb{R}$. Then the quantity

$$(2.1) \quad \int_X (\langle u, x \rangle - \alpha) \, dx$$

is called the *first moment* of K with respect to the hyperplane $H(u, \alpha)$.

We note that the quantity $\langle u, x \rangle - \alpha$ in (2.1) is equal to the signed distance of x from $H(u, \alpha)$, according to the orientation induced by u . From Definitions 2.1 and 2.2 it follows that the first moment of X with respect to $H(u, \alpha)$ is equal to $\text{vol}_d(X)c(X)$, that is, it is equal to that of a point mass at $c(X)$ with weight $\text{vol}_d(X)$. While the notion of centroid in non-Euclidean spaces is outside the scope of this paper, we remark that an axiomatic generalization of centroid in various manifolds including hyperbolic and spherical spaces, based partly on this observation, can be found in [46]. For more information on centroids in spherical or hyperbolic spaces, the interested reader is referred to [8, 40, 43, 45].

The geometric concept of centroid is partly motivated by the mechanics of rigid bodies. In this context the centroid of a convex body K coincides with the *center of mass* of the solid K in the case of homogeneous mass distribution, as well as with the *center of gravity* of K in a uniform gravitational field. However a physical object may also have inhomogeneous mass distribution, in which case the center of mass and the center of gravity of K are identical to each other but different from the centroid, as they can be any interior point of K depending on the mass distribution.

2.1. Grünbaum's centroid inequality. Consider a convex disk $K \subset \mathbb{R}^2$, and let L be a line through its centroid. This line dissects K into two convex disks K_1 and K_2 . It was observed by Winternitz in 1923 [12] that for any such disk K and line L , we have

$$\frac{4}{5} \leq \frac{\text{area}(K_1)}{\text{area}(K_2)} \leq \frac{5}{4},$$

with equality if K is a triangle.

This result was generalized by Grünbaum [53] for any dimension in 1960 as follows.

Theorem 2.3. *Let $K \subset \mathbb{R}^d$ be a centered convex body. For any $u \in \mathbb{S}^{d-1}$ let K_u denote the intersection of K with the closed half space bounded by u^\perp and containing u . Then, for any $u \in \mathbb{S}^{d-1}$, we have*

$$\frac{\text{vol}_d(K_u)}{\text{vol}_d(K_{-u})} \leq \left(1 + \frac{1}{d}\right)^d - 1.$$

Based on Theorem 2.3, let $\theta(K) = \sup \left\{ \frac{\text{vol}_d(K_u)}{\text{vol}_d(K_{-u})} : u \in \mathbb{S}^{d-1} \right\}$. This quantity can be regarded as a measure of symmetry of K : it satisfies $\theta(K) \geq 1$ for every convex body K , where we have equality for centrally symmetric bodies, and it attains its maximal value $\theta_d = \left(1 + \frac{1}{d}\right)^d - 1$ e.g. for convex cones.

In 2000, Groemer [52] proved a stability version of Grünbaum's result. Namely, he proved that for every $\varepsilon \geq 0$, there is some $\lambda_d > 0$ depending only on d such that if $\theta(K) \geq \theta_d - \varepsilon$ for some convex body K in \mathbb{R}^d , then there is a convex cone C in \mathbb{R}^d such that

$$\text{vol}_d(K \Delta C) \leq \lambda_d \text{vol}_d(K) \varepsilon^{1/(2d^2)},$$

where $X \Delta Y$ denotes the symmetric difference of X, Y .

Using the inequality $\left(1 + \frac{1}{d}\right)^d < e$, Theorem 2.3 yields a simple dimension-free estimate for the volumes of the intersections: for any convex body K in

\mathbb{R}^d , and for any closed half space H containing the centroid of K in its boundary, $\text{vol}_d(H \cap K) \geq \frac{1}{e} \text{vol}_d(K)$. This inequality was generalized by Bertsimas and Vempala [7] for a convex body K in *isotropic position*, defined as a special type of centered convex body: for any unit vector $u \in \mathbb{S}^{d-1}$, K satisfies the equality $1 = \frac{1}{\text{vol}_d(K)} \int_K (\langle u, x \rangle)^2 dx$. Their result states that if K is a convex body in \mathbb{R}^d in isotropic position, and $z \in K$ is at distance t from the origin, then for any closed half space H containing z , $\text{vol}_d(H \cap K) \geq (\frac{1}{e} - t) \text{vol}_d(K)$. Here we note that for any centered convex body K there is a linear transformation A such that $A(K)$ is in isotropic position (see e.g. [7]). Since linear transformations do not change the ratio of volumes and the lengths of parallel segments, it seems possible to restate the result in [7] for an arbitrary convex body K centered at o . This was done in [102] by Shyntar and Yaskin, who strengthened of the result in [7] for \mathbb{R}^2 in the following form: Let K be a plane convex body with o as its centroid. Let $-1 < \alpha < 2$ and $u \in \mathbb{S}^1$. Let H be the closed half plane defined as $H = \{x \in \mathbb{R}^2 : \langle x, u \rangle \geq \alpha h_K(-u)\}$. Then

$$(2.2) \quad C_1(\alpha) \text{area}(K) \leq \text{area}(K \cap H) \leq C_2(\alpha) \text{area}(K),$$

where

$$C_1(\alpha) = \begin{cases} \frac{1}{9}(2 - \alpha)^2, & \text{if } \alpha \in (-1, 0), \\ \frac{4}{9}(1 + \alpha)(1 - 2\alpha), & \text{if } \alpha \in (0, 1/2), \\ 0, & \text{if } \alpha \in (1/2, 2), \end{cases}$$

and

$$C_2(\alpha) = \begin{cases} 1 - \frac{4}{9}(1 + \alpha)^2, & \text{if } \alpha \in (-1, 0), \\ \frac{5 - 3\alpha}{9(1 + \alpha)}, & \text{if } \alpha \in (0, 1), \\ \frac{1}{9}(2 - \alpha)^2, & \text{if } \alpha \in (1, 2). \end{cases}$$

The authors of [102] also characterized the equality cases in (2.2).

Observe that if K is a centered convex body in \mathbb{R}^d , and H is a hyperplane through o , then o may not be the centroid of $K|H$. Thus, it is a meaningful question to examine Grünbaum's problem for projections of convex bodies. This problem was completely solved by Stephen and Zhang [107], who proved that for any centered convex body K in \mathbb{R}^d , any k -dimensional linear subspace E with $1 \leq k \leq d$, and any half space H bounded by u^\perp with $u \in E \cap \mathbb{S}^{d-1}$, we have

$$\text{vol}_k((K|E) \cap H) \geq \left(\frac{k}{d+1}\right)^k \text{vol}_k(K|E).$$

Furthermore, here there is equality for some K and E if and only if $K = \text{conv}\{y_1 + L_1, y_2 + L_2\}$, where

- $L_1 \subset u^\perp$ and $L_1|(E \cap u^\perp)$ are $(k-1)$ -dimensional convex bodies,
- $L_2 \subset E^\perp$ is a $(d-k)$ -dimensional convex body,
- $y_1, y_2 \in \mathbb{R}^d$ satisfy $y_1 \notin H$ and $y_2 \in \text{int}(H)$,
- the centroid of K is o .

More specifically, the authors of [107] prove a more general version of the above result for volumes, where the centroid of K is replaced by an arbitrary point in the interior of K , representing the center of mass of an inhomogeneous object.

In the same vein, one can examine Grünbaum's problem for sections of a convex body. This was proposed by Fradelizi, Meyer and Yaskin [44], who proved that for any centered convex body K , linear subspace V of dimension $d-k$ with $0 \leq k \leq$

$d - 1$, and closed half space H with boundary u^\perp for some $u \in V \cap \mathbb{S}^{d-1}$, we have

$$(2.3) \quad \text{vol}_{d-k}(K \cap V \cap H) \geq c \text{vol}_{d-k}(K \cap V),$$

where $c \geq \frac{c_0}{(k+1)^2} \left(1 + \frac{k+1}{d-k}\right)^{-(d-k-2)}$ for some absolute constant $c_0 > 0$. This result follows from a more general setting investigated in [44] for intersections of a convex body K with a convex cone, which the authors use to prove a conjecture of Meyer and Reisner [84] about the volumes of convex intersection bodies. Grünbaum's problem for sections was settled in [87] by Myroshnychenko, Stephen and Zhang, who proved the following: for any convex body K in \mathbb{R}^d , a k -dimensional linear subspace E of \mathbb{R}^d and a closed half space H bounded by u^\perp with $u \in \mathbb{S}^{d-1} \cap E$, if the centroid of K lies in $E \cap u^\perp$, then

$$\text{vol}_k(K \cap E \cap H) \geq \left(\frac{k}{d+1}\right)^k \text{vol}_k(K \cap E).$$

Furthermore, here we have equality if and only if $K = \text{conv} \left\{ -\left(\frac{d-k+1}{k}\right)z + D_0, z + D_1 \right\}$, where

- $z \in E \cap \text{int}(H)$,
- D_0 is a $(k-1)$ -dimensional convex body in $E \cap u^\perp$,
- D_1 is a centered $(d-k)$ -dimensional convex body in a $(d-k)$ -dimensional subspace F of \mathbb{R}^d such that $E \cup F$ spans \mathbb{R}^d .

The results about both the intersections and the projections were generalized for dual volumes by Stephen and Yaskin [106] as follows. Let K be a centered convex body in \mathbb{R}^d with o , and let E be a k -dimensional linear subspace of \mathbb{R}^d , where $1 \leq k \leq d$. Let H be a closed half space bounded by u^\perp with $u \in \mathbb{S}^{d-1} \cap E$. Then, for any $1 \leq i \leq k$,

$$\tilde{V}_i((K|E) \cap H) \geq \left(\frac{i}{d+1}\right)^i \tilde{V}_i(K|E),$$

and

$$\tilde{V}_i(K \cap E \cap H) \geq \left(\frac{i}{d+1}\right)^i \tilde{V}_i(K \cap E),$$

where $\tilde{V}_i(\cdot)$ denotes i th dual volume with respect to the k -dimensional subspace E . It is worth noting that the case $i = k$ of these statements coincides with the results in [107] and [87], respectively, mentioned in the previous two paragraphs.

An interesting discrete version of the theorem of Winternitz was proposed by Shyntar and Yaskin [102]. To state the problem, for any set $X \subseteq \mathbb{R}^2$ we denote the cardinality of the point set $X \cap \mathbb{Z}^2$ by $\#X$, and we call a convex polygon P an *integer polygon* if all the vertices of P have integer coordinates. In this setting, the problem of Winternitz asks for finding the largest positive number $C > 0$ such that for any integer polygon P , and any closed half plane H containing $c(P)$ in its boundary, we have $\frac{\#(K \cap H)}{\#K} \geq C$. The authors of [102] give an example showing that there is no positive number C that satisfies this property, and prove that, for any integer polygon P and any closed half plane containing $c(3P)$ in its boundary,

$$\frac{\#((3P) \cap H)}{\#(3P)} > \frac{1}{6}.$$

Furthermore, for any $t \geq 6$, any integer polygon P and any closed half space containing $c(tP)$ in its boundary, we have

$$(2.4) \quad \frac{\#((tP) \cap H)}{\#(tP)} \geq \frac{\frac{4}{9}t^2 - 2t - 3}{t^2 + 3t + 2}.$$

Here we note that for $t > 0$, the expression on the right-hand side of (2.4) is increasing, and tends to $\frac{4}{9}$ as $t \rightarrow \infty$.

A generalization of Grünbam's theorem for not necessarily convex sets was considered by Marín Sola and Yepes Nicolás [82]. To state their result, we recall that by Brunn's theorem, for any compact, convex set K in \mathbb{R}^d , and any $u \in \mathbb{S}^{d-1}$, the function $f_X : \mathbb{R} \rightarrow \mathbb{R}$, $f(t) = \text{vol}_{d-1}(K \cap (tu + u^\perp))$ is $\frac{1}{d-1}$ -concave on its support. The authors of [82] proved that if $X \subset \mathbb{R}^d$ is a centered compact set with nonempty interior, and H is a closed half space with u^\perp as its boundary for some $u \in \mathbb{S}^{d-1}$, and the function $f_X : \mathbb{R} \rightarrow \mathbb{R}$, $f(t) = \text{vol}_{d-1}(X \cap (tu + u^\perp))$ is p -concave on its support for some $p > 0$, then

$$\frac{\text{vol}_d(X \cap H)}{\text{vol}_d(X)} \geq \left(\frac{p+1}{2p+1} \right)^{(p+1)/p},$$

and if f_X is log-concave on its support, then

$$\frac{\text{vol}_d(X \cap H)}{\text{vol}_d(X)} \geq \frac{1}{e}.$$

They also characterize the equality case. The same problem, under the more general condition that the function f_X is φ -concave for some strictly increasing, continuous function $\varphi : [0, \infty) \rightarrow \mathbb{R} \cup \{-\infty\}$ satisfying some mild regularity conditions, was investigated by Marín Sola [81].

2.2. The Busemann-Petty centroid inequality. Let K be a centered convex body in \mathbb{R}^d . Consider the function $H_K : \mathbb{R}^d \rightarrow \mathbb{R}$, defined by

$$(2.5) \quad H_K(u) = \frac{1}{\text{vol}_d(K)} \int_K |\langle x, u \rangle| dx.$$

It is well known that this function is positively homogeneous, nonnegative, convex function which is zero only at o [95]. Thus, H_K is the support function of a convex body CK (see Section 1), called the *centroid body* of K . This body is strictly convex, and if K is symmetric to o , i.e. $-K = K$, then the hypersurface $\text{bd}(CK)$, called the *centroid surface* of K , coincides with the set of the centroids of the intersections of K with the closed half spaces containing o in their boundaries. Centroid bodies, attributed by Blaschke to Dupin (see e.g. [100]), were given their name in the seminal work of Petty [95], who proved the inequality that

$$(2.6) \quad \text{vol}_d(CK) \geq \left(\frac{(d+1)\kappa_d}{2\kappa_{d-1}} \right)^d \text{vol}_d(K),$$

with equality if and only if K is an o -symmetric ellipsoid. This inequality is called the *Busemann-Petty centroid inequality*. In 2016, Ivaki [63] proved a stability version of the equality case of this inequality in the plane. Namely, he proved that there are positive constants ε_0 and γ such that for any $0 < \varepsilon < \varepsilon_0$ and any plane convex body K satisfying $\frac{\text{area}(CK)}{\text{area}(K)} \leq \left(\frac{4}{3\pi}\right)^2 (1 + \varepsilon)$, the Banach-Mazur distance of K and an ellipsoid is at most $1 + \gamma\varepsilon^{1/8}$.

It was observed by Milman and Pajor [85] that all centroid bodies are zonoids; that is, they can be obtained as limits, in Hausdorff distance, of a sequence of convex bodies, called zonotopes, obtained as the Minkowski sums of finitely many closed segments.

The concept of centroid body, using a different normalization, was extended to L_p -Brunn-Minkowski theory by Lutwak and Zhang [74] in the following way. For any real $p \geq 1$, let

$$c_{d,p} = \frac{\kappa_{d+p}}{\kappa_2 \kappa_d \kappa_{p-1}}.$$

For any star body $K \subset \mathbb{R}^d$, let $\Gamma_p^* K$ be the o -symmetric convex body defining the norm

$$\|x\|_{\Gamma_p^* K} = \left(\frac{1}{c_{n,p} \text{vol}_d(K)} \int_K |\langle x, y \rangle|^p dy \right)^{1/p}.$$

This body is called the *polar L_p -centroid body* of K , and it can be extended to the case $p = \infty$ by taking the limit $p \rightarrow \infty$. It is worth noting that if K is an o -symmetric convex body and $p = \infty$, then $\Gamma_p^* K$ coincides with the Euclidean polar body K^* of K , defined as $K^* = \{y : \langle x, y \rangle \leq 1 \text{ for every } x \in K\}$. It was proved in [74] that for any star body K , we have

$$(2.7) \quad \text{vol}_d(K) \text{vol}_d(\Gamma_p^* K) \leq \kappa_d^2,$$

with equality if and only if K is an o -symmetric ellipsoid. This inequality, for the case $p = \infty$, coincides with the Blaschke-Santaló inequality. The polar body of $\Gamma_p^* K$ is called the *L_p -centroid body* of K ; if $p = 1$, this body coincides with a normalization of the centroid body CK of K . We note that the body $\Gamma_2 K$ also has a geometric meaning: this is the ellipsoid which has the same inertia about every axis as K . This body is called the *ellipsoid of inertia* or *Legendre ellipsoid* of K [15]. Verifying a conjecture in [74], Lutwak, Yang and Zhang [72] proved that

$$(2.8) \quad \text{vol}_d(K) \leq \text{vol}_d(\Gamma_p K),$$

with equality if and only if K is an o -symmetric ellipsoid. Note that (2.8), combined with the Blaschke-Santaló inequality, yields (2.7), and that (2.8) is a generalization of the Busemann-Petty centroid inequality in (2.6). A different proof of the inequality in (2.8) was given in [15], and a version, establishing a stronger inequality for nonsymmetric bodies, can be found in the paper [56] of Haberl and Schuster. The estimate in [56] was generalized for compact sets in [61]. A different stronger version of the L_p Busemann-Petty centroid inequality can be found in [76], and a variant for the first quermassintegrals of the bodies in [113].

Let $\phi : \mathbb{R} \rightarrow [0, \infty)$ be a convex function satisfying $\phi(0) = 0$ such that ϕ is strictly decreasing on $(-\infty, 0]$ or strictly increasing on $[0, \infty)$. The *Orlitz centroid body* $\Gamma_\phi K$ of a star body K is defined as the convex body with support function

$$(2.9) \quad h_{\Gamma_\phi K}(x) = \inf \left\{ \lambda > 0 : \frac{1}{\text{vol}_d(K)} \int_K \phi \left(\frac{\langle x, y \rangle}{\lambda} \right) \leq 1 \right\}.$$

We note that if $\phi(t) = |t|^p$ with $p \geq 1$, then $\Gamma_\phi K = \Gamma_p K$. This concept was introduced by Lutwak, Yang and Zhang in [73], who proved that for any function ϕ satisfying the above conditions, the ratio $\frac{\text{vol}_d(\Gamma_\phi K)}{\text{vol}_d(K)}$ is minimal on the family of convex bodies K with $o \in \text{int}(K)$ if and only if K is an o -symmetric ellipsoid. This result was extended for arbitrary star bodies by Zhu [119], and the equality case, under a weaker condition on ϕ , was studied in [114].

It is an interesting question to find a converse inequality to (2.6). This question was proposed by Bisztriczky and Böröczky Jr. [11] in 2001. They proved that in the family of o -symmetric plane convex bodies (resp. plane convex bodies containing o), the quantity $\frac{\text{area}(CK)}{\text{area}(K)}$ is maximal if K is a parallelogram (resp. a triangle with o as a vertex). Campi and Gronchi [16] extended these results to L_p -centroid bodies in the plane, and Chen, Zhou and Yang [19] to Orlicz centroid bodies in the plane.

The definition of Orlicz centroid bodies in (2.9) can be extended to that of Orlicz-Lorenz centroid bodies by adding a weight function to the integral in (2.9). The Busemann-Petty centroid inequality in this setting was investigated in [18, 41, 42, 91, 116, 117]. Variants of centroid bodies, and the Busemann-Petty centroid inequality, in non-Euclidean spaces (including complex, spherical and hyperbolic spaces) can be found in [8, 55, 64, 68, 75].

3. STATIC EQUILIBRIUM POINTS

Consider a convex body K in \mathbb{R}^3 . Clearly, if K is balanced on a horizontal plane H , then it is in static equilibrium if and only if the orthogonal projection of its center of mass onto H belongs to K . This leads to the following definition (see e.g. [67]).

Definition 3.1. Let K be a convex body in \mathbb{R}^d and $c \in \mathbb{R}^d$. A point $q \in \text{bd}(K)$ is called an *equilibrium point of K with respect to c* if the hyperplane passing through q and perpendicular to $[c, q]$ supports K .

We note that by convention, in Definition 3.1, every point $c \in \text{bd}(K)$ is an equilibrium point of K with respect to c . In the literature, c is often assumed to be the centroid of K . In this case we call q simply an *equilibrium point* of K . As we mentioned in the beginning of Section 2, any reference point $c \in \text{int}(K)$ is the center of mass of K equipped with some suitable, possibly inhomogeneous density function. Thus, unless we state it otherwise, we assume from now on that the reference point lies in the interior of the body.

For the case $c = o \in \text{int}(K)$, a simple description of equilibrium points can be given by means of the radial or the support function of K . More specifically, if ρ_K or h_K is C^1 -class, then q is an equilibrium point of K with respect to o if and only if $\frac{q}{|q|}$ is a critical point of ρ_K or h_K , respectively. Indeed, since the restriction of ρ_K to a plane coincides with the radial function of the intersection of K with the plane, and a similar observation holds for h_K and the projection of K onto the plane, it is sufficient to show our remark for plane convex bodies. For this case a simple geometric argument shows the statement (see also [2, 31]). We note that the properties that h_K or ρ_K are C^1 -class are not equivalent [100], but if both are C^1 -class, their critical points coincide.

A remarkable result of Zamfirescu [115] states that, in Baire category sense with respect to the topology induced by Hausdorff distance, a typical convex body has infinitely many equilibrium points with respect to a typical point. On the other hand, Montejano [86] showed that the Euclidean balls centered at o are the only convex bodies whose every boundary point is an equilibrium point with respect to o .

Next, we define non-degenerate equilibria.

Definition 3.2. Let K be a smooth convex body in \mathbb{R}^d and $c \in \mathbb{R}^d$. Assume that K has finitely many equilibrium points with respect to c , and its boundary is C^2 -class

in the neighborhood of each equilibrium point $q \in \text{bd}(K)$. Assume that the Hessian of the function $\delta_K : \text{bd}(K) \rightarrow \mathbb{R}$, $\delta_K(x) = |x - c|$ is not zero at any equilibrium point q . Then we say that K is *nondegenerate* with respect to c . Furthermore, the *index* of any equilibrium point is the number of negative eigenvalues of the Hessian of δ_K at q . Equilibrium points of index 0 and $d - 1$ are called *stable* and *unstable* points, respectively, and equilibrium points of all other indices are called *saddle* points.

We note that for the special case $c = o \in \text{int}(K)$, there is an equivalent formulation of the concepts in Definition 3.2 in terms of the radial function ρ_K . Indeed, the conditions for the nondegeneracy of K are equivalent to the properties that ρ_K is C^1 -class, it has finitely many critical points and ρ_K is C^2 -class in a neighborhood of every one of them, and at every critical point the Hessian of ρ_K is not zero. Furthermore, the index of the equilibrium point $q \in \text{bd}(K)$ is equal to the number of the negative eigenvalues of the Hessian of ρ_K at the corresponding critical point $\frac{q}{|q|}$.

Whereas there is no definition, in general, for a not necessarily smooth convex body being nondegenerate, the condition in Definition 3.2 can be replaced by a geometric one if K is a polytope. We note that by definition, every face (of any dimension) of a convex polytope contains at most one point of equilibrium in its relative interior.

Definition 3.3. Let P be a convex polytope in \mathbb{R}^d , and let $c \in \mathbb{R}^d$. For any equilibrium point q of P with respect to c , let H_q denote the supporting hyperplane of P at q perpendicular to $[c, q]$. Set $F_q = P \cap H_q$, and let $f_q = \dim F_q$. If, for every equilibrium point q , q lies in the relative interior of F_q , we say that P is *nondegenerate*. In this case the *index* of the equilibrium point q is the quantity $(d - 1) - f_q$.

Consider a C^1 -class function $f : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$ with finitely many critical points such that f is C^2 -class in a neighborhood of every critical point, and its Hessian is nonzero at every critical point. A standard convolution technique (see e.g. [48]) shows that there is a C^∞ -class function $g : \mathbb{S}^{d-1} \rightarrow \mathbb{R}$ arbitrarily close to f such that for every $0 \leq k \leq d - 1$, the number of critical points of f with index k coincide with the same quantity for g . The function g is a Morse function, and hence, applying this observation for the radial function ρ_K of a nondegenerate smooth convex body K with $c = o \in \text{int}(K)$, and combining it with the Poincaré-Hopf formula for the sphere \mathbb{S}^{d-1} , we have the following.

Theorem 3.4. Let K be a nondegenerate smooth convex body in \mathbb{R}^3 with respect to $c \in \text{int}(K)$, and let S, H, U denote the numbers of stable, saddle and unstable points of K with respect to c . Then

$$(3.1) \quad S - H + U = 2.$$

Similarly, if K is a nondegenerate plane convex body with respect to some $c \in \text{int}(K)$, with S stable and U unstable points, then $S = U$.

Finally, let P be a nondegenerate convex polytope in \mathbb{R}^d with respect to $c \in \text{int}(P)$. Then, if $\varepsilon > 0$ is sufficiently small, the convex body $K = P + \varepsilon \mathbf{B}^d$ is a nondegenerate convex body with respect to $c \in \text{int}(K)$, and for every $0 \leq k \leq d - 1$, the number of equilibrium points of P of index k coincide with the same quantity

for K . Thus, Theorem 3.4 is satisfied also for nondegenerate convex polyhedra and polygons. We also note that the notion of radial function and Theorem 3.4 can be extended to the case $c \notin \text{int}(K)$ in a natural way [47].

3.1. Primary classes of convex bodies based on their equilibria. To distinguish convex bodies based on their static equilibrium properties it is a straightforward way to use the numbers of their equilibrium points of different indices. We observe that by Theorem 3.4, for any nondegenerate plane convex body it is sufficient to know the number of either the stable or the unstable points, and in 3-dimensions two of the numbers of stable, saddle and unstable points. Thus, we define the set $\{S\}$ as the family of all nondegenerate convex bodies in \mathbb{R}^2 that have S stable points, and the set $\{S, U\}$ as the family of all nondegenerate convex bodies in \mathbb{R}^3 that have S stable and U unstable points. For later use, we decompose each class $\{S\}$ into two subclasses $\{S\}_s, \{S\}_p$, consisting of the nondegenerate smooth plane convex bodies and the convex polygons in this class, respectively, and we introduce the classes $\{S, U\}_s, \{S, U\}_p$ in the same way (Figure 1). Elements of class $\{1, U\}$ for some U are called *monostable* or *unistable*, elements of class $\{S, 1\}$ are called *mono-unstable* or *uni-unstable*, and a body that is both monostable and mono-unstable is called *mono-monostatic*.

We call this classification the primary classification system, and a class $\{S\}$ or $\{S, U\}$ the *primary class* of its elements [31, 111]. It is a natural question to ask what are the integers $S \geq 0$ (resp. the integers $S, U \geq 0$) such that the class $\{S\}$ (resp. the class $\{S, U\}$) is not empty. Our first step in answering this problem is the observation that if K is a nondegenerate convex body, then every boundary point of K closest to $c(K)$ is a stable, and every boundary point of K farthest from $c(K)$ is an unstable point of K . This implies that in every nonempty primary class, S, U are positive. The first nontrivial result in this direction is due to Domokos, Papadopoulos and Ruina [33], who proved that the class $\{1\}$ is empty, or in other words, every nondegenerate plane convex body has at least two stable and two unstable points. On the other hand, it is easy to see that the class $\{S\}$ is not empty for any $S \geq 2$. Indeed, the class $\{2\}$ contains all ellipses different from a circle, and for any $S \geq 3$, the class $\{S\}$ contains, e.g., the regular S -gons. This completes the planar case. As a final remark, we mention that the result that $\{1\}$ is empty is equivalent to the famous 4-vertex theorem in differential geometry [111].

The characterization of nonempty primary classes $\{S, U\}$ was carried out by Várkonyi and Domokos [111]. In particular, they presented a construction of a convex body with only one stable and one unstable point. For preciseness, we mention that the body constructed in [111] was not C^2 -class at its two equilibrium points, and thus, it is not nondegenerate as defined in Definition 3.2; nevertheless the radial function of the body has a global minimum and maximum at its equilibrium points. We also add that, as it was remarked in [31], a standard smoothing technique yields a C^∞ -class, nondegenerate approximation of this body, with the same numbers of stable and unstable points. For an explicit construction of nondegenerate convex bodies in class $\{1, 1\}$, with C^2 -class boundaries, the reader is also referred to [32]. A representative example of class $\{1, 1\}$ constructed by patching together elementary surfaces became known as ‘Gömböc’ (Figure 2). The existence of a convex polyhedron in class $\{1, 1\}$ was shown in [67]. Another remarkable simple shape in class $\{1, 1\}$ given by an explicit formula was recently published by Sloan [103], and possible symmetries of mono-monostatic bodies have been determined in [32].

$s \setminus u$	1	2	3	4	5	6	7	8	...
1									
2									
3									
4									
5									
6									
7									
8									
htb ...									

FIGURE 1. Equilibrium classes with some representative examples. Classes corresponding to polyhedral pairs (see Section 3.6, below) are highlighted by grey background. Unistable, uni-unstable, and mono-monostatic objects correspond to the first row, the first column, and the top-left corner of the table. The examples have been provided by Tímea Szabó and Gábor Domokos.

Once we have verified that the class $\{1, 1\}$ is not empty, to show that no class $\{S, U\}$, with $S, U \geq 1$ is empty it is sufficient to find a way to increase the number of equilibrium points of a body. This was also carried out in [111] by Domokos and Várkonyi, who showed that, in a neighborhood of a fixed stable point of a convex body K , it is possible to create a stable point by truncating K by a suitable plane, or an unstable point by truncating K by a suitable cone. These steps are called ‘Columbus steps’ in memory of the famous discoverer Christopher Columbus, who balanced an egg on its top by creating a stable point near it. Since the class $\{1, 1\}$ is not empty, subsequent applications of these steps show that no class $\{S, U\}$ with $S, U \geq 1$ is empty. We note that a more rigorous investigation of Columbus steps, involving also a construction of a smooth approximation of the resulting body, can be found in [31]. The fact that all these classes contain also nondegenerate convex polyhedra follows from [67].

3.2. Secondary and tertiary classes. In this subsection, we investigate nondegenerate smooth convex bodies in \mathbb{R}^3 . Consider the radial function $\rho_K : \mathbb{S}^2 \rightarrow \mathbb{R}$ of such a body. By the smoothing technique described in the beginning of Section 3, we may assume that ρ_K is a C^∞ -class function. Then ρ_K is a Morse function on




FIGURE 2. Left: illustration of the ömböc: a piecewise smooth body in class $\{1, 1\}$. The boundary is coloured according to level curves of the radial function, which has two critical points at the top and at the bottom of the surface. Right: the monostatic polyhedron of Guy [21], see Section 3.6, below. The illustration has been created with the aid of a 3D model available at <https://www.thingiverse.com/thing:90866>.

\mathbb{S}^2 , and the gradient of ρ_K is nonzero at q if and only if q is not critical, i.e. it is regular. By the Picard-Lindelöf theorem, for any regular point q of \mathbb{S}^2 there is a unique curve $\gamma: \mathbb{R} \rightarrow \mathbb{S}^2$ with the property that $\gamma(0) = q$, and $\gamma'(t)$ is equal to the gradient of ρ_K at $\gamma(t)$ for all $t \in \mathbb{R}$. Such a curve γ , called an *integral curve* of the gradient flow induced by ρ_K starts and ends at two different critical points of ρ_K , called the *origin* and the *destination* of γ . For any critical point x , the *descending* (resp. *ascending*) *manifold* of x , denoted by $D(x)$ (resp. $A(x)$) is the union of $\{x\}$ and all integral curves with x as their destination (resp. origin). The connected components of the nonempty sets, obtained as the intersection of an ascending and a descending manifold, are called *cells* of the *Morse-Smale complex generated by ρ_K* . The 2-, 1-, and 0-dimensional cells of this complex are called *faces*, *edges*, and *vertices*, respectively.

If $x \neq y$ are critical points of ρ_K , then $A(x) \cap D(x) = \{x\}$, and $A(x) \cap D(y)$ is the union of all integral curves with origin x and destination y . Thus, the vertices of the Morse-Smale complex of ρ_K are the critical points of ρ_K , and if x is a stable and y is an unstable point, then $A(x) \cap D(y)$ is an open set in \mathbb{S}^2 , implying that its connected components are faces of the complex (provided $A(x) \cap D(y) \neq \emptyset$). On the other hand, the same does not hold if x or y is a saddle point since every saddle point is the origin, and also the destination, of exactly two, *isolated* integral curves, corresponding to edges of the complex [39, 70].

The function ρ_K is called *Morse-Smale* if all ascending and descending manifolds of ρ_K intersect only transversally; or equivalently, if any pair of intersecting ascending and descending 1-dimensional manifolds cross. In this case the crossing point of these 1-manifolds is necessarily a saddle, since crossing at a regular point would contradict the property that any regular point belongs to exactly one integral curve. In the following we only deal convex bodies whose radial function ρ_K is Morse-Smale. In this case every face of the Morse-Smale complex is bounded by

a union of lower dimensional cells, either two disjoint vertices or a cycle of vertices and edges.

The existence of a face whose boundary is a pair of vertices implies that the body K has no saddle point, that is, its primary class is $\{1, 1\}$. In this case the Morse-Smale complex is made up of a unique face corresponding to the single connected component of $A(x) \cap D(y)$, where x is the stable and y is the unstable point of K , and its boundary consisting of the two critical points x and y . In addition, the function ρ_K has no saddle points, no isolated integral curves, therefore its Morse-Smale complex has no edges. In the opposite case the boundary of every face consists of a cycle of connecting edges and vertices, and it is known (see e.g. [39]) that it consists of four edges, and the four critical points in the boundary are a stable, a saddle, an unstable and a saddle point, in this cyclic order. The boundary is possibly glued to itself along vertices and edges.

The topological graph G on \mathbb{S}^2 , whose vertices are the critical points of ρ_K , and whose edges are the edges of the Morse-Smale complex, is called the *Morse-Smale graph* generated by ρ_K [31]. This graph is usually regarded as a 3-colored quadrangulation of \mathbb{S}^2 , where the ‘colors’ of the vertices are the three types of a critical point. We regard the topological graph G defined by ρ_K as the drawing of a 3-colored abstract graph \bar{G} on \mathbb{S}^2 . The graphs \bar{G} and G are called the *secondary* and the *tertiary class* of K , respectively. The secondary classification of convex bodies is a refinement of the primary one, as convex bodies whose Morse-Smale complexes induce the same abstract graph have the same numbers of stable, saddle and unstable points, whereas the statement is not necessarily true in the opposite direction. Similarly, tertiary classification is a refinement of the secondary one.

Let us denote by \mathcal{Q}_3^* the family of finite 3-colored quadrangulations of \mathbb{S}^2 , up to homeomorphism, where the color classes are denoted by $\mathcal{S}, \mathcal{H}, \mathcal{U}$, each element of \mathcal{H} has degree 4, and the cardinalities of the classes satisfy the equation $\text{card } \mathcal{S} - \text{card } \mathcal{H} + \text{card } \mathcal{U} = 2$. We remark that the Morse-Smale complex of every Morse-Smale function on \mathbb{S}^2 belongs to \mathcal{Q}_3^* . Furthermore, let us denote the Morse-Smale graph of a nondegenerate convex body K , if it exists, by $Q_3^*(K)$. We note that by convention, the Morse-Smale graphs of the elements of the class $\{1, 1\}$, consisting of two vertices with different colors and no edges, are regarded as elements of \mathcal{Q}_3^* . These graphs are called *path graphs* on two vertices, and are denoted by P_2 (see Figure 3).




FIGURE 3. The topological graphs of convex bodies with at most two stable and two unstable points.

As in the case of primary classes, it is a natural question to ask which elements of \mathcal{Q}_3^* appear as Morse-Smale graphs of nondegenerate convex bodies. The answer, namely that *every* such quadrangulation is realizable by a convex body, was given in [31] by Domokos, Lángi and Szabó. The idea of the proof is a refinement of Columbus steps, described in the previous subsection. More specifically, a generalization of a method of Batagelj [4], and also Negami and Nakamoto [90] shows that every 3-colored quadrangulation in \mathcal{Q}_3^* can be reduced to P_2 by finitely many applications of a graph operation, called *face contraction*, or equivalently, every such quadrangulation can be obtained from P_2 by finitely many subsequent *vertex splittings*, the dual operations of face contractions. Based on this result, the authors of [31] show that for any convex body K and any 3-colored quadrangulation Q in \mathcal{Q}_3^* , obtained from $Q_3^*(K)$ by an arbitrary vertex splitting, there is a convex body K' , obtained as a smoothed truncation of K by a plane or a cone, satisfying that $Q_3^*(K') = Q$.

We note that the above representation of the Morse-Smale complex of K is also called the *primal* Morse-Smale graph of K . Saddle points can be removed from the primal Morse-Smale graph without losing information: first we connect maxima and minima in the quadrangles, then cancel saddle points and edges incident to them (see Figure 4). This representation is called the *quasi-dual* Morse-Smale graph (see. [36]). Every face of a quasi-dual representation is a quadrangle, and the vertices on its boundary form an alternating sequence of two stable and two unstable points. Connecting the two stable points in each face, and removing all unstable points and their edges we obtain an even more compact representation of the Morse-Smale graph. This representation is a planar graph that might contain loops and multiple edges. The vertices of this graph are the stable points of the Morse-Smale complex, and its edges correspond to the saddle points of the complex (See Figure 4).

It is a natural question to ask the numbers of secondary and tertiary classes in each primary class. However, the answer to this question is known only for some small values of S and U .

The definitions of the secondary and tertiary classes of a smooth nondegenerate convex body K rely on the topological properties of the Morse-Smale complex of the gradient flow of the radial function ρ_K of the body. This task seems much more complicated if K is a nondegenerate convex polyhedron. Partial results in this direction can be found in the recent paper [70] of Ludmány, Lángi and Domokos.

3.3. Transition between different classes. The next topic in our survey is to examine how difficult it is to move a convex body from its class to another one. We start the investigation with primary classes. Following the paper [26] of Domokos and Lángi, the question we ask is this: For a given nondegenerate convex body K , what is the minimum fraction of the volume of K that is necessary to remove from K to obtain a nondegenerate convex body with a different number of equilibrium points? As we have already seen in Subsection 3.1, this number is equal to zero for any smooth convex body, because it is possible to use an arbitrarily small truncation of K to *increase* the number of equilibrium points of the body. This shows that we should aim at *decreasing* the number of equilibrium points of K . Thus, we define the (*downward*) *robustness* of the convex body K in \mathbb{R}^d as the quantity

$$\rho(K) = \frac{1}{\text{vol}_d(K)} \inf \{ \text{vol}_d(K \setminus K') : K' \in \mathcal{F}_<(K) \},$$




FIGURE 4. Different representations of a Morse-Smale complex. a) Primal Morse-Smale graph. b) Primal Morse-Smale graph with the stable and unstable points connected. c) Quasi-dual Morse-Smale graph. d) Quasi-dual graph with the stable points connected. e) The induced uncolored planar graph; vertices, edges and faces correspond to the stable, saddle and unstable points of the primal representation.

where the set $\mathcal{F}_{<}(K)$ denotes the family of convex bodies in $K' \subset K$ with fewer equilibrium points than K . Furthermore, we define the robustness of an equilibrium class as the supremum of the robustnesses of the elements of the class. To find the robustness of a convex body, or an equilibrium class, is challenging partly because truncating a body changes also the position of its centroid. This is the motivation behind defining the *internal* and *external robustness* of a convex body. As in [26], we do it for plane convex bodies with piecewise smooth boundaries, and remark that the nondegeneracy of such a body, and its stable and unstable points, can be defined for them by combining the conditions in Definitions 3.2 and 3.3. We denote the family containing such plane convex bodies by \mathcal{K}_2 .

Definition 3.5. Let $K \in \mathcal{K}_2$ and $c \in \text{int}(K)$. Assume that K has S stable points with respect to c . Let $R(K, c) \subseteq \text{int}(K)$ denote the set of the points such that K has S stable points with respect to any point of $R(K, c)$. The *internal robustness* of K with respect to c is

$$\rho^{in}(K, c) = \frac{\inf \{|q - c| : q \notin R(K, c)\}}{\text{perim}(K)},$$

where $\text{perim}(K)$ is the perimeter of K .

The existence of the quantity $\rho^{in}(K, c)$ follows from the observations that, first, for any $S > 0$, the set with respect to which K has S stable points is open, and second, Blaschke's rolling ball theorem implies that any $K \in \mathcal{K}_2$ contains two points with respect to which K has different numbers of stable points. The question of how

the number of equilibria depends on the location of the center of mass is discussed in detail in Section 3.5, and the concept of internal robustness is illustrated by Figure 5.

Definition 3.6. Let $K \in \mathcal{K}_2$ and $c \in \text{int}(K)$. Assume that K has S stable points with respect to c . We define the (*downward*) *external robustness* of K with respect to c as the quantity

$$\rho^{ex}(K, c) = \frac{\inf\{\text{area}(K \setminus K') : K' \in \mathcal{F}_<(K, c)\}}{\text{area}(K)},$$

where $\mathcal{F}_<(K, c)$ denotes the family of the nondegenerate plane convex bodies $K' \in \mathcal{K}_2$, satisfying $c \in \text{int}(K') \subset K$, which has strictly less than S stable points with respect to c . If the set in the numerator is empty, we let $\rho^{ex}(K, c) = 1$.

The internal and the external robustness of the equilibrium class $\{S\}$ is defined as the quantity $\rho_S^{in} = \sup\{\rho^{in}(K, c(K)) : K \in \{S\}\}$ and $\rho_S^{ex} = \sup\{\rho^{ex}(K, c(K)) : K \in \{S\}\}$, respectively. Regarding these quantities, the authors of [26] prove that for any $S \geq 3$, both $\rho^{in}(K, p)$ and $\rho^{ex}(K, p)$ are maximal over all $K \in \mathcal{K}_2$ and $c \in \text{int}(K)$ if K is a regular S -gon, and $c = c(K)$.

In [26], the authors extend the concepts of external and internal robustness to nondegenerate smooth convex bodies and nondegenerate convex polyhedra in \mathbb{R}^3 in the straightforward way, where, in case of internal robustness, they replace the normalizing factor $\text{perim}(K)$ by $\sqrt{\text{surf}(K)}$, where $\text{surf}(\cdot)$ denotes surface area. They show that any platonic solid in class $\{S, U\}$ has maximal internal robustness among the elements of (S, U) having the same numbers of faces, edges and vertices, respectively. They also show that the (full) robustness of the classes $\{1, 1\}$, $\{1, 2\}$ and $\{2, 1\}$ is equal to 1.

Transitions between equilibrium classes can be described also from the point of view of dynamical systems. The equivalence classes of nondegenerate, smooth convex bodies are generic in the sense that a sufficiently small C^2 -class deformation of the body does not change any of its classes. In particular, if $\{K(t) : t \in [0, 1]\}$ is a C^2 -class differentiable 1-parameter family of convex bodies, where $K(0)$ and $K(1)$ belong to different equilibrium classes, then there is some value $0 < t_0 < 1$, $K(t_0)$, where $K(t_0)$ is degenerate. If, in addition, for any $0 \leq t < t_0$ (resp. $t_0 < t \leq 1$) $K(t)$ belongs to the same equilibrium class as $K(0)$ (resp. $K(1)$), then there is a bifurcation of the corresponding gradient vector fields induced by $\rho_{K(t)}$. In a ‘generic’ case, this bifurcation has codimension 1 [54]. The codimension 1 bifurcations of gradient vector fields on \mathbb{S}^2 have two types, called *saddle-node* bifurcations and *saddle-saddle connections*.

In terms of topological graphs, saddle-node bifurcations correspond to vertex splittings/face contractions, while saddle-saddle connections correspond to another type graph transformations on 3-colored quadrangulations on \mathbb{S}^2 , called *diagonal slides*. These graph transformations establish a combinatorial connection between primary/secondary or tertiary equilibrium classes, namely we can say that the transition between two classes \mathcal{C}_0 and \mathcal{C}_1 induced by such a graph transformation T is *combinatorially realizable* if there are 3-colored quadrangulations Q_0 and Q_1 , belonging to the classes \mathcal{C}_0 and \mathcal{C}_1 , respectively, such that Q_1 is obtained from Q_0 by Q_1 . In other words, we regard the 3-colored quadrangulations of \mathbb{S}^2 , up to homeomorphism, as vertices of a ‘metagraph’ \mathcal{G} , and connect two vertices of \mathcal{G} by an edge if one of them is induced from the other one by a vertex splitting or a diagonal

slide. We call an edge *primary*, *secondary* or *tertiary* if the two quadrangulations are induced by convex bodies in different primary, secondary or tertiary classes, respectively.

It is a meaningful question to ask which edges of \mathcal{G} are realizable by a generic 1-parameter family of convex bodies. This problem was partially solved in [23], who proved that every primary and secondary edge of \mathcal{G} is realizable. The case of tertiary edges is still open.

3.4. The location and clustering of equilibria. Flattish objects like a coin tend to have two stable equilibrium points, and elongated ones like a needle tend to have two unstable ones. This simple observation can be turned into a more precise statement in various ways. For example, it was shown in [28] that for any convex body K in \mathbb{R}^d , with C^∞ -class boundary and strictly positive Gaussian curvature and for any orthogonal affinity $f_\lambda : \mathbb{R}^d \rightarrow \mathbb{R}^d$ with affinity ratio $\lambda > 0$, if λ is sufficiently large, then $f_\lambda(K)$ has exactly two unstable points, and if λ is sufficiently small, then $f_\lambda(K)$ has exactly two stable points.

Similarly, a convex body K with a significantly flattened or elongated *bounding box* may not have equilibrium points but in some small regions of $\text{bd } K$ including two small patches and a narrow ring as demonstrated by [34]. However, the number of equilibria within these areas can be arbitrary. Indeed, many objects have localized clusters of equilibria. For example, randomly generated fine polygonal/polyhedral discretizations of smooth surfaces often possess small groups of nearby equilibria, in place of isolated equilibria of the original surface. Based on this observation, global equilibria consisting of clusters of local equilibria have been defined in [34]. This concept was made precise in [30], where the authors gave simple formulas for the numbers of stable, saddle and unstable points on a fine polyhedral approximation of a C^2 -class convex surface S in \mathbb{R}^3 induced by an equidistant partition of the parameter range, based on the principal curvatures of the equilibrium points and their distances from the reference points. They called these quantities the *imaginary equilibrium indices* of the equilibrium points of S , and carried out a similar investigation of C^2 -class curves in \mathbb{R}^2 . In [29], the authors removed some technical assumptions from the results in [30], and proved a similar result for the number of equilibrium points of a polygonal curve generated by randomly chosen points on the curve.

3.5. The number of equilibria with respect to a general reference point.

Let K be a convex body in \mathbb{R}^d . Let the function $n_K(c) : \mathbb{R}^d \rightarrow \mathbb{N}$ be defined as the number of equilibrium points with respect to the point c . We note that if K has C^2 -class boundary and strictly positive Gaussian curvature, then this quantity is equal to the number of points of $\text{bd}(K)$ through which the line normal to $\text{bd}(K)$ contains c . Thus, in the literature, questions related to this function are often referred to as *problems of concurrent normals*. In this subsection we discuss results related to $n_K(\cdot)$.

First, we consider plane convex bodies. The function $n_K(\cdot)$ was first investigated by Apollonius of Perga [92] in the special case of an ellipse, and found that its value in a generic point is either 2 or 4. It is known that for any K in \mathbb{R}^2 with C^2 -class boundary and strictly positive curvature, $n_K(\cdot)$ is a piecewise constant function. Even more, it was proved by Allemann, Hungerbühler and Wasem [2] that $n_K(c) = 2 + 2|m|$ where m is the winding number of the evolute of $\text{bd}(K)$ with




FIGURE 5. Two examples of smooth, planar curves with their evolutes, and the corresponding number of equilibria. In the left panel, the dashed circle illustrates the internal robustness of a point P . The right one is an example with $n_{min,K} = 4$.

respect to c . This result makes it possible to have a nice illustration of the function $n_K(\cdot)$: we decompose the plane into connected regions by the evolute of $\text{bd}(K)$ (which in the following we call the evolute of K) and observe that within each region $n_K(\cdot)$ is a constant (Figure 5). As a consequence, $n_K(c)$ is discontinuous at the points of the evolute, and K is degenerate only with respect to the points of its evolute.

It is trivial that the number of equilibrium points of any convex body K with respect to any point c is at least two; this follows from the observation that the points of $\text{bd}(K)$ closest to and farthest from c are equilibrium points. On the other hand, for any convex body with C^2 -class boundary and strictly positive Gaussian curvature, $n_K(c) = 2$ for all points c sufficiently far from K . However, due to motivation by mechanics, it is often assumed that the reference point c lies in the interior of the convex body K . Thus, we define the quantity

$$n_{min,K} = \min\{n_K(c) : c \in \text{int}(K)\}.$$

We observe that there are plane convex bodies K satisfying $n_{min,K} \geq 4$. A simple example for this is a smoothed regular triangle (Figure 5). Even higher values of $n_{min,K}$ are possible if we drop the differentiability properties for $\text{bd}(K)$: for example, rectangles have eight equilibrium points with respect to all interior points or more generally, bricks in \mathbb{R}^d have $3^d - 1$ equilibrium points with respect to all their interior points. The highest possible value of $n_{min,K}$ among polytopes and among smooth bodies appear to be unsolved problems. For example, it is an intriguing question if there exist a plane convex body K with C^2 -class boundary and strictly positive curvature that satisfies $n_{min,K} > 4$. A related result by Grebennikov [51] is that almost every normal through a boundary point to such a convex body in any dimensions contains a point c with $n_K(c) \geq 6$.

In contrast, the quantity

$$n_{max,K} = \max\{n_K(c) : c \in \text{int}(K)\}$$

has been studied by several authors. More specifically, it has been proven under mild smoothness assumptions that $n_{max,K} \geq 2d$ for any convex body with C^1 -class

boundary in \mathbb{R}^d with $2 \leq d \leq 4$ [59, 94], and the same statement in arbitrary dimensions has been known for long as *concurrent normal conjecture*. In addition, the inequality $n_{max,K} \geq 6$ has been proven for any convex body K in \mathbb{R}^d with an arbitrary value of d [59]. In a recent paper, another lower bound for $n_{max,K}$, over the family of d -dimensional convex polytopes, was given by Nasonov, Panina and Siersma [89], namely they proved the inequality $n_{max,K} \geq 2d + 2$, and conjectured that the inequality $n_{max,K} \geq 10$ holds for any 3-dimensional convex polytope.

Several works have addressed the average number of equilibria

$$(3.2) \quad \bar{n}_K = \frac{1}{\text{vol}_d(K)} \int_{\text{int}(K)} n_K(x) dx$$

in a convex body K , for certain families of convex bodies. A list of bounds, established in the papers [27, 37, 57, 58, 62, 99], have been collected in Table 1.

Dimension	2	3	d
\mathcal{P}	$\overset{?}{> 4}$ [37], ≤ 12 [57]	$\overset{*}{> 2}$ [27]	$\overset{*}{> 2}$ [27]
\mathcal{F}	$\overset{*}{\geq 2}$, ≤ 12 [57, 62]; $\overset{?}{\leq 8}$ [58]	$\overset{*}{\geq 2}$ ≤ 62 ; $\overset{?}{\leq 26}$ [58]	$\overset{*}{\geq 2}$ $\leq \frac{3}{2} \binom{2d}{d} - 1$
\mathcal{F}_{cs}	$\overset{*}{\leq 8}$ [57, 62]	$\overset{*}{\leq 26}$ [57, 62]	$\overset{*}{\leq 3^d - 1}$ [57, 62]
\mathcal{F}_{cc}	≤ 6 [57, 62]	≤ 20 [27]	$\leq \binom{2d}{d}$ [27]
\mathcal{F}_{cc}	$\overset{*}{\leq} \frac{2\pi}{\pi - \sqrt{3}}$ [17, 99]	≤ 20 [27]	$\leq \binom{2d}{d}$ [27]

TABLE 1. A summary of known bounds of the average number of equilibria. Star means bounds verified to be sharp, and question marks denote published conjectures. The notation in the first row is as follows: \mathcal{P} and \mathcal{F} are the families of convex polytopes, convex bodies with C^2 -class boundary and strictly positive curvature, respectively, in the appropriate dimension, and for $x \in \{cs, cc, cw\}$, \mathcal{F}_x denotes the subfamily of \mathcal{F} consisting of centrally symmetric convex bodies, of the convex bodies containing all their centers of curvature, and of constant width bodies. The bounds established by [62] do not require C^2 -class boundary.

3.6. Equilibrium points of convex polyhedra. In modern times, the study of equilibrium points of convex polyhedra was probably started with a problem of Conway and Guy [21] in 1966 who conjectured that there is no monostable tetrahedron but there is a monostable convex polyhedron in \mathbb{R}^3 . These two questions were answered by Goldberg and Guy in [50] in 1969 (Figure 2). For a more detailed proof of the first problem, see also [77]. In addition, in [50] Guy presented some problems regarding monostable polyhedra, stating that three of them are due to Conway (for similar statements in the literature, see e.g. [22, 38, 80]). These three questions appear also in the problem collection of Croft, Falconer and Guy [22] as Problem B12. These problems were:

- (1) Can a monostable polyhedron in the Euclidean 3-space \mathbb{R}^3 have an n -fold axis of symmetry for $n > 2$?

- (2) What is the smallest possible ratio of diameter to girth for a monostable polyhedron?
- (3) What is the set of convex bodies uniformly approximable by monostable polyhedra, and does this contain the sphere?

Related to these problems we recall that the *girth* of a convex body in \mathbb{R}^3 is the minimum perimeter of an orthogonal projection of the body onto a plane [38]. It is worth noting that, as stated in [50], Conway showed that no body of revolution can be monostable, and also that the polyhedron constructed in [50] has a 2-fold rotational symmetry.

The first two questions have recently been answered by Lángi [67], but the third one is still open. The main tool of the proofs of the result in [67] is a theorem stating that a nondegenerate convex body, under some mild condition on its curvatures, can be approximated arbitrarily well by a nondegenerate convex polyhedron with the same numbers of equilibria and with the same symmetry group. This result can be regarded as the counterpoint of the one in [30] about approximations of a convex surface by equidistant partitions of its parameter range, mentioned in Subsection 3.4.

A problem proposed in [50] asks about the minimum dimension, if it exists, in which a d -simplex can be monostable. This problem has been investigated by Dawson et al. [77–80], who proved that there is no monostable d -simplex if $d \leq 8$ and there is a monostable 11-simplex. With regard to Problem XVI in [101], asking about the minimum number of faces of a monostable polyhedron in \mathbb{R}^3 , the original construction of Guy [50] with 19 faces (attributed also to Conway) was modified by Bezdek to obtain a monostable polyhedron with 18 faces [9], while a computer-aided search by Reshetov [97] yields a monostable polyhedron with 14 faces. The non-existence of mono-unstable homogeneous tetrahedra has been proved by [25] where a construction of a mono-unstable polyhedron was also given.

We have seen that all equilibrium classes $\{S, U\}$ contain nondegenerate convex polyhedra [67] and all classes $\{S\}$ contain convex polygons [67], as one can construct a fine polyhedral approximation of a smooth convex body from the same class. On the other hand, these representatives have much more faces, edges, and vertices than the number of equilibria. Recall that every vertex, edge or face of a nondegenerate convex polyhedron contains at most one equilibrium point in its relative interior. Thus, an interesting related problem is to find the ‘simplest’ convex polyhedron in each class in terms of the number of vertices, edges or faces. This leads to the following concept.

For an arbitrary nondegenerate, convex polyhedron P the total number of faces, edges, and vertices without equilibria is called the *complexity* of P , denoted by $C(P)$. Furthermore, for any $S, U \geq 1$, we denote the smallest complexity among polyhedra in the class $\{S, U\}_p$ by $C(S, U)$, and call it the *complexity of equilibrium class* $\{S, U\}_p$. In addition, recall a classical theorem of Steinitz [104, 105] stating that a triple of integers (v, e, f) are the numbers of the vertices, edges and faces of a convex polyhedron, respectively, if, and only if $f \geq 4$, $\frac{f}{2} + 2 \leq v \leq 2f - 4$, and $e = v + f - 2$. Based on this result, let us call a pair of integers (f, v) a *polyhedral pair* if they satisfy these inequalities, and let \mathcal{PP} denote the set of all polyhedral pairs.

It has been shown in [25] that for all equilibrium classes with $S, U > 1$,

$$C(S, U) = 2 \min\{f + v - S - U : (f, v) \in \mathcal{PP}\}.$$

In particular, this result implies that $C(S, U) = 0$ if and only if (S, U) is a polyhedral pair, or in other words, any primary equilibrium class $\{S, U\}_p$, where (S, U) is a polyhedral pair, can be represented by a *minimal polyhedron*, that is, by a polyhedron that contains an equilibrium point in the relative interior of every face, edge and vertex. An illustration of polyhedral pairs, and some examples of minimal polyhedra are given in Figure 1. There is no similar result for secondary and tertiary equilibrium classes, though the problem seems to be related to the problem of centering Koebe polyhedra (see [25], and for more information on centering Koebe polyhedra, see [66]).

While the above result answers our question for all classes except for monostable and mono-unstable ones, only estimates are known about the complexities of the classes of monostatic polyhedra. In particular, the monostable polyhedron P_{CG} of Conway and Guy in [21] is in class $\{1, 4\}_p$ and has complexity $C(P_{CG}) = 96$. The ones P_B and P_R , constructed by Bezdek and Reshetov, are in classes $\{1, 3\}_p$ and $\{1, 2\}_p$, respectively, having complexity $C(P_B) = 64$ and $C(P_R) = 70$. The authors of [25] extended these constructions for all classes $\{1, U\}_p$ with $U \geq 2$, and constructed polyhedra in classes $\{S, 1\}_p$ for all $S \geq 2$, yielding the estimates $C(2, 1) \leq 66$, $C(3, 1) \leq 64$, and for all $S, U \geq 4$, $C(S, 1) \leq 61 + 2 \lfloor \frac{S}{2} \rfloor$ and $C(1, U) \leq 92 + 2 \lfloor \frac{U}{2} \rfloor$. A lower bound $v \geq 7$ for the number v of vertices of mono-unstable polyhedra was established by [14] using a computational approach based on semi-definite programming. Presently there is no explicit estimate for the complexity of the class $\{1, 1\}_p$. However it has been conjectured that $C(1, 1)$ is at least approximately 1000 [111].

The picture outlined above changes significantly, if the centroid of the polyhedron is replaced by the center of mass of the vertices. In this case, mono-unstable polyhedra with less than eight vertices do not exist [13], but a mono-monostatic polyhedron with 21 faces and 21 vertices has been found [24]. If the assumption of homogeneity is relaxed, and the reference point is permitted to be an arbitrary interior point of the polyhedron, then even some (non-regular) tetrahedra may be monostable as first pointed out by Conway (see [80]) whereas other tetrahedra may be mono-unstable [3]. It is also shown in [3] that all monostable and mono-unstable tetrahedra belong to the equilibrium class $\{1, 2\}_p$ and $\{2, 1\}_p$, respectively. In higher dimensions, even some regular polytopes may be monostable [80].

4. RELATED PROBLEMS AND APPLICATIONS

The concepts of equilibrium and stability are closely related to the equilibria of floating bodies, which is the main focus of a review paper of Alfonseca et al. in preparation. The equilibrium positions of a convex body K floating in a liquid correspond to the equilibrium positions in the sense of Definition 3.1 of a related body $U_V(K)$, called the *Ulam floating body* of K [60]. The boundary of this body is also referred to as the *surface of buoyancy* (Alfonseca et al., in preparation). For any $0 < V < \text{vol}_d(K)$, the surface of buoyancy contains the centroids of the intersections of K with all closed half spaces H that satisfy the condition $\text{vol}_d(K \cap H) = V$. Thus, if K is symmetric and $V = \frac{1}{2} \text{vol}_d(K)$ coincides with the centroid body of K (see Subsection 2.2). Moreover, in the limit cases $V \rightarrow 0$ and $V \rightarrow \text{vol}_d(K)$, up to homothety, $U_V(K)$ tends to the original body K .

A possible generalization of the notion of equilibrium point of a convex body is the following [2]: given an angle $0 < \alpha < \frac{\pi}{2}$, we say that a point q in the boundary

of a convex body K is an *equilibrium point of K of parameter α* , with respect to c , if K has a supporting hyperplane at q with angle equal to $\frac{\pi}{2} - \alpha$ to $[c, q]$. This definition is motivated by physical objects that can be balanced at a point on planes with slope angle $-\frac{\pi}{2} < \alpha < \frac{\pi}{2}$ provided that there is sufficient friction to prevent downhill slip. The extension of stability is also straightforward in two dimensions in the case $c = o$ by noting that such points correspond to critical points of the function $u \mapsto \cos \alpha h_K(u) + \sin \alpha s(u)$, $u \in \mathbb{S}^1$, where $s(u)$ is the (signed) arclength of the arc of $\text{bd}(K)$ between $\rho_K(u)u \in \text{bd}(K)$ and an arbitrary fixed reference point in $\text{bd}(K)$. This function corresponds to the potential energy function of a heavy plane body rolling on a slope without slip. Allemann, Hungerbühler and Wasem [2] showed that the number of equilibria of parameter α of a plane convex body K is a piecewise constant function of α , which drops to zero outside a bounded interval (with 0 in its interior) of the slope angle α . In dimensions $d \geq 3$ there is no straightforward way of defining stability, as strictly convex bodies can always escape from an equilibrium by means of a spin motion.

The concepts outlined in this paper can also be extended to equilibria of objects enclosed in a sphere of radius α [110]; in this context the equilibrium points in Definition 3.1 correspond to the limit case $\alpha \rightarrow \infty$. The equilibria of a convex body K in a sphere of radius α are in one-to-one correspondence with the equilibrium points of the α -hull of K , defined as the intersection of all balls of radius α that contain K (for the definition of α -convex sets and a systematic investigation of these bodies, the reader is referred to [83] and [10], respectively). The number of equilibrium points of a plane convex body K with respect to an interior point c , defined in this way, is a piecewise constant, strictly increasing function of α , with minimum value of two for every c different from the center of the smallest ball containing K . In case of 3-dimensional convex bodies, this function is not necessarily monotonic, but it also drops to two if α is sufficiently close to the circumradius of the body. In other words, almost all objects become mono-monostatic inside a sufficiently small sphere [110].

A further generalization considers equilibria of objects with multiple contact points on an underlying surface with arbitrary geometry. A partial characterization of the positions of the centers of mass corresponding to equilibria has been given by [93, 98], and rich geometric structures have been uncovered. The stability analysis of these systems relies heavily on the theory of dynamical systems [96, 112] and is beyond the scope of this work.

We finish the paper by briefly mentioning a few applications of the above results outside mathematics. The theorem of Winternitz (see Subsection 2.1) has recently found an application to data depth in statistics [88]. Equilibrium classes (see Subsections 3.1, 3.2) are used as a shape classification scheme in geology [34, 65, 69]. Flocks of equilibria (see Subsection 3.4) help to understand fascinating geological formations called *balancing rocks* [5, 71].

Monostable objects play special roles in industrial object manipulation processes called part feeding [6], and the statistical behaviour of randomly dropped objects with multiple stable equilibria is also attracts much interest [49, 109]. Monostable shapes also appear in nature [20, 35] as well as in robotics [108, 118], and they allow spontaneous self-righting if an object topples. A further medical application of monostable objects concerns autonomous drug delivery [1].

REFERENCES

- [1] Alex Abramson, Ester Caffarel-Salvador, Minsoo Khang, David Dellal, David Silverstein, Yuan Gao, Morten Revsgaard Frederiksen, Andreas Vegge, František Hubálek, Jorrit J Water, et al. An ingestible self-orienting system for oral delivery of macromolecules. *Science*, 363(6427):611–615, 2019.
- [2] Jonas Allemann, Norbert Hungerbühler, and Micha Wasem. Equilibria of plane convex bodies. *J. Nonlinear Sci.*, 31(5):86, 2021.
- [3] Gergő Almádi, Robert J MacG. Dawson, Gábor Domokos, and Krisztina Regős. On equilibria of tetrahedra. *Math. Intelligencer*, pages 1–8, 2023.
- [4] Vladimir Batagelj. An inductive definition of the class of 3-connected quadrangulations of the plane. *Discrete Math.*, 78(1-2):45–53, 1989.
- [5] John W Bell, James N Brune, Tanzhuo Liu, Marek Zreda, and James C Yount. Dating precariously balanced rocks in seismically active parts of California and Nevada. *Geology*, 26(6):495–498, 1998.
- [6] Dina R Berkowitz and John Canny. Designing parts feeders using dynamic simulation. In *Proceedings of IEEE International Conference on Robotics and Automation*, volume 2, pages 1127–1132. IEEE, 1996.
- [7] Dimitris Bertsimas and Santosh Vempala. Solving convex programs by random walks. *J. ACM*, 51(4):540–556, 2004.
- [8] Florian Besau, Thomas Hack, Peter Pivovarov, and Franz E Schuster. Spherical centroid bodies. *Amer. J. Math.*, 145(2):515–542, 2023.
- [9] Andras Bezdek. On stability of polyhedra. In *Workshop on Discrete Geometry, Sep 13-16 2011, Fields Institute, Canada*, pages 2490—2491, 2011.
- [10] Károly Bezdek, Zsolt Lángi, Márton Naszódi, and Peter Papez. Ball-polyhedra. *Discrete Comput. Geom.*, 38(2):201–230, 2007.
- [11] Ted Bisztriczky and Károly Böröczky, Jr. About the centroid body and the ellipsoid of inertia. *Mathematika*, 48(1-2):1–13, 2001.
- [12] Wilhelm Blaschke. *Vorlesungen über Differentialgeometrie. II, Affine Differentialgeometrie*. Springer, 1923.
- [13] Sándor Bozóki, Gábor Domokos, Flórián Kovács, and Krisztina Regős. Mono-unstable polyhedra with point masses have at least 8 vertices. *Int. J. Solids Struct.*, 234:111276, 2022.
- [14] Sándor Bozóki, Gábor Domokos, Dávid Papp, and Krisztina Regős. The smallest mono-unstable, homogeneous convex polyhedron has at least 7 vertices. *arXiv preprint arXiv:2401.17906*, 2024.
- [15] Stefano Campi and Paolo Gronchi. The L^p -Busemann-Petty centroid inequality. *Adv. Math.*, 167(1):128–141, 2002.
- [16] Stefano Campi and Paolo Gronchi. On the reverse L^p -Busemann-Petty centroid inequality. *Mathematika*, 49(1-2):1–11, 2002.
- [17] Gulbank D Chakerian. The number of diameters through a point inside an oval. *Rev. Un. Mat. Argentina*, 29:282–290, 1984.
- [18] Fangwei Chen and Congli Yang. A new proof of the Orlicz-Lorentz centroid inequality. *J. Inequal. Appl.*, pages Paper No. 12, 12, 2019.
- [19] Fangwei Chen, Jiazou Zhou, and Congli Yang. On the reverse Orlicz Busemann-Petty centroid inequality. *Adv. in Appl. Math.*, 47(4):820–828, 2011.
- [20] Ylenia Chiari, Arie van Der Meijden, Adalgisa Caccone, Julien Claude, and Benjamin Gilles. Self-righting potential and the evolution of shell shape in Galápagos tortoises. *Sci. Rep.*, 7(1):15828, 2017.
- [21] John H Conway and Richard K Guy. Stability of polyhedra. *SIAM Rev.*, 8(3):381, 1966.
- [22] Hallard T Croft, Kenneth J Falconer, and Richard K Guy. *Unsolved problems in geometry*. Problem Books in Mathematics. Springer-Verlag, New York, 1994. Corrected reprint of the 1991 original [MR1107516 (92c:52001)], *Unsolved Problems in Intuitive Mathematics, II*.
- [23] Gábor Domokos, Philip Holmes, and Zsolt Lángi. A genealogy of convex solids via local and global bifurcations of gradient vector fields. *J. Nonlinear Sci.*, 26:1789–1815, 2016.
- [24] Gábor Domokos and Flórián Kovács. Conway’s spiral and a discrete Gömböc with 21 point masses. *Amer. Math. Monthly*, 130(9):795–807, 2023.
- [25] Gábor Domokos, Flórián Kovács, Zsolt Lángi, Krisztina Regős, and Péter T Varga. Balancing polyhedra. *Ars Math. Contemp.*, 19(1):95–124, 2020.

- [26] Gábor Domokos and Zsolt Lángi. The robustness of equilibria on convex solids. *Mathematika*, 60(1):237–256, 2014.
- [27] Gábor Domokos and Zsolt Lángi. On the average number of normals through points of a convex body. *Studia Sci. Math. Hungar.*, 52(4):457–476, 2015.
- [28] Gábor Domokos and Zsolt Lángi. The evolution of geological shape descriptors under distance-driven flows. *Math. Geosci.*, 50(3):337–363, 2018.
- [29] Gábor Domokos, Zsolt Lángi, and András A. Sipos. Tracking critical points on evolving curves and surfaces. *Exp. Math.*, 31(1):1–20, 2022.
- [30] Gábor Domokos, Zsolt Lángi, and Tímea Szabó. On the equilibria of finely discretized curves and surfaces. *Monatsh. Math.*, 168(3-4):321–345, 2012.
- [31] Gábor Domokos, Zsolt Lángi, and Tímea Szabó. A topological classification of convex bodies. *Geom. Dedicata*, 182:95–116, 2016.
- [32] Gábor Domokos, Zsolt Lángi, and Péter L Várkonyi. A characterization of the symmetry groups of mono-monostatic convex bodies. *Monatsh. Math.*, 201(3):703–724, 2023.
- [33] Gábor Domokos, Jim Papadopoulos, and Andy Ruina. Static equilibria of planar, rigid bodies: is there anything new? *J. Elasticity*, 36(1):59–66, 1994.
- [34] Gábor Domokos, András Sipos, Tímea Szabó, and Péter Várkonyi. Pebbles, shapes, and equilibria. *Math. Geosci.*, 42:29–47, 2010.
- [35] Gábor Domokos and Péter L Várkonyi. Geometry and self-righting of turtles. *P. Roy. Soc. B - Biol. Sci.*, 275(1630):11–17, 2008.
- [36] Shen Dong, Peer-Timo Bremer, Michael Garland, Valerio Pascucci, and John C Hart. Spectral surface quadrangulation. *ACM Trans. Graphics*, 25(3):1057–1066, 2006.
- [37] Stelian Dumitraşcu. Every convex polygon is swept by its inner normal more than 4 times. *An. Univ. Timișoara Ser. Mat.-Inform.*, 36(1):43–58, 1998.
- [38] Adrian Dumitrescu and Csaba D Tóth. On the cover of the rolling stone. In *Proceedings of the 2020 ACM-SIAM Symposium on Discrete Algorithms*, pages 2575–2586. SIAM, Philadelphia, PA, 2020.
- [39] Herbert Edelsbrunner, John Harer, and Afra Zomorodian. Hierarchical Morse—Smale complexes for piecewise linear 2-manifolds. *Discrete Comput. Geom.*, 30:87–107, 2003.
- [40] Frederick Fabricius-Bjerre. Centroids and medians in spherical space. II. *Mat. Tidsskr. B*, 1947:48–52, 1947.
- [41] Niufa Fang, Xiao Li, and Di Ma. L_p centroid bodies with respect to weights $|x|^\ell$. *J. Math. Anal. Appl.*, 526(1):Paper No. 127321, 17, 2023.
- [42] Y Feng and T Ma. On the reverse Orlicz-Lorentz Busemann-Petty centroid inequality. *Acta Math. Hungar.*, 159(1):211–228, 2019.
- [43] David Fog. Centroids and medians in spherical space. I. *Mat. Tidsskr. B*, 1947:41–47, 1947.
- [44] Matthieu Fradelizi, Mathieu Meyer, and Vlad Yaskin. On the volume of sections of a convex body by cones. *Proc. Amer. Math. Soc.*, 145(7):3153–3164, 2017.
- [45] Gregory A Gal’perin. The concept of the center of mass of a system of mass points in spaces of constant curvature. *Dokl. Akad. Nauk SSSR*, 302(5):1039–1044, 1988.
- [46] Gregory A Gal’perin. A concept of the mass center of a system of material points in the constant curvature spaces. *Comm. Math. Phys.*, 154(1):63–84, 1993.
- [47] Richard J Gardner. *Geometric tomography*, volume 58. Cambridge University Press Cambridge, 1995.
- [48] Mohammad Ghomi. The problem of optimal smoothing for convex functions. *Proc. Amer. Math. Soc.*, 130(8):2255–2259, 2002.
- [49] Ken Goldberg, Brian V Mirtich, Yan Zhuang, John Craig, Brian R Carlisle, and John Canny. Part pose statistics: Estimators and experiments. *IEEE T. Robot.*, 15(5):849–857, 1999.
- [50] Michael Goldberg and Richard K Guy. Stability of polyhedra (JH Conway and RK Guy). *SIAM Rev.*, 11(1):78–82, 1969.
- [51] Alexandr Grebennikov and Gaiane Panina. A note on the concurrent normal conjecture. *Acta Math. Hungar.*, 167(2):529–532, 2022.
- [52] Helmut Groemer. Stability theorems for two measures of symmetry. volume 24, pages 301–311. 2000. The Branko Grünbaum birthday issue.
- [53] Branko Grünbaum. Partitions of mass-distributions and of convex bodies by hyperplanes. *Pacific J. Math.*, 10:1257–1261, 1960.

- [54] John Guckenheimer and Philip Holmes. *Nonlinear oscillations, dynamical systems, and bifurcations of vector fields*, volume 42 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1990. Revised and corrected reprint of the 1983 original.
- [55] Christoph Haberl. Complex affine isoperimetric inequalities. *Calc. Var. Partial Differential Equations*, 58(5):Paper No. 169, 22, 2019.
- [56] Christoph Haberl and Franz E Schuster. General L_p affine isoperimetric inequalities. *J. Differential Geom.*, 83(1):1–26, 2009.
- [57] Kathy Hann. The average number of normals through a point in a convex body and a related Euler-type identity. *Geom. Dedicata*, 48:27–55, 1993.
- [58] Kathy Hann. What’s the bound on the average number of normals? *Amer. Math. Monthly*, 103(10):897–900, 1996.
- [59] Erhard Heil. Concurrent normals and critical points under weak smoothness assumptions. In *Discrete geometry and convexity (New York, 1982)*, volume 440 of *Ann. New York Acad. Sci.*, pages 170–178. New York Acad. Sci., New York, 1985.
- [60] Han Huang, Boaz A Slomka, and Elisabeth M Werner. Ulam floating bodies. *J. Lond. Math. Soc. (2)*, 100(2):425–446, 2019.
- [61] QingZhong Huang and BinWu He. An asymmetric Orlicz centroid inequality for probability measures. *Sci. China Math.*, 57(6):1193–1202, 2014.
- [62] Daniel Hug. On the mean number of normals through a point in the interior of a convex body. *Geom. Dedicata*, 55:319–340, 1995.
- [63] Mohammad N Ivaki. The planar Busemann-Petty centroid inequality and its stability. *Trans. Amer. Math. Soc.*, 368(5):3539–3563, 2016.
- [64] Tian Jin and Gangsong Leng. The properties of complex centroid body. *J. Shanghai Univ. Nat. Sci.*, 27(6):1144–1148, 2021.
- [65] Gábor Korvin. The shape of pebbles, grains and pores. In *Statistical Rock Physics*, pages 229–263. Springer, 2024.
- [66] Zsolt Lángi. Centering Koebe polyhedra via Möbius transformations. *Groups Geom. Dyn.*, 15(1):197–221, 2021.
- [67] Zsolt Lángi. A solution to some problems of Conway and Guy on monostable polyhedra. *Bull. Lond. Math. Soc.*, 54(2):501–516, 2022.
- [68] Chao Li and Weidong Wang. Inequalities for complex centroid bodies. *Acta Math. Hungar.*, 161(1):313–326, 2020.
- [69] Balázs Ludmány and Gábor Domokos. Pebbles, graphs and equilibria: Higher order shape descriptors for sedimentary particles. *Centr. Eur. Geol.*, 65(2):158–177, 2023.
- [70] Balázs Ludmány, Zsolt Lángi, and Gábor Domokos. Morse–Smale complexes on convex polyhedra. *Period. Math. Hungar.*, 2024.
- [71] Balázs Ludmány, Ignacio Pérez-Rey, Gábor Domokos, Mauro Muñoz-Menéndez, Leandro R Alejano, and András Á Sipos. A new insight into the stability of precariously balanced rocks. *Rock Mech. Rock Eng.*, 56(5):3539–3550, 2023.
- [72] Erwin Lutwak, Deane Yang, and Gaoyong Zhang. L_p affine isoperimetric inequalities. *J. Differential Geom.*, 56(1):111–132, 2000.
- [73] Erwin Lutwak, Deane Yang, and Gaoyong Zhang. Orlicz centroid bodies. *J. Differential Geom.*, 84(2):365–387, 2010.
- [74] Erwin Lutwak and Gaoyong Zhang. Blaschke-Santaló inequalities. *J. Differential Geom.*, 47(1):1–16, 1997.
- [75] Songjun Lv, Shengjie Zhang, and Shiyu Shen. L_p Busemann-Petty centroid inequality in hyperbolic and spherical spaces. *Arch. Math. (Basel)*, 120(6):651–663, 2023.
- [76] Tong-yi Ma and De-yan Zhang. L_p -centroid bodies and its characterizations. *Commun. Math. Res.*, 31(4):333–344, 2015.
- [77] Robert J MacG Dawson. Monostatic simplexes. *Amer. Math. Monthly*, 92(8):541–546, 1985.
- [78] Robert J MacG Dawson and Wendy Finbow. Monostatic simplexes iii. *Geom. Dedicata*, 84:101–113, 2001.
- [79] Robert J MacG Dawson, Wendy Finbow, and Philip Mak. Monostatic simplexes ii. *Geom. Dedicata*, 70:209–219, 1998.
- [80] Robert J MacG Dawson and Wendy A Finbow. What shape is a loaded die? *Math. Intelligencer*, 21:32–37, 1999.
- [81] Francisco Marín Sola. On general concavity extensions of Grünbaum type inequalities. *Bull. Braz. Math. Soc. (N.S.)*, 55(1):Paper No. 2, 17, 2024.

- [82] Francisco Marín Sola and Jesús Y Nicolás. On Grünbaum type inequalities. *J. Geom. Anal.*, 31(10):9981–9995, 2021.
- [83] Anton E Mayer. Eine Überkonvexität. *Math. Z.*, 39:511–531, 1935.
- [84] Mathieu Meyer and Shlomo Reisner. The convex intersection body of a convex body. *Glasg. Math. J.*, 53(3):523–534, 2011.
- [85] Vitali D Milman and Alain Pajor. Isotropic position and inertia ellipsoids and zonoids of the unit ball of a normed n -dimensional space. In *Geometric aspects of functional analysis (1987–88)*, volume 1376 of *Lecture Notes in Math.*, pages 64–104. Springer, Berlin, 1989.
- [86] Luis Montejano. On a problem of Ulam concerning a characterization of the sphere. *Stud. Appl. Math.*, 53(3):243–248, 1974.
- [87] Sergii Myroshnychenko, Matthew Stephen, and Ning Zhang. Grünbaum’s inequality for sections. *J. Funct. Anal.*, 275(9):2516–2537, 2018.
- [88] Stanislav Nagy, Carsten Schütt, and Elizabeth M Werner. Halfspace depth and floating body. *Stat. Surv.*, 13:52 – 118, 2019.
- [89] Ivan Nasonov, Gaiane Panina, and Dirk Siersma. Concurrent normals problem for convex polytopes and Euclidean distance degree. *arXiv preprint arXiv:2406.01773*, 2024.
- [90] Seiya Negami and Atsuhiko Nakamoto. Diagonal transformations of graphs on closed surfaces. *Sci. Rep. Yokohama Nat. Univ. Sect. I Math. Phys. Chem.*, (40):71–97, 1993.
- [91] Van Hoang Nguyen. Orlicz-Lorentz centroid bodies. *Adv. in Appl. Math.*, 92:99–121, 2018.
- [92] Apollonius of Perga. *Treatise on Conic Sections*. Cambridge University Press, 1896.
- [93] Yizhar Or and Elon Rimon. Computation and graphical characterization of robust multiple-contact postures in two-dimensional gravitational environments. *Int. J. Robot. Res.*, 25(11):1071–1086, 2006.
- [94] John Pardon. Concurrent normals to convex bodies and spaces of Morse functions. *Math. Ann.*, 352(1):55–71, 2012.
- [95] Clinton M Petty. Centroid surfaces. *Pacific J. Math.*, 11:1535–1547, 1961.
- [96] Michael Posa, Mark Tobenkin, and Russ Tedrake. Stability analysis and control of rigid-body systems with impacts and friction. *IEEE Trans. Automat. Control*, 61(6):1423–1437, 2015.
- [97] Alexander Reshetov. A unistable polyhedron with 14 faces. *Internat. J. Comput. Geom. Appl.*, 24(1):39–59, 2014.
- [98] Elon Rimon, Richard Mason, Joel W Burdick, and Yizhar Or. A general stance stability test based on stratified Morse theory with application to quasi-static locomotion planning. *IEEE T. Robot.*, 24(3):626–641, 2008.
- [99] Luis A Santaló. Note on convex spherical curves. *Bull. Amer. Math. Soc.*, 50(12):528–534, 1944.
- [100] Rolf Schneider. *Convex bodies: the Brunn-Minkowski theory*, volume 151 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, expanded edition, 2014.
- [101] Geoffrey C Shephard. Twenty problems on convex polyhedra. II. *Math. Gaz.*, 52:359–367, 1968.
- [102] Alexandra Shyntar and Vladyslav Yaskin. A generalization of Winternitz’s theorem and its discrete version. *Proc. Amer. Math. Soc.*, 149(7):3089–3104, 2021.
- [103] Millard L Sloan. Analytical Gomboc. *arXiv preprint arXiv:2306.14914*, 2023.
- [104] Ernst Steinitz. Polyeder und Raumeinteilungen. In *Enzykl. math. Wiss., Vol. 3 (Geometrie) Part 3AB12*, pages 1–139.
- [105] Ernst Steinitz. Über die Eulersche Polyederrelationen. *Arch. Math. Phys.*, 11:86–88, 1906.
- [106] Matthew Stephen and Vlad Yaskin. Applications of Grünbaum-type inequalities. *Trans. Amer. Math. Soc.*, 372(9):6755–6769, 2019.
- [107] Matthew Stephen and Ning Zhang. Grünbaum’s inequality for projections. *J. Funct. Anal.*, 272(6):2628–2640, 2017.
- [108] Edward Tunstel. Evolution of autonomous self-righting behaviors for articulated nanorovers. 1999.
- [109] Péter L Várkonyi. Estimating part pose statistics with application to industrial parts feeding and shape design: New metrics, algorithms, simulation experiments and datasets. *IEEE T. Autom. Sci. Eng.*, 11(3):658–667, 2014.
- [110] Péter L Várkonyi. Sensorless part feeding with round cages in two and three dimensions. *IEEE T. Robot. Autom.*, 1(2):724–731, 2016.

- [111] Péter L Várkonyi and Gábor Domokos. Static equilibria of rigid bodies: dice, pebbles, and the Poincaré-Hopf theorem. *J. Nonlinear Sci.*, 16:255–281, 2006.
- [112] Péter L Várkonyi and Yizhar Or. Lyapunov stability of a rigid body with two frictional contacts. *Nonlinear Dynam.*, 88:363–393, 2017.
- [113] Weidong Wang. A quermassintegral form of the L_p -Busemann-Petty centroid inequality. *J. Geom. Anal.*, 32(4):Paper No. 111, 9, 2022.
- [114] Denghui Wu and Jiazou Zhou. The LYZ centroid conjecture for star bodies. *Sci. China Math.*, 61(7):1273–1286, 2018.
- [115] Tudor Zamfirescu. How do convex bodies sit? *Mathematika*, 42(1):178–181, 1995.
- [116] Zengle Zhang. The Orlicz-Lorentz centroid inequality for star bodies*. *Monatsh. Math.*, 200(1):179–190, 2023.
- [117] Zengle Zhang and Niufa Fang. A new proof of the Orlicz-Lorentz Busemann-Petty centroid inequality. *J. Math. Inequal.*, 13(3):703–712, 2019.
- [118] Jianguo Zhao, Jing Xu, Bingtuan Gao, Ning Xi, Fernando J Cintron, Matt W Mutka, and Li Xiao. MSU jumper: A single-motor-actuated miniature steerable jumping robot. *IEEE T. Robot.*, 29(3):602–614, 2013.
- [119] Guangxian Zhu. The Orlicz centroid inequality for star bodies. *Adv. in Appl. Math.*, 48(2):432–445, 2012.

ZSOLT LÁNGI, DEPARTMENT OF ALGEBRA AND GEOMETRY, BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS,, MŰEGYETEM RKP. 3., H-1111 BUDAPEST, HUNGARY, AND, ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS,, REÁLTANODA UTCA 13-15, H-1053, BUDAPEST, HUNGARY
Email address: zlangi@math.bme.hu

PÉTER L. VÁRKONYI, DEPARTMENT OF MECHANICS, MATERIALS, AND STRUCTURES, BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS,, MŰEGYETEM RKP. 3., H-1111 BUDAPEST, HUNGARY
Email address: varkonyi.peter@epk.bme.hu