LOCAL EXACT LAGRANGIAN CONTROLLABILITY FOR 1D BAROTROPIC COMPRESSIBLE NAVIER–STOKES EQUATIONS

KAI KOIKE, FRANCK SUEUR, AND GASTÓN VERGARA-HERMOSILLA

ABSTRACT. We consider a viscous compressible barotropic flow in the interval $[0, \pi]$ with homogeneous Dirichlet boundary conditions for the flow velocity and a constant rest state as initial data. Given two sufficiently close subintervals $I = [\alpha_1, \alpha_2]$ and $J = [\beta_1, \beta_2]$ of (0, 1), a nonempty open set $\omega \subset (1, \pi)$, and T > 0, we construct an external force f supported in ω acting on the momentum equation such that the corresponding flow map moves the fluid particles initially occupying I exactly onto J in time T.

1. INTRODUCTION

Let a viscous compressible fluid occupy the interval $[0, \pi]$. Denote by $\rho = \rho(t, x)$ and u = u(t, x) the density and the velocity of the fluid, respectively. We assume that the flow is barotropic, that is, the pressure $p = p(\rho)$ is a function only of ρ . Moreover, we assume a physically natural condition that p is smooth around $\rho = 1$ and that p'(1) > 0. Assuming also that the fluid is initially at rest and cannot escape from both ends of $[0, \pi]$, the time evolution of (ρ, u) can be described by the following barotropic compressible Navier–Stokes equations:

(1)
$$\begin{cases} \rho_t + (\rho u)_x = 0 & \text{in } (0, T) \times (0, \pi), \\ \rho(u_t + uu_x) + p(\rho)_x = u_{xx} + f & \text{in } (0, T) \times (0, \pi), \\ u(t, 0) = 0, \quad u(t, \pi) = 0 & \text{for } t \in (0, T), \\ (\rho, u)(0, x) = (1, 0) & \text{for } x \in (0, \pi). \end{cases}$$

Here *T* is a positive real number and the viscous coefficient in front of u_{xx} is set to unity for simplicity. More importantly, $f \in L^2((0,T) \times (0,\pi))$ is an external force whose support in *x* is included in a nonempty open set $\omega \subset (1,\pi)$.

Associated to the solution (ρ, u) to (1), which is determined by f, we define the flow map

$$\varphi = \varphi[f] \colon [0,T] \times [0,\pi] \to [0,\pi]$$

by $\varphi(0, x) = x$ and $\varphi_t(t, x) = u(t, \varphi(t, x))$. That this ODE is uniquely solvable (in its integral form) is guaranteed by the regularity $u \in L^1(0, T; \operatorname{Lip}(0, \pi))$ from the following proposition, which can be proved similarly to [11, Theorem 7.1]; note however that we only consider smooth f in this paper, so that u is actually smooth.

Date: July 31, 2024.

Proposition 1.1. For $f \in L^2((0,T) \times (0,\pi))$, there exists $\varepsilon_0 > 0$ such that if $\varepsilon := \|f\|_{L^2((0,T) \times (0,\pi))} \le \varepsilon_0$, then (1) has a unique strong solution in the class

$$\rho \in C([0,T]; H^1(0,\pi)) \cap C^1([0,T]; L^2(0,\pi)),$$

$$u \in C([0,T]; H_0^1(0,\pi)) \cap L^2(0,T; H^2(0,\pi)), \quad u_t \in L^2(0,T; L^2(0,\pi)).$$

Moreover, there exists a positive constant C > 0 such that

$$\sup_{0 \le t \le T} \left(\|\rho(t, \cdot) - 1\|_{H^1(0,\pi)} + \|u(t, \cdot)\|_{H^1(0,\pi)} \right) + \left(\int_0^T \|u_x(t, \cdot)\|_{H^1(0,\pi)}^2 \, dt \right)^{1/2} \le C\varepsilon.$$

1.1. Local exact Lagrangian controllability. The purpose of this paper is to investigate the *local exact Lagrangian controllability* for the compressible Navier–Stokes equations (1). That is, we ask whether one can choose an appropriate external force f supported in $\omega \subset (1, \pi)$ so that the flow map $\varphi = \varphi[f]$ moves the fluid particles initially occupying an interval $I = [\alpha_1, \alpha_2] \subset (0, 1)$ exactly onto another interval $J = [\beta_1, \beta_2] \subset (0, 1)$ in a given time T > 0; see Figure 1.



FIGURE 1. Can we find a remote control f = f(t, x) whose support in *x* is contained in ω such that the corresponding flow map $\varphi = \varphi[f]$ moves the interval $I = [\alpha_1, \alpha_2]$ exactly onto $J = [\beta_1, \beta_2]$ in a given time T > 0?

Our main result, Theorem 1.1 below, answers this question affirmatively when I and J are sufficiently close. In fact, we can achieve this using a smooth control of the form $f = \varepsilon_1 f_1 + \varepsilon_2 f_2$ (where $\varepsilon_1, \varepsilon_2 \in \mathbb{R}$) with f_1 and f_2 defined by

(2)
$$f_i(t,x) = \chi_\eta(x)\xi_i(t,x)$$

Here χ_{η} ($\eta > 0$) is a non-negative smooth function supported in ω such that

$$\chi_{\eta}(x) = 1 \quad (\operatorname{dist}(x, \omega^c) \ge \eta)$$

and (ζ_i, ξ_i) is the solution to the linearized adjoint system

(3)
$$\begin{cases} -\partial_t \zeta_i - \partial_x \xi_i = 0 & \text{in } (0, T) \times (0, \pi), \\ -\partial_t \xi_i - c^2 \partial_x \zeta_i = \partial_{xx} \xi_i + \delta_{\alpha_i} & \text{in } (0, T) \times (0, \pi), \\ \xi_i(t, 0) = 0, \quad \xi_i(t, \pi) = 0 & \text{for } t \in (0, T), \\ (\zeta_i, \xi_i)(T, x) = (0, 0) & \text{for } x \in (0, \pi). \end{cases}$$

Here, $c = \sqrt{p'(1)} > 0$ and δ_{α_i} is the Dirac delta function centered at $x = \alpha_i$. We note that this definition is inspired by [8] dealing with an analogous problem for

viscous Burgers' equation; the role of f_i becomes clear in the proof of Lemma 2.2. In Section 2.2, we prove the following properties of ξ_1 and ξ_2 :

Proposition 1.2. Let $\omega \subset (1, \pi)$ be a nonempty open set and $0 < \alpha_1 < \alpha_2 < 1$. Then ξ_1 and ξ_2 defined through (3) are bounded and smooth on $(0, T) \times (1, \pi)$. In particular, f_1 and f_2 are bounded and smooth on $(0, T) \times (0, \pi)$. Moreover, ξ_1 and ξ_2 are linearly independent in $L^2((0, T) \times \omega)$.

In what follows, we fix $\eta > 0$ small so that $\chi_{\eta}^{1/2}\xi_1$ and $\chi_{\eta}^{1/2}\xi_2$ are also linearly independent in $L^2((0,T) \times \omega)$. With these preparations, our main theorem reads:

Theorem 1.1. Let T > 0 and $\omega \subset (1, \pi)$ be a nonempty open set. Then for any $0 < \alpha_1 < \alpha_2 < 1$, there exists $\delta = \delta(\alpha_1, \alpha_2, T, \omega) > 0$ and a neighborhood U of (0,0) such that if $0 < \beta_1 < \beta_2 < 1$ satisfies $\max_{i=1,2}(|\alpha_i - \beta_i|) < \delta$, there exists a unique pair of real numbers $(\varepsilon_1, \varepsilon_2)$ in U satisfying $\varphi[\varepsilon_1 f_1 + \varepsilon_2 f_2](T, \alpha_i) = \beta_i$ for i = 1, 2.

Since the flow map preserves order, that is, $\varphi(T, x_1) < \varphi(T, x_2)$ if $0 \le x_1 < x_2 \le \pi$, the mapping of the end points $\varphi(T, \alpha_i) = \beta_i$ (i = 1, 2) implies the transportation of the whole interval $\varphi(T, I) = J$. Thus our main theorem shows that a remote action *f* can move the fluid particles initially occupying the interval $I = [\alpha_1, \alpha_2]$ exactly onto $J = [\beta_1, \beta_2]$ if both are close enough.

Remark 1.1. We can generalize Theorem 1.1 to more than two points: Let $0 < \alpha_1 < \alpha_2 < \cdots < \alpha_d < 1$ and $0 < \beta_1 < \beta_2 < \cdots < \beta_d < 1$. Define f_i by (2) for $i = 1, 2, \ldots, d$. Then if $\max_{i=1,2,\ldots,d} |\alpha_i - \beta_i|$ is sufficiently small, we can find $(\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_d)$ such that the flow map associated to $f = \sum_{i=1}^d \varepsilon_i f_i$ maps α_i to β_i for every $i = 1, 2, \ldots, d$ in a given time T > 0. The proof is almost identical except for an additional difficulty for which we use Lemma B.1 for general d.

1.2. Comparison with Eulerian controllability results. It is interesting to compare our theorem with known *Eulerian* controllability results. In [5], Ervedoza, Glass, Guerrero, and Puel proved an exact Eulerian controllability result for (1) with initial data in a small neighborhood of $(\rho, u) = (\bar{\rho}, \bar{u})$ with $\bar{\rho} > 0$ and $\bar{u} \neq 0$. More precisely, they proved the following:

Theorem 1.2 ([5, Theorem 1.1]). Let L and T be positive numbers. Assume that $\bar{\rho} > 0$ and $\bar{u} \neq 0$ satisfy $T > L/|\bar{u}|$. Then there exists $\varepsilon_0 > 0$ such that for any $(\rho_0, u_0) \in H^3(0, L) \times H^3(0, L)$ with

$$\|u_0 - \bar{u}\|_{H^3(0,L)} + \|\rho_0 - \bar{\rho}\|_{H^3(0,L)} \le \varepsilon_0,$$

there exists a solution (ρ, u) to

(4)
$$\begin{cases} \rho_t + (\rho u)_x = 0 & \text{in } (0, T) \times (0, L), \\ \rho(u_t + uu_x) + p(\rho)_x = u_{xx} & \text{in } (0, T) \times (0, L), \\ (\rho, u)(0, x) = (\bar{\rho}, \bar{u}) & \text{for } x \in (0, L), \end{cases}$$

satisfying

$$(\rho, u)(T, \cdot) = (\bar{\rho}, \bar{u})$$
 in $(0, L)$.

Remark 1.2. Note that the missing boundary conditions in (4) play the role of controls. Thus this is a boundary controllability result. We remark that by restricting the domain from $(0, \pi)$ to (0, 1) in (1), Theorem 1.1 implies a boundary controllability result with controls acting only on x = 1.

In contrast with the theorem above, if $\bar{u} = 0$, it is proved that the linearization of (4) is not exactly Eulerian controllable [12, 2].¹ However, in these papers, it is also shown that the system is *approximately* Eulerian controllable even when $\bar{u} = 0$. Since Horsin showed in [8] that for viscous Burgers' equation, the local exact Lagrangian controllability can be deduced from the approximate Eulerian controllability—or rather its equivalent unique continuation property—it is natural to ask whether we can prove the local exact Lagrangian controllability for (1) as well. This is answered in the affirmative by Theorem 1.1.

1.3. Comparison with controllability results for incompressible flows. Controllability properties for incompressible flows are much well studied. Let us focus here on Lagrangian controllability results and we refer to [10] and the references therein for Eulerian ones.

First, the following global Lagrangian controllability result for 2D incompressible Euler equations are proved in [6]: Take two homotopic Jordan curves γ_1 and γ_2 bounding domains of the same area in a bounded domain $\Omega \subset \mathbb{R}^2$, and let Σ be a nonempty open subset of $\partial \Omega$. Then for any $\varepsilon > 0$ and smooth initial data, there exists a solution to the 2D incompressible Euler equations with the no-flux boundary condition on $\partial \Omega \setminus \Sigma$ such that the corresponding flow map moves γ_1 to γ_2 within an error smaller than ε in a given time. This result was later extended to the 3D case in [7] (see also [9] for another approach) and to incompressible viscous flows in [10].

All the works cited above use potential flows and the harmonic functions theory as basic building blocks of the proof. But since compressible potential flows are not directly related to harmonic functions and are hard to analyze, it seems difficult to employ these methods to compressible flows. Perhaps for this reason, at present, all the controllability results for compressible Navier–Stokes equations, including ours, are local ones. It remains an interesting future research direction to investigate global controllability theorems for compressible Euler or Navier–Stokes flows.

¹We should remark that this does not exclude the controllability for the *nonlinear* system as we might be able to control the nonlinear one using for example the return method of Coron [3]. We also note that the result in [1] does not contradict these non-controllability results as she considers the equations in a Lagrangian coordinate.

2. Proof

Let us briefly explain the strategy following [8]: Using f_1 and f_2 defined by (2), we introduce a map $\Theta: U' \to \mathbb{R}^2$ in a neighborhood U' of (0, 0); this map is designed so that if we can apply the inverse function theorem to Θ around (0, 0), Theorem 1.1 follows immediately. Therefore, the proof is reduced to proving the non-vanishing of the Jacobian of Θ at (0, 0). To prove this, the key point is the linear independence of ξ_1 and ξ_2 stated in Proposition 1.2, which is essentially a unique continuation property for the adjoint linear system (3); see Remark 2.2. For the viscous Burgers equation considered in [8], the adjoint linear system is the heat equation and the unique continuation property is easily obtained by Holmgren's theorem. For the compressible Navier–Stokes equations considered in this paper, we need a more subtle analysis. In the boundary controllability setting, this is showed in [12, 2] (in its equivalent approximate Eulerian controllability form) using Fourier analytic techniques. We adapt this technique to our distributed control setting. Let us pursue this strategy below.

2.1. Jacobian as a Gram determinant. First, let U' be a sufficiently small neighborhood of $(0,0) \in \mathbb{R}^2$ and set

$$\Theta \colon U' \ni (\varepsilon_1, \varepsilon_2) \mapsto (\varphi[\varepsilon_1 f_1 + \varepsilon_2 f_2](T, \alpha_1), \varphi[\varepsilon_1 f_1 + \varepsilon_2 f_2](T, \alpha_2)) \in \mathbb{R}^2$$

where f_1 and f_2 are defined by (2). We remind the reader that $\varphi[\varepsilon_1 f_1 + \varepsilon_2 f_2]$ is the flow map associated to the external force $f = \varepsilon_1 f_1 + \varepsilon_2 f_2$ and that the flow map is well-defined by Proposition 1.1. We also note that Θ is C^1 in a neighborhood of (0,0) which follows from smooth dependence on ε_1 and ε_2 of solutions to (1) with $f = \varepsilon_1 f_1 + \varepsilon_2 f_2$; here we need sufficient regularity of f_i , which is why we introduced a cutoff function in (2). So if we can show that the Jacobian of Θ at (0,0) is non-zero, that is,

(5)
$$J \coloneqq \det \begin{pmatrix} d\varphi[\varepsilon_1 f_1](T, \alpha_1)/d\varepsilon_1 \Big|_{\varepsilon_1 = 0} & d\varphi[\varepsilon_1 f_1](T, \alpha_2)/d\varepsilon_1 \Big|_{\varepsilon_1 = 0} \\ d\varphi[\varepsilon_2 f_2](T, \alpha_1)/d\varepsilon_2 \Big|_{\varepsilon_2 = 0} & d\varphi[\varepsilon_2 f_2](T, \alpha_2)/d\varepsilon_2 \Big|_{\varepsilon_2 = 0} \end{pmatrix} \neq 0,$$

it follows that there exists an open neighborhood U of (0,0) and V of $\Theta(0,0) = (\alpha_1, \alpha_2)$ such that Θ induces a one-to-one map from U to V. Note that this implies Theorem 1.1: If the interval $J = [\beta_1, \beta_2]$ is sufficient close to I, then $(\beta_1, \beta_2) \in V$, so we can find a unique $(\varepsilon_1, \varepsilon_2) \in U$ such that $\varphi[\varepsilon_1 f_1 + \varepsilon_2 f_2](T, \alpha_i) = \beta_i$ for i = 1, 2.

To show the non-vanishing of the Jacobian *J*, it turns out that it suffices to analyze the linearization of (1) around $(\rho, u) = (1, 0)$: We consider the solution (η, v) to

(6)
$$\begin{cases} \eta_t + v_x = g & \text{in } (0, T) \times (0, \pi), \\ v_t + c^2 \eta_x = v_{xx} + f & \text{in } (0, T) \times (0, \pi), \\ v(t, 0) = 0, \quad v(t, \pi) = 0 & \text{for } t \in (0, T), \\ (\eta, v)(0, x) = (0, 0) & \text{for } x \in (0, \pi), \end{cases}$$

where $c = \sqrt{p'(1)} > 0$. This claim is justified by Lemma 2.1 below. Before stating the lemma, let us note that by simple energy estimates, we can show that there exists $C_T > 0$ such that

(7)
$$\sup_{0 \le t \le T} \left(\|\eta(t, \cdot)\|_{L^2(0,\pi)} + \|v(t, \cdot)\|_{L^2(0,\pi)} \right) + \left(\int_0^T \|v_x(t, \cdot)\|_{L^2(0,\pi)}^2 \, dt \right)^{1/2} \\ \le C_T \left(\|f\|_{L^2((0,T) \times (0,\pi))} + \|g\|_{L^2((0,T) \times (0,\pi))} \right).$$

Lemma 2.1. Let (η_i, v_i) be the solution to (6) with $f = f_i$ and g = 0. Then we have

$$d\varphi[\varepsilon_i f_i](T, x)/d\varepsilon_i\Big|_{\varepsilon_i=0} = \int_0^T v_i(t, x) dt$$

for all $x \in (0, \pi)$.

Proof. Fix $x \in (0, \pi)$. Let (ρ_i, u_i) be the solution to (1) with $f = \varepsilon_i f_i$. Note that

(8)
$$\varphi[\varepsilon_i f_i](t, x) = x + \int_0^t u_i(s, \varphi[\varepsilon_i f_i](s, x)) \, ds \quad (0 \le t \le T)$$

Using this equality and applying the mean value theorem, with the help of Sobolev inequalities, we obtain

$$\int_0^T |u_i(t,\varphi[\varepsilon_i f_i](t,x)) - u_i(t,x)| dt$$

$$\leq \int_0^T |\varphi[\varepsilon_i f_i](t,x) - x| \cdot ||\partial_x u_i(t,\cdot)||_{L^{\infty}(0,\pi)} dt$$

$$\leq C \int_0^T \int_0^t |u_i(s,\varphi[\varepsilon_i f_i](s,x))| ds \cdot ||\partial_x u_i(t,\cdot)||_{H^1(0,\pi)} dt$$

$$\leq C \int_0^T \int_0^t ||\partial_x u_i(s,\cdot)||_{L^2(0,\pi)} ds \cdot ||\partial_x u_i(t,\cdot)||_{H^1(0,\pi)} dt.$$

Then by Hölder's inequality as well as Proposition 1.1, we get

$$\begin{split} &\int_{0}^{T} \int_{0}^{t} \|\partial_{x} u_{i}(s, \cdot)\|_{L^{2}(0,\pi)} \, ds \cdot \|\partial_{x} u_{i}(t, \cdot)\|_{H^{1}(0,\pi)} \, dt \\ &\leq \int_{0}^{T} t^{1/2} \left(\int_{0}^{t} \|\partial_{x} u_{i}(s, \cdot)\|_{L^{2}(0,\pi)}^{2} \, ds \right)^{1/2} \cdot \|\partial_{x} u_{i}(t, \cdot)\|_{H^{1}(0,\pi)} \, dt \\ &\leq T^{1/2} \|\partial_{x} u_{i}\|_{L^{2}(0,T; \, L^{2}(0,\pi))} \int_{0}^{T} \|\partial_{x} u_{i}(t, \cdot)\|_{H^{1}(0,\pi)} \, dt \\ &\leq T \|\partial_{x} u_{i}\|_{L^{2}(0,T; \, H^{1}(0,\pi))}^{2} \leq CT \varepsilon_{i}^{2}. \end{split}$$

Next, since $(\eta, v) = (\rho_i - \varepsilon_i \eta_i, u_i - \varepsilon_i v_i)$ satisfies (6) with

$$f = -u_i \partial_x u_i - \frac{p'(\rho_i) - c^2(\rho_i - 1)}{\rho_i} \partial_x \rho_i - \frac{\rho_i - 1}{\rho_i} \partial_{xx} u_i - \frac{\rho_i - 1}{\rho_i} \varepsilon_i f_i,$$

$$g = -(\rho_i - 1) \partial_x u_i - (\partial_x \rho_i) u_i,$$

Proposition 1.1 and inequality (7) implies

$$\int_0^T |u_i(t,x) - \varepsilon_i v_i(t,x)| \, dt \le CT^{1/2} \|\partial_x v\|_{L^2(0,T;\,L^2(0,\pi))}$$
$$\le C_T \left(\|f\|_{L^2((0,T)\times(0,\pi))} + \|g\|_{L^2((0,T)\times(0,\pi))} \right) \le C_T \varepsilon_i^2.$$

Now using these inequalities to (8) with t = T, we conclude that

$$\varphi[\varepsilon_i f_i](T, x) = x + \int_0^T u_i(t, \varphi[\varepsilon_i f_i](t, x)) dt = x + \varepsilon_i \int_0^T v_i(t, x) dt + O(\varepsilon_i^2)$$

as $\varepsilon_i \to 0$. This proves the lemma.

Now by a duality argument, we can write the Jacobian *J* as the Gram determinant of $(\chi_{\eta}^{1/2}\xi_1, \chi_{\eta}^{1/2}\xi_2) \in L^2((0,T) \times \omega)$:

Lemma 2.2. The identity

$$J = \det\left(\int_0^T \int_\omega \chi_\eta \xi_i \xi_j \, dx \, dt\right)_{i,j=1,2}$$

holds for J defined by (5).

Proof. Define v_i as in Lemma 2.1. Then (3) and integration-by-parts imply²

$$\int_0^T v_i(t, \alpha_j) dt = \int_0^T \int_0^\pi v_i \delta_{\alpha_j} dx dt$$

= $\int_0^T \int_0^\pi v_i (-\partial_t \xi_j - c^2 \partial_x \zeta_j - \partial_{xx} \xi_j) dx dt$
= $\int_0^T \int_0^\pi v_i \left(-\partial_t \xi_j - c^2 \partial_{xx} \int_t^T \xi_j ds - \partial_{xx} \xi_j \right) dx dt$
= $\int_0^T \int_0^\pi \left(\partial_t v_i - c^2 \partial_{xx} \int_0^t v_i ds - \partial_{xx} v_i \right) \xi_j dx dt$
= $\int_0^T \int_0^\pi \left(\partial_t v_i + c^2 \partial_x \eta_i - \partial_{xx} v_i \right) \xi_j dx dt$
= $\int_0^T \int_0^\pi f_i \xi_j dx dt = \int_0^T \int_0^\pi \chi_\eta \xi_i \xi_j dx dt.$

This computation and Lemma 2.1 yield the desired result.

Remark 2.1. Our Lagrangian controllability problem can be recast into that for a system of ODEs: Let $u_{\varepsilon_1 f_1 + \varepsilon_2 f_2}$ be the solution to (1) with $f = \varepsilon_1 f_1 + \varepsilon_2 f_2$ and

²We note that $\xi_j(t, \cdot)$ is continuous on $(0, \pi)$ but its first derivative is discontinuous at $x = \alpha_j$. This fact can be shown using the explicit formulas in Lemma 2.3; see also Lemma 2.4. So the calculation below should be understood in the sense of distributions.

consider the ODE system

$$\begin{cases} dx_1(t)/dt = u_{\varepsilon_1 f_1 + \varepsilon_2 f_2}(t, x_1(t)) & (0 \le t \le T), \\ dx_2(t)/dt = u_{\varepsilon_1 f_1 + \varepsilon_2 f_2}(t, x_2(t)) & (0 \le t \le T), \\ x_1(0) = \alpha_1, \quad x_2(0) = \alpha_2. \end{cases}$$

Then we try to find appropriate ε_1 and ε_2 so that $x_1(T) = \beta_1$ and $x_2(T) = \beta_2$ hold. If we linearize this system around $(\varepsilon_1, \varepsilon_2, x_1, x_2) = (0, 0, \alpha_1, \alpha_2)$, we get

(9)
$$\begin{cases} \frac{d}{dt} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} (t) = \begin{bmatrix} v_1(t,\alpha_1) & v_2(t,\alpha_1) \\ v_1(t,\alpha_2) & v_2(t,\alpha_2) \end{bmatrix} \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \end{bmatrix} \quad (0 \le t \le T), \\ x_1(0) = \alpha_1, \quad x_2(0) = \alpha_2 \end{cases}$$

with v_i defined in Lemma 2.1. The controllability Gramian (see [4, Definition 1.10]) of (9) is

$$\left(\int_0^T v_i(t,\alpha_j)\right)_{i,j=1,2} = \left(\int_0^T \int_\omega \chi_\eta \xi_i \xi_j \, dx \, dt\right)_{i,j=1,2}$$

by the proof of Lemma 2.2. Therefore, the controllability of (9) is equivalent to the non-vanishing of our Jacobian J. Note that by the linear test [4, Theorem 3.6], the controllability of the linearized system implies the nonlinear local controllability result as well.

2.2. Smoothness and linear independence of ξ_1 and ξ_2 . By Lemma 2.2 and the Cauchy–Schwarz inequality, Theorem 1.1 is proved if we can show that ξ_1 and ξ_2 are linearly independent in $L^2((0,T) \times \omega)$ and $\eta > 0$ is taken small enough. So it remains to prove Proposition 1.2.

For this purpose, we first need an explicit formula for ξ_i :

Lemma 2.3. Let (ζ_i, ξ_i) be the solution to (3). If $n_0 := 2c$ is not an integer, we have

$$\xi_i(t,x) = \frac{2}{\pi} \sum_{n \ge 1} \frac{\sin n\alpha_i}{\mu_n - \lambda_n} \left(e^{\mu_n(T-t)} - e^{\lambda_n(T-t)} \right) \sin nx$$

If n_0 is an integer, we have

$$\xi_i(t,x) = \frac{2}{\pi}q(t)\sin n_0\alpha_i \sin n_0x + \frac{2}{\pi}\sum_{n\ge 1, n\ne n_0}\frac{\sin n\alpha_i}{\mu_n - \lambda_n} \left(e^{\mu_n(T-t)} - e^{\lambda_n(T-t)}\right)\sin nx$$

where

$$q(t) = \frac{1}{c^2} \left(1 - e^{-2c^2(T-t)} \right) - (T-t)e^{-2c^2(T-t)}.$$

Proof. To calculate the solution to the non-homogeneous system (3) by Duhamel's principle, we first need to compute the associated semigroup. Define the operator $A: [L^2(0,\pi)]^2 \supset D(A) \rightarrow [L^2(0,\pi)]^2$ by

$$D(A) = \left\{ (\zeta, \xi)^T \in [L^2(0, \pi)]^2 \mid \xi \in H^1_0(0, \pi), \ c^2 \zeta + \xi' \in H^1(0, \pi) \right\}$$

and

$$A = \begin{pmatrix} 0 & \partial_x \\ c^2 \partial_x & \partial_x^2 \end{pmatrix}.$$

It can be shown that -A is a maximal monotone operator in $[L^2(0,\pi)]^2$; see [2, Lemma 2.1]. Let us compute the semigroup $(S(t))_{t\geq 0}$ generated by -A following [2]. For this purpose, we introduce

$$\Phi_0(x) = \frac{1}{\sqrt{c^2 \pi}} (1, 0)^T,$$

$$\Phi_{2n}(x) = \sqrt{\frac{2}{c^2 \pi}} (\cos nx, 0)^T, \quad \Phi_{2n-1}(x) = \sqrt{\frac{2}{\pi}} (0, \sin nx)^T \quad (n \ge 1).$$

1

These form an orthonormal basis of $[L^2(0,\pi)]^2$ with respect to the inner product

$$\left\langle \begin{pmatrix} \zeta_1\\ \xi_1 \end{pmatrix}, \begin{pmatrix} \zeta_2\\ \xi_2 \end{pmatrix} \right\rangle = c^2 \int_0^\pi \zeta_1 \zeta_2 \, dx + \int_0^\pi \xi_1 \xi_2 \, dx$$

Note that the spaces

$$V_0 = \text{span}\{\Phi_0\}, \quad V_n = \text{span}\{\Phi_{2n}, \Phi_{2n-1}\} \quad (n \ge 1)$$

are invariant under A, and the restriction of A to V_n ($n \ge 1$) has the matrix representation

$$A_n = \begin{pmatrix} 0 & cn \\ -cn & -n^2 \end{pmatrix}$$

with respect to the basis (Φ_{2n}, Φ_{2n-1}) . If $n \neq n_0$, it can be diagonalized as

$$A_n = P_n \begin{pmatrix} \lambda_n & 0\\ 0 & \mu_n \end{pmatrix} P_n^{-1}$$

with

(10)
$$\lambda_n = -\frac{n^2}{2} - \frac{n^2}{2}\sqrt{1 - 4c^2/n^2}, \quad \mu_n = -\frac{n^2}{2} + \frac{n^2}{2}\sqrt{1 - 4c^2/n^2},$$

and

$$P_n = \begin{pmatrix} c & c \\ \lambda_n/n & \mu_n/n \end{pmatrix}.$$

Therefore, S(t) has the matrix representation

$$S_n(t) = e^{-A_n t} = P_n \begin{pmatrix} e^{-\lambda_n t} & 0\\ 0 & e^{-\mu_n t} \end{pmatrix} P_n^{-1}$$

on the space V_n . If $n = n_0$ on the other hand, A_n can be triangulated as

$$A_n = Q_n \begin{pmatrix} \lambda_n & 1 \\ 0 & \lambda_n \end{pmatrix} Q_n^{-1}$$

with

$$Q_n = \begin{pmatrix} c & -c/\lambda_n \\ \lambda_n/n & 0 \end{pmatrix},$$

and S(t) has the matrix representation

$$S_n(t) = Q_n \begin{pmatrix} e^{-\lambda_n t} & t e^{-\lambda_n t} \\ 0 & e^{-\lambda_n t} \end{pmatrix} Q_n^{-1}.$$

Now, by Duhamel's principle, the solution (ζ_i, ξ_i) to (3) can be written as

$$\begin{pmatrix} \zeta_i \\ \xi_i \end{pmatrix}(t) = \int_t^T S(t-\tau) \begin{pmatrix} 0 \\ \delta_{\alpha_i} \end{pmatrix} d\tau.$$

So we get

$$\begin{pmatrix} \zeta_i \\ \xi_i \end{pmatrix} (t, x) = \int_t^T \left\langle \begin{pmatrix} 0 \\ \delta_{\alpha_i} \end{pmatrix}, \Phi_0 \right\rangle \Phi_0 \, d\tau + \sum_{n \ge 1} \int_t^T \left(\Phi_{2n}, \Phi_{2n-1} \right) S_n(t-\tau) \left\{ \left\langle \begin{pmatrix} 0 \\ \delta_{\alpha_i} \end{pmatrix}, \Phi_{2n} \right\rangle e_1 + \left\langle \begin{pmatrix} 0 \\ \delta_{\alpha_i} \end{pmatrix}, \Phi_{2n-1} \right\rangle e_2 \right\} \, d\tau = \sum_{n \ge 1} \int_t^T \left(\Phi_{2n}, \Phi_{2n-1} \right) S_n(t-\tau) \left\langle \begin{pmatrix} 0 \\ \delta_{\alpha_i} \end{pmatrix}, \Phi_{2n-1} \right\rangle e_2 \, d\tau.$$

Here $e_1 = (1,0)^T$ and $e_2 = (0,1)^T$ are the standard basis of \mathbb{R}^2 . Computing these terms explicitly, we get the desired formulas.

The smoothness of ξ_i on $(0, T) \times (1, \pi)$ can be proved using Lemma 2.3:

Lemma 2.4. Let (ζ_i, ξ_i) be the solution to (3). Set $X_i = (0, \pi) \setminus {\alpha_i}$. Then ξ_i is bounded and smooth on $(0, T) \times X_i$.

Proof. The boundedness is clear from Lemma 2.3 and the fact that $\mu_n - \lambda_n = O(n^2)$ as $n \to \infty$. Now note that λ_n and μ_n are both real for all $n \ge n_0 = 2c$. By Lemma 2.3 and $\lambda_n \le -n^2/2$ for $n \ge n_0$, it suffices to show that

$$\sum_{n=n_0}^{\infty} \frac{\sin n\alpha_i}{\mu_n - \lambda_n} e^{\mu_n (T-t)} \sin nx = \sum_{n=n_0}^{\infty} \frac{1}{\mu_n - \lambda_n} e^{\mu_n (T-t)} \frac{1}{2} \left\{ \cos n(x - \alpha_i) - \cos n(x + \alpha_i) \right\}$$

is smooth on $(0, T) \times X_i$. For this purpose, we introduce

$$u(t,x) = \sum_{n=n_0}^{\infty} \frac{1}{\mu_n - \lambda_n} e^{\mu_n(T-t)} \cos nx.$$

It is easy to see that *u* is infinitely differentiable in *t* on $(0, T) \times \mathbb{R}$ and that

$$\partial_t^l u(t,x) = \sum_{n=n_0}^{\infty} \frac{(-\mu_n)^l}{\mu_n - \lambda_n} e^{\mu_n (T-t)} \cos nx$$

for any $l \ge 0$. To prove the lemma, it suffices to show that $\partial_t^l u$ is infinitely differentiable in x on $(0, T) \times \{(-2\pi, 0) \cup (0, 2\pi)\}$. Fix $t_0 \in (0, T)$. By (10), for any

integer $K \ge 1$, there exist real numbers $a_k = a_k(t_0, l)$ (k = 1, 2, ..., K) such that

$$\frac{(-\mu_n)^l}{\mu_n - \lambda_n} e^{\mu_n (T - t_0)} = \sum_{k=1}^K \frac{a_k}{n^{2k}} + O(n^{-2K-2}) \quad (n \to \infty).$$

Note that

$$\sum_{n=n_0}^{\infty} O(n^{-2K-2}) \cos nx$$

is 2*K*-times continuously differentiable on \mathbb{R} . On the other hand, it is well-known that³

$$\sum_{n=1}^{\infty} \frac{\cos nx}{n^2} = \frac{x^2}{4} - \frac{\pi x}{2} + \frac{\pi^2}{6} \quad (0 \le x \le 2\pi).$$

Therefore, the sum

$$\sum_{n=n_0}^{\infty} \frac{a_1}{n^2} \cos nx$$

is infinitely differentiable on $(-2\pi, 0) \cup (0, 2\pi)$. By repeating termwise integration, we see that

$$\sum_{n=n_0}^{\infty} \frac{a_k}{n^{2k}} \cos nx$$

is also infinitely differentiable on $(-2\pi, 0) \cup (0, 2\pi)$ for any $1 \le k \le K$. Since $K \ge 1$ is arbitrary, the proof is complete.

We finally prove the independence of ξ_1 and ξ_2 in $L^2((0,T) \times \omega)$:

Lemma 2.5. The functions ξ_1 and ξ_2 defined through (3) are linearly independent in $L^2((0,T) \times \omega)$.

Proof. Suppose that there exist $c_1, c_2 \in \mathbb{R}$ such that $c_1\xi_1 + c_2\xi_2 = 0$ in $L^2((0, T) \times \omega)$. By Lemma 2.3, if 2c is not an integer, we have

$$\xi_i(t,x) = \frac{2}{\pi} \sum_{n \ge 1} \frac{\sin n\alpha_i}{\mu_n - \lambda_n} \left(e^{\mu_n(T-t)} - e^{\lambda_n(T-t)} \right) \sin nx$$

for $(t, x) \in (0, T) \times \omega$, where λ_n and μ_n are defined by (10). By the density of irrational numbers, there exists $\gamma \in \mathbb{R} \setminus \mathbb{Q}$ such that $\gamma \pi \in \omega$. Since f_i is continuous on $(0, T) \times \omega$ by Lemma 2.4, we have

$$\sum_{n\geq 1} \frac{c_1 \sin n\alpha_1 + c_2 \sin n\alpha_2}{\mu_n - \lambda_n} \left(e^{\mu_n (T-t)} - e^{\lambda_n (T-t)} \right) \sin n\gamma \pi = 0.$$

Note that $\sin n\gamma \pi \neq 0$ for $n \geq 1$. As $\mu_n - \lambda_n = O(n^2)$, the series is absolutely convergent, and we can apply Lemma A.1 to conclude that $c_1 \sin n\alpha_1 + c_2 \sin n\alpha_2 = 0$

³Extend the polynomial on the right-hand side from $[0, \pi]$ to $[-\pi, \pi]$ as an even function; the left-hand side is then its cosine expansion.

for $n \ge 1$. By Lemma B.1, this implies $c_1 = c_2 = 0$. The case when 2c is an integer can be treated in a similar manner.

Remark 2.2. By a similar reasoning to that in the proof of Proposition 1.2, we can prove the following unique continuation property: Let (ζ, ξ) be the solution to

$$\begin{cases} -\zeta_t - \xi_x = 0 & \text{in } (0, T) \times (0, \pi), \\ -\xi_t - c^2 \zeta_x = \xi_{xx} & \text{in } (0, T) \times (0, \pi), \\ \xi(t, 0) = 0, \quad \xi(t, \pi) = 0 & \text{for } t \in (0, T), \\ (\zeta, \xi)(T, x) = (\zeta^T, \xi^T)(x) & \text{for } x \in (0, \pi) \end{cases}$$

with $(\zeta^T, \xi^T) \in H^1(0, \pi) \times L^2(0, \pi)$ and $\int_0^{\pi} \zeta^T dx = 0$. Then if $\xi \equiv 0$ on $(0, T) \times \omega$, we must have $(\zeta, \xi) \equiv (0, 0)$ on $(0, T) \times (0, \pi)$. We note that the condition $\xi \equiv 0$ on $(0, T) \times \omega$ can be replaced by $\xi(t, \gamma \pi) = 0$ for 0 < t < T, where $\gamma \in (0, 1)$ is an arbitrary irrational number. Thus the independence property in Proposition 1.2, which is the key point in the proof of Theorem 1.1, is essentially a unique continuation property for the linearized adjoint system.

Proposition 1.2 is obtained by combining Lemmas 2.4 and 2.5. Then taking $\eta > 0$ small enough, Propositions 1.2 and 2.2 imply that the Jacobian *J* is non-zero, which in turn proves Theorem 1.1 thanks to the inverse function theorem.

Appendix A. Generalization of a Lemma by Rosier and Rouchon

We state here a generalization of [12, Lemma 1]. The purpose of the generalization is to take care of the case when $n_0 = 2c$ is an integer in the proof of Lemma 2.5.

Lemma A.1. Let $(c_n)_{n\geq -1}$ and $(\lambda_n)_{n\geq 0}$ be two sequences of complex numbers with $\lambda_n \neq \lambda_n$ for $n \neq m$. Assume that λ_0 is real and $(\lambda)_{n\geq 1}$ are real for all but finitely many $n \geq 0$; that $\sum_{n\geq -1} |c_n| < \infty$; and that there exists $\Lambda \in \mathbb{R}$ such that $\operatorname{Re} \lambda_n \leq \Lambda$ for all $n \geq 0$. Then if

$$(c_{-1}t + c_0)e^{\lambda_0 t} + \sum_{n \ge 1} c_n e^{\lambda_n t} = 0 \quad (0 < t < T)$$

for some positive number T > 0, then $c_n = 0$ for all $n \ge -1$.

Proof. Let

$$F(z) = (c_{-1}z + c_0)e^{\lambda_0 z} + \sum_{n \ge 1} c_n e^{\lambda_n z}.$$

By the assumptions of the lemma, *F* is analytic on the right half-plane. Since F(t) = 0 for 0 < t < T, by the unique continuation property for analytic functions, it follows that F(z) = 0 for every *z* in the right half-plane. Now let I_+ be the finite set of $n \ge 1$ such that the imaginary part of λ_n is positive. Then since $\sum_{n\ge -1} |c_n| < \infty$,

$$\sum_{n \in I_+} c_n e^{\lambda_n (1-it)} = F(1-it) - \sum_{n \ge 1; n \notin I_+} c_n e^{\lambda_n (1-it)} = -\sum_{n \ge 1; n \notin I_+} c_n e^{\lambda_n (1-it)}$$

is bounded as $t \to \infty$. Let $\lambda_n = \mu_n + i\nu_n$ ($\mu_n, \nu_n \in \mathbb{R}$) and set $\nu = \max_{n \in I_+} \nu_n$. Then we see that

$$\sum_{n \in I_+; \nu_n = \nu} c_n e^{\lambda_n (1 - it)} = \left(\sum_{n \in I_+; \nu_n = \nu} c_n e^{\lambda_n} e^{-i\mu_n t} \right) e^{\nu t} = 0$$

because otherwise $\sum_{n \in I_+} c_n e^{\lambda_n(1-it)}$ is unbounded as $t \to \infty$. Therefore, we obtain $c_n = 0$ for $n \in I_+$ with $v_n = v$. Arguing similarly, it follows that $c_n = 0$ for any $n \in I_+$. Considering also the behavior as $t \to -\infty$, we get $c_n = 0$ for all n such that the imaginary part of λ_n is negative.

From the argument above, we may assume that $(\lambda_n)_{n\geq 0}$ are real for all $n \geq 0$. Now write $c'_{-1} = e^{\lambda_0}c_{-1}$ and $c'_n = e^{\lambda_n}c_n$ for $n \geq 0$. By taking the average of the relation $tF(1+it)e^{-i\lambda_0 t} = 0$ over the interval [-T, T], we get

$$ic'_{-1}T^2/3 = \frac{1}{2T} \int_{-T}^{T} t\left\{c'_{-1}(1+it) + c'_0\right\} dt = -\sum_{n\geq 1} \frac{1}{2T} \int_{-T}^{T} c'_n t e^{i(\lambda_n - \lambda_0)t} dt.$$

By the assumption $\sum_{n\geq -1} |c_n| < \infty$, for any $\delta > 0$, there exists a sufficiently large integer M such that

$$|c'_{-1}| \leq \sum_{1 \leq n \leq M} \frac{3}{2T^3} \left| \int_{-T}^T c'_n t e^{i(\lambda_n - \lambda_0)t} dt \right| + \delta.$$

Since

r

$$\frac{3}{2T^3} \int_{-T}^{T} t e^{i\lambda t} dt = \frac{3\cos\lambda T}{i\lambda T^2} - \frac{3\sin\lambda T}{i\lambda^2 T^3}$$

for $\lambda \in \mathbb{R} \setminus \{0\}$, by taking T > 0 sufficient large, we get $|c'_{-1}| \leq 2\delta$. Hence $c_{-1} = 0$.

The rest of the proof is the same as that of [12, Lemma 1]; we reproduce it here for the sake of completeness. By taking the average of the relation $F(1 + it)e^{-i\lambda_N t} = 0$ over the interval [-T, T], we get

$$c'_{-N} = -\sum_{n\geq 0;\,n\neq N} \frac{1}{2T} \int_{-T}^{T} c'_n e^{i(\lambda_n - \lambda_N)t} dt$$

for any $N \ge 0$. For any $\delta > 0$, there exists a sufficiently large integer M such that

$$|c'_N| \le \sum_{0 \le n \le M; n \ne N} \frac{1}{2T} \left| \int_{-T}^T c'_n e^{i(\lambda_n - \lambda_N)t} dt \right| + \delta.$$

Since

$$\frac{1}{2T} \int_{-T}^{T} e^{i\lambda t} dt = \frac{\sin \lambda T}{\lambda T}$$

for $\lambda \in \mathbb{R} \setminus \{0\}$, by taking T > 0 sufficient large, we get $|c'_N| \le 2\delta$. Hence, $c_N = 0$. This completes the proof. Appendix B. Linear independence of some trigonometric vectors

In the proof of Lemma 2.5, we used the fact that $c_1 \sin n\alpha_1 + c_2 \sin n\alpha_2 = 0$ for $n \ge 1$ implies $c_1 = c_2 = 0$. We give a proof of this in the following. In fact, to prove the extension mentioned in Remark 1.1, we prove a more general lemma:

Lemma B.1. Let $0 < \alpha_1 < \alpha_2 < \cdots < \alpha_d < \pi$ and let c_1, c_2, \ldots, c_d be real numbers. If

$$c_1\sin(n\alpha_1) + c_2\sin(n\alpha_2) + \dots + c_d\sin(n\alpha_d) = 0$$

holds for n = 1, 2, ..., d, then $c_1 = c_2 = \cdots = c_d = 0$.

To prove this, it suffices to show that

14

$$D = D(\alpha_1, \alpha_2, \dots, \alpha_d) \coloneqq \det \begin{pmatrix} \sin \alpha_1 & \cdots & \sin \alpha_i & \cdots & \sin \alpha_d \\ \vdots & \ddots & & \vdots \\ \sin i\alpha_1 & & \sin i\alpha_i & & \sin i\alpha_d \\ \vdots & & \ddots & \vdots \\ \sin d\alpha_1 & \cdots & \sin d\alpha_i & \cdots & \sin d\alpha_d \end{pmatrix} \neq 0.$$

This is a consequence of the following lemma:

Lemma B.2. Let $0 < \alpha_1 < \alpha_2 < \cdots < \alpha_d < \pi$. Then the following identity holds:

$$D(\alpha_1, \alpha_2, \dots, \alpha_d) = 2^{d(d-1)} \left(\prod_{i=1}^d \sin \alpha_i \right) \prod_{1 \le i < j \le d} \sin \left(\frac{\alpha_i - \alpha_j}{2} \right) \sin \left(\frac{\alpha_i + \alpha_j}{2} \right).$$

Proof. Let $1 \le i \le d$. By the multiple-angle formula

$$\sin(i\theta) = \sin\theta \sum_{k=0}^{\lfloor (i-1)/2 \rfloor} (-1)^k \binom{i}{2k+1} (1-\cos^2\theta)^k \cos^{i-2k-1}\theta$$

and an elementary identity involving binomial coefficients, it follows that there exists a polynomial $S_i(x)$ that satisfies:

- (i) $\sin(i\theta) = S_i(\cos\theta)\sin\theta$,
- (ii) $\deg S_i(x) = i 1$,
- (iii) the leading coefficient of $S_i(x)$ is 2^{i-1} .

Now by (i)-(iii) and some elementary row operations,

$$D = \det \begin{pmatrix} \sin \alpha_1 & \cdots & \sin \alpha_i & \cdots & \sin \alpha_d \\ \vdots & \ddots & & \vdots \\ S_i(\cos \alpha_1) \sin \alpha_1 & S_i(\cos \alpha_i) \sin \alpha_i & S_i(\cos \alpha_d) \sin \alpha_d \\ \vdots & & \ddots & \vdots \\ S_d(\cos \alpha_1) \sin \alpha_1 & \cdots & S_d(\cos \alpha_i) \sin \alpha_i & \cdots & S_d(\cos \alpha_d) \sin \alpha_d \end{pmatrix}$$

$$= \left(\prod_{i=1}^{d} \sin \alpha_{i}\right) \det \begin{pmatrix} 1 & \cdots & 1 & \cdots & 1 \\ \vdots & \ddots & & \vdots \\ S_{i}(\cos \alpha_{1}) & S_{i}(\cos \alpha_{i}) & S_{i}(\cos \alpha_{d}) \\ \vdots & & \ddots & \vdots \\ S_{d}(\cos \alpha_{1}) & \cdots & S_{d}(\cos \alpha_{i}) & \cdots & S_{d}(\cos \alpha_{d}) \end{pmatrix}$$
$$= \left(\prod_{i=1}^{d} \sin \alpha_{i}\right) \det \begin{pmatrix} 1 & \cdots & 1 & \cdots & 1 \\ \vdots & \ddots & & & \vdots \\ 2^{i-1} \cos^{i-1} \alpha_{1} & 2^{i-1} \cos^{i-1} \alpha_{i} & 2^{i-1} \cos^{i-1} \alpha_{d} \\ \vdots & & \ddots & \vdots \\ 2^{d-1} \cos^{d-1} \alpha_{1} & \cdots & 2^{d-1} \cos^{d-1} \alpha_{i} & \cdots & 2^{d-1} \cos^{d-1} \alpha_{d} \end{pmatrix}.$$

The determinant in the last line is a Vandermonde determinant. So we have

$$D = \left(\prod_{i=1}^{d} \sin \alpha_{i}\right) 2^{d(d-1)/2} \prod_{1 \le i < j \le d} (\cos \alpha_{j} - \cos \alpha_{i})$$
$$= 2^{d(d-1)} \left(\prod_{i=1}^{d} \sin \alpha_{i}\right) \prod_{1 \le i < j \le n} \sin \left(\frac{\alpha_{i} - \alpha_{j}}{2}\right) \sin \left(\frac{\alpha_{i} + \alpha_{j}}{2}\right).$$

This proves the lemma.

Acknowledgements. K. Koike was supported by JSPS Grant-in-Aid for Early-Career Scientists (Grant Number 22K13938) and Grant-in-Aid for Pioneering Research (Organization for Fundamental Research, Tokyo Institute of Technology). F. Sueur was supported by the Bourgeons project, grant ANR-23-CE40-0014-01 of the French National Research Agency (ANR). G. Vergara-Hermosilla was supported by the ANID program BCH 2022 grant No. 4220003. We thank Kazuaki Miyatani for giving a nice proof of Lemma B.2. We also acknowledge Cavallina Lorenzo for his suggestion to consider the generalization to more than two points (Remark 1.1). Finally, we are grateful to Sylvain Ervedoza for several helpful discussions.

Data availability statement. Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

Declarations. The authors declare that they have no conflicts of interest.

References

- 1. E. V. Amosova, *Exact local controllability for the equations of viscous gas dynamics*, Differentsial'nye Uravneniya **47** (2011), 1776–1795.
- S. Chowdhury, M. Ramaswamy, and J.-P. Raymond, *Controllability and stabilizability of the linearized compressible Navier–Stokes system in one dimension*, SIAM J. Control Optim. **50** (2012), 2959–2987.
- J.-M. Coron, Contrôlabilité exacte frontière de l'équation d'Euler des fluides parfaits incompressibles bidimensionnels, C. R. Acad. Sci. Paris 317 (1993), 271–276.
- 4. _____, *Control and nonlinearity*, Mathematical Surveys and Monographs, vol. 136, American Mathematical Society, Providence, RI, 2007.

15

KAI KOIKE, FRANCK SUEUR, AND GASTÓN VERGARA-HERMOSILLA

- S. Ervedoza, O. Glass, S. Guerrero, and J.-P. Puel, Local exact controllability for the onedimensional compressible Navier–Stokes equation, Arch. Rational Mech. Anal. 206 (2012), 189–238.
- O. Glass and T. Horsin, Approximate Lagrangian controllability for the 2-D Euler equation. Application to the control of the shape of vortex patches, J. Math. Pures Appl. 93 (2010), 61–90.
- 7. _____, *Prescribing the motion of a set of particles in a three-dimensional perfect fluid*, SIAM J. Control Optim. **50** (2012), 2726–2742.
- T. Horsin, Local exact Lagrangian controllability of the Burgers viscous equation, Ann. I. H. Poincaré 25 (2008), 219–230.
- 9. T. Horsin and O. Kavian, Lagrangian controllability of inviscid incompressible fluids: A constructive approach, ESAIM: Control Optim. Calc. Var. 23 (2017), 1179–1200.
- J. Liao, F. Sueur, and P. Zhang, Smooth controllability of the Navier–Stokes equation with Navier conditions: Application to Lagrangian controllability, Arch. Rational Mech. Anal. 243 (2022), 869–941.
- 11. A. Matsumura and T. Nishida, *The initial value problem for the equations of motion of viscous and heat-conductive gases*, J. Math. Kyoto Univ. **20** (1980), 67–104.
- L. Rosier and P. Rouchon, On the controllability of a wave equation with structural damping, Int. J. Tomogr. Stat. 5 (2007), 79–84.

Department of Mathematics, Tokyo Institute of Technology, 2-12-1, Ookayama, Meguroku, Tokyo 152-8551, Japan

Email address: koike.k@math.titech.ac.jp

16

Institut de Mathématiques de Bordeaux, UMR CNRS 5251, Université de Bordeaux, 351 cours de la Libération, F33405 Talence Cedex, France *Email address*: franck.sueur@icloud.com

LAMME, UNIV. EVRY, CNRS, UNIVERSITÉ PARIS-SACLAY, 23 BOULEVARD DE FRANCE, 91037 EVRY CEDEX, FRANCE

Email address: gaston.v-h@outlook.com