

Protecting Quantum Information via Many-Body Dynamical Localization

Ling-Zhi Tang,¹ Dan-Wei Zhang,^{1,2,*} Hai-Feng Yu,³ and Z. D. Wang^{4,†}

¹Key Laboratory of Atomic and Subatomic Structure and Quantum Control (Ministry of Education), Guangdong Basic Research Center of Excellence for Structure and Fundamental Interactions of Matter, South China Normal University, Guangzhou 510006, China

²Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Frontier Research Institute for Physics, School of Physics, South China Normal University, Guangzhou 510006, China

³Beijing Academy of Quantum Information Sciences, Beijing 100193, China

⁴Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Department of Physics, and HK Institute of Quantum Science & Technology, The University of Hong Kong, Pokfulam Road, Hong Kong, China

(Dated: July 30, 2024)

Dynamically localized states in quantum many-body systems are fundamentally important in understanding quantum thermalization and have applications in quantum information processing. Here we explore many-body dynamical localization (MBDL) without disorders in a non-integrable quantum XY spin chain under periodical and quadratic kicks. We obtain the localization phase diagram with the MBDL and delocalization states and show dynamical observables to extract the phase diagram. For proper kick strengths in the MBDL regime, we reveal a local dynamical decoupling effect for persistent Rabi oscillation of certain spins. Furthermore, we propose the MBDL-protected quantum information at high temperatures, and present an analysis of the dynamical decoupling to obtain the required system parameters for quantum storage. Compared to other non-thermalized states, the disorder-free MBDL states require much fewer repetitions and resources, providing a promising way to protect and store quantum information robust against thermal noises.

Introduction.— Coherent and controllable many-body quantum dynamics plays an important role in quantum matter engineering and quantum information processing. As a consequence of thermalization, however, coherence in generic many-body quantum systems is fragile once they are excited to a finite energy density above ground states [1, 2]. There are several ways to avoid thermalization, apart from integrability. One is the many-body localized (MBL) states induced by sufficiently strong disorders [3–6], and the other is quantum many-body scars [7–11], which correspond to the strong and weak breaking of ergodicity, respectively. In MBL systems, all excitations are localized and thermalization is strongly suppressed, which can protect coherence of any initial states. The scar states also have a long coherence time as their dynamics are effectively restricted in a subspace. These localized states may provide resources for quantum information processing [12–22].

Floquet engineering provides another versatile toolbox to understand thermalization and non-equilibrium phases in periodically driven quantum systems [23–31]. As a paradigm of quantum chaos, the quantum kicked rotor (QKR) can avoid chaotic dynamics due to the dynamical localization [32–36], which can be viewed as Anderson localization in momentum space and has been observed in ultracold atomic gases [37–40]. Recently, increasing efforts are made to study the dynamically localized states under many-body interactions, such as interacting QKR at the few-body limit [41–47], integrable case [48], and mean-field level [49–51], which were experimentally observed using Bose-Einstein condensates

in periodically pulsed optical lattices [52, 53]. Recent theoretical works suggest that the many-body dynamical localization (MBDL) persists in periodically kicked 1D Bose gases in strongly interacting regimes [54–57]. The observation of MBDL in an interaction-tunable 1D Bose gas was just reported in Ref. [58]. Simultaneously, quantum spin chains with tunable disorders and driving fields have been proposed to realize MBL and scarred (confined) states [59–65]. The periodically kicked Ising chains with exactly solvable dynamics can serve as a minimal model to study many-body quantum chaos [66–70]. Nevertheless, the MBDL in clean driven spin systems, which are amenable in quantum processors, and its related coherence protection remain unexplored.

In this work, we investigate the disorder-free MBDL in a non-integrable kicked XY spin chain and its application in protecting quantum information. For the quantum XY chain with a longitudinal field under periodic transverse-field kicks, we obtain the phase diagram with the MBDL and delocalization phases and show dynamical observables to extract the localization phase diagram. We uncover a local dynamical decoupling (DD) effect under proper kick strengths in the MBDL regime, which enables persistent Rabi oscillation of certain spins. Moreover, we demonstrate the MBDL-protected quantum information at high temperatures, and present an analysis of the local DD to obtain the system parameters for robust quantum storage. Our proposal is realizable with superconducting and Rydberg atom quantum simulators.

Model.— We consider a kicked quantum spin chain of

length L described by the Hamiltonian:

$$H(t) = H_{XY} + KT \sum_{j=1}^L (j - j_0)^2 \sigma_j^z \sum_{n=1}^{N_T} \delta(t - nT), \quad (1)$$

where $H_{XY} = \sum_{j=1}^L J (\sigma_j^+ \sigma_{j+1}^- + \sigma_{j+1}^+ \sigma_j^-) + \Omega \sigma_j^x$ is the static Hamiltonian with XY interaction between nearest-neighbour spins and a longitudinal field, K and T denote the strength and period of the transverse-field kicks with a quadratic potential centered at j_0 , respectively. Here $\sigma_j^\pm = \sigma_j^x \pm i \sigma_j^y$ with $\sigma_j^{x,y,z}$ being the Pauli matrices of j -th spin, J is the interaction strength, and Ω is the strength of the longitudinal field, which leads to coherent spin flips and breaks $U(1)$ symmetry. Notably, we will emphasize in protection of quantum coherence raising from Ω via the MBDL in this kicked XY chain, which was not considered in kicked Ising chains [66–70]. To do this, we consider the high-frequency driving by setting $T = 1/16$ and $\Omega = \hbar = 1$ as the energy (frequency) unit.

For the kicked spin chain, the Floquet operator U_F is

$$U_F = e^{-iKT \sum_{j=1}^L (j-j_0)^2 \sigma_j^z} e^{-iH_{XY}T} \equiv e^{-iH_F T}, \quad (2)$$

where $H_F = \frac{i}{T} \ln U_F$ is the effective Floquet Hamiltonian. The wave function at time $t = nT$ after n kick cycles is $|\psi(t)\rangle = U_F^n |\psi(0)\rangle$. As the model is non-integrable and $U(1)$ -symmetry broken, we numerically study the static and dynamical properties using the exact diagonalization for system size $L \leq 14$ and density matrix renormalization group for larger L . Without loss of generality, we consider centrosymmetric spin chains by setting $j_0 = (L + 1)/2$ under open boundary condition.

Localization phase diagram.— We first study localization properties of many-body Floquet eigenstates $\{|\psi_\alpha^F\rangle\}$. They can be numerically obtained from the eigen equation $U_F |\psi_\alpha^F\rangle = e^{-i\epsilon_\alpha T} |\psi_\alpha^F\rangle$, where $\epsilon_\alpha \in [0, 2\pi]$ are the folded quasi-energies with $\alpha = \{1, 2, \dots, D\}$ and $D = 2^L$ as the Hilbert-space dimension. To probe localization, we compute the average inverse participation ratio (IPR) $I_F = \frac{1}{D} \sum_{\alpha=1}^D \sum_{p=1}^D |\langle p | \psi_\alpha^F \rangle|^4$, where $|p\rangle$ denote product states in the computational basis. The result of I_F in Fig. 1(a) shows the localization phase diagram of Floquet eigenstates in the K - J parameter space. For weak (no) kick and strong interaction, the Floquet system is in the delocalization phase with vanishing I_F when $L \rightarrow \infty$, which corresponds to thermalization and loss of quantum information of initial states. As K is increased, it can enter the MBDL (dynamical localization) phase with finite I_F in the interacting (noninteracting) case. In the localization regime, a proper kick strength can induce the DD of certain spins with persistent Rabi oscillation, such as two edge spins for $L = 11$ and $K/\Omega = 16\pi/25 \approx 2.01$ in Fig. 1(a), which will be discussed later. Here the local DD leads to the decrease of I_F (and O_F) with respect to K without transitions, as depicted in Fig. 1(c).

Another quantity to reveal localization in the kicked spin chain is the infinite-time staggered magnetization




FIG. 1. (Color online) Localization phase diagrams on the K - J parameter space obtained by (a) the IPR I_F and (b) the staggered magnetization O_F averaging over all Floquet eigenstates of $L = 11$. The dashed line in (a) corresponding to the values of $I_F \approx 0.009$ is plotted to guide the eye and as a rough estimation of the boundary between the delocalization and MBDL phase regions. A proper kick gives rise to the DD of certain spins. The dashed lines with triangles and circles for critical points in (b) represent the phase boundaries determined by finite-size scaling of O_F and half-chain entanglement S_F of Floquet eigenstates, respectively. (c) I_F and O_F as a function of K near the DD. (d) S_F versus L for various K . (e) \bar{r} as a function of K for various L . Other parameter is $J = 1$ in (c) and $J = 2$ in (d,e).

evolving from an initial Néel state $|\psi(0)\rangle = |\uparrow\downarrow\uparrow\downarrow\dots\rangle$, which can be calculated via the Floquet ensemble $\{|\psi_\alpha^F\rangle\}$ as $O_F = \frac{1}{L} \sum_{j=1}^L \sum_{\alpha=1}^D (-1)^j |c_\alpha|^2 \langle \psi_\alpha^F | \sigma_j^z | \psi_\alpha^F \rangle$ [1], with the coefficients $c_\alpha = \langle \psi_\alpha^F | \psi(0) \rangle$. Figure 1(b) shows the result of O_F consistent with the phase diagram in Fig. 1(a). The phase boundary can be estimated by the finite-size scaling of O_F (see Fig. 5 in the Supplementary Material (SM) [71]). We also compute the average half-chain entanglement entropy: $S_F = -\frac{1}{D} \sum_{\alpha=1}^D \text{Tr} [\rho_\alpha^F \log \rho_\alpha^F]$, where $\rho_\alpha^F = \text{Tr}_{j \leq L/2} [|\psi_\alpha^F\rangle \langle \psi_\alpha^F|]$ with even L . The finite-size scaling of S_F [71, 72] gives roughly consistent boundary the delocalization and MBDL phases in Fig. 1(b). Figure 1(c) shows the area-law entanglement in the deep MBDL phase since only the area of the bipartitioning contributes to the entanglement [73–76], which is different from the volume law in the delocalization phase.

Figure 1(e) indicates the crossover from delocalization to the MBDL by the mean gap-ratio parameter \bar{r} [23, 77]. The level-spacing statistics of Floquet eigenstates is characterized by $r_\alpha = \min\{\delta_\alpha, \delta_{\alpha+1}\} / \max\{\delta_\alpha, \delta_{\alpha+1}\}$, where $\delta_\alpha = \epsilon_{\alpha+1} - \epsilon_\alpha$ is the quasi-energy gap. We obtain \bar{r} up to $L = 14$ by averaging r_α over all Floquet eigenstates and over 20 realizations of spin chains with a center




FIG. 2. (Color online) (a) Time evolution of \mathcal{I} for typical parameters. (b) $\bar{\mathcal{I}}$ as a function of J for $L = 13, 15, 19$. (c) Phase diagram extracted by $\bar{\mathcal{I}}$. (d) Time evolution of subsystem entanglement $S_{A=L/2}$ for various parameters. (e) Saturation entanglement \bar{S} versus L . Other parameters are $L = 13$ in (a,e) and $K = 0.5$ in (b,d,e), and the initial Néel state is $|\psi(0)\rangle = |\uparrow\downarrow\uparrow\downarrow\dots\rangle$ in (a-e).

offset $j_0 \rightarrow j_0 + j_{\text{offset}}$ (to lift the level-spacing degeneracy), with j_{offset} uniformly distributed in the small region $[-0.02, 0.02]$. The MBDL phase is indicated by a Poissonian level-spacing distribution with $\bar{r} \approx 0.386$, whereas presents a Wigner-Dyson distribution with $\bar{r} \approx 0.53$ in the delocalized phase. It remains an important task to capture the delocalization-MBDL transition and criticality in the thermodynamics limit [78]. However, our results shown in Fig. 1 pinpoint the large parameter region for the MBDL phase in the kicked spin chain.

Dynamical observables.— The (de)localization can be observed via the stroboscopic dynamics of the kicked spin chain. For an initial product state in typical experiments, a direct observation is the spin-up (excitation) density distribution $P_{\uparrow}(j, t) = \langle \psi(t) | \frac{\sigma_j^z + \mathbb{I}}{2} | \psi(t) \rangle$ at the stroboscopic time $t = nT$, where \mathbb{I} is the 2×2 identity matrix. The results of $P_{\uparrow}(j, t)$ for different initial states show similar dynamics [71]: The spin chain tends to thermalize in the static case. Turning on periodical kicks up to the MBDL regime, the quantum coherence of initial states preserves without thermalization in the long-time evolution. This disorder-free non-thermalized dynamical phase is due to the kinetic constriction by the quadratic kick potential. The MBDL can be broken down and the thermalization dynamics dominates as J is increased.

To better characterize the MBDL and extract the localization phase diagram, one can prepare an initial Néel state and measure the evolution of the generalized imbalance $\mathcal{I}(t) = \frac{1}{L} \sum_j (-1)^{j+1} P_{\uparrow}(j, t)$. Typical results of $\mathcal{I}(t)$ are shown in Fig. 2(a). In the MBDL (delocalization) phase when $L \rightarrow \infty$, \mathcal{I} remains finite (becomes vanishing) after long time evolution, which is expected to tend to the value of O_F in the infinite-time limit. We can use the long-time-averaged imbalance $\bar{\mathcal{I}}$ (averaging over the

time interval from $t = 100T$ to $200T$ in simulations) to probe (de)localization and extract the phase diagram, as shown in Figs. 2(c) and 2(c), respectively.

Another observable is the dynamics of subsystem entanglement entropy $S_A(t) = -\text{Tr}[\rho_A(t) \log \rho_A(t)]$ [18, 78–80], where $\rho_A(t) = \text{Tr}_{j \leq A} [|\psi(t)\rangle \langle \psi(t)|]$ is the reduced density matrix of subsystem-A. By performing tomography, one can measure ρ_A after n kicks to obtain $S_A(t)$. Figure 2(d) shows distinct saturation behaviors of the half-chain entanglement $S_{A=L/2}$ for various L in the MBDL and delocalization phases. In finite localized (delocalized) systems, $S_{A=L/2}(t)$ approaches to saturate at late times to certain value independent (linearly dependent) of L [81–85]. The system-size dependence of the saturation entanglement \bar{S} , numerically obtained by averaging $S_{A=L/2}(t)$ over the time interval from $t = 400T$ to $600T$, is shown in Fig. 2(e). The result confirms the area- and volume-law of the half-chain entanglement in the MBDL and delocalization phases, respectively. Notably, the universal fluctuation in entanglement dynamics [85] in this kicked spin chain worth further studying.

MBDL-protected quantum information.— In the deep MBDL phase, arbitrary states in the energy spectrum are non-thermalized. Due to this strong breaking of ergodicity, the quantum information can be robustly encoded with initial MBDL states. For a generic initial state $|\psi(0)\rangle$, its effective inverse temperature β_{eff} can be obtained by solving $\text{Tr}[H_F(|\psi(0)\rangle \langle \psi(0)| - \rho_{\beta})] = 0$, where $\rho_{\beta} = e^{-\beta_{\text{eff}} H_F} / \text{Tr}(e^{-\beta_{\text{eff}} H_F})$ is the thermal density matrix [86, 87]. We plot β_{eff} as a function of quasi-energies of all product states $\epsilon_p = \langle \psi(0) | H_F | \psi(0) \rangle$ for the system of $L = 11$ in the MBDL phase and local DD case in Fig. 3(a) and 3(b), respectively. To show the protection of quantum information at high temperatures, we can consider the initial product states with $\beta_{\text{eff}} \approx 0$ near the center of the energy spectra. The non-thermalization dynamics of initial MBDL states can be observed from the long-time coherent evolution of spin density distributions, with an example shown in Fig. 3(d) (and the SM [71]). We can study additional information on the coherent dynamics beyond local observables. We compute the time evolution of fidelity $F_A(t) = \left[\text{Tr} \sqrt{\sqrt{\rho_A(t)} \rho_A(0) \sqrt{\rho_A(t)}} \right]^2$ and entanglement entropy $S_A(t)$ of subsystem-A with $A = 4$ for the high-temperature initial states in Figs. 3(e) and 3(f). The subsystem fidelity with high values and low entanglement entropy persists in the long-time evolution.

For the local DD case in Fig. 3(d), one can find the persistent Rabi oscillation in $P_{\uparrow}(j, t)$ of $j = 1, L$ for $L = 11$. In this case, the two edge spins of arbitrary initial states are effectively decoupled and behave as free spins with nearly the same Rabi oscillation, as long as the system is in the MBDL phase. Similar coherent dynamics and the DD of edge spins for other initial states are shown in Fig. 8. Note that the DD effect will be destroyed if J is increased up to the delocalization regime. Interestingly,




FIG. 3. (Color online) β_{eff} as a function of ϵ_p for (a) MBDL with $K = 1$; and (b) DD with $K = 16\pi/25$. The initial high-temperature states with $\beta_{\text{eff}} \approx 0$ are denoted in red. (c) Time evolution of F_A and S_A (with $A = 4$) for the initial state in (a). Time evolution of (d) P_j ; and (e) F_A , S_A and $\langle \sigma_1^z \rangle$ for the initial state in (b). (f) Lifetime τ of the $\langle \sigma_1^z \rangle$ -oscillation versus L for the initial vacuum ($|\downarrow\downarrow\dots\rangle$) and entangled states [71]. Other parameters are $J = 0.5$ and $L = 11$.

the revivals in the local density dynamics in Fig. 3(d), as well as $\langle \sigma_1^z(t) \rangle$ and $F_A(t)$ in Fig. 3(f) imply that the quantum information of initial states in the DD case can be well restored at high temperatures. Moreover, we estimate the lifetime τ of the oscillation of $\langle \sigma_1^z(t) \rangle$ under the DD effect [71]. The results of τ versus L for vacuum ($|\downarrow\downarrow\dots\rangle$) and entangled initial states shown in Fig. 3(d). Here the exponential increase of the lifetime with the system size indicates strong non-thermalization. Thus, the MBDL states with the DD effect provide a promising way of quantum storage robust against to thermal noises and even far away equilibrium. Notably, the long-time coherent dynamics is exhibited for generic initial MBDL states, such as entangled states shown in the SM [71]. Compared to the MBL and scar states, MBDL states in our kicked spin system do not require strong disorder average and special quantum state preparation.

Analysis of local DD.— We proceed to present an analysis of the local DD to obtain the required system parameters. For the j -th decoupled spin in the MBDL regime, its effective coupling strength is vanishing, corresponding to the local Floquet operator $U_F^{(j)} = e^{-i\tilde{K}\sigma_j^z} e^{-i\Omega T\sigma_j^x}$, where $\tilde{K} = KT(j - j_0)^2$ with m being positive integers. When $\tilde{K} = m\pi$ for certain kick strengths, the operator reduces to $U_F^{(j)} = (-1)^m e^{-i\Omega T\sigma_j^x}$. This implies the Rabi oscillation of the effectively decoupled spin with the frequency $\omega_d = \Omega T/\pi$. The analytical expression of the




FIG. 4. (Color online) (a) ω_d with respect to Ω for the DD of edge spins in the chain of $L = 11$. (b) $K_d^{(1,1)}$ with respect to L . (c) K versus N_d for $L = 21, 41, 61$. (d) Evolution of $P_j(j, t)$ in the DD case of $N_d = 4$ for $K = 16\pi/25$ and $L = 21$.

oscillation frequency is consistent of the numerical result, as shown in Fig. 4(a). The kick strength required for the DD of j -th spin is $K_d^{(j,m)} = m\pi/T(j - j_0)^2$. For the centrosymmetric chain of odd spins considered here, the local DD occurs with pairs of spins, such as $j = 1, L$ for $L = 11$ shown in Fig. 3(d) and Fig. 4. We numerically extract $K_d^{(1,1)}$ for various L from the local dynamics and find well agreement with the analytical expression $K_d^{(1,1)} = 4\pi/T(L - 1)^2$, as shown in Fig. 4(b).

To store more quantum information without thermalization, one needs more decoupled spins. Remarkably, we obtain the following relation between the kick strength and the DD spin number N_d of system size L [71]:

$$K(N_d, L) = \frac{\pi}{T \lfloor (L - 1)/N_d \rfloor^2}, \quad (3)$$

where $\lfloor \dots \rfloor$ is the floor function. The results for larger spin chain of $L = 21, 41, 61$ are shown in Fig. 4(c). One can see that the number of DD spins can be increased up to $N_d = L - 1$ by increasing the kick strength. We also numerically confirm this relation from the local dynamics of the spin chain of $L = 21$ and $N_d = 4$ in Fig. 4(d).

Discussions and conclusions.— Before concluding, we discuss experimental realization of the kicked XY quantum spin chain. The first system is the superconducting quantum simulators, consisting of 1D arrays of controllable qubits with tunable nearest-neighbour XY coupling [30, 88, 89]. One can individually and dynamically tune local potentials to realize the static and periodical kick terms along the x - and z -axis in Eq. (1), similar as those in experiments in Ref. [30]. Alternatively, one can realize the Floquet unitary U_F given by Eq. (2) in a digital way using tunable quantum gates on superconducting qubits [28, 29]. The second platform is the 1D driven Rydberg atom array [9, 90]. One can encode the effective spin-1/2 in a pair of opposite-parity Rydberg states and realize the XY spin Hamiltonian via the resonant dipole-dipole interactions [90]. The kicked potential may be produced

by applying pulsed laser beams with spatially dependent light shifts [9, 90]. Both superconducting and Rydberg quantum simulators are programmable with controllable many-body dynamics to detect the MBDL phase and the related coherence protection.

In summary, we have explored the disorder-free MBDL in the non-integrable quantum XY chain under periodical kicks. We have revealed the localization phase diagram with dynamical observables and large parameter regions for the MBDL, and uncovered the local DD effect for persistent Rabi oscillation of certain spins. We have proposed the MBDL-protected quantum information at high temperatures, and presented an analysis of the DD to obtain the required system parameters for quantum storage. The MBDL states in spin systems offer a promising platform for studying quantum thermalization and quantum information processing robust against thermal noises.

The authors acknowledge stimulating discussions with Chushun Tian, Dong-Ling Deng, Zhen-Yu Wang, and Shi-Liang Zhu. This work was supported by the National Natural Science Foundation of China (Grants No. 12174126 and No. 92365206), Innovation Program for Quantum Science and Technology (No. 2021ZD0301802), the Guangdong Basic and Applied Basic Research Foundation (Grant No. 2024B1515020018), the Science and Technology Program of Guangzhou (Grant No. 2024A04J3004), and Quantum Science Center of Guangdong-Hong Kong-Macau Great Bay Area.

* danweizhang@m.scnu.edu.cn

† zwang@hku.hk

- [1] M. Rigol, V. Dunjko, and M. Olshanii, *Nature* **452**, 854 (2008).
- [2] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, *Advances in Physics* **65**, 239 (2016).
- [3] P. W. Anderson, *Phys. Rev.* **109**, 1492 (1958).
- [4] R. Nandkishore and D. A. Huse, *Annual Review of Condensed Matter Physics* **6**, 15 (2015).
- [5] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, *Rev. Mod. Phys.* **91**, 021001 (2019).
- [6] P. Sierant, M. Lewenstein, A. Scardicchio, L. Vidmar, and J. Zakrzewski, *arXiv:2403.07111*.
- [7] H. Bernien et al., *Nature* **551**, 579 (2017).
- [8] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papić, *Nature Physics* **14**, 745 (2018).
- [9] D. Bluvstein, A. Omran, H. Levine, A. Keesling, G. Semeghini, S. Ebadi, T. T. Wang, A. A. Michailidis, N. Maskara, W. W. Ho, S. Choi, M. Serbyn, M. Greiner, V. Vuletić, and M. D. Lukin, *Science* **371**, 1355 (2021).
- [10] M. Serbyn, D. A. Abanin, and Z. Papić, *Nat. Phys.* **17**, 675 (2021).
- [11] P. Zhang, H. Dong, Y. Gao, L. Zhao, J. Hao, J.-Y. Desaulles, Q. Guo, J. Chen, J. Deng, B. Liu, W. Ren, Y. Yao, X. Zhang, S. Xu, K. Wang, F. Jin, X. Zhu, B. Zhang, H. Li, C. Song, Z. Wang, F. Liu, Z. Papić, L. Ying, H. Wang, and Y.-C. Lai, *Nat. Phys.* **19**, 120 (2022).
- [12] D. A. Huse, R. Nandkishore, V. Oganesyan, A. Pal, and S. L. Sondhi, *Phys. Rev. B* **88**, 014206 (2013).
- [13] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. Lett.* **111**, 127201 (2013).
- [14] M. Serbyn, M. Knap, S. Gopalakrishnan, Z. Papić, N. Y. Yao, C. R. Laumann, D. A. Abanin, M. D. Lukin, and E. A. Demler, *Phys. Rev. Lett.* **113**, 147204 (2014).
- [15] Y. Bahri, R. Vosk, E. Altman, and A. Vishwanath, *Nat. Commun.* **6**, 7341 (2015).
- [16] N. Y. Yao, C. R. Laumann, and A. Vishwanath, *arXiv:1508.06995*.
- [17] D.-W. Zhang, Y.-Q. Zhu, Y. X. Zhao, H. Yan, and S.-L. Zhu, *Adv. Phys.* **67**, 253 (2018).
- [18] H. Dong, J.-Y. Desaulles, Y. Gao, N. Wang, Z. Guo, J. Chen, Y. Zou, F. Jin, X. Zhu, P. Zhang, H. Li, Z. Wang, Q. Guo, J. Zhang, L. Ying, and Z. Papić, *arXiv:2312.10216*.
- [19] G. K. Naik, R. Singh, and S. K. Mishra, *Phys. Rev. A* **99**, 032321 (2019).
- [20] X. Mi and et al., *Science* **378**, 785 (2022).
- [21] H. Schmid, A.-G. Penner, K. Yang, L. Glazman, and F. von Oppen, *Phys. Rev. Lett.* **132**, 210401 (2024).
- [22] H. Yarloo, H.-C. Zhang, and A. E. B. Nielsen, *PRX Quantum* **5**, 020365 (2024).
- [23] L. D'Alessio and M. Rigol, *Phys. Rev. X* **4**, 041048 (2014).
- [24] A. Haldar and A. Das, *Annalen der Physik* **529** (2017).
- [25] R. Moessner and S. L. Sondhi, *Nat. Phys.* **13**, 424 (2017).
- [26] S. Geier, N. Thaicharoen, C. Hainaut, T. Franz, A. Salzinger, A. Tebben, D. Grimshandl, G. Zurn, and M. Weidemüller, *Science* **374**, 1149 (2021).
- [27] T. Shimasaki, M. Prichard, H. E. Kondakci, J. E. Pagett, Y. Bai, P. Dotti, A. Cao, A. R. Dardia, T.-C. Lu, T. Grover, and D. M. Weld, *Nat. Phys.* **20**, 409 (2024).
- [28] X. Mi and et. al., *Nature* **601**, 531 (2021).
- [29] X. Zhang, W. Jiang, J. Deng, K. Wang, J. Chen, P. Zhang, W. Ren, H. Dong, S. Xu, Y. Gao, F. Jin, X. Zhu, Q. Guo, H. Li, C. Song, A. V. Gorshkov, T. Iadecola, F. Liu, Z.-X. Gong, Z. Wang, D.-L. Deng, and H. Wang, *Nature* **607**, 468 (2022).
- [30] H. Dong, P. Zhang, C. B. Dag, Y. Gao, N. Wang, J. Deng, X. Zhang, J. Chen, S. Xu, K. Wang, Y. Wu, C. Zhang, F. Jin, X. Zhu, A. Zhang, Y. Zou, Z. Tan, Z. Cui, Z. Zhu, F. Shen, T. Li, J. Zhong, Z. Bao, H. Li, Z. Wang, Q. Guo, C. Song, F. Liu, A. Chan, L. Ying, and H. Wang, *arXiv:2403.16935*.
- [31] V. Tiwari, D. S. Bhakuni, and A. Sharma, *Phys. Rev. B* **109**, L161104 (2024).
- [32] S. Fishman, D. R. Grempel, and R. E. Prange, *Phys. Rev. Lett.* **49**, 509 (1982).
- [33] C. Tian, A. Kamenev, and A. Larkin, *Phys. Rev. Lett.* **93**, 124101 (2004).
- [34] C. Tian, A. Altland, and M. Garst, *Phys. Rev. Lett.* **107**, 074101 (2011).
- [35] S. Weidemann, M. Kremer, S. Longhi, and A. Szameit, *Nature Photonics* **15**, 576 (2021).
- [36] M. Santhanam, S. Paul, and J. B. Kannan, *Phys. Rep.* **956**, 1 (2022).
- [37] F. L. Moore, J. C. Robinson, C. F. Bharucha, B. Sundaram, and M. G. Raizen, *Phys. Rev. Lett.* **75**, 4598 (1995).
- [38] G. Lemarié, J. Chabé, P. Szriftgiser, J. C. Garreau, B. Grémaud, and D. Delande, *Phys. Rev. A* **80**, 043626 (2009).

- [39] G. Lemarié, H. Lignier, D. Delande, P. Szriftgiser, and J. C. Garreau, *Phys. Rev. Lett.* **105**, 090601 (2010).
- [40] I. Manai, J.-F. Clement, R. Chicireanu, C. Hainaut, J. C. Garreau, P. Szriftgiser, and D. Delande, *Phys. Rev. Lett.* **115**, 240603 (2015).
- [41] S. Adachi, M. Toda, and K. Ikeda, *Phys. Rev. Lett.* **61**, 659 (1988).
- [42] T. Boness, S. Bose, and T. S. Monteiro, *Phys. Rev. Lett.* **96**, 187201 (2006).
- [43] E. B. Rozenbaum and V. Galitski, *Phys. Rev. B* **95**, 064303 (2017).
- [44] P. Qin, A. Andreanov, H. C. Park, and S. Flach, *Scientific Reports* **7**, 41139 (2017).
- [45] S. Notarnicola, F. Iemini, D. Rossini, R. Fazio, A. Silva, and A. Russomanno, *Phys. Rev. E* **97**, 022202 (2018).
- [46] R. Chicireanu and A. Rançon, *Phys. Rev. A* **103**, 043314 (2021).
- [47] S. Paul, J. B. Kannan, and M. S. Santhanam, [arXiv:2405.06622](https://arxiv.org/abs/2405.06622).
- [48] A. C. Keser, S. Ganeshan, G. Refael, and V. Galitski, *Phys. Rev. B* **94**, 085120 (2016).
- [49] D. L. Shepelyansky, *Phys. Rev. Lett.* **70**, 1787 (1993).
- [50] G. Gligoric, J. D. Bodyfelt, and S. Flach, *EPL (Europhysics Letters)* **96**, 30004 (2011).
- [51] S. Lellouch, A. Rançon, S. De Bièvre, D. Delande, and J. C. Garreau, *Phys. Rev. A* **101**, 043624 (2020).
- [52] J. H. See Toh, K. C. McCormick, X. Tang, Y. Su, X.-W. Luo, C. Zhang, and S. Gupta, *Nat. Phys.* **18**, 1297 (2022).
- [53] A. Cao, R. Sajjad, H. Mas, E. Q. Simmons, J. L. Tanlimco, E. Nolasco-Martinez, T. Shimasaki, H. E. Kondakci, V. Galitski, and D. M. Weld, *Nat. Phys.* **18**, 1302 (2022).
- [54] C. Rylands, E. B. Rozenbaum, V. Galitski, and R. Konik, *Phys. Rev. Lett.* **124**, 155302 (2020).
- [55] V. Vuatelet and A. Rançon, *Phys. Rev. A* **104**, 043302 (2021).
- [56] V. Vuatelet and A. Rançon, *Quantum* **7**, 917 (2023).
- [57] M. Fava, R. Fazio, and A. Russomanno, *Phys. Rev. B* **101**, 064302 (2020).
- [58] Y. Guo, S. Dhar, A. Yang, Z. Chen, H. Yao, M. Horvath, L. Ying, M. Landini, and H.-C. Nägerl, [arXiv:2312.13880](https://arxiv.org/abs/2312.13880).
- [59] A. Lazarides, A. Das, and R. Moessner, *Phys. Rev. Lett.* **115**, 030402 (2015).
- [60] P. Ponte, A. Chandran, Z. Papić, and D. A. Abanin, *Ann. Phys.* **353**, 196 (2015).
- [61] P. Bordia, H. Lüschen, U. Schneider, M. Knap, and I. Bloch, *Nat. Phys.* **13**, 460 (2017).
- [62] P. Sierant, M. Lewenstein, A. Scardicchio, and J. Zakrzewski, *Phys. Rev. B* **107**, 115132 (2023).
- [63] M. Kormos, M. Collura, G. Takács, and P. Calabrese, *Nat. Phys.* **13**, 246 (2016).
- [64] S. Liu, S.-X. Zhang, C.-Y. Hsieh, S. Zhang, and H. Yao, *Phys. Rev. Lett.* **130**, 120403 (2023).
- [65] R. Verdel, G.-Y. Zhu, and M. Heyl, *Phys. Rev. Lett.* **131**, 230402 (2023).
- [66] T. Prosen, *Phys. Rev. E* **65**, 036208 (2002).
- [67] B. Bertini, P. Kos, and T. Prosen, *Phys. Rev. Lett.* **121**, 264101 (2018).
- [68] B. Bertini, P. Kos, and T. Prosen, *Phys. Rev. X* **9**, 021033 (2019).
- [69] R. Khasseh, J. Zhang, M. Heyl, and M. A. Rajabpour, *Phys. Rev. Lett.* **131**, 216701 (2023).
- [70] H. Sharma and U. T. Bhosale, *Phys. Rev. B* **109**, 014412 (2024).
- [71] See Supplementary Materials.
- [72] L.-Z. Tang, G.-Q. Zhang, L.-F. Zhang, and D.-W. Zhang, *Phys. Rev. A* **103**, 033325 (2021).
- [73] D. N. Page, *Phys. Rev. Lett.* **71**, 1291 (1993).
- [74] D. J. Luitz, N. Laflorencie, and F. Alet, *Phys. Rev. B* **91**, 081103 (2015).
- [75] J. A. Kjäll, J. H. Bardarson, and F. Pollmann, *Phys. Rev. Lett.* **113**, 107204 (2014).
- [76] A. Pal and D. A. Huse, *Phys. Rev. B* **82**, 174411 (2010).
- [77] G.-Q. Zhang, D.-W. Zhang, Z. Li, Z. D. Wang, and S.-L. Zhu, *Phys. Rev. B* **102**, 054204 (2020).
- [78] M. Rispoli, A. Lukin, R. Schittko, S. Kim, M. E. Tai, J. Léonard, and M. Greiner, *Nature* **573**, 385 (2019).
- [79] A. M. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, *Science* **353**, 794 (2016).
- [80] J. Choi, A. L. Shaw, I. S. Madjarov, X. Xie, R. Finkelstein, J. P. Covey, J. S. Cotler, D. K. Mark, H.-Y. Huang, A. Kale, H. Pichler, F. G. S. L. Brandão, S. Choi, and M. Endres, *Nature* **613**, 468 (2023).
- [81] M. Fagotti and P. Calabrese, *Phys. Rev. A* **78**, 010306 (2008).
- [82] J. H. Bardarson, F. Pollmann, and J. E. Moore, *Phys. Rev. Lett.* **109**, 017202 (2012).
- [83] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. Lett.* **110**, 260601 (2013).
- [84] O. Hart, S. Gopalakrishnan, and C. Castelnovo, *Phys. Rev. Lett.* **126**, 227202 (2021).
- [85] L.-K. Lim, C. Lou, and C. Tian, *Nat. Commun.* **15**, 1775 (2024).
- [86] M. C. Bañuls, J. I. Cirac, and M. B. Hastings, *Phys. Rev. Lett.* **106**, 050405 (2011).
- [87] F. Chen, Z.-H. Sun, M. Gong, Q. Zhu, Y.-R. Zhang, Y. Wu, Y. Ye, C. Zha, S. Li, S. Guo, H. Qian, H.-L. Huang, J. Yu, H. Deng, H. Rong, J. Lin, Y. Xu, L. Sun, C. Guo, N. Li, F. Liang, C.-Z. Peng, H. Fan, X. Zhu, and J.-W. Pan, *Phys. Rev. Lett.* **127**, 020602 (2021).
- [88] X. Li, H. Xu, J. Wang, L.-Z. Tang, D.-W. Zhang, C. Yang, T. Su, C. Wang, Z. Mi, W. Sun, X. Liang, M. Chen, C. Li, Y. Zhang, K. Linghu, J. Han, W. Liu, Y. Feng, P. Liu, G. Xue, J. Zhang, S. P. Zhao, Y. Jin, S.-L. Zhu, H. Yu, and Q.-K. Xue, [arXiv:2301.12138](https://arxiv.org/abs/2301.12138).
- [89] X. Tan, D.-W. Zhang, W. Zheng, X. Yang, S. Song, Z. Han, Y. Dong, Z. Wang, D. Lan, H. Yan, S.-L. Zhu, and Y. Yu, *Phys. Rev. Lett.* **126**, 017702 (2021).
- [90] C. Chen, G. Bornet, M. Bintz, G. Emperauger, L. Leclerc, V. S. Liu, P. Scholl, D. Barredo, J. Hauschild, S. Chatterjee, M. Schuler, A. M. Läuchli, M. P. Zaletel, T. Lahaye, N. Y. Yao, and A. Browaeys, *Nature* **616**, 691 (2023).

SUPPLEMENTAL MATERIALS

A. Finite-size scaling to obtain phase boundaries

We numerically determine the phase boundary between the MBDL and delocalization phase through the finite-size scaling analysis of infinite-time staggered magnetization O_F and half-chain entanglement entropy S_F defined in the main text. Figures 5(a1-e1) show the result of O_F versus the kick strength K for various system sizes L and interaction strengths J . For the case of $J = 0.5$ in Fig. 5(a1), O_F saturates to small finite value at weak kick strength, which indicates the thermalization of the systems. As the kick strength is increased, the curves of O_F is separated around $K = K_c$ and increases with L . The curves can collapse well by considering O_F as a function of $(K - K_c)L^{1/\nu}$ with the critical exponent $\nu \approx 0.70$ and $K_c \approx 0.2$, as shown in Fig. 5(a2), which indicates a delocalization-MBDL transition when approaching to the thermodynamic limit. Similar finite-size scaling analysis are taken for $J = \{1, 1.5, 2.0, 2.5\}$ in Figs. 5(b1-e1), which give the critical points at $K_c \approx \{0.38, 0.52, 0.62, 0.75\}$ in Figs. 5(b2-e2).

Half-chain entanglement entropy is a complementary quantity to distinguish thermal phase and the MBL phase [73–76]. For fully thermal Floquet states in our system, the entanglement entropy is given by the Page value $S_F^{\text{Page}} = (L \ln 2 - 1)/2 + O(L)$ [73], which indicates that the value of entanglement entropy increases linearly with system size L , i.e., the volume law. On the other hand, these eigenstates in MBL systems are partly entangled, hence the entanglement entropy $S_F \sim O(L^0)$ obeys the area law. Figures 6(a1-e1) show the site-averaged half-chain entanglement entropy S_F/L of Floquet eigenstates of U_F as functions of the kick strength K for different system sizes L and J . For $J = 0.5$ in Fig. 6(a1), S_F/L obeys the volume law at weak kick strength, which indicates the thermalization in the system. As K is increased, the curves of S_F/L for different L cross together at $K = K_c$, which indicates the delocalization-MBDL transition. To determine the MBDL transition point K_c , we collapse these curves in Fig. 6(a2) by considering S_F/L as a function of $(K - K_c)L^{1/\nu}$ with $K_c \approx 0.28$. For other values of $J = \{1, 1.5, 2.0, 2.5\}$ in Figs. 6(b1-e1), the corresponding critical points can be determined as $K_c \approx \{0.42, 0.51, 0.60, 0.70\}$ in Figs. 6(b2-e2), respectively. The result of MBDL transition point $\{K_c\}$ are in consistent with ones obtained from O_F , while the critical exponent $\nu \approx 0.65$ in Figs. 6(a2-e2) is slightly different with that in Figs. 5(a2-e2).

B. Dynamics of various initial states

The (de)localization of the many-body Floquet eigenstates $\{|\psi_\alpha^F\rangle\}$ can be observed through dynamics of

spin-up (excitation) density distribution $P_\uparrow(j, t)$ defined in main text. The results of $P_\uparrow(j, t)$ as functions of t/T for different initial states with different parameters $(K/\Omega, J/\Omega)$ are showed in Fig. 7. For Néel state $|\psi(0)\rangle = |\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\rangle$ with $(K/\Omega, J/\Omega) = (0, 0.5)$ in Fig. 7(a), Hamiltonian Eq. (1) reduces to H_{XY} . In this case, the system tends to thermalize with losing embedded quantum information [the upper panel]. Turning on kick strength up to MBDL regime with $K = J = 0.5$, the system turns to preserve the quantum coherence of initial states without thermalization in a long-time evolution up to $t = 200T$ [the middle panel]. This disorder-free non-thermalized dynamical phase comes from dynamical constraints induced by quadratic kick potential. Noted that the system slightly thermalize around the central spin, which is due to the zero kick strength on spin $j = 7$. Further increasing the interaction strength up to $J = 2.0$, the systems reenter the delocalized regime of phase diagram, which leads to delocalization dynamics of spin chains [the lower panel]. Similar phenomenons can be observed for other typical initial states in Fig. 1(b, c), which indicates the independence of the choice of initial states for observing MBDL and delocalization dynamics.

The edge spins of the spin chains exhibit persistent Rabi oscillation in $P_\uparrow(j, t)$ for specific value of kick strength. Such phenomenon is induced by DD of edge spins in MBDL regime, which is independent of the choice of initial state. Here we consider systems size $L = 11$ and kick strength $K = K_d^{(1,1)}$ to enable the DD of edge spins. The coherent dynamics and DD of edge spins of Néel state $|\psi(0)\rangle = |\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\rangle$ for $J = 0, 1$ are shown in upper and middle panels of Fig. 8(a), respectively. One can see the Rabi oscillation of $P_\uparrow(1, t)$ and $P_\uparrow(11, t)$ independent of interaction strength $J = 0, 1$. By applying the Fourier transformation of $P_\uparrow(1, t)$ for varying $J = 0, 1$, we obtain the corresponding Fourier spectrum in Fig. 8(a) [lower panel]. The Rabi-like oscillation frequency of $P_\uparrow(1, t)$ for $J = 1$ are the same as that for free spins with $J = 0$. This indicates that the spins of $j = 1, L$ are effectively decoupled and behavior as free spins with Rabi oscillation. Noted that Rabi oscillation also exhibits in central spin $j = 6$ for decoupled limit $J = 0$, which breaks up when interaction strength dominates the dynamics around central spins with $J = 1$. DD of other initial state can also be observed in Fig. 8(b-e).

The persistent coherent dynamics protected by the disorder-free MBDL is exhibited for generic initial states. We compute the time evolution of fidelity $F_A(t)$ and entanglement entropy $S_A(t)$ of the high-temperature Bell states and entangled states in Figs. 9(a,b) and (c,d), respectively. The parameters are $(K/\Omega, J/\Omega) = (1.0, 0.5)$ and $(16\pi/25, 1.0)$ in (a,c) and (b,d), respectively. The subsystem fidelity dynamics with low entanglement entropy persists in the long-time evolution up to $t = 500T$. Noted that $F_A(t)$ oscillates in (b,d) due to DD of edge spins.




FIG. 5. (Color online) Finite-size scaling of O_F to determine the critical points K_c . The lower column shows O_F after the data collapse for $J = \{0.5, 1.0, 1.5, 2.0, 2.5\}$ with $K_c \approx \{0.2, 0.38, 0.52, 0.62, 0.75\}$ and critical exponent $\nu \approx 0.70$ in (a-e), respectively.




FIG. 6. (Color online) Finite-size scaling of S_F to determine the critical points K_c . The lower column shows S_F after the data collapse for $J = \{0.5, 1.0, 1.5, 2.0, 2.5\}$ with $K_c \approx \{0.28, 0.42, 0.51, 0.60, 0.70\}$ and critical exponent $\nu \approx 0.65$ in (a-e), respectively.

We can also estimate the lifetime τ of the persist Rabi-like oscillation of edge spins from the long-time evolution of $\langle \sigma_1^z \rangle$ under the DD effect. For instance, we show the time evolution of $\langle \sigma_1^z \rangle$ in Figs. 10(a,b,c) starting from the vacuum states $|\psi(0)\rangle = |\downarrow\downarrow\cdots\rangle$ and (d,e,f) for entangled states $|\psi(0)\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle_4|\downarrow\rangle_5 + |\downarrow\rangle_4|\uparrow\rangle_5) \otimes \prod_{j \neq 4,5} |\downarrow\rangle_j$, for system sizes $L = 7, 8, 11$. The parameters $(K/\Omega, J/\Omega) = (K_d^{(1,1)}, 1)$ are set to enable the DD of edge spins for various L . The upper envelop of $\langle \sigma_1^z \rangle$ denoted by red dashed lines decrease to local minimum with increasing time. The lifetime are roughly determined by the first minimum point of the envelop. The obtained values of lifetime τ/T are plotted in the figure.

C. Derivation of the relation between K and N_d

There may exist more than two DD spins fulfill the condition $K_d^{(j,m)}$ at the same time. For example,

$$\begin{aligned} K_d^{(j_0+1,1)} &= K_d^{(j_0+2,4)} = \dots = K_d^{(j_0+j_1, j_1^2)} = \frac{\pi}{T}, \\ K_d^{(j_0+2,1)} &= K_d^{(j_0+4,4)} = \dots = K_d^{(j_0+j_2, j_2^2/4)} = \frac{\pi}{4T}, \end{aligned} \quad (4)$$

where $j_1 = (L-1)/2$ and $j_2 = 2\lfloor(L-1)/4\rfloor$. Noted that the total number of DD spins N_d depends on the number of the total terms in Eq. (4), one can obtain

$$\begin{aligned} N_d^{(1)} &= L-1, \\ N_d^{(2)} &= 2\lfloor(L-1)/4\rfloor, \end{aligned} \quad (5)$$

respectively. The general formalization of Eq. (4) is

$$K_d^{(j_0+l,1)} = K_d^{(j_0+2l,4)} = \dots = K_d^{(j_0+j_l, j_l^2/l^2)} = \frac{\pi}{l^2 T}, \quad (6)$$

where $l \leq (L-1)/2$, $j_l = l\lfloor(L-1)/(2l)\rfloor$, hence

$$N_d^{(l)} = 2j_l/l = 2\lfloor(L-1)/(2l)\rfloor. \quad (7)$$

By iterating l from 1 to $(L-1)/2$, one can obtain all the values of $\{N_d^{(l)}\}$ with corresponding kick strength $K = \pi/l^2 T$, which leads to a relation between N_d and K

$$K = \frac{\pi}{T[(L-1)/N_d]^2}. \quad (8)$$




FIG. 9. (Color online) Dynamics of the subsystem fidelity F_A and entanglement entropy S_A from (a,b) high-temperature Bell state $|\psi(0)\rangle = \frac{1}{\sqrt{2}}(|\downarrow\uparrow\uparrow\uparrow\uparrow\uparrow\downarrow\downarrow\downarrow\downarrow\rangle + |\uparrow\downarrow\downarrow\downarrow\downarrow\downarrow\uparrow\uparrow\uparrow\uparrow\rangle)$; and from (c,d) entangled state $|\psi(0)\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle_4 |\downarrow\rangle_5 + |\downarrow\rangle_4 |\uparrow\rangle_5) \otimes \prod_{j \neq 4,5} |\downarrow\rangle_j$. The system size is $L = 11$ and the parameters are $(K/\Omega, J/\Omega) = (1.0, 0.5)$ in (a,c) and $(16\pi/25, 1.0)$ in (b,d), respectively.




FIG. 10. (Color online) Dynamics of $\langle \sigma_1^z \rangle$ starting from (a1,a2,a3) vacuum states $|\psi(0)\rangle = |\downarrow\downarrow \dots\rangle$; and from (b1,b2,b3) entangled states $|\psi(0)\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle_4 |\downarrow\rangle_5 + |\downarrow\rangle_4 |\uparrow\rangle_5) \otimes \prod_{j \neq 4,5} |\downarrow\rangle_j$ under the local DD effect. Red dashed lines denote upper envelop of $\langle \sigma_1^z \rangle$. Lifetime τ/T are roughly determined by the first minimum point of the envelop, with the values being plotted in the figure. The system sizes are (a1,b1) $L = 7$, (a2,b2) $L = 9$, and (a3,b3) $L = 11$, and the parameters are $(K/\Omega, J/\Omega) = (K_d^{(1,1)}, 1)$ for ensuring the DD of edge spins.