

# DISTRIBUTION AND CONGRUENCES OF $(u, v)$ -REGULAR BIPARTITIONS

NABIN KUMAR MEHER

ABSTRACT. Let  $B_{u,v}(n)$  denote the number of  $(u, v)$ -regular bipartitions of  $n$ . In this article, we prove that  $B_{p,m}(n)$  is always almost divisible by  $p$ , where  $p \geq 5$  is a prime number and  $m = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_r^{\alpha_r}$ , where  $\alpha_i \geq 0$  and  $p_i \geq 5$  be distinct primes with  $\gcd(p, m) = 1$ . Further, we obtain an infinities families of congruences modulo 3 for  $B_{3,7}(n)$ ,  $B_{3,5}(n)$  and  $B_{3,2}(n)$  by using Hecke eigenform theory and a result of Newman [15]. Furthermore, we get many infinite families of congruences modulo 7, 11 and 13 respectively for  $B_{2,7}(n)$ ,  $B_{2,11}(n)$  and  $B_{2,13}(n)$ , by employing an identity of Newman [15]. In addition, we prove infinite families of congruences modulo 2 for  $B_{4,3}(n)$ ,  $B_{8,3}(n)$  and  $B_{4,5}(n)$  by applying another result of Newman [16].

## 1. INTRODUCTION

Let  $t$  and  $n$  be positive integers. A partition of  $n$  is called  $t$ -regular if none of its parts is divisible by  $t$ . Let  $b_t(n)$  denotes the number of  $t$ -regular partitions of  $n$ . The generating function for  $b_t(n)$  is given by

$$\sum_{n=0}^{\infty} b_t(n)q^n = \frac{f_t}{f_1},$$

where  $f_t$  is defined by  $f_t = \prod_{m=1}^{\infty} (1 - q^{tm})$ . Gordan and Ono[4] proved a density result on  $t$ -regular partitions. In particular, they proved that if  $p$  is a prime number and  $p^{\text{ord}_p(t)} \geq \sqrt{t}$ , then for any positive integer  $j$ , the arithmetic density of positive integers  $n$  such that  $b_t(n) \equiv 0 \pmod{p^j}$  is one.

Note that a bipartition  $(\alpha, \beta)$  of  $n$  is a pair of partitions  $(\alpha, \beta)$  such that the sum of all of the parts is  $n$ . For any positive integers  $u \geq 2, v \geq 2$ , an  $(u, v)$ -regular bipartition of  $n$  is a bipartition  $(\alpha, \beta)$  of  $n$  such that  $\alpha$  is a  $u$ -regular partition and  $\beta$  is a  $v$ -regular partition. Let  $B_{u,v}(n)$  denote the number of  $(u, v)$ -regular bipartitions of  $n$ .

---

2010 Mathematics Subject Classification: Primary 05A17, 11P83, Secondary 11F11  
 Keywords:  $(u, v)$ -regular bipartitions; Eta-quotients; modular forms; Hecke eigenform; Newman Identity; arithmetic density

Conventionally, assume that  $B_{u,v}(0) = 1$ . The generating function of  $B_{u,v}(n)$  is given by

$$\sum_{n=0}^{\infty} B_{u,v}(n)q^n = \frac{f_u f_v}{f_1^2}. \quad (1.1)$$

In order to state the following results, we introduce the Legendre symbol. Let  $p \geq 3$  be a prime and  $a$  be an integer. The Legendre symbol  $\left(\frac{a}{p}\right)_L$  is defined as

$$\left(\frac{a}{p}\right)_L := \begin{cases} 1, & \text{if } a \text{ is a quadratic residue modulo } p \text{ and } p \nmid a, \\ -1, & \text{if } a \text{ is a nonquadratic residue modulo } p, \\ 0, & \text{if } p|a. \end{cases}$$

In 2017, Xia and Yao [24] studied  $(k, \ell)$ -regular bipartitions and proved that for a positive integer  $s$  and a prime  $p \geq 5$  with  $\left(\frac{-s}{p}\right) = -1$ , we have

$$B_{3,s} \left( p^{2\alpha+1}n + \frac{(1+s)(p^{2\alpha+2} - 1)}{24} \right) \equiv 0 \pmod{3} \quad (1.2)$$

whenever  $n, \alpha \geq 0$  with  $p \nmid n$ . Motivated by the above results, we show that  $B_{p,m}(n)$  is almost always divisible by a prime number  $p \geq 5$ . In particular, we prove the following.

**Theorem 1.1.** *Let  $p \geq 5$  be a prime number. Let  $m = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_r^{\alpha_r}$ , where  $\alpha_i \geq 0$  be non-negative integers and  $p_i \geq 5$  be distinct primes with  $\gcd(p, m) = 1$ . Then the set*

$$\{n \in \mathbb{N} : B_{p,m}(n) \equiv 0 \pmod{p}\}$$

*has arithmetic density 1.*

In 2016, Dou [3] found that, for  $n \geq 0$  and  $\alpha \geq 2$ ,

$$B_{3,11} \left( 3^\alpha n + \frac{5 \cdot 3^{\alpha-1} - 1}{2} \right) \equiv 0 \pmod{11}. \quad (1.3)$$

Further, in 2017, L. Wang [22] studied the arithmetic properties of  $B_{3,\ell}(n)$  and  $B_{5,\ell}(n)$ . In particular, he proved that for any non-negative integer  $n$ , we have

$$B_{3,7} \left( 2^{2\alpha} n + \frac{5 \cdot 2^{2\alpha-1} - 1}{3} \right) \equiv 0 \pmod{3} \quad \text{for all } \alpha \geq 1, \quad (1.4)$$

$$B_{3,7}(4n+1) \equiv -B_{3,7}(n) \pmod{3}, \quad (1.5)$$

and

$$B_{3,7}(4n+3) \equiv 0 \pmod{3}. \quad (1.6)$$

In 2019, T. Kathiravan and K. Srilakshmi [8] proved infinite families of congruences modulo 5 for  $B_{2,15}(n)$ , modulo 11 for  $B_{7,11}(n)$  and modulo 17 for  $B_{243,17}(n)$ . Motivated from the above result, by using Hecke eigenform and modular form theory, we obtain the following congruences modulo 3 for  $B_{3,7}(n)$ ,  $B_{3,5}(n)$  and  $B_{3,2}(n)$ . In particular, it says the following.

**Theorem 1.2.** *Let  $k$  and  $n$  be non-negative integers. Then we have*

- (i)  $B_{3,7} \left( p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 n + \frac{p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 (3j + p_{k+1}) - 1}{3} \right) \equiv 0 \pmod{3}$ , where  $p_i$  is a prime number such that  $p_i \not\equiv 1 \pmod{3}$  for  $i \in \{1, 2, \dots, k+1\}$ , and  $p_{k+1} \nmid j$ .
- (ii)  $B_{3,5} \left( p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 n + \frac{p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 (4j + p_{k+1}) - 1}{4} \right) \equiv 0 \pmod{3}$ , where  $p_i$  is a prime number such that  $p_i \equiv 3 \pmod{4}$  for  $i \in \{1, 2, \dots, k+1\}$ , and  $p_{k+1} \nmid j$ .
- (iii)  $B_{3,2} \left( p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 n + \frac{p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 (8j + p_{k+1}) - 1}{8} \right) \equiv 0 \pmod{3}$ , where  $p_i$  is a prime number such that  $p_i \not\equiv 1 \pmod{8}$  for  $i \in \{1, 2, \dots, k+1\}$ , and  $p_{k+1} \nmid j$ .

If we substitute  $p_1 = p_2 = \cdots = p_{k+1} = p$  in Theorem 1.2, then we obtain the following result.

**Corollary 1.3.** *Let  $k$  and  $n$  be non-negative integers. Then we have*

- (i)  $B_{3,7} \left( p^{2k+2} n + p^{2k+1} j + \frac{p^{2k+2} - 1}{3} \right) \equiv 0 \pmod{3}$ , where  $p$  is an odd prime such that  $p \equiv 2 \pmod{3}$ , and  $p \nmid j$ .
- (ii)  $B_{3,5} \left( p^{2k+2} n + p^{2k+1} j + \frac{p^{2k+2} - 1}{4} \right) \equiv 0 \pmod{3}$ , where  $p$  is an odd prime such that  $p \equiv 3 \pmod{4}$ , and  $p \nmid j$ .
- (iii)  $B_{3,2} \left( p^{2k+2} n + p^{2k+1} j + \frac{p^{2k+2} - 1}{8} \right) \equiv 0 \pmod{3}$ , where  $p$  is an odd prime such that  $p \not\equiv 1 \pmod{8}$ , and  $p \nmid j$ .

**Remark 1.1.** (i) *If we substitute  $p = 2, k = 0$  and  $j = 1$  in 1.3(i), we obtain (1.6).*  
(ii) *If we put  $p = 2, k = \alpha - 1$  and  $j = 1$  in 1.3(i), we obtain (1.4).*

*In Particular, our results generalize some of the result obtained by Wang [22].*

Further, by employing a result of Newman [15], we obtain another infinite families of congruences modulo 3 for  $B_{3,7}(n), B_{3,5}(n)$  and  $B_{3,2}(n)$ .

**Theorem 1.4.** *Let  $\alpha$  be a non-negative integer.*

- (i) *Let  $p$  be a prime number with  $p \equiv 1 \pmod{6}$ . If  $p \nmid (3n + 1)$  and  $B_{3,7} \left( \frac{p-1}{3} \right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ ,*

$$B_{3,7} \left( p^{2\alpha+1} n + \frac{p^{2\alpha+1} - 1}{3} \right) \equiv 0 \pmod{3}. \quad (1.7)$$

- (ii) *Let  $p$  be a prime number with  $p \equiv 1 \pmod{4}$ . If  $p \nmid (4n + 1)$  and  $B_{3,5} \left( \frac{p-1}{4} \right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ , and*

$$B_{3,5} \left( p^{2\alpha+1} n + \frac{p^{2\alpha+1} - 1}{4} \right) \equiv 0 \pmod{3}. \quad (1.8)$$

- (iii) Let  $p$  be a prime number with  $p \equiv 1 \pmod{8}$ . If  $p \nmid (8n + 1)$  and  $B_{3,2} \left( \frac{p-1}{8} \right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ ,

$$B_{3,2} \left( p^{2\alpha+1}n + \frac{p^{2\alpha+1} - 1}{8} \right) \equiv 0 \pmod{3}. \quad (1.9)$$

Furthermore, we prove the following multiplicative formulae for  $B_{3,7}(n)$ ,  $B_{3,5}(n)$  and  $B_{3,2}(n)$  modulo 3.

**Theorem 1.5.** *Let  $k$  and  $n$  be positive integers.*

- (i) Let  $p$  be a prime number such that  $p \equiv 2 \pmod{3}$ . Let  $r$  be a non-negative integer such that  $p$  divides  $3r + 2$ , then

$$B_{3,7} \left( p^{k+1}n + pr + \frac{2p-1}{3} \right) \equiv - \left( \frac{-7}{p} \right)_L B_{3,7} \left( p^{k-1}n + \frac{3r+2-p}{3p} \right) \pmod{3}.$$

- (ii) Let  $p$  be a prime number such that  $p \equiv 3 \pmod{4}$ . Let  $r$  be a non-negative integer such that  $p$  divides  $4r + 3$ , then

$$B_{3,5} \left( p^{k+1}n + pr + \frac{3p-1}{4} \right) \equiv - \left( \frac{-5}{p} \right)_L B_{3,5} \left( p^{k-1}n + \frac{4r+3-p}{4p} \right) \pmod{3}.$$

- (iii) Let  $t \in \{3, 5, 7\}$ . Let  $p$  be a prime number such that  $p \equiv t \pmod{8}$ . Let  $r$  be a non-negative integer such that  $p$  divides  $8r + t$ , then

$$B_{3,2} \left( p^{k+1}n + pr + \frac{tp-1}{8} \right) \equiv - \left( \frac{-2}{p} \right)_L B_{3,2} \left( p^{k-1}n + \frac{8r+t-p}{8p} \right) \pmod{3}.$$

**Corollary 1.6.** *Let  $k$  and  $n$  be positive integers.*

- (i) Let  $p$  be a prime number such that  $p \equiv 2 \pmod{3}$ . Then

$$B_{3,7} \left( p^{2k}n + \frac{p^{2k}-1}{3} \right) \equiv (-1)^k \left( \frac{-7}{p} \right)_L^k B_{3,7}(n) \pmod{3}.$$

- (ii) Let  $p$  be a prime number such that  $p \equiv 3 \pmod{4}$ . Then

$$B_{3,5} \left( p^{2k}n + \frac{p^{2k}-1}{4} \right) \equiv (-1)^k \left( \frac{-5}{p} \right)_L^k B_{3,5}(n) \pmod{3}.$$

- (iii) Let  $t \in \{3, 5, 7\}$ . Let  $p$  be a prime number such that  $p \equiv t \pmod{8}$ . Then

$$B_{3,2} \left( p^{2k}n + \frac{p^{2k}-1}{4} \right) \equiv (-1)^k \left( \frac{-2}{p} \right)_L^k B_{3,2}(n) \pmod{3}.$$

**Remark 1.2.** *In particular, if we put  $p = 2, k = 1$  in the above corollary 1.6(i), we get (1.5).*

Next, by applying an identity due to Newman [15], we obtain the following congruences modulo 7, 11 and 13 respectively.

**Theorem 1.7.** (i) *Let  $p$  be a prime number with  $p \equiv 1 \pmod{24}$ . If  $p \nmid (24n + 7)$  and  $B_{7,2} \left( \frac{7(p-1)}{24} \right) \equiv 0 \pmod{7}$ , then for  $n \geq 0$ , and  $\alpha \geq 0$ ,*

$$B_{7,2} \left( p^{2\alpha+1}n + \frac{7(p^{2\alpha+1} - 1)}{24} \right) \equiv 0 \pmod{7}. \quad (1.10)$$

(ii) *Let  $p$  be a prime number with  $p \equiv 1 \pmod{24}$ . If  $p \nmid (24n+1)$  and  $B_{11,2} \left( \frac{11(p-1)}{24} \right) \equiv 0 \pmod{11}$ , then for  $n \geq 0$ , and  $\alpha \geq 0$ ,*

$$B_{11,2} \left( p^{2\alpha+1}n + \frac{11(p^{2\alpha+1} - 1)}{24} \right) \equiv 0 \pmod{11}. \quad (1.11)$$

(iii) *Let  $p$  be a prime number with  $p \equiv 1 \pmod{24}$ . If  $p \nmid (24n+13)$  and  $B_{13,2} \left( \frac{13(p-1)}{24} \right) \equiv 0 \pmod{13}$ , then for  $n \geq 0$ , and  $\alpha \geq 0$ ,*

$$B_{13,2} \left( p^{2\alpha+1}n + \frac{13(p^{2\alpha+1} - 1)}{24} \right) \equiv 0 \pmod{13}. \quad (1.12)$$

In addition, by employing another result of Newman [16], we obtain the following congruences of modulo 2 for  $B_{4,3}(n)$ .

**Theorem 1.8.** *Let  $a_1(n)$  be defined by*

$$\sum_{n=0}^{\infty} a_1(n)q^n := (q; q)_{\infty}^2 (q^3; q^3)_{\infty} \quad (1.13)$$

and let  $p \geq 5$  be a prime number. Define

$$w_1(p) := a_1 \left( \frac{5(p^2 - 1)}{24} \right) + \left( \frac{-6}{p} \right)_L \left( \frac{-5(p^2 - 1)}{24p} \right)_L \quad (1.14)$$

(i) *If  $w_1(p) \equiv 0 \pmod{2}$ , then for  $n, k \geq 0$ , if  $p \nmid n$ , we have*

$$B_{4,3} \left( p^{4k+3}n + \frac{5(p^{4k+4} - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (1.15)$$

(ii) *If  $w_1(p) \not\equiv 0 \pmod{2}$ , then for  $n, k \geq 0$ , if  $p \nmid n$ , we have*

$$B_{4,3} \left( p^{6k+5}n + \frac{5(p^{6k+6} - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (1.16)$$

(iii) If  $w_1(p) \not\equiv 0 \pmod{2}$ , then for  $n, k \geq 0$ , with  $w_1(p) \equiv \left(\frac{-6n-1+\frac{p^2-1}{4}}{p}\right)_L \pmod{2}$ , we have

$$B_{4,3} \left( p^{6k+2}n + \frac{5(p^{6k+2}-1)}{24} \right) \equiv 0 \pmod{2}. \quad (1.17)$$

**Remark 1.3.** (i) Note that (1.13) was considered in [Theorem 1.5, [23]] and [Theorem 1.1, [9]]. They derived infinite families of congruences modulo 3 and 2 respectively.

(ii) If we put  $p = 7$ , (1.14) we obtain that  $a_1(10) = 0$  and  $w_1(7) = 1$ . Note that  $w_1(7) \equiv \left(\frac{11-6n}{7}\right)_L \equiv \left(\frac{n-3}{7}\right)_L \equiv 1 \pmod{2}$  when  $n \not\equiv 3 \pmod{7}$ . Thus putting  $k = 0$ , in (1.4), we get

$$B_{4,3}(343n + 49r + 10) \equiv 0 \pmod{2},$$

where  $r \in \{0, 1, 2, 4, 5, 6\}$

Similarly, by employing the same result of Newman [16], we get another infinite families of congruences modulo 2 for  $B_{8,3}(n)$ , and  $B_{4,5}(n)$ .

**Theorem 1.9.** Let  $a_2(n)$  be defined by

$$\sum_{n=0}^{\infty} a_2(n)q^n := (q; q)_{\infty}^6 (q^3; q^3)_{\infty} \quad (1.18)$$

and let  $p \geq 5$  be a prime number. Define

$$w_2(p) := a_2 \left( \frac{3(p^2-1)}{8} \right) + p^2 \cdot \left( \frac{-6}{p} \right)_L \left( \frac{-3(p^2-1)}{8p} \right)_L \quad (1.19)$$

(i) If  $w_2(p) \equiv 0 \pmod{2}$ , then for  $n, k \geq 0$ , if  $p \nmid n$ , we have

$$B_{8,3} \left( p^{4k+3}n + \frac{3(p^{4k+4}-1)}{8} \right) \equiv 0 \pmod{2}. \quad (1.20)$$

(ii) If  $w_2(p) \not\equiv 0 \pmod{2}$ , then for  $n, k \geq 0$ , if  $p \nmid n$ , we have

$$B_{8,3} \left( p^{6k+5}n + \frac{3(p^{6k+6}-1)}{8} \right) \equiv 0 \pmod{2}. \quad (1.21)$$

(iii) If  $w_2(p) \not\equiv 0 \pmod{2}$ , then for  $n, k \geq 0$ , with  $w_2(p) \equiv p^2 \cdot \left(\frac{-6n-2+\frac{p^2-1}{4}}{p}\right)_L \pmod{2}$ , we have

$$B_{8,3} \left( p^{6k+2}n + \frac{3(p^{6k+2}-1)}{8} \right) \equiv 0 \pmod{2}. \quad (1.22)$$

**Theorem 1.10.** *Let  $a_3(n)$  be defined by*

$$\sum_{n=0}^{\infty} a_3(n)q^n := (q; q)_{\infty}^2 (q^5; q^5)_{\infty} \quad (1.23)$$

and let  $p \geq 5$  be a prime number. Define

$$w_3(p) := a_3\left(\frac{7(p^2-1)}{24}\right) + \left(\frac{-10}{p}\right)_L \left(\frac{-7(p^2-1)}{24p}\right)_L \quad (1.24)$$

(i) *If  $w_3(p) \equiv 0 \pmod{2}$ , then for  $n, k \geq 0$ , if  $p \nmid n$ , we have*

$$B_{4,5}\left(p^{4k+3}n + \frac{7(p^{4k+4}-1)}{24}\right) \equiv 0 \pmod{2}. \quad (1.25)$$

(ii) *If  $w_3(p) \not\equiv 0 \pmod{2}$ , then for  $n, k \geq 0$ , if  $p \nmid n$ , we have*

$$B_{4,5}\left(p^{6k+5}n + \frac{7(p^{6k+6}-1)}{24}\right) \equiv 0 \pmod{2}. \quad (1.26)$$

(iii) *If  $w_3(p) \not\equiv 0 \pmod{2}$ , then for  $n, k \geq 0$ , with  $w_3(p) \equiv \left(\frac{-10n-2+\frac{11(p^2-1)}{12}}{p}\right)_L \pmod{2}$ , we have*

$$B_{4,5}\left(p^{6k+2}n + \frac{7(p^{6k+2}-1)}{24}\right) \equiv 0 \pmod{2}. \quad (1.27)$$

## 2. PRELIMINARIES

We recall some basic facts and definition on modular forms. For more details, one can see [10], [17]. We start with some matrix groups. We define

$$\Gamma := \mathrm{SL}_2(\mathbb{Z}) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{Z}, ad - bc = 1 \right\},$$

$$\Gamma_{\infty} := \left\{ \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix} : n \in \mathbb{Z} \right\}.$$

For a positive integer  $N$ , we define

$$\Gamma_0(N) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) : c \equiv 0 \pmod{N} \right\},$$

$$\Gamma_1(N) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_0(N) : a \equiv d \equiv 1 \pmod{N} \right\}$$

and

$$\Gamma(N) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) : a \equiv d \equiv 1 \pmod{N}, b \equiv c \equiv 0 \pmod{N} \right\}.$$

A subgroup of  $\Gamma = \mathrm{SL}_2(\mathbb{Z})$  is called a congruence subgroup if it contains  $\Gamma(N)$  for some  $N$  and the smallest  $N$  with this property is called its level. Note that  $\Gamma_0(N)$  and  $\Gamma_1(N)$  are congruence subgroups of level  $N$ , whereas  $\mathrm{SL}_2(\mathbb{Z})$  and  $\Gamma_\infty$  are congruence subgroups of level 1. The index of  $\Gamma_0(N)$  in  $\Gamma$  is

$$[\Gamma : \Gamma_0(N)] = N \prod_{p|N} \left(1 + \frac{1}{p}\right)$$

where  $p$  runs over the prime divisors of  $N$ .

Let  $\mathbb{H}$  denote the upper half of the complex plane  $\mathbb{C}$ . The group

$$\mathrm{GL}_2^+(\mathbb{R}) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{R}, ad - bc > 0 \right\},$$

acts on  $\mathbb{H}$  by  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} z = \frac{az+b}{cz+d}$ . We identify  $\infty$  with  $\frac{1}{0}$  and define  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \frac{r}{s} = \frac{ar+bs}{cr+ds}$ , where  $\frac{r}{s} \in \mathbb{Q} \cup \{\infty\}$ . This gives an action of  $\mathrm{GL}_2^+(\mathbb{R})$  on the extended half plane  $\mathbb{H}^* = \mathbb{H} \cup \mathbb{Q} \cup \{\infty\}$ . Suppose that  $\Gamma$  is a congruence subgroup of  $\mathrm{SL}_2(\mathbb{Z})$ . A cusp of  $\Gamma$  is an equivalence class in  $\mathbb{P}^1 = \mathbb{Q} \cup \{\infty\}$  under the action of  $\Gamma$ .

The group  $\mathrm{GL}_2^+(\mathbb{R})$  also acts on functions  $h : \mathbb{H} \rightarrow \mathbb{C}$ . In particular, suppose that  $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{GL}_2^+(\mathbb{R})$ . If  $h(z)$  is a meromorphic function on  $\mathbb{H}$  and  $k$  is an integer, then define the slash operator  $|_k$  by

$$(h|_k\gamma)(z) := (\det \gamma)^{k/2} (cz + d)^{-k} h(\gamma z).$$

**Definition 2.1.** *Let  $\Gamma$  be a congruence subgroup of level  $N$ . A holomorphic function  $h : \mathbb{H} \rightarrow \mathbb{C}$  is called a modular form with integer weight  $k$  on  $\Gamma$  if the following hold:*

(1) *We have*

$$h\left(\frac{az+b}{cz+d}\right) = (cz+d)^k h(z)$$

*for all  $z \in \mathbb{H}$  and  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma$ .*

(2) *If  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ , then  $(h|_k\gamma)(z)$  has a Fourier expansion of the form*

$$(h|_k\gamma)(z) := \sum_{n \geq 0} a_\gamma(n) q_N^n$$

*where  $q_N := e^{2\pi iz/N}$ .*



For a positive integer  $k$ , the complex vector space of modular forms of weight  $k$  with respect to a congruence subgroup  $\Gamma$  is denoted by  $M_k(\Gamma)$ .

**Definition 2.2.** [17, Definition 1.15] *If  $\chi$  is a Dirichlet character modulo  $N$ , then we say that a modular form  $h \in M_k(\Gamma_1(N))$  has Nebentypus character  $\chi$  if*

$$h\left(\frac{az+b}{cz+d}\right) = \chi(d)(cz+d)^k h(z)$$

for all  $z \in \mathbb{H}$  and  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_0(N)$ . The space of such modular forms is denoted by  $M_k(\Gamma_0(N), \chi)$ .

The relevant modular forms for the results obtained in this article arise from eta-quotients. Recall that the Dedekind eta-function  $\eta(z)$  is defined by

$$\eta(z) := q^{1/24}(q; q)_\infty = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$$

where  $q := e^{2\pi iz}$  and  $z \in \mathbb{H}$ . A function  $h(z)$  is called an eta-quotient if it is of the form

$$h(z) := \prod_{\delta|N} \eta(\delta z)^{r_\delta}$$

where  $N$  and  $r_\delta$  are integers with  $N > 0$ .

**Theorem 2.3.** [17, Theorem 1.64] *If  $h(z) = \prod_{\delta|N} \eta(\delta z)^{r_\delta}$  is an eta-quotient such that*

$$k = \frac{1}{2} \sum_{\delta|N} r_\delta \in \mathbb{Z},$$

$$\sum_{\delta|N} \delta r_\delta \equiv 0 \pmod{24} \quad \text{and} \quad \sum_{\delta|N} \frac{N}{\delta} r_\delta \equiv 0 \pmod{24},$$

then  $h(z)$  satisfies

$$h\left(\frac{az+b}{cz+d}\right) = \chi(d)(cz+d)^k h(z)$$

for each  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_0(N)$ . Here the character  $\chi$  is defined by  $\chi(d) := \left(\frac{(-1)^k s}{d}\right)$  where  $s = \prod_{\delta|N} \delta^{r_\delta}$ .

**Theorem 2.4.** [17, Theorem 1.65] *Let  $c, d$  and  $N$  be positive integers with  $d|N$  and  $\gcd(c, d) = 1$ . If  $f$  is an eta-quotient satisfying the conditions of Theorem 2.3 for  $N$ , then the order of vanishing of  $f(z)$  at the cusp  $\frac{c}{d}$  is*

$$\frac{N}{24} \sum_{\delta|N} \frac{\gcd(d, \delta)^2 r_\delta}{\gcd(d, \frac{N}{d}) d \delta}.$$

Suppose that  $h(z)$  is an eta-quotient satisfying the conditions of Theorem 2.3 and that the associated weight  $k$  is a positive integer. If  $h(z)$  is holomorphic at all of the cusps of  $\Gamma_0(N)$ , then  $h(z) \in M_k(\Gamma_0(N), \chi)$ . Theorem 2.4 gives the necessary criterion for determining orders of an eta-quotient at cusps. In the proofs of our results, we use Theorems 2.3 and 2.4 to prove that  $h(z) \in M_k(\Gamma_0(N), \chi)$  for certain eta-quotients  $h(z)$  we consider in the sequel.

We shall now mention a result of Serre [19, P. 43] which will be used later.

**Theorem 2.5.** *Let  $h(z) \in M_k(\Gamma_0(N), \chi)$  has Fourier expansion*

$$h(z) = \sum_{n=0}^{\infty} b(n)q^n \in \mathbb{Z}[[q]].$$

*Then for a positive integer  $r$ , there is a constant  $\alpha > 0$  such that*

$$\#\{0 < n \leq X : b(n) \not\equiv 0 \pmod{r}\} = \mathcal{O}\left(\frac{X}{(\log X)^\alpha}\right).$$

*Equivalently*

$$\lim_{X \rightarrow \infty} \frac{\#\{0 < n \leq X : b(n) \not\equiv 0 \pmod{r}\}}{X} = 0. \quad (2.1)$$

We finally recall the definition of Hecke operators and a few relevant results. Let  $m$  be a positive integer and  $h(z) = \sum_{n=0}^{\infty} a(n)q^n \in M_k(\Gamma_0(N), \chi)$ . Then the action of Hecke operator  $T_m$  on  $h(z)$  is defined by

$$h(z)|T_m := \sum_{n=0}^{\infty} \left( \sum_{d|\gcd(n,m)} \chi(d) d^{k-1} a\left(\frac{mn}{d^2}\right) \right) q^n.$$

In particular, if  $m = p$  is a prime, we have

$$h(z)|T_p := \sum_{n=0}^{\infty} \left( a(pn) + \chi(p) p^{k-1} a\left(\frac{n}{p}\right) \right) q^n.$$

We note that  $a(n) = 0$  unless  $n$  is a non-negative integer.

### 3. PROOF OF THEOREM 1.1

Let  $m = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_r^{\alpha_r}$ , where  $p_i \geq 5$  be distinct prime numbers and  $\alpha_i \geq 0$  be non-negative integers. From (1.1), we get

$$\sum_{n=0}^{\infty} B_{p,m}(n)q^n = \frac{(q^p; q^p)_{\infty} (q^m; q^m)_{\infty}}{(q; q)_{\infty}^2}. \quad (3.1)$$

Note that for any prime  $p$  and positive integers  $j$ , we have

$$(q; q)_\infty^{p^j} \equiv (q^p; q^p)_\infty^{p^{j-1}} \pmod{p^j}. \quad (3.2)$$

Thus, we have

$$\sum_{n=0}^{\infty} B_{p,m}(n)q^n \equiv \frac{(q^p; q^p)_\infty (q^m; q^m)_\infty}{(q; q)_\infty^2} \equiv (q; q)_\infty^{p-2} (q^m; q^m)_\infty \pmod{p}. \quad (3.3)$$

We define

$$H_p(z) := \frac{\eta^p(24z)}{\eta(24pz)}.$$

Using the binomial theorem, we get

$$H_p^{p^j}(z) = \frac{\eta^{p^{j+1}}(24z)}{\eta^{p^j}(24pz)} \equiv 1 \pmod{p^{j+1}}.$$

Define

$$F_{p,m,j}(z) := \eta^{p-2}(24z)\eta(24mz)H_p^{p^j}(z) = \frac{\eta^{(p^{j+1}+p-2)}(24z)\eta(24mz)}{\eta^{p^j}(24pz)}.$$

On modulo  $p^{j+1}$ , we get

$$F_{p,m,j}(z) \equiv \eta^{p-2}(24z)\eta(24mz) \equiv q^{p-2+m}(q^{24}, q^{24})_\infty^{p-2}(q^{24m}, q^{24m})_\infty. \quad (3.4)$$

Combining (3.3) and (3.4) together, we obtain

$$F_{p,m,j}(z) \equiv q^{p-2+m}(q^{24}, q^{24})_\infty^{p-2}(q^{24m}, q^{24m})_\infty \equiv \sum_{n=0}^{\infty} B_{p,m}(n)q^{24n+p-2+m} \pmod{p}. \quad (3.5)$$

Next, we show that  $F_{p,m,j}(z)$  is a modular form for certain values of  $p, m$ , and  $j$ .

**Lemma 3.1.** *Let  $m = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_r^{\alpha_r}$ , where  $p_i \geq 5$  be distinct primes and  $\alpha_i \geq 0$  be integers. Then, we have  $F_{p,m,j}(z) \in M_k(\Gamma_0(N), \chi)$ , where  $k = \frac{p-1}{2}(p^j+1)$ ,  $N = 2^5 \cdot 3^2 \cdot p \cdot m$  and  $\chi = \left( \frac{2^{3 \cdot (p-1)(p^j+1)} 3^{(p-1)(p^j+1)} m p^{-p^j}}{\bullet} \right)$ .*

*Proof.* Applying Theorem 2.3, we first estimate the level of eta quotient  $F_{p,m,j}(z)$ . The level of  $F_{p,m,j}(z)$  is  $N = 3 \cdot 2^3 \cdot p \cdot m \cdot M$ , where  $M$  is the smallest positive integer which satisfies

$$\begin{aligned} 3 \cdot 2^3 \cdot p \cdot m \cdot M \left[ \frac{p^{j+1} + p - 2}{2^3 \cdot 3} + \frac{1}{2^3 \cdot 3 \cdot m} - \frac{p^j}{2^3 \cdot 3 \cdot p} \right] &\equiv 0 \pmod{24} \\ \implies M [p \cdot m \cdot (p^{j+1} + p - 2) + p - m p^j] &\equiv 0 \pmod{24} \\ \implies M [m \cdot p^j (p^2 - 1) + p \cdot m \cdot (p - 2) + p] &\equiv 0 \pmod{24} \\ \implies 2M \left[ m \cdot p^j \cdot \frac{(p^2 - 1)}{2} + p \cdot \frac{(m \cdot (p - 2) + 1)}{2} \right] &\equiv 0 \pmod{24}. \end{aligned}$$

Therefore  $M = 12$  and the level of  $F_{p,m,j}(z)$  is  $N = 2^5 \cdot 3^2 \cdot p \cdot m$ . The cusps of  $\Gamma_0(2^5 \cdot 3^2 \cdot p \cdot m)$  are given by fractions  $\frac{c}{d}$  where  $d|2^5 \cdot 3^2 \cdot p \cdot m$  and  $\gcd(c, d) = 1$ . By using Theorem 2.4, we obtain that  $F_{p,m,j}(z)$  is holomorphic at a cusp  $\frac{c}{d}$  if and only if

$$(p^{j+1} + p - 2) \frac{\gcd^2(d, 2^3 \cdot 3)}{2^3 \cdot 3} + \frac{\gcd^2(d, 24m)}{24m} - p^j \frac{\gcd^2(d, 2^3 \cdot 3p)}{2^3 \cdot 3p} \geq 0 \quad (3.6)$$

$$\iff L := (p^{j+1} + p - 2) \cdot pG_1 + \frac{p}{m} \cdot G_2 - p^j \geq 0,$$

where  $G_1 = \frac{\gcd^2(d, 2^3 \cdot 3)}{\gcd^2(d, 2^3 \cdot 3 \cdot p)}$  and  $G_2 = \frac{\gcd^2(d, 2^3 \cdot 3m)}{\gcd^2(d, 2^3 \cdot 3 \cdot p)}$ .

Let  $d$  be a divisor of  $2^5 \cdot 3^2 \cdot p \cdot m$ . We can write  $d = 2^{r_1} 3^{r_2} p^{r_3} t : 0 \leq r_1 \leq 5, 0 \leq r_2 \leq 2, 0 \leq r_3 \leq 1, t|m$ .

Next, we find all possible value of divisors of  $2^5 \cdot 3^2 \cdot p \cdot m$  to compute equation (3.6).

When  $d = 2^{r_1} 3^{r_2} p^{r_3} t : 0 \leq r_1 \leq 5, 0 \leq r_2 \leq 2, 0 \leq r_3 \leq 1, t|m$ . Then  $\frac{1}{p^2} \leq G_1 \leq 1$  and  $\frac{t^2}{p^2} \leq G_2 \leq 1$ . Then equation (3.6) will be

$$L \geq (p^{j+1} + p - 2) \cdot p \cdot \frac{1}{p^2} + \frac{p}{m} \cdot \frac{t^2}{p^2} - p^j \geq \left(1 - \frac{2}{p}\right) + \frac{t^2}{pm} \geq 0. \quad (3.7)$$

Therefore,  $F_{p,m,j}(z)$  is holomorphic at every cusp  $\frac{c}{d}$ . Using Theorem 2.3, we compute the weight of  $F_{p,m,j}(z)$  is  $k = \frac{(p-1)}{2} (p^j + 1)$  which is a positive integer. The associated character for  $F_{p,m,j}(z)$  is

$$\chi = \left( \frac{2^{3 \cdot (p-1)(p^j+1)} 3^{(p-1)(p^j+1)} m p^{-p^j}}{\bullet} \right).$$

Thus  $F_{p,m,j}(z) \in M_k(\Gamma_0(N), \chi)$  where  $k, N$  and  $\chi$  are as above □

**Proof of Theorem 1.1.** Using Lemma 3.1, we find  $F_{p,m,j}(z) \in M_k(\Gamma_0(N), \chi)$  where  $k = \frac{(p-1)}{2} (p^j + 1)$ ,  $N = 2^5 \cdot 3^2 \cdot p \cdot m$  and  $\chi = \left( \frac{2^{3 \cdot (p-1)(p^j+1)} 3^{(p-1)(p^j+1)} m p^{-p^j}}{\bullet} \right)$ . Employing Theorem 2.5, we obtain that the Fourier coefficients of  $F_{p,m,j}(z)$  satisfies (2.1) for  $r = p$  which implies that the Fourier coefficient of  $F_{p,m,j}(z)$  are almost always divisible by  $p$ . Hence, from (3.5), we conclude that  $B_{p,m}(n)$  are almost always divisible by  $p$ . This completes the proof of Theorem 1.1. □

4. CONGRUENCES FOR  $B_{3,7}(n)$ .

**Proof of Theorem 1.2(i).** From equation (1.1), we have

$$\sum_{n=0}^{\infty} B_{3,7}(n)q^n = \frac{(q^3; q^3)_{\infty}(q^7; q^7)_{\infty}}{(q; q)_{\infty}^2}. \quad (4.1)$$

From (4.1) and (3.2), we get

$$\sum_{n=0}^{\infty} B_{3,7}(n)q^n \equiv \frac{(q^3; q^3)_{\infty}(q^7; q^7)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty}(q^7; q^7)_{\infty} \pmod{3}. \quad (4.2)$$

Thus, we have

$$\sum_{n=0}^{\infty} B_{3,7}(n)q^{3n+1} \equiv q(q^3; q^3)_{\infty}(q^{21}; q^{21})_{\infty} \equiv \eta(3z)\eta(21z) \pmod{3}. \quad (4.3)$$

By using Theorem 2.3, we obtain  $\eta(3z)\eta(21z) \in S_1(\Gamma_0(63), (\frac{-63}{\bullet}))$ . Thus  $\eta(3z)\eta(21z)$  has a Fourier expansion i.e.

$$\eta(3z)\eta(21z) = q - q^4 - q^7 - \dots = \sum_{n=1}^{\infty} a(n)q^n \quad (4.4)$$

Thus,  $a(n) = 0$  if  $n \not\equiv 1 \pmod{3}$ , for all  $n \geq 0$ . From (4.3) and (4.4), comparing the coefficient of  $q^{3n+1}$ , we get

$$B_{3,7}(n) \equiv a(3n+1) \pmod{3}. \quad (4.5)$$

Since  $\eta(3z)\eta(21z)$  is a Hecke eigenform (see [14]), it gives

$$\eta(3z)\eta(21z)|T_p = \sum_{n=1}^{\infty} \left( a(pn) + \left( \frac{-63}{p} \right)_L a\left(\frac{n}{p}\right) \right) q^n = \lambda(p) \sum_{n=1}^{\infty} a(n)q^n.$$

Note that  $\left( \frac{-63}{p} \right)_L = \left( \frac{-7}{p} \right)_L$ . Comparing the coefficients of  $q^n$  on both sides of the above equation, we get

$$a(pn) + \left( \frac{-7}{p} \right)_L a\left(\frac{n}{p}\right) = \lambda(p)a(n). \quad (4.6)$$

Since  $a(1) = 1$  and  $a(\frac{1}{p}) = 0$ , if we put  $n = 1$  in the above expression, we get  $a(p) = \lambda(p)$ . As  $a(p) = 0$  for all  $p \not\equiv 1 \pmod{3}$  this implies that  $\lambda(p) = 0$  for all  $p \not\equiv 1 \pmod{3}$ . From (4.6) we get that for all  $p \not\equiv 1 \pmod{3}$

$$a(pn) + \left( \frac{-7}{p} \right)_L a\left(\frac{n}{p}\right) = 0. \quad (4.7)$$

Now, we consider two cases here. If  $p \nmid n$ , then replacing  $n$  by  $pn + r$  with  $\gcd(r, p) = 1$  in (4.7), we get

$$a(p^2n + pr) = 0. \quad (4.8)$$

Replacing  $n$  by  $3n - pr + 1$  in (4.8) and using (4.5), we get

$$B_{3,7} \left( p^2 n + pr \frac{(1-p^2)}{3} + \frac{p^2-1}{3} \right) \equiv 0 \pmod{3}. \quad (4.9)$$

Since  $p \equiv 2 \pmod{3}$ , we have  $3|(1-p^2)$  and  $\gcd\left(\frac{(1-p^2)}{3}, p\right) = 1$ , when  $r$  runs over a residue system excluding the multiples of  $p$ , so does  $r\frac{(1-p^2)}{3}$ . Thus for  $p \nmid j$ , (4.9) can be written as

$$B_{3,7} \left( p^2 n + pj + \frac{p^2-1}{3} \right) \equiv 0 \pmod{3}. \quad (4.10)$$

Now we consider the second case, when  $p|n$ . Here replacing  $n$  by  $pn$  in (4.7), we get

$$a(p^2 n) = - \left( \frac{-7}{p} \right)_L \cdot a(n). \quad (4.11)$$

Further substituting  $n$  by  $3n + 1$  in (4.11) we obtain

$$a(3p^2 n + p^2) = - \left( \frac{-7}{p} \right)_L \cdot a(3n + 1). \quad (4.12)$$

Using (4.5) in (4.12), we get

$$B_{3,7} \left( p^2 n + \frac{p^2-1}{3} \right) = - \left( \frac{-7}{p} \right)_L \cdot B_{3,7}(n). \quad (4.13)$$

Let  $p_i$  be primes such that  $p_i \equiv 2 \pmod{3}$ . Further note that

$$p_1^2 p_2^2 \cdots p_k^2 n + \frac{p_1^2 p_2^2 \cdots p_k^2 - 1}{3} = p_1^2 \left( p_2^2 \cdots p_k^2 n + \frac{p_2^2 \cdots p_k^2 - 1}{3} \right) + \frac{p_1^2 - 1}{3}.$$

Repeatedly using (4.13) and (4.10), we get

$$\begin{aligned} & B_{3,7} \left( p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 n + \frac{p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 (3j + p_{k+1}) - 1}{3} \right) \\ & \equiv - \left( \frac{-7}{p_1} \right)_L \cdot B_{3,7} \left( p_2^2 \cdots p_k^2 p_{k+1}^2 n + \frac{p_2^2 \cdots p_k^2 p_{k+1}^2 (3j + p_{k+1}) - 1}{3} \right) \equiv \cdots \\ & \equiv (-1)^k \left( \left( \frac{-7}{p_1} \right)_L \left( \frac{-7}{p_2} \right)_L \cdots \left( \frac{-7}{p_k} \right)_L \right) B_{3,7} \left( p_{k+1}^2 n + p_{k+1} j + \frac{p_{k+1}^2 - 1}{3} \right) \equiv 0 \pmod{3} \end{aligned}$$

when  $j \not\equiv 0 \pmod{p_{k+1}}$ . This completes the proof of the theorem.  $\square$

**Proof of Theorem 1.4(i).** From (4.1) and (3.2), we get

$$\sum_{n=0}^{\infty} B_{3,7}(n) q^n \equiv \frac{(q^3; q^3)_{\infty} (q^7; q^7)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty} (q^7; q^7)_{\infty} \pmod{3}. \quad (4.14)$$

Define

$$\sum_{n=0}^{\infty} c(n) q^n := f_1 f_7 = (q; q)_{\infty} (q^7; q^7)_{\infty}. \quad (4.15)$$

In 1959, Newman [15] proved that if  $p$  is a prime with  $p \equiv 1 \pmod{6}$ , then

$$c\left(pn + \frac{p-1}{3}\right) = c\left(\frac{p-1}{3}\right) c(n) - (-1)^{\frac{p-1}{2}} \left(\frac{7}{p}\right)_L c\left(\frac{n}{p} - \frac{p-1}{3p}\right). \quad (4.16)$$

Observe that, if  $p \nmid (3n+1)$ , then  $\left(\frac{n}{p} - \frac{p-1}{3p}\right)$  is not an integer and

$$c\left(\frac{n}{p} - \frac{p-1}{3p}\right) = 0. \quad (4.17)$$

From (4.16) and (4.17), we get that if  $p \nmid (3n+1)$ , then

$$c\left(pn + \frac{p-1}{3}\right) = c\left(\frac{p-1}{3}\right) c(n). \quad (4.18)$$

Thus, if  $p \nmid (3n+1)$  and  $c\left(\frac{p-1}{3}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ ,

$$c\left(pn + \frac{p-1}{3}\right) \equiv 0 \pmod{3}. \quad (4.19)$$

Replacing  $n$  by  $pn + \frac{p-1}{3}$  in (4.16), we get

$$c\left(p^2n + \frac{p^2-1}{3}\right) = c\left(\frac{p-1}{3}\right) c\left(pn + \frac{p-1}{3}\right) - (-1)^{\frac{p-1}{2}} \left(\frac{7}{p}\right)_L c(n). \quad (4.20)$$

Observe that from (4.20) if  $c\left(\frac{p-1}{3}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ , we get

$$c\left(p^2n + \frac{p^2-1}{3}\right) \equiv -(-1)^{\frac{p-1}{2}} \left(\frac{7}{p}\right)_L c(n) \pmod{3}. \quad (4.21)$$

From (4.21) and by using mathematical induction, we get that if  $c\left(\frac{p-1}{3}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ , and  $\alpha \geq 0$ , we get

$$c\left(p^{2\alpha}n + \frac{p^{2\alpha}-1}{3}\right) \equiv \left(-(-1)^{\frac{p-1}{2}} \left(\frac{7}{p}\right)_L\right)^\alpha c(n) \pmod{3}. \quad (4.22)$$

Replacing  $n$  by  $pn + \frac{p-1}{3}$  in (4.22) and using (4.19), we obtain that if  $p \nmid (3n+1)$  and  $c\left(\frac{p-1}{3}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$  and  $\alpha \geq 0$ , we get

$$c\left(p^{2\alpha+1}n + \frac{p^{2\alpha+1}-1}{3}\right) \equiv \pmod{3}. \quad (4.23)$$

From (4.15) and (4.14), we obtain that for  $n \geq 0$ ,

$$B_{3,7}(n) \equiv c(n) \pmod{3}. \quad (4.24)$$

Thus, combining (4.24) and (4.23), we obtain the result of Theorem.  $\square$

**Proof of Theorem 1.5(i).** From (4.7), we get that for any prime  $p \equiv 2 \pmod{3}$

$$a(pn) = -\left(\frac{-7}{p}\right)_L a\left(\frac{n}{p}\right). \quad (4.25)$$

Replacing  $n$  by  $3n + 2$ , we obtain

$$a(3pn + 2p) = - \left( \frac{-7}{p} \right)_L a \left( \frac{3n + 2}{p} \right). \quad (4.26)$$

Next substituting  $n$  by  $p^k n + r$  with  $p \nmid r$  in (4.26), we obtain

$$a \left( 3 \left( p^{k+1} n + pr + \frac{2p-1}{3} \right) + 1 \right) = - \left( \frac{-7}{p} \right)_L a \left( 3 \left( p^{k-1} n + \frac{3r+2-p}{3p} \right) + 1 \right). \quad (4.27)$$

Note that  $\frac{2p-1}{3}$  and  $\frac{3r+2-p}{3p}$  are integers. Using (4.27) and (4.5), we get

$$B_{3,7} \left( p^{k+1} n + pr + \frac{2p-1}{3} \right) \equiv - \left( \frac{-7}{p} \right)_L B_{3,7} \left( p^{k-1} n + \frac{3r+2-p}{3p} \right) \pmod{3}. \quad (4.28)$$

□

**Proof of Corollary 1.6(i).** Let  $p$  be a prime such that  $p \equiv 2 \pmod{3}$ . Choose a non negative integer  $r$  such that  $3r + 2 = p^{2k-1}$ . Substituting  $k$  by  $2k - 1$  in (4.28), we obtain

$$\begin{aligned} B_{3,7} \left( p^{2k} n + \frac{p^{2k} - 1}{3} \right) &\equiv - \left( \frac{-7}{p} \right)_L B_{3,7} \left( p^{2k-2} n + \frac{p^{2k-2} - 1}{3} \right) \\ &\equiv \dots \equiv (-1)^k \left( \frac{-7}{p} \right)_L^k B_{3,7}(n) \pmod{3}. \end{aligned}$$

□

## 5. CONGRUENCES FOR $B_{3,5}(n)$

**Proof of Theorem 1.2(ii).** From equation (1.1), we have

$$\sum_{n=0}^{\infty} B_{3,5}(n) q^n = \frac{(q^3; q^3)_{\infty} (q^5; q^5)_{\infty}}{(q; q)_{\infty}^2}. \quad (5.1)$$

Note that for any prime number  $p$ , we have

$$(q^p; q^p)_{\infty} \equiv (q; q)_{\infty}^p \pmod{p}. \quad (5.2)$$

From (5.1) and (5.2), we get

$$\sum_{n=0}^{\infty} B_{3,5}(n) q^n \equiv \frac{(q^3; q^3)_{\infty} (q^5; q^5)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty} (q^5; q^5)_{\infty} \pmod{3}. \quad (5.3)$$

Thus we have

$$\sum_{n=0}^{\infty} B_{3,5}(n) q^{4n+1} \equiv q(q^4; q^4)_{\infty} (q^{20}; q^{20})_{\infty} \equiv \eta(4z)\eta(20z) \pmod{3}. \quad (5.4)$$



By using Theorem 2.3, we obtain  $\eta(4z)\eta(20z) \in S_1(\Gamma_0(80), \left(\frac{-80}{\bullet}\right)_L)$ . Observe that  $\eta(4z)\eta(20z)$  has a Fourier expansion i.e.

$$\eta(4z)\eta(20z) = q - q^5 - q^9 + \cdots = \sum_{n=1}^{\infty} b(n)q^n \quad (5.5)$$

Thus,  $b(n) = 0$  if  $n \not\equiv 1 \pmod{4}$ , for all  $n \geq 0$ . From (5.4) and (5.5), comparing the coefficient of  $q^{4n+1}$ , we get

$$B_{3,5}(n) \equiv b(4n+1) \pmod{3}. \quad (5.6)$$

Since  $\eta(4z)\eta(20z)$  is a Hecke eigenform (see [14]), it gives

$$\eta(4z)\eta(20z)|T_p = \sum_{n=1}^{\infty} \left( b(pn) + \left(\frac{-80}{p}\right)_L b\left(\frac{n}{p}\right) \right) q^n = \lambda(p) \sum_{n=1}^{\infty} b(n)q^n.$$

Note that  $\left(\frac{-80}{p}\right)_L = \left(\frac{-5}{p}\right)_L$ . Comparing the coefficients of  $q^n$  on both sides of the above equation, we get

$$b(pn) + \left(\frac{-5}{p}\right)_L b\left(\frac{n}{p}\right) = \lambda(p)b(n). \quad (5.7)$$

Since  $b(1) = 1$  and  $b\left(\frac{1}{p}\right) = 0$ , if we put  $n = 1$  in the above expression, we get  $b(p) = \lambda(p)$ . As  $b(p) = 0$  for all  $p \not\equiv 1 \pmod{4}$  this implies that  $\lambda(p) = 0$  for all  $p \not\equiv 1 \pmod{4}$ . From (5.7) we get that for all  $p \not\equiv 1 \pmod{4}$

$$b(pn) + \left(\frac{-5}{p}\right)_L b\left(\frac{n}{p}\right) = 0. \quad (5.8)$$

Now, we consider two cases here. If  $p \nmid n$ , then replacing  $n$  by  $pn + r$  with  $\gcd(r, p) = 1$  in (5.8), we get

$$b(p^2n + pr) = 0. \quad (5.9)$$

Replacing  $n$  by  $4n - pr + 1$  in (5.9) and using (5.6), we get

$$B_{3,5} \left( p^2n + pr \frac{(1-p^2)}{4} + \frac{p^2-1}{4} \right) \equiv 0 \pmod{3}. \quad (5.10)$$

Since  $p \equiv 3 \pmod{4}$ , we have  $4|(1-p^2)$  and  $\gcd\left(\frac{(1-p^2)}{4}, p\right) = 1$ , when  $r$  runs over a residue system excluding the multiples of  $p$ , so does  $r \frac{(1-p^2)}{4}$ . Thus for  $p \nmid j$ , (5.10) can be written as

$$B_{3,5} \left( p^2n + pj + \frac{p^2-1}{4} \right) \equiv 0 \pmod{3}. \quad (5.11)$$

Now we consider the second case, when  $p|n$ . Here replacing  $n$  by  $pn$  in (5.8), we get

$$b(p^2n) = - \left(\frac{-5}{p}\right)_L b(n). \quad (5.12)$$

Further substituting  $n$  by  $4n + 1$  in (5.12) we obtain

$$b(4p^2n + p^2) = - \left( \frac{-5}{p} \right)_L b(4n + 1). \quad (5.13)$$

Using (5.6) in (5.13), we get

$$B_{3,5} \left( p^2n + \frac{p^2 - 1}{4} \right) \equiv - \left( \frac{-5}{p} \right)_L \cdot B_{3,5}(n) \pmod{3}. \quad (5.14)$$

Let  $p_i$  be primes such that  $p_i \equiv 3 \pmod{4}$ . Further note that

$$p_1^2 p_2^2 \cdots p_k^2 n + \frac{p_1^2 p_2^2 \cdots p_k^2 - 1}{4} = p_1^2 \left( p_2^2 \cdots p_k^2 n + \frac{p_2^2 \cdots p_k^2 - 1}{4} \right) + \frac{p_1^2 - 1}{4}.$$

Repeatedly using (5.14) and (5.11), we get

$$\begin{aligned} & B_{3,5} \left( p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 n + \frac{p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 (4j + p_{k+1}) - 1}{4} \right) \\ & \equiv (-1) \left( \frac{-5}{p_1} \right)_L B_{3,5} \left( p_2^2 \cdots p_k^2 p_{k+1}^2 n + \frac{p_2^2 \cdots p_k^2 p_{k+1}^2 (4j + p_{k+1}) - 1}{4} \right) \equiv \cdots \\ & \equiv (-1)^k \left( \left( \frac{-5}{p_1} \right)_L \left( \frac{-5}{p_2} \right)_L \cdots \left( \frac{-5}{p_k} \right)_L \right) B_{3,5} \left( p_{k+1}^2 n + p_{k+1} j + \frac{p_{k+1}^2 - 1}{4} \right) \equiv 0 \pmod{3} \end{aligned}$$

when  $j \not\equiv 0 \pmod{p_{k+1}}$ . This completes the proof of the theorem.  $\square$

**Proof of Theorem 1.4(ii).** From (4.1) and (3.2), we get

$$\sum_{n=0}^{\infty} B_{3,5}(n) q^n \equiv \frac{(q^3; q^3)_{\infty} (q^5; q^5)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty} (q^5; q^5)_{\infty} \pmod{3}. \quad (5.15)$$

Define

$$\sum_{n=0}^{\infty} t(n) q^n := f_1 f_5 = (q; q)_{\infty} (q^5; q^5)_{\infty}. \quad (5.16)$$

In 1959, Newman [15] proved that if  $p$  is a prime with  $p \equiv 1 \pmod{4}$ , then

$$t \left( pn + \frac{p-1}{4} \right) = t \left( \frac{p-1}{4} \right) t(n) - \left( \frac{5}{p} \right)_L t \left( \frac{n}{p} - \frac{p-1}{4p} \right). \quad (5.17)$$

Therefore, if  $p \nmid (4n + 1)$ , then  $\left( \frac{n}{p} - \frac{p-1}{4p} \right)$  is not an integer and

$$t \left( \frac{n}{p} - \frac{p-1}{4p} \right) = 0. \quad (5.18)$$

From (5.17) and (5.18) we get that if  $p \nmid (4n + 1)$ , then

$$t \left( pn + \frac{p-1}{4} \right) = t \left( \frac{p-1}{4} \right) t(n). \quad (5.19)$$

Thus, if  $p \nmid (4n + 1)$  and  $t\left(\frac{p-1}{4}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ ,

$$t\left(pn + \frac{p-1}{4}\right) \equiv 0 \pmod{3}. \quad (5.20)$$

Replacing  $n$  by  $pn + \frac{p-1}{4}$  in (5.17), we get

$$t\left(p^2n + \frac{p^2-1}{4}\right) = t\left(\frac{p-1}{4}\right) t\left(pn + \frac{p-1}{4}\right) - \left(\frac{5}{p}\right)_L t(n). \quad (5.21)$$

Observe that from (5.21) if  $t\left(\frac{p-1}{4}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ , we get

$$t\left(p^2n + \frac{p^2-1}{4}\right) \equiv -\left(\frac{5}{p}\right)_L t(n) \pmod{3}. \quad (5.22)$$

From (5.22) and by using mathematical induction, we get that if  $t\left(\frac{p-1}{4}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ , and  $\alpha \geq 0$ , we get

$$t\left(p^{2\alpha}n + \frac{p^{2\alpha}-1}{4}\right) \equiv \left(-\left(\frac{5}{p}\right)_L\right)^\alpha t(n) \pmod{3}. \quad (5.23)$$

Replacing  $n$  by  $pn + \frac{p-1}{4}$  in (5.23) and using (5.20), we obtain that if  $p \nmid (4n + 1)$  and  $t\left(\frac{p-1}{4}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$  and  $\alpha \geq 0$ , we get

$$t\left(p^{2\alpha+1}n + \frac{p^{2\alpha+1}-1}{4}\right) \equiv 0 \pmod{3}. \quad (5.24)$$

From (5.16) and (5.15), we obtain that for  $n \geq 0$ ,

$$B_{3,5}(n) \equiv t(n) \pmod{3}. \quad (5.25)$$

Thus, combining (5.25) and (5.24), we obtain the result of Theorem.  $\square$

**Proof of Theorem 1.5(ii).** From (5.8), we get that for any prime  $p \equiv 3 \pmod{4}$

$$b(pn) = -\left(\frac{-5}{p}\right)_L b\left(\frac{n}{p}\right). \quad (5.26)$$

Replacing  $n$  by  $4n + 3$ , we obtain

$$b(4pn + 3p) = -\left(\frac{-5}{p}\right)_L b\left(\frac{4n + 3}{p}\right). \quad (5.27)$$

Next substituting  $n$  by  $p^k n + r$  with  $p \nmid r$  in (4.26), we obtain

$$b\left(4\left(p^{k+1}n + pr + \frac{3p-1}{4}\right) + 1\right) = -\left(\frac{-5}{p}\right)_L b\left(4\left(p^{k-1}n + \frac{4r+3-p}{4p}\right) + 1\right). \quad (5.28)$$

Note that  $\frac{3p-1}{4}$  and  $\frac{4r+3-p}{4p}$  are integers. Using (5.28) and (5.6), we get

$$B_{3,5} \left( p^{k+1}n + pr + \frac{3p-1}{4} \right) \equiv - \left( \frac{-5}{p} \right)_L B_{3,5} \left( p^{k-1}n + \frac{4r+3-p}{4p} \right) \pmod{3}. \quad (5.29)$$

□

**Proof of Corollary 1.6(ii).** Let  $p$  be a prime such that  $p \equiv 3 \pmod{4}$ . Choose a non negative integer  $r$  such that  $4r+3 = p^{2k-1}$ . Substituting  $k$  by  $2k-1$  in (5.29), we obtain

$$\begin{aligned} B_{3,5} \left( p^{2k}n + \frac{p^{2k}-1}{4} \right) &\equiv - \left( \frac{-5}{p} \right)_L B_{3,5} \left( p^{2k-2}n + \frac{p^{2k-2}-1}{4} \right) \\ &\equiv \cdots \equiv \left( - \left( \frac{-5}{p} \right)_L \right)^k B_{3,5}(n) \pmod{3}. \end{aligned}$$

□

## 6. CONGRUENCES FOR $B_{3,2}(n)$

**Proof of Theorem 1.2(iii).** From equation (1.1), we have

$$\sum_{n=0}^{\infty} B_{3,2}(n)q^n = \frac{(q^3; q^3)_{\infty}(q^2; q^2)_{\infty}}{(q; q)_{\infty}^2}. \quad (6.1)$$

From (6.1) and (3.2), we obtain

$$\sum_{n=0}^{\infty} B_{3,2}(n)q^n = \frac{(q^3; q^3)_{\infty}(q^2; q^2)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty}(q^2; q^2)_{\infty} \pmod{3}. \quad (6.2)$$

Thus, we have

$$\sum_{n=0}^{\infty} B_{3,2}(n)q^{8n+1} \equiv q(q^8; q^8)_{\infty}(q^{16}; q^{16})_{\infty} \equiv \eta(8z)\eta(16z) \pmod{3}. \quad (6.3)$$

By using Theorem 2.3, we obtain  $\eta(8z)\eta(16z) \in S_1(\Gamma_0(128), (\frac{-128}{\bullet})_L)$ . Note that  $\eta(8z)\eta(16z)$  has a Fourier expansion i.e.

$$\eta(8z)\eta(16z) = q - q^9 - 2q^{17} - \cdots = \sum_{n=1}^{\infty} d(n)q^n \quad (6.4)$$

Thus,  $d(n) = 0$  if  $n \not\equiv 1 \pmod{8}$ , for all  $n \geq 0$ . From (6.3) and (6.4), comparing the coefficient of  $q^{8n+1}$ , we get

$$B_{3,2}(n) \equiv d(8n+1) \pmod{3}. \quad (6.5)$$

Since  $\eta(8z)\eta(16z)$  is a Hecke eigenform (see [14]), it gives

$$\eta(8z)\eta(16z)|T_p = \sum_{n=1}^{\infty} \left( d(pn) + \left( \frac{-128}{p} \right)_L d\left(\frac{n}{p}\right) \right) q^n = \lambda(p) \sum_{n=1}^{\infty} d(n)q^n.$$

Note that  $\left( \frac{-128}{p} \right)_L = \left( \frac{-2}{p} \right)_L$ . Comparing the coefficients of  $q^n$  on both sides of the above equation, we get

$$d(pn) + \left( \frac{-2}{p} \right)_L d\left(\frac{n}{p}\right) = \lambda(p)d(n). \quad (6.6)$$

Since  $d(1) = 1$  and  $d\left(\frac{1}{p}\right) = 0$ , if we put  $n = 1$  in the above expression, we get  $d(p) = \lambda(p)$ . As  $d(p) = 0$  for all  $p \not\equiv 1 \pmod{8}$  this implies that  $\lambda(p) = 0$  for all  $p \not\equiv 1 \pmod{8}$ . From (6.6) we get that for all  $p \not\equiv 1 \pmod{8}$

$$d(pn) + \left( \frac{-2}{p} \right)_L d\left(\frac{n}{p}\right) = 0. \quad (6.7)$$

Now, we consider two cases here. If  $p \nmid n$ , then replacing  $n$  by  $pn + r$  with  $\gcd(r, p) = 1$  in (6.7), we get

$$d(p^2n + pr) = 0. \quad (6.8)$$

Replacing  $n$  by  $8n - pr + 1$  in (6.8) and using (6.5), we get

$$B_{3,2} \left( p^2n + pr \frac{(1-p^2)}{8} + \frac{p^2-1}{8} \right) \equiv 0 \pmod{3}. \quad (6.9)$$

Since  $p \not\equiv 1 \pmod{8}$ , we have  $8 \mid (1-p^2)$  and  $\gcd\left(\frac{(1-p^2)}{8}, p\right) = 1$ , when  $r$  runs over a residue system excluding the multiples of  $p$ , so does  $r \frac{(1-p^2)}{8}$ . Thus for  $p \nmid j$ , (6.9) can be written as

$$B_{3,2} \left( p^2n + pj + \frac{p^2-1}{8} \right) \equiv 0 \pmod{3}. \quad (6.10)$$

Now we consider the second case, when  $p \mid n$ . Here replacing  $n$  by  $pn$  in (6.7), we get

$$d(p^2n) = - \left( \frac{-2}{p} \right)_L \cdot d(n). \quad (6.11)$$

Further substituting  $n$  by  $8n + 1$  in (6.11) we obtain

$$d(8p^2n + p^2) = - \left( \frac{-2}{p} \right)_L d(8n + 1). \quad (6.12)$$

Using (6.5) in (6.12), we get

$$B_{3,2} \left( p^2n + \frac{p^2-1}{8} \right) = - \left( \frac{-2}{p} \right)_L B_{3,2}(n). \quad (6.13)$$

Let  $p_i$  be primes such that  $p_i \not\equiv 1 \pmod{8}$ . Further note that

$$p_1^2 p_2^2 \cdots p_k^2 n + \frac{p_1^2 p_2^2 \cdots p_k^2 - 1}{8} = p_1^2 \left( p_2^2 \cdots p_k^2 n + \frac{p_2^2 \cdots p_k^2 - 1}{8} \right) + \frac{p_1^2 - 1}{8}.$$

Repeatedly using (6.13) and (6.10), we get

$$\begin{aligned}
& B_{3,2} \left( p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 n + \frac{p_1^2 p_2^2 \cdots p_k^2 p_{k+1}^2 (8j + p_{k+1}) - 1}{8} \right) \\
& \equiv - \left( \frac{-2}{p_1} \right)_L B_{3,2} \left( p_2^2 \cdots p_k^2 p_{k+1}^2 n + \frac{p_2^2 \cdots p_k^2 p_{k+1}^2 (8j + p_{k+1}) - 1}{8} \right) \equiv \cdots \\
& \equiv (-1)^k \left( \frac{-2}{p_1} \right)_L \left( \frac{-2}{p_2} \right)_L \cdots \left( \frac{-2}{p_k} \right)_L B_{3,2} \left( p_{k+1}^2 n + p_{k+1} j + \frac{p_{k+1}^2 - 1}{8} \right) \equiv 0 \pmod{3}
\end{aligned}$$

when  $j \not\equiv 0 \pmod{p_{k+1}}$ . This completes the proof of the theorem.  $\square$

**Proof of Theorem 1.4(iii).** From (4.1) and (3.2), we get

$$\sum_{n=0}^{\infty} B_{3,2}(n) q^n \equiv \frac{(q^3; q^3)_{\infty} (q^2; q^2)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty} (q^2; q^2)_{\infty} \pmod{3}. \quad (6.14)$$

Define

$$\sum_{n=0}^{\infty} u(n) q^n := f_1 f_2 = (q; q)_{\infty} (q^2; q^2)_{\infty}. \quad (6.15)$$

In 1959, Newman [15] proved that if  $p$  is a prime with  $p \equiv 1 \pmod{8}$ , then

$$u \left( pn + \frac{p-1}{8} \right) = u \left( \frac{p-1}{8} \right) u(n) - \left( \frac{2}{p} \right)_L u \left( \frac{n}{p} - \frac{p-1}{8p} \right). \quad (6.16)$$

Therefore, if  $p \nmid (8n+1)$ , then  $\left( \frac{n}{p} - \frac{p-1}{8p} \right)$  is not an integer and

$$u \left( \frac{n}{p} - \frac{p-1}{8p} \right) = 0. \quad (6.17)$$

From (6.16) and (6.17) we get that if  $p \nmid (8n+1)$ , then

$$u \left( pn + \frac{p-1}{8} \right) = u \left( \frac{p-1}{8} \right) u(n). \quad (6.18)$$

Thus, if  $p \nmid (8n+1)$  and  $u \left( \frac{p-1}{8} \right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ ,

$$u \left( pn + \frac{p-1}{8} \right) \equiv 0 \pmod{3}. \quad (6.19)$$

Replacing  $n$  by  $pn + \frac{p-1}{8}$  in (6.16), we get

$$u \left( p^2 n + \frac{p^2 - 1}{8} \right) = u \left( \frac{p-1}{8} \right) u \left( pn + \frac{p-1}{8} \right) - \left( \frac{2}{p} \right)_L u(n). \quad (6.20)$$

Observe that from (6.20) if  $u\left(\frac{p-1}{8}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ , we get

$$u\left(p^2n + \frac{p^2-1}{8}\right) \equiv -\left(\frac{2}{p}\right)_L u(n) \pmod{3}. \quad (6.21)$$

From (6.21) and by using mathematical induction, we get that if  $u\left(\frac{p-1}{8}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$ , and  $\alpha \geq 0$ , we get

$$u\left(p^{2\alpha}n + \frac{p^{2\alpha}-1}{8}\right) \equiv \left(\left(\frac{2}{p}\right)_L\right)^\alpha u(n) \pmod{3}. \quad (6.22)$$

Replacing  $n$  by  $pn + \frac{p-1}{8}$  in (5.23) and using (6.19), we obtain that if  $p \nmid (8n+1)$  and  $u\left(\frac{p-1}{8}\right) \equiv 0 \pmod{3}$ , then for  $n \geq 0$  and  $\alpha \geq 0$ , we get

$$u\left(p^{2\alpha+1}n + \frac{p^{2\alpha+1}-1}{8}\right) \equiv 0 \pmod{3}. \quad (6.23)$$

From (6.15) and (6.14), we obtain that for  $n \geq 0$ ,

$$B_{3,2}(n) \equiv u(n) \pmod{3}. \quad (6.24)$$

Thus, combining (6.24) and (6.23), we obtain the result of Theorem.  $\square$

**Proof of Theorem 1.5(iii).** From (6.7), we get that for any prime  $p \equiv t \pmod{8}$

$$d(pn) = -\left(\frac{-2}{p}\right)_L \cdot d\left(\frac{n}{p}\right). \quad (6.25)$$

Replacing  $n$  by  $8n+t$ , we obtain

$$d(8pn+tp) = -\left(\frac{-2}{p}\right)_L d\left(\frac{8n+t}{p}\right). \quad (6.26)$$

Next substituting  $n$  by  $p^k n + r$  with  $p \nmid r$  in (6.26), we obtain

$$d\left(8\left(p^{k+1}n + pr + \frac{tp-1}{8}\right) + 1\right) = -\left(\frac{-2}{p}\right)_L d\left(8\left(p^{k-1}n + \frac{8r+t-p}{8p}\right) + 1\right). \quad (6.27)$$

Note that  $\frac{tp-1}{8}$  and  $\frac{8r+t-p}{8p}$  are integers. Using (6.27) and (6.5), we get

$$B_{3,2}\left(p^{k+1}n + pr + \frac{tp-1}{8}\right) \equiv -\left(\frac{-2}{p}\right)_L B_{3,2}\left(p^{k-1}n + \frac{8r+t-p}{8p}\right) \pmod{3}. \quad (6.28)$$

$\square$

**Proof of Corollary 1.6(iii).** Let  $p$  be a prime such that  $p \equiv t \pmod{8}$ . Choose a non negative integer  $r$  such that  $8r+t = p^{2k-1}$ . Substituting  $k$  by  $2k-1$  in (6.28), we obtain

$$B_{3,2}\left(p^{2k}n + \frac{p^{2k}-1}{8}\right) \equiv -\left(\frac{2}{p}\right)_L B_{3,2}\left(p^{2k-2}n + \frac{p^{2k-2}-1}{4}\right)$$

$$\equiv \dots \equiv (-1)^k \left(\frac{2}{p}\right)_L^k B_{3,2}(n) \pmod{3}.$$

□

7. CONGRUENCES FOR  $B_{7,2}(n)$ 

**Proof of Theorem 1.7(i).** From (1.1) and (3.2), we get

$$\sum_{n=0}^{\infty} B_{7,2}(n)q^n \equiv \frac{(q^7; q^7)_{\infty}(q^2; q^2)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty}^5 (q^2; q^2)_{\infty} \pmod{7}. \quad (7.1)$$

Define

$$\sum_{n=0}^{\infty} c(n)q^n := f_1^5 f_2 = (q; q)_{\infty}^5 (q^2; q^2)_{\infty}. \quad (7.2)$$

In 1959, Newman [15] proved that if  $p$  is a prime with  $p \equiv 1 \pmod{24}$ , then

$$c\left(pn + \frac{7(p-1)}{24}\right) = c\left(\frac{7(p-1)}{24}\right) c(n) - p^2 \cdot c\left(\frac{n}{p} - \frac{7(p-1)}{24p}\right). \quad (7.3)$$

Therefore, if  $p \nmid (24n + 7)$ , then  $\left(\frac{n}{p} - \frac{7(p-1)}{24p}\right)$  is not an integer and

$$c\left(\frac{n}{p} - \frac{7(p-1)}{24p}\right) = 0. \quad (7.4)$$

From (7.3) and (7.4) we get that if  $p \nmid (24n + 7)$ , then

$$c\left(pn + \frac{7(p-1)}{24}\right) = c\left(\frac{7(p-1)}{24}\right) c(n). \quad (7.5)$$

Thus, if  $p \nmid (24n + 7)$  and  $c\left(\frac{7(p-1)}{24}\right) \equiv 0 \pmod{7}$ , then for  $n \geq 0$ ,

$$c\left(pn + \frac{7(p-1)}{24}\right) \equiv 0 \pmod{7}. \quad (7.6)$$

Replacing  $n$  by  $pn + \frac{7(p-1)}{24}$  in (7.3), we get

$$c\left(p^2n + \frac{7(p^2-1)}{24}\right) = c\left(\frac{7(p-1)}{24}\right) c\left(pn + \frac{7(p-1)}{24}\right) - p^2 \cdot c(n). \quad (7.7)$$

Observe that from (7.7) if  $c\left(\frac{7(p-1)}{24}\right) \equiv 0 \pmod{7}$ , then for  $n \geq 0$ , we get

$$c\left(p^2n + \frac{7(p^2-1)}{24}\right) \equiv -p^2 \cdot c(n) \pmod{7}. \quad (7.8)$$



From (7.8) and by using mathematical induction, we get that if  $c\left(\frac{7(p-1)}{24}\right) \equiv 0 \pmod{7}$ , then for  $n \geq 0$ , and  $\alpha \geq 0$ , we get

$$c\left(p^{2\alpha}n + \frac{7(p^{2\alpha} - 1)}{24}\right) \equiv (-p^{2\cdot})^\alpha c(n) \pmod{7}. \quad (7.9)$$

Replacing  $n$  by  $pn + \frac{7(p-1)}{24}$  in (7.9) and using (7.6), we obtain that if  $p \nmid (24n + 7)$  and  $c\left(\frac{7(p-1)}{24}\right) \equiv 0 \pmod{7}$ , then for  $n \geq 0$  and  $\alpha \geq 0$ , we get

$$c\left(p^{2\alpha+1}n + \frac{7(p^{2\alpha+1} - 1)}{24}\right) \equiv 0 \pmod{7}. \quad (7.10)$$

From (7.2) and (7.1), we obtain that for  $n \geq 0$ ,

$$B_{7,2}(n) \equiv c(n) \pmod{7}. \quad (7.11)$$

Thus, combining (7.11) and (7.10), we obtain the result of Theorem.  $\square$

## 8. CONGRUENCES FOR $B_{11,2}(n)$

**Proof of Theorem 1.7(ii).** From (1.1) and (3.2), we get

$$\sum_{n=0}^{\infty} B_{11,2}(n)q^n = \frac{(q^{11}; q^{11})_{\infty}(q^2; q^2)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty}^9 (q^2; q^2)_{\infty} \pmod{11}. \quad (8.1)$$

Define

$$f_1^9 f_2 = (q; q)_{\infty}^9 (q^2; q^2)_{\infty} := \sum_{n=0}^{\infty} f(n)q^n. \quad (8.2)$$

In 1959, Newman [15] proved that if  $p$  is a prime with  $p \equiv 1 \pmod{24}$ , then

$$f\left(pn + \frac{11(p-1)}{24}\right) = f\left(\frac{11(p-1)}{24}\right) f(n) - p^4 \cdot f\left(\frac{n}{p} - \frac{11(p-1)}{24p}\right). \quad (8.3)$$

Therefore, if  $p \nmid (24n + 1)$ , then  $\left(\frac{n}{p} - \frac{11(p-1)}{24p}\right)$  is not an integer and

$$f\left(\frac{n}{p} - \frac{11(p-1)}{24p}\right) = 0. \quad (8.4)$$

From (8.3) and (8.4) we get that if  $p \nmid (24n + 1)$ , then

$$f\left(pn + \frac{11(p-1)}{24}\right) = f\left(\frac{11(p-1)}{24}\right) f(n). \quad (8.5)$$

Thus, if  $p \nmid (24n + 11)$  and  $f\left(\frac{11(p-1)}{24}\right) \equiv 0 \pmod{11}$ , then for  $n \geq 0$ ,

$$f\left(pn + \frac{11(p-1)}{24}\right) \equiv 0 \pmod{11}. \quad (8.6)$$

Replacing  $n$  by  $pn + \frac{11(p-1)}{24}$  in (8.3), we get

$$f\left(p^2n + \frac{11(p^2-1)}{24}\right) = f\left(\frac{11(p-1)}{24}\right) f\left(pn + \frac{11(p-1)}{24}\right) - p^4 \cdot f(n). \quad (8.7)$$

Observe that from (8.7) if  $f\left(\frac{11(p-1)}{24}\right) \equiv 0 \pmod{11}$ , then for  $n \geq 0$ , we get

$$f\left(p^2n + \frac{11(p^2-1)}{24}\right) \equiv -p^4 \cdot f(n) \pmod{11}. \quad (8.8)$$

From (8.8) and by using mathematical induction, we get that if  $f\left(\frac{11(p-1)}{24}\right) \equiv 0 \pmod{11}$ , then for  $n \geq 0$ , and  $\alpha \geq 0$ , we get

$$f\left(p^{2\alpha}n + \frac{11(p^{2\alpha}-1)}{24}\right) \equiv (-p^4)^\alpha \cdot f(n) \pmod{11}. \quad (8.9)$$

Replacing  $n$  by  $pn + \frac{11(p-1)}{24}$  in (8.9) and using (8.6), we obtain that if  $p \nmid (24n + 11)$  and  $f\left(\frac{11(p-1)}{24}\right) \equiv 0 \pmod{11}$ , then for  $n \geq 0$  and  $\alpha \geq 0$ , we get

$$f\left(p^{2\alpha+1}n + \frac{11(p^{2\alpha+1}-1)}{24}\right) \equiv 0 \pmod{11}. \quad (8.10)$$

From (8.2) and (8.1), we obtain that for  $n \geq 0$ ,

$$B_{11,2}(n) \equiv f(n) \pmod{11}. \quad (8.11)$$

Thus, combining (8.11) and (8.10), we obtain the result of Theorem.  $\square$

## 9. CONGRUENCES FOR $B_{13,2}(n)$

**Proof of Theorem 1.7(iii).** From (1.1) and (3.2), we get

$$\sum_{n=0}^{\infty} B_{13,2}(n)q^n = \frac{(q^{13}; q^{13})_{\infty} (q^2; q^2)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty}^{11} (q^2; q^2)_{\infty} \pmod{13}. \quad (9.1)$$

Define

$$f_1^{11} f_2 = (q; q)_{\infty}^{11} (q^2; q^2)_{\infty} := \sum_{n=0}^{\infty} g(n)q^n. \quad (9.2)$$

In 1959, Newman [15] proved that if  $p$  is a prime with  $p \equiv 1 \pmod{24}$ , then

$$g\left(pn + \frac{13(p-1)}{24}\right) = g\left(\frac{13(p-1)}{24}\right)g(n) - p^5 \cdot g\left(\frac{n}{p} - \frac{13(p-1)}{24p}\right). \quad (9.3)$$

Therefore, if  $p \nmid (24n + 13)$ , then  $\left(\frac{n}{p} - \frac{13(p-1)}{24p}\right)$  is not an integer and

$$g\left(\frac{n}{p} - \frac{13(p-1)}{24p}\right) = 0. \quad (9.4)$$

From (9.3) and (9.4) we get that if  $p \nmid (24n + 13)$ , then

$$g\left(pn + \frac{13(p-1)}{24}\right) = g\left(\frac{13(p-1)}{24}\right)g(n). \quad (9.5)$$

Thus, if  $p \nmid (24n + 13)$  and  $g\left(\frac{13(p-1)}{24}\right) \equiv 0 \pmod{13}$ , then for  $n \geq 0$ ,

$$g\left(pn + \frac{13(p-1)}{24}\right) \equiv 0 \pmod{13}. \quad (9.6)$$

Replacing  $n$  by  $pn + \frac{13(p-1)}{24}$  in (9.3), we get

$$g\left(p^2n + \frac{13(p^2-1)}{24}\right) = g\left(\frac{13(p-1)}{24}\right)g\left(pn + \frac{13(p-1)}{24}\right) - p^5 \cdot g(n). \quad (9.7)$$

Observe that from (9.7) if  $g\left(\frac{11(p-1)}{24}\right) \equiv 0 \pmod{13}$ , then for  $n \geq 0$ , we get

$$g\left(p^2n + \frac{13(p^2-1)}{24}\right) \equiv -p^5 \cdot g(n) \pmod{13}. \quad (9.8)$$

From (9.8) and by using mathematical induction, we get that if  $g\left(\frac{13(p-1)}{24}\right) \equiv 0 \pmod{13}$ , then for  $n \geq 0$ , and  $\alpha \geq 0$ , we get

$$g\left(p^{2\alpha}n + \frac{11(p^{2\alpha}-1)}{24}\right) \equiv (-p^5)^\alpha g(n) \pmod{13}. \quad (9.9)$$

Replacing  $n$  by  $pn + \frac{13(p-1)}{24}$  in (9.9) and using (9.6), we obtain that if  $p \nmid (24n + 13)$  and  $g\left(\frac{13(p-1)}{24}\right) \equiv 0 \pmod{13}$ , then for  $n \geq 0$  and  $\alpha \geq 0$ , we get

$$g\left(p^{2\alpha+1}n + \frac{13(p^{2\alpha+1}-1)}{24}\right) \equiv 0 \pmod{13}. \quad (9.10)$$

From (9.2) and (8.1), we obtain that for  $n \geq 0$ ,

$$B_{13,2}(n) \equiv e(n) \pmod{13}. \quad (9.11)$$

Thus, combining (9.11) and (9.10), we obtain the result of Theorem.  $\square$

## 10. APPLICATION OF NEWMANN IDENTITY II

We are going to state a result of Newman [ [16], Theorem 3 ] which play an important role in the proof of our Theorems. We state his result as a lemma. We need the following notations to state the lemma. Let  $r, s$  be integers such that  $r, s \neq 0, r \not\equiv s \pmod{2}$ . Set

$$\phi(\tau) = \prod_{n=1}^{\infty} (1 - x^n)^r (1 - x^{nq})^s = \sum_{n=0}^{\infty} a(n)x^n$$

$$\epsilon = \frac{r+s}{2}, t = \frac{r+sq}{24}, \Delta = t(p^2 - 1) = \frac{(r+sq)(p^2-1)}{24}, \theta = (-1)^{\frac{1}{2}-\epsilon} 2q^s.$$

**Lemma 10.1.** *With the notations defined as above, the coefficients  $a(n)$  of  $\phi(\tau)$  satisfy*

$$a(np^2 + \Delta) = \gamma_0(n)a(n) - p^{2\epsilon-2} a\left(\frac{n-\Delta}{p^2}\right), \quad (10.1)$$

where

$$\gamma_0(n) = p^{2\epsilon-2} c - \left(\frac{\theta}{p}\right)_L p^{\epsilon-3/2} \left(\frac{n-\Delta}{p}\right)_L, \quad (10.2)$$

where  $c$  is a constant.

## 11. CONGRUENCES FOR $B_{4,3}(n)$

**Proof of Theorem 1.8.** Note that

$$\sum_{n=0}^{\infty} B_{4,3}(n)q^n = \frac{(q^4; q^4)_{\infty} (q^3; q^3)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty}^2 (q^3; q^3)_{\infty} \pmod{2}. \quad (11.1)$$

Let  $a_1(n)$  be defined by (1.13). Substituting  $q = 3, r = 2, s = 1$  in Lemma 10.1, we obtain

$$a_1\left(p^2 n + \frac{5(p^2-1)}{24}\right) = \gamma_1(n)a_1(n) - p \cdot a_1\left(\frac{n - \frac{5(p^2-1)}{24}}{p^2}\right), \quad (11.2)$$

where

$$\gamma_1(n) = p \cdot c - \left(\frac{-6}{p}\right)_L \left(\frac{n - \frac{5(p^2-1)}{24}}{p}\right)_L. \quad (11.3)$$

If we put  $n = 0$  in (11.2) and using the facts that  $a_1(0) = 1$  and  $a_1\left(\frac{-\frac{5(p^2-1)}{24}}{p^2}\right) = 0$ , we get

$$\gamma_1(0) = a_1\left(\frac{5(p^2-1)}{24}\right). \quad (11.4)$$

Further taking  $n = 0$  in (11.3) and using (11.4), we obtain

$$pc = w_1(p), \quad (11.5)$$

where  $w_1(p) := a_1 \left( \frac{5(p^2-1)}{24} \right) + \left( \frac{-6}{p} \right)_L \left( \frac{-5(p^2-1)}{24p} \right)_L$ . Combining (11.5), (11.2) and (11.3) together, we get

$$a_1 \left( p^2n + \frac{5(p^2-1)}{24} \right) = \left[ w_1(p) - \left( \frac{-6}{p} \right)_L \left( \frac{n - \frac{5(p^2-1)}{24}}{p} \right)_L \right] a_1(n) - p \cdot a_1 \left( \frac{n - \frac{5(p^2-1)}{24}}{p^2} \right). \quad (11.6)$$

Replacing  $n$  by  $pn + \frac{5(p^2-1)}{24}$  in (11.6), we obtain

$$a_1 \left( p^3n + \frac{5(p^4-1)}{24} \right) = w_1(p)a_1 \left( pn + \frac{5(p^2-1)}{24} \right) - p \cdot a_1 \left( \frac{n}{p} \right). \quad (11.7)$$

If  $w_1(p) \equiv 0 \pmod{2}$ , then

$$a_1 \left( p^3n + \frac{5(p^4-1)}{24} \right) \equiv -p \cdot a_1 \left( \frac{n}{p} \right) \pmod{2}. \quad (11.8)$$

Replacing  $n$  by  $pn$  we have,

$$a_1 \left( p^4n + \frac{5(p^4-1)}{24} \right) \equiv -p \cdot a_1(n) \pmod{2}. \quad (11.9)$$

Using mathematical induction in (11.9), we obtain for  $n, k \geq 0$ ,

$$a_1 \left( p^{4k}n + \frac{5(p^{4k}-1)}{24} \right) \equiv (-p)^k \cdot a_1(n) \pmod{2}. \quad (11.10)$$

Congruence (11.8) implies that If  $w_1(p) \equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$a_1 \left( p^3n + \frac{5(p^4-1)}{24} \right) \equiv 0 \pmod{2}. \quad (11.11)$$

Replacing  $n$  by  $p^3n + \frac{5(p^4-1)}{24}$  in (11.10) and using (11.11), we get that if  $w_1(p) \equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$a_1 \left( p^{4k+3}n + \frac{5(p^{4k+4}-1)}{24} \right) \equiv 0 \pmod{2}. \quad (11.12)$$

From (11.13), (11.1) and (11.12) we obtain that, if  $w_1(p) \equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$B_{4,3} \left( p^{4k+3}n + \frac{5(p^{4k+4}-1)}{24} \right) \equiv 0 \pmod{2}. \quad (11.13)$$

Further, replacing  $n$  by  $p^2n + \frac{5p(p^2-1)}{24}$  in (11.7), we get

$$a_1 \left( p^5n + \frac{5(p^6-1)}{24} \right) = w_1(p)a_1 \left( p^3n + \frac{5(p^4-1)}{24} \right) - p \cdot a_1 \left( pn + \frac{5(p^2-1)}{24} \right). \quad (11.14)$$

Combining (11.14) and (11.7) together, we get

$$a_1 \left( p^5 n + \frac{5(p^6 - 1)}{24} \right) = (w_1^2(p) - p) a_1 \left( pn + \frac{5(p^2 - 1)}{24} \right) - w_1(p) \cdot p \cdot a_1 \left( \frac{n}{p} \right). \quad (11.15)$$

Note that if  $w_1(p) \not\equiv 0 \pmod{2}$ , then  $w_1(p) \equiv 1 \pmod{2}$  and hence  $(w_1^2(p) - p) \equiv 0 \pmod{2}$ . Thus, if  $w_1(p) \not\equiv 0 \pmod{2}$ , then (11.15) will be

$$a_1 \left( p^5 n + \frac{5(p^6 - 1)}{24} \right) \equiv -w_1(p) \cdot p \cdot a_1 \left( \frac{n}{p} \right) \pmod{2}. \quad (11.16)$$

Therefore, if  $p \nmid n$ , then  $a_1 \left( \frac{n}{p} \right) = 0$  and

$$a_1 \left( p^5 n + \frac{5(p^6 - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (11.17)$$

Replacing  $n$  by  $pn$  in (11.16), we obtain

$$a_1 \left( p^6 n + \frac{5(p^6 - 1)}{24} \right) \equiv -w_1(p) \cdot p \cdot a_1(n) \pmod{2}. \quad (11.18)$$

By using mathematical induction in (11.18), we get

$$a_1 \left( p^{6k} n + \frac{5(p^{6k} - 1)}{24} \right) \equiv (-w_1(p) \cdot p)^k \cdot a_1(n) \pmod{2}. \quad (11.19)$$

Replacing  $n$  by  $p^5 n + \frac{5(p^6 - 1)}{24}$  in (11.19) and using (11.17), it follows that if  $w_1(p) \not\equiv 0 \pmod{2}$  and  $p \nmid n$ , we get

$$a_1 \left( p^{6k+5} n + \frac{5(p^{6k+6} - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (11.20)$$

From (11.20) and (11.1) we obtain that, if  $w_1(p) \not\equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$B_{4,3} \left( p^{6k+5} n + \frac{5(p^{6k+6} - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (11.21)$$

Note that  $\left( \frac{-6}{p} \right)_L \left( \frac{n - \frac{5(p^2 - 1)}{24}}{p} \right)_L = \left( \frac{-6n - 1 + \frac{(p^2 - 1)}{4}}{p} \right)_L$ . From (11.6) it follows that if  $w_1(p) \equiv \left( \frac{-6n - 1 + \frac{(p^2 - 1)}{4}}{p} \right)_L \pmod{2}$ , we obtain

$$a_1 \left( p^2 n + \frac{5(p^2 - 1)}{24} \right) \equiv -p \cdot a_1 \left( \frac{n - \frac{5(p^2 - 1)}{24}}{p^2} \right) \pmod{2}. \quad (11.22)$$

Observe that if  $\left(\frac{-6n-1+\frac{(p^2-1)}{4}}{p}\right)_L \not\equiv 0 \pmod{2}$ , then  $p \nmid (24n+5)$  and hence  $\left(\frac{n-\frac{5(p^2-1)}{24}}{p}\right)$  and  $\left(\frac{n-\frac{5(p^2-1)}{24}}{p^2}\right)$  are not integers. Thus, for  $n \geq 0$ , from (11.22) we have

$$a_1\left(p^2n + \frac{5(p^2-1)}{24}\right) \equiv 0 \pmod{2}. \quad (11.23)$$

Replacing  $n$  by  $p^2n + \frac{5(p^2-1)}{24}$  in (11.19) and using (11.23), we obtain that if  $w_1(p) \not\equiv 0 \pmod{2}$ , and  $w_1(p) \equiv \left(\frac{-6n-1+\frac{(p^2-1)}{4}}{p}\right)_L \pmod{2}$ , then

$$a_1\left(p^{6k+2}n + \frac{5(p^{6k+2}-1)}{24}\right) \equiv 0 \pmod{2}. \quad (11.24)$$

Combining (11.1) and (11.24) together, we get

$$B_{4,3}\left(p^{6k+2}n + \frac{5(p^{6k+2}-1)}{24}\right) \equiv 0 \pmod{2}. \quad (11.25)$$

□

## 12. CONGRUENCES FOR $B_{8,3}(n)$

**Proof of Theorem 1.9 .** Note that

$$\sum_{n=0}^{\infty} B_{8,3}(n)q^n = \frac{(q^8; q^8)_{\infty}(q^3; q^3)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty}^6 (q^3; q^3)_{\infty} \pmod{2}. \quad (12.1)$$

Let  $a_2(n)$  be defined by (1.13). Substituting  $q = 3, r = 6, s = 1$  in Lemma 10.1, we obtain

$$a_2\left(p^2n + \frac{3(p^2-1)}{8}\right) = \gamma_2(n)a_2(n) - p^5 \cdot a_2\left(\frac{n - \frac{3(p^2-1)}{8}}{p^2}\right), \quad (12.2)$$

where

$$\gamma_2(n) = p^5 \cdot c - p^2 \cdot \left(\frac{-6}{p}\right)_L \left(\frac{n - \frac{3(p^2-1)}{8}}{p}\right)_L. \quad (12.3)$$

If we put  $n = 0$  in (12.2) and using the facts that  $a_2(0) = 1$  and  $a_2\left(\frac{-\frac{3(p^2-1)}{8}}{p^2}\right) = 0$ , we get

$$\gamma_2(0) = a_2\left(\frac{3(p^2-1)}{8}\right). \quad (12.4)$$

Further taking  $n = 0$  in (12.3) and using (12.4), we obtain

$$p^5c = w_2(p), \quad (12.5)$$

where  $w_2(p) := a_2\left(\frac{3(p^2-1)}{8}\right) + p^2 \cdot \left(\frac{-6}{p}\right)_L \left(\frac{-3(p^2-1)}{8p}\right)_L$ . Combining (12.5), (12.2) and (12.3) together, we get

$$a_2\left(p^2n + \frac{3(p^2-1)}{8}\right) = \left[ w_2(p) - p^2 \cdot \left(\frac{-6}{p}\right)_L \left(\frac{n - \frac{3(p^2-1)}{8}}{p}\right)_L \right] a_2(n) - p^5 \cdot a_2\left(\frac{n - \frac{3(p^2-1)}{8}}{p^2}\right). \quad (12.6)$$

Replacing  $n$  by  $pn + \frac{3(p^2-1)}{8}$  in (12.6), we obtain

$$a_2\left(p^3n + \frac{3(p^4-1)}{8}\right) = w_2(p)a_2\left(pn + \frac{3(p^2-1)}{8}\right) - p^5 \cdot a_2\left(\frac{n}{p}\right). \quad (12.7)$$

If  $w_2(p) \equiv 0 \pmod{2}$ , then

$$a_2\left(p^3n + \frac{3(p^4-1)}{8}\right) \equiv -p^5 \cdot a_2\left(\frac{n}{p}\right) \pmod{2}. \quad (12.8)$$

Replacing  $n$  by  $pn$  we have,

$$a_2\left(p^4n + \frac{3(p^4-1)}{8}\right) \equiv -p^5 \cdot a_2(n) \pmod{2}. \quad (12.9)$$

Using mathematical induction in (12.9), we obtain for  $n, k \geq 0$ ,

$$a_2\left(p^{4k}n + \frac{3(p^{4k}-1)}{8}\right) \equiv (-p^5)^k \cdot a_2(n) \pmod{2}. \quad (12.10)$$

Congruence (12.8) implies that If  $w_2(p) \equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$a_2\left(p^3n + \frac{3(p^4-1)}{8}\right) \equiv 0 \pmod{2}. \quad (12.11)$$

Replacing  $n$  by  $p^3n + \frac{3(p^4-1)}{8}$  in (12.10) and using (12.11), we get that if  $w_2(p) \equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$a_2\left(p^{4k+3}n + \frac{3(p^{4k+4}-1)}{8}\right) \equiv 0 \pmod{2}. \quad (12.12)$$

From (12.1) and (12.12) we obtain that, if  $w_2(p) \equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$B_{8,3}\left(p^{4k+3}n + \frac{3(p^{4k+4}-1)}{8}\right) \equiv 0 \pmod{2}. \quad (12.13)$$

Replacing  $n$  by  $p^2n + \frac{3p(p^2-1)}{8}$  in (12.7), we get

$$a_2\left(p^5n + \frac{3(p^6-1)}{8}\right) = w_2(p)a_2\left(p^3n + \frac{3(p^4-1)}{8}\right) - p^5 \cdot a_2\left(pn + \frac{3(p^2-1)}{8}\right). \quad (12.14)$$



Combining (12.14) and (12.7) together, we get

$$a_2 \left( p^5 n + \frac{3(p^6 - 1)}{8} \right) = (w_2(p) - p^5) a_2 \left( pn + \frac{3(p^2 - 1)}{8} \right) - w_2(p) \cdot p^5 \cdot a_2 \left( \frac{n}{p} \right). \quad (12.15)$$

Note that if  $w_2(p) \not\equiv 0 \pmod{2}$ , then  $w_2(p) \equiv 1 \pmod{2}$  and hence  $(w_2(p) - p^5) \equiv 0 \pmod{2}$ . Thus, if  $w_2(p) \not\equiv 0 \pmod{2}$ , then (12.15) will be

$$a_2 \left( p^5 n + \frac{3(p^6 - 1)}{8} \right) \equiv -w_2(p) \cdot p^5 \cdot a_2 \left( \frac{n}{p} \right) \pmod{2}. \quad (12.16)$$

Therefore, if  $p \nmid n$ , then  $a_2 \left( \frac{n}{p} \right) = 0$  and

$$a_2 \left( p^5 n + \frac{3(p^6 - 1)}{8} \right) \equiv 0 \pmod{2}. \quad (12.17)$$

Replacing  $n$  by  $pn$  in (12.16), we obtain

$$a_2 \left( p^6 n + \frac{3(p^6 - 1)}{8} \right) \equiv -w_2(p) \cdot p^5 \cdot a_2(n) \pmod{2}. \quad (12.18)$$

By using mathematical induction in (12.18), we get

$$a_2 \left( p^{6k} n + \frac{3(p^{6k} - 1)}{8} \right) \equiv (-w_2(p) \cdot p^5)^k \cdot a_2(n) \pmod{2}. \quad (12.19)$$

Replacing  $n$  by  $p^5 n + \frac{3(p^6 - 1)}{8}$  in (12.19) and using (12.17), it follows that if  $w_2(p) \not\equiv 0 \pmod{2}$  and  $p \nmid n$ , we get

$$a_2 \left( p^{6k+5} n + \frac{3(p^{6k+6} - 1)}{8} \right) \equiv 0 \pmod{2}. \quad (12.20)$$

From (12.20) and (12.1) we obtain that, if  $w_2(p) \not\equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$B_{8,3} \left( p^{6k+5} n + \frac{3(p^{6k+6} - 1)}{8} \right) \equiv 0 \pmod{2}. \quad (12.21)$$

Note that  $\left( \frac{-6}{p} \right)_L \left( \frac{n - \frac{3(p^2 - 1)}{8}}{p} \right)_L = \left( \frac{-6n - 2 + \frac{(p^2 - 1)}{4}}{p} \right)_L$ . From (12.6) it follows that if  $w_2(p) \equiv p^2 \cdot \left( \frac{-6n - 2 + \frac{(p^2 - 1)}{4}}{p} \right)_L \pmod{2}$  we get

$$a_2 \left( p^2 n + \frac{3(p^2 - 1)}{8} \right) \equiv -p^5 \cdot a_2 \left( \frac{n - \frac{3(p^2 - 1)}{8}}{p^2} \right) \pmod{2}. \quad (12.22)$$

Observe that  $\left(\frac{-6n-2+\frac{(p^2-1)}{4}}{p}\right)_L \not\equiv 0 \pmod{2}$ , then  $p \nmid (8n+3)$  and hence  $\left(\frac{n-\frac{3(p^2-1)}{8}}{p}\right)$  and  $\left(\frac{n-\frac{3(p^2-1)}{8}}{p^2}\right)$  are not integers. Thus, for  $n \geq 0$ , from (12.22) we have

$$a_2\left(p^2n + \frac{3(p^2-1)}{8}\right) \equiv 0 \pmod{2}. \quad (12.23)$$

Replacing  $n$  by  $p^2n + \frac{3(p^2-1)}{8}$  in (12.19) and using (12.23), we obtain that if  $w_2(p) \not\equiv 0 \pmod{2}$ , and  $w_2(p) \equiv p^2 \cdot \left(\frac{-6n-2+\frac{(p^2-1)}{4}}{p}\right)_L \pmod{2}$ , then

$$a_2\left(p^{6k+2}n + \frac{3(p^{6k+2}-1)}{8}\right) \equiv 0 \pmod{2}. \quad (12.24)$$

Combining (12.1) and (12.24) together, we get

$$B_{8,3}\left(p^{6k+2}n + \frac{3(p^{6k+2}-1)}{8}\right) \equiv 0 \pmod{2}. \quad (12.25)$$

□

### 13. CONGRUENCES FOR $B_{4,5}(n)$

**Proof of Theorem 1.10.** Note that

$$\sum_{n=0}^{\infty} B_{4,5}(n)q^n = \frac{(q^4; q^4)_{\infty}(q^5; q^5)_{\infty}}{(q; q)_{\infty}^2} \equiv (q; q)_{\infty}^2 (q^5; q^5)_{\infty} \pmod{2}. \quad (13.1)$$

Let  $a_3(n)$  be defined by (1.23). Substituting  $q = 5, r = 2, s = 1$  in Lemma 10.1, we obtain

$$a_3\left(p^2n + \frac{7(p^2-1)}{24}\right) = \gamma_3(n)a_3(n) - p \cdot a_3\left(\frac{n - \frac{7(p^2-1)}{24}}{p^2}\right), \quad (13.2)$$

where

$$\gamma_3(n) = p \cdot c - \left(\frac{-10}{p}\right)_L \left(\frac{n - \frac{7(p^2-1)}{24}}{p}\right)_L. \quad (13.3)$$

If we put  $n = 0$  in (13.2) and using the facts that  $a_3(0) = 1$  and  $a_3\left(\frac{-\frac{7(p^2-1)}{24}}{p^2}\right) = 0$ , we get

$$\gamma_3(0) = a_3\left(\frac{7(p^2-1)}{24}\right). \quad (13.4)$$

Further taking  $n = 0$  in (13.3) and using (13.4), we obtain

$$pc = w_3(p), \quad (13.5)$$

where  $w_3(p) := a_3\left(\frac{7(p^2-1)}{24}\right) + \left(\frac{-10}{p}\right)_L \left(\frac{-7(p^2-1)}{24p}\right)_L$ . Combining (13.5), (13.2) and (13.3) together, we get

$$a_3\left(p^2n + \frac{7(p^2-1)}{24}\right) = \left[ w_3(p) - \left(\frac{-10}{p}\right)_L \left(\frac{n - \frac{7(p^2-1)}{24}}{p}\right)_L \right] a_3(n) - p \cdot a_3\left(\frac{n - \frac{7(p^2-1)}{24}}{p^2}\right). \quad (13.6)$$

Replacing  $n$  by  $pn + \frac{7(p^2-1)}{24}$  in (13.6), we obtain

$$a_3\left(p^3n + \frac{7(p^4-1)}{24}\right) = w_3(p)a_3\left(pn + \frac{7(p^2-1)}{24}\right) - p \cdot a_3\left(\frac{n}{p}\right). \quad (13.7)$$

If  $w_3(p) \equiv 0 \pmod{2}$ , then

$$a_3\left(p^3n + \frac{7(p^4-1)}{24}\right) \equiv -p \cdot a_3\left(\frac{n}{p}\right) \pmod{2}. \quad (13.8)$$

Replacing  $n$  by  $pn$  we have,

$$a_3\left(p^4n + \frac{7(p^4-1)}{24}\right) \equiv -p \cdot a_3(n) \pmod{2}. \quad (13.9)$$

Using mathematical induction in (13.9), we obtain for  $n, k \geq 0$ ,

$$a_3\left(p^{4k}n + \frac{7(p^{4k}-1)}{24}\right) \equiv (-p)^k \cdot a_3(n) \pmod{2}. \quad (13.10)$$

Congruence (13.8) implies that If  $w_3(p) \equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$a_3\left(p^3n + \frac{7(p^4-1)}{24}\right) \equiv 0 \pmod{2}. \quad (13.11)$$

Replacing  $n$  by  $p^3n + \frac{7(p^4-1)}{24}$  in (13.10) and using (13.11), we get that if  $w_2(p) \equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$a_3\left(p^{4k+3}n + \frac{7(p^{4k+4}-1)}{24}\right) \equiv 0 \pmod{2}. \quad (13.12)$$

From (13.1) and (13.12) we obtain that, if  $w_2(p) \equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$B_{4,5}\left(p^{4k+3}n + \frac{7(p^{4k+4}-1)}{24}\right) \equiv 0 \pmod{2}. \quad (13.13)$$

Replacing  $n$  by  $p^2n + \frac{7(p^2-1)}{24}$  in (13.7), we get

$$a_3\left(p^5n + \frac{7(p^6-1)}{24}\right) = w_3(p)a_3\left(p^3n + \frac{7(p^4-1)}{24}\right) - p \cdot a_3\left(pn + \frac{7(p^2-1)}{24}\right). \quad (13.14)$$

Combining (13.14) and (13.7) together, we get

$$a_3\left(p^5n + \frac{7(p^6-1)}{24}\right) = (w_3^2(p) - p) a_3\left(pn + \frac{7(p^2-1)}{24}\right) - w_3(p) \cdot p \cdot a_3\left(\frac{n}{p}\right). \quad (13.15)$$

Note that if  $w_3(p) \not\equiv 0 \pmod{2}$ , then  $w_3(p) \equiv 1 \pmod{2}$  and hence  $(w_3^2(p) - p) \equiv 0 \pmod{2}$ . Thus, if  $w_3(p) \not\equiv 0 \pmod{2}$ , then (13.15) will be

$$a_3 \left( p^5 n + \frac{7(p^6 - 1)}{24} \right) \equiv -w_3(p) \cdot p \cdot a_3 \left( \frac{n}{p} \right) \pmod{2}. \quad (13.16)$$

Therefore, if  $p \nmid n$ , then  $a_3 \left( \frac{n}{p} \right) = 0$  and

$$a_3 \left( p^5 n + \frac{7(p^6 - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (13.17)$$

Replacing  $n$  by  $pn$  in (13.16), we obtain

$$a_3 \left( p^6 n + \frac{7(p^6 - 1)}{24} \right) \equiv -w_3(p) \cdot p \cdot a_3(n) \pmod{2}. \quad (13.18)$$

By using mathematical induction in (13.18), we get

$$a_3 \left( p^{6k} n + \frac{7(p^{6k} - 1)}{24} \right) \equiv (-w_3(p) \cdot p)^k \cdot a_3(n) \pmod{2}. \quad (13.19)$$

Replacing  $n$  by  $p^5 n + \frac{7(p^6 - 1)}{24}$  in (13.19) and using (13.17), it follows that if  $w_3(p) \not\equiv 0 \pmod{2}$  and  $p \nmid n$ , we get

$$a_3 \left( p^{6k+5} n + \frac{7(p^{6k+6} - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (13.20)$$

From (13.20) and (13.1) we obtain that, if  $w_3(p) \not\equiv 0 \pmod{2}$  and  $p \nmid n$ , then

$$B_{4,5} \left( p^{6k+5} n + \frac{7(p^{6k+6} - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (13.21)$$

Note that  $\left( \frac{-10}{p} \right)_L \left( \frac{n - \frac{7(p^2 - 1)}{24}}{p} \right)_L = \left( \frac{-10n - 2 + \frac{11(p^2 - 1)}{12}}{p} \right)_L$ . From (13.6) it follows that if  $w_3(p) \equiv \left( \frac{-10n - 2 + \frac{11(p^2 - 1)}{12}}{p} \right)_L \pmod{2}$  we get

$$a_3 \left( p^2 n + \frac{7(p^2 - 1)}{24} \right) \equiv -p \cdot a_3 \left( \frac{n - \frac{7(p^2 - 1)}{24}}{p^2} \right) \pmod{2}. \quad (13.22)$$

Observe that  $\left( \frac{-10n - 2 + \frac{11(p^2 - 1)}{12}}{p} \right)_L \not\equiv 0 \pmod{2}$ , then  $p \nmid (24n + 7)$  and hence  $\left( \frac{n - \frac{7(p^2 - 1)}{24}}{p} \right)$  and  $\left( \frac{n - \frac{7(p^2 - 1)}{24}}{p^2} \right)$  are not integers. Thus, for  $n \geq 0$ , from (13.22) we have

$$a_3 \left( p^2 n + \frac{7(p^2 - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (13.23)$$

Replacing  $n$  by  $p^2n + \frac{7(p^2-1)}{24}$  in (13.19) and using (13.23), we obtain that if  $w_3(p) \not\equiv 0 \pmod{2}$ , and  $w_3(p) \equiv \left( \frac{-10n-2+\frac{11(p^2-1)}{12}}{p} \right)_L \pmod{2}$ , then

$$a_3 \left( p^{6k+2}n + \frac{7(p^{6k+2} - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (13.24)$$

Combining (13.1) and (13.24) together, we get

$$B_{4,5} \left( p^{6k+2}n + \frac{7(p^{6k+2} - 1)}{24} \right) \equiv 0 \pmod{2}. \quad (13.25)$$

□

**Data availability statement:** There is no data associated to our manuscript.

#### REFERENCES

- [1] J. P. Allouche and L. Goldmakher, *Mock character and Kronecker symbol*, J. Number Theory **192** (2018), 356-372.
- [2] R. Barman and A. Singh, *On mex-related partition functions of Andrews and Newman*, Res. Number Theory **7** (2021), Paper No. 53, 11 pp.
- [3] D.Q.J. Dou, *Congruences for  $(3, 11)$ -regular bipartitions modulo 11*, Ramanujan J. **40** (2016), 535-540.
- [4] B. Gordan and K. Ono, *Divisibility of certain partition functions by power of primes*, Ramanujan j. **1** (1997), 25-34.
- [5] A. Granville and K. Ono, *Defect zero  $p$ -blocks for finite simple groups*, Trans. Amer. Math. Soc. **348** (1996), no. 1, 331-347.
- [6] M. D. Hirschhorn and J. A. Sellers, *Parity results for partitions wherein each parts an odd number of times*, Bull. Aust. Math. Soc. **1** (2019), 51-55.
- [7] T. Kathiravan and S. N. Fathima, *On  $\ell$ -regular bipartitions modulo  $\ell$* , Ramanujan J. **44** (2017), 549-558.
- [8] T Kathiravan and K. Srilakshmi, *Some congruences for  $(s, t)$ -regular bipartitions modulo  $t$* , <https://arxiv.org/pdf/1908.06642>.
- [9] T. Kathiravan, K. Srinivas and U. K. Sangale, *Some congruences for  $(\ell, k)$  and  $(\ell, k, r)$ -regular partitions*, Preprint, <https://arxiv.org/abs/2303.13906>
- [10] N. Koblitz, *Introduction to elliptic curves and modular forms*, Springer-Verlag New York (1991).
- [11] B. L. S. Lin, *Arithmetic properties of bipartitions with even parts distinct*, Ramanujan J. **33** (2013), 269-279.
- [12] B. L. S. Lin, *Arithmetic of the 7-regular bipartition function modulo 3*, Ramanujan J. **37** (2015), 469-478.
- [13] B. L. S. Lin, *An infinite family of congruences modulo 3 for 13-regular bipartitions*, Ramanujan J. **39** (2016), 169-178.
- [14] Y. Martin, *Multiplicative  $\eta$ -quotients*, Trans. Am. Math. Soc. **348** (1996), 4825-4856.
- [15] M. Newman, *Modular forms whose coefficients possess multiplicative properties*, Ann. of Math. **70** (1959), 478-489.
- [16] M. Newman, *Modular forms whose coefficients possess multiplicative properties II*, Ann. of Math. **75** (1962), 242-250.

- [17] K. Ono, *The web of modularity: arithmetic of the coefficients of modular forms and  $q$ -series*, CBMS Regional Conference Series in Mathematics, 102, Amer. Math. Soc., Providence, RI, 2004.
- [18] The Sage Developers, *sagemath, the Sage Mathematics Software System (Version 8.1)*. <https://www.sagemath.org>
- [19] J. -P. Serre, *Divisibilité des coefficients des formes modularies de poids entier*, C. R. Acad. Sci. Paris (A), **279** (1974), 679-682.
- [20] J. -P. Serre, *Divisibilité de certaines fonctions arithmétiques*, in: Séminaire Delanga-Pisot-Poitou, Théorie Nr., **16** (1974), 1-28.
- [21] J. Tate, *Extensions of  $\mathbb{Q}$  un-ramified outside 2*, in: Arithmetic Geometry: Conference on Arithmetic Geometry with an Emphasis on Iwasawa Theory, Arizona State University, March 15 – 18, 1993, Vol. 174, No. 174, American Mathematical Society, Providence, 1994.
- [22] L. Wang, *Arithmetic properties of  $(k, l)$ -regular bipartitions*, Bull. Aust. Math. Soc. **95** (2017), 353-364.
- [23] E.X.W. Xia, *New congruence properties for Ramanujan's  $\phi$  function*, Proc. AMS. **149** (2021), 4985-4999.
- [24] E.X.W. Xia and O.X.M. Yao, *Arithmetic properties of  $(s, t)$ -regular bipartition functions*, J. Number Theory **171** (2017), 1-17.

NABIN KUMAR MEHER, DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF INFORMATION RAICHUR, GOVT. ENGINEERING COLLEGE CAMPUS, YERMARUS, RAICHUR, KARNATAKA, INDIA 584135.

*Email address:* mehernabin@gmail.com, nabinmeher@iiitr.ac.in