

Computing the degreewidth of a digraph is hard

Pierre Aboulker¹, Nacim Oijid², Robin Petit³, Mathis Rocton⁴, and Christopher-Lloyd Simon⁵

¹DIENS, École normale supérieure, CNRS, PSL University, Paris, France.

²Univ Lyon, CNRS, INSA Lyon, UCBL, Centrale Lyon, Univ Lyon 2, LIRIS, UMR5205, F-69622 Villeurbanne, France.

³Computer Science Department, Université Libre de Bruxelles, Brussels, Belgium.

⁴Algorithm and Complexity Group, TU Wien, Vienna, Austria.

⁵The Pennsylvania State University, Department of Mathematics

August 1, 2024

Abstract

Given a digraph, an ordering of its vertices defines a *backedge graph*, namely the undirected graph whose edges correspond to the arcs pointing backwards with respect to the order. The *degreewidth* of a digraph is the minimum over all ordering of the maximum degree of the backedge graph. We answer an open question by Keeney and Lokshantov [WG 2024], proving that it is NP-hard to determine whether an oriented graph has degreewidth at most 1, which settles the last open case for oriented graphs. We complement this result with a general discussion on parameters defined using backedge graphs and their relations to classical parameters.

Contents

1	Introduction	1
2	Complexity of Computing $\vec{\Delta}$	2
3	A Unifying Point of View	4
3.1	From Undirected to Directed Graphs	4
3.2	General Observations	5
3.3	Dichromatic Number and Directed Clique Number	6
3.4	Directed Vertex Cover is Feedback Vertex Set	6
3.5	Back to the Degreewidth	6
3.6	Optimal Linear Arrangement, Cutwidth and Bandwidth	8

1 Introduction

Throughout this paper, we consider digraphs D which, unless specified otherwise, will always be *simple*, namely they have no loops and for distinct vertices $u, v \in V(D)$ there is at most one arc from u to v (but D may contain an arc from u to v and an arc from v to u). The word *edge* always refers to an undirected graph whereas *arc* always refers to a directed one.

Given a digraph D , to a total ordering \prec on $V(D)$ we associate the undirected graph D^\prec on the vertex set $V(D)$ whose edges correspond to the arcs $uv \in A(D)$ such that $v \prec u$. We call it the *backedge graph* of D with respect to \prec .

The *degreewidth* $\vec{\Delta}(D)$ has been recently introduced in [11] as a parameter to solve problems on tournaments and is defined as follows:

$$\vec{\Delta}(D) = \min \{ \Delta(D^\prec) \mid \prec \text{ is a total ordering of } V(D) \}$$

An *oriented graph* (resp. *tournament*, *semi-complete digraph*) is a digraph in which there is at most (resp. exactly, at least) one arc between each pair of vertices.

The degreewidth of tournaments has been studied in [11], where it is proved that it is NP-hard to compute, that sorting by in-degree gives a 3-approximation, and that one can decide if an n -vertex tournament has degreewidth 1 in $O(n^3)$ steps. In a subsequent paper [14], it is proved that the degreewidth of a semi-complete digraph D can be computed in $\vec{\Delta}(D)^{O(\vec{\Delta}(D))}n + O(n^2)$, and that FEEDBACK ARC SET and CUTWIDTH are FPT parametrised by the degreewidth on semi-complete digraphs. Finally, it is proved [14] that deciding if an oriented graph has degreewidth 2 is NP-hard and the complexity of deciding if an oriented graph has degreewidth 1 is left as an open question. We solve this question by proving that it is NP-hard.

2 Complexity of Computing $\vec{\Delta}$

For an integer $k \in \mathbb{N}$, we define the problem k -DEGREewidth as follows:

- Input: a digraph $D = (V, A)$.
- Question: is $\vec{\Delta}(D)$ at most k ?

Observe that 0-DEGREewidth is exactly the problem of deciding whether a given digraph is an acyclic digraph and it is clearly in P since acyclic digraphs are recognisable in linear time using for example a depth first search. This is however not the case for $k > 0$ (unless $P = NP$) as we show in [Theorem 2.3](#).

Let D be a digraph. A digraph D' is a 1-subdivision of D if D' can be obtained from D by adding a vertex v_a for every arc $a = (x, y)$ of D , and by replacing a by the two arcs (x, v_a) and (v_a, y) .

Remark 2.1. For a digraph D , if D' is a 1-subdivision of D , the underlying graph G of D' is bipartite and 2-degenerate.

A feedback arc set (FAS) of a digraph D is a subset $F \subseteq A(D)$ such that $D - F$ contains no directed cycles. Given a digraph D and a set $F \subseteq A(D)$ of arcs, we define the undirected graph $D[F]$ with vertex set $V(D[F]) = V(D)$ and such that for all $u, v \in V(D[F])$, $uv \in E(D[F])$ if and only if $(u, v) \in A(D)$ or $(v, u) \in A(D)$.

The next lemma ensures that in order to decide whether a given digraph has $\vec{\Delta}(D) \leq k$ it is enough to check the existence of a feedback arc set F such that $\Delta(D[F]) \leq k$.

Lemma 2.2. Let D be a digraph. We have $\vec{\Delta}(D) \leq k$ if and only if D has a FAS F such that $D[F]$ has maximum degree at most k .

Proof: Assume first that $\vec{\Delta}(D) \leq k$. There exists an ordering \prec such that D^\prec has maximum degree k . Hence the set of arcs $F \subseteq A(D)$ corresponding to the edges of D^\prec is a feedback arc set such that $\Delta(D[F]) \leq k$.

Assume now that D has a feedback arc set F such that $\Delta(D[F]) \leq k$. The digraph $D - F$ is acyclic and thus admits a topological ordering \prec . Hence $E(D^\prec) \subseteq F$ and thus $\Delta(D^\prec) \leq k$. ■

Theorem 2.3

For every integer $k \geq 1$, k -DEGREewidth is NP-complete, even restricted to 1-subdivisions of multidigraphs.

Proof: Let $k \geq 1$. The problem k -DEGREewidth is in NP, as for a digraph D , any given vertex ordering \prec achieving $\Delta(D^\prec) \leq k$ can be provided as a certificate.

We prove the NP-hardness of k -DEGREewidth by reducing from 3-SAT.

Before starting the reduction, we define for any integer p the transfer digraph of size p from s to t as $T_p = (V_p^T, E_p^T)$ with $V_p^T = \{s, v_1, \dots, v_p, t\}$ and $E_p^T = \{(s, v_i) : 1 \leq i \leq p\} \cup \{(v_i, t) : 1 \leq i \leq p\}$. Informally, this digraph consists of p disjoint paths of length 2 from the source s to the sink t , as depicted in [Fig. 1](#).

The next claim follows from the pigeonhole principle.

Claim 2.4. Let $T_{2k} = (V, E)$ be a transfer digraph of size $2k$ from s to t and let $F \subset E$. If F disconnects s from t , then $\Delta(D[F]) \geq k$, with equality if and only if $d(s) = d(t) = k$ in $D[F]$.

Let $T_{2k+1} = (V, E)$ be a transfer digraph of size $2k + 1$ from s to t and let $F \subset E$. If F disconnects s from t , then $\Delta(D[F]) \geq k + 1$.




Figure 1: Transfer digraph.

Construction of the reduction.

On a set of boolean variables $X = \{x_1, \dots, x_n\}$, consider a 3-SAT formula $\varphi = \bigwedge_{1 \leq j \leq m} K_j$, where for $1 \leq j \leq m$ the clause $K_j = (\lambda_j^1 \vee \lambda_j^2 \vee \lambda_j^3)$ is a disjunction of literals λ_j^i for $1 \leq i \leq 3$. To the formula φ , we associate a digraph D as follows.

First to every clause $K_j = (\lambda_j^1 \vee \lambda_j^2 \vee \lambda_j^3)$ we associate a digraph D_j constructed as follows (see Figure 2).

- For $1 \leq i \leq 3$, we add the vertices ℓ_j^i and $\tilde{\ell}_j^i$ in D_j .
- For $1 \leq i \leq 3$, we add a transfer digraph of size $2k$ from ℓ_j^i to $\tilde{\ell}_j^i$.
- For $1 \leq i \neq i' \leq 3$, we add a transfer digraph of size $2k + 1$ from $\tilde{\ell}_j^i$ to $\ell_j^{i'}$.
- We finally add three vertices c_j^1, c_j^2, c_j^3 and create a directed cycle $C_j = (\tilde{\ell}_j^1, c_j^1, \tilde{\ell}_j^2, c_j^2, \tilde{\ell}_j^3, c_j^3, \tilde{\ell}_j^1)$.

We connect the digraphs D_j by adding transfer digraphs as follow: For each $1 \leq j \neq j' \leq m$, and each pairs $1 \leq i, i' \leq 3$, such that $\lambda_j^i = \neg \lambda_{j'}^{i'}$, we add a transfer digraph of size $2k + 1$ from $\tilde{\ell}_j^i$ to $\ell_{j'}^{i'}$.

The resulting digraph D has $O(km)$ vertices, so the reduction is polynomial. Note that D belongs to the desired class of digraphs: it is a 1-subdivision of the multi-digraph obtained by replacing every transfer digraph of size p by p parallel arcs and every directed path $(\tilde{\ell}_j^i, c_j^i, \tilde{\ell}_j^{i'})$ by an arc $(\tilde{\ell}_j^i, \tilde{\ell}_j^{i'})$.

Intuition behind the reduction.

1. The cycle C_j associated to the clause K_j guarantees that for every FAS F , at least one path joining some ℓ_j^i to the associated $\tilde{\ell}_j^i$ must not intersect F , and this index i corresponds to a literal λ_j^i that must be set to \top .
2. The gadget linking literals that are negation of one another ensures that the correspondence in the previous point only provides sound valuations (so that $\nu(\lambda) = \top \iff \nu(\neg\lambda) = \perp$ for every literal λ).

Proof of the equivalence.

We now prove that φ is satisfiable if and only if D has a feedback arc set F with $\Delta(D[F]) \leq k$.

(\implies) Suppose first that φ is satisfiable. To a valuation ν satisfying φ we associate a feedback arc sets F such that $\Delta(D[F]) = k$.

We construct the set F as follows:

- For each $1 \leq j \leq m$, we choose one index i_j such that $\nu(\lambda_j^{i_j}) = \top$ and we add the arc $(\ell_j^{i_j}, c_j^{i_j})$ to F . Denote by F_\top the set of arcs added to F during this step.
- For each $1 \leq j \leq m$, for each $1 \leq i \leq 3$ with $i \neq i_j$, we add $2k$ arcs of the transfer digraph from ℓ_j^i to $\tilde{\ell}_j^i$ so that these vertices are disconnected.

By construction, we have $\Delta(D[F]) = k$. We now prove that F is indeed a feedback arc set. Note that for $1 \leq j \leq m$ and $1 \leq i \leq 3$, a directed cycle going through a vertex ℓ_j^i must also go through $\tilde{\ell}_j^i$. Let C be a directed cycle of D .

- Suppose first that $C \subset D_j$ for some $1 \leq j \leq m$. If $C = C_j$ then $E(C) \cap F \neq \emptyset$ since one arc of C_j is in F_\top . Otherwise, C must contain at least one vertex ℓ_j^i , and we distinguish two cases.
 - If $i \neq i_j$ then C intersects F because all the out-arcs from ℓ_j^i are in the transfer digraph from ℓ_j^i to $\tilde{\ell}_j^i$, and these have been disconnected by construction of F .




Figure 2: Representation of a clause gadget.

- If $i = i_j$, then all out-arcs from $\ell_j^{i_j}$ lead to $\tilde{\ell}_j^{i_j}$. Hence every path from $\tilde{\ell}_j^{i_j}$ either contains $(\tilde{\ell}_j^{i_j}, c_j^{i_j})$ which is in $F_\top \subseteq F$, or leads to ℓ_j^i for some $i \neq i_j$, and we are back to the previous point. In any case we proved that that $E(C) \cap F \neq \emptyset$.
- If C is not included in any digraph D_j , then C goes through a transfer digraph from some $\tilde{\ell}_j^i$ to some $\ell_{j^\top}^{i^\top}$ with $1 \leq j \neq j^\top \leq m$, and $1 \leq i, i^\top \leq 3$. Since $\ell_{j^\top}^{i^\top}$ is not a sink of $D - F$, we know that $\nu(\lambda_{j^\top}^{i^\top}) = \top$, hence $(\tilde{\ell}_{j^\top}^{i^\top}, c_{j^\top}^{i^\top}) \in F_\top \subseteq F$. The directed cycle can thus either go (through a transfer digraph) to $\ell_{j^\top}^{i'}$ for $i' \neq i^\top$ but that vertex is a sink of $D - F$ by construction, or to $\ell_{j^\perp}^{i^\perp}$ with $\lambda_{j^\perp}^{i^\perp} = \neg \lambda_{j^\top}^{i^\top}$ but then $\nu(\lambda_{j^\perp}^{i^\perp}) = \perp$ and that vertex is a sink of $D - F$ as well. In any case, C intersects F .

(\Leftarrow) Suppose that D has a feedback arc set F with $\Delta(D[F]) \leq k$. We construct a valuation ν that satisfies φ as follows. For each directed cycle C_j , at least one arc needs to be in F and this arc is necessarily incident to exactly one $\tilde{\ell}_j^i$. We define ν such that $\nu(\lambda_j^i) = \top$ if an arc from $E(C_j)$ incident to $\tilde{\ell}_j^i$ is in F . After this operation, if some variables x_i have not been assigned a truth value, then we assign them arbitrarily. This construction ensures that for every clause, at least one of its literals is assigned to \top . It remains to prove that this valuation is well-defined, namely that there is no literal λ such that $\nu(\lambda) = \top$ and $\nu(\neg \lambda) = \top$. Suppose by contradiction that there exists some λ such that $\nu(\lambda) = \top$ and $\nu(\neg \lambda) = \top$. By definition of ν , there exist two pairs (j_1, i_1) and (j_2, i_2) with $\lambda_{j_1}^{i_1} = \neg \lambda_{j_2}^{i_2}$ such that an arc containing $\tilde{\ell}_{j_1}^{i_1}$ in C_{j_1} and an arc containing $\tilde{\ell}_{j_2}^{i_2}$ in C_{j_2} are in F . By construction of the transfer digraph, since one arc incident to $\tilde{\ell}_{j_1}^{i_1}$ (respectively $\tilde{\ell}_{j_2}^{i_2}$) is in F , there exists a path in the transfer digraph from $\ell_{j_1}^{i_1}$ to $\tilde{\ell}_{j_1}^{i_1}$ (respectively from $\ell_{j_2}^{i_2}$ to $\tilde{\ell}_{j_2}^{i_2}$) that does not intersect F , otherwise, one of these vertices would have degree at least $k + 1$ according to **Claim 2.4**. Therefore, since there is at least one path from $\tilde{\ell}_{j_1}^{i_1}$ to $\ell_{j_2}^{i_2}$ and one path from $\tilde{\ell}_{j_2}^{i_2}$ to $\ell_{j_1}^{i_1}$, there is a directed cycle that does not intersect F , contradicting the fact that F is a feedback arc set. Consequently, ν is well-defined and φ is satisfiable. \blacksquare

3 A Unifying Point of View

3.1 From Undirected to Directed Graphs

Given an (undirected) graph parameter γ , we define its directed version as:

$$\vec{\gamma}(D) = \min \{ \gamma(D^\prec) \mid \prec \text{ is a total ordering of } V(D) \}. \quad (1)$$

We say that an ordering \prec of a digraph D is a $\vec{\gamma}$ -ordering if \prec witnesses $\vec{\gamma}(D)$, that is $\vec{\gamma}(D) = \gamma(D^\prec)$.

As for the degreewidth, several digraph parameters are derived by applying (1) to an undirected parameter γ . Notably, for certain digraph parameters γ' , there exists a related undirected parameter γ such that $\vec{\gamma}$ retrieves γ' ; examples include the dichromatic number and the Feedback Vertex Number, as detailed below. We believe that exploring the perspective given by the backedge graphs can provide new insights into these parameters and potentially uncover novel digraph parameters of interest.

In this section, we give an overview of existing digraphs parameters defined using (1), namely dichromatic number, directed clique number, feedback vertex number and of course degreewidth, that we explore a bit further by considering it as a maximum degree concept, rather than merely a parameter for obtaining FPT algorithms. Next, we turn our attention to optimal linear arrangement, cutwidth, and bandwidth, which are graph parameters for which each of their directed analogues is defined via the backed graphs, but not by applying (1). We will detail this in the last subsection. The goal is not to prove new results, but rather to draw attention to this backedge graph perspective, which we believe is not very well known.

First, the backedge graph perspective provided by (1) offers a unified approach to presenting the decision problem related to a digraph parameter $\vec{\gamma}$, that is deciding if a given digraph D satisfies $\vec{\gamma}(D) \leq k$. Given a class of (undirected) graphs \mathcal{C} , we say that a FAS is a \mathcal{C} -FAS if the graph induced by the edges of F (forgetting their orientations) belongs to \mathcal{C} . Given a class of undirected graphs \mathcal{C} and a class of digraphs \mathcal{D} , the \mathcal{C} -FAS Problem is the problem of deciding whether a digraph $D \in \mathcal{D}$ has a \mathcal{C} -FAS.

In this formalism, the main result of this paper proves that, for every $k \geq 1$, the \mathcal{C} -FAS problem in oriented graphs (more precisely in 1-subdivision of oriented multigraphs) is NP-hard when \mathcal{C} is the class of graphs with maximum degree k .

More generally, for a graph parameter γ which is monotone (in the sense $\gamma(H) \leq \gamma(G)$ for any subgraph H of G), deciding whether a digraph D satisfies $\vec{\gamma}(D) \leq k$ is the same as the \mathcal{C} -FAS problem in \mathcal{D} where \mathcal{C} is the class of graphs G satisfying $\gamma(G) \leq k$ and \mathcal{D} is the class of all digraphs. This problem is often of particular interest when \mathcal{D} is taken to be the class of all tournaments. For more on this subject we refer to [3] (in particular the last section).

For an undirected graph G , the digraph \overleftrightarrow{G} is obtained from G by replacing each edge with a *digon* consisting of two opposite arcs. The adjacency relation of \overleftrightarrow{G} is symmetric, and the map $G \mapsto \overleftrightarrow{G}$ naturally injects graphs into digraphs. Observe that every ordering \prec of \overleftrightarrow{G} satisfies $\overleftrightarrow{G}^\prec = G$, hence for any graph parameter γ we have $\gamma(G) = \vec{\gamma}(\overleftrightarrow{G})$. Consequently $\vec{\gamma}$ generalises γ : any property satisfied by γ on undirected graphs will also be satisfied by $\vec{\gamma}$ in restriction to symmetric digraphs, and one can wonder how this property extends to all other digraphs.

3.2 General Observations

This generalisation of parameters $\gamma \mapsto \vec{\gamma}$ from graphs to digraphs also preserves inequalities. If graph parameters γ_1, γ_2 are such that for every graph G we have $\gamma_1(G) \leq \gamma_2(G)$, then for every digraph D and every ordering \prec of D we have $\gamma_1(D^\prec) \leq \gamma_2(D^\prec)$ hence $\vec{\gamma}_1(D) \leq \vec{\gamma}_2(D)$. Moreover, since $\gamma \mapsto \vec{\gamma}$ is defined by a minimum, for a non-decreasing function $f : \mathbb{N} \rightarrow \mathbb{R}$ we have $f \circ \gamma \leq f \circ \vec{\gamma}$. Hence monotone inequalities are also preserved: if there exists a non-decreasing function $f : \mathbb{N} \rightarrow \mathbb{R}$ such that for every graph G we have $\gamma_1(G) \leq f(\gamma_2(G))$, then for every digraph D we have $\vec{\gamma}_1(D) \leq f \circ \vec{\gamma}_2(D) \leq f(\vec{\gamma}_2(D))$.

Let $\mathcal{F}(D)$ be the set of inclusion-minimal feedback arc sets of D . For a set of arcs F of D , denote by $D[F]$ the undirected graph with vertices $V(D)$ and edges F (removing orientations). Observe that for every $F \in \mathcal{F}(D)$, there is a topological ordering \prec of $D - F$ such that $D^\prec = D[F]$.

Lemma 3.1. *If γ is a monotone graph parameter, then for every digraph D :*

$$\vec{\gamma}(D) = \min_{F \in \mathcal{F}(D)} \gamma(D[F]).$$

Proof: Consider a $\vec{\gamma}$ -ordering \prec of D . Let F be the set of arcs $(x, y) \in A(D)$ such that $y \prec x$ and note that $D[F] = D^\prec$. Since F is a feedback arc set of D , it contains a minimal feedback arc set $F' \in \mathcal{F}(D)$, and $D[F']$ is a subgraph of $D[F]$. Since γ is monotone, we have:

$$\vec{\gamma}(D) = \gamma(D^\prec) = \gamma(D[F]) \geq \gamma(D[F']) \geq \min_{\tilde{F} \in \mathcal{F}(D)} \gamma(D[\tilde{F}]).$$

Consider a minimal feedback arc set $F \in \mathcal{F}(D)$. Choose a topological ordering \prec_F of $D - F$. We have $D^\prec = D[F]$ hence $\vec{\gamma}(D) \leq \gamma(D[F])$, which ends the proof. ■

Proposition 3.2

For every undirected graph parameter γ and digraph D we have $\vec{\gamma}(D) = \vec{\gamma}(D^\dagger)$, where D^\dagger is obtained from D by reversing the direction of every arc.

Proof: If \prec is a $\vec{\gamma}$ -ordering of D then the opposite ordering \prec' defined by $v \prec' w \iff w \prec v$ satisfies $D^\prec = (D^\dagger)^{\prec'}$ hence $\vec{\gamma}(D) \geq \vec{\gamma}(D^\dagger)$ and the other inequality follows by symmetry. ■

3.3 Dichromatic Number and Directed Clique Number

Let us explain why the directed generalisation $\vec{\chi}$ obtained by applying (1) to the chromatic number χ the *dichromatic number* [16] χ_a , traditionally defined for a digraph D as the minimum k such that D can be partitioned into k acyclic induced subdigraph and denoted here by χ_a .

On the one hand, for every ordering \prec , a stable set of D^\prec induces an acyclic subdigraph in D , so a colouring of D^\prec is a dicolouring of D and we have $\chi_a(D) \leq \chi(D^\prec)$ for every \prec . On the other hand, it is enough to provide an ordering \prec such that $\chi(D^\prec) \leq \chi_a(D)$. This can be done by taking an ordering built from a dicolouring, choosing an ordering on the colour classes, and ordering the vertices of each colour class with a topological order.

This point of view on the dichromatic number has been used several times, for example in [6, 17, 4], but we believe it deserves more attention.

A recent development in this direction is the study of the *directed clique number* $\vec{\omega}$ defined by applying (1) to the clique number ω . It has been first introduced in [2] (see also [17, 5]). This gives a notion of clique number for digraphs that leads to a notion of $\vec{\chi}$ -bounded classes of digraphs, which we think deserve further exploration.

3.4 Directed Vertex Cover is Feedback Vertex Set

The *vertex cover number* $\tau(G)$ of an undirected graph G is the minimum size of a vertex set S such that $G - S$ is edgeless. Its directed version is therefore:

$$\vec{\tau}(D) = \min \{ \tau(D^\prec) \mid \prec \text{ is a total ordering of } V(D) \}$$

The Feedback Vertex number $\text{fvn}(D)$ of a digraph D is the size of a minimum set of vertices S such that $D - S$ is acyclic.

Interestingly, and maybe not surprisingly, fvn is the same as $\vec{\tau}$ as we show now. Let D be a digraph. Let S be a Feedback Vertex Set of D . The acyclic digraph $D - S$ has a topological ordering \prec' , hence $(D - S)^{\prec'}$ is edgeless. Hence, by defining \prec as any ordering of D placing the vertices in S first followed by the ordering \prec' , we get that S is a vertex cover of D^\prec . Hence $\vec{\tau}(D) \leq |S| = \text{fvn}(D)$. Let \prec be a $\vec{\tau}(D)$ -ordering of D . For every vertex cover S of D^\prec , we have that $D^\prec - S$ is edgeless, so $D - S$ is acyclic; hence $\text{fvn}(D) \leq |S| = \vec{\tau}(D)$.

3.5 Back to the Degreewidth

The maximum degree of a graph has several directed analogues. Let v be a vertex of a digraph D . We define the *maxdegree* of v as $d_{\max}(v) = \max\{d^+(v), d^-(v)\}$ and the *mindegree* of v as $d_{\min}(v) = \min\{d^+(v), d^-(v)\}$. We then define the corresponding maximum degrees of D : $\Delta_{\max}(D) = \max\{d_{\max}(v) \mid v \in V(D)\}$ and $\Delta_{\min}(D) = \max\{d_{\min}(v) \mid v \in V(D)\}$. The degreewidth $\vec{\Delta}$ can be seen as a third notion of maximum degree. With this point of view, the degreewidth can be studied by drawing inspiration from results involving the maximum degree in the undirected world. In this section, we first compare the three notions, and then have a look to a potential generalisation of Brooks' theorem, that relates the maximum degree and the chromatic number of undirected graphs.

Note that $\vec{\Delta}, \Delta_{\max}, \Delta_{\min}$ coincide on symmetric digraphs.

Lemma 3.3. For every digraph D , we have $\vec{\Delta}(D) \leq \Delta_{\min}(D) \leq \Delta_{\max}(D)$.

Proof: Let D be a digraph. It is clear that $\Delta_{\min}(D) \leq \Delta_{\max}(D)$. Let \prec be an ordering of $V(D)$ that minimises the number of edges of D^{\prec} and assume by contradiction that there exists $u \in V(D)$ such that $d_{D^{\prec}}(u) > \Delta_{\min}(D)$. Let α (respectively β) be the number vertices x such that $x \prec u$ and (u, x) is an arc of D (respectively $u \prec x$ and (x, u) is an arc of D). So $d_{D^{\prec}}(u) = \alpha + \beta \geq \Delta_{\min}(D) + 1$. Assume first that $d^-(u) = d_{\min}(u)$. Define the ordering \prec^* from \prec by keeping \prec on $V(D) \setminus \{u\}$ untouched and placing u at the beginning. Then

$$|E(D^{\prec^*})| = |E(D^{\prec})| - \alpha + (d^-(u) - \beta)$$

Since $d^-(u) \leq \Delta_{\min}(D) < \alpha + \beta$, we have $|E(D^{\prec^*})| < |E(D^{\prec})|$, contradicting the choice of \prec . The case $d^+(u) = d_{\min}(u)$ is treated similarly by placing u at the end of the ordering. ■

Corollary 3.4

If D is a k -regular digraph, namely for which every vertex v satisfies $d^-(v) = d^+(v) = k$, then $\vec{\Delta}(D) = \Delta_{\min}(D) = \Delta_{\max}(D) = k$.

Proof: Let D be a k -regular digraph. It is clear that $\Delta_{\min}(D) = \Delta_{\max}(D) = k$. By Lemma 3.3, we know that $\vec{\Delta}(D) \leq k$. Moreover, for every ordering \prec , the smallest vertex u satisfies $d^-(u) = d_{D^{\prec}}(u) = k$, so $\vec{\Delta}(D) \geq k$. ■

Let D be a digraph, and \prec a $\vec{\Delta}$ -ordering of D . By greedily colouring D^{\prec} , we get that $\vec{\chi}(D) \leq \vec{\Delta}(D) + 1$. Hence, for every digraph D , we have:

$$\vec{\chi}(D) \leq \vec{\Delta}(D) + 1 \leq \Delta_{\min}(D) + 1 \leq \Delta_{\max}(D) + 1.$$

For $k \in \mathbb{N}$, the k -DICOLOURABILITY decision problem takes as input a digraph D and decides if it is k -dicolourable. It is known to be NP-complete [7] as soon as $k \geq 2$.

Brooks' theorem [8] is a fundamental result in graph colouring, stating that every graph G satisfies $\chi(G) \leq \Delta(G) + 1$ with equality if and only if G is an odd cycle or a complete graph. A directed version of Brooks' theorem for Δ_{\max} , giving a full characterisation of digraphs D satisfying $\vec{\chi}(D) = \Delta_{\max}(D) + 1$, has been proved in [15] and a polynomial time algorithm can be deduced to decide if a given digraph D satisfies $\vec{\chi}(D) = \Delta_{\max}(D) + 1$. In [1], it is proved that it is NP-complete to decide if a given digraph D satisfies $\vec{\chi}(D) = \Delta_{\min}(D) + 1$.

Theorem 3.5 ([1])

For $k \geq 2$, k -DICOLOURABILITY is NP-complete even when restricted to digraphs D with $\Delta_{\min}(D) = k$.

Using a very similar proof, we now show that the same holds after replacing Δ_{\min} by $\vec{\Delta}$, hence a directed version of Brooks' theorem using $\vec{\Delta}$ is unfortunately very unlikely.

Given a digraph D and a vertex v of D , we denote by $\text{dig}_D(v)$ the number of digons incident to v in D :

$$\text{dig}_D(v) = |\{w \in V(D) \mid (v, w), (w, v) \in A(D)\}|.$$

If the digraph D is clear, we omit the subscript and write $\text{dig}(v)$.

Lemma 3.6. *Every digraph D satisfies $\vec{\Delta}(D) \geq \max\{\text{dig}(v) \mid v \in V(D)\}$.*

Proof: For every $v \in V(D)$, let $w_1, \dots, w_{\text{dig}(v)}$ be the "digon-neighbours" of v . Now fix any ordering \prec of $V(D)$ and observe that for every $1 \leq i \leq \text{dig}(v)$ either $w_i \prec v$, in which case $(v, w_i) \in E(D^{\prec})$, or $v \prec w_i$ in which case $(w_i, v) \in E(D^{\prec})$. Hence $d_{D^{\prec}}(v) \geq \text{dig}_D(v)$ which proves the result. ■

Theorem 3.7

For $k \geq 2$, k -DICOLOURABILITY is NP-complete even when restricted to digraphs D with $\vec{\Delta}(D) = k$.

Proof: Let $k \geq 2$. In [1] the proof of Theorem 3.5 associates to any digraph D a digraph D' satisfying $\Delta_{\min}(D') = k$ and $\vec{\chi}(D) = k$ if and only if $\vec{\chi}(D') = k$. We prove that the constructed D' also satisfies $\vec{\Delta}(D) = k$, implying the theorem. We start, for sake of completeness, by giving the construction of D' .

Start with a given digraph D , and construct the digraph D' as follows. For each vertex v , D' has $k + 1$ vertices $v^-, v^+, v_1, \dots, v_{k-1}$ (so D' has $|V(D)|(k + 1)$ vertices). For each vertex v of D , create arcs so that $D'[\{v^-, v_1, \dots, v_{k-1}\}]$ and $D'[\{v^+, v_1, \dots, v_{k-1}\}]$ are complete symmetric digraphs, and then add the arc (v^-, v^+) . Finally, for every arc uv of D , add the arc (u^+, v^-) to D' .

As proved in [1], we have $\Delta_{\min}(D') = k$. Indeed, for every $v \in V(D)$ and $1 \leq i < k$, we have $\text{dig}_{D'}(v_i) = d^+(v_i) = d^-(v_i) = k$, therefore $\Delta_{\min}(D') \geq k$. Moreover, for every $v \in V(D)$ we have $d^+(v^-) = d^-(v^+) = k$, so $d_{\min}(v^+) \leq k$ and $d_{\min}(v^-) \leq k$. Therefore $\Delta_{\min}(D') = k$.

We now prove that $\vec{\Delta}(D) = k$. Combining [Lemma 3.3](#) and [Lemma 3.6](#), one gets:

$$k \leq \max_{v \in V(D')} \text{dig}(v) \leq \vec{\Delta}(D') \leq \Delta_{\min}(D') = k.$$

Finally, the proof concludes the same way as for [Theorem 3.7](#) since D is k -dicolourable if and only if D' is. \blacksquare

Even though we cannot hope for a complete characterisation of digraphs D satisfying $\vec{\chi}(D) = \vec{\Delta}(D) + 1$, we may still show the following partial result.

Observation 3.8

Let D be a digraph with $\vec{\Delta}(D) = k$. If $\vec{\chi}(D) = k + 1$ then for every $\vec{\Delta}$ -ordering \prec of D , the graph D^\prec has a connected component equal to an odd cycle if $k = 2$ or to the clique K_{k+1} if $k \neq 2$.

Proof: Let \prec be a $\vec{\Delta}$ -ordering. If no connected component of D^\prec is one of Brooks' exception, then $\chi(D^\prec) \leq \Delta(D^\prec) = \vec{\Delta}(D)$, and the result follows from $\vec{\chi}(D) \leq \chi(D^\prec)$. \blacksquare

3.6 Optimal Linear Arrangement, Cutwidth and Bandwidth

Given an ordered undirected graph (G, \prec) , the *length* of an edge (u, v) such that $u \prec v$ is 1 plus the number of vertices x such that $u \prec x \prec v$. One can define three undirected parameters by minimising certain functions of the lengths over all ordering of the graph: Optimal Linear Arrangement, cutwidth, and bandwidth. Each of these parameters has a directed version defined similarly, but for a given digraph D and an ordering \prec , only the edges of D^\prec are considered. These directed versions differ from those obtained by applying (1) to the undirected parameters. The sole purpose of this section is to highlight this difference. The authors did not explore the relationship between these parameters.

Let us provide a detailed explanation for the Optimal Linear Arrangement. For an undirected graph G with an ordering \prec , we denote by $\text{cost}(G, \prec)$ the sum of the lengths of its edges. An *Optimal Linear Arrangement* of G is an ordering \prec that minimises $\text{cost}(G, \prec)$, and its value is denoted $\text{OLA}(G)$, thus:

$$\text{OLA}(G) = \min \{ \text{cost}(G, \prec) \mid \prec \text{ is a total ordering of } V(G) \}.$$

The directed version diOLA of OLA is defined similarly [9, 12], but only the length of backward arcs are summed. More precisely,

$$\text{diOLA}(D) = \min \{ \text{cost}(D^\prec, \prec) \mid \prec \text{ is a total ordering of } V(D) \}.$$

This is not the same as $\overrightarrow{\text{OLA}}$ defined according to the formula (1) by:

$$\begin{aligned} \overrightarrow{\text{OLA}}(D) &= \min \{ \text{OLA}(D^\prec) \mid \prec \text{ is a total ordering of } V(D) \} \\ &= \min \{ \text{cost}(D^\prec, \prec') \mid \prec, \prec' \text{ are total orderings of } V(D) \}. \end{aligned}$$

Hence, in $\overrightarrow{\text{OLA}}$, we can choose the Optimal Linear Arrangement over each backedge graph D^\prec , while in diOLA we only consider the cost of (D^\prec, \prec) . Thus, for every digraph D , we have $\overrightarrow{\text{OLA}}(D) \leq \text{diOLA}(D)$.

The *bandwidth* of an ordered graph (G, \prec) is the maximum length of an edge, and the bandwidth of a graph G is $\text{bw}(G) = \min \{ \text{bw}(G, \prec) \mid \prec \text{ is a total ordering of } V(G) \}$. The *cutwidth* $\text{cw}(G)$ of an undirected graph G is the smallest integer k with the following property: there is an ordering of the vertices of the graph, such that every cut obtained by partitioning the vertices into earlier and later subsets of the ordering is crossed by at most k edges. As with OLA , the directed bandwidth [13] and directed cutwidth [10] of a digraph D are defined similarly to their undirected counterparts, but for each ordering \prec , only the edges of D^\prec are considered. Therefore, as with OLA , applying (1) to these parameters does not yield the same results.

Acknowledgement

This research was partially supported by the french *Agence Nationale de la Recherche* under contract *DAGDigDec (JCJC) ANR-21-CE48-0012*, the group *Casino/ENS Chair on Algorithmics and Machine Learning*, the *Austrian Science Foundation (FWF, Project 10.55776/Y1329)* and the  *European Union's Horizon 2020 COFUND programme (LogiCS@TUWien, No. 101034440)*.

References

- [1] Pierre Aboulker and Guillaume Aubian. *Four proofs of the directed Brooks' Theorem*. *Discrete Mathematics*, 346(11):113193, 2023.
- [2] Pierre Aboulker, Guillaume Aubian, Pierre Charbit, and Raul Lopes. *Clique number of tournaments*. arXiv preprint arXiv:2310.04265, 2023.
- [3] Pierre Aboulker, Guillaume Aubian, and Raul Lopes. *Finding forest-orderings of tournaments is NP-complete*. arXiv preprint arXiv:2402.10782, 2024.
- [4] Pierre Aboulker, Frédéric Havet, François Pirot, and Juliette Schabanel. *Minimum acyclic number and maximum dichromatic number of oriented triangle-free graphs of a given order*. arXiv preprint arXiv:2403.02298, 2024.
- [5] Guillaume Aubian. *Computing the clique number of tournaments*. arXiv preprint arXiv:2401.07776, 2024.
- [6] Eli Berger, Krzysztof Choromanski, Maria Chudnovsky, Jacob Fox, Martin Loebl, Alex Scott, Paul Seymour, and Stéphan Thomassé. *Tournaments and colouring*. *Journal of Combinatorial Theory, Series B*, 103:1–20, 01 2013.
- [7] Drago Bokal, Gasper Fijavz, Martin Juvan, P Mark Kayll, and Bojan Mohar. *The circular chromatic number of a digraph*. *Journal of Graph Theory*, 46(3):227–240, 2004.
- [8] R. L. Brooks. *On colouring the nodes of a network*. *Math. Proc. Cambridge Philos. Soc.*, 37:194–197, 1941.
- [9] P. Z. Chinn, J. Chvátalová, A. K. Dewdney, and N. E. Gibbs. *The bandwidth problem for graphs and matrices—a survey*. *Journal of Graph Theory*, 6(3):223–254, 1982.
- [10] Maria Chudnovsky, Alexandra Fradkin, and Paul Seymour. *Tournament immersion and cutwidth*. *Journal of Combinatorial Theory, Series B*, 102(1):93–101, 2012.
- [11] Tom Davot, Lucas Isenmann, Sanjukta Roy, and Jocelyn Thiebaut. *Degreewidth: A New parameter for Solving Problems on Tournaments*. In *International Workshop on Graph-Theoretic Concepts in Computer Science*, pages 246–260. Springer, 2023.
- [12] Fedor V. Fomin and Michał Pilipczuk. *On width measures and topological problems on semi-complete digraphs*. *Journal of Combinatorial Theory, Series B*, 138:78–165, 2019.
- [13] Pallavi Jain, Lawqueen Kanesh, William Lochet, Saket Saurabh, and Roohani Sharma. *Exact and approximate digraph bandwidth*. In *Arkadev Chattopadhyay and Paul Gustin, editors, 39th IARCS Annual Conference on Foundations of Software Technology and Theoretical Computer Science, FSTTCS 2019, December 11-13, 2019, Bombay, India, volume 150 of LIPIcs, pages 18:1–18:15. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019*.
- [14] Ryan Keeney and Daniel Lokshantov. *Degreewidth on semi-complete digraphs*. In *International Workshop on Graph-Theoretic Concepts in Computer Science*, page to be published, 2024.
- [15] B. Mohar. *Eigenvalues and colourings of digraphs*. *Linear Algebra and its Applications*, 432(9):2273–2277, 2010.
- [16] Victor Neumann-Lara. *The dichromatic number of a digraph*. *Journal of Combinatorial Theory, Series B*, 33(3):265–270, 1982.
- [17] Tung Nguyen, Alex Scott, and Paul Seymour. *Some results and problems on tournament structure*, 2023.