

QUANTITATIVE STABILITY IN OPTIMAL TRANSPORT FOR GENERAL POWER COSTS

OCTAVE MISCHLER AND DARIO TREVISAN

ABSTRACT. We establish novel quantitative stability results for optimal transport problems with respect to perturbations in the target measure. We provide explicit bounds on the stability of optimal transport potentials and maps, which are relevant for both theoretical and practical applications. Our results apply to a wide range of costs, including all Wasserstein distances with power cost exponent strictly larger than 1 and leverage mostly assumptions on the source measure, such as log-concavity and bounded support. Our work provides a significant step forward in the understanding of stability of optimal transport problems, as previous results were mostly limited to the case of the quadratic cost.

1. INTRODUCTION

A well-posed mathematical problem in the Hadamard sense is one for which existence and uniqueness of solutions hold, but also stability with respect to perturbations in the data. Although qualitative stability in most situations emerges as a by-product of existence and uniqueness arguments, quantitative stability becomes particularly crucial in applications, for it directly relates to the convergence of numerical methods. This applies in particular to optimal transport problems, which in the last decades have had significant impact across various fields of applied mathematics, most notably PDEs of evolution type [AGS08] and machine learning [PC+19]. In this work, we establish novel quantitative stability results for optimal transport problems with respect to a large variety of costs, including Wasserstein distances.

The Wasserstein distance of order $p \geq 1$ between two probability measures ρ, μ on \mathbb{R}^d , with finite p -th moment, is defined as the quantity

$$(1.1) \quad \mathcal{W}_p(\rho, \mu) := \inf_{\pi \in \Pi(\rho, \mu)} \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} |y - x|^p d\pi(x, y) \right)^{1/p},$$

where $\Pi(\rho, \mu)$ denotes the set of couplings between ρ and μ , i.e., (joint) probability measures on $\mathbb{R}^d \times \mathbb{R}^d$ with marginal distributions ρ, μ , and $|\cdot|$ denotes the Euclidean norm. Under mild assumptions, in particular if ρ is absolutely continuous with respect to the Lebesgue measure and $p > 1$, one can argue [AGS08, Theorem 6.2.4] that the optimizer is unique and given by a optimal transport map T for $\mathcal{W}_p(\rho, \mu)$, i.e., one has $y = T(x)$ for π -a.e. (x, y) (with necessarily $T_{\#}\rho = \mu$) and therefore

$$(1.2) \quad \mathcal{W}_p(\rho, \mu)^p = \int_{\mathbb{R}^d} |T(x) - x|^p d\rho(x).$$

For the quadratic cost case $p = 2$, the optimal transport map is often referred to as Brenier's map.

Even when an optimal transport map exists, it is useful to define \mathcal{W}_p as a minimum over couplings, as one can exploit convex duality and obtain the equivalent formulation

$$(1.3) \quad \mathcal{W}_p(\rho, \mu)^p = \sup_{(\phi, \psi)} \left\{ \int_{\mathbb{R}^d} \phi d\rho + \int_{\mathbb{R}^d} \psi d\mu \right\},$$

where the supremum runs among upper semi-continuous functions (ϕ, ψ) such that $\phi(x) + \psi(y) \leq |x - y|^p$ for every $x, y \in \mathbb{R}^d$. An optimizing pair (ϕ, ψ) is called an optimal

Date: July 30, 2024.

Key words and phrases. Optimal transport, quantitative stability, Prékopa-Leindler inequality, fractional Sobolev spaces.

(or Kantorovich) pair of potentials for $\mathcal{W}_p(\rho, \mu)$. Under minimal assumptions, one can always argue that an optimizer exists, together with a qualitative stability result stating convergence of the potentials or the maps, see e.g. [Vil+09, Theorem 5.20, Corollary 5.23].

1.1. Main results. Aim of this work is to establish new quantitative stability results for optimal transport potentials and maps with respect to perturbations of the “target” measure μ , keeping one “source” measure ρ fixed. In a slightly simplified form, to ease the notation, our main contributions can be summarized in the following two results. The first one concerns stability of Kantorovich potentials and applies to any log-concave ρ with bounded support.

Theorem 1.1. *Let $d \geq 1$, λ be a log-concave probability measure on \mathbb{R}^d with bounded support, ρ be a probability measure on \mathbb{R}^d absolutely continuous with respect to λ , with density bounded from above and below by strictly positive constants, let $\mathcal{Y} \subseteq \mathbb{R}^d$ be compact and $p > 1$. Then, there exists $C = C(\lambda, \rho, \mathcal{Y}, p) < \infty$ such that, for any μ, ν , probability measures supported on \mathcal{Y} , it holds*

$$(1.4) \quad \|\phi_\mu - \phi_\nu\|_{L^2(\rho)} \leq C \mathcal{W}_1(\mu, \nu)^\theta,$$

with

$$(1.5) \quad \theta = \begin{cases} 1 - \frac{1}{p} & \text{if } 1 < p < 2, \\ \frac{1}{2} & \text{if } p \geq 2, \end{cases}$$

and where (ϕ_μ, ψ_μ) denotes the Kantorovich potentials for $\mathcal{W}_p(\rho, \mu)$, i.e., the optimizer in (1.3), with zero ρ -mean, i.e., $\int \phi_\mu d\rho = 0$ (and similarly (ϕ_ν, ψ_ν) , for $\mathcal{W}_p(\rho, \nu)$).

Our second main result deals instead with optimal transport maps: in this case, we require ρ to have a density bounded from above and below with respect to the Lebesgue measure on a bounded convex set.

Theorem 1.2. *Let $d \geq 1$, ρ be a probability measure on \mathbb{R}^d with bounded and convex support, absolutely continuous with respect to the Lebesgue measure, with density bounded from above and below by strictly positive constants, let $\mathcal{Y} \subseteq \mathbb{R}^d$ be compact and $p > 1$. Then,*

$$(1.6) \quad \begin{cases} \text{for every } \theta \in \left(0, \frac{(p-1)^2}{p(p+1)}\right) & \text{if } 1 < p < 2, \\ \text{for } \theta = \frac{1}{6(p-1)} & \text{if } p \geq 2, \end{cases}$$

there exists $C = C(\rho, \mathcal{Y}, p, \theta) < \infty$ such that, for any μ, ν , probability measures supported on \mathcal{Y} , it holds

$$(1.7) \quad \|T_\mu - T_\nu\|_{L^2(\rho)} \leq C \mathcal{W}_1(\mu, \nu)^\theta,$$

where T_μ denotes the optimal transport map for $\mathcal{W}_p(\rho, \mu)$, i.e., (1.2) holds (and similarly T_ν , for $\mathcal{W}_p(\rho, \nu)$).

1.2. Related literature. To our knowledge, most of the existing quantitative results concern quadratic optimal transport only. The first of these results is due to Ambrosio, reported by Gigli in [Gig11] and reformulated in form closer to our setting in [MDC20]. It states stability of Brenier’s map around measures μ such that T_μ is Lipschitz.

Theorem 1.3 (Ambrosio and Gigli). *Let ρ be a probability density with bounded support $\mathcal{X} \subseteq \mathbb{R}^d$, let $\mathcal{Y} \subset \mathbb{R}^d$ be compact, and given $\mu, \nu \in \mathcal{P}(\mathcal{Y})$, write T_μ for the Brenier map from ρ to μ , i.e., (1.2) holds with $p = 2$ and $T = T_\mu$, similarly write T_ν . Then,*

$$(1.8) \quad \|T_\mu - T_\nu\|_{L^2(\rho)} \leq 2 \text{Lip}(T_\mu) \text{diam}(\mathcal{X}) \mathcal{W}_1(\mu, \nu)^{1/2},$$

where $\text{Lip}(\cdot)$ denotes the Lipschitz constant and $\text{diam}(\cdot)$ the diameter.

When compared with Theorem 1.2, we see that the assumptions on ρ are weaker, but are compensated by the fact that T_μ must be Lipschitz continuous, otherwise the bound is vacuous. In applications, regularity for the Brenier map is usually obtained as a consequence

of Caffarelli's theory [Caf92], which requires stronger assumptions than [Theorem 1.2](#). As a recent extension to more general costs of this type of results, we mention [GMT22].

Other results in this direction have been obtained, notably [NZ18; Ber21; LN21], but to keep this part short, we only state one other result, which summarizes the results [Del22, Theorem 5.14, Theorem 5.12, Corollary 5.8], first exposed in [MDC20; DM23].

Theorem 1.4 (Mérigot, Delalande and Chazal). *Let $\mathcal{X} \subseteq \mathbb{R}^d$ be a compact convex set, ρ be a probability density supported on \mathcal{X} , bounded from above and below by strictly positive constants. Let $p > \max\{d, 3\}$ and assume that $\mu, \nu \in \mathcal{P}(\mathbb{R}^d)$ have bounded p -th moment,*

$$(1.9) \quad \int_{\mathbb{R}^d} |x|^p d\mu(x) + \int_{\mathbb{R}^d} |x|^p d\nu(x) =: m_p < \infty.$$

Then, there exists $C = C(d, p, \mathcal{X}, \rho, m_p) < \infty$ such that

$$(1.10) \quad \|\phi_\mu - \phi_\nu\|_{L^2(\rho)} \leq C\mathcal{W}_1(\mu, \nu)^{1/2} \quad \text{and} \quad \|T_\mu - T_\nu\|_{L^2(\rho, \mathbb{R}^d)} \leq C\mathcal{W}_1(\mu, \nu)^{1/(6+16d/p)},$$

where ϕ_μ, T_μ denote respectively the Kantorovich potential with zero ρ -mean and the optimal transport map for $\mathcal{W}_2(\rho, \mu)$ (and similarly ϕ_ν, T_ν for $\mathcal{W}_2(\rho, \nu)$).

If moreover μ, ν are supported on \mathcal{Y} compact one can replace the exponent $1/(6+16d/p)$ above with $1/6$, for some constant $C = C(d, \mathcal{X}, \rho, \mathcal{Y}) < \infty$.

Let us finally mention [DNWP24] which provides stronger estimates than the previous result in the semi-discrete setting.

Technically, our work is to improve upon the above results by extending the quantitative stability for optimal transport to larger families of costs, paving the way for a wider range of application. Our analysis, in particular [Theorem 1.1](#), allows for a source probability ρ that is merely log-concave, and actually this assumption is relaxed by allowing for a density with respect to a given log-concave measure that is bounded from above and below by strictly positive constants; in a similar way, in [Theorem 1.2](#) the bounded support assumption for the target measures could be replaced by moment bounds – see also extensions and open problems below.

1.3. Comments on the proof technique. The main lines of our argument follow [Del22], where the stability inequality for potentials is first established for the entropic semi-discrete optimal transport problem, and then extended to the Wasserstein distance by an approximation argument. However, the proofs in [Del22] rely heavily on the fact that the cost is quadratic, so that up to a shift by removing the second moments, it is linear, hence both concave and convex.

Our first key realization is that the convexity assumption can actually be circumvented, and only concavity of the cost plays an essential role in the stability inequality for the Kantorovich potentials (1.4). This leads to a more straightforward argument for the case $p = 2$. The extension to general costs however requires further ideas. A crucial one is a generalization of the Prekopa-Leindler inequality to log-concave measures, which is proved using a modification of McCann's optimal transport argument [McC94]. Actually, for $p \geq 2$, also the argument by Brascamp and Lieb [BL76] can be used to obtain Prékopa-Leindler's inequality, however for $p < 2$ McCann's argument seems necessary, since concavity of the cost fails (even allowing for a quadratic shift). Curiously, our argument yields therefore a quantitative stability in optimal transport through an optimal transport derived inequality.

Finally, to obtain stability for the optimal transport maps (1.7), a Gagliardo-Nirenberg type inequality is used in [Del22]. For $p \geq 2$ the argument goes through similarly, however for $p < 2$ this requires some modifications, in particular the use of fractional Sobolev spaces.

1.4. Extensions and open problems. We mention some extensions of our results that may be worth exploring, possibly using the techniques we developed.

(1) One notices immediately that [Theorem 1.2](#) holds for target measures with bounded support, while [Theorem 1.4](#) requires only bounds on their moments. It seems that this gap can be filled simply following the arguments from [Del22].

(2) It may be also worth exploring whether quantitative stability results for potentials hold when the reference measure has unbounded support. An interesting example may be that of a Gaussian density ρ . Let us remark however that extending the stability result for optimal transport maps to general log-concave measures appears to be a more open-ended problem, as it would require establishing suitable “weighted” Gagliardo-Nirenberg type inequalities.

(3) We conjecture that quantitative stability bounds on Riemannian manifolds or even more general settings could be obtained using suitable versions of our arguments, at least under bounded curvature assumptions – see e.g. [CEMS06] for an extension of Prékopa–Leindler type inequalities on Riemannian manifolds.

1.5. Structure of the paper. Section 2 contains definitions and general well-known results. Section 3 is devoted to the proof of Theorem 1.1, while Section 4 focuses on the derivation of Theorem 1.2. In both sections, we naturally split the arguments in three cases $p = 2$, which we believe is particularly useful to understand the key points in our argument, $p > 2$ which is a relatively easy variation of the former, and then the hardest case $1 < p < 2$, which requires substantially new tools.

2. NOTATION AND BASIC FACTS

2.1. General notation. Throughout the paper, we let $d \geq 1$ and write $|\cdot|$ for the Euclidean norm on \mathbb{R}^d and $\langle \cdot, \cdot \rangle$ for the scalar product. Given a vector $v \in \mathbb{R}^d$ and $\alpha > 0$, we write $v^{(\alpha)} := |v|^{\alpha-1}v$. We let $\mathcal{X}, \mathcal{Y} \subseteq \mathbb{R}^d$ denote compact sets and write $\mathcal{C}(\mathcal{X})$ for the set of continuous functions on \mathcal{X} , $\mathcal{P}(\mathcal{X})$ for the set of Borel probability measures on \mathcal{X} , $R_{\mathcal{X}}$ the smallest positive real such that $\mathcal{X} \subseteq B(0, R_{\mathcal{X}})$, and $\text{diam}(\mathcal{X})$ for the diameter of \mathcal{X} . Unless otherwise stated, we always tacitly assume that $\rho \in \mathcal{P}(\mathcal{X})$ and $\mu, \nu \in \mathcal{P}(\mathcal{Y})$. Given $\psi \in \mathcal{C}(\mathcal{Y})$ and $\mu \in \mathcal{P}(\mathcal{Y})$, we write $\langle \mu | \psi \rangle = \int_{\mathcal{Y}} \psi d\mu$ for the duality pairing (which we also use for signed measures). When \mathcal{Y} is finite, $\mathcal{Y} = \{y_i\}_{i=1}^n$, we naturally identify functions $\psi \in \mathcal{C}(\mathcal{Y})$ with vectors $(\psi_i)_{i=1}^n = (\psi(y_i))_{i=1}^n \in \mathbb{R}^n$ and probabilities $\mathcal{P}(\mathcal{Y})$ with the simplex, i.e., $(\mu_i)_{i=1}^n = \{\mu(y_i)\}_{i=1}^n \in [0, 1]^n$ with $\sum_{i=1}^n \mu_i = 1$. We write $\nabla = (\partial_i)_{i=1}^n$ for the gradient operator on $\mathcal{C}(\mathcal{Y})$ induced by such identification with \mathbb{R}^n , and similarly $\nabla^2 = (\partial_{i,j}^2)_{i,j=1}^n$ for the Hessian. Let us stress that these are in fact functional derivatives, i.e., $\partial_i = \frac{\partial}{\partial \psi_i}$ and not partial derivatives with respect to the variable y_i .

We always tacitly identify any absolutely continuous measure (with respect to the Lebesgue measure) with its density. In particular, we say that λ is log-concave if it has a density $\lambda = e^{-V}$ for some convex lower-semicontinuous $V : \mathbb{R} \rightarrow [0, \infty]$. The support of a log-concave measure is necessarily convex. Our main results apply as well to singular log-concave measures, arguing by approximations via convolutions, but we omit the details to keep the exposition simple.

We write $\|\cdot\|_{L^p(\rho)}$ for the L^p norm with respect to the measure ρ and simply $\|\cdot\|_p$ if ρ is understood. We occasionally use the notation $\|\cdot\|_{\infty}$ for the uniform norm. Similarly, we write

$$(2.1) \quad \mathfrak{m}_{\rho}(f) = \int_{\mathcal{X}} f d\rho$$

$$(2.2) \quad \text{Var}_{\rho}(f) = \int_{\mathcal{X}} \left(f - \int_{\mathcal{X}} f d\rho \right)^2 d\rho$$

for the mean and variance of $f : \mathcal{X} \rightarrow \mathbb{R}$ – assuming that the ρ -mean of f is well-defined. We recall the characterization

$$(2.3) \quad \text{Var}_{\rho}(f) = \min_{\lambda \in \mathbb{R}} \int_{\mathcal{X}} (f - \lambda)^2 d\rho,$$

which easily yields that if $\tilde{\rho} \in \mathcal{P}(\mathcal{X})$ is absolutely continuous with respect to ρ , with a uniformly bounded density $d\tilde{\rho}/d\rho$, then

$$(2.4) \quad \text{Var}_{\tilde{\rho}}(f) \leq \left(\sup_{\mathcal{X}} \frac{d\tilde{\rho}}{d\rho} \right) \text{Var}_{\rho}(f).$$

Moreover, $f \mapsto \text{Var}_{\rho}(f)$ is easily seen to be convex. We write

$$(2.5) \quad \text{osc}(f) := \sup_{x \in \mathcal{X}} f(x) - \inf_{x \in \mathcal{X}} f(x)$$

for the oscillation of f . We notice the elementary inequality

$$(2.6) \quad \text{osc}(f) \leq \text{diam}(\mathcal{X}) \text{Lip}(f),$$

where $\text{Lip}(f)$ denotes the Lipschitz constant of f (possibly ∞ if f is not Lipschitz continuous).

2.2. Optimal transport and its entropic regularization. Given a continuous cost function $c : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ we define the optimal transport problem with cost c as the following minimization problem:

$$(OT_c) \quad \mathcal{T}^c(\rho, \mu) := \inf_{\pi \in \Pi(\rho, \mu)} \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi(x, y)$$

where $\Pi(\rho, \mu)$ denotes the set of couplings between ρ and μ . For any $\varepsilon > 0$, we introduce the ε -entropic regularized optimal transport problem, with cost c , as the following minimization problem:

$$(EOT_{c, \varepsilon}) \quad \mathcal{T}^{c, \varepsilon}(\rho, \mu) := \inf_{\pi \in \Pi(\rho, \mu)} \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi(x, y) + \varepsilon \text{Ent}(\pi || \rho \otimes \mu)$$

where Ent denotes the relative entropy functional.

Note also that for $\sigma \in \mathcal{P}(\mathcal{Y})$ such that $\mu \ll \sigma$ and $\sigma \ll \mu$, it admits the following rewriting

$$(2.7) \quad \mathcal{T}^{c, \varepsilon}(\rho, \mu) := \inf_{\pi \in \Pi(\rho, \mu)} \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi(x, y) + \varepsilon \text{Ent}(\pi || \rho \otimes \sigma) - \varepsilon \text{Ent}(\mu || \sigma),$$

which will be quite useful in our derivations.

Since for simplicity we assume that \mathcal{X}, \mathcal{Y} are compact and c is continuous, it always holds $\mathcal{T}^c(\rho, \mu) < \infty$ for every $\rho \in \mathcal{P}(\mathcal{X}), \mu \in \mathcal{P}(\mathcal{Y})$. When $p \geq 1$ and c is the so-called p -cost, $c(x, y) = \frac{1}{p}|x - y|^p$ for $x \in \mathcal{X}, y \in \mathcal{Y}$, we write \mathcal{T}^p instead of \mathcal{T}^c and notice that

$$(2.8) \quad \mathcal{T}^p(\rho, \mu) = \frac{1}{p} \mathcal{W}_p(\rho, \mu)^p.$$

When $p = 2$, writing $|x - y|^2 = |x|^2 + |y|^2 - 2\langle x, y \rangle$, we collect the identity

$$(2.9) \quad \mathcal{T}^2(\rho, \mu) = \frac{1}{2} \int_{\mathcal{X}} |x|^2 d\rho(x) + \frac{1}{2} \int_{\mathcal{Y}} |y|^2 d\mu(y) + \mathcal{T}'(\rho, \mu),$$

where $'$ denotes the linear cost $(x, y) \mapsto -\langle x, y \rangle$.

We will make crucial use of the following result on the so-called displacement interpolation between two absolutely continuous probabilities, which is well-known for the quadratic case (see e.g. McCann's proof of Prékopa-Leindler [McC94, appendix D]), while for general power cost a proof can be found as a byproduct of the proof of [AGS08, proposition 9.3.9] applied to the (opposite of) the differential entropy functional $\mathcal{F}(\rho) = \int \rho \log(\rho)$.

Lemma 2.1. *Let $p > 1$, $\rho_0, \rho_1 \in \mathcal{P}(\mathcal{X})$ be absolutely continuous and let T be the optimal transport map for $\mathcal{W}_p(\rho_0, \rho_1)$. For $t \in [0, 1]$, let $T_t(x) = (1-t)x + tT(x)$ and set $\rho_t = (T_t)_\# \rho_0$ the displacement interpolation between ρ_0 and ρ_1 . Then ρ_t is absolutely continuous and*

$$(2.10) \quad \rho_t(T_t(x)) \leq \rho_0(x)^{1-t} \rho_1(T(x))^t \quad \text{for } \rho_0\text{-a.e. } x \in \mathcal{X}.$$

We also recall the following upper bound for the quadratic Wasserstein distance between two densities, in terms of the L^2 norm of their difference, with respect to a log-concave reference measure ρ .

Lemma 2.2. *Let $\rho \in \mathcal{P}(\mathbb{R}^d)$ be log-concave. Then, there exists $C = C(\rho) < \infty$ such that the following holds. For every $\mu_0 = f_0\rho$, $\mu_1 = f_1\rho \in \mathcal{P}(\mathbb{R}^d)$, both absolutely continuous with respect to ρ ,*

$$(2.11) \quad \mathcal{W}_2(\mu_0, \mu_1)^2 \leq \frac{C}{\text{ess-inf } f_1} \|f_1 - f_0\|_{L^2(\rho)}^2.$$

Inequalities of this kind have been used by many authors, see e.g. [Pey18; AST19; GHO21; Led17] and can be proved combining the Benamou-Brenier formula for \mathcal{W}_2 and the Poincaré-Wirtinger inequality with respect to the measure ρ .

2.3. Dual formulations and Kantorovich functional. Since the “source” measure $\rho \in \mathcal{P}(\mathcal{X})$ plays a distinguished role in our results, we write the dual formulation of the optimal transport problem in the following asymmetric way (also called semi-dual in [Del22]):

$$(2.12) \quad \mathcal{T}^c(\rho, \mu) = \sup_{\psi \in \mathcal{C}(\mathcal{Y})} \left\{ \int_{\mathcal{X}} \psi^c d\rho + \int_{\mathcal{Y}} \psi d\mu \right\},$$

where

$$(2.13) \quad \psi^c(x) := \inf_{y \in \mathcal{Y}} \{c(x, y) - \psi(y)\}$$

denotes the c -transform of ψ . For the ε -entropic regularized version, duality reads [MG20; NW22]

$$(2.14) \quad \mathcal{T}^{c, \varepsilon}(\rho, \mu) = \sup_{\psi \in \mathcal{C}(\mathcal{Y})} \left\{ \int_{\mathcal{X}} \psi^{c, \varepsilon} d\rho + \int_{\mathcal{Y}} \psi d\mu \right\} - \varepsilon \text{Ent}(\mu || \sigma),$$

where the (c, ε) -transform of ψ is given by

$$(2.15) \quad \psi^{c, \varepsilon}(x) := -\varepsilon \log \left(\int_{\mathcal{Y}} \exp \left(-\frac{c(x, y) - \psi(y)}{\varepsilon} \right) d\sigma \right).$$

We notice that this does not depend anymore upon μ . Being able to use the same definition of (c, ε) -transform for different target measure μ is a key element of the proof. Symmetrically, c -transform and (c, ε) -transform can be defined for continuous functions on \mathcal{X} . When a function in $\mathcal{C}(\mathcal{Y})$ can be written as a c -transform (resp. (c, ε) -transform) we say it is c -concave (resp. (c, ε) -concave function). In this formulation, a c -concave or (c, ε) -concave optimizer $\psi \in \mathcal{C}(\mathcal{Y})$ is called a Kantorovich potential (for $\mathcal{T}^c(\rho, \mu)$ or $\mathcal{T}^{c, \varepsilon}(\rho, \mu)$), and $(\phi, \psi) = (\psi^c, \psi) \in \mathcal{C}(\mathcal{X}) \times \mathcal{C}(\mathcal{Y})$ is called a pair of Kantorovich potentials (and similarly for $\varepsilon > 0$). We notice that such transforms define monotone decreasing maps, i.e., for $\psi, \tilde{\psi} \in \mathcal{C}(\mathcal{Y})$

$$(2.16) \quad \psi(y) \leq \tilde{\psi}(y) \quad \text{for every } y \in \mathcal{Y} \quad \Rightarrow \quad \psi^{c, \varepsilon}(x) \geq \tilde{\psi}^{c, \varepsilon}(x) \quad \text{for every } x \in \mathcal{X},$$

and are concave,

$$(2.17) \quad ((1-t)\tilde{\psi} + t\psi)^{c, \varepsilon}(x) \geq (1-t)\tilde{\psi}^{c, \varepsilon}(x) + t\psi^{c, \varepsilon}(x) \quad \text{for every } t \in [0, 1], x \in \mathcal{X},$$

as a consequence of Hölder’s inequality. Moreover, $(\psi + \lambda)^{c, \varepsilon} = \psi^{c, \varepsilon} - \lambda$ for every constant $\lambda \in \mathbb{R}$.

We introduce, for $\phi_0, \phi_1 \in \mathcal{C}(\mathcal{X})$, the quantity

$$(2.18) \quad M_{\phi_0, \phi_1}^c = \sup_{t \in [0, 1]} \{\text{osc}(\phi_t)\},$$

where $\phi_t := ((1-t)\phi_0^c + t\phi_1^c)^c$.

Example 2.3. If the cost c is Lipschitz continuous, then any function $\phi = \psi^c$ that is a c -transform is Lipschitz continuous. In the case of the p -cost, we obtain that $\text{Lip}(\phi) \leq (R_{\mathcal{X}} + R_{\mathcal{Y}})^{p-1}$, hence $\text{osc}(\phi) \leq 2R_{\mathcal{X}}(R_{\mathcal{X}} + R_{\mathcal{Y}})^{p-1}$. In particular, we can always bound from above

$$(2.19) \quad M_{\phi_0, \phi_1}^c \leq 2R_{\mathcal{X}}(R_{\mathcal{X}} + R_{\mathcal{Y}})^{p-1}.$$

As in [Del22] we introduce the c -Kantorovich functional

$$(2.20) \quad \psi \in \mathcal{C}(\mathcal{Y}) \quad \mapsto \quad \mathcal{K}^c(\psi) = - \int_{\mathcal{X}} \psi^c(x) d\rho(x)$$

and similarly the (c, ε) -Kantorovich functional

$$(2.21) \quad \psi \in \mathcal{C}(\mathcal{Y}) \quad \mapsto \quad \mathcal{K}^{c, \varepsilon}(\psi) = - \int_{\mathcal{X}} \psi^{c, \varepsilon}(x) d\rho(x).$$

We remark that the notation does not highlight the dependence upon ρ , which will be clear from the context. Both functionals $\mathcal{K}^c, \mathcal{K}^{c, \varepsilon}$ are convex, and the duality formulas read

$$(2.22) \quad \begin{aligned} \mathcal{T}^c(\rho, \mu) &= \sup_{\psi \in \mathcal{C}(\mathcal{Y})} \{ \langle \mu | \psi \rangle - \mathcal{K}^c(\psi) \} \\ \mathcal{T}^{c, \varepsilon}(\rho, \mu) &= \sup_{\psi \in \mathcal{C}(\mathcal{Y})} \{ \langle \mu | \psi \rangle - \mathcal{K}^{c, \varepsilon}(\psi) \} - \varepsilon \text{Ent}(\mu | \sigma). \end{aligned}$$

Remark 2.4. With the above definitions, we have convergence of the regularized problems to the usual transport problem as $\varepsilon \downarrow 0$, as discussed in [NW22]. More precisely, since both \mathcal{X} and \mathcal{Y} are bounded and the cost function is (uniformly) continuous, any family of functions that are (c, ε) -transforms has a uniform modulus of continuity, also uniformly with respect to $\varepsilon > 0$. If we require that optimal potential $\phi = \psi^{c, \varepsilon}$ have e.g. zero mean with respect to ρ , we get by Arzelà-Ascoli that the families are relatively compact, hence up to a subsequence they converge uniformly. Reversing the roles of ϕ and ψ leads to compactness also for the optimizers ψ 's. If moreover uniqueness of the (zero mean) optimal potentials holds, as is the case of p -costs for $p > 1$ in our setting, one obtains uniform convergence of the optimizers (and also of the costs), otherwise one can still extract a uniformly converging subsequence.

2.4. Semi-discrete problem. It will be convenient to further approximate by assuming that \mathcal{Y} is a finite set. The following result extends [Del22, Lemma 2.7] and allows us to pass from such semi-discrete case to the general case of \mathcal{Y} compact.

Lemma 2.5. *Let $\mathcal{X} \subset \mathbb{R}^d$ be compact and convex and $\mathcal{Y} \subset \mathbb{R}^d$ be compact, $\rho \in \mathcal{P}(\mathcal{X})$ be absolutely continuous, $\mu_0, \mu_1 \in \mathcal{P}(\mathcal{Y})$ and $c : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$ be continuous. For $k \in \{0, 1\}$, let $\phi_k \in \mathcal{C}(\mathcal{X})$ assume that there exists a unique Kantorovich potential with zero ρ -mean for $\mathcal{T}^c(\rho, \mu_k)$ and ψ_k its c -transform. Then, there exists sequences of finite sets $(\mathcal{Y}^n)_{n=1}^{\infty}$, discrete measures $(\mu_k^n)_{n=1}^{\infty}$, Kantorovich potentials $(\phi_k^n, \psi_k^n)_{n=1}^{\infty}$ such that, for $k \in \{0, 1\}$,*

- (1) $\text{supp}(\mu_k^n) = \mathcal{Y}^n$ for every $n \geq 1$,
- (2) $\|\phi_k^n - \phi_k\|_{\infty} \rightarrow 0$ and $\|\psi_k^n - \psi_k\|_{\infty} \rightarrow 0$ as $n \rightarrow \infty$
- (3) it holds $\langle \mu_k^n | \psi_1^n - \psi_0^n \rangle \rightarrow \langle \mu_k | \psi_1 - \psi_0 \rangle$ as $n \rightarrow \infty$.

Without the uniqueness assumption, one may still extract a converging subsequence to a pair of Kantorovich potentials.

Proof. For any $n \geq 1$, consider a finite partition $\mathcal{Y} = \bigsqcup_{1 \leq i \leq n} \mathcal{Y}_i^n$ with $\varepsilon_n := \max_{1 \leq i \leq n} \text{diam}(\mathcal{Y}_{i, N})$ infinitesimal as $n \rightarrow \infty$. Define $\mathcal{Y}^n = \{y_i^n\}_{i=1}^n$, with $y_i^n \in \mathcal{Y}_i^n$ and set

$$(2.23) \quad \mu_k^n = \sum_{i=1}^n \left[\left(1 - \frac{1}{n}\right) \mu_k(\mathcal{Y}_i^n) + \frac{1}{n^2} \right] \delta_{y_i^n},$$

so that (1) is verified.

Letting ϕ_k^n be the Kantorovich potential for $\mathcal{T}^c(\rho, \mu_k^n)$ with zero ρ -mean, then (2) holds by Arzelà-Ascoli and the fact that the limiting potential is unique.

To show (3), notice that

$$(2.24) \quad \mathcal{W}_1(\mu_k, \mu_k^n) \leq \varepsilon_n + \frac{\text{diam}(\mathcal{Y})}{n},$$

so that $\mu_k^n \rightarrow \mu_k$ weakly and therefore the limit holds, since the functions ψ_k^n converge uniformly. \square

In the case where \mathcal{Y} is finite, and fixing σ as the uniform measure on \mathcal{Y} , we recall some formulas for the derivatives of $\mathcal{K}^{c,\varepsilon}$ which is easily seen to be a smooth map from $\mathcal{C}(\mathcal{Y})$ (identified with \mathbb{R}^n) to \mathbb{R} . These are proved in [Del22, lemma 4.2, proposition 3.6] for the case of the “linear” cost $'$ defined above (2.9), but the proof can straightforwardly be generalized to any cost.

$$(2.25) \quad \partial_i(\psi^{c,\varepsilon})(x) = -\frac{\exp\left(\frac{\psi_i - c(x, y_i)}{\varepsilon}\right) \sigma_i}{\sum_{j=1}^n \exp\left(\frac{\psi_j - c(x, y_j)}{\varepsilon}\right) \sigma_j}, \quad \text{for } i = 1 \dots, n,$$

which we notice defines for every $x \in \mathcal{X}$ a probability vector, i.e., $-\partial_i(\psi^{c,\varepsilon})(x) \in [0, 1]$ and $-\sum_i \partial_i(\psi^{c,\varepsilon})(x) = 1$, which we naturally identify with a probability on \mathcal{Y} , and plays the role of a generalized optimal transport map in this context. We write therefore

$$(2.26) \quad \pi_{\varepsilon, \psi}^c(y_i | x) := -\partial_i(\psi^{c,\varepsilon})(x).$$

which in turn play a role in the expression for the gradient and Hessian of $\mathcal{K}^{c,\varepsilon}$:

$$(2.27) \quad \begin{aligned} \langle \nabla \mathcal{K}^{c,\varepsilon}(\psi), v \rangle &= \int_{\mathcal{X}} \mathbf{m}_{\pi_{\varepsilon, \psi}^c(\cdot | x)}(v) d\rho(x), \\ \langle v, \nabla^2 \mathcal{K}^{c,\varepsilon}(\psi)v \rangle &= \frac{1}{\varepsilon} \int_{\mathcal{X}} \text{var}_{\pi_{\varepsilon, \psi}^c(\cdot | x)}(v) d\rho(x). \end{aligned}$$

We also introduce the following “partition function”

$$(2.28) \quad \psi \in \mathcal{C}(\mathcal{Y}) \mapsto \mathcal{Z}_\beta(\psi) := \int_{\mathcal{X}} e^{\beta \psi^{c,\varepsilon}} d\rho.$$

As with the function \mathcal{K} , the notation does not highlight the dependence upon ρ , which will be clear from the context. The function \mathcal{Z} is smooth and in particular the following formula

$$(2.29) \quad \langle v, \nabla^2 \log \mathcal{Z}_\beta(\psi)v \rangle = -\frac{\beta}{\varepsilon} \int_{\mathcal{X}} \text{Var}_{\pi_{\varepsilon, \psi}^c(\cdot | x)}(v) d\rho_\psi^\varepsilon(x) + \beta^2 \text{Var}_{\rho_\psi^\varepsilon} \left(\mathbf{m}_{\pi_{\varepsilon, \psi}^c(\cdot | x)}(v) \right),$$

holds with

$$(2.30) \quad \rho_{\beta, \psi}^\varepsilon := \frac{1}{\mathcal{Z}_\beta(\psi)} e^{\beta \psi^{c,\varepsilon}} \rho.$$

Remark 2.6. In most proofs, to keep notation simple we will fix $\beta = 1$ and write $\mathcal{Z} := \mathcal{Z}_1$.

Finally, recalling that we denote with $'$ the cost $(x, y) \mapsto -\langle x, y \rangle$, we obtain

$$(2.31) \quad \psi'(x) = -\sup_{y \in \mathcal{Y}} \{ \langle x, y \rangle + \psi(y) \}, \quad \psi'^{\varepsilon}(x) = -\varepsilon \log \left(\int_{\mathcal{Y}} \exp \left(\frac{\langle x, y \rangle + \psi(y)}{\varepsilon} \right) d\sigma(y) \right).$$

Remark 2.7. Let $f \in L^1(\rho)$ and $\tilde{c}(x, y) = c(x, y) + f(x)$. Then the following equalities hold, with equality of the set of minimizers.

$$(2.32) \quad \mathcal{T}_c(\rho, \mu) = \mathcal{T}_{\tilde{c}}(\rho, \mu) + \mathbf{m}_\rho(f)$$

$$(2.33) \quad \mathcal{T}_{c,\varepsilon}(\rho, \mu) = \mathcal{T}_{\tilde{c},\varepsilon}(\rho, \mu) + \mathbf{m}_\rho(f)$$

Moreover, for any $\psi \in \mathcal{C}(\mathcal{Y})$,

$$(2.34) \quad \mathcal{K}^{c,\varepsilon}(\psi) = \mathcal{K}^{\tilde{c},\varepsilon}(\psi) + \mathbf{m}_\rho(f)$$

In particular, if (ϕ_c, ψ_c) (resp. $(\phi_{c,\varepsilon}, \psi_{c,\varepsilon})$) denotes a couple of optimal potentials for the cost c for (OT $_c$), (resp. for (EOT $_{c,\varepsilon}$)),

$$(2.35) \quad (\phi_{\tilde{c}}, \psi_{\tilde{c}}) = (\phi_c + f, \psi_c)$$

$$(2.36) \quad (\phi_{\tilde{c},\varepsilon}, \psi_{\tilde{c},\varepsilon}) = (\phi_{c,\varepsilon} + f, \psi_{c,\varepsilon})$$

2.5. Fractional Sobolev spaces. We start by recalling a definition of the spaces $W^{s,p}(\Omega)$, when the order s is not a integer and $\Omega \subseteq \mathbb{R}^d$ is a bounded domain with Lipschitz boundary. For $0 < s < 1$ and $1 \leq p < \infty$, we set

$$(2.37) \quad [u]_{s,p}^p := \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{d+sp}} dx dy$$

and

$$(2.38) \quad \|u\|_{W^{s,p}} := \|u\|_{L^p(\Omega)} + [u]_{s,p} \quad W^{s,p}(\Omega) := \{u \in L^p(\Omega) \text{ and } \|u\|_{W^{s,p}} < \infty\}.$$

When $s > 1$ is not an integer, we write $s = m + \sigma$, with $m \in \mathbb{N}$, and $\sigma \in (0, 1)$ and let

$$(2.39) \quad W^{s,p}(\Omega) := \{u \in W^{m,p}(\Omega); D^m u \in W^{\sigma,p}(\Omega)\}$$

normed with

$$(2.40) \quad \|u\|_{W^{s,p}} := \|u\|_{W^{m,p}} + [D^m u]_{W^{\sigma,p}}.$$

When $p = 2$ we usually note $H^s(\Omega) := W^{s,2}(\Omega)$.

We recall the fractional Gagliardo-Nirenberg inequality (see [BM18, Theorem 1]): for any $0 \leq s_1 < s_2$, $p, p_1, p_2 \in [1, \infty]$, with $\theta \in (0, 1)$,

$$(2.41) \quad s := \theta s_1 + (1 - \theta)s_2, \quad \text{and} \quad \frac{1}{p} = \theta \frac{1}{p_1} + (1 - \theta) \frac{1}{p_2},$$

then

$$(2.42) \quad \|u\|_{W^{s,p}(\Omega)} \leq C \|u\|_{W^{s_1,p_1}(\Omega)}^{\theta} \|u\|_{W^{s_2,p_2}(\Omega)}^{1-\theta}$$

provided that the following condition *fails*:

$$(2.43) \quad s_2 \text{ is an integer, } p_2 = 1 \text{ and } s_1 - \frac{1}{p_1} \geq s_2 - \frac{1}{p_2}.$$

By iterations, we may also consider $\theta_1, \theta_2, \theta_3 \in (0, 1)$ with $\theta_1 + \theta_2 + \theta_3 = 1$, $0 \leq s_1 < s_2 < s_3$ and $p, p_1, p_2, p_3 \in [1, \infty]$ such that

$$(2.44) \quad s := \sum_{i=1}^3 \theta_i s_i, \quad \frac{1}{p} = \sum_{i=1}^3 \frac{\theta_i}{p_i},$$

provided that both (2.43) and the following condition *fail*:

$$(2.45) \quad s_3 \text{ is an integer, } p_3 = 1 \text{ and } s - \frac{1}{p} \geq s_3 - \frac{1}{p_3}.$$

In our application, we need the following special case: $s_1 = 0 < s_2 = 1 < s_3 < 2$ and $p_1 = 2, p_2 = \infty, p_3 = 1$, and $s = 1, p = 2$, so that $\theta_1, \theta_2, \theta_3$ must satisfy $\theta_1 + \theta_2 + \theta_3 = 1$ and

$$(2.46) \quad 1 = \theta_2 + \theta_3 s_3 \quad \text{and} \quad \frac{1}{2} = \frac{\theta_1}{2} + \theta_3.$$

These conditions yield

$$(2.47) \quad \theta_1 = 1 - \frac{2}{1 + s_3}, \quad \theta_2 = \theta_3 = \frac{1}{1 + s_3}.$$

which must be also complemented with the failure of both (2.43), which fails because $p_2 = \infty$ and (2.44), which fails because s_3 is not an integer. In conclusion, writing $r \in (1, 2)$ instead of s_3 , we have the following inequality:

$$(2.48) \quad \|u\|_{H^1(\Omega)} \leq C \|u\|_{L^2(\Omega)}^{1-\frac{2}{1+r}} \|u\|_{W^{1,\infty}(\Omega)}^{\frac{1}{1+r}} \|u\|_{W^{r,1}(\Omega)}^{\frac{1}{1+r}}.$$

3. QUANTITATIVE STABILITY OF POTENTIALS

Aim of this section is to establish our first main result, [Theorem 1.1](#). To simplify our argument and highlight the differences with the results in [\[Del22\]](#), we assume that $\rho = \lambda$ is a log-concave probability measure with bounded (convex) support and point out in [Remark 3.5](#) below the minor modifications for the general case of ρ equivalent to λ . We also split our argument into three subsections. We deal first with the quadratic case, in particular recovering and slightly extending the results in [\[Del22\]](#), but mainly in order to point out where our arguments differ and allow for extensions respectively to the case $p \geq 2$ ([Section 3.2](#)) and $1 < p < 2$ ([Section 3.3](#)).

3.1. The quadratic case. In this section, we prove [Theorem 1.1](#) in the case $p = 2$. In view of the identity [\(2.9\)](#), it is sufficient to argue in the case of the linear cost $c(x, y) = -\langle x, y \rangle$ that we denote with $'$. Therefore, we establish the following result.

Theorem 3.1. *Let ρ be a log-concave probability measure with bounded convex support $\mathcal{X} \subseteq \mathbb{R}^d$ and let $\mathcal{Y} \subseteq \mathbb{R}^d$ be compact. Given $\mu_0, \mu_1 \in \mathcal{P}(\mathcal{Y})$, let $(\phi_{\mu_0}, \psi_{\mu_0}), (\phi_{\mu_1}, \psi_{\mu_1}) \in \mathcal{C}(\mathcal{X}) \times \mathcal{C}(\mathcal{Y})$ be pairs of Kantorovich potentials respectively for $\mathcal{T}'(\rho, \mu_0)$ and $\mathcal{T}'(\rho, \mu_1)$, then*

$$(3.1) \quad \text{Var}_\rho(\phi_{\mu_0} - \phi_{\mu_1}) \leq 2M'_{\phi_{\mu_0}, \phi_{\mu_1}} \langle \mu_0 - \mu_1 | \psi_{\mu_0} - \psi_{\mu_1} \rangle.$$

Recall that, by [\(2.19\)](#) for $p = 2$, we have $M'_{\phi_{\mu_0}, \phi_{\mu_1}} \leq 2R_{\mathcal{X}}(R_{\mathcal{X}} + R_{\mathcal{Y}})$.

Proof of [Theorem 1.1](#) for $p = 2$. Let us now prove [Theorem 1.1](#) from [Theorem 3.1](#). Being a Kantorovich potential for the $'$ -cost, ψ_{μ_0} is the $'$ -transform of $\phi_{\mu_0} \in \mathcal{C}(\mathcal{X})$. As a consequence, ψ_{μ_0} is $R_{\mathcal{X}}$ -Lipschitz. Similarly, ψ_{μ_1} is $R_{\mathcal{X}}$ -Lipschitz. Then, by Kantorovich-Rubinstein duality formula, one gets

$$(3.2) \quad \langle \mu_0 - \mu_1 | \psi_{\mu_0} - \psi_{\mu_1} \rangle \leq R_{\mathcal{X}} W_1(\mu_0, \mu_1)$$

which, put along with [Theorem 3.1](#) yields [Theorem 1.1](#). □

We collect also as a corollary the following slightly different bound, where we change the point of view by taking a single target measure $\mu \in \mathcal{P}(\mathcal{Y})$ and consider any $\psi \in \mathcal{C}(\mathcal{Y})$ (not necessarily a potential).

Corollary 3.2. *Let ρ be a log-concave probability measure with bounded convex support $\mathcal{X} \subseteq \mathbb{R}^d$ and let $\mathcal{Y} \subseteq \mathbb{R}^d$ be compact. Given $\mu \in \mathcal{P}(\mathcal{Y})$, let $(\phi_\mu, \psi_\mu) \in \mathcal{C}(\mathcal{X}) \times \mathcal{C}(\mathcal{Y})$ be a pair of Kantorovich potentials for $\mathcal{T}'(\rho, \mu)$, let $\psi \in \mathcal{C}(\mathcal{Y})$ and $\phi = \psi'$, it holds*

$$(3.3) \quad \text{Var}_\rho(\phi_\mu - \phi) \leq 2M_{\phi_\mu, \phi} [\mathcal{T}'(\rho, \mu) - (\langle \mu | \psi \rangle + \langle \rho | \phi \rangle)].$$

Proof of [Corollary 3.2](#). \mathcal{K}' is a convex functional defined on $\mathcal{C}(\mathcal{Y})$ and therefore differentiable in duality with $\mathcal{M}(\mathcal{Y})$ the space of Radon measures. Moreover, one can easily check that for any $\mu \in \mathcal{P}_2(\mathcal{Y})$, if ψ_μ denotes its Kantorovich potential, then $\psi_\mu \in D(\partial \mathcal{K}')$ and $\partial \mathcal{K}'(\psi_\mu) = \mu$. One concludes by monotonicity of the slopes. □

We now turn to the proof of [Theorem 3.1](#). It is known or easy to check that while strict convexity yields uniqueness in a minimization problem, strong convexity yields explicit stability estimates. Following the idea developed in [\[Del22\]](#), we aim at a suitable convexity property for the functional \mathcal{K}' , which we obtain first in the ε -entropic semi-discrete (i.e., for finite \mathcal{Y}) case, then via an approximation argument for the general case of \mathcal{Y} compact. However, to highlight the differences (and the improvements) with respect to [\[Del22\]](#), we split the proof of [Theorem 3.1](#) into three steps:

Step 1 (log-concavity of \mathcal{Z}). Arguing for fixed $\varepsilon > 0$ and \mathcal{Y} finite, we establish log-concavity of the “partition function” \mathcal{Z} introduced in [\(2.28\)](#). Unlike [\[Del22, proposition 4.6\]](#), where this is obtained as a “black box” application of the Prékopa-Leindler inequality when ρ is the Lebesgue measure on a convex bounded \mathcal{X} , we obtain this property for a general log-concave measure ρ through an optimal transport argument ([Proposition 3.3](#)), which itself

can be used to prove the Prékopa-Leindler inequality, but is more amenable to further generalizations.

Step 2 (modified convexity of \mathcal{K}'^ε). By direct computation, we go from the log-concavity of \mathcal{Z} to a strong convexity-like inequality on \mathcal{K}'^ε , that reads for $\psi_0, \psi_1 \in \mathcal{C}(\mathcal{Y})$,

$$(3.4) \quad \text{Var}_\rho(\psi_1'^\varepsilon - \psi_0'^\varepsilon) \leq C \langle \nabla \mathcal{K}'^\varepsilon(\psi_1) - \nabla \mathcal{K}'^\varepsilon(\psi_0) | \psi_1 - \psi_0 \rangle,$$

for a suitable constant C . Let us stress that the inequality above provides quantitative convexity, but not strong convexity in the usual sense, for the dual potentials appear in the left hand side. Still, our use of integration against ρ in the first step allows us to yield a constant independent of upper and lower bounds on the density of ρ with respect to the Lebesgue measure, going beyond the case covered in [Del22]. Moreover, when compared to [Del22], where the (stronger) inequality with $\text{Var}_{(\mu+\nu)/2}(\psi_\mu - \psi_\nu)$ in the left hand side is established, our proof does not rely on linearity of the cost (while [Del22, lemma 2.3] apparently does).

Step 3 (approximation and scaling) This final step consists in moving from the ε -entropic regularization to the classical optimal transport by letting $\varepsilon \downarrow 0$, as briefly discussed in Remark 2.4, and then via an approximation argument to go from the discrete case of finite \mathcal{Y} to the general case of compact \mathcal{Y} using Lemma 2.5. Finally, thanks to a scaling argument we improve the constant in the stability inequality: indeed, it is sufficient to introduce an “inverse temperature” parameter $\beta > 0$, defining

$$(3.5) \quad \mathcal{Z}_\beta(\psi) := \int_{\mathcal{X}} e^{\beta \psi'^\varepsilon} d\rho$$

and, after repeating the first two steps, we optimize upon β . This step does not differ substantially from [Del22, section 4.5, section 2.4, proposition 3.3]. The scaling argument, also in [Del22], is actually of great importance for the case $1 < p < 2$ and will be detailed in Section 3.3, while adapting it to the present setting is straightforward.

In view of the scheme detailed above, we establish the following proposition (settling Step 1).

Proposition 3.3 (log-concavity of \mathcal{Z}). *Let ρ be a log-concave measure with (convex) bounded support \mathcal{X} , let $\mathcal{Y} \subseteq \mathbb{R}^d$ be finite, $\varepsilon > 0$ and $\beta > 0$. For every $\psi_0, \psi_1 \in \mathcal{C}(\mathcal{Y})$ and $t \in [0, 1]$, it holds*

$$(3.6) \quad \log \mathcal{Z}_\beta((1-t)\psi_0 + t\psi_1) \geq (1-t) \log \mathcal{Z}_\beta(\psi_0) + t \log \mathcal{Z}_\beta(\psi_1).$$

Proof. Recall that we identify $\mathcal{C}(\mathcal{Y})$ with \mathbb{R}^n , since $\mathcal{Y} = \{y_i\}_{i=1}^n$, and write for brevity $\psi_i := \psi(y_i)$, $\mu_i := \mu(y_i)$. To simplify the notation, we also simply write ψ' instead of ψ'^ε and argue in the case $\beta = 1$ only. Thus, given $\psi_0, \psi_1 \in \mathbb{R}^n$ and $x_0, x_1 \in \mathcal{X}$ and setting $\psi_t = (1-t)\psi_0 + t\psi_1$, we have

$$(3.7) \quad \begin{aligned} -(\psi_t)'(x_t) &= \varepsilon \log \left(\sum_{i=1}^n \exp \left(\frac{\langle x_t, y_i \rangle + (\psi_t)_i}{\varepsilon} \right) \sigma_i \right) \\ &= \varepsilon \log \left(\sum_{i=1}^n \exp \left(\frac{(1-t)\langle x_0, y_i \rangle + t\langle x_1, y_i \rangle + (1-t)(\psi_0)_i + t(\psi_1)_i}{\varepsilon} \right) \sigma_i \right). \end{aligned}$$

Using Hölder’s inequality, we get

$$(3.8) \quad -(\psi_t)'(x_t) \leq -(1-t)(\psi_0)'(x_0) - t(\psi_1)'(x_1)$$

Therefore, taking the exponential and setting $h_t(x) = \exp((\psi_t)'(x))$ we obtain the inequality

$$(3.9) \quad h_t(x_t) \geq h_0(x_0)^{1-t} h_1(x_1)^t$$

As mentioned earlier, one can now use as a black box the Prékopa-Leindler inequality for log-concave measures which yields $\|h_t\|_{L^1(\rho)} \geq \|h_0\|_{L^1(\rho)}^{1-t} \|h_1\|_{L^1(\rho)}^t$. For completeness we recall the optimal transport version of the proof. Up to renormalization, we can always

assume $\|h_0\|_{L^1(\rho)} = \|h_1\|_{L^1(\rho)} = 1$, so that h_0 and h_1 are probability densities with respect to ρ .

By [Lemma 2.1](#) applied with $p = 2$, $\rho_0 := h_0\rho$, $\rho_1 := h_1\rho$, using also the log-concavity of ρ and writing ρ_t the displacement interpolation between $h_0\rho$ and $h_1\rho$, we get for $h_0\rho$ -a.e. $x \in \mathcal{X}$ the inequality

$$\begin{aligned} \rho_t(T_t(x)) &\leq (h_0(x)\rho(x))^{1-t}(h_1(T(x))\rho(T(x)))^t \\ &\leq h_0(x)^{1-t}(h_1(T(x)))^t \rho(x)^{1-t} \rho(T(x))^t \\ (3.10) \quad &\leq h_0(x)^{1-t}(h_1(T(x)))^t \rho(T_t(x)) \\ &\leq h_t(T_t(x))\rho(T_t(x)). \end{aligned}$$

Therefore, we deduce that

$$(3.11) \quad \rho_t(x) \leq h_t(x)\rho(x) \quad \text{holds } \rho_t \text{ a.e.}$$

Integrating thus yields $\|h_t\|_{L^1(\rho)} \geq 1$. Introducing back $\|h_0\|_{L^1(\rho)}$ and $\|h_1\|_{L^1(\rho)}$, we get the Prékopa-Leindler inequality :

$$(3.12) \quad \|h_t\|_{L^1(\rho)} \geq \|h_0\|_{L^1(\rho)}^{1-t} \|h_1\|_{L^1(\rho)}^t.$$

Taking the log and noticing that $\|h_t\|_{L^1(\rho)} = \mathcal{Z}(\psi_t)$ yields the thesis. \square

In order to address *Step 2* with lighter notations, fixing $\psi_0, \psi_1 \in \mathcal{C}(\mathcal{Y})$ and $\psi_t = (1-t)\psi_0 + t\psi_1$, we write $\pi_t(\cdot|x) := \pi'_{\varepsilon, \psi_t}(\cdot|x)$, so that [\(2.29\)](#) reads

$$(3.13) \quad \langle v, \nabla^2 \log \mathcal{Z}(\psi_t)v \rangle = -\frac{1}{\varepsilon} \int_{\mathcal{X}} \text{Var}_{\pi_t(\cdot|x)}(v) d\rho_{\psi_t}^{\varepsilon}(x) + \text{Var}_{\rho_{\psi_t}^{\varepsilon}} \left(\mathbf{m}_{\pi_t(\cdot|x)}(v) \right),$$

Proposition 3.4. *Let ρ be a log-concave probability measure with compact support $\mathcal{X} \subset \mathbb{R}^d$, let $\mu \in \mathcal{P}(\mathcal{Y})$ with $\mathcal{Y} \subseteq \mathbb{R}^d$ finite and let $\varepsilon > 0$. Given $\psi_0, \psi_1 \in \mathcal{C}(\mathcal{Y})$, set*

$$(3.14) \quad \phi_t^{\varepsilon} := ((1-t)\psi_0 + t\psi_1)^{\varepsilon} \quad \text{for } t \in [0, 1],$$

Then,

$$(3.15) \quad \text{Var}_{\rho}(\phi_1^{\varepsilon} - \phi_0^{\varepsilon}) \leq C \langle \nabla \mathcal{K}'^{\varepsilon}(\psi_1) - \nabla \mathcal{K}'^{\varepsilon}(\psi_0) | \psi_1 - \psi_0 \rangle$$

with $C := \exp(2 \sup_{t \in [0,1]} \text{osc}(\phi_t^{\varepsilon}))$.

Proof. Again, we identify $\mathcal{C}(\mathcal{Y})$ with \mathbb{R}^n . From [Proposition 3.3](#), we get that $\nabla^2 \log \mathcal{Z}(\psi_t) \in \mathbb{R}^{n \times n}$ is a symmetric negative semi-definite matrix, which yields for every $v \in \mathbb{R}^n$,

$$(3.16) \quad \langle v, \nabla^2 \log \mathcal{Z}(\psi_t)v \rangle \leq 0 \quad \forall 0 \leq t \leq 1.$$

By the identity [\(2.29\)](#), we get

$$(3.17) \quad \text{Var}_{\rho_{\psi_t}^{\varepsilon}} \left(\mathbf{m}_{\pi_t(\cdot|x)}(v) \right) \leq \frac{1}{\varepsilon} \int_{\mathcal{X}} \text{Var}_{\pi_t(\cdot|x)}(v) d\rho_{\psi_t}^{\varepsilon} \quad \forall 0 \leq t \leq 1.$$

Our goal is now to change the integration against $\rho_{\psi_t}^{\varepsilon}$ to integration against ρ . Recalling the the definition of $\rho_{\psi_t}^{\varepsilon}$ in [\(2.30\)](#), we deduce

$$(3.18) \quad e^{-\text{osc}(\phi_t^{\varepsilon})} \rho_{\psi_t}^{\varepsilon} \leq \rho \leq e^{\text{osc}(\phi_t^{\varepsilon})} \rho_{\psi_t}^{\varepsilon}.$$

We remark that here is where the choice of ρ as a ‘‘reference’’ measure in the definition of \mathcal{Z} , instead e.g. of the Lebesgue measure on \mathcal{X} , turns out to be useful, as it frees us from making assumptions on the density of ρ (besides its log-concavity). Using [\(2.4\)](#) in both sides of [\(3.17\)](#) thus find, for every $0 \leq t \leq 1$,

$$(3.19) \quad \text{Var}_{\rho} \left(\mathbf{m}_{\pi_t(\cdot|x)}(v) \right) \leq \frac{e^{2\text{osc}(\phi_t^{\varepsilon})}}{\varepsilon} \int_{\mathcal{X}} \text{Var}_{\pi_t(\cdot|y)}(v) d\rho = e^{2\text{osc}(\phi_t^{\varepsilon})} \langle v, \nabla^2 \mathcal{K}'^{\varepsilon}(\psi_t)v \rangle$$

where the second identity follows from [\(2.27\)](#).

Recalling that $\phi_t^\varepsilon := (\psi_t)'^{\varepsilon}$, we have by the chain rule

$$(3.20) \quad \frac{d}{dt}\phi_t^\varepsilon(x) = \langle \nabla \psi'^{\varepsilon}(y_t|x) | \psi_1 - \psi_0 \rangle = -\mathbf{m}_{\pi_t(\cdot|x)}(\psi_1 - \psi_0)$$

hence, by convexity of the variance,

$$(3.21) \quad \begin{aligned} \text{Var}_\rho(\phi_1^\varepsilon - \phi_0^\varepsilon) &= \text{Var}_\rho\left(\int_0^1 \frac{d}{dt}\phi_t^\varepsilon dt\right) \\ &\leq \int_0^1 \text{Var}_\rho\left(\frac{d}{dt}\phi_t^\varepsilon\right) dt \\ &\leq \int_0^1 \text{Var}_\rho\left(\mathbf{m}_{\pi_t(\cdot|x)}(\psi_1 - \psi_0)\right) dt. \end{aligned}$$

Using (3.19) with $v = \psi_1 - \psi_0$, we thus obtain

$$(3.22) \quad \begin{aligned} \text{Var}_\rho(\phi_1^\varepsilon - \phi_0^\varepsilon) &\leq \int_0^1 e^{2\text{osc}(\phi_t^\varepsilon)} \langle \psi_1 - \psi_0, \nabla^2 \mathcal{K}'^\varepsilon(\psi_t)(\psi_1 - \psi_0) \rangle dt \\ &\leq C \int_0^1 \langle \psi_1 - \psi_0, \nabla^2 \mathcal{K}'^\varepsilon(\psi_t)(\psi_1 - \psi_0) \rangle dt, \\ &\leq C \langle \psi_1 - \psi_0, \int_0^1 \nabla^2 \mathcal{K}'^\varepsilon(\psi_t)(\psi_1 - \psi_0) dt \rangle \end{aligned}$$

with $C = \exp(2\sup_{t \in [0,1]} \text{osc}(\phi_t^\varepsilon))$. Finally, using that

$$(3.23) \quad \nabla^2 \mathcal{K}'^\varepsilon(\psi_t)(\psi_1 - \psi_0) = \frac{d}{dt} \nabla \mathcal{K}'^\varepsilon(\psi_t)$$

we obtain the thesis. \square

Let us recall here that we refer to Remark 2.4 and Lemma 2.5 for the approximation Step 3 and to [Del22, section 4.5, section 2.4, proposition 3.3] for detailed proofs.

Remark 3.5. The only point where log-concavity of ρ is used is in the proof of Proposition 3.3. Therefore, in order to cover the case of ρ that admits a density uniformly bounded from above and below by strictly positive constants with respect to a log-concave measure λ with bounded support \mathcal{X} ,

$$(3.24) \quad m_\rho \lambda(x) \leq \rho(x) \leq M_\rho \lambda(x) \quad \text{for } \lambda\text{-a.e. } x \in \mathcal{X},$$

we need to define \mathcal{Z} using λ instead of the measure ρ . This leads to an additional step in the argument, as log-concavity of \mathcal{Z} leads to a modified version of (3.17), which reads

$$(3.25) \quad \text{Var}_{\lambda_{\psi_t}^\varepsilon}\left(\mathbf{m}_{\pi_t(\cdot|x)}(v)\right) \leq \frac{1}{\varepsilon} \int_{\mathcal{X}} \text{Var}_{\pi_t(\cdot|x)}(v) d\lambda_{\psi_t}^\varepsilon \quad \forall 0 \leq t \leq 1.$$

It is then sufficient to replace the variance and integration with respect to $\lambda_{\psi_t}^\varepsilon$ with those with respect to $\rho_{\psi_t}^\varepsilon$, so that the proof may then proceed without further changes: by the very definition of $\lambda_{\psi_t}^\varepsilon$, we find the inequalities

$$(3.26) \quad \frac{m_\rho}{M_\rho} \lambda_{\psi_t}^\varepsilon(x) \leq \rho_{\psi_t}^\varepsilon(x) \leq \frac{M_\rho}{m_\rho} \lambda_{\psi_t}^\varepsilon(x) \quad \text{for } \lambda\text{-a.e. } x \in \mathcal{X}.$$

Therefore, recalling (2.4) we obtain that (3.17) holds with an additional factor $(M_\rho/m_\rho)^2$ in the right hand side. Similar arguments apply in the general cases we consider below, so we only focus on the case of a log-concave measure ρ to keep the exposition simple.

3.2. The case $p \geq 2$. In this section, we prove Theorem 1.1 in the case $p \geq 2$. The previous proof does not fully use the linearity of the cost but only its concavity, hence an appropriate change of cost allows to generalize it with minimum effort.

Theorem 3.6. *Let ρ be a log-concave probability measure with bounded convex support $\mathcal{X} \subseteq \mathbb{R}^d$ and let $\mathcal{Y} \subseteq \mathbb{R}^d$ be compact finally let $c : \mathcal{X} \times \mathcal{Y} \rightarrow [0, +\infty)$ be a C^2 cost function. Given $\mu_0, \mu_1 \in \mathcal{P}(\mathcal{Y})$, let $(\phi_{\mu_0}, \psi_{\mu_0}), (\phi_{\mu_1}, \psi_{\mu_1}) \in \mathcal{C}(\mathcal{X}) \times \mathcal{C}(\mathcal{Y})$ be pairs of Kantorovich*

potentials respectively for $\mathcal{T}^c(\rho, \mu_0)$ and $\mathcal{T}^c(\rho, \mu_1)$ then, there exists $C = C(c, \rho, \mathcal{Y}) < \infty$ such that

$$(3.27) \quad \text{Var}_\rho(\phi_{\mu_0} - \phi_{\mu_1}) \leq C \langle \mu_0 - \mu_1 | \psi_{\mu_0} - \psi_{\mu_1} \rangle.$$

Moreover, for the p -cost one can take $C(p, \rho, \mathcal{Y}) = 4pR_{\mathcal{X}}(R_{\mathcal{X}} + R_{\mathcal{Y}})^{p-1}$.

Proof of Theorem 1.1 for $p \geq 2$. The thesis follows from Theorem 3.6 in the same way as Theorem 1.1 for $p = 2$ followed from Theorem 3.1. \square

As in the case $p = 2$, we collect as a corollary the following slightly different bound, where we change the point of view by taking a single target measure $\mu \in \mathcal{P}(\mathcal{Y})$ and consider any $\psi \in \mathcal{C}(\mathcal{Y})$ (not necessarily a potential).

Corollary 3.7. *Let ρ be a log-concave probability measure with bounded (convex) support $\mathcal{X} \subseteq \mathbb{R}^d$, let $\mathcal{Y} \subseteq \mathbb{R}^d$ be compact and let $c : \mathcal{X} \times \mathcal{Y} \rightarrow [0, +\infty)$ be a \mathcal{C}^2 cost function. Given $\mu \in \mathcal{P}(\mathcal{Y})$, let $(\phi_\mu, \psi_\mu) \in \mathcal{C}(\mathcal{X}) \times \mathcal{C}(\mathcal{Y})$ be a pair Kantorovich potentials for $\mathcal{T}^c(\rho, \mu)$, let $\psi \in \mathcal{C}(\mathcal{Y})$ and write $\phi := \psi^c$. Then it holds, with $C = C(c, \rho, \mathcal{Y})$ the constant in Theorem 3.6,*

$$(3.28) \quad \text{Var}_\rho(\phi_\mu - \phi) \leq C [\mathcal{T}^c(\rho, \mu) - (\langle \mu | \psi \rangle + \langle \rho | \phi \rangle)].$$

This theorem follows mainly from the observation that concavity in the first variable is a sufficient condition on the cost for Step 1 to hold. Step 2 and Step 3 will then apply as they hold for c continuous and \mathcal{X} and \mathcal{Y} compact. We conclude thanks to an appropriate modification of the cost.

Proposition 3.8 (log-concavity of \mathcal{Z}). *Let \mathcal{Y} be finite. Then, the concavity of $x \rightarrow c(x, y)$ for all $y \in \mathcal{Y}$ is a sufficient condition for (3.6) to hold, for every $\psi_0, \psi_1 : \mathcal{Y} \rightarrow \mathbb{R}$.*

Proof. Assuming that $\mathcal{Y} = \{y_i\}_{i=1}^n$, and recalling that

$$(3.29) \quad -(\psi_t)^{c, \epsilon}(x_t) = \epsilon \log \left(\sum_{i=1}^n \exp \left(\frac{(\psi_t)_i - c(x_t, y_i)}{\epsilon} \right) \right),$$

with $x_t = (1-t)x_0 + tx_1$, one simply uses first the concavity of c and then Holder's inequality. \square

Definition 3.9. For any \mathcal{C}^2 cost function c , we introduce the shifted cost $\tilde{c} : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}_+$, defined by

$$(3.30) \quad \tilde{c}(x, y) = c(x, y) - \frac{\gamma}{2}|x|^2$$

where

$$(3.31) \quad \gamma := \sup_{x \in \mathcal{X}, y \in \mathcal{Y}} \left\| \nabla_x^2 c(x, y) \right\|_{op},$$

and $\|\cdot\|_{op}$ denotes the operator norm. When \mathcal{X} and \mathcal{Y} are two compact subsets of \mathbb{R}^d , the shifted cost is well defined and concave in the variable x .

In particular, this definition holds for the p -cost when $p \geq 2$, with $\gamma = (p-1)(R_{\mathcal{X}} + R_{\mathcal{Y}})^{p-2}$. Note also that this is actually the same exact strategy than going from the quadratic optimal transport problem to the maximum correlation problem where we go from a convex cost to a linear (and thus concave) one. In the current case, using the modified convexity (3.15) and not usual strong convexity is crucial, because the cost is no longer linear.

Proof of Theorem 3.6. Letting \mathcal{X}, c, ρ as in the theorem and setting \tilde{c} as in Definition 3.9, Remark 2.7 shows that proving Theorem 3.6 for \tilde{c} is equivalent to proving it with c . We can therefore suppose c concave. Then for \mathcal{Y} finite Step 1 applies thanks to Proposition 3.8. Step 2 as proved in Proposition 3.4 still holds. For Step 3 we argue letting first $\epsilon \downarrow 0$ by Remark 2.4 and then use Lemma 2.5 to extend from the semi-discrete to the general case. Note finally that the only requirement on C is $C(p, \rho, \mathcal{Y}) \geq 2M_{\phi_0, \phi_1}^{\tilde{c}}$, for all ϕ_0, ϕ_1 Kantorovitch potentials and where $M^{\tilde{c}}$ means that it is computed with the \tilde{c} cost. It is then easy to obtain a bound similar to (2.19) and thus completing the thesis. \square

Remark 3.10. The entire argument straightforwardly extends to cost functions c that are Lipschitz continuous and such that, for some $\gamma \in \mathbb{R}$, the modified cost (3.30) is concave with respect to the variable x , for every $y \in \mathcal{Y}$. This larger family includes for example the “boundary cost” from [FG10],

$$(3.32) \quad b_{\Omega}^2(x, y) := \min(|x - y|^2, d(x, \Omega^c)^2 + d(y, \Omega^c)^2),$$

where $\Omega \subseteq \mathbb{R}^d$ and $d(\cdot, \Omega^c) = \inf_{z \in \Omega^c} |\cdot - z|$. Quantitative stability with respect to such cost in the form of Corollary 3.7 plays a key role in [GHT24], where asymptotic limits for the quadratic transportation between random measures is investigated.

3.3. The case $1 < p < 2$. We finally move to the most challenging case of exponents $p \in (1, 2)$ for which in particular, the smoothness or even the concavity condition of the above remark fail. We prove the following inequality.

Theorem 3.11. *Let $p \in (1, 2)$, set $q = p/(p-1)$, let ρ be a log-concave probability measure with bounded (convex) support $\mathcal{X} \subseteq \mathbb{R}^d$ and let $\mathcal{Y} \subseteq \mathbb{R}^d$ be compact. Given $\mu_0, \mu_1 \in \mathcal{P}(\mathcal{Y})$, let $(\phi_{\mu_0}, \psi_{\mu_0}), (\phi_{\mu_1}, \psi_{\mu_1}) \in \mathcal{C}(\mathcal{X}) \times \mathcal{C}(\mathcal{Y})$ be pairs of Kantorovich potentials respectively for $\mathcal{W}_p(\rho, \mu_0)$ and $\mathcal{W}_p(\rho, \mu_1)$. Then, there exists $C = C(p, \rho, \mathcal{Y}) < \infty$ such that*

$$(3.33) \quad \text{Var}_{\rho}(\phi_1 - \phi_0) \leq C \langle \mu_1 - \mu_0 | \psi_1 - \psi_0 \rangle^{\frac{2}{q}}.$$

We also obtain the following corollary.

Corollary 3.12. *Let $p \in (1, 2)$, set $q = p/(p-1)$, let ρ be a log-concave probability measure with bounded (convex) support $\mathcal{X} \subseteq \mathbb{R}^d$ and let $\mathcal{Y} \subseteq \mathbb{R}^d$ be compact. Given $\mu \in \mathcal{P}(\mathcal{Y})$, let $(\phi_{\mu}, \psi_{\mu}) \in \mathcal{C}(\mathcal{X}) \times \mathcal{C}(\mathcal{Y})$ be Kantorovich potentials for $\mathcal{W}_p(\rho, \mu)$, let $\psi \in \mathcal{C}(\mathcal{Y})$ and write $\phi := \psi^c$. Then, there exists $C = C(p, \rho, \mathcal{Y}) < \infty$ such that it holds*

$$(3.34) \quad \text{Var}_{\rho}(\phi_{\mu} - \phi) \leq C [\mathcal{W}_p(\rho, \mu)^p - (\langle \mu | \psi \rangle + \langle \rho | \phi \rangle)]^{\frac{2}{q}}.$$

As already noticed, we cannot restrict ourselves to the previous \mathcal{C}^2 case as the cost does not have bounded derivatives on the diagonal. We will derive a weaker version of Step 1, before performing Step 2 and Step 3, where the scaling argument will be more important to control the additional term.

Proposition 3.13. *Let ρ be log-concave with (convex) bounded support $\mathcal{X}, \mathcal{Y} \subseteq \mathbb{R}^d$ be finite, $\varepsilon > 0$ and $\beta > 0$. For every $\psi_0, \psi_1 \in \mathcal{C}(\mathcal{Y})$ and $t \in [0, 1]$, it holds*

$$(3.35) \quad \begin{aligned} \log \mathcal{Z}_{\beta}((1-t)\psi_0 + t\psi_1) &\geq (1-t) \log \mathcal{Z}_{\beta}(\psi_0) + t \log \mathcal{Z}_{\beta}(\psi_1) \\ &\quad - \beta t(1-t) \gamma \mathcal{W}_p(\rho_{\beta, \psi_0}^{\varepsilon}, \rho_{\beta, \psi_1}^{\varepsilon})^p. \end{aligned}$$

where $\gamma = \gamma(p) < \infty$ and for $k \in \{0, 1\}$, $\rho_{\beta, \psi_k}^{\varepsilon}$ is defined as in (2.30) with ψ_k instead of ψ .

Proof of Proposition 3.13. Recall that we identify $\mathcal{C}(\mathcal{Y})$ with \mathbb{R}^n and again for simplicity we prove the result only for $\beta = 1$.

We first note that, for every $x_0, x_1 \in \mathbb{R}^d$, with the notation $x_t = (1-t)x_0 + tx_1$, there exists a constant $\gamma = \gamma(p)$ such that the following inequality holds:

$$(3.36) \quad |x_t|^p \geq (1-t)|x_0|^p + t|x_1|^p - \gamma t(1-t)|x_0 - x_1|^p \quad \text{for every } t \in [0, 1].$$

Therefore, letting $\psi_0, \psi_1 \in \mathbb{R}^n$, it holds

$$(3.37) \quad \begin{aligned} -(\psi_t)^{c, \varepsilon}(x_t) &= \varepsilon \log \left[\sum_{i=1}^n \exp \left(\frac{(\psi_t)_i - |x_t - y_i|^p}{\varepsilon} \right) \sigma_i \right] \\ &\leq \varepsilon \log \left[\sum_{i=1}^n \exp \left(\frac{(1-t)(\psi_0)_i + t(\psi_1)_i - (1-t)|x_0 - y_i|^p - t|x_1 - y_i|^p}{\varepsilon} \right) \right. \\ &\quad \left. \cdot \exp \left(\frac{t(1-t)|x_0 - x_1|^p}{\varepsilon} \right) \sigma_i \right] \end{aligned}$$

and we thus get, applying Holder’s inequality,

$$(3.38) \quad (\psi_t)^{c, \varepsilon}(x_t) \geq (1-t)(\psi_0)^{c, \varepsilon}(x_0) + t(\psi_1)^{c, \varepsilon}(x_1) - t(1-t)\gamma|x_0 - x_1|^p.$$

Therefore, taking the exponential and setting $h_t(x) = \exp((\psi_t)^{c;\varepsilon}(x))$ we have

$$(3.39) \quad h_t(x_t) \geq h_0(x_0)^{1-t} h_1(x_1)^t \exp(-t(1-t)\gamma|x_0 - x_1|^p).$$

We next obtain from this bound an integral inequality, in a similar fashion than the Prékopa-Leindler inequality. Up to renormalization, we can always assume $\|h_0\|_{L^1(\rho)} = \|h_1\|_{L^1(\rho)} = 1$, so that h_0 and h_1 are probability densities with respect to ρ .

By [Lemma 2.1](#) applied to the p -cost, $\rho_0 := h_0\rho$, $\rho_1 := h_1\rho$, using also the log-concavity of ρ and writing ρ_t for the displacement interpolation between $h_0\rho$ and $h_1\rho$, we get, for $h_0\rho$ -a.e. $x \in \mathcal{X}$, the following inequality :

$$(3.40) \quad \begin{aligned} \rho_t(T_t(x)) &\leq (h_0(x)\rho(x))^{1-t} (h_1(T(x))\rho(T(x)))^t \\ &\leq h_0(x)^{1-t} (h_1(T(x)))^t \rho(x)^{1-t} \rho(T(x))^t \\ &\leq h_0(x)^{1-t} (h_1(T(x)))^t \rho(T_t(x)) \\ &\leq h_t(T_t(x))\rho(T_t(x)) \exp(t(1-t)\gamma|x - T(x)|^p). \end{aligned}$$

so that

$$(3.41) \quad \rho_t(x) \exp(-t(1-t)\gamma|(T_t)^{-1}(x) - T((T_t)^{-1}(x))|^p) \leq h_t(x)\rho(x)$$

holds ρ_t a.e., where the inverse of T_t also makes sense ρ_t a.e.. Integrating over \mathcal{X} and using that $\rho_t = (T_t)_\#(h_0\rho)$,

$$(3.42) \quad \begin{aligned} \|h_t\|_1 &\geq \int_{\mathcal{X}} \exp(-\gamma t(1-t)|((T_t)^{-1}(x) - T((T_t)^{-1}(x))|^p) \rho_t(x) dx \\ &\geq \int_{\mathcal{X}} \exp(-\gamma t(1-t)|x - T(x)|^p) h_0(x) d\rho(x). \end{aligned}$$

Applying Jensen's inequality with the exponential function we get, introducing back the norms of h_0 and h_1 ,

$$(3.43) \quad \|h\|_{L^1(\rho)} \geq \|h_0\|_{L^1(\rho)}^{1-t} \|h_1\|_{L^1(\rho)}^t \exp\left(-\gamma t(1-t) \int_{\mathcal{X}} |x - T(x)|^p h_0(x) d\rho(x)\right).$$

Finally, taking the log yields the thesis:

$$\begin{aligned} \log \mathcal{Z}((1-t)\psi^0 + t\psi^1) &\geq (1-t) \log \mathcal{Z}(\psi^0) + t \log \mathcal{Z}(\psi^1) \\ &\quad - t(1-t)\gamma \mathcal{W}_p(\rho_{\psi^0}^\varepsilon, \rho_{\psi^1}^\varepsilon)^p. \end{aligned} \quad \square$$

Using the established bound for $\log \mathcal{Z}_\beta$, we prove the main result for this section.

Proof of [Theorem 3.11](#). From [Proposition 3.13](#), writing $I_\beta := \log \mathcal{Z}_\beta$ for brevity, one gets for all $t \in (0, 1)$,

$$(3.44) \quad \begin{aligned} \beta\gamma \mathcal{W}_p(\rho_{\beta, \psi_0}^\varepsilon, \rho_{\beta, \psi_1}^\varepsilon)^p &\geq \frac{(1-t)I_\beta(\psi^0) + tI_\beta(\psi^1) - I_\beta((1-t)\psi^0 + t\psi^1)}{t(1-t)} \\ &\geq - \int_0^1 \langle \nabla I_\beta(\psi_{st}) | \psi_1 - \psi_0 \rangle ds + \int_0^1 \langle \nabla I_\beta(\psi_{1-(1-t)s}) | \psi_1 - \psi_0 \rangle ds. \end{aligned}$$

Letting successively $t \rightarrow 0$ and $t \rightarrow 1$ we find

$$(3.45) \quad \beta\gamma \mathcal{W}_p(\rho_{\beta, \psi_0}^\varepsilon, \rho_{\beta, \psi_1}^\varepsilon)^p \geq -\langle \nabla I_\beta(\psi_0) | \psi_1 - \psi_0 \rangle + \int_0^1 \langle \nabla I_\beta(\psi_{1-s}) | \psi_1 - \psi_0 \rangle ds$$

and

$$(3.46) \quad \beta\gamma \mathcal{W}_p(\rho_{\beta, \psi_0}^\varepsilon, \rho_{\beta, \psi_1}^\varepsilon)^p \geq - \int_0^1 \langle \nabla I_\beta(\psi_s) | \psi_1 - \psi_0 \rangle ds + \langle \nabla I_\beta(\psi_1) | \psi_1 - \psi_0 \rangle$$

so that, by summing the two inequalities,

$$(3.47) \quad 2\beta\gamma \mathcal{W}_p(\rho_{\beta, \psi_0}^\varepsilon, \rho_{\beta, \psi_1}^\varepsilon)^p \geq \langle \nabla I_\beta(\psi_1) - \nabla I_\beta(\psi_0) | \psi_1 - \psi_0 \rangle.$$

Following the proof of *Step 2* as proposed in [Proposition 3.4](#) and letting for $\phi_t^\varepsilon = \psi_t^{c,\varepsilon}$, for $t \in [0, 1]$ where $\psi_t = (1-t)\psi_0 + t\psi_1$, $M^\varepsilon := \sup_{t \in [0,1]} \{\text{osc}(\phi_t^\varepsilon)\}$, and for all $\phi \in \mathcal{C}(\mathcal{X})$,

$$(3.48) \quad \rho_\phi := \frac{e^\phi}{\int_{\mathcal{X}} e^\phi d\rho} \rho,$$

we find

$$(3.49) \quad \beta e^{-\beta M^\varepsilon} \text{Var}_\rho(\phi_1^\varepsilon - \phi_0^\varepsilon) \leq e^{\beta M^\varepsilon} \langle \nabla \mathcal{K}^{c,\varepsilon}(\psi_1) - \nabla \mathcal{K}^{c,\varepsilon}(\psi_0) | \psi_1 - \psi_0 \rangle + 2\gamma \mathcal{W}_p(\rho_{\beta\phi_0^\varepsilon}, \rho_{\beta\phi_1^\varepsilon})^p.$$

If we then let $\varepsilon \downarrow 0$ minding [Remark 2.4](#), by choosing for simplicity the potentials so that ϕ_k have zero ρ -mean, we get that for all $\beta > 0$ the following bound holds:

$$(3.50) \quad \beta e^{-\beta M_{\phi_0, \phi_1}} \|\phi_1 - \phi_0\|_{L^2(\rho)}^2 \leq e^{\beta M_{\phi_0, \phi_1}} \langle \mu_0 - \mu_1 | \psi_1 - \psi_0 \rangle + 2\gamma \mathcal{W}_p(\rho_{\beta\phi_0}, \rho_{\beta\phi_1})^p,$$

The next step is to derive an estimate of $\mathcal{W}_p(\rho_{\beta\phi_0}, \rho_{\beta\phi_1})$ in terms of $\|\phi_1 - \phi_0\|_{L^2(\rho)}$. Again, for simplicity we argue in the case $\beta = 1$. First, we notice that by the monotonicity of Wasserstein distances, we have $\mathcal{W}_p(\rho_{\phi_0}, \rho_{\phi_1}) \leq \mathcal{W}_2(\rho_{\phi_0}, \rho_{\phi_1})$. Then, we apply [Lemma 2.2](#), with $\mu_k = \rho_{\phi_k}$, $k \in \{0, 1\}$, so that $f_k \geq e^{-M_{\phi_0, \phi_1}}$ and we get

$$(3.51) \quad \mathcal{W}_2^2(\rho_{\phi_1}, \rho_{\phi_0}) \leq C_\rho e^{M_{\phi_0, \phi_1}} \left\| \frac{e^{\phi_1}}{\int_{\mathcal{X}} e^{\phi_1} d\rho} - \frac{e^{\phi_0}}{\int_{\mathcal{X}} e^{\phi_0} d\rho} \right\|_{L^2(\rho)}^2.$$

For any $x \in \mathcal{X}$, we have

$$(3.52) \quad \begin{aligned} \left| \frac{e^{\phi_1(x)}}{\int_{\mathcal{X}} e^{\phi_1} d\rho} - \frac{e^{\phi_0(x)}}{\int_{\mathcal{X}} e^{\phi_0} d\rho} \right| &\leq \left| \frac{e^{\phi_1(x)} - e^{\phi_0(x)}}{\int_{\mathcal{X}} e^{\phi_1} d\rho} \right| + e^{\phi_0(x)} \left| \frac{1}{\int_{\mathcal{X}} e^{\phi_1} d\rho} - \frac{1}{\int_{\mathcal{X}} e^{\phi_0} d\rho} \right| \\ &\leq e^{M_{\phi_0, \phi_1}} |e^{\phi_1(x)} - e^{\phi_0(x)}| + e^{2M_{\phi_0, \phi_1}} \left| \int_{\mathcal{X}} (e^{\phi_1} - e^{\phi_0}) d\rho \right| \\ &\leq e^{2M_{\phi_0, \phi_1}} (|\phi_1(x) - \phi_0(x)| + \|\phi_1 - \phi_0\|_{L^1(\rho)}) \end{aligned}$$

where the zero ρ -mean property of ϕ_k assures that

$$(3.53) \quad -M_{\phi_0, \phi_1} \leq \min \phi_k \leq \max \phi_k \leq M_{\phi_0, \phi_1}.$$

Integrating with respect to ρ yields

$$(3.54) \quad \left\| \frac{e^{\phi_1}}{\int_{\mathcal{X}} e^{\phi_1} d\rho} - \frac{e^{\phi_0}}{\int_{\mathcal{X}} e^{\phi_0} d\rho} \right\|_{L^2(\rho)}^2 \leq C_\rho e^{4M_{\phi_0, \phi_1}} \|\phi_1 - \phi_0\|_{L^2(\rho)}^2.$$

From this estimation, [\(3.50\)](#) and [\(3.51\)](#) we get

$$(3.55) \quad \beta e^{-\beta M_{\phi_0, \phi_1}} \|\phi_1 - \phi_0\|_{L^2(\rho)}^2 \leq e^{\beta M_{\phi_0, \phi_1}} \langle \mu_0 - \mu_1 | \psi_1 - \psi_0 \rangle + \gamma C_\rho e^{\frac{5p\beta}{2} M_{\phi_0, \phi_1}} \beta^p \|\phi_1 - \phi_0\|_{L^2(\rho)}^p$$

so,

$$(3.56) \quad \beta e^{-2\beta M_{\phi_0, \phi_1}} \|\phi_1 - \phi_0\|_{L^2(\rho)}^2 - \gamma C_\rho e^{\frac{(5p-2)\beta}{2} M_{\phi_0, \phi_1}} \beta^p \|\phi_1 - \phi_0\|_{L^2(\rho)}^p \leq \langle \mu_0 - \mu_1 | \psi_1 - \psi_0 \rangle.$$

Setting $C = \gamma C_\rho$, $\alpha = 2M_{\phi_0, \phi_1}$ and $\alpha' = \frac{(5p-2)}{2} M_{\phi_0, \phi_1}$ so that $C, \alpha, \alpha' > 0$ and $h(\beta) := \beta e^{-\alpha\beta} - C e^{\alpha'\beta} \beta^p$, finally optimizing in β we get

$$(3.57) \quad \min(\|\phi_1 - \phi_0\|_{L^2(\rho)}^2, \|\phi_1 - \phi_0\|_{L^2(\rho)}^{\frac{p}{p-1}}) \leq \frac{1}{\sup_{\beta \in \mathbb{R}_+} h(\beta)} \langle \mu_0 - \mu_1 | \psi_1 - \psi_0 \rangle.$$

Now, deriving the absolute bound

$$(3.58) \quad \begin{aligned} \|\phi_1 - \phi_0\|_{L^2(\rho)} &\leq |\mathcal{X}| (\|\phi_0\|_{L^\infty} + \|\phi_1\|_{L^\infty}) \\ &\leq 2|\mathcal{X}| M_{\phi_0, \phi_1} \leq C = C(\rho, p, \mathcal{Y}) \end{aligned}$$

we get $\|\phi_1 - \phi_0\|_{L^2(\rho)}^{\frac{p}{p-1}} \leq C\|\phi_1 - \phi_0\|_{L^2(\rho)}^2$. From this last inequality and (3.57) we can conclude

$$(3.59) \quad \|\phi_1 - \phi_0\|_{L^2(\rho)}^{\frac{p}{p-1}} \leq C\langle \mu_0 - \mu_1 | \psi_1 - \psi_0 \rangle$$

for some $C = C(p, \rho, \mathcal{X}, \mathcal{Y}) > 0$. One final application of Lemma 2.5 extends the bounds from the semi-discrete case to the general case of compact \mathcal{Y} . \square

Remark 3.14. One can easily check that the entire proof above relies only on (Lipschitz continuity and) the curvature condition (3.36) satisfied by the p -cost. Thus, if the cost c is such that, for some $p > 1$ and $\gamma < \infty$, it holds

$$(3.60) \quad c(x_t, y) \geq (1-t)c(x_0, y) + tc(x_1, y) - \gamma t(1-t)|x_0 - x_1|^p \quad \text{for all } x_0, x_1 \in \mathcal{X} \text{ and } y \in \mathcal{Y}$$

then Proposition 3.13 still holds. We notice that such property, which we may refer to as a (p, γ) -curvature condition for the cost function c , is stable by taking sums and infima. As an example, it holds for the boundary cost analogue to (3.32) with general exponent p .

Remark 3.15. Again, we notice that the only point where log-concavity of ρ is used is in the proof of Proposition 3.13. Therefore, in order to cover the case of ρ that admits a density uniformly bounded from above and below by strictly positive constants with respect to a log-concave measure λ , we need to replace ρ with λ in the definition of \mathcal{Z} . This leads to a version of (3.35) with $\lambda_{\beta, \psi_k}^\varepsilon$ instead of $\rho_{\beta, \psi_k}^\varepsilon$. However, this changes very little in the proof of Theorem 3.11: we obtain (3.49) arguing as in Proposition 3.4 and taking into account Remark 3.5; subsequently, we apply Lemma 2.2 with respect to the measure λ ; and finally we replace $L^2(\lambda)$ with $L^2(\rho)$, up to a multiplicative constant.

4. QUANTITATIVE STABILITY OF MAPS

In this section, we focus on the stability of optimal transport maps by establishing Theorem 1.2. We make the following standing assumption: ρ is a probability measure with bounded and convex support $\mathcal{X} \subseteq \mathbb{R}^d$, with density bounded below and above by two constants:

$$(4.1) \quad 0 < m_\rho \leq \rho(x) \leq M_\rho < +\infty \quad \text{for every } x \in \mathcal{X}.$$

Our goal is now to use the results from the previous section to yield a bound on the L^2 distance between two optimal maps T_0 and T_1 from the source ρ to two target distributions μ_0, μ_1 , supported on a bounded set \mathcal{Y} . By the extension of Brenier theorem to general p -power costs [AGS08, Theorem 6.2.4] ($p > 1$), the optimal transport map T for $\mathcal{W}_p(\rho, \mu)$ is given by

$$(4.2) \quad T(x) = x - (\nabla\phi(x))^{(q/p)} \quad \text{for a.e. } x \in \mathcal{X},$$

where ϕ is the Kantorovitch potential, $q = p/(p-1)$ and we recall the notation $v^{(\alpha)} = |v|^{\alpha-1}v$. Thus, our strategy is to establish first upper bounds on $\|\nabla\phi_1 - \nabla\phi_0\|_{L^2}$ in terms of $\|\phi_1 - \phi_0\|_{L^2}$, whenever $\phi_k, k \in \{0, 1\}$ are c -concave functions. In the following section, we derive such bounds, first for the quadratic case – recalling the argument from [Del22] – and then more generally for c -concave functions.

4.1. The quadratic case. We report in this section the argument from [Del22]. Let us recall that the Brenier map is the optimal transport map for the p -cost when $p = 2$.

Theorem 4.1. *Let \mathcal{X}, \mathcal{Y} be compact subsets of \mathbb{R}^d with \mathcal{X} convex, let ρ be a probability measure with density w.r.t. the Lebesgue measure bounded below and above by strictly positive constants as in (4.1) and let $\mu_0, \mu_1 \in \mathcal{P}(\mathcal{Y})$. Then, there exists a constant $C = C(d, \mathcal{X}, \mathcal{Y}, m_\rho, M_\rho) < \infty$ such that*

$$(4.3) \quad \|T_0 - T_1\|_{L^2(\rho)} \leq C\mathcal{W}_1(\mu_0, \mu_1)^{\frac{1}{6}}.$$

where T_k denotes the Brenier map from ρ to μ_k , for $k \in \{0, 1\}$.

This theorem is an immediate consequence of [Theorem 3.1](#), [\(4.2\)](#) which reads

$$(4.4) \quad T(x) = x - \nabla\phi(x) = \nabla \left(\frac{|x|^2}{2} - \phi(x) \right), \quad \text{for a.e. } x \in \mathcal{X},$$

so that $T(x)$ is the gradient of the function $x \mapsto |x|^2/2 - \phi(x)$ which is convex and Lipschitz, and the following “reverse” Poincaré inequality, [[Del22](#), proposition 5.11] – but see also [[SS05](#); [HPS08](#); [PSSURNH15](#)] for similar results – whose statement we report here for the sake of completeness and for comparison with our extension on fractional spaces, [Proposition 4.2](#) below.

Proposition 4.2. *Let $\mathcal{X} \subseteq \mathbb{R}^d$ be a bounded domain with rectifiable boundary and let $u, v : \mathcal{X} \rightarrow \mathbb{R}$ be convex Lipschitz functions. Then, there exists a constant $C = C(d) < \infty$ depending on d only such that*

$$(4.5) \quad \|\nabla u - \nabla v\|_{L^2(\mathcal{X})}^2 \leq C \mathcal{H}^{d-1}(\partial\mathcal{X})^{\frac{2}{3}} (\|\nabla u\|_{L^\infty(\mathcal{X})} + \|\nabla v\|_{L^\infty(\mathcal{X})})^{\frac{4}{3}} \|u - v\|_{L^2(\mathcal{X})}^{\frac{2}{3}},$$

where \mathcal{H}^{d-1} denotes the $(d-1)$ -dimensional Hausdorff measure.

4.2. The case $p > 2$. This is a minor modification of the quadratic case, minding that the optimal transport map depends on the gradient of the Kantorovich potential in a Hölder continuous way. We prove the following result.

Theorem 4.3. *Let \mathcal{X}, \mathcal{Y} be compact subsets of \mathbb{R}^d with \mathcal{X} convex, let ρ be a probability measure, with density w.r.t. the Lebesgue measure bounded below and above by strictly positive constants as in [\(4.1\)](#). For $k \in \{0, 1\}$, let $\mu_k, \mu_1 \in \mathcal{P}(\mathcal{Y})$ and denote with T_k the optimal transport map for $\mathcal{W}_p(\rho, \mu_k)$. Then, there exists a constant $C = C(p, d, \mathcal{X}, \mathcal{Y}, m_\rho, M_\rho) < \infty$ such that*

$$(4.6) \quad \|T_0 - T_1\|_{L^2(\rho)} \leq C \mathcal{W}_1(\mu_0, \mu_1)^{\frac{1}{6(p-1)}}.$$

Proof. We start from with the inequality $|v^{(\alpha)} - w^{(\alpha)}| \leq C(\alpha)|v - w|^\alpha$, valid for all $0 < \alpha < 1$ and $v, w \in \mathbb{R}^d$ with $C = C(\alpha, d) < \infty$. We find

$$(4.7) \quad \begin{aligned} \|T_0 - T_1\|_{L^2(\rho)}^2 &= \int_{\mathcal{X}} |\nabla\phi_0^{(q/p)} - \nabla\phi_1^{(q/p)}|^2 d\rho \\ &\leq C \int_{\mathcal{X}} |\nabla\phi_0 - \nabla\phi_1|^{2q/p} d\rho \\ &\leq C \|\nabla\phi_0 - \nabla\phi_1\|_{L^2(\rho)}^{2q/p}, \end{aligned}$$

Having used Hölder inequality and the fact that $q/p = 1/(p-1) < 1$ for $p > 2$. Next, we notice that there exists $\gamma = \gamma(p, \mathcal{X}, \mathcal{Y}) < \infty$ such that, for any function ϕ that is c -concave with respect to the p -cost, the function $x \mapsto \phi(x) - \gamma|x|^2$ is concave and Lipschitz (with a bound on the Lipschitz constant uniform on ϕ). We then proceed with the upper bounds (where the constant C may change from line to line)

$$(4.8) \quad \begin{aligned} \|\nabla\phi_0 - \nabla\phi_1\|_{L^2(\rho)}^2 &\leq C \|\nabla\phi_0 - \nabla\phi_1\|_{L^2(\mathcal{X})}^2 = C \left\| \nabla(\phi_0 - \gamma|\cdot|^2) - \nabla(\phi_1 - \gamma|\cdot|^2) \right\|_{L^2(\mathcal{X})}^2 \\ &\leq C \|\phi_0 - \phi_1\|_{L^2(\mathcal{X})}^{\frac{2}{3}} \text{ by } \text{Proposition 4.2} \\ &\leq C \|\phi_0 - \phi_1\|_{L^2(\rho)}^{\frac{2}{3}} \leq C \mathcal{W}_1(\mu_0, \mu_1)^{\frac{1}{3}}. \end{aligned}$$

where the last inequality follows by [Theorem 1.1](#) assuming without loss of generality that ϕ_0, ϕ_1 have zero ρ -mean. \square

4.3. The case $1 < p < 2$. This case is instead less obvious, again because there is no shift reducing a c -concave function to a concave function when c is the p -cost for $1 < p < 2$. We prove the following result.

Theorem 4.4. *Let $p \in (1, 2)$, set $q = p/(p - 1)$, let $\mathcal{X}, \mathcal{Y} \subseteq \mathbb{R}^d$ be compact subsets with \mathcal{X} convex, let ρ be a log-concave probability measure on \mathcal{X} such that (4.1) holds, let $\mu_0, \mu_1 \in \mathcal{P}(\mathcal{Y})$ and denote with T_k the optimal transport map for $\mathcal{W}_p(\rho, \mu_k)$. Then, for every $\theta \in (0, (p - 1)/(p + 1))$ there exists a constant $C = C(\theta, p, d, \mathcal{X}, \mathcal{Y}, m_\rho, M_\rho) < \infty$ such that*

$$(4.9) \quad \|T_0 - T_1\|_{L^2(\rho)} \leq C \mathcal{W}_1(\mu_0, \mu_1)^{\frac{\theta}{q}}.$$

To establish this theorem we need an equivalent of Proposition 4.2 for c -concave functions with respect to the p -cost.

Proposition 4.5. *Let $p \in (1, 2)$, $\mathcal{X} \subseteq \mathbb{R}^d$ be a bounded domain with rectifiable boundary and let $u, v : \mathcal{X} \rightarrow \mathbb{R}$ be c -concave with respect to the p -cost. Then, for all $\theta \in (0, (p - 1)/(p + 1))$ there exists a constant $C = C(\theta, \mathcal{X}, p) < \infty$ such that*

$$(4.10) \quad \|\nabla u - \nabla v\|_{L^2(\mathcal{X})} \leq C \|u - v\|_{L^2(\mathcal{X})}^\theta.$$

For comparison with our argument, let us recall that the proof of Proposition 4.2 relies on two ingredients. First, one establishes the one-dimensional inequality [Del22, Proposition 5.24]

$$(4.11) \quad \int_0^1 |u' - v'|^2 \leq 8 (\text{Lip}(u) + \text{Lip}(v))^{\frac{4}{3}} \left(\int_0^1 |u - v|^2 \right)^{\frac{1}{3}}$$

that settles the case $\mathcal{X} = [0, 1] \subseteq \mathbb{R}$ (and by scaling any other interval). Then, for the general case, one applies an integral-geometric argument [Del22, Lemma 5.25] decomposing the d -dimensional integrals into integral over segments. In our context, the integral-geometric argument formula will be identical, up to harmless variations in the exponents. Our aim is therefore to establish a suitable analogue of (4.11) for functions that are c -concave with respect to the p -cost. Before we do this, it is convenient to notice the following consequence of c -concavity with respect to the p -cost.

Lemma 4.6. *Let $p \in (1, 2)$, $\mathcal{X} \subseteq \mathbb{R}^d$ be open and bounded and $\phi : \mathcal{X} \rightarrow \mathbb{R}$ be c -concave with respect to the p -cost. Then, at every $x, y \in \mathcal{X}$ that are differentiability points for ϕ , it holds*

$$(4.12) \quad \langle \nabla \phi(x) - \nabla \phi(y), x - y \rangle \leq \frac{2\gamma}{p^2} |x - y|^p,$$

where $\gamma = \gamma(p)$ is the same constant as in (3.36).

Proof. By c -concavity, the graph of the associated transport map

$$(4.13) \quad x \rightarrow T(x) := x - (\nabla \phi(x))^{(q/p)}$$

where $q = p/(p - 1)$, that is defined at every differentiability point of x , is c -cyclically monotone, i.e.,

$$(4.14) \quad |T(x) - x|^p - |T(x) - y|^p \leq |T(y) - x|^p - |T(y) - y|^p.$$

Next, we recall the inequality, valid for any $v, h, z \in \mathbb{R}^d$,

$$(4.15) \quad \begin{aligned} |v + h - z|^p - |v - z|^p - p \langle (v - z)^{(p-1)}, h \rangle \\ = p \int_0^1 \langle (v + th - z)^{(p-1)} - (v - z)^{(p-1)}, h \rangle dt \\ \leq \gamma \int_0^1 t^{p-1} |h|^p dt = \frac{\gamma}{p} |h|^p, \end{aligned}$$

where we used the inequality [Lin19, Chapter 12, (III)]

$$(4.16) \quad p \langle (a - z)^{(p-1)} - (b - z)^{(p-1)}, a - b \rangle \leq \gamma |a - b|^p,$$

valid for all $a, b, z \in \mathbb{R}^d$, since $1 < p < 2$. Applying (4.15) in both sides of (4.14), i.e., first with $z = T(x)$, $h = y - x$ and then for $z = T(y)$, $h = x - y$, we get

$$(4.17) \quad -p \langle (x - T(x))^{(p-1)}, y - x \rangle - \frac{\gamma}{p} |y - x|^p \leq p \langle (y - T(y))^{(p-1)}, x - y \rangle + \frac{\gamma}{p} |y - x|^p$$

so that, noticing that $(x - T(x))^{(p-1)} = \nabla\phi(x)$ by the very definition of T , we obtain the thesis. \square

In the proof below we only use that c -concave functions for the p -cost are Lipschitz (with uniform bounds on their Lipschitz constant) and satisfy inequality (4.12). Thus, we introduce the following notation: for $p \in (1, 2)$, $\lambda \in \mathbb{R}$, we say that a Lipschitz function $u : \mathcal{X} \rightarrow \mathbb{R}$ is (p, λ) -concave if, for every x, y that are differentiability point of u it holds

$$(4.18) \quad \langle \nabla u(x) - \nabla u(y), x - y \rangle \leq \lambda |x - y|^p.$$

We are now in a position to state and prove the following analogue of (4.11).

Lemma 4.7. *Let $p \in (1, 2)$ and $\lambda, L \geq 0$. For every $\theta \in (0, (p-1)/(p+1))$, there exists $C = C(p, \lambda, L, \theta) < \infty$ such that for $u, v : [0, 1] \rightarrow \mathbb{R}$ both (p, λ) -concave and L -Lipschitz, it holds*

$$(4.19) \quad \int_0^1 |u' - v'|^2 dx \leq C \left(\int_0^1 |u - v|^2 dx \right)^\theta.$$

Proof of Lemma 4.7. Without loss of generality, we may assume throughout the proof that $\int_0^1 u dx = \int_0^1 v dx = 0$.

For comparison with the proof in our case, we recall that a key point in the proof of (4.11) in [Del22] is the following integration by parts (assuming by approximation that the functions are C^2 smooth):

$$(4.20) \quad \begin{aligned} \int_0^1 |u' - v'|^2 &= (u - v)(u' - v') \Big|_0^1 - \int_0^1 (u - v)(u'' - v'') \\ &\leq 2(\text{Lip}(u) + \text{Lip}(v)) \|u - v\|_\infty + \|u - v\|_\infty \int_0^1 (|u''| + |v''|) \\ &= 2(\text{Lip}(u) + \text{Lip}(v)) \|u - v\|_\infty + \|u - v\|_\infty \int_0^1 (u'' + v'') \\ &\leq \left[4(\|u'\|_{L^\infty} + \|v'\|_{L^\infty}) + (u' + v') \Big|_0^1 \right] \|u - v\|_\infty \\ &\leq 8(\|u'\|_{L^\infty} + \|v'\|_{L^\infty}) \|u - v\|_\infty, \end{aligned}$$

where the convexity assumption on u and v allowed us to remove the absolute values in the terms $|u''|$ and $|v''|$. The above argument can be interpreted as an interpolation bound on the $H^{1,2}$ norm in terms of the L^∞ norm of the function and the L^1 norm of its second derivative, that together with the positivity condition reduces to a L^∞ norm of the derivative. Finally, the standard Gagliardo-Nirenberg inequality allows to replace the L^∞ norm in the right hand side with the L^2 norm (which gives the exponents $4/3$ and $2/3$).

In our case of (p, λ) -concave functions, the positivity of second derivatives is replaced by the weaker condition (4.18), so the idea is to interpolate with fractional derivatives of order between 1 and 2. We start from the assumption, which in the one dimensional case reads

$$(4.21) \quad (u'(x) - u'(y))(x - y) - \lambda |x - y|^p \leq 0$$

for every $x, y \in (0, 1)$ where u is differentiable, which is a set of full Lebesgue measure, because u is Lipschitz continuous. Without loss of generality, we may also assume that $\lambda \geq 0$. Therefore, we have the inequality (assuming $x \neq y$):

$$(4.22) \quad \begin{aligned} \frac{|u'(x) - u'(y)|}{|x - y|} &= \frac{|(u'(x) - u'(y))(x - y)|}{|x - y|^2} \\ &\leq \frac{|(u'(x) - u'(y))(x - y) - \lambda |x - y|^p| + \lambda |x - y|^p}{|x - y|^2} \\ &= \frac{(u'(y) - u'(x))}{x - y} + \frac{2\lambda}{|x - y|^{2-p}}. \end{aligned}$$

Given $\alpha \in (0, 1)$, we bound from above

(4.23)

$$\begin{aligned} [u']_{\alpha,1} &= \int_0^1 \int_0^1 \frac{|u'(x) - u'(y)|}{|x - y|^{1+\alpha}} dy dx \\ &\leq \int_0^1 \int_0^1 \frac{u'(y) - u'(x)}{(x - y)|x - y|^\alpha} dy dx + \int_0^1 \int_0^1 \frac{2\lambda dx dy}{|x - y|^{2-p+\alpha}} \\ &= - \int_0^1 \int_{-1}^1 \chi_{\{0 \leq x+h \leq 1\}} \frac{u'(x+h) - u'(x)}{h|h|^\alpha} dh dx + 2\lambda \int_0^1 \int_{-1}^1 \chi_{\{0 \leq x+h \leq 1\}} \frac{dh dx}{|h|^{2-p+\alpha}}. \end{aligned}$$

where χ denotes the indicator function. We bound separately the two terms. For the second one, we simply use the fact that $\int_0^1 h^{-\beta} ds < \infty$ if $\beta < 1$, hence

$$(4.24) \quad 2\lambda \int_0^1 \int_{-1}^1 \chi_{\{0 \leq x+h \leq 1\}} \frac{dh dx}{|h|^{2-p+\alpha}} \leq C,$$

for some $C = C(p, \alpha, \lambda) < \infty$, provided that $2 - p + \alpha < 1$, i.e., $\alpha < p - 1$. For the first term, we integrate first with respect to $x \in [0, 1]$, so that for fixed $h \in [-1, 1]$, we find

$$(4.25) \quad \int_0^1 \chi_{\{0 \leq x+h \leq 1\}} (u'(x+h) - u'(x)) dx = u(\min\{1+h, 1\}) - u(\max\{h, 0\}) \\ - u(\min\{1-h, 1\}) + u(\max\{-h, 0\}).$$

When we integrate also with respect to $h \in [-1, 1]$ it is convenient to consider separately the two cases $h > 0$ and $h < 0$. In the first case, we find

$$(4.26) \quad \int_0^1 \int_0^1 \chi_{\{0 \leq x+h \leq 1\}} \frac{u'(x+h) - u'(x)}{|h|^\alpha h} dx dh = \int_0^1 \frac{u(1) - u(1-h) - u(h) + u(0)}{h^{1+\alpha}} dh \\ \leq 2 \text{Lip}(u) \int_0^1 \frac{1}{h^\alpha} dh \leq C,$$

where $C = C(\alpha, L) < \infty$, having used that $\alpha < 1$. In the case $h < 0$, we find similarly

$$(4.27) \quad \int_{-1}^0 \int_0^1 \chi_{\{0 \leq x+h \leq 1\}} \frac{u'(x+h) - u'(x)}{|h|^\alpha h} dx dh = \int_{-1}^0 \frac{u(1+h) - u(0) - u(1) + u(-h)}{|h|^\alpha h} dh \\ \leq 2 \text{Lip}(u) \int_0^1 \frac{1}{h^\alpha} dh = C.$$

Thus, we collect the bound

$$(4.28) \quad [u']_{\alpha,1} \leq C = C(p, \alpha, \lambda, L) < \infty,$$

provided that $0 < \alpha < p - 1$. Since u is Lipschitz (and has zero mean), it also holds

$$(4.29) \quad \|u\|_{W^{1,1}((0,1))} \leq \|u\|_{W^{1,\infty}((0,1))} \leq C.$$

In conclusion, the thesis follows as an application of (2.48) on $\Omega = (0, 1)$ and $r := 1 + \alpha \in (1, p)$ and letting $\theta := 1 - 2/(1 + r)$, which can be any real in the interval $(0, (p - 1)/(p + 1))$. \square

We now indicate how to complete the proof of [Proposition 4.5](#) following the steps of [\[Del22, Proposition 5.11\]](#), for which we refer for more details.

Proof of Proposition 4.5. The first step is to notice the good scaling properties of inequality (4.19). Through an affine change of variable, one easily proves that if $p \in (1, 2)$, $\lambda, L, l \geq 0$ and $\theta \in (0, (p - 1)/(p + 1))$, there exists $C = C(p, \lambda, L, l, \theta) < \infty$ such that for any interval $I \subset \mathbb{R}$ with $|I| \leq l$, $u, v : I \rightarrow \mathbb{R}$ both (p, λ) -concave and L -Lipschitz, it holds

$$(4.30) \quad \int_0^1 |u' - v'|^2 dx \leq C \left(\int_0^1 |u - v|^2 dx \right)^\theta.$$

To pursue the proof we need to introduce some new notation that we try to keep as light as possible, referring to [\[Del22\]](#) and the references therein for more details.

Let \mathcal{L}^d be the set of affine lines in \mathbb{R}^d identified with $\{\ell = (e, p) \in \mathbb{R}^d \times \mathbb{R}^d \mid e \in \mathbb{S}^{d-1}, p \in e^\perp\}$ equipped with the Riemannian metric. We write $d\mathcal{L}^d$ the volume measure, so that for any $\phi : \mathcal{L}^d \rightarrow \mathbb{R}$

$$(4.31) \quad \int_{\ell \in \mathcal{L}^d} \phi(\ell) d\mathcal{L}^d(\ell) = \int_{e \in \mathbb{S}^{d-1}} \int_{p \in e^\perp} \phi((e, p)) d\mathcal{H}^{d-1}(p) d\mathcal{H}^{d-1}(e)$$

where \mathcal{H}^{d-1} denotes the $(d-1)$ -dimensional Hausdorff measure.

Back to our setting, by [Del22, Lemma 5.25], one obtains

$$(4.32) \quad \|u\|_{L^2(\mathcal{X})}^2 = C \int_{\ell \in \mathcal{L}^d} \int_{y \in \ell \cap \mathcal{X}} u(y)^2 dy d\mathcal{L}^d(\ell)$$

and

$$(4.33) \quad \|\nabla u\|_{L^2(\mathcal{X})}^2 = C' \int_{\ell \in \mathcal{L}^d} \int_{y \in \ell \cap \mathcal{X}} \langle \nabla u(y), e(\ell) \rangle^2 dy d\mathcal{L}^d(\ell)$$

where $C = C(d), C' = C'(d) \in (0, \infty)$ are dimensional constants. Since \mathcal{X} is convex, we can write $\ell \cap \mathcal{X} = I_\ell$ for a suitable line segment (that we identify with a real interval). Then we use (4.30) with $t \in I_\ell \rightarrow u(p + te)$ and $l = \text{diam}(\mathcal{X})$ for a.e. $\ell \in \mathcal{L}^d$, (4.32), (4.33) and Holder's inequality obtaining the thesis,

$$(4.34) \quad \|\nabla u - \nabla v\|_{L^2(\mathcal{X})} \leq C \|u - v\|_{L^2(\mathcal{X})}^\theta. \quad \square$$

We end this section with the proof of [Theorem 4.4](#), which in turn concludes the proof of [Theorem 1.2](#).

Proof of [Theorem 4.4](#). Naming ϕ_0 and ϕ_1 the zero-mean (with respect to ρ) Kantorovich potentials associated to T_0 and T_1 , by (4.2) we have

$$(4.35) \quad \|T_0 - T_1\|_{L^2(\mathcal{X})} = \left\| (\nabla \phi_0)^{(q/p)} - (\nabla \phi_1)^{(q/p)} \right\|_{L^2(\mathcal{X})},$$

where $q = p/(p-1)$. Using the inequality

$$(4.36) \quad |v^{(\alpha)} - w^{(\alpha)}| \leq C |v - w| (|v|^{\alpha-1} + |w|^{\alpha-1}),$$

valid for $v, w \in \mathbb{R}^d, \alpha > 1$ with $C = C(\alpha, d) < \infty$, we find for $\alpha := q/p > 1$,

$$(4.37) \quad \|T_0 - T_1\|_{L^2(\mathcal{X})} \leq C \left(\text{Lip}(\phi_0)^{q/p-1} + \text{Lip}(\phi_1)^{q/p-1} \right) \|\nabla \phi_0 - \nabla \phi_1\|_{L^2(\mathcal{X})}.$$

Since the Lipschitz constant of ϕ_0 and ϕ_1 are uniformly bounded by c -concavity, it follows that

$$(4.38) \quad \begin{aligned} \|T_0 - T_1\|_{L^2(\mathcal{X})} &\leq C \|\nabla \phi_0 - \nabla \phi_1\|_{L^2(\mathcal{X})} \leq C \|\phi_0 - \phi_1\|_{L^2(\mathcal{X})}^\theta \\ &\leq C \|\phi_0 - \phi_1\|_{L^2(\rho)}^\theta \end{aligned}$$

where the constant C may change from line to line, and we applied [Proposition 4.5](#). To conclude, we apply [Theorem 3.11](#) and obtain

$$(4.39) \quad \|T_0 - T_1\|_{L^2(\rho)} \leq C \mathcal{W}_1(\mu_0, \mu_1)^{\frac{\theta}{q}},$$

which is the thesis. \square

ACKNOWLEDGEMENTS AND FUNDING

Both authors thank A. Pratelli, L. Ambrosio, S. Di Marino and A. Gerolin for useful discussions and comments during the development of this work.

D.T. acknowledges the MUR Excellence Department Project awarded to the Department of Mathematics, University of Pisa, CUP I57G22000700001, the HPC Italian National Centre for HPC, Big Data and Quantum Computing - Proposal code CN1 CN00000013, CUP I53C22000690001, the PRIN 2022 Italian grant 2022WHZ5XH - "understanding the LLearning process of QUantum Neural networks (LeQun)", CUP J53D23003890006, the INdAM-GNAMPa project 2024 "Tecniche analitiche e probabilistiche in informazione quantistica" and the project G24-202 "Variational methods for geometric and optimal matching problems" funded by Università Italo Francese. Research also partly funded by PNRR - M4C2 - Investimento 1.3, Partenariato Esteso PE00000013 - "FAIR - Future

Artificial Intelligence Research" - Spoke 1 "Human-centered AI", funded by the European Commission under the NextGeneration EU programme.

REFERENCES

- [AGS08] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient flows in metric spaces and in the space of probability measures*. Second edition. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2008. ISBN: 978-3-7643-8721-1 (cit. on pp. [1](#), [5](#), [18](#)).
- [AST19] L. Ambrosio, F. Stra, and D. Trevisan. “A PDE approach to a 2-dimensional matching problem”. In: *Probability Theory and Related Fields* 173 (2019), pp. 433–477 (cit. on p. [6](#)).
- [Ber21] R. J. Berman. “Convergence rates for discretized Monge–Ampère equations and quantitative stability of optimal transport”. In: *Foundations of Computational Mathematics* 21.4 (2021), pp. 1099–1140 (cit. on p. [3](#)).
- [BL76] H. J. Brascamp and E. H. Lieb. “On extensions of the Brunn–Minkowski and Prékopa–Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation”. In: *Journal of functional analysis* 22.4 (1976), pp. 366–389 (cit. on p. [3](#)).
- [BM18] H. Brezis and P. Mironescu. “Gagliardo–Nirenberg inequalities and non-inequalities: the full story”. In: *Ann. Inst. H. Poincaré C Anal. Non Linéaire* 35.5 (2018), pp. 1355–1376. ISSN: 0294-1449,1873-1430. URL: <https://doi.org/10.1017/S001309151800009> (cit. on p. [9](#)).
- [Caf92] L. A. Caffarelli. “The regularity of mappings with a convex potential”. In: *Journal of the American Mathematical Society* 5.1 (1992), pp. 99–104 (cit. on p. [3](#)).
- [CEMS06] D. Cordero-Erausquin, R. J. McCann, and M. Schmuckenschläger. “Prékopa–Leindler type inequalities on Riemannian manifolds, Jacobi fields, and optimal transport”. In: *Annales de la Faculté des sciences de Toulouse: Mathématiques*. Vol. 15. 4. 2006, pp. 613–635 (cit. on p. [4](#)).
- [Del22] A. Delalande. “Quantitative Stability in Quadratic Optimal Transport”. PhD thesis. Université Paris-Saclay, 2022 (cit. on pp. [3](#), [6–8](#), [10](#), [11](#), [13](#), [18–23](#)).
- [DM23] A. Delalande and Q. Merigot. “Quantitative stability of optimal transport maps under variations of the target measure”. In: *Duke Mathematical Journal* 172.17 (2023), pp. 3321–3357 (cit. on p. [3](#)).
- [DNWP24] V. Divol, J. Niles-Weed, and A.-A. Pooladian. “Tight stability bounds for entropic Brenier maps”. In: *arXiv preprint arXiv:2404.02855* (2024) (cit. on p. [3](#)).
- [FG10] A. Figalli and N. Gigli. “A new transportation distance between non-negative measures, with applications to gradients flows with Dirichlet boundary conditions”. In: *Journal de mathématiques pures et appliquées* 94.2 (2010), pp. 107–130 (cit. on p. [15](#)).
- [GHO21] M. Goldman, M. Huesmann, and F. Otto. “Quantitative Linearization Results for the Monge–Ampère Equation”. In: *Comm. Pure Appl. Math.* 74.12 (2021), pp. 2483–2560 (cit. on p. [6](#)).
- [GHT24] M. Goldman, M. Huesmann, and D. Trevisan. “Asymptotics for Random Quadratic Transportation Costs”. In: *in preparation* (2024) (cit. on p. [15](#)).
- [Gig11] N. Gigli. “On Hölder continuity-in-time of the optimal transport map towards measures along a curve”. In: *Proc. Edinb. Math. Soc. (2)* 54.2 (2011), pp. 401–409. ISSN: 0013-0915,1464-3839. URL: <https://doi.org/10.1017/S001309151100009> (cit. on p. [2](#)).
- [GMT22] A. Gallouët, Q. Mérigot, and B. Thibert. “Strong c-concavity and stability in optimal transport”. In: *arXiv preprint arXiv:2207.11042* (2022) (cit. on p. [3](#)).

- [HPS08] S. Hussain, J. Pečarić, and M Shashiashvili. “The weighted square integral inequalities for the first derivative of the function of a real variable”. In: *Journal of inequalities and applications* 2008 (2008), pp. 1–14 (cit. on p. 19).
- [Led17] M. Ledoux. “On optimal matching of Gaussian samples”. In: *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 457. Veroyatnost’ i Statistika. 25 (2017), pp. 226–264. ISSN: 0373-2703 (cit. on p. 6).
- [Lin19] P. Lindqvist. *Notes on the stationary p-Laplace equation*. Springer, 2019 (cit. on p. 20).
- [LN21] W. Li and R. H. Nochetto. “Quantitative stability and error estimates for optimal transport plans”. In: *IMA Journal of Numerical Analysis* 41.3 (2021), pp. 1941–1965 (cit. on p. 3).
- [McC94] R. McCann. “A convexity theory for interacting gases and equilibrium crystals”. In: *PhD Thesis, Princeton Univ.* (1994) (cit. on pp. 3, 5).
- [MDC20] Q. Mérigot, A. Delalande, and F. Chazal. “Quantitative stability of optimal transport maps and linearization of the 2-Wasserstein space”. In: *Proceedings of the Twenty Third International Conference on Artificial Intelligence and Statistics*. Ed. by S. Chiappa and R. Calandra. Vol. 108. Proceedings of Machine Learning Research. PMLR, 2020, pp. 3186–3196. URL: <https://proceedings.mlr.press/v108/merigot20a.html> (cit. on pp. 2, 3).
- [MG20] S. D. Marino and A. Gerolin. “An optimal transport approach for the Schrödinger bridge problem and convergence of Sinkhorn algorithm”. In: *Journal of Scientific Computing* 85.2 (2020), p. 27 (cit. on p. 6).
- [NW22] M. Nutz and J. Wiesel. “Entropic optimal transport: Convergence of potentials”. In: *Probability Theory and Related Fields* 184.1 (2022), pp. 401–424 (cit. on pp. 6, 7).
- [NZ18] M. Neilan and W. Zhang. “Rates of Convergence in Norm for the Monge–Ampère Equation”. In: *SIAM Journal on Numerical Analysis* 56.5 (2018), pp. 3099–3120 (cit. on p. 3).
- [PC+19] G. Peyré, M. Cuturi, et al. “Computational Optimal Transport: With Applications to Data Science”. In: *Foundations and Trends® in Machine Learning* 11.5-6 (2019), pp. 355–607 (cit. on p. 1).
- [Pey18] R. Peyre. “Comparison between W_2 distance and \dot{H}^{-1} norm, and localization of Wasserstein distance”. English. In: *ESAIM, Control Optim. Calc. Var.* 24.4 (2018), pp. 1489–1501. ISSN: 1292-8119 (cit. on p. 6).
- [PSSURNH15] J. Pečarić, M. Shoaib Saleem, H. Ur Rehman, A. M. Nizami, and A. Hussain. “Reverse Poincaré-type inequalities for the difference of superharmonic functions”. In: *Journal of Inequalities and Applications* 2015 (2015), pp. 1–11 (cit. on p. 19).
- [SS05] K. Shashiashvili and M. Shashiashvili. “Estimation of the derivative of the convex function by means of its uniform approximation”. In: *Journal of Inequalities in Pure and Applied Mathematics* 6.4 (2005), pp. 1–10 (cit. on p. 19).
- [Vil+09] C. Villani et al. *Optimal transport: old and new*. Vol. 338. Springer, 2009 (cit. on p. 2).

(Octave Mischler) DMA, ECOLE NORMALE SUPÉRIEURE PSL, 45 RUE D’ULM, 75005 PARIS 5, FRANCE
Email address: octave.mischler@ens.psl.eu

(Dario Trevisan) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127 - PISA (PI) ITALY
Email address: dario.trevisan@unipi.it