

# Rule 60 cellular automaton, Mersenne numbers, and the Newman-Moore spin lattice

Jonás Carmona-Pérez<sup>1</sup>, Adrian J. Peguero<sup>2</sup>, Vanja Dunjko<sup>2</sup>, Maxim Olshanii<sup>2</sup>, Joanna Ruhl<sup>2\*</sup>

**1** Technical Advisory Subdirectorate of Information Management (STAGI), Andalusian Health Service (SAS), 41071 Seville, Spain

**2** Department of Physics, University of Massachusetts Boston, Boston Massachusetts 02125, USA

\* joanna.ruhl001@umb.edu

July 30, 2024

## Abstract

The goal of this paper is to review the properties of a Rule 60 cellular automaton on a ring with a Mersenne number circumference and to use this knowledge to explicitly construct all the ground state configurations of the classical Newman-Moore model (a particular two-dimensional spin lattice model with a specific three-spin interaction) on a square lattice of the same size. In this particular case, the number of ground states is equal to half of the available spin configurations in any given row of the lattice.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Rule 60 cellular automaton</b>	<b>2</b>
2.1	Periodic trajectories	2
2.2	Configurations and their characterization	2
2.3	Useful lemmas	3
<b>3</b>	<b>Fundamental periods and a propagator for the Rule 60 on a ring of a Mersenne number circumference</b>	
3.1	Fundamental periods	5
3.2	Propagator	9
<b>4</b>	<b>Application to the Newman-Moore model</b>	<b>11</b>
4.1	Connection between the Rule 60 automaton and the Newman-Moore model	11
4.2	Newman-Moore results inferred from the properties of the Rule 60 automaton	12
<b>5</b>	<b>Summary of results</b>	<b>12</b>
	<b>References</b>	<b>13</b>

---

# 1 Introduction

Both the classical Newman-Moore model (a particular two-dimensional spin lattice model with a specific three-spin interaction) [1] and its quantum counterpart [2] play an important role in studies of systems with glassy dynamics (see, for example, [3], [4], [5] and references therein). In this paper, we review the properties of a Rule 60 cellular automaton on a ring with Mersenne number circumference and use this knowledge to explicitly construct all ground state configurations of the classical Newman-Moore model on a square lattice of the same size.

## 2 Rule 60 cellular automaton

For the one-dimensional cellular automata [6], the state of the system at the instant of time  $j$  is given by a length  $L$  string:

$$\vec{x}_j = [x_{0,j}, x_{1,j}, \dots, x_{i,j}, \dots, x_{L-1,j}], \quad (1)$$

where  $x_{i,j}$  is the state of the  $i$ 'th bit at the instance of time  $j$ . Each  $x_{i,j}$  is either 0, if the site is unoccupied, or 1, if the site is occupied. Consider a Rule 60 automaton [6] on a ring of a circumference  $L$ . According to the rule, the state of the automaton at the instance of time  $j + 1$  is related to its state at time  $j$  by

$$\begin{aligned} x_{i,j+1} &= \text{Xor}(x_{i-1,j}, x_{i,j}) & \text{for } i = 1, 2, \dots, L-1 \\ x_{0,j+1} &= \text{Xor}(x_{L-1,j}, x_{0,j}) \\ j &= 0, 1, \dots \end{aligned} \quad (2)$$

### 2.1 Periodic trajectories

In what follows, we will be interested in periodic trajectories generated by the rule (2).

**Definition 2.1.** An initial condition  $\vec{x}_0$  initiates a  **$M$ -periodic trajectory**  $\vec{x}_j$  if

$$\vec{x}_M = \vec{x}_0 .$$

**Definition 2.2.** An initial condition  $\vec{x}_0$  initiates a  **$M$ -fundamental-periodic trajectory**  $\vec{x}_j$ . if

$$\begin{aligned} \vec{x}_M &= \vec{x}_0 \\ \text{and} \\ \text{for any } 0 < j < M, \quad \vec{x}_j &\neq \vec{x}_0 . \end{aligned}$$

### 2.2 Configurations and their characterization

The state of the system at a given instant of time is called a **configuration**. E.g. for the trajectory

$$\begin{aligned} j = 0 & & [100] \\ j = 1 & & [110] \\ j = 2 & & [011] \\ j = 3 & & [101] \\ & & \dots \end{aligned} ,$$

[011] is the state of the size  $L = 3$  lattice at the instance of time  $j = 2$ , given that the initial condition at  $j = 0$  was [100].

**Definition 2.3.** Let  ${}_L N_{\text{occupied}}$ -type state indicate a configuration that features  $N_{\text{occupied}}$  occupied sites total on a length  $L$  lattice.

E.g. [110] will be characterised as a configuration of a  ${}_3 2$ -type.

### 2.3 Useful lemmas

**Lemma 2.1.** For any  $L$ , there are as many  ${}_L(\text{even})$ -type as there are  ${}_L(\text{odd})$ -type states.

*Proof.* One can establish a one-to-one correspondence between  ${}_L(\text{even})$  and  ${}_L(\text{odd})$  states simply by controlling the state of the first bit, leaving other bits intact.  $\square$

**Lemma 2.2.** For any  $L$ , the state [00...0] is a stationary state (a 1-periodic trajectory).

*Proof.* This directly follows from the rule (2).  $\square$

**Lemma 2.3.** For any ring of circumference  $L$ , both  ${}_L(\text{odd})$ -type and  ${}_L(\text{even})$ -type states become an  ${}_L(\text{even})$ -type state after one time step:

$$\begin{aligned} {}_L(\text{odd}) &\rightarrow {}_L(\text{even}) \\ {}_L(\text{even}) &\rightarrow {}_L(\text{even}) . \end{aligned}$$

*Proof.* One proof is presented in [6] as Lemma 3.1. We present an alternative proof here. Assume that the state of the system at an instance  $j$  is

$$[x_{0,j}, x_{1,j}, \dots, x_{i,j}, \dots, x_{L-1,j}] .$$

According to the rule (2), at the next step, 1's will be marking points where the site state changes from 0 to 1 or vice versa, along the lattice. For example, in

$$\begin{aligned} \vec{x}_j &= [011\mathbf{0}11001] \\ \vec{x}_{j+1} &= [110110101] , \end{aligned}$$

the boldface “1” at  $j + 1$  marks a lateral switch from “0” to “1”, in bold, at  $j$ . Likewise in

$$\begin{aligned} \vec{x}_j &= [\mathbf{0}11011001] \\ \vec{x}_{j+1} &= [110110101] , \end{aligned}$$

the boldface “1” at  $j + 1$  marks a lateral switch from “1” to “0” (in boldface), as the ring closes.

Observe now that on a ring, the site state can only change laterally an *even* number of times. This proves the Lemma.  $\square$

**Corollary 2.3.1.** Type  ${}_L(\text{odd})$  can not initiate a periodic motion.

*Proof.* For an  $M$ -periodic motion, the state must return to itself after  $M$  steps. But according to the Lemma 2.3, an  ${}_L(\text{odd})$ -type state can never become an  ${}_L(\text{odd})$ -type state.  $\square$

**Lemma 2.4.** *Let  $M_{\max}$  be the maximum fundamental period for a given  $L$ . Let  $M_{\mathcal{P}}$  be the period which the  $L1$  system converges to. Then*

$$M_{\max} = M_{\mathcal{P}} .$$

*Other periods divide  $M_{\max}$ .*

*Proof.* The Lemma 2.4 appears in [7] as Theorem 1. An alternative formulation related to cyclotomic polynomials is given in [8], Proposition 3.6.  $\square$

An explicit form of a propagator, for  $L = 2^n - 1$  ( $n = 2, 3, \dots$ ) is given by equation 9 in Theorem 3.6.

**Lemma 2.5.** *Let  $k$  and  $n$  be positive integers. Then, the binomial coefficient  $\binom{2^n}{k}$  is even for all  $0 < k < 2^n$ .*

Several proofs for this property are known, the following is perhaps the simplest one, adapted from [9] with minor changes.

*Proof.* According to the binomial theorem,

$$(x + 1)^{2^n} = \sum_{k=0}^{2^n} \binom{2^n}{k} x^k .$$

Therefore the lemma will be established if we prove that for all positive integer  $n$  there is some integer  $a_k$  with  $0 < k < 2^n$  such that

$$(x + 1)^{2^n} = x^{2^n} + 1 + \sum_{k=1}^{2^n-1} 2a_k x^k . \quad (3)$$

We will prove this by induction on  $n$ . First we prove the base case where  $n = 1$ , so that

$$(x + 1)^{2^n} = (x + 1)^2 = x^2 + 1 + 2x .$$

which has the form of 3 with  $a_1 = 1$ .

The induction hypothesis is that this property holds for  $n = m$ , for a given  $m \geq 1$ , so that

$$(x + 1)^{2^m} = x^{2^m} + 1 + \sum_{k=1}^{2^m-1} 2a_k x^k .$$

We now prove the property holds for  $m + 1$ :

$$(x + 1)^{2^{m+1}} = ((x + 1)^{2^m})^2$$

By the induction hypothesis this is

$$\begin{aligned} ((x + 1)^{2^m})^2 &= \left( x^{2^m} + 1 + \sum_{k=1}^{2^m-1} 2a_k x^k \right)^2 \\ &= \left\{ (x^{2^m} + 1)^2 \right\} + \left\{ 2(x^{2^m} + 1) \left( 2 \sum_{k=1}^{2^m-1} a_k x^k \right) \right\} + \left\{ \left( 2 \sum_{k=1}^{2^m-1} a_k x^k \right)^2 \right\} \\ &= x^{2^{m+1}} + 1 + 2x^{2^m} \\ &\quad + 4 \left\{ (x^{2^m} + 1) \sum_{k=1}^{2^m-1} a_k x^k + \left( \sum_{k=1}^{2^m-1} a_k x^k \right)^2 \right\} \end{aligned}$$

The first two terms to the right of the final equality are the first two terms of 3 with  $n = m + 1$ . The binomial theorem applied to  $(x + 1)^{2^{m+1}}$  produces these same two terms, and then a polynomial in  $x$  with powers running from 1 to  $2^{m+1} - 1$ , inclusive. Because all terms in this particular remainder are multiplied either by two or by four, they are all explicitly even, and the resulting polynomial will be sums of even numbers, which must therefore also be even. So it is possible to find numbers  $b_k$  such that

$$(x + 1)^{2^{m+1}} = x^{2^{m+1}} + 1 + \sum_{k=1}^{2^{m+1}-1} 2b_k x^k .$$

In particular,

$$b_k = \begin{cases} 2a_1, & \text{for } k = 1; \\ 2a_k + \sum_{m=1}^{k-1} a_{k-m}a_m, & \text{for } k = 2, \dots, 2^m - 1; \\ 1 + 2 \sum_{m=1}^{2^m-1} a_{2^m-m}a_m, & \text{for } k = 2^m; \\ 2a_{k-2^m} + \sum_{m=k-2^m+1}^{2^m-1} a_m a_{k-m}, & \text{for } k = 2^m + 1, \dots, 2^{m+1} - 2; \text{ and} \\ 2a_{2^m-1}, & \text{for } k = 2^{m+1} - 1. \end{cases}$$

□

**Lemma 2.6.** *For the Rule 60 automaton on a ring of circumference  $L = 2^n$  ( $n = 2, 3, \dots$ ), the only allowed fundamental period is  $M = 1$ .*

This result is well known [5–7]. However, in what follows, we will be proving a statement concerning temporal periods on a  $L = 2^n - 1$  ring, and we are going to use the proof of the Lemma above as a model.

*Proof.* [7] Lemma 1, Corollary 1; [5] Eq. 13. □

### 3 Fundamental periods and a propagator for the Rule 60 on a ring of a Mersenne number circumference

#### 3.1 Fundamental periods

**Definition 3.1.** A domain is the maximal contiguous set of sites having the same occupation state bounded by a right domain boundary and a left domain boundary.

**Definition 3.2.** A site  $x_{i,j}$  is said to be a right domain boundary (RDB) if the site  $x_{i+1,j}$  has a different occupation than  $x_{i,j}$ . Similarly, a site  $x_{i,j}$  is said to be a left domain boundary (LDB) if the site  $x_{i-1,j}$  has a different occupation. If  $x_{i,j}$  is both a RDB and a LDB, then it is a single-site domain. If  $x_{i,j}$  is neither a RDB or a LDB, then it is an interior site of a domain.

**Definition 3.3.** A bit flip on  $x_{i,j}$  is an operation which changes the occupation of that site, either from 1 to 0 or from 0 to 1, and leaves all other sites in  $\vec{x}_j$  unchanged.

**Definition 3.4.** A maximal state is a state of  $L$  with the maximum number of domains.

**Lemma 3.1.** *Every ring of odd circumference  $L$  has an even number of domains for all occupation states.*

*Proof.* Since  $L$  is odd, it is not possible to have only single-site domains on a ring. For all states of odd  $L$ , at least one domain must contain at least two sites. For the maximal state, only one domain will contain two sites, all others will be a single-site domain. Thus with periodic boundary conditions, when  $L$  is odd, the maximum number of domains is  $L - 1$ , which is even.

Any other state can be made from a maximal state by a series of bit flips on successive sites, up to  $L$  sites. On each site there are four possible cases:

Case 1: The site  $x_{i,j}$  is a single-site domain. By definition, this means the occupation states of  $x_{i-1,j}$  and  $x_{i+1,j}$  are the same. Therefore when the  $x_{i,j}$  site is bit flipped, its single-site domain will merge with the two domains to the right and to the left, reducing the total number of domains by two (three original domains become one). In a maximal state, there are an even number of domains, and bit flipping any of the single site domains results in a net loss of two domains, so the total number of domains remains even.

Case 2: The site  $x_{i,j}$  is an interior site of a domain. This is essentially the reverse of Case 1; when this site is bit flipped, it will then become a single site domain, with two new domains on either side. The original single domain becomes three, for a net gain of two domains. Since the number of domains in a maximal state is even, if a site interior to a domain is bit flipped, two domains are added and the total remains even.

Case 3: The site  $x_{i,j}$  is a LDB, but not a RDB. By definition, since  $x_{i,j}$  is a LDB but not a RDB it cannot be a single-site domain. Bit flipping this site will therefore shift the domain boundary to the right one site, merging the  $x_{i,j}$  site with the domain that previously had the RDB at  $x_{i-1,j}$  and creating a new LDB at  $x_{i+1,j}$ . The total number of domains remains the same. Since the maximal state has an even number of domains, bit flipping a site which is a LDB but not a single-site domain results in the same even number of domains.

Case 4: The site  $x_{i,j}$  is a RDB, but not a LDB. Again by definition, this site cannot be a single-site domain. Similarly to Case 3, bit flipping this site will result in the domain boundary shifting left one site, making  $x_{i,j}$  the RDB previously at  $x_{i+1,j}$ . The total number of domains again remains the same. Therefore all possible cases of bit flips from the maximal state result in a state with an even number of domains.  $\square$

**Remark:** The above proof can be extended to even  $L$  to formulate an alternate proof of Lemma 2.3.

**Definition 3.5.** A predecessor state of  $\vec{x}_{j+1}$  is any state  $\vec{x}_j$  that evolves to  $\vec{x}_{j+1}$  under the time evolution specified in (2)

**Lemma 3.2.** A state on a ring of circumference  $L$  has a predecessor state if, and only if, it is even.

*Proof.* We have already shown in Lemma 2.3 that all Rule 60 time evolution results in even states, so no  $L(\text{odd})$  state has a predecessor. If a predecessor does exist, it must be constructible by the following inverse of Rule 60. Here we construct an  $\vec{x}_j$  configuration

from the knowledge of the  $\vec{x}_{j+1}$  configuration. This inverse rule is

$$\begin{aligned} x_{0,j} &= 1 \\ x_{i,j} &= \text{Xor}(x_{i-1,j}, x_{i-1,j+1}) \quad \text{for } i = 1, 2, \dots, L-1. \end{aligned} \quad (4)$$

Because of the periodic boundary condition, there is also the self-consistency condition

$$x_{0,j} = \text{Xor}(x_{L-1,j}, x_{L-1,j+1}). \quad (5)$$

Note that there is a freedom in the initial choice of  $x_{0,j}$ , we have chosen 1 for this proof, but one can equally choose 0 and the rest of the construction goes through as above.

We will now prove that all even states have a predecessor by induction on  $L$ .

For the base case, we take  $L = 2$  at some time  $j + 1$ . There are four possible configurations:

$$\begin{aligned} &[00] \\ &[01] \\ &[10] \\ &[11] \end{aligned}$$

Applying the inverse rule (4), we find:

$$\begin{aligned} [00] &\text{ has predecessor } [11] \\ [11] &\text{ has predecessor } [10] \end{aligned}$$

The configurations [01] and [10] do not have predecessors, as application of the inverse rule (4) can not satisfy the self-consistency condition in both cases. Thus, we have proven by demonstration that all even states for  $L = 2$  have a predecessor state, and no odd states have a predecessor.

The induction hypothesis is that this holds for some  $L = N$ , so that given an  $\vec{x}_{j+1}$  of length  $N > 2$

$$\begin{aligned} x_{i,j} &= \text{Xor}(x_{i-1,j}, x_{i-1,j+1}) && \text{for } i = 1, 2, \dots, N-1 \\ \text{Xor}(x_{N-1,j}, x_{N-1,j+1}) &= 1 && \text{for}_N(\text{even}) \\ \text{Xor}(x_{N-1,j}, x_{N-1,j+1}) &= 0 && \text{for}_N(\text{odd}) \end{aligned}$$

We now prove the property holds for  $N + 1$ . There are two cases:

Case 1: The ring of circumference  $N$  had an even occupation, so by the induction hypothesis

$$\text{Xor}(x_{N-1,j}, x_{N-1,j+1}) = 1.$$

Extending the ring to  $N + 1$  means adding another site,  $x_{N,j+1}$ , which can be either occupied or unoccupied. If the site is occupied then the total occupation of  $N + 1$  is odd. By (4)

$$x_{N,j} = \text{Xor}(x_{N-1,j}, x_{N-1,j+1}) = 1$$

but by the self-consistency condition (5)

$$x_{0,j} = \text{Xor}(x_{N,j}, x_{N,j+1}) = 0$$

which is a contradiction, and so there can be no predecessor state.

If  $x_{N,j+1}$  is unoccupied, then again

$$x_{N,j} = \text{Xor}(x_{N-1,j}, x_{N-1,j+1}) = 1$$

but now

$$x_{0,j} = \text{Xor}(x_{N,j}, x_{N,j+1}) = \text{Xor}(1, 0) = 1$$

and the predecessor state has been successfully constructed using rule (4).

Case 2: The ring of circumference  $N$  had an odd occupation, so by the induction hypothesis

$$\text{Xor}(x_{N-1,j}, x_{N-1,j+1}) = 0.$$

If the additional site is occupied, so  $x_{N,j+1} = 1$  then the occupation of  $N + 1$  is now even, and

$$x_{0,j} = \text{Xor}(x_{N,j}, x_{N,j+1}) = \text{Xor}(1, 0) = 1$$

and the predecessor state has been successfully constructed.

If  $x_{N,j+1} = 0$ , then the  $N + 1$  state remains odd and

$$x_{0,j} = \text{Xor}(x_{N,j}, x_{N,j+1}) = \text{Xor}(0, 0) = 0$$

in contradiction to (5) and a predecessor state can not be constructed. Therefore, the ring of circumference  $N + 1$  has a predecessor state only if the occupation is even.  $\square$

**Lemma 3.3.** *Every  $L$ (even) state on a ring has two possible predecessor states, which are related by bit flipping every site.*

*Proof.* In Lemma 3.2 we demonstrated that every  $L$ (even) state has a predecessor, but noted that there is a freedom in choosing the occupation of the initial site of the predecessor state. Since the Xor operation only considers relationships between pairs of sites,  $x_{i,j}$  and  $x_{i-1,j}$ , bit flipping every site preserves the outcome of the Xor operation and generates a second predecessor. Since this is the only freedom in constructing the predecessor states, these two states are the only predecessor states.  $\square$

**Lemma 3.4.** *On a ring of odd circumference  $L$ , every  $L$ (even) state has one  $L$ (odd) precursor.*

*Proof.* Construct an  $\vec{x}_j$  state from a given  $\vec{x}_{j+1}$  by applying (4). If the occupation is odd, then the proof is complete. If the occupation is even, then by Lemma 3.3 the second state can be found by bit flipping every site of the even state. Since  $L$  is odd, this flipped state will be odd.  $\square$

**Theorem 3.5.** *For a ring whose circumference is a Mersenne number,  $L = 2^n - 1$  ( $n = 2, 3, \dots$ ), every  $(L=2^n-1)$ (even)-type initial state ( $2^{L-1}$  of them) initiates an  $L$ -periodic trajectory. These periodic trajectories exhaust the set of periodic trajectories.*

**Remark:** The proof that follows is not new. The most elegant one can be found in [7] Lemma 1, Corollary 3. Our proof differs in several details from the one in [7]. In particular, it does not rely on the properties of the propagators, and as such is technically close to the one given in [5] for  $L = 2^n$ .

*Proof.* Regard the state of the system at an instance of time  $j$  as a vector  $\vec{x}_j$ . Observe that

$$\vec{x}_j \stackrel{\text{mod } 2}{=} (\hat{I} + \hat{S}_L)^j \vec{x}_0,$$



where  $\hat{I}$  is a unit matrix,  $\hat{S}_L$  is a left shift operator:

$$\begin{aligned} (\hat{S}_L \vec{x})_i &= (\vec{x})_{i-1} \quad \text{for } i = 1, 2, \dots, L-1 \\ (\hat{S}_L \vec{x})_0 &= (\vec{x})_{L-1} \end{aligned} .$$

Below we will use a particular obvious property of the shift operator:

$$\left(\hat{S}_L\right)^L = \hat{I} . \tag{6}$$

Now, let us consider  $L = 2^n - 1$  and apply the Rule 60  $L + 1 = 2^n$  times. We get

$$\begin{aligned} (\hat{I} + \hat{S}_L)^{L+1} &= \hat{I} + \sum_{r=1}^L \binom{L+1}{r} (\hat{S}_L)^r + (\hat{S}_L)^{L+1} \\ &= \hat{I} + \sum_{r=1}^{2^n-1} \binom{2^n}{r} (\hat{S}_L)^r + \hat{S}_L \\ &\stackrel{\text{mod } 2}{=} \hat{I} + \hat{S}_L . \end{aligned}$$

Above, we used the properties the property (6) and Lemma 2.5. In the other words, for any initial state  $\vec{x}_0$ ,

$$\vec{x}_{L+1} = \vec{x}_1 .$$

Now, thanks to the Lemma 3.4,  $\vec{x}_1$  can be represented by *any*  $L$ (even) state. Shifting the time by one step back, we get that for *any*  $(L=2^n-1)$ (even) initial state  $\vec{x}_0$ ,

$$\vec{x}_L = \vec{x}_0 .$$

This is exactly the property we need.

Finally, according to the Corollary 2.3.1, no other periodic motions are possible.  $\square$

**Corollary 3.5.1.** *For a ring whose circumference  $L = 2^n - 1 = \text{prime}$  is a Mersenne prime, every  $L$ (even)-type initial state different from all zeros ( $2^{L-1} - 1$  of them) initiates a  $L$ -fundamental-periodic trajectory.*

*Proof.* According to the Theorem 3.5,  $L$  is always one of the periods of *every* periodic trajectory. Thus, it is the maximal fundamental period Lemma 2.4 is devoted to:

$$M_{\max} = L .$$

The second part of the Lemma 2.4 says that all the smaller periods divide  $M_{\max}$ . But in the case of  $L$  being a Mersenne prime,  $L$  is prime and as such has no factors besides unity (and the “all zeros” initial state is the only one that is capable of producing a unit period) and itself. Hence  $L$  is indeed the only available fundamental period.

Finally, while *any*  $(L=2^n-1)$ (even)-type state initiates a periodic trajectory, these exhaust the list. Indeed, according to the Corollary 2.3.1, the  $L$ (odd)-type states cannot initiate periodic trajectories.  $\square$

### 3.2 Propagator

To aid in the aim of explicitly constructing all ground states of the Newman-Moore model, let us first introduce a propagator, i.e. a solution of (2) that has only one site occupied at  $j = 0$ . Namely let us introduce

$$\mathcal{P}_{(i,j) \leftarrow (i_0,0)} \tag{7}$$

that obeys

$$\begin{aligned}
\mathcal{P}_{(i,0)\leftarrow(i_0,0)} &= \delta_{i,i_0} \\
\mathcal{P}_{(i,j+1)\leftarrow(i_0,0)} &= \text{Xor}(\mathcal{P}_{(i-1,j)\leftarrow(i_0,0)}, \mathcal{P}_{(i,j)\leftarrow(i_0,0)}) \quad \text{for } i = 1, 2, \dots, L-1 \\
\mathcal{P}_{(0,j+1)\leftarrow(i_0,0)} &= \text{Xor}(\mathcal{P}_{(L-1,j)\leftarrow(i_0,0)}, \mathcal{P}_{(0,j)\leftarrow(i_0,0)}) \\
j &= 0, 1, \dots \\
i_0 &= 0, 1, L-1
\end{aligned} \tag{8}$$

**Theorem 3.6.** For  $L = 2^n - 1$  ( $n = 2, 3, \dots$ ),

$$\mathcal{P}_{(i,j)\leftarrow(0,0)} = \begin{cases} \binom{j}{i}_L \bmod 2, & 0 \leq j \leq L-1 \\ (1 - \delta_{i,0}), & j = L \\ \mathcal{P}_{(i,(j-1 \bmod L)+1)\leftarrow(0,0)}, & j > L \end{cases} \tag{9}$$

where  $\binom{j}{i}_L$  is the q-binomial coefficient defined as

$$\binom{n}{m}_q = \prod_{k=0}^{m-1} \frac{1 - q^{n-k}}{1 - q^{k+1}} \tag{10}$$

*Proof.* For the initial state, note that

$$\binom{0}{i}_L = \begin{cases} 1 & i = 0 \\ 0 & \text{otherwise} \end{cases} \tag{11}$$

so the  $j = 0$  state has a single occupied site at  $i = 0$  as desired.

As stated in (2), the time evolution is given by

$$x_{i,j+1} = \text{Xor}(x_{i-1,j}, x_{i,j}), \quad 0 < j \leq L-1. \tag{12}$$

This is equivalent to addition mod 2, so

$$x_{i,j+1} = [x_{i-1,j} + x_{i,j}] \bmod 2. \tag{13}$$

We make use of the known recursion relation for the q-binomial coefficients

$$\binom{j+1}{i}_L = \binom{j}{i-1}_L + L^i \binom{j}{i}_L. \tag{14}$$

Note that for  $L = 2^n - 1$

$$L^i \bmod 2 = 1 \quad \forall i \tag{15}$$

since all terms in the expansion of  $(2^n - 1)^i$  will be powers of two and their multiples, except the  $\binom{i}{i} (2^n)^0 (-1)^i$  term. Thus

$$\binom{j+1}{i}_L \bmod 2 = \left[ \binom{j}{i-1}_L + \binom{j}{i}_L \right] \bmod 2, \tag{16}$$

which is equivalent to (13).  $\square$

For other propagators, the translational invariance dictates

**Theorem 3.7.**

$$\mathcal{P}_{(i,j)\leftarrow(i_0,0)} = \begin{cases} \mathcal{P}_{(i+i_0,j)\leftarrow(i_0,0)} & \text{for } i \leq 2(L-1) - i_0 \\ \mathcal{P}_{(i+i_0-(L-1),j)\leftarrow(i_0,0)} & \text{for } i > 2(L-1) - i_0 \end{cases}$$

$$i_0 = 0, 1 \dots, L-1$$

$$i = 0, 1 \dots, L-1$$

$$j = 0, 1 \dots, L-1$$
(17)

*Proof.* This follows directly from the definition of the propagator. □

The propagator (17) can be used to propagate the Rule 60 automaton with  $L = 2^n - 1$ :

$$x_{i,j} = \sum_{i_0=0}^{L-1} \mathcal{P}_{(i,j)\leftarrow(i_0,0)} x_{i_0,j}$$

$$i = 0, 1 \dots, L-1$$

$$j = 0, 1 \dots, L-1$$

Note that the formulas ((9), (17)) are not directly applicable for  $L \neq 2^n - 1$ .

## 4 Application to the Newman-Moore model

### 4.1 Connection between the Rule 60 automaton and the Newman-Moore model

To connect the Rule 60 automaton and the two-dimensional spin lattices, one considers the so-called Triangular Plaquette Model or Newman-Moore model [1, 3, 10]:

$$H = -J \sum_{i=0}^{L-1} \sum_{j=0}^{M-1} \sigma_{i,j} \sigma_{i+1,j} \sigma_{i+1,j+1}$$
(18)

with periodic boundary conditions. Notice that according to the rule 60, it is always the case that

$$(1 - 2x_{i,j})(1 - 2x_{i+1,j})(1 - 2x_{i+1,j+1}) = 1 .$$

Hence, every  $M$ -periodic trajectory of (2) will produce a ground state of the Hamiltonian (18), whose energy is

$$E_{\text{ground}} = E_{\uparrow \dots \uparrow} = -JLM ,$$

if one uses the following association:

$$\sigma_{i,j} = (1 - 2x_{i,j}) .$$

Notice that “all spins up” is one of the ground states.

## 4.2 Newman-Moore results inferred from the properties of the Rule 60 automaton

We are now ready to address the principal application of our cellular automata result: the Newman-Moore model. In what follows, we will be able to explicitly list all the ground state configurations of the Newman-Moore model (18) with

$$\begin{aligned} M = L = 2^n - 1 \\ n = 2, 3, \dots, \end{aligned}$$

i.e. for a square Newman-Moore lattice with the sides equal to one of the Mersenne numbers.

We assert the following.

**Theorem 4.1.** *The ground state of the Newman-Moore model (18) with  $M = L = 2^n - 1$  is  $2^{L-1}$  degenerate.*

*Proof.* The proof follows directly from the Theorem 3.5.  $\square$

We will now provide explicit expressions for the ground state configurations of the Newman-Moore model (18) on an  $L \times L$  square lattice with  $L$  being equal to one of the Mersenne numbers  $2^n - 1$ .

The ground state configurations will be labeled by the state of the 0-th row:

$$[\sigma_{0,0}, \sigma_{1,0}, \dots, \sigma_{L-1,0}],$$

where number of spin-down sites,  $\sigma = -1$  is even: there are  $2^{L-1}$  such configurations, consistent with the  $2^{L-1}$ -fold degeneracy.

**Theorem 4.2.** *The ground states of the Newman-Moore model (18) on an  $L \times L$  square lattice with  $L = (2^n - 1)$  ( $n = 2, 3, \dots$ ) read*

$$\begin{aligned} \sigma_{i,j} &= \sum_{i_0=0}^{L-1} \left( 1 - 2\mathcal{P}_{(i,j) \leftarrow (i_0,0)} \frac{1 - \sigma_{i_0,j}}{2} \right) \\ i &= 0, 1 \dots, 2^n - 2 \\ j &= 0, 1 \dots, 2^n - 2 \end{aligned} \tag{19}$$

where the propagator  $\mathcal{P}_{(i,j) \leftarrow (i_0,0)}$  is given by (17) and  $[\sigma_{0,0}, \sigma_{1,0}, \dots, \sigma_{2^n-2,0}]$  is any of the  $2^{L-1}$  spin configurations characterised by the number of the  $\sigma = -1$  sites being even.

## 5 Summary of results

The Theorem 3.5 plays the central role in our paper. It allows to identify all the initial conditions that lead to periodic trajectories in a Rule 60 cellular automaton on a ring whose circumference  $L$  is given by one of the Mersenne numbers,  $L = 2^n - 1$ . Furthermore, all periodic trajectories are shown to share the period  $L$ . The Theorem 3.6 constructs the propagators for this model.

The above results allow us to list all the ground states of the Newman-Moore model explicitly. These results are summarised in Theorems 4.1 and 4.2

## Acknowledgements

The authors wish to thank Konstantinos Sfairopoulos for bringing this problem to their attention, for providing invaluable guidance on references, and for many useful discussions.

A significant portion of this work was produced during the thematic trimester on “Quantum Many-Body Systems Out-of-Equilibrium”, at the Institut Henri Poincaré (Paris): MO is immeasurably grateful to the organizers of the trimester, Rosario Fazio, Thierry Giamarchi, Anna Minguzzi, and Patrizia Vignolo, for an opportunity to be a part of it. MO and JC-P wish to express their gratitude to the International High IQ Society for the networking opportunities that it offers.

**Funding information** MO was supported by the NSF Grant No. PHY-1912542. MO would like to thank the Institut Henri Poincaré (UAR 839 CNRS-Sorbonne Université) and the LabEx CARMIN (ANR-10-LABX-59-01) for their support.

## References

- [1] M. E. J. Newman and C. Moore, *Glassy dynamics and aging in an exactly solvable spin model*, Phys. Rev. E **60**, 5068 (1999), doi:[10.1103/PhysRevE.60.5068](https://doi.org/10.1103/PhysRevE.60.5068).
- [2] B. Yoshida and A. Kubica, *Quantum criticality from ising model on fractal lattices* (2014), [1404.6311](https://arxiv.org/abs/1404.6311).
- [3] R. L. Jack, L. Berthier and J. P. Garrahan, *Static and dynamic length scales in a simple glassy plaquette model*, Phys. Rev. E **72**, 016103 (2005), doi:[10.1103/PhysRevE.72.016103](https://doi.org/10.1103/PhysRevE.72.016103).
- [4] A. Lipowski, D. Johnston and D. Espriu, *Slow dynamics of ising models with energy barriers*, Phys. Rev. E **62**, 3404 (2000), doi:[10.1103/PhysRevE.62.3404](https://doi.org/10.1103/PhysRevE.62.3404).
- [5] K. Sfairopoulos, L. Causer, J. F. Mair and J. P. Garrahan, *Boundary conditions dependence of the phase transition in the quantum Newman-Moore model*, Phys. Rev. B **108**, 174107 (2023), doi:[10.1103/PhysRevB.108.174107](https://doi.org/10.1103/PhysRevB.108.174107).
- [6] O. Martin, A. M. Odlyzko and S. Wolfram, *Algebraic properties of cellular automata*, Commun. Math. Phys. **93**, 219 (1984).
- [7] A. Ehrlich, *Periods in Ducci’s n-number game of differences*, The Fibonacci Quarterly **28**, 302 (1990).
- [8] F. Breuer, E. Lötter and B. van der Merwe, *Ducci-sequences and cyclotomic polynomials*, Finite Fields and Their Applications **13**(2), 293 (2007), doi:<https://doi.org/10.1016/j.ffa.2005.11.003>.
- [9] D. Schepler, *How to prove binomial coefficient  $\binom{2^n}{k}$  is even number?*, Mathematics Stack Exchange, URL:<https://math.stackexchange.com/q/3087725> (version: 2019-01-25), <https://math.stackexchange.com/q/3087725>.
- [10] *Newman-Moore code*, In V. V. Albert and P. Faist, eds., *The Error Correction Zoo* (2023).