

The Hadamard Product of Moment Sequences

Philipp J. di Dio

Department of Mathematics and Statistics, University of Konstanz, Universitätsstraße 10,
D-78464 Konstanz, Germany

Zukunftskolleg, University of Konstanz, Universitätsstraße 10, D-78464 Konstanz,
Germany

philipp.didio@uni-konstanz.de

Abstract

We define the (Hadamard) product $s \odot t := (s_\alpha \cdot t_\alpha)_{\alpha \in \mathbb{N}_0^n}$ of two sequences $s = (s_\alpha)_{\alpha \in \mathbb{N}_0^n}$ and $t = (t_\alpha)_{\alpha \in \mathbb{N}_0^n}$. If s is represented by a measure μ and t by a measure ν , i.e., both are moment sequences, then we give an explicit measure $\mu \odot \nu$ that represents $s \odot t$, i.e., $s \odot t$ is a moment sequence. We pick a result from *Moment-sequence transforms* (J. Eur. Math. Soc. **24** (2022), 3109–3160) and show how the multiplication $s \odot t$ of moment sequences simplifies the proof and also gives deeper insight by providing the representing measure $\mu \odot \nu$.

Keywords: moment sequence, moment, Hadamard, product, entry-wise
2020 MSC: Primary 44A60; Secondary 30E05, 26C05.

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function and $n \in \mathbb{N}$. Entry-wise maps

$$f : \mathbb{R}^{\mathbb{N}_0^n} \rightarrow \mathbb{R}^{\mathbb{N}_0^n}, \quad s \mapsto f(\odot s) := (f(s_\alpha))_{\alpha \in \mathbb{N}_0^n} \quad (1)$$

have intensively been studied in various connections, see e.g. [Sch42, KT21, BGKP22, BGKP23] and reference therein. Especially in [BGKP22] maps (1) were studied which preserve moment sequences, i.e., $f(\odot s)$ is a moment sequence for any moment sequence s . For a closed set $K \subseteq \mathbb{R}^n$ recall that a sequence $s = (s_\alpha)_{\alpha \in \mathbb{N}_0^n}$ is a *K-moment sequence* if there exists a measure μ on K such that $s_\alpha = \int_K x^\alpha d\mu(x)$ holds for all $\alpha \in \mathbb{N}_0^n$.

Despite the long history of the theory of moments and its extensive literature [Sti94, Akh65, KN77, Lau09, Mar08, Sch17] the (Hadamard) product of moment sequences is not contained in the standard literature. It has been studied for a long time but it is surprising that the connection to [Bor11, Cor. 4.3] was not made before. We apply this to [BGKP22, Thm. 3.1] and not only simplify the proof and arguments but also provide a deeper understanding what maps (1) do with moment sequences and their representing measures.

We thank Greg Blekherman for valuable remarks and reference [BRS⁺22].

Our work is based on the following result of Borcea [Bor11]. Let

$$T : \mathbb{R}[x_1, \dots, x_n] \rightarrow \mathbb{R}[x_1, \dots, x_n] \quad \text{with} \quad Tx^\alpha = t_\alpha x^\alpha \quad \text{for all } \alpha \in \mathbb{N}_0^n \quad (2)$$

be a linear map for some real sequence $t = (t_\alpha)_{\alpha \in \mathbb{N}_0^n}$. Operators T as in (2) are called *diagonal operators* and $(t_\alpha)_{\alpha \in \mathbb{N}_0^n}$ is its *diagonal sequence*. Define $\text{Pos}(\mathbb{R}^n) := \{p \in \mathbb{R}[x_1, \dots, x_n] \mid p \geq 0 \text{ on } \mathbb{R}^n\}$. By [Bor11, Cor. 4.3]¹ the following are equivalent:

- (a) T is a positivity preserver, i.e., $T\text{Pos}(\mathbb{R}^n) \subseteq \text{Pos}(\mathbb{R}^n)$.
- (b) $t = (t_\alpha)_{\alpha \in \mathbb{N}_0^n}$ is a moment sequence.

Additionally, if μ is a representing measure of the moment sequence t then

$$(Tp)(x) = \int_{\mathbb{R}^n} p(x_1 y_1, \dots, x_n y_n) \, d\mu(y_1, \dots, y_n) \quad (3)$$

holds for all $p \in \mathbb{R}[x_1, \dots, x_n]$.

Its proof is $L(p) := (Tp)(1, \dots, 1) \geq 0$ for all $p \in \text{Pos}(\mathbb{R}^n)$ and $L(x^\alpha) = \lambda_\alpha$.

Let T and S be operators of the form (2) with moment sequences $t = (t_\alpha)_{\alpha \in \mathbb{N}_0^n}$ and $s = (s_\alpha)_{\alpha \in \mathbb{N}_0^n}$ represented by measures μ and ν , respectively. Then the operator $ST = TS$ is a diagonal operator which fulfills

$$(TS)(p)(x) = (ST)(p)(x) = \int_{\mathbb{R}^n \times \mathbb{R}^n} p(x_1 y_1 z_1, \dots, x_n y_n z_n) \, d\mu(y_1, \dots, y_n) \, d\nu(z_1, \dots, z_n) \quad (4)$$

for all $p \in \mathbb{R}[x_1, \dots, x_n]$, i.e., $TSx^\alpha = t_\alpha s_\alpha x^\alpha$ for all $\alpha \in \mathbb{N}_0^n$ and therefore with diagonal sequence $(t_\alpha \cdot s_\alpha)_{\alpha \in \mathbb{N}_0^n}$. This motivates the following.

Definition 1. Let $n \in \mathbb{N}$, let $s = (s_\alpha)_{\alpha \in \mathbb{N}_0^n}$ and $t = (t_\alpha)_{\alpha \in \mathbb{N}_0^n} \in \mathbb{R}^{\mathbb{N}_0^n}$ be real sequences, let μ and ν be measures on \mathbb{R}^n , and let $k \in \mathbb{N}$. We define

$$s \odot t := (s_\alpha \cdot t_\alpha)_{\alpha \in \mathbb{N}_0^n}, \quad s^{\odot 0} := (1)_{\alpha \in \mathbb{N}_0^n}, \quad \text{and} \quad s^{\odot k} := \underbrace{s \odot \dots \odot s}_{k\text{-times}} = (s_\alpha^k)_{\alpha \in \mathbb{N}_0^n}.$$

Additionally, we define $\mu \odot \nu$ by

$$(\mu \odot \nu)(A) := \int_{\mathbb{R}^n \times \mathbb{R}^n} \chi_A(x_1 y_1, \dots, x_n y_n) \, d\mu(x_1, \dots, x_n) \, d\nu(y_1, \dots, y_n)$$

for all Borel sets $A \subseteq \mathbb{R}^n$. χ_A is the characteristic function of A . We define

$$\mu^{\odot 0} := \delta_1 \quad \text{and} \quad \mu^{\odot k} := \underbrace{\mu \odot \dots \odot \mu}_{k\text{-times}}$$

where δ_x is the Dirac measure supported at $x \in \mathbb{R}^n$.

¹Submitted: 17th of October 2008; revised: 12th of January 2009; published online: 7th of January 2011, i.e., after the death of the author Julius Borcea (8th of June 1968 – 8th of April 2009). Because of the untimely death of J. Borcea the publication was significantly delayed.

Clearly, $\mu \odot \nu$ is a measure on \mathbb{R}^n . For two sets $A, B \subseteq \mathbb{R}^n$ we define

$$A \cdot B := \{(a_1 b_1, \dots, a_n b_n) \mid (a_1, \dots, a_n) \in A, (b_1, \dots, b_n) \in B\}. \quad (5)$$

Then (4) implies the following.

Theorem 2. *Let $n \in \mathbb{N}$ and let $s, t \in \mathbb{R}^{\mathbb{N}_0^n}$ be moment sequences with representing measures μ and ν . Then $s \odot t$ is a moment sequence with representing measure $\mu \odot \nu$ which fulfills $\text{supp } \mu \odot \nu = \text{supp } \mu \cdot \text{supp } \nu$.*

Proof. Follows immediately from (4) and Definition 1. \square

Remark 3. Theorem 2 not only states that the map $f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2$ induces a moment sequence preserving map f as in (1). By induction any map $f(x) = \sum_{k=0}^d a_k \cdot x^k$ with $d \in \mathbb{N}_0$ and $a_k \geq 0$ for all $k = 0, \dots, d$ preserves moment sequences entry-wise. If s is represented by μ then $f(\odot s) = \sum_{k=0}^d a_k \cdot s^{\odot k}$ is represented by the measure $f(\odot \mu) := \sum_{k=0}^d a_k \cdot \mu^{\odot k}$. Under appropriate conditions on the coefficients a_k the limit $d \rightarrow \infty$ exists for the sequence $f(\odot s)$ and the (signed) measure $f(\odot \mu)$. That f is entire is sufficient. \circ

Theorem 2 and Remark 3 are simple observations from [Bor11, Cor. 4.3] but have neither been formulated nor used before. It reduces the long studied problem of the Hadamard product to triviality. Let us collect properties of \odot .

Lemma 4. *Let $n \in \mathbb{N}$, let μ, ν , and ω be measures on \mathbb{R}^n , let A be a Borel set of \mathbb{R}^n , let $u = (u_1, \dots, u_n), v = (v_1, \dots, v_n) \in \mathbb{R}^n$, and let $a, b \geq 0$ and $c > 0$ be constants. Then the following hold:*

- (i) $\mu \odot \nu = \nu \odot \mu$.
- (ii) $(a\mu + b\nu) \odot \omega = a \cdot \mu \odot \omega + b \cdot \nu \odot \omega$.
- (iii) $(\mu \odot \nu) \odot \omega = \mu \odot (\nu \odot \omega)$.
- (iv) $(\mu \odot \nu)(\mathbb{R}^n) = \mu(\mathbb{R}^n) \cdot \nu(\mathbb{R}^n)$.
- (v) $\delta_0 \odot \mu = \mu(\mathbb{R}^n) \cdot \delta_0$.
- (vi) $\delta_{\mathbf{1}} \odot \mu = \mu$ with $\mathbf{1} := (1, \dots, 1) \in \mathbb{R}^n$.
- (vii) $(\delta_{c \cdot \mathbf{1}} \odot \mu)(A) = \mu(c^{-1} \cdot A)$.
- (viii) $\delta_u \odot \delta_v = \delta_{u \odot v}$ with $u \odot v := (u_1 v_1, \dots, u_n v_n)$.

Proof. (i) – (viii) follow all immediately from Definition 1 and Theorem 2. We prove only (v) – (viii) explicitly.

(v): Let $A \subseteq \mathbb{R}^n$ be a Borel set. We have

$$(\delta_0 \odot \mu)(A) = \int_{\mathbb{R}^n \times \mathbb{R}^n} \chi_A(x_1 y_1, \dots, x_n y_n) d\mu(x_1, \dots, x_n) d\delta_0(y_1, \dots, y_n)$$

$$\begin{aligned}
&= \int_{\mathbb{R}^n} \chi_A(x_1 \cdot 0, \dots, x_n \cdot 0) \, d\mu(x_1, \dots, x_n) = \mu(\mathbb{R}^n) \cdot \chi_A(0, \dots, 0) \\
&= \mu(\mathbb{R}^n) \cdot \delta_0(A).
\end{aligned}$$

(vi): Let $A \subseteq \mathbb{R}^n$ be a Borel set. We have

$$\begin{aligned}
(\delta_{\mathbf{1}} \odot \mu)(A) &= \int_{\mathbb{R}^n \times \mathbb{R}^n} \chi_A(x_1 y_1, \dots, x_n y_n) \, d\mu(x_1, \dots, x_n) \, d\delta_{\mathbf{1}}(y_1, \dots, y_n) \\
&= \int_{\mathbb{R}^n} \chi_A(x_1 \cdot 1, \dots, x_n \cdot 1) \, d\mu(x_1, \dots, x_n) = \mu(A).
\end{aligned}$$

(vii): Let $A \subseteq \mathbb{R}^n$ be a Borel set. We have

$$\begin{aligned}
(\delta_{c \cdot \mathbf{1}} \odot \mu)(A) &= \int_{\mathbb{R}^n \times \mathbb{R}^n} \chi_A(x_1 y_1, \dots, x_n y_n) \, d\delta_{c \cdot \mathbf{1}}(x_1, \dots, x_n) \, d\mu(y_1, \dots, y_n) \\
&= \int_{\mathbb{R}^n} \chi_A(c y_1, \dots, c y_n) \, d\mu(y_1, \dots, y_n) \\
&= \int_{\mathbb{R}^n} \chi_{c^{-1} \cdot A}(y_1, \dots, y_n) \, d\mu(y_1, \dots, y_n) = \mu(c^{-1} \cdot A).
\end{aligned}$$

(viii): Let $A \subseteq \mathbb{R}^n$ be a Borel set. We have

$$\begin{aligned}
(\delta_u \odot \delta_v)(A) &= \int_{\mathbb{R}^n \times \mathbb{R}^n} \chi_A(x_1 y_1, \dots, x_n y_n) \, d\delta_u(x_1, \dots, x_n) \, d\delta_v(y_1, \dots, y_n) \\
&= \chi_A(u_1 v_1, \dots, u_n v_n) = \delta_{u \odot v}(A). \quad \square
\end{aligned}$$

As an application we look at the following. The equivalence (i) \Leftrightarrow (ii) is [BGKP22, Thm. 3.1]. [BGKP22, Thm. 9.1] is the multi-dimensional version. We only use [BGKP22, Thm. 3.1] (i) \Rightarrow (ii). Recall that an *absolutely monotonic entire function* has the form $f(x) = \sum_{k \in \mathbb{N}_0} a_k \cdot x^k$ for all $x \in \mathbb{R}$ with Taylor coefficients $a_k \geq 0$ for all $k \in \mathbb{N}_0$.

Corollary 5. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$. Then the following are equivalent:*

- (i) *f maps $[-1, 1]$ -moment sequences to $[-1, 1]$ -moment sequence.*
- (ii) *f is an absolutely monotonic entire function.*
- (iii) *f is entire and f maps \mathbb{R} -moment sequence to \mathbb{R} -moment sequences.*
- (iv) *f is entire and $(f(2^{-k}))_{k \in \mathbb{N}_0}$ is a moment sequence.*
- (v) *f maps $[-1, 1]^n$ -moment sequences to $[-1, 1]^n$ -moment sequences for a fixed $n \in \mathbb{N}$.*
- (vi) *f maps $[-1, 1]^n$ -moment sequences to $[-1, 1]^n$ -moment sequences for all $n \in \mathbb{N}$.*
- (vii) *f is entire and f maps \mathbb{R}^n -moment sequences to \mathbb{R}^n -moment sequences for a fixed $n \in \mathbb{N}$.*

(viii) f is entire and f maps \mathbb{R}^n -moment sequences to \mathbb{R}^n -moment sequences for all $n \in \mathbb{N}$.

Proof. (i) \Rightarrow (ii): That is [BGKP22, Thm. 3.1].

(ii) \Rightarrow (i): Let s be a $[-1, 1]$ -moment sequence with representing measure μ . Then $f(\odot s)$ is represented by $f(\odot \mu) = \sum_{k \in \mathbb{N}_0} a_k \cdot \mu^{\odot k}$ and

$$\text{supp } f(\odot \mu) = \bigcup_{k \in \mathbb{N}_0} \text{supp } \mu^{\odot k} \subseteq \bigcup_{k \in \mathbb{N}} \underbrace{[-1, 1] \cdot \dots \cdot [-1, 1]}_{k\text{-times, see (5)}} = [-1, 1] \quad (6)$$

by Theorem 2. That $f(\odot \mu)$ represents $f(\odot s)$, i.e., the limit $d \rightarrow \infty$ in Remark 3 exists, follows from (6) and

$$(f(\odot \mu))(\mathbb{R}) = \sum_{k \in \mathbb{N}_0} a_k \cdot (\mu^{\odot k})(\mathbb{R}) = \sum_{k \in \mathbb{N}_0} a_k \cdot (\mu(\mathbb{R}))^k = f(\mu(\mathbb{R})) < \infty. \quad (7)$$

(ii) \Rightarrow (iii): Eq. (7) does not require $\text{supp } \mu \subseteq [-1, 1]$ but holds for arbitrary $\text{supp } \mu \subseteq \mathbb{R}$ since f is entire.

(iii) \Rightarrow (iv): Clear since $(2^{-k})_{k \in \mathbb{N}_0}$ is a moment sequence with representing measure $\delta_{1/2}$.

(iv) \Rightarrow (ii): The sequence $(2^{-k})_{k \in \mathbb{N}_0}$ is a moment sequence with (unique) representing measure $\delta_{1/2}$. Hence, by Theorem 2 and Lemma 4 (viii) we have that $(f(2^{-k}))_{k \in \mathbb{N}_0}$ is represented by the (possibly signed) measure

$$f(\odot \delta_{2^{-1}}) = \sum_{k \in \mathbb{N}_0} a_k \cdot \delta_{2^{-k}}$$

with $\text{supp } f(\odot \delta_{2^{-1}}) \subseteq \{2^{-k} \mid k \in \mathbb{N}_0\} \subseteq [0, 1]$ and $|f(\odot \mu)|(\mathbb{R}) \leq \sum_{k \in \mathbb{N}_0} |a_k| \cdot (\mu(\mathbb{R}))^k < \infty$. Assume there exists a $k \in \mathbb{N}_0$ such that $a_k < 0$. Since $\mathbb{R}[x]$ is dense in $C([0, 1], \mathbb{R})$ there exists a $p \in \mathbb{R}[x]$ such that

$$p \geq 0 \text{ on } [0, 1], \quad p(2^{-k}) = 1, \quad \text{and} \quad p(2^{-m}) \leq \frac{|a_k|}{2 \sum_{l \in \mathbb{N}_0 \setminus \{k\}} |a_l|}$$

for all $m \in \mathbb{N}_0 \setminus \{k\}$. Then with the Riesz functional $L_{f(\odot s)} : \mathbb{R}[x] \rightarrow \mathbb{R}$ defined by $L_{f(\odot s)}(x^m) = f(s_m)$ for all $m \in \mathbb{N}_0$ we have

$$L_{f(\odot s)}(p) = \sum_{l \in \mathbb{N}_0} a_l \cdot p(2^{-l}) \leq a_k + \sum_{m \in \mathbb{N}_0 \setminus \{k\}} |a_m| \cdot p(2^{-m}) \leq a_k + \frac{|a_k|}{2} < 0,$$

i.e., $f(\odot s)$ is not a moment sequence. Hence, $a_k \geq 0$ holds for all $k \in \mathbb{N}_0$.

(ii) \Rightarrow (v), ..., (viii): Follows with the same arguments from (ii) \Rightarrow (i) for general $n \in \mathbb{N}$ from Theorem 2 and Lemma 4.

(v) or (vi) \Rightarrow (i): For the multi-moment sequence $s = (s_\alpha)_{\alpha \in \mathbb{N}_0^n}$ take the marginal moment sequence $s^{[1]} := (s_{(k, 0, \dots, 0)})_{k \in \mathbb{N}_0}$.

(vii) or (viii) \Rightarrow (iii): For the multi-moment sequence $s = (s_\alpha)_{\alpha \in \mathbb{N}_0^n}$ take the marginal moment sequence $s^{[1]} := (s_{(k, 0, \dots, 0)})_{k \in \mathbb{N}_0}$. \square

The Hankel matrix arguments in [BGKP22] were used to show that $f(\odot s)$ is a moment sequence. Theorem 2 provides a representing measure of $f(\odot s)$ and therefore removes the Hankel matrix arguments. It also simplifies [BRS⁺22].

Funding

The author and this project are supported by the Deutsche Forschungsgemeinschaft DFG with the grant DI-2780/2-1 and his research fellowship at the Zukunftskolleg of the University of Konstanz, funded as part of the Excellence Strategy of the German Federal and State Government.

References

- [Akh65] N. I. Akhiezer, *The classical moment problem and some related questions in analysis*, Oliver & Boyd, Edinburgh, 1965.
- [BGKP22] A. Belton, D. Guillot, A. Khare, and M. Putinar, *Moment-sequence transforms*, *J. Eur. Math. Soc.* **24** (2022), 3109–3160.
- [BGKP23] ———, *Totally positive kernels, Pólya frequency functions, and their transforms*, *J. D’Anal. Math.* **150** (2023), 83–158.
- [Bor11] J. Borcea, *Classification of linear operators preserving elliptic, positive and non-negative polynomials*, *J. reine angew. Math.* **650** (2011), 67–82.
- [BRS⁺22] G. Blekherman, F. Rincón, R. Sinn, C. Vinzant, and J. Yu, *Moments, sums of squares, and tropicalization*, arXiv:2203.06291v1.
- [KN77] M. G. Kreĭn and A. A. Nudel’man, *The Markow Moment Problem and Extremal Problems*, American Mathematical Society, Providence, Rhode Island, 1977, translation of the Russian original from 1973.
- [KT21] A. Khare and T. Tao, *On the sign patterns of entrywise positivity preservers in fixed dimension*, *Am. J. Math.* **143** (2021), 1863–1929.
- [Lau09] M. Laurent, *Sums of squares, moment matrices and optimization over polynomials*, *Emerging application of algebraic geometry*, IMA Vol. Math. Appl., vol. 149, Springer, New York, 2009, pp. 157–270.
- [Mar08] M. Marshall, *Positive Polynomials and Sums of Squares*, *Mathematical Surveys and Monographs*, no. 146, American Mathematical Society, Rhode Island, 2008.
- [Sch42] I. J. Schoenberg, *Positive definite functions on spheres*, *Duke Math. J.* **9** (1942), 96–108.
- [Sch17] K. Schmüdgen, *The Moment Problem*, Springer, New York, 2017.
- [Sti94] T. J. Stieltjes, *Recherches sur les fractions continues*, *Ann. Fac. Sci. Toulouse* **8** (1894), no. 4, J1–J122.