

# Bi-Lipschitz embedding metric triangles in the plane

Xinyuan Luo, Matthew Romney, Alexandria L. Tao

## Abstract

A *metric polygon* is a metric space comprised of a finite number of closed intervals joined cyclically. The second-named author and Ntalampikos recently found a method to bi-Lipschitz embed an arbitrary metric triangle in the Euclidean plane with uniformly bounded distortion, which we call here the *tripodal embedding*. In this paper, we prove the sharp distortion bound  $4\sqrt{7/3}$  for the tripodal embedding. We also give a detailed analysis of four representative examples of metric triangles: the intrinsic circle, the three-petal rose, tripods and the twisted heart. In particular, our examples show the sharpness of the tripodal embedding distortion bound and give a lower bound for the optimal distortion bound in general. Finally, we show the triangle embedding theorem does not generalize to metric quadrilaterals by giving a family of examples of metric quadrilaterals that are not bi-Lipschitz embeddable in the plane with uniform distortion.

## 1 Introduction

The *bi-Lipschitz embedding problem* asks one to given an intrinsic characterization of those metric spaces that can be embedded in some Euclidean space under a bi-Lipschitz map. Moreover, if a metric space does embed in some Euclidean space, one may then attempt to find or estimate the optimal target dimension and bi-Lipschitz distortion. The bi-Lipschitz embedding problem is wide open in full generality and has produced a rich body of research from a variety of mathematical points of view. We refer the reader to [5], [6], [7, Chapter 15], [8], [10] and [11] for general background.

In this article, we study a specific class of metric spaces, namely, *metric polygons*, and their embeddings in Euclidean spaces, with a focus on *metric triangles*. We say that a metric polygon is a metric space comprised of finitely many closed intervals (i.e., metric spaces each isometric to  $[0, a]$  for some  $a > 0$ ), called edges, joined in cyclic fashion. For two points lying on different edges, there is no constraint on the distance between them beyond what follows from the triangle inequality. A metric polygon with three edges is called a metric triangle, with four edges is called a metric quadrilateral, and so forth. See Definition 2.4 below for a formal definition.

To our knowledge, metric polygons first appear explicitly in the recent paper [9], in which it is shown that any metric triangle embeds in  $\mathbb{R}^2$  (equipped with the Euclidean metric) under a bi-Lipschitz map with distortion at most 12. The fact is then applied to the problem of finding good polyhedral approximations of metric surfaces under minimal geometric assumptions. We refer to the embedding found in [9] as the *tripodal embedding*; it will be reviewed in Section 3 below. Our goal is to explore the topic of embedding metric polygons more systematically and extend the embedding result from [9] in several directions.

As our main objective, we find the sharp bound on the distortion of the tripodal embedding in the general case.

**Theorem 1.1.** Every metric triangle embeds in  $\mathbb{R}^2$  under the tripodal embedding with distortion at most  $D = 4\sqrt{7/3} \approx 6.11$ .

In Section 4.4, we use an example that we call the *twisted heart* to show that the constant  $4\sqrt{7/3}$  cannot be improved. This leaves open the question of the best possible value for the distortion  $D$  with an arbitrary embedding. Let  $\text{lip}(\Delta)$  denote the minimum value for which any metric triangle embeds in  $\mathbb{R}^2$  with distortion at most  $\text{lip}(\Delta)$ . A simple example, called the *three-petal rose*, shows that  $\text{lip}(\Delta) \geq 2$ ; see Section 4.2. The example of the twisted heart suggests an even larger value for  $\text{lip}(\Delta)$ , perhaps roughly the value 4. However, we have not been able to find any general embedding method that improves upon the tripodal embedding.

---

2020 Mathematics Subject Classification. 51F30, 30L05.

M. Romney was partially supported by the National Science Foundation under grant DMS-2413156. The main results of this paper form a part of the Senior Research Project of X. Luo and A. L. Tao.

**Question 1.2.** What is the value of  $\text{lip}(\Delta)$ ?

In Section 4, we study the previously mentioned examples, along with two other representative examples: tripods and the intrinsic circle. Specifically, we consider the problem of finding distortion-minimizing bi-Lipschitz embeddings for these triangles. We can think of such a distortion-minimizing embedding as an optimal “Euclidean realization” of the original triangle. Note, however, that such an embedding is typically not unique; since the distortion of an embedding depends only on the worst pair of points in the space, one can typically perturb a given embedding elsewhere without affecting the distortion.

In the final main section this paper, Section 5, we consider the situation for arbitrary metric polygons. Note that it follows immediately from general properties of bi-Lipschitz embeddings (see Theorem 3.2 in [6]) that every metric  $n$ -gon bi-Lipschitz embeds in  $\mathbb{R}^{2n+1}$  with uniformly bounded distortion depending only on  $n$ . Thus our main question is, for a given  $n$ , to determine the smallest dimension for which one obtains an embedding of every  $n$ -gon with uniformly bounded distortion. We show that the analogue of Theorem 1.1 is false for metric quadrilaterals.

**Proposition 1.3.** There is a sequence of metric quadrilaterals  $X_n$  such that  $\text{lip}(X_n, \mathbb{R}^2)$  is unbounded as  $n \rightarrow \infty$ .

This proposition is related to the fact that a simple closed curve separates the plane. On the other hand, we expect that there is no obstruction to embedding arbitrary metric  $n$ -gons in  $\mathbb{R}^3$  with distortion bound depending only on  $n$ .

**Question 1.4.** For each  $n \geq 4$ , is there a value  $D_n$  such that every metric  $n$ -gon embeds in  $\mathbb{R}^3$  with distortion  $D_n$ ?

We also expect that every metric quadrilateral is individually bi-Lipschitz embeddable in  $\mathbb{R}^2$ , i.e., without a uniform bound on the bi-Lipschitz constant.

**Question 1.5.** Is every metric quadrilateral bi-Lipschitz embeddable in  $\mathbb{R}^2$ ?

On the other hand, the two fundamental examples of non-planar graphs— $K_5$ , the complete graph on five vertices, and  $K_{3,3}$ , the complete bipartite graph on two sets of three vertices—can each be realized as a metric pentagon. See Example 5.3 below.

In addition to our work, there have been other investigations into the metric geometry of classes of one-dimensional metric spaces. For example, bi-Lipschitz embeddings of so-called *quasiconformal trees* have also been studied in [3] and [4].

## Acknowledgement

We thank Guy C. David for feedback on a draft of our paper.

## 2 Background

### 2.1 Metric spaces and bi-Lipschitz maps

We review some of the definitions and concepts related to metric spaces. For a detailed account of the subject, see the monographs of Bridson–Haefliger [1] and Burago–Burago–Ivanov [2].

**Definition 2.1.** Given a set  $X$ , a *metric* (or *distance function*) on  $X$  is a function  $d: X \times X \rightarrow [0, \infty)$  satisfying the following properties for all  $x, y, z \in X$ :

1.  $d(x, y) = 0$  if and only if  $x = y$  (positive definiteness)
2.  $d(x, y) = d(y, x)$  (symmetry)
3.  $d(x, y) \leq d(x, z) + d(z, y)$  (triangle inequality)

A *metric space* is a pair  $(X, d)$ , where  $X$  is a set and  $d$  is a metric on  $X$ .

Perhaps the most familiar example of a metric space is  $n$ -dimensional Euclidean space, which is the set  $\mathbb{R}^n$  equipped with the Euclidean metric  $d_{\text{Euc}}(x, y) = \|x - y\| = \sqrt{(x_1 - y_1)^2 + \cdots + (x_n - y_n)^2}$ . For an arbitrary metric space  $(X, d)$ , a natural question is how different the metric  $d$  is from the Euclidean metric, quantitatively. To make this precise, we use the following definition.

**Definition 2.2.** For each  $L > 0$ , a map  $f: X \rightarrow Y$  between metric spaces  $(X, d_X)$  and  $(Y, d_Y)$  is *L-Lipschitz* if  $d_Y(f(x), f(y)) \leq Ld_X(x, y)$  for all  $x, y \in X$ . The map  $f$  is *L-bi-Lipschitz* if

$$L^{-1}d_X(x, y) \leq d_Y(f(x), f(y)) \leq Ld_X(x, y)$$

for all  $x, y \in X$ . The map  $f$  is *Lipschitz* (resp. *bi-Lipschitz*) if it is *L-Lipschitz* (resp. *L-bi-Lipschitz*) for some  $L > 0$ .

The *bi-Lipschitz embedding problem* then asks one to characterize when a given metric space can be mapped onto a subset of Euclidean space of some dimension under a bi-Lipschitz map. As discussed in the introduction, this is a difficult open problem in full generality. To refine this problem, we give the following definition.

**Definition 2.3.** Let  $f: X \rightarrow Y$  be a bi-Lipschitz map between metric spaces  $(X, d_X)$  and  $(Y, d_Y)$ . The *distortion* of  $f$  is  $\text{lip}(f) = L_0 \cdot L_1$ , where  $L_0, L_1$  are the minimum values such that  $L_0^{-1}d_X(x, y) \leq d_Y(f(x), f(y)) \leq L_1d_X(x, y)$ .

For metric spaces  $(X, d_X)$  and  $(Y, d_Y)$ , we define

$$\text{lip}(X, Y) = \inf \text{lip}(f),$$

where the infimum is taken over all bi-Lipschitz embeddings  $f: X \rightarrow Y$ . Note that  $\text{lip}(X, Y) = \infty$  if no such embedding exists.

A *curve* is a continuous function from an interval into a metric space. Let  $(X, d)$  denote this target metric space. A curve  $\Gamma: [a, b] \rightarrow X$  is *closed* if  $\Gamma(a) = \Gamma(b)$  and *simple* if it is injective (that is, it has no self-intersections) except that possibly  $\Gamma(a) = \Gamma(b)$ .

Given a curve  $\Gamma: [a, b] \rightarrow X$ , the *length* of  $\Gamma$  is

$$\ell(\Gamma) = \sup \sum_{j=1}^n d(\Gamma(t_{j-1}), \Gamma(t_j)),$$

where the supremum is taken over all finite partitions  $a = t_0 < t_1 < \dots < t_n = b$ . A curve  $\Gamma$  between two points  $x$  and  $y$  is a *geodesic* if  $\ell(\Gamma) = d(x, y)$ . A metric space  $(X, d)$  is a *length space* if

$$d(x, y) = \inf(\Gamma)$$

for all  $x, y \in X$ , where the infimum is taken over all curves  $\Gamma$  whose image contains  $x$  and  $y$ . If  $X$  is a compact length space, then by the Hopf–Rinow Theorem (see [1, Proposition I.3.7]) every two points are connected by at least one geodesic. Given two points  $x, y \in X$ , we write  $[xy]$  to denote some choice of geodesic between the two points.

The examples of metric polygons we consider in this paper are all either length spaces or subsets of length spaces. Topologically, they are graphs, so a length metric can be defined simply by specifying the length of each edge. See [2, Chapter 2] for a more detailed overview of length spaces.

Finally, the *concatenation* of two curves  $\Gamma_1: [a, b] \rightarrow X$ ,  $\Gamma_2: [c, d] \rightarrow X$  satisfying  $\Gamma_1(b) = \Gamma_2(c)$  is denoted by  $\Gamma_1 * \Gamma_2$ . This is the curve from  $[a, b + d - c]$  to  $X$  defined by the formula  $\Gamma_1(t)$  if  $t \leq b$  and  $\Gamma_2(t - b + c)$  if  $t \geq b$ .

## 2.2 Metric polygons

Our interest in this paper is in a specific class of metric spaces called metric polygons.

**Definition 2.4.** Let  $n \geq 2$ . A *metric n-gon* is a metric space that is the union of  $n$  subspaces  $I_1, \dots, I_n$  that are each isometric to a closed interval  $[0, \ell_n]$  and connected at the endpoints cyclically. More precisely, if we let  $\varphi_j: [0, \ell_j] \rightarrow I_j$  denote the required isometry onto the interval  $I_j$ , then it holds that  $\varphi_j(\ell_j) = \varphi_{j+1}(0)$  for all  $j \in \{1, \dots, n\}$ , with the identification  $n + 1 = 1$ . A metric 3-gon is called a *metric triangle*, and a metric 4-gon is called a *metric quadrilateral*, and so forth. More generally, a *metric polygon* is a metric  $n$ -gon for some  $n$ .

Each subspace  $I_j$  is called an *edge* of the metric polygon. Each endpoint of one of the edges  $I_j$  is called a *vertex* of the metric polygon.

Note that this definition allows the sets  $I_j$  to intersect at points other than the endpoints, although the nature of such additional intersection points is constrained by the triangle inequality. Also, note that the terms “edge” and “vertex” here are distinct from the topological meaning of these terms for a graph. For example, graphs (such as  $K_5$  or  $K_{3,3}$ , as discussed in Example 5.3) can be represented as metric polygons. In these cases, the topological edges and vertices of the graph need not coincide with its edges and vertices as a metric polygon.

Next, we introduce some main examples of metric triangles that will be studied further in Section 4. These are illustrated in Figure 1. In each of these examples, the space is a length space or subset of a length space.

**Example 2.5.** (a) **The intrinsic circle.** Let  $S$  be the unit circle in  $\mathbb{R}^2$ , which we give the arc length or angular distance. More precisely, each point in  $S$  can be represented by some  $\theta \in [0, 2\pi)$ , with the correspondence  $\theta \mapsto (\cos(\theta), \sin(\theta))$ . Then  $d(\theta, \tau) = \min\{|\theta - \tau|, |\theta - \tau - 2\pi|, |\theta - \tau + 2\pi|\}$ .

To view this space as a metric triangle, we pick any three points that do not all belong to the same semicircle to be vertices. For example, we can take  $\theta_1 = 0$ ,  $\theta_2 = 2\pi/3$  and  $\theta_3 = 4\pi/3$ .

(b) **The three-petal rose.** Let  $S_1, S_2, S_3$  be three copies of the intrinsic circle as just defined. Let  $R$  be the wedge sum of  $S_1, S_2, S_3$  with basepoint 0 in each  $S_i$  (that is, the quotient space of the disjoint union of  $S_1, S_2, S_3$  given by identifying the point 0 in each  $S_i$ ). We define  $d$  to be the length metric on  $R$ . More precisely,  $d$  is defined as follows. If  $\theta, \tau \in S_i$  for the same  $i$ , then  $d(\theta, \tau)$  agrees with the metric in (a). If  $\theta \in S_i$  and  $\tau \in S_j$  for  $i \neq j$ , then  $d(\theta, \tau) = d(\theta, 0) + d(0, \tau)$ .

To view this space as a metric triangle, we take the antipodal point on each circle as vertices. That is, let  $p = \pi \in S_1$ ,  $q = \pi \in S_2$  and  $r = \pi \in S_3$ . The points 0 and  $p$  divide  $S_1$  into two semicircles  $S_1^1, S_1^2$ . Likewise,  $S_2$  and  $S_3$  are divided into semicircles  $S_2^1, S_2^2$  and  $S_3^1, S_3^2$ , respectively. Then the edges of  $R$  are  $[pq] = S_1^2 \cup S_2^1$ ,  $[qr] = S_2^2 \cup S_3^1$  and  $[rp] = S_3^2 \cup S_1^1$ .

(c) **Tripods.** Choose values  $l_i \geq 0$ ,  $i = 1, 2, 3$ , and consider the three intervals  $T_i = [0, l_i]$ ,  $i = 1, 2, 3$ . We define  $T$  to be the wedge sum of  $T_1, T_2, T_3$  with basepoint 0 in each  $T_i$ . We define  $d$  to be the length metric on  $T$ . That is, if  $x, y \in T_i$  for the same  $i$ , then  $d(x, y) = |x - y|$ . If  $x \in T_i$  and  $y \in T_j$  for  $i \neq j$ , then  $d(x, y) = d(x, 0) + d(0, y) = x + y$ . Such a metric space is called a *tripod*, or a *metric tripod* for emphasis. We call the point 0 the *center* of the tripod.

Let  $p = l_1 \in T_1$ ,  $q = l_2 \in T_2$ ,  $r = l_3 \in T_3$ . We can view  $T$  as a metric triangle by taking the points  $p, q, r$  as vertices and  $[pq] = T_1 \cup T_2$ ,  $[qr] = T_2 \cup T_3$ ,  $[rp] = T_3 \cup T_1$  as edges.

(d) **The twisted heart.** The example is less obvious but turns out to be the extremal example for the tripod embedding. We start with the graph consisting of fourteen edges of unit length as shown in Figure 1d. The distance between two points is the length of the shortest curve between them. We obtain a triangle by taking  $p, q, r$  as vertices and removing the edges from  $a$  to  $c$  and from  $b$  to  $e$  (the red edges in Figure 1d) without changing the metric on the remaining points. Observe that each pair of vertices is at distance 4 apart. Moreover,  $d(a, b) = 4$  while  $d(a, c) = d(b, e) = 1$ . We denote this example by  $H$ .

Our final definition, used in the proof of Theorem 1.1, is the Gromov product, which measures the deficit in the triangle inequality for a triple of points.

**Definition 2.6.** Let  $X$  be a metric space and  $p, q, r \in X$ . The *Gromov product* of  $p$  and  $q$  at  $r$  is

$$(p \cdot q)_r = \frac{1}{2}(d(p, r) + d(q, r) - d(p, q)).$$

The usefulness of the Gromov product for our purpose is that it allows us to associate to any metric triangle a tripod with matching edge lengths. This idea forms the basis of our proof that any metric triangle is bi-Lipschitz embeddable in  $\mathbb{R}^2$ . The following simple lemma was proved as Lemma 3.1 in [9].

**Lemma 2.7.** Let  $\Delta$  be a metric triangle with vertices  $p, q, r$ . Let  $T$  be the tripod formed as in Example 2.5 by taking  $l_1 = (q \cdot r)_p$ ,  $l_2 = (p \cdot r)_q$  and  $l_3 = (p \cdot q)_r$ . Define a map  $P: \Delta \rightarrow T$  by setting  $P(p) = l_1 \in T_1$ ,  $P(q) = l_2 \in T_2$ ,  $P(r) = l_3 \in T_3$ , and mapping each edge of  $\Delta$  by arc length onto the corresponding edge of  $T$ . Then  $P$  is 1-Lipschitz.

Each tripod  $T$  can in turn be embedded into the plane in a natural way. For a given tripod  $T$  with vertices  $p, q, r$  and center  $o$ , let  $r_1 = d(o, p)$ ,  $r_2 = d(o, q)$ ,  $r_3 = d(o, r)$ . Let  $u_1 = (r_1, 0)$ ,  $u_2 = r_2(-\frac{1}{2}, \frac{\sqrt{3}}{2})$ ,




Figure 1: Examples of metric polygons.

and  $u_3 = r_3(-\frac{1}{2}, -\frac{\sqrt{3}}{2})$ . For each  $i = 1, 2, 3$ , let  $J_i$  be the union of the straight line segment from  $(0, 0)$  to  $u_i$ . Then let  $Y = J_1 \cup J_2 \cup J_3$ , equipped with the Euclidean metric from  $\mathbb{R}^2$ . We define a map  $\varphi: T \rightarrow Y$  by mapping the segment  $[op]$  onto  $J_1$  by arc length,  $[oq]$  onto  $J_2$  by arc length and  $[or]$  onto  $J_3$  by arc length.

### 3 Proof of optimal distortion bound

In this section, we give the proof of our main result, Theorem 1.1, giving the optimal distortion bound for the tripodal embedding of an arbitrary metric triangle.

We start by reviewing the definition of the tripodal embedding as given in [9]. Let  $\Delta$  be an arbitrary metric triangle with edges  $I_1, I_2, I_3$ . Denote the metric on  $\Delta$  by  $d$ . Let  $p$  be the common vertex of  $I_1$  and  $I_2$ ,  $q$  be the common vertex of  $I_2$  and  $I_3$ , and  $r$  be the common vertex of  $I_3$  and  $I_1$ . For each edge  $I_j$ , let  $\widehat{I}_j$  denote the union of the other two edges of  $\Delta$ . We can associate to the triangle  $\Delta$  a metric tripod  $T$  with matching edge lengths and corresponding map  $P: \Delta \rightarrow T$  as in Lemma 2.7. To this  $T$  we associate an embedded tripod  $Y$  and map  $\varphi: T \rightarrow Y$  following the notation of the previous section. For conciseness, we write  $\bar{x}$  in place of  $(\varphi \circ P)(x)$ . Let  $v_1 = (\frac{1}{2}, \frac{\sqrt{3}}{2})$ ,  $v_2 = (-1, 0)$  and  $v_3 = (\frac{1}{2}, -\frac{\sqrt{3}}{2})$ . Then  $F: \Delta \rightarrow \mathbb{R}^2$  is defined by the formula

$$F(x) = \bar{x} + \text{dist}(x, \widehat{I}_j)v_j \text{ if } x \in I_j.$$

Here, given a point  $x \in X$  and a set  $A \subset X$ , we write  $\text{dist}(x, A) = \inf_{a \in A} d(x, a)$ .

We fix some additional notation. Let  $I_1^p$  denote the subarc of  $I_1$  consisting of those points  $x \in I_1$  for which  $d(x, p) \leq (q \cdot r)_p$ . Define  $I_1^q, I_2^q, I_2^r, I_3^p$  and  $I_3^r$  similarly. For each edge  $I_j$ , let  $o_j$  denote the unique point such that  $\bar{o}_j$  is the origin. Moreover, we let  $z \in \Delta$  denote a point satisfying  $d(x, z) = \text{dist}(x, \widehat{I}_1)$ . This point may fail to be uniquely determined, so we fix a choice of  $z$ . Similarly, we let  $w \in \Delta$  denote a point satisfying  $d(y, w) = \text{dist}(y, \widehat{I}_1)$ , where  $j$  is such that  $y \in I_j$ .

In this proof, we consider two arbitrary distinct points  $x, y \in \Delta$  and show that  $\|F(x) - F(y)\|$  satisfies appropriate bounds. By symmetry, we may consider five individual cases in our proof of Theorem 1.1: (1)  $x, y \in I_1^p$ ; (2)  $x \in I_1^p, y \in I_1^q$ ; (3)  $x \in I_1^p, y \in I_3^p$ ; (4)  $x \in I_1^p, y \in I_3^r$ ; (5)  $x \in I_1^q, y \in I_3^r$ . See Figure 2. Each of these cases is represented by one of the lemmas in this section.

**Lemma 3.1.** Suppose that  $x, y \in I_1^p$ . Then

$$\frac{\sqrt{3}}{2}d(x, y) \leq \|F(x) - F(y)\| \leq \sqrt{3}d(x, y).$$

*Proof.* Assume that  $d(x, p) < d(y, p)$ . We first verify the lower bound. Note that both  $\bar{x}$  and  $\bar{y}$  lie on the positive real axis, with  $\|\bar{x} - \bar{y}\| = d(x, y)$ . Moreover,  $F(x)$  and  $F(y)$  lie on the lines  $\{\bar{x} + tv_1 : t \in \mathbb{R}\}$




Figure 2: The five cases in the proof of Theorem 1.1.

and  $\{\bar{y} + tv_1 : t \in \mathbb{R}\}$ , respectively. The distance between these lines is  $\frac{\sqrt{3}}{2}d(x, y)$ . See Figure 3a. This verifies the lower bound.

Next, we show the upper bound. Observe that  $|d(x, \hat{I}_1) - d(y, \hat{I}_2)| \leq d(x, y)$ . Write  $F(y)$  as  $F(y) = \bar{y} + t(y)v_1$  for some  $t(y) \geq 0$ . Then  $F(x)$  belongs to the set  $\bar{x} + tv_j$ , where  $t \in [t(y) - d(x, y), t(y) + d(x, y)]$ . The point in this set furthest from  $F(y)$  is  $\bar{x} + (t(y) + d(x, y))v_j$ , which is at distance  $\sqrt{3}d(x, y)$  from  $F(y)$ . See Figure 3b. This verifies the upper bound.  $\square$

**Lemma 3.2.** Suppose that  $x \in I_1^p$  and  $y \in I_1^q$ . Then

$$\frac{\sqrt{3}}{2}d(x, y) \leq \|F(x) - F(y)\| \leq \sqrt{3}d(x, y).$$

*Proof.* Assume without loss of generality that  $d(o_1, y) \leq d(o_1, x)$ . We first verify the lower bound. As in the previous case, observe that  $F(x)$  and  $F(y)$  lie on the lines  $\{\bar{x} + tv_1 : t \in \mathbb{R}\}$  and  $\{\bar{y} + tv_1 : t \in \mathbb{R}\}$ , respectively. The distance between these lines is  $\frac{\sqrt{3}}{2}d(o_1, x) + \frac{\sqrt{3}}{2}d(o_1, y)$ . See Figure 4a. Since  $d(x, y) = d(x, o_1) + d(o_1, y)$ , we obtain the lower bound.

We next verify the upper bound. Write  $F(y)$  as  $F(y) = \bar{y} + t(y)v_1$  for some  $t(y) \geq 0$ . As in the previous case,  $F(x)$  belongs to the set  $\bar{x} + tv_j$ , where  $t \in [t(y) - d(x, y), t(y) + d(x, y)]$ . Since  $d(o_1, y) \leq d(o_1, x)$ , the point in this set furthest from  $F(y)$  is  $\bar{x} + (t(y) + d(x, y))v_j$ . Denote the latter point by  $\tilde{x}$ . Let  $\tilde{y} = -|\bar{y}|u_1 + t(y)v_1$ , that is, the point along the negative horizontal axis at the same distance from the origin as  $\bar{y}$ . The point  $\tilde{y}$  belongs to the line  $\{\bar{y} + tv_1 : t \in \mathbb{R}\}$  but is further away from  $\tilde{x}$  than  $F(y)$ . Thus  $\|F(x) - F(y)\| \leq \|\tilde{x} - \tilde{y}\|$ . But  $\|\tilde{x} - \tilde{y}\| \leq \sqrt{3}d(x, y)$  by the same analysis as in Lemma 3.1. This establishes the upper bound.  $\square$

In the remaining lemmas, it is more convenient to phrase the inequalities in terms of the square of the distortion rather than the distortion itself. We introduce some more notation. Note that  $x \in I_1$  and  $y \in I_3$  in all the remaining cases. We let  $a = d(x, z)$ ,  $b = d(y, w)$ ,  $s = d(x, o_1)$  and  $t = d(y, o_3)$ . Recall that  $z$  be a point in  $\hat{I}_1$  such that  $d(x, z) = d(x, \hat{I}_1)$ , and  $w$  a point in  $\hat{I}_3$  such that  $d(y, w) = d(y, \hat{I}_3)$ .

**Lemma 3.3.** Suppose that  $x \in I_1^p$  and  $y \in I_3^p$ . Then

$$\frac{3}{16} \leq \frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq 4.$$

*Proof.* Observe that  $\bar{x}$  and  $\bar{y}$  belong to the same leg of the embedded tripod  $Y$ . First we show the upper bound. Note that  $d(x, y) \geq |t - s|$ , since by Lemma 2.7 the map  $\varphi \circ P$  is 1-Lipschitz. It follows that  $d(x, y) \geq \max\{a, b, |t - s|\}$ . Let  $e = |t - s|$ . A computation shows that

$$\|F(x) - F(y)\|^2 = \frac{1}{4}(a - b + 2s - 2t)^2 + \frac{3}{4}(a + b)^2.$$




Figure 3

By rescaling and symmetry, we assume now that  $a = 1$  and  $b \leq 1$ . If  $|e| \leq 1$ , then  $d(x, y) \geq \max\{a, b\} = 1$  and hence

$$\frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq M_1(b, e)$$

for

$$M_1(b, e) = \frac{1}{4}(1 - b + 2e)^2 + \frac{3}{4}(1 + b)^2.$$

The function  $M_1(b, e)$  is maximized on the set  $\{(b, e) : 0 \leq b, e \leq 1\}$  by taking  $b = e = 1$ , which gives a value of 4. Since this statement is plausible enough (for example, by plotting the function), and checking it is unrelated to the rest of the proof, we defer the verification of this claim to the Appendix. See Lemma 6.1 for the details. We do the same with similar inequalities arising in this section.

If  $|e| \geq 1$ , then

$$\frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq \frac{1}{e^2} M_1(b, e).$$

On the set  $\{(b, e) : 0 \leq b \leq 1, 1 \leq e < \infty\}$ , the function  $e^{-2} M_1(b, e)$  is maximized by again taking  $b = e = 1$ , which gives a value of 4. See Lemma 6.2. This establishes the upper bound.

Now we show the lower bound. At this point, we separate into two cases.

*Case 1.* Assume first that  $z \in I_3$  or  $w \in I_1$ . By symmetry, we may assume that  $z \in I_3$ . If  $d(p, z) \leq d(p, y)$ , the triangle inequality implies that

$$d(p, o_1) \leq d(p, z) + a + s = d(p, o_3) - t - d(y, z) + a + s.$$

Since  $d(p, o_1) = d(p, o_3)$ , this rearranges to  $d(y, z) \leq a + s - t$ , from which we conclude that

$$d(x, y) \leq d(y, z) + d(z, x) \leq 2a + s - t.$$

Assume without loss of generality that  $a = 1$ . Set  $e = s - t$ . Then  $d(x, y)^2 / \|F(x) - F(y)\|^2$  is bounded by the function

$$M_2(b, e) = \frac{(2 + e)^2}{\frac{1}{4}(1 - b + 2e)^2 + \frac{3}{4}(1 + b)^2}.$$

In Lemma 6.3, we show that  $M_2(b, e)$  is bounded on the set  $\{(b, e) : b \geq 0\}$  by 4.

On the other hand, if  $d(p, y) \leq d(p, z)$ , the triangle inequality implies that

$$d(p, o_3) \leq d(o_3, z) + a + d(x, p) = t - d(z, y) + a + d(p, o_1) - s,$$

which gives  $d(z, y) \leq a + t - s$  and hence

$$d(x, y) \leq d(y, z) + d(z, x) \leq 2a + t - s.$$

In this case,  $d(x, y)^2 / \|F(x) - F(y)\|^2$  is bounded by the function

$$\widetilde{M}_2(b, e) = \frac{(2 - e)^2}{\frac{1}{4}(1 - b + 2e)^2 + \frac{3}{4}(1 + b)^2}.$$




Figure 4

If  $s \geq t$ , then  $e \geq 0$  and we have the relation  $\widetilde{M}_2(b, e) \leq M_2(b, e)$ . If  $s < t$ , we claim that one of two possibilities necessarily holds: either  $b \geq t - s$ , or  $w \in I_1$  with  $d(p, w) \leq d(p, x)$ . If  $w \in I_2$ , then necessarily  $b \geq t$  and hence the first alternative holds. If  $w \in I_1$  with  $d(p, w) > d(p, x)$ , then the triangle inequality gives

$$d(p, o_1) \leq d(p, y) + d(y, w) + d(w, o_1) = d(p, o_3) - t + b + d(w, o_1),$$

and hence  $t - d(w, o_1) \leq b$ . Since  $d(w, o_1) < s$ , this establishes the claim.

By Lemma 6.4, the function  $\widetilde{M}_2(b, e)$  is bounded on the set  $\{(b, e) : b \geq 0, -b \leq e \leq 0\}$  by  $13/3$ . This covers the case where  $b \geq t - s$ .

Assume next that  $w \in I_1$  with  $d(p, w) \leq d(p, x)$ . Since  $w \in I_1$ , we can follow the previous argument with roles of  $x$  and  $y$  reversed. Since  $d(p, w) \leq d(p, x)$ , we get the inequality  $d(x, y) \leq 2b + t - s$  analogously to the first alternative above, which is handled by Lemma 6.3.

*Case 2.* As the second case, assume now that both  $w, z \in I_2$ . Let  $d_1 = d(r, w) - d(r, o_2)$  and let  $d_2 = d(q, z) - d(q, o_2)$ . Note that  $d_1$  and  $d_2$  may be negative. The triangle inequality yields the inequalities

$$d(p, q) \leq d(p, y) + d(y, w) + d(w, q) = d(p, o_3) - t + b + d(o_2, q) - d_1$$

and

$$d(p, r) \leq d(p, y) + d(y, w) + d(w, r) = d(p, o_3) - t + b + d(o_2, r) + d_1.$$

If  $d_1 \geq 0$ , we use the first inequality and the relation  $d(p, q) = d(p, o_3) + d(o_2, q)$  to conclude that  $d_1 \leq b - t$ . If  $d_1 \leq 0$ , we use the second inequality and the relation  $d(p, r) = d(p, o_3) + d(o_2, r)$  to conclude that  $-d_1 \leq b - t$ . Together, these show that  $|d_1| \leq b - t$ . Similarly, we have  $|d_2| \leq a - s$ .

This gives

$$d(z, w) \leq |d_1| + |d_2| \leq a + b - s - t$$

and hence

$$d(x, y) \leq d(x, z) + d(z, w) + d(w, y) \leq a + b + |d_1| + |d_2| \leq 2a + 2b - s - t.$$

We compute

$$\frac{d(x, y)^2}{\|F(x) - F(y)\|^2} \leq \frac{(2a + 2b - s - t)^2}{\frac{1}{4}(a - b + 2s - 2t)^2 + \frac{3}{4}(a + b)^2}.$$

Assume without loss of generality that  $a = 1$  and  $b \leq 1$ . Then  $d(x, y)^2 / \|F(x) - F(y)\|^2$  is bounded by the function

$$M_3(b, s, t) = \frac{(2 + 2b - s - t)^2}{\frac{1}{4}(1 - b + 2s - 2t)^2 + \frac{3}{4}(1 + b)^2}.$$

Note as well that  $s \leq a = 1$  and  $t \leq b \leq 1$ . Thus we must maximize the function  $M_3(b, s, t)$  on the set  $0 \leq b, s, t \leq 1$ . We show in Lemma 6.5 that the maximum value is  $16/3$ , which is attained when  $b = 1$  and  $s = t = 0$ .  $\square$

The remaining two cases are handled similarly to Lemma 3.3, and so our proofs omit some details.



**Lemma 3.4.** Suppose that  $x \in I_1^p$  and  $y \in I_3^r$ . Then

$$\frac{3}{16} \leq \frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq 7.$$

*Proof.* We begin with the upper bound. Observe as before that  $d(x, y) \geq \max\{a, b, s + t\}$ . On the other hand, a computation shows that

$$\|F(x) - F(y)\|^2 = \frac{1}{4}(a - b + 2s + t)^2 + \frac{3}{4}(a + t + b)^2.$$

Assume now without loss of generality that  $a = 1$ . If  $\max\{a, b, s + t\} = a = 1$ , then  $b \leq 1$  and  $s + t \leq 1$  and so

$$\frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq M_4(b, s, t),$$

where

$$M_4(b, s, t) = \frac{1}{4}(1 - b + 2s + t)^2 + \frac{3}{4}(1 + t + b)^2.$$

Over the region  $\{(b, s, t) : 0 \leq b, s, t \leq 1, s + t \leq 1\}$ , the expression on the right of the previous inequality is maximized by taking  $s = 0, b = t = 1$  by Lemma 6.6. This gives the desired upper bound  $1/4 + 3/4 \cdot 3^2 = 7$ .

If  $\max\{a, b, s + t\} = s + t$ , then

$$\frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq \frac{M_4(b, s, t)}{(s + t)^2}.$$

Over the region  $\{(b, s, t) : 0 \leq s, t \leq 1, s + t \geq 1, 0 \leq b \leq s + t\}$ , the expression on the right of the previous inequality is again maximized by taking  $s = 0, b = t = 1$  by Lemma 6.7.

If  $\max\{a, b, s + t\} = b$ , then

$$\frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq \frac{M_4(b, s, t)}{b^2}.$$

Over the region  $\{(b, s, t) : 0 \leq s, t \leq 1, b \geq 1, s + t \leq b\}$ , the expression on the right of the previous inequality is again maximized by taking  $s = 0, b = t = 1$  by Lemma 6.8.

Next, we show the lower bound. We separate into three cases.

*Case 1.* Assume first that  $z \in I_3$ . We claim that  $d(x, y) \leq 2a + s + t$ . If  $d(z, p) \leq d(y, p)$ , then we bound  $d(x, y)$  as follows. By the triangle inequality,

$$d(o_1, p) = s + d(x, p) \leq s + a + d(z, p).$$

Since  $d(z, p) = d(o_3, p) - d(o_3, z) = d(o_1, p) - d(o_3, z)$ , this gives

$$d(o_3, z) \leq a + s.$$

Thus  $d(x, y) \leq t + d(o_3, z) + a \leq 2a + s + t$ . If  $d(z, p) > d(y, p)$ , we obtain the same bound as follows. By the triangle inequality,

$$d(r, p) = d(r, z) + d(z, y) + t + d(o_3, p) \leq d(r, z) + a + d(x, p).$$

Since  $d(o_3, p) = d(o_1, p) = d(x, p) + s$ , the previous inequality implies that  $d(z, y) \leq a - s - t$  and hence  $d(x, y) \leq 2a - s - t \leq 2a + s + t$ . It follows that

$$\frac{d(x, y)^2}{\|F(x) - F(y)\|^2} \leq \frac{(2a + s + t)^2}{\frac{1}{4}(a - b + 2s + t)^2 + \frac{3}{4}(a + t + b)^2}.$$

A bound on the right-hand side quantity will be found together with the next case.

*Case 2.* Assume in the second case that  $w \in I_1$ . We claim now that  $d(x, y) \leq 2b + s + t$ . If  $d(w, q) \leq d(o_1, q)$ , then we have

$$d(q, r) \leq d(y, q) + b + d(w, r) = d(r, o_3) - t + b + d(q, o_1) - d(w, o_1).$$

This implies that  $t + d(w, o_1) \leq b$ . Then

$$d(x, y) \leq b + d(w, o_1) + s \leq 2b + s - t.$$

Next, if  $w$  lies between  $o_1$  and  $x$ , then  $d(x, y) \leq b + s \leq 2b + s - t$ , since  $b \geq t$ . If  $w$  lies between  $x$  and  $p$ , then

$$d(p, r) \leq d(p, o_3) - t + b + d(w, p) \leq d(p, o_3) - t + b + d(o_1, p) - s - d(x, w),$$

which implies  $d(x, y) \leq d(x, w) + d(w, y) \leq 2b - s - t \leq 2b - s + t$ .

It follows in each case that  $d(x, y) \leq 2b + s + t$  and hence that

$$\frac{d(x, y)^2}{\|F(x) - F(y)\|^2} \leq \frac{(2b + s + t)^2}{\frac{1}{4}(a - b + 2s + t)^2 + \frac{3}{4}(a + t + b)^2}.$$

Without loss of generality, assume that  $a = 1$ . Note as well that  $s \leq a = 1$ . In both of the first two cases,  $d(x, y)^2/\|F(x) - F(y)\|^2$  is bounded by

$$M_5(b, s, t) = \frac{(\max\{2, 2s\} + s + t)^2}{\frac{1}{4}(1 - b + 2s + t)^2 + \frac{3}{4}(1 + t + b)^2}.$$

By Lemma 6.9, the function  $M_5(b, s, t)$  is bounded by 4 on the set  $\{(b, s, t) : 0 \leq b, s, t, 1 \geq s\}$ .

*Case 3.* In the final case, we assume that  $z \in I_2$  and  $w \in I_2$ . Proceeding similarly to Case 2 of Lemma 3.3, we obtain the relation  $d(x, y) \leq 2a + 2b - s + t$ . Assume without loss of generality that  $a = 1$ . Then  $d(x, y)^2/\|F(x) - F(y)\|^2 \leq M_6(b, s, t)$ , where

$$M_6(b, s, t) = \frac{(2 + 2b + 2t)^2}{\frac{1}{4}(1 - b + 2s + t)^2 + \frac{3}{4}(1 + t + b)^2}.$$

Note that the numerator of this formula uses the weaker bound  $2 + 2b + 2t$  instead of  $2 + 2b - s + t$ , but this turns out to simplify the calculations. By Lemma 6.10, this is bounded on the set  $\{(b, s, t) : 0 \leq b, s, t\}$  by  $16/3$ .  $\square$

**Lemma 3.5.** Suppose that  $x \in I_1^p$  and  $y \in I_3^r$ . Then

$$\frac{3}{16} \leq \frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq 7.$$

*Proof.* We prove the upper bound. This is similar to the proof of Lemma 3.4. As before, we have the lower bound  $d(x, y) \geq \max\{a, b, s + t\}$ . However, we have the new computation

$$\|F(x) - F(y)\|^2 = \frac{1}{4}(a - b - s + t)^2 + \frac{3}{4}(a + s + t + b)^2.$$

Assume without loss of generality that  $a = 1$  and  $b \leq 1$ . If  $s + t \leq 1$ , then

$$\frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq M_7(b, s, t),$$

where

$$M_7(b, s, t) = \frac{1}{4}(1 - b - s + t)^2 + \frac{3}{4}(1 + s + t + b)^2.$$

Over the region  $\{(b, s, t) : 0 \leq b, s, t \leq 1, s + t \leq 1\}$ ,  $M_7(b, s, t)$  is maximized by taking  $s = 0, t = 1, b = 1$  or  $s = 1, t = 0, b = 1$ . By Lemma 6.11, this gives the value  $1/4 + 3/4 \cdot 3^2 = 7$ , which gives the desired bound.

If  $s + t \geq 1$ , then

$$\frac{\|F(x) - F(y)\|^2}{d(x, y)^2} \leq \frac{M_7(b, s, t)}{(s + t)^2}.$$

Over the region  $\{(b, s, t) : 0 \leq b, s, t, 1 \leq s + t\}$ , the function  $M_7(b, s, t)/(s + t)^2$  is maximized by taking  $s = 0, t = 1, b = 1$  or  $s = 1, t = 0, b = 1$ , which by Lemma 6.12 again gives the desired bound.

Next, we show the lower bound. Assume first that  $z \in I_3$ . Then we can show similarly to in Lemma 3.4 that  $d(y, z) \leq a + t - s$ . Actually, for the remainder of the proof it is slightly more convenient




Figure 5: The image of  $F_{s_0}$ , which embeds the intrinsic circle in  $\mathbb{R}^2$

to use the weaker bound  $d(y, z) \leq a + t + s$ . From this we conclude that  $d(x, y) \leq 2a + s + t$ . Assume without loss of generality that  $a = 1$ . Then  $d(x, y)^2 / \|F(x) - F(y)\|^2$  is bounded by the function

$$M_8(b, s, t) = \frac{(2 + s + t)^2}{\frac{1}{4}(1 - b - s + t)^2 + \frac{3}{4}(1 + s + t + b)^2},$$

where  $b, s, t \geq 0$ . By Lemma 6.13, this function is bounded on the given region by 4.

Next, the case that  $w \in I_1$  is symmetric to the case that  $z \in I_3$ , and so we obtain the same bound.

Finally, assume that both  $z$  and  $w$  are in  $I_2$ . We can show similarly to Case 2 in Lemma 3.3 that  $d(w, z) \leq a + b + s + t$ . From this we conclude that  $d(x, y) \leq 2a + 2b + s + t$ . Assume without loss of generality that  $a = 1$ . Then  $d(x, y)^2 / \|F(x) - F(y)\|^2$  is bounded by the function

$$M_9(b, s, t) = \frac{(2 + 2b + 2s + 2t)^2}{\frac{1}{4}(1 - b - s + t)^2 + \frac{3}{4}(1 + s + t + b)^2},$$

where  $b, s, t \geq 0$ . As in Case 3 of Lemma 3.4, we use a weaker bound in the numerator in order to simplify the calculations. By Lemma 6.14, this function is bounded above by  $16/3$ .  $\square$

## 4 Examples

We give a detailed analysis of the examples introduced in Example 2.5: the intrinsic circle, the three-petal rose, tripods and the twisted heart. These are denoted respectively by  $S$ ,  $R$ ,  $T$  and  $H$ .

### 4.1 The intrinsic circle

This example represents the ‘‘fattest’’ possible metric triangle.

The obvious embedding from  $S$  into  $\mathbb{R}^2$  is the identity map, which has distortion  $\pi/2 \approx 1.5708$ . Interestingly, the identity map is not distortion minimizing. To improve this, for each  $s \in [0, \pi/3)$ , we define the map  $F_s: S \rightarrow \mathbb{R}^2$  by the formula

$$F_s(\theta) = \left( \frac{3\pi \cos(\tau(\theta)) - \sqrt{3}\pi \sin(s)}{3\pi - 9s}, \frac{\pi(-\sin(s) + \sin(\tau(\theta)))}{\pi - 3s} \right)$$

for  $\theta \in [0, 2\pi/3)$ . Here,  $\tau(\theta) = s + (1 - 3s/\pi)\theta$ . We extend the definition to all  $\theta \in [0, 2\pi)$  by symmetry. Namely, let  $A: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be the linear map representing rotation by  $\pi/3$ . For  $\theta \in [2\pi/3, 2\pi)$ , we define  $F_s(\theta) = A \circ F_s(\theta - 2\pi/3)$ . The map  $F_s$  takes  $S$  to a ‘‘fattened triangle’’ as shown in Figure 5 (with the value  $s_0$  found below). Note that  $F_0$  is the identity map and that the limit of  $F_s$  as  $s \rightarrow \pi/3$  maps  $S$  onto an equilateral triangle. The intuition behind the family  $\{F_s\}$  is that, beginning from the identity map, one can lower the maximal distortion by adding corners, which increases the distortion on small scales (where the identity map is close to an isometry) but lowers distortion between antipodal points (where the maximal distortion is attained).

The map  $F_s$  is length-preserving, and in particular is 1-Lipschitz but not  $L$ -Lipschitz for any  $L < 1$ . Thus the distortion of  $F_s$  is given by finding two points  $x, y \in S$  that maximize  $d(x, y) / \|F_s(x) - F_s(y)\|$ . A calculation shows that the distortion of  $F_s$  is attained by the pair of antipodal points  $F_s(0)$  and  $F_s(\pi)$  if  $s$  is not too large. For such  $s$  we calculate the distortion of  $F_s$  to be

$$\text{lip}(F_s) = \frac{\pi - 3s}{1 + \cos(s) - \sqrt{3}\sin(s)}.$$

The function  $\text{lip}(F_s)$  attains a minimum value at some point  $s_0$ . It can be shown numerically that  $s_0 \approx .2627$ , and that  $\text{lip}(F_{s_0}) \approx 1.5525$ , which is a mild improvement over the value 1.5708 above. We leave as a question whether the embedding  $F_{s_0}$  is indeed optimal.

**Question 4.1.** What is  $\text{lip}(S, \mathbb{R}^2)$ , and what embedding attains it?

In the other direction, an easy argument shows that  $\text{lip}(S, \mathbb{R}^2) \geq \sqrt{2}$ . Let  $x, y, z, w \in S$  be four evenly spaced points, so that  $d(x, y) = d(y, z) = d(z, w) = d(w, x) = \pi/2$  and  $d(x, z) = d(y, w) = \pi$ . Let  $f: S \rightarrow \mathbb{R}^2$  be an arbitrary embedding. By rescaling, we may assume that  $f$  is 1-Lipschitz. A consequence of Euler's quadrilateral theorem is the inequality

$$\|f(x) - f(z)\|^2 + \|f(y) - f(w)\|^2 \leq \|f(x) - f(y)\|^2 + \|f(y) - f(z)\|^2 + \|f(z) - f(w)\|^2 + \|f(w) - f(x)\|^2.$$

The right-hand side of this inequality is bounded by  $\pi^2$ . This implies that either  $\|f(x) - f(z)\| \leq \pi/\sqrt{2}$  or  $\|f(y) - f(w)\| \leq \pi/\sqrt{2}$ . In the first case we have

$$\text{lip}(f) \geq \frac{d(x, z)}{\|f(x) - f(z)\|} \geq \sqrt{2}.$$

The second case likewise gives  $\text{lip}(f) \geq \sqrt{2}$ .

## 4.2 The three-petal rose

This example provides a lower bound on  $\text{lip}(\Delta)$ , the minimal distortion required to embed every metric triangle in  $\mathbb{R}^2$ .

**Proposition 4.2.** Let  $G$  be an arbitrary embedding of  $R$  into  $\mathbb{R}^2$ . Then  $\text{lip}(G) \geq 2$ .

*Proof.* Without loss of generality, we may assume that  $G(o) = 0$  and that  $G$  is non-expanding, but that any rescaling of  $G$  by a factor  $r > 1$  is no longer non-expanding. In particular, for all  $\varepsilon > 0$ , we can find points  $x, y \in R$  such that  $d(x, y)/\|G(x) - G(y)\| \leq 1 + \varepsilon$ .

The triangle  $R$  is the union of six arcs  $S_i^1, S_i^2, i \in \{1, 2, 3\}$  with one endpoint  $o$  and other endpoint the antipodal point of  $S_i$ , which we denote by  $y_i$ .

Choose a value  $\varepsilon \in (0, 1/(2 \text{lip}(G)))$ . Observe that  $d(G(y_i), 0) \geq \varepsilon$  for each  $i$ . In particular, since each set  $G(S_i^j)$  is connected, we can find a point  $x_i^j \in S_i^j$  such that  $d(f(x_i^j), 0) = \varepsilon$ .

Enumerate the points  $x_i^j$  based on the order of  $G(x_i^j)$  along the circle  $S(0, \varepsilon)$  (with the counterclockwise orientation) from the positive  $x$ -axis as  $x_1, x_2, \dots, x_6$ . We can find consecutive points  $x_i, x_{i+1}$  (with the identification  $x_7 = x_1$ ) such that the angle  $\angle(0; G(x_i), G(x_{i+1})) \leq \pi/3$ . In particular,  $\|G(x_i) - G(x_{i+1})\| \leq \varepsilon$ . On the other hand, since  $G$  is non-expanding,  $d(x_i, x_{i+1}) = d(x_i, o) + d(o, x_{i+1}) \geq 2\varepsilon$ .

We conclude that

$$\frac{d(x_i, x_{i+1})}{\|G(x_i) - G(x_{i+1})\|} \geq \frac{2\varepsilon}{\varepsilon} = 2,$$

and hence that the distortion of  $G$  is at least  $2\varepsilon/\varepsilon = 2$ .  $\square$

The tripodal embedding  $F$  for the three-petal rose has distortion  $\text{lip}(F) = 2$ . Alternatively, let  $X$  be the union of three equilateral triangles with the same side length and sharing a common vertex, each separated by an angle of  $\pi/3$ . Then any arc length-preserving homeomorphism from  $R$  onto  $X$  also has distortion 2.

## 4.3 Tripods

As we have seen, tripods play a key role in the proof of Theorem 1.1. The class of tripods represents the "thinnest" possible metric triangles. We verify the natural expectation that the embedding  $F$  from the previous section is distortion minimizing for this class of triangles.

**Proposition 4.3.** Let  $T$  be a tripod that is non-degenerate, meaning that  $p_i > 0$  for all  $i \in \{1, 2, 3\}$ . Then  $\text{lip}(T, \mathbb{R}^2) = 2/\sqrt{3}$ , and this is obtained by the tripodal embedding.

*Proof.* We show that the tripodal embedding  $F$  has distortion  $2/\sqrt{3}$ . If  $x, y \in T_i$  for the same  $i \in \{1, 2, 3\}$ , then the tripodal embedding satisfies  $\|F(x) - F(y)\| = d(x, y)$ . Next, assume that  $x \in T_i$  and  $y \in T_j$  for some  $i \neq j$ . Assume that  $i = 1$  and  $j = 2$ . Let  $s = d(x, o)$  and  $t = d(y, o)$ . Then

$$\frac{\|F(x) - F(y)\|}{d(x, y)} \leq \frac{\|F(x) - F(o)\|}{d(x, y)} + \frac{\|F(o) - F(y)\|}{d(x, y)} = \frac{s}{s+t} + \frac{t}{s+t} = 1,$$

and

$$\frac{d(x, y)}{\|F(x) - F(y)\|} = \frac{s + t}{\|(s + \frac{t}{2}, \frac{\sqrt{3}t}{2})\|} = \frac{s + t}{\sqrt{(s + \frac{t}{2})^2 + \frac{3}{4}t^2}}.$$

For fixed  $t$ , the expression on the right-hand side attains a maximum when  $s = t$ . To check this, assume without loss of generality that  $t = 1$  and let

$$M(s) = \frac{(s + 1)^2}{(s + \frac{1}{2})^2 + \frac{3}{4}},$$

defined for  $s \geq 0$ . We compute  $M'(s) = (1 - s^2)/(1 + s + s^2)^2$ . Thus we have a single critical point at  $s = 1$ . We evaluate  $M(1) = 4/3$ . Also,  $M(0) = 1$  and  $\lim_{s \rightarrow \infty} M(s) = 1$ , which establishes that  $M(1)$  is a maximum.

The proof that no embedding can have smaller distortion is similar to the proof of Proposition 4.2 but with three arcs emanating from the center point  $o$  rather than six. We omit the details.  $\square$

#### 4.4 The twisted heart

This example shows the sharpness of Theorem 1.1. Observe first that  $d(b, e) = d(c, a) = 1$  and  $d(a, b) = 4$ . On the other hand, we compute that  $F(a) = (\frac{1}{2}, \frac{\sqrt{3}}{2})$ ,  $F(b) = (\frac{1}{2}, -\frac{\sqrt{3}}{2})$  and  $F(c) = (-\frac{3}{2}, -\frac{\sqrt{3}}{2})$ . Then

$$\frac{\|F(a) - F(b)\|}{d(a, b)} = \frac{\|(0, \sqrt{3})\|}{4} = \frac{\sqrt{3}}{4}$$

and

$$\frac{\|F(a) - F(c)\|}{d(a, c)} = \|(2, \sqrt{3})\| = \sqrt{7}.$$

Thus the tripod embedding on the twisted heart has distortion  $4\sqrt{7/3}$ .

The natural follow-up is to what extent this value can be reduced by using a different embedding. One candidate is to map  $H$  to an equilateral triangle in the obvious way; this gives a distortion of  $2\sqrt{7} \approx 5.29$ . By perturbing the image points, it seems possible from computations to reduce the distortion to roughly 4. We leave this example as an area for further exploration.

**Question 4.4.** What is  $\text{lip}(H, \mathbb{R}^2)$ , and what embedding attains it?

## 5 Metric polygons

In this section, we prove Proposition 1.3, which shows that the embedding result for metric triangles does not extend to the case of metric quadrilaterals.

We define for all  $\varepsilon > 0$  a quadrilateral  $Q = Q(\varepsilon)$  as follows. First we define a corresponding superset  $\tilde{Q}$ . Start with a circle of length 4 divided into four subarcs of length 1, enumerated in cyclic order as  $I_1, I_2, I_3, I_4$ . For each  $j = 1, \dots, 4$ , let  $p_j$  denote the initial point of  $I_j$  and  $m_j$  denote the midpoint. Let  $I_5$  be an interval of length  $\varepsilon$  connecting the point  $p_1$  to  $p_3$ , and let  $I_6$  be an interval of length  $\varepsilon$  connecting the point  $p_2$  to  $p_4$ . Let  $I_7$  be an interval of length  $\varepsilon$  connecting  $m_1$  and  $m_3$ . We now define  $\tilde{Q}$  to be the union of  $I_1, \dots, I_7$ . We equip  $\tilde{Q}$  with the length metric, denoted by  $d$ . We then take  $Q$  to be the subspace  $I_1 \cup \dots \cup I_4$ . One checks that  $Q$  is a metric quadrilateral with vertices  $p_1, \dots, p_4$ .


Proposition 1.3 is a consequence of the following proposition.

**Proposition 5.1.** Fix  $L \geq 1$  and choose  $0 < \varepsilon < 1/(6L^4)$ . Then the quadrilateral  $Q = Q(\varepsilon)$  is not  $L$ -bi-Lipschitz embeddable in the plane.

*Proof.* Assume that there is an  $L$ -bi-Lipschitz embedding  $f: Q \rightarrow \mathbb{R}^2$ . Given a point  $x \in Q$ , we write  $x'$  in place of  $f(x)$ .


Observe that  $f(Q)$  is a simple closed curve that, by the Jordan curve theorem, separates the plane and has two complementary components.

Let  $J_1$  be the straight line segment connecting  $p'_1$  to  $p'_3$ , which has length at most  $L\varepsilon$ . We can find a maximal closed subinterval  $\tilde{J}_1$  with interior contained in  $\mathbb{R}^2 \setminus f(Q)$  such that one endpoint  $j_1^1 \in \tilde{J}_1$  belongs to  $f(I_1 \cup I_4)$  and the other endpoint  $j_1^2 \in \tilde{J}_1$  belongs to  $f(I_2 \cup I_3)$ . Let  $\tilde{J}_1^1$  be the minimal subarc




(a) The quadrilateral  $Q(\varepsilon)$ .  
Each red edge has length  $\varepsilon$ .

(b) An embedding  
of  $Q(\varepsilon)$  in  $\mathbb{R}^2$



(c) The graph  $K_5$



(d) The graph  $K_{3,3}$

Figure 6: The metric polygons in Section 5.

of  $f(I_1 \cup I_4)$  connecting  $p'_1$  to  $j_1^1$ , and let  $\tilde{J}_1^2$  be the minimal subarc of  $f(I_2 \cup I_3)$  connecting  $j_1^2$  to  $p'_3$ . Observe that  $\tilde{J}_1^1$  and  $\tilde{J}_1^2$  each have length at most  $L^3\varepsilon$ , since

$$\ell(\tilde{J}_1^1) \leq L\ell(f^{-1}(\tilde{J}_1^1)) = Ld(p_1, f^{-1}(j_1^1)) \leq L^2\|p'_1 - j_1^1\| \leq L^3\varepsilon,$$

and similarly for  $\tilde{J}_1^2$ . Then the concatenation  $K_1 = \tilde{J}_1^1 * \tilde{J}_1^2 * \tilde{J}_1^3$  is a simple curve having length at most  $3L^3\varepsilon$ .

We define  $K_2$  in the same way as a simple curve with length at most  $3L^3\varepsilon$  connecting  $p'_2$  to  $p'_4$ .

If  $K_1$  and  $K_2$  were to have a point of intersection, it would follow that  $\|p'_1 - p'_2\| \leq 6L^3\varepsilon$ , and hence that  $d(p_1, p_2) \leq 6L^4\varepsilon < 1$ . Since  $d(p_1, p_2) = 1$ , we conclude that  $K_1$  and  $K_2$  do not have a point of intersection. In particular, it follows that  $\tilde{J}_1$  and  $\tilde{J}_2$  lie in different complementary components of  $f(Q)$ .

Let  $\tilde{S}$  be the union of  $K_1$ ,  $K_2$ ,  $f(I_2)$  and  $f(I_4)$ . We observe that  $\tilde{S}$  separates  $m'_1$  and  $m'_3$ . Then  $\tilde{S}$  contains a simple closed curve  $S$  also separating  $m'_1$  and  $m'_3$  (see Theorem IV.6.7 in [12]). Now  $\|m'_1 - m'_3\| \leq L\varepsilon$ , which implies that  $m'_1$  and  $m'_3$  lie at distance at most  $L\varepsilon < (6L^3)^{-1} \leq (6L)^{-1}$  from  $S$ .

However, the distance from  $m'_1$  to  $K_1 \cup K_2$  is at least

$$\frac{1}{2L} - 3L^2\varepsilon \geq \frac{1}{2L} - \frac{1}{6L^4} \geq \frac{1}{4L}.$$

Likewise, the distance from  $m'_1$  to  $f(I_2) \cup f(I_4)$  is at least  $(2L)^{-1}$ . Thus  $\text{dist}(m'_1, S) \geq (4L)^{-1}$ , which gives a contradiction.  $\square$

**Remark 5.2.** One might initially assume that taking  $\varepsilon = 0$  in the previous construction gives a metric quadrilateral that cannot be bi-Lipschitz embedded in  $\mathbb{R}^2$ . While indeed we do get a metric quadrilateral in this way, it is possible to bi-Lipschitz embed it in  $\mathbb{R}^2$ . The previous proof relies on the fact that a simple closed curve separates the plane. However, if  $\varepsilon = 0$ , the corresponding quadrilateral is not a simple closed curve, and there is more flexibility for where to map each edge.

**Example 5.3.** A metric pentagon may fail to even be topologically embeddable in  $\mathbb{R}^2$ . There are two basic examples of non-planar graphs:  $K_5$ , the complete graph on five vertices, and  $K_{3,3}$ , the complete bipartite graph on two sets of three vertices. Each of these can be realized as a metric pentagon.

We provide the details for  $K_5$ . Denote its topological vertices in cyclic order by  $p_1, p_2, \dots, p_5$ . Denote the topological edge from  $p_i$  to  $p_j$  by  $I_{ij}$ . We make  $K_5$  into a metric space by assigning each edge length 1 and taking the length metric. For each  $i = 1, \dots, 5$ , let  $m_i$  be the midpoint of  $I_{i(i+1)}$  (with

the usual identification of  $I_{56}$  with  $I_{51}$ ). The space  $K_5$  can be viewed as a metric pentagon by taking  $m_1, m_3, m_5, m_2, m_4$  as its vertices (as a metric polygon) in cyclic order. The edges (as a metric polygon) are  $[m_1p_1]*I_{14}*[p_4m_3]$ ,  $[m_3p_3]*I_{31}*[p_1m_5]$ ,  $[m_5p_5]*I_{53}*[p_3m_2]$ ,  $[m_2p_2]*I_{25}*[p_5m_4]$ ,  $[m_4p_4]*I_{42}*[p_2m_1]$ . Each of these edges has length 2.

Note that, for both metric spaces  $K_5$  and  $K_{3,3}$  (with the standard length metric just described), any two points are at distance at most 2 apart. This implies that any edge (as a metric polygon) has length at most 2. In particular, these spaces cannot be realized as metric quadrilaterals.

## 6 Appendix

In this appendix, we provide the proofs of the various inequalities used in Section 3. These are mostly straightforward calculus optimization problems, but we include the details for completeness. We recall the following functions defined in Section 3:

$$\begin{aligned} M_1(b, e) &= \frac{1}{4}(1-b+2e)^2 + \frac{3}{4}(1+b)^2; \\ M_2(b, e) &= \frac{(2+e)^2}{\frac{1}{4}(1-b+2e)^2 + \frac{3}{4}(1+b)^2}; \\ \widetilde{M}_2(b, e) &= \frac{(2-e)^2}{\frac{1}{4}(1-b+2e)^2 + \frac{3}{4}(1+b)^2}; \\ M_3(b, s, t) &= \frac{(2+2b-s-t)^2}{\frac{1}{4}(1-b+2s-2t)^2 + \frac{3}{4}(1+b)^2}; \\ M_4(b, s, t) &= \frac{1}{4}(1-b+2s+t)^2 + \frac{3}{4}(1+t+b)^2; \\ M_5(b, s, t) &= \frac{(\max\{2, 2b\} + s + t)^2}{\frac{1}{4}(1-b+2s+t)^2 + \frac{3}{4}(1+t+b)^2}; \\ M_6(b, s, t) &= \frac{(2+2b+2t)^2}{\frac{1}{4}(1-b+2s+t)^2 + \frac{3}{4}(1+t+b)^2}; \\ M_7(b, s, t) &= \frac{1}{4}(1-b-s+t)^2 + \frac{3}{4}(1+s+t+b)^2; \\ M_8(b, s, t) &= \frac{(2+s+t)^2}{\frac{1}{4}(1-b-s+t)^2 + \frac{3}{4}(1+s+t+b)^2}; \\ M_9(b, s, t) &= \frac{(2+2b+2s+2t)^2}{\frac{1}{4}(1-b-s+t)^2 + \frac{3}{4}(1+s+t+b)^2}. \end{aligned}$$

**Lemma 6.1.** The function  $M_1(b, e)$  is maximized on the set  $\{(b, e) : 0 \leq b, e \leq 1\}$  by taking  $b = e = 1$ , which gives a value of 4.

*Proof.* We compute  $\frac{\partial M_1}{\partial b}(b, e) = 1 + 2b - e$  and  $\frac{\partial M_1}{\partial e}(b, e) = 1 - b + 2e$ . These are both non-negative on the set  $\{(b, e) : 0 \leq b, e \leq 1\}$ . Thus  $M_1$  attains a maximum value by making  $b, e$  as large as possible, that is,  $b = e = 1$ . We compute  $M_1(1, 1) = 4$ .  $\square$

**Lemma 6.2.** The function  $\frac{1}{e^2}M_1(b, e)$  is maximized on the set  $\{(b, e) : 0 \leq b \leq 1, 1 \leq e < \infty\}$  by taking  $b = e = 1$ , which gives a value of 4.

*Proof.* We first compute

$$\frac{\partial M_1}{\partial e}(b, e) = -\frac{2 + 2b + 2b^2 + e - be}{e^3}.$$

Observe that since  $b \leq 1$ , the numerator is always positive, and we conclude that  $\frac{\partial M_1}{\partial e}(b, e) < 0$ . This implies that  $\frac{1}{e^2}M_1(b, e)$  attains a maximum when  $e = 1$ . The previous lemma now implies that a maximum value of 4 is attained when  $b = 1$ .  $\square$

**Lemma 6.3.** The function  $M_2(b, e)$  is maximized on the set  $\{(b, e) : b \geq 0\}$  by taking  $b = e = 0$ , which gives a value of 4.

*Proof.* We compute

$$\frac{\partial M_2}{\partial b}(b, e) = -\frac{(2+e)^2(1+2b-e)}{\left(\frac{1}{4}(1-b+2e)^2 + \frac{3}{4}(1+b)^2\right)^2}.$$

Observe that  $\frac{\partial M_2}{\partial b}(b, e) > 0$  if  $2b < e - 1$  and  $\frac{\partial M_2}{\partial b}(b, e) < 0$  if  $2b > e - 1$ . This implies that  $M_2(b, e)$  is greatest if  $2b + 1 = e$ , or if  $b = 0$ .

Now

$$\frac{\partial}{\partial b}M_2(b, 2b+1) = -\frac{2(3+2b)}{3(1+b)^3},$$

which is negative. Thus  $M_2(b, 2b+1)$  is maximized when  $b = 0$ , which gives a value of 4.

Similarly, if  $b = 0$ , then we compute

$$\frac{\partial}{\partial e}M_2(0, e) = -\frac{3e(2+e)}{(1+e+e^2)^2},$$

which is negative if  $e > 0$  or  $e < -2$  and positive if  $-2 < e < 0$ . Thus  $M_2(0, e)$  has a local maximum at  $e = 0$ . We also check that  $\lim_{e \rightarrow -\infty} M_2(0, e) = 1$ , and so  $e = 0$  is a global maximum. This completes the proof.

$$M_2(-e, e) = \frac{(2-e)^2}{\frac{1}{4}(1+3e)^2 + \frac{3}{4}(1-e)^2} \cdot \frac{2(e-2)(1+6e)}{(1+3e^2)^2}.$$

□

**Lemma 6.4.** The function  $\widetilde{M}_2(b, e)$  is maximized on the set  $\{(b, e) : b \geq 0, -b \leq e \leq 0\}$  by taking  $b = -e = 1/6$ , which gives a value of  $13/3$ .

*Proof.* We compute

$$\frac{\partial M_2}{\partial b}(b, e) = -\frac{(1+2b-e)(e-2)^2}{\left(\frac{1}{4}(1-b+2e)^2 + \frac{3}{4}(1+b)^2\right)^2}.$$

This is strictly negative over the given region. It follows that  $M_2(b, e)$  is maximized if  $b = -e$ . Now

$$\frac{\partial}{\partial e}M_2(-e, e) = \frac{2(e-2)(1+6e)}{(1+3e^2)^2}.$$

This is positive if  $e < -1/6$  and negative if  $e > -1/6$ . Thus there is a maximum at  $e = -1/6$ . We evaluate  $M_2(1/6, -1/6) = 13/3$ . □

**Lemma 6.5.** The function  $M_3(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t \leq 1\}$  by taking  $b = 1, s = t = 0$ , which gives a value of  $16/3$ .

*Proof.* We compute

$$\frac{\partial M_3}{\partial s}(b, s, t) = -\frac{(2+2b-s-t)(6+7b+3b^2+s-3t)}{2\left(\frac{1}{4}(1-b+2s-2t)^2 + \frac{3}{4}(1+b)^2\right)^2},$$

which is negative for all  $0 < b, s, t < 1$ . It follows that  $M_3(b, s, t)$  is maximized if  $s = 0$ . Next, we evaluate

$$\frac{\partial}{\partial t}M_3(b, 0, t) = -\frac{(2+2b-t)(2+3b+3b^2-t)}{2\left(\frac{1}{4}(1-b-2t)^2 + \frac{3}{4}(1+b)^2\right)^2},$$

which is also negative for all  $0 < b, t < 1$ . Thus  $M_3(b, 0, t)$  is maximized if  $t = 0$ . Finally, we evaluate

$$\frac{\partial M_3}{\partial b}(b, 0, 0) = \frac{(3-b)(1+b)}{\left(\frac{1}{4}(1-b)^2 + \frac{3}{4}(1+b)^2\right)^2},$$

which is positive if  $0 < b < 1$ . We conclude that  $M_3(b, s, t)$  is maximized when  $b = 1, s = t = 0$ . Evaluate  $M_3(1, 0, 0) = 16/3$ . □

**Lemma 6.6.** The function  $M_4(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t \leq 1, s+t \leq 1\}$  by taking  $s = 0, b = t = 1$ , which gives a value of 7.



*Proof.* Since  $1 - b + 2s + t \geq 0$ , we see that  $M_4(b, s, t)$  is increasing in  $s$  and  $t$ . Thus the maximum is attained when  $s + t = 1$ . Now

$$M_4(b, s, 1 - s) = \frac{3}{4}(2 + b - s)^2 + \frac{1}{4}(2 - b + s)^2.$$

Let  $u = b - s$ , so the previous expression becomes  $h(u) = \frac{3}{4}(2 + u)^2 + \frac{1}{4}(2 - u)^2$ . Then  $h'(u) = 2u + 2$ . So  $M_4$  is maximized when  $b - s$  is greatest, that is,  $b = 1$  and  $s = 0$ . Then  $t = 1$ . We evaluate  $M_4(1, 0, 1) = 7$ .  $\square$

**Lemma 6.7.** The function  $\frac{1}{(s+t)^2}M_4(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t, 1 \leq s + t, b \leq s + t\}$  by taking  $s = 0, b = t = 1$ , which gives a value of 7.

*Proof.* We evaluate

$$\frac{\partial}{\partial b} \frac{M_4(b, s, t)}{(s+t)^2} = \frac{1 + 2b - s + t}{(s+t)^2},$$

which is positive when  $2b > s - t - 1$ . If  $2b > s - t - 1$ , then  $\frac{1}{(s+t)^2}M_4(b, s, t)$  increases as  $b$  increases. If  $2b < s - t - 1$ , then  $\frac{1}{(s+t)^2}M_4(b, s, t)$  increases as  $b$  decreases. From this we conclude that  $\frac{1}{(s+t)^2}M_4(b, s, t)$  is greatest when either  $b = s + t$  or  $b = 0$ . When analyzing the latter case, note that  $t + 1 < s$ . Then

$$M_4(s + t, s, t) = \frac{s^2 + 2s + 3ts + 3t^2 + 3t + 1}{(s+t)^2}$$

and we compute

$$\frac{\partial}{\partial s} \frac{M_4(s + t, s, t)}{(s+t)^2} = -\frac{2 + 2s + st + 4t + 3t^2}{(s+t)^3},$$

which is negative over the same region. Thus the function is greatest when  $s + t = 1$ . The previous lemma now implies that a maximum value of 7 is attained when  $s = 0$  and  $b = t = 1$ .

On the other hand,

$$M_4(0, s, t) = \frac{s^2 + 2s + 3ts + 3t^2 + 3t + 1}{(s+t)^2}.$$

Let  $u = (\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}})$ . We compute

$$\frac{\partial}{\partial u} \frac{M_4(0, s, t)}{(s+t)^2} = \frac{s - t - 1}{\sqrt{2}(s+t)^2},$$

which is positive. From this, we can assume that  $t = 0$ . Then  $M_4(0, s, 0) = 1 + \frac{1}{s} + \frac{1}{s^2}$ , which is at most 3.  $\square$

**Lemma 6.8.** The function  $\frac{1}{b^2}M_4(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t, 1 \leq b, s + t \leq b\}$  by taking  $s = 0, b = t = 1$ , which gives a value of 7.

*Proof.* Let  $u = (0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ . The directional derivative of  $\frac{1}{b^2}M_4(b, s, t)$  in the direction of  $u$  is  $3(1 + s + t)/(\sqrt{2}b^2)$ , which is positive. We conclude that  $\frac{1}{b^2}M_4(b, s, t)$  is greatest when  $s + t = b$ . This reduces to Lemma 6.7, and so we obtain the same conclusion.  $\square$

**Lemma 6.9.** The function  $M_5(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t, s \leq 1\}$  by taking  $b = 6, s = 1, t = 0$ , which gives a value of 13/3.

*Proof.* Let  $M_5^a(b, s, t)$  denote the function defined like  $M_5$  but replacing “ $\min\{2, 2b\}$ ” with “2”, and similarly for  $M_5^b$ . It suffices to consider the functions  $M_5^a$  and  $M_5^b$  separately.

Let  $u = (0, \frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}})$ . Then the directional derivative of  $M_5^a$  in the  $u$ -direction is

$$D_u M_5^a(b, s, t) = \frac{(1 + 2b - s + t)(2 + s + t)^2}{\sqrt{2}(\frac{1}{4}(1 - b + 2s + t)^2 + \frac{3}{4}(1 + b + t)^2)}.$$

If  $s < 1 + 2b + t$ , then  $D_u M_5^a$  is positive; if  $s > 1 + 2b + t$ , then  $D_u M_5^a$  is negative. Thus along a given curve in the  $u$ -direction,  $M_5^a(b, s, t)$  is greatest when  $s = 1 + 2b + t$ . Note that this point lies outside the interior of the target region  $\{(b, s, t) : 0 \leq b, s, t, s \leq 1\}$ . From this we deduce that the maximum necessarily occurs when  $t = 0$  or when  $s = 1$ .

$$M_5^a(b, 1 + 2b + t, t) = \frac{(3 + 2b + 2t)^2}{3(1 + b + t)^2}.$$

In the case that  $t = 0$ , we evaluate

$$M_5^a(b, s, 0) = \frac{(2 + s)^2}{\frac{1}{4}(1 - b + 2s)^2 + \frac{3}{4}(1 + b)^2}.$$

Compute

$$\frac{\partial}{\partial b} \frac{(2 + s)^2}{\frac{1}{4}(1 - b + 2s)^2 + \frac{3}{4}(1 + b)^2} = -\frac{(1 + 2b - s)(2 + s)^2}{(\frac{1}{4}(1 - b + 2s)^2 + \frac{3}{4}(1 + b)^2)^2}.$$

Since  $s \leq 1$ , this is negative, so we conclude that the maximum occurs when  $b = 0$ . Now

$$M_5^a(0, s, 0) = \frac{(2 + s)^2}{\frac{1}{4}(1 + 2s)^2 + \frac{3}{4}}.$$

This function is decreasing in  $s$  for  $s > 0$ , so we conclude that  $s = 0$ . We compute  $M_5^a(0, 0, 0) = 4$ , so  $M_5^a$  attains a maximum value for  $b = s = t = 0$ .

In the case that  $s = 1$ , we evaluate

$$M_5^a(b, 1, t) = \frac{(3 + t)^2}{\frac{1}{4}(3 - b + t)^2 + \frac{3}{4}(1 + b + t)^2}.$$

A computation shows that the  $b$ -derivative of the right-hand side is negative, and thus any maximum occurs when  $b = 0$ . We then deduce that  $t = 0$ . Evaluate  $M_5^a(0, 1, 0) = 3$ .

The function  $M_5^b$  is handled similarly and we highlight the differences. We compute

$$D_u M_5^b(b, s, t) = \frac{(1 + 2b - s + t)(2b + s + t)^2}{\sqrt{2}(\frac{1}{4}(1 - b + 2s + t)^2 + \frac{3}{4}(1 + b + t)^2)^2}.$$

From this, we conclude that the maximum occurs when  $s = 1 + 2b + t$ , when  $t = 0$  or when  $s = 1$ . As before, the line  $s = 1 + 2b + t$  lies outside the target domain, so we assume that  $t = 0$  or  $s = 1$ .

If  $t = 0$ , we compute

$$\frac{\partial}{\partial s} M_5^b(b, s, 0) = \frac{(2b + s)(2 + 4b^2 + s - 5bs)}{(\frac{1}{4}(1 - b + 2s + t)^2 + \frac{3}{4}(1 + b + t)^2)^2}.$$

This is positive for  $b \geq 0$ ,  $0 < s < 1$ , so we conclude that  $s = 1$ .

Thus we can consider the second case that  $s = 1$ . We compute

$$\frac{\partial}{\partial b} M_5^b(b, 1, t) = -\frac{(1 + 2b + t)(-12 + 2b - 11t - 3t^2)}{(\frac{1}{4}(1 - b + 2s + t)^2 + \frac{3}{4}(1 + b + t)^2)^2}$$

and

$$\frac{\partial}{\partial t} M_5^b(b, 1, t) = -\frac{(1 + 2b + t)(-3 + 7b - t + 3bt)}{(\frac{1}{4}(1 - b + 2s + t)^2 + \frac{3}{4}(1 + b + t)^2)^2}.$$

The first of these is zero if  $b = -(11t + 3t^2)/12$ , while the second of these is zero if  $b = (3 + t)/(7 + 3t)$ . There is no  $t > 0$  satisfying both these relations, and so we conclude that  $M_5^b(b, 1, t)$  does not have any critical points when  $0 < b, t$ . Thus any maximum must occur when  $t = 0$  or  $b = 0$ .

Next,

$$\frac{\partial}{\partial b} M_5^b(b, 1, 0) = -\frac{2(-6 + b)(1 + 2b)}{(3 + b^2)^2},$$

so  $M_5^b(b, 1, 0)$  has a maximum at  $b = 6$ . Compute  $M_5^b(6, 1, 0) = 13/3$ . Similarly,

$$\frac{\partial}{\partial b} M_5^b(0, 1, t) = -\frac{(1 + t)(3 + t)}{(3 + 3t + t^2)^2},$$

which is positive for  $t > 0$ . So  $\lim M_5^b(0, 1, t) = 1$  is a bound on  $M_5^b(0, 1, t)$  for  $t > 0$ .

Finally,  $\frac{\partial}{\partial t} M_5^b(b, 1, t)$  is negative whenever  $b \geq 1$ , which implies  $M_5^b(b, 1, t) \leq M_5^b(b, 1, 0)$  for all  $b, t$  satisfies  $b \geq 1$ . Also,  $\frac{\partial}{\partial b} M_5^b(b, 1, t)$  is positive whenever  $b \leq 1$ . These facts allow us to conclude that  $M_5^b(b, 1, t)$  does in fact attain a maximum.  $\square$

**Lemma 6.10.** The function  $M_6(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t\}$  by taking  $s + t = 1 + b$ , which gives a value of  $16/3$ .

*Proof.* Compute

$$\frac{\partial M_6}{\partial s}(b, s, t) = \frac{4(-1 + b - 2s - t)(1 + b + t)^2}{(\frac{1}{4}(1 - b + 2s + t)^2 + \frac{3}{4}(1 + t + b)^2)^2}.$$

This is positive if  $2s < -1 + b - t$  and negative if  $2s > -1 + b - t$ . Thus any maximum of  $M_6$  occurs when  $2s = -1 + b - t$ , if  $-1 + b - t \geq 0$ , or  $-1 + b - t < 0$  and  $s = 0$ . In the first case, we evaluate  $M_6(b, (-1 + b - t)/2, t) = 16/3$ .

In the second case, evaluate

$$\frac{\partial M_6}{\partial b}(b, 0, t) = -\frac{4(-1 + b - t)(1 + t)(1 + b + t)}{(b^2 + b(1 + t) + (1 + t)^2)^2}.$$

If  $-1 + b - t < 0$ , then this derivative is positive. We conclude that any maximum for  $M_6(b, 0, t)$  occurs when  $-1 + b - t = 0$ , which returns us to the previous case.  $\square$

**Lemma 6.11.** The function  $M_7(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t \leq 1, s + t \leq 1\}$ ,  $M_7(b, s, t)$  by taking  $s = 0$ ,  $b = t = 1$ , or  $t = 0$ ,  $b = s = 1$ . This gives a value of 7.

*Proof.* We evaluate

$$\frac{\partial}{\partial b} M_7(b, s, t) = \frac{\partial}{\partial s} M_7(b, s, t) = 1 + 2b + 2s + t,$$

which is positive. Thus  $M_7$  is greatest when  $b = 1$  and  $s + t = 1$ . Now

$$M_7(1, s, 1 - s) = \frac{27}{4} + \frac{1}{4}(1 - 2s)^2,$$

which attains a maximum of 7 when  $s = 0$  or  $s = 1$ .  $\square$

**Lemma 6.12.** The function  $\frac{1}{(s+t)^2} M_7(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t \leq 1, s + t \geq 1\}$ ,  $M_7(b, s, t)$  by taking  $s = 0$ ,  $b = t = 1$ , or  $t = 0$ ,  $b = s = 1$ . This gives a value of 7.

*Proof.* We evaluate

$$\frac{\partial}{\partial b} \frac{M_7(b, s, t)}{(s+t)^2} = \frac{1 + 2b + 2s + t}{(s+t)^2},$$

which is positive for  $0 < b, s, t < 1$ . This implies the function is greatest when  $b = 1$ . We now compute

$$\frac{\partial}{\partial s} \frac{M_7(1, s, t)}{(s+t)^2} = \frac{-6 + ts - 3s - 3t - t^2}{(s+t)^3},$$

which is negative over the same region. Thus the function is greatest when  $s + t = 1$ . The previous lemma now implies that a maximum value of 7 is attained when  $s = 0$  and  $b = t = 1$ , or  $t = 0$  and  $b = s = 1$ .  $\square$

**Lemma 6.13.** The function  $M_8(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t\}$  by taking  $b = s = t = 0$ . This gives a value of 4.

*Proof.* Let  $u = (0, \frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}})$ . Then the directional derivative of  $M_8$  in the  $u$ -direction is

$$D_u M_8(b, s, t) = \frac{(b - 1 + s - t)(2 + s + t)^2}{\sqrt{2}(\frac{1}{4}(1 - b - s + t)^2 + \frac{3}{4}(1 + b + s + t)^2)^2}.$$

If  $s - t < 1 - b$ , then  $D_u M_8$  is positive; if  $s - t > 1 - b$ , then  $D_u M_8$  is negative. Thus along a given curve in the  $u$ -direction,  $M_8(b, s, t)$  is maximized when  $s = 1 - b + t$ . Note that this point may lie outside the target region  $\{(b, s, t) : 0 \leq b, s, t\}$ , in which case the maximum necessarily occurs when  $t = 0$ . We examine each case separately. First, we take  $s = 1 - b + t$ . Then

$$M_8(b, 1 - b + t, t) = \frac{(-3 + b - 2t)^2}{3(1 + t)^2}.$$

This function is decreasing in  $b$  if  $b < 3 + 2t$  and increasing in  $b$  if  $b > 3 + 2t$ . Moreover, for fixed  $t$ , since  $s \geq 0$ , it follows that  $b$  satisfies the relation  $b \leq 1 + t$ . Thus the maximum must occur when  $b = 0$  or  $b = 1 + t$ . This gives

$$M_8(0, 1 + t, t) = \frac{4(3 + 2t)^2}{3(2 + 2t)^2}$$

and

$$M_8(1 + t, 0, t) = \frac{4(2 + t)^2}{3(2 + 2t)^2}.$$

Each of these functions is decreasing. So each has a maximum when  $t = 0$ . We evaluate  $M_8(0, 1, 0) = 3$  and  $M_8(1, 0, 0) = 4/3$ .

Next, we take  $t = 0$ . Then

$$M_8(b, s, 0) = \frac{(2 + s)^2}{\frac{1}{4}(1 - b - s)^2 + \frac{3}{4}(1 + b + s)^2}.$$

Compute

$$\frac{\partial}{\partial b} \frac{(2 + s)^2}{\frac{1}{4}(1 - b - s)^2 + \frac{3}{4}(1 + b + s)^2} = -\frac{((2 + s)^2(1 + 2b + 2s))}{1 + b + b^2 + s + 2bs + s^2}.$$

This is negative, so we conclude that the maximum occurs when  $b = 0$ . Now

$$M_8(0, s, 0) = \frac{(2 + s)^2}{\frac{1}{4}(1 - s)^2 + \frac{3}{4}(1 + s)^2}.$$

This function is decreasing in  $s$  for  $s > 0$ , so we conclude that  $s = 0$ . We compute  $M_8(0, 0, 0) = 4$ , so  $M_8$  attains a maximum value for  $b = s = t = 0$ .  $\square$

**Lemma 6.14.** The function  $M_9(b, s, t)$  is maximized on the set  $\{(b, s, t) : 0 \leq b, s, t\}$  by taking  $t = b + s - 1$ , which gives a value of  $16/3$ .

*Proof.* We compute

$$\frac{\partial}{\partial t} M_9(b, s, t) = \frac{4(b + s)(-1 + b + s - t)(1 + b + s + t)}{(\frac{1}{4}(1 - b - s + t)^2 + \frac{3}{4}(1 + b + s + t)^2)^2}.$$

This is positive if  $t + 1 < b + s$  and negative if  $t + 1 > b + s$ , so we conclude that  $M_9(b, s, t)$  is largest when  $t = b + s - 1$ . Now  $M_9(b, s, b + s - 1)$  evaluates to  $16/3$ .  $\square$

## References

- [1] M. R. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [2] D. Burago, Y. Burago, and S. Ivanov. *A course in metric geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.
- [3] G. C. David, S. Eriksson-Bique, and V. Vellis. Bi-Lipschitz embeddings of quasiconformal trees. *Proc. Amer. Math. Soc.*, 151(5):2031–2044, 2023.
- [4] G. C. David and V. Vellis. Bi-Lipschitz geometry of quasiconformal trees. *Illinois J. Math.*, 66(2):189–244, 2022.
- [5] J. Heinonen. *Geometric embeddings of metric spaces*, volume 90 of *Report. University of Jyväskylä Department of Mathematics and Statistics*. University of Jyväskylä, Jyväskylä, 2003.
- [6] U. Lang and C. Plaut. Bilipschitz embeddings of metric spaces into space forms. *Geom. Dedicata*, 87(1-3):285–307, 2001.
- [7] J. Matoušek. *Lectures on discrete geometry*, volume 212 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2002.

- [8] A. Naor and O. Neiman. Assouad's theorem with dimension independent of the snowflaking. *Rev. Mat. Iberoam.*, 28(4):1123–1142, 2012.
- [9] D. Ntalampekos and M. Romney. Polyhedral approximation of metric surfaces and applications to uniformization. *Duke Math. J.*, 172(9):1673–1734, 2023.
- [10] M. I. Ostrovskii. *Metric embeddings*, volume 49 of *De Gruyter Studies in Mathematics*. De Gruyter, Berlin, 2013. Bilipschitz and coarse embeddings into Banach spaces.
- [11] S. Semmes. Bilipschitz embeddings of metric spaces into Euclidean spaces. *Publ. Mat.*, 43(2):571–653, 1999.
- [12] R. L. Wilder. *Topology of manifolds*, volume 32 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1963.

Department of Mathematical Sciences, Stevens Institute of Technology, 1 Castle Point Terrace, Hoboken NJ 07030, USA.

*Email address:* xluo15@stevens.edu, mromney@stevens.edu, atao@stevens.edu