THE HI EXTENSION OF THE STANDARD HI SPACES

SPIROS A. ARGYROS, ANTONIS MANOUSSAKIS, AND PAVLOS MOTAKIS

ABSTRACT. A Hereditarily Indecomposable (HI) Banach space X admits an HI extension if there exists an HI space Z such that X is isomorphic to a subspace Y of Z and Z/Y is of infinite dimension. The problem whether or not every HI space admits an HI extension is attributed to A. Pelczynski. In this paper we present a method to define HI-extensions of the standard HI spaces, a class which includes the Gowers-Maurey space, asymptotic ℓ_p -HI space and others.

1. INTRODUCTION

The problem of the HI extension of an HI Banach space can be considered as part of a general problem which concerns the embedding of Banach spaces into a space with tight structure. There are several results related to this problem with the more representative being the following: The first one stated and proved in [7] asserts that every separable reflexive Banach space is isomorphic to a subspace of a reflexive indecomposable Banach space. The second one is similar to the previous result where the superspace admits very few operators. In particular every separable reflexive Banach space is isomorphic to a subspace of a \mathcal{L}_{∞} space which satisfies the "scalar-plus-compact" property. This result is presented in [8] for uniformly convex spaces and the general case is proved in an unpublished paper of the same authors. In this direction it remains open whether every separable Banach space not containing c_0 is embedded into a \mathcal{L}_{∞} space with very few operators.

Returning to the problem of the HI extension of an HI space, attributed to A. Pelczynski, we point out that this is a delicate and hard problem. This is mainly due to the quasi-minimality of HI Banach spaces [14]. To explain the obstacle a little further, assume that X is an HI space and Z is also HI containing isomorphically the space X. Then every subspace Y of X and every subspace E of Z contain further subspaces D_1 of Y and D_2 of E which are isomorphic. This means that for a possible construction of an HI superspace Z of X we ought to extract the structure of the space X in the space Z and this is hard, in particular when we do not have a precise description of the structure of the space X.

As it stands, there exists a specific type of HI space for which the answer to Pelczynski's problem is known to be affirmative. In [11], J. M. F. Castillo, V. Ferenczi and M. Gonzalez proved that Ferenczi's uniformly convex HI space \mathcal{F} from [13] admits an HI extension \mathcal{F}_2 . The space \mathcal{F} is the outcome of a complex interpolation scheme, and this is an unavoidable component of the definition of \mathcal{F}_2 . As such, the method from [11] is not applicable to the more standard HI spaces defined via the method of norming sets. The aim of this paper is to exhibit a method that permits us to define HI extensions of the latter class of HI spaces,

²⁰¹⁰ Mathematics Subject Classification: Primary 46B03, 46B06.

which includes the first generation of HI spaces such as the Gowers-Maurey HI space, [15], the Argyros-Deliyanni asymptotic ℓ_1 HI space, [3], and the Deliyanni-Manoussakis asymptotic ℓ_p -HI spaces, [12]. We have chosen to apply the method on a variant of Gowers-Maurey space, denoted by X_{hi} , with its norming set W_{hi} being a subset of the norming set of the mixed Tsirelson space $T[(\mathcal{A}_{n_i}, m_i^{-1})_{j \in \mathbb{N}}]$.

The norming set of any HI space which has been defined in the literature has two inevitable characteristics. The first one is some unconditional structure which is usually imposed by a mixed Tsirelson space. For example in Gowers-Maurey space, Schlumprecht space [20] plays that role. The second one is the conditional structure which is responsible for the tight structure of the HI space and it is traced in the earlier work of B. Maurey and H. P. Rosenthal [19]. In the case of X_{hi} the norming set W_{hi} is defined as follows: The set W_{hi} is a subset of $c_{00}(\mathbb{N})$ which is minimal with respect to the following properties.

- (i) It is rationally convex, symmetric, closed on the projections on intervals of \mathbb{N} and contains $\pm e_n^*$, $n \in \mathbb{N}$.
- (ii) It is closed in all $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})$ operations. That means that for every family $(f_i)_{i=1}^{n_{2j}}$ of elements of W_{hi} with successive supports the functional $f = m_{2j}^{-1} \sum_{i=1}^{n_{2j}} f_i \in W_{hi}$. In this case we say that f is a weighted functional with weight w(f) of f equal to m_{2j} . (Note that this is the unconditional structure mentioned above).
- (iii) It includes all $f = m_{2j-1}^{-1} \sum_{i=1}^{n_{2j-1}} f_i$ where $(f_i)_{i=1}^{n_{2j-1}}$ is a (2j-1)-special sequence. For such a functional we set $w(f) = m_{2j-1}$.

(2j-1)-special sequences are sequences $(f_i)_{i=1}^{n_{2j-1}}$ of successive weighted functionals which are defined through a coding function. Their important property is a tree-like property, namely for every i > 1 the weight $w(f_i) = m_{2j_i}$ of f_i uniquely determines the previous elements of the sequence. For a precise definition of the (2j-1)-special sequences see Definition 4.1. This type of functionals imposes the HI structure of the space X_{hi} .

We now pass to describe the HI extension of the space X_{hi} mentioned above. Our approach is the natural one. Namely, we shall define a new norming set W_{ex} such that the induced norm on $c_{00}(\mathbb{N})$ is an HI one and also the completion of the subspace generated by the even coordinates $(e_{2n})_{n\in\mathbb{N}}$ is isometric to the initial space X_{hi} . For this, we let $\widehat{W}_{hi} = \{\widehat{g} : g \in W_{hi}\}$ where for $g \in W_{hi}$ and $n \in \mathbb{N}$, $\widehat{g}(2n) = g(n)$ and $\widehat{g}(2n-1) = 0$. Then, W_{ex} is the norming set that satisfies the following two conditions:

- (a) The norm induced by W_{ex} is an HI one.
- (b) $W_{ex}|_{\mathbb{N}_{ev}} = \widehat{W}_{hi}$, where \mathbb{N}_{ev} is the set of even positive integers and $g|_{\mathbb{N}_{ev}} = g \cdot 1_{\mathbb{N}_{ev}}$.

We set X_{ex} the completion of $(c_{00}(\mathbb{N}), \|\cdot\|_{W_{ex}})$. Condition (b) yields that the subspace $X_{ev} = \overline{\langle e_{2n} : n \in \mathbb{N} \rangle}$ of X_{ex} is isometric to X_{hi} . In the sequel we denote the subspace X_{ev} by \widehat{X}_{hi} . Let us also observe that since the space X_{ex} is an HI space, the set \widehat{W}_{hi} cannot be a subset of W_{ex} . Otherwise, the subspace \widehat{X}_{hi} would be complemented in X_{ex} . Moreover, since $\|\cdot\|_{W_{ex}}$ is HI, the norming set W_{ex} must include special functionals which are entirely responsible for the HI structure. At the same time, for every special functional $f \in W_{ex}$, $f|_{\mathbb{N}_{ev}} = \hat{g}$ with $g \in W_{hi}$. This means that the added special functional in W_{ex} does not contribute to \widehat{X}_{hi} norm beyond the elements of \widehat{W}_{hi} . To satisfy the previously explained two necessary requirements, we introduce paired special functionals that will be described below. The norming set W_{ex} is defined to satisfy the following properties.

- (i) It contains $(e_n^*)_{n \in \mathbb{N}}$, it is symmetric, closed under rational convex combinations, and the restrictions on intervals of \mathbb{N} .
- (ii) It is closed under the even operations $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})_j$.
- (iii) $W_{ex}|_{\mathbb{N}_{ev}} = \widehat{W}_{hi}$.
- (iv) It contains the paired special functionals.

The paired special functionals are of the form $f = m_{2j+1}^{-1} \sum_{k=1}^{n_{2j+1}} (f_k + \hat{g}_k)$ such that $\sup f_k \subset \mathbb{N}_{od}$, the set of the odd positive integers, $w(f_k) = w(g_k)$, and $m_{2j+1}^{-1} \sum_{k=1}^{n_{2j+1}} g_k$ is a special functional in the norming set W_{hi} . The definition of the paired special sequence $f_1, \hat{g}_1, \ldots, f_{n_{2j+1}}, \hat{g}_{n_{2j+1}}$ uses a new coding function so that the tree-like property is also satisfied. Notice that the paired special functionals satisfy the requirement that $f|_{\mathbb{N}_{ev}} = \hat{g}$ with $g \in W_{hi}$. The HI property of the space X_{ex} follows from the following criterion from [11, Lemma 7.1] (see Proposition 6.1): the subspace \hat{X}_{hi} is HI and the quotient map $Q: X_{ex} \to X_{ex}/\hat{X}_{hi}$ is a strictly singular operator. The latter is a consequence of the next proposition (Proposition 7.7).

Proposition 1.1. For every infinite dimensional subspace Y of X_{ex} we have that $\operatorname{dist}(S_Y, S_{\widehat{X}_{b,i}}) = 0.$

The key ingredients for proving this result are paired special functionals.

The paper is organized into nine sections. The first three sections, after the introduction, are devoted to a brief presentation of basic ingredients and results for defining the norming set W_{hi} and proving its properties. The next section contains the definition of the norming set W_{ex} of the space X_{ex} . In the following two sections the proof that the space X_{ex} is HI is given. In the last two sections we present two further HI extensions. The first one concerns a norming set W which is not closed in rational convex combinations and the second one refers to HI extensions of asymptotic ℓ_1 HI spaces.

2. Mixed Tsirelson spaces

In this section we recall the definition of a class of mixed Tsirelson spaces and some of their properties that we shall use. We refer to [9] for a detailed presentation. Also we follow standard terminology as in [1],[18].

Let $(m_j)_{j=0}^{+\infty}$, $(n_j)_{j=0}^{+\infty}$ be two strictly increasing sequences satisfying the following:

- (1) $m_0 = 2$ and $m_{j+1} = m_j^5$.
- (2) $n_0 = 4$ and $n_{j+1} = (12n_j)^{s_j}$ where $2^{s_j} \ge m_{j+1}^3$.

By \mathcal{A}_n we denote the family of the finite subsets of \mathbb{N} with at most n elements, i.e., $\mathcal{A}_n = \{F \subset \mathbb{N} : \#F \leq n\}.$

Definition 2.1. 1) Let $n \in \mathbb{N}$. A sequence $(f_i)_{i=1}^k$, $k \leq n$, of elements of $c_{00}(\mathbb{N})$ is said to be \mathcal{A}_n -admissible if $\operatorname{supp}(f_1) < \operatorname{supp}(f_2) < \cdots < \operatorname{supp}(f_k)$.

2) The $(\mathcal{A}_{n_j}, m_j^{-1})$ -operation on $c_{00}(\mathbb{N})$ is the operation which assigns to each n_j -admissible sequence $(f_i)_{i=1}^k$ the vector $m_j^{-1}(f_1 + f_2 + \cdots + f_k)$.

For $f \in c_{00}(\mathbb{N})$ we shall write $w(f) = m_j$ if f is the result of an (n_j, m_j^{-1}) -operation.

We present the norming set of the mixed Tsirelson space defined by the families \mathcal{A}_{n_j} and the "weights" $(m_j)_j$ which is a discretization of the norming set of Schlumprecht space \mathcal{S} , [20].

Definition 2.2. Let W_{mT} be the smallest subset of $c_{00}(\mathbb{N})$ with the following properties

- (1) $\pm e_n^* \in W_{mT}$ for every $n \in \mathbb{N}$.
- (2) W_{mT} is closed in the $(\mathcal{A}_{n_j}, m_j^{-1})$ -operations, i.e., for every \mathcal{A}_{n_j} -admissible sequence $(f_i)_{i=1}^k$ of elements of W_{mT} we have that $m_j^{-1} \sum_{i=1}^k f_i \in W_{mT}$.
- (3) W_{mT} is symmetric and closed under the restriction of its elements to intervals of \mathbb{N} .
- (4) W_{mT} is closed in rational convex combinations.

The set W_{mT} can also be defined inductively as $W_{mT} = \bigcup_{n=0}^{\infty} W_{mT}^n$ where $W_{mT}^0 = \{\pm e_n^* : n \in \mathbb{N}\}$ and W_{mT}^n is the smallest subset of $c_{00}(\mathbb{N})$ that contains W_{mT}^{n-1} and satisfies properties (2)-(3) of the above definition with respect to the functionals of W_{mT}^{n-1} .

We denote by $X_{mT} = T[(\mathcal{A}_{n_j}, m_j^{-1})_j]$ the completion of $c_{00}(\mathbb{N})$ under the norm $\|\cdot\|_{W_{mT}}$ induced by the set W_{mT} , i.e.,

$$||x||_{W_{mT}} = \sup\{f(x) : f \in W_{mT}\}$$

Definition 2.3. Let $f \in W_{mT}$. For a finite tree \mathcal{A} , a family $(f_{\alpha})_{\alpha \in \mathcal{A}}$ is said to be a tree analysis of f if the following are satisfied.

- (1) \mathcal{A} has a unique root denoted by 0 and $f_0 = f$.
- (2) For every maximal node $\alpha \in \mathcal{A}$ we have that $f_{\alpha} = \pm e_{n_{\alpha}}^{*}$.
- (3) Let α be a non-maximal node of \mathcal{A} and denote by S_{α} the set of the immediate successors of α . Then either there exists $j \in \mathbb{N}$ such that $(f_{\beta})_{\beta \in S_{\alpha}}$ is a n_j admissible subset of W_{mT} and $f_{\alpha} = m_j^{-1} \sum_{\beta \in S_{\alpha}} f_{\beta}$ or there exists a rational convex combination $(r_{\beta})_{\beta \in S_{\alpha}}$ such that $f_{\alpha} = \sum_{\beta \in S_{\alpha}} r_{\beta} f_{\beta}$.

Proposition 2.4. Every $f \in W_{mT}$ admits a tree analysis.

The proof is easy. We simple consider all $f \in W_{mT}$ admitting a tree analysis and then we show that this set satisfies (1),(2) and (3) of the above definition. The result follows from the minimality of the set W_{mT} .

Definition 2.5. Let $f \in W_{mT}$ and $(f_{\alpha})_{\alpha \in \mathcal{A}}$ be a tree analysis of f. For $\alpha \in \mathcal{A}$ the functional f_{α} is said to be weighted with weight $w(f_{\alpha}) = m_j$ if f_{α} is the result of an $(\mathcal{A}_{n_j}, m_j^{-1})$ -operation of its successors.

It was proved in [10], Lemma 3.15, that every $f \in W_{mT}$ admits a tree analysis $(f_{\alpha})_{\alpha \in \mathcal{A}}$ such that whenever f_{α} is rational convex combination of its successors, i.e., $f_{\alpha} = \sum_{\beta \in S_{\alpha}} r_{\beta} f_{\beta}$, then every f_{β} is a weighted functional and $\operatorname{ran}(f_{\beta}) \subset \operatorname{ran}(f_{\alpha})$.

3. Even operation complete subsets of W_{mT}

Definition 3.1. A subset W of W_{mT} is said to be even operation complete (EO-complete) if

- (1) $\pm e_n^* \in W$ for every $n \in \mathbb{N}$.
- (2) W is closed in the $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})$ -operation for every $j \in \mathbb{N}$.
- (3) W is symmetric and closed under the restriction of its elements to intervals of N.

(4) Every $f \in W$ admits a tree analysis $(f_{\alpha})_{\alpha \in \mathcal{A}}$ such that $f_{\alpha} \in W$ for every

In the rest of this section by W we will denote any EO-complete subset of W_{mT} . We will denote by X_W the completion of $c_{00}(\mathbb{N})$ under the norm $\|\cdot\|_W$ induced by W.

Definition 3.2. A vector $x \in X_W$ is said to be a $C - \ell_1^k$ - average if there exists $x_1 < x_2 < \cdots < x_k$ such that $x = \frac{1}{k} \sum_{i=1}^k x_i$, and $||x_i|| \le C ||x||$ for each *i*. Moreover if ||x|| = 1 then x is called a normalized $C - \ell_1^k$ -average.

Lemma 3.3. Let Y be a block subspace of X_W . Then for every $j \in \mathbb{N}$ there exists a block sequence $(y_i)_{i=1}^{n_{2j}}$ such that $y_i \in Y$ and $n_{2j}^{-1} \sum_{i=1}^{n_{2j}} y_i$ is a normalized $2 - \ell_1^{n_{2j}} - \ell_1^{n_{2j}} - \ell_1^{n_{2j}}$. average

The existence of normalized ℓ_1^k -averages was first established by Th. Schlumprecht in his space S ([20]), with the use of Krivine theorem, since ℓ_1 is finitely block representable in S. Later W. T. Gowers and B. Maurey, ([15] Lemma 3), provided a James type argument to construct $2 - \ell_1^k$ -averages in S. For a proof in the setting of the space X_W we refer the reader to [9] Lemma II.22, [4] Lemma 4.6

The following lemma concerning ℓ_1^{nj} -averages is an essential tool for providing upper bounds on the norm of certain vectors.

Lemma 3.4. Let x be an $C - \ell_1^{n_j}$ -average in X_W and $f \in W$ with $w(f) = m_i$, i < j. Then

$$\left|f(x)\right| \le \frac{3C}{2m_i}.$$

We refer to [15] Lemma 4, or [9] Lemma II.23, for a proof.

Definition 3.5. A block sequence $(x_k)_{k\in\mathbb{N}}$ in X_W is said to be a (C,ϵ) -rapidly increasing sequence (RIS), if there exists a strictly increasing sequence $(j_k)_k$ of positive integers such that

- (1) $||x_k|| \leq C$ for all k. (2) $\frac{1}{m_{j_{k+1}}} \cdot \# \operatorname{supp}(x_k) \leq \epsilon$ for every k.
- (3) For every k = 1, 2, ... and every $f \in W$ with $w(f) = m_i, i < j_k$ we have that $|f(x_k)| \leq C/m_i$.

Proposition 3.6. Let Y be a block subspace of X_W . Then for $\epsilon > 0$ there exists a normalized $(3, \epsilon)$ -RIS sequence $(x_k)_k$ with $x_k \in Y$ for every k.

Indeed, Lemma 3.3 yields that we can get a block sequence $(x_j)_j$ such that x_j is a normalized 2- $\ell_1^{n_{2j}}$ -average and $x_j \in Y$ for every k. Choose a subsequence $(x_{j_k})_k$ such that $\# \operatorname{supp}(x_{j_k}) \leq \epsilon m_{2j_{k+1}}$. Lemma 3.4 yields that $(x_{j_k})_k$ is a $(3, \epsilon)$ -RIS.

3.1. The auxiliary space. In order to get upper bounds on the action of weighted functionals on certain vectors we shall use an auxiliary space to reduce the high complexity of the calculations (see subsection 3.2). The auxiliary space X_{aux} is the mixed Tsirelson space $T[(\mathcal{A}_{4n_j}, m_j^{-1})_{j \in \mathbb{N}}].$

The result we shall use from the auxiliary space are the upper estimations of the functionals of its norming set W_{aux} on the averages of its basis, and in particular the following lemma.

Lemma 3.7. Let $x = \frac{m_j}{n_j} \sum_{k=1}^{n_j} e_k \in X_{aux}$. Then for every $f \in W_{aux}$ with $w(f) = m_i$,

$$\left| f(x) \right| \le \begin{cases} \frac{2}{m_i} & \text{if } i < j \\ \frac{m_j}{m_i} & \text{if } i \ge j. \end{cases}$$

Moreover if f has a tree analysis $(f_{\alpha})_{\alpha \in \mathcal{A}}$ such that for every weighted f_{α} , $w(f_{\alpha}) \neq m_j$ then $|f(x)| \leq \frac{2}{m_i^2}$.

For a proof of this result see [9], Lemma II.9, (or [4], Lemma 4.2).

Remark 3.8. Let us note that the above estimations hold also for n_j -averages of the basis of the space X_W .

3.2. The basic inequality. The basic inequality is the fundamental tool which permits us to show that the action of a weighted functional $f \in W$ on a n_j -average of a RIS admits similar upper bound as the action of a functional g in W_{aux} with the same weight with f, acting on a n_j -average of the basis of X_{aux} . In particular the following holds.

Proposition 3.9. Let $(x_k)_k$ be a (C, ϵ) -RIS in X_W and $(\lambda_k)_k \in c_{00}(\mathbb{N})$ be a sequence of scalars. Then for every $f \in W$ of type I we can find g_1 such that $g_1 = h_1$ or $g_1 = e_t^* + h_1$ with $t \notin \operatorname{supp}(h_1)$ where $h_1 \in W_{aux}$ with $w(h_1) = w(f)$ and $g_2 \in c_{00}(\mathbb{N})$ with $||g_2||_{\infty} \leq \epsilon$ with g_1, g_2 having nonnegative coordinates and such that

$$\left| f(\sum_{k} \lambda_k x_k) \right| \le C(g_1 + g_2) (\sum_{k} |\lambda_k| e_k).$$
(3.1)

If we additionally assume that there exists $j_0 \in \mathbb{N}$ such that for every $h \in X_W$ with $w(h) = m_{j_0}$ and every interval E of \mathbb{N} we have that

T

$$\left| h(\sum_{k \in E} \lambda_k x_k) \right| \le C(\max_{k \in E} |\lambda_k| + \epsilon \sum_{k \in E} |\lambda_k|), \tag{3.2}$$

then if $w(f) \neq m_{j_0}$, we may select h_1 to have a tree analysis $(h_t)_{t \in \mathcal{T}}$ with $w(h_t) \neq m_{j_0}$ for all $t \in \mathcal{T}$ with h_t a weighted functional.

We refer the reader to [9] Proposition II.14 or [4] Proposition 4.3 for a proof. As a first application of the basic inequality we get the following result concerning the action of weighted functionals on RIS.

Corollary 3.10. Let $(x_k)_{k=1}^{n_j}$ be a (C, m_j^{-2}) -RIS and $x = n_j^{-1} \sum_{k=1}^{n_j} x_k$. Then for $f \in W$ with $w(f) = m_i$ we have that

$$\left| f(x) \right| \le \begin{cases} \frac{3C}{m_i m_j} & \text{if } i < j \\ \frac{C}{m_i} + \frac{2C}{n_j} & \text{if } i \ge j \end{cases}.$$

$$(3.3)$$

In particular for every $j \in \mathbb{N}$, $m_{2j}^{-1} \le ||n_{2j}^{-1} \sum_{k=1}^{n_{2j}} x_k|| \le 2Cm_{2j}^{-1}$.

Т

Definition 3.11. A pair (x, f) with $x \in X_W$ and $f \in W$ is said to be a (C, j)-exact pair if the following conditions are satisfied:

- (1) $1 \leq ||x|| \leq C$, for every weighted functional $g \in W$ with $w(g) = m_i$, $i \neq j$ we have $|g(x)| \leq \frac{3C}{m_i}$ if i < j, while $|g(x)| \leq \frac{C}{m_i^2}$ if i > j.
- (2) f is weighted and $w(f) = m_j$.
- (3) f(x) = 1 and $\operatorname{ran}(f) \subset \operatorname{ran}(x)$.

The primitive example of a (2, 2j)-exact pair is

$$\left(\frac{m_{2j}}{n_{2j}}\sum_{k=1}^{n_{2j}}e_k, m_{2j}^{-1}\sum_{k=1}^{n_{2j}}e_k^*\right).$$
(3.4)

Indeed using Lemma 3.7 and Remark 3.8 it readily follows that the pair given by (3.4) is a (2,2j)-exact pair. Also for every block subspace Y of X_W we can find a (6,2j)-exact pair (y,f) with $y \in Y$. Indeed let $(x_k)_{k=1}^{n_{2j}}$ be a $(3,m_{2j}^{-3})$ -RIS where every $x_k \in Y$ is a $2 - \ell_1^{n_{2jk}}$ -average with $||x_k|| > 1$. Choose for every k a $f_k \in W$, such that $f_k(x_k) = 1$ and $\operatorname{ran}(f_k) \subset \operatorname{ran}(x_k)$. Then from Proposition 3.10 we get that that pair (y,f) where $y = \frac{m_{2j}}{n_{2j}} \sum_{k=1}^{n_{2j}} x_k$ and $f = m_{2j}^{-1} \sum_{k=1}^{n_{2j}} f_k$, is a (6,2j)-exact pair.

Definition 3.12. Let $C \ge 1$ and $j_0 \in \mathbb{N}$, $j_0 \ge 2$. We call a pair (x, f) for $x \in X_W$ and $f \in W$ a (C, j_0) -standard exact pair (SEP) if there exists a $(C, m_{j_0}^{-2})$ -RIS $(x_k)_{k=1}^{n_{j_0}}$ with

(1) $x = \frac{m_{j_0}}{n_{j_0}} \sum_{k=1}^{n_{j_0}} x_k.$ (2) $f = m_{j_0}^{-1} \sum_{k=1}^{n_{j_0}} f_k$ where $f_k \in W$ with $f_k(x_k) = 1$ and $\operatorname{ran}(f_k) = \operatorname{ran}(x_k).$ (3) x_k is a $\ell_1^{n_k}$ -average and $n_{j_0}^2 < m_{n_1}.$

Proposition 3.13. Let $j \in \mathbb{N}$ and $(x_i, f_i)_{i=1}^{n_{2j-1}} \subset X_W \times W$ such that $(x_i)_{i=1}^{n_{2j-1}}$ is a RIS and (x_i, f_i) is standard exact pair for every *i*. Let $x = n_{2j-1}^{-1} \sum_{i=1}^{n_{2j-1}} x_i \in X_W$ and assume that for every $g \in W$ with $w(g) = m_{2j-1}$ we have $|g(x)| \leq \frac{C}{m_{2j-1}^2}$. Then

$$\|x\|_W \le \frac{4C}{m_{2j-1}^2}.\tag{3.5}$$

This result follows from the basic inequality and especially the moreover part. We refer the reader to [4] Proposition 4.9 or [9] Proposition II.19 for a proof.

A consequence of the basic inequality is the reflexivity of the space X_W .

Proposition 3.14. The space X_W is reflexive.

We refer to [9] Proposition II.28, for a proof. As a consequence of the reflexivity of the space X_W and the rational convexity of the set W we get the following corollary.

Corollary 3.15. For the space X_W we have that $B_{X_{W}^*} = \overline{W}^{\|\cdot\|}$.

4. Standard HI spaces

In this section we show how we can refine W in order to get a norming set W_{hi} which will be the norming set of an HI-space. We start with the definition of the special sequences.

Let $\mathbb{Q}_s = \{(f_1, f_2, \dots, f_l) : f_i \in c_{00}(\mathbb{N}), f_i \neq 0, f_i(n) \in \mathbb{Q} \ \forall n \in \mathbb{N}, \forall i \leq l, f_1 < f_2 < \dots < f_l\}$. We fix a pair Ω_1, Ω_2 of disjoint infinite subsets of \mathbb{N} . From the fact

that \mathbb{Q}_s is countable there exists an injective coding function $\sigma : \mathbb{Q}_s \to \{2j : j \in \Omega_2\}$ such that

$$m_{\sigma(f_1, f_2, \dots, f_l)} > \max\{\frac{1}{|f_i(e_k)|} : k \in \operatorname{supp} f_i, i \le l\} \cdot \operatorname{maxsupp} f_l.$$
(4.1)

Definition 4.1. For $j \in \mathbb{N}$ and $k \leq n_{2j-1}$ a sequence (f_1, \ldots, f_k) of elements of W is said to be a (2j-1) special sequence if $w(f_1) = m_{2j_1}$ with $j_1 \in \Omega_1$, $m_{2j_1} \geq n_{2j-1}^2$ and $w(f_{i+1}) = m_{\sigma(f_1,\ldots,f_i)}$ for every $i \geq 1$.

Remark 4.2. The significant property of special sequences is the tree like behavior, i.e., if $j_1, j_2 \in \mathbb{N}$, $(f_i)_{i=1}^{n_{2j_1}-1}$ is an $(2j_1-1)$ -special sequence, $(g_i)_{i=1}^{n_{2j_2}-1}$ is an $(2j_2-1)$ -special sequence and if for some k, l we have $w(f_k) = w(g_l)$, then k = l and $f_i = g_i$ for every i < k. Moreover if $f_k \neq g_k$ then $w(f_{k+l}) \neq w(g_{k+i})$ for every $i, l \geq 1$.

4.1. The norming set of a standard HI-space. We consider the minimal subset W_{hi} of $c_{00}(\mathbb{N})$ satisfying the following

- (1) $\pm e_n^* \in W_{hi}$ for every $n \in \mathbb{N}$.
- (2) W_{hi} is closed in the $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})$ -operations.
- (3) W_{hi} is closed in the $(\mathcal{A}_{n_{2j-1}}, m_{2j-1}^{-1})$ -operations on special sequences, i.e., for every (2j-1)-special sequence (f_1, \ldots, f_k) with $f_i \in W_{hi}$ for every i, the functional $m_{2j-1}^{-1} \sum_{i=1}^k f_i \in W_{hi}$.
- (4) W_{hi} is symmetric and closed under the restriction of its elements to intervals of \mathbb{N} .
- (5) W_{hi} is closed under rational convex combinations.

The set W_{hi} is defined inductively as follows: Set $W_{hi}^0 = \{\pm e_n^* : n \in \mathbb{N}\}$ and assume that W_{hi}^n has been defined. To get W_{hi}^{n+1} , first we close the set of the functionals $\bigcup_{k=1}^n W_{hi}^k$ in the $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})$ -operations, and then we close the above set of functionals in the $(\mathcal{A}_{n_{2j-1}}, m_{2j-1}^{-1})$ operations on special sequences. The set W_{hi}^{n+1} is the set of the rational convex combinations of restrictions to intervals of the functionals we defined in the above two steps.

We denote by X_{hi} the completion of $c_{00}(\mathbb{N})$ under the norm induced by W_{hi} .

Note that by the definition of W_{hi} we get that it is an EO-complete subset of W_{mT} and hence all the results of the previous section hold for the space X_{hi} .

Definition 4.3. Let $j \in \mathbb{N}$. A sequence $(x_1, f_1, x_2, f_2, \ldots, x_{n_{2j-1}}, f_{n_{2j-1}})$ with $x_i \in X_{hi}$, $f_i \in W_{hi}$ is said to be (C, 2j - 1)-dependent if

- (1) $(f_1, \ldots, f_{n_{2j-1}})$ is a $W_{hi} (2j-1)$ special sequence
- (2) each pair (x_i, f_i) is a $(C, 2j_i)$ -SEP where $m_{2j_i} = w(f_i)$.

It follows from the definition of a (C, 2j - 1)-dependent sequence $(x_i, f_i)_{i=1}^{n_{2j-1}}$ that the block sequence $(x_i)_{i=1}^{n_{2j-1}}$ is a 3C-RIS.

Remark 4.4. From the results of the previous section we get that in every block subspace Y of X_{hi} we can find for every $j \in \mathbb{N}$ an (3, 2j)-SEP (x, f) with $x \in Y$. Hence if Y, Z are any two block subspaces of X_{hi} we can chooce a a (3, 2j - 1)dependent sequence $(x_i, f_i)_{i=1}^{n_{2j-1}}$ with $x_{2i-1} \in Y$ and $x_{2i} \in Z$ for every i.

Regarding the norm of the average $n_{2j-1}^{-1} \sum_{i=1}^{n_{2j-1}} x_i$ of the vectors of a (C, 2j-1)-dependent sequence we have the following result.

Proposition 4.5. Let $(x_1, f_1, x_2, f_2, ..., x_{n_{2j-1}}, f_{n_{2j-1}})$ be (C, 2j - 1)-dependent sequence. Then

$$m_{2j-1}^{-1} \le \|\sum_{i=1}^{n_{2j-1}} \frac{x_i}{n_{2j-1}}\| \quad and \quad \|\sum_{i=1}^{n_{2j-1}} (-1)^{i+1} \frac{x_i}{n_{2j-1}}\| \le 4Cm_{2j-1}^{-2}.$$
(4.2)

The first inequality in (4.2) follows easily. Indeed if $(x_1, f_1, x_2, f_2, \ldots, x_{n_{2j-1}}, f_{n_{2j-1}})$ is a (C, 2j - 1)-dependent sequence, the properties of the set W_{hi} yield that $F = m_{2j-1}^{-1} \sum_{i=1}^{n_{2j-1}} f_i \in W_{hi}$. Using this functional the first inequality in (4.2) readily follows. Note also $F(\sum_{i=1}^{n_{2j-1}} (-1)^{i+1})x_i) = 0$. It is proved also that

$$\left| g(\sum_{i=1}^{n_{2j-1}} (-1)^{i+1} x_i) \right| \le 2C(1+2m_{2j-1}^{-2}) \quad \text{for every } g \in W_{hi} \text{ with } w(g) = m_{2j-1}.$$

This inequality is a consequence of the following two facts:

The first one is the tree structure of the special sequences (see Remark 4.2). The second one is that every pair (x_k, f_k) of a dependent sequence $(x_i, f_i)_{i=1}^{n_{2j-1}}$ is a $(C, 2j_k)$ -SEP. This yields that for every $g \in W_{hi}$ with $w(g) \neq m_{2j_k}$ we have $|g(x_k)| \leq 3C \max\{w(g)^{-1}, m_{2j_k}^{-2}\}$ Combining the above two facts we get that the assumption of Proposition 3.13 holds. This gives us the second inequality in (4.2).

As a corollary of the previous proposition we get the following corollary.

Corollary 4.6. The space X_{hi} is an HI space.

Indeed let Y, Z be any two block subspaces of X_0 and $j \in \mathbb{N}$. From Remark 4.4 we get a (3, 2j - 1)-dependent sequence $(x_i, f_i)_{i=1}^{n_{2j-1}}$ with $x_{2i-1} \in Y$ and $x_{2i} \in Z$ for every *i*. From (4.2) we get that

$$\left\|\sum_{i=1}^{n_{2j-1}} (-1)^{i+1} x_i\right\| \le \frac{12}{m_{2j-1}} \left\|\sum_{i=1}^{n_{2j-1}} x_i\right\|.$$
(4.3)

Since $j \in \mathbb{N}$ the above inequality yields that $dist(S_Y, S_Z) = 0$ and this implies that X_0 is an HI space.

5. The definition of the space X_{ex} .

In this section we define the HI space X_{ex} which contains as an isometric subspace the initial HI space X_{hi} . For this we will define a norming set W_{ex} , a subset of $c_{00}(\mathbb{N})$, that induces the norm of the space. Actually the set W_{ex} is a subset of the set W_{mT} which is the norming set of the space $X_{mT} = T[(\mathcal{A}_{n_{2j}}, m_{2j}^{-1}) \cup (\mathcal{A}_{2n_{2j-1}}, m_{2j-1}^{-1})]$. Note that X_{mT} uses in the odd operations the families $\mathcal{A}_{2n_{2j-1}}$ instead of $\mathcal{A}_{n_{2j-1}}$ which were used in the definition of X_{hi} .

5.1. The definition of W_{ex} . Let $\mathbb{N} = \mathbb{N}_{od} \cup \mathbb{N}_{ev}$, \mathbb{N}_{od} the odd and \mathbb{N}_{ev} the even positive integers.

We shall denote also by $K = \bigcup_j K^{2j}$ the subset of W_{mT} , where

$$K^{2j} = \{m_{2j}^{-1} \sum_{i \in F} e_i^* : \#F = n_{2j}\}.$$

For every j, let $(K_k^{2j})_{k\in\mathbb{N}}$, be disjoint subsets of K^{2j} such that $\sup\{\min(f) : f \in K_k^{2j}\} = +\infty$ for every k. For $g = \sum_{i=1}^{+\infty} a_i e_i^* \in c_{00}(\mathbb{N})$ we denote by \widehat{g} the vector

 $\widehat{g} = \sum_{i} a_i e_{2i}^* \in c_{00}(\mathbb{N})$. We shall denote by $\widehat{K_k^{2j}}$ the set $\{\widehat{f} : f \in K_k^{2j}\}$ and by \widehat{W}_{hi} the set $\{\widehat{f}: f \in W_{hi}\}$.

Definition 5.1 (The coding functions $\varrho_k^{2i}, k \in \mathbb{N}, i = 0, 1, ...$). Let $(D_k)_{k \in \mathbb{N}}$ be a sequence of disjoint subsets of W_{mT} , to be determined later. For every $i, k \in \mathbb{N}$ we shall denote by L_k^{2i} the following subset of D_k ,

$$L_k^{2i} = \{ f \in D_k : w(f) = m_{2i} \text{ and } \operatorname{supp}(f) \subset \mathbb{N}_{od} \}.$$

For every $k, i \in \mathbb{N}$ let $\rho_k^{2i} : L_k^{2i} \to K_k^{2i}$ be an injection such that

10

for every
$$f \in L_k^{2i}$$
 we have $f < \widehat{\varrho_k^{2i}}(\widehat{f})$. (5.1)

Remark 5.2. From the above definitions it follows that if $g \in K_k^{2i}$ for some $i, k \in \mathbb{N}$ then there is no $n \neq k$ and $f \in L_n^{2i}$ such that $g = \varrho_n^{2i}(f) \in K_n^{2i}$. This follows from the fact that the domains of $\varrho_k^{2i}, \varrho_n^{2i}$ are disjoint as well as their

ranges K_k^{2i}, K_n^{2i} .

Our goal in this subsection is to define an EO-complete subset W_{ex} of W_{mT} such that the restriction of every element $f \in W_{ex}$ to the set \mathbb{N}_{ev} belongs to W_{hi} and in particular $\widehat{W}_{hi} = \{f_{|\mathbb{N}_{ev}} : f \in W_{ex}\}$. Denoting by X_{ex} the space defined by the set W_{ex} , the aforementioned property yields that its subspace \hat{X}_{hi} generated by the subsequence $(e_{2n})_{n \in \mathbb{N}}$ of the basis is isometric to the initial HI space X_{hi} . We shall also include in W_{ex} some new special functionals, named paired special functionals which will permit us to export the HI structure of the space X_{hi} to the whole space X_{ex} .

We proceed to give three definitions of special functionals which will be used in the definition of the norming W_{ex} . We assume the existence of a family $(D_k)_{k\in\mathbb{N}}$ of disjoint subsets of W_{mT} that will be specified later and we consider the corresponding functions $\varrho_k^{2i}: L_k^{2i} \to K_k^{2i}, k \in \mathbb{N}, i \geq 0$ as they were defined in Definition 5.1

Definition 5.3. Let $j, n \in \mathbb{N}$ and $V \subset W_{mT}$. We call a $(2j-1, V, (\varrho_k^{2i})_{k=1}^n, i \geq 0)$ paired special sequence any block sequence $(f_1, \hat{g}_1, \ldots, f_p, \hat{g}_p)$ satisfying the following:

- (1) p ≤ n_{2j-1} and for every k ≤ p there exist j_k ∈ N and i_k ≤ n such that f_k ∈ V ∩ L^{2j_k}_{i_k} and g_k = ϱ^{2j_k}_{i_k}(f_k).
 (2) The finite sequence (g₁, g₂, ..., g_p) is a (2j 1)-special sequence of W_{hi}.

Definition 5.4. Let $j, n \in \mathbb{N}$ and $V \subset W_{mT}$. We call a $(2j-1, V, (\varrho_k^{2i})_{k=1}^n), i \geq 0)$ semi paired special sequence any block sequence of the form $(f_i, \widehat{g}_i)_{i=1}^l \cap (\widehat{g}_i)_{i=l+1}^p$ satisfying the following:

- (1) $p \leq n_{2j-1}, l \geq 0$ and $(f_1, \hat{g}_1 f_2, \hat{g}_2, \dots, f_l, \hat{g}_l)$ is a $(2j-1, V, (\varrho_i^{2k})_{i=1}^n, k =$ $0, 1, \ldots$)-paired special sequence.
- (2) For every m = l + 1, ..., p we have that $g_m \in K_{i_m}^{2j_m}$ for some $i_m \leq n$, and $(g_1, g_2, ..., g_p)$ is a (2j 1)- special sequence of W_{hi}^1 .
- (3) There is no $\phi \in V$ such that $(f_1, \hat{g}_1 f_2, \hat{g}_2, \dots, f_l, \hat{g}_l, \phi, \hat{g}_{l+1})$ is a (2j i)1, $V, (\varrho_k^{2i})_{k=1}^n, i = 0, 1, ...)$ paired special sequence.

The third definition is the following.

Definition 5.5. Let $j, n \in \mathbb{N}$ and $V \subset W_{mT}$. We call a $(2j-1, V, (\varrho_k^{2i})_{k=1}^n, i \geq 0)$ special sequence any block sequence $(f_1, \hat{g}_1, \ldots, f_l, \hat{g}_l, \hat{g}_{l+1}, \ldots, \hat{g}_m, h_{m+1}, h_{m+2}, \ldots, h_p)$ satisfying the following:

- (1) $p \leq n_{2j-1}, 0 \leq l \leq m$ and $(f_1, \hat{g}_1, \ldots, f_l, \hat{g}_l, \hat{g}_{l+1}, \ldots, \hat{g}_m)$ is a $(2j 1, V, (g_k^{2i})_{k=1}^n, i = 0, 1, \ldots)$ -semi paired special sequence.
- (2) $h_{m+1|\mathbb{N}_{ev}} \in V \setminus \bigcup_{j=0}^{+\infty} \bigcup_{i=1}^{n} \widehat{K}_{i}^{2j}$ and for every $q \ge 1$, $h_{m+q|\mathbb{N}_{ev}} = \widehat{\phi}_{m+q} \in \widehat{W}_{hi}$ and $(g_1, \ldots, g_m, \phi_{m+1}, \phi_{m+2}, \ldots, \phi_p)$ is a (2j-1)-special sequence of W_{hi} .

Remark 5.6. It follows from the above definitions that the paired special sequences are semi paired special sequences and the later are special sequences.

5.2. Definition of the norming set W_{ex} . We proceed by induction to define an increasing family of sets W_n , n = 0, 1, 2, ... which will be subsets of W_{mT} such that $W_{ex} = \bigcup_{n=0}^{\infty} W_n$. For every $n \ge 1$ we shall choose a set $D_n \subset W_n$ such that the sets $(D_n)_{n \in \mathbb{N}}$ are disjoint and for every $n \in \mathbb{N}$, $i \ge 0$, we will consider the injective function $\varrho_n^{2i} : L_n^{2i} \to K_n^{2i}$ as it was defined in Definition 5.1.

Let $W_0 = \{\pm e_n^* : n \in \mathbb{N}\}$. Let $n \in \mathbb{N}$ and assume that for every k < n the sets $D_k \subset W_k$ and the functions ϱ_k^{2i} , $i \ge 0$ have been defined.

We set

$$U_n = \bigcup_j \{ m_{2j}^{-1} \sum_{i=1}^k f_i : (f_i)_{i=1}^k \text{ is an } \mathcal{A}_{n_{2j}} \text{-admissible sequence of } W_{n-1} \},$$

and $V_n = (U_n \setminus \bigcup_{j=0}^{\infty} \bigcup_{i=n+1}^{+\infty} \widehat{K_i^{2j}}) \cup W_{n-1}.$ We set $D_n = V_n \setminus W_{n-1}$ and for $i \ge 0$ we define

$$L_n^{2i} = \{ f \in D_n : w(f) = m_{2i} \text{ and } \operatorname{supp}(f) \subset \mathbb{N}_{od} \}.$$

For every $i \in \mathbb{N}$ we a choose an injective function $\varrho_n^{2i} : L_n^{2i} \to K_n^{2i}$ satisfying $f < \widehat{\varrho_n^{2i}(f)}$ for every $f \in L_n^{2i}$.

It is clear that $D_n \cap W_{n-1} = \emptyset$ and hence D_n is disjoint from every W_k , k < n. Also we have that $L_n^{2i} \subset V_n$ for every $i \ge 0$.

We define now for every $j \in \mathbb{N}$ the $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i \ge 0)$ special functionals. For $j \in \mathbb{N}$ we define

$$W_n^{2j-1} = \left\{ \frac{1}{m_{2j-1}} \left(\sum_{i=1}^l (f_i + \hat{g}_i) + \sum_{i=l+1}^m \hat{g}_i + \sum_{i=m+1}^p h_i \right) :$$
(5.2)

 $(f_1, \widehat{g}_1, \dots, f_l, \widehat{g}_l, \widehat{g}_{l+1}, \dots, \widehat{g}_m, h_{m+1}, \dots, h_p) \text{ is a } (2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i=0, 1, \dots)$ special sequence.}

We set

$$W_n = \operatorname{conv}_{\mathbb{Q}} \{ Ef : f \in V_n \cup \bigcup_{j=1}^{\infty} W_n^{2j-1} : E \text{ interval of } \mathbb{N} \}.$$

The norming set W_{ex} is the set $W_{ex} = \bigcup_{n=0}^{+\infty} W_n$.

From the inductive assumption we have that W_{n-1} is a subset of set W_{mT} . It follows that the set U_n is also subset of W_{mT} . Hence for every $j \in \mathbb{N}$ the set W_n^{2j-1} is a subset of W_{mT} and finally the set W_n is subset of W_{mT} . It is clear also that the sets W_{n-1} and D_n are subsets of W_n . Remarks 5.7.

a) If $(f_1, \hat{g}_1, f_2, \hat{g}_2, f_l, \hat{g}_l, \hat{g}_{l+1}, \hat{g}_{l+2}, \dots, \hat{g}_p)$ is a $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i = 0, 1, \dots)$ semi-paired special and $l \ge 1$, then for the functional \hat{g}_{l+1} we have that i) either $g_{l+1} \notin \bigcup_{k=1}^n \varrho_k^{2j_{l+1}}(L_k^{2j_{l+1}}), w(g_{l+1}) = m_{2j_{l+1}},$ SPIROS A. ARGYROS, ANTONIS MANOUSSAKIS, AND PAVLOS MOTAKIS

ii) or $g_{l+1} = \rho_{k_{l+1}}^{2j_{l+1}}(f_{l+1})$ for an $f_{l+1} \in L_{k_{l+1}}^{2j_{l+1}}, k_{l+1} \le n$ and minsupp $f_{l+1} \le \max \rho_{k_{l+1}}$

Note that in both cases the definition of the coding functions ϱ_k^{2i} and the disjointness of the sets K_k^{2i} implies that there is no m > n and a $(2j-1, V_m, (\varrho_k^{2i})_{k=0}^m, i \ge 0)$ - paired special sequence having as an initial segment the sequence $(f_i, \hat{g}_i)_{i=1}^{l+1}$.

- segment the sequence $(f_i, \hat{g}_i)_{i=1}^{l+1}$. b) For every $g_1 \in \varrho_{i_1}^{2j_1}(L_{i_1}^{2j_1})$ with $w(g_1) = m_{2j_1} \ge n_{2j-1}^2$ for some $j \in \mathbb{N}$ and $i_1 \le n$, we have that $((\varrho_{i_1}^{2j_1})^{-1}(g_1), \hat{g}_1)$ is a $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i = 0, 1, ...)$ paired sequence.
- c) If (g_1, \ldots, g_p) is a (2j-1)-special sequence in W_{hi} , such that for every $l \leq p, g_l \in K^{2j_l}$ for some $4j_l \in \mathbb{N}$ then if $\widehat{g}_1 \notin \bigcup_{k=1}^{\infty} K_k^{2j_1}$ it follows that $(\widehat{g}_1, \ldots, \widehat{g}_p)$ is a $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i=0,1,\ldots)$ special sequence where $n = \max\{1, k : \exists l \leq p \text{ with } g_l \in K_k^{2j_l}\}.$

If $g_1 \in K_{k_1}^{2j_1} \setminus \varrho_{k_1}^{2j_1}(L_{k_1}^{2j_1})$ then $(\widehat{g}_1, \ldots, \widehat{g}_p)$ is a $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i = 0, 1, \ldots)$ -semi paired special sequence where $n = \max\{k : \exists l \leq p \text{ with } g_l \in K_k^{2j_l}\}$.

It follows form the definition of W_{ex} that it is an EO-complete subset of W_{mt} . Hence from the results of Section 3 we have the following.

Proposition 5.8. The space X_{ex} is reflexive. Moreover $B_{X_{ex}^*} = \overline{W_{ex}}^{\parallel\parallel}$.

We show now that the semi paired special sequences have the tree property, a result that will be essential in the next sections.

Notation 5.9. If $(f_1, \hat{g}_1, f_2, \hat{g}_2, f_l, \hat{g}_l, \hat{g}_{l+1}, \ldots, \hat{g}_m)$ is a $(2j - 1, V_q, (\varrho_k^{2i})_{k=1}^q, i = 0, 1, \ldots)$ -semi paired special sequence, the pair defined by g_i , $i \leq m$ is defined to be the pair $\{f_i, g_i\}$ if $i \leq l$ and the singleton $\{g_i\}$ if $i = l + 1, \ldots, m$. The weight of the pair is defined to be the weight of g_i .

Lemma 5.10. The set of the semi paired special sequences has a tree structure, i.e., if $(f_i^1, \hat{g}_i^1)_{i=1}^{p_1} \frown (\hat{g}_i^1)_{i=p_1+1}^{m_1}$ and $(f_i^2, \hat{g}_i^2)_{i=1}^{p_2} \frown (\hat{g}_i^2)_{i=p_2+1}^{m_2}$ are two semi-paired special sequences and the couples determined by some g_i^1, g_j^2 have the same weight then i = j and then couples determined by g_l^1 and g_l^2 are equal for all $l \leq i$. Moreover if $g_i^1 \neq g_i^2$ then all the couples determined by $g_k^1, g_p^2, k, p > i$ have different weights.

Proof. By the definition of the semi-paired special sequences it follows that $(g_l^1)_{l=1}^i$ and $(g_k^2)_{k=1}^j$ are special sequences of W_{hi}^1 . Since the special sequences of W_{hi} have the tree property and $w(g_i^1) = w(g_j^2)$ it follows that i = j and $g_k^1 = g_k^2$ for all k < i = j. Also from the definition of the (semi)-paired special sequences and the injectivity of the functions $g_n^{2j}, n \in \mathbb{N}, j \ge 0$ it follows that the couples determined by $g_k^1, g_k^2, k < i$, are equal and in particular they have the same cardinality.

The same holds of $g_i^1 = g_i^2$. In the case $g_i^1 \neq g_i^2$ again from the tree property of the special sequences of W_{hi} it follows that $w(g_k^1) \neq w(g_l^2)$ for all k, l > i.

In the case $i = j = l_1$ (or $i = j = l_2$) we have that $l_1 - 1 \le l_2$ ($l_2 - 1 \le l_1$).

We proceed to show that $\widehat{W}_{hi} = \{f_{|\mathbb{N}_{ev}} : f \in W_{ex}\}$. First we show that $W_{ex|\mathbb{N}_{ev}}$ is a subset of \widehat{W}_{hi} . This will be an immediate consequence of the following lemma. Lemma 5.11. For every $n \in \mathbb{N}$ and every $f \in W_n$ we have that $f_{|\mathbb{N}_{ev}} \in \widehat{W}_{hi}$.

12

Proof. We prove the lemma by induction. For every $n \in \mathbb{N}$, $e_{n|\mathbb{N}_{ev}}^* = 0$ if n is odd otherwise if n = 2k, $e_{n|\mathbb{N}_{ev}}^* = e_n^* = \widehat{e_k^*} \in \widehat{W}_{hi}$. Assume that the result holds for all $k \leq n$ and $f \in W_{n+1}$.

Let f be the result of an $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})$ - operation of elements of W_{n-1} , i.e., f = $m_{2i}^{-1} \sum_{i=1}^{d} f_i$. Then from the inductive hypothesis we have that $f_{i|\mathbb{N}_{ev}} = \widehat{\phi}_i \in \widehat{W}_{hi}$ for every *i*. Since W_{hi} is closed in the $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})$ -operations it follows that $\phi =$ $m_{2j}^{-1} \sum_{i=1}^{d} \phi_i \in W_{hi}$ and $f_{|\mathbb{N}_{ev}} = \widehat{\phi} \in \widehat{W}_{hi}$. Assume now that there exists a $(2j - 1, V_n, (\varrho_k^{2i})_{k=1}^n, i = 0, 1, \dots)$ -special se-

quence $(f_1, \hat{g}_1, f_l, \hat{g}_l, \hat{g}_{l+1}, \dots, \hat{g}_m, h_{m+1}, \dots, h_p)$ such that $f = m_{2j-1}^{-1} (\sum_{i=1}^l (f_i + \hat{g}_i) + \sum_{i=l+1}^m \hat{g}_i + \sum_{i=m+1}^p h_i)$. By the definition of the special sequences we have that $f_{i|\mathbb{N}_{ev}} = 0$ for every $i \leq l$ and $h_{i|\mathbb{N}_{ev}} = \widehat{\phi}_i \in \widehat{W}_{hi}$ for every $i \geq m+1$. Moreover $(g_1, \ldots, g_m, \phi_{m+1}, \cdots \phi_p)$ is a (2j-1)-special sequence of W_{hi} . It follows that

$$\phi = m_{2j-1}^{-1} (\sum_{i=1}^{m} g_i + \sum_{i=m+1}^{p} \phi_i) \in W_{hi} \text{ and } f_{|\mathbb{N}_{ev}} = m_{2j-1}^{-1} (\sum_{i=1}^{m} \widehat{g}_i + \sum_{i=m+1}^{p} h_{i|\mathbb{N}_{ev}}) = \widehat{\phi}_{i}$$

The case f is rational convex combination of functionals of the first two cases is immediate consequence of the fact that W_{hi} is closed in rational convex combinations. This completes the proof. \square

The next lemma will be used to prove the converse inclusive relation.

Lemma 5.12. Let $g_i \in K_{n_i}^{2j_i}$, $i \leq m$ be such that (g_1, \ldots, g_m) is a (2j - 1)-special sequence of W_{hi} . Then there exist $0 \leq l \leq m$ and $f_1, \ldots, f_l \in W_{ex}$ such that $(f_1, \hat{g}_1, \dots, f_l, \hat{g}_l, \hat{g}_{l+1}, \dots, \hat{g}_m)$ is a $(2j - 1, V_n, (\varrho_k^n)_{k=1}^n, i = 0, \dots)$ semi-paired special sequence where $n = \max\{n_i : i \leq m\}$.

Proof. Let $l \leq m$ be maximal with the properties

- (1) $g_i \in \varrho_{n_i}^{2j_i}(L_{n_i}^{2j_i})$ for every $i \le l$. (2) $F = ((\varrho_{n_i}^{2j_i})^{-1}(g_i), \widehat{g}_i)_{i=1}^l$ is a $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i = 0, 1...)$ -paired special sequence, where $n = \max\{n_i : i \le m\}$.

If l = 0, Remark 5.7b) yields that $g_1 \notin \varrho_{n_1}^{2j_1}(L_{n_1}^{2j_1})$ and from Remark 5.7c) we get that $(\widehat{g}_1, \widehat{g}_2, \dots, \widehat{g}_m)$ is $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i=0, 1, \dots,)$ -semi-paired special sequence.

If l = m we have that $((\varrho_{n_i}^{2j_i})^{-1}(g_i), \widehat{g}_i)_{i=1}^m$ is $(2j - 1, V_n, (\varrho_k^{2i})_{k=1}^n, i = 0, 1...)$ paired special sequence.

Let l < m. Then for the functional g_{l+1} we have that either does not belong to $p_{n_{l+1}}^{2j_{l+1}}(L_{n_{l+1}}^{2j_{l+1}})$ or belongs to $p_{n_{l+1}}^{2j_{l+1}}(L_{n_{l+1}}^{2j_{l+1}})$ and $\min(p_{n_{l+1}}^{2i_{l+1}})^{-1}(g_{l+1}) \leq \max \widehat{g}_l$. In both cases we get that

$$((\varrho_1^{2j_1})^{-1}(g_1), \widehat{g}_1, \dots, (\varrho_l^{2j_1})^{-1}(g_l), \widehat{g}_l, \widehat{g}_{l+1}, \dots, \widehat{g}_m)$$

is $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i=0,1...)$ semi-paired special sequence.

Lemma 5.13. For every $g \in W_{hi}$ there exists $\phi \in W_{ex}$ such that $\phi_{|\mathbb{N}_{ey}} = \hat{g}$.

Proof. We shall prove by induction on n that for every $g \in W_{hi}^n$ there exists $\phi \in W_{hi}^n$ $W_{ex}^{n_0}, n_0 = n_0(g) \ge n$, such that $\phi_{|\mathbb{N}_{ev}} = \widehat{g}$.

The case $g = \pm e_k^* \in W_{hi}^0$ it trivial. Let $g \in W_{hi}^1$. We shall consider the following two cases.

Case 1. $g = m_{2j}^{-1} \sum_{i \in F} \pm e_i^*$ with $\#F \le n_{2j}$.

We have to consider the following two subcases.

Subcase a). $g \notin \bigcup_{n=2}^{\infty} K_n^{2j}$. In this subcase we get that $\phi = m_{2j}^{-1} \sum_{i \in F} \pm e_{2i}^* \in V_1$ and $\phi = \phi_{|\mathbb{N}_{ev}} = \widehat{g}$.

Subcase b). $g \in K_n^{2j}$ for some $n \ge 2$. In this subcase we get that $\phi = m_{2j}^{-1} \sum_{i \in F} e_{2i}^* \in V_n \subset W_n$ and $\phi = \widehat{g}$. We pass now to the next case.

Case 2. Let $g = m_{2j-1}^{-1} \sum_{i=1}^{p} g_i$ where $g_i \in W_{hi}^1$ and (g_1, \ldots, g_p) is a (2j-1)special sequence.

Let $m \leq p$ be maximal with the property $g_k \in K_{i_k}^{2j_k}$ for every $k \leq m$. If m = 0 from 5.7c) we get that $(\widehat{g}_1, \ldots, \widehat{g}_m)$ is $(2j-1, V_{n_0}, (\varrho_k^{2i})_{k=1}^{n_0}, i = 0, \ldots)$ special sequence for some n.

If $m \geq 1$, Lemma 5.12 yields that there exist $l \leq m, f_1, \ldots, f_l \in W_{ex}$ and $n_0 \in \mathbb{N}$ such that $(f_1, \hat{g}_1, \dots, f_l, \hat{g}_l, \hat{g}_{l+1}, \dots, \hat{g}_m)$ is a $(2j - 1, V_{n_0}, (\varrho_k^{2i})_{k=1}^{n_0}, i = 0, \dots)$ semipaired special sequence. Clearly by the definition of the $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i =$ $(0,\ldots)$ special sequences, we get that $(f_1,\widehat{g}_1,\ldots,f_l,\widehat{g}_l,\widehat{g}_{l+1},\ldots,\widehat{g}_p)$ is $(2j-1,V_n,(\varrho_k^{2i})_{k=1}^n,i=1,\ldots,\widehat{g}_p)$ $(0,\ldots)$ special sequence where $n = \max\{r_i : g_i \in K_{r_i}^{2j_i}, i \in \{1,\ldots,p\}\}$. It follows

$$\phi = m_{2j-1}^{-1} \left(\sum_{i=1}^{l} (f_i + \widehat{g}_i) + \sum_{i=l+1}^{p} \widehat{g}_i \right) \in W_{ex} \text{ and } \phi_{|\mathbb{N}_{ev}} = m_{2j-1}^{-1} \sum_{i=1}^{d} \widehat{g}_i = \widehat{g}.$$

The case q is rational convex combination of functional of the above two cases follows readily. This complete the first inductive step.

Assume that we have prove the result for every $k \leq n$ and let $g \in W_{hi}^{n+1}$.

The case g is that result of an $(\mathcal{A}_{n_{2i}}, m_{2i}^{-1})$ -operation of elements of W_n^{hi} follows easily from the inductive hypothesis. Indeed if $g = m_{2j}^{-1} \sum_{i=1}^{d} g_i$ then for every i we have that there exists $\phi_i \in W_{ex}$ such that $\phi_{i|\mathbb{N}_{ev}} = \widehat{g}_i$. Since W_{ex} is closed in projections on intervals we may assume that $ran(\phi_i) = ran(\hat{g}_i)$. Also using that W_{ex} is closed in $(\mathcal{A}_{n_{2j}}, m_{2j}^{-1})$ -operations we get

$$\phi = m_{2j}^{-1} \sum_{i=1}^{d} \phi_i \in W_{ex} \text{ and } \phi_{|\mathbb{N}_{ev}} = \widehat{g}$$

Assume now that $g = m_{2j-1}^{-1} \sum_{i=1}^{p} g_i$ where (g_1, \ldots, g_p) is a (2j-1)-special sequence of W_{hi}^{n+1}

Let $m \leq p$ be maximal with respect the property $g_k \in K_{i_k}^{2j_k}$ for some i_k , for every $k \leq m$. Then from Lemma 5.12 we get that there exist $l \leq m, f_1, \ldots, f_l \in W_{ex}$ and $n_0 \in \mathbb{N}$ such that $(f_1, \widehat{g}_1, \ldots, f_l, \widehat{g}_l, \widehat{g}_{l+1}, \ldots, \widehat{g}_m)$ is a $(2j-1, V_{n_0}, (\varrho_k^{2i})_{k=1}^{n_0}, i=0, \ldots)$ semi-paired special sequence. For every i > m from the inductive hypothesis we have that there exists $h_i \in W_{ex}^{n_i}$ such that $h_{i|\mathbb{N}_{ev}} = \widehat{g}_i$. Since W_{ex} is closed in projections on intervals we may assume that $ran(h_i) = ran(\hat{g}_i)$ for every *i*. Moreover from the choice of m we have also that $h_{m+1|\mathbb{N}_{ev}} = \widehat{g}_{m+1} \notin \bigcup_{k=1}^{\infty} \widehat{K}_k^{2j_{m+1}}$.

It follows that $(f_1, \widehat{g}_1, \ldots, f_l, \widehat{g}_l, \widehat{g}_{l+1}, \ldots, \widehat{g}_m, h_{m+1}, \cdots, h_p)$ is a $(2j-1, V_{\overline{n}_0}, (\varrho_k^{2i})_{k=1}^{\overline{n}_0}, i = 0$ $0,\ldots$) special sequence for some $\bar{n}_0 \geq n$. Hence

$$\phi = m_{2j-1}^{-1} \left(\sum_{i=1}^{l} (f_i + \widehat{g}_i) + \sum_{i=l+1}^{m} \widehat{g}_i + \sum_{i=m+1}^{p} h_i \right) \in W_{ex} \text{ and } \phi_{|\mathbb{N}_{ev}} = \widehat{g}.$$

14

As a corollary we get the following.

Proposition 5.14. For the norming sets W_{hi} and W_{ex} of the spaces X_{hi} and X_{ex} we have that

$$W_{ex|\mathbb{N}_{ev}} = \widehat{W}_{hi}.$$

Morever the space X_{hi} embeds isometrically into X_{ex} .

6. A CRIRETION FOR THE HI PROPERTY

In this section we provide a criterion for a Banach space to be an HI space. It is equivalent to [11, Lemma 7.1], but we provide a proof for completeness.

Proposition 6.1. Let X be a Banach space and X_0 be an HI subspace of X. If for every subspace Z of X we have $dist(S_{X_0}, S_Z) = 0$ then the space X is HI.

Proof. We show that for every two infinite dimensional closed subspaces Y_1, Y_2 of X we have $dist(S_{Y_1}, S_{Y_2}) = 0$. Let $\epsilon \in (0, 1/10)$. We start with the following observation.

In every subspace Y of X we can find a Schauder basic sequence $(y_n)_{n\in\mathbb{N}} \subset Y$ and a normalized sequence $(x_n)_{n\in\mathbb{N}} \subset S_{X_0}$ such that $||y_n - x_n|| < \frac{\epsilon}{2^{n+3}}$. Indeed using Mazur Theorem [18] we get a normalized Schauder basic sequence $(u_n)_n$ in Y with basis constant $K \leq 2$. Using the hypothesis by a standard inductive argument we get a block sequence $(y_n)_{n\in\mathbb{N}}$ of $(u_n)_n$ and a normalized sequence $(x_n)_{n\in\mathbb{N}} \subset S_{X_0}$ such that $||y_n - x_n|| < \frac{\epsilon}{2^{n+3}}$. This implies that $(x_n)_n$ is also Schauder basic sequence.

By the above observation we find Schauder basic sequences $(y_n^1)_n \,\subset Y_1, (y_n^2) \,\subset Y_2$ and normalized Schauder basic sequences $(x_n^1)_n, (x_n^2)_n$ in X_0 such that $||y_n^i - x_n^i|| < \epsilon/2^{n+3}$ for every n and i = 1, 2. Since the subspace X_0 is HI, there exist normalized vectors $x_1 \in [x_n^1 : n \in \mathbb{N}]$ and $x_2 \in [x_n^2 : n \in \mathbb{N}]$ such that $||x_1 - x_2|| < \epsilon/3$. Since $(x_n^1)_n$ and $(x_n^2)_n$ are Schauder basic sequences by $\epsilon/3$ perturbation we may assume that there exists $n_1 \in \mathbb{N}$ such that $x_1 \in [x_n^1 : n \le n_1]$ and $x_2 \in [x_n^2 : n \le n_1]$. Let $x_1 = \sum_{i=1}^{n_1} a_n x_n^1$ and $x_2 = \sum_{i=1}^{n_1} b_n x_n^2$. Then

$$\begin{split} \|\sum_{i=1}^{n_1} a_n y_n^1 - \sum_{i=1}^{n_1} b_n y_n^2 \| &\leq \|x_1 - x_2\| + \|\sum_{i=1}^{n_1} a_n y_n^1 - \sum_{i=1}^{n_1} a_n x_n^1 \| + \|\sum_{i=1}^{n_1} b_n x_n^2 - \sum_{i=1}^{n_1} b_n y_n^2 \| \\ &\leq 3\frac{\epsilon}{3} + \max_n |a_n| \frac{\epsilon}{2^3} + \max_n |b_n| \frac{\epsilon}{2^3} < 3\epsilon. \end{split}$$

Using once more that $||x_n^i - y_n^i|| < \epsilon/2^{n+3}$ we get that $\operatorname{dist}(S_{Y_1}, S_{Y_2}) < 10\epsilon$. This completes the proof that $\operatorname{dist}(S_{Y_1}, S_{Y_2}) = 0$ and therefore X is HI.

7. The space X_{ex} is HI

To show that the space X_{ex} is HI we will use Proposition 6.1 and as X_0 the space \hat{X}_{hi} . In order to show that the assumption of the lemma holds we will use the RISs and the standard exact pairs as we did for the space X_{hi} .

7.1. Dependent paired sequences. In this subsection we shall define the dependent paired sequences which is the analogue of the dependent sequences defined for the original space X_{ex} and we will provide upper estimations for the vectors defined by them. The definition of the (C, ϵ) -RIS is the same as in Definition 3.5 with the obvious modifications. Similar the definition of the (C, j_0) -standard exact pair (SEP) is as in Definition 3.12.

To get upper bounds for the estimation of a weighted functional on a (C, ϵ) -RIS we shall use the basic inequality and a slight modification of the auxiliary space. The new auxiliary space is the mixed Tsirelson space $X_{aux2} = T[(\mathcal{A}_{12n_j}, m_j^{-1})]_{j \in \mathbb{N}}$. We denote by W_{aux2} its norming set. The choice of the sequences $(m_j)_{j \in \mathbb{N}}$, $(n_j)_{j \in \mathbb{N}}$ gives us that on the averages of the basis we have similar estimations as in Lemma 3.7. Namely the following holds.

Lemma 7.1. Let $x = \frac{m_j}{n_j} \sum_{k=1}^{n_j} e_k \in X_{aux2}$. Then for every $f \in W_{aux2}$ with $w(f) = m_i.$

$$\left| f(x) \right| \le \begin{cases} \frac{2}{m_i} & \text{if } i < j \\ \frac{m_j}{m_i} & \text{if } i \ge j. \end{cases}$$

Moreover if f has a tree analysis $(f_{\alpha})_{\alpha \in \mathcal{A}}$ such that for every weighted f_{α} we have $w(f_{\alpha}) \neq m_j \ then \left| f(x) \right| \leq \frac{2}{m_j^2}.$

For a proof of this result see [9], Lemma II.9, (or [4], Lemma 4.2).

The upper bounds of the action of a weighted functional on averages of a (C, ϵ) -RIS in the space X_{ex} are the same as in Proposition 3.10. For sake of completeness we state it again.

Proposition 7.2. Let $(x_k)_{k=1}^{n_j}$ be a (C, m_j^{-2}) -RIS in X_{ex} and $x = n_j^{-1} \sum_{k=1}^{n_j} x_k$. Then for every $f \in W_{ex}$ with $w(f) = m_i$ we have

$$\left|f(x)\right| \le \begin{cases} \frac{3C}{m_i m_j} & \text{if } i < j\\ \frac{C}{m_i} + \frac{2C}{n_j} & \text{if } i \ge j \end{cases}.$$
(7.1)

To show that the assumptions of the Lemma 6.1 hold we shall use the notion of dependent paired sequences.

Definition 7.3. A double sequence $(x_k, f_k)_{k=1}^{2n_{2j-1}}$ is said to be a (C, 2j-1)-dependent paired sequence if there exist $j_k, k \leq n_{2j-1}$ such that

- (1) $x_{2k} = \frac{m_{2j_k}}{n_{2j_k}} \sum_{i \in F_k} e_i, \#F_k = n_{2j_k} \text{ and } F_k \subset \mathbb{N}_{ev}, \text{ and } f_{2k} = m_{2j_k}^{-1} \sum_{i \in F_k} e_i^*.$ (2) (x_{2k-1}, f_{2k-1}) is a $(C, 2j_k)$ -standard exact pair with $\operatorname{supp}(f_{2k-1}) \subset \mathbb{N}_{od}$ for
- every k. (3) $(f_i)_{i=1}^{2n_{2j-1}}$ is a (2j-1)-special sequence of W_{ex} with $w(f_{2k-1}) = m_{2j_k} = w(f_{2k})$ for every k.

Let us note that if $(x_k, f_k)_{k=1}^{2n_{2j-1}}$ is a (C, 2j-1)-dependent paired sequence then the functional $F = m_{2j-1}^{-1} \sum_{k=1}^{n_{2n-1}} (f_{2k-1} + f_{2k}) \in W_{ex}$ and

$$\left\|\frac{m_{2j-1}}{n_{2j-1}}\sum_{k=1}(x_{2k-1}+x_{2k})\right\| \ge F\left(\frac{m_{2j-1}}{n_{2j-1}}\sum_{k=1}(x_{2k-1}+x_{2k})\right) = 2$$

while $F(n_{2j-1}^{-1}\sum_{k=1}(x_{2k-1}-x_{2k})) = 0$. Our aim is to show that there exists a K such that for every j and (C, 2j-1)-dependent paired sequence we have

$$\left\|\frac{m_{2j-1}}{n_{2j-1}}\sum_{k=1}^{n_{2j-1}} (x_{2k-1} - x_{2k})\right\| \le KCm_{2j-1}^{-1}.$$

For this we employ a variant of the basic inequality.

Proposition 7.4 (A variant of the basic inequality). Let $(x_k, f_k)_{k=1}^{2n_{2j-1}}$ be a (C, 2j-1)-dependent paired sequence and $\lambda_k, k \leq 2n_{2j-1}$ be scalars. Then for every weighted functional $f \in W_{ex}$ we can find $g_1, g_2 \in c_{00}(\mathbb{N})$ having non-negative coordinates such that $||g_2||_{\infty} < m_{2j-1}^{-3}$ and $g_1 = h_1$ or $g_1 = e_k^* + h_1$ or $g_1 = e_k^* + e_{k+1}^* + h_1$ for some $h_1 \in W_{aux2}$ with $w(f) = w(h_1)$ and $k \notin \operatorname{supp}(h_1)$ if $g_1 = e_k^* + h_1$ or $k, k+1 \notin \operatorname{supp}(h_1)$ if $g_1 = e_k^* + e_{k+1}^* + h_1$, satisfying

$$\left| f\left(\sum_{k=1}^{2n_{2j-1}} \lambda_k x_k\right) \right| \le C(g_1 + g_2) \left(\sum_{k=1}^{2n_{2j-1}} \lambda_k e_k\right).$$
(7.2)

If we additionally assume that there exist $j_0 \in \mathbb{N}$ such that for all weighted functionals $g \in W_{ex}$ with $w(g) = m_{j_0}$ and every interval E of \mathbb{N} we have that

$$\left|g(\sum_{k\in E}\lambda_k x_k)\right| \le C(\max_k |\lambda_k| + m_{2j-1}^{-3}\sum_{k\in E} |\lambda_k|)$$
(7.3)

then if $w(f) \neq m_{j_0}$ we may select h_1 to have a tree analysis $(h_t)_{t \in \mathcal{T}}$ with $w(h_t) \neq m_{j_0}$ for all $t \in \mathcal{T}$.

The proof is identical to the original one with the only exception that a functional with weight in the interval $[m_{2j_k}, m_{2j_{k+1}})$ "can norm" at most two vectors, namely x_{2k-1}, x_{2k} , instead of one in the case of the original basic inequality.

As a corollary of the basic inequality we get the following.

Corollary 7.5. Let $(x_k, f_k)_{k=1}^{2n_{2j-1}}$ be a (C, 2j-1)-dependent paired sequence. Then

$$\left\|\frac{m_{2j-1}}{n_{2j-1}}\sum_{k=1}^{n_{2j-1}} (x_{2k-1} - x_{2k})\right\| \le \frac{12C}{m_{2j-1}}.$$
(7.4)

The proof of the corollary follows from Proposition 7.4 and Lemma 7.1 once it is proved that for every (2j-1)-special sequence $(g_k)_{k=1}^d$ of W_{ex} we have

$$m_{2j-1}^{-1} \sum_{k=1}^{a} g_k (\sum_{k \in E} (-1)^{k+1} x_k) \le 6C(1 + (\#E)m_{2j-1}^{-2}).$$
(7.5)

The proof of this inequality is based on two key ingredients which are the same as in the proof of Proposition 4.5. The first one is that for every k the estimation of a weighted functional on x_{2k-1} or x_{2k} satisfies upper bounds which depend on the weight of the functional (see Lemma 7.1 and Proposition 7.2) and the second one is that the family of paired special sequences has a tree structure (see Remark 5.6b).

Using these two ingredients it is shown that for every (C, 2j - 1)-dependent paired sequence $(x_k, f_k)_{k=1}^{2n_{2j-1}}$ and for every (2j - 1)-special sequence $(g_k)_{k=1}^d$ of W_{ex} (7.5) holds. This implies that the additional part of the variant of the basic inequality holds and using Lemma 7.1 we get (7.4). 7.2. The HI property of X_{ex} . We start with the following lemma.

18

Lemma 7.6. Let $\epsilon > 0$ and $x \in X_{ex}$ be a normalized block vector such that $\operatorname{dist}(x, \widehat{X}_{hi}) > \epsilon$. Then for every $\delta \in (0, \epsilon)$ there exists $f \in W_{ex}$ such that

a)
$$\operatorname{supp}(f) \subset \mathbb{N}_{od},$$
 b) $f(x) > \epsilon - \delta$

Proof. Since dist $(x, \hat{X}_{hi}) > \epsilon$ the Hahn-Banach theorem yields that there exists $g \in S_{X_{ex}^*}$ such that $g(x) > \epsilon$ and $g_{|\hat{X}_{hi}|} = 0$. Denoting by E the range of x we get that the functional F = Eg satisfies also that $F(x) > \epsilon$ and $F(e_{2n}) = 0$ for every $n \in \mathbb{N}$. Using that $B_{X_{ex}^*} = \overline{W_{ex}}^{\parallel \parallel}$ we get $x^* \in W_{ex}$ such that

$$||F - x^*|| < \delta_1 \quad \text{where } \delta_1 \text{ satisfies } \delta_1 < \frac{\delta(1 - \epsilon)}{(1 - \delta)(1 + \#E)}. \tag{7.6}$$

The basis of X_{ex} is bimonotone hence $|x^*(e_{2n})| = |x^*(e_{2n}) - F(e_{2n})| < \delta_1$ and also we may assume that $x^* = Ex^*$. Assuming by a small perturbation if necessary, that δ is rational we set

$$f = (1 - \delta)x^* + (1 - \delta)\sum_{2n \in E} |x^*(e_{2n})| \epsilon_{2n} e_{2n}^*$$
(7.7)

where $\epsilon_{2n} = -sign(x^*(e_{2n}))$. Note that $f(e_{2n}) = 0$ for every n. Indeed for every n such that $2n \notin E$ we have $x^*(e_{2n}) = Ex^*(e_{2n}) = 0$ while if $2n \in E$,

$$x^*(e_{2n}) + |x^*(e_{2n})| \epsilon_n e_{2n}^*(e_{2n}) = x^*(e_{2n}) - x^*(e_{2n}) = 0.$$

Also f belongs to the rational convex hull of W_{ex} . This follows from the fact that $x^*(e_n) \in \mathbb{Q}$ for every $n \in \mathbb{N}$ and that

$$\sum_{2n\in E} |Ex^*(e_{2n})| \le \sum_{2n\in E} ||Ex^* - F|| \le \#E\,\delta_1 < \delta.$$
(7.8)

We conclude that $f \in W_{ex}$ since W_{ex} is rationally convex. To show that f is the functional we are seeking it remains to show that $f(x) > \epsilon - \delta$. Using (7.8) we get

$$f(x) = (1 - \delta)x^*(x) + (1 - \delta)\sum_{2n \in E} |x^*(e_{2n})| \epsilon_{2n} e_{2n}^*(x)$$

$$\geq (1 - \delta)F(x) - (1 - \delta)||F - x^*|| \cdot ||x|| - (1 - \delta)\delta_1 \# E$$

$$\geq (1 - \delta)\epsilon - (1 - \delta) \cdot \delta_1(1 + \# E) > \epsilon - \delta.$$

Proposition 7.7. For every closed infinite dimensional subspace Y of X_{ex} we have that $dist(S_Y, S_{\widehat{X}_{h,i}}) = 0$.

Proof. On the contrary assume that $dist(S_Y, S_{\widehat{X}_{hi}}) > \epsilon$ for an infinite dimensional subspace Y. We may assume that Y is a block subspace.

We first show that for every $j, p \in \mathbb{N}$ there exists a $(6/\epsilon, 2j)$ - SEP (y, f) with $y \in Y$, min y > p and supp $(f) \subset \mathbb{N}_{od}$.

Since dist $(S_Y, S_{\widehat{X}_{hi}}) > \epsilon$ we may choose a RIS $(y_k)_{k \in \mathbb{N}}$ of normalized $2 - \ell_1^{n_{2j_k}}$ averages in Y such that dist $(y_k, S_{\widehat{X}_{hi}}) > \epsilon$ for every k. Form Lemma 7.6 we get a sequence of functionals $(f_k)_k \subset W_{ex}$ such that

$$f_k(y_k) > \frac{\epsilon}{2}, \quad \operatorname{ran}(f_k) \subset \operatorname{ran}(y_k) \text{ and } f_{k|\widehat{X}_{hi}} = 0.$$
 (7.9)

For every k choose $c_k \in [1, 2/\epsilon)$ such that $f_k(c_k y_k) = 1$. Then for every $j \in \mathbb{N}$ and $F_j \subset \mathbb{N}$ with $\#F_j = n_{2j}$ and $\min F_j > \max\{n_{2j}, p\}$ it follows that the pair (y, f) where

$$y = \frac{m_{2j}}{n_{2j}} \sum_{k \in F_j} c_k y_k \quad \text{and} \quad f = m_{2j}^{-1} \sum_{k \in F_j} f_k \tag{7.10}$$

is a $(6/\epsilon, 2j)$ - SEP (see Definition 3.12) with $y \in Y$, min supp(y) > p and supp $(f) \subset \mathbb{N}_{od}$.

Next for every $j \in \mathbb{N}$ we defined a $(6\epsilon^{-1}, 2j - 1)$ -dependent paired sequence $(x_k, f_k)_{k=1}^{2n_{2j-1}}$ as follows:

Choose j_1 such that $n_{2j-1}^2 < m_{2j_1}$. Let (x_1, f_1) be a $(6\epsilon^{-1}, 2j_1)$ -SEP as in (7.10). Since $f_1 \in W_{ex}$ we have that $f_1 \in V_{n_1}$ for some n_1 . Using the coding function $\varrho_{n_1}^{2j_1}$ we get the functional $g_2 = \varrho_{n_1}^{2j_1}(f_1) = m_{2j_1}^{-1} \sum_{k \in B_1} e_k^* \in K_{n_1}^{2j_1}$. We set

$$f_2 = \hat{g}_2$$
 and $x_2 = \frac{m_{2j_1}}{n_{2j_1}} \sum_{k \in B_1} e_{2k}$

Let $2j_2 = \sigma(g_2)$ where σ is the coding function used to define the special sequences in the space X_{hi} . Choose (x_3, f_3) a $(6\epsilon^{-1}, 2j_2)$ -SEP as in (7.10) satisfying $\operatorname{supp}(x_2) < \operatorname{supp}(x_3)$. This yields that also $\operatorname{supp}(f_2) < \operatorname{supp}(f_3)$. Let n_3 be such that $f_3 \in V_{n_3}$. Using the coding function $\varrho_{n_3}^{2j_2}$ we get the functional $g_4 = \varrho_{n_3}^{2j_2}(f_3) = m_{2j_2}^{-1} \sum_{k \in B_3} e_k^* \in K_{n_3}^{2j_2}$. We set

$$f_4 = \widehat{g}_4$$
 and $x_4 = \frac{m_{2j_2}}{n_{2j_2}} \sum_{k \in B_3} e_{2k}.$

It is clear now how we proceed to get a $(6\epsilon^{-1}, 2j-1)$ -dependent paired sequence. Set $y = \frac{m_{2j-1}}{n_{2j-1}} \sum_{k=1}^{n_{2j-1}} x_{2k-1}$ and $x = \frac{m_{2j-1}}{n_{2j-1}} \sum_{k=1}^{n_{2j-1}} x_{2k}$. From the choice of the

sequence we have that $y \in Y$ and $x \in \widehat{X}_{hi}$. Corollary 7.5 yields that $||y - x|| < 72\epsilon^{-1}m_{2j-1}^{-1}$. Also since $(x_{2k-1})_{k=1}^{n_{2j-1}}$ is a $(2, m_{2j-1}^{-2})$ -RIS it follows that $1 \le ||x|| \le 4$. This yields $\operatorname{dist}(S_Y, S_{\widehat{X}_{hi}}) < 2 \cdot 72C\epsilon^{-1}m_{2j-1}^{-1}$. It follows that

$$\frac{\epsilon}{2} < \frac{72}{\epsilon \cdot m_{2j-1}} \Leftrightarrow m_{2j-1}\epsilon^2 < 2 \cdot 72$$

a contradiction for large enouge j. Therefore $dist(S_Y, S_{\widehat{X}_{hi}}) = 0$ for every infinite dimensional subspace Y of X_{ex} .

The above yields the following.

Theorem 7.8. The HI space X_{hi} is isometric to an infinite codimensional subspace of the HI space X_{ex} .

Proof. From Proposition 5.14 we have that $W_{ex|\mathbb{N}_{ev}} = \widehat{W}_{hi}$ and this shows that the space $\widehat{X_{hi}} = [e_{2n} : n \in \mathbb{N}]$ is isometric to X_{hi} . Propositions 6.1 and 7.7 yield that the space X_{ex} is HI and this completes the proof.

8. The HI extension of a variant of X_{hi} .

There are several examples of HI spaces where the norming set W does not include convex combinations and this is necessary for the restricted saturation method applied for the definition of W, see [5],[6]. Also the norming set in the Argyros-Deliyanni asymptotic ℓ_1 -HI space is defined by a norming set W which does not include rational convex combinations. Since the closedness in rational convex combinations is essential to show that the extended space is HI, naturally this raises the question if the spaces defined by a norming set not closed in rational convex combinations admit an HI extension. In this section we provide a positive answer to this problem for a variant of the space X_{hi} used before, which is defined by a norming set W_{vhi} which is similar to W_{hi} but not closed in rational convex combinations. More precisely we consider the following norming set. Let $\sigma : \mathbb{Q}_s \to \{2j : j \in \Omega_2\}$ be a coding function satisfying (4.1).

Definition 8.1. The set W_{vhi} is the minimal subset of c_{00} satisfying the following conditions:

- (1) $\{\pm e_n : n \in \mathbb{N}\} \subset W_{vhi}$.
- (2) W_{vhi} is symmetric and is closed under the restriction of its elements on intervals of \mathbb{N} .
- (3) W_{vhi} is closed under the (A_{n₂j}, m_{2j}⁻¹)-operations, i.e., m_{2j}⁻¹∑_{i=1}^d f_i ∈ W_{vhi} for all n_{2j}-admissible sequences (f_i)_{i=1}^d of W_{vhi}.
 (4) W_{vhi} is closed under the (A_{n₂j-1}, m_{2j-1})-operations on special sequences,
- (4) W_{vhi} is closed under the $(\mathcal{A}_{n_{2j-1}}, m_{2j-1}^{-1})$ -operations on special sequences, i.e., $m_{2j-1}^{-1} \sum_{i=1}^{d} f_i \in W_{vhi}$ for all n_{2j-1} -admissible sequences $(f_i)_{i=1}^{d}$ of W_{vhi} such that $w(f_1) = m_{2j_1} > n_{2j-1}^2$ and $w(f_{i+1}) = m_{\sigma(f_1,\dots,f_i)}$ for $1 \leq i \leq d-1$.

The difference between W_{vhi} and W_{hi} is the rational convexity. It follows by the same arguments as in section 4 that the space X_{vhi} with the set W_{vhi} as its norming set is also HI.

We define a variant W_{vex} of the extended norming set W_{ex} . The definition of W_{vex} is very similar to the previous one W_{ex} and goes as follows:

Let $W_0 = \{\pm e_n^* : n \in \mathbb{N}\}$ and assume that W_{n-1} has been defined. We set

$$\widetilde{U}_n = \bigcup_j \{ m_{2j}^{-1} \sum_{i=1}^k f_i : (f_i)_{i=1}^k \text{ is an } \mathcal{A}_{n_{2j}} \text{-admissible sequence of } \widetilde{W}_{n-1} \},\$$

and $\widetilde{V}_n = (\widetilde{U}_n \setminus \bigcup_{j=0}^{\infty} \bigcup_{k=n+1}^{+\infty} K_k^{2j}) \cup \widetilde{W}_{n-1}.$ For $j \in \mathbb{N}$ we define

$$\widetilde{W}_{n}^{2j-1} = \left\{ \frac{1}{m_{2j-1}} \left(\sum_{i=1}^{l} (f_{i} + \widehat{g}_{i}) + \sum_{i=l+1}^{m} \widehat{g}_{i} + \sum_{i=m+1}^{p} h_{i} \right) :$$

$$(f_{1}, \widehat{g}_{1}, \dots, f_{l}, \widehat{g}_{l}, \widehat{g}_{l+1}, \dots, \widehat{g}_{m}, h_{m+1}, \dots, h_{p}) \text{ is a } (2j-1, \widetilde{V}_{n}, (\varrho_{i}^{2k})_{i=1}^{n}, k = 0, 1, \dots)$$
special sequence}.
$$(8.1)$$

where the definition of a $(2j - 1, \tilde{V}_n, (\varrho_i^{2k})_{i=1}^n, k = 0, 1, ...)$ special sequence is identical to the corresponding definition in W_{ex} . We set

$$\widetilde{W}_n = \bigcup_j \widetilde{W}_n^{2j-1} \cup \widetilde{V}_n \cup \{\sum_{i=1}^d r_i f_i : (r_i)_{i=1}^d \text{ is a rational convex combination}, \\ f_i \in \bigcup_j \widetilde{W}_n^{2j-1} \cup \widetilde{V}_n \text{ and } \operatorname{supp}(\sum_{i=1}^d r_i f_i) \subset \mathbb{N}_{od}\}$$

The norming set W_{vex} is the set $W_{vex} = \bigcup_{n=0}^{+\infty} \widetilde{W}_n$.

Lemmas 5.11 and 5.13 remain valid for the norming sets W_{vhi} and W_{vex} and the proofs are identical. These yield that X_{vhi} is isometric to the subspace $\hat{X}_{vhi} = [e_{2n} : n \in \mathbb{N}]$ of X_{vex} . We have to prove that X_{vex} is HI. To prove this first we note that the variant of the basic inequality, Proposition 7.4 and Corollary 7.5 hold also for the space X_{vex} . Moreover the following holds.

Proposition 8.2. The space X_{vex} is reflexive and $B_{X_{ver}^*} = \overline{\text{conv}}^{\parallel \parallel}(W_{vex})$

To prove that X_{vex} is HI we need a slight modification in the proof of Lemma 7.6 since in that proof we use that $B_{X_{ex}^*} = \overline{W_{ex}}^{\|\cdot\|}$ which does not hold for the space X_{vex} .

Lemma 8.3. Let $\epsilon > 0$ and $x \in X_{vex}$ be a normalized block vector such that $\operatorname{dist}(x, \widehat{X}_{vhi}) > \epsilon$. Then for every $\delta \in (0, \epsilon)$ there exists $f \in W_{vex}$ such that

a)
$$\operatorname{supp}(f) \subset \mathbb{N}_{od},$$
 b) $f(x) > \epsilon - \delta$

Proof. Since dist $(x, \hat{X}_{vhi}) > \epsilon$ there exists $f \in S_{X_{vex}^*}$ such that $g(x) > \epsilon$ and $g_{|\hat{X}_{vhi}|} = 0$. Denoting by E the range of x we get that the functional F = Eg satisfies also that $F(x) > \epsilon$ and $F(e_{2n}) = 0$ for every $n \in \mathbb{N}$. Using that $B_{X_{vex}^*} = \overline{\operatorname{conv}}(W_{vex})^{\|\|}$ we get $x^* \in \operatorname{conv}(W_{vex})$ such that

$$||F - x^*|| < \delta_1 \quad \text{where } \delta_1 \text{ satisfies } \delta_1 < \frac{\delta(1 - \epsilon)}{(1 - \delta)(1 + \#E)}. \tag{8.2}$$

The basis of X_{vex} is bimonotone hence $|x^*(e_{2n})| = |x^*(e_{2n}) - F(e_{2n})| < \delta_1$ and also we may assume that x^* is supported by the set E, i.e., $x^* = Ex^*$.

Assuming by a small perturbation if necessary that δ is rational we set

$$f = (1 - \delta)x^* + (1 - \delta)\sum_{2n \in E} |x^*(e_{2n})| \epsilon_{2n} e_{2n}^* \text{ where } \epsilon_{2n} = -sign(x^*(e_{2n})).$$
(8.3)

As in Lemma 7.6 we get that that $f \in \operatorname{conv}_{\mathbb{Q}}(W_{vex})$ and $\operatorname{supp}(f) \subset \mathbb{N}_{od}$.

Since W_{vex} is closed in the rational convex combinations whose support is subset of \mathbb{N}_{od} it follows that $f \in W_{vex}$. The rest of the proof is identical with the proof of Lemma 7.6.

Using the above lemma we get the following.

Proposition 8.4. The space X_{vex} is HI.

The proof of the proposition is identical to the proof of the Proposition 7.7.

Remark 8.5. The extension method applied for the HI spaces defined by the families A_n can also be applied with identical proofs for the Gowers-Maurey space [15].

9. The case of asymptotic ℓ_1 -HI Banach spaces

In this section we show that the extension method can be applied also to asymptotic ℓ_1 -HI Banach spaces, see [3]. The method is identical to the one for the spaces defined by the families \mathcal{A}_{n_j} with some slight modifications due to the usage of the Schreier families.

Let us recall the definition of the norming set of a standard asymptotic $\ell_1\text{-HI}$ space.

Let $(n_j)_{j=0}^{\infty}, (m_j)_{j=0}^{\infty}$ be two sequences of positive integers such that $m_0 = 2$, $m_{j+1} = m_j^{m^j}$, $n_0 = 1$ and $n_{j+1} = m_1^{2m_{j+1}} n_j$. Let also σ be a coding function defined as in Section 4.

The norming set W_{hi} of a standard asymptotic ℓ_1 -HI space is the minimal subset of $c_{00}(\mathbb{N})$ satisfying the following properties:

- i) It is rationally convex, symmetric, closed on the projections on intervals of \mathbb{N} and contains $\pm e_n^*, n \in \mathbb{N}$.
- ii) It is closed in (S_{n2j}, m⁻¹_{2j})-operations, i.e., if (f_i)_{i∈F} is an S_{n2j}-admissible sequence of W_{hi} then m⁻¹_{2j} ∑_{i∈F} f_i ∈ W_{hi}.
 iii) It is closed in (S_{n2j-1}, m⁻¹_{2j-1})-operations on special sequences, i.e., if (f_i)_{i∈F} is an S admissible sequence of W_i = (f_i) = (f
- is an $S_{n_{2j}}$ -admissible sequence of W_{hi} $w(f_i) = m_{2j_i} = \sigma(f_1, \ldots, f_{i-1})$ and $m_{2j_1} \ge n_{2j-1}^2$ then $m_{2j-1}^{-1} \sum_{i \in F} f_i \in W_{hi}$.

To define the extended HI space we shall use a variation of the Schreier families $(\mathcal{S}_n)_{n \in \mathbb{N}}$, (see [10] for some of their properties).

Definition 9.1. We set
$$\mathcal{S}_1^f = \{\{1\}\} \cup \{F \subset \mathbb{N} : 2\#F \leq minF\}$$
 and

$$\mathcal{S}_{n}^{f} = \mathcal{S}_{1}^{f}[\mathcal{S}_{n-1}^{f}] = \{ \cup_{i=1}^{k} F_{i} : F_{1} < \dots < F_{k} \in \mathcal{S}_{n-1}^{f}, \{\min F_{i} : i \leq k\} \in \mathcal{S}_{1}^{f} \}.$$

We also the define the families

$$\mathcal{S}_n^f \odot \mathcal{A}_2 = \{ \bigcup_{i=1}^d F_i : F_1 < \dots < F_d, \ \#F_i \le 2 \text{ and } \{ \max F_i \}_{i=1}^d \in \mathcal{S}_n^f \}.$$

It is easy to see that for every n the families $\mathcal{S}_m^f, \mathcal{S}_n^f \odot \mathcal{A}_2$ are regular.

For a finite subset $G = \{n_1 < \cdots < n_d\}$ of \mathbb{N} we denote by \widehat{G} the set $\widehat{G} = \{2n_1 < \cdots < n_d\}$ $\cdots < \cdots < 2n_d$. We also denote by $\widehat{\mathcal{S}}_n$ the family $\{\widehat{F} : F \in \mathcal{S}_n\}$.

Lemma 9.2. For every $n \in \mathbb{N}$ we have

$$\widehat{\mathcal{S}_n} = \mathcal{S}_n^f \cap [\mathbb{N}_{ev}]^{<\omega}.$$

Proof. First we show by induction on n that

$$\mathcal{S}_n^f \cap [\mathbb{N}_{ev}]^{<\omega} \subset \widehat{\mathcal{S}_n}.$$

Let $F = \{2n_1 < 2n_2 < \cdots < 2n_d\} \in \mathcal{S}_1^f$. Then $2d \leq 2n_1 \Rightarrow d \leq n_1$. It follows that $G = \{n_1 < n_2 < \dots < n_d\} \in \mathcal{S}_1 \text{ and } F = \widehat{G}.$

Assume that the result holds for every $k \leq n$ and $F \in \mathcal{S}_{n+1}^f \cap [\mathbb{N}_{ev}]^{<\omega}$. Then $F = \bigcup_{i=1}^{d} F_i$ with $F_i \in \mathcal{S}_n^f$ and $\{\min F_i : i \leq d\} \in \mathcal{S}_1^f$. By the inductive hypothesis we have for every $i \leq d$ there exits $G_i \in S_n$ such that $F_i = \widehat{G}_i$. From the case n = 1we get that $\{\min G_i : i \leq d\} \in S_1$ and therefore $G = \bigcup_{i=1}^d G_i \in S_{n+1}$. Also $\widehat{G} = F$ and this completes the proof of the induction.

The inclusion $\widehat{\mathcal{S}_n} \subset \mathcal{S}_n^f \cap [\mathbb{N}_{ev}]^{<\omega}$ is proved similarly.

We give the definition of the admissibility with respect the families \mathcal{S}_n and $\mathcal{S}_n^f \odot \mathcal{A}_2.$

Definition 9.3. We call a block sequence $(x_i)_{i=1}^d$ in $c_{00}(\mathbb{N})$

- (1) S_n -admissible if $\min\{\operatorname{supp}(x_i) : i \leq d\} \in S_n$.
- (2) \mathcal{S}_n^f -admissible if $\min\{\operatorname{supp}(x_i) : i \leq d\} \in \mathcal{S}_n^f$. (3) $\mathcal{S}_n^f \odot \mathcal{A}_2$ -admissible if $\{\min \operatorname{supp}(x_i)_{i=1}^d\} = \bigcup_{i=1}^m F_i, F_i < F_{i+1}, \#F_i \leq 2$ for every i and $\{\max F_i : i \leq m\} \in \mathcal{S}_n^f$.

Let us mention that admissibility with respect the families $\mathcal{S}_n, \mathcal{S}_n^f, n \in \mathbb{N}$ is the standard one, see [9], [10], while the admissibility with respect the families $\mathcal{S}_n^f \odot \mathcal{A}, n \in \mathbb{N}$ is a variation of the standard one, suitable for our applications.

Remark 9.4. Let $G = \{m_1 < m_2 < \cdots < m_d\} \in S_n$. For every $0 \le j \le j$ d and every $k_1 < 2m_1 < k_2 < 2m_2 < \cdots < k_j < 2m_j$ we have that the set $\{k_1, 2m_1, \ldots, k_j, 2m_j, 2m_{j+1}, 2m_{j+2}, \ldots, 2m_d\}$ is $\mathcal{S}_n^f \odot \mathcal{A}_2$ -admissible.

Indeed if $F_i = \{k_i, 2m_i\}$ for $i \leq j$ and $F_i = \{2m_i\}$ for i > j, then Lemma 9.2 yields that $\{\max F_i : i \leq d\}$ is S_n^f -admissible and hence $\bigcup_{i=1}^d F_i$ is $S_n^f \odot \mathcal{A}_2$ -admissible (see Definition 9.3).

The norming set of the extended space will be an EO-complete subset of the norming set W_{mT} of the mixed Tsirelson space $T[(\mathcal{S}_{n_{2j}}^f, m_{2j}^{-1})_{j \in \mathbb{N}}, (\mathcal{S}_{n_{2j-1}}^f \odot \mathcal{A}_2, m_{2j-1}^{-1})_{j \in \mathbb{N}}]$. In order to define the extended norming set as in Subsection 5.1 we consider the

sets

$$K^{2j} = \{m_{2j}^{-1} \sum_{i \in F} e_i^* : F \text{ is maximal } \mathcal{S}_{n_{2j}}\text{-set}\}$$

and a partition $(K_i^{2j})_i$ of them such that for every i, $\sup\{\min(f) : f \in K_i^{2j}\} = +\infty$. The sets W_{mT}^{2j} and the coding functions $\varrho_i^{2j} : W_{mT}^{2j} \to K_i^{2j}$ are also defined as in Subsection 5.1.

The definition of the paired special, semi paired special and special sequences are the same using $\mathcal{S}_{n_{2i-1}}^f \odot \mathcal{A}_2$ -admissibility instead of the $\mathcal{A}_{n_{2j}}$ -admissibility. Let us give the definition of the $(2j-1, V, (\varrho_k^{2i})_{k=1}^n, i=0, 1, ...)$ - paired special sequence. **Definition 9.5.** Let $j, n \in \mathbb{N}$ and $V \subset W_{mT}$. We call $(2j - 1, V, (\varrho_k^{2i})_{k=1}^n, i =$ $(0,1,\ldots)$ - paired special sequence any block sequence $(f_1,\widehat{g}_1,\ldots,f_p,\widehat{g}_p)$ satisfying the following:

(1) it is $\mathcal{S}_{n_{2i-1}}^f \odot \mathcal{A}_2$ -admissible and for every k there exists $j_k \in \mathbb{N}$ such that $f_k \in V \cap W_{mT}^{2j_k}$ and $\widehat{g}_k = \widehat{\varrho_{i_k}^{2j_k}(f_k)}$ for some $i_k \leq n$. (2) (g_1, g_2, \dots, g_p) is a (2j-1)-special sequence of W_{hi} .

Remark 9.6. Let us point out that the $S_{n_{2j-1}}$ -admissibility of (g_1, g_2, \ldots, g_p) implies the $\mathcal{S}_{n_{2i-1}}^f \odot \mathcal{A}_2$ -admissibility of the sequence $(f_1, \widehat{g}_1, \ldots, f_p, \widehat{g}_p)$ (see Remark 9.4).

The definition of the $(2j - 1, V, (\varrho_k^{2i})_{k=1}^n, i = 0, 1, ...)$ - paired special sequences and $(2j - 1, V, (\varrho_k^{2i})_{k=1}^n, i = 0, 1, ...)$ - special sequence are as in Subsection 5.1.

Next we prove that $W_{ex|N_{ev}} = \widehat{W}_{hi}$ proving the analogues of Lemmas 5.11,5.13.

Lemma 9.7. For every $n \in \mathbb{N}$ and every $f \in W_n$ we have that $f_{|\mathbb{N}_{ev}} \in \widehat{W}_{hi}$.

Proof. As in Lemma 5.11 we use induction. The case n = 0 is identical. Assume that the result holds for all $k \leq n$ and let $f \in W_{n+1}$.

If f is the result of an $(\mathcal{S}_{n_{2j}}^f, m_{2j}^{-1})$ operation of elements of W_n , i.e., f = $m_{2i}^{-1} \sum_{i=1}^{d} f_i$, from the inductive hypothesis we have that for every *i* there exists $\phi_i \in W_{hi}$ such that $f_{i|\mathbb{N}_{ev}} = \widehat{\phi_i}$. Lemma 9.2 yields that $\{\text{minsupp}(\phi_i) : i \leq d\}$ is $\mathcal{S}_{n_{2j}}$ -admissible. Since W_{hi} is closed in $(\mathcal{S}_{n_{2j}}, m_{2j}^{-1})$ -operations we get that $f_{|\mathbb{N}_{ev}} =$ $m_{2j}^{-1} \widehat{\sum_{i=1}^d} \phi_i \in \widehat{W}_{hi}.$

The result in the case where f is the result of an $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i = 0, 1, ...)$ special sequence is analogue to the previous one using Remark 9.6. The case is fis a rational convex combination follows readily. **Lemma 9.8.** For every $f \in W_{hi}$ there exists $\phi \in W_{ex}$ such that $\phi_{|\mathbb{N}_{ey}} = \widehat{f}$.

Proof. Similar to the proof of Lemma 5.13 we proceed by induction.

For n = 1 and $f = m_{2i}^{-1} \sum_{i \in G} \pm e_i^*$, $G \in \mathcal{S}_{n_{2j}}$ we observe from Lemma 9.2 that $\widehat{G} \in \mathcal{S}_n^f$. Using this observation, as in Lemma 5.13 we conclude that $\widehat{f} \in W_{ex}$ We pass now to the case where $f = m_{2j-1}^{-1} \sum_{i=1}^p g_i$ and (g_1, \ldots, g_p) is a (2j-1)-

special sequence of W_{hi}^1 . As in Lemma 5.13 the proof is based on the following claim.

Claim. Let $g_i \in K_{n_i}^{2j_i}$, $i \leq m$ be such that (g_1, \ldots, g_m) is a (2j - 1)-special sequence of W_{hi}^1 . Then there exist $0 \leq l \leq m$ and $f_1, \ldots, f_l \in W_{ex}$ such that $(f_1, \widehat{g}_1, \ldots, f_l, \widehat{g}_l, \widehat{g}_{l+1}, \ldots, \widehat{g}_m)$ is a $(2j-1, V_n, (\varrho_k^n)_{k=1}^n, i=0, \ldots)$ semi-paired special sequence where $n = \max\{n_i : i \leq m\}$.

Sketch of the proof. Let $l \leq m$ be maximal with the properties

- (1) $g_i \in \varrho_{n_i}^{2j_i}(V_n \cap W_{mT}^{2j_i})$ for every $i \le l$. (2) $F = ((\varrho_{n_i}^{2j_i})^{-1}(g_i), \widehat{g}_i)_{i=1}^l$ is a $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i = 0, 1...)$ -paired special sequence, where $n = \max\{n_i : i \le m\}$.

For every i, as in the first case we have $\hat{g}_i \in W_{ex}$, and from Remark 9.4 we get $(\varrho_{n_i}^{2j_1})^{-1}(g_1), \widehat{g}_1, \ldots, (\varrho_{n_i}^{2j_l})^{-1}(g_i), \widehat{g}_l, \widehat{g}_{l+1}, \ldots, \widehat{g}_m)$ is $\mathcal{S}_{n_{2j-1}}^f \odot \mathcal{A}_2$ -admissible in W_{ex} . Considering the cases l = 0, l = m and 0 < l < m, as in the proof of Lemma 5.13 we get that $(\varrho_{n_i}^{2j_1})^{-1}(g_1), \widehat{g}_1, \ldots, (\varrho_{n_i}^{2j_l})^{-1}(g_i), \widehat{g}_l, \widehat{g}_{l+1}, \ldots, \widehat{g}_m)$ is a $(2j-1, V_n, (\varrho_k^{2i})_{k=1}^n, i=0, 1, \dots)$ -semi paired special sequence. \Box

Returning back to the proof, let $f = m_{2j-1}^{-1} \sum_{i=1}^{p} g_i$ where (g_1, \ldots, g_p) is a (2j-1)-special sequence of W_{hi}^1 , and $m \leq p$ be maximal with the property $g_i \in K_{2n_i}^{2j_i}$ for every $i \leq m$. It follows from the Claim and the definition of the special sequences that $((\varrho_{n_1}^{2j_1})^{-1}(g_1), \hat{g}_1, \dots, (\varrho_{n_m}^{2j_l})^{-1}(g_m), \hat{g}_m, \hat{g}_{m+1}, \dots, \hat{g}_p)$ is a $(2j - 1, V_n, (\varrho_k^{2i})_{k=1}^n, i = 0, 1, \dots)$ -special sequence. Hence $\phi = m_{2j-1}^{-1}(\sum_{i=1}^m ((\varrho_{n_i}^{2j_i})^{-1}(g_i) + ((\varphi_{n_i}^{2j_i})^{-1}(g_i)))$ $\widehat{g}_i) + \sum_{m+1}^p \widehat{g}_i) \in W_{ex} \text{ and } \widehat{f} = \phi_{|\mathbb{N}_{ev}}.$

Assume that we have proved the result for every $k \leq n$ and let $f \in W_{hi}^{n+1}$.

In the case f is the result of an $(\mathcal{S}_{n_{2j}}, m_{2j}^{-1})$ -operation of elements of W_{hi}^n , using Lemma 9.2 and the property W_{ex} is closed in the $(\mathcal{S}_{n_{2i}}^f, m_{2i}^{-1})$ -operations, we show, as in Lemma 5.13, that $\hat{f} = \phi_{|\mathbb{N}_{ev}}$ for some $\phi \in W_{ex}$. In the case $f = m_{2j-1}^{-1} \sum_{i=1}^{p} g_i$ where (g_1, \ldots, g_p) is a (2j-1)-special sequence

of W_{hi}^{n+1} , is proved as in Lemma 5.13 using the above Claim.

As a corollary we get the following.

Proposition 9.9. For the norming sets W_{hi} and W_{ex} of the spaces X_{hi} and X_{ex} we have that

$$W_{ex|\mathbb{N}_{ev}} = \widehat{W}_{hi}.$$

As a consequence the space X_{hi} is isometric to the subspace of X_{ex} generated by the sequence $(e_{2n})_{n \in \mathbb{N}}$.

The results of the Sections 4,5,7 with the obvious modifications imposed by the usage of the Schreier families in place of the \mathcal{A}_{n_i} families, hold also for the new spaces with similar proofs (see [10] for a detailed exposition of the related results).

The above yield the following.

Theorem 9.10. A standard asymptotic ℓ_1 HI Banach space is isometric to an infinite codimensional subspace of a reflexive asymptotic ℓ_1 HI Banach space.

Remark 9.11. Applying the analogue modifications we get that any standard asymptotic ℓ_p HI space, 1 , (see [12]), admits an HI extension.

10. FINAL DISCUSSION

We discuss some open problems that are related to this article. We begin with the one that motivated this paper and remains open in its full generality.

Problem 1 (Pełczyński). Does every HI space X embed into an HI space Y such that Y/X is infinite dimensional?

Problem 1 was posed by Pełczynski at a conference where the first author was present, however we were unable to find it in the literature. Yet, he posed a strongly related one in [17, Problem 3.3, page 348].

Problem 2 (Pełczynski). *Does every separable HI space X embed into an HI space Y with a Schauder basis?*

It is remarkable that in the same paper [17] Pełczyński posed the problem of the existence of a \mathcal{L}_{∞} -space that is HI, a fact that the first author was unaware of. This is why this paper was not cited in [2], where the scalar-plus-compact problem was solved using such a space. Let us point out that the methods developed in the current paper show promise in constructing HI extensions of \mathcal{L}_{∞} -HI spaces, such as the Argyros-Haydon space.

After sharing a preliminary version of this article to W. B. Johnson, he pointed out to us a more general form of Pełczyński's HI extension Problem 1, based on the following definition (see also [11, page 4701] for a formulation in the setting of twisted sums).

Definition 10.1. Let X be a Banach space Y be a superspace of X. We call Y a singular extension of X if Y/X is infinite dimensional and the quotient map $Q: Y \to Y/X$ is strictly singular.

Problem 3 (W. B. Johnson). Does every infinite dimensional separable Banach space X that is not isomorphic to c_0 admit a singular extension Y?

For example, as proved by Kalton and Peck, the space $X = \ell_p$, 1 , $admits a singular extension, namely the Banach space <math>Z_p$ (see [16, Theorem 6.4, page 27]). It is not hard to see that if X is HI and it admits a singular extension Y then Y must be HI as well. Therefore, a positive answer to Problem 3 would imply a positive answer of Problem 1, at least in the separable setting (whether every separable HI space admits a singular extension was also asked by Castillo, Ferenczi, and Gonzalez in [11, page 4701]).

In [11] it is proved that the uniformly convex HI space \mathcal{F} constructed by V. Ferenczi in [13] admits an HI short exact sequence $0 \to \mathcal{F} \to \mathcal{F}_2 \to \mathcal{F} \to 0$, such that \mathcal{F}_2 is HI, i.e., \mathcal{F}_2 is an HI twisted sum of \mathcal{F} . This is achieved by using the fact that \mathcal{F} is a space resulting from complex interpolation. Note that in particular \mathcal{F}_2 is an HI extension of \mathcal{F} .

It has long been an open question whether it is possible to construct a uniformly convex HI space by directly defining a norming set instead of using complex interpolation. It is therefore relevant to ask whether it is possible to find an HI space X

and an HI twisted sum Y of X by defining a norming set W. The natural approach would be to try to construct a rationally convex norming set W that induces an HI space Y such that

$$\{f|_{\mathbb{N}_{ev}} : f \in W\} = \{f \circ \pi : f \in W \text{ with } \operatorname{supp}(f) \subset \mathbb{N}_{od}\},\$$

where $\pi : \mathbb{N}_{ev} \to \mathbb{N}_{od}$ is the order preserving bijection. Then, by taking $X = \overline{\langle \{e_{2n} : n \in \mathbb{N}\} \rangle}$, the Hahn-Banach theorem would yield that Y/X is isometrically isomorphic to X.

References

- Albiac, F., Kalton, Nigel J. Topics in Banach space theory. Second edition. With a foreword by Gilles Godefory. Graduate Texts in Mathematics, 233. Springer, 2016. xx+508 pp.
- [2] Argyros, S. A.; Haydon, R. G., A hereditarily indecomposable \mathcal{L}_{∞} -space that solves the scalar-plus-compact problem. Acta Math. 206 (2011), no. 1, 1-54.
- [3] Argyros, S. A., Deliyanni, I., Examples of asymptotic l₁ Banach spaces. Trans. Amer. Math. Soc. 349 (1997), no. 3, 973–995.
- [4] Argyros, S. A.; Manoussakis, A., An indecomposable and unconditionally saturated Banach space. Dedicated to Professor Aleksander Pełczyński on the occasion of his 70th birthday. Studia Math. 159 (2003), no. 1, 1–32.
- [5] Argyros, S. A.; Motakis, P., A dual method of constructing hereditarily indecomposable Banach spaces. Positivity 20 (2016), no. 3, 625–662.
- [6] Argyros, S. A.; Motakis, P., On the complete separation of asymptotic structures in Banach spaces, Adv. in Math., 362 (2020), 106962, 51 pp.
- [7] Argyros, S. A., Raikoftsalis, Th., The cofinal property of the reflexive indecomposable Banach spaces, Ann. Inst. Fourier, 62 (2012) no 1, 1-45.
- [8] Argyros, S. A., Freeman, D., Haydon, R., Odell, E., Raikoftsalis, Th., Schlumprecht Th., Zisimopoulou, D., Embedding uniformly convex spaces into spaces with very few operators, Journal of Functional Analysis 262 (2012), no 3, 825-849.
- [9] Argyros, S. A.; Todorcevic, S. Ramsey methods in analysis. Advanced Courses in Mathematics. CRM Barcelona. Birkhäuser Verlag, Basel, 2005. viii+257 pp
- [10] Argyros, S. A.; Tolias, A., Methods in the theory of hereditarily indecomposable Banach spaces. Mem. Amer. Math. Soc. 170 (2004), no. 806, vi+114 pp.
- [11] Castillo, J. M. F.; Ferenczi, V.; González, M., Singular twisted sums generated by complex interpolation. Trans. Amer. Math. Soc. 369 (2017), no. 7, 4671-4708.
- [12] Deliyanni I., Manoussakis A., Asymptotic ℓ_p hereditarily indecomposable Banach spaces. Illinois J. Math. 51 (2007), no. 3, 767-803.
- [13] Ferenczi, V., A uniformly convex hereditarily indecomposable Banach space. Israel J. Math. 102 (1997), 199-225.
- [14] Gowers, W.T. An infinite Ramsey theorem and some Banach- space dichotomies, Annals of Mathematics, 156 (2002), 797–833.
- [15] Gowers, W. T.; Maurey, B. The unconditional basic sequence problem, J. Amer. Math. Soc. 6 (1993), no. 4, 851–874
- [16] Kalton, N. J.; Peck, N. T., Twisted sums of sequence spaces and the three space problem. Trans. Amer. Math. Soc. 255 (1979), 1-30.
- [17] Pełczyński, A., Selected problems on the structure of complemented subspaces of Banach spaces. Methods in Banach space theory, 341-354, London Math. Soc. Lecture Note Ser., 337, Cambridge Univ. Press, Cambridge, 2006.
- [18] Lindenstrauss, J., Tzafriri, L, Classical Banach spaces. I. Sequence spaces. Ergebnisse der Mathematik und ihrer Grenzgebiete, Vol. 92. Springer-Verlag, Berlin-New York, 1977.
- [19] Maurey, B., Rosenthal H. P., Normalized weakly null sequences with no unconditional subsequence, Studia Math., 61 (1977), 77-98.
- [20] Schlumprecht, Th. An arbitrarily distortable Banach space. Israel J. Math. 76 (1991), no. 1-2, 81–95.

NATIONAL TECHNICAL UNIVERSITY OF ATHENS, FACULTY OF APPLIED SCIENCES, DEPARTMENT OF MATHEMATICS, ZOGRAFOU CAMPUS, 157 80, ATHENS, GREECE. Email address: sargyros@math.ntua.gr

School of Environmental Engineering, Technical University of Crete, University Campus, 73100 Chania, Greece

Email address: amanousakis@isc.tuc.gr

Department of Mathematics and Statistics, York University, 4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada *Email address:* pmotakis@yorku.ca