

ON BEST COAPPROXIMATIONS IN SUBSPACES OF DIAGONAL MATRICES

DEBMALYA SAIN, SHAMIM SOHEL, SOUVIK GHOSH AND KALLOL PAUL

ABSTRACT. We characterize the best coapproximation(s) to a given matrix T out of a given subspace \mathbb{Y} of the space of diagonal matrices \mathcal{D}_n , by using Birkhoff-James orthogonality techniques and with the help of a newly introduced property, christened the $*$ -Property. We also characterize the coproximal subspaces and the co-Chebyshev subspaces of \mathcal{D}_n in terms of the $*$ -Property. We observe that a complete characterization of the best coapproximation problem in ℓ_∞^n follows directly as a particular case of our approach.

1. INTRODUCTION.

The study of best coapproximations in Banach spaces was initiated by Franchetti and Furi in [3], as a complementary notion to the classical theory of best approximations. Unlike the Hilbert space case, the existence and the uniqueness of best coapproximations are known to be difficult problems in the setting of Banach spaces, especially from a computational point of view. We refer the readers to [3, 7] for more information on this topic. The purpose of the present article is to study the best coapproximation problem in subspaces of diagonal matrices, from the perspective of orthogonality. Indeed, we obtain a complete characterization of the best coapproximations to any given square matrix, in the said subspaces of matrices. It should be noted that recently in [13], the concept of orthogonality has been utilized to address best approximation problems in Banach spaces. Although our present exploration is motivated in spirit by the said approach, the techniques used in this study are completely different from those used in [13]. Let us now introduce the relevant notations and the terminologies to be used throughout the article.

We use the symbol \mathbb{H} to denote a Hilbert space, along with its usual inner product $\langle \cdot, \cdot \rangle$ and its usual norm $\|\cdot\|_2$. In this article, we will only work with *real* Hilbert spaces. Let \perp denote the usual orthogonality relation on \mathbb{H} . $\mathbb{L}(\mathbb{H})$ ($\mathbb{K}(\mathbb{H})$) denotes the Banach space of all bounded (compact) linear operators on \mathbb{H} , endowed with the usual operator norm. Given $T \in \mathbb{L}(\mathbb{H})$, let M_T denote the norm attainment set of

2010 *Mathematics Subject Classification.* Primary 46B20, Secondary 47L05.

Key words and phrases. best coapproximations; coproximal subspace; co-Chebyshev subspace; diagonal matrices; Birkhoff-James orthogonality .

The research of Dr. Debmalya Sain is sponsored by SERB N-PDF fellowship under the mentorship of Professor Apoorva Khare. Dr. Sain feels grateful to have the opportunity to acknowledge the loving friendship of his childhood friend Dr. Subhamoy Neogi, an accomplished physician with an empathetic heart. The second and third author would like to thank CSIR, Govt. of India, for the financial support in the form of Junior Research Fellowship under the mentorship of Prof. Kallol Paul. The research of Prof. Paul is supported by project MATRICS (MTR/2017/000059) of SERB, DST, Govt. of India.

T , i.e. $M_T = \{x \in H : \|x\|_2 = 1, \|Tx\|_2 = \|T\|\}$. We note that $M_T \neq \phi$ whenever $T \in \mathbb{L}(\mathbb{H})$. In case \mathbb{H} is finite-dimensional, given any $T \in \mathbb{L}(\mathbb{H})$, we identify T with its matrix representation with respect to the canonical basis of \mathbb{H} . Let \mathcal{M}_n denote the space of all $n \times n$ real matrices and let \mathcal{D}_n be the subspace of \mathcal{M}_n , consisting of diagonal matrices. Given $T \in \mathcal{M}_n$, let T^t denotes the transpose of T . Given any $A \in \mathcal{D}_n$ with diagonal entries a_{ii} , $1 \leq i \leq n$, we write $A = ((a_{11}, a_{22}, \dots, a_{nn}))$, for the sake of brevity. The zero element of \mathbb{R}^n is denoted by θ , whenever $n > 1$. The following definition is of fundamental importance in our entire study:

Definition 1.1. Let $(\mathbb{X}, \|\cdot\|)$ be a Banach space and let \mathbb{Y} be a subspace of \mathbb{X} . Given any $x \in \mathbb{X}$, we say that $y_0 \in \mathbb{Y}$ is a best coapproximation to x out of \mathbb{Y} if $\|y_0 - y\| \leq \|x - y\|$ for all $y \in \mathbb{Y}$.

In general, neither the existence nor the uniqueness of best coapproximation(s) is guaranteed, even in the finite-dimensional case. A subspace \mathbb{Y} of the Banach space \mathbb{X} is said to be coproximal if a best coapproximation to any element of \mathbb{X} out of \mathbb{Y} exists. A coproximal subspace \mathbb{Y} is said to be co-Chebyshev if the best coapproximation is unique in each case. Given $x \in \mathbb{X}$ and a subspace \mathbb{Y} of \mathbb{X} , the (possibly empty) set of all best coapproximations to x out of \mathbb{Y} is denoted by $\mathcal{R}_{\mathbb{Y}}(x)$. Our aim in this article is to explore the problem of finding the best coapproximation(s) to any given $T \in \mathcal{M}_n$ out of any given subspace \mathbb{Y} of \mathcal{D}_n , provided the best coapproximation(s) exist. We employ Birkhoff-James orthogonality techniques and the concept of numerical range of an operator $T \in \mathbb{L}(\mathbb{H})$, to obtain a complete solution to the above problem, which is also computationally effective. Let us recall from the pioneering articles [1, 5] that given any two elements x, y in a Banach space \mathbb{X} , we say that x is Birkhoff-James orthogonal to y , written as $x \perp_B y$, if $\|x + \lambda y\| \geq \|x\|$ for all $\lambda \in \mathbb{R}$. It should be noted that given any subspace \mathbb{Y} of a Banach space \mathbb{X} and an element $x \in \mathbb{X}$, $y_0 \in \mathbb{Y}$ is a best coapproximation to x out of \mathbb{Y} if and only if $\mathbb{Y} \perp_B (x - y_0)$ i.e., $y \perp_B (x - y_0)$ for all $y \in \mathbb{Y}$. Using Theorem 1.1 of [2], also known as the Bhatia-Šemrl Theorem, we study the best coapproximation problem from the perspective of Birkhoff-James orthogonality. We also recall that given any $T \in \mathbb{L}(\mathbb{H})$, the numerical range of T is defined as $W(T) := \{\langle Tx, x \rangle : \|x\|_2 = 1\}$. We refer the readers to [4], for a comprehensive study and possible applications of the numerical range of an operator in $\mathbb{L}(\mathbb{H})$.

In order to apply the above concepts in our designated study, we need to introduce the following definitions whose importance will be self-evident in due course of time.

Definition 1.2. Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be a set of linearly independent elements in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Considering the diagonal matrices A_1, A_2, \dots, A_m as column vectors, we form the $n \times m$ matrix $\tilde{A} = (\tilde{a}_{ij})_{1 \leq i \leq n, 1 \leq j \leq m}$, where $\tilde{a}_{ij} = a_{ii}^j$.

- (i) For each $i \in \{1, 2, \dots, n\}$, the i -th component of \mathcal{A} is defined as the i -th row of \tilde{A} , i.e., $(a_{ii}^1, a_{ii}^2, \dots, a_{ii}^m)$. Whenever the context is clear we simply say the i -th component of \mathcal{A} as the i -th component.
- (ii) The i -th component and the j -th component are said to be equivalent if $(a_{ii}^1, a_{ii}^2, \dots, a_{ii}^m) = \pm (a_{jj}^1, a_{jj}^2, \dots, a_{jj}^m)$.

(iii) The positively associated set $P_i^+(\mathcal{A})$ of the i -th component is defined as

$$P_i^+(\mathcal{A}) = \left\{ j \in \{1, 2, \dots, n\} : (a_{jj}^1, a_{jj}^2, \dots, a_{jj}^m) = (a_{ii}^1, a_{ii}^2, \dots, a_{ii}^m) \right\}.$$

Similarly, the negatively associated set $P_i^-(\mathcal{A})$ is defined as

$$P_i^-(\mathcal{A}) = \left\{ j \in \{1, 2, \dots, n\} : (a_{jj}^1, a_{jj}^2, \dots, a_{jj}^m) = - (a_{ii}^1, a_{ii}^2, \dots, a_{ii}^m) \right\}.$$

For simplicity, we write $P_i^+(\mathcal{A}) = P_i^+$ and $P_i^-(\mathcal{A}) = P_i^-$, if the context is clear.

(iv) The i -th component is said to satisfy the $*$ -Property with respect to \mathcal{A} if there exist $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$ such that

$$\left| \sum_{k=1}^m \beta_k a_{ii}^k \right| > \max \left\{ \left| \sum_{k=1}^m \beta_k a_{jj}^k \right| : 1 \leq j \leq n, j \notin P_i^+ \cup P_i^- \right\}.$$

Definition 1.3. Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Suppose that the i -th component satisfies the $*$ -Property with $|P_i^+ \cup P_i^-| = k_i$. Given $T = (b_{pq})_{1 \leq p, q \leq n}$, we define the $*$ -associated matrix of T corresponding to the i -th component as a square matrix of order k_i , given by ${}^i T_* = (c_{rs})_{1 \leq r, s \leq k_i}$, where

$$\begin{aligned} c_{rs} &= b_{rs}, (r, s) \in P_i^+ \times (P_i^+ \cup P_i^-) \\ &= -b_{rs}, (r, s) \in P_i^- \times (P_i^+ \cup P_i^-) \end{aligned}$$

In this paper, we obtain a complete characterization of the best coapproximation to an element of \mathcal{M}_n out of a given subspace of \mathcal{D}_n . We emphasize that our method is computationally convenient and it is possible to present a tractable algorithmic solution to the above problem by using it. We further illustrate this by presenting explicit numerical examples in support of our claim. The first step in this direction is to obtain a theoretical characterization of the best coapproximation problem in $\mathbb{K}(\mathbb{H})$. The second step is to explore some fundamental attributes of the newly introduced $*$ -Property in connection with the best coapproximation problem. In the final step, we assimilate the previously obtained results to present the desired algorithm to study the best coapproximation problem in any given subspace of \mathcal{D}_n . We also characterize the coproximal subspaces and co-Chebyshev subspaces of \mathcal{D}_n in \mathcal{M}_n . As another important application of the present study, we observe that a particular case of our method gives a complete solution to the best coapproximation problem in ℓ_∞^m , for any given $m \in \mathbb{N}$.

2. MAIN RESULTS

We begin with a theoretical characterization of best coapproximations in $\mathbb{K}(\mathbb{H})$, that will play a crucial role in the computational approach towards finding best coapproximation(s) (provided it exists) in any given subspace of \mathcal{D}_n , as adopted in the present article.

Theorem 2.1. *Let \mathbb{H} be a Hilbert space and let $T, A_1, A_2, \dots, A_m \in \mathbb{K}(\mathbb{H})$. Given any $\alpha_1, \alpha_2, \dots, \alpha_m \in \mathbb{R}$, $\sum_{i=1}^m \alpha_i A_i$ is a best coapproximation to T out of $\text{span}\{A_1, A_2, \dots, A_m\}$ if and only if given any $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$, there exists $x \in M_{\sum_{i=1}^m \beta_i A_i}$ such that $\langle \sum_{i=1}^m \beta_i A_i x, (T - \sum_{i=1}^m \alpha_i A_i)x \rangle = 0$.*

Proof. It follows from the definitions of Birkhoff-James orthogonality and best coapproximation that $\sum_{i=1}^m \alpha_i A_i$ is a best coapproximation to T out of $\text{span}\{A_1, A_2, \dots, A_m\}$ if and only if $A \perp_B (T - \sum_{i=1}^m \alpha_i A_i)$, for all $A \in \text{span}\{A_1, A_2, \dots, A_m\}$. Clearly, this is equivalent to the following:

$$\sum_{i=1}^m \beta_i A_i \perp_B (T - \sum_{i=1}^m \alpha_i A_i) \forall \beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}.$$

It follows from Theorem 2.2 of [10] that for any $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$, $M_{\sum_{i=1}^m \beta_i A_i}$ is the unit sphere of some subspace of \mathbb{H} . Now applying Theorem 2.2 of [11], we conclude that the above condition is equivalent to the existence of $x = x(\beta_1, \dots, \beta_m) \in M_{\sum_{i=1}^m \beta_i A_i}$ such that $\langle \sum_{i=1}^m \beta_i A_i x, (T - \sum_{i=1}^m \alpha_i A_i) x \rangle = 0$, for any $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$. This completes the proof of the theorem. \square

We establish some fundamental attributes of the newly introduced $*$ -Property which also plays an important role in our scheme. To begin with, we establish the basis invariance of equivalent components and the $*$ -Property.

Proposition 2.2. *Let Y be a subspace of \mathcal{D}_n and let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$, $\mathcal{B} = \{B_1, B_2, \dots, B_m\}$ be two bases of Y , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$ and $B_k = ((b_{11}^k, b_{22}^k, \dots, b_{nn}^k))$, for each $1 \leq k \leq m$. Then*

- (i) $P_i^+(\mathcal{A}) = P_i^+(\mathcal{B})$ and $P_i^-(\mathcal{A}) = P_i^-(\mathcal{B})$, $i \in \{1, 2, \dots, n\}$.
- (ii) For any $i \in \{1, 2, \dots, n\}$, the i -th component satisfies the $*$ -Property with respect to \mathcal{A} if and only if the i -th component satisfies the $*$ -Property with respect to \mathcal{B} .

Proof. (i) Consider the two matrices \tilde{A} and \tilde{B} as constructed in Definition 1.2. Since \mathcal{A} and \mathcal{B} are two bases of Y , so there exists an invertible matrix $Q = (q_{ij})_{1 \leq i, j \leq m}$ such that $\tilde{A} = \tilde{B}Q$, where $\tilde{a}_{ij} = \sum_{k=1}^m \tilde{b}_{ik} q_{kj}$, for any $1 \leq i \leq n, 1 \leq j \leq m$. The desired result then follows easily.

(ii) We first prove the necessary part. As before let $Q = (q_{ij})_{1 \leq i, j \leq m}$ be the invertible matrix such that $\tilde{A} = \tilde{B}Q$, where $\tilde{a}_{ij} = \sum_{k=1}^m \tilde{b}_{ik} q_{kj}$, for any $1 \leq i \leq n, 1 \leq j \leq m$. Since the i -th component satisfies the $*$ -Property with respect to \mathcal{A} , there exist $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$ such that

$$\left| \sum_{k=1}^m \beta_k a_{ii}^k \right| > \max \left\{ \left| \sum_{k=1}^m \beta_k a_{jj}^k \right| : 1 \leq j \leq n, j \notin P_i^+ \cup P_i^- \right\}.$$

Observe that $\tilde{A}\tilde{\beta} = \tilde{B}Q\tilde{\beta}$, where $\tilde{\beta} = (\beta_1 \ \beta_2 \ \dots \ \beta_m)^t$. Considering $\tilde{\gamma} = (\gamma_1 \ \gamma_2 \ \dots \ \gamma_m)^t = Q\tilde{\beta}$, it is easy to see that for any $r \in \{1, 2, \dots, n\}$,

$$\left| \sum_{k=1}^m \gamma_k b_{rr}^k \right| = \left| \sum_{k=1}^m \left(\sum_{j=1}^m q_{kj} \beta_j \right) \tilde{b}_{rk} \right| = \left| \sum_{j=1}^m \left(\sum_{k=1}^m \tilde{b}_{rk} q_{kj} \right) \beta_j \right| = \left| \sum_{j=1}^m \tilde{a}_{rj} \beta_j \right| = \left| \sum_{j=1}^m \beta_j a_{rr}^j \right|.$$

This immediately shows that the i -th component satisfies the $*$ -Property with respect to \mathcal{B} . This completes the necessary part. The sufficient part follows similarly. \square

In light of the above theorem, from now onwards we will not explicitly mention the choice of basis in the description of the $*$ -Property. Our next theorem essentially guarantees the existence of the $*$ -Property.

Theorem 2.3. *Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Then there exists $1 \leq i \leq n$ such that the i -th component satisfies the $*$ -Property.*

Proof. Let the i_1 -th, i_2 -th, \dots , i_p -th components represent all the nonequivalent components. For any $\tilde{w} = (\gamma_1, \gamma_2, \dots, \gamma_m) \in \mathbb{R}^m$, consider the set of scalars

$$S_{\tilde{w}} := \left\{ \left| \sum_{k=1}^m \gamma_k a_{i_1 i_1}^k \right|, \left| \sum_{k=1}^m \gamma_k a_{i_2 i_2}^k \right|, \dots, \left| \sum_{k=1}^m \gamma_k a_{i_p i_p}^k \right| \right\}.$$

Case 1 : If $S_{\tilde{w}}$ attains its maximum at a unique point, say at $|\sum_{k=1}^m \gamma_k a_{i_r i_r}^k|$, where $r \in \{1, 2, \dots, p\}$, then clearly the i_r -th component satisfies the $*$ -Property.

Case 2 : Let us assume that the maximum of $S_{\tilde{w}}$ is attained at exactly two points. Suppose that for $i_s, i_t \in \{i_1, i_2, \dots, i_p\}$ and $i_s \neq i_t$,

$$\left| \sum_{k=1}^m \gamma_k a_{i_s i_s}^k \right| = \left| \sum_{k=1}^m \gamma_k a_{i_t i_t}^k \right| > \max \left\{ \left| \sum_{k=1}^m \gamma_k a_{qq}^k \right| : q \in \{i_1, i_2, \dots, i_p\} \setminus \{i_s, i_t\} \right\}.$$

Let us define functions $f_s, f_t : \mathbb{R}^m \rightarrow \mathbb{R}$ given by

$$f_s(\tilde{u}) := |\langle \tilde{u}, \tilde{a}_{i_s} \rangle| \text{ and } f_t(\tilde{u}) := |\langle \tilde{u}, \tilde{a}_{i_t} \rangle|,$$

where $\tilde{u} = (\alpha_1, \alpha_2, \dots, \alpha_m) \in \mathbb{R}^m$ and $\tilde{a}_{i_s} = (a_{i_s i_s}^1, a_{i_s i_s}^2, \dots, a_{i_s i_s}^m)$, $\tilde{a}_{i_t} = (a_{i_t i_t}^1, a_{i_t i_t}^2, \dots, a_{i_t i_t}^m)$ are the i_s -th and the i_t -th component, respectively.

Let us also define another function, $g : \mathbb{R}^m \rightarrow \mathbb{R}$ given by

$$g(\tilde{u}) := \max \{ |\langle \tilde{u}, \tilde{a}_q \rangle| : q \in \{i_1, i_2, \dots, i_p\} \setminus \{i_s, i_t\} \},$$

where $\tilde{a}_q = (a_{qq}^1, a_{qq}^2, \dots, a_{qq}^m)$ is the q -th component. Since f_i, g are continuous function on \mathbb{R}^m , $\phi_i = f_i - g$ is also continuous and $\phi_i(\tilde{w}) > 0$, for all $i \in \{s, t\}$, where $\tilde{w} = (\gamma_1, \gamma_2, \dots, \gamma_m) \in \mathbb{R}^m$. It is easy to observe that there exists an open ball $\mathcal{B}_\delta(\tilde{w})$, with radius $\delta > 0$ and centered at \tilde{w} , such that $\phi_i(\tilde{y}) > 0$, for all $\tilde{y} \in \mathcal{B}_\delta(\tilde{w})$. Consider the hyperspaces H_1, H_2 of \mathbb{R}^m given by

$$H_1 = \{ \tilde{x} \in \mathbb{R}^m : \langle \tilde{x}, (\tilde{a}_{i_s} + \tilde{a}_{i_t}) \rangle = 0 \},$$

$$H_2 = \{ \tilde{x} \in \mathbb{R}^m : \langle \tilde{x}, (\tilde{a}_{i_s} - \tilde{a}_{i_t}) \rangle = 0 \}.$$

We note that $\{ \tilde{x} \in \mathbb{R}^m : f_s(\tilde{x}) = f_t(\tilde{x}) \} = H_1 \cup H_2$, which is a nowhere dense set in \mathbb{R}^m . Therefore, by choosing $\tilde{v} := (\beta_1, \beta_2, \dots, \beta_m) \in \mathcal{B}_\delta(\tilde{w}) \setminus (H_1 \cup H_2)$, we obtain that $f_s(\tilde{v}) \neq f_t(\tilde{v})$. Without loss of generality, assume $f_s(\tilde{v}) > f_t(\tilde{v})$. It is now easy to observe that

$$\left| \sum_{k=1}^m \beta_k a_{i_s i_s}^k \right| > \max \left\{ \left| \sum_{k=1}^m \beta_k a_{qq}^k \right| : 1 \leq q \leq n, q \notin P_{i_s}^+ \cup P_{i_s}^- \right\}.$$

Therefore, the i_s -th component satisfies the $*$ -Property.

Case 3 : Suppose that the maximum of $S_{\tilde{w}}$ is attained at $r (> 2)$ number of points and let the i_1 -th, i_2 -th, \dots , i_r -th components satisfy

$$\left| \sum_{k=1}^m \gamma_k a_{i_1 i_1}^k \right| = \dots = \left| \sum_{k=1}^m \gamma_k a_{i_r i_r}^k \right| > \max \left\{ \left| \sum_{k=1}^m \gamma_k a_{qq}^k \right| : q \in \{i_1, \dots, i_p\} \setminus \{i_1, \dots, i_r\} \right\}.$$

By similar argument as given in Case 2, it can be shown that at least one of the i_l -th components satisfies the *-Property, where $l \in \{1, 2, \dots, r\}$. This completes the theorem. \square

Remark 2.4. Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. In particular, for any $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$, there exists an i -th component such that $|\sum_{k=1}^m \beta_k A_k| = |\sum_{k=1}^m \beta_k a_{ii}^k|$, where the i -th component satisfies the *-Property.

Our next aim is to obtain a tractable necessary and sufficient condition for the *-Property. In this context, we first recall the definition of a normal cone. A non-empty set $K \subset \mathbb{R}^n$ is said to be a normal cone if the following three conditions are satisfied:

$$(i) u, v \in K \Rightarrow u + v \in K, (ii) u \in K, \alpha \geq 0 \Rightarrow \alpha u \in K, (iii) K \cap (-K) = \{\theta\}.$$

We define interior of the normal cone K , denoted by $int(K)$, as the collection of all interior points of the normal cone K . We refer the readers to [12], for an application of the notion of normal cones in studying approximate Birkhoff-James orthogonality in Banach spaces. We also require the following lemma for our purpose.

Lemma 2.5. Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Given any $i, j \in \{1, 2, \dots, n\}$, where $j \notin P_i^+ \cup P_i^-$, there exist a pair of normal cones whose interiors are the collection of all the $(\beta_1, \beta_2, \dots, \beta_m) \in \mathbb{R}^m$ such that $|\sum_{k=1}^m \beta_k a_{ii}^k| > |\sum_{k=1}^m \beta_k a_{jj}^k|$.

Proof. Let us consider the set

$$C := \left\{ (\beta_1, \beta_2, \dots, \beta_m) \in \mathbb{R}^m : \left| \sum_{k=1}^m \beta_k a_{ii}^k \right| > \left| \sum_{k=1}^m \beta_k a_{jj}^k \right| \right\}.$$

Let us also construct two sets K_1 and K_2 such that

$$K_1 := \left\{ (\beta_1, \beta_2, \dots, \beta_m) \in C : \sum_{k=1}^m \beta_k a_{ii}^k > 0 \right\} \cup \{\theta\},$$

$$K_2 := \left\{ (\beta_1, \beta_2, \dots, \beta_m) \in C : \sum_{k=1}^m \beta_k a_{ii}^k < 0 \right\} \cup \{\theta\}.$$

From the definition of K_1 and K_2 , it is evident that $K_1 = -K_2$. Now, it is immediate that $\tilde{x} \in K_1$ implies that $\alpha \tilde{x} \in K_1$, for all $\alpha \geq 0$. Therefore, to prove that K_1 is a normal cone, we only need to show $\tilde{u}, \tilde{v} \in K_1$ implies $\tilde{u} + \tilde{v} \in K_1$. Suppose that $\tilde{u} = (\alpha_1, \alpha_2, \dots, \alpha_m)$ and $\tilde{v} = (\gamma_1, \gamma_2, \dots, \gamma_m) \in K_1$. Then, $\sum_{k=1}^m (\alpha_k + \gamma_k) a_{ii}^k > 0$ and for any $j \in \{1, 2, \dots, n\}$ such that $j \notin P_i^+ \cup P_i^-$, it follows that

$$\begin{aligned} \left| \sum_{k=1}^m (\alpha_k + \gamma_k) a_{jj}^k \right| &\leq \left| \sum_{k=1}^m \alpha_k a_{jj}^k \right| + \left| \sum_{k=1}^m \gamma_k a_{jj}^k \right| < \left| \sum_{k=1}^m \alpha_k a_{ii}^k \right| + \left| \sum_{k=1}^m \gamma_k a_{ii}^k \right| \\ &= \left| \sum_{k=1}^m (\alpha_k + \gamma_k) a_{ii}^k \right|. \end{aligned}$$

This proves that K_1 (and therefore, K_2) is a normal cone. It is rather straightforward to verify that $int(K_1) = K_1 \setminus \{\theta\}$ and $int(K_2) = K_2 \setminus \{\theta\}$. Therefore, $int(K_1) \cup int(K_2) = C$, as desired. This completes the proof of the lemma. \square

Next we introduce the notion of associated pair of cones, which turns out to be useful in characterizing the $*$ -Property.

Definition 2.6. Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Given any $i \in \{1, 2, \dots, n\}$, we define the pair of normal cones $K_j^i, -K_j^i$ as the associated pair of cones of the i -th component with respect to the j -th component, given by

$$K_j^i := \left\{ (\beta_1, \beta_2, \dots, \beta_m) \in \mathbb{R}^m : \left| \sum_{k=1}^m \beta_k a_{ii}^k \right| > \left| \sum_{k=1}^m \beta_k a_{jj}^k \right| \text{ and } \sum_{k=1}^m \beta_k a_{ii}^k > 0 \right\} \cup \{\theta\},$$

for all $j \notin P_i^+ \cup P_i^-$.

Finally, we are in a position to characterize the $*$ -Property from a geometric perspective.

Theorem 2.7. Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Then for any $i \in \{1, 2, \dots, n\}$, the i -th component satisfies the $*$ -Property if and only if

$$\bigcap \{ \text{int}(K_j^i) \cup \text{int}(-K_j^i) : 1 \leq j \leq n, j \notin P_i^+ \cup P_i^- \} \neq \phi.$$

Proof. Suppose that the i -th component satisfies the $*$ -Property, i.e., there exists $\tilde{x} = (\beta_1, \beta_1, \dots, \beta_m) \in \mathbb{R}^m$ such that

$$\left| \sum_{k=1}^m \beta_k a_{ii}^k \right| > \max \left\{ \left| \sum_{k=1}^m \beta_k a_{jj}^k \right| : j \notin P_i^+ \cup P_i^- \right\}.$$

This is equivalent to $\tilde{x} \in \text{int}(K_j^i) \cup \text{int}(-K_j^i)$, for all $1 \leq j \leq n$ and $j \notin P_i^+ \cup P_i^-$, where $K_j^i, -K_j^i$ are the pair of associated cones of the i -th component with respect to the j -th component. Therefore,

$$\bigcap \{ \text{int}(K_j^i) \cup \text{int}(-K_j^i) : 1 \leq j \leq n, j \notin P_i^+ \cup P_i^- \} \neq \phi.$$

This completes the proof of the necessary part of the theorem. We note that the sufficient part of the theorem also follows from similar arguments and the definition of pair of associated cones. This establishes the theorem. \square

We next obtain a simple and useful sufficient condition for the $*$ -Property. It should be noted that in practise, the following result can be readily applied in most cases of the computations involving the $*$ -Property, since checking the linear independence of a given set of vectors is not complicated at all by virtue of the well-known method of row reduction of matrices.

Proposition 2.8. Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Suppose that the i -th component $(a_{ii}^1, a_{ii}^2, \dots, a_{ii}^m) \notin \text{span}\{(a_{jj}^1, a_{jj}^2, \dots, a_{jj}^m) : 1 \leq j \leq n, j \notin P_i^+ \cup P_i^-\}$, where $(a_{jj}^1, a_{jj}^2, \dots, a_{jj}^m)$ is the j -th component. Then the i -th component satisfies the $*$ -Property.

Proof. Let $Y_1 = \text{span}\{(a_{jj}^1, a_{jj}^2, \dots, a_{jj}^m) : 1 \leq j \leq n, j \notin P_i^+ \cup P_i^-\}$ and let $Y_2 = \text{span}\{(a_{jj}^1, a_{jj}^2, \dots, a_{jj}^m) : 1 \leq j \leq n\}$. Clearly, $Y_1 \subsetneq Y_2 = \mathbb{R}^m$, which implies that $Y_2^\perp \subsetneq Y_1^\perp$. Therefore, there exists $(\gamma_1, \gamma_2, \dots, \gamma_m) \in Y_1^\perp \setminus Y_2^\perp$ such

that $|\sum_{k=1}^m \gamma_k a_{ii}^k| > \max \{|\sum_{k=1}^m \gamma_k a_{jj}^k| : 1 \leq j \leq n, j \notin P_i^+ \cup P_i^-\} = 0$. In other words, the i -th component satisfies the $*$ -Property, as desired. \square

Remark 2.9. Suppose that \mathcal{T}_i is the collection of all those j such that the j -th component is a scalar multiple of the i -th component, where $i, j \in \{1, 2, \dots, n\}$. Let us assume that the i -th component $(a_{ii}^1, a_{ii}^2, \dots, a_{ii}^m) = c_j (a_{jj}^1, a_{jj}^2, \dots, a_{jj}^m)$, where $(a_{jj}^1, a_{jj}^2, \dots, a_{jj}^m)$ is the j -th component and $|c_j| \geq 1$, for all $j \in \mathcal{T}_i$. Also assume that $(a_{ii}^1, a_{ii}^2, \dots, a_{ii}^m) \notin \text{span}\{(a_{kk}^1, a_{kk}^2, \dots, a_{kk}^m) : 1 \leq k \leq n, k \notin \mathcal{T}_i\}$. Following similar argument from Proposition 2.8, the i -th component satisfies the $*$ -Property.

We are now ready to present a computationally convenient characterization of the best coapproximation to an element of \mathcal{M}_n out of a given subspace of \mathcal{D}_n .

Theorem 2.10. *Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Suppose that the j_1 -th, j_2 -th, \dots , j_r -th nonequivalent components satisfy the $*$ -Property. Then given any $T \in \mathcal{M}_n$, $\sum_{k=1}^m \alpha_k A_k$ is a best coapproximation to T out of $\text{span}\{A_1, A_2, \dots, A_m\}$ if and only if $\alpha_1, \alpha_2, \dots, \alpha_m \in \mathbb{R}$ satisfy the following relations:*

$$a_{j_p j_p}^1 \alpha_1 + a_{j_p j_p}^2 \alpha_2 + \dots + a_{j_p j_p}^m \alpha_m \in W(j_p T_*),$$

for all $p \in \{1, 2, \dots, r\}$, where $W(j_p T_*)$ is the numerical range of the $*$ -associated matrix of T corresponding to the j_p -th component.

Proof. Let us first prove the necessary part of the theorem. Assume that the j_s -th component satisfies the $*$ -Property, where $s \in \{1, 2, \dots, r\}$. Then there exists $\tilde{v} = (\beta_1, \beta_2, \dots, \beta_m) \in \mathbb{R}^m$ such that $|\sum_{k=1}^m \beta_k a_{j_s j_s}^k| > \max\{|\sum_{k=1}^m \beta_k a_{qq}^k| : 1 \leq q \leq n, q \notin P_{j_s}^+ \cup P_{j_s}^-\}$. Therefore, $\|\sum_{k=1}^m \beta_k A_k\| = |\sum_{k=1}^m \beta_k a_{j_s j_s}^k|$ and the norm attainment set of $\sum_{k=1}^m \beta_k A_k$ is

$$M_{\sum_{k=1}^m \beta_k A_k} = \{x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n : \|x\|_2 = 1, x_h = 0 \forall h \notin P_{j_s}^+ \cup P_{j_s}^-\}.$$

Let $T = (b_{uv})_{1 \leq u, v \leq n} \in \mathcal{M}_n$. Since $\sum_{k=1}^m \alpha_k A_k$ is a best coapproximation to T out of $\text{span}\{A_1, A_2, \dots, A_m\}$, it follows from Theorem 2.1 that $\langle \sum_{k=1}^m \beta_k A_k x, (T - \sum_{k=1}^m \alpha_k A_k)x \rangle = 0$, i.e., $\langle \sum_{k=1}^m \beta_k A_k x, T x \rangle = \langle \sum_{k=1}^m \beta_k A_k x, \sum_{k=1}^m \alpha_k A_k x \rangle$, for some $x \in M_{\sum_{k=1}^m \beta_k A_k}$. By a straight forward calculation, the previous equation can be expressed as

$$\begin{aligned} a_{j_s j_s}^1 \alpha_1 + \dots + a_{j_s j_s}^m \alpha_m &= \sum_{u \in P_{j_s}^+, v \in P_{j_s}^+ \cup P_{j_s}^-} b_{uv} x_u x_v - \sum_{u \in P_{j_s}^-, v \in P_{j_s}^+ \cup P_{j_s}^-} b_{uv} x_u x_v \\ &\in W(j_s T_*). \end{aligned}$$

Similarly, we can observe that for all $p \in \{1, 2, \dots, r\}$,

$$(1) \quad a_{j_p j_p}^1 \alpha_1 + a_{j_p j_p}^2 \alpha_2 + \dots + a_{j_p j_p}^m \alpha_m \in W(j_p T_*),$$

completing the proof of the necessary part.

We now prove the sufficient part of the theorem. For any $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$, not all zero, by virtue of Remark 2.4, there exists $t \in \{1, 2, \dots, r\}$ such that the j_t -th component satisfies $\|\sum_{k=1}^m \beta_k A_k\| = |\sum_{k=1}^m \beta_k a_{j_t j_t}^k|$. Let $P_{j_t}^+ = \{j_t, j_{t_2}, \dots, j_{t_w}\}$

and $P_{j_t}^- = \{j_{t_{w+1}}, j_{t_{w+2}}, \dots, j_{t_v}\}$, so that $|P_{j_t}^+ \cup P_{j_t}^-| = v$. From the hypothesis, $\alpha_1, \alpha_2, \dots, \alpha_m \in \mathbb{R}$ satisfy the following relations:

$$a_{j_t j_t}^1 \alpha_1 + a_{j_t j_t}^2 \alpha_2 + \dots + a_{j_t j_t}^m \alpha_m \in W(j_t T_*).$$

Therefore, there exists $y = (y_t, y_{t_2}, \dots, y_{t_v}) \in \mathbb{R}^v$ with $\|y\|_2 = 1$ such that

$$(2) \quad a_{j_t j_t}^1 \alpha_1 + a_{j_t j_t}^2 \alpha_2 + \dots + a_{j_t j_t}^m \alpha_m = \langle j_t T_* y, y \rangle.$$

Now by taking $\hat{y} = (\tilde{y}_1, \tilde{y}_2, \dots, \tilde{y}_n) \in \mathbb{R}^n$ such that $\tilde{y}_h = 0 \forall j_h \notin P_{j_t}^+ \cup P_{j_t}^-$ and $\tilde{y}_h = y_h \forall j_h \in P_{j_t}^+ \cup P_{j_t}^-$, it is easy to observe that $\hat{y} \in M_{\sum_{k=1}^m \beta_k A_k}$. By some easy calculations and by using the equation (2), we conclude that

$$\left\langle \sum_{k=1}^m \beta_k A_k \hat{y}, \left(T - \sum_{k=1}^m \alpha_k A_k \right) \hat{y} \right\rangle = \left(\sum_{k=1}^m \beta_k a_{j_t j_t}^k \right) \left[\langle y, j_t T_* y \rangle - \sum_{k=1}^m \alpha_k a_{j_t j_t}^k \right] = 0.$$

The sufficient part of the theorem now follows directly from Theorem 2.1. This establishes the theorem. □

Remark 2.11. Suppose that $A_1, A_2, \dots, A_m \in \mathcal{M}_n$, where $1 \leq m \leq n$, are such that $A_i A_j^t, A_i^t A_j$ are symmetric, for all $i, j \in \{1, 2, \dots, m\}$. Then from Corollary 9 of [6], there exist orthogonal matrices P and Q such that $P^t A_i Q = D_i$, where $D_i \in \mathcal{D}_n$, for all $i \in \{1, 2, \dots, m\}$. Moreover, since P and Q are orthogonal matrices, it is easy to see that $\|\sum_{i=1}^m \beta_i A_i\| = \|\sum_{i=1}^m \beta_i D_i\|$, for all $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$.

The above remark allows us to present the following strengthened version of Theorem 2.10.

Theorem 2.12. Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{M}_n such that $A_i A_j^t, A_i^t A_j$ are symmetric, for all $i, j \in \{1, 2, \dots, m\}$. Let $D_1, D_2, \dots, D_m \in \mathcal{D}_n$ be such that $D_i = P^t A_i Q$, where $D_i = ((d_{11}^i, d_{22}^i, \dots, d_{nn}^i))$, for all $i \in \{1, 2, \dots, m\}$ and $P, Q \in \mathcal{M}_n$ are orthogonal matrices. Suppose that the j_1 -th, j_2 -th, \dots, j_r -th nonequivalent components satisfy the $*$ -Property (with respect to $\text{span}\{D_1, D_2, \dots, D_m\}$). Then given any $T \in \mathcal{M}_n$, $\sum_{i=1}^m \alpha_i A_i$ is a best coapproximation to T out of $\text{span}\{A_1, A_2, \dots, A_m\}$ if and only if $\alpha_1, \alpha_2, \dots, \alpha_m \in \mathbb{R}$ satisfy the following relations:

$$d_{j_p j_p}^1 \alpha_1 + d_{j_p j_p}^2 \alpha_2 + \dots + d_{j_p j_p}^m \alpha_m \in W(j_p (P^t T Q)_*),$$

for all $p \in \{1, 2, \dots, r\}$, where $W(j_p (P^t T Q)_*)$ is the numerical range of the $*$ -associated matrix of $P^t T Q$ corresponding to the j_p -th component.

Proof. $\sum_{i=1}^m \alpha_i A_i$ is a best coapproximation to T out of $\text{span}\{A_1, A_2, \dots, A_m\}$ if and only if given any $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$, there exists $x \in M_{\sum_{i=1}^m \beta_i A_i}$ such that $\langle (\sum_{i=1}^m \beta_i A_i) x, (T - \sum_{i=1}^m \alpha_i A_i) x \rangle = 0$, i.e.,

$$\left\langle \left(P \sum_{i=1}^m \beta_i D_i Q^t \right) x, \left(T - P \sum_{i=1}^m \alpha_i D_i Q^t \right) x \right\rangle = 0.$$

So, for $y = Q^t x$, it is immediate that $\langle (\sum_{i=1}^m \beta_i D_i) y, (P^t T Q - \sum_{i=1}^m \alpha_i D_i) y \rangle = 0$. We also note that $x \in M_{\sum_{i=1}^m \beta_i A_i}$ if and only if $y = Q^t x \in M_{\sum_{i=1}^m \beta_i D_i}$. Therefore, $\sum_{i=1}^m \alpha_i A_i$ is a best coapproximation to T out of $\text{span}\{A_1, A_2, \dots, A_m\}$ if and only if $\sum_{i=1}^m \alpha_i D_i$ is a best coapproximation to $P^t T Q$ out of $\text{span}\{D_1, D_2, \dots, D_m\}$. Now the desired result follows directly from Theorem 2.10. This completes the proof of the theorem. □

To illustrate the applicability of Theorem 2.10 from a computational point of view, we next present a series of explicit numerical examples elaborating the different features of the best coapproximation problem, related to the existence and the uniqueness. In each case, an algorithmic approach is presented which further underlines the usefulness of the *-Property in studying best coapproximation problems in subspaces of \mathcal{D}_n .

Example 2.13. Let $A_1 = ((7, -5, 2, 6, -7, -5, 1))$, $A_2 = ((1, 3, 4, 3, -1, 3, 2))$, $A_3 = ((3, -7, -4, 5, -3, -7, -2))$ be linearly independent matrices in \mathcal{D}_7 . Our aim is to find the best coapproximation(s) to any given T out of $\mathbb{Y} = \text{span}\{A_1, A_2, A_3\}$. In view of the Theorem 2.10, we proceed in the following steps.

Step 1 : For $i \in \{1, 2, \dots, 7\}$, the i -th components are respectively

$$(7, 1, 3), (-5, 3, -7), (2, 4, -4), (6, 3, 5), (-7, -1, -3), (-5, 3, -7), (1, 2, -2).$$

Step 2 : $P_1^+ = \{1\}, P_1^- = \{5\}; P_2^+ = \{2, 6\}, P_2^- = \phi; P_3^+ = \{3\}, P_3^- = \phi; P_4^+ = \{4\}, P_4^- = \phi; P_5^+ = \{5\}, P_5^- = \{1\}; P_6^+ = \{2, 6\}, P_6^- = \phi; P_7^+ = \{7\}, P_7^- = \phi$, respectively, where P_i^+ and P_i^- are the positively associated set and the negatively associated set of the i -th component, respectively, for all $i \in \{1, 2, \dots, 7\}$.

Step 3 : Here the nonequivalent components satisfying the *-Property may be taken as the 1-st component, the 2-nd component, the 3-rd component and the 4-th component.

Step 4 : In this final step, we consider a given $T \in \mathcal{M}_7$ and apply Theorem 2.10 to obtain the best coapproximation to T out of \mathbb{Y} . In order to illustrate the various possibilities arising in the best coapproximation problem in \mathcal{D}_7 , it suffices to consider the following three particular cases.

Case 1 : Let $T_1 \in \mathcal{M}_7$ be given by $T_1 = (b_{ij})_{1 \leq i, j \leq 7}$, where $b_{11} = 2, b_{15} = 4, b_{22} = 1, b_{26} = 3, b_{33} = 4, b_{44} = 1, b_{51} = -7, b_{55} = -2, b_{62} = 2, b_{66} = 1$ and the other b_{ij} 's can be chosen arbitrarily.

$$\text{Therefore, } {}^1T_{1*} = \begin{pmatrix} 2 & 4 \\ 7 & 2 \end{pmatrix}, {}^2T_{1*} = \begin{pmatrix} 1 & 3 \\ 2 & 1 \end{pmatrix}, {}^3T_{1*} = (4), {}^4T_{1*} = (1).$$

Then from Theorem 2.10, $\sum_{i=1}^3 \alpha_i A_i$ is a best coapproximation to T_1 out of \mathbb{Y} if and only if $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ satisfies the following relations:

$$\begin{aligned} 7\alpha_1 + \alpha_2 + 3\alpha_3 &\in W({}^1T_{1*}) = [-7/2, 15/2] \\ -5\alpha_1 + 3\alpha_2 - 7\alpha_3 &\in W({}^2T_{1*}) = [-3/2, 7/2] \\ 2\alpha_1 + 4\alpha_2 - 4\alpha_3 &\in W({}^3T_{1*}) = \{4\} \\ 6\alpha_1 + 3\alpha_2 - 5\alpha_3 &\in W({}^4T_{1*}) = \{1\}. \end{aligned}$$

Since there are infinitely many $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ satisfying the above relations, there are infinitely many best coapproximation to T_1 out of \mathbb{Y} . Moreover,

$$\mathcal{R}_{\mathbb{Y}}(T_1) = \{((x, 4-x, 4, 1, -x, 4-x, 2)) : 1/2 \leq x \leq 11/2\}.$$

Case 2 : Let $T_2 \in \mathcal{M}_7$ be given by $T_1 = (c_{ij})_{1 \leq i, j \leq 7}$, where $c_{11} = 3$, $c_{15} = -5$, $c_{22} = 1$, $c_{26} = 3$, $c_{33} = 4$, $c_{44} = 1$, $c_{51} = -5$, $c_{55} = -3$, $c_{62} = 2$, $c_{66} = 1$ and the other c_{ij} 's can be chosen arbitrarily.

$$\text{Therefore, } {}^1T_{2*} = \begin{pmatrix} 3 & -5 \\ 5 & 3 \end{pmatrix}, {}^2T_{2*} = \begin{pmatrix} 1 & 3 \\ 2 & 1 \end{pmatrix}, {}^3T_{2*} = (4), {}^4T_{2*} = (1).$$

Then from Theorem 2.10,

$$\begin{aligned} 7\alpha_1 + \alpha_2 + 3\alpha_3 \in W({}^1T_{2*}) &= \{3\} \\ -5\alpha_1 + 3\alpha_2 - 7\alpha_3 \in W({}^2T_{2*}) &= [-3/2, 7/2] \\ 2\alpha_1 + 4\alpha_2 - 4\alpha_3 \in W({}^3T_{2*}) &= \{4\} \\ 6\alpha_1 + 3\alpha_2 - 5\alpha_3 \in W({}^4T_{2*}) &= \{1\}. \end{aligned}$$

Since there exist unique $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ satisfying the above relations, the best coapproximation to T out of \mathbb{Y} is unique. Moreover,

$$\mathcal{R}_{\mathbb{Y}}(T_2) = \{((3, 1, 4, 1, -3, 1, 2))\}.$$

Case 3 : Let $T_3 \in \mathcal{M}_7$ be given by $T_3 = (d_{ij})_{1 \leq i, j \leq 7}$, where $d_{11} = 14$, $d_{15} = 1$, $d_{22} = 1$, $d_{26} = 3$, $d_{33} = 4$, $d_{44} = 1$, $d_{51} = 1$, $d_{55} = -14$, $d_{62} = 2$, $d_{66} = 1$ and the other d_{ij} 's can be chosen arbitrarily.

$$\text{Therefore, } {}^1T_{3*} = \begin{pmatrix} 14 & 1 \\ -1 & 14 \end{pmatrix}, {}^2T_{3*} = \begin{pmatrix} 1 & 3 \\ 2 & 1 \end{pmatrix}, {}^3T_{3*} = (4), {}^4T_{3*} = (1).$$

Then from Theorem 2.10,

$$\begin{aligned} 7\alpha_1 + \alpha_2 + 3\alpha_3 \in W({}^1T_{3*}) &= \{14\} \\ -5\alpha_1 + 3\alpha_2 - 7\alpha_3 \in W({}^2T_{3*}) &= [-3/2, 7/2] \\ 2\alpha_1 + 4\alpha_2 - 4\alpha_3 \in W({}^3T_{3*}) &= \{4\} \\ 6\alpha_1 + 3\alpha_2 - 5\alpha_3 \in W({}^4T_{3*}) &= \{1\}. \end{aligned}$$

Since there exists no such $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ satisfying the above relations, it follows that

$$\mathcal{R}_{\mathbb{Y}}(T_3) = \phi.$$

Our next goal is to obtain a tractable characterization of the coproximal subspaces of \mathcal{D}_n with respect to \mathcal{M}_n . The following lemma is crucial for that purpose, besides being interesting in its own right by providing a lower bound on the number of nonequivalent components satisfying the *-Property.

Lemma 2.14. *Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Let the total number of nonequivalent components satisfying the *-Property be p . Then $p \geq m$.*

Proof. Suppose that the j_1 -th, j_2 -th, \dots , j_p -th nonequivalent components satisfy the *-Property. Suppose on the contrary that $p < m$. Let $Y_1 = \text{span}\{(a_{j_s j_s}^1, a_{j_s j_s}^2, \dots, a_{j_s j_s}^m) : 1 \leq s \leq p\}$ and let $Y_2 = \text{span}\{(a_{ii}^1, a_{ii}^2, \dots, a_{ii}^m) : 1 \leq i \leq n\}$. Clearly, $Y_1 \subsetneq Y_2 = \mathbb{R}^m$, which implies that $Y_2^\perp \subsetneq Y_1^\perp$. Therefore, there exists $(\gamma_1, \gamma_2, \dots, \gamma_m) \in Y_1^\perp \setminus Y_2^\perp$ such that $|\sum_{k=1}^m \gamma_k a_{ii}^k| > \max\{|\sum_{k=1}^m \gamma_k a_{j_s j_s}^k| : 1 \leq s \leq p\} = 0$, for some $i \notin \{j_1, j_2, \dots, j_p\}$. Following Theorem 2.3, we obtain that the i -th component, which is nonequivalent to the j_1 -th, j_2 -th, \dots , j_p -th components, satisfies the *-Property. This contradiction completes the proof of the lemma.

□

We now obtain a characterization of the coproximal subspaces of \mathcal{D}_n in terms of the *-Property.

Theorem 2.15. *Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Then $\text{span}\{A_1, A_2, \dots, A_m\}$ is a coproximal subspace of \mathcal{M}_n if and only if there exist exactly m number of nonequivalent components satisfying the *-Property.*

Proof. Let us first prove the sufficient part of the theorem. Let the j_1 -th, j_2 -th, \dots , j_m -th components be chosen as the nonequivalent m number of components satisfying the *-Property. Let us consider $C \in \mathcal{M}_m$ given by $C = (c_{st})_{1 \leq s, t \leq m}$ such that $c_{st} = a_{j_s j_s}^t$, where $(a_{j_s j_s}^1, a_{j_s j_s}^2, \dots, a_{j_s j_s}^m)$ is the j_s -th component. We claim that $\text{rank}(C) = m$. Suppose on the contrary $\text{rank}(C) < m$. Let $Y_1 = \text{span}\{(a_{j_s j_s}^1, a_{j_s j_s}^2, \dots, a_{j_s j_s}^m) : 1 \leq s \leq m\}$ and let $Y_2 = \text{span}\{(a_{ii}^1, a_{ii}^2, \dots, a_{ii}^m) : 1 \leq i \leq n\}$. Clearly, $Y_1 \subsetneq Y_2 = \mathbb{R}^m$, which implies that $Y_2^\perp \subsetneq Y_1^\perp$. Therefore, there exists $(\gamma_1, \gamma_2, \dots, \gamma_m) \in Y_1^\perp \setminus Y_2^\perp$ such that $|\sum_{k=1}^m \gamma_k a_{ii}^k| > \max\{|\sum_{k=1}^m \gamma_k a_{j_s j_s}^k| : 1 \leq s \leq m\} = 0$, for some $i \notin \{j_1, j_2, \dots, j_m\}$. Following Theorem 2.3, there exists an i -th component, which is nonequivalent to the j_1 -th, j_2 -th, \dots , j_m -th components, that satisfies the *-Property. This contradiction establishes our claim. Therefore, C is invertible and hence onto. So, for any $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$, there always exist $\alpha_1, \alpha_2, \dots, \alpha_m \in \mathbb{R}$ such that $a_{j_s j_s}^1 \alpha_1 + a_{j_s j_s}^2 \alpha_2 + \dots + a_{j_s j_s}^m \alpha_m = \beta_s$, for all $s \in \{1, 2, \dots, m\}$. Noting that for any $T \in \mathcal{M}_n$, $W(j^s T_*) \subset \mathbb{R}$, therefore we conclude that

$$(3) \quad a_{j_s j_s}^1 \alpha_1 + a_{j_s j_s}^2 \alpha_2 + \dots + a_{j_s j_s}^m \alpha_m \in W(j^s T_*),$$

for all $s \in \{1, 2, \dots, m\}$. Following Theorem 2.10, it is now evident that $\sum_{k=1}^m \alpha_k A_k$ is the best coapproximation to T out of $\text{span}\{A_1, A_2, \dots, A_m\}$. This shows that $\text{span}\{A_1, A_2, \dots, A_m\}$ is a coproximal subspace of \mathcal{M}_n .

Let us now prove the necessary part of the theorem. Suppose that the j_1 -th, j_2 -th, \dots , j_p -th nonequivalent components satisfy the *-Property. Then from Lemma 2.14, we get that $p \geq m$. Let us now take the $p \times m$ matrix $D = (d_{st})_{1 \leq s \leq p, 1 \leq t \leq m}$ such that $d_{st} = a_{j_s j_s}^t$, where $(a_{j_s j_s}^1, a_{j_s j_s}^2, \dots, a_{j_s j_s}^m)$ is the j_s -th component. Let $T_D \in \mathbb{L}(\mathbb{H}_1, \mathbb{H}_2)$ be the linear operator corresponding to the matrix D with respect to the standard ordered bases of $\mathbb{H}_1, \mathbb{H}_2$, where $\mathbb{H}_1 = \mathbb{R}^m$ and $\mathbb{H}_2 = \mathbb{R}^p$. Since $\text{span}\{A_1, A_2, \dots, A_m\}$ is a coproximal subspace of \mathcal{M}_n , for $T \in \mathcal{M}_n$, there exist $\alpha_1, \alpha_2, \dots, \alpha_m \in \mathbb{R}$ satisfying the following relations:

$$(4) \quad a_{j_s j_s}^1 \alpha_1 + a_{j_s j_s}^2 \alpha_2 + \dots + a_{j_s j_s}^m \alpha_m \in W(j^s T_*),$$

for all $s \in \{1, 2, \dots, p\}$. We now claim that T_D is onto. Let $\beta = (\beta_1, \beta_2, \dots, \beta_p) \in \mathbb{R}^p$. We note that for any $T = (b_{ij})_{1 \leq i, j \leq n}$, $j^s T_*$ is a $h \times h$ matrix whose entries are $(\pm b_{ij})$ depending on $P_{j_s}^+$ and $P_{j_s}^-$, where $|P_{j_s}^+ \cup P_{j_s}^-| = h$. So we can choose T suitably so that $W(j^s T_*) = \{\beta_s\}$ for each $s \in \{1, 2, \dots, p\}$. This shows that for each $s \in \{1, 2, \dots, p\}$, we get

$$a_{j_s j_s}^1 \alpha_1 + a_{j_s j_s}^2 \alpha_2 + \dots + a_{j_s j_s}^m \alpha_m = \beta_s$$

and so $T_D(\alpha) = \beta$, where $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_m) \in \mathbb{R}^m$. Thus T_D is onto and therefore, $m \geq p$. This along with Lemma 2.14 completes the proof. □

We now obtain a characterization of the co-Chebyshev subspaces of \mathcal{D}_n in terms of the *-Property.

Theorem 2.16. *Let $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ be linearly independent in \mathcal{D}_n , where $A_k = ((a_{11}^k, a_{22}^k, \dots, a_{nn}^k))$, for each $1 \leq k \leq m$. Suppose that the i_1 -th, i_2 -th, \dots , i_p -th nonequivalent components satisfy the *-Property. Then $\text{span}\{A_1, A_2, \dots, A_m\}$ is a co-Chebyshev subspace of \mathcal{M}_n if and only if $p = m$ and $|P_{i_s}^+ \cup P_{i_s}^-| = 1$ for all $s \in \{1, 2, \dots, p\}$.*

Proof. We first prove the sufficient part of the theorem. Since $p = m$, we note from Theorem 2.15 that $\text{span}\{A_1, A_2, \dots, A_m\}$ is a coproximal subspace of \mathcal{M}_n . Therefore, for any given $T \in \mathcal{M}_n$, there exist $\alpha_1, \alpha_2, \dots, \alpha_m \in \mathbb{R}$ satisfying the following relations:

$$(5) \quad a_{i_s i_s}^1 \alpha_1 + a_{i_s i_s}^2 \alpha_2 + \dots + a_{i_s i_s}^m \alpha_m \in W({}^{i_s}T_*),$$

for all $s \in \{1, 2, \dots, m\}$. Since $|P_{i_s}^+ \cup P_{i_s}^-| = 1$, it follows that ${}^{i_s}T_*$ is of order 1. Moreover, $W({}^{i_s}T_*) = (b_{i_s i_s})$, for every $s \in \{1, 2, \dots, m\}$, where $T = (b_{ij})_{1 \leq i, j \leq n}$. Therefore, relations (5) represent a system of linear equation with coefficient matrix $C = (c_{st})_{1 \leq s, t \leq m}$, where $c_{st} = a_{i_s i_s}^t$. Following the arguments given in the proof of Theorem 2.15, we conclude that C is invertible. Hence for any given $T \in \mathcal{M}_n$, there exists a unique $(\alpha_1, \alpha_2, \dots, \alpha_m) \in \mathbb{R}^m$ satisfying the relations (5). Therefore, $\text{span}\{A_1, A_2, \dots, A_m\}$ is a co-Chebyshev subspace of \mathcal{M}_n .

We now prove the necessary part of the theorem. Let us assume that $\text{span}\{A_1, A_2, \dots, A_m\}$ is a co-Chebyshev subspace of \mathcal{M}_n . In particular, $\text{span}\{A_1, A_2, \dots, A_m\}$ is a coproximal subspace of \mathcal{M}_n . Therefore, from Theorem 2.15, we get $p = m$. Suppose on the contrary $|P_{i_s}^+ \cup P_{i_s}^-| = k_s > 1$, for some $s \in \{1, 2, \dots, m\}$. Therefore, ${}^{i_s}T_*$ is of order k_s . Let us consider $Q = (q_{rt})_{1 \leq r \leq p, 1 \leq t \leq m}$, where $q_{rt} = a_{i_r i_r}^t$. Since $p = m$, following the arguments given in the proof of Theorem 2.15, Q is invertible. Now, for any two scalars c and d , where $c \neq d$, we can choose a suitable $T \in \mathcal{M}_n$ such that $c, d \in W({}^{i_s}T_*)$. Therefore, we can conclude that there exist at least two different sets of $(\alpha_1, \alpha_2, \dots, \alpha_m) \in \mathbb{R}^m$ satisfying the relations:

$$a_{i_l i_l}^1 \alpha_1 + a_{i_l i_l}^2 \alpha_2 + \dots + a_{i_l i_l}^m \alpha_m \in W({}^{i_l}T_*),$$

for all $l \in \{1, 2, \dots, m\}$. This contradicts that $\text{span}\{A_1, A_2, \dots, A_m\}$ is a co-Chebyshev subspace of \mathcal{M}_n . Hence the theorem. \square

As an immediate application of the above theorem, we record the following interesting observation.

Corollary 2.17. \mathcal{D}_n is a co-Chebyshev subspace of \mathcal{M}_n .

Proof. Clearly, $\{E_k : 1 \leq k \leq n\}$ is a basis of \mathcal{D}_n , where $E_k = (e_{ij}^k)_{1 \leq i, j \leq n}$ is given by

$$\begin{aligned} e_{ij}^k &= 1, \text{ whenever } i = j = k \\ &= 0, \text{ otherwise.} \end{aligned}$$

It is trivial to observe that for all $1 \leq i \leq n$, the i -th component satisfies the *-Property and $|P_i^+ \cup P_i^-| = 1$. Therefore, the desired result follows directly from Theorem 2.16. \square

As another important application of the theories developed in the present article, it is possible to characterize the best coapproximation problem in the setting of ℓ_∞^n , for any given $n \in \mathbb{N}$. This in turn is equivalent to the following optimization problem:

Problem: Let $a_{ij}, \alpha_k \in \mathbb{R}$ be fixed, where $1 \leq i \leq m, 1 \leq j, k \leq n$. Find a necessary and sufficient condition for the existence of $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$ such that for any $c_1, c_2, \dots, c_m \in \mathbb{R}$, the following inequality holds true:

$$\max \left\{ \left| \alpha_k - \sum_{i=1}^m c_i a_{ik} \right| : 1 \leq k \leq n \right\} \geq \max \left\{ \left| \sum_{i=1}^m \beta_i a_{ik} - \sum_{i=1}^m c_i a_{ik} \right| : 1 \leq k \leq n \right\}.$$

Moreover, in case existence is guaranteed, find all such $\beta_1, \beta_2, \dots, \beta_m \in \mathbb{R}$.

We emphasize that the above problem is not entirely trivial, most notably because the existence of a desired solution is not a priori guaranteed. However, it is possible to completely solve the problem (from both theoretical and computational perspectives), by applying the methodology already developed in this article. It is well-known that ℓ_∞^n (endowed with its usual maximum norm) is isometrically isomorphic to \mathcal{D}_n endowed with the usual operator norm. Indeed, the natural choice map $\Psi : \ell_\infty^n \rightarrow \mathcal{D}_n$, taking $(a_1, a_2, \dots, a_n) \in \ell_\infty^n$ to $((a_1, a_2, \dots, a_n)) \in \mathcal{D}_n$ is easily seen to be the desired isometric isomorphism. This connection allows us to obtain an algorithmic approach to the best coapproximation problem in any given subspace \mathbb{Y} of ℓ_∞^n via the methods already developed to treat the corresponding best coapproximation problem in the subspace $\Psi(\mathbb{Y})$ of \mathcal{D}_n . It should be noted in this context that our theory essentially translates into characterizing the best coapproximation(s) to any given $T \in \mathcal{M}_n$ out of any given subspace of ℓ_∞^n , and therefore, the best coapproximation problem in subspaces of ℓ_∞^n is only a particular case of the results developed so far. To illustrate this further, we make note of the following two remarks pertaining to the best coapproximation problem in ℓ_∞^n :

- ℓ_∞^n is a coproximal subspace of \mathcal{M}_n for each $n \in \mathbb{N}$. This is simply a reformulation of Corollary 2.17.
- By using the concept of the *-Property, and the above mentioned isometric isomorphism $\Psi : \ell_\infty^n \rightarrow \mathcal{D}_n$, it is quite straightforward to construct subspaces of ℓ_∞^n which are (not) coproximal. Indeed, in light of the theories developed in this article, any such construction essentially reduces to controlling the number (nonequivalent) of i -th components satisfying the *-Property, for $1 \leq i \leq n$. As an explicit example, it can be readily verified that \mathbb{Y}_1 is a coproximal subspace of ℓ_∞^7 , whereas \mathbb{Y}_2 is not, where

$$\begin{aligned} \mathbb{Y}_1 &= \text{span} \{ (6, 1, 4, 3, 3, 1, 1), (2, 5, 2, 3, 1, 5, 1), (4, 3, 8, 6, 2, 3, 2), (2, 1, 4, 9, 1, 1, 3) \}, \\ \mathbb{Y}_2 &= \text{span} \{ (2, -5, 3, 1, -2, -5, 2), (-4, 2, 2, -2, -4, 2, -4) \}. \end{aligned}$$

In view of our treatment of the theory of coapproximations in ℓ_∞^n spaces, it seems appropriate to end the present article with the following concluding remark:

Remark 2.18. The theory of best coapproximations in Banach spaces remains a much less exposed area of research, especially from a computational point of view, in comparison to the theory of best approximations. Certainly, this is in part due to the inherently complicated non-linear nature of the best coapproximation problem and the difficulty of the corresponding computations involved in the process. In this context, the reader is encouraged to look up the literature, including [7, 8]. Our main focus in this article is to illustrate the following principle in the setting of ℓ_∞^n (or, more generally, for matrices in \mathcal{M}_n out of subspaces of \mathcal{D}_n):

It is possible to essentially reduce the much harder “best coapproximation problem” to the well-known and way more simpler “existence and uniqueness problem corresponding to a particular system of linear problems”, by applying the concept of orthogonality.

Indeed, using the methodology developed so far, it is now very easy to explicitly produce examples of coproximinal and co-Chebyshev subspaces in the setting of ℓ_∞^n . Therefore, in light of the above fact, a natural query would be to test the validity of such a nicety, in the setting of classical Banach spaces other than ℓ_∞^n .

REFERENCES

- [1] Birkhoff, G., *Orthogonality in linear metric spaces*, Duke Math. J. **1** (1935), 169–172.
- [2] Bhatia, R., Šemrl, P., *Orthogonality of matrices and distance problem*, Linear Alg. Appl., **287** (1999), 77–85.
- [3] Franchetti, C., Furi, M., *Some characteristic properties of real Hilbert spaces*, Rev. Roumaine Math. Pures Appl., **17** (1972), 1045–1048.
- [4] Gustafson, K. E., Rao, D. K. M., *Numerical range*, Springer-Verlag New York, DOI: 10.1007/978-1-4613-8498-4.
- [5] James, R. C., *Orthogonality and linear functionals in normed linear spaces*, Trans. American Math. Soc., **61** (1947b), 265–292.
- [6] Maehara, T., Murota, K., *Simultaneous singular value decomposition*, Linear Alg. Appl., **435** (2011), 106–116.
- [7] Papini, P. L., Singer, I., *Best coapproximation in normed linear spaces*, Mh. Math., **88** (1979) 27–44.
- [8] Rao, G.S., Swaminathan, M., *Best coapproximation and schauder bases in Banach spaces*, Acta Sci. Math., **54** (1990), 339–354.
- [9] Sain, D., *Birkhoff-James orthogonality of linear operators on finite dimensional Banach spaces*, J. Math. Anal. Appl., **447** (2017), 860–866.
- [10] Sain, D., Paul, K., *Operator norm attainment and inner product spaces*, Linear Alg. Appl., **439** (2013), 2448–2452.
- [11] Sain, D., Mal, A., Paul, K., *Some remarks on Birkhoff-James orthogonality of linear operators*, Expo. Math., **38** (2020), 138–147.
- [12] Sain, D., Paul, K., Mal, A., *On approximate Birkhoff-James orthogonality and normal cones in a normed space*, J. Convex Anal., **26** (2019), 341–351.
- [13] Sain, D., *On best approximations to compact operators*, Proc. Amer. Math. Soc., **149** (2021), 4273–4286.

(Sain) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BENGALURU 560012, KARNATAKA, INDIA

Email address: saindebmalya@gmail.com

(Sohel) DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, WEST BENGAL, INDIA

Email address: shamimsohel11@gmail.com

(Ghosh) DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, WEST BENGAL, INDIA

Email address: `sghosh0019@gmail.com`

(Paul) DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, WEST BENGAL, INDIA

Email address: `kalloldada@gmail.com`