Quantum group coproducts and universality under scalar extensions

Alexandru Chirvasitu

Abstract

We characterize the families of bialgebras or Hopf algebras over fields for which the product in the corresponding category is finite-dimensional, answering a question of M. Lorenz: if the ground field is infinite then bialgebra or Hopf products are finite-dimensional precisely when the factors are, with at most one of dimension > 1; over finite fields the necessary and sufficient condition is instead that factors be finite-dimensional with at most finitely many of dimension > 1; finally, these statements hold for coalgebras as well, provided the family is finite. We also characterize (a) finite field extensions as precisely those whose underlying scalar extension functor preserves coalgebra, or bialgebra, or Hopf algebra products (correcting an error in the literature); (b) algebraic field extensions as those along which finite coalgebra (bialgebra, Hopf algebra) products are preserved; and (c) again algebraic field extensions as precisely those which intertwine cofree coalgebras on vector spaces, or cofree bialgebras (Hopf algebras) on algebras.

Key words: Hopf algebra; Tannaka reconstruction; adjoint functor; affine algebra; algebraic extension; bialgebra; coalgebra; cofree; comonadic; coproduct; coradical; cosemisimple; finite dual; free; geometric quotient; locally presentable category; monadic; product; pseudo-pullback; sink

MSC 2020: 16T05; 16T15; 18A30; 12F05; 14L30; 20G15; 16T20; 18N10; 18C15; 18C20

Contents

1	Small coalgebra (bialgebra, Hopf algebra) products	3
2	Transporting universal constructions along field extensions	10
References		24

Introduction

The present paper is in part motivated by the natural question [24] of when and how often products $\prod_{\text{HALG}} H_i$ in the category \mathbb{R} HALG of Hopf algebras over a field \mathbb{R} are finite-dimensional. The quantum

groups in the title are those attached to the Hopf algebras in question *contra*variantly, as common in non-commutative geometry (e.g. [10, Chapters 7 and 13]); it is in this sense that we are here concerned with *co*products: dual counterparts to Hopf (and more generally, bialgebra or coalgebra) products.

That the categories $_{\Bbbk}$ HALG (along with those of bialgebras and coalgebras, $_{\Bbbk}$ BIALG and $_{\Bbbk}$ COALG) admit arbitrary (co)limits and afford a wealth of universal constructions ((co)free objects, etc.) is

well familiar, and the literature on category-theoretic aspects of the subject abounds: [3, 4, 11, 12, 17, 31, 32, 33, 39], [38, §§6.4, 12.2 and so on], [14, §1.6], etc. etc. All categories mentioned are in fact (e.g. by [33, Lemmas 1 and 2 and Theorem 6]) *locally presentable* [2, Definition 1.17] and are thus amenable to various adjoint functor theorems [2, §§0.7 and 1.66].

Devoted to variations on that initial motivating question, Section 1 confirms (in Theorem 1.4, Corollary 1.8 and Theorem 1.9) that it is only in the self-evident, trivial cases that a coalgebra (bialgebra, Hopf algebra) product can be small. The focus is specifically on the three categories COALG, BIALG and HALG for definiteness: the reader can easily supply analogues for Hopf algebras with bijective antipode and cousins (e.g. Hopf algebras with antipode S satisfying $S^{2d} = \text{id}$ for some fixed d).

Theorem A Let \Bbbk be a field.

(1) If k is infinite, a product $\prod_i H_i$ in the category of k-bialgebras or Hopf k-algebras is finitedimensional if and only if all H_i are, with at most one of dimension > 1.

(2) If k is finite, such a product is infinite-dimensional all H_i are, with only finitely many of dimension > 1.

(3) All of the above applies to finite families of k-coalgebras.

In part prompted by the natural need to adjust the ground field in addressing Theorem A, and in part in order to correct an error (of the author's) in the literature, Section 2 turns to permanence properties for universal constructions (products, cofree objects) in categories of coalgebras or bialgebras under extending scalars via the functor $(-)_{\Bbbk'} := - \otimes_{\Bbbk} \Bbbk'$ attached to a field extension $\Bbbk \leq \Bbbk'$.

The mistake alluded to above is the claim [12, Theorem 4.1(2)] that $(-)_{k'}$ preserves arbitrary products of coalgebras, or bialgebras, or Hopf algebras: Examples 2.1 and 2.2 each give instances of failure (or rather a families of them). Allowing for variations in taste, the corrected statement(s) may well be more interesting:

- that specific permanence property in fact characterizes the *finite* field extensions (Theorem 2.6);
- and furthermore, there is an analogue characterizing *algebraic* extensions by relaxing the product-preservation property appropriately (Theorem 2.13).

Theorem B (1) A field extension $\mathbb{k} \leq \mathbb{k}'$ is finite if and only if the corresponding scalar extension functor $(-)_{\mathbb{k}'}$

• is a right adjoint between the corresponding categories of coalgebras (equivalently, bialgebras or Hopf algebras);

• or preserves products in COALG (again equivalently, in BIALG or HALG).

(2) Similarly, a field extension $\mathbb{k} \leq \mathbb{k}'$ is finite if and only if the corresponding scalar extension functor $(-)_{\mathbb{k}'}$ preserves finite limits (equivalently, finite products) in any one of the categories COALG, BIALG or HALG.

That discussion in turn generates a bit of downstream momentum: it is clear from the above statement that, in analyzing ways in which $(-)_{k'}$ might fail to preserve various universal constructions, the distinction between algebraic and transcendental field extensions $k \leq k'$ is crucial. The following characterization (Theorem 2.9 below) of algebraic extensions is an upshot of that analysis, and in the same spirit as Theorem B.

Theorem C A field extension $\mathbb{k} \leq \mathbb{k}'$ is algebraic if and only if the corresponding scalar extension functor $(-)_{\mathbb{k}'}$

• preserves cofree coalgebras on arbitrary vector spaces or equivalently, cofree bialgebras (or Hopf algebras) on arbitrary algebras;

• equivalently, any of the above for only the 1-dimensional vector spaces or algebras.

Acknowledgements

I am grateful for insightful comments, questions and suggestions from A. Agore, M. Lorenz and G. Militaru. This work is partially supported by NSF grant DMS-2001128.

1 Small coalgebra (bialgebra, Hopf algebra) products

All (co)algebras are assumed (co)unital, along with their respective morphisms. Everything in sight is linear over a field k, and we frequently denote scalar extensions along $\mathbb{k} \to \mathbb{k}'$ by $(-)_{\mathbb{k}'}$ (mostly fields, but the notation applies generally, for ring morphisms). For background on coalgebras, bialgebras and Hopf algebras we refer the reader to [14, 27, 34, 38]; more specific citations are scattered throughout the text, where needed.

As is customary in the category-theoretic literature (e.g. [1, Definition 19.3]), (\top) symbols occasionally indicate adjunctions by having the narrow end point towards the left adjoint. $\mathcal{C}(-,-)$ stands for morphisms in the category \mathcal{C} . Some common categories in use below include

• $_{\bullet}\mathcal{M}$ and \mathcal{M}_{\bullet} , left and right modules respectively and similarly, \mathcal{M}^{\bullet} and $^{\bullet}\mathcal{M}$ for comodules;

• additional 'f' subscripts indicate finite-dimensional objects therein, as in \mathcal{M}_f^C for finite-dimensional right C-comodules;

• VECT, ALG, COALG, BIALG and HALG denote categories of vector spaces, algebras, coalgebras, bialgebras and Hopf algebras respectively, occasionally decorated with the ground field for clarity or emphasis (e.g. _kCOALG for k-coalgebras).

We need some 2-categorical (or bicategorical) background, as the reader can find covered in [16, 20, 21] and numerous other sources. In particular, extend the notion of a *pseudo-pullback* of [21, §6.10, Example 15] from pairs of functors to families of co-terminal functors (i.e. *sinks* [1, Definition 10.62]).

Definition 1.1 A pseudo-pullback PPLB($F_i, i \in I$) is the pseudo-limit [21, §6.10] of the diagram consisting of a family of functors $C_i \xrightarrow{F_i} C$ (a sink with codomain C and domain $(C_i)_i$). Concretely, it is the category with

• objects consisting of tuples $(c_i \in C_i, i \in I; c \in C)$ and C-isomorphisms $F_i c_i \xrightarrow{\varphi_i} c_i;$

• and morphisms $(c_i; c; \varphi_i) \to (c'_i; c'; \varphi'_i)$ consisting of \mathcal{C}_i -morphisms $c_i \to c'_i$ and a \mathcal{C} -morphism $c \to c'$ making the obvious diagrams (involving φ_i and φ'_i) commute.

PPLB(F_i), in words, is the universal category equipped with functors to the C_i and C and natural isomorphisms



the top left-hand arrows are $(c_i; c; \varphi_i) \mapsto c_i$, the bottom arrow selects the *c* instead, and the natural isomorphisms are what the φ_i respectively aggregate to.

That in place, note the following representation-theoretic description of the product $\prod_i C_i$ of a family of coalgebras in the category thereof.

Theorem 1.2 Let $(C_i)_{i \in I}$ be a family of coalgebras and $C := \prod C_i$ the product in the category $_{\Bbbk}COALG$.

(1) The corestriction functors $\mathcal{M}^C \to \mathcal{M}^{C_i}$ induced by the product structure morphisms $C \xrightarrow{\pi_i} C_i$ realize \mathcal{M}^C as the pseudo-pullback of the forgetful functors $\mathcal{M}^{C_i} \to VEC$.

- (2) The same goes for categories $\mathcal{M}_{f}^{\bullet}$ of finite-dimensional comodules.
- (3) The statements hold also for categories of bialgebras or Hopf algebras.

Proof This follows straightforwardly from standard *Tannaka reconstruction* [37, Theorem 2.1.12 and Lemma 2.2.1] of coalgebras and morphisms from their respective categories of comodules. In one formulation [26, Proposition 3.3] particularly convenient here,

$$\operatorname{COALG} \ni C \longmapsto \left(\mathcal{M}_{f}^{C} \xrightarrow{\text{forgetful functor } \operatorname{Fgt}} \operatorname{Vect}_{f} \right)$$

is a right *biadjoint* [16, Definition 9.8] from

• the category of coalgebras, enhanced to a bicategory by equipping it with 2-morphisms



consisting of functionals $\psi \in C^* = \text{Vect}(C, \Bbbk)$ intertwining the algebra morphisms f^* and g^* , in the sense that

 $\psi \cdot f^*(\theta) = g^*(\theta) \cdot \psi$ in the dual algebra [34, §2.3] C^* , $\forall \theta \in D^*$.

• and a certain bicategory consisting of categories equipped with functors $\rightarrow \text{VECT}_f$ and *actions* [26, post Example 2.3] by the monoidal category VECT_f .

That right biadjoint will in particular preserve products, and checking that these are precisely the usual coalgebra products in the former bicategory and pseudo-pullbacks in the latter is straightforward.

The statement for \mathcal{M}^C (arbitrary, rather than finite-dimensional comodules) follows by realizing \mathcal{M}^C as a *cocompletion* [2, Definition 1.44] of \mathcal{M}_f^C (every comodule being canonically the *filtered* [2, Definition 1.4] union of its finite-dimensional subcomodules [27, Finiteness Theorem 5.1.1]).

As for the bialgebra- and Hopf-algebra-flavored versions, simply observe that products in either category coincide with those of the underlying algebras. For the functors

 ${}_{\Bbbk}\mathrm{HAlg} \xleftarrow{\mathrm{Incl}}{}_{\Bbbk}\mathrm{BiAlg} \xrightarrow{\mathrm{Fgt}}{}_{\Bbbk}\mathrm{Coalg}$

are right adjoints [32, Proposition 47 and Theorem 54] (and indeed, by [32, Proposition 47 3.(b)] and [33, Theorem 10 2.], *monadic* in the sense of [7, Definition 4.4.1]), so by [8, Proposition 3.2.2] also *continuous* (i.e. [25, $\SV.4$] preserve small limits). In particular, said functors preserve products.

The following bit of language (and notation) will occasionally be useful.

Definition 1.3 Let $(C_i)_i$ be a family of coalgebras. The corresponding category $\mathcal{M}^{(C_i)}$ of (C_i) -*comodules* is by definition that of *C*-comodules for the product $C := \prod C_i$ in the category COALG.
Per Theorem 1.2, it consists of vector spaces

- carrying separate C_i -comodule structures for all i;
- with those structures leaving invariant an exhaustive filtration by finite-dimensional subspaces.

Given the product coincidence for coalgebras, bialgebras and Hopf algebras (remarked upon in the course of the proof of Theorem 1.2), the terminology just introduced will be used unambiguously for bialgebras and Hopf algebras as well (as in speaking of (H_i) -comodules for $H_i \in \text{HALG}$, etc.).

Theorem 1.4 For a family $(H_i)_{i \in I}$ of bialgebras over an infinite field the following conditions are equivalent.

- (a) The H_i are finite-dimensional, with at most one of dimension > 1.
- (b) The product $H := \prod_i H_i$ in the category of bialgebras is finite-dimensional.
- (c) H has finitely many (isomorphism classes of) simple comodules.

(d) For every $d \in \mathbb{Z}_{\geq 0}$, H has finitely many (isomorphism classes of) simple d-dimensional comodules.

For both Theorem 1.4 and later in Theorem 1.9, when discussing coalgebras, it will be convenient to have to handle binary families of objects only.

Lemma 1.5 Let $(C_i)_{i \in I}$ be a family of \Bbbk -coalgebras.

(1) Suppose I is finite and the coalgebra product $\prod_{I} C_{i}$

- either is finite dimensional;
- or has finitely many simple comodules;
- or has finitely many simple comodules in any given dimension.

Then the same, respectively, goes for every product $\prod_J C_j$ over every subfamily $J \subseteq I$.

(2) If the C_i are bialgebras (or Hopf algebras) then the above holds for arbitrary, possibly infinite families.

Proof Both remarks are simple enough, but the second is particularly straightforward. As in all categories C with zero objects (both initial and final [8, pre Proposition 4.5.16]), in both BIALG and HALG the canonical morphisms $\prod_{I} c_i \to \prod_{J} c_j$, $J \subseteq I$ to products of subfamilies split:



for any subset $J \subseteq I$, with

$$\iota_i := \begin{cases} \text{product structure map } \pi_i & \text{if } i \in J \\ \text{zero morphism} & \text{otherwise.} \end{cases}$$

 \prod_{J} is thus a retract of \prod_{I} , and the properties in the statement are plainly inherited by retracts. We thus turn to (1).

Fix a non-zero finite-dimensional $(C_l)_{l \in I \setminus J}$ -comodule W (in the sense of Definition 1.3); one such exists, given the finiteness of I (the finiteness of $I \setminus J$ suffices). If $(V_t)_t$ are infinitely many simple $(C_j)_{j \in J}$ -comodules (of bounded dimension) then $(V_t \otimes W)_t$ carry obvious $(C_i)_{i \in I}$ -comodule (respectively of bounded dimension). They are semisimple and *isotypic* (i.e. sums of copies of single simple comodules) over $(C_j)_{j \in J}$, so some infinite family (respectively of bounded dimension) of mutually distinct simple $(C_i)_I$ -comodules can be extracted from among their subquotients.

This, so far, takes care of the two last bullet points in (1). The first can be handled similarly, noting that infinite-dimensionality will provide an infinite family (V_t) of mutually non-isomorphic $(C_j)_J$ -comodules that are either

- simple;
- or non-split extensions of two fixed simple $(C_i)_J$ -comodules.

Either way, the argument above applies.

As Remark 1.6(2) below observes, one cannot, generally, drop the finiteness requirement in Lemma 1.5(1).

Proof of Theorem 1.4 The implications

(1-1) (a)
$$\xrightarrow{1-\text{dimensional object is terminal}}$$
 (b) \implies (c) \implies (d)

being obvious, we focus, for the duration of the proof, on $(d) \Rightarrow (a)$.

It is enough, by Lemma 1.5(2), to consider binary products $H := H_{\alpha} \times H_{\beta}$ of finite-dimensional bialgebras H_i , $i = \alpha, \beta$. Throughout, we use the description of (finite-dimensional) $(H_i)_i$ -comodules implicit in Theorem 1.2: vector spaces equipped with separate H_i -comodule structures. More specifically, this means H_i -comodules V_i together with linear isomorphisms $H_i \xrightarrow{\varphi_i} V$ to a common vector space V, morphisms $(V, V_i, \varphi) \to (V', V'_i, \varphi'_i)$ being commutative diagrams

$$V_{i} \xrightarrow{\varphi_{i}} V \xrightarrow{\psi \in \operatorname{VEC}} V'_{i}$$
$$\xrightarrow{\cong} V'_{i}$$
$$\xrightarrow{\varphi_{i}'} V',$$

(same ψ for both $i = \alpha, \beta$). There are several cases to consider, along with a more broadly-applicable preamble.

(I) A general argument. We will construct $(H_i)_{i=\alpha,\beta}$ -comodules W equipped with isomorphisms $W_i \xrightarrow{\varphi_i} W$, $W_i \in \mathcal{M}_f^{H_i}$, $i = \alpha, \beta$ varying the choices involved. Specifically, in each case considered below the W_i will be fixed and the φ_i will vary. Those choices will always ensure that W is simple over $H := H_{\alpha} \times H_{\beta}$ (bialgebra product), so it will suffice to argue that there are infinitely many mutually non-isomorphic choices of φ_i .

Upon fixing a pair $(\varphi_{i,0})i = \alpha, \beta$, arbitrary pairs $(\varphi_i)_i$ are in bijection with the group $GL(W)^2$: simply compose the selected $\varphi_{i,0}$ separately with elements of the general linear group of GL(W). We will regard GL(W) as the group of \Bbbk -points of a linear algebraic (\Bbbk) -group [9, §I.1.1]: the A-points, for commutative \Bbbk -algebras A, are defined as $GL(W_A)$. Bold fonts will distinguish between schemes (as in $\mathbf{GL}(W)$) and the corresponding groups of k-points (such as GL(W)). We similarly have groups $GL_i(W_i)$, $i = \alpha, \beta$ of comodule automorphisms, as k-points of algebraic groups $\mathbf{GL}_i(W_i)$.

We will furthermore assume \Bbbk is large enough for Jordan-Hölder filtrations of the W_{α} involved to be stable (i.e. the simple subquotients stay simple after extending scalars to $\overline{\Bbbk}$). This is always achievable by extending scalars along some *finite* field extension $\Bbbk \leq \Bbbk'$, and such extensions do not affect products (Theorem 2.6 below, where we discuss the compatibility between field extensions and universal constructions).

The fixed $\varphi_{i,0}$ transport comodule automorphisms over to W, realizing embeddings $\mathbf{GL}_i(W_i) \to \mathbf{GL}(W)$. The set of isomorphism classes of $(\varphi_{\alpha,\beta}) \in \mathcal{M}^{(H_{\alpha,\beta})}$ is then identifiable with the double coset space

(1-2)
$$GL(W)\backslash GL(W)^2/GL_{\alpha}(W_{\alpha}) \times GL_{\beta}(W_{\beta}),$$

with the left-hand GL acting on $(\varphi_{\alpha,\beta})$ by simultaneous composition (i.e. via the diagonal embedding $GL \to GL^2$) and the right-hand $GL_{\alpha} \times GL_{\beta}$ acting by precomposition in the two GL(W) factors. To compress the notation, set

$$\mathbf{X} := \mathbf{GL}(W)^2 \quad \text{(the scheme in (1-2) being acted upon)} \\ \mathbf{G} := \mathbf{GL}(W) \times \mathbf{GL}_{\alpha}(W_{\alpha}) \times \mathbf{GL}_{\beta}(W_{\beta}) \quad \text{(the algebraic group in (1-2) doing the acting)}$$

Orbit spaces such as the

$$X/G, \quad X := \mathbf{X}(\Bbbk), \quad G := \mathbf{G}(\Bbbk)$$

of (1-2) might not be the space of k-points of a scheme (the familiar issues attending the formation of geometric quotient [9, §§6.1 and 6.16], [28, Definition 0.6], etc.). This can nevertheless be remedied [35, Theorem and opening paragraph] upon passing to a dense open **G**-invariant $\mathbf{U} \subseteq \mathbf{X}$. Because furthermore the resulting quotient \mathbf{U}/\mathbf{G} will be unirational [9, §13.7] (being surjected upon by a dense open subspace **U** of affine space), the infinitude of k implies [9, §13.7, concluding sentence] that $\mathbf{U}(\mathbf{k})/\mathbf{G}(\mathbf{k})$ is infinite (and hence so is (1-2)) provided it has positive dimension. Addressing that positivity hinges on (what we refer to as) the crude dimension (or size) estimate

(1-3)
$$\dim \mathbf{X} > \dim \mathbf{G} - 1 \quad \text{or} \quad \dim GL(W) > \dim GL_{\alpha}(W_{\alpha}) + \dim GL_{\beta}(W_{\beta}) - 1,$$

the -1 to account for the fact that diagonal scaling by the left-hand GL in (1-2) coincides with scaling by the diagonal copy of \Bbbk in $GL_{\alpha} \times GL_{\beta}$ on the right. We next tackle the aforementioned cases.

(II) Each H_i , $i = \alpha, \beta$ has a simple comodule V_i of respective dimension $d_i \ge 2$. The W_i of the above sketch will be $V_i^{\oplus d/d_i}$ respectively, with $d = \operatorname{lcm}(d_\alpha, d_\beta)$. The common vector space receiving the isomorphisms $W_i \xrightarrow{\varphi_i} W$ is thus *d*-dimensional.

The resulting (H_i) -comodule structures are indeed simple, for any non-zero subcomodule must restrict to a sum of copies of each V_i , and hence have dimension divisible by both d_i so also by d. The crude estimate (1-3) reads

$$d^{2} > \left(\frac{d}{d_{\alpha}}\right)^{2} + \left(\frac{d}{d_{\beta}}\right)^{2} - 1$$
 i.e. $1 > \frac{1}{d_{\alpha}^{2}} + \frac{1}{d_{\beta}^{2}} - \frac{1}{d^{2}}$

indeed valid by the assumption that $d_{\alpha,\beta} \geq 2$ (note that the -1 correction of (1-3) is not even needed in this case).

(III) H_{α} has a simple comodule V_{α} of dimension $d_{\alpha} \geq 2$ and H_{β} does not. H_{β} , in other words, is *pointed* [34, Definition 3.4.4]: its simple right comodules are all 1-dimensional. Set $d := \operatorname{lcm}(d_{\alpha}, 2)$ (so $d = d_{\alpha}$ if the latter is even and $2d_{\alpha}$ if not). $W_{\alpha} \cong V_{\alpha}^{\oplus d/d_{\alpha}}$ again, while W_{β} will be a sum of $\frac{d}{2}$ copies of a single 2-dimensional H_{β} -comodule

 $(1-4) V_{\beta} := \begin{cases} V_{\beta,0} \oplus V_{\beta,1} & \dim V_{\beta,s} = 1, \ V_{\beta,s} \text{ non-isomorphic if they exist} \\ a & \text{non-split} & \text{self-} \\ extension & \text{of} & a & 1- \\ \dim extensional & \text{comodule} & \text{otherwise} \\ V_{\beta,0} \end{cases}$

If d_{α} is even then W is certainly simple as an $H_{\alpha,\beta}$ -comodule, for over H_{α} it is W_{α} . Otherwise (assuming d_{α} odd, that is), we can force simplicity as follows.

Having fixed a specific decomposition $W \cong W_{\alpha} \cong V_{\alpha}^2$, all other copies of V_{α} embedded therein as H_{α} -subcomodules are precisely the graphs $\Gamma_{\lambda}, \lambda \in \mathbb{k}^{\times}$ of the scaling maps from one distinguished V_{α} summand to the other. Any non-zero $H_{\alpha,\beta}$ -subcomodule of W will of course intersect its socle (i.e. [14, §3.1] largest semisimple subobject) $W_{\beta} = V_{\beta,0}^{\frac{d}{2}}$ as an H_{β} -comodule. Restricting to an open dense choice of $\varphi_{\alpha,\beta}$, we can assume said socle is similarly the graph Γ_T of an isomorphism

$$V_{\alpha} \xrightarrow{T = T_{\varphi_{\alpha,\beta}}} V_{\alpha}$$

The elements of that socle belonging to some H_{α} -subcomodule V_{α} , then, are (identifiable with) the eigenspaces of T. Now, because $d_{\alpha} \geq 3$ (being ≥ 2 and odd), a non-zero $H_{\alpha,\beta}$ -subcomodule, if proper, must

- be isomorphic to V_{α} as an H_{α} -comodule, coinciding with one of the $\Gamma_{\lambda}, \lambda \in \mathbb{k}^{\times}$;
- and also intersect Γ_T along at least a *two*-dimensional space.

To avoid this, it is enough to range over those choices of $\varphi_{\alpha,\beta}$ for which the corresponding T has no repeated eigenvalues (an open dense condition).

Having thus ensured that W is simple, it remains to verify the estimate (1-3); here, it translates to

(1-5)
$$\begin{cases} d_{\alpha}^2 > 1 + 2\left(\frac{d_{\alpha}}{2}\right)^2 - 1 = \frac{d_{\alpha}^2}{2} \iff d_{\alpha} > 0 & \text{if } d_{\alpha} \text{ is even} \\ (2d_{\alpha})^2 > 4 + 2d_{\alpha}^2 - 1 = 3 + 2d_{\alpha}^2 \iff d_{\alpha} > 1 & \text{if } d_{\alpha} \text{ is odd} \end{cases}$$

Both inequalities, of course, hold. And once more, the -1 terms on the right-hand side is not needed, on either line of (1-5).

(IV) $H_{\alpha,\beta}$ are both pointed. W will now be 2-dimensional, and its comodule structures W_{α} and W_{β} are as in (1-4) (for α , now, as well as β).

W will be simple as soon as the socles $V_{i,0} \subset V_i$, $i = \alpha, \beta$ are distinct, an open dense condition in the space of choices of $(\varphi_{\alpha,\beta})$. As for the size comparison (1-3), here it is 4 > 2 + 2 - 1; indeed valid (with the -1 term needed, for once). **Remarks 1.6** (1) The infinitely many simple (H_i) -comodules provided by Theorem 1.4 for binary (say) families dim $H_{\alpha,\beta} > 1$ certainly need not have unbounded dimensions, in general: see Example 1.7 below.

(2) Despite COALG having "the same" products as BIALG and HALG (proof of Theorem 1.2), the statement of Theorem 1.4 will not go through in that generality for coalgebras: if a (necessarily infinite) coalgebra family C_i is such that

$$\sup_{i} \inf_{0 \neq V \in \mathcal{M}^{C_i}} \dim V = \infty$$

(e.g. if C_i is a family of matrix coalgebras $M_{n_i}^*$ with $\{n_i\}_i$ unbounded) then no finite-dimensional vector space can support C_i -comodule structures simultaneously for all *i*. By the *comodule finiteness theorem* [27, 5.1.1] the coalgebra product $\prod C_i$ must be trivial (i.e. the 0 coalgebra).

(3) Note the contrast between the situation of (2) above and the dual version: algebras over a field always embed into their coproduct in $_{k}$ ALG, for instance by the standard normal-form description of that coproduct [6, Corollary 8.1].

(4) To pursue the duality issue, the counterpart to (2) is the remark that the coproducts of bialgebras or Hopf algebras are the same as those of the underlying algebras. The dual version of Theorem 1.4, moreover, is easily checked (again standard, via normal forms).

Having observed (in the proof of Lemma 1.5) that BIALG and HALG have zero objects, there are canonical morphisms

(1-6)
$$\coprod_i H_i \longrightarrow \prod_i H_i.$$

These do not appear to be, however, especially useful in deducing Theorem 1.4 from its dual variant: (1-6) are not, generally, embeddings (so that even if the domain is infinite-dimensional, it is not clear, a priori, that this entails the infinite dimensionality of the codomain).

To see this, consider what happens when all bialgebras H_i happen to be commutative: the subcategories

<sup>**BIALG**
$$\subset$$
 ^{**BIALG** and ^{**BIALG** \subset ^{**BIALG**}}}</sup>

of commutative objects being *reflective* by [33, diagram (9)] (i.e. [8, Definition 3.5.2] full, with the inclusions being right adjoints), they are automatically closed under limits [8, Proposition 3.5.3]. In particular, the bialgebra product of commutative bialgebras is automatically commutative. Naturally, their coproduct will not be (save for trivial particular cases).

Example 1.7 Take for both $H_{\alpha,\beta}$ the function algebra on the group $\mathbb{Z}/2$:

$$H_{\alpha} := \mathbb{k}^{\mathbb{Z}/2} =: H_{\beta},$$

with the pointwise multiplication and the coalgebra structure dualizing the multiplication on $\mathbb{Z}/2$. A finite-dimensional (H_i) -comodule is then nothing but a finite-dimensional representation (over \Bbbk) of the *infinite dihedral group* [36, Exercise 11.63]

group coproduct
$$(\mathbb{Z}/2) * (\mathbb{Z}/2) \cong \mathbb{Z} \rtimes \mathbb{Z}/2$$

(semidirect product [36, Theorem 7.2 and Definition preceding it]) for the inversion $\mathbb{Z}/2$ -action on \mathbb{Z} .

For algebraically closed k irreducible finite-dimensional $(\mathbb{Z} \rtimes \mathbb{Z}/2)$ -modules are at most 2dimensional: a generator for \mathbb{Z} will have a λ -eigenvector for some $\lambda \in \mathbb{k}^{\times}$, and the $\mathbb{Z}/2$ -factor will swap such an eigenvector with one with eigenvalue λ^{-1} .

Theorem 1.4 cannot quite go through in precisely the same form over finite fields, but the needed adjustments are not overly drastic.

Corollary 1.8 If the ground field k is finite, the conditions of Theorem 1.4 satisfy

$$(a) \iff (b) \iff (c) \Longrightarrow (d).$$

The last condition is strictly weaker, holding whenever all H_i are finite-dimensional, with at most finitely many of dimension > 1.

Proof We of course still have (1-1) in any case, and the last claim is simple: a *d*-dimensional vector space over a finite field is of course finite, and supports finitely many joint comodule structures over the finitely many (> 1)-dimensional, finite-dimensional H_i .

It thus remains to argue that the implication (c) \Rightarrow (a) (equivalently, its contrapositive) still holds over finite fields, restricting the discussion, once again, to pairs of bialgebras.

If $H_{\alpha,\beta}$ both have dimension > 1, Theorem 1.4 shows that for some $d \in \mathbb{Z}_{\geq 0}$ a *d*-dimensional \overline{k} -vector space V over the algebraic closure $\overline{k} \supset \overline{k}$ supports infinitely many mutually non-isomorphic pairs of $(H_{\alpha,\beta})_{\overline{k}}$ -comodule structures (i.e. *H*-comodule structures V_t). Observe that in general, for any \overline{k} -coalgebra C, every finite-dimensional (or simple) $C_{\overline{k}}$ -comodule arises as a subquotient of a comodule of the form

 $W_{\overline{k}}, \quad W \in {}_{\mathbb{k}}\mathcal{M}^C$ finite-dimensional (respectively simple).

Applying this to $C = H_{\alpha,\beta}$, there must be *H*-comodules (necessarily mutually non-isomorphic, and possibly of differing, possibly unbounded dimensions) recovering the V_t .

As to coalgebras (as opposed to bialgebras), the phenomenon noted in Remark 1.6(2), necessarily requiring *infinite* families, turns out to be the only impediment to Theorem 1.4 (and Corollary 1.8) going through in full generality.

Theorem 1.9 The conclusions of Theorem 1.4 and Corollary 1.8 hold in $_{\Bbbk}$ COALG for finite families of coalgebras.

Proof Having reduced the matter to binary families by Lemma 1.5(1), the proofs of Theorem 1.4 and Corollary 1.8 apply verbatim.

2 Transporting universal constructions along field extensions

The proof of Theorem 1.4 remarks in passing on the behavior of coalgebra (bialgebra, Hopf algebra) products under extending the ground field; it is perhaps apposite then, at this stage, to point out an error in the literature. [12, Theorem 4.1(2)] claims that scalar extension $(-)_{k'} = - \otimes_k k'$ along a field extension $k \leq k'$ is a right adjoint between the respective categories COALG, BIALG and HALG. This is not so: the functor preserves equalizers ([12, Lemma 4.2] appears to be correct), but not, in general, arbitrary products (there is a flaw in the proof of [12, Lemma 4.3]).

That in general infinite coalgebra products cannot be preserved by infinite field extensions can be seen as follows.

Example 2.1 Recall the *cofree coalgebra* construction

$$_{\Bbbk}$$
VECT $\ni V \longmapsto C_{\rm CF}(V) \in _{\Bbbk}$ COALG

of [14, §1.6] (or [38, §6.4]): the right adjoint to the forgetful functor COALG \rightarrow VECT. Being a right adjoint it preserves limits, and hence also products. In the commutative diagram

$$(\prod_{i} C_{\rm CF}(V_i))_{\Bbbk'} \xrightarrow{\qquad} \prod_{i} C_{\rm CF}(V_i)_{\Bbbk'} \xrightarrow{\qquad} \prod_{i} V_{i,\Bbbk'}$$

the double-headed arrows are indeed surjections [14, Exercise 1.6.2], whereas the right-hand bottom arrow *never* is for infinite $\mathbb{k} \leq \mathbb{k}'$ and infinitely many non-vanishing V_i . The top left-hand arrow thus fails to be an isomorphism (indeed, surjective) in all of those cases (infinite field extensions, infinite vector-space families).

Even more drastically, it is possible, for infinite field extensions $\mathbb{k} \leq \mathbb{k}'$, for the coalgebra product of C_i to vanish while that of $(C_i)_{\mathbb{k}'}$ does not (Example 2.2).

Example 2.2 Suppose the algebraic closure $\overline{\Bbbk} \supset \Bbbk$ is infinite (equivalently, by the celebrated Artin-Schreier theorem [19, Theorem 11.14], \Bbbk is neither algebraically closed nor real closed in the sense of [18, §5.1], [22, p.87], etc.).

Let $\mathbb{k}_i \supset \mathbb{k}$ be finite extensions with $\{[\mathbb{k}_i : \mathbb{k}]\}_i$ unbounded, and set $C_i := \mathbb{k}_i^*$, the dual \mathbb{k} -coalgebra. No finite-dimensional \mathbb{k} -vector space carries \mathbb{k}_i -linear structures for all i simultaneously, so the coalgebra product $\prod_i C_i$ vanishes as in Remark 1.6(2). On the other hand, $\mathbb{k}_i \otimes_{\mathbb{k}} \overline{\mathbb{k}}$ all have 1-dimensional modules (by the *Hilbert Nullstellensatz* [5, Corollary 7.10], for they are all finite-dimensional commutative algebras over an algebraically closed field). The coalgebra product $\prod_i (C_i)_{\overline{\mathbb{k}}}$ thus has at least a 1-dimensional comodule, so cannot vanish.

Remark 2.3 It will become clear later (Theorem 2.13) that even *finite* coalgebra products are not preserved by arbitrary field extensions.

The following simple strengthening of [14, Exercise 1.6.2] recalled in Example 2.1 above (the surjectivity of the canonical map $C_{\rm CF}(V) \to V$ from the cofree coalgebra) will be useful later. The cofree Hopf algebra $H_{\rm CF}(A) \to A$ on an algebra, referred to in the statement, is (the image of A through) the right adjoint to the forgetful functor HALG \to ALG. That forgetful functor is a left adjoint by, say, [4, Theorem 2.5], and indeed (by [33, Theorem 10 1.]) comonadic (the notion categorically dual to [7, Definition 4.4.1]).

Lemma 2.4 For any k-algebra A the cofree Hopf algebra structure map $H_{CF}(A) \to A$ is onto.

Proof Were it not, all algebra maps into A with Hopf domain would factor through some proper subspace of A (one subspace, valid for all Hopf algebras). On the other hand though, every $a \in A$ is the image of the algebra morphism

$$\Bbbk[x] \ni x \longmapsto a \in A.$$

The polynomial ring can be made into a Hopf algebra by declaring x primitive (i.e. [27, Definition 1.3.4(b)] $\Delta(x) = x \otimes 1 + 1 \otimes x$), and we are done.

Remark 2.5 The existence of the cofree Hopf algebra $H_{\rm CF}$ (and its bialgebra analogue $B_{\rm CF}(A)$ for an algebra A: nothing but $C_{\rm CF}(A)$, equipped with a natural bialgebra structure [38, §VI, pp.134-135]) is also an instance of *monadic adjunction lifting*: apply [7, Theorem 4.5.6] to the forgetful diagram(s)



with monadic [32, Proposition 47, points 1. and 3.] and [33, Theorem 10 2.] south-eastward functors.

It is no accident that in Example 2.2 and Example 2.1 the field extensions had to be infinite, for [12, Theorem 4.1(2)] does, at least, hold for finite extensions.

Theorem 2.6 The following conditions on a field extension $\mathbb{k} \leq \mathbb{k}'$ are equivalent.

- (a) The extension is finite.
- (b) The corresponding

(2-1)
$${}_{\Bbbk}COALG \xrightarrow{(-)_{\Bbbk'} = -\otimes_{\Bbbk} \Bbbk'} {}_{\Bbbk'}COALG$$

is a right adjoint.

- (c) (2-1) is continuous or, equivalently, preserves products.
- (d) (2-1) preserves powers (i.e. products of copies of a single coalgebra).
- (e) The analogue of (b) for categories of bialgebras and/or Hopf algebras.
- (f) The analogue of (c) for categories of bialgebras and/or Hopf algebras.
- (g) The analogue of (d) for categories of bialgebras and/or Hopf algebras.

(h) $(-)_{k'}$ preserves the \aleph_0 -power of the cofree Hopf algebra on the ground field, regarded as an algebra.

Proof We tackle the various implications in turn.

(b) \Leftrightarrow (c): Regardless of finiteness, (2-1) is in any case a *left* adjoint or, equivalently (by *Freyd's special adjoint functor theorem* [2, §0.7], for COALG is a locally presentable [2, Definition 1.17] category by [31, §2.7, Proposition, item 1.] or [33, Lemma 1 2.]), cocontinuous. Indeed, scalar extension is left adjoint to scalar restriction at the level of vector spaces, and colimits of coalgebras are those of underlying vector spaces (COALG $\xrightarrow{\text{forget}}$ VECT being left adjoint [31, §2.7, Proposition, item 2.]).

As a cocontinuous functor between locally presentable categories, (2-1) will be a right adjoint as soon as it is also continuous by [2, Theorem 1.66]. And continuity is in turn equivalent to product preservation (by [1, Proposition 13.4]), for equalizers are preserved in any case, by possibly infinite field extensions, by [12, Lemma 4.2 and discussion pertaining to equalizers in the proof of Theorem 4.1].

(e) \Leftrightarrow (f): The argument just given works here too, substituting

 ${\sf BIALG} \quad {\rm or} \quad {\sf HALG} \quad \xrightarrow{{\rm forget}} \quad {\sf ALG}$

for the forgetful functors COALG \rightarrow VECT. $(-)_{k'}$ is left adjoint at the level of algebras, all categories in sight are locally presentable [33, Lemma 2 and Theorem 6], etc.

The two unlabeled implications in

$$\begin{array}{c} \text{obvious} \\ (c) & & \\$$

hold because products (and more generally, limits) of bialgebras or Hopf algebras are computed at the coalgebra level, by the continuity (in fact, *monadicity* [7, Definition 4.4.1], per [31, §4.1, Proposition] and [33, Theorem 10 2.]) of the corresponding forgetful functors.

(a) \Rightarrow (c): We break up the construction of a coalgebra product $C := \prod_i C_i$ into a handful of steps, each of which is in turn preserved by extending scalars along *finite* field extensions. The recipe:

• First form the vector-space product $W := \prod_{i=1}^{V_{\text{ECT}}} C_i$. The construction is indeed preserved by $(-)_{\mathbb{k}'}$ by the assumed finiteness.

• Then form the cofree coalgebra (Example 2.1) $C_{\rm CF}(W)$, naturally equipped with maps

cofree structure map

$$C_{\rm CF}(W) \xrightarrow{\pi_i} C_i$$

(linear only; not, in general, coalgebra morphisms). We relegate the invariance of $W \mapsto C_{CF}(W)$ under $(-)_{k'}$ to Proposition 2.7 below.

• Finally, $C \leq C_{\rm CF}(W)$ will be the subspace defined as the common equalizer of all parallel pairs

(2-2)
$$C_{\rm CF}(W) \xrightarrow{\Delta^{(n)}} C_{\rm CF}(W) \xrightarrow{\otimes(n+1)} C_i \xrightarrow{\pi_i^{\otimes(n+1)}} C_i^{\otimes(n+1)}$$

where $\Delta^{(\bullet)}$ denote the iterates of the comultiplication. Tensoring with *any* overfield preserves both equalizers and arbitrary subspace intersections, finishing the proof of the present implication.

(h) \Rightarrow (a): The idea, in its essence, is that driving Example 2.1. Denote cofree Hopf algebras by $H_{\rm CF}(\bullet)$, as in Lemma 2.4. Given the surjectivity, by the latter, of the map

$$H_{\rm CF}(\Bbbk^S) \longrightarrow \Bbbk^S$$
, arbitrary set S

(for arbitrary k, hence also for k'), the argument of Example 2.1 carries over: if S is infinite then

$$\left(\Bbbk^{S}\right)_{\Bbbk'} \hookrightarrow \left(\Bbbk'\right)^{S}$$

is not onto.

This suffices to connect the implication graph.

The following simple remark is perhaps worth setting out on its own; it is auxiliary to the proof of Theorem 2.6 above (implication (a) \Rightarrow (c)).

Proposition 2.7 The scalar extension functor $(-)_{k'}$ along a finite field extension $k \leq k'$ intertwines all of the following right adjoints:

- (1) the cofree coalgebra $C_{CF}(V) \to V$ on a vector space;
- (2) the cofree bialgebra $B_{CF}(A) \rightarrow A$ on an algebra;
- (3) the cofree Hopf algebra $H_{CF}(B) \rightarrow B$ on a bialgebra;
- (4) the cofree Hopf algebra $H_{CF}(A) \to A$ on an algebra.
- of [14, §1.6] over \Bbbk and \Bbbk' respectively.

Proof (1): First, the finiteness of $\mathbb{k} \leq \mathbb{k}'$ ensures that $(-)_{\mathbb{k}'}$ intertwines the formation of vector-space duals:

(2-3)
$$(V^*)_{\mathbb{k}'} \xrightarrow{\text{canonically}} (V_{\mathbb{k}'})^*, \quad \forall \ V \in {}_{\mathbb{k}} \text{VECT}.$$

Secondly, it also preserves the *finite dual* construction

$${}_{\Bbbk} ALG \ni A \longmapsto A^{\circ} := \{ f \in A^* \mid \ker f \text{ contains a cofinite-dimensional ideal} \}$$
$$= \bigcup_{A \to \overline{A}} \overline{A}^* \le A^* \quad (\text{ranging over finite-dimensional quotient algebras } \overline{A})$$

of [14, §1.5], in the sense that (2-3) holds with $A \in ALG$ and $(-)^{\circ}$ in place of V and $(-)^{*}$ respectively. It follows in particular that the scalar extension functor intertwines the cofree coalgebra construction

VECT
$$\ni V \longmapsto C_{CF}(V^{**}) \xrightarrow{[14, \text{ Lemma 1.6.4}]} T(V^*)^{\circ}, \quad T(-) := \text{tensor algebra}$$

To conclude, recall the following description of $C_{CF}(V)$; it can be extracted from the proof of [14, Theorem 1.6.6] (via [14, Lemma 1.6.5]):

(2-4)
$$C_{\rm CF}(V) = \left\{ x \in C_{\rm CF}(V^{**}) \mid \left(\pi_{V^{**}}^{\otimes (n+1)} \circ \Delta^{(n)} \right)(x) \in V^{\otimes (n+1)} \subset (V^{**})^{\otimes (n+1)} \right\},$$

where $C_{\rm CF}(\bullet) \xrightarrow{\pi_{\bullet}} \bullet$ denotes the structure map of the cofree coalgebra and, as in (2-2), $\Delta^{(n)}$ are the iterates of Δ . Everything on the right-hand side of (2-4) is preserved by $(-)_{\Bbbk'}$ (even for *arbitrary* field extensions; finiteness is not needed here), hence the conclusion.

(2): As observed in [38, §VI, discussion following Theorem 6.4.8], $B_{\rm CF}(A)$ is nothing but the cofree coalgebra $C_{\rm CF}(A)$: the algebra structure on A lifts automatically along the canonical map $C_{\rm CF}(A) \to A$.

(3): This follows from the description of the cofree Hopf algebra $H_{CF}(B) \to B$ on a bialgebra B given in [11, Theorem 3.1(a)]:

• Form the product

$$P := \prod_{n \in \mathbb{Z}_{\geq 0}}^{\text{BiAlg}} B_n \cong \prod_{n \in \mathbb{Z}_{\geq 0}}^{\text{Coalg}} B_n, \quad B_n := \begin{cases} B & \text{if } n \text{ is even} \\ B^{op, cop} & \text{otherwise,} \end{cases}$$

where 'op' superscripts indicate opposite algebra structures and dually (per standard notation [34, Definition 2.1.4]), 'cop' means opposite coalgebra structures. That product construction is compatible with $(-)_{k'}$ by Theorem 2.6.

• Consider the bialgebra morphism $P^{op,cop} \xrightarrow{S} P$ that "shifts" factor indices, defined factor-wise by $B_n \xrightarrow{id} B_{n-1}$.

• Finally, $H_{\rm \scriptscriptstyle CF}(B)\subseteq P$ is then defined as the largest subcoalgebra contained in

$$\{x \in P \mid x_1 S(x_2) = \eta \varepsilon(x) = S(x_1) x_2\}, \qquad \begin{array}{l} \varepsilon, \eta := (\text{co}) \text{unit of the bialgebra } P, \\ x \xrightarrow[\text{comultiplication } \Delta]{\text{Sweedler notation } [34, p.23]} x_1 \otimes x_2 \end{array}$$

(automatically a sub-bialgebra, Hopf with antipode S).

This last step thus first forms an equalizer (in VECT) of the three self-maps



and then cuts out the largest subcoalgebra of P contained therein. Both of those procedures are preserved by *arbitrary* (not just finite) field extensions: the former obviously, and the latter by [12, Lemma 4.2].

(4): This right adjoint is the composition of the preceding two:

$$ALG \xrightarrow{B_{CF}} BIALG \xrightarrow{H_{CF}} HALG$$

(4) thus follows from (2) + (3).

The construction of $C_{\rm CF}(V)$ just recalled in the proof of Proposition 2.7, via the embedding $V \leq V^{**}$, can in a sense be turned around: $C_{\rm CF}$ can also be built up (rather than down) from

Lemma 2.8 The cofree coalgebra construction

$$VECT \ni V \longmapsto C_{CF}(V) \in COALG$$

. .

preserves filtered colimits. In particular, we have

(2-5)
$$\lim_{fin-dim} V' \leq V \qquad C_{CF}(V') \cong \lim_{fin-dim} T(V'^*)^\circ \xrightarrow{canonical map} C_{CF}(V).$$

Proof This is an application of a familiar general principle applicable to adjunctions between locally presentable categories.

In both COALG and VECT every object is a *filtered colimit* [2, Definition 1.4] of finite-dimensional subobjects: obvious for vector spaces, and standard [27, Theorem 5.1.1] for coalgebras. These, in turn, are precisely the *finitely presentable* [2, Definition 1.1] objects • in each respective category: their corresponding representable (covariant) functors $\mathcal{C}(\bullet, -)$ preserve filtered colimits.

The forgetful functor COALG $\xrightarrow{\text{FGT}}$ VECT is thus a left adjoint between locally finitely presentable categories which respects object finite presentability; per [2, Theorem 1.66, part (2) of the proof], its right adjoint C_{CF} respects filtered colimits.

The finiteness requirement in Proposition 2.7 is also, it turns out, not accidental, but it is suboptimal: that statement cannot quite be reversed. In fact, preservation by $(-)_{\mathbb{k}'}$ of the functor C_{CF} singles out precisely the *algebraic* field extensions.

Theorem 2.9 The following conditions on a field extension $\mathbb{k} \leq \mathbb{k}'$ are equivalent.

- (a) The extension is algebraic.
- (b) The corresponding functor $(-)_{k'}$ intertwines either (equivalently, both) of the adjunctions

$$\begin{array}{ccc} COALG & BIALG \\ & & & \\ & &$$

(c) The functor $(-)_{\mathbb{k}'}$ preserves the cofree coalgebra on the 1-dimensional vector space (equivalently, cofree bialgebra on the 1-dimensional algebra).

(d) The functor $(-)_{k'}$ intertwines the adjunction

(e) $(-)_{k'}$ preserves the cofree Hopf algebra on the 1-dimensional algebra.

Proof (a) \Rightarrow (b): We prove the coalgebra version, for the corresponding statement for bialgebras follows. Indeed, the bialgebra adjunction carries the same underlying coalgebra structure: the cofree coalgebra on an algebra, regarded as a vector space, automatically carries a natural bialgebra structure; this is noted explicitly in [38, §VI, post Theorem 6.4.8].

Lemma 2.8 reduces the problem to finite-dimensional V, whereupon

$$C_{\rm CF}(V) \cong C_{\rm CF}(V^{**}) \xrightarrow{[38, \text{ Theorem 6.4.1}]} T(V^*)^{\circ} \cong \Bbbk \langle \mathcal{S} \rangle^{\circ}$$

 $\Bbbk \langle \mathcal{S} \rangle$ denoting the free \Bbbk -algebra on a basis S for V^* .

The functor $(-)_{k'}$ (for any $k \leq k'$ at all, algebraic or not) certainly intertwines the formation of duals for finite-dimensional V as well as T(-), so the crux of the matter is the preservation of finite duals, specifically for free finitely-generated algebras $k \langle S \rangle$:

$$\mathbb{k} \leq \mathbb{k}' \text{ algebraic} \Longrightarrow (\mathbb{k} \langle \mathcal{S} \rangle^{\circ})_{\mathbb{k}'} \xrightarrow{\text{canonical map}}{\text{ is onto }} \mathbb{k}' \langle \mathcal{S} \rangle^{\circ}, \quad \forall S \text{ finite.}$$

This, though, follows from Lemma 2.11 below (implication (a) \Rightarrow (b)).

(b) \Rightarrow (c) and (d) \Rightarrow (e) are formal.

(a) \Rightarrow (d): Recall [27, Definition 4.2.3] that finitely-generated k-algebras are sometimes also termed (k-) affine. By analogy, we call a Hopf algebra (k-)S-affine or (k-)-antipode-affine if it is generated as an algebra by the iterates $S^{\bullet}F$ under the antipode of a finite subset (or subspace) F. We observe that an arbitrary Hopf algebra H can be recovered as

(2-6)
$$H = \varinjlim_{\substack{\text{S-affine } H' \le H}}^{\text{ALG}} H'.$$

What is more, in the present context of a given algebraic extension $\mathbb{k} \leq \mathbb{k}'$, an S-affine Hopf \mathbb{k}' algebra H will be of the form $H \cong K_{\mathbb{k}'}$ for some finite intermediate $\mathbb{k}(H)/\mathbb{k}$ and S-affine Hopf $\mathbb{k}(H)$ -algebra $K \leq H$: fix a finite-dimensional \mathbb{k}' -coalgebra C generating H as an algebra, take $\mathbb{k}(H)$ large enough to contain the structure constants c_{ij}^k in

$$e_i \cdot e_j = \sum_k c_{ij}^k e_k$$
, (e_i) a k'-basis for the algebra C^* ,

and take for K the \mathbb{k}'' -algebra generated by the antipode-iterates of the dual basis (e_i^*) .

Consider a Hopf \Bbbk' -algebra H and an \Bbbk -algebra A. Relegating one crucial step to Lemma 2.10, the proof functions as follows:

(c) \Rightarrow (a): Since $(-)_{\Bbbk'}$ in any case respects duality for finite-dimensional vector spaces (such as the $V \cong \Bbbk$ under consideration), the issue is whether it respects the finite duality in $C_{CF}(V) \cong \Bbbk[x]^{\circ}$ (for x spanning V^*). For that, we can fall back on the implication (c) \Rightarrow (a) of Lemma 2.11.

(e) \Rightarrow (a): The selfsame implication (c) \Rightarrow (a) of Lemma 2.11 also handles this claim, given that $H_{\rm CF}(\Bbbk) \cong \Bbbk[x^{\pm 1}]^{\circ}$ (Remark 2.17(2)).

Lemma 2.10 Let H be an S-affine Hopf \Bbbk -algebra, A a \Bbbk -algebra, and $\Bbbk \leq \Bbbk'$ a field extension. An algebra morphism $H \to A_{\Bbbk'}$ factors through $A_{\Bbbk''}$ for some intermediate $\Bbbk \leq \Bbbk'' \leq \Bbbk'$ finitely-generated over \Bbbk .

Proof The assumption is that H is generated as an algebra by the iterates $S^{\bullet C}$ for some finitedimensional coalgebra $C \leq H$, S being the antipode of H. Coalgebra morphisms $C \to H$ factor as

$$C \xrightarrow{H_{\rm F}(C)} H^{\rm HALG morphism} H$$

through the free Hopf algebra [39, §1] $H_{\rm F}(C)$ on C. There is no loss in assuming $H = H_{\rm F}(C)$, which we henceforth do.

k-algebra morphisms $H_{\mathbb{F}}(C) \to A_{\mathbb{K}'}$ are in bijection [39, Corollary 10] with the sequences

$$(x_n)_{n \in \mathbb{Z}_{\geq 0}} \subset C^* \otimes A_{\Bbbk'}, \quad x_{n+1} = \begin{cases} \text{the inverse } x_n^{-1} \text{ in } C^* \otimes A_{\Bbbk'} & \text{if } n \text{ is even} \\ x_n^{-1} \text{ in } C^* \otimes (A_{\Bbbk'})^{op} & \text{otherwise} \end{cases}$$

The first term x_0 of that sequence belongs to some $A_{\Bbbk''}$ for finitely-generated intermediate \Bbbk'' as in the statement. That the inverse $x_1 = x_0^{-1}$ then also belongs to $A_{\Bbbk''}$ (as do the other x_n afterwards, inductively) follows: multiplication by x_0 , regarded as a \Bbbk'' -linear endomorphism, preserves the direct-sum decomposition

$$A_{\Bbbk'} \cong A_{\Bbbk''} \oplus \left(A \otimes_{\Bbbk} \Bbbk''^{\perp} \right) \text{ in }_{\Bbbk''} \text{VECT}, \quad \Bbbk' = \Bbbk'' \oplus \Bbbk''^{\perp}$$

for some complementary subspace \Bbbk''^{\perp} of $\Bbbk'' \leq \Bbbk'$ (i.e. leaves invariant both summands); its inverse must do so as well.

Lemma 2.11 The following conditions on a field extension $\mathbb{k} \leq \mathbb{k}'$ are equivalent.

- (a) The extension is algebraic.
- (b) $(-)_{k'}$ preserves the formation of finite duals for affine algebras.
- (c) $(-)_{\Bbbk'}$ preserves either one of the finite duals $\Bbbk[x]^{\circ}$ or $\Bbbk[x^{\pm 1}]^{\circ}$.

Proof (a) \Rightarrow (b): Let A be an affine k-algebra. Recast, the claim is that any cofinite ideal of $A_{k'}$ contains one such ideal for $A \leq A_{k'}$ (provided $k \leq k'$ is algebraic). Or again: for a finite-dimensional k'-algebra B, a finite subset $F \subseteq B$ generates a finite-dimensional k-algebra.

In this last formulation, simply note that finite dimensionality means that for some ℓ all words of length $> \ell$ in F are k'-linear combinations of words of length $\leq \ell$. The coefficients appearing in such expansions for length- $(\ell + 1)$ words are all contained in some *finite* intermediate extension $k \leq \tilde{k} \leq k'$, so the \tilde{k} -algebra $\tilde{k}(F) \leq B$ generated by F is certainly finite-dimensional over \tilde{k} . But then it is also finite-dimensional over k, and along with it its k-subspace $k(F) \leq \tilde{k}(F)$.

(b) \Rightarrow (c) is self-evident.

(c) \Rightarrow (a): Consider $t \in \mathbb{k}'$ transcendental over \mathbb{k} and observe that the algebra morphism

$$\Bbbk'[x] \cong \Bbbk[x]_{\Bbbk'} \ni x \longmapsto t \in \Bbbk'$$

does not annihilate any cofinite ideals in $\Bbbk[x]$. For that reason, regarded as an element of $\Bbbk'[x]^{\circ}$, that character cannot belong to $(\Bbbk[x]^{\circ})_{\Bbbk'}$. The argument works just as well for $\Bbbk[x^{\pm 1}]^{\circ}$: pick an *invertible* transcendental element $t \in \Bbbk'$.

Affineness is essential in Lemma 2.11, in the proof of the implication (a) \Rightarrow (b):

Example 2.12 No infinite extension $\Bbbk \leq \Bbbk'$ (algebraic or not) preserves the formation of finite duals for (commutative or non-commutative) polynomial rings on infinite sets S: an algebra morphism $\Bbbk'[S] \to \Bbbk'$ for which the \Bbbk -span of S is infinite over \Bbbk cannot annihilate a cofinite ideal of $\Bbbk[S]$.

A little additional work will bring us full circle back to products, in the following analogue of Theorem 2.6 characterizing *algebraic* (as opposed to finite) field extensions.

Theorem 2.13 The following conditions on a field extension $\mathbb{k} \leq \mathbb{k}'$ are equivalent.

- (a) The extension is algebraic.
- (b) (2-1) preserves finite (or binary) products of finite-dimensional coalgebras.
- (c) (2-1) is finitely continuous or, equivalently, preserves finite (or binary) products.
- (d) (2-1) preserves finite (or binary) powers.
- (e) The analogue of (c) for categories of bialgebras and/or Hopf algebras.
- (f) The analogue of (d) for categories of bialgebras and/or Hopf algebras.
- (g) $(-)_{k'}$ preserves the binary power of some Hopf algebra of finite dimension > 1.
- (h) $(-)_{k'}$ preserves the binary power of some coalgebra of finite dimension > 1.

Proof 'Finite' and 'binary' are interchangeable throughout by induction, and the conditions in (c) are equivalent by [1, Proposition 13.3] given that, as noted previously in the proof of Theorem 2.6, equalizers are preserved regardless of the field extension. Immediate implications include

$$\begin{array}{c} (b) & & \text{obvious} \\ \text{obvious} & (c) & & \\ \text{same products} & (e) & & \\ \end{array} \\ \begin{array}{c} (f) & \xrightarrow{\text{obvious}} \\ (g) & \xrightarrow{\text{obvious}} \\ (h) \end{array} \end{array}$$

A few others will complete the proof.

(b) \Rightarrow (c) follows from the fact [27, Theorem 5.1.1] that every coalgebra *C* is a filtered colimit $C \cong \varinjlim C'$ of its finite-dimensional subcoalgebras, along with the commutation [2, Proposition 1.59], given the local finite presentability of COALG recalled in the proof of Lemma 2.8 of finite limits and filtered colimits:

$$\varinjlim_{\substack{C'_i \leq C_i \\ \dim C'_i < \infty}} \prod_i^{\text{finite}} C'_i \xrightarrow{\text{canonical}} \prod_i \left(\varinjlim C'_i \right) \cong \prod_i C_i.$$

(a) \Rightarrow (b): For finite-dimensional coalgebras C the unit morphism $C \rightarrow C^{*\circ}$ attached to the contravariant right-hand adjunction ([14, Theorem 1.5.22], [38, Theorem 6.0.5])

$$\operatorname{COALG} \xleftarrow{*}_{\circ} \operatorname{ALG}$$

is an isomorphism. Contravariant right adjoints convert coproducts into products, so that

(2-7)
$$\dim C_i < \infty \Longrightarrow \prod_i^{\text{COALG}} C_i \cong \left(\coprod_i^{\text{ALG}} C_i^* \right)^{\circ}.$$

It remains to observe that arbitrary scalar extensions preserve algebra coproducts and duality for finite-dimensional vector spaces, whereas *algebraic* scalar extensions preserve finite duals of affine algebras by Lemma 2.11.

(h) \Rightarrow (a): (2-7) applies to the binary power of a finite-dimensional $C \in \text{COALG}$. The claim is thus that, given

$$(C^* =) A \in ALG, \quad 1 < \dim A < \infty,$$

preservation by $(-)_{\Bbbk}$ of the finite dual $(A \coprod A)^{\circ}$ forces $\Bbbk \leq \Bbbk'$ to be algebraic. Rephrased, the goal is to show that whenever $\Bbbk \leq \Bbbk'$ is transcendental there is a morphism

(2-8)
$$A \coprod A \xrightarrow{\varphi_{\ell,r} \text{ for 'left' and 'right'}}_{\text{ in }_{\Bbbk} \text{ALG}} B_{\Bbbk'} \text{ for } B \in {}_{\Bbbk} \text{ALG}, \quad \dim B < \infty$$

which does *not* factor through a finite-dimensional k-subalgebra of $B_{k'}$. Take for B the matrix algebra $\operatorname{End}(A) \cong M_{\dim A}(k)$. The argument bifurcates, referring to a transcendental $t \in k'$ fixed throughout.

(I) A contains, possibly after an appropriate (harmless) finite extension of k, a nonzero nilpotent element. It then also contains a square-zero $0 \neq x \in A$. Fix a square-zero $0 \neq x \in A$. φ_{ℓ} will simply be the left regular self-action

$$A \ni a \longmapsto (a \cdot) \in B = \operatorname{End}(A),$$

whereas φ_r will be the same embedding followed by a conjugation in $B_{\Bbbk(t)} \cong M_{\dim A}(\Bbbk(t))$. Choosing that conjugation so that $\varphi_r(x) = x't$ with

$$x' \in B = M_{\dim A}(\mathbb{k}), \quad xx' \text{ is not nilpotent},$$

the images through the corresponding (2-8) of the powers of

(left-hand copy of x) \cdot (right-hand copy of x) $\in A \coprod A$

have infinite-dimensional k-span span $\{(xx')^n t^n\}_n$.

(II) No finite extensions of \Bbbk produce non-zero nilpotent elements in A. We can then assume (again, perhaps after a scalar extension) that A is a finite product of (at least two) copies

of k. Focusing on $\varphi_{\ell,r}$ that factor through some quotient $A \twoheadrightarrow k^2$, we may as well take $A = k^2$; algebra morphisms (2-8), then, simply pick out idempotents $\varphi_{\ell,r}(x), x := (1,0) \in k^2$. Define

$$x \xrightarrow{\varphi_{\ell}} \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}$$

$$x \xrightarrow{\varphi_{r}} \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \cdot \begin{pmatrix} 1 & -t \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1+t & -t+t^{2} \\ 1 & -t \end{pmatrix}$$

We thus have

(right-hand x) – (left-hand x)
$$\xrightarrow{(2-8)} \begin{pmatrix} 1+t & -t+t^2 \\ 1 & -t \end{pmatrix} - \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} t & -t+t^2 \\ 0 & -t \end{pmatrix},$$

an element whose powers are plainly have infinite k-span.

We record the following description of the cofree Hopf algebra on a finite-dimensional algebra.

Proposition 2.14 In the diagram

$$H_{CF}(A) \xrightarrow{B_{CF}(A) \cong T(A^*)^{\circ}} \xrightarrow{A,} A,$$

relating the cofree bialgebra and Hopf algebra on a finite-dimensional \Bbbk -algebra A the left-hand arrow is the finite dual

$$H_F(T(A^*))^\circ \longrightarrow T(A^*)^\circ$$

of the canonical map $T(A^*) \to H_F(T(A^*))$ of the bialgebra $T(A^*)$ into its Hopf envelope.

Proof The bialgebra $T(A^*)$ in the statement carries the tensor algebra structure and the coalgebra structure induced by that of A^* . That the cofree bialgebra is indeed $T(A^*)^\circ$ follows from [38, pp.134-135], given that by (the proof of) [38, Theorem 6.4.1] this is also the cofree coalgebra on the finite-dimensional vector space $A \cong A^{**}$.

Note next that $(-)^{\circ}$ induces a contravariant self-adjoint on the right on both categories BIALG and HALG: the natural bijection

$$BIALG(H, K^{\circ}) \cong BIALG(K, H^{\circ}), \quad H, K \in BIALG \text{ or } HALG$$

follows essentially from [34, Proposition 7.7.5], which identifies both spaces with that of *bialgebra (or Hopf) pairings* ([34, Definition 7.7.6], [10, §4.1 D, pp.114-115]) $H \times K \to \Bbbk$. Said self-adjoint $(-)^{\circ}$ also intertwines the forgetful functor from Hopf algebras to bialgebras, hence the commutativity up to natural isomorphism of (the northwestward branch of)



More elaborately (but perhaps also more transparently), given [25, §IV.8, Theorem 1] that the adjoint of a composition of functors is the composition of the corresponding adjoints, we have adjunctions



so that the top and bottom must be naturally isomorphic; $(-)^{\circ}$, in other words, intertwines the free and cofree Hopf algebra constructions. This, though, is precisely the sought-after claim in our particular setup: that $(-)^{\circ}$ turns the free Hopf algebra structure map $B \to H_F(B)$ into the cofree counterpart $H_F(B)^{\circ} \to B^{\circ}$ thereof with $B = T(A^*)^{\circ}$.

Remarks 2.15 (1) The principle underlying Proposition 2.14 also fits the isomorphism $C_{\rm CF}(V^{**}) \cong T(V^*)^\circ$ of [14, Lemma 1.6.4] into the broader observation that $C_{\rm CF}(W^*) \cong T(W)^\circ$: the latter is (the image of W through) the composition

$$\operatorname{VECT} \xrightarrow{T} \operatorname{ALG} \xrightarrow{\circ} \operatorname{COALG}^{op}$$

of left adjoints; the former instead chains together

VECT
$$\xrightarrow{*}$$
 VECT^{op} $\xrightarrow{C_{CF}}$ COALG^{op},

again left adjoints. The common right adjoint to the two compositions is

(2) Dual to the phenomenon recalled in Remark 2.5, of the cofree coalgebra $C_{CF}(A)$ on an algebra A inheriting a natural (bi)algebra structure, the tensor algebra T(C) on a coalgebra is nothing but the free bialgebra on C ([34, Theorem 5.3.1], [38, Proposition 3.2.4 and Exercise (1) following it]): the result of applying to C the left adjoint of BIALG \rightarrow COALG.

In light of this, the free Hopf algebra $H_{\rm F}(T(A^*))$ of Proposition 2.14 is also the free Hopf algebra on the coalgebra A^* (introduced in [39]; see also [34, §7.5]): the image of C through chaining together the left adjoints of the two composable functors in

$$\begin{array}{c} \stackrel{\mathrm{FGT}}{\longrightarrow} & \mathrm{BIALG} \xrightarrow{\mathrm{FGT}} \\ \mathrm{HALG} \xrightarrow{\mathrm{FGT}} & \mathrm{COALG} \end{array}$$

producing the left adjoint of the composite.

An illustration of Proposition 2.14:

۲

Example 2.16 Set $A := \Bbbk^S$ for a finite set S. The coalgebra A^* is then the span of |S| grouplikes [27, Definition 1.3.4(a)] and $T(A^*) \cong \Bbbk \langle S \rangle$, the free algebra on said grouplike elements (with the bialgebra structure resulting therefrom). The free Hopf algebra on $\Bbbk \langle S \rangle$ is easily seen to be what we will denote by $\Bbbk \langle S^{\pm 1} \rangle$: the non-commutative Laurent polynomial algebra, obtained by formally adjoining inverses to every $s \in S$.

The cofree Hopf algebra map $\Bbbk \langle S^{\pm 1} \rangle^{\circ} \to \Bbbk \langle S \rangle^{\circ}$ of Proposition 2.14 is an embedding: if a functional $\Bbbk \langle S^{\pm 1} \rangle \to \Bbbk$ annihilates all of $\Bbbk \langle S \rangle$ and also a cofinite ideal, it must annihilate arbitrary monomials

(2-9)
$$s_1^{e_1} \cdots s_n^{e_n}, \quad s_k \in \mathcal{S}, \quad e_k \in \mathbb{Z}$$

(and hence vanish identically) by induction on the absolute value of the smallest exponent $e_k < 0$. Suppose, say, that $e_k < 0$, f's vanishing on a cofinite ideal implies that it annihilates elements of the form

•
$$p(s) \cdot \bullet', \quad \bullet, \bullet' \in \mathbb{kS}^{\pm 1}$$
 and some polynomial $p \in \mathbb{k}[x].$

(2-9) can thus be expressed as a linear combination of analogous terms with strictly larger exponents in place of e_k , completing the induction step.

Remarks 2.17 (1) The isomorphism $B_{CF}(\mathbb{k}^{S}) \cong \mathbb{k} \langle S \rangle^{\circ}$ is valid generally, regardless of whether S is finite (as it was in Example 2.16): $\mathbb{k}^{S} \cong (\mathbb{k}^{\oplus S})^{*}$, and Remark 2.15(1) applies.

(2) The case $|\mathcal{S}| = 1$ of Example 2.16 features prominently in [23]. Switching to the more common notation $\mathbb{k}[x]$ for the single $x \in \mathcal{S}$, [23, §2] identifies $\mathbb{k}[x^{\pm 1}]^{\circ}$ as a Hopf subalgebra of the bialgebra $\mathbb{k}[x]^{\circ}$ of *linearly recursive sequences* [15, §1.1.1]: a functional $\mathbb{k}[x] \xrightarrow{f} \mathbb{k}$ belongs to the finite dual precisely when the corresponding sequence

$$(f_n)_{n \in \mathbb{Z}_{>0}}, \quad f_n := f(x^n)$$

satisfies a linear recurrence. The Hopf subalgebra $k[x^{\pm 1}]^{\circ}$ consists of those sequences that extend across all of \mathbb{Z} , satisfying the linear recurrence throughout. Per Proposition 2.14, that Hopf subalgebra is in fact the *universal* bialgebra morphism received by $k[x]^{\circ}$ from a Hopf algebra, so is in particular also the *largest* Hopf subalgebra of the bialgebra of linearly recursive sequences.

A small subtlety present in the proof of Theorem 2.9 (specifically, the implication (a) \Rightarrow (a)) suggests a side-note:

• On the one hand, Lemma 2.11 handles, for algebraic $\mathbb{k} \leq \mathbb{k}'$, preservation by $(-)_{\mathbb{k}'}$ of finite duals for *affine* algebras.

• On the other hand, affineness cannot simply be dropped in general (Example 2.12).

• The issue of applying Lemma 2.11 in the course of the proof of the aforementioned implication suggests itself, and with it so does the question of when the free Hopf algebra $H_{\rm F}(C)$ on a finite-dimensional coalgebra is affine. Such C, it turns out, can be classified.

Proposition 2.18 The free Hopf algebra $H_F(C)$ on a finite-dimensional k-coalgebra C is k-affine if and only if the scalar extension $C_{\overline{k}}$ to the algebraic closure is pointed.

Proof As a consequence [30, §3.5, Corollary b] of the Wedderburn-Artin theorem [30, §3.5, Theorem], the coradical $(C_{\overline{k}})_0$, being finite-dimensional and cosemisimple over an algebraically closed field, must be a finite sum

$$(C_{\overline{\Bbbk}})_0 \cong \bigoplus_{i=1}^{\ell} M_{n_i}(\overline{\Bbbk})^*$$

of matrix coalgebras; the analogous isomorphism then holds for sufficiently large (in the sense of inclusion) finite $\Bbbk'' \ge \Bbbk$. Affineness being preserved and reflected by finite field extensions we can, upon substituting for \Bbbk' a composite field [13, §11.7] of $\Bbbk', \Bbbk'' \ge \Bbbk$ (thus finite over \Bbbk'), assume that $C_0 \cong \bigoplus M_{n_i}^*$ to begin with. Under that assumption, the claim is that $H_{\rm F}(C)$ is affine precisely when all n_i are 1.

Choosing a (plain, vector-space) decomposition $C \cong C_0 \oplus V$, [39, Theorem 32] gives the algebra structure

$$H_{\mathrm{F}}(C) \cong T(V) \sqcup \coprod_{i} H_{\mathrm{F}}(M_{n_{i}}^{*}) \quad \text{(coproduct in ALG)}.$$

Affineness for $H_{\rm F}(C)$ as a whole thus reduces to affineness for each individual $H_{\rm F}(M_{n_i}^*)$, and hence the desired equivalence again reduces:

$$H := H_{\rm F}(M_n^*)$$
 is affine $\xleftarrow{\text{claim}} n = 1.$

One implication (\Leftarrow) is plain enough: $H_{\rm F}(\Bbbk) \cong \Bbbk[x^{\pm 1}]^{\circ}$ [39, Lemma 34], certainly affine. For the converse, the basis for H described in [29, Theorem 5] makes it clear that while generated as an algebra by the iterates

$$(S^{\bullet}M_n^*)_{\bullet\in\mathbb{Z}_{>0}}, \quad S := \text{antipode of } H,$$

H is not generated as an algebra by only finitely many of those iterates.

Remark 2.19 The coalgebras featuring in Proposition 2.18, with pointed extensions to the algebraic closure, are also precisely those whose free Hopf algebra happens to have bijective antipode [39, Theorem 18].

References

- Jiří Adámek, Horst Herrlich, and George E. Strecker. Abstract and concrete categories: the joy of cats. *Repr. Theory Appl. Categ.*, 2006(17):1–507, 2006. 3, 12, 19
- [2] Jiří Adámek and Jiří Rosický. Locally presentable and accessible categories, volume 189 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1994. 2, 4, 12, 16, 19
- [3] A. L. Agore. Monomorphisms of coalgebras. Colloq. Math., 120(1):149–155, 2010. 2
- [4] A. L. Agore. Categorical constructions for Hopf algebras. Commun. Algebra, 39(4):1476–1481, 2011. 2, 11

- [5] Michael F. Atiyah and I. G. Macdonald. Introduction to commutative algebra. Reading, Mass.-Menlo Park, Calif.-London-Don Mills, Ont.: Addison-Wesley Publishing Company (1969)., 1969. 11
- [6] George M. Bergman. The diamond lemma for ring theory. Adv. Math., 29:178–218, 1977. 9
- [7] Francis Borceux. Handbook of categorical algebra. 2: Categories and structures, volume 51 of Encycl. Math. Appl. Cambridge: Univ. Press, 1994. 4, 11, 12, 13
- [8] Francis Borceux. Handbook of categorical algebra. Volume 1: Basic category theory, volume 50 of Encycl. Math. Appl. Cambridge: Cambridge Univ. Press, 1994. 4, 5, 9
- [9] Armand Borel. *Linear algebraic groups*, volume 126 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991. 6, 7
- [10] Vyjayanthi Chari and Andrew Pressley. A guide to quantum groups. Cambridge: Cambridge University Press, 1995. 1, 21
- [11] Alexandru Chirvăsitu. On epimorphisms and monomorphisms of Hopf algebras. J. Algebra, 323(5):1593–1606, 2010. 2, 14
- [12] Alexandru Chirvasitu. Epimorphic quantum subgroups and coalgebra codominions. Algebr. Represent. Theory, 27(1):219–244, 2024. 2, 10, 12, 15
- [13] P. M. Cohn. Basic algebra. Groups, rings and fields. London: Springer, 2003. 24
- [14] Sorin Dăscălescu, Constantin Năstăsescu, and Şerban Raianu. Hopf algebras, volume 235 of Monographs and Textbooks in Pure and Applied Mathematics. Marcel Dekker, Inc., New York, 2001. An introduction. 2, 3, 8, 11, 14, 20, 22
- [15] Graham Everest, Alf van der Poorten, Igor Shparlinski, and Thomas Ward. Recurrence sequences, volume 104 of Math. Surv. Monogr. Providence, RI: American Mathematical Society (AMS), 2003. 23
- [16] Thomas M. Fiore. Pseudo limits, biadjoints, and pseudo algebras: categorical foundations of conformal field theory, volume 860 of Mem. Am. Math. Soc. Providence, RI: American Mathematical Society (AMS), 2006. 3, 4
- [17] Paul Großkopf and Joost Vercruysse. Free and co-free constructions for Hopf categories. J. Pure Appl. Algebra, 228(10):32, 2024. Id/No 107704. 2
- [18] Nathan Jacobson. Basic algebra I. 2nd ed. New York: W. H. Freeman and Company. XVIII, 499 p. £ 19.95 (1985)., 1985. 11
- [19] Nathan Jacobson. Basic algebra II. New York, NY: W. H. Freeman and Company, 2nd ed. edition, 1989. 11
- [20] Niles Johnson and Donald Yau. 2-dimensional categories. Oxford: Oxford University Press, 2021. 3
- [21] Stephen Lack. A 2-categories companion. In *Towards higher categories*, pages 105–191. Berlin: Springer, 2010. 3

- [22] T. Y. Lam. A first course in noncommutative rings., volume 131 of Grad. Texts Math. New York, NY: Springer, 2nd ed. edition, 2001. 11
- [23] Richard G. Larson and Earl J. Taft. The algebraic structure of linearly recursive sequences under Hadamard product. Isr. J. Math., 72(1-2):118–132, 1990. 23
- [24] Martin Lorenz. personal communication, 2021-01-08. 1
- [25] Saunders Mac Lane. Categories for the working mathematician, volume 5 of Graduate Texts in Mathematics. Springer-Verlag, New York, second edition, 1998. 4, 22
- [26] Paddy McCrudden. Tannaka duality for Maschkean categories. J. Pure Appl. Algebra, 168(2-3):265–307, 2002. 4
- [27] S. Montgomery. Hopf algebras and their actions on rings, volume 82 of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1993. 3, 4, 9, 11, 16, 17, 19, 23
- [28] D. Mumford, J. Fogarty, and F. Kirwan. Geometric invariant theory, volume 34 of Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)]. Springer-Verlag, Berlin, third edition, 1994. 7
- [29] Warren D. Nichols. Quotients of Hopf algebras. Comm. Algebra, 6(17):1789–1800, 1978. 24
- [30] Richard S. Pierce. Associative algebras, volume 88 of Grad. Texts Math. Springer, Cham, 1982.
 24
- [31] Hans-E. Porst. On categories of monoids, comonoids, and bimonoids. Quaest. Math., 31(2):127– 139, 2008. 2, 12, 13
- [32] Hans-E. Porst. The formal theory of Hopf algebras. Part I: Hopf monoids in a monoidal category. Quaest. Math., 38(5):631–682, 2015. 2, 4, 12
- [33] Hans-E. Porst. The formal theory of Hopf algebras. Part II: The case of Hopf algebras. Quaest. Math., 38(5):683–708, 2015. 2, 4, 9, 11, 12, 13
- [34] David E. Radford. Hopf algebras, volume 49 of Series on Knots and Everything. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2012. 3, 4, 8, 15, 21, 22
- [35] Maxwell Rosenlicht. A remark on quotient spaces. Anais Acad. Brasil. Ci., 35:487–489, 1963.
 7
- [36] Joseph J. Rotman. An introduction to the theory of groups, volume 148 of Graduate Texts in Mathematics. Springer-Verlag, New York, fourth edition, 1995. 9, 10
- [37] Peter Schauenburg. Tannaka duality for arbitrary Hopf algebras, volume 66 of Algebra-Ber. München: R. Fischer, 1992. 4
- [38] Moss E. Sweedler. Hopf algebras. Mathematics Lecture Note Series. W. A. Benjamin, Inc., New York, 1969. 2, 3, 11, 12, 14, 16, 20, 21, 22
- [39] Mitsuhiro Takeuchi. Free Hopf algebras generated by coalgebras. J. Math. Soc. Japan, 23:561– 582, 1971. 2, 18, 22, 24

Department of Mathematics, University at Buffalo Buffalo, NY 14260-2900, USA *E-mail address*: achirvas@buffalo.edu