Deformation of moduli spaces of meromorphic G-connections on \mathbb{P}^1 via unfolding of irregular singularities

Kazuki Hiroe*

Department of Mathematics and Informatics, Chiba University 1-33, Yayoi-cho, Inage-ku, Chiba-shi, Chiba, 263-8522 JAPAN email: kazuki@math.s.chiba-u.ac.jp

Abstract

The unfolding of singular points of linear differential equations is a classical technique to study the properties of irregular singular points from those of regular singular points. In this paper, we propose a general framework for the unfolding of unramified irregular singularities of meromorphic G-connections on \mathbb{P}^1 . As one of main results, we give a description for the unfolding of singularities of connections as the deformation of moduli spaces of them, and show that every moduli space of irreducible meromorphic G-connections with unramified irregular singularities on \mathbb{P}^1 has a deformation to a moduli space of irreducible Fuchsian G-connections on \mathbb{P}^1 . Furthermore, we can consider the unfolding of additive Delinge-Simpson problems, namely, the unfolding of irregular singularities generates a family of additive Deligne-Simpson problems. Then as an application of our main result, we show that a Deligne-Simpson problem for G-connections with unramified irregular singularities has a solution if and only if every other unfolded Deligne-Simpson problem simultaneously has a solution. Also we give a combinatorial and diagrammatic description of the unfolding of irregular singularities via spectral types and unfolding diagrams, and then consider a conjecture proposed by Oshima which asks the existence of irreducible G-connections realizing the given spectral types and their unfolding. Our main result gives an affirmative answer of this conjecture.

Contents

1

Intr	oduction	3
1.1	Classical and recent known examples	3
1.2	Unfolding of the irregular singular point of canonical forms for meromorphic	
	G -connections on the formal punctured disk $\ldots \ldots \ldots \ldots \ldots \ldots \ldots$	4
1.3	Statement of the main theorem	6
1.4	Spectral types and unfolding diagrams	7
1.5	Additive Deligne-Simpson problem	9
1.6	A conjecture by Oshima	11

*The author is supported by JSPS KAKENHI Grant Number 20K03648.

2	Meromorphic G-connections on the formal punctured disk2.1Hukuhara-Turrittin theory for \mathfrak{g} -valued connections2.2Spectral types of canonical forms2.3 δ -invariants of canonical forms	13 13 15 17
3	Truncated orbit of an unramified canonical form3.1Affine algebraic groups over the ring of formal power series3.2Truncated orbit as coadjoint orbit3.3 δ -invariant and dimension of truncated orbit	19 19 22 23
4	Meromorphic G-connections on \mathbb{P}^1 with unramified canonical forms4.1Meromorphic G-connections on \mathbb{P}^1 and index of rigidity4.2Example I: Heun equation and its confluent equations4.3Example II	25 25 27 29
5	δ-constant deformation of canonical forms5.1Complement of hypersurfaces associated with H 5.2Stratification of \mathbb{C}^{k+1} associated to partitions5.3Unfolding of canonical forms and decomposition of the spectral types5.4δ-constant deformation of canonical form5.5rig-constant deformation of a family of canonical forms5.6Examples	 32 32 32 33 35 36 37
6	Additive Deligne-Simpson problem and a conjecture by Oshima6.1Additive Deligne-Simpson problem for unfolding families of canonical forms6.2Unfolding of spectral types6.3A conjecture by Oshima	41 41 43 45
7	Moduli spaces of meromorphic connections on \mathbb{P}^1 with unramified irreg- ular sigularities 7.1 Lie groupoids, Orbifolds, and symplectic stratified spaces 7.2 Moduli spaces of meromorphic connections on a trivial bundle over $\mathbb{P}^1(\mathbb{C})$. 7.3 Stability of G/Z -action on the space of irreducible connections 7.4 Symplectic orbifold $\mathcal{M}_{\mathbf{H}}^{i\mathbf{r}}$	47 47 49 50 51
8	Triangular decompositions of truncated orbits8.1A fiber bundle with the fiber $\mathbb{O}_{H}^{L_1 \ltimes G(\mathbb{C}[z]_k)_1}$ and the truncated orbit8.2Decompositions of $G(\mathbb{C}[z]_l)_1$	53 54 55 56 57 59 60 61
9	Deformation of $N_{l,l+1}(\mathbb{C}[z]_l)_1$ and $\mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1})$ 9.1 Deformation of $\mathbb{C}[z]_l$ and $\mathbb{C}[z^{-1}]_l$	64 67 68 70

	9.5	Lie subgroupoid $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)_1$	73
	9.6	Subbundle $\mathfrak{u}_{l,l+1}(\widehat{L}(\mathbb{B})^{rev}_{[l-1]})$	74
10) Defe	ormation of truncated orbit	75
	10.1	Deformation of truncated orbit	75
	10.2	Deformation of the moment map $\mu_{\mathbb{O}_H \downarrow G}$	77
	10.3	Fibers on each stratum of \mathbb{B}_H	78
	D 4		

11 Deformation of moduli spaces of G-connections on \mathbb{P}^1 via unfolding of irregular singularities 82

1 Introduction

The unfolding of singular points of linear differential equations is a classical technique to study the properties of irregular singular points from those of regular singular points. Hence it will be a natural expectation that moduli spaces of meromorphic connections with irregular singularities should have a deformation to moduli spaces of meromorphic Fuchsian connections, i.e., connections only with regular singularities. In this paper, we consider moduli spaces of meromorphic *G*-connections with unramified irregular singularities on \mathbb{P}^1 and show that every such moduli space has the deformation whose generic fibers are moduli spaces of Fuchsian connections on \mathbb{P}^1 . Also we apply the deformation to several existence problems of irreducible linear differential equations on \mathbb{P}^1 .

1.1 Classical and recent known examples

There are many classical and recent examples of the confluence and the unfolding of singular points of differential equations on the Riemann sphere \mathbb{P}^1 . As the first example, let us recall the differential equation for the Gauss hypergeometric function which has regular singular points at $0, 1, \infty$. Then it is classically known that by the confluence of the singular points 1 and ∞ , we obtain the differential equation for the Kummer confluent hypergeometric function which has regular singular point at 0 and irregular singular point at ∞ . Further we can consider the confluence of singular points 0 and ∞ and then obtain the differential equation for the Hermite-Weber function which has the irregular singular point at ∞ . We can illustrate this procedure by the following diagram in which the arrows represent the confluences of singular points.



As well as for the Heun differential equation it is known that several differential equations with irregular singular points are derived by the confluence of singular points: the confluent Heun differential equation, the biconfluent Heun differential equation, the doublyconfluent Heun differential equation, and the triconfluent Heun differential equation.



In the more recent papers [37] and [38], Oshima constructed versal unfolding families of unramified irregular singularities of differential equations without accessory parameters. Also in [25], Kawakami-Nakamura-Sakai considered the linear Fuchsian differential equations on \mathbb{P}^1 which relate to the 4-dimensional Painlevé-type equations and made s lists of linear differential equations arising from these Fuchsian differential equations by the confluence of singular points. Here is the one example from the lists in [25].



The collections of partitions of the integer 4 with parentheses in the diagram stand for spectral types at singular points of differential equations of rank 4. For more detail we refer the original paper [25], and also we shall introduce a generalization of spectral types for meromorphic *G*-connections later in this paper, see Sections 1.4, 2.2 and 4.3.

The purpose of this paper is to propose a general framework for the unfolding of singular points of linear differential equations on \mathbb{P}^1 including these known classical and recent examples. Furthermore, since in these above-mentioned examples we notice that the number of accessory parameters is preserved by the unfolding, it will be expected that the unfolding gives rise to deformation of the spaces of accessory parameters, i.e., moduli spaces of meromorphic connections. Actually in this paper, we shall give a description of the unfolding of singular points of differential equations as the deformation of their moduli spaces.

1.2 Unfolding of the irregular singular point of canonical forms for meromorphic *G*-connections on the formal punctured disk

Before the discussion on the deformation of moduli spaces of connections, let us introduce the deformation of canonical forms of meromorphic connections on the formal punctured disk.

Let G be a connected linear reductive algebraic group defined over \mathbb{C} . As we recall in Section 2.1, according to the formal classification theory by Hukuhara-Turrittin-Levelt-Babbit-Varadarajan, meromorphic G-connections on the 1-dimensional formal punctured disk are classified by canonical forms, i.e. \mathfrak{g} -valued meromorphic 1-forms

$$H dz = \left(\frac{H_k}{z^k} + \dots + \frac{H_1}{z} + H_{\rm res}\right) \frac{dz}{z}$$

defined on ramified coverings of the punctured disk, where H_i are mutually commuting semisimple elements and also H_{res} commutes with all H_i . Namely, for any meromorphic *G*-connection ∇ on the formal punctured disk, there exists a \mathfrak{g} -valued 1-form as above such that ∇ is reduced to d + H dz on a ramified covering of the punctured disk under the gauge transformation. In this paper, we especially focus only on unramified cases. Now we explain the unfolding of irregular singularities of unramified canonical forms. Let us consider a canonical form H as above. For $\mathbf{c} = (c_0, c_1, \ldots, c_k) \in \mathbb{C}^{k+1}$, let

$$H(\mathbf{c}) = \left(\frac{H_k}{(z-c_1)(z-c_2)\cdots(z-c_k)} + \cdots + \frac{H_1}{(z-c_1)} + H_{\text{res}}\right) \frac{1}{(z-c_0)}$$

and call this function on \mathbb{C}^{k+1} the *unfolding* of H. Since obviously we have $H(\mathbf{0}) = H$, it is a deformation of H. Moreover we have the partial fraction decomposition

$$H(\mathbf{c}) = \frac{A^{[0]}}{z - c_0} + \frac{A^{[1]}}{z - c_1} + \dots + \frac{A^{[k]}}{z - c_k}$$

which is the sum of regular singular canonical forms for generic \mathbf{c} , i.e., \mathbf{c} satisfying $c_i \neq c_j$, $i \neq j$. Thus $H(\mathbf{c})$ gives a deformation of the unramified irregular singular canonical form H to the sum of regular singular canonical forms $H(\mathbf{c})$.

Let us look at this deformation $H(\mathbf{c})$ in more detail. Now we introduce a stratification on \mathbb{C}^{k+1} associated to the partition of the index set $\{0, 1, \ldots, k\}$ as follows. For a partition $\mathcal{I} = I_0 \sqcup I_2 \sqcup \cdots \sqcup I_r = \{0, 1, \ldots, k\}$, we consider the subset of \mathbb{C}^{k+1}

$$C(\mathcal{I}) = \left\{ (c_0, c_1, \dots, c_k) \in \mathbb{C}^{k+1} \middle| \begin{array}{c} c_i = c_j & \text{if } i, j \in I_l \text{ for some } l, \\ c_i \neq c_j & \text{otherwise} \end{array} \right\}$$

and these subsets define the stratification of \mathbb{C}^{k+1} ,

$$\mathbb{C}^{k+1} = \bigsqcup_{\mathcal{I}: \text{ partitions of } \{0,1,\ldots,k\}} \mathcal{C}(\mathcal{I}),$$

see Section 5.2. Then by setting $I_j = \{i_{[j,0]}, i_{[j,1]}, \ldots, i_{[j,k_j]}\}$, we obtain the partial fraction decomposition

$$H(\mathbf{c}) = \sum_{j=0}^{r} \left(\frac{A_{k_j}^{[j]}}{(z - c_{i_{[j,0]}})^{k_j}} + \dots + \frac{A_1^{[j]}}{z - c_{i_{[j,0]}}} + A_0^{[j]} \right) \frac{1}{z - c_{i_{[j,0]}}}$$

for $\mathbf{c} \in \mathcal{C}(\mathcal{I})$ which is again the sum of unramified canonical forms whose pole orders are the cardinalities $\sharp I_i$ of the components in the partition \mathcal{I} .

Thus for each fixed $\mathbf{c} \in \mathbb{C}^{k+1}$, there exists the unique stratum $\mathcal{C}(\mathcal{I})$ containing \mathbf{c} and we may regard the deformed canonical form $H(\mathbf{c})$ as the collection

$$H(\mathbf{c}) = \left(\left(\frac{A_{k_j}^{[j]}}{(z - c_{i_{[j,0]}})^{k_j}} + \dots + \frac{A_1^{[j]}}{z - c_{i_{[j,0]}}} + A_0^{[j]} \right) \frac{1}{z - c_{i_{[j,0]}}} \right)_{j=0,\dots,r}$$

of canonical forms at $z = c_{i[j,0]}$, j = 0, ..., r. Therefore $(H(\mathbf{c}))_{\mathbf{c}\in\mathbb{C}^{k+1}}$ can be seen as a family of collections of unramifed canonical forms associated to the stratification of \mathbb{C}^{k+1} . On the unique closed stratum $\mathcal{C}(\{0, ..., k\})$ corresponding to the trivial partition $\{0, ..., k\} = \{0, ..., k\}, H(\mathbf{c})$ are of the forms

$$H(\mathbf{c}) = \left(\frac{H_k}{(z-c_0)^k} + \dots + \frac{H_1}{(z-c_0)} + H_{\text{res}}\right) \frac{1}{(z-c_0)}$$

which are essentially same as the original canonical form H, and also on the unique open stratum $\mathcal{C}(\{0\} \sqcup \cdots \sqcup \{k\})$ corresponding to the finest partition $\{0, \ldots, k\} = \{0\} \sqcup \cdots \sqcup \{k\}$, $H(\mathbf{c})$ are collections of regular singular canonical forms. And on the other strata, $H(\mathbf{c})$ are collections canonical forms with intermediate singularities.

1.3 Statement of the main theorem

Now we explain the main theorem of this paper. Let G be a connected reductive linear algebraic group defined over \mathbb{C} . Let $|D| = \{a_1, a_2, \ldots, a_d\}$ be a finite subset of \mathbb{P}^1 and $D = \sum_{i=1}^d (k_{a_i} + 1)a_i$ be an effective divisor with non negative integers k_{a_i} . Let us consider a collection $\mathbf{H} = (H^{(a)})_{a \in |D|}$ of unramified canonical forms

$$H^{(a)} dz_a = \left(\frac{H_{k_a}^{(a)}}{z_a^{k_a}} + \dots + \frac{H_1^{(a)}}{z_a} + H_{\rm res}^{(a)}\right) \frac{dz_a}{z_a}$$

for $a \in |D|$, where z_a are coordinate functions on \mathbb{P}^1 centered at a. Then by following the paper [5] by Boalch, we can define the moduli space $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}}$ of irreducible meromorphic G-connections with local canonical form $H^{(a)}$ at each singular points $a \in |D|$, see Definition 7.12. Note that $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}}$ is a holomorphic symplectic orbifold if it is nonempty, see Proposition 7.16.

For each $a \in |D|$ we can consider the deformation $H^{(a)}(\mathbf{c}^{(a)})$ of the canonical form $H^{(a)}$ and the parameter space \mathbb{C}^{k_a+1} with the stratification as in the previous subsection. Then the stratifications on \mathbb{C}^{k_a+1} define the product stratification on $\prod_{a\in|D|}\mathbb{C}^{k_a+1}$. As we explained before, for a fixed $\mathbf{c} = (\mathbf{c}^{(a)})_{a\in|D|} \in \prod_{a\in|D|}\mathbb{C}^{k_a+1}$, there exists the unique stratum $\prod_{a\in|D|} \mathcal{C}(\mathcal{I}^{(a)})$ containing \mathbf{c} and we can regard $\mathbf{H}(\mathbf{c}) = (H^{(a)}(\mathbf{c}^{(a)}))_{a\in|D|}$ as a collection of canonical forms.

Therefore for each fixed $\mathbf{c} \in \prod_{a \in |D|} \mathbb{C}^{k_a+1}$, we can consider the moduli space $\mathcal{M}_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}}$ of irreducible meromorphic *G*-connections associated to the collection $\mathbf{H}(\mathbf{c})$ of canonical forms. Then in the following main theorem we construct a holomorphic family of these moduli spaces.

Theorem 1.1 (Theorem 11.5). Suppose $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}} \neq \emptyset$. Then there exists an open neighborhood $\widetilde{\mathbb{B}}_{\mathbf{H}}$ of **0** in $\prod_{a \in |D|} \mathbb{C}^{k_a+1}$ and a complex orbifold $\mathcal{M}_{\mathbf{H},\widetilde{\mathbb{B}}_{\mathbf{H}}}^{\mathrm{ir}}$ with a holomorphic map $\pi_{\widetilde{\mathbb{B}}_{\mathbf{H}}} : \mathcal{M}_{\mathbf{H},\widetilde{\mathbb{B}}_{\mathbf{H}}}^{\mathrm{ir}} \to \widetilde{\mathbb{B}}_{\mathbf{H}}$ satisfying the following.

1. The orbifold $\mathcal{M}_{\mathbf{H},\widetilde{\mathbb{B}}_{\mathbf{H}}}^{\mathrm{ir}}$ is a deformation of $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}}$, i.e., the map $\pi_{\widetilde{\mathbb{B}}_{\mathbf{H}}} : \mathcal{M}_{\mathbf{H},\widetilde{\mathbb{B}}_{\mathbf{H}}}^{\mathrm{ir}} \to \widetilde{\mathbb{B}}_{\mathbf{H}}$ is a surjective submersion and we have

$$\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}} \cong \pi_{\mathbb{B}_{\mathbf{H}}}^{-1}(\mathbf{0}).$$

- 2. Every stratum $\prod_{a \in |D|} C(\mathcal{I}^{(a)})$ of $\prod_{a \in |D|} \mathbb{C}^{k_a+1}$ has the nonempty intersection with the base space $\widetilde{\mathbb{B}}_{\mathbf{H}}$.
- 3. For each $\mathbf{c} \in \widetilde{\mathbb{B}}_{\mathbf{H}}$ the fiber $\pi_{\widetilde{\mathbb{B}}_{\mathbf{H}}}^{-1}(\mathbf{c})$ is isomorphic to an open dense subspace of $\mathcal{M}_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}}$.

As we explained before, if **c** is in the open stratum $\prod_{a \in |D|} \mathcal{C}(\{0\} \sqcup \cdots \sqcup \{k_a\}) \cap \mathbb{B}_{\mathbf{H}}$, then $\mathcal{M}_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}}$ is a moduli space of Fuchsian *G*-connections. Therefore this theorem says that every (non-empty) moduli space $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}}$ of *G*-connections with unramified irregular singularities has the deformation to a moduli space of Fuchsian *G*-connections. Then the question now arises:

What kind of moduli spaces $\mathcal{M}_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}}$ of meromorphic connections appear on the other strata of $\widetilde{\mathbb{B}}_{\mathbf{H}}$?

Looking at the examples in Section 1.1, we notice that many different types of differential equations appear during the unfolding of irregular singular points. For instance, as the unfolding of the triconfluent Heun equation, one can obtain the biconfluent Heun, doubly-confluent Heun, the confluent Heun, and the Heun equation. Therefore the above question for the deformation of moduli spaces of connections will be translated to the following natural question for the unfolding of irregular singularities of differential equations:

What kind of differential equations appear from a given equation through the unfolding of its irregular singularities?

In the next subsection, with regard to these questions, we shall extract a discrete data from the canonical forms **H** which is called the spectral types, and also we introduce the unfolding diagram of the spectral types. Then the spectral types and the unfolding diagram give us a combinatorial and diagrammatic explanation of how the singularities and local canonical forms of G-connections change and relate with each other during the deformation of the moduli space $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}}$.

Now let us give some comments for the related previous works. The relationship between the confluence/unfolding of singular points of linear differential equations and the deformation of their moduli spaces have already been studied by several researchers, especially in the case of $G = \operatorname{GL}_n$. In [22], Inaba constructed a 1-parameter deformation of moduli spaces of meromorphic connections with unramified irregular singularities and also discussed applications to their isomonodromic deformations. The above main theorem is a generalization of Theorem 2.11 in [22] where Inaba dealt only with a special unramified irregular singularities which he called *generic* unramified irregular singularities. Also in [13] Gaiur-Mazzocco-Rubtsov discussed the confluence of singular points of linear differential equations and a deformation of phase spaces of their isomonodromic deformations in a different point of view from our approach, and they moreover discussed their quantizations.

Analyzing Stokes structures is an important application of the unfolding which is not dealt with in this paper. There are many classical and recent studies around this area, see [41] by Schäfke, [39] by Ramis, [48] by Zhang, [12] by Glutsuk, [30] by Lambert-Rousseau, [21] by Hurtubise-Rousseau, [27] by Klimeš, and their references.

1.4 Spectral types and unfolding diagrams

Let us take an unramified canonical form $H dz = \left(\sum_{i=1}^{k} H_i/z^i + H_{\text{res}}\right) dz/z$. Let $H_{\text{res}} = H_0 + J_0$ be the Jordan decomposition of H_{res} with the semisimple H_0 and the nilpotent J_0 . Let us fix a set of simple roots Π of G. Then under the action of the Weyl group of G, we can associate to H the sequence of subsets of Π ,

$$\Pi_i := \{ \alpha \in \Phi \mid \alpha(H_k) = \dots = \alpha(H_{i+1}) = \alpha(H_i) = 0 \}$$

for i = 0, 1, ..., k. Let L_i be the Levi subgroup of G associated to Π_i . Then the pair

$$\operatorname{sp}(H) := (\Pi_k \supset \cdots \supset \Pi_1 \supset \Pi_0; [J_0])$$

of the sequence of subsets of Π and the L_1 -orbit of J_0 is called the *spectral type* of the canonical form H. For a collection $\mathbf{H} = (H^{(a)})_{a \in |D|}$ of canonical forms we can also consider the collection $\operatorname{sp}(\mathbf{H}) := (\operatorname{sp}(H^{(a)}))_{a \in |D|}$ of spectral types. On the other hand, without referring H, we can consider the pair $(\Pi_k \supset \cdots \supset \Pi_1 \supset \Pi_0; [J_0])$ of a sequence of subsets

of Π and the L_1 -orbit of a nilpotent element J_0 in the Lie algebra \mathfrak{l}_0 of L_0 . We call this pair an *abstract spectral type*. For more details, see Section 2.2.

Now we explain that the unfolding $H(\mathbf{c})$ of H is well captured by the spectral type $\mathrm{sp}(H)$. First we introduce the unfolding of spectral types. Let $S = (\prod_k \supset \cdots \supset \prod_1 \supset \prod_0; [J_0])$ be an abstract spectral type. We consider the set $\mathcal{P}_{[k+1]}$ of all partitions of the index set $\{0, 1, \ldots, k\}$ which has the natural partial order by the refinement of partitions. Let us take a partition $\mathcal{I} = I_0 \sqcup \cdots \sqcup I_r = \{0, 1, \ldots, k\}$ and write $I_j = \{i_{[j,0]} < i_{[j,1]} < \cdots < i_{[j,k_j]}\}, j = 0, 1, \ldots, r$. We may assume that $0 \in I_0$. Then to the partition \mathcal{I} we associate the collection of spectral types $S^{\mathcal{I}} = (S^{I_j})_{j=0,1,\ldots,r}$ defined by

$$S^{I_0} := (\Pi_{i_{[0,k_0]}} \supset \cdots \supset \Pi_{i_{[0,1]}} \supset \Pi_0; [J_0]),$$

$$S^{I_j} := (\Pi_{i_{[j,k_j]}} \supset \cdots \supset \Pi_{i_{[j,1]}} \supset \Pi_{i_{[j,0]}}; [0]), \ j = 1, \dots, r,$$

which corresponds to the decomposition of the sequence $\Pi_k \supset \cdots \prod_1 \supset \Pi_0$ with regard to the partition \mathcal{I} . Then the following says that this decomposition actually describes the unfolding $\mathbf{H}(\mathbf{c})$ of the collection $\mathbf{H} = (H^{(a)})_{a \in |D|}$ of canonical forms.

Proposition 1.2 (Corollary 5.5). Let $\widetilde{\mathbb{B}}_{\mathbf{H}}$ be the open neighborhood of $\mathbf{0} \in \prod_{a \in |D|} \mathbb{C}^{k_a+1}$ in Theorem 1.1. Then for each stratum $\prod_{a \in |D|} \mathcal{C}(\mathcal{I}^{(a)})$ of $\prod_{a \in |D|} \mathbb{C}^{k_a+1}$ and each $\mathbf{c} \in \widetilde{\mathbb{B}}_{\mathbf{H}} \cap \prod_{a \in |D|} \mathcal{C}(\mathcal{I}^{(a)})$, we have

$$\operatorname{sp}(\mathbf{H}(\mathbf{c})) = (\operatorname{sp}(H^{(a)})^{\mathcal{I}^{(a)}})_{a \in |D|}.$$

Namely, on each stratum $\prod_{a \in |D|} C(\mathcal{I}^{(a)})$, spectral types of $\mathbf{H}(\mathbf{c})$ are all the same and the spectral type on the stratum is the decomposition of the original spectral type $\operatorname{sp}(\mathbf{H})$ with respect to the partitions $\mathcal{I}^{(a)}$, $a \in |D|$.

Next we introduce a diagrammatic description of the unfolding of spectral types. Let

$$\mathbf{S} = (S_i)_{i=1,\dots,d} = (\Pi_{k_i}^{(i)} \supset \dots \supset \Pi_1^{(i)} \supset \Pi_0^{(i)}; [J_0^{(i)}])_{i=1,\dots,d}$$

be a collection of abstract spectral types. Under the poset structures of $\mathcal{P}_{[k_i+1]}$, the product $\prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$ naturally becomes a poset as well, i.e., for $(\mathcal{I}^{(i)})_{i=1,...,d}, (\mathcal{J}^{(i)})_{i=1,...,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$, we have $(\mathcal{I}^{(i)})_{i=1,...,d} \leq (\mathcal{J}^{(i)})_{i=1,...,d}$ if and only if $\mathcal{I}^{(i)} \leq \mathcal{J}^{(i)}$ for all $i = 1, \ldots, d$. Thus we obtain the Hasse diagram of the poset $\prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$. Then we attach to each vertex $(\mathcal{I}^{(i)})_{i=1,...,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$ the corresponding unfolding $\mathbf{S}^{(\mathcal{I}^{(i)})_{i=1,2,...,d}}$ of \mathbf{S} . Then we call the resulting diagram the *unfolding diagram* of \mathbf{S} . We notice that there may appear same collections of abstract spectral types in several vertices, namely, it may happen $\mathbf{S}^{(\mathcal{I}^{(i)})_{i=1,...,d}} = \mathbf{S}^{(\mathcal{I}^{(i)})_{i=1,...,d}}$ for some $(\mathcal{I}^{(i)})_{i=1,...,d}, (\mathcal{J}^{(i)})_{i=1,...,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$. Therefore by identifying these same collections of abstract spectral types we can obtain the smaller diagram, which we call the *reduced unfolding diagram* of \mathbf{S} . See Section 6.2 for the detail.

Let us see some examples. Let $G = \operatorname{GL}_2$ and we consider the canonical form Hwith $\operatorname{sp}(H) = (\emptyset \supset \emptyset \supset \emptyset \supset \emptyset; [0])$ which is the canonical form for the triconfluent Heun equation, see Section 4.2. Then the deformation of the moduli space $\mathcal{M}_{(H)}^{\operatorname{ir}}$ given in Theorem 1.1 is depicted by the following reduced unfolding diagram of the spectral type,



We notice that this diagram coincides with the diagram in Section 1.1 which describes the confluence of singular points of Heun equations. Therefore we can say that the deformation of $\mathcal{M}_{(H)}^{\mathrm{ir}}$ recovers the confluence procedure of the Heun equation.

Also let $G = GL_4$ and we consider the canonical form H with

$$\operatorname{sp}(H) = (\{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\} \supset \{e_{12}\}; [0]).$$

Here e_{ij} for $1 \leq i < j \leq 4$ are standard simple roots of GL_4 . Then the deformation of the moduli space $\mathcal{M}_{(H)}^{ir}$ is depicted by the reduced unfolding diagram,



This diagram also recovers the one in Section 1.1 which describes the confluence of singular points of the differential equation appeared in [25]. We can also check that the other diagrams describing the confluence of singular points in the lists obtained by Kawakami-Nakamura-Sakai in [25] can be recovered from our reduced unfolding diagrams.

Moreover moduli spaces $\mathcal{M}_{\mathbf{H}}^{ir}$ with dim $\mathcal{M}_{\mathbf{H}}^{ir} = 0$ correspond to differential equations without accessory parameters, and we can check that the deformations of $\mathcal{M}_{\mathbf{H}}^{ir}$ in Theorem 1.1 recover Oshima's versal unfolding families of differential equations without accessory parameters.

Therefore one can conclude that the deformation constructed in Theorem 1.1 gives a natural generalization of many classical and recent known examples of confluent/unfolding families of linear differential equations on \mathbb{P}^1 , and also a natural translation of the theory for the unfolding of singular points of differential equations to the theory for deformation of their moduli spaces of meromorphic connections.

1.5 Additive Deligne-Simpson problem

Let us explain an application of Theorem 1.1 to an existence problem for irreducible G-connections, so-called the additive Deligne-Simpson problem.

The additive Deligne-Simpson problem asks the existence of irreducible Fuchsian differential equations

$$\frac{dY}{dz} = \sum_{i=1}^{r} \frac{A_i}{z - a_i} Y \quad (A_i \in M_n(\mathbb{C}))$$

on \mathbb{P}^1 whose residue matrices A_i at $z = a_i$ for $i = 1, \ldots, r$ and $A_{\infty} := -\sum_{i=1}^r A_i$ at $z = \infty$ are contained in the prescribed conjugacy classes $C_i \subset M_n(\mathbb{C})$ for $i \in \{1, \ldots, r, \infty\}$. This problem is studied by Deligne, Simpson, Kostov and many other researchers, then finally solved in [10] by Crawley-Boevey by using the theory of quiver representations. In [28] by Kostov, the historical background for the Deligne-Simpson problem is explained in detail. There are many generalizations of this problem, for instance, in [6] by Boalch, [19] by

Hiroe-Yamakawa, and [15] by Hiroe, similar problems for linear differential equations with unramified irregular singularities are discussed, also in [29] by Kulkarni-Livesay-Matherne-Nguyen-Sage and [32] by Livesay-Sage-Nguyen similar problems for linear differential equations with ramified irregular singularities are discussed, and furthermore in [23] by Jakob-Yun a similar problem for G-connections with ramified irregular singularities is discussed.

Now we propose the following problem which is a natural generalizations of the problems considered in [6], [19], and [15] to *G*-connections. We may assume $|D| = \{a_1, \ldots, a_d\} \subset \mathbb{C}$ under the projective transformation.

Problem 1.3 (Additive Deligne-Simpson problem for **H**). Let us take a collection of unramified canonical forms $\mathbf{H} = (H^{(a)})_{a \in |D|}$ satisfying that the sum of residues $\sum_{a \in |D|} H_{\text{res}}^{(a)}$ is contained in the semisimple part \mathfrak{g}_{ss} of \mathfrak{g} . Then find an irreducible meromorphic *G*-connection

$$d_A = d + \sum_{a \in |D|} \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^i} \frac{dz}{z-a} \quad (A_i^{(a)} \in \mathfrak{g})$$

with singularities only on |D| in \mathbb{P}^1 such that

$$\sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}} \in \mathbb{O}_{H^{(a)}}$$

for all $a \in |D|$.

Here $\mathbb{O}_{H^{(a)}}$ stands for the $G(\mathbb{C}[[z_a]])$ -orbit through $H^{(a)} \in \mathfrak{g}(\mathbb{C}((z_a^{-1})))$, see Section 2.1 for the detail.

As in the previous subsections, we can consider an unfolding family of canonical forms $\mathbf{H}(\mathbf{c})_{\mathbf{c}\in U} = ((H^{(a)}(\mathbf{c}^{(a)}))_{a\in |D|})_{\mathbf{c}\in U}$ over an open neighborhood U of $\mathbf{0} \in \prod_{a\in |D|} \mathbb{C}^{k_a+1}$. Then the fibers $(H^{(a)}(\mathbf{c}^{(a)}))_{a\in |D|}$ on each stratum of $\prod_{a\in |D|} \mathbb{C}^{k_a+1}$ can be seen as collections of canonical forms. Then we can extend the additive Deligne-Simpson problem to the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c}\in U}$ as below.

Problem 1.4 (Additive Deligne-Simpson problem for an unfolding family of canonical forms). Let U be an open neighborhood of $\mathbf{0} \in \prod_{a \in |D|} \mathbb{C}^{k_a+1}$. Let $(\mathbf{H}(\mathbf{c}))_{\mathbf{c} \in U}$ be the unfolding of a collection of canonical forms $\mathbf{H} = (H^{(a)})_{a \in |D|}$ as above. Then find an meromorphic G-connection on \mathbb{P}^1 ,

$$d_{A(\mathbf{c})} = d + A(\mathbf{c}) \, dz$$

satisfying all the following conditions.

- 1. The g-valued 1-form $A(\mathbf{c}) dz$ depends holomorphically on $\mathbf{c} \in U$.
- 2. For each $\mathbf{c} \in U$, $d_{A(\mathbf{c})}$ is a solution of the additive Deligne-Simpson problem for $\mathbf{H}(\mathbf{c})$.

Namely, we would like to find a holomorphic family $d_{A(\mathbf{c})} = d + A(\mathbf{c}) dz$ of irreducible *G*-connections which gives simultaneous solutions for all the Deligne-Simpson problems associated to $\mathbf{H}(\mathbf{c})$ for $\mathbf{c} \in U$.

Then as a direct consequence of Theorem 1.1, it follows that if one can solve the additive Deligne-Simpson problem for \mathbf{H} , then one can simultaneously solve the every other additive Deligne-Simpson problem for the deformed collection of canonical forms $\mathbf{H}(\mathbf{c})$.

Theorem 1.5 (Theorem 6.4). Let **H** be a collection of unramified canonical forms as above. Let $d_A = d + A dz$ be a solution of the additive Deligne-Simpson problem for **H**. Then there exists an open neighborhood U of $\mathbf{0} \in \prod_{a \in |D|} \mathbb{C}^{k_a+1}$ and a holomorphic family of G-connections $d_{A(\mathbf{c})} = d + A(\mathbf{c}) dz$ ($\mathbf{c} \in U$) on \mathbb{P}^1 such that

- $d_{A(\mathbf{0})} = d_A$,
- $d_{A(\mathbf{c})} = d + A(\mathbf{c}) dz$ is a solution of the additive Deligne-Simpson for the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c}\in U}$.

In particular, the following are equivalent.

- 1. The additive Deligne-Simpson problem for H has a solution.
- 2. There exists an open neighborhood U of $\mathbf{0} \in \prod_{a \in |D|} \mathbb{C}^{k_a+1}$ and the additive Deligne-Simpson problem for the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c} \in U}$ has a solution.

1.6 A conjecture by Oshima

Let us explain another existence problem proposed by Oshima which is closely related to the additive Deligne-Simpson problem.

Let us consider a collection $\mathbf{S} = (S_i)_{i=1,\dots,d}$ of abstract spectral types. We say that the collection \mathbf{S} is *irreducibly realizable* if there exists an irreducible meromorphic *G*connection which has \mathbf{S} as its spectral type. Precisely to say, \mathbf{S} is irreducibly realizable if and only if there exists a collection $\mathbf{H} = (H^{(a_i)})_{i=1,\dots,d}$ of unramified canonical forms such that

- 1. $\mathbf{S} = \operatorname{sp}(\mathbf{H}),$
- 2. the additive Deligne-Simpson problem for **H** has a solution.

Here a solution $d_A = d + A dz$ of this additive Deligne-Simpson problem is called a *realization* of **S**.

Furthermore, a collection $\mathbf{S} = (S_i)_{i=1,2,\dots,d}$ of spectral types is said to be *versally* realizable when there exists a collection $\mathbf{H} = (H^{(a_i)})_{i=1,2,\dots,d}$ of unramified canonical forms satisfying

- 1. S = sp(H),
- 2. there exists an open neighborhood U of $\mathbf{0} \in \prod_{i=1}^{d} \mathbb{C}^{k_i+1}$ and the additive Deligne-Simpson problem for the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c}\in U}$ has a solution.

Then in [37] and [38], Oshima proposed the following conjecture which states that the realizability of \mathbf{S} is equivalent to the versal realizability of it, and moreover equivalent to the realizability of the *finest unfolding*

$$\mathbf{S}^{\text{reg}} := \mathbf{S}^{(\{0\} \sqcup \cdots \sqcup \{k_i\})_{i=1,\dots,d}} = \left((\Pi_0^{(1)}; [J_0^{(1)}]), (\Pi_1^{(1)}; [0]), \dots, (\Pi_{k_1}^{(1)}; [0]), \dots, (\Pi_{k_d}^{(d)}; [J_0^{(d)}]), (\Pi_1^{(d)}; [0]), \dots, (\Pi_{k_d}^{(d)}; [0]) \right)$$

of **S**. Here we note that the finest unfolding \mathbf{S}^{reg} is the collection of spectral types for regular singular canonical forms associated to the finest partitions $\{0\} \sqcup \cdots \sqcup \{k_i\} = \{0, \ldots, k_i\}, i = 1, \ldots, d$.

Conjecture 1.6 (Conjecture 6.6, Oshima [37], [38]). Let $\mathbf{S} = (S_i)_{i=1,...,d}$ be a collection of abstract spectral types. Then the following are equivalent.

- 1. The collection \mathbf{S} is irreducibly realizable.
- 2. The collection **S** is versally realizable.
- 3. The finest unfolding \mathbf{S}^{reg} of \mathbf{S} is irreducibly realizable.

In [37] and [38], Oshima showed that this conjecture is true when $G = \operatorname{GL}_n$ and $-2 \leq \operatorname{rig}(\mathbf{S}) \leq 2$. Here $\operatorname{rig}(\mathbf{S})$ is the index of rigidity of \mathbf{S} which will be explained in Section 4.1.

With regard to this conjecture, we show the following as a corollary of Theorems 1.1 and 1.5.

Theorem 1.7 (Theorem 6.5). Let $\mathbf{S} = (S_i)_{i=1,...,d}$ be a collection of abstract spectral types. Suppose that \mathbf{S} is irreducibly realizable and let $d_A = d + A dz$ be a realization of \mathbf{S} . Then there exists an open neighborhood U of $\mathbf{0} \in \prod_{i=1}^d \mathbb{C}^{k_{a_i}+1}$ and a holomorphic family of G-connections $d_{A(\mathbf{c})} = d + A(\mathbf{c}) dz$ ($\mathbf{c} \in U$) on \mathbb{P}^1 such that

- $d_{A(\mathbf{0})} = d_A$,
- for each $(\mathcal{I}_i)_{i=1,\dots,d} \in \prod_{i=1}^d \mathcal{P}_{[k_i+1]}$ and $\mathbf{c} \in U \cap \prod_{i=1}^d \mathcal{C}(\mathcal{I}^{(i)}), d_{A(\mathbf{c})} = d + A(\mathbf{c}) dz$ is a realization of $\mathbf{S}^{(\mathcal{I}_i)_{i=1,\dots,d}}$.

In particular, the following are equivalent.

- 1. The collection \mathbf{S} of abstract spectral types is irreducibly realizable.
- 2. For every $(\mathcal{I}^{(i)})_{i=1,\ldots,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$, the collection $\mathbf{S}^{(\mathcal{I}^{(i)})_{i=1,\ldots,d}}$ is irreducibly realizable.

As a result we obtain the following.

Corollary 1.8. In Conjecture 1.6, 1 and 2 are equivalent and the implication $1 \Rightarrow 3$ holds true.

In the case of $G = \operatorname{GL}_n$, the implication $3 \Rightarrow 1$ in Conjecture 1.6 is shown in [16]. Therefore combining these results, we obtain the affirmative solution of the conjecture by Oshima in the case of $G = \operatorname{GL}_n$.

Acknowledgements

This project has been started since around 2018 when the conference in honor of Toshio Oshima was held at Josai University. Since then the author had many opportunities to give talks in conferences and seminars, discuss with many researchers, and receive many valuable comments. The author would like to thank all of them, Philip Boalch, Masahiro Futaki, Sampei Hirose, Michiaki Inaba, Kohei Iwaki, Shingo Kamimoto, Tatsuki Kuwagaki, Frank Loray, Takuro Mochizuki, Hiraku Nakajima, Vladimir Rubtsov, Masa-Hiko Saito, Shinji Sasaki, Yoshitsugu Takei, and Daisuke Yamakawa. Especially the author express his gratitude to Hiroshi Kawakami, Akane Nakamura, and Hidetaka Sakai for their work breaking a new ground for the unfolding of singularities of differential equations. The author thanks Shunya Adachi for his careful reading of the draft of this article. Finally the author express his deepest gratitude of thanks to Toshio Oshima for his inspiring works and many valuable discussions.

2 Meromorphic *G*-connections on the formal punctured disk

In this section, we shall recall the local and formal classification theory of meromorphic \mathfrak{g} -valued connections following the work by Babbitt and Varadarajan [4] which is a natural generalization of the classical classification theory given by Hukuhara [20], Turrittin [45], and also by Levelt [31]. Also we shall introduce some numerical invariants associated with the canonical forms of meromorphic connections on the formal punctured disk: δ -invariants and spectral types.

Let us fix some notations. For an indeterminate z, $\mathbb{C}[z]$, $\mathbb{C}[[z]]$, and $\mathbb{C}((z))$ denote the ring of polynomials, formal power series, and field of formal Laurent series of z with coefficients in \mathbb{C} respectively. The *order* of $f = \sum_{i=m}^{\infty} a_i z^i \in \mathbb{C}((z))$ is the smallest exponent of z^n which has non-zero coefficient, i.e., $\operatorname{ord}(f) := \min\{n \in \mathbb{Z} \mid a_n \neq 0\}$.

We denote the ideal of a commutative ring \mathcal{R} generated by $r_1, r_2, \ldots, r_l \in \mathcal{R}$ by $\langle r_1, r_2, \ldots, r_l \rangle_{\mathcal{R}}$ or $\langle r_1, r_2, \ldots, r_l \rangle_{\mathcal{R}}$.

In this paper, G denotes a fixed connected reducitve affine algebraic group defined over \mathbb{C} , and \mathfrak{g} denotes its Lie algebra. Let H be a closed subgroup of G and \mathfrak{h} its Lie algebra. Let \mathcal{R} be a unitial \mathbb{C} -algebra. Then $H(\mathcal{R})$ denotes the group of \mathcal{R} -valued points of H. Namely, for the coordinate ring $\mathcal{O}(H)$ of H, $H(\mathcal{R})$ is defined by $H(\mathcal{R}) := \operatorname{Hom}_{\mathbb{C}\text{-alg}}(\mathcal{O}(H), \mathcal{R})$ whose group multiplication comes from the comultiplication of the Hopf algebra $\mathcal{O}(H)$. We also define the corresponding Lie algebra by $\mathfrak{h}(\mathcal{R}) := \mathfrak{h} \otimes_{\mathbb{C}} \mathcal{R}$. The Lie bracket on $\mathfrak{h}(\mathcal{R})$ is defined by $[X_1 \otimes r_1, X_2 \otimes r_2] := [X_1, X_2] \otimes r_1 r_2$ for $X_1, X_2 \in \mathfrak{h}$ and $r_1, r_2 \in \mathcal{R}$. Here we notice that in the definition of $\mathfrak{h}(\mathcal{R})$ the \mathbb{C} -algebra \mathcal{R} need not to be unitial. Thus we can consider the Lie algebra $\mathfrak{h}(\mathcal{R})$ even for non-unitial \mathbb{C} -algebra \mathcal{R} .

We fix an embedding $G \hookrightarrow \operatorname{GL}_N$ throughout the remaining of this paper. We frequently write $G(\mathbb{C}) = G$.

For a group H acting on a set M, the stabilizer subgroup for $m \in M$ is denoted by $\operatorname{Stab}_H(m) := \{h \in H \mid h \cdot m = m\}.$

For a Lie algebra \mathfrak{h} and $X \in \mathfrak{h}$, \mathfrak{h}_X denotes the kernel of the linear map $\operatorname{ad}(X) \in \operatorname{End}_{\mathbb{C}}(\mathfrak{h})$.

2.1 Hukuhara-Turrittin theory for g-valued connections

Let us recall covariant exterior derivatives of principal G-bundles on the formal punctured disk.

Definition 2.1. Let \mathcal{R} be either $\mathbb{C}[\![z]\!]$ or $\mathbb{C}(\!(z)\!)$. Let us take $A \in \mathfrak{g}(\mathbb{C}(\!(z)\!))$ and define the \mathbb{C} -linear map $d_A \colon \mathfrak{g}(\mathcal{R}) \to \mathfrak{g}(\mathcal{R}) \otimes_{\mathcal{R}} \mathbb{C}(\!(z)\!) dz$ by $d_A := d + A dz$, namely, d_A is define by the formula

 $d_A X := dX + [A, X] dz, \quad X \in \mathfrak{g}(\mathcal{R}).$

Here $\mathbb{C}((z)) dz := \{f(z)dz \mid f(z) \in \mathbb{C}((z))\}$ is the space of formal meromorphic 1-forms and $d: \mathcal{R} \to \mathbb{C}((z))dz$ is the formal exterior derivative defined by $df(z) := (\frac{d}{dz}f(z)) dz$ for $f(z) \in \mathcal{R}$.

Note that d_A satisfies the Leibniz rule,

$$d_A(f \cdot X) = X \otimes df + f \cdot d_A(X), \quad f \in \mathcal{R}, \ X \in \mathfrak{g}(\mathcal{R}).$$

We call d_A a formal meromorphic exterior covariant derivative or formal meromorphic G-connection over \mathcal{R} .

In particular when $\operatorname{ord} A = -1$, we say that the formal connection d_A has the regular singularity of first kind at z = 0. Sometimes we omit the word, first kind, and just call regular singularity.

Definition 2.2 (Gauge transformation). Let \mathcal{R} be either $\mathbb{C}[\![z]\!]$ or $\mathbb{C}(\!(z)\!)$. Let us consider a *G*-connection $d_A: \mathfrak{g}(\mathcal{R}) \to \mathfrak{g}(\mathcal{R}) \otimes_{\mathcal{R}} \mathbb{C}(\!(z)\!) dz$ over \mathcal{R} . Then for $g \in G(\mathcal{R})$, we can define a new connection $\mathrm{Ad}(g) \circ d_A \circ \mathrm{Ad}(g^{-1})$. We call this transformation for *G*-connection over \mathcal{R} the gauge transformation over \mathcal{R} .

Remark 2.3. Let \mathcal{R} be as above. Let $\rho: G \hookrightarrow \operatorname{GL}_N$ be the fixed embedding. Then for each $g \in G(\mathcal{R})$, there uniquely exists $\delta_G(g) \in \mathfrak{g}(\mathcal{R})$ such that

$$d\rho(\delta_G(g)) = \left(\frac{d}{dz}\rho(g)\right)\rho(g)^{-1},$$

Then the transformed connection $\operatorname{Ad}(g) \circ d_A \circ \operatorname{Ad}(g^{-1})$ is of the form $d_{g[A]}$ with

$$g[A] := \operatorname{Ad}(g)(A) + \delta_G(g).$$

See 1.6 in [4] for the detail.

Let us recall the classification theory of G-connections under the gauge transformation.

Definition 2.4 (Canonical form). We say that $H \in \mathfrak{g}(\mathbb{C}((z)))$ is a *canonical form* if H dz is of the form

$$H dz = \left(\sum_{i=1}^{k} H_i z^{-i} + H_{\rm res}\right) \frac{dz}{z}$$

with semisimple elements H_i satisfying $[H_i, H_j] = 0$ and $[H_i, H_{res}] = 0$ for $1 \le i, j \le k$.

In particular $H_{irr} := \sum_{i=1}^{k} H_i z^{-(i+1)}$ is called the *irregular part* of the canonical form H. If the irregular part is zero, then we call H a *regular singular* canonical form

If the irregular part is zero, then we call ${\cal H}$ a $regular\ singular\ canonical\ form.$

As a natural generalization of the classical results established by Hukuhara, Turrittin, and Levelt, Babbit-Varadarajan showed that any meromorphic G-connections are reduced to canonical forms under the gauge transformation.

Theorem 2.5 (Babbitt-Varadarajan [4], cf. Hukukara [20], Turrittin [45], and Levelt [31]). Let us consider a meromorphic G-connection $d_A = d + A dz$ with $A \in \mathfrak{g}(\mathbb{C}((z)))$. Then there exists a field extension $\mathbb{C}((t))$ of $\mathbb{C}((z))$ satisfying $t^q = z$ with $q \in \mathbb{Z}_{\geq 1}$, and d_A is transformed to $d_H = d + H dt$ with a canonical form $H \in \mathfrak{g}(\mathbb{C}((t)))$ under the gauge transformation over $\mathbb{C}((t))$.

In this case d_H is called a canonical form of d_A . The minimal $q \in \mathbb{Z}_{\geq 1}$ to transform d_A to a canonical form is called the ramification index of d_A . If in particular q = 1, d_A is called an unramified connection.

Now we take an unramified canonical form

$$H dz = \left(\sum_{i=1}^{k} H_i z^{-i} + H_{\rm res}\right) \frac{dz}{z}$$

and the associated G-connection $d_H = d + H dz$ over $\mathbb{C}[\![z]\!]$. Then the isomorphic class of d_H under the gauge transformation of $G(\mathbb{C}[\![z]\!])$ is parametrized by the orbit

 $O_H := \left\{ \operatorname{Ad}(g)(H) + \delta_G(g) \in \mathfrak{g}(\mathbb{C}((z))) \mid g \in G(\mathbb{C}[[z]]) \right\}.$

As an analogue of O_H , Boalch introduced the truncated orbit of H in [5] which can be regarded as a coadjoint orbit of a finite dimensional complex Lie group as it will be explained later.

Definition 2.6 (Truncated orbit of canonical form). Let us take an unramified canonical form H and regard it as an element in $\mathfrak{g}(\mathbb{C}((z))/\mathbb{C}[[z]])$. Then the orbit of H under the adjoint action of $G(\mathbb{C}[[z]])$,

$$\mathbb{O}_H := \{ \mathrm{Ad}(g) H \in \mathfrak{g}(\mathbb{C}((z))/\mathbb{C}[\![z]\!]) \mid g \in G(\mathbb{C}[\![z]\!]) \}$$

is called the *truncated orbit* of H.

We can compare the orbits O_H and \mathbb{O}_H as follows.

Proposition 2.7. Let $\pi: \mathbb{C}((z)) \to \mathbb{C}((z))/\mathbb{C}[[z]]$ be the natural projection. Let us take an unramified canonical form $H \in \mathfrak{g}(\mathbb{C}((z)))$ as above. Let us consider the Lie subalgebra $\mathfrak{g}_{H_{irr}} = \{X \in \mathfrak{g} \mid [X, H_i] = 0 \text{ for all } i = 1, 2, \ldots, k\}$ of \mathfrak{g} .

Suppose that $\operatorname{ad}_{\mathfrak{g}_{H_{irr}}}(H_0) \in \operatorname{End}_{\mathbb{C}}(\mathfrak{g}_{H_{irr}})$ has no nonzero integers as its eigenvalues. Then $X \in O_H$ if and only if $\pi(X) \in \mathbb{O}_H$.

Proof. Lemma 6.2.1 and Proposition 9.3.2 in [4] show that if $\pi(X) \in \mathbb{O}_H$, then our assumption implies that there exists $g \in G(\mathbb{C}[\![z]\!])$ such that $\operatorname{Ad}(g)X + \delta_G(g) = H$, i.e., $X \in O_H$.

Conversely, we assume that $X \in O_H$. Then there exists $g \in G(\mathbb{C}[\![z]\!])$ such that $X = \operatorname{Ad}(g)H + \delta_G(g)$. On the other hand, we have $\operatorname{Ad}(g)H + \delta_G(g) \equiv \operatorname{Ad}(g)H \pmod{\mathbb{C}[\![z]\!]}$ since $\delta_G(g) \in \mathfrak{g} \otimes \mathbb{C}[\![z]\!]$. Thus $\pi(X) \in \mathbb{O}_H$.

Definition 2.8 (Truncated canonical form). Let H dz be an unramified canonical form. Let $d_A = d + A dz$ with $A \in \mathfrak{g}(\mathbb{C}((z)))$ be a meromorphic *G*-connection. If $\pi(A) \in \mathbb{O}_H$, then we call *H* the *truncated canonical form* of d_A . In this case, we say that d_A has the unramified truncated canonical form *H*.

If H satisfies the condition in Proposition 2.7, the truncated canonical form of d_A coincides with the usual canonical form appears in Theorem 2.5.

2.2 Spectral types of canonical forms

To an unramified canonical form

$$H dz = \left(\sum_{i=1}^{k} H_i z^{-i} + H_{\rm res}\right) \frac{dz}{z},$$

we associate the following discrete data, called a spectral type.

Let $H_{\text{res}} = H_0 + J_0$ be the Jordan decomposition with the semisimple H_0 and nilpotent J_0 . Then since H_0, H_1, \ldots, H_k are all semisimple and mutually commuting, $\bigoplus_{i=0}^k \mathbb{C}H_i$ becomes a Lie subalgebra of \mathfrak{g} which is toral, i.e., consists of semisimple elements. Thus

we can choose a maximal toral subalgebra, that is, a Cartan subalgebra \mathfrak{t} containing this toral subalgebra. Then we obtain the root space decomposition of \mathfrak{g} ,

$$\mathfrak{g}=\mathfrak{t}\oplus\bigoplus_{\alpha\in\Phi}\mathfrak{g}_{\alpha},$$

where $\mathfrak{g}_{\alpha} := \{X \in \mathfrak{g} \mid \operatorname{Ad}(t)X = \alpha(t)X, t \in \mathfrak{t}\}$ are root spaces with respect to $\alpha \in \mathfrak{t}^*$, and Φ is the set of roots, i.e., $\Phi := \{\alpha \in \mathfrak{t}^* \mid \mathfrak{g}_{\alpha} \neq 0\}$. Also we fix a simple system Π of the set of roots Φ and denote the associated positive and negative systems by Φ^+ and Φ^- respectively.

Recall that \mathfrak{t} has a fundamental domain

$$\mathcal{D} := \{ X \in \mathfrak{t} \mid \mathfrak{Re} \, \alpha(X) \ge 0 \text{ and if } \mathfrak{Re} \, \alpha(X) = 0 \text{ then } \mathfrak{Im} \, \alpha(X) \ge 0 \text{ for any } \alpha \in \Pi \}$$

under the action of the Weyl group W of G. Thus the W-action allows us to assume that $H_k \in \mathcal{D}$. Then we define the subset

$$\Pi_k := \{ \alpha \in \Phi \mid \alpha(H_k) = 0 \}$$

of Π . Similarly under the action of $W_k := \operatorname{Stab}_W(H_k)$, we may assume that H_{k-1} belongs to

$$\mathcal{D}_k := \{ X \in \mathfrak{t} \mid \mathfrak{Re}\,\alpha(X) \ge 0 \text{ and if } \mathfrak{Re}\,\alpha(X) = 0 \text{ then } \mathfrak{Im}\,\alpha(X) \ge 0 \text{ for any } \alpha \in \Pi_k \}$$

and then the we define the subset of Π_k by

$$\Pi_{k-1} := \{ \alpha \in \Pi_k \mid \alpha(H_{k-1}) = 0 \} = \{ \alpha \in \Phi \mid \alpha(H_k) = \alpha(H_{k-1}) = 0 \}.$$

Inductively for i = 0, 1, ..., k - 2, we may assume that H_i belongs to

 $\mathcal{D}_{i+1} := \{ X \in \mathfrak{t} \mid \mathfrak{Re}\,\alpha(X) \ge 0 \text{ and if } \mathfrak{Re}\,\alpha(X) = 0 \text{ then } \mathfrak{Im}\,\alpha(X) \ge 0 \text{ for any } \alpha \in \Pi_{i+1} \}$

under the action of $W_{i+1} := \operatorname{Stab}_{W_{i+2}}(H_{i+1})$ and then we define the subset of Π_{i+1} by

$$\Pi_i := \{ \alpha \in \Pi_{i+1} \mid \alpha(H_i) = 0 \}$$

= $\{ \alpha \in \Pi_{i+2} \mid \alpha(H_{i+1}) = \alpha(H_i) = 0 \}$
= \cdots
= $\{ \alpha \in \Pi \mid \alpha(H_k) = \cdots = \alpha(H_{i+1}) = \alpha(H_i) = 0 \}$

Therefore we obtain the sequence

$$\Pi = \Pi_{k+1} \supset \Pi_k \supset \cdots \supset \Pi_0$$

of the set of simple roots associated to the canonical form H.

For each Π_i , we can consider the subset of roots $\Phi_i := (\bigoplus_{\alpha \in \Pi_i} \mathbb{Z}\alpha) \cap \Phi$ generated by Π_i and then define the Lie subalgebra

$$\mathfrak{l}_i := \mathfrak{t} \bigoplus \bigoplus_{\alpha \in \Phi_i} \mathfrak{g}_\alpha = \{ X \in \mathfrak{l}_{i+1} \mid [X, H_i] = 0 \}.$$

Thus we also obtain the sequence of Lie subalgebras

$$\mathfrak{g} = \mathfrak{l}_{k+1} \supset \mathfrak{l}_k \supset \cdots \supset \mathfrak{l}_0$$

of \mathfrak{g} . For each pair l, m of positive integers with $0 \leq l < m \leq k$, \mathfrak{l}_l can be seen as the Levi subalgebra of the parabolic subalgebra of \mathfrak{l}_m ,

$$\mathfrak{p}_{l,m} := \mathfrak{l}_l \bigoplus \bigoplus_{\Phi_m^+ \setminus \Phi_l} \mathfrak{g}_\alpha.$$

Let us denote the nilpotent radical of $\mathfrak{p}_{l,m}$ and its opposite by

$$\mathfrak{u}_{l,m} := igoplus_{\Phi_{m}^{+} \setminus \Phi_{l}} \mathfrak{g}_{lpha}, \qquad \qquad \mathfrak{n}_{l,m} := igoplus_{\Phi_{m}^{-} \setminus \Phi_{l}} \mathfrak{g}_{lpha}$$

respectively. Here we put $\Phi_m^{\pm} := \Phi_m \cap \Phi^{\pm}$. Then we have the decomposition

$$\mathfrak{l}_m = \mathfrak{n}_{l,m} \oplus \mathfrak{l}_l \oplus \mathfrak{u}_{l,m}.$$

Especially when m = k we simply write $\mathfrak{p}_l := \mathfrak{p}_{l,k}$, $\mathfrak{u}_l := \mathfrak{u}_{l,k}$, $\mathfrak{n}_l := \mathfrak{n}_{l,k}$. Also we denote the analytic subgroups of $G = G(\mathbb{C})$ corresponding to \mathfrak{l}_l , $\mathfrak{p}_{l,m}$, $\mathfrak{u}_{l,m}$, $\mathfrak{n}_{l,m}$, \mathfrak{u}_l , and \mathfrak{n}_l by L_i , $P_{l,m}$, $U_{l,m}$, $N_{l,m}$, U_l , and N_l respectively.

Definition 2.9 (Spectral type). For the unramified canonical form H dz, the spectral type of H is the pair $sp(H) := (\Pi_H; [J_0])$ of the above sequence

$$\Pi_H \colon \Pi_k \supset \Pi_{k-1} \supset \cdots \supset \Pi_0$$

of subsets of Π , and the orbit $[J_0]$ of J_0 under the adjoint action of L_1 . Here we note that $J_0 \in \mathfrak{l}_0$ since it commute with all H_i for $i = 0, 1, \ldots, k$.

If a meromorphic G-connection $d_A = d + A dz$ has the above H as the unramified truncated canonical form, then we also call $(\Pi_H, [J_0])$ the spectral type of d_A .

Without referring a canonical form H, we can consider the pair $(\Pi_k \supset \Pi_{k-1} \supset \cdots \supset \Pi_0; [J_0])$ of a sequence of subsets of Π and L_1 -orbit of a nilpotent element $J_0 \in \mathfrak{l}_0$. We call this pair an *abstract spectral type*.

2.3 δ -invariants of canonical forms

We introduce some more invariants of a canonical form

$$H dz = \left(\sum_{i=1}^{k} H_i z^{-i} + H_{\rm res}\right) \frac{dz}{z}.$$

Definition 2.10 (Irregularities of canonical forms). For the above canonical form H dz, the quantity

$$\operatorname{Irr}(H) := \sum_{i=1}^{k} (\dim G - \dim L_i)$$

is called the *irregularity* of H dz.

This quantity is an analogue of the Komatsu-Malgrange irregularity of connections on the punctured disk. **Remark 2.11.** Let us consider the case $G = \operatorname{GL}_n$. We consider a GL_n -connection $d_A = d + A dz : \mathfrak{gl}_n(\mathbb{C}((z))) \to \mathfrak{gl}_n(\mathbb{C}((z))) \otimes_{\mathbb{C}((z))} \mathbb{C}((z)) dz$ with the above unramified canonical form H dz. We may rewrite

$$H dz = \left(\begin{pmatrix} h_1(z^{-1}) & & \\ & h_2(z^{-1}) & \\ & & \ddots & \\ & & & h_n(z^{-1}) \end{pmatrix} + H_{\text{res}} \right) \frac{dz}{z}$$

with polynomials $h_i(t) \in t \mathbb{C}[t]$ for i = 1, 2, ..., n. Then it is known that the Komatsu-Malgrange irregularity of d_A equals to the sum of degrees of the polynomials,

$$\sum_{i=1}^{n} \sum_{j=1}^{n} \deg (h_i(t) - h_j(t)).$$

Then easy computation deduces the equations

$$\sharp\{(i,j)\in\{1,2,\ldots,n\}^{\times 2}\mid \deg\left(h_i(t)-h_j(t)\right)=l\}=\dim L_{l+1}-\dim L_l$$

for l = 1, 2, ..., k. Here $\sharp A$ denotes the cardinality of a set A. Thus we obtain

$$\sum_{i=1}^{n} \sum_{j=1}^{n} \deg \left(h_i(t) - h_j(t) \right) = \sum_{l=1}^{k} l \cdot \left(\dim L_{l+1} - \dim L_l \right) = \sum_{i=1}^{k} (\dim G - \dim L_i).$$

This allows us to regard the irregularity Irr(H) defined above as an analogue of the Komatsu-Malgrange irregularity.

Let us introduce another quantity.

Definition 2.12 (δ -invariants of canonical forms). For the above canonical form H dz, the quantity

$$\delta(H) := \dim G + \operatorname{Irr}(H) - \dim \operatorname{Stab}_G(H)$$

is called the δ -invariant of H dz.

Definition 2.13 (Truncated δ -invariant). Let $d_A = d + A dz$ be a meromorphic *G*connection with the unramified truncated canonical form *H*, namely, $A \in \mathbb{O}_H$. Then define the *truncated* δ -invariant of d_A by $\delta^{tr}(d_A) := \delta(H)$.

Remark 2.14. In the case $G = GL_n$, the quantity

$$\delta(d_A) := \operatorname{rk}(d_A) + \operatorname{Irr}(d_A) - \dim_{\mathbb{C}} \operatorname{Ker}(d_A)$$

appears as an important invariant for a connection $d_A: \mathfrak{gl}_n(\mathbb{C}((z))) \to \mathfrak{gl}_n(\mathbb{C}((z))) \otimes_{\mathbb{C}((z))} \mathbb{C}((z)) dz$. Since we can check that this invariant coincides with our δ -invariant under the assumption in Proposition 2.7 in the case of $G = \operatorname{GL}_n$, we may regard our δ is an analogue of this classical quantity.

Indeed, this classical quantity appears as a local part of Euler characteristic of a meromorphic connection on a compact Riemann surfacea, see Deligne [11], Arinkin [2], and Malgrange [34] for the detail. Furthermore, a relevance of this quantity and the δ -invariant of singularities of a plane curve germ is pointed out in [17].

Let us notice that the invariants Irr(H) and $\delta(H)$ depend only on the spectral type $(\Pi_H, [J_0])$ of H dz. Therefore we can also define the δ -invariant $\delta(S)$ for an abstract spectral type S as well.

3 Truncated orbit of an unramified canonical form

The truncated orbit \mathbb{O}_H of an unramified canonical form

$$H dz = \left(\frac{H_k}{z^k} + \dots + \frac{H_1}{z} + H_{\rm res}\right) \frac{dz}{z}$$

was introduced in the paper [5] by Boalch and its many fundamental properties were studied there, see also a subsequent paper [19] by Hiroe and Yamakawa, and for the case of *G*-connections, we also refer Yamakawa's paper [47]. In this section, among these properties, we shall recall that \mathbb{O}_H can been seen as a coadjoint orbit of a finite dimensional complex Lie group. We shall also give a formula for the dimension of \mathbb{O}_H in terms of δ invariant of *H* introduced in the previous section.

3.1 Affine algebraic groups over the ring of formal power series

Let us recall some fundamental structures of $G(\mathbb{C}[\![z]\!])$ following Babbitt and Varadarajan [4]. Recall that $\mathbb{C}[\![z]\!]$ is a local ring with the maximal ideal $\mathfrak{m}_z := \langle z \rangle$. The projection map $\pi : \mathbb{C}[\![z]\!] \to \mathbb{C}[\![z]\!]/\mathfrak{m}_z \cong \mathbb{C}$ has the inclusion $\iota : \mathbb{C} \hookrightarrow \mathbb{C}[\![z]\!]$ as a section. Thus the induced group homomorphisms $\pi_* : G(\mathbb{C}[\![z]\!]) \to G$ and $\iota_* : G \to G(\mathbb{C}[\![z]\!])$ also satisfy $\pi_* \circ \iota_* = \mathrm{id}_G$, and we have the short exact sequence

$$1 \to \operatorname{Ker} \pi_* \to G(\mathbb{C}\llbracket z \rrbracket) \xrightarrow{\pi_*} G \to 1$$

with the right splitting ι_* . Therefore by denoting $G(\mathbb{C}[[z]])_1 := \operatorname{Ker} \pi_*$, we obtain the semidirect product decomposition

$$G(\mathbb{C}\llbracket z \rrbracket) \cong G \ltimes G(\mathbb{C}\llbracket z \rrbracket)_1.$$

Since $\operatorname{GL}_N(\mathbb{C}[\![z]\!])$ is written by

$$\operatorname{GL}_{N}(\mathbb{C}[\![z]\!]) = \left\{ \sum_{i=0}^{\infty} g_{i} z^{i} \, \middle| \, g_{0} \in \operatorname{GL}_{N}(\mathbb{C}), g_{i} \in M_{N}(\mathbb{C}), i = 1, 2, \ldots \right\},\$$

the normal subgroup $\operatorname{GL}_N(\mathbb{C}[\![z]\!])_1$ has the explicit description

$$\operatorname{GL}_{N}(\mathbb{C}\llbracket z \rrbracket)_{1} = \left\{ \sum_{i=0}^{\infty} g_{i} z^{i} \in \operatorname{GL}(\mathbb{C}\llbracket z \rrbracket) \middle| g_{0} = I_{N} \right\}.$$

Here I_N denotes the identity matrix of size N. Therefore if we fix an embedding $G \hookrightarrow GL_N$, we can regard

$$G(\mathbb{C}\llbracket z \rrbracket)_1 = \mathrm{GL}_N(\mathbb{C}\llbracket z \rrbracket)_1 \cap G(\mathbb{C}\llbracket z \rrbracket).$$

Let us consider the Lie algebra $\mathfrak{g}(\mathbb{C}[\![z]\!])_1$ of $G(\mathbb{C}[\![z]\!])_1$. Then we have the semidirect product decomposition

$$\mathfrak{g}(\mathbb{C}\llbracket z \rrbracket) \cong \mathfrak{g} \oplus \mathfrak{g}(\mathbb{C}\llbracket z \rrbracket)_1$$

Since $\mathfrak{g}(\mathbb{C}[\![z]\!])_1$ is the kernel of the projection

$$\mathfrak{g}(\mathbb{C}[\![z]\!]) \ni \sum_{i=0}^{\infty} X_i z^i \longmapsto X_0 \in \mathfrak{g},$$

we have

$$\mathfrak{g}(\mathbb{C}[\![z]\!])_1 = \left\{ \sum_{i=0}^{\infty} X_i z^i \in \mathfrak{g}(\mathbb{C}[\![z]\!]) \, \middle| \, X_0 = 0 \right\},\$$

and we can identify

$$\mathfrak{g}(\mathbb{C}[\![z]\!])_1 \cong \mathfrak{g} \otimes_{\mathbb{C}} \mathfrak{m}_z.$$

Here we regard \mathfrak{m}_z as a (non-unitial) \mathbb{C} -algebra.

Let us give an explicit description of elements in $G(\mathbb{C}[[z]])_1$. For a non-negative integer $l \in \mathbb{Z}_{\geq 0}$, we consider the quotient ring

$$\mathbb{C}[z]_l := \mathbb{C}[[z]] / \langle z^{l+1} \rangle \cong \mathbb{C}[z] / \langle z^{l+1} \rangle,$$

which is a finite dimensional \mathbb{C} -algebra with the unique maximal ideal $\mathfrak{m}_{z}^{(l)} := \langle z \rangle_{\mathbb{C}[z]_{l}}$. We may identify $\mathbb{C}[z]_{l}$ with the space of polynomials $\{\sum_{i=0}^{l} a_{i} z^{i} \mid a_{i} \in \mathbb{C}\} \cong \mathbb{C}^{l+1}$ of degree at most l as \mathbb{C} -vector spaces. Then the projection map $\pi_{l} : \mathbb{C}[z]_{l} \to \mathbb{C}[z]_{l}/\mathfrak{m}_{z}^{(l)} \cong \mathbb{C}$ gives the split short exact sequence

$$1 \to \operatorname{Ker}(\pi_l)_* \to G(\mathbb{C}[z]_l) \xrightarrow{(\pi_l)_*} G \to 1$$

and the semidirect product decomposition

$$G(\mathbb{C}[z]_l) = G \ltimes G(\mathbb{C}[z]_l)_1$$

by denoting $G(\mathbb{C}[z]_l)_1 := \text{Ker}(\pi_l)_*$. Similar argument as above shows that the Lie algebra $\mathfrak{g}(\mathbb{C}[z]_l)_1$ of $G(\mathbb{C}[z]_l)_1$ is isomorphic to $\mathfrak{g} \otimes_{\mathbb{C}} \mathfrak{m}_z^{(l)}$.

Since the ring of formal power series can be obtained as the projective limit of $\mathbb{C}[z]_l$, i.e., $\mathbb{C}[\![z]\!] = \varprojlim_l \mathbb{C}[z]_l$, we have

$$G(\mathbb{C}[\![z]\!]) = \operatorname{Hom}_{\mathbb{C}\text{-alg}}(\mathcal{O}(G), \varprojlim_{l} \mathbb{C}[z]_{l}) \cong \varprojlim_{l} \operatorname{Hom}_{\mathbb{C}\text{-alg}}(\mathcal{O}(G), \mathbb{C}[z]_{l}) = \varprojlim_{l} G(\mathbb{C}[z]_{l})$$

On the other hand, the induced sequence

$$1 \longrightarrow \varprojlim_{l} \operatorname{Ker} (\pi_{l})_{*} \longrightarrow \varprojlim_{l} G(\mathbb{C}[z]_{l}) \longrightarrow \varprojlim_{l} G = G$$

is also exact since the projective limit functor is left exact. Also since the right most arrow coincides with π_* , this is moreover a short exact sequence. Thus we have $G(\mathbb{C}[[z]])_1 = \text{Ker } \pi_* \cong \varprojlim_l (\pi_l)_* = \varprojlim_l G(\mathbb{C}[z]_l)_1.$

Lemma 3.1. The finite dimensional complex Lie group $G(\mathbb{C}[z]_l)_1$ is a connected and simply connected nilpotent Lie group.

Proof. Since $\mathfrak{m}_z^{(l)}$ is a nilpotent ideal, the Lie algebra $\mathfrak{g}(\mathbb{C}[z]_l)_1 \cong \mathfrak{g} \otimes_{\mathbb{C}} \mathfrak{m}_z^{(l)}$ is also nilpotent. Thus $G(\mathbb{C}[z]_l)_1$ is a nilpotent Lie group. Also since

$$\operatorname{GL}_N(\mathbb{C}[z]_l)_1 \cong \left\{ I_N + X_1 z + \cdots + X_l z^l \, \big| \, X_i \in M_N(\mathbb{C}) \right\} \cong M_N(\mathbb{C})^{\oplus l},$$

is connected and simply connected nilpotent Lie group, it follows from Corollary 1.2.2 in [9] that the Lie subgroup $G(\mathbb{C}[z]_l)_1$ is connected and simply connected nilpotent Lie group as well.

The nilpotent Lie algebra $\mathfrak{g}(\mathbb{C}[z]_l)_1$ is equipped with the gradation with respect to the degree of z in $\mathbb{C}[z]_l$. Namely if we set

$$\mathfrak{g}^{[i]} := \{ X z^i \in \mathfrak{g}(\mathbb{C}[z]_l)_1 \mid X \in \mathfrak{g} \} = \mathfrak{g} \otimes_{\mathbb{C}} \left((\mathfrak{m}_z^{(l)})^i / (\mathfrak{m}_z^{(l)})^{i-1} \right)$$

for i = 1, 2, ..., l, then we have the natural decomposition

$$\mathfrak{g}(\mathbb{C}[z]_l)_1 = \bigoplus_{i=1}^l \mathfrak{g}^{[i]}$$

satisfying

$$[\mathfrak{g}^{[i]},\mathfrak{g}^{[j]}] \subset \mathfrak{g}^{[i+j]}$$

for i, j = 1, 2, ..., l.

Proposition 3.2. The exponential map $\exp: \mathfrak{g}(\mathbb{C}[z]_l)_1 \to G(\mathbb{C}[z]_l)_1$ is biholomorphic. Furthermore the map

$$\mathfrak{g}(\mathbb{C}[z]_l)_1 = \bigoplus_{i=1}^l \mathfrak{g}^{[i]} \ni (Z_i)_{i=1,2,\dots,l} \longmapsto e^{Z_l} \cdots e^{Z_2} e^{Z_1} \in G(\mathbb{C}[z]_l)_1$$

is biholomorphic.

Proof. The first assertion follows from the well-known fact for nilpotent Lie groups, see Theorem 1.2.1 in [9].

The second assertion follows from the graded structure of $\mathfrak{g}(\mathbb{C}[z]_l)_1$ as follows. Let us take $g \in G(\mathbb{C}[z]_l)_1$. Then the first assertion tells us that there uniquely exists $Z \in \mathfrak{g}(\mathbb{C}[z]_l)_1$ such that $g = e^Z$. Then according to the grading, we can uniquely write $Z = Z_1 + Z^{(1)}$ by $Z_1 \in \mathfrak{g}^{[1]}$ and $Z^{(1)} \in \bigoplus_{i=2}^l \mathfrak{g}^{[i]}$. Thus the Campbell-Baker-Hausdorff formula shows that there uniquely exists $Z^{(2)} \in \bigoplus_{i=2}^l \mathfrak{g}^{[i]}$ such that

$$e^{-Z_1}e^Z = e^{Z^{(2)}}.$$

Similarly for $Z^{(2)}$ we have the unique decomposition $Z^{(2)} = Z_2 + Z^{(3)}$ by $Z_2 \in \mathfrak{g}^{[2]}$ and $X^{(3)} \in \bigoplus_{i=3}^l \mathfrak{g}^{[i]}$. Then moreover there uniquely exists $Z^{(3)} \in \bigoplus_{i=3}^l \mathfrak{g}^{[i]}$ such that

$$e^{-Z_2}e^{-Z_1}e^Z = e^{Z^{(3)}}.$$

This procedure allows us to find $(Z_i)_{i=1,2,\dots,l} \in \bigoplus_{i=1}^l \mathfrak{g}^{[i]}$ uniquely from $g \in G(\mathbb{C}[z]_l)_1$, and thus we obtain the well-defined map

$$G(\mathbb{C}[z]_l)_1 \ni g \longmapsto \sum_{i=1}^l Z_i \in \mathfrak{g}(\mathbb{C}[z]_l)_1$$

which is holomorphic by the construction. And obviously this map gives the inverse map of the above map. $\hfill \Box$

From this Proposition, we obtain the following explicit description of $G(\mathbb{C}[[z]])_1$.

Proposition 3.3 (Proposition 9.3.1 in [4]). Each $g(z) \in G(\mathbb{C}[[z]])_1$ has the unique infinite product expansion,

$$g(z) = \prod_{l=1}^{\infty} e^{X_l z^l} := \lim_{l \to \infty} e^{X_l z^l} \cdots e^{X_2 z^2} e^{X_1 z}, \quad (X_l \in \mathfrak{g}).$$

Proof. This follows from Proposition 3.2 and the isomorphism $G(\mathbb{C}[\![z]\!])_1 \cong \varprojlim_l G(\mathbb{C}[\![z]\!]_l)_1$.

Let us notice that the natural projection $\pi_l \colon G(\mathbb{C}[\![z]\!])_1 \to G(\mathbb{C}[\![z]\!])_1$ is surjective. Indeed, Proposition 3.2 says that every element in $G(\mathbb{C}[\![z]\!]_l)_1$ is of the form $e^{X_l z^l} \cdots e^{X_2 z^2} e^{X_1 z}$ for $X_i \in \mathfrak{g}$ which is the projection image of $e^{X_l z^l} \cdots e^{X_2 z^2} e^{X_1 z} \in G(\mathbb{C}[\![z]\!])_1$. Thus since $G(\mathbb{C}[\![z]\!]) = G \ltimes G(\mathbb{C}[\![z]\!])_1$ and $G(\mathbb{C}[\![z]\!]_l) = G \ltimes G(\mathbb{C}[\![z]\!]_l)_1$, the projection $\pi_l \colon G(\mathbb{C}[\![z]\!]) \to G(\mathbb{C}[\![z]\!]_l)$ is surjective as well.

3.2 Truncated orbit as coadjoint orbit

The truncated orbit \mathbb{O}_H of an unramified canonical form $H \in \mathfrak{g}(\mathbb{C}((z))/\mathbb{C}[[z]])$ is defined as the orbit under the action of $G(\mathbb{C}[[z]])$. We shall explain that \mathbb{O}_H can be seen as a coadjoint orbit of a finite dimensional complex Lie group. Before explaining this fact, we prepare some notations. For a positive integer l, we consider a $\mathbb{C}[[z]]$ -module

$$\mathbb{C}[z^{-1}]_l := z^{-(l+1)} \mathbb{C}[\![z]\!] / \mathbb{C}[\![z]\!]$$

which can be also seen as a module of the quotient ring $\mathbb{C}[z]_l$. Then for a Lie subalgebra $\mathfrak{h} \subset \mathfrak{g}$, we can define the \mathbb{C} -vector space

$$\mathfrak{h}(\mathbb{C}[z^{-1}]_l) := \mathfrak{h} \otimes_{\mathbb{C}} \mathbb{C}[z^{-1}]_l$$

on which the action of the Lie algebra $\mathfrak{h}(\mathbb{C}[z]_l)$ is naturally defined via the Lie algebra structure of \mathfrak{h} and the $\mathbb{C}[z]_l$ -module structure of $\mathbb{C}[z^{-1}]_l$.

We shall frequently use the following basis and identifications as \mathbb{C} -vector spaces,

$$\mathbb{C}[z^{-1}]_l \cong \left\{ \sum_{i=0}^l \frac{a_i}{z^{i+1}} \, \middle| \, a_i \in \mathbb{C} \right\}, \qquad \mathbb{C}[z]_l \cong \left\{ \sum_{i=0}^l a_i z^i \, \middle| \, a_i \in \mathbb{C} \right\}.$$

According to these identifications, we set $f(0) := a_0$ for $f(z) = \sum_{i=0}^l a_i z^i \in \mathbb{C}[z]_l$ and $\underset{z=0}{\operatorname{res}} g(z) := b_0$ for $g(z) = \sum_{i=0}^l \frac{b_i}{z^{i+1}} \in \mathbb{C}[z^{-1}]_l$.

Let us introduce a non-degenerate bilinear pairing on $\mathfrak{g}(\mathbb{C}[z]_l) \times \mathfrak{g}(\mathbb{C}[z^{-1}]_l)$ as follows. Recall that the trace form

$$M_N(\mathbb{C}[z]_l) \times M_N(\mathbb{C}[z^{-1}]_l) \ni (A, B) \mapsto \operatorname{res}_{z=0}(\operatorname{tr}(AB)) \in \mathbb{C}$$
(1)

is non-degenerate. Since we have the embeddings $\mathfrak{g}(\mathbb{C}[z]_l) \hookrightarrow \mathfrak{gl}_N(\mathbb{C}[z]_l) = M_N(\mathbb{C}[z]_l)$ and $\mathfrak{g}(\mathbb{C}[z^{-1}]_l) \hookrightarrow \mathfrak{gl}_N(\mathbb{C}[z^{-1}]_l) = M_N(\mathbb{C}[z^{-1}]_l)$ induced by $G \hookrightarrow \operatorname{GL}_N$, this trace pairing defines the non-degenerate pairing on $\mathfrak{g}(\mathbb{C}[z]_l) \times \mathfrak{g}(\mathbb{C}[z^{-1}]_l)$. This non-degenerate pairing moreover gives us the identification

$$\mathfrak{g}(\mathbb{C}[z]_l)^* \cong \mathfrak{g}(\mathbb{C}[z^{-1}]_l).$$

We can consider the canonical form H as an element in $\mathfrak{g}(\mathbb{C}[z^{-1}]_l)$ by regarding $\mathfrak{g}(\mathbb{C}[z^{-1}]_l)$ as a subspace of $\mathfrak{g}(\mathbb{C}((z))/\mathbb{C}[[z]])$. Then we compare the orbits of H under the actions of $G(\mathbb{C}[[z]])$ and $G(\mathbb{C}[[z]]_l)$ as follows. The natural projection map $p_l \colon \mathbb{C}[[z]] \to \mathbb{C}[[z]]_l$ induces the short exact sequence

$$1 \to \operatorname{Ker}(p_l)_* \to G(\mathbb{C}[\![z]\!]) \xrightarrow{(p_l)_*} G(\mathbb{C}[\![z]\!]) \to 1.$$

As we previously remarked, the projection map $(p_l)_* \colon G(\mathbb{C}[\![z]\!]) \to G(\mathbb{C}[\![z]\!])$ is surjective. Since Ker $(p_k)_*$ is contained in the stabilizer group $\operatorname{Stab}_{G(\mathbb{C}[\![z]\!])}(H)$ of H, we also have the exact sequence

$$1 \to \operatorname{Ker}(p_k)_* \to \operatorname{Stab}_{G(\mathbb{C}[[z]])}(H) \xrightarrow{(p_k)_*|_{\operatorname{Stab}_{G(\mathbb{C}[[z]])}(H)}} \operatorname{Stab}_{G(\mathbb{C}[[z]_k)}(H) \to 1.$$

Therefore we obtain the isomorphism

$$G(\mathbb{C}[\![z]\!])/\mathrm{Stab}_{G(\mathbb{C}[\![z]\!])}(H) \cong G(\mathbb{C}[\![z]\!]_k)/\mathrm{Stab}_{G(\mathbb{C}[\![z]\!]_k)}(H),$$

which allows us to regard the truncated orbit \mathbb{O}_H as the $G(\mathbb{C}[z]_k)$ -orbit through $H \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k)$. Namely, \mathbb{O}_H can be regarded as the $G(\mathbb{C}[z]_k)$ -coadjoint orbit through $H \in \mathfrak{g}(\mathbb{C}[z]_k)^*$ under the identification $\mathfrak{g}(\mathbb{C}[z]_k)^* \cong \mathfrak{g}(\mathbb{C}[z^{-1}]_k)$.

3.3 δ -invariant and dimension of truncated orbit

We shall show that δ -invariants of unramified canonical forms represent dimensions of corresponding truncated orbits.

Let us consider the stabilizer of H.

Proposition 3.4 (Lemma 3.3 in [47]). The stabilizer of $H \in \mathfrak{g}(\mathbb{C}[z]_k)^*$ in $G(\mathbb{C}[z]_k)_1$ is of the following form,

$$\operatorname{Stab}_{(G(\mathbb{C}[z]_k)_1)}(H) = \left\{ e^{X_k z^k} \cdots e^{X_2 z^2} e^{X_1 z} \in G(\mathbb{C}[z]_k)_1 \, \middle| \, X_i \in \mathfrak{l}_i, \ i = 1, \dots, k \right\}.$$

We moreover have

$$\operatorname{Stab}_{G(\mathbb{C}[z]_k)}(H) = \operatorname{Stab}_G(H) \ltimes \operatorname{Stab}_{G(\mathbb{C}[z]_k)_1}(H).$$

Proof. Let us recall that for $g = e^{z^k X_k} \cdots e^{zX_1} \in G(\mathbb{C}[z]_k)_1$ and $B = \sum_{i=0}^k B_i z^{-i-1} \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k) = \mathfrak{g}(\mathbb{C}[z]_k)^*$, we have $\operatorname{Ad}^*(g)(B) = \sum_{i=0}^k C_i z^{-i-1}$ with

$$C_{i} = \sum_{l_{1}, l_{2}, \dots, l_{k} \ge 0} \frac{\operatorname{ad}(X_{k})^{l_{k}} \circ \dots \circ \operatorname{ad}(X_{2})^{l_{2}} \circ \operatorname{ad}(X_{1})^{l_{1}}(B_{i+l_{1}+2l_{2}+\dots+kl_{k}})}{l_{1}!l_{2}! \cdots l_{k}!}.$$

Let us apply this formula to the case $g \in \operatorname{Stab}_{G(\mathbb{C}[z]_k)_1}(H)$ and B = H. Then the equation $C_{k-1} = H_{k-1}$ implies

$$H_{k-1} = H_{k-1} + \operatorname{ad}(X_1)(H_k)$$

which gives us $X_1 \in \mathfrak{l}_k$. Secondly, the equation $C_{k-2} = H_{k-2}$ implies

$$H_{k-2} = H_{k-2} + \operatorname{ad}(X_1)(H_{k-1}) + \operatorname{ad}(X_2)(H_k).$$

Let us recall the decomposition $\mathfrak{g} = \mathfrak{g}_{H_k} \oplus \operatorname{ad}(H_k)(\mathfrak{g})$ and the equation $\mathfrak{g}_{H_k} = \mathfrak{l}_k$. Here we recall that \mathfrak{g}_{H_k} denotes the kernel of $\operatorname{ad}(H_k) \in \operatorname{End}_{\mathbb{C}}(\mathfrak{g})$. Then since $\operatorname{ad}(X_1)(H_{k-1}) \in \mathfrak{l}_k$ and $\operatorname{ad}(X_2)(H_k) \in \operatorname{ad}(H_k)(\mathfrak{g})$, we have $\operatorname{ad}(X_1)(H_{k-1}) = \operatorname{ad}(X_2)(H_k) = 0$. This implies $X_1 \in \mathfrak{l}_{k-1}$ and $X_2 \in \mathfrak{l}_k$.

Similarly, for i > 2 under the assumption $X_j \in \mathfrak{l}_{k-i+j+1}$ for $j = 1, 2, \ldots, i-1$ the equation $C_{k-i} = H_{k-i}$ implies

$$\sum_{l_1, l_2, \dots, l_{i-1} > 0} \frac{\operatorname{ad}(X_{i-1})^{l_{i-1}} \circ \dots \circ \operatorname{ad}(X_2)^{l_2} \circ \operatorname{ad}(X_1)^{l_1} (H_{k-i+l_1+2l_2+\dots+(i-1)l_{i-1}})}{l_1! l_2! \cdots l_{i-1}!} + \operatorname{ad}(X_i) (H_k) = 0$$

From the assumption, the first term belongs to \mathfrak{l}_k . Thus the above argument shows $\operatorname{ad}(X_i)(H_k) = 0$, i.e., $X_i \in \mathfrak{l}_k$. Further this implies that

$$\sum_{l_1, l_2, \dots, l_{i-2} > 0} \frac{\operatorname{ad}(X_{i-1})^{l_{i-1}} \circ \dots \circ \operatorname{ad}(X_2)^{l_2} \circ \operatorname{ad}(X_1)^{l_1} (H_{k-i+l_1+2l_2+\dots+(i-2)l_{i-2}})}{l_1! l_2! \cdots l_{i-2}!} + \operatorname{ad}(X_{i-1}) (H_{k-1}) = 0.$$

Here we used the equation $\operatorname{ad}(X_{i-1}) \circ \operatorname{ad}(X_1)(H_k) = \operatorname{ad}(X_{i-1})(H_k) = 0$ which follows from the assumption $X_1 \in \mathfrak{l}_{k-i+2}$ and $X_{i-1} \in \mathfrak{l}_k$. Then since first term belongs to \mathfrak{l}_{k-1} , the decomposition $\mathfrak{l}_k = (\mathfrak{l}_k)_{H_{k-1}} \oplus \operatorname{ad}(H_{k-1})(\mathfrak{l}_k)$ and the equation $(\mathfrak{l}_k)_{H_{k-1}} = \mathfrak{l}_{k-1}$ imply that $\operatorname{ad}(X_{i-1})(H_{k-1})$, i.e., $X_{i-1} \in \mathfrak{l}_{k-1}$. Iterating this procedure, we obtain $X_j \in \mathfrak{l}_{k-i+j}$ for $j = 1, 2, \ldots, i$.

Thus inductively we obtain $X_j \in l_j$ for j = 1, 2, ..., k which shows the desired equation for $\operatorname{Stab}_{G(\mathbb{C}[z]_k)_1}(H)$.

Let us see the second assertion. Take $g \in \operatorname{Stab}_{G(\mathbb{C}[z]_k)}(H)$ and decompose $g = g_0 \cdot g_1$ by $g_0 \in G$ and $g_1 \in G(\mathbb{C}[z]_k)_1$. Then since the constant term of g_1 is identity, we have

$$\operatorname{Ad}^*(g_1)(H) - H \equiv 0 \pmod{z^{-k}},$$

and moreover $g_0 \in \operatorname{Stab}_G(H_k) = L_k$. This implies that

$$\operatorname{Ad}^*(g_1)(H) = \operatorname{Ad}^*(g_0^{-1})(H) \in \mathfrak{l}_k(\mathbb{C}[z^{-1}]_k)$$

which further shows

$$g_1 \in \left\{ e^{X_k z^k} \cdots e^{X_2 z^2} e^{X_1 z} \in G(\mathbb{C}[z]_k)_1 \, \middle| \, X_i \in \mathfrak{l}_k, \ i = 1, \dots, k \right\}.$$

Then since H_k is in the center of l_k , we have

$$\operatorname{Ad}^{*}(g_{1})(H) - H \equiv 0 \pmod{z^{-(k-1)}},$$

and moreover $g_0 \in \operatorname{Stab}_G(H_k) \cap \operatorname{Stab}_G(H_{k-1}) = L_{k-1}$. Therefore we have

$$\operatorname{Ad}^{*}(g_{1})(H) = \operatorname{Ad}^{*}(g_{0}^{-1})(H) \in \mathfrak{l}_{k-1}(\mathbb{C}[z^{-1}]_{k})$$

which further shows

$$g_1 \in \left\{ e^{X_k z^k} \cdots e^{X_2 z^2} e^{X_1 z} \in G(\mathbb{C}[z]_k)_1 \, \middle| \, X_k \in \mathfrak{l}_k, \, X_i \in \mathfrak{l}_{k-1}, \, i = 1, \dots, k-1 \right\}.$$

Here we recall that $e^{X_k z^k}$ with $X_k \in \mathfrak{l}_k$ is in $\operatorname{Stab}_{G(\mathbb{C}[z]_k)_1}(H)$ by the first assertion. Iterating this procedure, we obtain the second assertion.

Corollary 3.5. Let $H dz \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k) dz$ be an unramified canonical form. Then we have

$$\dim \mathbb{O}_H = \delta(H).$$

Proof. Recall that

 $\dim G(\mathbb{C}[z]_k) = \dim \mathfrak{g}(\mathbb{C}[z]_k) = \dim \mathfrak{g} \times \dim \mathbb{C}[z]_k = (k+1) \cdot \dim G.$

Therefore from Proposition 3.4 we have

$$\dim \mathbb{O}_H = \dim G(\mathbb{C}[z]_k) - \dim \operatorname{Stab}_{G(\mathbb{C}[z]_k)}(H)$$
$$= (k+1) \cdot \dim G - \left(\sum_{i=1}^k \dim L_i + \dim \operatorname{Stab}_G(H)\right)$$
$$= \dim G + \sum_{i=1}^k (\dim G - \dim L_i) - \dim \operatorname{Stab}_G(H)$$
$$= \dim G + \operatorname{Irr}(H) - \dim \operatorname{Stab}_G(H) = \delta(H).$$

4 Meromorphic G-connections on \mathbb{P}^1 with unramified canonical forms

Let us consider meromorphic G-connections on \mathbb{P}^1 and define the invariant of them, called the index of rigidity.

4.1 Meromorphic G-connections on \mathbb{P}^1 and index of rigidity

Let $|D| := \{a_1, a_2, \ldots, a_d\}$ be a finite set of points in \mathbb{P}^1 , and $D := \sum_{a \in \{a_1, \ldots, a_d\}} (k_a + 1) \cdot a$ an effective divisor with multiplicities $k_a \in \mathbb{Z}_{\geq 0}$. Let z_a for $a \in |D|$ are coordinate functions on \mathbb{P}^1 centered at a. Let $\mathcal{O}_{\mathbb{P}^1}$ be the sheaf of regular function on \mathbb{P}^1 . Let $\mathcal{O}_{\mathbb{P}^1,D}$ and $\Omega_{\mathbb{P}^1,D}$ denote the sheaves of rational function and of rational 1-forms associated to D respectively.

Let us take a \mathfrak{g} -valued meromorphic algebraic 1-form on \mathbb{P}^1 , $A dz \in \mathfrak{g} \otimes_{\mathbb{C}} \Omega_{\mathbb{P}^1,D}(\mathbb{P}^1)$. Then the covariant exterior derivative

$$d_A := d + A \, dz$$

is a \mathbb{C} -linear morphism $\mathcal{O}_{\mathbb{P}^1}^{\mathfrak{g}} := \mathfrak{g} \otimes_{\mathbb{C}} \mathcal{O}_{\mathbb{P}^1} \to \Omega_{\mathbb{P}^1,D}^{\mathfrak{g}} := \mathfrak{g} \otimes_{\mathbb{C}} \Omega_{\mathbb{P}^1,D}$ satisfying the Leibniz rule. We call this d_A a *meromorphic G-connection* on \mathbb{P}^1 . By the projective transformation on \mathbb{P}^1 , we may assume that $|D| \subset \mathbb{C}$. Then we can write

$$A \, dz = \sum_{a \in |D|} \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^i} \frac{dz}{z-a}, \quad (A_i^{(a)} \in \mathfrak{g})$$

Here we note that the residues satisfy the relation

$$\sum_{a \in |D|} A_0^{(a)} = 0 \tag{2}$$

since A dz is regular at ∞ . Now we moreover suppose that d_A has unramified truncated canonical form $H^{(a)} \in \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})$ at each singular point $a \in |D|$. Then the collection $\mathbf{H} = (H^{(a)})_{a \in |D|}$ of canonical forms should satisfy the following relation which is an analogue of the Fuchs relation.

Proposition 4.1. Let us take a meromorphic G-connection

$$d + \sum_{a \in |D|} \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^i} \frac{dz}{z-a}$$

on \mathbb{P}^1 as above. Let us suppose that there exists a collection of canonical forms $\mathbf{H} = (H^{(a)})_{a \in |D|} \in \prod_{a \in |D|} \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})$ such that

$$\sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}} \in \mathbb{O}_{H^{(a)}}, \quad a \in |D|.$$

Then we have

$$\sum_{a \in |D|} H_{\rm res}^{(a)} \in \mathfrak{g}_{\rm ss}$$

Here $\mathfrak{g}_{ss} := [\mathfrak{g}, \mathfrak{g}]$ is the semisimple part of the reductive Lie algebra \mathfrak{g} .

Proof. Firstly, note that computations in this proof are carried out in $\operatorname{GL}_N(\mathbb{C}[z_a]_{k_a})$, $M_N(\mathbb{C}[z_a]_{k_a}) = \mathfrak{gl}_N(\mathbb{C}[z_a]_{k_a})$, and $M_N(\mathbb{C}[z_a^{-1}]_{k_a}) = \mathfrak{gl}_N(\mathbb{C}[z_a^{-1}]_{k_a})$ through the embedding $G \hookrightarrow \operatorname{GL}_N$.

Since $\sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}} \in \mathbb{O}_{H^{(a)}}$, there exists $g_a \in G(\mathbb{C}[z_a]_{k_a})$ such that $\operatorname{Ad}^*(g_a)(H^{(a)}) = \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}}$ for each $a \in |D|$.

Let \mathfrak{z} be the center of \mathfrak{g} . Let $\mathfrak{g}_{ss}^{\perp}$ be the orthogonal complement of \mathfrak{g}_{ss} with respect to the trace pairing. Then $\mathfrak{g}_{ss}^{\perp} = \mathfrak{z}$. Indeed, for $X \in \mathfrak{g}_{ss}^{\perp}$ and $Y_1, Y_2 \in \mathfrak{g}$, we have

$$tr([X, Y_1] \cdot Y_2) = tr(X \cdot [Y_1, Y_2]) = 0$$

since $[Y_1, Y_2] \in \mathfrak{g}_{ss} = [\mathfrak{g}, \mathfrak{g}]$. Thus since the trace pairing is non-degenerate for \mathfrak{g} , we have $[X, Y_1] = 0$, i.e., $\mathfrak{g}_{ss}^{\perp} \subset \mathfrak{z}$. Conversely, recalling that $\mathfrak{g} = \mathfrak{g}_{ss}^{\perp} \oplus \mathfrak{g}_{ss} \subset \mathfrak{z} \oplus \mathfrak{g}_{ss} = \mathfrak{g}$, we obtain $\mathfrak{g}_{ss}^{\perp} = \mathfrak{z}$. Then for $X \in \mathfrak{g}_{ss}^{\perp} = \mathfrak{z}$, we have

$$0 = \operatorname{tr}(X \cdot 0) = \operatorname{tr}\left(X \cdot \sum_{a \in |D|} A_0^{(a)}\right) = \operatorname{tr}\left(X \cdot \sum_{a \in |D|} \operatorname{res}_{z=a}\left(\sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}}\right)\right)$$
$$= \operatorname{tr}\left(X \cdot \sum_{a \in |D|} \operatorname{res}_{z=a}(\operatorname{Ad}(g_a)(H^{(a)}))\right) = \sum_{a \in |D|} \operatorname{res}_{z=a}\operatorname{tr}(X \cdot \operatorname{Ad}(g_a)(H^{(a)}))$$
$$= \sum_{a \in |D|} \operatorname{res}_{z=a}\operatorname{tr}(\operatorname{Ad}(g_a^{-1})(X) \cdot H^{(a)}) = \sum_{a \in |D|} \operatorname{res}_{z=a}\operatorname{tr}(X \cdot H^{(a)}) = \operatorname{tr}\left(X \cdot \sum_{a \in |D|} \operatorname{res}_{z=a}(H^{(a)})\right).$$

This equation implies that $\sum_{a \in |D|} \operatorname{res}_{z=a}(H^{(a)}) \in \mathfrak{z}^{\perp} = \mathfrak{g}_{ss}$ as desired.

Definition 4.2 (Index of rigidity of **H**). Let $\mathbf{H} = (H^{(a)})_{a \in |D|}$ be a collection of canonical forms satisfying $\sum_{a \in |D|} \underset{z=a}{\operatorname{res}} (H^{(a)}) \in \mathfrak{g}_{ss}$. The integer defined by

$$\operatorname{rig}(\mathbf{H}) := 2 \dim G - \sum_{a \in |D|} \delta(H^{(a)})$$

is called the *index of rigidity of* \mathbf{H} .

Here we note that rig (\mathbf{H}) depends only on the spectral type sp (\mathbf{H}) of \mathbf{H} . Thus we can define the index of rigidity rig (\mathbf{S}) for a collection \mathbf{S} of abstract spectral types as well.

We can also associate this quantity to meromorphic G-connections on \mathbb{P}^1 .

Definition 4.3 (Truncated index of rigidity). Let $\mathbf{H} = (H^{(a)})_{a \in |D|}$ be a collection of canonical forms as above. Let $d_A = d + A dz$, $A dz \in \Omega^{\mathfrak{g}}_{\mathbb{P}^1,D}(\mathbb{P}^1)$, be a meromorphic *G*-connection on \mathbb{P}^1 with the truncated canonical form $H^{(a)}$ at each $a \in |D|$. Then the truncated index of rigidity of d_A is defined by

$$\operatorname{rig}^{\operatorname{tr}}(d_A) := \operatorname{rig}(\mathbf{H}).$$

This is an analogue of the index of rigidity originally defined by Katz in [24] for local systems on $\mathbb{P}^1 \setminus \{n\text{-points}\}$. Under the Riemann-Hilbert correspondence, one can also define the index of rigidity for meromorphic connections, see Arinkin's paper [2] and its references for instance, and also see the paper [23] by Jakob and Yun for *G*-connections.

4.2 Example I: Heun equation and its confluent equations

In this and the next subsections, we restrict our interest to the case $G = GL_n$ and consider some explicit differential equations on \mathbb{P}^1 , and then see their canonical forms and spectral types and indices of rigidity.

The first example is the Heun-type equation

$$\frac{dY}{dz} = \left(\frac{A_0^{(0)}}{z} + \frac{A_0^{(1)}}{z-1} + \frac{A_0^{(t)}}{z-t}\right)Y, \quad (A_0^{(a)} \in \mathfrak{gl}_2 = M_2(\mathbb{C}), \ a \in \{0, 1, t\})$$

This is an analogue of the usual Heun differential equation which is a 2nd order scalar differential equation with 4 regular singular points, $z = 0, 1, t, \infty$. We regard this differential equation as a GL₂-connection on \mathbb{P}^1 and see its canonical forms at singular points. For simplicity we may assume that eigen-values of each residue matrices $A_0^{(0)}$, $A_0^{(1)}$, $A_0^{(t)}$, $A_0^{(\infty)} := -(A_0^{(0)} + A_0^{(1)} + A_0^{(t)})$ does not differ by integers. Then the canonical forms at these singular points are of the forms

$$H^{(a)} = \begin{pmatrix} \theta_1^{(a)} & \\ & \theta_2^{(a)} \end{pmatrix} z_a^{-1} \quad (a \in \{0, 1, t, \infty\})$$

where $\theta_1^{(a)}$ and $\theta_2^{(a)}$ are distinct eigen-values of A_a .

Similarly we consider the confluent Heun-type equation

$$\frac{dY}{dz} = \left(\frac{A_0^{(0)}}{z} + \frac{A_0^{(1)}}{z-1} - A_1^{(\infty)}\right)Y$$

with the canonical forms

$$H^{(a)} = \begin{pmatrix} \theta_1^{(a)} & \\ & \theta_2^{(a)} \end{pmatrix} z_a^{-1} \quad (a \in \{0, 1\}),$$
$$H^{(\infty)} = \begin{pmatrix} \alpha_1^{(\infty)} & \\ & \alpha_2^{(\infty)} \end{pmatrix} \zeta^{-2} + \begin{pmatrix} \theta_1^{(\infty)} & \\ & \theta_2^{(\infty)} \end{pmatrix} \zeta^{-1}$$

where $\zeta := 1/z$ and we assume $\theta_1^{(a)} - \theta_2^{(a)} \notin \mathbb{Z}$ for $a \in \{0, 1\}$ and $\alpha_1^{(\infty)} \neq \alpha_2^{(\infty)}$. Also consider the doubly-confluent Heun-type equation

$$\frac{dY}{dz} = \left(\frac{A_1^{(0)}}{z^2} + \frac{A_0^{(0)}}{z} - A_1^{(\infty)}\right)Y$$

with the canonical forms

$$H^{(0)} = \begin{pmatrix} \alpha_1^{(0)} & \\ & \alpha_2^{(0)} \end{pmatrix} z^{-2} + \begin{pmatrix} \theta_1^{(0)} & \\ & \theta_2^{(0)} \end{pmatrix} z^{-1},$$
$$H^{(\infty)} = \begin{pmatrix} \alpha_1^{(\infty)} & \\ & \alpha_2^{(\infty)} \end{pmatrix} \zeta^{-2} + \begin{pmatrix} \theta_1^{(\infty)} & \\ & \theta_2^{(\infty)} \end{pmatrix} \zeta^{-1}$$

where $\alpha_1^{(a)} \neq \alpha_2^{(a)}$ for $a = 0, \infty$. Further consider the biconfluent Heun-type equation

$$\frac{dY}{dz} = \left(\frac{A_0^{(0)}}{z} - A_1^{(\infty)} - A_2^{(\infty)}z\right)Y$$

with the canonical forms

$$H^{(0)} = \begin{pmatrix} \theta_1^{(0)} & \\ & \theta_2^{(0)} \end{pmatrix} z^{-1},$$

$$H^{(\infty)} = \begin{pmatrix} \alpha_1^{(\infty)} & \\ & \alpha_2^{(\infty)} \end{pmatrix} \zeta^{-3} + \begin{pmatrix} \beta_1^{(\infty)} & \\ & \beta_2^{(\infty)} \end{pmatrix} \zeta^{-2} + \begin{pmatrix} \theta_1^{(\infty)} & \\ & \theta_2^{(\infty)} \end{pmatrix} \zeta^{-1}$$

where we assume $\theta_1^{(0)} - \theta_2^{(0)} \notin \mathbb{Z}$ and $\alpha_1^{(\infty)} \neq \alpha_2^{(\infty)}$. Finally consider the triconfluent Heun-type equation

$$\frac{dY}{dz} = \left(-A_3^{(\infty)}z^2 - A_2^{(\infty)}z - A_1^{(\infty)}\right)Y$$

with the canonical form

$$H^{(\infty)} = \begin{pmatrix} \alpha_1^{(\infty)} & \\ & \alpha_2^{(\infty)} \end{pmatrix} \zeta^{-4} + \begin{pmatrix} \beta_1^{(\infty)} & \\ & \beta_2^{(\infty)} \end{pmatrix} \zeta^{-3} + \begin{pmatrix} \gamma_1^{(\infty)} & \\ & \gamma_2^{(\infty)} \end{pmatrix} \zeta^{-2} + \begin{pmatrix} \theta_1^{(\infty)} & \\ & \theta_2^{(\infty)} \end{pmatrix} \zeta^{-1}$$

where we assume $\alpha_1^{(\infty)} \neq \alpha_2^{(\infty)}$.

Let us consider the spectral types and indices of rigidity of these canonical forms. Since we are in the case $G = GL_2$, as a set of simple roots Π we can take $\Pi = \{e_{12} := e_1 - e_2\}$ where e_i are *i*-th projection of $\mathbb{C}^2 \cong \mathfrak{t}$. Then for the Heun-type equation,

$$sp(H^{(a)}) = (\emptyset; [0]) \quad (a \in \{0, 1, t, \infty\}),$$

for the confluent Heun-type equation,

$$\operatorname{sp}(H^{(a)}) = (\emptyset; [0]) \ (a \in \{0, 1\}), \ \operatorname{sp}(H^{(\infty)}) = (\emptyset \supset \emptyset; [0]),$$

for the doubly-confluent Heun-type equation,

$$\operatorname{sp}(H^{(0)}) = (\emptyset \supset \emptyset; [0]), \quad \operatorname{sp}(H^{(\infty)}) = (\emptyset \supset \emptyset; [0]),$$

for the biconfluent Heun-type equation,

$$\operatorname{sp}(H^{(0)}) = (\emptyset; [0]), \quad \operatorname{sp}(H^{(\infty)}) = (\emptyset \supset \emptyset \supset \emptyset; [0]),$$

and finally for the triconfluent Heun-type equation,

$$\operatorname{sp}(H^{(\infty)}) = (\emptyset \supset \emptyset \supset \emptyset \supset \emptyset; [0]).$$

It can be checked that all these cases have the same index of rigidity

$$\operatorname{rig}\left(\mathbf{H}\right)=0.$$

4.3 Example II

The Heun-type equations in the previous subsection have the spectral types of quite simple forms since they are GL_2 -connections, i.e., the set of simple root Π is the singleton set. Here we shall consider the following little more complicated example

$$\frac{dY}{dz} = \left(\frac{A_0^{(0)}}{z} + \frac{A_0^{(1)}}{z-1} + \frac{A_0^{(t)}}{z-t}\right)Y, \quad (A_0^{(a)} \in \mathfrak{gl}_4 = M_4(\mathbb{C}), \ a \in \{0, 1, t\})$$

with the truncated canonical forms,

$$H^{(a)} = \begin{pmatrix} \theta_1^{(a)} I_2 & \\ & \theta_2^{(a)} I_2 \end{pmatrix} z_a^{-1} \quad (a \in \{0, 1, t\}), \qquad H^{(\infty)} = \begin{pmatrix} \theta_1^{(\infty)} I_2 & & \\ & \theta_2^{(\infty)} & \\ & & \theta_3^{(\infty)} \end{pmatrix} \zeta^{-1}$$

where I_2 is the identity matrix of size 2 and we assume $\theta_i^{(a)} \neq \theta_j^{(a)}$ for $i \neq j$ and $a \in \{0, 1, t, \infty\}$. This differential equation appears in [7] by Boalch and [40] by Sakai to construct a higher dimensional analogue of the Painlevé VI equation, called the matrix Painlevé VI system in [25] by Kawakami-Nakamura-Sakai. Furthermore in the same paper Kawakami-Nakamura-Sakai classified confluent type equations with unramified irregular singularities arising from the above Fuchsian differential equation. Now we recall these confluent type equations.

The confluent equation of type I:

$$\frac{dY}{dz} = \left(\frac{A_1^{(0)}}{z^2} + \frac{A_0^{(0)}}{z} + \frac{A_0^{(1)}}{z-1}\right)Y, \quad (A_i^{(a)} \in \mathfrak{gl}_4 = M_4(\mathbb{C}), \ a \in \{0,1\})$$

with the truncated canonical forms,

$$\begin{split} H^{(0)} &= \begin{pmatrix} \alpha_1^{(0)} I_2 & \\ & \alpha_2^{(0)} I_2 \end{pmatrix} z^{-2} + \begin{pmatrix} \theta_1^{(0)} I_2 & \\ & \theta_2^{(0)} I_2 \end{pmatrix} z^{-1}, \\ H^{(1)} &= \begin{pmatrix} \theta_1^{(1)} I_2 & \\ & & \theta_2^{(1)} I_2 \end{pmatrix} (z-1)^{-1}, \quad H^{(\infty)} = \begin{pmatrix} \theta_1^{(\infty)} I_2 & \\ & & \theta_2^{(\infty)} & \\ & & & \theta_3^{(\infty)} \end{pmatrix} \zeta^{-1} \end{split}$$

where we assume $\alpha_1^{(0)} \neq \alpha_2^{(0)}$ and $\theta_i^{(a)} \neq \theta_j^{(a)}$ for $i \neq j$ and $a \in \{1, \infty\}$.

The confluent equation of type II:

$$\frac{dY}{dz} = \left(\frac{A_0^{(0)}}{z} + \frac{A_0^{(1)}}{z-1} - A_1^{(\infty)}\right)Y, \quad (A_i^{(a)} \in \mathfrak{gl}_4 = M_4(\mathbb{C}), \ a \in \{0, 1, \infty\})$$

with the truncated canonical forms,

$$\begin{aligned} H^{(a)} &= \begin{pmatrix} \theta_1^{(a)} I_2 & \\ & \theta_2^{(a)} I_2 \end{pmatrix} z_a^{-1} \quad (a \in \{0, 1\}), \\ H^{(\infty)} &= \begin{pmatrix} \alpha_1^{(\infty)} I_2 & \\ & \alpha_2^{(\infty)} I_2 \end{pmatrix} \zeta^{-2} + \begin{pmatrix} \theta_1^{(\infty)} I_2 & \\ & \theta_2^{(\infty)} & \\ & & \theta_3^{(\infty)} \end{pmatrix} \zeta^{-1} \end{aligned}$$

where we assume $\alpha_1^{(\infty)} \neq \alpha_2^{(\infty)}$, $\theta_1^{(a)} \neq \theta_2^{(a)}$ for $a \in \{0, 1\}$, and $\theta_2^{(\infty)} \neq \theta_3^{(\infty)}$. The doubly-confluent equation:

$$\frac{dY}{dz} = \left(\frac{A_1^{(0)}}{z^2} + \frac{A_0^{(0)}}{z} - A_1^{(\infty)}\right)Y, \quad (A_i^{(a)} \in \mathfrak{gl}_4 = M_4(\mathbb{C}), \ a \in \{0, \infty\})$$

with the truncated canonical forms,

$$H^{(0)} = \begin{pmatrix} \alpha_1^{(0)} I_2 & \\ & \alpha_2^{(0)} I_2 \end{pmatrix} z^{-2} + \begin{pmatrix} \theta_1^{(0)} I_2 & \\ & \theta_2^{(0)} I_2 \end{pmatrix} z^{-1},$$
$$H^{(\infty)} = \begin{pmatrix} \alpha_1^{(\infty)} I_2 & \\ & \alpha_2^{(\infty)} I_2 \end{pmatrix} \zeta^{-2} + \begin{pmatrix} \theta_1^{(\infty)} I_2 & \\ & \theta_2^{(\infty)} & \\ & & \theta_3^{(\infty)} \end{pmatrix} \zeta^{-1}$$

where we assume $\alpha_1^{(a)} \neq \alpha_2^{(a)}$, for $a \in \{0, \infty\}$ and $\theta_2^{(\infty)} \neq \theta_3^{(\infty)}$. The biconfluent equation of type I:

$$\frac{dY}{dz} = \left(\frac{A_2^{(0)}}{z^3} + \frac{A_1^{(0)}}{z^2} + \frac{A_0^{(0)}}{z}\right)Y, \quad (A_i^{(a)} \in \mathfrak{gl}_4 = M_4(\mathbb{C}), \ a \in \{0, 1\})$$

with the truncated canonical forms,

$$\begin{split} H^{(0)} &= \begin{pmatrix} \alpha_1^{(0)} I_2 & \\ & \alpha_2^{(0)} I_2 \end{pmatrix} z^{-3} + \begin{pmatrix} \beta_1^{(0)} I_2 & \\ & \beta_2^{(0)} I_2 \end{pmatrix} z^{-2} + \begin{pmatrix} \theta_1^{(0)} I_2 & \\ & \theta_2^{(0)} I_2 \end{pmatrix} z^{-1}, \\ H^{(\infty)} &= \begin{pmatrix} \theta_1^{(\infty)} I_2 & \\ & \theta_2^{(\infty)} \\ & & \theta_3^{(\infty)} \end{pmatrix} \zeta^{-1} \end{split}$$

where we assume $\alpha_1^{(0)} \neq \alpha_2^{(0)}$ and $\theta_i^{(\infty)} \neq \theta_j^{(\infty)}$ for $i \neq j$. The biconfluent equation of type II:

$$\frac{dY}{dz} = \left(\frac{A_0^{(0)}}{z} - A_2^{(\infty)}z - A_1^{(\infty)}\right)Y, \quad (A_i^{(a)} \in \mathfrak{gl}_4 = M_4(\mathbb{C}), \ a \in \{0, \infty\})$$

with the truncated canonical forms,

$$H^{(0)} = \begin{pmatrix} \theta_1^{(0)} I_2 & \\ & \theta_2^{(0)} I_2 \end{pmatrix} z^{-1},$$

$$H^{(\infty)} = \begin{pmatrix} \alpha_1^{(\infty)} I_2 & \\ & \alpha_2^{(\infty)} I_2 \end{pmatrix} \zeta^{-2} + \begin{pmatrix} \beta_1^{(\infty)} I_2 & \\ & \beta_2^{(\infty)} I_2 \end{pmatrix} \zeta^{-2} + \begin{pmatrix} \theta_1^{(\infty)} I_2 & \\ & \theta_2^{(\infty)} & \\ & & \theta_3^{(\infty)} \end{pmatrix} \zeta^{-1}$$

where we assume $\alpha_1^{(\infty)} \neq \alpha_2^{(\infty)}$, $\theta_1^{(0)} \neq \theta_2^{(0)}$ and $\theta_2^{(\infty)} \neq \theta_3^{(\infty)}$. The triconfluent equation:

$$\frac{dY}{dz} = \left(-A_3^{(\infty)}z^2 - A_2^{(\infty)}z - A_1^{(\infty)}\right)Y, \quad (A_i^{(a)} \in \mathfrak{gl}_4 = M_4(\mathbb{C}), \ a \in \{0, \infty\})$$

with the truncated canonical form,

$$H^{(\infty)} = \begin{pmatrix} \alpha_1^{(\infty)} I_2 \\ \alpha_2^{(\infty)} I_2 \end{pmatrix} \zeta^{-3} + \begin{pmatrix} \beta_1^{(\infty)} I_2 \\ \beta_2^{(\infty)} I_2 \end{pmatrix} \zeta^{-2} \\ + \begin{pmatrix} \gamma_1^{(\infty)} I_2 \\ \gamma_2^{(\infty)} I_2 \end{pmatrix} \zeta^{-2} + \begin{pmatrix} \theta_1^{(\infty)} I_2 \\ \theta_2^{(\infty)} \\ \theta_3^{(\infty)} \end{pmatrix} \zeta^{-1}$$

where we assume $\alpha_1^{(\infty)} \neq \alpha_2^{(\infty)}$ and $\theta_2^{(\infty)} \neq \theta_3^{(\infty)}$.

Let us consider the spectral types and indices of rigidity of these canonical forms. Since we are in the case $G = GL_4$, as a set of simple roots Π we can take $\Pi = \{e_{12}, e_{23}, e_{34}\}$ where $e_{ij} := e_i - e_j$ and e_i are *i*-th projection of $\mathbb{C}^4 \cong \mathfrak{t}$. Then for the first Fuchsian equation, the collection of spectral types is

$$\operatorname{sp}(H^{(a)}) = (\{e_{12}, e_{34}\}; [0]) \quad (a \in \{0, 1, t\}), \quad \operatorname{sp}(H^{(\infty)}) = (\{e_{12}\}; [0]),$$

also for the confluent equation of type I,

$$sp(H^{(0)}) = (\{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\}; [0]), \qquad sp(H^{(1)}) = (\{e_{12}, e_{34}\}; [0]),$$
$$sp(H^{(\infty)}) = (\{e_{12}\}; [0]),$$

for the confluent equation of type II,

$$sp(H^{(a)}) = (\{e_{12}, e_{34}\}; [0]) \ (a \in \{0, 1\}), \qquad sp(H^{(\infty)}) = (\{e_{12}, e_{34}\} \supset \{e_{12}\}; [0]),$$
for the doubly-confluent equation,

 $\operatorname{sp}(H^{(0)}) = (\{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\}; [0]), \qquad \operatorname{sp}(H^{(\infty)}) = (\{e_{12}, e_{34}\} \supset \{e_{12}\}; [0]),$

for the biconfluent equation of type I,

 $sp(H^{(0)}) = (\{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\}; [0]), \qquad sp(H^{(\infty)}) = (\{e_{12}\}; [0]),$ for the biconfluent equation of type II,

 $sp(H^{(0)}) = (\{e_{12}, e_{34}\}; [0]), \qquad sp(H^{(\infty)}) = (\{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\} \supset \{e_{12}\}; [0]),$ and finally for the triconfluent equation,

$$\operatorname{sp}(H^{(\infty)}) = (\{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\} \supset \{e_{12}\}; [0]).$$

It can be checked that all these cases have the same index of rigidity

$$\operatorname{rig}\left(\mathbf{H}\right) = -2.$$

5 δ -constant deformation of canonical forms

In this section, we shall introduce a deformation of unramified canonical forms which preserves their δ -invariants.

Let us consider an unramified canonical form

$$H dz = \left(\frac{H_k}{z^k} + \dots + \frac{H_1}{z} + H_{\rm res}\right) \frac{dz}{z} \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k) dz.$$

Let $H_{\text{res}} = H_0 + J_0$ be the Jordan decomposition with the semisimple H_0 and nilpotent J_0 . We may assume H satisfies the assumptions in Section 2.2.

5.1 Complement of hypersurfaces associated with H

To each simple root $\alpha \in \Pi \setminus \Pi_0$, we associate the positive integer

$$d(\alpha) := \max\{i \in \{1, \dots, k\} \mid \alpha \notin \Pi_i\}.$$

Then it follows that $\alpha(H_{d(\alpha)}) \neq 0$, and $\alpha(H_i) = 0$ for all $d(\alpha) < j \leq k$. We also define the following polynomials

$$f_{\alpha}^{(i)}(x_0, x_1, \dots, x_k) := \sum_{j=i}^{d(\alpha)} \left(\alpha(H_j) \prod_{\nu=j+1}^{d(\alpha)} (x_i - x_{\nu}) \right)$$

for $i = 0, 1, ..., d(\alpha)$. Here we formally put $\prod_{\nu=d(\alpha)+1}^{d(\alpha)} (x_i - x_{\nu}) := 1$. Then we denote the complement of hypersurfaces defined by these polynomials $f_{\alpha}^{(i)}(x)$ by

$$\mathbb{B}_{H} := \left\{ \mathbf{c} \in \mathbb{C}^{k+1} \, \middle| \, f_{\alpha}^{(i)}(\mathbf{c}) \neq 0 \text{ for all } i = 0, 1, \dots, d(\alpha), \, \alpha \in \Pi \backslash \Pi_{0} \right\}.$$

Since $f_{\alpha}^{(i)}(0,0,\ldots,0) = \alpha(H_{d(\alpha)}) \neq 0$, we have $\mathbf{0} \in \mathbb{B}_H$, thus in particular, $\mathbb{B}_H \neq \emptyset$.

5.2 Stratification of \mathbb{C}^{k+1} associated to partitions

Let $\mathcal{I}: I_1, I_2, \ldots, I_r$ be a partition of $\{0, 1, 2, \ldots, k\}$, i.e., the direct sum decomposition

$$I_0 \sqcup I_1 \sqcup \cdots \sqcup I_r = \{0, 1, 2, \dots, k\}$$

with nonempty direct summands I_i , i = 0, 1, ..., r. We may assume that $0 \in I_0$. Let us introduce an embedding of \mathbb{C}^{r+1} into \mathbb{C}^{k+1} associated to \mathcal{I} ,

$$\iota_{\mathcal{I}} \colon \mathbb{C}^{r+1} \ni \mathbf{a} = (a_0, a_1, \dots, a_r) \mapsto (\iota_{\mathcal{I}}(\mathbf{a})_0, \iota_{\mathcal{I}}(\mathbf{a})_1, \dots, \iota_{\mathcal{I}}(\mathbf{a})_k) \in \mathbb{C}^{k+1}$$

by setting $\iota_{\mathcal{I}}(\mathbf{a})_i := a_l$ if $i \in I_l$. Let us denote the configuration space of r + 1 points in \mathbb{C} by

$$C_{r+1}(\mathbb{C}) := \{ (a_0, a_1, \dots, a_r) \in \mathbb{C}^{r+1} \mid a_i \neq a_j \text{ for } i \neq j \}.$$

We consider the subspace of \mathbb{C}^{k+1} ,

$$C(\mathcal{I}) := \iota_{\mathcal{I}}(C_{r+1}(\mathbb{C})) = \left\{ (a_0, a_1, \dots, a_k) \in \mathbb{C}^{k+1} \middle| \begin{array}{c} a_i = a_j & \text{if } i, j \in I_l \text{ for some } l, \\ a_i \neq a_j & \text{otherwise} \end{array} \right\}.$$

Then we obtain the direct sum decomposition of \mathbb{C}^{k+1} ,

$$\mathbb{C}^{k+1} = \bigsqcup_{\mathcal{I} \in \mathcal{P}_{[k+1]}} C(\mathcal{I}),$$

where $\mathcal{P}_{[k+1]}$ is the set of all partitions of $\{0, 1, \ldots, k\}$.

Definition 5.1 (\mathcal{P} -decomposed space). Let \mathcal{P} be a poset. An \mathcal{P} -decomposition of a Hausdorff paracompact topological space X is a locally finite correction $(S_i)_{i \in \mathcal{I}}$ of disjoint locally closed manifolds such that

1.
$$X = \bigsqcup_{i \in \mathcal{I}} S_i,$$

2. $S_i \cap \overline{S_j} \neq \emptyset \iff S_i \subset \overline{S_j} \iff i \leq j.$

We call the space X an \mathcal{P} -decomposed space and each S_i a piece of X.

According to the natural ordering of $\mathcal{P}_{[k+1]}$ defined by the refinement of partitions, the above decomposition defines a $\mathcal{P}_{[k+1]}$ -decomposed space structure on \mathbb{C}^{k+1} . This decomposed space structure on \mathbb{C}^{k+1} moreover defines the stratification of which we shall recall the definition in Section 7.1. Thus we call the pieces of \mathbb{C}^{k+1} the *strata* of the stratified space \mathbb{C}^{k+1} .

The lemma below assures that the open subset $\mathbb{B}_H \subset \mathbb{C}^{k+1}$ defined above intersects with every stratum $\mathcal{C}(\mathcal{I})$. Therefore the decomposition $\mathbb{B}_H = \bigsqcup_{\mathcal{I} \in \mathcal{P}_{[k+1]}} \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$ defines the stratification on \mathbb{B}_H .

Lemma 5.2. Let U be an open neighborhood of $\mathbf{0} \in \mathbb{C}^{k+1}$. Then $U \cap \mathcal{C}(\mathcal{I}) \neq \emptyset$ for any $\mathcal{I} \in \mathcal{P}_{[k+1]}$.

Proof. Let us take $\mathcal{I} \in \mathcal{P}_{[k+1]}$. Then the inclusion map $\iota_{\mathcal{I}} \colon \mathbb{C}^{r+1} \hookrightarrow \mathbb{C}^{k+1}$ defined above satisfies $\mathbf{0} \in \operatorname{Im} \iota_{\mathcal{I}}$. Thus $\iota_{\mathcal{I}}^{-1}(U) \neq \emptyset$. The configuration space $C_{r+1}(\mathbb{C})$ of r+1 points in \mathbb{C} is a Zariski open subset of the connected space \mathbb{C}^{r+1} and thus it is dense in \mathbb{C}^{r+1} . Therefore we have

$$\emptyset \neq C_{r+1}(\mathbb{C}) \cap \iota_{\mathcal{I}}^{-1}(U) = \iota_{\mathcal{I}}^{-1}(\mathcal{C}(\mathcal{I})) \cap \iota_{\mathcal{I}}^{-1}(U) = \iota_{\mathcal{I}}^{-1}(\mathcal{C}(\mathcal{I}) \cap U)$$

which shows $\mathcal{C}(\mathcal{I}) \cap U \neq \emptyset$ as desired.

5.3 Unfolding of canonical forms and decomposition of the spectral types

Definition 5.3 (Deformation of unramified canonical forms). Let H dz be the canonical form as above. Then the *unfolding* of H is the function from \mathbb{C}^{k+1} to $\mathfrak{g}(\mathbb{C}(z))$ of the form,

$$H(c_0, c_1, \dots, c_k) := \left(\frac{H_k}{(z - c_1)(z - c_2)\cdots(z - c_k)} + \frac{H_{k-1}}{(z - c_1)(z - c_2)\cdots(z - c_{k-1})} + \dots + H_{\text{res}}\right) \frac{1}{z - c_0}.$$

Let us take a partition $\mathcal{I}: I_0, I_1, \ldots, I_r \in \mathcal{P}_{[k+1]}$ and write $I_j = \{i_{[j,0]} < i_{[j,1]} < \cdots < i_{[j,k_j]}\}$. Then we consider $H(\mathbf{c})$ for $\mathbf{c} \in C(\mathcal{I})$. Since $H(\mathbf{c})$ has poles at $c_{i_{[0,0]}}, c_{i_{[1,0]}}, \ldots, c_{i_{[r,0]}}$ as the function of z, we obtain the partial fraction decomposition of $H(\mathbf{c})$,

$$H(\mathbf{c}) = \sum_{j=0}^{r} \sum_{\nu=0}^{k_j} \frac{A_{\nu}^{[j]}}{(z - c_{i_{[j,0]}})^{\nu+1}} \quad (A_{\nu}^{[j]} \in \mathfrak{g}).$$

Then obviously, each $H(\mathbf{c})_j := \sum_{\nu=0}^{k_j} \frac{A_{\nu}^{[j]}}{(z-c_{i_{[j,0]}})^{\nu+1}}$ becomes a canonical form at $z = c_{i_{[j,0]}}$ for $j = 0, 1, \ldots, r$. The coefficients $A_{\nu}^{[j]} \in \mathfrak{g}$ are linear combinations of H_i and H_{res} which might be complicated to write down explicitly. However, the spectral types of $H(\mathbf{c})_j$ are rather easily described in terms of the partition \mathcal{I} as follows.

Proposition 5.4. Let $(\Pi_k \supset \Pi_{k-1} \supset \cdots \supset \Pi; [J_0])$ be the spectral type of H. Suppose that $\mathbf{c} \in \mathbb{B}_H \cap C(\mathcal{I})$. Then we have

$$\Pi_{i_{[j,\nu]}} = \left\{ \alpha \in \Pi \, \big| \, \alpha((A_{\mu}^{[j]})_{\rm ss}) = 0 \text{ for all } \mu = \nu, \nu + 1, \dots, k_j \right\}$$

for $\nu = 0, 1, \ldots, k_j$, $j = 0, 1, \ldots, r$. Here X_{ss} denotes the semisimple part in the Jordan decomposition of $X \in \mathfrak{g}$.

Proof. Recalling that $A^{[j]}_{\mu} = \underset{z=c_{i_{[j,0]}}}{\operatorname{res}} \left((z - c_{i_{[j,0]}})^{\mu} H(\mathbf{c}) \right)$, we have

$$A_{\mu}^{[j]} = \sum_{i=i_{[j,\mu]}}^{i_{[j,\mu+1]}-1} \frac{H_i}{\prod_{\substack{0 \le l \le i \\ l \notin I_j}} (c_{i_{[j,0]}} - c_l)} + (\text{linear combination of } H_i \text{ for } i_{[j,\mu+1]} \le i \le k)$$

for $(\mu, j) \neq (0, 0)$ and

$$A_0^{[0]} = H_{\text{res}} + \sum_{i=1}^{i_{[0,1]}-1} \frac{H_i}{\prod_{\substack{0 \le l \le i \\ l \notin I_0}} (c_{i_{[0,0]}} - c_l)} + \left(\text{linear combination of } H_i \text{ for } i_{[0,1]} \le i \le k\right).$$
(3)

Here we formally put $i_{[j,k_j+1]} := k + 1$. Thus, in any cases, semisimple parts are written by

$$(A_{\mu}^{[j]})_{\rm ss} = \sum_{i=i_{[j,\mu]}}^{i_{[j,\mu+1]}-1} \frac{H_i}{\prod_{\substack{0 \le l \le i \\ l \notin I_j}} (c_{i_{[j,0]}} - c_l)} + (\text{linear combination of } H_i \text{ for } i_{[j,\mu+1]} \le i \le k)$$

for all $\mu = 0, 1, ..., k$ and j = 0, 1, ..., r.

Therefore if a simple root α satisfies $\alpha(H_i) = 0$ for all $i \ge i_{[j,\mu]}$, then $\alpha((A^{[j]}_{\mu})_{ss}) = 0$. Namely, we have the inclusion

$$\Pi_{i_{[j,\nu]}} \subset \left\{ \alpha \in \Pi \mid \alpha((A_{\mu}^{[j]})_{ss}) = 0 \text{ for all } \mu = \nu, \nu + 1, \dots, k_j \right\}$$

To obtain the converse inclusion, we take a simple root $\alpha \notin \prod_{i_{[j,\nu]}}$. Then we can find $\mu \in \{\nu, \nu + 1, \ldots, k_j\}$ such that $i_{[j,\mu]} \leq d(\alpha) < i_{[j,\mu+1]}$. Since we have $\alpha(H_i) = 0$ for all $d(\alpha) < i \leq k$, we obtain

$$\alpha((A_{\mu}^{[j]})_{ss}) = \sum_{i=i_{[j,\mu]}}^{i_{[j,\mu+1]}-1} \frac{\alpha(H_i)}{\prod_{\substack{0 \le l \le i \ l \le i}} (c_{i_{[j,0]}} - c_l)} = \sum_{i=i_{[j,\mu]}}^{d(\alpha)} \frac{\alpha(H_i)}{\prod_{\substack{0 \le l \le i \ l \le i}} (c_{i_{[j,0]}} - c_l)}$$
$$= \sum_{i=i_{[j,\mu]}}^{d(\alpha)} \frac{\alpha(H_i) \prod_{\substack{l=i+1 \ l \le i}}^{d(\alpha)} (c_{i_{[j,0]}} - c_l)}{\prod_{\substack{0 \le l \le d(\alpha) \ l \le l \le i}} (c_{i_{[j,0]}} - c_l)} = \frac{f_{\alpha}^{(i_{[j,\mu]})}(\mathbf{c})}{\prod_{\substack{0 \le l \le d(\alpha) \ l \le l \le j}} (c_{i_{[j,0]}} - c_l)} \neq 0.$$

Here we used the assumption $\mathbf{c} \in \mathbb{B}_H$. Therefore it follows that

$$\alpha \notin \left\{ \alpha \in \Pi \mid \alpha((A_{\mu}^{[j]})_{ss}) = 0 \text{ for all } \mu = k_j, k_j - 1, \dots, \nu \right\}$$

as desired.

Therefore we obtain the following decomposition of the spectral type of H. Corollary 5.5. Suppose that $\mathbf{c} \in \mathbb{B}_H \cap C(\mathcal{I})$. Then the spectral types of $H(\mathbf{c})_j$ are

$$\begin{array}{ll} (\Pi^{I_0}; [J_0]) & if \ j = 0, \\ (\Pi^{I_j}; [0]) & otherwise \end{array}$$

Here Π_{I_i} denotes the sequence

$$\Pi_{i_{[j,k_j]}} \supset \cdots \supset \Pi_{i_{[j,1]}} \supset \Pi_{i_{[j,0]}}$$

of subset of Π for each $j = 0, 1, \ldots, r$.

5.4 δ -constant deformation of canonical form

Let us show that the unfolding $H(\mathbf{c})$ of the canonical form H preserves its δ -invariant. **Proposition 5.6.** Suppose that $\mathbf{c} \in \mathbb{B}_H \cap C(\mathcal{I})$.

1. For j = 0, 1, ..., r we have

$$\begin{aligned} \operatorname{Stab}_{G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_{j}})_{1}}(H(\mathbf{c})_{j}) \\ &= \left\{ e^{X_{k_{j}} z_{c_{i_{[j,0]}}}^{k_{j}}} \cdots e^{X_{1} z_{c_{i_{[j,0]}}}} \in G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_{j}})_{1} \middle| X_{\nu} \in \mathfrak{l}_{i_{[j,\nu]}}, \ \nu = 1, \dots, k_{j} \right\}. \end{aligned}$$

2. For j = 1, 2, ..., r, we have

$$\operatorname{Stab}_G(H(\mathbf{c})_j) = L_{i_{[j,0]}}.$$

For j = 0, we also have

$$\operatorname{Stab}_G(H(\mathbf{c})_0) = \operatorname{Stab}_G(H).$$

Proof. The first assertion follows directly from Propositions 3.4 and 5.4.

For $j = 1, 2, \ldots, r$, Proposition 5.4 shows

$$\left\{ \alpha \in \Pi \, \middle| \, \alpha(A_{\mu}^{[j]}) = 0 \text{ for all } \mu = 0, 1, \dots, k_j \right\} = \Pi_{i_{[j,0]}}$$

which implies $\operatorname{Stab}_G(H(\mathbf{c})_j) = L_{i_{[j,0]}}$.

For j = 0, let us recall the equation (3). Then it follows that the stabilizer of $A_0^{[0]} - J_0$ in $L_{i_{[0,1]}}$ is $L_1 \subset L_{i_{[0,1]}}$. Here we recall that $H_{res} = J_0 + H_0$ is the Jordan decomposition. This implies that $\operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]}) = \operatorname{Stab}_{L_1}(H_{res})$ of which we postpone to give a proof. Then we obtain

$$\operatorname{Stab}_{G}(H(\mathbf{c})_{0}) = \operatorname{Stab}_{G}((H(\mathbf{c})_{0})_{\operatorname{res}}) \cap \operatorname{Stab}_{G}((H(\mathbf{c})_{0})_{\operatorname{irr}})$$
$$= \operatorname{Stab}_{G}(A_{0}^{[0]}) \cap L_{i_{[0,1]}} = \operatorname{Stab}_{L_{1}}(H_{\operatorname{res}}) = \operatorname{Stab}_{G}(H),$$

which shows the second assertion for j = 0.

Thus we finally show the equation $\operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]}) = \operatorname{Stab}_{L_1}(H_{\operatorname{res}})$. Firstly, we can obtain the inclusion $\operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]}) \subset \operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]} - H_{\operatorname{res}})$ as follows. The equation (3) shows that the nilpotent part of the Jordan decomposition of $A_0^{[0]}$ equals J_0 . Thus for $l \in \operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]})$, we obtain

$$Ad(l)(A_0^{[0]} - H_{res}) = Ad(l)(A_0^{[0]} - J_0 - H_0)$$

= Ad(l)(A_0^{[0]} - J_0) - Ad(l)(H_0)
= A_0^{[0]} - J_0 - H_0 = A_0^{[0]} - H_{res}.

Here the third equation follows from the uniqueness of the Jordan decomposition and the fact that H_0 is stabilized by $L_{i_{[0,1]}}$. Thus we have the desired inclusion $\operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]}) \subset \operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]} - H_{\operatorname{res}}).$

On the other hand, the equation (3) tells us that $\operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]} - H_{\operatorname{res}}) = L_1$. Thus obviously we have $\operatorname{Stab}_{L_1}(H_{\operatorname{res}}) \subset \operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]})$. Moreover the relation $\operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]}) \subset \operatorname{Stab}_{L_{i_{[0,1]}}}(A_0^{[0]} - H_{\operatorname{res}})$ gives the converse inclusion.

Corollary 5.7. Let us take $\mathcal{I}: I_0, I_1, \ldots, I_r \in \mathcal{P}_{[k+1]}$ and consider $H(\mathbf{c})$ for $\mathbf{c} \in C(\mathcal{I}) \cap \mathbb{B}_H$. Let $H(\mathbf{c}) = \sum_{j=0}^r H(\mathbf{c})_j$ be the partial fraction decomposition of $H(\mathbf{c})$ as above. Then we have the equation

$$\sum_{i=0}^{r} \delta(H(\mathbf{c})_j) = \delta(H).$$

Proof. By Corollary 5.5 and Proposition 5.6, we have

$$\sum_{i=0}^{r} \delta(H(\mathbf{c})_{j}) = \sum_{i=1}^{r} (\dim G + \sum_{j=1}^{k_{i}} (\dim G - L_{i_{[i,j]}}) - \dim L_{i_{[i,0]}}) + (\dim G + \sum_{j=1}^{k_{0}} (\dim G - L_{i_{[0,j]}}) - \dim \operatorname{Stab}_{G}(H)) = \dim G - \sum_{l=1}^{k} (\dim G - \dim L_{l}) - \operatorname{Stab}_{G}(H) = \dim G - \operatorname{Irr}(H) - \operatorname{Stab}_{G}(H) = \delta(H).$$

5.5 rig-constant deformation of a family of canonical forms

As well as in Section 4, let us take an effective divisor $D = \sum_{a \in \{a_1, \dots, a_d\}} (k_a + 1) \cdot a$ in \mathbb{C} and a collection of canonical forms $\mathbf{H} = (H^{(a)})_{a \in |D|} \in \prod_{a \in |D|} \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})$ with the relation $\sum_{a \in |D|} H^{(a)}_{\text{res}} \in \mathfrak{g}_{\text{ss}}$. Then we can consider the unfolding

$$\mathbf{H}(\mathbf{c}) := (H^{(a)}(\mathbf{c}_a))_{\mathbf{c} = (\mathbf{c}_a)_{a \in |D|} \in \prod_{a \in |D|} \mathbb{C}^{k_a + 1}}$$

of the collection **H**.

Recall that each \mathbb{C}^{k_a+1} is equipped with the stratification $\mathbb{C}^{k_a+1} = \bigsqcup_{\mathcal{I}^{(a)} \in \mathcal{P}_{k_a+1}} C(\mathcal{I}^{(a)})$, and thus the product $\prod_{a \in |D|} \mathbb{C}^{k_a+1}$ also has the stratification with the strata $\prod_{a \in |D|} C(\mathcal{I}^{(a)})$ for $(\mathcal{I}^{(a)}) \in \prod_{a \in |D|} \mathcal{P}_{k_a+1}$.

Let us take a stratum $\prod_{a \in |D|} C(\mathcal{I}^{(a)})$ of $\prod_{a \in |D|} \mathbb{C}^{k_a+1}$ associated with the collection of partitions $\mathcal{I}^{(a)} \colon I_0^{(a)}, \ldots, I_{r_a}^{(a)} \in \mathcal{P}_{k_a+1}$ for $a \in |D|$. Then for $\mathbf{c} = (\mathbf{c}_a)_{a \in |D|} \in \prod_{a \in |D|} C(\mathcal{I}_a)$, we can write $H^{(a)}(\mathbf{c}_a) = \sum_{i=0}^{r_a} H^{(a)}(\mathbf{c}_a)_i$ according to the partial fraction decomposition and regard $\mathbf{H}(\mathbf{c})$ as the collection of canonical forms

$$\mathbf{H}(\mathbf{c}) = (H^{(a)}(\mathbf{c}_a)_i)_{\substack{i=0,1,\dots,r_a\\a\in |D|}}.$$

Lemma 5.8. Let $(\mathcal{I}^{(a)})_{a\in|D|} \in \prod_{a\in|D|} \mathcal{P}_{[k_{a}+1]}$ be a collection of partitions. We write $\mathcal{I}^{(a)}: I_{0}^{(a)}, \ldots, I_{r_{a}}^{(a)}$ and $I_{j}^{(a)} = \{i_{[j,0]}^{(a)}, \ldots, i_{[j,k_{j}^{(a)}]}^{(a)}\}$. Let us take $\mathbf{c} = (\mathbf{c}_{a})_{a\in|D|} \in \prod_{a\in|D|} \mathcal{C}(\mathcal{I}^{(a)})$ and write $\mathbf{c}^{(a)} = (c_{0}^{(a)}, \ldots, c_{k_{a}}^{(a)})$.

Then the collection $\mathbf{H}(\mathbf{c}) = (H^{(a)}(\mathbf{c}_a)_i)_{i=0,1,\dots,r_a}$ of canonical forms satisfies the residue condition

$$\sum_{a \in |D|} \sum_{j=0}^{r_a} \operatorname{res}_{z_a - c_{i_{[j,0]}}^{(a)}} \left(H^{(a)}(\mathbf{c}_a)_i \right) \in \mathfrak{g}_{\mathrm{ss}}.$$
Proof. Lemma 9.3 in the latter section tells us that

$$\sum_{a \in |D|} \sum_{j=0}^{r_a} \operatorname{res}_{z_a = c_{i_{[j,0]}}^{(a)}} H^{(a)}(\mathbf{c}_a)_i = \sum_{a \in |D|} \operatorname{res}_{z=a}(H^{(a)}) \in \mathfrak{g}_{ss}.$$

Now we can consider the index of rigidity of $\mathbf{H}(\mathbf{c})$,

$$\operatorname{rig}(\mathbf{H}(\mathbf{c})) = 2 \dim G - \sum_{a \in |D|} \sum_{i=0}^{r_a} \delta(H^{(a)}(\mathbf{c}_a)_i).$$

For each $a \in |D|$, let $\mathbb{B}_{H^{(a)}}$ be the complement of hypersurfaces associated with the canonical form $H^{(a)}$ defined in Section 5.1. Then we consider the product of them,

$$\mathbb{B}_{\mathbf{H}} := \prod_{a \in |D|} \mathbb{B}_{H^{(a)}}.$$

Corollary 5.9. For any $\mathbf{c} \in \prod_{a \in |D|} \mathbb{B}_{H^{(a)}}$, we have

$$\operatorname{rig}\left(\mathbf{H}(\mathbf{c})\right) = \operatorname{rig}\left(\mathbf{H}\right).$$

Proof. This is a direct consequence of Corollary 5.7.

5.6 Examples

Let us consider the canonical form

$$H = \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix} z^{-4} + \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} z^{-3} + \begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix} z^{-2} + \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix} z^{-1} \in \mathfrak{gl}_2(\mathbb{C}[z^{-1}]_3) \, dz$$

with $\alpha_1 \neq \alpha_2$ which is the canonical form of the triconfluent Heun-type equation in Section 4.2, and also consider its unfolding

$$H(\mathbf{c}) = \frac{\begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}}{(z - c_0)(z - c_1)(z - c_2)(z - c_3)} + \frac{\begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}}{(z - c_0)(z - c_1)(z - c_2)} + \frac{\begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix}}{(z - c_0)(z - c_1)} + \frac{\begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}}{z - c_0}.$$

On the original setting in Section 4.2, H is the canonical form at $z = \infty$. Here we however consider at z = 0 under some projective transformation of \mathbb{P}^1 for simplicity. Let us see the partial fraction decomposition of $H(\mathbf{c})$ on each stratum of $\mathbb{B}_H \subset \mathbb{C}^4$.

Let us take the trivial partition $\mathcal{I}: \{0, 1, 2, 3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

$$H(\mathbf{c}) = \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix} z_{c_0}^{-4} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} z_{c_0}^{-3} + \begin{pmatrix} \gamma_1 \\ \gamma_2 \end{pmatrix} z_{c_0}^{-2} + \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix} z_{c_0}^{-1},$$

which is the canonical form of the triconfluent Heun-type equation.

Let us consider the partition $\mathcal{I}: \{0\} \sqcup \{1, 2, 3\}$. Then Proposition 5.4 and its proof imply that for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$, we have

$$H(\mathbf{c}) = \begin{pmatrix} \theta_1^{(c_0)}(\mathbf{c}) & \\ & \theta_2^{(c_0)}(\mathbf{c}) \end{pmatrix} z_{c_0}^{-1} + \\ \begin{pmatrix} \alpha_1^{(c_1)}(\mathbf{c}) & \\ & \alpha_2^{(c_1)}(\mathbf{c}) \end{pmatrix} z_{c_1}^{-3} + \begin{pmatrix} \beta_1^{(c_1)}(\mathbf{c}) & \\ & \beta_2^{(c_1)}(\mathbf{c}) \end{pmatrix} z_{c_1}^{-2} + \begin{pmatrix} \theta_{(c_1)}^{(1)}(\mathbf{c}) & \\ & \theta_2^{(c_1)}(\mathbf{c}) \end{pmatrix} z_{c_1}^{-1},$$

with some $\alpha_j^{(c_i)}(\mathbf{c}), \beta_j^{(c_i)}(\mathbf{c}), \theta_j^{(c_i)}(\mathbf{c}) \in \mathbb{C}$ satisfying $\theta_1^{(c_0)}(\mathbf{c}) \neq \theta_2^{(c_0)}(\mathbf{c})$ and $\alpha_1^{(c_1)}(\mathbf{c}) \neq \alpha_2^{(c_1)}(\mathbf{c})$. The canonical forms in the right hand side are those of the biconfluent Heuntype equation. We obtain the same canonical forms for the partitions $\{1\} \sqcup \{0, 2, 3\}, \{2\} \sqcup \{0, 1, 3\}, \text{ and } \{3\} \sqcup \{0, 1, 2\}$ as well.

Let us consider the partition $\mathcal{I}: \{0, 1\} \sqcup \{2, 3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

$$H(\mathbf{c}) = \begin{pmatrix} \alpha_1^{(c_0)}(\mathbf{c}) & & \\ & \alpha_2^{(c_0)}(\mathbf{c}) \end{pmatrix} z_{c_0}^{-2} + \begin{pmatrix} \beta_{(c_0)}^{(1)}(\mathbf{c}) & & \\ & \beta_2^{(c_0)}(\mathbf{c}) \end{pmatrix} z_{c_0}^{-1} + \\ \begin{pmatrix} \alpha_1^{(c_2)}(\mathbf{c}) & & \\ & \alpha_2^{(c_2)}(\mathbf{c}) \end{pmatrix} z_{c_2}^{-2} + \begin{pmatrix} \beta_{(c_2)}^{(1)}(\mathbf{c}) & & \\ & \beta_2^{(c_2)}(\mathbf{c}) \end{pmatrix} z_{c_2}^{-1},$$

with $\alpha_1^{(c_i)}(\mathbf{c}) \neq \alpha_2^{(c_i)}(\mathbf{c})$ for i = 0, 2. The canonical forms in the right hand side are those of the doubly-confluent Heun-type equation. We can obtain the same canonical forms for the partitions $\{0, 2\} \sqcup \{1, 3\}$, and $\{0, 3\} \sqcup \{1, 2\}$ as well.

Let us consider the partition $\mathcal{I}: \{0\} \sqcup \{1\} \sqcup \{2,3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

$$H(\mathbf{c}) = \begin{pmatrix} \theta_1^{(c_0)}(\mathbf{c}) & \\ & \theta_2^{(c_0)}(\mathbf{c}) \end{pmatrix} z_{c_0}^{-1} + \\ \begin{pmatrix} \theta_1^{(c_1)}(\mathbf{c}) & \\ & & \theta_2^{(c_1)}(\mathbf{c}) \end{pmatrix} z_{c_1}^{-1} + \\ \begin{pmatrix} \alpha_1^{(c_2)}(\mathbf{c}) & \\ & & & \alpha_2^{(c_2)}(\mathbf{c}) \end{pmatrix} z_{c_2}^{-2} + \begin{pmatrix} \theta_{(c_2)}^{(1)}(\mathbf{c}) & \\ & & & \theta_2^{(c_2)}(\mathbf{c}) \end{pmatrix} z_{c_2}^{-1},$$

with $\theta_1^{(c_i)}(\mathbf{c}) \neq \theta_2^{(c_i)}(\mathbf{c})$ for i = 0, 1 and $\alpha_1^{(c_2)}(\mathbf{c}) \neq \alpha_2^{(c_2)}(\mathbf{c})$. The canonical forms in the right are those of the confluent Heun-type equation. We can obtain the same canonical forms for the partitions $\{0\} \sqcup \{2\} \sqcup \{1,3\}, \{0\} \sqcup \{3\} \sqcup \{1,2\}, \{0,1\} \sqcup \{2\} \sqcup \{3\}, \{0,2\} \sqcup \{1\} \sqcup \{3\}, \text{ and } \{0,3\} \sqcup \{1\} \sqcup \{2\} \text{ as well.}$

Finally consider the finest partition $\mathcal{I}: \{0\} \sqcup \{1\} \sqcup \{2\} \sqcup \{3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

$$H(\mathbf{c}) = \sum_{i=0}^{3} \begin{pmatrix} \theta_1^{(c_i)}(\mathbf{c}) & \\ & \theta_2^{(c_i)}(\mathbf{c}) \end{pmatrix} z_{c_i}^{-1},$$

with $\theta_1^{(c_i)}(\mathbf{c}) \neq \theta_2^{(c_i)}(\mathbf{c})$ for i = 0, 1, 2, 3. The canonical forms in the right hand side are those of the Heun-type equation. Therefore all canonical forms for confluent Heun-type equations in Section 4.2 can be obtained from the unfolding of the canonical form of the triconfluent Heun-type equation.

Let us see another example,

$$\begin{split} H &= \begin{pmatrix} \alpha_1 I_2 \\ \alpha_2 I_2 \end{pmatrix} z^{-4} + \begin{pmatrix} \beta_1 I_2 \\ \beta_2 I_2 \end{pmatrix} z^{-3} \\ &+ \begin{pmatrix} \gamma_1 I_2 \\ \gamma_2 I_2 \end{pmatrix} z^{-2} + \begin{pmatrix} \theta_1 I_2 \\ \theta_2 \\ \theta_3 \end{pmatrix} z^{-1} \in \mathfrak{gl}_4(\mathbb{C}[z^{-1}]_3) \, dz \end{split}$$

with $\alpha_1 \neq \alpha_2$ and $\theta_2 \neq \theta_3$ which is the canonical form of the triconfluent equation in Section 4.3, and also consider its unfolding

$$H(\mathbf{c}) = \frac{\begin{pmatrix} \alpha_1 I_2 \\ \alpha_2 I_2 \end{pmatrix}}{(z - c_0)(z - c_1)(z - c_2)(z - c_3)} + \frac{\begin{pmatrix} \beta_1 I_2 \\ \beta_2 I_2 \end{pmatrix}}{(z - c_0)(z - c_1)(z - c_2)} + \frac{\begin{pmatrix} \gamma_1 I_2 \\ \gamma_2 I_2 \end{pmatrix}}{(z - c_0)(z - c_1)} + \frac{\begin{pmatrix} \theta_1 I_2 \\ \theta_2 \\ \theta_3 \end{pmatrix}}{z - c_0}.$$

Take the trivial partition \mathcal{I} : $\{0, 1, 2, 3\}$, then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

$$H(\mathbf{c}) = \begin{pmatrix} \alpha_1 I_2 \\ \alpha_2 I_2 \end{pmatrix} (z - c_0)^{-4} + \begin{pmatrix} \beta_1 I_2 \\ \beta_2 I_2 \end{pmatrix} (z - c_0)^{-3} \\ + \begin{pmatrix} \gamma_1 I_2 \\ \gamma_2 I_2 \end{pmatrix} (z - c_0)^{-2} + \begin{pmatrix} \theta_1 I_2 \\ \theta_2 \\ \theta_3 \end{pmatrix} (z - c_0)$$

which is the canonical form of the triconfluent equation in Section 4.3.

Let us consider the partition $\mathcal{I}: \{0\} \sqcup \{1, 2, 3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$, we have

with $\alpha_1^{(c_1)}(\mathbf{c}) \neq \alpha_2^{(c_1)}(\mathbf{c})$ and $\theta_i^{(c_0)}(\mathbf{c}) \neq \theta_j^{(c_0)}(\mathbf{c})$ for $i \neq j$. The canonical forms in the right hand side are those of the biconfluent equation of type I in Section 4.3.

Let us consider the partition $\mathcal{I}: \{0, 1, 2\} \sqcup \{3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

$$H(\mathbf{c}) =$$

$$\frac{\begin{pmatrix} \alpha_1^{(c_0)}(\mathbf{c})I_2 & \\ & \alpha_2^{(c_0)}(\mathbf{c})I_2 \end{pmatrix}}{z_{c_0}^3} + \frac{\begin{pmatrix} \beta_1^{(c_0)}(\mathbf{c})I_2 & \\ & \beta_2^{(c_0)}(\mathbf{c})I_2 \end{pmatrix}}{z_{c_0}^2} + \frac{\begin{pmatrix} \theta_1^{(c_0)}(\mathbf{c})I_2 & \\ & \theta_2^{(c_0)}(\mathbf{c}) \end{pmatrix}}{z_{c_0}} + \frac{\begin{pmatrix} \theta_1^{(c_0)}(\mathbf{c})I_2 & \\ & \theta_2^{(c_0)}(\mathbf{c})I_2 \end{pmatrix}}{z_{c_0}} + \frac{\begin{pmatrix} \theta_1^{(c_0)}(\mathbf{c})I_2 & \\ & \theta_$$

with $\alpha_1^{(c_0)}(\mathbf{c}) \neq \alpha_2^{(c_0)}(\mathbf{c})$, $\theta_2^{(c_0)}(\mathbf{c}) \neq \theta_3^{(c_0)}(\mathbf{c})$, and $\theta_1^{(c_3)}(\mathbf{c}) \neq \theta_2^{(c_3)}(\mathbf{c})$. The canonical forms in the right are those of the biconfluent equation of type II in Section 4.3. We can obtain the same canonical forms for the partitions $\{0, 2, 3\} \sqcup \{1\}$ and $\{0, 1, 3\} \sqcup \{2\}$ as well.

Let us consider the partition $\mathcal{I}: \{0,1\} \sqcup \{2,3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

$$H(\mathbf{c}) = \frac{\begin{pmatrix} \alpha_1^{(c_0)}(\mathbf{c})I_2 & \\ & \alpha_2^{(c_0)}(\mathbf{c})I_2 \end{pmatrix}}{z_{c_0}^2} + \frac{\begin{pmatrix} \theta_1^{(c_0)}(\mathbf{c})I_2 & \\ & \theta_2^{(c_0)}(\mathbf{c}) \\ & & \\ & \frac{z_{c_0}}{z_{c_0}} \end{pmatrix}}{z_{c_0}^{(c_2)}(\mathbf{c})I_2} + \frac{\begin{pmatrix} \theta_1^{(c_2)}(\mathbf{c})I_2 & \\ & \theta_2^{(c_2)}(\mathbf{c})I_2 \\ & & \\ & & \\ & & \\ & & \\ & & z_{c_2} \end{pmatrix}},$$

with $\alpha_1^{(c_i)}(\mathbf{c}) \neq \alpha_2^{(c_i)}(\mathbf{c})$ for i = 0, 2 and $\theta_2^{(c_0)}(\mathbf{c}) \neq \theta_3^{(c_0)}(\mathbf{c})$. The canonical forms in the right hand side are those of the doubly-confluent equation in Section 4.3. We can obtain the same canonical forms for the partitions $\{0, 2\} \sqcup \{1, 3\}$, and $\{0, 3\} \sqcup \{1, 2\}$ as well.

Let us consider the partition $\mathcal{I}: \{0\} \sqcup \{1\} \sqcup \{2,3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

with $\theta_j^{(c_i)}(\mathbf{c}) \neq \theta_k^{(c_i)}(\mathbf{c})$ for i = 0, 1 and $j \neq k$ and $\alpha_1^{(c_2)}(\mathbf{c}) \neq \alpha_2^{(c_2)}(\mathbf{c})$. The canonical forms in the right are those of the confluent equation of type I in Section 4.3. We can

obtain the same canonical forms for the partitions $\{0\} \sqcup \{2\} \sqcup \{1,3\}, \{0\} \sqcup \{3\} \sqcup \{1,2\}$ as well.

Let us consider the partition $\mathcal{I}: \{0,1\} \sqcup \{2\} \sqcup \{3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

$$\begin{split} H(\mathbf{c}) &= \\ \frac{\left(\alpha_{1}^{(c_{0})}(\mathbf{c})I_{2} \\ \alpha_{2}^{(c_{0})}(\mathbf{c})I_{2} \right)}{z_{c_{0}}^{2}} + \frac{\left(\begin{array}{c} \theta_{1}^{(c_{0})}(\mathbf{c})I_{2} \\ \theta_{2}^{(c_{0})}(\mathbf{c}) \\ z_{c_{0}} \end{array} \right)}{z_{c_{0}}} \\ \frac{\left(\theta_{1}^{(c_{2})}(\mathbf{c})I_{2} \\ \theta_{2}^{(c_{2})}(\mathbf{c})I_{2} \end{array} \right)}{z_{c_{2}}} + \\ \frac{\left(\theta_{1}^{(c_{3})}(\mathbf{c})I_{2} \\ \theta_{2}^{(c_{3})}(\mathbf{c})I_{2} \right)}{z_{c_{3}}}, \end{split}$$

with $\alpha_1^{(c_0)}(\mathbf{c}) \neq \alpha_2^{(c_0)}(\mathbf{c}), \ \theta_2^{(c_0)}(\mathbf{c}) \neq \theta_3^{(c_0)}(\mathbf{c}), \ \text{and} \ \theta_1^{(c_i)}(\mathbf{c}) \neq \theta_2^{(c_i)}(\mathbf{c}) \text{ for } i = 2, 3.$ The canonical forms in the right hand side are those of the confluent equation of type II in Section 4.3. We can obtain the same canonical forms for the partitions $\{0, 2\} \sqcup \{1\} \sqcup \{3\}$ and $\{0, 3\} \sqcup \{1\} \sqcup \{2\}$ as well.

Finally consider the finest partition $\mathcal{I}: \{0\} \sqcup \{1\} \sqcup \{2\} \sqcup \{3\}$. Then for $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}_H$,

$$H(\mathbf{c}) = \frac{\begin{pmatrix} \theta_1^{(c_0)}(\mathbf{c})I_2 & & \\ & \theta_2^{(c_0)}(\mathbf{c}) \\ & & \theta_3^{(c_0)}(\mathbf{c}) \end{pmatrix}}{z_{c_0}} + \sum_{i=1}^3 \frac{\begin{pmatrix} \theta_1^{(c_i)}(\mathbf{c})I_2 & \\ & \theta_2^{(c_i)}(\mathbf{c})I_2 \\ & & z_{c_i} \end{pmatrix}}{z_{c_i}},$$

with $\theta_j^{(c_i)}(\mathbf{c}) \neq \theta_k^{(c_i)}(\mathbf{c})$ for i = 0, 1, 2, 3 and $j \neq k$. The canonical forms in the right hand side are those of the first equation in Section 4.3.

Therefore as well as the Heun-type equations, the unfolding of H recovers the canonical forms for all the confluent equations appeared in Section 4.3.

6 Additive Deligne-Simpson problem and a conjecture by Oshima

In the previous section, we introduced a deformation of unramified canonical forms and showed that this deformation preserves some important invariants of the canonical forms. In this section, we propose a problem to find a deformation of irreducible meromorphic G-connection defined over \mathbb{P}^1 which realizes the given deformation of canonical forms.

6.1 Additive Deligne-Simpson problem for unfolding families of canonical forms

Let us consider an effective divisor $D = \sum_{a \in \{a_1, \dots, a_d\}} (k_a + 1) \cdot a$ as in Section 4.

Definition 6.1 (Irreducible connection, cf. Arinkin [3]). For a \mathfrak{g} -valued meromorphic 1-form $A dz \in \Omega^{\mathfrak{g}}_{\mathbb{P}^1,D}(\mathbb{P}^1)$, we say the associated *G*-connection $d_A = d + A dz$ is *irreducible* if there is no proper parabolic subalgebra \mathfrak{p} of \mathfrak{g} such that $A dz \in \mathfrak{p} \otimes_{\mathbb{C}} \Omega_{\mathbb{P}^1,D}(\mathbb{P}^1)$

We now consider the following problem which asks the existence of an irreducible Gconnection with the prescribed local isomorphic classes. We assume $|D| \subset \mathbb{C}$ under the
projective transformation for simplicity.

Problem 6.2 (Additive Deligne-Simpson problem for **H**). Let us take a collection of unramified canonical forms $\mathbf{H} = (H^{(a)})_{a \in |D|} \in \prod_{a \in |D|} \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})$ satisfying the residue condition $\sum_{a \in |D|} H^{(a)}_{\text{res}} \in \mathfrak{g}_{\text{ss}}$. Then find an irreducible meromorphic *G*-connection

$$d_A = d + \sum_{a \in |D|} \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^i} \frac{dz}{z-a}$$

with singularities only on $|D| \subset \mathbb{P}^1$ such that

$$\sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}} \in \mathbb{O}_{H^{(a)}}$$

for all $a \in |D|$.

As it is explained in Section 1.5, this problem was considered by Deligne, Simpson, and Kostov, and after their pioneering works there are many subsequent developments around this problem, see [10], [28], [6], [19], [15], [29], [32], and [23].

Now we also consider a generalization of this problem as follows. For a collection $\mathbf{H} = (H^{(a)})_{a \in |D|} \in \prod_{a \in |D|} \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})$ of unramified canonical forms satisfying the residue condition $\sum_{a \in |D|} H^{(a)}_{\text{res}} \in \mathfrak{g}_{\text{ss}}$, let $\mathbf{H}(\mathbf{c}) = (H^{(a)}(\mathbf{c}_a))_{\mathbf{c}=(\mathbf{c}_a)\in\prod_{a \in |D|} \mathbb{C}^{k_a+1}}$ be the unfolding of \mathbf{H} . Then as in Section 5.5, for each $\mathbf{c} \in \prod_{a \in |D|} \mathbb{C}^{k_a+1}$ we regard $\mathbf{H}(\mathbf{c})$ as the collection of the canonical forms $\mathbf{H}(\mathbf{c}) = (H^{(a)}(\mathbf{c}_a)_i)_{i=0,1,\dots,r_a}$.

Problem 6.3 (Additive Deligne-Simpson problem for an unfolding family of canonical forms). Let U be an open neighborhood of $\mathbf{0} \in \prod_{a \in |D|} \mathbb{C}^{k_a+1}$. Let $(\mathbf{H}(\mathbf{c}))_{\mathbf{c} \in U}$ be the unfolding of a collection of canonical forms $\mathbf{H} = (H^{(a)})_{a \in |D|}$ as above. Then find an meromorphic G-connection over \mathbb{P}^1 ,

$$d_{A(\mathbf{c})} = d + A(\mathbf{c}) \, dz$$

satisfying all the following conditions.

- 1. The g-valued 1-form $A(\mathbf{c}) dz$ depends holomorphically on $\mathbf{c} \in U$.
- 2. For each $\mathbf{c} \in U$, $d_{A(\mathbf{c})}$ is a solution of the additive Deligne-Simpson problem for $\mathbf{H}(\mathbf{c})$.

In regard to Problems 6.2 and 6.3, we obtain the following whose proof is postponed until Section 11.

Theorem 6.4. Let **H** be a collection of unramified canonical forms as above. Let $d_A = d + A dz$ be a solution of the additive Deligne-Simpson problem for **H**. Then there exists an open neighborhood U of $\mathbf{0} \in \prod_{a \in |D|} \mathbb{C}^{k_a+1}$ and a holomorphic family of G-connections $d_{A(\mathbf{c})} = d + A(\mathbf{c}) dz$ ($\mathbf{c} \in U$) on \mathbb{P}^1 such that

- $d_{A(\mathbf{0})} = d_A$,
- $d_{A(\mathbf{c})} = d + A(\mathbf{c}) dz$ is a solution of the additive Deligne-Simpson for the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c}\in U}$.

In particular, the following are equivalent.

- 1. The additive Deligne-Simpson problem for H has a solution.
- 2. There exists an open neighborhood U of $\mathbf{0} \in \prod_{a \in |D|} \mathbb{C}^{k_a+1}$ and the additive Deligne-Simpson problem for the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c} \in U}$ has a solution.

6.2 Unfolding of spectral types

As we saw in Corollary 5.5, the unfolding $H(\mathbf{c})$ of an unramified canonical form H induces decompositions of the spectral type of H. We give a diagrammatic description of these decompositions.

According to the decomposition in Corollary 5.5, we define the unfolding of abstract spectral types as follows. Let S be an abstract spectral type as above. For a partition $\mathcal{I}: I_0, \ldots, I_r \in \mathcal{P}_{[k+1]}$ of the index set $\{0, 1, \ldots, k\}$, write $I_j = \{i_{[j,0]} < i_{[j,1]} < \cdots < i_{[j,k_j]}\}$ for $j = 1, 2, \ldots, r$ and assume $0 \in I_0$. Then for each $\mathcal{I} \in \mathcal{P}_{[k+1]}$, define the collection $S^{\mathcal{I}} = (S^{I_j})_{j=1,\ldots,r}$ of spectral types by

$$S^{I_0} := (\Pi_{i_{[0,k_0]}} \supset \cdots \supset \Pi_{i_{[0,1]}} \supset \Pi_0; [J_0]),$$

$$S^{I_j} := (\Pi_{i_{[j,k_j]}} \supset \cdots \supset \Pi_{i_{[j,1]}} \supset \Pi_{i_{[j,0]}}; [0]), \ j = 1, \dots, r,$$

which corresponds to the decomposition of the sequence $\Pi_k \supset \cdots \supset \Pi_1 \supset \Pi_0$ with regard to the partition \mathcal{I} . We call this collection $S^{\mathcal{I}} = (S^{I_j})_{j=1,\dots,r}$ an *unfolding* of S associated to \mathcal{I} .

Also for a collection

$$\mathbf{S} = (S_i)_{i=1,2,\dots,d} = \left((\Pi_{k_i}^{(i)} \supset \Pi_{k_i-1}^{(i)} \supset \dots \supset \Pi_0^{(i)}; [J_0^{(i)}]) \right)_{i=1,2,\dots,d}$$

of abstract spectral types, we can define the *unfolding*

$$\mathbf{S}^{(\mathcal{I}^{(i)})_{i=1,2,\dots,d}} := (S_i^{I_j^{(i)}})_{\substack{i=1,2,\dots,d,\\j=1,2,\dots,r_i}}$$

of **S** associated to $(\mathcal{I}^{(i)})_{i=1,2,\dots,d} = (I_0^{(i)} \sqcup \cdots \sqcup I_{r_i}^{(i)})_{i=1,2,\dots,d} \in \prod_{i=1}^d \mathcal{P}_{[k_i+1]}.$ In particular for the finest partitions $(\{0\} \sqcup \cdots \{k_i\})_{i=1,\dots,d} \in \prod_{i=1}^d \mathcal{P}_{[k_i+1]}$, the corre-

In particular for the finest partitions $(\{0\} \sqcup \cdots \{k_i\})_{i=1,\dots,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$, the corresponding unfolding

$$\mathbf{S}^{\text{reg}} := \mathbf{S}^{(\{0\} \sqcup \dots \{k_i\})_{i=1,\dots,d}} = \left((\Pi_0^{(1)}; [J_0^{(1)}]), (\Pi_1^{(1)}; [0]), \dots, (\Pi_{k_1}^{(1)}; [0]), \dots, (\Pi_{k_d}^{(d)}; [J_0^{(d)}]), (\Pi_1^{(d)}; [0]), \dots, (\Pi_{k_d}^{(d)}; [0]) \right)$$

of \mathbf{S} which is a collection of spectral types for regular singular canonical forms is called the *finest unfolding* of \mathbf{S} .

Let us introduce the unfolding diagram of a collection **S** of spectral types. Recall that $\prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$ becomes a product poset, i.e., for $(\mathcal{I}^{(i)})_{i=1,\dots,d}, (\mathcal{J}^{(i)})_{i=1,\dots,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$, we have $(\mathcal{I}^{(i)})_{i=1,\dots,d} \leq (\mathcal{J}^{(i)})_{i=1,\dots,d}$ if and only if $\mathcal{I}^{(i)} \leq \mathcal{J}^{(i)}$ for all $i = 1,\dots,d$. Thus

we obtain the Hasse diagram of the poset $\prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$. Then we attach to each vertex $(\mathcal{I}^{(i)})_{i=1,\dots,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$ the corresponding unfolding $\mathbf{S}^{(\mathcal{I}^{(i)})_{i=1,2,\dots,d}}$ of \mathbf{S} . Then we call the resulting diagram the *unfolding diagram* of \mathbf{S} .

For example let us recall the spectral type of the triconfluent Heun-type equation $(\emptyset \supset \emptyset \supset \emptyset \supset \emptyset; [0])$. Then we obtain the following unfolding diagram.



In this diagram we notice that same collections of spectral types appear in several vertices. This implies that we can reduce the diagram to a smaller one as follows.

Let $\mathbf{S} = (S_i)_{i=1,2,\dots,d}$ be a collection of abstract spectral types as above. Then there exist positive integers $0 \leq l_1^{(i)} < l_2^{(i)} < \cdots < l_{t_i}^{(i)} = k_i$ such that

$$\Pi_{k_i}^{(i)} = \dots = \Pi_{l_{t_i-1}^{(i)}+1}^{(i)} \supseteq \dots \supseteq \Pi_{l_2^{(i)}}^{(i)} \dots = \Pi_{l_1^{(i)}+1}^{(i)} \supseteq \Pi_{l_1^{(i)}}^{(i)} = \dots = \Pi_0^{(i)}$$

for i = 1, 2, ..., d. Let us consider the subgroup $\mathfrak{S}_{l_0^{(i)}+1} \times \mathfrak{S}_{l_1^{(i)}-l_0^{(i)}} \times \cdots \mathfrak{S}_{l_{t_i}^{(i)}-l_{t_i-1}^{(i)}}$ of the symmetric group \mathfrak{S}_{k_i+1} of degree $k_i + 1$ which acts on $\{0, 1, ..., k_i\}$ as the permutation. Then we can define quotient posets

$$\mathcal{P}^{S_i}_{[k_i+1]} := \mathcal{P}_{[k_i+1]} / (\mathfrak{S}_{l_0^{(i)}+1} \times \mathfrak{S}_{l_1^{(i)}-l_0^{(i)}} \times \cdots \mathfrak{S}_{l_{t_i}^{(i)}-l_{t_i-1}^{(i)}})$$

and the product $\prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}^{S_i}$ of them. If we take $(\mathcal{I}_i)_{i=1,\dots,d}, (\mathcal{J}_i)_{i=1,\dots,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$ and they are equal in the quotient $\prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}^{S_i}$, then we have $\mathbf{S}^{(\mathcal{I}_i)_{i=1,\dots,d}} = \mathbf{S}^{(\mathcal{J}_i)_{i=1,\dots,d}}$. Thus the following diagram is well-defined. Namely, let us consider the Hasse diagram of the poset $\prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}^{S_i}$ and attach to the each vertex $[(\mathcal{I}_i)_{i=1,\dots,d}]$ the unfolding $\mathbf{S}^{(\mathcal{I}_i)_{i=1,\dots,d}}$ of \mathbf{S} . We call this diagram the *reduced unfolding diagram* of \mathbf{S} .

For the above spactral type $(\emptyset \supset \emptyset \supset \emptyset \supset \emptyset; [0])$, the reduced unfolding diagram is drawn as follows.



Let us also consider the spectral type $(\{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\} \supset \{e_{12}, e_{34}\} \supset \{e_{12}\}; [0])$. of the triconfluent equation in Section 4.3. Then we obtain the reduced unfolding diagram,



6.3 A conjecture by Oshima

Let us explain a conjecture proposed by Oshima in the paper [38] which asks the existence of differential equations on \mathbb{P}^1 with a prescribed collection of spectral types. A collection

$$\mathbf{S} = (S_i)_{i=1,2,\dots,d} = \left((\Pi_{k_i}^{(i)} \supset \Pi_{k_i-1}^{(i)} \supset \dots \supset \Pi_0^{(i)}; [J_0^{(i)}]) \right)_{i=1,2,\dots,d}$$

of abstract spectral types are said to be *irreducibly realizable* when there exists an irreducible meromorphic G-connections over \mathbb{P}^1 with poles only at $\{a_1, a_2, \ldots, a_d\} \subset \mathbb{P}^1$ which has the spectral type $(\prod_{k=1}^{(i)} \supset \prod_{k=1}^{(i)} \supset \cdots \supset \prod_{0}^{(i)}; [J_0^{(i)}])$ at $z = a_i$ for $i = 1, 2, \ldots, d$. This can be restated as follows. A collection $\mathbf{S} = (S_i)_{i=1,2,\ldots,d}$ of abstract spectral types is irreducibly realizable if and only if there exists a collection $\mathbf{H} = (H^{(a_i)})_{i=1,2,\ldots,d}$ of unramified canonical forms satisfying that

- sp (**H**) = **S**, where sp (**H**) := (sp ($H^{(a_i)}$))_{i=1,2,\dots,d},
- the additive Deligne-Simpson problem for **H** has a solution.

Furthermore, a collection $\mathbf{S} = (S_i)_{i=1,2,\dots,d}$ of spectral types is said to be *versally realizable* when there exists a collection $\mathbf{H} = (H^{(a_i)})_{i=1,2,\dots,d}$ of unramified canonical forms satisfying that

- $\operatorname{sp}(\mathbf{H}) = \mathbf{S},$
- there exists an open neighborhood U of $\mathbf{0} \in \prod_{i=1}^{d} \mathbb{C}^{k_i+1}$ and the additive Deligne-Simpson problem for the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c}\in U}$ has a solution.

As we saw in the previous subsection, for a collection $\mathbf{S} = (S_i)_{i=1,2,...,d}$ of spectral types we can consider the reduced unfolding diagram of \mathbf{S} as in the previous section. Then we say that the unfolding diagram is *irreducibly realizable* if there exists a collection $\mathbf{H} = (H^{(a_i)})_{i=1,2,...,d}$ of unramified canonical forms satisfying

• there exists an open neighborhood U of $\mathbf{0} \in \prod_{i=1}^{d} \mathbb{C}^{k_i+1}$ such that the deformation $\mathbf{H}(\mathbf{c})_{\mathbf{c}\in U}$ realizes the unfolding diagram, namely, for each $(\mathcal{I}^{(i)})_{i=1,\dots,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$ and $\mathbf{c} \in U \cap (\prod_{i=1}^{d} \mathcal{C}(\mathcal{I}^{(i)}))$, we have

$$\operatorname{sp}(\mathbf{H}(\mathbf{c})) = \mathbf{S}^{(\mathcal{I}^{(i)})_{i=1,\dots,d}}$$

• there exists an open neighborhood of $\mathbf{0} \in \mathbb{B}_{\mathbf{H}}$ and the additive Deligne-Simpson problem for the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c}\in U}$ has a solution.

Then as a corollary of Theorem 6.4 we obtain the following.

Theorem 6.5. Let $\mathbf{S} = (S_i)_{i=1,...,d}$ be a collection of abstract spectral types. Suppose that \mathbf{S} is irreducibly realizable and let $d_A = d + A dz$ be a realization of \mathbf{S} . Then there exists an open neighborhood U of $\mathbf{0} \in \prod_{i=1}^d \mathbb{C}^{k_i+1}$ and a holomorphic family of G-connections $d_{A(\mathbf{c})} = d + A(\mathbf{c}) dz$ ($\mathbf{c} \in U$) on \mathbb{P}^1 such that

- $d_{A(\mathbf{0})} = d_A$,
- for each $(\mathcal{I}_i)_{i=1,\dots,d} \in \prod_{i=1}^d \mathcal{P}_{[k_i+1]}$ and $\mathbf{c} \in U \cap \prod_{i=1}^d \mathcal{C}(\mathcal{I}^{(i)}), d_{A(\mathbf{c})} = d + A(\mathbf{c}) dz$ is a realization of $\mathbf{S}^{(\mathcal{I}_i)_{i=1,\dots,d}}$.

In particular, the following are equivalent.

- 1. The collection \mathbf{S} of abstract spectral types is irreducibly realizable.
- 2. For every $(\mathcal{I}^{(i)})_{i=1,\dots,d} \in \prod_{i=1}^{d} \mathcal{P}_{[k_i+1]}$, the collection $\mathbf{S}^{(\mathcal{I}^{(i)})_{i=1,\dots,d}}$ is irreducibly realizable.
- 3. The unfolding diagram of \mathbf{S} is irreducibly realizable.

Proof. Conditions 2 and 3 are obviously equivalent. From 2 or 3, 1 immediately follows. Suppose 1 holds. Then there exists a collection $\mathbf{H} = (H^{(i)})_{i=1,...,d}$ of unramified canonical forms such that $\operatorname{sp}(\mathbf{H}) = \mathbf{S}$ and the additive Deligne-Simpson problem for \mathbf{H} has a solution $d_A = d + A \, dz$. Then Theorem 6.4 shows that there exists an open neighborhood U of $\mathbf{0} \in \prod_{i=1}^d \mathbb{C}^{k_i+1}$ and a holomorphic family of G-connections $d_{A(\mathbf{c})} = d + A(\mathbf{c}) \, dz$ ($\mathbf{c} \in U$) on \mathbb{P}^1 such that $d_z A(\mathbf{0}) = d_A$, and $d_{A(\mathbf{c})} = d + A(\mathbf{c}) \, dz$ is a solution of the additive Deligne-Simpson for the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c}\in U}$. Let $\mathbb{B}_{\mathbf{H}} = \prod_{i=1}^d \mathbb{B}_{H^{(i)}}$ be open neighborhood of $\mathbf{0} \in \prod_{i=1}^d \mathbb{C}^{k_i+1}$ defined in Section 5.5 and set $V := U \cap \mathbb{B}_{\mathbf{H}} \neq \emptyset$. Then since V is an open neighborhood of $\mathbf{0} \in \prod_{i=1}^d \mathbb{C}^{k_i+1}$ Lemma 5.2 assures that $V \cap \prod_{i=1}^d \mathcal{C}(\mathcal{I}^{(i)}) \neq \emptyset$ for any $(\mathcal{I}^{(i)})_{i=1,...,d} \in \prod_{i=1}^d \mathcal{P}_{[k_i+1]}$. Then by Corollary 5.5 we have $\operatorname{sp}(\mathbf{H}(\mathbf{c})) = \mathbf{S}^{(\mathcal{I}^{(i)})_{i=1,...,d}}$ for all $\mathbf{c} \in V \cap \prod_{i=1}^n \mathcal{C}(\mathcal{I}^{(i)})$ and $(\mathcal{I}^{(i)})_{i=1,...,d} \in \prod_{i=1}^d \mathcal{P}_{[k_i+1]}$. Namely the unfolding diagram of \mathbf{S} is realizable.

Now we are ready to state the conjecture by Oshima.

Conjecture 6.6 (Oshima [38]). Let $\mathbf{S} = (S_i)_{i=1,\dots,d}$ be a collection of abstract spectral types. Then the following are equivalent.

- 1. The collection \mathbf{S} is irreducibly realizable.
- 2. The collection \mathbf{S} is versally realizable.
- 3. The finest unfolding \mathbf{S}^{reg} of \mathbf{S} is irreducibly realizable.

As a direct consequence of Theorems 6.4 and 6.5, we obtain the following.

Corollary 6.7. In Conjecture 6.6, 1 and 2 are equivalent and the implication $1 \Rightarrow 3$ holds true.

Proof. The first assertion directly follows from Theorem 6.4. The direction $1 \Rightarrow 3$ follows from Theorem 6.5.

7 Moduli spaces of meromorphic connections on \mathbb{P}^1 with unramified irregular signarities

We recall the definition of moduli spaces of algebraic meromorphic connections on a trivial bundle on the projective line $\mathbb{P}^1 = \mathbb{P}^1(\mathbb{C})$ with unramified irregular singularities.

7.1 Lie groupoids, Orbifolds, and symplectic stratified spaces

Let us recall the notion of Lie groupoids, see [33] by Mackenzie and its references for more detail.

Definition 7.1 (Complex Lie groupoid). Let us consier a complex manifold M and its submanifold M_0 with the inclusion map $i: M_0 \hookrightarrow M$. Let us also consider holomorphic surjective submersions $s, t: M \to M_0$ satisfying $t \circ i = s \circ i = \mathrm{id}_{M_0}$. Further, we consider a holomorphic map $m: M^{(2)} := \{(g_1, g_2) \in M \times M \mid s(g_1) = t(g_2)\} \to M$ satisfying $s \circ m(g_1, g_2) = s(g_2)$ and $t \circ m(g_1, g_2) = t(g_1)$. Then if the following conditions are satisfied, the tuple (M, M_0, s, t, m) is called a *complex Lie groupoid*.

- 1. Associativity: $m(m(g_1, g_2), g_3) = m(g_1, m(g_2, g_3))$ for all $(g_1, g_2, g_3) \in M^3$ satisfying $s(g_1) = t(g_2)$ and $s(g_2) = t(g_3)$.
- 2. Units: $m(i \circ t(g), g) = g = m(g, i \circ s(g))$ for all $g \in M$.
- 3. Inverces: For all $g \in M$, there exists $h \in M$ such that s(h) = t(g), t(h) = s(g) and $m(g,h), m(h,g) \in M_0$.

Here the maps s and t are called the *source map* and *target map* respectively, and m is called the *multiplication map*. Also elements of the submanifold $i: M_0 \hookrightarrow M$ are called *units*.

Definition 7.2 (Bisection). Let $M = (M, M_0, s, t, m)$ be a complex Lie groupoid. A *bisection* of M is a submanifold $S \subset M$ such that both s and t restrict to biholomorphic map $S \to M_0$.

Let us recall the notion of orbifold groupoids. We refer [1] by Adem-Leida-Ruan, for more details.

Definition 7.3 (Orbit space of a Lie groupoid). Let $M = (M, M_0, s, t, m)$ a complex Lie groupoid. For $m \in M_0$, the *orbit* of m is the subset of M_0 defined by

$$M(m) := \{t(g) \in M_0 \mid s(g) = m\} = t(s^{-1}(m)).$$

Then we can define the equivalent relation $m_1 \sim m_2$ on M_0 by $m_1 \in M(m_2)$. The quotient space by this equivalent relation $|M| := M_0 / \sim$ is called the *orbit space* of M.

As a typical example of Lie groupoids, we introduce action groupoids.

Definition 7.4 (Action groupoids). Let X be a complex manifold and H a complex Lie group acting holomorphically on X from the left. Then let $M := H \times X$ and $M_0 = X$. We define the source map $s \colon H \times X \to X$ by the second projection pr_2 and the target map by

$$t \colon H \times X \ni (h, x) \longmapsto h \cdot x \in X.$$

Also define the inclusion by $i: X \ni x \mapsto (e, x) \in H \times X$. Then the multiplication map

$$m: M^{(2)} \ni ((h_1, x), (h_2, h_2^{-1} \cdot x)) \longmapsto (h_1 h_2, h_2^{-1} \cdot x) \in M$$

defines the groupoid structure on $(H \times X, X, s, t, m)$. We call this Lie groupoid an *action* groupoid. In this case, the orbit space of the action groupoid coincides with the usual orbit space $H \setminus X$.

Let us recall the proper, étale and foliation groupoids.

Definition 7.5 (Proper, étale and foliation groupoids). Let $M = (M, M_0, s, t, m)$ be a Lie groupoid. Then M is called a *proper groupoid* if the product map $s \times t \colon M \to M_0 \times M_0$ is a proper map. If both s and t are locally homeomorphisms, then M is called an *étale groupoid*. If *isotropy groups* $M_m := s^{-1}(m) \cap t^{-1}(m)$ are discrete for all $m \in M_0$, then M is called a *foliation groupoid*.

Then we can define orbifold groupoids.

Definition 7.6 (Orbifold groupoids). We call a proper étale foliation groupoid an *orbifold* groupoid.

If a Lie group H acts on a complex manifold X almost freely and properly, then one can check that the action groupoid $H \times X$ is an example of orbifold groupoids.

Definition 7.7 (Orbifolds). An orbifold structure on a paracompact Hausdorff space X is given by an orbifold groupoid M and a homeomorphism $|M| \to X$. Equivalent classes of orbifold structures are defined by Morita equivalence, see [1]. An orbifold \mathcal{X} is a paracompact Hausdorff space with an equivalence class of orbifold structures.

Therefore if H acts on X almost freely and properly as above, the quotient space $H \setminus X$ becomes an orbifold with the orbifold structure coming from the action groupoid $H \times X$.

Next let us recall symplectic stratified spaces and symplectic reductions. Firstly, according to the paper [14] by Goresky and MacPherson, recall topological stratified spaces.

Definition 7.8 (Stratified space). Let X be a \mathcal{P} -decomposed space with the pieces $(S_i)_{i \in \mathcal{P}}$. Then X is called a *stratified space* if the pieces of X, called *strata*, satisfies the following *local normal triviality*. Namely, for each point $x \in S_i$, there is a compact stratified space L, called the *link* of x, and a homeomorphism h of an open neighborhood U of x in X on the product $B \times cL$. Here B is a open ball in S_i and cL is the open cone $L \times [0, 1)/L \times \{0\}$ over L. Moreover h preserves the decomposition.

As it is also well-known, a quotient orbifold $H \setminus X$ has the following natural stratification. For a subgroup $\Gamma \subset H$, we consider the subset $X_{(\Gamma)}$ of X consisting of all points whose stabilizer group is conjugate to Γ . Then the collection of the quotient spaces of these subset $(H \setminus X_{(\Gamma)})_{\Gamma \subset H}$ defines $H \setminus X$ as a stratified space.

The quotient orbifold $H \setminus X$ has the following natural holomorphic structure,

$$\mathcal{O}(H \setminus X) := \{ f \in C^0(H \setminus X) \mid f \circ \pi \text{ is holomorphic} \},\$$

where $\pi: X \to H \setminus X$ is the quotient map. In general, for a stratified space X with the strata $(S_i)_{i \in \mathcal{I}}$ which are complex manifolds, a *holomorphic structure* $\mathcal{O}(X)$ is a subalgebra of $C^0(X)$ having the property that for any $f \in \mathcal{O}(X)$ the restriction on a stratum S_i is holomorphic, $f|_{S_i} \in \mathcal{O}(S_i)$.

Definition 7.9 (Stratified holomorphic symplectic space). Let X be a stratified space with a holomorphic structure $\mathcal{O}(X)$. Then X is called a *stratified holomorphic symplectic space* if the following are satisfied,

- 1. each stratum S_i is a holomorphic symplectic manifolds,
- 2. $\mathcal{O}(X)$ is a Poisson algebra,
- 3. the embeddings $S_i \hookrightarrow X$ are Poisson.

Then the following is a generalization of the symplectic reduction theory by Marsden and Weinstein [35] to stratified spaces given by Sjammar and Lerman [43].

Theorem 7.10 (Marsden-Weinstein and Sjamaar-Lerman). Let X be a holomorphic symplectic manifold on which a complex Lie group H acts properly and almost freely. Let $\mu: X \to \mathfrak{h}^*$ be a moment map. Then the quotient space $H \setminus \mu^{-1}(0)$ becomes an orbifold whose canonical stratification gives the structure of stratified holomorphic symplectic space.

Remark 7.11. This is a consequence of Theorem 2.1 in [43] in which the Lie group H is assumed to be compact. However as it is stated in the introduction of the same paper [43], Theorem 2.1 is valid for proper actions of arbitrary Lie groups.

7.2 Moduli spaces of meromorphic connections on a trivial bundle over $\mathbb{P}^1(\mathbb{C})$

Let $D = \sum_{a \in \{a_1, a_2, \dots, a_d\}} (k_a + 1) \cdot a$ be an effective divisor on \mathbb{P}^1 . Under the projective transformation, we assume $|D| \subset \mathbb{C}$ for simplicity. Let us consider the space of meromorphic *G*-connections

$$\overline{\mathcal{M}}_D := \left\{ d_A = d + A \, dz \, \middle| \, A \, dz \in \Omega^{\mathfrak{g}}_{\mathbb{P}^1, D}(\mathbb{P}^1) \right\}.$$

Since each $d + A dz \in \overline{\mathcal{M}}_D$ is of the form

$$A \, dz = \sum_{a \in |D|} \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^i} \frac{dz}{z-a} \quad (A_i^{(a)} \in \mathfrak{g})$$

with the relation $\sum_{a \in |D|} A_0^{(a)} = 0$, we can identify $\overline{\mathcal{M}}_D$ with the space

$$\left\{ \left(X_i^{(a)}\right)_{\substack{a\in|D|,\\i=0,1,\dots,k_a}} \in \prod_{a\in|D|} \mathfrak{g}^{k_a+1} \left| \sum_{a\in|D|} X_0^{(a)} = 0 \right\} \right\}.$$

The G-action on $\overline{\mathcal{M}}_D$ by the gauge transformation is translated into the diagonal adjoint action, i.e.,

$$G \times \overline{\mathcal{M}}_D \ni \left(g, \left(X_i^{(a)}\right)_{\substack{a \in |D|, \\ i=0,1,\dots,k_a}}\right) \longmapsto \left(\operatorname{Ad}(g)(X_i^{(a)})\right)_{\substack{a \in |D|, \\ i=0,1,\dots,k_a}} \in \overline{\mathcal{M}}_D.$$

Under this identification, we regard $\overline{\mathcal{M}}_D$ as a complex manifold with the algebraic *G*-action.

Then following the paper [5] by Boalch we define the moduli space of meromorphic G-connections on \mathbb{P}^1 with given local canonical forms, see also [19] by Hiroe-Yamakawa and [46] by Yamakawa, and also see [8] by Bremer-Sage for ramified cases.

Definition 7.12 (Moduli spaces of meromorphic *G*-connections). For a collection of canonical forms $\mathbf{H} = (H^{(a)})_{a \in |D|} \in \prod_{a \in |D|} \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})$ with $\sum_{a \in |D|} H^{(a)}_{\text{res}} \in \mathfrak{g}_{\text{ss}}$, we consider quotient spaces

$$\mathcal{M}_{\mathbf{H}} := G \left\{ d_A = d + \sum_{a \in |D|} \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^i} \frac{dz}{z-a} \in \overline{\mathcal{M}}_D \left| \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}} \in \mathbb{O}_{H_a} \text{ for all } a \in |D| \right\} \right\}$$

and call this space the moduli spaces of connections associated with **H**. We also consider the subspace $\mathcal{M}_{\mathbf{H}}^{ir}$ of $\mathcal{M}_{\mathbf{H}}$ consists of irreducible connections and call this subspace the moduli spaces of irreducible connections associated with **H**.

Although the moduli space $\mathcal{M}_{\mathbf{H}}$ itself is a naive quotient space, we shall later explain that the subspace $\mathcal{M}_{\mathbf{H}}^{ir}$ of $\mathcal{M}_{\mathbf{H}}$ has a structure of holomorphic symplectic orbifold.

7.3 Stability of G/Z-action on the space of irreducible connections

Let us also consider

$$\left(\prod_{i=1}^{n} \mathfrak{g}\right)^{\mathrm{ir}} := \left\{ (X_i)_{i=1,\dots,n} \in \prod_{i=1}^{n} \mathfrak{g} \left| (X_i)_{i=1,\dots,n} \text{ is irreducible} \right\},\right.$$

the subspace of $\prod_{i=1}^{n} \mathfrak{g}$ consisting of all irreducible elements, where $(X_i)_{i=1,\dots,n} \in \prod_{i=1}^{n} \mathfrak{g}$ is said to be *irreducible* if there exists no proper parabolic subalgebra of \mathfrak{g} containing all X_i for $i = 1, \dots, n$.

Since the center Z of G acts trivially on $\prod_{i=1}^{n} \mathfrak{g}$, we can consider the action of G/Z on $\prod_{i=1}^{n} \mathfrak{g}$. Let us explain that $(\prod_{i=1}^{n} \mathfrak{g})^{\text{ir}}$ coincides with the space of stable points in $\prod_{i=1}^{n} \mathfrak{g}$ with respect to the G/Z-action.

We follow the formulations in [26] by Kempf. Let $\lambda : \mathbb{G}_m \to G/Z$ be a one-parameter subgroup of the algebraic group G/Z. Let χ be a character of the image of λ . Then we can write

$$\chi(\lambda(t)) = t^{\chi(\lambda)} \quad (t \in \mathbb{G}_m)$$

by some integer $\chi(\lambda)$ which characterizes χ .

Let V be a \mathbb{C} -vector space with a linear G/Z-action. Then we consider the weight decomposition $V = \bigoplus V^{\chi}$ of V where χ runs through the set of characters of the image of λ , namely, V^{χ} are χ -eigenspaces of V under the action of \mathbb{G}_m through λ . Then we can associate the integer to each $v \in V$ defined by

$$m(v, \lambda) := \min\{\chi(\lambda) \mid \text{projection of } v \text{ onto } V^{\chi} \text{ is non-zero}\}.$$

Then we can see that the limit $\lim_{t\to 0} \lambda(t) \cdot v$ exists if and only if $m(v, \lambda) \ge 0$.

On the other hand, to a one-parameter subgroup λ , we can associate the parabolic subalgebra of $\mathfrak{g}_{ss} = \operatorname{Lie}(G/Z)$ defined by

$$\mathfrak{p}(\lambda) := \{ X \in \mathfrak{g}_{ss} \mid m(X, \lambda) \ge 0 \}.$$

Proposition 7.13. For $X \in \prod_{i=1}^{n} \mathfrak{g}$, X is irreducible if and only if it is stable under the action of G/Z, i.e., the G/Z-orbit of X is Zariski closed and the stabilizer $\operatorname{Stab}_{G/Z}(X)$ of X has the finite cardinality.

Proof. Obviously any element X in the center \mathfrak{z} of \mathfrak{g} satisfies $m(X,\lambda) = 0$. Thus by Proposition 8.4.5 in [44], any parabolic subalgebra of \mathfrak{g} is of the form $\mathfrak{z} \oplus \mathfrak{p}(\lambda)$. Therefore we can conclude that $X \in \prod_{i=1}^{n} \mathfrak{g}$ is irreducible if and only if X satisfies $m(X,\lambda) < 0$ for any one-parameter subgroup λ of G/Z. Then the desired result follows from the Hilbert-Mumford criterion (see Theorem 2.1 in [36]).

It follows from this proposition that $(\prod_{i=1}^{n} \mathfrak{g})^{\text{ir}}$ coincides with the space of stable points in $\prod_{i=1}^{n} \mathfrak{g}$ as desired. In particular, $(\prod_{i=1}^{n} \mathfrak{g})^{\text{ir}}$ is a Zariski open subset of $\prod_{i=1}^{n} \mathfrak{g}$.

7.4 Symplectic orbifold \mathcal{M}_{H}^{ir}

Let us consider the injective immersion

$$\iota_{\mathbf{H}} \colon \prod_{a \in |D|} \mathbb{O}_{H_a} \longrightarrow \prod_{a \in |D|} \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})$$

defined as the product of the natural immersions

$$\iota_{\mathbb{O}_{H_a}} \colon \mathbb{O}_{H_a} \cong G(\mathbb{C}[z_a]_{k_a})/G(\mathbb{C}[z_a]_{k_a})_{H_a} \ni [g] \longmapsto \mathrm{Ad}^*(g)(H_a) \in \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a}) \quad (a \in |D|).$$

We consider the open subset of $\prod_{a \in |D|} \mathbb{O}_{H_a}$ defined by

$$\left(\prod_{a\in |D|} \mathbb{O}_{H_a}\right)^{\mathrm{ir}} := \iota_{\mathbf{H}}^{-1} \left(\left(\prod_{a\in |D|} \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})\right)^{\mathrm{ir}} \right).$$

Under the identification $\mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a}) \cong \mathfrak{g}(\mathbb{C}[z_a]_{k_a})^*$ via the trace pairing, the injective immersion

$$\mu_{\mathbb{O}_{H_a}} = \iota_{\mathbb{O}_{H_a}} \colon \mathbb{O}_{H_a} \longmapsto \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})$$

is considered as the moment map with respect to the coadjoint action of $G(\mathbb{C}[z]_k)$. Furthermore, since the residue map $\mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a}) \ni X(z_a) \mapsto \underset{z_a=0}{\operatorname{res}} X(z_a) \in \mathfrak{g}$ is the dual map of the inclusion $\mathfrak{g} \hookrightarrow \mathfrak{g}(\mathbb{C}[z_a]_{k_a})$, the map

$$\mu_{\mathbb{O}_{H_a}\downarrow G} := \operatorname{res}_{z_a=0} \circ \iota_{\mathbb{O}_{H_a}} \colon \mathbb{O}_{H_a} \longrightarrow \mathfrak{g}$$

is the moment map with respect to the coadjoint action of G under the identification $\mathfrak{g} \cong \mathfrak{g}^*$ via the trace pairing. As the product of these moment maps, we obtain the moment map

$$\mu_{\mathbf{H}} \colon \prod_{a \in |D|} \mathbb{O}_{H_a} \longrightarrow \mathfrak{g}^* \\ (X_a)_{a \in |D|} \longmapsto \sum_{a \in |D|} \mu_{\mathbb{O}_{H_a} \downarrow G}(X_a)$$

with respect to the diagonal action of G. Also we denote the restriction of this moment map $\mu_{\mathbf{H}}$ on $\left(\prod_{a \in |D|} \mathbb{O}_{H_a}\right)^{\text{ir}}$ by $\mu_{\mathbf{H}}^{\text{ir}}$ which is still a moment map since $\left(\prod_{a \in |D|} \mathbb{O}_{H_a}\right)^{\text{ir}}$ is a open submanifold of $\prod_{a \in |D|} \mathbb{O}_{H_a}$ and closed under the *G*-action. Then under the injective immersion $\iota_{\mathbf{H}}$, we obtain the following identification,

$$\begin{cases} d_A = d + \sum_{a \in |D|} \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^i} \frac{dz}{z-a} \in \overline{\mathcal{M}}_D \middle| \begin{array}{c} d_A \text{ is irreducible,} \\ \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}} \in \mathbb{O}_{H_a} \text{ for all } a \in |D| \end{array} \right\} \\ = \left\{ \left(\sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}} \right)_{a \in |D|} \in \left(\prod_{a \in |D|} \mathbb{O}_{H_a} \right)^{\text{ir}} \middle| \sum_{a \in |D|} \sum_{z_a=0}^{res} \sum_{i=0}^{k_a} \frac{A_i^{(a)}}{(z-a)^{i+1}} = 0 \right\} \\ = \left\{ (\xi_a)_{a \in |D|} \in \left(\prod_{a \in |D|} \mathbb{O}_{H_a} \right)^{\text{ir}} \middle| \mu_{\mathbf{H}}^{\text{ir}}((\xi_a)_{a \in |D|}) = 0 \right\} = (\mu_{\mathbf{H}}^{\text{ir}})^{-1}(0). \end{cases}$$

Namely, the moduli space $\mathcal{M}_{\mathbf{H}}^{ir}$ of irreducible connections is regarded as the symplectic reduction

$$\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}} = G \setminus (\mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0).$$

Now let us look at the *G*-action on the level 0 set $(\mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0)$.

Lemma 7.14. Let us consider the decomposition $\mathfrak{g}^* = \mathfrak{z}^* \oplus \mathfrak{g}^*_{ss}$ induced by the decomposition $\mathfrak{g} = \mathfrak{z} \oplus \mathfrak{g}_{ss}$. Then we have $\operatorname{Im} \mu_{\mathbf{H}} \subset \mathfrak{g}^*_{ss}$.

Proof. As we saw in Proposition 4.1, for $X \in \mathfrak{g}_{ss}^{\perp} = \mathfrak{z}$ and $(g_a) \in \prod_{a \in |D|} G(\mathbb{C}[z_a]_{k_a})$, we have

$$\operatorname{tr}(X \cdot \sum_{a \in |D|} \operatorname{res}_{z=a}(\operatorname{Ad}(g_a)(H^{(a)})))$$

=
$$\operatorname{tr}\sum_{a \in |D|} \operatorname{res}_{z=a}(X \cdot \operatorname{Ad}(g_a)(H^{(a)})) = \operatorname{tr}\sum_{a \in |D|} \operatorname{res}_{z=a}(\operatorname{Ad}(g_a^{-1})(X)H^{(a)})$$

=
$$\operatorname{tr}\sum_{a \in |D|} \operatorname{res}_{z=a}(X \cdot H^{(a)}) = \operatorname{tr}(X \sum_{a \in |D|} \operatorname{res}_{z=a}(H^{(a)})) = 0,$$

since $\sum_{a \in |D|} \operatorname{res}_{z=a}(H^{(a)}) \in \mathfrak{g}_{ss}$. Thus $\sum_{a \in |D|} \operatorname{res}_{z=a}(\operatorname{Ad}(g_a)(H_a)) \in \mathfrak{g}_{ss}$. Then the correspondence between the residue map and the moment map $\mu_{\mathbf{H}}$ show the result.

Lemma 7.15. Let L be a locally compact topological group. Let us consider locally compact topological spaces V and W with continuous L-actions. Suppose that L acts on W properly and almost freely, and there exists a L-equivariant continuous map $\phi: V \to W$. Then the L-action on V is also proper and almost free. In particular if L-action on W is free, so is that on V.

Proof. Let us consider the stabilizer group $\operatorname{Stab}_L(v)$ of $v \in V$. If $g \in \operatorname{Stab}_L(v)$, then $g \cdot \phi(v) = \phi(g \cdot v) = \phi(v)$. Thus we have $\operatorname{Stab}_L(v) \subset \operatorname{Stab}_L(\phi(v))$. Then since the *L*-action on *W* is almost free, $\operatorname{Stab}_L(v)$ is a finite set.

Let us take a compact subset $K \subset V$ and consider $L_K := \{g \in L \mid g \cdot K \cap K \neq \emptyset\}$. Since ϕ is continuous, the image $\phi(K) \subset W$ is a compact subset. Let us take $g \in L_K$. Then we have $g \cdot \phi(K) \cap \phi(K) = \phi(g \cdot K) \cap \phi(K) \supset \phi(g \cdot K \cap K) \neq \emptyset$ since $g \cdot K \cap K \neq \emptyset$. Hence L_K is a subset of $L_{\phi(K)} := \{g \in G \mid g \cdot \phi(K) \cap \phi(K) \neq \emptyset\}$ which has the compact closure. Therefore L_K has the compact closure as well. **Proposition 7.16.** The moduli space $\mathcal{M}_{\mathbf{H}}^{ir}$ has a structure of complex symplectic orbifold of dimension

$$\mathcal{M}_{\mathbf{H}}^{ir} = 2 \dim Z - \operatorname{rig} (\mathbf{H}),$$

if it is nonempty.

Proof. Let us identify $\mathfrak{g}^* \cong \mathfrak{g}$ via the trace pairing and denote the dual map of the inclusion $\iota_{ss} \colon \mathfrak{g}_{ss} \hookrightarrow \mathfrak{g} = \mathfrak{z} \oplus \mathfrak{g}_{ss}$ by $\iota_{ss}^* \colon \mathfrak{g}^* \to \mathfrak{g}_{ss}^*$. Then we obtain the moment map

$$\iota_{\mathrm{ss}}^* \circ \mu_{\mathbf{H}}^{\mathrm{ir}} \colon \left(\prod_{a \in |D|} \mathbb{O}_{H_a} \right)^{\mathrm{ir}} \to \mathfrak{g}_{\mathrm{ss}}^*$$

of G/Z-action. Since we know that $\iota_{ss}^* \circ \mu_{\mathbf{H}}^{ir} = \mu_{\mathbf{H}}^{ir}$ by Lemma 7.14, we obtain

$$G \setminus (\mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0) = (G/Z) \setminus (\iota_{\mathrm{ss}}^* \circ \mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0).$$

Thus it suffices to show that the right hand side has a structure of complex symplectic orbifold.

Proposition 7.13 and Lemma 7.15 show that G/Z action on $\left(\prod_{a\in |D|} \mathbb{O}_{H_a}\right)^{\mathrm{ir}}$ is proper and almost free. Therefore since $\iota_{\mathrm{ss}}^* \circ \mu_{\mathbf{H}}^{\mathrm{ir}}$ is a moment map, the almost free-ness of G/Zaction assures that 0 is a regular value of $\iota_{\mathrm{ss}}^* \circ \mu_{\mathbf{H}}^{\mathrm{ir}}$. Namely, $(\iota_{\mathrm{ss}}^* \circ \mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0)$ becomes a submanifold of $\left(\prod_{a\in |D|} \mathbb{O}_{H_a}\right)^{\mathrm{ir}}$. Thus $(G/Z)\setminus(\iota_{\mathrm{ss}}^* \circ \mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0)$ is the quotient space of the complex manifold $(\iota_{\mathrm{ss}}^* \circ \mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0)$ by the proper and almost free action of G/Z, namely, it has the complex symplectic orbifold structure by Theorem 7.10.

Let us consider the dimension of $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}}$ under the assumption $(\iota_{\mathrm{ss}}^* \circ \mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0) \neq \emptyset$. Since the G/Z-action is almost free, we have

$$\dim \mathcal{M}_{\mathbf{H}}^{\mathrm{ir}} = \dim \left(G/Z \right) \setminus (\iota_{\mathrm{ss}}^* \circ \mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0) = \sum_{a \in |D|} \dim \mathbb{O}_{H^{(a)}} - 2 \dim G/Z$$
$$= \sum_{a \in |D|} \delta(H^{(a)}) - 2 \dim G + 2 \dim Z = 2 \dim Z - \mathrm{rig}(\mathbf{H})$$

as desired.

8 Triangular decompositions of truncated orbits

We shall recall the triangular decompositions of truncated orbits. This will play an important role to construct the deformation of truncated orbits in latter sections. Similar decomposition can also be found in [19] by Hiroe-Yamakawa, [15] by Hiroe, and [47] by Yamakawa.

Let us fix an unramified canonical form

$$H = \left(\frac{H_k}{z^k} + \dots + \frac{H_1}{z} + H_{\text{res}}\right) \frac{1}{z} \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k)$$

throughout this section.

8.1 A fiber bundle with the fiber $\mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}}$ and the truncated orbit

For a positive integer l with $0 \leq l \leq k$ and a closed subgroup $S \subset G(\mathbb{C}[z]_l)$, we denote the S-orbit of $\xi \in \mathfrak{g}(\mathbb{C}[z]_l)^*$ by

$$\mathbb{O}^S_{\xi} := \mathrm{Ad}^*(S)(\xi).$$

Now let us consider the orbit $\mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}}$ of H under the $L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}$ -action. Let us also consider the product manifold $G \times \mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}}$ with the L_{1} -action defined by $h \cdot (g, \xi) := (gh^{-1}, \mathrm{Ad}^{*}(h)(\xi))$ for $h \in L_{1}, g \in G$, and $\xi \in \mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}}$. Then the quotient space

$$G \times_{L_1} \mathbb{O}_H^{L_1 \ltimes G(\mathbb{C}[z]_k)_1} := L_1 \setminus (G \times \mathbb{O}_H^{L_1 \ltimes G(\mathbb{C}[z]_k)_1})$$

is the fiber bundle on G/L_1 with the fiber $\mathbb{O}_H^{L_1 \ltimes G(\mathbb{C}[z]_k)_1}$ which is associated with the principal L_1 -bundle $G \to G/L_1$. Then we shall show that the total space $G \times_{L_1} \mathbb{O}_H^{L_1 \ltimes G(\mathbb{C}[z]_k)_1}$ is isomorphic to the truncated orbit \mathbb{O}_H .

Consider the multiplication map $\tilde{\psi}: G \times (L_1 \ltimes G(\mathbb{C}[z]_k)_1) \ni (g, (h, f)) \mapsto ghf \in G(\mathbb{C}[z]_k)$. This is obviously $\operatorname{Stab}_{L_1 \ltimes G(\mathbb{C}[z]_k)_1}(H)$ -equivariant under the multiplications from the right. Therefore it induces the map

$$\overline{\psi} \colon G \times \mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}} \ni (g, \xi) \longmapsto \mathrm{Ad}^{*}(g)(\xi) \in \mathbb{O}_{H}$$

Moreover since this $\overline{\psi}$ is invariant under the L_1 -action, it factors through the map

$$\psi \colon G \times_{L_1} \mathbb{O}_H^{L_1 \ltimes G(\mathbb{C}[z]_k)_1} \ni [(g,\xi)] \longmapsto \mathrm{Ad}^*(g)(\xi) \in \mathbb{O}_H$$

which does not depend on the choice of representatives of $[(g,\xi)]$. Namely, we obtain the commutative diagram

Here vertical arrows are natural projections.

Proposition 8.1 (cf. Proposition 4.9 in [15], and also Proposition 2.12 in [19]). The above map $I = \mathcal{L}(\mathbb{C}[1])$

$$\psi \colon G \times_{L_1} \mathbb{O}_H^{L_1 \ltimes G(\mathbb{C}[z]_k)_1} \xrightarrow{\sim} \mathbb{O}_H$$

is an isomorphism as complex manifolds.

Proof. Obviously the map $\overline{\psi} \colon G \times \mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}} \to \mathbb{O}_{H}$ is surjective, and thus ψ is surjective as well. Then let us see the injectivity. Suppose that $(g_{1},\xi_{1}), (g_{2},\xi_{2}) \in G \times \mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}}$ are sent to the same image by $\overline{\psi}$, i.e., $\operatorname{Ad}^{*}(g_{1})(\xi_{1}) = \operatorname{Ad}^{*}(g_{2})(\xi_{2})$. Since there exist $(h_{i},f_{i}) \in L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}, i = 1, 2$, such that $\operatorname{Ad}^{*}(h_{i}f_{i})(H) = \xi_{i}$, we obtain

$$H = \mathrm{Ad}^*((h_1 f_1)^{-1}) \circ \mathrm{Ad}^*(g_1^{-1}) \circ \mathrm{Ad}^*(g_2) \circ \mathrm{Ad}^*(h_2 f_2)(H).$$

This implies that

$$(h_1^{-1}g_1^{-1}g_2h_2, \operatorname{Ad}(h_2^{-1}g_2^{-1}g_1h_1)(f_1^{-1}) \cdot f_2) \in G \ltimes G(\mathbb{C}[z]_k)_1$$

stabilizes H. Then Proposition 3.4 shows that $h_1^{-1}g_1^{-1}g_2h_2 \in \operatorname{Stab}_G(H) \subset L_1$. Thus we have $h := g_1^{-1}g_2 \in L_1$ and $(g_2, \xi_2) = (g_1h, \operatorname{Ad}^*(h^{-1})(\xi_1))$ which show the injectivity of ψ .

In the remaining of this section, we shall give a proof of the following triangular decomposition theorem of the fiber $\mathbb{O}_{H}^{L_1 \ltimes G(\mathbb{C}[z]_k)_1}$.

Theorem 8.2 (cf. Theorem 3.6 [19], Proposition 4.15 [15], and Corollary 3.2 [47]). There exists an isomorphism

$$\Psi \colon \left(\prod_{l=1}^{k} \left(N_{l,l+1}(\mathbb{C}[z]_{l})_{1} \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}) \right) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_{1}} \longrightarrow \mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{2}}$$

as complex manifolds.

Here for the notations $N_{l,l+1}$ and $\mathfrak{n}_{l,l+1}$, see Section 2.2.

8.2 Decompositions of $G(\mathbb{C}[z]_l)_1$

Recall that $G(\mathbb{C}[z]_l)_1$ is a connected and simply connected nilpotent Lie group and moreover it has the grading with respect to the degrees of z^m . Therefore $G(\mathbb{C}[z]_l)_1$ admits the following decomposition property.

Proposition 8.3. Let \mathfrak{h}_1 , \mathfrak{h}_2 , and \mathfrak{l} be Lie subalgebras of \mathfrak{g} , which satisfy $\mathfrak{l} = \mathfrak{h}_1 \oplus \mathfrak{h}_2$. Let us define the subgroups of $G(\mathbb{C}[z]_l)_1$ by $H_i(\mathbb{C}[z]_l)_1 := \exp(\mathfrak{h}_i(\mathbb{C}[z]_l)_1)$ for i = 1, 2, and $L(\mathbb{C}[z]_l)_1 := \exp(\mathfrak{l}(\mathbb{C}[z]_l)_1)$. Then we obtain the following decomposition,

$$L(\mathbb{C}[z]_{l})_{1} = H_{1}(\mathbb{C}[z]_{l})_{1} \cdot H_{2}(\mathbb{C}[z]_{l})_{1}.$$

Proof. Take $g \in L(\mathbb{C}[z]_l)_1$ and write $g = e^{X_l z^l} \cdots e^{X_2 z^2} e^{X_1 z}$ by $X_i \in \mathfrak{l}, i = 1, 2, \ldots, l$ according to Proposition 3.2. First let us decompose $X_1 = H_1^{(1)} + H_1^{(2)}$ by $H_1^{(i)} \in \mathfrak{h}_i$. Then the Campbell-Baker-Hausdorff formula shows that

$$e^{X_1 z} = e^{(H_1^{(1)} + H_1^{(2)})z} \equiv e^{H_1^{(1)} z} e^{H_1^{(2)} z} \pmod{z^2}.$$

Thus we can write

$$g \cdot (e^{H_1^{(1)}z} e^{H_1^{(2)}z})^{-1} = e^{X_l' z^l} \cdots e^{X_2' z^2}.$$

Then we decompose $X'_2 = H_2^{(1)} + H_2^{(2)}$ by $H_2^{(i)} \in \mathfrak{h}_i$ and we have

$$e^{X'_2 z^2} = e^{(H_2^{(1)} + H_2^{(2)})z^2} \equiv e^{H_2^{(1)} z^2} e^{H_2^{(2)} z^2} \pmod{z^3}$$

as well. Since $e^{H_1^{(i)}z}$ and $e^{H_2^{(i')}z^2}$ are commutative modulo z^3 by the Campbell-Baker-Hausdorff formula, we obtain

$$g \cdot (e^{H_2^{(1)}z^2} e^{H_1^{(1)}z} e^{H_2^{(2)}z^2} e^{H_1^{(2)}z})^{-1} = e^{X_l''z^l} \cdots e^{X_3''z^3}$$

Repeating this procedure, we obtain $h_i \in H_i(\mathbb{C}[z]_l)_1$, i = 1, 2, such that $g = h_1 h_2$.

Let us notice that $H_1(\mathbb{C}[z]_l)_1 \cap H_2(\mathbb{C}[z]_l)_1 = \{e\}$ since $\mathfrak{h}_1(\mathbb{C}[z]_l)_1 \cap \mathfrak{h}_2(\mathbb{C}[z]_l)_1 = \{0\}$ and the exponential map is an analytic diffeomorphism. Thus the description $g = h_1h_2$ is unique.

8.3 Filtration of $\mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}}$

Let us consider the sequence of subgroups

$$L_1 \ltimes L_1(\mathbb{C}[z]_k)_1 \subset L_1 \ltimes L_2(\mathbb{C}[z]_k)_1 \subset \cdots \subset L_1 \ltimes L_{k+1}(\mathbb{C}[z]_k)_1 = L_1 \ltimes G(\mathbb{C}[z]_k)_1$$

of $L_1 \ltimes G(\mathbb{C}[z]_k)_1$, and the induced filtration of $\mathbb{O}_H^{L_1 \ltimes G(\mathbb{C}[z]_k)_1}$,

$$\mathbb{O}_{H}^{L_{1}\ltimes L_{1}(\mathbb{C}[z]_{k})_{1}} \subset \mathbb{O}_{H}^{L_{1}\ltimes L_{2}(\mathbb{C}[z]_{k})_{1}} \subset \cdots \subset \mathbb{O}_{H}^{L_{1}\ltimes L_{k+1}(\mathbb{C}[z]_{k})_{1}} = \mathbb{O}_{H}^{L_{1}\ltimes G(\mathbb{C}[z]_{k})_{1}}.$$

Here we notice that Lemma 3.3 in [47] or computations carried out in the proof of Proposition 3.4 shows that

$$\mathbb{O}_{H}^{L_{1} \ltimes L_{1}(\mathbb{C}[z]_{k})_{1}} = H_{\mathrm{irr}} + \mathrm{Ad}^{*}(L_{1})(H_{\mathrm{res}}) \cong \mathbb{O}_{H_{\mathrm{res}}}^{L_{1}}$$

see also Lemma 4.10 in [15].

Recall that for a pair of positive integers $0 \le l < l' \le k$, there is the natural projection map

$$\pi_{l',l} \colon \mathbb{C}[z]_{l'} \to \mathbb{C}[z]_{l'}/(\mathfrak{m}_z^{(l')})^{l+1} = \mathbb{C}[z]_l$$

where $\mathfrak{m}_{z}^{(l')} = \langle z \rangle_{\mathbb{C}[z]_{l'}}$ is the unique maximal ideal of $\mathbb{C}[z]_{l'}$. Then we obtain exact sequences of Lie algebras

$$0 \to \mathfrak{h}((\mathfrak{m}_{z}^{(l')})^{l+1}) \to \mathfrak{h}(\mathbb{C}[z]_{l'}) \xrightarrow{\pi_{l',l}} \mathfrak{h}(\mathbb{C}[z]_{l}) \to 0$$

and nilpotent Lie groups

$$1 \to \exp\left(\mathfrak{h}((\mathfrak{m}_{z}^{(l')})^{l+1})\right) \to H(\mathbb{C}[z]_{l'})_{1} \xrightarrow{\pi_{l',l}} H(\mathbb{C}[z]_{l})_{1} \to 1$$

for a Lie subgroup $H \subset G$ and its Lie algebra \mathfrak{h} .

For a positive integer $0 \leq l \leq k$, let us define

$$H^{[l]} := \left(\frac{H_l}{z^l} + \dots + H_{\text{res}}\right) \frac{1}{z} \in \mathfrak{g}(\mathbb{C}[z^{-1}]_l).$$

Proposition 8.4. Let us take a pair of positive integers l, l' with $0 \leq l \leq l' \leq k$. Then the projection map $\pi_{l',l}: L_1 \ltimes L_{l+1}(\mathbb{C}[z]_{l'})_1 \to L_1 \ltimes L_{l+1}(\mathbb{C}[z]_l)_1$ induces the isomorphism

$$\pi^{\mathbb{O}}_{l',l} \colon \mathbb{O}_{H^{[l']}}^{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_{l'})_1} \xrightarrow{\sim} \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_{l})_1}.$$

Proof. From Proposition 3.4, we have

 $\operatorname{Stab}_{L_{l+1}(\mathbb{C}[z]_{l'})_1}(H^{[l']}) = \left\{ e^{X_{l'}z^{l'}} \cdots e^{X_2z^2} e^{X_1z} \in G(\mathbb{C}[z]_{l'})_1 \, \middle| \, X_i \in \mathfrak{l}_i \cap \mathfrak{l}_{l+1}, \ i = 1, \dots, l' \right\},$ $\operatorname{Stab}_{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_{l'})_1}(H^{[l']}) = \operatorname{Stab}_{L_1}(H_{\operatorname{res}}) \ltimes \operatorname{Stab}_{L_{l+1}(\mathbb{C}[z]_{l'})_1}(H^{[l']}).$

Here we note that the above condition $X_i \in l_i \cap l_{l+1}$, $i = 1, \ldots, l'$ is equivalent to

$$X_i \in \begin{cases} \mathfrak{l}_i & i = 1, 2, \dots, l, \\ \mathfrak{l}_{l+1} & i = l+1, \dots, l' \end{cases}$$

Therefore since we moreover have

$$\operatorname{Stab}_{L_{l+1}(\mathbb{C}[z]_{l})_{1}}(H^{[l]}) = \left\{ e^{X_{l}z^{l}} \cdots e^{X_{2}z^{2}} e^{X_{1}z} \in G(\mathbb{C}[z]_{l})_{1} \, \middle| \, X_{i} \in \mathfrak{l}_{i}, \ i = 1, \dots, l \right\},$$

$$\operatorname{Stab}_{L_{1} \ltimes L_{l+1}(\mathbb{C}[z]_{l})_{1}}(H^{[l]}) = \operatorname{Stab}_{L_{1}}(H_{\operatorname{res}}) \ltimes \operatorname{Stab}_{L_{l+1}(\mathbb{C}[z]_{l})_{1}}(H^{[l]}),$$

we obtain the commutative diagram

whose vertical arrows are natural inclusions and horizontal sequences are exact. Then by a diagram chase shows $\pi_{l,l'}$ induces the bijective holomorphic map

$$\mathbb{O}_{H^{[l']}}^{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_{l'})_1} = L_{l+1}(\mathbb{C}[z]_{l'})_1 / \operatorname{Stab}_{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_{l'})_1}(H^{[l']}) \xrightarrow{\pi_{l',l}} L_1 \ltimes L_{l+1}(\mathbb{C}[z]_l)_1 / \operatorname{Stab}_{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_l)_1}(H^{[l]}) = \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_{l'})_1}.$$

By the isomorphisms in this proposition, the above filtration is replaced by

$$\mathbb{O}_{H_{\mathrm{res}}}^{L_1 \ltimes L_1(\mathbb{C}[z]_0)_1} \subset \mathbb{O}_{H^{[1]}}^{L_1 \ltimes L_2(\mathbb{C}[z]_1)_1} \subset \cdots \subset \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_l)_1} \subset \cdots \subset \mathbb{O}_{H}^{L_1 \ltimes G(\mathbb{C}[z]_k)_1}$$

8.4 LU-type decomposition and Levi-type decomposition

For each pair $1 \leqslant l < l' \leqslant k$ of positive integers, Proposition 8.3 gives us the Levi-type decomposition

$$P_{l,l'}(\mathbb{C}[z]_l)_1 = L_l(\mathbb{C}[z]_l)_1 \ltimes U_{l,l'}(\mathbb{C}[z]_l)_1$$

and the LU-type decomposition

$$L_{l'}(\mathbb{C}[z]_l)_1 = N_{l,l'}(\mathbb{C}[z]_l)_1 \cdot P_{l,l'}(\mathbb{C}[z]_l)_1.$$

Since L_1 normalizes $N_{l,l'}(\mathbb{C}[z]_l)_1$, we further obtain

$$L_{1} \ltimes L_{l'}(\mathbb{C}[z]_{l})_{1} = N_{l,l'}(\mathbb{C}[z]_{l})_{1} \cdot (L_{1} \ltimes P_{l,l'}(\mathbb{C}[z]_{l})_{1})$$

= $N_{l,l'}(\mathbb{C}[z]_{l})_{1} \cdot ((L_{1} \ltimes L_{l}(\mathbb{C}[z]_{l})_{1}) \ltimes U_{l,l'}(\mathbb{C}[z]_{l})_{1}).$

Proposition 8.5. Let us take a pair $1 \le l \le l' \le k$ of positive integers. The multiplication map

$$\overline{\phi} \colon N_{l,l'}(\mathbb{C}[z]_l)_1 \times (L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1) \longrightarrow L_1 \ltimes L_{l'}(\mathbb{C}[z]_l)_1$$

induces the isomorphism

$$\phi \colon N_{l,l'}(\mathbb{C}[z]_l)_1 \times \mathbb{O}_{H^{[l]}}^{L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1} \longrightarrow \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_{l'}(\mathbb{C}[z]_l)_1}$$

as complex manifolds.

Proof. Let $\pi_1: L_1 \ltimes L_{l'}(\mathbb{C}[z]_l)_1 \to \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_{l'}(\mathbb{C}[z]_l)_1}, \pi_2: L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1 \to \mathbb{O}_{H^{[l]}}^{L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1}$ be natural quotient maps. Since we obviously have

$$\operatorname{Stab}_{L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1}(H^{[l]}) \subset \operatorname{Stab}_{L_1 \ltimes L_{l'}(\mathbb{C}[z]_l)_1}(H^{[l]}),$$

the composition map $\pi_1 \circ \overline{\phi} \colon N_{l,l'}(\mathbb{C}[z]_l)_1 \times (L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1) \to \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_{l'}(\mathbb{C}[z]_l)_1}$ factors through the quotient map $\mathrm{id}_{N_{l,l'}(\mathbb{C}[z]_l)_1} \times \pi_2 \colon N_{l,l'}(\mathbb{C}[z]_l)_1 \times (L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1) \to N_{l,l'}(\mathbb{C}[z]_l)_1 \times \mathbb{O}_{H^{[l]}}^{L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1}$. Namely, we obtain the commutative diagram

Since $\overline{\phi}$ is isomorphism, ϕ is surjective. Let us notice that Proposition 3.4 shows that $\operatorname{Stab}_{L_1 \ltimes L_{l'}(\mathbb{C}[z]_l)_1}(H^{[l]}) \subset L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1$. Thus the above inclusion becomes the equation $\operatorname{Stab}_{L_1 \ltimes P_{l,l'}(\mathbb{C}[z]_l)_1}(H^{[l]}) = \operatorname{Stab}_{L_1 \ltimes L_{l'}(\mathbb{C}[z]_l)_1}(H^{[l]})$ which shows the injectivity of ϕ by a diagram chase.

Let us focus on the case l' = l + 1 and consider the multiplication map

$$\chi \colon (L_1 \ltimes L_l(\mathbb{C}[z]_l)_1) \times U_{l,l+1}(\mathbb{C}[z]_l)_1 \longrightarrow L_1 \ltimes P_{l,l+1}(\mathbb{C}[z]_l)_1$$

Also we define the map

$$\begin{array}{cccc} a_{H^{[l]}} \colon & (L_1 \ltimes L_l(\mathbb{C}[z]_l)_1) \times U_{l,l+1}(\mathbb{C}[z]_l)_1 & \longrightarrow & (L_1 \ltimes L_l(\mathbb{C}[z]_l)_1) \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}) \\ & (g,u) & \longmapsto & (g, \operatorname{Ad}^*(gu)(H^{[l]}) - \operatorname{ad}^*(g)(H^{[l]})) \end{array}$$

Lemma 8.6. The above map $a_{H^{[l]}}$ is an isomorphism as complex manifolds.

Proof. First we show that $a_{H^{[l]}}$ is well-defined. Let us recall that for $u = e^{z^l X_l} \cdots e^{zX_1} \in U_{l,l+1}(\mathbb{C}[z]_l)_1$ and $B = \sum_{i=0}^l B_i z^{-i-1} \in \mathfrak{l}_l(\mathbb{C}[z^{-1}]_l)$, we have $\operatorname{Ad}^*(u)(B) = \sum_{i=0}^l C_i z^{-i-1}$ with

$$C_{i} = B_{i} + \sum_{\substack{m_{1}, m_{2}, \dots, m_{l} \ge 0 \\ (m_{1}, m_{2}, \dots, m_{l}) \neq \mathbf{0}}} \frac{\operatorname{ad}(X_{l})^{m_{l}} \circ \dots \circ \operatorname{ad}(X_{2})^{m_{2}} \circ \operatorname{ad}(X_{1})^{m_{1}}(B_{i+m_{1}+2m_{2}+\dots+lm_{l}})}{m_{1}!m_{2}! \cdots m_{l}!}.$$

Then we notice that that the second term in the right hand side is contained in $\mathfrak{u}_{l,l+1}$, and $C_l = B_l$. Thus $a_{H^{[l]}}$ is well-defined.

Next show that $a_{H^{[l]}}$ is injective. Suppose that (g, u) and (g', u') have the same image by $a_{H^{[l]}}$. Then obviously g = g' and also $\operatorname{Ad}^*(gu)(H^{[l]}) = \operatorname{Ad}^*(gu')(H^{[l]})$. Thus $u^{-1}u' \in U_{l,l+1}(\mathbb{C}[z]_l)_1$ belongs to $\operatorname{Stab}_{L_{l+1}(\mathbb{C}[z]_l)_1}(H^{[l]})$ which means that $u^{-1}u' = e$ by Proposition 3.4, i.e., u = u' as desired.

Finally let us show that $a_{H^{[l]}}$ is surjective. Recall that $\mathfrak{u}_{l,l+1} = \operatorname{Ker} (\operatorname{ad}(H_m)|_{\mathfrak{u}_{l,l+1}}) \oplus \operatorname{ad}(H_m)(\mathfrak{u}_{l,l+1}) = \operatorname{ad}(H_m)(\mathfrak{u}_{l,l+1})$ for $m = 0, 1, \ldots, l$, since $\operatorname{Ker} (\operatorname{ad}(H_m)|_{\mathfrak{u}_{l,l+1}}) = 0$. Then we can see that the above equations for given $C_l = B_l$ and $C_i \in B_i + \mathfrak{u}_{l,l+1}$ for $i = 0, 1, \ldots, l-1$ can be solved inductively from i = l-1 to i = 0 and we find $u \in U_{l,l+1}(\mathbb{C}[z]_l)_1$ such that $\operatorname{Ad}^*(u)(B) = \sum_{i=0}^l C_i z^{-i-1}$. In particular when $B \in \mathbb{O}_H^{L_1 \ltimes L_l(\mathbb{C}[z]_l)_1}$, there exists $g \in L_1 \ltimes L_l(\mathbb{C}[z]_l)_1$ such that $\operatorname{Ad}^*(g)(H^{[l]}) = B$. Thus in this case we have

$$\operatorname{Ad}^*(g(g^{-1}ug))(H^{[l]}) = \operatorname{Ad}^*(ug)(H^{[l]}) = \operatorname{Ad}^*(u)(B) = C$$

This shows that $a_{H^{[l]}}$ is surjective.

Proposition 8.7. The map

$$\begin{array}{cccc} X \colon & \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_l(\mathbb{C}[z]_l)_1} \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}) & \longrightarrow & \mathbb{O}_{H^{[l]}}^{L_1 \ltimes P_{l,l+1}(\mathbb{C}[z]_l)_1} \\ & & (\eta, \xi) & \longmapsto & \eta + \xi \end{array}$$

is an isomorphism of complex manifolds.

Proof. One can check that the following diagram is commutative,

$$\begin{array}{cccc} (L_1 \ltimes L_l(\mathbb{C}[z]_l)_1) \times U_{l,l+1}(\mathbb{C}[z]_l)_1 & \xrightarrow{\chi} & L_1 \ltimes P_{l,l+1}(\mathbb{C}[z]_l)_1 \\ & \downarrow^{a_H[l]} & & \downarrow^{a_H[l]} & & \downarrow^{\pi_P} \\ (L_1 \ltimes L_l(\mathbb{C}[z]_l)_1) \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}) & & \downarrow^{\pi_P} \\ & \downarrow^{\pi_L \times \mathrm{id}} & & \downarrow^{\pi_P} \\ \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_l(\mathbb{C}[z]_l)_1} \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}) & \xrightarrow{X} & \mathbb{O}_{H^{[l]}}^{L_1 \ltimes P_{l,l+1}(\mathbb{C}[z]_l)_1} \end{array}$$

Here $\pi_L \colon L_1 \ltimes L_l(\mathbb{C}[z]_l)_1 \to \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_l(\mathbb{C}[z]_l)_1}$ and $\pi_P \colon L_1 \ltimes P_{l,l+1}(\mathbb{C}[z]_l)_1 \to \mathbb{O}_{H^{[l]}}^{L_1 \ltimes P_{l,l+1}(\mathbb{C}[z]_l)_1}$ are natural projections. Since χ and $a_{H^{[l]}}$ are isomorphisms, and $\pi_L \times \text{id}$ and π_P are surjective, it follows that X is surjective.

Let us equip the action of $L_1 \ltimes L_l(\mathbb{C}[z]_l)_1$ with $(L_1 \ltimes L_l(\mathbb{C}[z]_l)_1) \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1})$ by the multiplication on the first component. Then the isomorphism $a_{H^{[l]}}$ becomes $(L_1 \ltimes L_l(\mathbb{C}[z]_l)_1)_H$ -equivariant and also the isomorphism $\chi \circ a_{H^{[l]}}^{-1}$ becomes $(L_1 \ltimes L_l(\mathbb{C}[z]_l)_1)_H$ -equivariant. Therefore since $(L_1 \ltimes L_l(\mathbb{C}[z]_l)_1)_H = (L_1 \ltimes P_{l,l+1}(\mathbb{C}[z]_l)_1)_H$, we can see that X is injective.

8.5 A proof of Theorem 8.2

Now we are ready to give a proof of Theorem 8.2.

Proof of Theorem 8.2. As a combination of Propositions 8.5 and 8.7, we obtain isomorphisms

$$\overline{\psi}_l \colon N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}) \times \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_l(\mathbb{C}[z]_l)_1} \longrightarrow \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_l)_1}$$

for l = 1, 2, ..., k. Since $\mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_l(\mathbb{C}[z]_l)_1} \cong \mathbb{O}_{H^{[l-1]}}^{L_1 \ltimes L_l(\mathbb{C}[z]_{l-1})_1}$ by Proposition 8.4, above isomorphisms moreover give us following isomorphisms

$$\psi_l \colon (N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1})) \times \mathbb{O}_{H^{[l-1]}}^{L_1 \ltimes L_l(\mathbb{C}[z]_{l-1})_1} \longrightarrow \mathbb{O}_{H^{[l]}}^{L_1 \ltimes L_{l+1}(\mathbb{C}[z]_l)_1}$$

for l = 1, 2, ..., k. Therefore as the combination of these isomorphisms ψ_l we obtain the desired isomorphism

$$\Psi \colon \left(\prod_{l=1}^{k} \left(N_{l,l+1}(\mathbb{C}[z]_{l})_{1} \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}) \right) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_{1}} \longrightarrow \mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}}$$

by recalling $\mathbb{O}_{H^{[0]}}^{L_1 \ltimes L_1(\mathbb{C}[z]_0)_1} = \mathbb{O}_{H_{\mathrm{res}}}^{L_1}$.

59

8.6 Triangular decomposition of \mathbb{O}_H

As a consequence of Theorem 8.2, we moreover obtain the triangular decomposition of the truncated orbit \mathbb{O}_H as follows.

Define an L_1 -action on $\left(\prod_{l=1}^k (N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}))\right) \times \mathbb{O}_{H_{\text{res}}}^{L_1}$ as follows,

$$g \cdot ((n_{l,l+1},\xi_l)_{l=1,\dots,k},\xi) := ((\mathrm{Ad}(g)(n_l),\mathrm{Ad}^*(g)(\xi_l))_{l=1,2,\dots,k},\mathrm{Ad}^*(g)(\xi))$$

for $g \in L_1$ and $((n_l, \xi_l)_{l=1,\dots,k}, \xi) \in \left(\prod_{l=1}^k (N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}))\right) \times \mathbb{O}_{H_{\text{res}}}^{L_1}$. Then

$$\mathrm{id}_G \times \Psi \colon G \times \left(\left(\prod_{l=1}^k (N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1})) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_1} \right) \longrightarrow G \times \mathbb{O}_H^{L_1 \ltimes G(\mathbb{C}[z]_k)_1}$$

is L_1 -equivariant isomorphism, and therefore Proposition 8.1 shows that the map

$$G \times \left(\left(\prod_{l=1}^{k} (N_{l,l+1}(\mathbb{C}[z]_{l})_{1} \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1})) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_{1}} \right) \xrightarrow{\mathrm{id}_{G} \times \Psi} G \times \mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}} \xrightarrow{\psi} \mathbb{O}_{H}$$

factors through the isomorphism

$$L_1 \setminus \left(G \times \left(\left(\prod_{l=1}^k (N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1})) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_1} \right) \right) \cong \mathbb{O}_H.$$

According to this triangular decomposition of \mathbb{O}_H , let us give a description of the moment map

$$\mu_{\mathbb{O}_H \downarrow G} \colon \mathbb{O}_H \longrightarrow \mathfrak{g}$$

defined in Section 7.4, which will play an important role in the latter sections. Let us take $\Xi \in \mathbb{O}_H$ and choose its representative

$$(g, (n_l, \nu_l)_{l=1,...,k}, \eta) \in G \times \prod_{l=1}^k \left(N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_1}.$$

Then under the injective immersion $\iota_{\mathbb{O}_H} \colon \mathbb{O}_H \hookrightarrow \mathfrak{g}(\mathbb{C}[z^{-1}]_k)$, the image $\iota_{\mathbb{O}_H}(\Xi)$ is computed by the following steps. Step 1. Let us define

Step 1. Let us define

$$\iota_{\mathbb{O}_H}^{(1)}(\Xi) := \mathrm{Ad}^*(n_1)(\nu_1 + \eta) \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k).$$

Step l for $1 < l \leq k$. Let us define

$$\iota_{\mathbb{O}_H}^{(l)}(\Xi) := \mathrm{Ad}^*(n_l)(\nu_l + \iota_{\mathbb{O}_H}^{(l-1)}(\Xi)) \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k).$$

Then we obtain

$$\iota_{\mathbb{O}_H}(\Xi) = \mathrm{Ad}^*(g)(\iota_{\mathbb{O}_H}^{(k)}(\Xi)) \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k).$$

Therefore we finally obtain the inductive formula for the moment map,

$$\mu_{\mathbb{O}_H \downarrow G}(\Xi) = \operatorname{res}_{z=0} \circ \operatorname{Ad}^*(g)(\iota_{\mathbb{O}_H}^{(k)}(\Xi)) \in \mathfrak{g}.$$

Another representative of Ξ leads to the same result. Indeed for another representative there exists $h \in L_1$ and it is written by $(gh^{-1}, (\mathrm{Ad}(h)(n_l), \mathrm{Ad}^*(h)(\nu_l))_{l=1,\dots,k}, \mathrm{Ad}^*(h)(\eta))$. Then we can also inductively define

$$\begin{aligned} \widetilde{\iota}_{\mathbb{O}_{H}}^{(1)}(\Xi) &:= \mathrm{Ad}^{*}(\mathrm{Ad}(h)(n_{1}))(\mathrm{Ad}^{*}(h)(\nu_{1}) + \mathrm{Ad}^{*}(h)(\eta)) \\ &= \mathrm{Ad}^{*}(\mathrm{Ad}(h)(n_{1})h)(\nu_{1} + \eta) \\ &= \mathrm{Ad}^{*}(h \cdot n_{1})(\nu_{1} + \eta), \end{aligned}$$

$$\begin{aligned} \hat{\iota}_{\mathbb{O}_{H}}^{(2)}(\Xi) &:= \mathrm{Ad}^{*}(\mathrm{Ad}(h)(n_{2}))(\mathrm{Ad}^{*}(h)(\nu_{2}) + \hat{\iota}_{\mathbb{O}_{H}}^{(1)}(\Xi)) \\ &= \mathrm{Ad}^{*}(\mathrm{Ad}(h)(n_{2}))(\mathrm{Ad}^{*}(h)(\nu_{2}) + \mathrm{Ad}^{*}(h \cdot n_{1})(\nu_{1} + \eta)) \\ &= \mathrm{Ad}^{*}(\mathrm{Ad}(h)(n_{2}) \cdot h)(\nu_{2} + \mathrm{Ad}^{*}(n_{1})(\nu_{1} + \eta)) \\ &= \mathrm{Ad}^{*}(h \cdot n_{2})(\nu_{2} + \iota_{\mathbb{O}_{H}}^{(1)}(\Xi)), \end{aligned}$$

$$\begin{aligned} \hat{\iota}_{\mathbb{O}_{H}}^{(l)}(\Xi) &:= \mathrm{Ad}^{*}(\mathrm{Ad}(h)(n_{l}))(\mathrm{Ad}^{*}(h)(\nu_{l}) + \hat{\iota}_{\mathbb{O}_{H}}^{(l-1)}(\Xi)) \\ &= \mathrm{Ad}^{*}(\mathrm{Ad}(h)(n_{l}))(\mathrm{Ad}^{*}(h)(\nu_{l}) + \mathrm{Ad}^{*}(h \cdot n_{l-1})(\nu_{l-2} + \iota_{\mathbb{O}_{H}}^{(l-2)}(\Xi))) \\ &= \mathrm{Ad}^{*}(h \cdot n_{l})(\nu_{l} + \iota_{\mathbb{O}_{H}}^{(l-1)}(\Xi)) \end{aligned}$$

for $l \leq k$, and finally we obtain

$$\operatorname{Ad}^{*}(gh^{-1})(\widetilde{\iota}_{\mathbb{O}_{H}}^{(l)}(\Xi)) = \operatorname{Ad}^{*}(gh^{-1})(\operatorname{Ad}^{*}(h \cdot n_{k})(\nu_{k} + \iota_{\mathbb{O}_{H}}^{(k-1)}(\Xi)))$$
$$= \operatorname{Ad}^{*}(g)(\operatorname{Ad}^{*}(n_{k})(\nu_{k} + \iota_{\mathbb{O}_{H}}^{(k-1)}(\Xi)))$$
$$= \operatorname{Ad}^{*}(g)(\iota_{\mathbb{O}_{H}}^{(l)}(\Xi)) = \iota_{\mathbb{O}_{H}}(\Xi)$$

as well.

8.7 A Zariski open subset of the truncated orbit of a semisimple canonical form

In this subsection, we particularly consider a semisimple canonical form

$$H dz = \left(\frac{H_k}{z^k} + \dots + \frac{H_1}{z} + H_0\right) \frac{dz}{z} \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k) dz,$$

 $H_i \in \mathfrak{t}, i = 0, 1, \dots, k$. Then we have $\operatorname{Stab}_G(H) = L_0$ and $\operatorname{Stab}_{G(\mathbb{C}[z]_k)}(H) = L_0 \ltimes \operatorname{Stab}_{G(\mathbb{C}[z]_k)_1}(H)$. Let us consider the Zariski open subset of G defined by

$$G_0 := N_0 \cdot P_0$$

which is the open cell of the Bruhat decomposition of G with respect to P_0 . Then for a non negative integer j, we can also define the Zariski open subset of $G(\mathbb{C}[z]_j)$ by

$$G_0(\mathbb{C}[z]_j) := \{ g_0 \cdot g_1 \in G \ltimes G(\mathbb{C}[z]_j)_1 \mid g_0 \in G_0 \}.$$

Then we can consider the Zariski open subset

$$\mathbb{O}_{H}^{G_{0}(\mathbb{C}[z]_{k})} := \{ \mathrm{Ad}^{*}(g)(H) \in \mathfrak{g}(\mathbb{C}[z]_{k})^{*} \mid g \in G_{0}(\mathbb{C}[z]_{k}) \}$$

of the truncated orbit \mathbb{O}_H and we shall show an analogue of Theorem 8.2 for $\mathbb{O}_H^{G_0(\mathbb{C}[z]_k)}$.

Recalling that $G_0(\mathbb{C}[z]_k)$ contains $\operatorname{Stab}_{G(\mathbb{C}[z]_k)}(H)$ and moreover $\operatorname{Stab}_{G(\mathbb{C}[z]_k)}(H)$ acts on $G_0(\mathbb{C}[z]_k)$ via the multiplication from the right, we have

$$\mathbb{O}_{H}^{G_{0}(\mathbb{C}[z]_{k})} \cong G_{0}(\mathbb{C}[z]_{k})/\mathrm{Stab}_{G(\mathbb{C}[z]_{k})}(H).$$

Proposition 8.8. Let *j* be a non negative integer. We have the following decomposition

$$G_0(\mathbb{C}[z]_j) = N_k(\mathbb{C}[z]_j) \cdot N_{k-1,k}(\mathbb{C}[z]_j) \cdot \cdots \cdot N_{0,1}(\mathbb{C}[z]_j) \cdot L_0(\mathbb{C}[z]_j) \\ \cdot U_{0,1}(\mathbb{C}[z]_j) \cdot \cdots \cdot U_{k-1,k}(\mathbb{C}[z]_j) \cdot U_k(\mathbb{C}[z]_j).$$

Proof. If j = 0, this follows from the decompositions of N_0 and $P_0 = L_0 \cdot U_0$. Let us consider the case $j \ge 1$. Let us take $g \in G_0(\mathbb{C}[z]_j)$ and write $g = g_0g_1$ by $g_0 \in G_0$ and $g_1 \in G(\mathbb{C}[z]_j)_1$. Then along the decomposition $G_0 = N_0 \cdot P_0$ we can write $g_0 = n_0p_0$ and thus we have $g = n_0g'_1p_0$ with $g'_1 := p_0g_1p_0^{-1} \in G(\mathbb{C}[z]_j)_1$. Proposition 8.3 gives us the decomposition $g'_1 = n_1p_1$ by $n_1 \in N_0(\mathbb{C}[z]_j)_1$ and $p_1 \in P_0(\mathbb{C}[z]_j)_1$. Then by putting $n := n_0n_1 \in N_0(\mathbb{C}[z]_j)$ and $p := p_0(p_0^{-1}p_1p_0) \in P_0(\mathbb{C}[z]_j)$ we obtain the decomposition g = np. Namely we obtain that the multiplication map $N_0(\mathbb{C}[z]_k) \times P_0(\mathbb{C}[z]_j) \ni (n,p) \mapsto$ $np \in G_0(\mathbb{C}[z]_j) \cap P_0(\mathbb{C}[z]_j) = \{e\}.$

Similarly from the decompositions $N_0 = N_k \cdot N_{k-1,k} \cdot \cdots \cdot N_{0,1}$ and $P_0 = L_0 \cdot U_{0,1} \cdot \cdots \cdot U_{k-1,k} \cdot U_k$ together with Proposition 8.3, we obtain the decompositions

$$N_0(\mathbb{C}[z]_j) = N_k(\mathbb{C}[z]_j) \cdot N_{k-1,k}(\mathbb{C}[z]_j) \cdots N_{0,1}(\mathbb{C}[z]_j),$$

$$P_0(\mathbb{C}[z]_j) = L_0(\mathbb{C}[z]_j) \cdot U_{0,1}(\mathbb{C}[z]_j) \cdots U_{k-1,k}(\mathbb{C}[z]_j) \cdot U_k(\mathbb{C}[z]_j).$$

Thus we obtain the desired decomposition of $G_0(\mathbb{C}[z]_i)$.

Let us set $L_l^{G_0} := L_l \cap G_0$ and define

$$L_{l}^{G_{0}}(\mathbb{C}[z]_{j}) := \{g_{0} \cdot g_{1} \in L_{l} \ltimes L_{l}(\mathbb{C}[z]_{j})_{1} \mid g_{0} \in L_{l}^{G_{0}}\}$$

for each $0 \leq l \leq k + 1$. Also define

$$P_{l,m}^{G_0}(\mathbb{C}[z]_j) := N_{l-1,l}(\mathbb{C}[z]_j) \cdot N_{l-2,l-1}(\mathbb{C}[z]_j) \cdot \cdots \cdot N_{0,1}(\mathbb{C}[z]_j) \cdot L_0(\mathbb{C}[z]_j) \\ \cdot U_{0,1}(\mathbb{C}[z]_j) \cdot \cdots \cdot U_{l'-2,l'-1}(\mathbb{C}[z]_j) \cdot U_{l'-1,l'}(\mathbb{C}[z]_j)$$

for each pair $0 \leq l < l' \leq k$. Then we obtain the Levi-type decomposition

$$P_{l,l'}^{G_0}(\mathbb{C}[z]_j) = L_l^{G_0}(\mathbb{C}[z]_j) \cdot U_{l,l'}(\mathbb{C}[z]_j)$$

and also the LU-type decomposition

$$L_{l'}^{G_0}(\mathbb{C}[z]_j) = N_{l,l'}(\mathbb{C}[z]_j) \cdot P_{l,l'}^{G_0}(\mathbb{C}[z]_j)$$

by Proposition 8.8.

Put $\mathbb{O}_{H}^{L_{l}^{G_{0}}(\mathbb{C}[z]_{k})} := \{ \operatorname{Ad}^{*}(g)(H) \in \mathfrak{g}(\mathbb{C}[z]_{k})^{*} \mid g \in L_{l}^{G_{0}}(\mathbb{C}[z]_{l'}) \}, \text{ and then we obtain a filtration} \}$

$$\{H\} = \mathbb{O}_{H}^{L_{0}^{G_{0}}(\mathbb{C}[z]_{k})} \subset \mathbb{O}_{H}^{L_{1}^{G_{0}}(\mathbb{C}[z]_{k})} \subset \dots \subset \mathbb{O}_{H}^{L_{k+1}^{G_{0}}(\mathbb{C}[z]_{k})} = \mathbb{O}_{H}^{G_{0}(\mathbb{C}[z]_{k})}$$

of $\mathbb{O}_{H}^{G_{0}(\mathbb{C}[z]_{k})}$ as well as that of $\mathbb{O}_{H}^{L_{1} \ltimes G(\mathbb{C}[z]_{k})_{1}}$ in Section 8.3.

Proposition 8.9. For each pair of integers $0 \leq l \leq l' \leq k$, the projection $\pi_{l',l} \colon \mathbb{C}[z]_{l'} \to \mathbb{C}[z]_l$ induces the isomorphism

$$\pi^{\mathbb{O}}_{l',l} \colon \mathbb{O}_{H^{[l']}}^{L^{G_0}_{l+1}(\mathbb{C}[z]_{l'})} \xrightarrow{\sim} \mathbb{O}_{H^{[l]}}^{L^{G_0}_{l+1}(\mathbb{C}[z]_l)}$$

Proof. This directly follows from the same argument as in Proposition 8.4.

Thus the above filtration becomes as follows,

$$\{H_{\rm res}\} = \mathbb{O}_{H^{[0]}}^{L_1^{G_0}(\mathbb{C}[z]_0)} \subset \mathbb{O}_{H^{[1]}}^{L_2^{G_0}(\mathbb{C}[z]_1)} \subset \cdots \subset \mathbb{O}_{H^{[l]}}^{L_{l+1}^{G_0}(\mathbb{C}[z]_l)} \subset \cdots \subset \mathbb{O}_{H^{[k]}}^{L_{k+1}^{G_0}(\mathbb{C}[z]_k)} = \mathbb{O}_{H}^{G_0(\mathbb{C}[z]_k)}.$$

As we saw in Section 8.4, we can obtain decompositions of $\mathbb{O}_{H^{[l]}}^{L_l^{G_0}(\mathbb{C}[z]_l)}$ along the Levitype decomposition of $P_{l,l'}^{G_0}(\mathbb{C}[z]_l)$ and the LU-type one of $L_l^{G_0}(\mathbb{C}[z]_l)$ as follows.

Proposition 8.10. Let us take a pair $0 \le l \le l' \le k$ of positive integers. The multiplication map

$$\overline{\phi} \colon N_{l,l'}(\mathbb{C}[z]_l) \times P_{l,l'}^{G_0}(\mathbb{C}[z]_l) \longrightarrow L_{l'}^{G_0}(\mathbb{C}[z]_l)$$

induces the isomorphism

$$\phi \colon N_{l,l'}(\mathbb{C}[z]_l)_1 \times \mathbb{O}_{H^{[l]}}^{P^{G_0}_{l,l'}(\mathbb{C}[z]_l)} \longrightarrow \mathbb{O}_{H^{[l]}}^{L^{G_0}_{l'}(\mathbb{C}[z]_l)}$$

as complex manifolds. Here $\mathbb{O}_{H^{[l]}}^{P_{l,l'}^{G_0}(\mathbb{C}[z]_l)} := \{ \mathrm{Ad}^*(g)(H^{[l]}) \in \mathfrak{g}(\mathbb{C}[z]_l)^* \mid g \in P_{l,l'}^{G_0}(\mathbb{C}[z]_l) \}.$

Proof. This directly follows from the same argument as in Proposition 8.5.

Proposition 8.11. The map

$$X: \quad \mathbb{O}_{H^{[l]}}^{L_{l}^{G_{0}}(\mathbb{C}[z]_{l})} \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l}) \longrightarrow \begin{array}{c} \mathbb{O}_{H^{[l]}}^{P_{l,l+1}^{G_{0}}(\mathbb{C}[z]_{l})} \\ (\eta,\xi) \longmapsto \eta + \xi \end{array}$$

is an isomorphism of complex manifolds.

Proof. Let us define the map

$$\begin{array}{rcl} a_{H^{[l]}} \colon & L_l^{G_0}(\mathbb{C}[z]_l) \times U_{l,l+1}(\mathbb{C}[z]_l) & \longrightarrow & L_l^{G_0}(\mathbb{C}[z]_l) \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_l) \\ & (g,u) & \longmapsto & (g, \operatorname{Ad}^*(gu)(H^{[l]}) - \operatorname{ad}^*(g)(H^{[l]})) \end{array}$$

and the argument in Lemma 8.6 tells us that this $a_{H^{[l]}}$ is an isomorphism. Then as well as in the proof of Proposition 8.7, the multiplication map $\chi \colon L_l^{G_0}(\mathbb{C}[z]_l) \times U_{l,l+1}(\mathbb{C}[z]) \to P_{l,l+1}^{G_0}(\mathbb{C}[z]_l)$ induces the commutative diagram

$$\begin{array}{cccc} L_{l}^{G_{0}}(\mathbb{C}[z]_{l})_{1} \times U_{l,l+1}(\mathbb{C}[z]_{l}) & \xrightarrow{\chi} & P_{l,l+1}^{G_{0}}(\mathbb{C}[z]_{l}) \\ & \downarrow^{a_{H}[l]} & & \downarrow^{a_{H}[l]} & & \downarrow^{\pi_{P}} \\ & \downarrow^{\pi_{L} \times \mathrm{id}} & & \downarrow^{\pi_{P}} \\ & & \bigcirc_{H^{[l]}}^{L_{l}^{G_{0}}(\mathbb{C}[z]_{l})} \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l}) & \xrightarrow{X} & \bigcirc_{H^{[l]}}^{P_{l,l+1}^{G_{0}}(\mathbb{C}[z]_{l})} \end{array}$$

which shows the surjectivity of X.

Moreover since $\operatorname{Stab}_{L_l^{G_0}(\mathbb{C}[z]_l)}(H^{[l]}) = \operatorname{Stab}_{P_{l,l+1}^{G_0}(\mathbb{C}[z]_l)}(H^{[l]}), X$ is injective by the same argument as in the proof of Proposition 8.7.

Then we finally obtain an analogue of Theorem 8.2.

Theorem 8.12. There exists an isomorphism

$$\Psi \colon \prod_{l=1}^{k} \left(N_{l,l+1}(\mathbb{C}[z]_l) \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_l) \right) \xrightarrow{\sim} \mathbb{O}_H^{G_0(\mathbb{C}[z]_k)}$$

as complex manifolds.

Proof. As a combination of Propositions 8.9, 8.10, and 8.11, we obtain isomorphisms

$$\psi_l \colon N_{l,l+1}(\mathbb{C}[z]_l) \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_l) \times \mathbb{O}_{H^{[l-1]}}^{L_l^{G_0}(\mathbb{C}[z]_l)} \longrightarrow \mathbb{O}_{H^{[l]}}^{L_{l+1}^{G_0}(\mathbb{C}[z]_l)}$$

for l = 1, 2, ..., k. Therefore as the combination of these isomorphisms ψ_l we obtain the desired isomorphism

$$\Psi \colon \prod_{l=1}^{k} \left(N_{l,l+1}(\mathbb{C}[z]_l) \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_l) \right) \longrightarrow \mathbb{O}_H^{G_0(\mathbb{C}[z]_k)_1}$$

by recalling $\mathbb{O}_{H^{[0]}}^{L_1^{G_0}(\mathbb{C}[z]_0)} = \{H_{\mathrm{res}}\}.$

9 Deformation of $N_{l,l+1}(\mathbb{C}[z]_l)_1$ and $\mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1})$

We saw in Section 8 that $N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}(\mathbb{C}[z^{-1}]_{l-1})$ are main building blocks of truncated orbit \mathbb{O}_H . In this section, we shall introduce deformations of these $N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}(\mathbb{C}[z^{-1}]_{l-1})$ which will define the deformation of the truncated orbit \mathbb{O}_H later.

9.1 Deformation of $\mathbb{C}[z]_l$ and $\mathbb{C}[z^{-1}]_l$

To each point $\mathbf{c} = (c_0, c_1, \dots, c_k) \in \mathbb{C}^{k+1}$, we associate the effective divisor of $\mathbb{A}^1(\mathbb{C}) = \mathbb{C}$ defined by

$$D(\mathbf{c}) := c_0 + c_1 + \dots + c_k.$$

Let $|D(\mathbf{c})|$ be the underlying set of points in \mathbb{C} . Then we can consider the subspace of rational functions

$$\mathcal{L}(D(\mathbf{c})) := \{ f \in \mathbb{C}(z) \, | \, \mathrm{ord}_P(f) \ge -n_P \text{ for all } P \in \mathbb{C} \}$$

where n_P is the coefficient of $P \in \mathbb{C}$ in the formal sum $D(\mathbf{c}) = \sum_{P \in \mathbb{C}} n_P P$, and $\operatorname{ord}_P(f)$ is the order of f at P. We can regard $L(D(\mathbf{c}))$ as $\mathbb{C}[z]$ -submodule of $\mathbb{C}(z)$ containing $\mathbb{C}[z]$. Thus we can consider the quotient module

$$\widehat{\mathcal{L}}(\mathbf{c}) := \mathcal{L}(D(\mathbf{c})) / \mathbb{C}[z].$$

We also consider

$$\mathcal{L}(-D(\mathbf{c})) := \{ f \in \mathbb{C}(z) \mid \operatorname{ord}_P(f) \ge n_P \text{ for all } P \in \mathbb{C} \}.$$

which may be regarded as the ideal of $\mathbb{C}[z]$ generated by $\prod_{i=0}^{k} (z - c_i)$. Thus we can consider the quotient ring

$$\widehat{\Gamma}(\mathbf{c}) := \mathbb{C}[z]/\mathrm{L}(-D(\mathbf{c})) = \mathbb{C}[z]/\langle \prod_{i=0}^{k} (z-c_i) \rangle_{\mathbb{C}[z]}$$

Then $\widehat{\Gamma}(\mathbf{c})$ is a finite dimensional \mathbb{C} -algebra with maximal ideals $\mathfrak{m}_{z_{c_i}}^{\widehat{\Gamma}(\mathbf{c})} := \langle z - c_i \rangle_{\widehat{\Gamma}(\mathbf{c})}$ for $i = 0, 1, \ldots, k$. Let us note that these maximal ideals $\mathfrak{m}_{z_{c_i}}^{\widehat{\Gamma}(\mathbf{c})}$ are not nilpotent in general, unlike the maximal ideal $\mathfrak{m}_z^{(l)}$ of the local ring $\mathbb{C}[z]_l$. We notice that since $L(-D(\mathbf{c})) \subset \mathbb{C}[z]$ acts trivially on $\widehat{L}(\mathbf{c})$, $\widehat{L}(\mathbf{c})$ has the $\widehat{\Gamma}(\mathbf{c})$ -module structure. Also we note that if $\mathbf{c} = \mathbf{0}$, then

$$\widehat{\Gamma}(\mathbf{0}) \cong \mathbb{C}[z]_k, \qquad \widehat{\mathrm{L}}(\mathbf{0}) \cong \mathbb{C}[z^{-1}]_k.$$

Therefore $\widehat{\Gamma}(\mathbf{c})$ and $\widehat{\mathcal{L}}(\mathbf{c})$ can be regarded as deformations of $\mathbb{C}[z]_k$ and $\mathbb{C}[z^{-1}]_k$ respectively.

Let us define a filtration on $\widehat{\mathcal{L}}(\mathbf{c})$ and cofiltration on $\widehat{\Gamma}(\mathbf{c})$ as follows. Consider projection maps $\operatorname{pr}^{(l)} : \mathbb{C}^{k+1} \ni (x_0, x_1, \dots, x_k) \mapsto (x_0, x_1, \dots, x_l) \in \mathbb{C}^{l+1}$ and define divisors $D(\mathbf{c})_l := D(\operatorname{pr}^{(l)}(\mathbf{c}))$ for $l = 0, 1, \dots, k$. Then we obtain the filtration

$$L(D(\mathbf{c})_0) \subset L(D(\mathbf{c})_1) \subset \cdots \subset L(D(\mathbf{c})_k) = L(D(\mathbf{c})).$$

Definition 9.1 (Standard filtration and basis of $\hat{L}(\mathbf{c})$). Let

$$\widehat{\mathcal{L}}(\mathbf{c})_l := \mathcal{L}(D(\mathbf{c})_l) / \mathbb{C}[z] \text{ for } l = 0, 1, \dots, k.$$

The filtration

$$\widehat{\mathrm{L}}(\mathbf{c})_0 \subset \widehat{\mathrm{L}}(\mathbf{c})_1 \subset \cdots \subset \widehat{\mathrm{L}}(\mathbf{c})_k = \widehat{\mathrm{L}}(\mathbf{c})$$

is called the *standard filtration* of $\widehat{L}(\mathbf{c})$. The associated basis of $\widehat{L}(\mathbf{c})$,

$$\frac{1}{z-c_0}, \frac{1}{(z-c_0)(z-c_1)}, \dots, \frac{1}{(z-c_0)(z-c_1)\cdots(z-c_k)}$$

as \mathbb{C} -vector space is called the *standard basis* of $\widehat{L}(\mathbf{c})$.

Notice that

$$\frac{1}{z-c_0}, \frac{1}{(z-c_0)(z-c_1)}, \dots, \frac{1}{(z-c_0)(z-c_1)\cdots(z-c_l)}$$

becomes a basis of the *l*-th component $\hat{\mathbf{L}}(\mathbf{c})_l$ of the standard filtration for each $l = 0, 1, \ldots, k$.

Similarly, we can consider the filtration

$$L(-D(\mathbf{c})_0) \supset L(-D(\mathbf{c})_1) \supset \cdots \supset L(-D(\mathbf{c})_k) = L(-D(\mathbf{c})).$$

Definition 9.2 (Standard cofiltration and basis of $\widehat{\Gamma}(\mathbf{c})$). Let

$$\widehat{\Gamma}(\mathbf{c})_l := \mathbb{C}[z]/L(-D(\mathbf{c})_l) = \mathbb{C}[z]/\langle \prod_{i=0}^l (z-c_i) \rangle_{\mathbb{C}[z]} \text{ for } l = 0, 1, \dots, k.$$

Then the sequence of projection maps

$$\widehat{\Gamma}(\mathbf{c})_0 \to \widehat{\Gamma}(\mathbf{c})_1 \to \cdots \to \widehat{\Gamma}(\mathbf{c})_k = \widehat{\Gamma}(\mathbf{c})$$

induced by the above filtration is called the *standard cofiltration* of $\widehat{\Gamma}(\mathbf{c})$. The basis of $\widehat{\Gamma}(\mathbf{c})$,

$$1, (z - c_0), (z - c_0)(z - c_1), \dots, (z - c_0)(z - c_1) \cdots (z - c_{k-1})$$

as \mathbb{C} -vector space is called the *standard basis* of $\widehat{\Gamma}(\mathbf{c})$.

We can see that

$$1, (z - c_0), (z - c_0)(z - c_1), \dots, (z - c_0)(z - c_1) \cdots (z - c_{l-1})$$

becomes a basis of $\widehat{\Gamma}(\mathbf{c})_l$ for each $l = 0, 1, \dots, k$.

Next we introduce the pairing of $\widehat{\Gamma}(\mathbf{c})_l$ and $\widehat{L}(\mathbf{c})_l$ as \mathbb{C} -vector spaces defined by

$$\langle , \rangle_{\mathbf{c},l} \colon \widehat{\Gamma}(\mathbf{c})_l \times \widehat{\mathcal{L}}(\mathbf{c})_l \longrightarrow \mathbb{C} (f(z), g(z)) \longmapsto \sum_{c \in |D(\mathbf{c})|} \operatorname{res}_{z=c} (f(z)g(z))$$

for each l = 0, 1, ..., k.

Lemma 9.3 (Residue formula). Let us take $f(z) \in \widehat{L}(\mathbf{c})_l$ and write

$$f(z) = \sum_{i=0}^{l} \frac{f_i}{\prod_{\nu=0}^{i} (z - c_{\nu})}$$

under the standard basis. Then we have

$$\sum_{c \in |D(\mathbf{c})|} \operatorname{res}_{z=c}(f(z)) = f_0$$

Proof. It follows from a direct computation.

Proposition 9.4. The pairing $\langle , \rangle_{\mathbf{c},l}$ is non-degenerate. Moreover, the bases

$$(z - c_0), (z - c_0)(z - c_1), \dots, (z - c_0)(z - c_1) \cdots (z - c_{l-1})$$

of $\widehat{\Gamma}(\mathbf{c})_l$ and

$$\frac{1}{z-c_0}, \frac{1}{(z-c_0)(z-c_1)}, \dots, \frac{1}{(z-c_0)(z-c_1)\cdots(z-c_l)}$$

of $\widehat{L}(\mathbf{c})_l$ are dual bases with respect to this pairing. Proof. It follows that

$$\left\langle 1, \frac{1}{(z-c_0)(z-c_1)\cdots(z-c_{j'})} \right\rangle_{\mathbf{c},l} = \delta_{0,j'}$$

for $0 \leq j' \leq l$ and

$$\left\langle (z-c_0)(z-c_1)\cdots(z-c_{j-1}), \frac{1}{(z-c_0)(z-c_1)\cdots(z-c_{j'})} \right\rangle_{\mathbf{c},l} = \delta_{j,j'},$$

for $0 < j \leq l, 0 \leq j' \leq l$, from Lemma 9.3. Here $\delta_{j,j'}$ is Kronecker delta.

Corollary 9.5. Let us take $\mathbf{c} = (c_i)_{i=0,1,...,l}$, $\mathbf{x} = (x_i)_{i=0,1,...,l}$, and $\mathbf{y} = (y_i)_{i=0,1,...,l}$ from $\mathbb{C}^{\oplus (l+1)}$. Define

$$f(\mathbf{x}, \mathbf{c}) := x_0 + x_1(z - c_0) + \dots + x_l(z - c_0)(z - c_1) \cdots (z - c_{l-1}),$$

$$g(\mathbf{x}, \mathbf{c}) := \frac{x_0}{z - c_0} + \frac{x_1}{(z - c_0)(z - c_1)} + \dots + \frac{x_l}{(z - c_0)(z - c_1) \cdots (z - c_l)}.$$

Then

$$\langle f(\mathbf{x},\mathbf{c}),g(\mathbf{x},\mathbf{c})
angle_{\mathbf{c},l}$$

is constant as a function of $\mathbf{c} \in \mathbb{C}^{\oplus (l+1)}$.

Proof. This is a direct consequence of the above proposition.

9.2 Partial fraction decomposition and Chinese remainder theorem

Let us take a partition $\mathcal{I}: I_0, I_1, \ldots, I_r$ of $\{0, 1, \ldots, k\}$ and write $I_j = \{i_{[j,0]} < i_{[j,1]} < \ldots < i_{[j,k_j]}\}$. For each $l = 0, 1, \ldots, k$, we define a partition $\mathcal{I}^{(l)}: I_0^{(l)}, I_1^{(l)}, \ldots, I_r^{(l)}$ of $\{0, 1, \ldots, l\} \subset \{0, 1, \ldots, k\}$ by $I_j^{(l)} := I_j \cap \{0, 1, \ldots, l\}$. Here we allow the case $I_j^{(l)} = \emptyset$. Let us set $l_j := |I_j^{(l)}| - 1$. Then for $\mathbf{c} = (c_0, c_1, \ldots, c_k) \in C(\mathcal{I})$, we can write $D(\mathbf{c})_l = \sum_{\nu=0}^r (l_j + 1) \cdot c_{i_{[j,0]}}$.

Then for $\mathbf{c} = (c_0, c_1, \dots, c_k) \in C(\mathcal{I})$, we can write $D(\mathbf{c})_l = \sum_{\nu=0}^r (l_j + 1) \cdot c_{i_{[j,0]}}$. The algorithm of the partial fractional decomposition of rational functions gives us the direct sum decomposition

$$\widehat{\mathcal{L}}(\mathbf{c})_l = \bigoplus_{j=0}^r \mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{l_j}.$$

Here we formally put $\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{l_j} := \{0\}$ if $l_j < 0$, i.e., $I_j^{(l)} = \emptyset$. We denote natural projection maps by $\operatorname{pr}_{c_{i_{[j,0]}}}: \widehat{\operatorname{L}}(\mathbf{c})_l \to \mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{l_j}$ for $j = 0, 1, \ldots, r$.

We also have the following decomposition of $\widehat{\Gamma}(\mathbf{c})_l$. Since $c_{i_{[j,0]}} \neq c_{i_{[j',0]}}$ for $j \neq j'$, it follows that

$$\langle (z - c_{i_{[j,0]}})^{l_j} \rangle_{\mathbb{C}[z]} + \langle (z - c_{i_{[j',0]}})^{l_{j'}} \rangle_{\mathbb{C}[z]} = \mathbb{C}[z], \quad j \neq j' \text{ with } I_j \neq \emptyset, I_{j'} \neq \emptyset,$$
$$\bigcap_{j=0}^r \langle (z - c_{i_{[j,0]}})^{l_j} \rangle_{\mathbb{C}[z]} = \mathcal{L}(-D(\mathbf{c})_l).$$

Therefore the Chinese remainder theorem implies that natural projection maps

$$\mathrm{pr}_{c_{i_{[j,0]}}} \colon \widehat{\Gamma}(\mathbf{c})_{l} \longrightarrow \mathbb{C}[z] / \langle (z - c_{i_{[j,0]}})^{l_{j}+1} \rangle_{\mathbb{C}[z]} = \mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}}$$

define the algebra isomorphism

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon \quad \widehat{\Gamma}(\mathbf{c})_{l} \longrightarrow \prod_{j=0}^{r} \mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}}$$
$$f(z) \longmapsto (\operatorname{pr}_{c_{i_{[j,0]}}}(f(z)))_{j=0,1,\dots,r} \ .$$

Here we formally put $\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j} := \{0\}$ if $l_j < 0$, i.e., $I_j^{(l)} = \emptyset$.

Under this isomorphism, maximal ideals $\mathfrak{m}_{z_{c_{i_{[j,0]}}}}^{\widehat{\Gamma}(\mathbf{c})_{l}}$ for $j = 0, 1, \ldots, r$ correspond to

$$\mathbb{C}[z_{c_{i_{[0,0]}}}]_{l_{0}} \times \cdots \times \mathbb{C}[z_{c_{i_{[j-1,0]}}}]_{l_{j-1}} \times \mathfrak{m}_{z_{c_{i_{[j,0]}}}}^{(l_{j})} \times \mathbb{C}[z_{c_{i_{[j+1,0]}}}]_{l_{j+1}} \times \cdots \times \mathbb{C}[z_{c_{i_{[r,0]}}}]_{l_{r}}$$

in $\prod_{j=0}^{r} \mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}$.

These decompositions of $\widehat{\Gamma}(\mathbf{c})_l$ and $\widehat{L}(\mathbf{c})_l$ are compatible with the pairing $\langle , \rangle_{\mathbf{c},l}$. Namely, for each $j = 0, 1, \ldots, r$, we define the non-degenerate pairing

$$\langle , \rangle_{c_{i_{[j,0]}},l} \quad \mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}} \times \mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{l_{j}} \longrightarrow \mathbb{C} \\ (f(z_{c_{i_{[j,0]}}}), g(z_{c_{i_{[j,0]}}})) \longmapsto \operatorname{res}_{z_{c_{i_{[j,0]}}}=0}(f(z_{c_{i_{[j,0]}}})g(z_{c_{i_{[j,0]}}})) + C_{c_{i_{[j,0]}}})$$

Then a direct computation shows the following.

Proposition 9.6. Let us take $\mathcal{I} \in \mathcal{P}_{[k+1]}$, $\mathbf{c} \in C(\mathcal{I})$, and an integer $0 \leq l \leq k$ as above. Then for $(f(z), g(z)) \in \widehat{\Gamma}(\mathbf{c})_l \times \widehat{L}(\mathbf{c})_l$, we have

$$\langle f(z), g(z) \rangle_{\mathbf{c},l} = \sum_{j=0}^{r} \langle \operatorname{pr}_{c_{i_{[j,0]}}}(f(z)), \operatorname{pr}_{c_{i_{[j,0]}}}(g(z)) \rangle_{c_{i_{[j,0]}},l}.$$

Let us notice that the decomposition $\widehat{\mathcal{L}}(\mathbf{c})_l = \bigoplus_{j=0}^r \mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{l_j}$ is not just as a vector space but as a $\widehat{\Gamma}(\mathbf{c})_l$ -module. Furthermore this is compatible with the decomposition of the algebra $\widehat{\Gamma}(\mathbf{c})_l = \prod_{j=0}^r \mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}$ in the following sense. Namely, we can easily check that the diagrams

$$\widehat{\Gamma}(\mathbf{c})_{l} \times \widehat{\mathbf{L}}(\mathbf{c})_{l} \xrightarrow{\cdot} \widehat{\mathbf{L}}(\mathbf{c})_{l} \xrightarrow{\cdot} \widehat{\mathbf{L}}(\mathbf{c})_{l} \\
\downarrow^{\operatorname{pr}_{c_{i_{[j,0]}}} \times \operatorname{pr}_{c_{i_{[j,0]}}}} \downarrow^{\operatorname{pr}_{c_{i_{[j,0]}}}} \\
\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}} \times \mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{l_{j}} \xrightarrow{\cdot} \mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{l_{j}}$$
(5)

are commutative for j = 0, 1, ..., r and l = 0, 1, ..., k. Here horizontal arrows are scalar multiplications of left modules.

9.3 Lie groupoid $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$

Let us recall the notion of Lie algebroids.

Definition 9.7 (Complex Lie algebroid). A complex Lie algebroid over a complex manifold M is a holomorphic vector bundle $A \to M$, together with a Lie bracket $[\cdot, \cdot]$ on its space of holomorphic sections, such that there exists a vector bundle map $\mathbf{a}: A \to TM$ called the *anchor map* satisfying the Leibniz rule

$$[\sigma, f\tau] = f[\sigma, \tau] + (\mathbf{a}(\sigma)f)\tau$$

for all global sections σ, τ of $A \to M$ and holomorphic functions f on M.

As well as the theory of Lie groups, we can associate the Lie algebroid to a Lie groupoid as follows.

Definition 9.8 (Lie algebroid of a Lie groupoid). For a complex Lie groupoid $M = (M, M_0, s, t, m)$, we consider the vector bundle

$$\operatorname{Lie}(M) := TM|_{M_0}/TM_0$$

defined as the normal bundle to M_0 in M. As the anchor map, we consider the unique map \mathbf{a} : Lie $(M) \to TM_0$ which makes the following diagram commutative,



The Lie bracket is defined as follows. Let (S_t) and (R_t) be families of bisections of M such that $\sigma = \frac{\partial}{\partial t}|_{t=0}S$ of bisections such that $\tau = \frac{\partial}{\partial t}|_t R_t$. Then the Lie bracket of σ and τ is defined by

$$[\sigma,\tau] := \frac{\partial}{\partial t_2}\Big|_{t_2=0} \left(\frac{\partial}{\partial t_1}\Big|_{t_1=0} (S_{t_2}R_{t_1}S_{t_2}^{-1})\right).$$

Now let us fix a positive integer $1 \leq l \leq k$ and an open neighborhood $\mathbb{B} \subset \mathbb{C}^{k+1}$ of **0**. Then we consider the trivial vector bundle

$$\mathfrak{g}^{\oplus(l+1)} \times \mathbb{B} \xrightarrow{\mathrm{pr}_2} \mathbb{B}.$$

This can be identified with the family $\bigsqcup_{\mathbf{c}\in\mathbb{B}}\mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_l)$ of Lie algebras by the following bijection,

$$\begin{array}{cccc}
\mathfrak{g}^{\oplus (l+1)} \times \mathbb{B} & \longmapsto & \bigsqcup_{\mathbf{c} \in \mathbb{B}} \mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_l) \\
((X_i)_{i=0,1,\dots,l}, \mathbf{c}) & \longmapsto & X_0 + X_1(z-c_0) + \dots + X_l(z-c_0) \cdots (z-c_{l-1})
\end{array}$$

Then the Lie algebra structure of each direct summand $\mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_l)$ allows us to regard the trivial bundle $\mathfrak{g}^{\oplus (l+1)} \times \mathbb{B} \to \mathbb{B}$ as a Lie algebroid with the trivial anchor map. We denote this Lie algebroid by

 $\mathfrak{g}(\widehat{\Gamma}(\mathbb{B})_l).$

Furthermore, let us consider the nilpotent Lie subalgebra $\mathfrak{n}_{l,l+1} \subset \mathfrak{g}$ and the subbundle $(\mathfrak{n}_{l,l+1})^{\oplus (l+1)} \times \mathbb{B} \xrightarrow{\operatorname{pr}_2} \mathbb{B}$. Then it has the structure of Lie algebroid as well. Then we denote the Lie algebroid $(\mathfrak{n}_{l,l+1})^{\oplus (l+1)} \times \mathbb{B} \xrightarrow{\operatorname{pr}_2} \mathbb{B}$ by

$$\mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l).$$

Since each fiber of the Lie algebroid $\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$ is nilpotent Lie algebra $\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)$, we can moreover equip this trivial bundle with the Lie groupoid structure as follows. As the source s and target t, we take the projection pr_2 : $(\mathbf{n}_{l,l+1})^{\oplus (l+1)} \times \mathbb{B} \to \mathbb{B}$, i.e., $s = t = \mathrm{pr}_2$. As the unit i, we take the zero section $\mathbf{0} \colon \mathbb{B} \ni \mathbf{c} \mapsto (\mathbf{0}, \mathbf{c}) \in (\mathbf{n}_{l,l+1})^{\oplus (l+1)} \times \mathbb{B}$. Then for each $\mathbf{c} \in \mathbb{B}$ we identify the fiber $(\mathbf{n}_{l,l+1})^{\oplus (l+1)} \times \{\mathbf{c}\}$ with the nilpotent Lie algebra $\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c}))$, and define the multiplication of $X_{\mathbf{c}}, Y_{\mathbf{c}} \in (\mathbf{n}_{l,l+1})^{\oplus (l+1)} \times \{\mathbf{c}\}$ by Dynkin's formula

$$m(X_{\mathbf{c}}, Y_{\mathbf{c}}) := \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \sum_{(p_1, q_1) \in (\mathbb{Z}_{\geq 0})^2 \setminus \{0\}} \cdots \sum_{(p_n, q_n) \in (\mathbb{Z}_{\geq 0})^2 \setminus \{0\}} \frac{[X_{\mathbf{c}}^{p_1} Y_{\mathbf{c}}^{q_1} \cdots X_{\mathbf{c}}^{p_n} Y_{\mathbf{c}}^{q_n}]}{(\sum_{i=1}^n (p_i + q_i)) \prod_{i=1}^n p_i! q_i!},$$

where

$$[X_{\mathbf{c}}^{p_1}Y_{\mathbf{c}}^{q_1}\cdots X_{\mathbf{c}}^{p_n}Y_{\mathbf{c}}^{q_n}] := (\mathrm{ad}X_{\mathbf{c}})^{p_1}(\mathrm{ad}Y_{\mathbf{c}})^{q_1}\cdots (\mathrm{ad}X_{\mathbf{c}})^{p_n}(\mathrm{ad}Y_{\mathbf{c}})^{q_n-1}Y_{\mathbf{c}}$$

Since $\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)$ is nilpotent, Dynkin's formula becomes a finite sum, namely, the multiplication is well-defined. This multiplication map obviously depends holomorphically on $\mathbf{c} \in \mathbb{B}$, and therefore the Campbell-Baker-Hausdorff formula shows that three properties in Definition 7.1 are satisfied. We denote this Lie groupoid by

$$N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l).$$

By definition, each fiber at $\mathbf{c} \in \mathbb{B}$ is isomorphic to the nilpotent Lie group $N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)$. Obviously the Lie algebroid of this Lie groupoid is $\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$. We denote the identity map by exp: $\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l) \to N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$ for the later convenience.

The adjoint action of the Lie group $N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)$ on the Lie algebra $\mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_l)$ at each $\mathbf{c} \in \mathbb{B}$ defines the adjoint action of the Lie groupoid $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$ on the Lie algebroid $\mathfrak{g}(\widehat{\Gamma}(\mathbb{B})_l)$. Namely we obtain the bundle map

$$\begin{array}{rcl} \mathrm{Ad} \colon & N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l) \times \mathfrak{g}(\widehat{\Gamma}(\mathbb{B})_l) & \longrightarrow & \mathfrak{g}(\widehat{\Gamma}(\mathbb{B})_l) \\ & & (n_{\mathbf{c}}, X_{\mathbf{c}}) & \longmapsto & \mathrm{Ad}(n_{\mathbf{c}})(X_{\mathbf{c}}) \end{array}$$

where $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l) \times \mathfrak{g}(\widehat{\Gamma}(\mathbb{B})_l)$ denotes the product bundle over \mathbb{B} . Let us introduce another realization of the trivial bundle $\mathfrak{g}^{\oplus(l+1)} \times \mathbb{B} \to \mathbb{B}$. We identify it with the family $\bigsqcup_{\mathbf{c}\in\mathbb{B}}\mathfrak{g}(\widehat{\mathcal{L}}(\mathbf{c})_l)$ by

$$\mathfrak{g}^{\oplus(l+1)} \times \mathbb{B} \longmapsto \underbrace{X_0}_{z-c_0} + \underbrace{X_1}_{(z-c_0)(z-c_1)} \oplus \underbrace{X_l}_{(z-c_0)\cdots(z-c_l)} + \cdots + \underbrace{X_l}_{(z-c_0)\cdots(z-c_l)}$$

Then we denote this trivial bundle by

$$\mathfrak{g}(\mathcal{L}(\mathbb{B})_l).$$

At each $\mathbf{c} \in \mathbb{B}$, we have the non-degenerate trace pairing

$$\begin{array}{ccc} \langle \,, \, \rangle_{\mathbf{c},l} \colon & \mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_l) \times \mathfrak{g}(\widehat{\mathbf{L}}(\mathbf{c})_l) & \longrightarrow & \mathbb{C} \\ & & (f(z), g(z)) & \longmapsto & \mathrm{tr} \left(\sum_{c \in |D(\mathbf{c})|} \mathop{\mathrm{res}}_{z=c} (f(z)g(z)) \right) \end{array}$$

Thus we can regard $\mathfrak{g}(\widehat{L}(\mathbb{B})_l)$ as the dual bundle of the Lie algebroid $\mathfrak{g}(\widehat{\Gamma}(\mathbb{B})_l)$ through these pairings.

The residue map

res:
$$\mathfrak{g}(\widehat{\mathbf{L}}(\mathbf{c})_l) \longrightarrow \mathfrak{g}$$

 $f(z) \longmapsto \sum_{c \in |D(\mathbf{c})|} \operatorname{res}_{z=c}(f(z))$

at each $\mathbf{c} \in \mathbb{B}$ defines the residue map for the family

res: $\mathfrak{q}(\widehat{\mathcal{L}}(\mathbb{B})_l) \longrightarrow \mathfrak{q}$.

The fiber-wise coadjoint action defines the action of $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$ on $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B})_l) \cong$ $\mathfrak{g}(\widehat{\Gamma}(\mathbb{B})_l)^*,$

$$\operatorname{Ad}^* \colon N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l) \times \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B})_l) \longrightarrow \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B})_l) \\ (n_{\mathbf{c}}, \xi_{\mathbf{c}}) \longmapsto \operatorname{Ad}^*(n_{\mathbf{c}})(\xi_{\mathbf{c}})$$

Here $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l) \times \mathfrak{g}(\widehat{L}(\mathbb{B})_l)$ denotes product bundle over \mathbb{B} .

Moreover take the opposite $\mathfrak{u}_{l,l+1}$ of $\mathfrak{n}_{l,l+1}$ and consider the subbundle $(\mathfrak{u}_{l,l+1})^{\oplus (l+1)} \times$ $\mathbb{B} \to \mathbb{B}$ of $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B})_l)$ and denote it by

$$\mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B})_l).$$

Decompositions of fibers of $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$ 9.4

Let us take a partition $\mathcal{I}: I_0, I_1, \ldots, I_r$ of $\{0, 1, \ldots, k\}$ as in Section 5.2. Then as we saw in Section 9.2, we can define the partition $\mathcal{I}^{(l)}: I_0^{(l)}, I_1^{(l)}, \ldots, I_r^{(l)}$ of $\{0, 1, \ldots, l\}$ from \mathcal{I} .

Let $\mathbf{c} \in C(\mathcal{I}) \cap \mathbb{B}$. Then the isomorphism $\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} : \widehat{\Gamma}(\mathbf{c})_{l} \xrightarrow{\sim} \prod_{j=0}^{r} \mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}}$ given in Section 9.2 induces the isomorphism

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon \mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_{l}) \xrightarrow{\sim} \prod_{j=0}^{r} \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}})$$

of Lie algebras which restricts to the isomorphism

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon \mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_{l}) \xrightarrow{\sim} \prod_{j=0}^{r} \mathfrak{n}_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}}).$$

Under the natural isomorphisms

$$\mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l) \cong N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l), \quad \mathfrak{n}_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}) \cong N_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j})$$

by exponential maps, we also obtain the isomorphism

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_{l}) \xrightarrow{\sim} \prod_{j=0}^{r} N_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}})$$

of complex manifolds.

Proposition 9.9. The above map $\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}}$ is an isomorphism as Lie groups.

Proof. It suffices to check that the above map is a group homomorphism which follows from the fact that linear algebraic groups defined over \mathbb{C} are functors form the category of \mathbb{C} -algebras to that of groups. This also can be checked directly in the following way.

Let us take $X, Y \in \mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)$. Then Dynkin's formula gives us

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}}(\exp\left(X\right) \cdot \exp\left(Y\right)) = \left(\exp\left(\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}}\left(\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \sum_{\substack{(p_{1},q_{1}) \in (\mathbb{Z}_{\geq 0})^{2} \setminus \{0\}}} \frac{\left[X^{p_{1}}Y^{q_{1}} \cdots X^{p_{n}}Y^{q_{n}}\right]}{\left(\sum_{i=1}^{n} (p_{i}+q_{i})\right) \prod_{i=1}^{n} p_{i}!q_{i}!}\right) \right).$$

Then since $\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}}$ is a Lie algebra isomorphism, we moreover have

$$\begin{split} \prod_{j=0}^{r} \mathrm{pr}_{c_{i[j,0]}} \left(\sum_{\substack{(p_{1},q_{1}) \in (\mathbb{Z}_{\geq 0})^{2} \setminus \{0\} \\ \vdots \\ (p_{n},q_{n}) \in (\mathbb{Z}_{\geq 0})^{2} \setminus \{0\}}} \frac{[X^{p_{1}}Y^{q_{1}} \cdots X^{p_{n}}Y^{q_{n}}]}{(\sum_{i=1}^{n} (p_{i}+q_{i})) \prod_{i=1}^{n} p_{i}!q_{i}!} \right) = \\ \left(\sum_{\substack{(p_{n},q_{n}) \in (\mathbb{Z}_{\geq 0})^{2} \setminus \{0\} \\ \vdots \\ (p_{n},q_{n}) \in (\mathbb{Z}_{\geq 0})^{2} \setminus \{0\}}} \frac{[\mathrm{pr}_{c_{i[j,0]}}(X)^{p_{1}}\mathrm{pr}_{c_{i[j,0]}}(Y)^{q_{1}} \cdots \mathrm{pr}_{c_{i[j,0]}}(X)^{p_{n}}\mathrm{pr}_{c_{i[j,0]}}(Y)^{q_{n}}]}{(\sum_{i=1}^{n} (p_{i}+q_{i})) \prod_{i=1}^{n} p_{i}!q_{i}!} \right)_{j=0,\dots,r}. \end{split}$$

Therefore we obtain

$$\begin{split} \prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}}(\exp\left(X\right) \cdot \exp\left(Y\right)) &= \left(\exp\left(\operatorname{pr}_{c_{i_{[j,0]}}}(X)\right) \cdot \exp\left(\operatorname{pr}_{c_{i_{[j,0]}}}(Y)\right)\right)_{j=0,\dots,r} \\ &= \left(\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}}(\exp\left(X\right))\right) \cdot \left(\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}}(\exp\left(X\right))\right) \\ \text{esired.} \end{split}$$

as desired.

Let us also recall the isomorphism $\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} : \widehat{\mathcal{L}}(\mathbf{c})_{l} \xrightarrow{\sim} \prod_{j=0}^{r} \mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]$ given in Section 9.2. Then it induces the isomorphism

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon \mathfrak{g}(\widehat{\mathcal{L}}(\mathbf{c})_{l}) \xrightarrow{\sim} \prod_{j=0}^{r} \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}])$$

which restricts to the isomorphism

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon \mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbf{c})_{l}) \xrightarrow{\sim} \prod_{j=0}^{r} \mathfrak{u}_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]).$$

Proposition 9.10. The following diagram is commutative.

Proof. Let us take $X \in \mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)$ and $Y \in \mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_l)$. Then since

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} : \mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_{l}) \to \prod_{j=0}^{r} \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}}),$$
$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} : \mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_{l}) \xrightarrow{\sim} \prod_{j=0}^{r} \mathfrak{n}_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}})$$

are Lie algebra isomorphisms, we have

$$\begin{split} \prod_{j=0}^{r} \mathrm{pr}_{c_{i_{[j,0]}}} \left(\mathrm{Ad}(\exp{(X)})(Y) \right) &= \prod_{j=0}^{r} \mathrm{pr}_{c_{i_{[j,0]}}} \left(\sum_{n=0}^{\infty} \frac{(\mathrm{ad}(X))^{n}(Y)}{n!} \right) \\ &= \left(\sum_{n=0}^{\infty} \frac{(\mathrm{ad}(\mathrm{pr}_{c_{i_{[j,0]}}}(X)))^{n} (\mathrm{pr}_{c_{i_{[j,0]}}}(Y))}{n!} \right)_{j=0,\dots,r} \\ &= \left(\mathrm{Ad} \left(\mathrm{pr}_{c_{i_{[j,0]}}} (\exp{(X)}) \right) \left(\mathrm{pr}_{c_{i_{[j,0]}}}(Y) \right) \right)_{j=0,\dots,r}. \end{split}$$

Namely, the following diagram is commutative,

$$\begin{split} N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l) &\times \mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_l) \xrightarrow{\mathrm{Ad}} \mathfrak{g}(\widehat{\Gamma}(\mathbf{c})_l) \\ & \downarrow^{\prod_{j=0}^r (\mathrm{pr}_{c_{i_{[j,0]}}} \times \mathrm{pr}_{c_{i_{[j,0]}}})} & \downarrow^{\prod_{j=0}^r \mathrm{pr}_{c_{i_{[j,0]}}}} \\ & \prod_{j=0}^r \left(N_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}) \times \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}) \right) \xrightarrow{\prod_{j=0}^r \mathrm{Ad}} \prod_{j=0}^r \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}) \end{split}$$
Then since vertical arrows are isomorphisms, the dual of this diagram gives our desired commutative diagram. $\hfill \Box$

Finally we look at the compatibility with the residue map res: $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbf{c})_l) \to \mathfrak{g}$ and the partial fraction decomposition. Since $\prod_{j=0}^r \operatorname{pr}_{c_{i_{[j,0]}}} : \mathfrak{g}(\widehat{\mathcal{L}}(\mathbf{c})_l) \to \prod_{j=0}^r \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}])$ is defined by the partial fractional decomposition, we obviously have

$$\operatorname{res}(X) = \sum_{j=0}^{r} \operatorname{res}_{z=c_{i_{[j,0]}}}(\operatorname{pr}_{c_{i_{[j,0]}}}(X))$$
(6)

for $X \in \mathfrak{g}(\widehat{\mathcal{L}}(\mathbf{c})_l)$

9.5 Lie subgroupoid $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)_1$

Let us consider the subbundle of the Lie algebroid $\mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l) = \bigsqcup_{\mathbf{c} \in \mathbb{B}} \mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)$ defined by

$$\mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)_1 := \left\{ X_0 + X_1(z-c_0) + \dots + X_l(z-c_0) \cdots (z-c_{l-1}) \in \bigsqcup_{\mathbf{c} \in \mathbb{B}} \mathfrak{g}(\widehat{\Gamma}(\mathbf{c})) \, \middle| \, X_0 = 0 \right\}$$

which has the natural Lie algebroid structure induce from that of $\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$. Then as well as for the Lie groupoid $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$, Dynkin's formula defines the groupoid structure on this vector bundle $\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)_1$, and we denote this Lie groupoid by

$$N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)_1$$

which is a Lie subgroupoid of $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)$.

By definition, each fiber of the Lie algebroid $\mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)_1$ at $\mathbf{c} \in \mathbb{B}$ is isomorphic to $\mathfrak{n}_{l,l+1} \otimes_{\mathbb{C}} \mathfrak{m}_{z_{c_0}}^{\widehat{\Gamma}(\mathbf{c})}$ where $\mathfrak{m}_{z_{c_0}}^{\widehat{\Gamma}(\mathbf{c})} = \langle z - c_0 \rangle_{\widehat{\Gamma}(\mathbf{c})}$ is a maximal ideal of $\widehat{\Gamma}(\mathbf{c})$. We denote this nilpotent Lie algebra by

$$\mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)_1 := \mathfrak{n}_{l,l+1} \otimes_{\mathbb{C}} \mathfrak{m}_{z_{c_0}}^{\widehat{\Gamma}(\mathbf{c})}$$

and also denote the corresponding nilpotent Lie group by

$$N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)_1.$$

Then $\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)_1$ and $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B})_l)_1$ can be regarded as the holomorphic families of nilpotent Lie algebras $\bigsqcup_{\mathbf{c}\in\mathbb{B}} \left(\mathbf{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)_1\right)$ and groups $\bigsqcup_{\mathbf{c}\in\mathbb{B}} \left(N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)_1\right)$ respectively.

Now we take a partition $\mathcal{I}: I_0, \ldots, I_r$ of $\{0, 1, \ldots, k\}$ and an element $\mathbf{c} \in \mathcal{C}(\mathcal{I}) \cap \mathbb{B}$ of \mathbb{B} . Then by recalling that the algebra isomorphism $\prod_{j=0}^r \operatorname{pr}_{c_{i_{[j,0]}}}: \widehat{\Gamma}(\mathbf{c})_l \xrightarrow{\sim} \prod_{j=0}^r \mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}$ restricts to the isomorphism

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon \mathfrak{m}_{z_{c_{0}}}^{\widehat{\Gamma}(\mathbf{c})} \xrightarrow{\sim} \mathfrak{m}_{z_{c_{0}}}^{(l_{0})} \times \prod_{j=1}^{r} \mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}}$$

of non-unitial \mathbb{C} -algebras, we can see that the isomorphism $\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} : \mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l) \xrightarrow{\sim} \prod_{j=0}^{r} \mathfrak{n}_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j})$ of Lie algebras restricts to the isomorphism

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon \mathfrak{n}_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_{l})_{1} \xrightarrow{\sim} \mathfrak{n}_{l,l+1}(\mathbb{C}[z_{c_{0}}]_{l_{0}})_{1} \times \prod_{j=1}^{r} \mathfrak{n}_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}}).$$

Then through the exponential maps we obtain the isomorphism as complex manifolds,

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_{l})_{1} \xrightarrow{\sim} N_{l,l+1}(\mathbb{C}[z_{c_{0}}]_{l_{0}})_{1} \times \prod_{j=1}^{r} N_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}})_{1}$$

Proposition 9.11. The map

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} \colon N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_{l})_{1} \xrightarrow{\sim} N_{l,l+1}(\mathbb{C}[z_{c_{0}}]_{l_{0}})_{1} \times \prod_{j=1}^{r} N_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_{j}})_{1}$$

is a Lie group isomorphism.

Proof. As well as Proposition 9.9, we can see that this map is compatible with the group multiplications. \Box

9.6 Subbundle $\mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B})^{\mathrm{rev}}_{\lceil l-1\rceil})$

In section 9.3 we gave a realization of the family $\bigsqcup_{\mathbf{c}\in\mathbb{B}}\mathfrak{u}_{l,l+1}(\widehat{\mathbf{L}}(\mathbf{c}))$ as the trivial vector bundle $\mathfrak{u}_{l,l+1}^{\oplus(l+1)}\times\mathbb{B}\to\mathbb{B}$ with respect to the standard basis

$$\frac{1}{z-c_0}, \frac{1}{(z-c_0)(z-c_1)}, \dots, \frac{1}{(z-c_0)(z-c_1)\cdots(z-c_l)}$$

of $\widehat{\mathcal{L}}(\mathbf{c})$. This vector bundle was denoted by $\mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B})_l)$. Now by reversing the order of c_0, c_1, \ldots, c_l , we obtain another basis

$$\frac{1}{(z-c_l)}, \frac{1}{(z-c_l)(z-c_{l-1})}, \dots, \frac{1}{(z-c_l)(z-c_{l-1})\cdots(z-c_1)}$$

of $\hat{\mathbf{L}}(\mathbf{c})$ for $\mathbf{c} \in \mathbb{B}$ and consider another realization of the family $\bigsqcup_{\mathbf{c}\in\mathbb{B}} \mathfrak{u}_{l,l+1}(\hat{\mathbf{L}}(\mathbf{c}))$ as the vector bundle $\mathfrak{u}_{l,l+1}^{\oplus(l+1)} \times \mathbb{B} \to \mathbb{B}$ with respect to this new basis. We denote this vector bundle by $\mathfrak{u}_{l,l+1}(\hat{\mathbf{L}}(\mathbb{B})_l)^{\text{rev}}$. There uniquely exists base change matrix $M_{\mathbf{c}} \in \mathrm{GL}_{l+1}(\mathbb{C})$ between above two basis of $\hat{\mathbf{L}}(\mathbf{c})$ for each $\mathbf{c} \in \mathbb{B}$, and $M_{\mathbf{c}}$ depends holomorphically on $\mathbf{c} \in \mathbb{B}$. Thus we obtain the holomorphic map $\mathbb{B} \ni \mathbf{c} \mapsto M_{\mathbf{c}} \in \mathrm{GL}_{l+1}(\mathbb{C})$ and then it defines the standard isomorphism

$$\phi_M \colon \mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B})_l)^{\operatorname{rev}} \xrightarrow{\sim} \mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B})_l)$$

as vector bundles over \mathbb{B} .

Let us consider the subbundle of $\mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B}))^{\mathrm{rev}}$ defined by

$$\mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B})_{l-1})^{\text{rev}} := \left\{ \frac{X_l}{z-c_l} + \frac{X_{l-1}}{(z-c_l)(z-c_{l-1})} + \dots + \frac{X_0}{(z-c_l)\cdots(z-c_0)} \in \bigsqcup_{\mathbf{c}\in\mathbb{B}} \mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbf{c})_l) \, \middle| \, X_0 = 0 \right\}.$$

Then we denote the corresponding subbundle of $\mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B}))$ under the isomorphism ϕ_M by

$$\mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B})^{\operatorname{rev}}_{[l-1]}) := \phi_M(\mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbb{B})_{l-1})^{\operatorname{rev}}).$$

Proposition 9.12. Let us take a partition $\mathcal{I}: I_0, \ldots, I_r$ of $\{0, 1, \ldots, k\}$. Also take $\mathbf{c} \in C(\mathcal{I}) \cap \mathbb{B}$. Let $\prod_{j=0}^r \operatorname{pr}_{c_{i_{[j,0]}}}: \mathfrak{g}(\widehat{\mathbf{L}}(\mathbf{c})_l) \xrightarrow{\sim} \prod_{j=0}^r \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{l_j})$ be the isomorphism introduced in Section 9.4. Let us denote the fiber of the vector bundle $\mathfrak{u}_{l,l+1}(\widehat{\mathbf{L}}(\mathbb{B})_{[l-1]}^{\operatorname{rev}})$ at \mathbf{c} by $\mathfrak{u}_{l,l+1}(\widehat{\mathbf{L}}(\mathbf{c})_{[l-1]}^{\operatorname{rev}})$.

Then by restricting the map $\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}}$ to $\mathfrak{u}_{l,l+1}(\widehat{L}(\mathbf{c})_{[l-1]}^{\operatorname{rev}})$, we obtain the isomorphism

$$\prod_{j=0}^{r} \operatorname{pr}_{c_{i_{[j,0]}}} : \mathfrak{u}_{l,l+1}(\widehat{\mathcal{L}}(\mathbf{c})_{[l-1]}^{\operatorname{rev}}) \xrightarrow{\sim} \mathfrak{u}_{l,l+1}(\mathbb{C}[z_{c_{0}}^{-1}]_{l_{0}-1}) \times \prod_{j=1}^{r} \mathfrak{u}_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{l_{j}}).$$

Proof. Let us consider the vector subspace of $\widehat{L}(\mathbf{c})_l$ defined by

$$\widehat{\mathcal{L}}(\mathbf{c})_{[l-1]}^{\mathrm{rev}} := \left\{ \frac{x_{;}}{z - c_{l}} + \frac{x_{l-1}}{(z - c_{l})(z - c_{l-1})} + \dots + \frac{x_{0}}{(z - c_{l}) \cdots (z - c_{0})} \in \widehat{\mathcal{L}}(\mathbf{c})_{l} \, \middle| \, x_{0} = 0 \right\}.$$

Since the zero-th component c_0 of **c** does not appear in this definition, then by setting $\mathbf{c}_{\geq 1} := (c_1, c_2, \ldots, c_k) \in \mathbb{C}^k$, we can regard

$$\widehat{\mathrm{L}}(\mathbf{c})_{[l-1]}^{\mathrm{rev}} = \widehat{\mathrm{L}}(\mathbf{c}_{\geq 1})_{l-1}.$$

For the partition $\mathcal{I}: I_0, I_1, \ldots, I_r$, let $I_j^{\geq 1} := I_j \cap \{1, 2, \ldots, k\}$ for $j = 0, 1, \ldots, r$ and define the partition $\mathcal{I}_{\geq 1}: I_0^{\geq 1}, I_1^{\geq 1}, \ldots, I_r^{\geq 1}$ of $\{1, 2, \ldots, k\}$ where we allow $I_j^{\geq 1} = \emptyset$. Recall that in our convention we always assume $0 \in I_0$. Thus $I_0^{\geq 1} = I_0 \setminus \{0\}$ and $I_j^{\geq 1} = I_j$ for $j \geq 1$, and we can write $\mathcal{I}_{\geq 1}: I_0^{\geq 1}, I_1, \ldots, I_r$. Then we can regard $\mathbf{c}_{\geq 1} \in C(\mathcal{I}_{\geq 1}) \subset \{(a_1, a_2, \ldots, a_k) \in \mathbb{C}^k\}$. Therefore as we saw in Section 9.2, we have the direct sum decomposition

$$\widehat{\mathcal{L}}(\mathbf{c})_{[l-1]}^{\mathrm{rev}} = \widehat{\mathcal{L}}(\mathbf{c}_{\geq 1})_{l-1} = \mathbb{C}[z_{c_0}]_{l_j-1} \oplus \bigoplus_{j=1}^r \mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}.$$

This decomposition induces our desired isomorphism.

10 Deformation of truncated orbit

Let us consider an unramified canonical normal form

$$H dz = \left(\frac{H_k}{z^k} + \dots + \frac{H_1}{z} + H_{\text{res}}\right) \frac{dz}{z} \in \mathfrak{g}(\mathbb{C}[z^{-1}]_k) dz.$$

Let $\mathbb{B}_H \subset \mathbb{C}^{k+1}$ be the complement of hypersurfaces defined in Section 5.1. In this section we shall construct a deformation of the truncated orbit \mathbb{O}_H over the parameter space \mathbb{B}_H .

10.1 Deformation of truncated orbit

Let us consider the product

$$\prod_{l=1}^{k} \left(N_{l,l+1}(\widehat{\Gamma}(\mathbb{B}_{H})_{l})_{1} \times \mathfrak{u}_{l,l+1}(\widehat{\mathrm{L}}(\mathbb{B}_{H})_{[l-1]}^{\mathrm{rev}}) \right)$$

of vector bundles over \mathbb{B}_H defined in the previous section. Let us define a left L_1 -action on this bundle. Recall that each component of this product is a subbundle of the trivial bundles $(\mathbf{n}_{l,l+1})^{\oplus (l+1)} \times \mathbb{B}_H$ or $(\mathbf{u}_{l,l+1})^{\oplus (l+1)} \times \mathbb{B}_H$. Then since $\mathbf{n}_{l,l+1}$ and $\mathbf{u}_{l,l+1}$ have the adjoint L_1 -actions, we can extend these adjoint actions diagonally on $(\mathbf{n}_{l,l+1})^{\oplus (l+1)}$ and $(\mathbf{u}_{l,l+1})^{\oplus (l+1)}$. Thus we obtain the L_1 -actions on $(\mathbf{n}_{l,l+1})^{\oplus (l+1)} \times \mathbb{B}_H$ and $(\mathbf{u}_{l,l+1})^{\oplus (l+1)} \times \mathbb{B}_H$ where L_1 acts trivially on \mathbb{B}_H . It can be directly checked that subbundles $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B}_H)_l)_1$ and $\mathbf{u}_{l,l+1}(\widehat{\Gamma}(\mathbb{B}_H)_{l-1}^{\mathrm{rev}})$ are preserved under these L_1 -actions. Therefore we can extend these L_1 -actions to the whole product bundle on which L_1 acts diagonally on each component through the above action.

Let us also consider the trivial fiber bundle $\mathbb{O}_{H_{\text{res}}}^{L_1} \times \mathbb{B}_H \to \mathbb{B}_H$ with the left L_1 -action defined by multiplication from the left on $L_1/\text{Stab}_{L_1}(H_{\text{res}}) \cong \mathbb{O}_{H_{\text{res}}}^{L_1}$. We regard this bundle as a subspace of the vector bundle $\mathfrak{g}(\hat{L}(\mathbb{B}_H))$ by the injective immersion

$$(L_1/\operatorname{Stab}_{L_1}(H_{\operatorname{res}})) \times \mathbb{B}_H \ni ([g], \mathbf{c}) \longmapsto \frac{H_k}{\prod_{j=0}^k (z - c_j)} + \frac{H_{k-1}}{\prod_{j=0}^{k-1} (z - c_j)} + \dots + \frac{\operatorname{Ad}(g)(H_{\operatorname{res}})}{z - c_0} \in \bigsqcup_{\mathbf{c} \in \mathbb{B}_H} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbf{c})) = \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H)).$$

Here we notice that $\operatorname{Ad}(g)(H_i) = H_i$ for $g \in L_1$ and $i = 1, 2, \ldots, k$.

Let us moreover consider the trivial bundle $G \times \mathbb{B}_H \to \mathbb{B}_H$. By the multiplication from the right, we define the left L_1 -action on G, i.e., $l \cdot g := gl^{-1}$ for $g \in G$ and $l \in L_1$, and thus the $G \times \mathbb{B}_H \to \mathbb{B}_H$ is equipped with the left L_1 -action as well.

Now we define the product of all these fiber bundles over \mathbb{B}_H with the left L_1 -action by

$$G \times \prod_{l=1}^{k} \left(N_{l,l+1}(\widehat{\Gamma}(\mathbb{B}_{H})_{l})_{1} \times \mathfrak{u}_{l,l+1}(\widehat{\mathrm{L}}(\mathbb{B}_{H})_{[l-1]}^{\mathrm{rev}}) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_{1}}.$$

We shall consider the quotient of the total space of this bundle under the L_1 -action. Since L_1 acts freely and properly on G, Lemma 7.15 assures that the L_1 -action on the total space of this fiber bundle is free and proper. Thus we can consider the quotient manifold

$$\mathbb{O}_{H,\mathbb{B}_{H}} := L_{1} \setminus \left(G \times \prod_{l=1}^{k} \left(N_{l,l+1}(\widehat{\Gamma}(\mathbb{B}_{H})_{l})_{1} \times \mathfrak{u}_{l,l+1}(\widehat{\mathrm{L}}(\mathbb{B}_{H})_{[l-1]}^{\mathrm{rev}}) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_{1}} \right)$$

with the projection map

 $\pi_{\mathbb{B}_H}\colon \mathbb{O}_{H,\mathbb{B}_H}\longrightarrow \mathbb{B}_H$

induced by the bundle projection.

Proposition 10.1. The fiber at $\mathbf{0} \in \mathbb{B}_H$ is isomorphic to the truncated orbit \mathbb{O}_H , i.e.,

$$\pi_{\mathbb{B}_H}^{-1}(\mathbf{0}) \cong \mathbb{O}_H$$

Proof. From the definitions of $N_{l,l+1}(\widehat{\Gamma}(\mathbb{B}_H)_l)_1$ and $\mathfrak{u}_{l,l+1}(\widehat{L}(\mathbb{B}_H)_{l-1}^{\mathrm{rev}})$, we obtain

$$\pi_{\mathbb{B}_{H}}^{-1}(\mathbf{0}) = L_{1} \setminus \left(G \times \prod_{l=1}^{k} \left(N_{l,l+1}(\mathbb{C}[z]_{l})_{1} \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1}) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_{1}} \right).$$

As we saw in Section 8.5, the right hand side is isomorphic to \mathbb{O}_H .

As a fiber bundle, $\mathbb{O}_{H,\mathbb{B}_H}$ is trivial, and no deformation occurs to each fiber. However group structures of $N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l)$ and their actions on $\mathfrak{u}_{l,l+1}(\widehat{L}(\mathbf{c})_{[l-1]}^{\mathrm{rev}})$ are deformed according to the change of parameter $\mathbf{c} \in \mathbb{B}_H$.

10.2 Deformation of the moment map $\mu_{\mathbb{O}_H \downarrow G}$

We shall define the map

$$\mu_{\mathbb{O}_H \downarrow G, \mathbb{B}_H} \colon \mathbb{O}_{H, \mathbb{B}_H} \longrightarrow \mathfrak{g}$$

which gives a deformation of the moment map $\mu_{\mathbb{O}_H \downarrow G} \colon \mathbb{O}_H \to \mathfrak{g}$. For this purpose we shall define the bundle map

$$\iota_{\mathbb{O}_{H,\mathbb{B}_{H}}}:\mathbb{O}_{H,\mathbb{B}_{H}}\longrightarrow\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H}))$$

which will be a deformation of the immersion $\iota_{\mathbb{O}_H} \colon \mathbb{O}_H \hookrightarrow \mathfrak{g}(\mathbb{C}[z^{-1}]_k)$, and the map $\mu_{\mathbb{O}_H \downarrow G, \mathbb{B}_H}$ will be defined as the composition of $\iota_{\mathbb{O}_{H, \mathbb{B}_H}}$ and res : $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H)) \to \mathfrak{g}$ defined in Section 9.3.

Let us take $\Xi \in \mathbb{O}_{H,\mathbb{B}_H}$ and choose its representative $(g, (n_l, \nu_l)_{l=1,\dots,k}, \eta) \in G \times \prod_{l=1}^k (N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_{l-1})) \times \mathbb{O}_{H_{res}}^{L_1}$. Then let us define $\iota_{\mathbb{O}_{H,\mathbb{B}_H}}(\Xi)$ inductively as follows.

Step 1. Let us define

$$\iota_{\mathbb{O}_{H,\mathbb{B}_{H}}}^{(1)}(\Xi) := \mathrm{Ad}^{*}(n_{1})(\nu_{1} + \eta) \in \mathfrak{g}(\widehat{\mathrm{L}}(\mathbb{B}_{H}))$$

where we regard $\nu_1, \eta \in \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H))$ and Ad^* is the coadjoint action of $N_1(\widehat{\Gamma}(\mathbb{B}_H))$ on $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H))$.

Step l for $1 < l \leq k$. Let us define

$$\iota_{\mathbb{O}_{H,\mathbb{B}_{H}}}^{(l)}(\Xi) := \mathrm{Ad}^{*}(n_{l})(\nu_{l} + \iota_{\mathbb{O}_{H,\mathbb{B}_{H}}}^{(l-1)}(\Xi)) \in \mathfrak{g}(\widehat{\mathrm{L}}(\mathbb{B}_{H})).$$

where we regard $\nu_l \in \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H))$ and Ad^* is the coadjoint action of $N_l(\widehat{\Gamma}(\mathbb{B}_H))$ on $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H))$. <u>Step k + 1</u>. Let us regard the trivial bundle $G \times \mathbb{B}_H \to \mathbb{B}_H$ as a Lie groupoid in the trivial way. Then the natural G-action on the each fiber $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbf{c}))$ at $\mathbf{c} \in \mathbb{B}_H$ defines the coadjoint action Ad^* of $G \times \mathbb{B}_H$ on $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H))$.

Then we define

$$\iota_{\mathbb{O}_{H,\mathbb{B}_{H}}}^{(k+1)}(\Xi) := \mathrm{Ad}^{*}(g)(\iota_{\mathbb{O}_{H,\mathbb{B}_{H}}}^{(k)}(\Xi)) \in \mathfrak{g}(\widehat{\mathrm{L}}(\mathbb{B}_{H})).$$

We can see that the definition of $\iota_{\mathbb{O}_{H,\mathbb{B}_{H}}}^{(k+1)}(\Xi)$ does not depend on the choice of representatives of Ξ as we saw in Section 8.6. Thus we obtain the bundle map

$$\iota_{\mathbb{O}_{H,\mathbb{B}_{H}}} := \iota_{\mathbb{O}_{H,\mathbb{B}_{H}}}^{(k+1)} : \mathbb{O}_{H,\mathbb{B}_{H}} \longrightarrow \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H}))$$

and the desired map

$$\mu_{\mathbb{O}_H \downarrow G, \mathbb{B}_H} \colon \mathbb{O}_{H, \mathbb{B}_H} \xrightarrow{\iota_{\mathbb{O}_{H, \mathbb{B}_H}}} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H)) \xrightarrow{\operatorname{res}} \mathfrak{g}.$$

Let us consider the left G-action on $\mathbb{O}_{H,\mathbb{B}_H}$ defined by the multiplication from the left on the first component of $G \times \left(\left(\prod_{l=1}^k (N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_l)_1) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_1} \right)$. Then we can see that the map $\mu_{\mathbb{O}_H \downarrow G,\mathbb{B}_H}$ is G-equivariant.

Proposition 10.2. The map $\mu_{\mathbb{O}_H \downarrow G, \mathbb{B}_H}$ is a deformation of the moment map $\mu_{\mathbb{O}_H \downarrow G}$, namely we have

$$\mu_{\mathbb{O}_H \downarrow G, \mathbb{B}_H}|_{\pi_{\mathbb{B}_H}^{-1}(\mathbf{0})} = \mu_{\mathbb{O}_H \downarrow G}$$

under the identification $\pi_{\mathbb{B}_H}^{-1}(\mathbf{0}) \cong \mathbb{O}_H$ given in Proposition 10.1.

Proof. By recalling the description of $\iota_{\mathbb{O}_H}$ given in Section 8.6, we can see that the restriction map $\iota_{\mathbb{O}_{H,\mathbb{B}_H}}|_{\pi_{\mathbb{B}_H}^{-1}(\mathbf{0})}$ coincides with the isomorphism

$$L_1 \setminus \left(G \times \left(\left(\prod_{l=1}^k (N_{l,l+1}(\mathbb{C}[z]_l)_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z^{-1}]_l)_1) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_1} \right) \right) \xrightarrow{\sim} \mathbb{O}_H \subset \mathfrak{g}(\mathbb{C}[z^{-1}]_k).$$

Also the map res : $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H)) \to \mathfrak{g}$ coincides with res : $\mathfrak{g}(\mathbb{C}[z^{-1}]_k) \to \mathfrak{g}$ on the fiber at $\mathbf{0} \in \mathbb{B}_H$. Thus we obtain the desired equation.

10.3 Fibers on each stratum of \mathbb{B}_H

Let us take a partition $\mathcal{I}: I_0, I_1, \ldots, I_r$ of $\{0, 1, \ldots, k\}$ as in Section 5.2 and write $I_j = \{i_{[j,0]} < i_{[j,1]} < \cdots < i_{[j,k_j]}\}$. Also take $\mathbf{c} \in C(\mathcal{I}) \cap \mathbb{B}_H$ and consider $H(\mathbf{c})$. Then we obtain the partial fraction decomposition

$$H(\mathbf{c}) = H(\mathbf{c})_0 + H(\mathbf{c})_1 + \dots + H(\mathbf{c})_r \text{ with } H(\mathbf{c})_j \in \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{k_j})$$

and $H(\mathbf{c})$ is regarded as the collection of canonical forms $(H(\mathbf{c})_j)_{j=0,\dots,r}$.

Proposition 10.3. Let $\pi_{\mathbb{B}_H} : \mathbb{O}_{H,\mathbb{B}_H} \to \mathbb{B}_H$ be the deformation of \mathbb{O}_H . Then the fiber $\pi_{\mathbb{B}_H}^{-1}(\mathbf{c})$ at $\mathbf{c} \in C(\mathcal{I}) \cap \mathbb{B}_H$ is isomorphic to an open dense subset of $\prod_{i=0}^r \mathbb{O}_{H(\mathbf{c})_i}$.

Before giving a proof of this proposition, we make some preparations. Let us consider the direct product of Lie groups $\prod_{j=0}^{r} G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})$. which naturally acts on the product space $\prod_{j=0}^{r} \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{k_j})$. Then by regarding $\mathbf{H}(\mathbf{c}) = (H(\mathbf{c})_j)_{j=0,\dots,r} \in \prod_{j=0}^{r} \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{k_j})$, we obtain

$$\prod_{j=0}^{r} \mathbb{O}_{H(\mathbf{c})_{j}} = \mathbb{O}_{\mathbf{H}(\mathbf{c})}^{\prod_{j=0}^{r} G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_{j}})}$$

where the right hand side is the $\prod_{j=0}^{r} G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_1})$ -orbit through $\mathbf{H}(\mathbf{c})$. We shall give a similar description of the orbit $\mathbb{O}_{\mathbf{H}(\mathbf{c})}^{\prod_{j=0}^{r} G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})}$ as that of the truncated orbit \mathbb{O}_H given in Proposition 8.1.

Let us consider the normal subgroup

$$G(\mathbb{C}[z_{c_0}]_{k_0})_1 \times \prod_{j=1}' G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})$$

of $\prod_{j=0}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})$ and the short exact sequence

$$1 \to G(\mathbb{C}[z_{c_0}]_{k_0})_1 \times \prod_{j=1}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j}) \to \prod_{j=0}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j}) \to G \to 1$$

induced by the inclusion. Then the diagonal embedding

$$G \ni g \longmapsto (g, \dots, g) \in \prod_{j=0}^{r} G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})$$

gives a right splitting of this exact sequence and we obtain the semidirect product decomposition

$$\prod_{j=0}^{r} G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_{1}}) = G \ltimes \left(G(\mathbb{C}[z_{c_{0}}]_{k_{0}})_{1} \times \prod_{j=1}^{r} G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_{j}}) \right).$$

Lemma 10.4. Let us consider the subgroup $L_1 \ltimes \left(G(\mathbb{C}[z_{c_0}]_{k_0})_1 \times \prod_{j=1}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j}) \right)$ of $\prod_{j=0}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})$. Then the orbit of $\mathbf{H}(\mathbf{c})$ under the action of this subgroup is

$$\mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}])_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j}.$$

Proof. Recall that $L_1 \subset \operatorname{Stab}_{G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})}(H(\mathbf{c})_j)$ for $j = 1, \ldots, r$ by Proposition 5.6. Thus the orbit of $\mathbf{H}(\mathbf{c})$ under the action of $L_1 \ltimes \left(G(\mathbb{C}[z_{c_0}]_{k_0})_1 \times \prod_{j=1}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})\right)$ coincides with that of $(L_1 \ltimes G(\mathbb{C}[z_{c_0}]_{k_0})_1) \times \prod_{j=1}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})$, that is, $\mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}])_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j}$.

Let us consider the left L_1 -action on G by the multiplication from the right. Also on $\mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}])_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j}$ we can consider the L_1 -action by the diagonal coadjoint action. Then we can consider

$$G \times_{L_1} \left(\mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}])_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j} \right) := L_1 \setminus \left(G \times \left(\mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}])_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j} \right) \right),$$

the fiber bundle over G/L_1 . Then as well as the diagram (4) we obtain the commutative diagram

where $\tilde{\psi}$, $\bar{\psi}$ and ψ are similarly defined as them in the diagram (4), and the upper left vertical arrow is the quotient map

$$\pi \colon L_1 \ltimes \left(G(\mathbb{C}[z_{c_0}]_{k_0})_1 \times \prod_{j=1}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j}) \right) \longrightarrow \mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}])_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j}$$

defined by Lemma 10.4.

Proposition 10.5. The above map

$$\psi \colon G \times_{L_1} \left(\mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}])_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j} \right) \longrightarrow \mathbb{O}_{\mathbf{H}(\mathbf{c})}^{\prod_{j=0}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})}$$

is an isomorphism as complex manifolds.

Proof. We can prove similarly as Proposition 8.1. The surjectivity comes from the definition. The injectivity comes from the fact $\operatorname{Stab}_G(\mathbf{H}(\mathbf{c})) = \operatorname{Stab}_G(H) \subset L_1$.

Now we are ready to give a proof of Proposition 10.3.

Proof of Proposition 10.3. By regarding the trivial bundle $\mathbb{O}_{H_{\text{res}}}^{L_1} \times \mathbb{B}_H$ as a subbundle of $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_H))$ as in Section 10.1, we can see that under the partial fraction decomposition $\mathfrak{g}(\widehat{\mathcal{L}}(\mathbf{c})) \cong \prod_{j=0}^r \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{k_j})$, the fiber $\mathbb{O}_{H_{\text{res}}}^{L_1} \times \{\mathbf{c}\} \subset \mathfrak{g}(\widehat{\mathcal{L}}(\mathbf{c}))$ corresponds to

$$\mathbb{O}_{H(\mathbf{c})_0}^{L_1} \times \prod_{j=1}^r \{H(\mathbf{c})_j\} \subset \prod_{j=0}^r \mathfrak{g}(\mathbb{C}[z_{c_{i_{[j,0]}}}^{-1}]_{k_j})$$

since $L_1 \subset \operatorname{Stab}_{G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})}(H(\mathbf{c})_j)$ for $j = 1, \ldots, r$ by Proposition 5.6.

Also consider the fiber $\prod_{l=1}^{k} (N_{l,l+1}(\hat{\Gamma}(\mathbf{c})_l) \times \mathfrak{u}_{l,l+1}(\hat{L}(\mathbf{c})_{[l-1]}^{rev}))$ of the vector bundle $\prod_{l=1}^{k} (N_{l,l+1}(\hat{\Gamma}(\mathbb{B}_H)_l) \times \mathfrak{u}_{l,l+1}(\hat{L}(\mathbb{B}_H)_{[l-1]}^{rev}))$. Then by combining the isomorphisms in Propositions 9.11 and 9.12, we obtain the isomorphism

$$\begin{split} \prod_{l=1}^{k} (N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_{l}) \times \mathfrak{u}_{l,l+1}(\widehat{\mathbf{L}}(\mathbf{c})_{[l-1]}^{\mathrm{rev}})) & \xrightarrow{\sim} \prod_{l=1}^{k_{0}} (N_{l,l+1}(\mathbb{C}[z_{c_{0}}]_{l_{0}})_{1} \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z_{c_{0}}^{-1}]_{l_{0}-1})) \\ & \times \prod_{j=1}^{r} \left(\prod_{l=1}^{k_{j}} (N_{l,l+1}(\mathbb{C}[z_{c_{i}_{[j,0]}}]_{l_{j}}) \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z_{c_{i}_{[j,0]}}]_{l_{j}})) \right). \end{split}$$

Therefore the fiber $\prod_{l=1}^{k} (N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_l) \times \mathfrak{u}_{l,l+1}(\widehat{\mathbf{L}}(\mathbf{c})_{[l-1]}^{\mathrm{rev}})) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_1}$ of the fiber bundle $\prod_{l=1}^{k} (N_{l,l+1}(\widehat{\Gamma}(\mathbb{B}_H)_l) \times \mathfrak{u}_{l,l+1}(\widehat{\mathbf{L}}(\mathbb{B}_H)_{[l-1]}^{\mathrm{rev}})) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_1}$ is isomorphic to

$$\left(\prod_{l=1}^{k_0} (N_{l,l+1}(\mathbb{C}[z_{c_0}]_{l_0})_1 \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z_{c_0}^{-1}]_{l_0-1})) \times \mathbb{O}_{H(\mathbf{c})_0}^{L_1}\right) \times \prod_{j=1}^r \left(\left(\prod_{l=1}^{k_j} (N_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}) \times \mathfrak{u}_{l,l+1}(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{l_j}))\right) \times \{H(\mathbf{c})_j\}\right).$$

By Theorems 8.2 and 8.12, this is isomorphic to the dense open subset

$$\mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}]_{k_0})_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j}^{G_0(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})}$$

of $\mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}]_{k_0})_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j}$.

As a conclusion, we obtain the diagram

which is checked to be commutative by the L_1 -equivariance of the upper horizontal arrow. Here the horizontal arrows are open embeddings and the vertical arrows are quotient maps. Therefore recalling the isomorphism

$$G \times_{L_1} \left(\mathbb{O}_{H(\mathbf{c})_0}^{L_1 \ltimes G(\mathbb{C}[z_{c_0}]_{k_0})_1} \times \prod_{j=1}^r \mathbb{O}_{H(\mathbf{c})_j} \right) \cong \mathbb{O}_{\mathbf{H}(\mathbf{c})}^{\prod_{j=0}^r G(\mathbb{C}[z_{c_{i_{[j,0]}}}]_{k_j})}$$

in Proposition 10.5, we obtain the desired open embedding

$$\pi_{H}^{-1}(\mathbf{c}) = L_{1} \setminus \left(G \times \left(\prod_{l=1}^{k} \left(N_{l,l+1}(\widehat{\Gamma}(\mathbf{c})_{l}) \times \mathfrak{u}_{l,l+1}(\widehat{L}(\mathbf{c})_{[l-1]}^{\mathrm{rev}}) \right) \times \mathbb{O}_{H_{\mathrm{res}}}^{L_{1}} \right) \right) \xrightarrow{\sim} L_{1} \setminus \left(G \times \left(\mathbb{O}_{H(\mathbf{c})_{0}}^{L_{1} \ltimes G(\mathbb{C}[z_{c_{0}}]_{k_{0}})_{1}} \times \prod_{j=1}^{r} \mathbb{O}_{H(\mathbf{c})_{j}}^{G_{0}(\mathbb{C}[z_{c_{i}}]_{j,0}]^{-1}_{k_{j}})} \right) \right) \hookrightarrow \mathbb{O}_{\mathbf{H}(\mathbf{c})}^{\prod_{j=0}^{r} G(\mathbb{C}[z_{c_{i}}]_{k_{j}})_{j}}.$$

Let us discuss the compatibility of the map $\mu_{\mathbb{O}_{H,\mathbb{B}_{H}}}$ and the embedding $\iota: \pi_{\mathbb{B}_{H}}^{-1}(\mathbf{c}) \hookrightarrow \prod_{i=0}^{r} \mathbb{O}_{H(\mathbf{c})_{i}}$ in Proposition 10.3. Define the map

$$\mu_{\mathbf{H}(\mathbf{c})} \colon \prod_{j=0}^{r} \mathbb{O}_{H(\mathbf{c})_{j}} \longrightarrow \mathfrak{g} (\xi_{j})_{j=0,\dots,r} \longmapsto \sum_{j=0}^{r} \mu_{\mathbb{O}_{H(\mathbf{c})_{j}} \downarrow G}(\xi_{j})$$

which is considered as the moment map with respect to the diagonal G-action under the identification $\mathfrak{g} \cong \mathfrak{g}^*$ via the trace pairing as we explained in Section 7.4.

Proposition 10.6. Let $\iota: \pi_{\mathbb{B}_H}^{-1}(\mathbf{c}) \hookrightarrow \prod_{j=0}^r \mathbb{O}_{H(\mathbf{c})_j}$ be the open embedding in Proposition 10.3. Then the following diagram is commutative,



Proof. By Proposition 9.10 and the definition of $\iota_{\mathbb{O}_H,\mathbb{B}_H}$, we obtain the commutative diagram,

Then since $\mu_{\mathbb{O}_H \downarrow G, \mathbb{B}_H} = \text{res} \circ \iota_{\mathbb{O}_{H, \mathbb{B}_H}}$ and $\mu_{\mathbf{H}(\mathbf{c})} = \sum_{j=0}^r \underset{z=c_{i_{[j,0]}}}{\text{res}} \circ \iota_{\mathbb{O}_{H(\mathbf{c})_j}}$, this diagram and the equation (6) give us the desired commutative diagram.

11 Deformation of moduli spaces of G-connections on \mathbb{P}^1 via unfolding of irregular singularities

Now we are ready to provide the precise statement of our main theorem and give a proof of it.

Let $D = \sum_{i=1}^{d} (k_{a_i} + 1) \cdot a_i$ be an effective divisor of \mathbb{P}^1 and we may assume that the underlying set |D| is contained in \mathbb{C} under the projective transformation. Let $\mathbf{H} = (H^{(a)})_{a \in |D|} \in \prod_{a \in |D|} \mathfrak{g}(\mathbb{C}[z_a^{-1}]_{k_a})$ be a collection of canonical forms with $\sum_{a \in |D|} \operatorname{res}_{z=a}(H^{(a)}) \in \mathfrak{g}_{ss}$. Let $\mathbb{B}_{\mathbf{H}} = \prod_{a \in |D|} \mathbb{B}_{H^{(a)}}$. We consider the product manifold $\prod_{a \in |D|} \mathbb{O}_{H^{(a)}, \mathbb{B}_{H^{(a)}}}$ with the projection

$$\pi_{\mathbb{B}_{\mathbf{H}}} \colon \prod_{a \in |D|} \mathbb{O}_{H^{(a)}, \mathbb{B}_{H^{(a)}}} \longrightarrow \prod_{a \in |D|} \mathbb{B}_{H^{(a)}}.$$

Let

$$\iota_{\mathbf{H},\mathbb{B}_{\mathbf{H}}} \colon \prod_{a \in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}} \longrightarrow \prod_{a \in |D|} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}}))$$

be the product map of the immersions $\iota_{\mathbb{O}_{H_a},\mathbb{B}_{H^{(a)}}} : \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}} \to \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}}))$ for $a \in |D|$. Let

$$\left(\prod_{a\in |D|}\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}}))\right)^{\mathrm{t}}$$

be the totality of stable points under the diagonal coadjoint action of G. Here we notice that under the trivialization $\prod_{a \in |D|} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}})) \cong \left(\bigoplus_{a \in |D|} \mathfrak{g}^{\oplus(k_a+1)}\right) \times \mathbb{B}_{\mathbf{H}}$, we have

$$\left(\prod_{a\in|D|}\mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}}))\right)^{\mathrm{ir}}\cong\left(\bigoplus_{a\in|D|}\mathfrak{g}^{\oplus(k_a+1)}\right)^{\mathrm{ir}}\times\mathbb{B}_{\mathbf{H}}$$

where $\left(\bigoplus_{a\in |D|} \mathfrak{g}^{\oplus(k_a+1)}\right)^{\text{ir}}$ is the set of irreducible elements defined in Section 7.3. We define the open subset of $\prod_{a\in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}^{H_{(a)}}}$ by

$$\left(\prod_{a\in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}}\right)^{\mathrm{ir}} := \iota_{\mathbf{H},\mathbb{B}_{H}}^{-1} \left(\left(\prod_{a\in |D|} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}}))\right)^{\mathrm{ir}} \right).$$

For the moment map

$$\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}} \colon \prod_{a \in |D|} \mathbb{O}_{H_{a},\mathbb{B}_{H^{(a)}}} \ni (X_{a})_{a \in |D|} \longmapsto \sum_{a \in |D|} \mu_{\mathbb{O}_{H^{(a)} \downarrow G,\mathbb{B}_{H^{(a)}}}}(X_{a}) \in \mathfrak{g}^{*},$$

we denote the restriction of the map on $\left(\prod_{a\in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}}\right)^{\mathrm{ir}}$ by $\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}}$.

Definition 11.1 (Deformation of $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}}$). We denote the quotient spaces of level zero sets of $\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}$ and $\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}}$ under the *G*-action by

$$\mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}} := G \setminus \mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{-1}(0), \qquad \qquad \mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}} := G \setminus (\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0).$$

The projection $\pi_{\mathbb{B}_{\mathbf{H}}} \colon \prod_{a \in |D|} \mathbb{O}_{H^{(a)}, \mathbb{B}_{H^{(a)}}} \to \mathbb{B}_{\mathbf{H}}$ induces projections $\pi_{\mathbb{B}_{\mathbf{H}}} \colon \mathcal{M}_{\mathbf{H}, \mathbb{B}_{\mathbf{H}}} \to \mathbb{B}_{\mathbf{H}}$ and $\pi_{\mathbb{B}_{H}} \colon \mathcal{M}_{\mathbf{H}, \mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}} \to \mathbb{B}_{\mathbf{H}}$

Proposition 11.2. Suppose $\mathcal{M}_{\mathbf{H}}^{ir} \neq \emptyset$. Then $\mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{ir}$ is a complex orbifold and the projection map $\pi_{\mathbb{B}_{\mathbf{H}}} : \mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{ir} \to \mathbb{B}_{\mathbf{H}}$ is a submersion.

We make some preparations for the proof of this proposition.

Lemma 11.3. The projection $\pi_{\mathbb{B}_{\mathbf{H}}}$: $\left(\prod_{a \in |D|} \mathbb{O}_{H^{(a)}, \mathbb{B}_{H^{(a)}}}\right)^{ir} \to \mathbb{B}_{\mathbf{H}}$ is a submersion.

Proof. Since the fiber bundle $\mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}}$ has the trivialization $\mathbb{O}_{H^{(a)}} \times \mathbb{B}_{H^{(a)}}$ for each $a \in |D|$, we can identify $\prod_{a \in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}} \cong \left(\prod_{a \in |D|} \mathbb{O}_{H^{(a)}}\right) \times \mathbb{B}_{\mathbf{H}}$ under which the projection $\pi_{\mathbb{B}_{\mathbf{H}}} \colon \prod_{a \in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}} \to \mathbb{B}_{\mathbf{H}}$ corresponds to the second projection that is obviously a submersion. Then since $\left(\prod_{a \in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}}\right)^{\mathrm{ir}}$ is an open submanifold of $\prod_{a \in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}}$, the restriction map $\pi_{\mathbb{B}_{\mathbf{H}}} \colon \left(\prod_{a \in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}}\right)^{\mathrm{ir}} \to \mathbb{B}_{\mathbf{H}}$ is still a submersion. \Box

Lemma 11.4. Suppose that $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}} \neq \emptyset$. Then $0 \in \mathfrak{g}^*$ is a regular value of $\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}}$.

Proof. Note that $(\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0) \neq \emptyset$ by the assumption. Let us take $m \in (\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0)$ and put $\mathbf{c} = \pi_{\mathbb{B}_{\mathbf{H}}}(m) \in \mathbb{B}_{\mathbf{H}}$. Then we have

$$\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}}|_{\pi_{\mathbb{B}_{\mathbf{H}}}^{-1}(\mathbf{c})} = \mu_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}}$$

by Proposition 10.6. Since dim $\operatorname{Stab}_{G}(m) = 0$ and $\mu_{\mathbf{H}(\mathbf{c})}^{\operatorname{ir}}$ is a moment map with respect to the *G*-action, $m \in (\mu_{\mathbf{H}(\mathbf{c})}^{\operatorname{ir}})^{-1}(0)$ is a regular point of $\pi_{\mathbb{B}_{\mathbf{H}}}^{-1}(\mathbf{c})$ with respect to $\mu_{\mathbf{H}(\mathbf{c})}^{\operatorname{ir}}$. Since $\pi_{\mathbb{B}_{\mathbf{H}}}^{-1}(\mathbf{c})$ is a closed submanifold of $\left(\prod_{a \in |D|} \mathbb{O}_{H_{a},\mathbb{B}_{H_{a}}}\right)^{\operatorname{ir}}$ by Lemma 11.3, *m* is also a regular point of $\left(\prod_{a \in |D|} \mathbb{O}_{H_{a},\mathbb{B}_{H_{a}}}\right)^{\operatorname{ir}}$ with respect to $\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\operatorname{ir}}$.

Proof of Proposition 11.2. The first assertion follows from Lemma 11.4 and the fact that the G-action on $\left(\prod_{a\in |D|} \mathbb{O}_{H_a,\mathbb{B}_{H_a}}\right)^{\text{ir}}$ is proper and almost free by Lemma 7.15. For the second assertion, it suffices to show that the projection $\pi := \pi_{\mathbb{B}_{\mathbf{H}}}|_{(\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\text{ir}})^{-1}(0)} : (\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\text{ir}})^{-1}(0) \rightarrow \mathbb{B}_{\mathbf{H}}$ is a submersion. Let us take $m \in (\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\text{ir}})^{-1}(0)$ and put $\mathbf{c} = \pi(m)$. Then by Proposition 10.6, we obtain the commutative diagram

$$\begin{aligned} (\mu_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}})^{-1}(0) & \stackrel{\iota_{\mathbf{c}}}{\longrightarrow} \pi_{\mathbb{B}_{\mathbf{H}}}^{-1}(\mathbf{c}) \\ & \downarrow & \downarrow \\ (\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0) & \stackrel{\iota_{\mathbb{B}_{\mathbf{H}}}}{\longrightarrow} \prod_{a \in |D|} \mathbb{O}_{H^{(a)},\mathbb{B}_{H^{(a)}}} \end{aligned}$$

where all the arrows are natural inclusions. This diagram induces the following commu-

tative diagram,

Here the horizontal sequences are exact, since as we saw in Lemma 11.4, m is a regular point of $\pi_{\mathbb{B}_{\mathbf{H}}}^{-1}(\mathbf{c})$ with respect to the map $\mu_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}}$ and also of $\left(\prod_{a \in |D|} \mathbb{O}_{H_a, \mathbb{B}_{H_a}}\right)^{\mathrm{ir}}$ with respect to the map $\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}$. Also the middle vertical sequence is exact by Lemma 11.3. The right vertical sequence is obviously exact. Then by the snake lemma, we obtain $T_{\mathbf{c}}\mathbb{B}_{H}\cong$ $\begin{array}{l} \operatorname{Coker}\left(T_{m}(\mu_{\mathbf{H}(\mathbf{c})}^{\operatorname{ir}})^{-1}(0) \to T_{m}(\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\operatorname{ir}})^{-1}(0)\right).\\ \operatorname{Since} T_{m}\pi = T_{m}\pi_{\mathbb{B}_{\mathbf{H}}} \circ T_{m}\iota_{\mathbb{B}_{\mathbf{H}}}, \text{ the above diagram shows that the composition map} \end{array}$

$$T_m(\mu_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}})^{-1}(0) \to T_m(\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0) \xrightarrow{T_m \pi} T_{\mathbf{c}} \mathbb{B}_{\mathbf{H}}$$

is the zero map. Thus the map $T_m \pi \colon T_m(\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0) \to T_{\mathbf{c}}\mathbb{B}_{\mathbf{H}}$ factors through the natural projection

$$T_m(\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0) \to \mathrm{Coker}\left(T_m(\mu_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}})^{-1}(0) \to T_m(\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0)\right).$$

Namely, we have the commutative diagram



where the arrow from the bottom to the upper right is the above isomorphism. Thus $T_m \pi \colon T_m(\mu_{\mathbf{H}, \mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0) \to T_{\mathbf{c}} \mathbb{B}_{\mathbf{H}} \text{ is a surjection.}$ Thus we conclude that $\pi \colon (\mu_{\mathbf{H}, \mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0) \to \mathbb{B}_{\mathbf{H}}$ is a submersion.

Suppose $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}} \neq \emptyset$. Let $\pi_{\mathbb{B}_{\mathbf{H}}} \colon \mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}} \to \mathbb{B}_{\mathbf{H}}$ be the projection map. Let $\widetilde{\mathbb{B}}_{\mathbf{H}} := \mathrm{Im} \pi \neq \mathbb{C}$ \emptyset and $\pi_{\widetilde{\mathbb{B}}_{\mathbf{H}}} \colon \mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}} \to \widetilde{\mathbb{B}}_{\mathbf{H}}$ be the projection map induced from $\pi_{\mathbb{B}_{\mathbf{H}}}$.

Theorem 11.5. Suppose $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}} \neq \emptyset$.

- 1. The reduced space $\mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}}$ is a complex orbifold.
- 2. The subspace $\widetilde{\mathbb{B}}_{\mathbf{H}}$ is an open submanifold of \mathbb{C}^{k+1} containing **0**.

- 3. The projection map $\pi_{\widetilde{\mathbb{B}}_{\mathbf{H}}} \colon \mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}} \to \widetilde{\mathbb{B}}_{\mathbf{H}}$ is a surjective submersion.
- 4. Every stratum $\prod_{a \in |D|} C(\mathcal{I}^{(a)})$ of $\prod_{a \in |D|} \mathbb{C}^{k_a+1}$ has the nonempty intersection with the base space $\widetilde{\mathbb{B}}_{\mathbf{H}}$.
- 5. The orbifold $\mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}}$ is a deformation of $\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}}$, i.e., we have

$$\mathcal{M}_{\mathbf{H}}^{\mathrm{ir}} \cong \pi_{\widetilde{\mathbb{B}}_{\mathbf{H}}}^{-1}(\mathbf{0}).$$

6. For each $\mathbf{c} \in \widetilde{\mathbb{B}}_{\mathbf{H}}$ the fiber $\pi_{\widetilde{\mathbb{B}}_{\mathbf{H}}}^{-1}(\mathbf{c})$ is isomorphic to an open dense subspace of $\mathcal{M}_{\mathbf{H}(\mathbf{c})}^{\mathrm{ir}}$.

Proof. The assertion 1 is the first assertion of Proposition 11.2.

By Proposition 11.2, the projection $\pi_{\mathbb{B}_{\mathbf{H}}} : \mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}} \to \mathbb{B}_{\mathbf{H}}$ is a submersion. Thus $\mathbb{B}_{\mathbf{H}} = \mathrm{Im} \pi \neq \emptyset$ is open since a submersion is an open map. Thus we have the assertion 2. The assertion 3 follows from the assertion 2 and Proposition 11.2. The assertion 4 is a consequence from the assertion 2 and Lemma 5.2. The assertion 6 and 7 directly follow from the definition of $\mathcal{M}_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}}$ and Propositions 10.2, 10.3, and 10.6.

Now we can give a proof of Theorem 6.4 as a corollary of Theorem 11.5.

Proof of Theorem 6.4. The implication $2 \Rightarrow 1$ is obvious.

Thus we assume that there exists a solution $d_A = d + A dz$ of the additive Deligne-Simpson problem for **H**. Then there exists the corresponding point $m_A \in (\mu_{\mathbf{H}}^{\mathrm{ir}})^{-1}(0) \subset (\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0)$. Then since $\pi_{\mathbb{B}_{\mathbf{H}}} : (\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0) \to \mathbb{B}_{\mathbf{H}}$ is a submersion, there exists an open neighborhood U of $\mathbf{0} \in \mathbb{B}_{\mathbf{H}}$ and a local section $s : U \to (\mu_{\mathbf{H},\mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0)$ of $\pi_{\mathbb{B}_{\mathbf{H}}}$ such that $s(\mathbf{0}) = m_A$. Then as the composition map

$$U \xrightarrow{s} (\mu_{\mathbf{H}, \mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0) \xrightarrow{\iota_{\mathbf{H}|(\mu_{\mathbf{H}, \mathbb{B}_{\mathbf{H}}}^{\mathrm{ir}})^{-1}(0)}} \prod_{a \in |D|} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}}))$$

we obtain the section $U \to \prod_{a \in |D|} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}}))$ of $\prod_{a \in |D|} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}})) \to \mathbb{B}_{\mathbf{H}}$ and write this section as the $\prod_{a \in |D|} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}}))$ -valued function on U,

$$\left(\frac{A_i^{(a)}(\mathbf{c})}{\prod_{j=0}^i (z_a - c_j^{(a)})}\right)_{a \in |D|} \in \prod_{a \in |D|} \mathfrak{g}(\widehat{\mathcal{L}}(\mathbb{B}_{H^{(a)}})) \quad (\mathbf{c} = ((c_0^{(a)}, c_1^{(a)}, \dots, c_{k_a}^{(a)}))_{a \in |D|} \in U).$$

Then the holomorphic family of G-connections

$$d_{A(\mathbf{c})} = d + \sum_{a \in |D|} \sum_{i=0}^{k_a} \frac{A_i^{(a)}(\mathbf{c})}{\prod_{j=0}^i (z_a - c_j^{(a)})} dz$$

is the desired solution of the additive Deligne-Simpson problem for the family $(\mathbf{H}(\mathbf{c}))_{\mathbf{c}\in U}$ with $d_{A(\mathbf{0})} = d_A$ by the assertion 6 in Theorem 11.5. As a result, we obtain the implication $1 \Rightarrow 2$.

References

- A. Adem, J. Leida, Y. Ruan, Orbifolds and stringy topology, Cambridge Tracts in Math., 171 Cambridge University Press, Cambridge, 2007. xii+149 pp.
- [2] D. Arinkin, Rigid irregular connections on \mathbb{P}^1 , Compos. Math. **146** (2010), no.5, 1323–1338.
- [3] D. Arinkin, *Irreducible connections admit generic oper structures*, preprint (2016) arXiv:1602.08989.
- [4] D. Babbitt, V. Varadarajan, Formal reduction theory of meromorphic differential equations: a group theoretic view, Pacific J. Math. **109** (1983), no.1, 1–80.
- [5] P. Boalch, Symplectic manifolds and isomonodromic deformations, Adv. Math. 163
 (2) (2001), 137–205.
- [6] P. Boalch, Irregular connections and Kac-Moody root systems, 2008, arXiv:0806.1050.
- [7] P. Boalch, Simply-laced isomonodromy systems, Publ. Math. IHES, 116, no. 1 (2012), 1–68.
- [8] C. Bremer, D. Sage, Moduli Spaces of Irregular Singular Connections, IMRN, 2013 Issue 8 (2013), 1800–1872.
- [9] L. Corwin, F. Greenleaf, Representations of nilpotent Lie groups and their applications. Part I. Basic theory and examples, Cambridge Stud. Adv. Math., 18, Cambridge University Press, Cambridge, 1990. viii+269 pp.
- [10] W. Crawley-Boevey, On matrices in prescribed conjugacy classes with no common invariant subspace and sum zero, Duke Math. J. 118, no. 2 (2003), 339–352.
- [11] P. Deligne, Équations différentielles à points singuliers réguliers, Lecture Notes in Math., Vol. 163 Springer-Verlag, Berlin-New York, 1970. iii+133 pp.
- [12] A. Glutsuk, Stokes operators via limit monodromy of generic perturbation, J. Dynam. Control Systems 5 (1999), 101–135.
- [13] I. Gaiur, M. Mazzocco, V. Rubtsov, Isomonodromic deformations: Confluence, Reduction and Quantisation, Comm. in Math. Phys., 400 (2023), 1385–1461.
- [14] M. Goresky, R. MacPherson, Intersection homology theory, Topology 19 (1980), no. 2, 135–162.
- [15] K. Hiroe, Linear differential equations on the Riemann sphere and representations of quivers, Duke Math. J., 166, 855–935, (2017).
- [16] K. Hiroe, Unfolding of spectral types, Josai Math. Monographs 12 (2020), 53–67.
- [17] K. Hiroe, Index of rigidity of differential equations and Euler characteristic of their spectral curves, J. Geom. Phys. 162 (2021), Paper No. 104060, 16 pp.
- [18] K. Hiroe, H. Kawakami, A. Nakamura, H. Sakai, 4-dimensional Painlevé type equations, MSJ memoirs, 37, 2018.

- [19] K. Hiroe, D. Yamakawa, Moduli spaces of meromorphic connection and quiver varieties, Adv. Math., 266 (2014), 120–151.
- [20] M. Hukuhara, Sur les points singuliers des équations diférentielles linéaires. III, Mem. Fac. Sci. Kyusyu Imp. Univ. A 2 (1942) 125–137.
- [21] J. Hurtubise, C. Rousseau, Moduli space for generic unfolded differential linear systems, Adv. Math. 307 (2017), 1268–1323.
- [22] M. Inaba, Unfolding of the unramified irregular singular generalized isomonodromic deformation, Bull. Sci. Math. 157 (2019), 102795, 121 pp.
- [23] K. Jakob, Z. Yun, A Deligne-Simpson problem for irregular G-connections over P¹, preprint (2023) arXiv:2301.10967.
- [24] N. Katz, *Rigid local systems*, Princeton University Press, Princeton, NJ, 1996. viii+223 pp.
- [25] H. Kawakami, A. Nakamura, H. Sakai, Degeneration Scheme of 4-dimensional Painlevé-type Equations in [18].
- [26] G. Kempf, *Instability in invariant theory* (preprint version), preprint (1976), arXiv:1807.02890.
- [27] M. Klimeš, Analytic classification of families of linear differential systems unfolding a resonant irregular singularity, SIGMA 16 (2020), 006, 46 pp.
- [28] V. Kostov, The Deligne-Simpson problem-a survey, J. Algebra, **281** (2004), 83–108.
- [29] M. Kulkarni, N. Livesay, J. Matherne, B. Nguyen, and D. Sage, *The Deligne-Simpson problem for connections on G_m with a maximally ramified singularity*, Adv. Math. 408 (2022), 108596.
- [30] C. Lambert, C. Rousseau, The Stokes phenomenon in the confluence of the hypergeometric equation using Riccati equation, J. Differential Equations 244 (2008), 2641– 2664.
- [31] G. Levelt, Jordan decomposition for a class of singular differential operators, Ark. Mat. 13 (1975) 1–27.
- [32] N. Livesay, D. Sage, B. Nguyen, *Explicit constructions of connections on the projective line with a maximally ramified irregular singularity*, preprint (2023), arXiv:2303.06581.
- [33] K. Mackenzie, General theory of Lie groupoids and Lie algebroids, London Mathematical Society Lecture Note Series, vol. 213, Cambridge University Press, Cambridge, 2005.
- [34] B. Malgrange, Équations différentielles à coefficients polynomiaux, Progr. Math., 96
 Birkhä user Boston, Inc., Boston, MA, 1991. vi+232 pp.
- [35] J. Marsden, A. Weinstein, Reduction of symplectic manifolds with symmetry, Rep. Mathematical Phys. 5 (1974), 121–130.

- [36] D. Mumford, J. Fogarty, F. Kirwan, Geometric invariant theory. Third edition, Ergeb. Math. Grenzgeb. (2), 34 Springer-Verlag, Berlin, 1994. xiv+292 pp.
- [37] T. Oshima, Confluence and versal unfolding of Pfaffian equations, Josai Mathematical Monographs 12 (2020), 117–151.
- [38] T. Oshima, Versal unfolding of irregular singurarities of a linear differential equation on the Riemann sphere, Publ. RIMS Kyoto Uinv. 57 (2021), 893–920.
- [39] J.-P. Ramis, Confluence et résurgence, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 36 (1989), 703–716.
- [40] H. Sakai, Isomonodromic Deformation and 4-dimensional Painlevé-type Equations in [18].
- [41] R. Schäfke, Confluence of several regular singular points into an irregular singular one, J. Dynam. Control Systems 4 (1998), 401–424.
- [42] C. Simpson, Products of matrices, in Differential Geometry, Global Analysis, and Topology (Halifax, 1990), CMS Conf. Proc. 12, Amer. Math. Soc. Providence, (1991), 157–185.
- [43] R. Sjamaar, E. Lerman Stratified symplectic spaces and reduction, Ann. Math. 134 (1991), 375–422.
- [44] T. Springer, Linear algebraic groups. Second edition, Progr. Math., 9 Birkhäuser Boston, Inc., Boston, MA, 1998. xiv+334 pp.
- [45] H. Turrittin, Convergent solutions of ordinary linear homogeneous differential equations in the neighborhood of an irreg- ular singular point, Acta Math. 93 (1955) 27-66.
- [46] D. Yamakawa, Quiver Varieties with Multiplicities, Weyl Groups of Non-Symmetric Kac-Moody Algebras, and Painlevé Equations, SIGMA 6 (2010), 087.
- [47] D. Yamakawa, Fundamental two-forms for isomonodromic deformations, J. Integrable Syst. 4 (2019), no.1, 1–35.
- [48] C. Zhang, Confluence et phénomène de Stokes, J. Math. Sci. Univ. Tokyo 3 (1996), 91–107.