

Hölder regularity of harmonic functions on metric measure spaces

Jin Gao and Meng Yang

Abstract

We introduce the Hölder regularity condition for harmonic functions on metric measure spaces and prove that under mild volume regular condition and upper heat kernel estimate, the Hölder regularity condition, the weak Bakry-Émery non-negative curvature condition, the heat kernel Hölder continuity with or without exponential terms and the heat kernel near-diagonal lower bound are equivalent. As applications, firstly, we prove the validity of the so-called generalized reverse Hölder inequality on the Sierpiński *carpet* cable system, which was left open by Devyver, Russ, Yang (Int. Math. Res. Not. IMRN (2023), no. 18, 15537–15583). Secondly, we prove that two-sided heat kernel estimates *alone* imply gradient estimate for the heat kernel on strongly recurrent fractal-like cable systems, which improves the main results of the aforementioned paper. Thirdly, we obtain Hölder (Lipschitz) estimate for heat kernel on general metric measure spaces, which extends the classical Li-Yau gradient estimate for heat kernel on Riemannian manifolds.

1 Introduction

Let us recall the following classical de Giorgi-Nash-Moser theorem from PDE theory. Let D be a bounded domain in \mathbb{R}^d . Consider the following divergence form elliptic operator

$$Lu = \sum_{i,j=1}^d \frac{\partial}{\partial x_i} \left(a_{ij} \frac{\partial u}{\partial x_j} \right),$$

where a_{ij} , $i, j = 1, \dots, d$, are measurable functions in D satisfying that $a_{ij} = a_{ji}$ for $i, j = 1, \dots, d$ and there exist some positive constants λ, Λ such that

$$\lambda |\xi|^2 \leq \sum_{i,j=1}^d a_{ij}(x) \xi_i \xi_j \leq \Lambda |\xi|^2 \text{ for any } x \in D, \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d.$$

Then any weak solution $u \in W_{loc}^{1,2}(D)$ of $Lu = 0$ in D , or (L -)harmonic function $u \in W_{loc}^{1,2}(D)$, is locally Hölder continuous in D or $u \in C^\alpha(D)$, where $\alpha \in (0, 1)$ is the Hölder exponent. One only needs the *measurability* of the coefficients a_{ij} to obtain the *Hölder regularity* of the weak solution u . To prove this theorem, very powerful iteration techniques were developed, including de Giorgi iteration [24] and Moser iteration [49], and were applied in many areas later, see [18, 36, 21, 32, 13, 29] and references therein for analysis and heat kernel on metric measure spaces. However, iteration techniques give the existence of some Hölder exponent α which depends only on d, λ, Λ , but may not give an explicit value of α .

The main purpose of this paper is to consider Hölder regularity of harmonic functions on metric measure spaces, where certain *explicit* Hölder exponent can be obtained. These metric measure spaces will originate from certain *strongly recurrent* fractals.

Date: August 20, 2024

MSC2010: 28A80, 35K08

Keywords: Hölder regularity, gradient estimate, harmonic function, heat kernel, Sierpiński carpet.

Jin Gao was supported by National Natural Science Foundation of China (12271282). Jin Gao acknowledged Jiaxin Hu and Eryan Hu for reading the preliminary draft. Part of the work was carried out while Meng Yang was participating a Mini-Symposia in *the 9th European Congress of Mathematics (9ECM)* in Seville (Spain) between 15 and 19 July 2024, Meng Yang was very grateful to Fabrice Baudoin and Li Chen for the invitation to the symposia.

Fractals are new examples with very different phenomena from Riemannian manifolds. On a complete non-compact Riemannian manifold, it was independently discovered by Grigor'yan [27] and Saloff-Coste [50, 51] that the following two-sided Gaussian estimate of the heat kernel

$$\frac{C_1}{V(x, \sqrt{t})} \exp\left(-C_2 \frac{d(x, y)^2}{t}\right) \leq p_t(x, y) \leq \frac{C_3}{V(x, \sqrt{t})} \exp\left(-C_4 \frac{d(x, y)^2}{t}\right), \quad \text{HK(2)}$$

is equivalent to the conjunction of the volume doubling condition and the scale-invariant L^2 -Poincaré inequality on balls. However, one important estimate on fractals is the following two-sided *sub-Gaussian* estimate of the heat kernel

$$\frac{C_1}{V(x, t^{1/\beta})} \exp\left(-C_2 \left(\frac{d(x, y)}{t^{1/\beta}}\right)^{\frac{\beta}{\beta-1}}\right) \leq p_t(x, y) \leq \frac{C_3}{V(x, t^{1/\beta})} \exp\left(-C_4 \left(\frac{d(x, y)}{t^{1/\beta}}\right)^{\frac{\beta}{\beta-1}}\right),$$

where β is a new parameter called the walk dimension, which is always strictly greater than 2 on fractals. Moreover, we say that a fractal is strongly recurrent if its Hausdorff dimension α is strictly less than its walk dimension β . For example, on the Sierpiński gasket (see Figure 1), we have $\alpha = \log 3 / \log 2 < \beta = \log 5 / \log 2$, see [14, 41], on the Sierpiński carpet (see Figure 2), we have $\alpha = \log 8 / \log 3 < \beta \approx 2.09697$, see [6, 7, 11, 8, 35, 46].




Figure 1: The Sierpiński gasket




Figure 2: The Sierpiński carpet

Another purpose of this paper is to improve some recent results by Devyver, Russ and the second named author [25], where gradient estimate for heat kernel was considered. Gradient estimate for heat kernel plays an important role in the L^p -boundedness of the Riesz transform for $p > 2$, see [3, 22]. To obtain pointwise upper *sub-Gaussian* estimate for the gradient of the heat kernel on fractal-like cable systems, they introduced the so-called generalized reverse Hölder inequality and verified this condition on the Vicsek and the Sierpiński gasket cable systems, where the proof can be generalized to the cable systems corresponding to a class of p.c.f. self-similar sets. Since the harmonic extension algorithm was intrinsically needed, their proof can not apply on the Sierpiński *carpet* cable system. In this paper, we introduce the Hölder regularity condition, which is stronger than the generalized reverse Hölder inequality, and, a little bit surprisingly, can be ensured by pointwise two-sided heat kernel estimates. Therefore, on the one hand, we prove the validity of the generalized reverse Hölder inequality on the Sierpiński *carpet* cable system, which was left open in [25, Page 15545, Line 2]. On the other hand, to obtain gradient estimate for the heat kernel, we do not need to assume gradient type conditions, such as the generalized reverse Hölder inequality, but *only* pointwise two-sided heat kernel estimates and some mild volume condition are enough, thus we improve main results of [25, Theorem 1.1, Corollary 1.3].

The next purpose of this paper is to generalize the recent result by Baudoin, Chen [15, Theorem 3.3] about Lipschitz estimate for heat kernel on the unbounded Vicsek set. Here we obtain Hölder (Lipschitz) estimate for heat kernel on general unbounded metric

measure spaces, which may behave differently at small scale and large scale. We follow a concise analytical approach, thus avoid to going into the probabilistic framework of fractional diffusions by Barlow [5].

To end the Introduction, let us state briefly our main result, more detailed statement will be given in Section 2. We introduce firstly two scaling functions. Let $\alpha_1, \alpha_2, \beta_1, \beta_2 \in (0, +\infty)$ satisfy

$$2 \leq \beta_i \leq \alpha_i + 1, \alpha_i < \beta_i \text{ for } i = 1, 2.$$

Let $\Phi, \Psi : (0, +\infty) \rightarrow (0, +\infty)$ be given by

$$\begin{aligned} \Phi(r) &= r^{\alpha_1} 1_{(0,1)}(r) + r^{\alpha_2} 1_{[1,+\infty)}(r), \\ \Psi(r) &= r^{\beta_1} 1_{(0,1)}(r) + r^{\beta_2} 1_{[1,+\infty)}(r). \end{aligned}$$

In an unbounded metric measure Dirichlet space, we introduce the Hölder regularity condition $\text{HR}(\Phi, \Psi)$ and prove that under mild volume regular condition $\text{V}(\Phi)$ and upper heat kernel estimate $\text{UHK}(\Psi)$, the following conditions are indeed equivalent.

- The Hölder regularity condition $\text{HR}(\Phi, \Psi)$.
- The weak Bakry-Émery non-negative curvature condition $\text{wBE}(\Phi, \Psi)$.
- The heat kernel Hölder continuity $\text{HHK}(\Phi, \Psi)$.
- The heat kernel Hölder continuity with exponential terms $\text{HHK}_{\text{exp}}(\Phi, \Psi)$.
- The heat kernel near-diagonal lower bound $\text{NLE}(\Psi)$.

Since we do not need to assume the existence of any gradient operator associated with the metric measure Dirichlet space, in particular, a “*carré du champ*”, the above equivalence generalizes the corresponding result by Coulhon, Jiang, Koskela, Sikora [22, Theorem 1.2].

In addition, we also extend the classical result on Riemannian manifolds to metric measure Dirichlet spaces, that is, the conjunction of $\text{UHK}(\Psi)$ and $\text{HHK}_{\text{exp}}(\Phi, \Psi)$ implies $\text{NLE}(\Psi)$ on metric measure Dirichlet spaces, which extends that the conjunction of *upper* and *gradient* estimates of heat kernel implies *lower* estimate of heat kernel on Riemannian manifolds, see Cheeger, Yau [19] and Li, Yau [47].

Throughout the paper, the letters C, C_1, C_2, C_A, C_B will always refer to some positive constants and may change at each occurrence. The sign \asymp means that the ratio of the two sides is bounded from above and below by positive constants. The sign \lesssim (\gtrsim) means that the LHS is bounded by positive constant times the RHS from above (below).

2 Statement of the main results

Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded metric measure Dirichlet (MMD) space, that is, (X, d) is a locally compact separable unbounded metric space, m is a positive Radon measure on X with full support, $(\mathcal{E}, \mathcal{F})$ is a strongly local regular Dirichlet form on $L^2(X; m)$. Throughout this paper, we always assume that all metric balls are relatively compact and $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$ is conservative.

For any $x \in X$, for any $r \in (0, +\infty)$, denote the (metric) ball $B(x, r) = \{y \in X : d(x, y) < r\}$, denote $V(x, r) = m(B(x, r))$. If $B = B(x, r)$, then we denote $\delta B = B(x, \delta r)$ for any $\delta \in (0, +\infty)$. Let $C(X)$ denote the space of all real-valued continuous functions on X and let $C_c(X)$ denote the space of all real-valued continuous functions on X with compact support.

Consider the strongly local regular Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$. Let Δ be the corresponding generator which is a non-negative definite self-adjoint operator. Let Γ be the corresponding energy measure. Denote $\mathcal{E}_1(\cdot, \cdot) = \mathcal{E}(\cdot, \cdot) + (\cdot, \cdot)$, where (\cdot, \cdot) is the inner product in $L^2(X; m)$. We refer to [26] for related results about Dirichlet forms.

We need to introduce some conditions to state our main results.

We say that the volume doubling condition VD holds if there exists $C_{VD} \in (0, +\infty)$ such that

$$V(x, 2r) \leq C_{VD} V(x, r) \text{ for any } x \in X, r \in (0, +\infty). \quad \text{VD}$$

We say that the volume regular condition $V(\Phi)$ holds if there exists $C_{VR} \in (0, +\infty)$ such that

$$\frac{1}{C_{VR}}\Phi(r) \leq V(x, r) \leq C_{VR}\Phi(r) \text{ for any } x \in X, r \in (0, +\infty). \quad V(\Phi)$$

Consider the regular Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$. Let $\{P_t\}$ be the corresponding heat semi-group. Let $\{X_t, t \geq 0, \mathbb{P}_x, x \in X \setminus \mathcal{N}_0\}$ be the corresponding Hunt process, where \mathcal{N}_0 is a properly exceptional set, that is, $m(\mathcal{N}_0) = 0$ and $\mathbb{P}_x(X_t \in \mathcal{N}_0 \text{ for some } t > 0) = 0$ for any $x \in X \setminus \mathcal{N}_0$. For any bounded Borel function f , we have $P_t f(x) = \mathbb{E}_x f(X_t)$ for any $t > 0$, for any $x \in X \setminus \mathcal{N}_0$.

The heat kernel $p_t(x, y)$ associated with the heat semi-group $\{P_t\}$ is a measurable function defined on $(0, +\infty) \times (X \setminus \mathcal{N}_0) \times (X \setminus \mathcal{N}_0)$ satisfying that:

- For any bounded Borel function f , for any $t > 0$, for any $x \in X \setminus \mathcal{N}_0$, we have

$$P_t f(x) = \int_{X \setminus \mathcal{N}_0} p_t(x, y) f(y) m(dy).$$

- For any $t, s > 0$, for any $x, y \in X \setminus \mathcal{N}_0$, we have

$$p_{t+s}(x, y) = \int_{X \setminus \mathcal{N}_0} p_t(x, z) p_s(z, y) m(dz).$$

- For any $t > 0$, for any $x, y \in X \setminus \mathcal{N}_0$, we have $p_t(x, y) = p_t(y, x)$.

See [32] for more details.

We say that the heat kernel upper bound $\text{UHK}(\Psi)$ holds if there exist a properly exceptional set \mathcal{N} and $C_1, C_2 \in (0, +\infty)$ such that for any $t \in (0, +\infty)$, for any $x, y \in X \setminus \mathcal{N}$, we have

$$p_t(x, y) \leq \frac{C_1}{V(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(x, y), t)), \quad \text{UHK}(\Psi)$$

where

$$\Upsilon(R, t) = \sup_{s \in (0, +\infty)} \left(\frac{R}{s} - \frac{t}{\Psi(s)} \right) \asymp \begin{cases} \left(\frac{R}{t^{1/\beta_1}} \right)^{\frac{\beta_1}{\beta_1-1}}, & \text{if } t < R, \\ \left(\frac{R}{t^{1/\beta_2}} \right)^{\frac{\beta_2}{\beta_2-1}}, & \text{if } t \geq R. \end{cases}$$

Then $\text{UHK}(\Psi)$ can also be re-written as follows:

$$p_t(x, y) \leq \begin{cases} \frac{C_1}{V(x, \Psi^{-1}(t))} \exp \left(-C_2 \left(\frac{d(x, y)}{t^{1/\beta_1}} \right)^{\frac{\beta_1}{\beta_1-1}} \right), & \text{if } t < d(x, y), \\ \frac{C_1}{V(x, \Psi^{-1}(t))} \exp \left(-C_2 \left(\frac{d(x, y)}{t^{1/\beta_2}} \right)^{\frac{\beta_2}{\beta_2-1}} \right), & \text{if } t \geq d(x, y). \end{cases}$$

If a lower bound, similar to $\text{UHK}(\Psi)$, with different constants C_i also holds, then we say that $\text{HK}(\Psi)$ holds.

We say that the heat kernel near-diagonal lower bound $\text{NLE}(\Psi)$ holds if there exist a properly exceptional set \mathcal{N} and $C, \varepsilon \in (0, +\infty)$ such that for any $t \in (0, +\infty)$, for any $x, y \in X \setminus \mathcal{N}$ with $d(x, y) < \varepsilon \Psi^{-1}(t)$, we have

$$p_t(x, y) \geq \frac{C}{V(x, \Psi^{-1}(t))}. \quad \text{NLE}(\Psi)$$

If the metric d is furthermore assumed to satisfy the chain condition, then the conjunction of $\text{UHK}(\Psi)$ and $\text{NLE}(\Psi)$ is equivalent to $\text{HK}(\Psi)$, see [32, Section 6].

Let D be an open subset of X . Let

$$\mathcal{F}_D = \text{the } \mathcal{E}_1\text{-closure of } \mathcal{F} \cap C_c(D).$$

Let $f \in L^1_{loc}(D)$. We say that $u \in \mathcal{F}$ is a solution of the Poisson equation $\Delta u = f$ in D if

$$\mathcal{E}(u, \varphi) = \int_D f \varphi dm \text{ for any } \varphi \in \mathcal{F} \cap C_c(D).$$

If $\Delta u = f$ in D with $f \in L^2(D)$, then the above equation also holds for any $\varphi \in \mathcal{F}_D$. We say that $u \in \mathcal{F}$ is harmonic in D if $\Delta u = 0$ in D .

Denote $\gamma_i = \beta_i - \alpha_i \in (0, +\infty)$ for $i = 1, 2$, then

$$\left(\frac{\Psi}{\Phi}\right)(r) = r^{\gamma_1} 1_{(0,1)}(r) + r^{\gamma_2} 1_{[1,+\infty)}(r).$$

We say that the Hölder regularity condition $\text{HR}(\Phi, \Psi)$ holds if there exists $C_H \in (0, +\infty)$ such that for any ball $B = B(x_0, r)$, for any $u \in \mathcal{F}$ which is harmonic in $2B$, for m -a.e. $x, y \in B$ with $x \neq y$, we have

$$\frac{\Phi(d(x, y))}{\Psi(d(x, y))} |u(x) - u(y)| \leq C_H \frac{\Phi(r)}{\Psi(r)} \int_{2B} |u| dm. \quad \text{HR}(\Phi, \Psi)$$

The above condition is some kind of generalization of the so-called generalized reversed Hölder inequality $\text{GRH}(\Phi, \Psi)$ introduced in [25].

We say that the Hölder estimate $\text{HHK}(\Phi, \Psi)$ for the heat kernel holds if there exists $C_{\text{HHK}} \in (0, +\infty)$ such that for any $t \in (0, +\infty)$, for any $x_1, x_2, y_1, y_2 \in X$, we have

$$|p_t(x_1, y_1) - p_t(x_2, y_2)| \leq C_{\text{HHK}} \frac{\left(\frac{\Psi}{\Phi}\right)(d(x_1, x_2)) + \left(\frac{\Psi}{\Phi}\right)(d(y_1, y_2))}{t}. \quad \text{HHK}(\Phi, \Psi)$$

The above condition implies that for any $t \in (0, +\infty)$, for any $x \in X$, $p_t(x, \cdot)$ is γ_1 -Hölder continuous in the small scale and γ_2 -Hölder continuous in the large scale.

We say that the Hölder estimate $\text{HHK}_{\text{exp}}(\Phi, \Psi)$ with exponential terms for the heat kernel holds if there exist $C_1, C_2 \in (0, +\infty)$ such that for any $t \in (0, +\infty)$, for any $x, y_1, y_2 \in X$, we have

$$\begin{aligned} & |p_t(x, y_1) - p_t(x, y_2)| \\ & \leq C_1 \frac{\left(\frac{\Psi}{\Phi}\right)(d(y_1, y_2))}{t} (\exp(-\Upsilon(C_2 d(x, y_1), t)) + \exp(-\Upsilon(C_2 d(x, y_2), t))). \end{aligned} \quad \text{HHK}_{\text{exp}}(\Phi, \Psi)$$

We say that the weak Bakry-Émery non-negative curvature condition $\text{wBE}(\Phi, \Psi)$ holds if there exists $C_{\text{wBE}} \in (0, +\infty)$ such that for any $t \in (0, +\infty)$, for any $f \in L^\infty(X; m)$, for any $x, y \in X$, we have

$$\frac{\Phi(d(x, y))}{\Psi(d(x, y))} |P_t f(x) - P_t f(y)| \leq C_{\text{wBE}} \frac{\Phi(\Psi^{-1}(t))}{\Psi(\Psi^{-1}(t))} \|f\|_{L^\infty(X; m)}. \quad \text{wBE}(\Phi, \Psi)$$

The above condition is some kind of generalization of the weak Bakry-Émery curvature condition introduced in [1, Definition 3.1].

The main result of this paper is as follows.

Theorem 2.1. *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$ and $\text{UHK}(\Psi)$. Then the followings are equivalent.*

- (1) $\text{HR}(\Phi, \Psi)$.
- (2) $\text{wBE}(\Phi, \Psi)$.
- (3) $\text{HHK}(\Phi, \Psi)$.
- (4) $\text{HHK}_{\text{exp}}(\Phi, \Psi)$.
- (5) $\text{NLE}(\Psi)$.

The key novelty of the above result is that the *pointwise* two-sided heat kernel estimates are enough to give some proper *explicit* Hölder regularity of harmonic functions. In previous literatures (for example [36, Section 5.3], [21, Proposition 4.5], [32, Section 5.3], [13, Section 4.2.1]), using Moser iteration technique or de Giorgi iteration technique and oscillation inequality, one can obtain only some Hölder exponent depending *implicitly* on other constants.

Our proof will follow the following diagrams.

$$\text{HR}(\Phi, \Psi) \Rightarrow \text{wBE}(\Phi, \Psi) \Rightarrow \text{HHK}(\Phi, \Psi) \Rightarrow \text{NLE}(\Psi) \Rightarrow \text{HR}(\Phi, \Psi),$$

and

$$\text{HR}(\Phi, \Psi) \Rightarrow \text{HHK}_{\text{exp}}(\Phi, \Psi) \Rightarrow \text{HHK}(\Phi, \Psi).$$

The implication “ $\text{HHK}_{\text{exp}}(\Phi, \Psi) \Rightarrow \text{HHK}(\Phi, \Psi)$ ” is trivial by neglecting exponential terms. The proof of “ $\text{wBE}(\Phi, \Psi) \Rightarrow \text{HHK}(\Phi, \Psi) \Rightarrow \text{NLE}(\Psi)$ ” is easy and standard and will be given for completeness. The proofs of “ $\text{HR}(\Phi, \Psi) \Rightarrow \text{wBE}(\Phi, \Psi)$ ” and “ $\text{HR}(\Phi, \Psi) \Rightarrow \text{HHK}_{\text{exp}}(\Phi, \Psi)$ ” will use a similar technique of gradient estimates as in [25, 22], see also [38, 39]. The proof of “ $\text{NLE}(\Psi) \Rightarrow \text{HR}(\Phi, \Psi)$ ” is new and will use a recent technique of resistance estimates in [40, Theorem 6.27].

Remark 2.2. *In the setting of fractional metric spaces with fractional diffusions introduced in [5, Section 3], when the dimension α of the state space is smaller than the walk dimension β , or $\alpha_1 = \alpha_2 = \alpha < \beta_1 = \beta_2 = \beta$ in our setting, it was proved in [1, Lemma 3.4, Remark 3.6, Theorem 3.7] that*

$$\text{wBE}(\Phi, \Psi) \Rightarrow \text{HHK}(\Phi, \Psi) \Rightarrow \text{NLE}(\Psi) \Rightarrow \text{wBE}(\Phi, \Psi),$$

respectively. To prove “ $\text{NLE}(\Psi) \Rightarrow \text{wBE}(\Phi, \Psi)$ ”, the Hölder regularity result [5, Theorem 3.40] for resolvents obtained within probabilistic framework was intrinsically needed.

Remark 2.3. *In the fractal setting, $\text{HHK}(\Phi, \Psi)$ was also proved on the unbounded Sierpiński gasket [14, Theorem 1.5 (c)] and on the unbounded Sierpiński carpet [8, Theorem 1.1 (e)].*

Remark 2.4. *The above equivalence can be regarded as some “Hölder” version generalization of the “gradient” version result [22, Theorem 1.2]. It was proved that in a non-compact doubling Dirichlet metric measure space (X, d, μ, \mathcal{E}) endowed with a “carré du champ”. Assume the upper Gaussian bound (UE) for the heat kernel and a local L^∞ -Poincaré inequality $(P_{\infty, \text{loc}})$. Then the following conditions are equivalent.*

- *The quantitative reverse L^∞ -Hölder inequality (RH_∞) for gradients of harmonic functions: There exists $C \in (0, +\infty)$ such that for any ball $B = B(x_0, r)$, for any u which is harmonic in $2B$, we have*

$$\|\nabla u\|_{L^\infty(B)} \leq \frac{C}{r} \int_{2B} |u| d\mu. \quad (\text{RH}_\infty)$$

- *The pointwise Li-Yau gradient estimate (GLY_∞) for heat kernel: There exist $C, c \in (0, +\infty)$ such that*

$$|\nabla_x h_t(x, y)| \leq \frac{C}{\sqrt{t}V(y, \sqrt{t})} \exp\left\{-c \frac{d(x, y)^2}{t}\right\} \quad (\text{GLY}_\infty)$$

for any $t \in (0, +\infty)$, for μ -a.e. $x, y \in X$.

- *The L^∞ -boundedness (G_∞) of the gradient of the heat semi-group $|\nabla H_t|$: There exists $C \in (0, +\infty)$ such that for any $t \in (0, +\infty)$, we have*

$$\|\nabla H_t\|_{\infty \rightarrow \infty} \leq \frac{C}{\sqrt{t}},$$

that is, for any $f \in L^\infty(X)$, we have

$$\|\nabla H_t f\|_{L^\infty(X)} \leq \frac{C}{\sqrt{t}} \|f\|_{L^\infty(X)}. \quad (\text{G}_\infty)$$

Our conditions are “Hölder” version of the above “gradient” conditions as in Figure 3.

In their proof to obtain (RH_∞) , they established a reproducing formula for harmonic functions using the finite propagation speed property, which may not be available on general MMD spaces. We leave it as an open question to generalize the reproducing formula technique to prove $\text{HR}(\Phi, \Psi)$ on MMD spaces or the following $\text{GRH}(\Phi, \Psi)$ on cable systems, in particular, to develop the corresponding finite propagation speed theory.

“Hölder” conditions	“gradient” conditions
HR(Φ, Ψ)	(RH $_{\infty}$)
HHK $_{\text{exp}}$ (Φ, Ψ)	(GLY $_{\infty}$)
wBE(Φ, Ψ)	(G $_{\infty}$)

Figure 3: The correspondence between “Hölder” and “gradient” conditions

Remark 2.5. *Let us compare the stability of “gradient” and “Hölder” conditions as follows.*

By a standard technique, it is easy to see that under (UE), any one of the equivalent “gradient” conditions implies the heat kernel near-diagonal lower Gaussian bound (NLE). In particular, under (UE), we have (GLY $_{\infty}$) implies (NLE). However, the converse is usually not true. Indeed, the conjunction of (UE) and (NLE), or equivalently, HK(2), is equivalent to the conjunction of the volume doubling condition and the scale-invariant L^2 -Poincaré inequality on balls, hence is quasi-isometry stable. However, to obtain (GLY $_{\infty}$), some curvature assumptions are usually needed, see [47, 16], hence (GLY $_{\infty}$) is quasi-isometry unstable.

For comparison, by our main result, Theorem 2.1, under UHK(Ψ), all the “Hölder” conditions are equivalent to NLE(Ψ). By the quasi-isometry stability of UHK(Ψ) (see [2]) and the conjunction of UHK(Ψ) and NLE(Ψ) (see [31]), our “Hölder” conditions are quasi-isometry stable. Roughly speaking, the stability originates from the strongly recurrent condition, which requires our MMD spaces to behave like \mathbb{R}^1 in the classical Riemannian setting.

Let (V, E) be an infinite, locally bounded, connected (undirected) graph. Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be the corresponding unbounded cable system with an elementary gradient operator ∇ , see [25, Section 3] for more details. Let $\alpha_1 = 1$, $\beta_1 = 2$, $\alpha_2 = \alpha$, $\beta_2 = \beta$.

We say that the generalized reverse Hölder inequality GRH(Φ, Ψ) holds if there exists $C_H \in (0, +\infty)$ such that for any ball $B = B(x_0, r)$, for any $u \in \mathcal{F}$ which is harmonic in $2B$, we have

$$\|\nabla u\|_{L^\infty(B)} \leq C_H \frac{\Phi(r)}{\Psi(r)} \int_{2B} |u| dm. \quad \text{GRH}(\Phi, \Psi)$$

It is obvious that on an unbounded cable system, HR(Φ, Ψ) implies GRH(Φ, Ψ). Indeed, assume HR(Φ, Ψ), then the LHS of HR(Φ, Ψ) becomes $|u(x) - u(y)|/d(x, y)$ for $d(x, y) \in (0, 1)$. Letting $y \rightarrow x$ and using the Lebesgue density theorem, we obtain the limit $|\nabla u(x)|$. Taking essential supremum with respect to $x \in B$, we have GRH(Φ, Ψ). Roughly speaking, HR(Φ, Ψ) encodes more Hölder continuity information than GRH(Φ, Ψ) about the behavior of harmonic functions at *large* scale.

We give some results as important applications of our main result, Theorem 2.1, on unbounded cable systems. Firstly, due to the fact that two-sided heat kernel estimates have already obtained on a large class of fractals (see [5, 34, 42, 43]), including the Sierpiński carpet (see [8, 9, 35]), we have GRH(Φ, Ψ) holds on a large class of fractal-like cable systems, including the Sierpiński *carpet* cable system, whose validity was left open in [25].

Corollary 2.6. *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded cable system satisfying $V(\Phi)$ and $HK(\Psi)$ with $\alpha < \beta$. Then GRH(Φ, Ψ) holds. In particular, GRH(Φ, Ψ) holds on the Sierpiński carpet cable system with $\alpha = \log 8 / \log 3 < 2 < \beta \approx 2.09697$.*

Remark 2.7. *In [25], the authors verified directly GRH(Φ, Ψ) on the Vicsek cable system and the Sierpiński gasket cable system using certain Hölder regularity¹ of harmonic functions on the corresponding fractals [52], which is also available on some class of p.c.f. self-similar sets [54] without any technical difficulty, but not available on general fractals, especially on the Sierpiński carpet.*

Secondly, we improve the results about gradient estimate for heat kernel as follows.

Corollary 2.8. *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded cable system satisfying $V(\Phi)$ and $HK(\Psi)$ with $\alpha < \beta$.*

¹The Hölder regularity follows indeed from certain extension algorithm for harmonic functions.

(1) Then $\text{HHK}_{\text{exp}}(\Phi, \Psi)$ holds. In particular, we have the gradient estimate $\text{GHK}(\Phi, \Psi)$ for the heat kernel as follows: there exist $C_1, C_2 \in (0, +\infty)$ such that for any $t \in (0, +\infty)$, for m -a.e. $x, y \in X$, we have

$$|\nabla_y p_t(x, y)| \leq \frac{C_1 \Phi(\Psi^{-1}(t))}{tV(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(x, y), t)). \quad \text{GHK}(\Phi, \Psi)$$

(2) There exist $C_3, C_4 \in (0, +\infty)$ such that for any $t \in (0, +\infty)$, for m -a.e. $x, y \in X$, we have

$$|\nabla_y p_t(x, y)| \leq C_3 \frac{\Phi(\Psi^{-1}(t))}{t} p_{C_4 t}(x, y).$$

Remark 2.9. (1) is certain improvement of [25, Theorem 1.1] and (2) is certain improvement of [25, Corollary 1.3], where $\text{GRH}(\Phi, \Psi)$ was needed. Now we can obtain gradient estimate result $\text{GHK}(\Phi, \Psi)$ without assuming gradient type conditions like $\text{GRH}(\Phi, \Psi)$.

As a direct consequence of $\text{HHK}_{\text{exp}}(\Phi, \Psi)$, we also obtain the following Hölder estimate for the heat kernel on unbounded MMD spaces.

Corollary 2.10. Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$ and $\text{HK}(\Psi)$. Then there exist $C, c \in (0, +\infty)$ such that for any $t \in (0, +\infty)$, for any $x, y_1, y_2 \in X$, we have

$$|p_t(x, y_1) - p_t(x, y_2)| \leq C \frac{\left(\frac{\Psi}{\Phi}\right)(d(y_1, y_2))}{\left(\frac{\Psi}{\Phi}\right)(\Psi^{-1}(t))} (p_{ct}(x, y_1) + p_{ct}(x, y_2)).$$

Remark 2.11. If $\beta_1 = \alpha_1 + 1$, then the above Hölder estimate gives a local Lipschitz estimate, which will imply the existence and some estimate of the gradient for the heat kernel. We give some examples as follows.

- Unbounded cable systems satisfying $V(\Phi)$ and $\text{HK}(\Psi)$. We have $\alpha_1 = 1$ and $\beta_1 = 2$ and this gives indeed also Corollary 2.8.
- The unbounded Vicsek set. We have $\alpha_1 = \alpha_2 = \log 5 / \log 3$, $\beta_1 = \beta_2 = \log 15 / \log 3$ and the above Hölder estimate is indeed a Lipschitz estimate, which was also given by Baudoin, Chen [15, Theorem 3.3]. However, a resolvent kernel estimate [5, Theorem 3.40] was intrinsically needed in their proof.
- The blowups of the Vicsek set by another fractal. We have $\alpha_1 = \log 5 / \log 3$ and $\beta_1 = \log 15 / \log 3$. α_2 and β_2 will be the Hausdorff dimension and walk dimension of another fractal, say the Sierpiński carpet or the unit interval, see Figure 7 or Figure 8, respectively. The estimates of the gradient for heat kernel can behave differently at small scale and large scale. We will give a detailed discussion in Section 9 about these two examples.

This paper is organized as follows. In Section 3, we list some characterizations of heat kernel estimates for later use. In Section 4, we give the existence, the uniqueness and Hölder estimate for the solutions of Poisson equation. In Section 5, we prove “ $\text{HR}(\Phi, \Psi) \Rightarrow \text{wBE}(\Phi, \Psi)$ ”. In Section 6, we prove “ $\text{HR}(\Phi, \Psi) \Rightarrow \text{HHK}_{\text{exp}}(\Phi, \Psi)$ ”. In Section 7, we prove “ $\text{wBE}(\Phi, \Psi) \Rightarrow \text{HHK}(\Phi, \Psi) \Rightarrow \text{NLE}(\Psi)$ ”. In Section 8, we prove “ $\text{NLE}(\Psi) \Rightarrow \text{HR}(\Phi, \Psi)$ ”. In Section 9, we give some examples related to the Sierpiński carpet and the Vicsek set including the blowups of a fractal by another different fractal.

3 Heat kernel estimates

In this section, we list some characterizations of heat kernel estimates for later use.

Let D be an open subset of X . Denote by $\lambda_1(D)$ the smallest Dirichlet eigenvalue for D , that is,

$$\lambda_1(D) = \inf \left\{ \frac{\mathcal{E}(u, u)}{\|u\|_2^2} : u \in \mathcal{F}_D \setminus \{0\} \right\}.$$

We say that the relative Faber-Krahn inequality $\text{FK}(\Psi)$ holds if there exist $C_F \in (0, +\infty)$, $\nu \in (0, 1)$ such that for any ball $B = B(x, r)$, for any open subset D of B , we have

$$\lambda_1(D) \geq \frac{C_F}{\Psi(r)} \left(\frac{m(B)}{m(D)} \right)^\nu. \quad \text{FK}(\Psi)$$

We will sometimes use the notation $\text{FK}(\Psi, \nu)$ to emphasize the role of the value of ν .

We say that the Poincaré inequality $\text{PI}(\Psi)$ holds if there exists $C_P \in (0, +\infty)$ such that for any ball $B = B(x, r)$, for any $u \in \mathcal{F}$, we have

$$\int_B |u - u_B|^2 dm \leq C_P \Psi(r) \int_{2B} d\Gamma(u, u), \quad \text{PI}(\Psi)$$

where u_A is the mean value of u on a measurable set A with $m(A) \in (0, +\infty)$, that is,

$$u_A = \int_A u dm = \frac{1}{m(A)} \int_A u dm.$$

Let U, V be two open subsets of X satisfying $U \subseteq \bar{U} \subseteq V$. We say that $\varphi \in \mathcal{F}$ is a cutoff function for $U \subseteq V$ if $0 \leq \varphi \leq 1$ m -a.e., $\varphi = 1$ m -a.e. in an open neighborhood of \bar{U} and $\text{supp}(\varphi) \subseteq V$, where $\text{supp}(f)$ refers to the support of the measure of $|f| dm$ for any given function f .

We say that the cutoff Sobolev inequality $\text{CS}(\Psi)$ holds if there exists $C_S \in (0, +\infty)$ such that for any $x \in X$, for any $R, r \in (0, +\infty)$, there exists a cutoff function $\varphi \in \mathcal{F}$ for $B(x, R) \subseteq B(x, R+r)$ such that for any $f \in \mathcal{F}$, we have

$$\begin{aligned} & \int_{B(x, R+r) \setminus \overline{B(x, R)}} f^2 d\Gamma(\varphi, \varphi) \\ & \leq \frac{1}{8} \int_{B(x, R+r) \setminus \overline{B(x, R)}} \varphi^2 d\Gamma(f, f) + \frac{C_S}{\Psi(r)} \int_{B(x, R+r) \setminus \overline{B(x, R)}} f^2 dm. \end{aligned} \quad \text{CS}(\Psi)$$

We have the characterisations of heat kernel estimates as follows.

Proposition 3.1. ([2, Theorem 1.12]) *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying VD. Then the followings are equivalent.*

- (1) $\text{UHK}(\Psi)$.
- (2) $\text{FK}(\Psi)$ and $\text{CS}(\Psi)$.

Proposition 3.2. ([31, THEOREM 1.2]) *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$. Then the followings are equivalent.*

- (1) $\text{UHK}(\Psi)$ and $\text{NLE}(\Psi)$.
- (2) $\text{PI}(\Psi)$ and $\text{CS}(\Psi)$.

Remark 3.3. *On any complete non-compact Riemannian manifold, $\text{CS}(\Psi)$ with $\beta_1 = \beta_2 = 2$, or equivalently, $\Psi(r) = r^2$ for any $r \in (0, +\infty)$, holds automatically, for example, one can simply take*

$$\varphi = 1 - \left(\frac{d(x, \cdot) - (R + \frac{r}{3})}{\frac{r}{3}} \vee 0 \right) \wedge 1,$$

so that the above equivalences hold without $\text{CS}(\Psi)$ and are classical, see [27, 50, 28]. However, on a general MMD space, $\text{CS}(\Psi)$ does not always hold and is involved in the formulation of the previous equivalences.

4 Poisson equation on unbounded MMD spaces

In this section, we give the existence, the uniqueness and Hölder estimate for the solutions of Poisson equation. We follow a similar technique of gradient estimate for the solutions of Poisson equation as in [25, 22], see also [38, 39]. The main result of this section is Proposition 4.3, which will play a central role in the proofs of “ $\text{HR}(\Phi, \Psi) \Rightarrow \text{wBE}(\Phi, \Psi)$ ” and “ $\text{HR}(\Phi, \Psi) \Rightarrow \text{HHK}_{\text{exp}}(\Phi, \Psi)$ ”.

First, we have the existence, the uniqueness and the regularity for the solutions of Poisson equation as follows.

Lemma 4.1. ([25, Lemma 2.2, Lemma 2.6]) Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $FK(\Psi, \nu)$. Let $q = \frac{2}{1-\nu}$. Then for any $p \in \left[\frac{q}{q-1}, +\infty\right)$, for any ball $B = B(x_0, r)$, for any $f \in L^p(B)$, there exists a unique $u \in \mathcal{F}_B$ such that $\Delta u = f$ in B . There exists $C \in (0, +\infty)$ depending only on ν, C_F such that

$$\int_B |u| dm \leq C \sqrt{\Psi(r)} \left(\frac{1}{m(B)} \mathcal{E}(u, u) \right)^{1/2} \leq C \Psi(r) \left(\int_B |f|^p dm \right)^{1/p}.$$

Second, we have the pointwise estimate for the solutions of Poisson equation as follows.

Lemma 4.2. ([25, Lemma 2.7]) Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $VD, FK(\Psi, \nu)$ and $CS(\Psi)$. Let $q = \frac{2}{1-\nu}$. Then for any $p \in \left[\frac{q}{q-1}, +\infty\right)$, there exists $C \in (0, +\infty)$ such that for any ball $B = B(x_0, r)$, for any $f \in L^\infty(2B)$, if $u \in \mathcal{F}$ satisfies $\Delta u = f$ in $2B$, then for m -a.e. $x \in B$, we have

$$|u(x)| \leq C \left(\int_{2B} |u| dm + F_1(x) \right),$$

where

$$F_1(x) = \sum_{j \leq \lfloor \log_2 r \rfloor} \Psi(2^j) \left(\int_{B(x, 2^j)} |f|^p dm \right)^{1/p}.$$

The proof of the above two results in [25] was given for the case $\beta_1 = 2$. It is easy to see that the same proof holds for more general Ψ , including the case in our paper.

Third, we have Hölder estimate for the solutions of Poisson equation as follows. The proof is similar to [25, Proposition 5.1] and [22, Theorem 3.2] to do gradient estimate. Some technical modifications will be needed.

Proposition 4.3. Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$, $FK(\Psi, \nu)$, $CS(\Psi)$ and $HR(\Phi, \Psi)$. Let $q = \frac{2}{1-\nu}$. Then for any $p \in \left[\frac{q}{q-1}, +\infty\right)$, there exists $C \in (0, +\infty)$ such that for any ball $B = B(x_0, r)$, for any $f \in L^\infty(2B)$, if $u \in \mathcal{F}$ satisfies $\Delta u = f$ in $2B$, then for m -a.e. $x, y \in \frac{1}{16}B$ with $x \neq y$, we have

$$\frac{\Phi(d(x, y))}{\Psi(d(x, y))} |u(x) - u(y)| \leq C \left(\frac{\Phi(r)}{\Psi(r)} \int_{2B} |u| dm + F_2(x) + F_2(y) \right),$$

where

$$F_2(x) = \sum_{j \leq \lfloor \log_2 r \rfloor} \Phi(2^j) \left(\int_{B(x, 2^j)} |f|^p dm \right)^{1/p}.$$

Proof. Let $k_0 = \lfloor \log_2 d(x, y) \rfloor$ and $k_1 = \lfloor \log_2 r \rfloor$, then $2^{k_0} \leq d(x, y) < 2^{k_0+1}$ and $2^{k_1} \leq r < 2^{k_1+1}$. Since $x, y \in \frac{1}{16}B$, we have $d(x, y) < \frac{1}{8}r$. Hence $2^{k_0} \leq d(x, y) < \frac{1}{8}r < \frac{1}{8}2^{k_1+1} = 2^{k_1-2}$, which implies $k_0 + 3 \leq k_1$.

For any $k = k_0 + 3, \dots, k_1$, by Lemma 4.1, there exists a unique $u_k \in \mathcal{F}_{B(x, 2^k)}$ such that $\Delta u_k = f$ in $B(x, 2^k)$ and

$$\int_{B(x, 2^{k-1})} |u_k| dm \lesssim \int_{B(x, 2^k)} |u_k| dm \lesssim \Psi(2^k) \left(\int_{B(x, 2^k)} |f|^p dm \right)^{1/p}.$$

Then

$$\begin{aligned} & |u(x) - u(y)| \\ & \leq |(u - u_{k_1})(x) - (u - u_{k_1})(y)| \\ & \quad + \sum_{k=k_0+4}^{k_1} |(u_k - u_{k-1})(x) - (u_k - u_{k-1})(y)| \end{aligned}$$

²in the sense that if $u_1, u_2 \in \mathcal{F}_B$ satisfy $\Delta u_1 = \Delta u_2 = f$ in B , then $u_1 = u_2$ m -a.e..

$$+ |u_{k_0+3}(x)| + |u_{k_0+3}(y)|.$$

For any $k = k_0 + 3, \dots, k_1$, we have $d(x, y) < 2^{k_0+1} \leq 2^{k-2}$, that is, $y \in B(x, 2^{k-2})$.

Since $\Delta(u - u_{k_1}) = 0$ in $B(x, 2^{k_1})$, by HR(Φ, Ψ), we have

$$\begin{aligned} & \frac{\Phi(d(x, y))}{\Psi(d(x, y))} |(u - u_{k_1})(x) - (u - u_{k_1})(y)| \\ & \lesssim \frac{\Phi(2^{k_1-1})}{\Psi(2^{k_1-1})} \int_{B(x, 2^{k_1})} |u - u_{k_1}| dm \\ & \leq \frac{\Phi(2^{k_1-1})}{\Psi(2^{k_1-1})} \left(\int_{B(x, 2^{k_1})} |u| dm + \int_{B(x, 2^{k_1})} |u_{k_1}| dm \right) \\ & \lesssim \frac{\Phi(2^{k_1-1})}{\Psi(2^{k_1-1})} \left(\int_{2B} |u| dm + \Psi(2^{k_1}) \left(\int_{B(x, 2^{k_1})} |f|^p dm \right)^{1/p} \right) \\ & \lesssim \frac{\Phi(r)}{\Psi(r)} \int_{2B} |u| dm + \Phi(2^{k_1}) \left(\int_{B(x, 2^{k_1})} |f|^p dm \right)^{1/p}. \end{aligned}$$

Similarly, for any $k = k_0 + 4, \dots, k_1$, since $\Delta(u_k - u_{k-1}) = 0$ in $B(x, 2^{k-1})$, by HR(Φ, Ψ), we have

$$\begin{aligned} & \frac{\Phi(d(x, y))}{\Psi(d(x, y))} |(u_k - u_{k-1})(x) - (u_k - u_{k-1})(y)| \\ & \lesssim \frac{\Phi(2^{k-2})}{\Psi(2^{k-2})} \int_{B(x, 2^{k-1})} |u_k - u_{k-1}| dm \\ & \leq \frac{\Phi(2^{k-2})}{\Psi(2^{k-2})} \left(\int_{B(x, 2^{k-1})} |u_k| dm + \int_{B(x, 2^{k-1})} |u_{k-1}| dm \right) \\ & \lesssim \frac{\Phi(2^{k-2})}{\Psi(2^{k-2})} \left(\Psi(2^k) \left(\int_{B(x, 2^k)} |f|^p dm \right)^{1/p} + \Psi(2^{k-1}) \left(\int_{B(x, 2^{k-1})} |f|^p dm \right)^{1/p} \right) \\ & \lesssim \Phi(2^k) \left(\int_{B(x, 2^k)} |f|^p dm \right)^{1/p} + \Phi(2^{k-1}) \left(\int_{B(x, 2^{k-1})} |f|^p dm \right)^{1/p}. \end{aligned}$$

Since $\Delta u_{k_0+3} = f$ in $B(x, 2^{k_0+3})$, by Lemma 4.2, we have

$$\begin{aligned} & \frac{\Phi(d(x, y))}{\Psi(d(x, y))} |u_{k_0+3}(x)| \\ & \lesssim \frac{\Phi(d(x, y))}{\Psi(d(x, y))} \left(\int_{B(x, 2^{k_0+3})} |u_{k_0+3}| dm + \sum_{j \leq k_0+2} \Psi(2^j) \left(\int_{B(x, 2^j)} |f|^p dm \right)^{1/p} \right) \\ & \lesssim \frac{\Phi(d(x, y))}{\Psi(d(x, y))} \left(\Psi(2^{k_0+3}) \left(\int_{B(x, 2^{k_0+3})} |f|^p dm \right)^{1/p} + \sum_{j \leq k_0+2} \Psi(2^j) \left(\int_{B(x, 2^j)} |f|^p dm \right)^{1/p} \right) \\ & \lesssim \frac{\Phi(2^{k_0+3})}{\Psi(2^{k_0+3})} \Psi(2^{k_0+3}) \left(\int_{B(x, 2^{k_0+3})} |f|^p dm \right)^{1/p} \\ & + \sum_{j \leq k_0+2} \frac{\Phi(2^{k_0+2})}{\Psi(2^{k_0+2})} \frac{\Psi(2^j)}{\Phi(2^j)} \Phi(2^j) \left(\int_{B(x, 2^j)} |f|^p dm \right)^{1/p} \\ & \leq \Phi(2^{k_0+3}) \left(\int_{B(x, 2^{k_0+3})} |f|^p dm \right)^{1/p} + \sum_{j \leq k_0+2} \Phi(2^j) \left(\int_{B(x, 2^j)} |f|^p dm \right)^{1/p}, \end{aligned}$$

where in the last inequality, we use the fact that $\gamma_1, \gamma_2 > 0$, which implies that

$$\frac{\Phi(2^{k_0+2})}{\Psi(2^{k_0+2})} \frac{\Psi(2^j)}{\Phi(2^j)} \leq 1 \text{ for any } j \leq k_0 + 2.$$

Similarly, since $y \in B(x, 2^{k_0+1}) \subseteq B(x, 2^{k_0+2})$, by Lemma 4.2, we also have

$$\begin{aligned} & \frac{\Phi(d(x, y))}{\Psi(d(x, y))} |u_{k_0+3}(y)| \\ & \lesssim \frac{\Phi(d(x, y))}{\Psi(d(x, y))} \left(\int_{B(x, 2^{k_0+3})} |u_{k_0+3}| dm + \sum_{j \leq k_0+2} \Psi(2^j) \left(\int_{B(y, 2^j)} |f|^p dm \right)^{1/p} \right) \\ & \lesssim \Phi(2^{k_0+3}) \left(\int_{B(x, 2^{k_0+3})} |f|^p dm \right)^{1/p} + \sum_{j \leq k_0+2} \Phi(2^j) \left(\int_{B(y, 2^j)} |f|^p dm \right)^{1/p}. \end{aligned}$$

In summary, we have

$$\begin{aligned} & \frac{\Phi(d(x, y))}{\Psi(d(x, y))} |u(x) - u(y)| \\ & \lesssim \frac{\Phi(r)}{\Psi(r)} \int_{2B} |u| dm + \Phi(2^{k_1}) \left(\int_{B(x, 2^{k_1})} |f|^p dm \right)^{1/p} \\ & + \sum_{k=k_0+4}^{k_1} \left(\Phi(2^k) \left(\int_{B(x, 2^k)} |f|^p dm \right)^{1/p} + \Phi(2^{k-1}) \left(\int_{B(x, 2^{k-1})} |f|^p dm \right)^{1/p} \right) \\ & + \Phi(2^{k_0+3}) \left(\int_{B(x, 2^{k_0+3})} |f|^p dm \right)^{1/p} + \sum_{j \leq k_0+2} \Phi(2^j) \left(\int_{B(x, 2^j)} |f|^p dm \right)^{1/p} \\ & + \Phi(2^{k_0+3}) \left(\int_{B(x, 2^{k_0+3})} |f|^p dm \right)^{1/p} + \sum_{j \leq k_0+2} \Phi(2^j) \left(\int_{B(y, 2^j)} |f|^p dm \right)^{1/p} \\ & \lesssim \frac{\Phi(r)}{\Psi(r)} \int_{2B} |u| dm + \sum_{k=k_0+3}^{k_1} \Phi(2^k) \left(\int_{B(x, 2^k)} |f|^p dm \right)^{1/p} \\ & + \sum_{k \leq k_0+3} \Phi(2^k) \left(\int_{B(x, 2^k)} |f|^p dm \right)^{1/p} + \sum_{k \leq k_0+2} \Phi(2^k) \left(\int_{B(y, 2^k)} |f|^p dm \right)^{1/p} \\ & \lesssim \frac{\Phi(r)}{\Psi(r)} \int_{2B} |u| dm + \sum_{k \leq k_1} \Phi(2^k) \left(\int_{B(x, 2^k)} |f|^p dm \right)^{1/p} + \sum_{k \leq k_1} \Phi(2^k) \left(\int_{B(y, 2^k)} |f|^p dm \right)^{1/p}. \end{aligned}$$

□

5 Proof of “HR(Φ, Ψ) \Rightarrow wBE(Φ, Ψ)”

The idea is to consider the heat equation “ $\Delta(P_t f) = \frac{\partial}{\partial t}(P_t f)$ ” for fixed t as a Poisson equation.

Proof of “HR(Φ, Ψ) \Rightarrow wBE(Φ, Ψ)”. For any $t \in (0, +\infty)$, for any $f \in L^\infty(X; m)$, we have $P_t f$ is a solution of the Poisson equation $\Delta(P_t f) = \frac{\partial}{\partial t}(P_t f)$. For any $x, y \in X$ with $x \neq y$. If $d(x, y) < \Psi^{-1}(t)$, then taking $r = 16\Psi^{-1}(t) > 16d(x, y)$, by Proposition 4.3, we have

$$\begin{aligned} & \frac{\Phi(d(x, y))}{\Psi(d(x, y))} |P_t f(x) - P_t f(y)| \\ & \lesssim \frac{\Phi(16\Psi^{-1}(t))}{\Psi(16\Psi^{-1}(t))} \int_{B(x, 32\Psi^{-1}(t))} |P_t f| dm \\ & + \sum_{k \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^k) \left(\int_{B(x, 2^k)} \left| \frac{\partial}{\partial t}(P_t f) \right|^p dm \right)^{1/p} \end{aligned}$$

$$+ \sum_{k \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^k) \left(\int_{B(y, 2^k)} \left| \frac{\partial}{\partial t} (P_t f)^p dm \right| \right)^{1/p}.$$

For the first term, we have

$$\begin{aligned} & \frac{\Phi(16\Psi^{-1}(t))}{\Psi(16\Psi^{-1}(t))} \int_{B(x, 32\Psi^{-1}(t))} |P_t f| dm \\ & \lesssim \frac{\Phi(\Psi^{-1}(t))}{\Psi(\Psi^{-1}(t))} \int_{B(x, 32\Psi^{-1}(t))} \|P_t f\|_{L^\infty(X; m)} dm \\ & \leq \frac{\Phi(\Psi^{-1}(t))}{\Psi(\Psi^{-1}(t))} \|f\|_{L^\infty(X; m)}. \end{aligned}$$

For the second term, by UHK(Ψ) and [23, THEOREM 4], we have the following estimate of the time derivative of the heat kernel

$$\left| \frac{\partial}{\partial t} p_t(z, w) \right| \leq \frac{C_1}{tV(z, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(z, w), t)),$$

hence

$$\begin{aligned} & \left| \frac{\partial}{\partial t} (P_t f)(z) \right| \\ & \leq \int_X \left| \frac{\partial}{\partial t} p_t(z, w) \right| \cdot |f(w)| m(dw) \\ & \leq \|f\|_{L^\infty(X; m)} \int_X \frac{C_1}{tV(z, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(z, w), t)) m(dw). \end{aligned}$$

By writing

$$\int_X = \int_{B(z, \Psi^{-1}(t))} + \sum_{n=0}^{\infty} \int_{B(z, 2^{n+1}\Psi^{-1}(t)) \setminus B(z, 2^n\Psi^{-1}(t))},$$

and using $V(\Phi)$, we have the integral

$$\int_X \frac{1}{V(z, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(z, w), t)) m(dw)$$

is bounded by some constant depending only on C_2 and Φ , see also [3, Page 944, Line -5], [20, Equation (2.5)], hence

$$\left| \frac{\partial}{\partial t} (P_t f)(z) \right| \lesssim \frac{\|f\|_{L^\infty(X; m)}}{t}.$$

Hence

$$\begin{aligned} & \sum_{k \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^k) \left(\int_{B(x, 2^k)} \left| \frac{\partial}{\partial t} (P_t f)^p dm \right| \right)^{1/p} \\ & \lesssim \sum_{k \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^k) \left(\int_{B(x, 2^k)} \left(\frac{\|f\|_{L^\infty(X; m)}}{t} \right)^p dm \right)^{1/p} \\ & = \frac{\|f\|_{L^\infty(X; m)}}{t} \sum_{k \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \frac{\Phi(2^k)}{\Phi(16\Psi^{-1}(t))} \frac{\Phi(16\Psi^{-1}(t))}{\Phi(\Psi^{-1}(t))} \Phi(\Psi^{-1}(t)) \\ & \lesssim \frac{\|f\|_{L^\infty(X; m)}}{t} \Phi(\Psi^{-1}(t)) \\ & = \frac{\Phi(\Psi^{-1}(t))}{\Psi(\Psi^{-1}(t))} \|f\|_{L^\infty(X; m)}. \end{aligned}$$

For the third term, similarly, we have

$$\sum_{k \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^k) \left(\int_{B(y, 2^k)} \left| \frac{\partial}{\partial t} (P_t f)^p dm \right| \right)^{1/p} \lesssim \frac{\Phi(\Psi^{-1}(t))}{\Psi(\Psi^{-1}(t))} \|f\|_{L^\infty(X; m)}.$$

In summary, we have

$$\frac{\Phi(d(x, y))}{\Psi(d(x, y))} |P_t f(x) - P_t f(y)| \lesssim \frac{\Phi(\Psi^{-1}(t))}{\Psi(\Psi^{-1}(t))} \|f\|_{L^\infty(X; m)}.$$

If $d(x, y) \geq \Psi^{-1}(t)$, then obviously

$$\begin{aligned} & \frac{\Phi(d(x, y))}{\Psi(d(x, y))} |P_t f(x) - P_t f(y)| \\ & \leq 2 \frac{\Phi(d(x, y))}{\Psi(d(x, y))} \|P_t f\|_{L^\infty(X; m)} \leq 2 \frac{\Phi(d(x, y))}{\Psi(d(x, y))} \|f\|_{L^\infty(X; m)} \\ & = 2 \frac{\frac{\Psi(\Psi^{-1}(t))}{\Phi(\Psi^{-1}(t))} \Phi(\Psi^{-1}(t))}{\frac{\Psi(d(x, y))}{\Phi(d(x, y))} \Psi(\Psi^{-1}(t))} \|f\|_{L^\infty(X; m)} \leq 2 \frac{\Phi(\Psi^{-1}(t))}{\Psi(\Psi^{-1}(t))} \|f\|_{L^\infty(X; m)}. \end{aligned}$$

□

6 Proof of “HR(Φ, Ψ) \Rightarrow HHK_{exp}(Φ, Ψ)”

The idea is to consider the heat equation “ $\Delta p_t(x, \cdot) = \frac{\partial}{\partial t} p_t(x, \cdot)$ ” for fixed t, x as a Poisson equation. Comparing with the proof of “HR(Φ, Ψ) \Rightarrow wBE(Φ, Ψ)” in Section 5, we need to pay special attention to exponential terms.

Proof of “HR(Φ, Ψ) \Rightarrow HHK_{exp}(Φ, Ψ)”. By UHK(Ψ) and [23, THEOREM 4], we have the following estimates of the heat kernel

$$p_t(x, y) \leq \frac{C_1}{V(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(x, y), t)),$$

and its time derivative

$$\left| \frac{\partial}{\partial t} p_t(z, w) \right| \leq \frac{C_1}{tV(z, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(z, w), t)).$$

Take any $t \in (0, +\infty)$, $x, y_1, y_2 \in X$. We now distinguish the following four cases.

- (i) $d(y_1, y_2) \geq \Psi^{-1}(t)$.
- (ii) $d(y_1, y_2) < \Psi^{-1}(t)$, $d(x, y_1) < 128\Psi^{-1}(t)$ and $d(x, y_2) < 128\Psi^{-1}(t)$.
- (iii) $d(y_1, y_2) < \Psi^{-1}(t)$ and $d(x, y_1) \geq 128\Psi^{-1}(t)$.
- (iv) $d(y_1, y_2) < \Psi^{-1}(t)$ and $d(x, y_2) \geq 128\Psi^{-1}(t)$.

For the case (i), we obtain the result directly from UHK(Ψ) as follows.

$$\begin{aligned} & |p_t(x, y_1) - p_t(x, y_2)| \leq p_t(x, y_1) + p_t(x, y_2) \\ & \leq \frac{C_1}{V(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(x, y_1), t)) + \frac{C_1}{V(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(x, y_2), t)) \\ & \asymp \frac{1}{\Phi(\Psi^{-1}(t))} (\exp(-\Upsilon(C_2 d(x, y_1), t)) + \exp(-\Upsilon(C_2 d(x, y_2), t))) \\ & = \frac{\Psi(\Psi^{-1}(t))}{\Phi(\Psi^{-1}(t))} \frac{1}{t} (\exp(-\Upsilon(C_2 d(x, y_1), t)) + \exp(-\Upsilon(C_2 d(x, y_2), t))) \\ & \leq \frac{\left(\frac{\Psi}{\Phi}\right)(d(y_1, y_2))}{t} (\exp(-\Upsilon(C_2 d(x, y_1), t)) + \exp(-\Upsilon(C_2 d(x, y_2), t))). \end{aligned}$$

For the remaining three cases, by Proposition 4.3, letting $r = 16\Psi^{-1}(t) > 16d(y_1, y_2)$, we have

$$\begin{aligned} & \frac{\Phi(d(y_1, y_2))}{\Psi(d(y_1, y_2))} |p_t(x, y_1) - p_t(x, y_2)| \\ & \lesssim \frac{\Phi(16\Psi^{-1}(t))}{\Psi(16\Psi^{-1}(t))} \int_{B(y_1, 32\Psi^{-1}(t))} p_t(x, z) m(dz) \end{aligned}$$

$$\begin{aligned}
& + \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \left(\int_{B(y_1, 2^j)} \left| \frac{\partial}{\partial t} p_t(x, z) \right|^p m(dz) \right)^{1/p} \\
& + \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \left(\int_{B(y_2, 2^j)} \left| \frac{\partial}{\partial t} p_t(x, z) \right|^p m(dz) \right)^{1/p}. \tag{6.1}
\end{aligned}$$

For the case (ii), we have $d(x, y_1) < 128\Psi^{-1}(t)$ and $d(x, y_2) < 128\Psi^{-1}(t)$. By Equation (6.1), we have

$$\begin{aligned}
& \frac{\Phi(d(y_1, y_2))}{\Psi(d(y_1, y_2))} |p_t(x, y_1) - p_t(x, y_2)| \\
& \lesssim \frac{\Phi(16\Psi^{-1}(t))}{\Psi(16\Psi^{-1}(t))} \int_{B(y_1, 32\Psi^{-1}(t))} \frac{C_1}{V(x, \Psi^{-1}(t))} m(dz) \\
& + \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \left(\int_{B(y_1, 2^j)} \left(\frac{C_1}{tV(x, \Psi^{-1}(t))} \right)^p m(dz) \right)^{1/p} \\
& + \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \left(\int_{B(y_2, 2^j)} \left(\frac{C_1}{tV(x, \Psi^{-1}(t))} \right)^p m(dz) \right)^{1/p} \\
& = \frac{\Phi(16\Psi^{-1}(t))}{\Psi(16\Psi^{-1}(t))} \frac{C_1}{V(x, \Psi^{-1}(t))} + 2 \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \frac{C_1}{tV(x, \Psi^{-1}(t))} \\
& \asymp \frac{\Phi(\Psi^{-1}(t))}{\Psi(\Psi^{-1}(t))} \frac{1}{\Phi(\Psi^{-1}(t))} + \frac{1}{t\Phi(\Psi^{-1}(t))} \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \\
& = \frac{1}{t} + \frac{1}{t\Phi(\Psi^{-1}(t))} \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j).
\end{aligned}$$

For any $r \in (0, +\infty)$. If $r < 1$, then

$$\sum_{j \leq \lfloor \log_2 r \rfloor} \Phi(2^j) = \sum_{j \leq \lfloor \log_2 r \rfloor} (2^j)^{\alpha_1} \asymp 2^{\lfloor \log_2 r \rfloor \alpha_1} \asymp r^{\alpha_1} = \Phi(r).$$

If $r \geq 1$, then

$$\sum_{j \leq \lfloor \log_2 r \rfloor} \Phi(2^j) = \sum_{j \leq -1} (2^j)^{\alpha_1} + \sum_{j=0}^{\lfloor \log_2 r \rfloor} (2^j)^{\alpha_2} \asymp 1 + 2^{\lfloor \log_2 r \rfloor \alpha_2} \asymp 1 + r^{\alpha_2} \asymp r^{\alpha_2} = \Phi(r).$$

Hence

$$\sum_{j \leq \lfloor \log_2 r \rfloor} \Phi(2^j) \asymp \Phi(r) \text{ for any } r \in (0, +\infty).$$

Then

$$\begin{aligned}
& \frac{\Phi(d(y_1, y_2))}{\Psi(d(y_1, y_2))} |p_t(x, y_1) - p_t(x, y_2)| \lesssim \frac{1}{t} + \frac{1}{t\Phi(\Psi^{-1}(t))} \Phi(16\Psi^{-1}(t)) \\
& \asymp \frac{1}{t} + \frac{1}{t\Phi(\Psi^{-1}(t))} \Phi(\Psi^{-1}(t)) = \frac{2}{t}.
\end{aligned}$$

By assumption, for $i = 1, 2$, we have

$$\Upsilon(C_2 d(x, y_i), t) = \sup_{s \in (0, +\infty)} \left(\frac{C_2 d(x, y_i)}{\Psi^{-1}(s)} - \frac{t}{s} \right) \leq \sup_{s \in (0, +\infty)} \left(\frac{128C_2 \Psi^{-1}(t)}{\Psi^{-1}(s)} - \frac{t}{s} \right).$$

By the following elementary result, Lemma 6.1, there exists $C_3 \in (0, +\infty)$ depending only on C_2, β_1, β_2 such that

$$\sup_{t, s \in (0, +\infty)} \left(\frac{128C_2 \Psi^{-1}(t)}{\Psi^{-1}(s)} - \frac{t}{s} \right) \leq C_3.$$

Hence

$$\begin{aligned} & \frac{\Phi(d(y_1, y_2))}{\Psi(d(y_1, y_2))} |p_t(x, y_1) - p_t(x, y_2)| \lesssim \frac{e^{C_3}}{t} (2e^{-C_3}) \\ & \leq \frac{e^{C_3}}{t} (\exp(-\Upsilon(C_2 d(x, y_1), t)) + \exp(-\Upsilon(C_2 d(x, y_2), t))). \end{aligned}$$

For the case (iii), we have $d(x, y_1) \geq 128\Psi^{-1}(t)$. Then in Equation (6.1), for any $z \in B(y_1, 32\Psi^{-1}(t))$, we have

$$\begin{aligned} d(x, z) & \geq d(x, y_1) - d(y_1, z) \geq d(x, y_1) - 32\Psi^{-1}(t) \\ & \geq d(x, y_1) - \frac{1}{4}d(x, y_1) = \frac{3}{4}d(x, y_1) \geq \frac{1}{2}d(x, y_1). \end{aligned}$$

For any $j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor$, for any $z \in B(y_1, 2^j)$, we have

$$\begin{aligned} d(x, z) & \geq d(x, y_1) - d(y_1, z) \geq d(x, y_1) - 16\Psi^{-1}(t) \\ & \geq d(x, y_1) - \frac{1}{8}d(x, y_1) = \frac{7}{8}d(x, y_1) \geq \frac{1}{2}d(x, y_1), \end{aligned}$$

for any $z \in B(y_2, 2^j)$, we have

$$\begin{aligned} d(x, z) & \geq d(x, y_1) - d(y_1, y_2) - d(y_2, z) \geq d(x, y_1) - \Psi^{-1}(t) - 16\Psi^{-1}(t) \\ & \geq d(x, y_1) - \frac{1}{4}d(x, y_1) = \frac{3}{4}d(x, y_1) \geq \frac{1}{2}d(x, y_1). \end{aligned}$$

Hence Equation (6.1) gives

$$\begin{aligned} & \frac{\Phi(d(y_1, y_2))}{\Psi(d(y_1, y_2))} |p_t(x, y_1) - p_t(x, y_2)| \\ & \lesssim \frac{\Phi(16\Psi^{-1}(t))}{\Psi(16\Psi^{-1}(t))} \int_{B(y_1, 32\Psi^{-1}(t))} \frac{C_1}{V(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(x, z), t)) m(dz) \\ & + \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \left(\int_{B(y_1, 2^j)} \left(\frac{C_1}{tV(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(x, z), t)) \right)^p m(dz) \right)^{1/p} \\ & + \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \left(\int_{B(y_2, 2^j)} \left(\frac{C_1}{tV(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(x, z), t)) \right)^p m(dz) \right)^{1/p} \\ & \leq \frac{\Phi(16\Psi^{-1}(t))}{\Psi(16\Psi^{-1}(t))} \int_{B(y_1, 32\Psi^{-1}(t))} \frac{C_1}{V(x, \Psi^{-1}(t))} \exp\left(-\Upsilon\left(\frac{C_2}{2}d(x, y_1), t\right)\right) m(dz) \\ & + \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \left(\int_{B(y_1, 2^j)} \left(\frac{C_1}{tV(x, \Psi^{-1}(t))} \exp\left(-\Upsilon\left(\frac{C_2}{2}d(x, y_1), t\right)\right) \right)^p m(dz) \right)^{1/p} \\ & + \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \left(\int_{B(y_2, 2^j)} \left(\frac{C_1}{tV(x, \Psi^{-1}(t))} \exp\left(-\Upsilon\left(\frac{C_2}{2}d(x, y_1), t\right)\right) \right)^p m(dz) \right)^{1/p} \\ & = \frac{\Phi(16\Psi^{-1}(t))}{\Psi(16\Psi^{-1}(t))} \frac{C_1}{V(x, \Psi^{-1}(t))} \exp\left(-\Upsilon\left(\frac{C_2}{2}d(x, y_1), t\right)\right) \\ & + 2 \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \frac{C_1}{tV(x, \Psi^{-1}(t))} \exp\left(-\Upsilon\left(\frac{C_2}{2}d(x, y_1), t\right)\right) \\ & \asymp \frac{\Phi(\Psi^{-1}(t))}{\Psi(\Psi^{-1}(t))} \frac{1}{\Phi(\Psi^{-1}(t))} \exp\left(-\Upsilon\left(\frac{C_2}{2}d(x, y_1), t\right)\right) \\ & + \frac{1}{t\Phi(\Psi^{-1}(t))} \exp\left(-\Upsilon\left(\frac{C_2}{2}d(x, y_1), t\right)\right) \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \\ & = \frac{1}{t} \left(1 + \frac{1}{\Phi(\Psi^{-1}(t))} \sum_{j \leq \lfloor \log_2 16\Psi^{-1}(t) \rfloor} \Phi(2^j) \right) \exp\left(-\Upsilon\left(\frac{C_2}{2}d(x, y_1), t\right)\right) \end{aligned}$$

$$\begin{aligned}
&\asymp \frac{1}{t} \left(1 + \frac{1}{\Phi(\Psi^{-1}(t))} \Phi(16\Psi^{-1}(t)) \right) \exp \left(-\Upsilon \left(\frac{C_2}{2} d(x, y_1), t \right) \right) \\
&\asymp \frac{1}{t} \exp \left(-\Upsilon \left(\frac{C_2}{2} d(x, y_1), t \right) \right).
\end{aligned}$$

For the case (iv), similar to the case (iii), we also have

$$\frac{\Phi(d(y_1, y_2))}{\Psi(d(y_1, y_2))} |p_t(x, y_1) - p_t(x, y_2)| \lesssim \frac{1}{t} \exp \left(-\Upsilon \left(\frac{C_2}{2} d(x, y_2), t \right) \right).$$

In summary, for any $t \in (0, +\infty)$, $x, y_1, y_2 \in X$, we have

$$\begin{aligned}
&|p_t(x, y_1) - p_t(x, y_2)| \\
&\lesssim \frac{1}{t} \frac{\Psi(d(y_1, y_2))}{\Phi(d(y_1, y_2))} \left(\exp \left(-\Upsilon \left(\frac{C_2}{2} d(x, y_1), t \right) \right) + \exp \left(-\Upsilon \left(\frac{C_2}{2} d(x, y_2), t \right) \right) \right).
\end{aligned}$$

□

Lemma 6.1. ([25, Lemma 5.2]) *Let $A \in (0, +\infty)$. Then there exists $C \in (0, +\infty)$ depending only on A, β_1, β_2 such that*

$$\sup_{t, s \in (0, +\infty)} \left(A \frac{\Psi^{-1}(t)}{\Psi^{-1}(s)} - \frac{t}{s} \right) \leq C.$$

Proof. Let $f : (0, +\infty) \times (0, +\infty) \rightarrow \mathbb{R}$ be given by

$$f(t, s) = A \frac{\Psi^{-1}(t)}{\Psi^{-1}(s)} - \frac{t}{s}.$$

By considering separately the cases $t, s \in (0, 1)$, $t, s \in [1, +\infty)$, $0 < t < 1 \leq s$ and $0 < s < 1 \leq t$, we easily obtain that

$$f(t, s) \leq A \max \left\{ \left(\frac{t}{s} \right)^{1/\beta_1}, \left(\frac{t}{s} \right)^{1/\beta_2} \right\} - \frac{t}{s}.$$

Since the function $(0, +\infty) \rightarrow \mathbb{R}$, $x \mapsto A \max\{x^{1/\beta_1}, x^{1/\beta_2}\} - x$ is bounded from above by some positive constant C depending only on A, β_1, β_2 , we have

$$\sup_{t, s \in (0, +\infty)} f(t, s) \leq C.$$

□

7 Proof of “wBE(Φ, Ψ) \Rightarrow HHK(Φ, Ψ) \Rightarrow NLE(Ψ)”

The proof of “wBE(Φ, Ψ) \Rightarrow HHK(Φ, Ψ)” is direct by using the semi-group property, see also [1, Lemma 3.4].

Proof of “wBE(Φ, Ψ) \Rightarrow HHK(Φ, Ψ)”. For any $t \in (0, +\infty)$, for any $x, y_1, y_2 \in X$, we have

$$\begin{aligned}
p_t(x, y_1) &= p_t(y_1, x) = \int_X p_{t/2}(y_1, z) p_{t/2}(z, x) m(dz) \\
&= \int_X p_{t/2}(y_1, z) p_{t/2}(x, z) m(dz) = P_{t/2}(p_{t/2}(x, \cdot))(y_1),
\end{aligned}$$

and $p_t(x, y_2) = P_{t/2}(p_{t/2}(x, \cdot))(y_2)$. By wBE(Φ, Ψ) and UHK(Ψ), we have

$$\begin{aligned}
&\frac{\Phi(d(y_1, y_2))}{\Psi(d(y_1, y_2))} |p_t(x, y_1) - p_t(x, y_2)| \\
&= \frac{\Phi(d(y_1, y_2))}{\Psi(d(y_1, y_2))} |P_{t/2}(p_{t/2}(x, \cdot))(y_1) - P_{t/2}(p_{t/2}(x, \cdot))(y_2)|
\end{aligned}$$

$$\begin{aligned}
&\leq C_{wBE} \frac{\Phi(\Psi^{-1}(\frac{t}{2}))}{\Psi(\Psi^{-1}(\frac{t}{2}))} \|p_{t/2}(x, \cdot)\|_{L^\infty(X; m)} \\
&\leq C_{wBE} \frac{\Phi(\Psi^{-1}(\frac{t}{2}))}{\frac{t}{2}} \frac{C_1}{V(x, \Psi^{-1}(t/2))} \\
&\leq 2C_{wBE} \frac{\Phi(\Psi^{-1}(\frac{t}{2}))}{t} \frac{C_1 C_{VR}}{\Phi(\Psi^{-1}(t/2))} \\
&= (2C_1 C_{VR} C_{wBE}) \frac{1}{t},
\end{aligned}$$

that is,

$$|p_t(x, y_1) - p_t(x, y_2)| \lesssim \frac{1}{t} \frac{\Psi(d(y_1, y_2))}{\Phi(d(y_1, y_2))}.$$

Hence for any $x_1, x_2 \in X$, we have

$$\begin{aligned}
&|p_t(x_1, y_1) - p_t(x_2, y_2)| \\
&\leq |p_t(x_1, y_1) - p_t(x_1, y_2)| + |p_t(x_1, y_2) - p_t(x_2, y_2)| \\
&\lesssim \frac{\left(\frac{\Psi}{\Phi}\right)(d(y_1, y_2)) + \left(\frac{\Psi}{\Phi}\right)(d(x_1, x_2))}{t}.
\end{aligned}$$

□

The proof of “ $\text{HHK}(\Phi, \Psi) \Rightarrow \text{NLE}(\Psi)$ ” is classical and standard, see also [21, Theorem 3.1], [32, Section 5].

Proof of “ $\text{HHK}(\Phi, \Psi) \Rightarrow \text{NLE}(\Psi)$ ”. First, we give a lower bound of $p_t(x, x)$. For any $N \geq 0$, by $\text{UHK}(\Psi)$, we have

$$\begin{aligned}
&\int_{X \setminus B(x, 2^N \Psi^{-1}(t))} p_t(x, y) m(dy) \\
&= \sum_{n=N}^{\infty} \int_{B(x, 2^{n+1} \Psi^{-1}(t)) \setminus B(x, 2^n \Psi^{-1}(t))} p_t(x, y) m(dy) \\
&\leq \sum_{n=N}^{\infty} \int_{B(x, 2^{n+1} \Psi^{-1}(t)) \setminus B(x, 2^n \Psi^{-1}(t))} \frac{C_1}{V(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 d(x, y), t)) m(dy) \\
&\leq \sum_{n=N}^{\infty} C_1 \frac{V(x, 2^{n+1} \Psi^{-1}(t))}{V(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_2 2^n \Psi^{-1}(t), t)).
\end{aligned}$$

By $V(\Phi)$, we have

$$\frac{V(x, 2^{n+1} \Psi^{-1}(t))}{V(x, \Psi^{-1}(t))} \leq C_{VR}^2 (2^{n+1})^{\alpha_1 \vee \alpha_2}.$$

Note that

$$\begin{aligned}
\Upsilon(C_2 2^n \Psi^{-1}(t), t) &= \sup_{s \in (0, +\infty)} \left(\frac{C_2 2^n \Psi^{-1}(t)}{s} - \frac{t}{\Psi(s)} \right) \\
&= \sup_{s \in (0, +\infty)} \left(\frac{C_2 2^n \Psi^{-1}(t)}{\Psi^{-1}(s)} - \frac{t}{s} \right) \geq C_2 2^n - 1.
\end{aligned}$$

Hence

$$\int_{X \setminus B(x, 2^N \Psi^{-1}(t))} p_t(x, y) m(dy) \leq C_1 C_{VR}^2 \sum_{n=N}^{\infty} (2^{n+1})^{\alpha_1 \vee \alpha_2} \exp(1 - C_2 2^n).$$

Since

$$\sum_{n=0}^{\infty} (2^{n+1})^{\alpha_1 \vee \alpha_2} \exp(1 - C_2 2^n) < +\infty,$$

there exists $N \geq 0$ depending only on $C_1, C_2, C_{VR}, \alpha_1, \alpha_2$ sufficiently large such that

$$\int_{X \setminus B(x, 2^N \Psi^{-1}(t))} p_t(x, y) m(dy) \leq C_1 C_{VR}^2 \sum_{n=N}^{\infty} (2^{n+1})^{\alpha_1 \vee \alpha_2} \exp(1 - C_2 2^n) \leq \frac{1}{2}.$$

Since $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$ is conservative, that is, $\int_X p_t(x, y) m(dy) = 1$, we have

$$\int_{B(x, 2^N \Psi^{-1}(t))} p_t(x, y) m(dy) = 1 - \int_{X \setminus B(x, 2^N \Psi^{-1}(t))} p_t(x, y) m(dy) \geq \frac{1}{2}.$$

On the other hand

$$\begin{aligned} & \int_{B(x, 2^N \Psi^{-1}(t))} p_t(x, y) m(dy) \\ & \leq \left(\int_{B(x, 2^N \Psi^{-1}(t))} p_t(x, y)^2 m(dy) \right)^{1/2} V(x, 2^N \Psi^{-1}(t))^{1/2} \\ & \leq \left(\int_X p_t(x, y)^2 m(dy) \right)^{1/2} V(x, 2^N \Psi^{-1}(t))^{1/2} \\ & = p_{2t}(x, x)^{1/2} V(x, 2^N \Psi^{-1}(t))^{1/2}, \end{aligned}$$

where we use the Cauchy-Schwarz inequality in the first inequality and the semi-group property in the last equality, hence

$$p_{2t}(x, x) \geq \frac{1}{4V(x, 2^N \Psi^{-1}(t))},$$

hence

$$p_t(x, x) \geq \frac{1}{4V(x, 2^N \Psi^{-1}(\frac{t}{2}))} \geq \frac{\tilde{C}_1}{\Phi(\Psi^{-1}(t))},$$

where \tilde{C}_1 is a positive constant depending only on $C_1, C_2, C_{VR}, N, \Phi, \Psi$.

Second, we prove NLE(Ψ). Indeed, for any $\varepsilon \in (0, \frac{1}{2})$, for any $y \in B(x, \varepsilon \Psi^{-1}(t))$, by HHK(Φ, Ψ), we have

$$\begin{aligned} & |p_t(x, x) - p_t(x, y)| \\ & \leq C_{HHK} \frac{\left(\frac{\Psi}{\Phi}\right)(d(x, y))}{t} \\ & \leq C_{HHK} \frac{\left(\frac{\Psi}{\Phi}\right)(\varepsilon \Psi^{-1}(t)) \left(\frac{\Psi}{\Phi}\right)(\Psi^{-1}(t))}{\left(\frac{\Psi}{\Phi}\right)(\Psi^{-1}(t)) t} \\ & \leq C_{HHK} \varepsilon^{\gamma_1 \wedge \gamma_2} \frac{1}{\Phi(\Psi^{-1}(t))}. \end{aligned}$$

There exists $\varepsilon \in (0, \frac{1}{2})$ depending only on $\tilde{C}_1, C_{HHK}, \gamma_1, \gamma_2$ sufficiently small such that

$$C_{HHK} \varepsilon^{\gamma_1 \wedge \gamma_2} \leq \frac{\tilde{C}_1}{2},$$

hence

$$|p_t(x, x) - p_t(x, y)| \leq \frac{\tilde{C}_1}{2\Phi(\Psi^{-1}(t))} \leq \frac{1}{2} p_t(x, x).$$

Thus we have

$$p_t(x, y) \geq p_t(x, x) - |p_t(x, x) - p_t(x, y)| \geq p_t(x, x) - \frac{1}{2} p_t(x, x) = \frac{1}{2} p_t(x, x) \geq \frac{\tilde{C}_1}{2\Phi(\Psi^{-1}(t))}$$

for any $y \in B(x, \varepsilon \Psi^{-1}(t))$, that is, we have NLE(Ψ). \square

8 Proof of “NLE(Ψ) \Rightarrow HR(Φ, Ψ)”

Recall that a Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$ is recurrent if for any non-negative function $f \in L^1(X; m)$, we have either $Gf = +\infty$ or $Gf = 0$ m -a.e., where G is the Green operator, see [26, Equation (1.6.2)]. In our setting, the Dirichlet form is recurrent as follows.

Lemma 8.1. *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$ and $NLE(\Psi)$. Then $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$ is recurrent.*

Proof. Taking any non-zero non-negative function $f \in L^1(X; m)$, we show that

$$Gf(x) := \int_0^{+\infty} \left(\int_X p_t(x, y) f(y) m(dy) \right) dt = +\infty \text{ for } m\text{-a.e. } x \in X.$$

Indeed, by $V(\Phi)$ and $NLE(\Psi)$, we have

$$\begin{aligned} \int_0^{+\infty} p_t(x, y) dt &\geq \int_{\Psi(\frac{d(x, y)}{\varepsilon})}^{+\infty} p_t(x, y) dt \geq \int_{\Psi(\frac{d(x, y)}{\varepsilon})}^{+\infty} \frac{C}{V(x, \Psi^{-1}(t))} dt \\ &\asymp \int_{\Psi(\frac{d(x, y)}{\varepsilon})}^{+\infty} \frac{1}{\Phi(\Psi^{-1}(t))} dt \geq \int_{\Psi(\frac{d(x, y)}{\varepsilon}) \vee 1}^{+\infty} \frac{dt}{t^{\alpha_2/\beta_2}} = +\infty. \end{aligned}$$

Since f is non-zero non-negative, there exists $a > 0$ such that $m(\{f > a\}) > 0$, then

$$Gf(x) = \int_X \left(\int_0^{+\infty} p_t(x, y) dt \right) f(y) m(dy) \geq a \int_{\{f > a\}} \left(\int_0^{+\infty} p_t(x, y) dt \right) m(dy) = +\infty$$

for m -a.e. $x \in X$. Hence $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$ is recurrent. \square

We use some techniques from resistance forms. The theory of resistance forms was developed by Kigami [44]. To begin with, let us introduce effective resistances. Let A, B be two non-empty subsets of X . The effective resistance between A and B is defined as follows.

$$R(A, B) = (\inf \{ \mathcal{E}(u, u) : u \in \mathcal{F}, u = 0 \text{ on } A, u = 1 \text{ on } B \})^{-1},$$

here we use the convention that $\inf \emptyset = +\infty$, $0^{-1} = +\infty$ and $(+\infty)^{-1} = 0$. For $x, y \in X$, write

$$R(x, A) = R(\{x\}, A), R(x, y) = R(\{x\}, \{y\}).$$

By definition, it is obvious that

$$R(x, y) = \sup \left\{ \frac{|u(x) - u(y)|^2}{\mathcal{E}(u, u)} : u \in \mathcal{F}, \mathcal{E}(u, u) > 0 \right\},$$

which implies that

$$|u(x) - u(y)|^2 \leq R(x, y) \mathcal{E}(u, u) \text{ for any } u \in \mathcal{F}, x, y \in X,$$

and if $A_1 \subseteq A_2, B_1 \subseteq B_2$, then

$$R(A_1, B_1) \geq R(A_2, B_2).$$

Proposition 8.2. *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$, $UHK(\Psi)$ and $NLE(\Psi)$. Then we have two-sided resistance estimates $R(\Phi, \Psi)$ as follows. There exists $C_R \in (0, +\infty)$ such that for any $x, y \in X$ with $x \neq y$, we have*

$$\frac{1}{C_R} \frac{\Psi(d(x, y))}{\Phi(d(x, y))} \leq R(x, y) \leq C_R \frac{\Psi(d(x, y))}{\Phi(d(x, y))}. \quad R(\Phi, \Psi)$$

Remark 8.3. *In the strongly recurrent setting, two-sided heat kernel estimates and two-sided resistance estimates are indeed equivalent. This equivalence was proved in [12] on graphs, in [45] on graphs and resistance forms and in [37] on metric spaces using an analytical approach.*

Under the assumptions of the above result, by Proposition 3.2, we have $\text{PI}(\Psi)$ and $\text{CS}(\Psi)$. The following result will play an important role in this section.

Lemma 8.4. (*Morrey-Sobolev inequality*) *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$ and $\text{PI}(\Psi)$. Then we have the following Morrey-Sobolev inequality $\text{MS}(\Phi, \Psi)$. There exists $C_{MS} \in (0, +\infty)$ such that for any $u \in \mathcal{F}$, for m -a.e. $x, y \in X$ with $x \neq y$, we have*

$$|u(x) - u(y)|^2 \leq C_{MS} \frac{\Psi(d(x, y))}{\Phi(d(x, y))} \mathcal{E}(u, u). \quad \text{MS}(\Phi, \Psi)$$

Hence any function in \mathcal{F} has a continuous version, or equivalently, $\mathcal{F} \subseteq C(X)$.

The proof is standard using telescopic technique, see also [12, Page 1654].

Proof. Let x, y be two different Lebesgue points of $u \in \mathcal{F} \subseteq L^2(X; m)$. Denote $r = d(x, y)$. Then

$$|u(x) - u(y)| \leq |u(x) - u_{B(x, r)}| + |u_{B(x, r)} - u_{B(y, r)}| + |u(y) - u_{B(y, r)}|.$$

For any $n \in \mathbb{Z}$, denote $B_n = B(x, 2^{-n}r)$. By $V(\Phi)$ and $\text{PI}(\Psi)$, we have

$$\begin{aligned} |u_{B_{n+1}} - u_{B_n}| &\leq \int_{B_{n+1}} |u - u_{B_n}| dm \lesssim \int_{B_n} |u - u_{B_n}| dm \leq \left(\int_{B_n} |u - u_{B_n}|^2 dm \right)^{1/2} \\ &\lesssim \left(\frac{1}{\Phi(2^{-n}r)} \Psi(2^{-n}r) \int_{B_{n-1}} d\Gamma(u, u) \right)^{1/2} \leq \left(\frac{\Psi(2^{-n}r)}{\Phi(2^{-n}r)} \mathcal{E}(u, u) \right)^{1/2}, \end{aligned}$$

hence

$$\begin{aligned} |u(x) - u_{B(x, r)}| &= |u(x) - u_{B_0}| = \lim_{n \rightarrow +\infty} |u_{B_n} - u_{B_0}| \leq \lim_{n \rightarrow +\infty} \sum_{k=0}^{n-1} |u_{B_{k+1}} - u_{B_k}| \\ &= \sum_{n=0}^{\infty} |u_{B_{n+1}} - u_{B_n}| \lesssim \sum_{n=0}^{\infty} \left(\frac{\Psi(2^{-n}r)}{\Phi(2^{-n}r)} \mathcal{E}(u, u) \right)^{1/2} \\ &= \left(\frac{\Psi(r)}{\Phi(r)} \mathcal{E}(u, u) \right)^{1/2} \sum_{n=0}^{\infty} \left(\frac{(\frac{\Psi}{\Phi})(2^{-n}r)}{(\frac{\Psi}{\Phi})(r)} \right)^{1/2} \leq \left(\frac{\Psi(r)}{\Phi(r)} \mathcal{E}(u, u) \right)^{1/2} \sum_{n=0}^{\infty} 2^{-\frac{(\gamma_1 \wedge \gamma_2)n}{2}} \\ &\asymp \left(\frac{\Psi(r)}{\Phi(r)} \mathcal{E}(u, u) \right)^{1/2}. \end{aligned}$$

Similarly, we have

$$|u(y) - u_{B(y, r)}| \lesssim \left(\frac{\Psi(r)}{\Phi(r)} \mathcal{E}(u, u) \right)^{1/2}.$$

Note that

$$\begin{aligned} |u_{B(x, r)} - u_{B(y, r)}| &\leq \int_{B(x, r)} \int_{B(y, r)} |u(z_1) - u(z_2)| m(dz_1) m(dz_2) \\ &\leq \left(\int_{B(x, r)} \int_{B(y, r)} |u(z_1) - u(z_2)|^2 m(dz_1) m(dz_2) \right)^{1/2} \\ &\asymp \frac{1}{\Phi(r)} \left(\int_{B(x, r)} \int_{B(y, r)} |u(z_1) - u(z_2)|^2 m(dz_1) m(dz_2) \right)^{1/2} \\ &\leq \frac{1}{\Phi(r)} \left(\int_{B(x, 2r)} \int_{B(x, 2r)} |u(z_1) - u(z_2)|^2 m(dz_1) m(dz_2) \right)^{1/2} \\ &= \frac{1}{\Phi(r)} \left(2V(x, 2r) \int_{B(x, 2r)} (u - u_{B(x, 2r)})^2 dm \right)^{1/2} \end{aligned}$$

$$\lesssim \frac{1}{\Phi(r)^{1/2}} \left(\Psi(2r) \int_{B(x,4r)} d\Gamma(u, u) \right) \lesssim \left(\frac{\Psi(r)}{\Phi(r)} \mathcal{E}(u, u) \right)^{1/2}.$$

In summary, we have

$$|u(x) - u(y)| \lesssim \left(\frac{\Psi(r)}{\Phi(r)} \mathcal{E}(u, u) \right)^{1/2}.$$

□

Proof of Proposition 8.2. The upper bound is obvious by $\text{MS}(\Phi, \Psi)$. We use the idea of an analytical proof of [37, Theorem 3.4] to give the lower bound. By the spectral calculus, it is easy to see that

$$\mathcal{E}(P_t u, P_t u) \leq \frac{1}{2et} (u, u) \text{ for any } t \in (0, +\infty), u \in L^2(X; m),$$

where $\{P_t\}$ is the heat semi-group corresponding to the Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$.

For any $t \in (0, +\infty)$, $x \in X$, take $u = p_t(x, \cdot) \in L^2(X; m)$. By the semi-group property, we have $P_t u = p_{2t}(x, \cdot)$ and $(u, u) = p_{2t}(x, x)$, hence

$$\mathcal{E}(p_{2t}(x, \cdot), p_{2t}(x, \cdot)) \leq \frac{1}{2et} p_{2t}(x, x),$$

clearly

$$\mathcal{E}(p_t(x, \cdot), p_t(x, \cdot)) \leq \frac{1}{et} p_t(x, x).$$

For any $y \in X$, we have

$$p_t(x, x) - p_t(x, y) \leq R(x, y)^{1/2} \mathcal{E}(p_t(x, \cdot), p_t(x, \cdot))^{1/2} \leq R(x, y)^{1/2} \left(\frac{1}{et} p_t(x, x) \right)^{1/2}. \quad (8.1)$$

By $\text{NLE}(\Psi)$, we have

$$p_t(x, x) \geq \frac{C_1}{V(x, \Psi^{-1}(t))}.$$

By $\text{UHK}(\Psi)$, we have

$$p_t(x, y) \leq \frac{C_2}{V(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_3 d(x, y), t)),$$

and

$$p_t(x, x) \leq \frac{C_2}{V(x, \Psi^{-1}(t))}.$$

Hence

$$\begin{aligned} & p_t(x, x) - p_t(x, y) \\ & \geq \frac{C_1}{V(x, \Psi^{-1}(t))} - \frac{C_2}{V(x, \Psi^{-1}(t))} \exp(-\Upsilon(C_3 d(x, y), t)) \\ & = \frac{1}{V(x, \Psi^{-1}(t))} (C_1 - C_2 \exp(-\Upsilon(C_3 d(x, y), t))), \end{aligned}$$

where

$$\begin{aligned} \Upsilon(C_3 d(x, y), t) &= \sup_{s \in (0, +\infty)} \left(\frac{C_3 d(x, y)}{s} - \frac{t}{\Psi(s)} \right) \\ &= \sup_{s \in (0, +\infty)} \left(\frac{C_3 d(x, y)}{\Psi^{-1}(s)} - \frac{t}{s} \right) \geq \frac{C_3 d(x, y)}{\Psi^{-1}(t)} - 1. \end{aligned}$$

Take $\varepsilon \in (0, 1)$, let $t = \Psi(\varepsilon d(x, y))$, then

$$p_t(x, x) - p_t(x, y) \geq \frac{1}{V(x, \Psi^{-1}(t))} \left(C_1 - C_2 \exp\left(1 - \frac{C_3}{\varepsilon}\right) \right) \geq \frac{C_1}{2V(x, \Psi^{-1}(t))}, \quad (8.2)$$

where $\varepsilon \in (0, 1)$ depending only on C_1, C_2, C_3 is sufficiently small such that

$$C_2 \exp\left(1 - \frac{C_3}{\varepsilon}\right) \leq \frac{C_1}{2}.$$

Plugging Equation (8.2) into Equation (8.1), we have

$$\frac{C_1}{2V(x, \Psi^{-1}(t))} \leq R(x, y)^{1/2} \left(\frac{1}{et} p_t(x, x)\right)^{1/2} \leq R(x, y)^{1/2} \left(\frac{1}{et} \frac{C_2}{V(x, \Psi^{-1}(t))}\right)^{1/2},$$

therefore

$$\begin{aligned} R(x, y)^{1/2} &\geq \frac{C_1}{2V(x, \Psi^{-1}(t))} \left(\frac{etV(x, \Psi^{-1}(t))}{C_2}\right)^{1/2} \\ &\asymp \left(\frac{t}{\Phi(\Psi^{-1}(t))}\right)^{1/2} = \left(\frac{\Psi(\varepsilon d(x, y))}{\Phi(\varepsilon d(x, y))}\right)^{1/2} \asymp \left(\frac{\Psi(d(x, y))}{\Phi(d(x, y))}\right)^{1/2}, \end{aligned}$$

that is,

$$R(x, y) \gtrsim \frac{\Psi(d(x, y))}{\Phi(d(x, y))}.$$

□

Let us recall the definition of resistance forms as follows.

Definition 8.5. ([44, DEFINITION 3.1]) *Let X be a set. A pair $(\mathcal{E}, \mathcal{G})$ is called a resistance form on X if the following five conditions are satisfied.*

- (RF1) \mathcal{G} is a linear subspace of $l(X)$ containing constant functions, where $l(X)$ is the space of all real-valued functions on X . \mathcal{E} is a non-negative symmetric quadratic form on \mathcal{G} . For $u \in \mathcal{G}$, $\mathcal{E}(u, u) = 0$ if and only if u is a constant function.
- (RF2) Let \sim be the equivalence relation on \mathcal{G} given by $u, v \in \mathcal{G}$, $u \sim v$ if and only if $u - v$ is a constant function. Then $(\mathcal{G}/\sim, \mathcal{E})$ is a Hilbert space.
- (RF3) For any $x, y \in X$ satisfying $x \neq y$, there exists $u \in \mathcal{G}$ such that $u(x) \neq u(y)$.
- (RF4) For any $x, y \in X$, we have

$$\sup \left\{ \frac{|u(x) - u(y)|^2}{\mathcal{E}(u, u)} : u \in \mathcal{G}, \mathcal{E}(u, u) > 0 \right\}$$

is finite, where we use the convention that $\sup \emptyset = 0$.

- (RF5) For any $u \in \mathcal{G}$, let $\bar{u} = (u \vee 0) \wedge 1$, then $\bar{u} \in \mathcal{G}$ and $\mathcal{E}(\bar{u}, \bar{u}) \leq \mathcal{E}(u, u)$.

Recall that for a Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$, its extended Dirichlet space is defined as the family of all m -measurable functions u which is finite m -a.e. and there exists an \mathcal{E} -Cauchy sequence $\{u_n\} \subseteq \mathcal{F}$ such that $u_n \rightarrow u$ m -a.e.. The sequence $\{u_n\} \subseteq \mathcal{F}$ is called an approximation sequence of $u \in \mathcal{F}_e$. We can extend \mathcal{E} to \mathcal{F}_e and $\mathcal{F} = \mathcal{F}_e \cap L^2(X; m)$. See [26, Section 1.5] for more details. We can also define the harmonicity of a function in the extended Dirichlet space as follows. Let D be a non-empty open subset of X . We say that $u \in \mathcal{F}_e$ is harmonic in D if $\mathcal{E}(u, \varphi) = 0$ for any $\varphi \in \mathcal{F} \cap C_c(D)$.

We collect some results about extended Dirichlet spaces as follows.

Lemma 8.6. *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$, $UHK(\Psi)$ and $NLE(\Psi)$.*

- (1) *The Morrey-Sobolev inequality $MS(\Phi, \Psi)$ holds for any $u \in \mathcal{F}_e$, hence $\mathcal{F}_e \subseteq C(X)$.*
- (2) *$(\mathcal{E}, \mathcal{F}_e)$ is a resistance form on X .*
- (3) *For any $x, y \in X$, we have*

$$R(x, y) = \sup \left\{ \frac{|u(x) - u(y)|^2}{\mathcal{E}(u, u)} : u \in \mathcal{F}_e, \mathcal{E}(u, u) > 0 \right\}.$$

(4) Let $u \in \mathcal{F}_e$ be harmonic in a non-empty open subset D of X . Then $\mathcal{E}(u, \varphi) = 0$ for any $\varphi \in \mathcal{F}_e \subseteq C(X)$ with $\varphi = 0$ on $X \setminus D$.

Proof. (1) It is obvious by the definition of \mathcal{F}_e and Lemma 8.4.

(2) Since the Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$ is recurrent, by [26, Theorem 1.6.3], we have $1 \in \mathcal{F}_e$ and $\mathcal{E}(1, 1) = 0$. For $u \in \mathcal{F}_e$, if $\mathcal{E}(u, u) = 0$, then by MS(Φ, Ψ), we have u is a constant function. Hence we have (RF1).

By (RF1), we have $(\mathcal{F}_e / \sim, \mathcal{E})$ is an inner product space. We only need to prove the completeness. Indeed, let $\{u_n\} \subseteq \mathcal{F}_e / \sim$ be an \mathcal{E} -Cauchy sequence, then $\{u_n\}$ is \mathcal{E} -bounded. Fix arbitrary $x_0 \in X$. By replacing u_n by $u_n - u_n(x_0)$ in \mathcal{F}_e / \sim , we may assume that $u_n(x_0) = 0$ for any $n \geq 1$.

For any $N \geq 1$, for any $x \in B(x_0, N)$, for any $n \geq 1$, by MS(Φ, Ψ), we have

$$|u_n(x)|^2 = |u_n(x) - u_n(x_0)|^2 \leq C_{MS} \frac{\Psi(d(x, x_0))}{\Phi(d(x, x_0))} \mathcal{E}(u_n, u_n) \leq C_{MS} \frac{\Psi(N)}{\Phi(N)} \sup_{n \geq 1} \mathcal{E}(u_n, u_n).$$

By MS(Φ, Ψ) again, using the Arzelà-Ascoli lemma and the diagonal argument, there exists a subsequence, still denoted by $\{u_n\}$, and $u \in C(X)$ such that u_n converges uniformly to u on any relatively compact open subset of X . By the definition of \mathcal{F}_e , we have $u \in \mathcal{F}_e$ and $\{u_n\}$ is \mathcal{E} -convergent to u . Hence $(\mathcal{F}_e / \sim, \mathcal{E})$ is complete, hence we have (RF2).

(RF3) follows from the regularity of the Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$.

(RF4) follows from MS(Φ, Ψ).

(RF5) follows from [26, Corollary 1.6.3].

(3) We need to prove that

$$\sup \left\{ \frac{|u(x) - u(y)|^2}{\mathcal{E}(u, u)} : u \in \mathcal{F}, \mathcal{E}(u, u) > 0 \right\} = \sup \left\{ \frac{|u(x) - u(y)|^2}{\mathcal{E}(u, u)} : u \in \mathcal{F}_e, \mathcal{E}(u, u) > 0 \right\}.$$

Indeed, “ \leq ” is trivial by $\mathcal{F} \subseteq \mathcal{F}_e$. “ \geq ”: For any $u \in \mathcal{F}_e$ with $\mathcal{E}(u, u) > 0$, there exists $\{u_n\} \subseteq \mathcal{F}$ which is \mathcal{E} -Cauchy and converges to u m -a.e.. Since $\{u_n\}$ is \mathcal{E} -convergent to u , we may assume that $\mathcal{E}(u_n, u_n) > 0$ for any $n \geq 1$. Take $x_0 \in X$ such that $\lim_{n \rightarrow +\infty} u_n(x_0) = u(x_0) \in \mathbb{R}$, then $\{u_n(x_0)\}$ is bounded. For any $N \geq 1$, for any $x \in B(x_0, N)$, for any $n \geq 1$, by MS(Φ, Ψ), we have

$$\begin{aligned} |u_n(x)|^2 &\leq 2 (|u_n(x) - u_n(x_0)|^2 + |u_n(x_0)|^2) \\ &\leq 2 \left(C_{MS} \frac{\Psi(d(x, x_0))}{\Phi(d(x, x_0))} \mathcal{E}(u_n, u_n) + \sup_{n \geq 1} |u_n(x_0)|^2 \right) \\ &\leq 2 \left(C_{MS} \frac{\Psi(N)}{\Phi(N)} \sup_{n \geq 1} \mathcal{E}(u_n, u_n) + \sup_{n \geq 1} |u_n(x_0)|^2 \right). \end{aligned}$$

By MS(Φ, Ψ) again, using the Arzelà-Ascoli lemma and the diagonal argument, there exists a subsequence, still denoted by $\{u_n\}$, and $v \in C(X)$ such that u_n converges uniformly to v on any relatively compact open subset of X . Since u_n also converges to u m -a.e. and $u, v \in C(X)$, we have $u = v$. Hence

$$\frac{|u(x) - u(y)|^2}{\mathcal{E}(u, u)} = \lim_{n \rightarrow +\infty} \frac{|u_n(x) - u_n(y)|^2}{\mathcal{E}(u_n, u_n)} \leq \text{LHS}.$$

Taking supremum with respect to $u \in \mathcal{F}_e$ with $\mathcal{E}(u, u) > 0$, we have “ \geq ”.

(4) Since $(\mathcal{E}, \mathcal{F}_e)$ is a resistance form on X , by [44, PROPOSITION 9.13], we have the continuity and the quasi-continuity coincide. Then $\mathcal{F}_{e,D}$ given by [26, Equation (2.3.14)] can be re-written as

$$\mathcal{F}_{e,D} := \{u \in \mathcal{F}_e : \tilde{u} = 0 \text{ q.e. on } X \setminus D\} = \{u \in \mathcal{F}_e \subseteq C(X) : u = 0 \text{ on } X \setminus D\},$$

where \tilde{u} is a quasi-continuous modification of $u \in \mathcal{F}_e$.

Since the Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$ is regular, by [26, Exercise 1.4.1], we have $\mathcal{F} \cap C_c(X)$ is a special standard core of \mathcal{E} . Then (4) follows from the equivalence in [26, Theorem 2.3.3 (ii)]. \square

By the resistance form theory, we can solve “boundary value problems” or take a trace of resistance forms as follows.

Lemma 8.7. ([44, Section 8]) *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$, $UHK(\Psi)$ and $NLE(\Psi)$. Let $Y \subsetneq X$ be a non-empty closed subset and $\mathcal{F}_e|_Y = \{u|_Y : u \in \mathcal{F}_e\}$. Then for any $u \in \mathcal{F}_e$, there exists a unique $v \in \mathcal{F}_e$ which is harmonic in $X \setminus Y$ such that $v = u$ on Y . Moreover, v is the unique function in \mathcal{F}_e that minimizes the following variational problem*

$$\inf \{ \mathcal{E}(v, v) : v \in \mathcal{F}_e, v = u \text{ on } Y \}.$$

Let $h_Y : \mathcal{F}_e|_Y \rightarrow \mathcal{F}_e$ be given by $u|_Y \mapsto v$ for $u \in \mathcal{F}_e$ as above. Let

$$\mathcal{E}|_Y(u, v) = \mathcal{E}(h_Y(u), h_Y(v))$$

for $u, v \in \mathcal{F}_e|_Y$. Then $(\mathcal{E}|_Y, \mathcal{F}_e|_Y)$ is a resistance form on Y .

We have the following maximum principle.

Corollary 8.8. (Maximum principle) *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$, $UHK(\Psi)$ and $NLE(\Psi)$. Let $D \subsetneq X$ be a non-empty open subset of X . Let $u, v \in \mathcal{F}_e$ be harmonic in D and satisfy $u \leq v$ on $X \setminus D$, then $u \leq v$ in D . In particular, $\inf_{X \setminus D} u \leq u \leq \sup_{X \setminus D} u$.*

Proof. We only need to show that if $u \in \mathcal{F}_e$ is harmonic in D and satisfies $u \leq 0$ on $X \setminus D$, then $u \leq 0$ in D . Indeed, $u \wedge 0 \in \mathcal{F}_e$ satisfies $u \wedge 0 = u$ on $X \setminus D$, by Lemma 8.7, we have

$$\mathcal{E}(u, u) \geq \mathcal{E}(u \wedge 0, u \wedge 0) \geq \inf \{ \mathcal{E}(v, v) : v \in \mathcal{F}_e, v = u \text{ on } X \setminus D \} = \mathcal{E}(u, u).$$

Hence the above inequalities are indeed equalities. By the uniqueness of $u \in \mathcal{F}_e$ from Lemma 8.7, we have $u \wedge 0 = u$, hence $u \leq 0$ in D . \square

The effective resistance between certain two sets can be achieved by the energy of some functions as follows.

Corollary 8.9. *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$, $UHK(\Psi)$ and $NLE(\Psi)$. Let A, B be two non-empty closed subsets of X satisfying $A \cap B = \emptyset$ and $A \cup B \subsetneq X$. If one of the following conditions holds.*

- (a) $X \setminus A$ is relatively compact and B is compact.
- (b) A, B are compact.

Then there exists a unique $v \in \mathcal{F}_e$ with $v = 0$ on A and $v = 1$ on B such that

$$R(A, B)^{-1} = \mathcal{E}(v, v) = \inf \{ \mathcal{E}(v, v) : v \in \mathcal{F}_e, v = 0 \text{ on } A, v = 1 \text{ on } B \}.$$

Moreover, $1_A, 1_B \in \mathcal{F}_e|_{A \cup B}$, $v = h_{A \cup B}(1_B) \in \mathcal{F}_e$ and

$$R(A, B)^{-1} = \mathcal{E}(h_{A \cup B}(1_A), h_{A \cup B}(1_A)) = \mathcal{E}(h_{A \cup B}(1_B), h_{A \cup B}(1_B)).$$

Proof. Since B is compact, $X \setminus A$ is open and $B \subseteq X \setminus A$, by the regularity of $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$, there exists $u \in \mathcal{F} \cap C_c(X) \subseteq \mathcal{F}_e$ such that $0 \leq u \leq 1$ in X , $u = 1$ on B and $\text{supp}(u) \subseteq X \setminus A$. Hence $u|_{A \cup B} = 1_B \in \mathcal{F}_e|_{A \cup B}$, $(1 - u)|_{A \cup B} = 1_A \in \mathcal{F}_e|_{A \cup B}$.

Since $A \cup B \subsetneq X$ is a non-empty closed subset, by Lemma 8.7, there exists a unique $v = h_{A \cup B}(u|_{A \cup B}) = h_{A \cup B}(1_B) \in \mathcal{F}_e$ which is harmonic in $X \setminus (A \cup B)$ and $v = u$ on $A \cup B$, that is, $v|_{A \cup B} = 1_B$, or $v = 0$ on A and $v = 1$ on B . Moreover, $v \in \mathcal{F}_e$ also minimizes the variational problem

$$\inf \{ \mathcal{E}(v, v) : v \in \mathcal{F}_e, v = 0 \text{ on } A, v = 1 \text{ on } B \},$$

and

$$\begin{aligned} \mathcal{E}(v, v) &= \mathcal{E}|_{A \cup B}(1_B, 1_B) = \mathcal{E}|_{A \cup B}(1_A, 1_A) \\ &= \mathcal{E}(h_{A \cup B}(1_B), h_{A \cup B}(1_B)) = \mathcal{E}(h_{A \cup B}(1_A), h_{A \cup B}(1_A)). \end{aligned}$$

It remains to prove that if either (a) or (b) holds, then

$$\mathcal{E}(v, v) = \inf \{ \mathcal{E}(v, v) : v \in \mathcal{F}, v = 0 \text{ on } A, v = 1 \text{ on } B \},$$

where the RHS is equal to $R(A, B)^{-1}$ by definition. Since $\mathcal{F} \subseteq \mathcal{F}_e$, we only need to show “ \geq ”. By the maximum principle Corollary 8.8, we have $0 \leq v \leq 1$ in X .

For (a), since $0 \leq v \leq 1$ in X , $v = 0$ on A and $X \setminus A$ is a relatively compact open set, we have $v \in \mathcal{F}_e \cap L^2(X; m) = \mathcal{F}$, “ \geq ” is obvious.

For (b), since A, B are both compact, by a similar proof to Lemma 8.6 (2), there exists an approximation sequence $\{v_n\} \subseteq \mathcal{F} \subseteq C(X)$ of $v \in \mathcal{F}_e$ that converges uniformly to v on $A \cup B$. For any $\varepsilon \in (0, 1/4)$, there exists $N \geq 1$ such that for any $n \geq N$, we have $|v_n - v| < \varepsilon$ on $A \cup B$, hence $-\varepsilon < v_n < \varepsilon$ on A and $1 - \varepsilon < v_n < 1 + \varepsilon$ on B . Let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be given by

$$\varphi(t) = \frac{\frac{1}{2}}{\frac{1}{2} - \varepsilon} \left((t \wedge (1 - \varepsilon)) \vee \varepsilon - \frac{1}{2} \right) + \frac{1}{2}, t \in \mathbb{R},$$

that is, $\varphi \in C(\mathbb{R})$ satisfies $\varphi = 0$ on $(-\infty, \varepsilon]$, $\varphi = 1$ on $[1 - \varepsilon, +\infty)$ and φ is linear on $[\varepsilon, 1 - \varepsilon]$. Then $\varphi(0) = 0$ and $|\varphi(t) - \varphi(s)| \leq \frac{\frac{1}{2}}{\frac{1}{2} - \varepsilon} |t - s|$ for any $t, s \in \mathbb{R}$. For any $n \geq N$, let $\tilde{v}_n = \varphi(v_n)$, then $\tilde{v}_n \in \mathcal{F}$ satisfies $\tilde{v}_n = 0$ on A , $\tilde{v}_n = 1$ on B and $\mathcal{E}(\tilde{v}_n, \tilde{v}_n) \leq \left(\frac{\frac{1}{2}}{\frac{1}{2} - \varepsilon} \right)^2 \mathcal{E}(v_n, v_n)$, hence

$$\inf \{ \mathcal{E}(v, v) : v \in \mathcal{F}, v = 0 \text{ on } A, v = 1 \text{ on } B \} \leq \mathcal{E}(\tilde{v}_n, \tilde{v}_n) \leq \left(\frac{\frac{1}{2}}{\frac{1}{2} - \varepsilon} \right)^2 \mathcal{E}(v_n, v_n).$$

Letting $n \rightarrow +\infty$, we have

$$\inf \{ \mathcal{E}(v, v) : v \in \mathcal{F}, v = 0 \text{ on } A, v = 1 \text{ on } B \} \leq \left(\frac{\frac{1}{2}}{\frac{1}{2} - \varepsilon} \right)^2 \mathcal{E}(v, v).$$

Since $\varepsilon \in (0, \frac{1}{4})$ is arbitrary, we have

$$\inf \{ \mathcal{E}(v, v) : v \in \mathcal{F}, v = 0 \text{ on } A, v = 1 \text{ on } B \} \leq \mathcal{E}(v, v),$$

that is, we have “ \geq ”. □

Remark 8.10. We will be interested in the case $A = X \setminus B(x, r)$, $B = \{y\}$ for (a), and the case $A = \{x\}$, $B = \{y\}$ for (b), where $x, y \in X$, $r \in (0, +\infty)$.

Lemma 8.11. Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$, $UHK(\Psi)$ and $NLE(\Psi)$. Then there exists $C \in (0, +\infty)$ such that for any $x_0 \in X$, for any $r \in (0, +\infty)$, we have

$$R(x_0, X \setminus B(x_0, r)) \geq C \frac{\Psi(r)}{\Phi(r)}.$$

The proof is essentially the same as the proof of [12, LEMMA 2.4], [45, Lemma 4.1] and [37, Proposition 5.3].

Proof. Denote $B = B(x_0, r)$. For any $x \in B \setminus (\frac{1}{2}B)$, by Corollary 8.9, there exists $h_x \in \mathcal{F}_e$ with $h_x(x_0) = 1$, $h_x(x) = 0$ and $0 \leq h_x \leq 1$ in X such that

$$\mathcal{E}(h_x, h_x) = R(x_0, x)^{-1} \leq \left(\frac{1}{C_R} \frac{\Psi(d(x_0, x))}{\Phi(d(x_0, x))} \right)^{-1} = C_R \frac{\Phi(d(x_0, x))}{\Psi(d(x_0, x))} \leq C_R \frac{\Phi(\frac{r}{2})}{\Psi(\frac{r}{2})},$$

where C_R is the constant in $R(\Phi, \Psi)$. Take $\varepsilon \in (0, \frac{1}{4})$ to be determined. For any $y \in B(x, \varepsilon r)$, we have

$$\begin{aligned} |h_x(y)| &= |h_x(x) - h_x(y)| \leq R(x, y)^{1/2} \mathcal{E}(h_x, h_x)^{1/2} \\ &\leq \left(C_R \frac{\Psi(d(x, y))}{\Phi(d(x, y))} C_R \frac{\Phi(\frac{r}{2})}{\Psi(\frac{r}{2})} \right)^{1/2} \leq C_R \left(\frac{\Psi(\varepsilon r)}{\Phi(\varepsilon r)} \frac{\Phi(\frac{r}{2})}{\Psi(\frac{r}{2})} \right)^{1/2} \leq C_R (2\varepsilon)^{\frac{\gamma_1 \wedge \gamma_2}{2}}. \end{aligned}$$

Then there exists $\varepsilon \in (0, \frac{1}{4})$ depending only on C_R, γ_1, γ_2 sufficiently small such that

$$|h_x(y)| \leq C_R(2\varepsilon)^{\frac{\gamma_1 \wedge \gamma_2}{2}} \leq \frac{1}{2} \text{ for any } y \in B(x, \varepsilon r).$$

By $V(\Phi)$, there exists $N \geq 1$ depending only on $C_{VR}, \alpha_1, \alpha_2, \varepsilon$, hence only on $C_{VR}, C_R, \alpha_1, \alpha_2, \beta_1, \beta_2$, such that $B \setminus (\frac{1}{2}B)$ can be covered by at most N balls with center in $B \setminus (\frac{1}{2}B)$ and radius εr . Take $x_1, \dots, x_N \in B \setminus (\frac{1}{2}B)$ such that $B \setminus (\frac{1}{2}B) \subseteq \cup_{1 \leq i \leq N} B(x_i, \varepsilon r)$, let $h = \min_{1 \leq i \leq N} h_{x_i}$. Then $h \in \mathcal{F}_e$ satisfies $h(x_0) = 1$, $0 \leq h \leq 1$ in X , $h \leq \frac{1}{2}$ in $B \setminus (\frac{1}{2}B)$ and

$$\mathcal{E}(h, h) \leq \sum_{i=1}^N \mathcal{E}(h_{x_i}, h_{x_i}) \leq \sum_{i=1}^N C_R \frac{\Phi(\frac{r}{2})}{\Psi(\frac{r}{2})} = C_R N \frac{\Phi(\frac{r}{2})}{\Psi(\frac{r}{2})}.$$

Let $g = 2((h - \frac{1}{2}) \vee 0)$, then $g \in \mathcal{F}_e$ satisfies $g(x_0) = 1$, $0 \leq g \leq 1$ in X , $g = 0$ in $B \setminus (\frac{1}{2}B)$ and

$$\mathcal{E}(g, g) = 4\mathcal{E}((h - \frac{1}{2}) \vee 0, (h - \frac{1}{2}) \vee 0) \leq 4\mathcal{E}(h, h).$$

By the regularity of the Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(X; m)$, there exists $\varphi \in \mathcal{F} \cap C_c(X)$ satisfying $0 \leq \varphi \leq 1$ in X , $\varphi = 1$ in an open neighborhood of $\frac{1}{2}\bar{B}$ and $\text{supp}(\varphi) \subseteq B$. Then $\varphi g \in \mathcal{F}_e$ satisfies $(\varphi g)(x_0) = 1$, $0 \leq \varphi g \leq 1$ in X and $\varphi g = g1_B = g1_{\frac{1}{2}B}$. By the strongly local property and Corollary 8.9, we have

$$\begin{aligned} R(x_0, X \setminus B)^{-1} &\leq \mathcal{E}(\varphi g, \varphi g) = \Gamma(\varphi g, \varphi g)(X) = \Gamma(\varphi g, \varphi g)(B) \\ &= \Gamma(g, g)(B) \leq \Gamma(g, g)(X) = \mathcal{E}(g, g) \leq 4\mathcal{E}(h, h) \leq 4C_R N \frac{\Phi(\frac{r}{2})}{\Psi(\frac{r}{2})}, \end{aligned}$$

hence

$$R(x_0, X \setminus B) \geq \frac{1}{4C_R N} \frac{\Psi(\frac{r}{2})}{\Phi(\frac{r}{2})} = \frac{1}{4C_R N} \frac{\Psi(\frac{r}{2})}{\Phi(\frac{r}{2})} \frac{\Psi(r)}{\Phi(r)} \geq \frac{1}{4 \cdot 2^{\gamma_1 \vee \gamma_2} C_R N} \frac{\Psi(r)}{\Phi(r)}.$$

□

Proposition 8.12. *Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying $V(\Phi)$, $UHK(\Psi)$ and $NLE(\Psi)$. Then for any ball $B = B(x_0, r)$, for any bounded $u \in \mathcal{F}_e$ which is harmonic in B , for any $x, y \in B$, we have*

$$|u(x) - u(y)| \leq \frac{R(x, y)}{R(x, X \setminus B)} \text{osc}_{X \setminus B} u,$$

where

$$\text{osc}_A u = \sup_A u - \inf_A u$$

is the oscillation of the function $u \in l(A)$ on the set A .

The idea of the proof is from a recent result [40, Theorem 6.27], where a general result in the setting of p -energy forms was given to prove certain p -resistance defines indeed a metric.

Proof. Let $Y = X \setminus B$, then $Y \subsetneq X$ is a non-empty closed subset of X . For fixed $x, y \in B$, we have $u - u(x) \in \mathcal{F}_e$ is harmonic in $B \setminus \{x\}$, hence $u - u(x) = h_{Y \cup \{x\}}(u - u(x))$. By the maximum principle Corollary 8.8, we have $u - u(x) \leq (\text{osc}_Y u)1_Y$ on $Y \cup \{x\}$, hence

$$u - u(x) = h_{Y \cup \{x\}}(u - u(x)) \leq h_{Y \cup \{x\}}((\text{osc}_Y u)1_Y) = (\text{osc}_Y u)h_{Y \cup \{x\}}(1_Y).$$

It remains to estimate $h_{Y \cup \{x\}}(1_Y)(y)$. Indeed, by Corollary 8.9, we have

$$\begin{aligned} R(x, X \setminus B)^{-1} &= R(x, Y)^{-1} = \mathcal{E}(h_{Y \cup \{x\}}(1_Y), h_{Y \cup \{x\}}(1_Y)) \\ &= \mathcal{E}(h_{Y \cup \{x\}}(1_Y), h_{Y \cup \{x\}}(1_Y) - 1) = -\mathcal{E}(h_{Y \cup \{x\}}(1_Y), h_{Y \cup \{x\}}(1_{\{x\}})). \end{aligned}$$

Since $\mathcal{F}_e \ni h_{Y \cup \{x,y\}}(1_{\{x\}}) - h_{Y \cup \{x\}}(1_{\{x\}}) = 0$ on $Y \cup \{x\}$ and $h_{Y \cup \{x\}}(1_Y) \in \mathcal{F}_e$ is harmonic in $B \setminus \{x\}$, by Lemma 8.6 (4), we have

$$\mathcal{E}(h_{Y \cup \{x\}}(1_Y), h_{Y \cup \{x,y\}}(1_{\{x\}}) - h_{Y \cup \{x\}}(1_{\{x\}})) = 0,$$

that is,

$$\mathcal{E}(h_{Y \cup \{x\}}(1_Y), h_{Y \cup \{x\}}(1_{\{x\}})) = \mathcal{E}(h_{Y \cup \{x\}}(1_Y), h_{Y \cup \{x,y\}}(1_{\{x\}})).$$

Let

$$\begin{aligned} u_1 &= h_{Y \cup \{x\}}(1_Y) - h_{Y \cup \{x\}}(1_Y)(y)h_{\{x,y\}}(1_{\{y\}}), \\ v_1 &= h_{Y \cup \{x,y\}}(1_{\{x\}}), \end{aligned}$$

then $u_1, v_1 \in \mathcal{F}_e$ are harmonic in $B \setminus \{x, y\}$. Let $u_2 = u_1|_{Y \cup \{x,y\}}$, $v_2 = v_1|_{Y \cup \{x,y\}}$, then $u_2, v_2 \in \mathcal{F}_e|_{Y \cup \{x,y\}}$ and

$$\mathcal{E}(u_1, v_1) = \mathcal{E}|_{Y \cup \{x,y\}}(u_2, v_2),$$

where $(\mathcal{E}|_{Y \cup \{x,y\}}, \mathcal{F}_e|_{Y \cup \{x,y\}})$ is a resistance form on $Y \cup \{x, y\}$, which follows from Lemma 8.7. On Y , we have $u_2 \in [0, 1]$, $v_2 = 0$, $u_2 \wedge v_2 = 0$, $u_2 \vee v_2 = u_2 = u_2 + v_2$. On $\{x\}$, we have $u_2 = 0$, $v_2 = 1$, $u_2 \wedge v_2 = 0$, $u_2 \vee v_2 = 1 = v_2 = u_2 + v_2$. On $\{y\}$, we have $u_2 = 0$, $v_2 = 0$, $u_2 \wedge v_2 = 0$, $u_2 \vee v_2 = 0 = u_2 + v_2$. Hence $u_2 \wedge v_2 = 0$, $u_2 \vee v_2 = u_2 + v_2$. By the following easy result Lemma 8.13, we have

$$\mathcal{E}(u_1, v_1) = \mathcal{E}|_{Y \cup \{x,y\}}(u_2, v_2) \leq 0,$$

that is,

$$\mathcal{E}(h_{Y \cup \{x\}}(1_Y), h_{Y \cup \{x,y\}}(1_{\{x\}})) \leq \mathcal{E}(h_{Y \cup \{x\}}(1_Y)(y)h_{\{x,y\}}(1_{\{y\}}), h_{Y \cup \{x,y\}}(1_{\{x\}})),$$

hence

$$\begin{aligned} R(x, X \setminus B)^{-1} &= -\mathcal{E}(h_{Y \cup \{x\}}(1_Y), h_{Y \cup \{x\}}(1_{\{x\}})) = -\mathcal{E}(h_{Y \cup \{x\}}(1_Y), h_{Y \cup \{x,y\}}(1_{\{x\}})) \\ &\geq -\mathcal{E}(h_{Y \cup \{x\}}(1_Y)(y)h_{\{x,y\}}(1_{\{y\}}), h_{Y \cup \{x,y\}}(1_{\{x\}})) \\ &= -h_{Y \cup \{x\}}(1_Y)(y)\mathcal{E}(h_{\{x,y\}}(1_{\{y\}}), h_{Y \cup \{x,y\}}(1_{\{x\}})). \end{aligned}$$

Since $\mathcal{F}_e \ni h_{Y \cup \{x,y\}}(1_{\{x\}}) - h_{\{x,y\}}(1_{\{x\}}) = 0$ on $\{x, y\}$ and $h_{\{x,y\}}(1_{\{y\}}) \in \mathcal{F}_e$ is harmonic in $X \setminus \{x, y\}$, by Lemma 8.6 (4), we have

$$\mathcal{E}(h_{\{x,y\}}(1_{\{y\}}), h_{Y \cup \{x,y\}}(1_{\{x\}}) - h_{\{x,y\}}(1_{\{x\}})) = 0,$$

that is,

$$\mathcal{E}(h_{\{x,y\}}(1_{\{y\}}), h_{Y \cup \{x,y\}}(1_{\{x\}})) = \mathcal{E}(h_{\{x,y\}}(1_{\{y\}}), h_{\{x,y\}}(1_{\{x\}})).$$

Hence

$$\begin{aligned} R(x, X \setminus B)^{-1} &\geq -h_{Y \cup \{x\}}(1_Y)(y)\mathcal{E}(h_{\{x,y\}}(1_{\{y\}}), h_{Y \cup \{x,y\}}(1_{\{x\}})) \\ &= -h_{Y \cup \{x\}}(1_Y)(y)\mathcal{E}(h_{\{x,y\}}(1_{\{y\}}), h_{\{x,y\}}(1_{\{x\}})) \\ &= h_{Y \cup \{x\}}(1_Y)(y)\mathcal{E}(1 - h_{\{x,y\}}(1_{\{y\}}), h_{\{x,y\}}(1_{\{x\}})) \\ &= h_{Y \cup \{x\}}(1_Y)(y)\mathcal{E}(h_{\{x,y\}}(1_{\{x\}}), h_{\{x,y\}}(1_{\{x\}})) \\ &= h_{Y \cup \{x\}}(1_Y)(y)R(x, y)^{-1}, \end{aligned}$$

which implies

$$h_{Y \cup \{x\}}(1_Y)(y) \leq \frac{R(x, y)}{R(x, X \setminus B)}.$$

Thus we have

$$u(y) - u(x) \leq (\text{osc}_Y u) \frac{R(x, y)}{R(x, X \setminus B)}.$$

Similarly, since $u - u(x) \geq -(\text{osc}_Y u)1_Y$ on $Y \cup \{x\}$, we have

$$u - u(x) = h_{Y \cup \{x\}}(u - u(x)) \geq -h_{Y \cup \{x\}}((\text{osc}_Y u)1_Y) = -(\text{osc}_Y u)h_{Y \cup \{x\}}(1_Y),$$

which implies

$$u(y) - u(x) \geq -(\operatorname{osc}_Y u) h_{Y \cup \{x\}}(1_Y)(y) \geq -(\operatorname{osc}_Y u) \frac{R(x, y)}{R(x, X \setminus B)}.$$

Therefore, we have

$$|u(x) - u(y)| \leq \frac{R(x, y)}{R(x, X \setminus B)} \operatorname{osc}_Y u.$$

□

Lemma 8.13. *Let X be a set. Let $(\mathcal{E}, \mathcal{G})$ be a resistance form on X . If $u, v \in \mathcal{G}$ satisfy $u \wedge v = 0$ and $u \vee v = u + v$, then $\mathcal{E}(u, v) \leq 0$.*

Proof. By the parallelogram identity and (RF5), we have

$$\mathcal{E}(u \wedge v, u \wedge v) + \mathcal{E}(u \vee v, u \vee v) \leq \mathcal{E}(u, u) + \mathcal{E}(v, v).$$

By assumption, we have

$$\mathcal{E}(u, u) + \mathcal{E}(v, v) \geq \mathcal{E}(u + v, u + v) = \mathcal{E}(u, u) + \mathcal{E}(v, v) + 2\mathcal{E}(u, v),$$

hence $\mathcal{E}(u, v) \leq 0$. □

We need the following L^1 -mean value inequality.

Lemma 8.14. *([31, THEOREM 6.3, LEMMA 9.2]) Let $(X, d, m, \mathcal{E}, \mathcal{F})$ be an unbounded MMD space satisfying VD, $FK(\Psi)$ and $CS(\Psi)$. Then there exists $C \in (0, +\infty)$ such that for any ball $B = B(x_0, r)$, for any $u \in \mathcal{F}$ which is harmonic in $2B$, we have*

$$\|u\|_{L^\infty(B)} \leq C \int_{2B} |u| dm.$$

We now give the proof of “ $NLE(\Psi) \Rightarrow HR(\Phi, \Psi)$ ” as follows.

Proof of “ $NLE(\Psi) \Rightarrow HR(\Phi, \Psi)$ ”. Note that we need to prove $HR(\Phi, \Psi)$ from $UHK(\Psi)$ and $NLE(\Psi)$. By Proposition 8.2, we have $R(\Phi, \Psi)$. Let $B = B(x_0, r)$ and $u \in \mathcal{F}$ harmonic in $2B$. By VD and Lemma 8.14, we have

$$M := \|u\|_{L^\infty(\frac{3}{2}B)} \lesssim \int_{2B} |u| dm.$$

Let $v = (u \vee (-M)) \wedge M$, then $v \in \mathcal{F} \cap L^\infty(X; m)$, $\|v\|_{L^\infty(X)} \leq M$ and $v = u$ in $\frac{3}{2}B$, hence $v \in \mathcal{F} \subseteq \mathcal{F}_e$ is bounded in X and harmonic in $\frac{3}{2}B$. By Proposition 8.12, for any $x, y \in B \subseteq \frac{3}{2}B$, we have

$$\begin{aligned} |v(x) - v(y)| &\leq \frac{R(x, y)}{R(x, X \setminus (\frac{3}{2}B))} \operatorname{osc}_{X \setminus (\frac{3}{2}B)} v \\ &\leq 2 \frac{R(x, y)}{R(x, X \setminus B(x, \frac{r}{2}))} \|v\|_{L^\infty(X)} \leq 2M \frac{R(x, y)}{R(x, X \setminus B(x, \frac{r}{2}))}, \end{aligned}$$

hence

$$|u(x) - u(y)| \leq 2\|u\|_{L^\infty(\frac{3}{2}B)} \frac{R(x, y)}{R(x, X \setminus B(x, \frac{r}{2}))} \lesssim \frac{R(x, y)}{R(x, X \setminus B(x, \frac{r}{2}))} \int_{2B} |u| dm.$$

By $R(\Phi, \Psi)$ and Lemma 8.11, we have

$$|u(x) - u(y)| \lesssim \frac{\binom{\Psi}{\Phi}(d(x, y))}{\binom{\Psi}{\Phi}(\frac{r}{2})} \int_{2B} |u| dm \asymp \frac{\binom{\Psi}{\Phi}(d(x, y))}{\binom{\Psi}{\Phi}(r)} \int_{2B} |u| dm,$$

hence $HR(\Phi, \Psi)$ holds. □

Remark 8.15. *A natural question is to consider the sharpness of the Hölder exponents $\beta_1 - \alpha_1$ and $\beta_2 - \alpha_2$ from $HR(\Phi, \Psi)$. Indeed, on a class of unbounded p.c.f. self-similar sets (see [54]) and the corresponding cable systems, these exponents are sharp, see the proof of [25, Proposition 4.1, Proposition 4.3] for the Vicsek and the SG cable systems. However, on the unbounded SC and the SC cable system, it was conjectured in [1, Conjecture 5.4, Open Question 5.5] that these exponents are not sharp, and a new fractal dimension, that is, the so-called topological Hausdorff dimension (see [4]), should be involved.*

9 Some examples related to the Sierpiński carpet and the Vicsek set

In this section, we give some examples related to the Sierpiński carpet and the Vicsek set including the blowups of a fractal by another different fractal. In \mathbb{R}^2 , let

$$\begin{aligned} p_1 &= (0, 0), p_2 = \left(\frac{1}{2}, 0\right), p_3 = (1, 0), p_4 = \left(1, \frac{1}{2}\right), \\ p_5 &= (1, 1), p_6 = \left(\frac{1}{2}, 1\right), p_7 = (0, 1), p_8 = \left(0, \frac{1}{2}\right), \end{aligned} \quad (9.1)$$

and

$$g_i(x) = \frac{1}{3}(x - p_i) + p_i, x \in \mathbb{R}^2, i = 1, \dots, 8.$$

Then the Sierpiński carpet is the unique non-empty compact set K in \mathbb{R}^2 satisfying $K = \cup_{i=1}^8 g_i(K)$, see Figure 2. Let $\alpha = \log 8 / \log 3$. By [42, Theorem 1.5.7, Proposition 1.5.8], the Hausdorff dimension of K is α and the α -dimensional Hausdorff measure $\mathcal{H}^\alpha(K) \in (0, +\infty)$. The construction of Brownian motion on the Sierpiński carpet was given by [6] and [46] using approximation of stochastic processes. See also [33] for another construction using Γ -convergence of quadratic forms, [9] for the construction on higher dimensional Sierpiński carpets, [17] for the construction on unconstrained Sierpiński carpets. Let $\beta = \log(8\rho) / \log 3 \approx 2.09697$, where $\rho \approx 1.25147 \in [\frac{7}{6}, \frac{3}{2}]$ is some parameter from resistance estimates in [7, 11].

For $a \in \mathbb{R}$, $A \subseteq \mathbb{R}^2$, denote $aA = \{ax : x \in A\}$.

Firstly, we consider the unbounded Sierpiński carpet $K^\infty = \cup_{n \geq 0} 3^n K$ as follows, where $\{3^n K\}_{n \geq 0}$ is an increasing sequence of subsets of \mathbb{R}^2 . Let $\alpha_1 = \alpha_2 = \alpha$, $\beta_1 = \beta_2 = \beta$. It is obvious that $V(\Phi)$ holds. It was proved in [8, Theorem 1.1] that $\text{HK}(\Psi)$ and $\text{HHK}(\Phi, \Psi)$ hold. In a general setting of fractional diffusions introduced in [5, Section 3], it was proved in [1, Theorem 3.7, Lemma 3.4] that $\text{wBE}(\Phi, \Psi)$ and $\text{HHK}(\Phi, \Psi)$ hold. It was proved in [8, Theorem 8.3] that all functions in the domain of the generator of the Dirichlet form are $(\beta - \alpha)$ -Hölder continuous. By the main result, Theorem 2.1, we now have the Hölder regularity for harmonic functions as $\text{HR}(\Phi, \Psi)$ and the Hölder estimate $\text{HHK}_{\text{exp}}(\Phi, \Psi)$ with exponential terms for the heat kernel.

Secondly, we consider the Sierpiński carpet cable system. Let $V_0 = \{p_1, \dots, p_8\}$ and $V_{n+1} = \cup_{i=1}^8 g_i(V_n)$ for any $n \geq 0$, then $\{V_n\}_{n \geq 0}$ is an increasing sequence of finite subsets of K and the closure of $\cup_{n \geq 0} V_n$ with respect to the Euclidean topology is K . For any $n \geq 0$, let $V^{(n)} = 3^n V_n$, see Figure 4 for $V^{(0)}$, $V^{(1)}$ and $V^{(2)}$.




Figure 4: $V^{(0)}$, $V^{(1)}$ and $V^{(2)}$

Obviously, $\{V^{(n)}\}_{n \geq 0}$ is an increasing sequence of finite sets. Let $V = \cup_{n \geq 0} V^{(n)}$ and

$$E = \left\{ \{p, q\} : p, q \in V, |p - q| = \frac{1}{2} \right\},$$

then (V, E) is an infinite, locally bounded, connected (undirected) graph, whose corresponding cable system is the Sierpiński carpet cable system, see [25, Section 3] for more detailed construction. Let $\alpha_1 = 1$, $\beta_1 = 2$, $\alpha_2 = \alpha$, $\beta_2 = \beta$. It is obvious that $V(\Phi)$ holds. Using the stability result [10] for heat kernel estimates or parabolic Harnack inequalities, we have $\text{HK}(\Psi)$ also holds, which is similar to the case on the unbounded Sierpiński carpet. By the main result, Theorem 2.1, we have $\text{HHK}(\Phi, \Psi)$, $\text{HHK}_{\text{exp}}(\Phi, \Psi)$, $\text{wBE}(\Phi, \Psi)$ and $\text{HR}(\Phi, \Psi)$ all hold. In particular, the validity of $\text{HR}(\Phi, \Psi)$ implies the validity of $\text{GRH}(\Phi, \Psi)$, which

was left open in [25]. Moreover, we also have $\text{GHK}(\Phi, \Psi)$, which can also be given directly by $\text{HHK}_{\text{exp}}(\Phi, \Psi)$.

Thirdly, we consider the blowups of fractals related to the Sierpiński carpet and the Vicsek set. However, we blowup a fractal by another different fractal. We consider the setting in \mathbb{R}^2 .

Let K, M be one of the following fractals in \mathbb{R}^2 : the unit interval $[0, 1] \times \{0\}$, the Vicsek set and the Sierpiński carpet. Let α_1 and β_1 be the Hausdorff dimension and walk dimension of K , α_2 and β_2 the Hausdorff dimension and walk dimension of M . Here, K serves as the “cell” fractal and M provides the “model” to blowup. Let $(\mathcal{E}^K, \mathcal{F}^K)$ be the strongly local regular Dirichlet form on $L^2(K; \mu)$ with two-sided sub-Gaussian heat kernel estimates with walk dimension β_1 , where μ is the normalized α_1 -Hausdorff measure on K , see [48, 42, 53] for the case K is the Vicsek set.

Let $\{g_i^{(1)}\}_{i=1, \dots, N^{(1)}}$, $\{g_i^{(2)}\}_{i=1, \dots, N^{(2)}}$ be the collections of contraction mappings generating K, M , respectively, that is, for $k = 1, 2$, $N^{(k)}, L^{(k)}$ are two positive integers, $p_1^{(k)} = (0, 0), \dots, p_{N^{(k)}}^{(k)} \in \mathbb{R}^2$,

$$g_i^{(k)}(x) = \frac{1}{L^{(k)}}x + \frac{L^{(k)} - 1}{L^{(k)}}p_i, x \in \mathbb{R}^2, i = 1, \dots, N^{(k)},$$

and K, M are the unique non-empty compact sets in \mathbb{R}^2 satisfying $K = \cup_{i=1}^{N^{(1)}} g_i^{(1)}(K)$, $M = \cup_{i=1}^{N^{(2)}} g_i^{(2)}(M)$, respectively.

- For the unit interval, we have $N^{(k)} = 2, L^{(k)} = 2$ and

$$p_1^{(k)} = (0, 0), p_2^{(k)} = (1, 0).$$

- For the Vicsek set, we have $N^{(k)} = 5, L^{(k)} = 3$ and

$$p_1^{(k)} = (0, 0), p_2^{(k)} = (1, 0), p_3^{(k)} = (1, 1), p_4^{(k)} = (0, 1), p_5^{(k)} = \left(\frac{1}{2}, \frac{1}{2}\right).$$

- For the Sierpiński carpet, we have $N^{(k)} = 8, L^{(k)} = 3$ and $p_1^{(k)}, \dots, p_8^{(k)}$ are p_1, \dots, p_8 from Equation (9.1).

We assume that for $i, j = 1, \dots, N^{(2)}$, if $g_i^{(2)}([0, 1]^2) \cap g_j^{(2)}([0, 1]^2) \neq \emptyset$, then $g_i^{(2)}(K) \cap g_j^{(2)}(K) \neq \emptyset$. Let $K_0 = K$. For any $n \geq 0$, let $K_{n+1} = \cup_{i=1}^{N^{(2)}} g_i^{(2)}(K_n)$, $K^{(n)} = (L^{(2)})^n K_n$, then

$$\begin{aligned} K^{(n+1)} &= (L^{(2)})^{n+1} K_{n+1} = (L^{(2)})^{n+1} \left(\cup_{i=1}^{N^{(2)}} g_i^{(2)}(K_n) \right) \\ &= (L^{(2)})^{n+1} \left(g_1^{(2)}(K_n) \cup \bigcup_{i=2}^{N^{(2)}} g_i^{(2)}(K_n) \right) \\ &= \left((L^{(2)})^n K_n \right) \cup \left((L^{(2)})^{n+1} \left(\bigcup_{i=2}^{N^{(2)}} g_i^{(2)}(K_n) \right) \right) \\ &\supseteq (L^{(2)})^n K_n = K^{(n)}, \end{aligned}$$

hence $\{K^{(n)}\}_{n \geq 0}$ is an increasing sequence of subsets of \mathbb{R}^2 .

Let $X = \cup_{n \geq 0} K^{(n)}$. We say that $W \subseteq X$ is a cell if W is a translation of $K^{(0)} = K_0 = K$. Let d be the Euclidean metric in \mathbb{R}^2 . Let m be the measure on X satisfying that for any cell W , $m|_W$ is the normalized α_1 -Hausdorff measure on W . For any cell W , by translation, there exists a strongly local regular Dirichlet form $(\mathcal{E}^W, \mathcal{F}^W)$ on $L^2(W; m|_W)$. Let

$$\mathcal{K} = \{u \in C_c(X) : u|_W \in \mathcal{F}^W \text{ for any cell } W\}.$$

Define

$$\mathcal{E}(u, u) = \sum_{W \text{ is a cell}} \mathcal{E}^W(u|_W, u|_W),$$

$$\mathcal{F} = \mathcal{E}_1\text{-closure of } \mathcal{K},$$

then $(\mathcal{E}, \mathcal{F})$ is a strongly local regular Dirichlet form on $L^2(X; m)$. Obviously, $(X, d, m, \mathcal{E}, \mathcal{F})$ is an unbounded MMD space satisfying $V(\Phi)$. Since two-sided heat kernel estimates hold on K, M , by [30, Theorem 3.14], we have the elliptic Harnack inequalities and the resistance estimates both hold on K, M . Combining these two conditions on K, M , we obtain easily the corresponding elliptic Harnack inequalities and the resistance estimates on X . Then by [30, Theorem 3.14] again, we have $\text{HK}(\Psi)$ holds on X . Indeed, in the strongly recurrent setting, only resistance estimates are enough for the above argument, see [12, 45, 37]. By the main result, Theorem 2.1, we have $\text{HHK}(\Phi, \Psi)$, $\text{HHK}_{\text{exp}}(\Phi, \Psi)$, $\text{wBE}(\Phi, \Psi)$ and $\text{HR}(\Phi, \Psi)$ all hold.

We give some examples as follows.

- (1) Let K be the Sierpiński carpet and M the unit interval, see Figure 5.




Figure 5: Blow up the Sierpiński carpet by the unit interval

- (2) Let K be the Sierpiński carpet and M the Vicsek set, see Figure 6 for $K^{(2)}$.




Figure 6: Blow up the Sierpiński carpet by the Vicsek set

- (3) Let K be the Vicsek set and M the Sierpiński carpet, see Figure 7 for $K^{(2)}$.

Since $(\Psi/\Phi)(r) = r$ for $r \in (0, 1)$, $\text{HHK}_{\text{exp}}(\Phi, \Psi)$ gives

$$\begin{aligned} & |p_t(x, y_1) - p_t(x, y_2)| \\ & \leq C_1 \frac{d(y_1, y_2)}{t} (\exp(-\Upsilon(C_2 d(x, y_1), t)) + \exp(-\Upsilon(C_2 d(x, y_2), t))) \end{aligned} \quad (9.2)$$

for any $x, y_1, y_2 \in X$ with $d(y_1, y_2) < 1$.

Let $S_0 = [(0, 0), (1, 1)] \cup [(0, 1), (1, 0)]$ be the union of the diagonals of the unit square $[0, 1]^2 \subseteq \mathbb{R}^2$. For any $n \geq 0$, let $S_{n+1} = \cup_{i=1}^{N^{(1)}} g_i^{(1)}(S_n)$. Then $\{S_n\}_{n \geq 0}$ is an increasing sequence of subsets of K and the closure of $\cup_{n \geq 0} S_n$ is K . Let $\mathcal{S}_0 = \cup_{n \geq 0} S_n$, $\nu^{(0)}$ the measure on \mathcal{S}_0 such that for any $n \geq 0$, $\nu^{(0)}|_{S_n}$ is the normalized 1-dimensional Hausdorff measure satisfying

$$\nu^{(0)}([(0, 0), (1, 1)]) = \nu^{(0)}([(0, 1), (1, 0)]) = 1.$$




Figure 7: Blow up the Vicsek set by the Sierpiński carpet

For any $n \geq 0$, let $\mathcal{S}_{n+1} = \cup_{i=1}^{N^{(2)}} g_i^{(2)}(\mathcal{S}_n)$, $\mathcal{S}^{(n)} = (L^{(2)})^n \mathcal{S}_n$, then similar to $\{K^{(n)}\}_{n \geq 0}$, $\{\mathcal{S}^{(n)}\}_{n \geq 0}$ is an increasing sequence of subsets of \mathbb{R}^2 , let $\mathcal{S} = \cup_{n \geq 0} \mathcal{S}^{(n)}$. For any cell $W \subseteq X$, we have $\mathcal{S}_W := \mathcal{S} \cap W$ is a translation of $\mathcal{S}^{(0)} = \mathcal{S}_0$. For any two cells $W_1, W_2 \subseteq X$, we have either the cardinal $\#(\mathcal{S}_{W_1} \cap \mathcal{S}_{W_2}) \in \{0, 1\}$ or $\mathcal{S}_{W_1} \cap \mathcal{S}_{W_2}$ is countably infinite, hence there exists a unique ν on \mathcal{S} such that for any cell W , $\nu|_{\mathcal{S}_W}$ is the push forward of $\nu^{(0)}$ under the translation from $\mathcal{S}^{(0)} = \mathcal{S}_0$ to \mathcal{S}_W . Note that $\nu(B(x_0), r) = +\infty$ for any ball $B(x_0, r)$ and $\nu \perp m$, that is, ν and m are mutually singular.

By the Lipschitz estimate, Equation (9.2), we have the weak gradient $\partial p_t(x, \cdot)$ exists ν -a.e. for any $t \in (0, +\infty)$, for any $x \in X$, and has the following estimate. For any $t \in (0, +\infty)$, for any $x \in X$, for ν -a.e. $y \in \mathcal{S}$, we have

$$|\partial_y p_t(x, y)| \leq \frac{C_1}{t} \exp(-\Upsilon(C_2 d(x, y), t)) \leq \begin{cases} \frac{C_3}{t} \exp\left(-C_4 \left(\frac{d(x, y)}{t^{1/\beta_1}}\right)^{\frac{\beta_1}{\beta_1-1}}\right), & \text{if } t < d(x, y), \\ \frac{C_3}{t} \exp\left(-C_4 \left(\frac{d(x, y)}{t^{1/\beta_2}}\right)^{\frac{\beta_2}{\beta_2-1}}\right), & \text{if } t \geq d(x, y). \end{cases} \quad (9.3)$$

Note that $\partial p_t(x, \cdot)$ exists and has the above estimate ν -a.e. instead of m -a.e.. At small scale, $\beta_1 = \log 15 / \log 3$, the gradient estimate has the sub-Gaussian behavior corresponding to the Vicsek set, while at large scale, $\beta_2 \approx 2.09697$, the gradient estimate has the sub-Gaussian behavior corresponding to the Sierpiński carpet. See [15, Section 3] for the results about Lipschitz and gradient estimates for heat kernel on unbounded Vicsek set, that is, the blowup of the Vicsek set by the Vicsek set itself.

- (4) Let K be the Vicsek set and M the unit interval, see Figure 8.




Figure 8: Blow up the Vicsek set by the unit interval

Similar to the blowup of the Vicsek set by the Sierpiński carpet, we also have the Lipschitz estimate, Equation (9.2) and the gradient estimate, Equation (9.3). However,

at small scale, $\beta_1 = \log 15 / \log 3$, the gradient estimate has the sub-Gaussian behavior corresponding to the Vicsek set, while at large scale, $\beta_2 = 2$, the gradient estimate has the Gaussian behavior corresponding to the unit interval.

References

- [1] Patricia Alonso-Ruiz, Fabrice Baudoin, Li Chen, Luke Rogers, Nageswari Shanmugalingam, and Alexander Teplyaev. Besov class via heat semigroup on Dirichlet spaces III: BV functions and sub-Gaussian heat kernel estimates. *Calc. Var. Partial Differential Equations*, 60(5):Paper No. 170, 38, 2021.
- [2] Sebastian Andres and Martin T. Barlow. Energy inequalities for cutoff functions and some applications. *J. Reine Angew. Math.*, 699:183–215, 2015.
- [3] Pascal Auscher, Thierry Coulhon, Xuan Thinh Duong, and Steve Hofmann. Riesz transform on manifolds and heat kernel regularity. *Ann. Sci. École Norm. Sup. (4)*, 37(6):911–957, 2004.
- [4] Richárd Balka, Zoltán Buczolich, and Márton Elekes. A new fractal dimension: the topological Hausdorff dimension. *Adv. Math.*, 274:881–927, 2015.
- [5] Martin T. Barlow. Diffusions on fractals. In *Lectures on probability theory and statistics (Saint-Flour, 1995)*, volume 1690 of *Lecture Notes in Math.*, pages 1–121. Springer, Berlin, 1998.
- [6] Martin T. Barlow and Richard F. Bass. The construction of Brownian motion on the Sierpiński carpet. *Ann. Inst. H. Poincaré Probab. Statist.*, 25(3):225–257, 1989.
- [7] Martin T. Barlow and Richard F. Bass. On the resistance of the Sierpiński carpet. *Proc. Roy. Soc. London Ser. A*, 431(1882):345–360, 1990.
- [8] Martin T. Barlow and Richard F. Bass. Transition densities for Brownian motion on the Sierpiński carpet. *Probab. Theory Related Fields*, 91(3-4):307–330, 1992.
- [9] Martin T. Barlow and Richard F. Bass. Brownian motion and harmonic analysis on Sierpinski carpets. *Canad. J. Math.*, 51(4):673–744, 1999.
- [10] Martin T. Barlow, Richard F. Bass, and Takashi Kumagai. Stability of parabolic Harnack inequalities on metric measure spaces. *J. Math. Soc. Japan*, 58(2):485–519, 2006.
- [11] Martin T. Barlow, Richard F. Bass, and John D. Sherwood. Resistance and spectral dimension of Sierpiński carpets. *J. Phys. A*, 23(6):L253–L258, 1990.
- [12] Martin T. Barlow, Thierry Coulhon, and Takashi Kumagai. Characterization of sub-Gaussian heat kernel estimates on strongly recurrent graphs. *Comm. Pure Appl. Math.*, 58(12):1642–1677, 2005.
- [13] Martin T. Barlow, Alexander Grigor’yan, and Takashi Kumagai. On the equivalence of parabolic Harnack inequalities and heat kernel estimates. *J. Math. Soc. Japan*, 64(4):1091–1146, 2012.
- [14] Martin T. Barlow and Edwin A. Perkins. Brownian motion on the Sierpiński gasket. *Probab. Theory Related Fields*, 79(4):543–623, 1988.
- [15] Fabrice Baudoin and Li Chen. Heat kernel gradient estimates for the Vicsek set. *arXiv e-prints*, page arXiv:2311.03499, November 2023.
- [16] Fabrice Baudoin and Nicola Garofalo. A note on the boundedness of Riesz transform for some subelliptic operators. *Int. Math. Res. Not. IMRN*, 2013(2):398–421, 2013.
- [17] Shiping Cao and Hua Qiu. Dirichlet forms on unconstrained Sierpinski carpets. *Probab. Theory Related Fields*, 189(1-2):613–657, 2024.
- [18] Eric A. Carlen, Shigeo Kusuoka, and Daniel W. Stroock. Upper bounds for symmetric Markov transition functions. *Ann. Inst. H. Poincaré Probab. Statist.*, 23(2, suppl.):245–287, 1987.
- [19] Jeff Cheeger and Shing Tung Yau. A lower bound for the heat kernel. *Comm. Pure Appl. Math.*, 34(4):465–480, 1981.

- [20] Li Chen, Thierry Coulhon, Joseph Feneuil, and Emmanuel Russ. Riesz transform for $1 \leq p \leq 2$ without Gaussian heat kernel bound. *J. Geom. Anal.*, 27(2):1489–1514, 2017.
- [21] Thierry Coulhon. Off-diagonal heat kernel lower bounds without Poincaré. *J. London Math. Soc. (2)*, 68(3):795–816, 2003.
- [22] Thierry Coulhon, Renjin Jiang, Pekka Koskela, and Adam Sikora. Gradient estimates for heat kernels and harmonic functions. *J. Funct. Anal.*, 278(8):108398, 67, 2020.
- [23] E. Brian Davies. Non-Gaussian aspects of heat kernel behaviour. *J. London Math. Soc. (2)*, 55(1):105–125, 1997.
- [24] Ennio De Giorgi. Sulla differenziabilità e l’analiticità delle estremali degli integrali multipli regolari. *Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat. (3)*, 3:25–43, 1957.
- [25] Baptiste Devyver, Emmanuel Russ, and Meng Yang. Gradient Estimate for the Heat Kernel on Some Fractal-Like Cable Systems and Quasi-Riesz Transforms. *International Mathematics Research Notices*, 2023(18):15537–15583, September 2023.
- [26] Masatoshi Fukushima, Yoichi Oshima, and Masayoshi Takeda. *Dirichlet forms and symmetric Markov processes*, volume 19 of *De Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, extended edition, 2011.
- [27] Alexander Grigor’yan. The heat equation on noncompact Riemannian manifolds. *Mat. Sb.*, 182(1):55–87, 1991.
- [28] Alexander Grigor’yan. Heat kernel upper bounds on a complete non-compact manifold. *Rev. Mat. Iberoamericana*, 10(2):395–452, 1994.
- [29] Alexander Grigor’yan, Eryan Hu, and Jiaxin Hu. Parabolic mean value inequality and on-diagonal upper bound of the heat kernel on doubling spaces. *Math. Ann.*, 389(3):2411–2467, 2024.
- [30] Alexander Grigor’yan and Jiaxin Hu. Heat kernels and Green functions on metric measure spaces. *Canad. J. Math.*, 66(3):641–699, 2014.
- [31] Alexander Grigor’yan, Jiaxin Hu, and Ka-Sing Lau. Generalized capacity, Harnack inequality and heat kernels of Dirichlet forms on metric measure spaces. *J. Math. Soc. Japan*, 67(4):1485–1549, 2015.
- [32] Alexander Grigor’yan and Andras Telcs. Two-sided estimates of heat kernels on metric measure spaces. *Ann. Probab.*, 40(3):1212–1284, 2012.
- [33] Alexander Grigor’yan and Meng Yang. Local and non-local Dirichlet forms on the Sierpiński carpet. *Trans. Amer. Math. Soc.*, 372(6):3985–4030, 2019.
- [34] Ben M. Hambly and Takashi Kumagai. Transition density estimates for diffusion processes on post critically finite self-similar fractals. *Proc. London Math. Soc. (3)*, 78(2):431–458, 1999.
- [35] Ben M. Hambly, Takashi Kumagai, Shigeo Kusuoka, and Xian Yin Zhou. Transition density estimates for diffusion processes on homogeneous random Sierpinski carpets. *J. Math. Soc. Japan*, 52(2):373–408, 2000.
- [36] Waldemar Hebisch and Laurent Saloff-Coste. On the relation between elliptic and parabolic Harnack inequalities. *Ann. Inst. Fourier (Grenoble)*, 51(5):1437–1481, 2001.
- [37] Jiaxin Hu. An analytical approach to heat kernel estimates on strongly recurrent metric spaces. *Proc. Edinb. Math. Soc. (2)*, 51(1):171–199, 2008.
- [38] Renjin Jiang. The Li-Yau inequality and heat kernels on metric measure spaces. *J. Math. Pures Appl. (9)*, 104(1):29–57, 2015.
- [39] Renjin Jiang, Pekka Koskela, and Dachun Yang. Isoperimetric inequality via Lipschitz regularity of Cheeger-harmonic functions. *J. Math. Pures Appl. (9)*, 101(5):583–598, 2014.
- [40] Naotaka Kajino and Ryosuke Shimizu. Contraction properties and differentiability of p -energy forms with applications to nonlinear potential theory on self-similar sets. *arXiv e-prints*, page arXiv:2404.13668, April 2024.
- [41] Jun Kigami. A harmonic calculus on the Sierpiński spaces. *Japan J. Appl. Math.*, 6(2):259–290, 1989.

- [42] Jun Kigami. *Analysis on fractals*, volume 143 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2001.
- [43] Jun Kigami. Volume doubling measures and heat kernel estimates on self-similar sets. *Mem. Amer. Math. Soc.*, 199(932):viii+94, 2009.
- [44] Jun Kigami. Resistance forms, quasisymmetric maps and heat kernel estimates. *Mem. Amer. Math. Soc.*, 216(1015):vi+132, 2012.
- [45] Takashi Kumagai. Heat kernel estimates and parabolic Harnack inequalities on graphs and resistance forms. *Publ. Res. Inst. Math. Sci.*, 40(3):793–818, 2004.
- [46] Shigeo Kusuoka and Xianyin Zhou. Dirichlet forms on fractals: Poincaré constant and resistance. *Probab. Theory Related Fields*, 93(2):169–196, 1992.
- [47] Peter Li and Shing-Tung Yau. On the parabolic kernel of the Schrödinger operator. *Acta Math.*, 156(3-4):153–201, 1986.
- [48] Tom L. Lindstrøm. Brownian motion on nested fractals. *Mem. Amer. Math. Soc.*, 83(420):iv+128, 1990.
- [49] Jürgen Moser. On Harnack’s theorem for elliptic differential equations. *Comm. Pure Appl. Math.*, 14:577–591, 1961.
- [50] Laurent Saloff-Coste. A note on Poincaré, Sobolev, and Harnack inequalities. *Internat. Math. Res. Notices*, 1992(2):27–38, 1992.
- [51] Laurent Saloff-Coste. Parabolic Harnack inequality for divergence-form second-order differential operators. *Potential Anal.*, 4(4):429–467, 1995. Potential theory and degenerate partial differential operators (Parma).
- [52] Robert S. Strichartz. Taylor approximations on Sierpinski gasket type fractals. *J. Funct. Anal.*, 174(1):76–127, 2000.
- [53] Robert S. Strichartz. *Differential equations on fractals*. Princeton University Press, Princeton, NJ, 2006. A tutorial.
- [54] Donglei Tang, Rui Hu, and Chunwei Pan. Hölder estimate of harmonic functions on a class of p.c.f. self-similar sets. *Anal. Theory Appl.*, 30(3):296–305, 2014.

DEPARTMENT OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY,
HANGZHOU 310036, CHINA
E-mail address: gaojin@hznu.edu.cn

INSTITUT FÜR ANGEWANDTE MATHEMATIK, UNIVERSITÄT BONN, ENDENICHER ALLEE 60,
53115 BONN, GERMANY
E-mail address: yangmengqh@gmail.com