

ON THE BOUNDEDNESS OF CERTAIN OPERATORS OF CONVOLUTION TYPE ON FOCK SPACES

SUNDARAM THANGAVELU

ABSTRACT. We find necessary and sufficient conditions on the convolution kernels φ so that certain operators on twisted Fock spaces $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ are bounded.

1. INTRODUCTION

In this article we are concerned with the boundedness of certain operators of convolution type on the Fock spaces $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ indexed by $\lambda \in \mathbb{R}$. These spaces are weighted Bergman spaces consisting of entire functions $F(\zeta)$ on \mathbb{C}^{2n} that are square integrable with respect to certain positive weight functions $w_\lambda(\zeta)$. Given $F, \varphi \in \mathcal{F}^\lambda(\mathbb{C}^{2n})$ we define the convolution

$$(1.1) \quad F *_\lambda \varphi(\zeta) = \int_{\mathbb{C}^{2n}} F(\zeta') \varphi(\zeta - \bar{\zeta}') \overline{K_\lambda(\zeta, \zeta')} w_\lambda(\zeta') d\zeta'$$

where $K_\lambda(\zeta, \zeta')$ is the reproducing kernel associated to $\mathcal{F}^\lambda(\mathbb{C}^{2n})$. The above integral need not converge for all $F, \varphi \in \mathcal{F}^\lambda(\mathbb{C}^{2n})$ but when φ is fixed, the convolution makes sense for all F from a dense class of functions in $\mathcal{F}^\lambda(\mathbb{C}^{2n})$. Thus we have a densely defined linear operator $S_\varphi^\lambda F = F *_\lambda \varphi$ whose boundedness on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ received considerable attention recently, see the following papers [4, 3, 2, 6].

In these works the authors have identified a subspace $\mathcal{A}^\lambda(\mathbb{C}^{2n}) \subset \mathcal{F}^\lambda(\mathbb{C}^{2n})$ with the property that S_φ^λ , initially defined on a dense subspace of $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ extends to the whole space as a bounded linear operator if and only if $\varphi \in \mathcal{A}^\lambda(\mathbb{C}^{2n})$. These subspaces turn out to be the image of certain Banach algebras \mathbf{A} under a map G_λ . Thus S_φ^λ is bounded on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ if and only if $\varphi = G_\lambda(\mathbf{a})$ for some $\mathbf{a} \in \mathbf{A}$. It has also been shown that there is a unitary map \mathcal{G}_λ setting up a one to one correspondence between a Hilbert space \mathcal{H}_λ and $\mathcal{F}^\lambda(\mathbb{C}^{2n})$. This map has the property that $\mathcal{G}_\lambda^*(F *_\lambda \varphi) = \mathcal{G}_\lambda^*(F) \mathbf{a}$ if $\varphi = G_\lambda(\mathbf{a})$ which justifies the definition of the convolution.

Though this characterisation is interesting in itself, there is no easy way to determine when a given $\varphi \in \mathcal{F}^\lambda(\mathbb{C}^{2n})$ belongs to the subspace $\mathcal{A}^\lambda(\mathbb{C}^{2n})$. It is desirable to have a necessary and sufficient condition on φ which are easy to verify, e.g. certain pointwise estimates/integrability conditions. In this article we address this problem and come up with a partial solution. More precisely, for each $0 < t \leq 1/2$ we introduce the subspaces

$$(1.2) \quad \mathcal{A}_{1/2}^\lambda(\mathbb{C}^{2n}) \subset \mathcal{A}^\lambda(\mathbb{C}^{2n}) \subset \mathcal{A}_t^\lambda(\mathbb{C}^{2n})$$

leading to the necessary condition, $\varphi \in \mathcal{A}_t^\lambda(\mathbb{C}^{2n})$ for all $0 < t < 1/2$ and a slightly stronger sufficient condition, $\varphi \in \mathcal{A}_{1/2}^\lambda(\mathbb{C}^{2n})$ for the operator S_φ^λ to be bounded on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$. All these

spaces $\mathcal{A}_t^\lambda(\mathbb{C}^{2n})$ are weighted Bergman spaces and hence their membership is defined in terms of integrability conditions/pointwise estimates.

We now give some details. Let us consider the case $\lambda = 0$ first which corresponds to analysis on the classical Fock space $\mathcal{F}^0(\mathbb{C}^{2n})$. These spaces are defined for any dimension and hence we consider $\mathcal{F}^0(\mathbb{C}^n) = \mathcal{F}(\mathbb{C}^n)$ consisting of entire functions F that are square integrable with respect to the Gaussian measure $w_0(z) = c_0 e^{-\frac{1}{2}|z|^2}$. For this case, we delete all subscripts and superscripts, e.g we simply write S_φ instead of S_φ^0 and so on. Boundedness of the operators

$$(1.3) \quad S_\varphi(z) = \int_{\mathbb{C}^n} F(w)\varphi(z - \bar{w}) e^{\frac{1}{2}z \cdot \bar{w}} w_0(w) dw$$

have been studied in [4]. Let $d\gamma(\xi) = e^{-|\xi|^2} d\xi$ and consider the Gauss-Bargmann transform

$$(1.4) \quad G : L^2(\mathbb{R}^n, d\gamma) \rightarrow \mathcal{F}(\mathbb{C}^n), \quad Gf(z) = e^{\frac{1}{4}z^2} \int_{\mathbb{R}^n} f(\xi) e^{iz \cdot \xi} d\gamma$$

which is unitary. By taking $\mathbf{A} = L^\infty(\mathbb{R}^n)$ and $\mathcal{A}(\mathbb{C}^n) = G(\mathbf{A})$ it has been proved that S_φ is bounded on $\mathcal{F}(\mathbb{C}^n)$ if and only if $\varphi = G(m)$ for some $m \in L^\infty(\mathbb{R}^n)$.

It is easy to see that the space $\mathcal{A}(\mathbb{C}^n)$ is invariant under all translations in the imaginary direction. That is for any $\varphi \in \mathcal{A}(\mathbb{C}^n)$ and $a \in \mathbb{R}^n$, the function $\tau(ia)\varphi(z) = \varphi(z + ia)$ belongs to $\mathcal{A}(\mathbb{C}^n)$. This leads a very simple necessary condition for S_φ to be bounded. A slight strengthening of the same condition also turns out to be sufficient.

Theorem 1.1. *Suppose $\varphi \in \mathcal{F}(\mathbb{C}^n)$ is such that S_φ is bounded. Then φ satisfies*

$$(i) \sup_{a \in \mathbb{R}^n} \int_{\mathbb{C}^n} |\varphi(z + ia)|^2 e^{-\frac{1}{2}|z|^2} dz < \infty, \quad (ii) \sup_{y \in \mathbb{R}^n} |\varphi(x + iy)| e^{-\frac{1}{4}|x|^2} < \infty.$$

Conversely, if we assume that φ satisfies (i) and the slightly stronger condition

$$\sup_{y \in \mathbb{R}^n} \int_{\mathbb{R}^n} |\varphi(x + iy)| e^{-\frac{1}{4}|x|^2} dx < \infty,$$

then $\varphi = Gm$ for some $m \in L^\infty(\mathbb{R}^n)$ and consequently S_φ is bounded.

Another necessary condition can be obtained by considering the weighted Bergman space $\mathcal{A}_t(\mathbb{C}^n)$, $t \geq 0$ defined by weight

$$w_t(x, y) = e^{-\frac{1}{2}(|x|^2 + |y|^2)} e^{\frac{2t}{(1+2t)}|y|^2}.$$

By considering a variant of the Gauss-Bargmann transform we can show that $\mathcal{A}(\mathbb{C}^n) \subset \mathcal{A}_t(\mathbb{C}^n)$ for $0 \leq t < 1/2$ leading to the following necessary condition.

Theorem 1.2. *Given $\varphi \in \mathcal{F}(\mathbb{C}^n)$, a necessary condition for S_φ to be bounded on $\mathcal{F}(\mathbb{C}^n)$ is that for $0 \leq t < 1/2$*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |\varphi(z)|^2 w_t(x, y) dx dy < \infty.$$

Let us consider the Hilbert space \mathcal{H} consisting of all tempered distributions f for which $f * q_{1/2} \in L^2(\mathbb{R}^n)$ where

$$q_t(x) = (4\pi t)^{-n/2} e^{-\frac{1}{4t}|x|^2}$$

is the heat kernel associated to the standard Laplacian Δ on \mathbb{R}^n . We equip \mathcal{H} with the obvious norm : $\|f\| = \|f * q_{1/2}\|_2$. Note that the Fourier transform $f \rightarrow \widehat{f}$ embeds \mathcal{H} isometrically into a subspace of $L^2(\mathbb{R}^n, d\gamma)$. For $f \in \mathcal{H}$ we define $\mathcal{G}(f) = G(\widehat{f})$ so that the image of \mathcal{H} under \mathcal{G} is a subspace $\mathcal{F}_0(\mathbb{C}^n)$ of the Fock space $\mathcal{F}(\mathbb{C}^n)$. Instead of G we work with \mathcal{G} and note that $\varphi = \mathcal{G}(f)$ defines a bounded operator S_φ if and only if $\widehat{f} \in L^\infty(\mathbb{R}^n)$. The usefulness of \mathcal{G} stems from the fact that it allows us to define the spaces $\mathcal{A}_t(\mathbb{C}^n)$ mentioned above, see Section 2.

We now consider the case $\lambda \neq 0$. The spaces $\mathcal{F}^\lambda(\mathbb{C}^{2n})$, known as twisted Fock spaces, are weighted Bergman spaces corresponding to the weight function

$$w_\lambda(\zeta) = c_\lambda e^{-\frac{1}{2}\lambda(\coth \lambda)|\zeta|^2} e^{\lambda[\operatorname{Re} \zeta, \operatorname{Im} \zeta]}$$

where $[\cdot, \cdot]$ is the symplectic form on \mathbb{R}^{2n} defined by $[(x, u), (y, v)] = (u \cdot y - v \cdot x)$. The reproducing kernel for this space is given by

$$K_\lambda(\zeta, \zeta') = e^{\frac{1}{2}\lambda(\coth \lambda)\zeta' \cdot \bar{\zeta}} e^{-\frac{i}{2}\lambda[\zeta', \bar{\zeta}]}$$

where $[(z, w), (a, b)] = (w \cdot a - z \cdot b)$ is the natural extension of the symplectic form to \mathbb{C}^{2n} . On $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ we consider operators of the form

$$(1.5) \quad S_\varphi^\lambda F(\zeta) = \int_{\mathbb{C}^{2n}} F(\zeta') \varphi(\zeta - \bar{\zeta}') e^{\frac{1}{2}\lambda(\coth \lambda)(\zeta \cdot \bar{\zeta}')} e^{-\frac{i}{2}\lambda[\zeta, \bar{\zeta}']} w_\lambda(\zeta') d\zeta'.$$

In a recent joint work with Rahul Garg [6] we have obtained a necessary and sufficient condition on φ so that S_φ^λ is bounded. The condition is in terms of a non-commutative analogue G_λ of the Gauss-Bargmann transform G .

In order to describe the known results we need to set up some notation. More details will be given in later sections. We let $\mathbf{A} = B(L^2(\mathbb{R}^n))$, the Banach algebra of all bounded linear operators on $L^2(\mathbb{R}^n)$. We let $e^{-tH(\lambda)}$, $t > 0$ stand for the Hermite semigroup generated by the scaled Hermite operator $H(\lambda) = -\Delta + \lambda^2|x|^2$ on \mathbb{R}^n . For $(x, u) \in \mathbb{R}^{2n}$, let $\pi_\lambda(x, u)$ be the Schrödinger representation of the Heisenberg group \mathbb{H}^n realised on $L^2(\mathbb{R}^n)$. Then it is known that $\pi_\lambda(z, w)$ can be defined as densely defined operators for $(z, w) \in \mathbb{C}^{2n}$. Finally we let

$$(1.6) \quad p_t^\lambda(x, u) = c_n \lambda^n (\sinh t\lambda)^{-n} e^{-\frac{1}{4}\lambda(\coth t\lambda)(|x|^2 + |u|^2)}$$

which also has a holomorphic extension $p_t^\lambda(z, w)$ to \mathbb{C}^{2n} . With these notations we define the non-commutative Gauss-Bargmann transform $G_\lambda : \mathbf{A} \rightarrow \mathcal{F}^\lambda(\mathbb{C}^{2n})$ by

$$(1.7) \quad G_\lambda(M)(z, w) = p_1^\lambda(z, w)^{-1} \operatorname{tr} \left(\pi_\lambda(-z, -w) e^{-\frac{1}{2}H(\lambda)} M e^{-\frac{1}{2}H(\lambda)} \right).$$

Let $\mathcal{A}^\lambda(\mathbb{C}^{2n})$ denote the image of $B(L^2(\mathbb{R}^n))$ under the Gauss-Bargmann transform. In [6], we have proved that the operator S_φ^λ is bounded on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ if and only if $\varphi \in \mathcal{A}^\lambda(\mathbb{C}^{2n})$. In other words, S_φ^λ is bounded if and only if $\varphi = G_\lambda(M)$ for some $M \in B(L^2(\mathbb{R}^n))$. However, given $\varphi \in \mathcal{F}^\lambda(\mathbb{C}^{2n})$ it is not easy to determine when it belongs to $\mathcal{A}^\lambda(\mathbb{C}^{2n})$.

For each $0 < t \leq 1/2$, let the weight function $w_t^\lambda(\xi, \eta)$ be defined on $\mathbb{R}^{2n} \times \mathbb{R}^{2n}$ by

$$(1.8) \quad w_t^\lambda(\xi, \eta) = e^{-\frac{\lambda}{2}(\coth \lambda)(|\xi|^2 + |\eta|^2)} e^{\lambda(\tanh 2t\lambda)|\xi|^2} e^{\lambda(\coth \lambda - \coth 2t\lambda)|\eta|^2}.$$

We then define $\mathcal{A}_t^\lambda(\mathbb{C}^{2n})$ as the weighted Bergman space corresponding to $w_t^\lambda(\xi, \eta)$. In other words, $\mathcal{A}_t^\lambda(\mathbb{C}^{2n})$ is the Hilbert space of all entire functions F on \mathbb{C}^{2n} for which

$$\int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} |F(\xi + i\eta)|^2 w_t^\lambda(\xi, \eta) d\xi d\eta < \infty.$$

Then it can be shown that (see Section 3.5) $\mathcal{A}_{1/2}^\lambda(\mathbb{C}^{2n}) \subset \mathcal{A}^\lambda(\mathbb{C}^{2n}) \subset \mathcal{A}_t^\lambda(\mathbb{C}^{2n})$ for $0 < t < 1/2$. In view of the result of [6] mentioned above this gives a necessary condition and also a sufficient condition on φ so that S_φ^λ is bounded.

Theorem 1.3. *Given $\varphi \in \mathcal{F}^\lambda(\mathbb{C}^{2n})$, a necessary condition for S_φ^λ to be bounded on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ is that for all $0 < t < 1/2$*

$$\int_{\mathbb{R}^{2n}} \int_{\mathbb{R}^{2n}} |\varphi(\xi + i\eta)|^2 w_t^\lambda(\xi, \eta) d\xi d\eta < \infty.$$

Conversely, if the above integral is finite for $t = 1/2$, then S_φ^λ is bounded on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$.

This theorem will be proved in Section 3.5. The proof depends on a curious relation between Hermite and special Hermite semigroups and the Gutzmer's formula for the Hermite expansions. In the course of the proof we will show that $\mathcal{A}_{1/2}^\lambda(\mathbb{C}^{2n})$ is precisely the image of \mathcal{S}_2 , the space of Hilbert-Schmidt operators on $L^2(\mathbb{R}^n)$, under the Gauss-Bargmann transform G_λ .

Thus if $\varphi \in \mathcal{A}_{1/2}^\lambda(\mathbb{C}^{2n})$ then there exists a Hilbert-Schmidt operator T on $L^2(\mathbb{R}^n)$ so that

$$\varphi(z, w) = p_1^\lambda(z, w)^{-1} \operatorname{tr} \left(\pi_\lambda(-z, -w) e^{-\frac{1}{2}H(\lambda)} T e^{-\frac{1}{2}H(\lambda)} \right).$$

As $T = \pi_\lambda(f)$ for a unique $f \in L^2(\mathbb{R}^{2n})$, by appealing to the inversion formula for the Weyl transform we get

$$\varphi(z, w) = p_1^\lambda(z, w)^{-1} p_{1/2}^\lambda *_\lambda f *_\lambda p_{1/2}^\lambda(z, w).$$

This suggests that we consider the semigroup $T_t f(x, u) = p_t^\lambda *_\lambda f *_\lambda p_t^\lambda(x, u)$ which turns out to be none other than the Hermite semigroup generated by $H(\lambda)$ on \mathbb{R}^{2n} . Thus the function $p_1^\lambda \varphi$ belongs to the image of $L^2(\mathbb{R}^{2n})$ under the Segal-Bargmann transform taking f into the holomorphic function $e^{-\frac{1}{2}H(\lambda)} f(z, w)$. The image of $L^2(\mathbb{R}^{2n})$ under the semigroup $e^{-tH(1)}$ is known to be a weighted Bergman space corresponding to the weight function

$$U_t(z, w) = c_n (\sinh 4t)^{-2n} e^{-(\coth 2t)(|y|^2 + |v|^2) + (\tanh 2t)(|x|^2 + |u|^2)}.$$

Therefore, it follows that φ is square integrable with respect to $|p_1^\lambda(z, w)|^2 U_{\lambda/2}(\sqrt{\lambda}(z, w))$ which after simplification reduces to $w_{1/2}^\lambda(\xi, \eta)$ proving the sufficiency part of the theorem.

We conclude this introduction with a brief description of the plan of this paper. In the next section we deal with operators S_φ on the classical Fock space $\mathcal{F}(\mathbb{C}^n)$ which corresponds to the case $\lambda = 0$. As one can see the proofs are quite easy in this case. In Section 3 we consider the operators S_φ^λ on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ in which case the proof requires quite a bit of preparation. After recalling the required preliminary material the main result is proved in Section 3.5.

2. THE OPERATORS S_φ ON THE FOCK SPACES $\mathcal{F}(\mathbb{C}^n)$

In this section we prove the results on operators of the form S_φ stated in the introduction. We recall that these operators studied by several authors [4, 3, 2] are defined by

$$S_\varphi F(z) = \int_{\mathbb{C}^n} F(w) \varphi(z - \bar{w}) e^{\frac{1}{2}z \cdot \bar{w}} e^{-\frac{1}{2}|w|^2} dw$$

where $\varphi \in \mathcal{F}(\mathbb{C}^n)$. As stated in the introduction, S_φ is bounded on the Fock space if and only if $\varphi = Gm$ for some $m \in L^\infty(\mathbb{R}^n)$. Here G is the Gauss-Bargmann transform defined by the equation (1.4). The proofs of Theorems 1.1 and 1.2 are rather easy which we present now.

Proof of Theorem 1.1 As mentioned above, the operator S_φ is bounded on $\mathcal{F}(\mathbb{C}^n)$ if and only if

$$\varphi(z) = e^{\frac{1}{4}z^2} \int_{\mathbb{R}^n} m(\xi) e^{-|\xi|^2} e^{iz \cdot \xi} d\xi$$

for some $m \in L^\infty(\mathbb{R}^n)$. A simple calculation shows that

$$\varphi(z + ia) = e^{\frac{1}{4}z^2} \int_{\mathbb{R}^n} m(\xi - \frac{1}{2}a) e^{-|\xi|^2} e^{iz \cdot \xi} d\xi$$

for any $a \in \mathbb{R}^n$. In particular,

$$\varphi(x + ia) = e^{\frac{1}{4}|x|^2} \int_{\mathbb{R}^n} m(\xi - \frac{1}{2}a) e^{-|\xi|^2} e^{ix \cdot \xi} d\xi.$$

As m is bounded we get two necessary conditions on φ : for any $a \in \mathbb{R}^n$

$$\int_{\mathbb{C}^n} |\varphi(z + ia)|^2 e^{-\frac{1}{2}|z|^2} dz \leq C, \quad |\varphi(x + ia)| e^{-\frac{1}{4}|x|^2} \leq C$$

where C is independent of a . Conversely, if we assume that φ satisfies the first of the above conditions, then for some $m_a \in L^2(\mathbb{R}^n, d\gamma)$ we have

$$\varphi(z + ia) = e^{\frac{1}{4}z^2} \int_{\mathbb{R}^n} m_a(\xi) e^{-|\xi|^2} e^{iz \cdot \xi} d\xi.$$

Since $\varphi(z) = Gm(z)$, we have the equality

$$e^{\frac{1}{4}x^2} \int_{\mathbb{R}^n} m_a(\xi) e^{-|\xi|^2} e^{ix \cdot \xi} d\xi = e^{\frac{1}{4}(x+ia)^2} \int_{\mathbb{R}^n} m(\xi) e^{-|\xi|^2} e^{i(x+ia) \cdot \xi} d\xi$$

which after a change of variables in the right hand side integral yields

$$\int_{\mathbb{R}^n} m_a(\xi) e^{-|\xi|^2} e^{ix \cdot \xi} d\xi = \int_{\mathbb{R}^n} m(\xi - \frac{1}{2}a) e^{-|\xi|^2} e^{ix \cdot \xi} d\xi.$$

Hence it follows that $m_a(\xi) = m(\xi - \frac{1}{2}a)$ and consequently φ satisfies the equation

$$\varphi(x + ia) = e^{\frac{1}{4}|x|^2} \int_{\mathbb{R}^n} m(\xi - \frac{1}{2}a) e^{-|\xi|^2} e^{ix \cdot \xi} d\xi.$$

If we impose the stronger condition on φ , namely

$$\int_{\mathbb{R}^n} |\varphi(x + ia)| e^{-\frac{1}{2}|x|^2} dx \leq C$$

for all $a \in \mathbb{R}^n$, then by Fourier inversion formula

$$m(\xi - \frac{1}{2}a) e^{-|\xi|^2} = \int_{\mathbb{R}^n} \varphi(x + ia) e^{-\frac{1}{4}|x|^2} e^{-ix \cdot \xi} dx.$$

Therefore, by choosing $\xi = 0$ we can conclude that m is bounded proving the theorem.

Remark 2.1. Note that $L^\infty(\mathbb{R}^n)$ is a translation invariant subspace of $L^2(\mathbb{R}^n, d\gamma)$ whose image under G is $\mathcal{A}(\mathbb{C}^n)$. Let $L_0^2(\mathbb{R}^n, d\gamma)$ be any translation invariant subspace which contains $L^\infty(\mathbb{R}^n)$. As shown above, if $\varphi = Gm$ with $m \in L_0^2(\mathbb{R}^n, d\gamma)$, then $\varphi_a(z) = \varphi(z + ia)$ corresponds to $m_a(\xi) = m(\xi - a/2)$.

Remark 2.2. Since $G : L^2(\mathbb{R}^n, d\gamma) \rightarrow \mathcal{F}(\mathbb{C}^n)$ is unitary we can define an action of \mathbb{C}^n on $L^2(\mathbb{R}^n, d\gamma)$ by transferring the unitary operator $\rho(w)$ which acts on the Fock space as

$$\rho(w)\varphi(z) = e^{-\frac{1}{4}|w|^2} \varphi(z + w) e^{-\frac{1}{2}z \cdot \bar{w}}.$$

A simple calculation shows that

$$\rho(u + iv)\varphi(z) = e^{-\frac{1}{2}(|v|^2 - iu \cdot v)} e^{iz \cdot v} e^{\frac{1}{4}z^2} \int_{\mathbb{R}^n} m(\xi) e^{iw \cdot \xi} e^{-|\xi|^2} e^{iz \cdot \xi} d\xi.$$

Making the change of variables $\xi \rightarrow \xi - v$ in the above integral and simplifying

$$\rho(u + iv)\varphi(z) = e^{-\frac{1}{2}|v|^2} e^{\frac{1}{4}z^2} \int_{\mathbb{R}^n} m(\xi - v) e^{i\bar{w} \cdot \xi} e^{-|\xi|^2} e^{iz \cdot \xi} d\xi.$$

Thus $\rho(w)G(m) = G(\pi(w)m)$ where $\pi(w)m(\xi) = e^{-\frac{1}{2}|v|^2} m(\xi - v) e^{i\xi \cdot \bar{w}}$.

We can get a different necessary condition for S_φ to be bounded by looking at another version of the Gauss-Bargmann transform. We let \mathcal{H} stand for the space of all tempered distributions f such that $f * q_{1/2}$ is in $L^2(\mathbb{R}^n)$ equipped with the norm $\|f\| = \|f * q_{1/2}\|_2$. Then it is clear that $f \in \mathcal{H}$ if and only if $\hat{f} \in L^2(\mathbb{R}^n, d\gamma)$. Hence $f \rightarrow G(\hat{f})$ is a unitary map between \mathcal{H} and $\mathcal{F}(\mathbb{C}^n)$. If $\varphi = G(f)$ and S_φ is bounded, then $\hat{f} \in L^\infty(\mathbb{R}^n)$. We can now prove Theorem 1.2.

Proof of Theorem 1.2 If φ is as above, then from the definition of G we see that

$$\varphi(z) e^{-\frac{1}{4}z^2} = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{-|\xi|^2} e^{iz \cdot \xi} d\xi.$$

By Plancherel, this immediately gives us the identity

$$\int_{\mathbb{R}^n} |\varphi(x + iy)|^2 e^{-\frac{1}{2}(|x|^2 - |y|^2)} dx = c_n \int_{\mathbb{R}^n} |\hat{f}(\xi)|^2 e^{-2|\xi|^2} e^{-2y \cdot \xi} d\xi.$$

Integrating the above with respect to $q_{t/2}(y)dy$ we obtain

$$(2\pi t)^{-n/2} \int_{\mathbb{C}^n} |\varphi(z)|^2 e^{-\frac{1}{2}|z|^2} e^{-\frac{(1-2t)}{2t}|y|^2} dx dy = c_n \int_{\mathbb{R}^{2n}} |\hat{f}(\xi)|^2 e^{-2(1-t)|\xi|^2} d\xi.$$

Under the assumption that φ is bounded, we have $\hat{f} \in L^\infty(\mathbb{R}^n)$ and hence from the above we get the necessary condition

$$\int_{\mathbb{C}^n} |\varphi(z)|^2 e^{-\frac{1}{2}|z|^2} e^{-\frac{(1-2t)}{2t}|y|^2} dx dy \leq C_t$$

for all $0 < t < 1$. Replacing $2t$ by $1 + 2t$ we can write the above as

$$\int_{\mathbb{C}^n} |\varphi(z)|^2 e^{-\frac{1}{2}|z|^2} e^{\frac{2t}{1+2t}|y|^2} dx dy \leq C_t$$

for $0 \leq t < 1/2$. This completes the proof of the theorem.

3. OPERATORS S_φ^λ ON THE TWISTED FOCK SPACES $\mathcal{F}^\lambda(\mathbb{C}^{2n})$.

In this section we consider the operators S_φ^λ on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ defined by the convolution

$$(3.1) \quad S_\varphi^\lambda F(\zeta) = \int_{\mathbb{C}^{2n}} F(\zeta') \varphi(\zeta - \bar{\zeta}') e^{\frac{1}{2}\lambda(\coth \lambda)(\zeta \cdot \bar{\zeta}')} e^{-\frac{1}{2}\lambda[\zeta, \bar{\zeta}']} w_\lambda(\zeta') d\zeta'.$$

With G_λ defined in (1.7) a necessary and sufficient condition on φ so that S_φ^λ is bounded on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ is that $\varphi = G_\lambda(M)$ for some bounded linear operator M on $L^2(\mathbb{R}^n)$. We are interested in finding certain integrability conditions on φ which does not involve the operator M . In order to do so we make use of the connection between Hermite and special Hermite operators. Recall that the definition of G_λ involves the Hermite as well as the special Hermite semigroup. We will be very sketchy in recalling the basic facts about these semigroups referring to the standard literature for more details, see e.g. [8, 9]. In what follows we assume $\lambda \neq 0$.

3.1. Heisenberg group and the Schrödinger representations. We begin by recalling some well known facts about the Heisenberg groups \mathbb{H}^n and their representations. The standard references for this section are [5, 9]. The group \mathbb{H}^n is just $\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}$ (or equivalently $\mathbb{C}^n \times \mathbb{R}$) equipped with the group law

$$(x, y, t)(u, v, s) = (x + u, y + v, t + s + \frac{1}{2}(u \cdot y - x \cdot v)).$$

For each $\lambda \in \mathbb{R}, \lambda \neq 0$, there is an irreducible unitary representation π_λ of \mathbb{H}^n on $L^2(\mathbb{R}^n)$ explicitly given by

$$\pi_\lambda(x, y, t)f(\xi) = e^{i\lambda t} e^{i\lambda(x \cdot \xi + \frac{1}{2}x \cdot y)} f(\xi + y), \quad f \in L^2(\mathbb{R}^n).$$

By setting $\pi_\lambda(x, y) = \pi_\lambda(x, y, 0)$ we observe that the irreducibility of π_λ has the following consequence: if a bounded linear operator T on $L^2(\mathbb{R}^n)$ commutes with $\pi_\lambda(x, y)$ for all $x, y \in \mathbb{R}^n$ then $T = cI$ for a constant c . However, there are non-trivial operators that commute either with the family $\pi_\lambda(x, 0)$ or with $\pi_\lambda(0, y)$.

The Fourier transform of a function $f \in L^1 \cap L^2(\mathbb{H}^n)$ is defined to be the operator valued function

$$\widehat{f}(\lambda) = \int_{\mathbb{H}^n} f(x, y, t) \pi_\lambda(x, y, t) dx dy dt = \int_{\mathbb{R}^{2n}} f^\lambda(x, y) \pi_\lambda(x, y) dx dy$$

where $f^\lambda(x, y)$ is the inverse Fourier transform of $f(x, y, t)$ in the central variable:

$$f^\lambda(x, y) = \int_{-\infty}^{\infty} f(x, y, t) e^{i\lambda t} dt.$$

Thus we see that $\widehat{f}(\lambda) = \pi_\lambda(f^\lambda)$ where for any $g \in L^1 \cap L^2(\mathbb{R}^{2n})$ the operator

$$\pi_\lambda(g) = \int_{\mathbb{R}^{2n}} g(x, y) \pi_\lambda(x, y) dx dy$$

is known as the Weyl transform of g . It is known that $\pi_\lambda : g \rightarrow \pi_\lambda(g)$ is a constant multiple of a unitary operator from $L^2(\mathbb{R}^{2n})$ onto the space \mathcal{S}_2 consisting of all Hilbert-Schmidt operators on $L^2(\mathbb{R}^n)$. The convolution between two functions f and g on \mathbb{H}^n is defined by

$$f * g(x, y, t) = \int_{\mathbb{H}^n} f((x, y, t)(u, v, s)^{-1}) g(u, v, s) du dv ds.$$

This gives rise to a family of convolution structures on $L^1(\mathbb{R}^{2n})$ defined by the relation $(f * g)^\lambda(x, y) = f^\lambda *_\lambda g^\lambda(x, y)$. These are known as λ -twisted convolutions and we make use of the relation $\pi_\lambda(f^\lambda *_\lambda g^\lambda) = \pi_\lambda(f^\lambda)\pi_\lambda(g^\lambda)$ in this article.

3.2. Hermite and special Hermite semigroups. The spectral decomposition of the Hermite operator $H(\lambda) = -\Delta + \lambda^2|x|^2$ is given by

$$H(\lambda) = \sum_{k=0}^{\infty} (2k + n)|\lambda| P_k(\lambda)$$

where the spectral projection $P_k(\lambda)$ onto the k -th eigenspace associated to the eigenvalue $(2k + n)|\lambda|$ is given by

$$P_k(\lambda)f = \sum_{|\alpha|=k} (f, \Phi_\alpha^\lambda) \Phi_\alpha^\lambda.$$

Here Φ_α^λ , $\alpha \in \mathbb{N}^n$ are the normraised, scaled Hermite functions which are eigenfunctions of $H(\lambda)$ with eigenvalues $(2|\alpha| + n)|\lambda|$. It is well known that they form an orthonormal basis for $L^2(\mathbb{R}^n)$. The Hermite semigroup is defined on $L^p(\mathbb{R}^n)$ by

$$e^{-tH(\lambda)} f = \sum_{k=0}^{\infty} e^{-t(2k+n)|\lambda|} P_k(\lambda) f.$$

The semigroup $e^{-tH(\lambda)}$ is an integral operator with an explicit kernel from which it follows that it is a contraction on $L^p(\mathbb{R}^n)$ for any $1 \leq p \leq \infty$.

The special Hermite operator denoted by L_λ is the operator on \mathbb{R}^{2n} given by

$$L_\lambda = -\Delta_{\mathbb{R}^{2n}} + \frac{\lambda^2}{4}(|x|^2 + |u|^2) + i\lambda \sum_{j=1}^n \left(x_j \frac{\partial}{\partial u_j} - u_j \frac{\partial}{\partial x_j} \right).$$

There is an orthonormal basis for $L^2(\mathbb{R}^{2n})$ consisting of joint eigenfunctions of L_λ and $H(\lambda)$ on \mathbb{R}^{2n} . These are defined by

$$\Phi_{\alpha\beta}^\lambda(x, u) = (2\pi)^{-n/2} |\lambda|^{n/2} (\pi_\lambda(x, u) \Phi_\alpha^\lambda, \Phi_\beta^\lambda).$$

$$(3.2) \quad L_\lambda \Phi_{\alpha\beta}^\lambda = (2|\beta| + n)|\lambda| \Phi_{\alpha\beta}^\lambda, \quad H(\lambda) \Phi_{\alpha\beta}^\lambda = (2|\alpha| + 2|\beta| + 2n)|\lambda| \Phi_{\alpha\beta}^\lambda.$$

The special Hermite functions satisfy another interesting orthogonality relation under twisted convolution (see [9]):

$$(3.3) \quad \Phi_{\alpha\beta}^\lambda *_\lambda \Phi_{\mu\nu}^\lambda = \delta_{\beta\mu} \Phi_{\alpha\nu}^\lambda.$$

The spectrum of L_λ consists of $(2k + n)|\lambda|$, $k \in \mathbb{N}$ but in view of the above the eigenspaces are infinite dimensional. However, the spectral decomposition of L_λ has a compact form in terms of Laguerre functions $\varphi_k^\lambda(x, u)$.

The λ -twisted convolution between two functions f and g on \mathbb{R}^{2n} defined earlier is explicitly given by the integral

$$(3.4) \quad f *_\lambda g(x, u) = \int_{\mathbb{R}^{2n}} f(a, b) g(x - a, u - b) e^{-i\frac{\lambda}{2}(u \cdot a - x \cdot b)} da db.$$

With this notation, the spectral decomposition of L_λ is given explicitly by

$$L_\lambda f(x, u) = (2\pi)^{-n} |\lambda|^n \sum_{\alpha \in \mathbb{N}^n} (2|\alpha| + n) |\lambda| f *_\lambda \Phi_{\alpha\alpha}^\lambda(x, u)$$

which can be further simplified. It is known that

$$\sum_{|\alpha|=k} \Phi_{\alpha\alpha}^\lambda(x, u) = \varphi_{k,\lambda}^{n-1}(x, u) = L_k^{n-1}\left(\frac{1}{2}|\lambda|(|x|^2 + |u|^2)\right) e^{-\frac{1}{4}|\lambda|(|x|^2 + |u|^2)}$$

where $L_k^{n-1}(t), t \geq 0$ stand for the Laguerre polynomials of type $(n-1)$. In terms of these functions the spectral decomposition of L_λ can be written as

$$L_\lambda f(x, u) = (2\pi)^{-n} |\lambda|^n \sum_{k=0}^{\infty} (2k + n) |\lambda| f *_\lambda \varphi_{k,\lambda}^{n-1}(x, u).$$

Consequently, the special Hermite semigroup generated by L_λ is given by the expansion

$$e^{-tL_\lambda} f(x, u) = (2\pi)^{-n} |\lambda|^n \sum_{k=0}^{\infty} e^{-t(2k+n)|\lambda|} f *_\lambda \varphi_{k,\lambda}^{n-1}(x, u).$$

From the above definition it is clear that $e^{-tL_\lambda} f = f *_\lambda p_t^\lambda$ where the kernel p_t^λ is given by the expansion

$$(2\pi)^{-n} |\lambda|^n \sum_{\alpha \in \mathbb{N}^n} e^{-t(2|\alpha|+n)|\lambda|} \Phi_{\alpha\alpha}^\lambda(x, u) = (2\pi)^{-n} |\lambda|^n \sum_{k=0}^{\infty} e^{-t(2k+n)|\lambda|} \varphi_{k,\lambda}^{n-1}(x, u).$$

The above series can be summed and p_t^λ is given explicitly by the formula

$$p_t^\lambda(x, u) = (4\pi)^{-n} \lambda^n (\sinh t\lambda)^{-n} e^{-\frac{1}{4}\lambda(\coth t\lambda)(|x|^2 + |u|^2)}.$$

3.3. Connection between Hermite and special Hermite semigroups. The relation between the Hermite semigroup $e^{-tH(\lambda)}$ on $L^2(\mathbb{R}^n)$ and the special Hermite semigroup e^{-tL_λ} is given by the Weyl transform: $e^{-tH(\lambda)} = \pi_\lambda(p_t^\lambda)$. This is a well known relation but there is yet another curious relation between Hermite and special Hermite semigroups both acting $L^2(\mathbb{R}^{2n})$ which plays an important role in the proof of our main theorem. In the next proposition $H(\lambda)$ stands for the Hermite operator on \mathbb{R}^{2n} .

Proposition 3.1. *For any $t > 0$ we have $e^{-tH(\lambda)} f = p_t^\lambda *_\lambda f *_\lambda p_t^\lambda$ for all $f \in L^2(\mathbb{R}^{2n})$.*

Proof. As the special Hermite functions $\Phi_{\alpha\beta}^\lambda$ form an orthonormal basis for $L^2(\mathbb{R}^{2n})$ it enough to check that

$$e^{-tH(\lambda)} \Phi_{\alpha\beta}^\lambda = p_t^\lambda *_\lambda \Phi_{\alpha\beta}^\lambda *_\lambda p_t^\lambda$$

for all α and β . Since p_t^λ is given by the expansion

$$p_t^\lambda(x, u) = \sum_{\mu \in \mathbb{N}^n} e^{-t(2|\mu|+n)|\lambda|} \Phi_{\mu\mu}^\lambda(x, u),$$

using the orthogonality relations (3.3) we get

$$p_t^\lambda *_\lambda \Phi_{\alpha\beta}^\lambda = e^{-t(2|\alpha|+n)|\lambda|} \Phi_{\alpha\beta}^\lambda, \quad \Phi_{\alpha\beta}^\lambda *_\lambda p_t^\lambda = e^{-t(2|\beta|+n)|\lambda|} \Phi_{\alpha\beta}^\lambda.$$

Consequently,

$$p_t^\lambda *_\lambda \Phi_{\alpha\beta}^\lambda *_\lambda p_t^\lambda = e^{-t(2|\alpha|+2|\beta|+2n)|\lambda|} \Phi_{\alpha\beta}^\lambda.$$

As $\Phi_{\alpha\beta}^\lambda$ are eigenfunctions of $H(\lambda)$ with eigenvalues $(2|\alpha| + 2|\beta| + 2n)|\lambda|$ we also have

$$e^{-tH(\lambda)} \Phi_{\alpha\beta}^\lambda = e^{-t(2|\alpha|+2|\beta|+2n)|\lambda|} \Phi_{\alpha\beta}^\lambda.$$

This completes the proof of the proposition. \square

The above result allows us to extend the domain of definition of the Hermite semigroup from $L^2(\mathbb{R}^{2n})$ to a bigger space. Let $L^2(\mathbb{R}^{2n}, p_t^\lambda)$ stand for the space of all tempered distributions f for which $p_t^\lambda *_\lambda f \in L^2(\mathbb{R}^{2n})$. For such functions

$$e^{-tH(\lambda)} f = p_t^\lambda *_\lambda f *_\lambda p_t^\lambda$$

is well defined and extends to \mathbb{C}^{2n} as an entire function.

3.4. On the image of $L^2(\mathbb{R}^{2n})$ under Hermite and special Hermite semigroups. For any $f \in L^2(\mathbb{R}^{2n})$ the functions $e^{-tH(\lambda)} f$ and e^{-tL_λ} both extend to \mathbb{C}^{2n} as entire functions. Thus the image of $L^2(\mathbb{R}^{2n})$ under either of these semigroups consist of entire functions which are square integrable with respect to certain explicit positive weight functions. For the case of the Hermite semigroup the weight function is given by

$$(3.5) \quad U_t^\lambda(z, w) = c_n (\sinh 4t\lambda)^{-2n} e^{-\lambda(\coth 2t\lambda)(|y|^2+|v|^2)+\lambda(\tanh 2t\lambda)(|x|^2+|u|^2)}.$$

Thus for any $f \in L^2(\mathbb{R}^{2n})$ the function $F(z, w) = e^{-tH(\lambda)} f(z, w)$ satisfies the identity

$$(3.6) \quad \int_{\mathbb{C}^n} \int_{\mathbb{C}^n} |F(z, w)|^2 U_t^\lambda(z, w) dz dw = c_{n,\lambda} \int_{\mathbb{R}^{2n}} |f(x, u)|^2 dx du.$$

We have stated the above result for $H(\lambda)$ on \mathbb{R}^{2n} but the result is true in any dimension. This was first proved by Byun [1] but there are many other proofs available in the literature, see [11]. The proof given in [11] relies on the so called Gutzmer's formula for the Hermite expansions which also plays an important role in getting a necessary condition on φ so that S_φ^λ is bounded on the twisted Fock space.

In order to state Gutzmer's formula we need to set up some notation. We make use of several facts about the Schrödinger representation π_λ of the Heisenberg group \mathbb{H}^n . The operator $\pi_\lambda(x, u)$ can also be defined for complex values of the arguments. Thus for $z, w \in \mathbb{C}^n$, $\pi_\lambda(z, w)$ can be defined on $L^2(\mathbb{R}^n)$ as densely defined unbounded operators. If we let $U(n)$ denote the group of $n \times n$ unitary matrices, then for any $\sigma \in U(n)$, the map $(z, t) \rightarrow (\sigma z, t)$ is an automorphism of \mathbb{H}^n . The Laguerre function $\varphi_{k,\lambda}^{n-1}(x, u)$ can also be extended to $\mathbb{C}^n \times \mathbb{C}^n$ as an entire function. With these notations we have

Theorem 3.2 (Gutzmer). *Let $f \in L^2(\mathbb{R}^n)$ has a holomorphic extension F to \mathbb{C}^n . Then we have*

$$\int_{U(n)} \int_{\mathbb{R}^n} |\pi_\lambda(i\sigma(y, v))F(\xi)|^2 d\xi d\sigma = c_n \sum_{k=0}^{\infty} \frac{k!(n-1)!}{(k+n-1)!} \varphi_{k,\lambda}^{n-1}(2iy, 2iv) \|P_k(\lambda)f\|_2^2.$$

We refer to [11] for a proof of this result. This formula along with the following lemma leads to a proof of the identity (3.6) in the case of $e^{-tH(\lambda)}$ on $L^2(\mathbb{R}^n)$.

Lemma 3.3. *For any $t > 0$ we have*

$$\frac{k!(n-1)!}{(k+n-1)!} \int_{\mathbb{R}^{2n}} p_{2t}^\lambda(2y, 2v) \varphi_{k,\lambda}^{n-1}(2iy, 2iv) dy dv = c_\lambda e^{2t(2k+n)|\lambda|}.$$

It is interesting to compare the above with the following identity which is easy to verify:

$$\frac{k!(n-1)!}{(k+n-1)!} \int_{\mathbb{R}^{2n}} p_{2t}^\lambda(2y, 2v) \varphi_{k,\lambda}^{n-1}(2y, 2v) dy dv = c_\lambda e^{-2t(2k+n)|\lambda|}$$

In fact, this is an immediate consequence of the expansion of p_t^λ in terms of Laguerre functions $\varphi_{k,\lambda}^{n-1}$ and their orthogonality in $L^2(\mathbb{R}^{2n})$.

In the case of the special Hermite semigroup we have a similar result. For $f \in L^2(\mathbb{R}^{2n})$ the function $F(x, u) = e^{-tL_\lambda} f(x, u)$ has a holomorphic extension to \mathbb{C}^{2n} and we have the identity

$$(3.7) \quad \int_{\mathbb{C}^n} \int_{\mathbb{C}^n} |F(z, w)|^2 W_t^\lambda(z, w) dz dw = c_{\lambda,n} \int_{\mathbb{R}^{2n}} |f(x, u)|^2 dx du$$

where the weight function $W_t^\lambda(z, w)$ is given by

$$W_t^\lambda(z, w) = e^{\lambda \operatorname{Im}(z \cdot \bar{w})} p_{2t}^\lambda(2y, 2v), \quad z = x + iy, w = u + iv.$$

Thus the image of $L^2(\mathbb{R}^{2n})$ under the special Hermite semigroup is precisely the weighted Bergman space consisting of entire functions on \mathbb{C}^{2n} for which (3.7) is finite. This was first proved in [7] but there is another proof using Gutzmer's formula for special Hermite expansion. We refer the reader to the paper [10] for details. We conclude this subsection with the following remarks.

Remark 3.4. Note that in view of Proposition 3.1 and (3.7), for any $f \in L^2(\mathbb{R}^{2n}, p_t^\lambda)$ we have the identity

$$\int_{\mathbb{C}^{2n}} |e^{-tH(\lambda)} f(z, w)|^2 W_t^\lambda(z, w) dz dw = \int_{\mathbb{R}^{2n}} |p_t^\lambda *_\lambda f(x, u)|^2 dx du.$$

As both $H(\lambda)$ and p_t^λ are even functions of λ , recalling the definition of W_t^λ and using the fact that $p_t^\lambda *_{-\lambda} f = f *_\lambda p_t^\lambda$ we have

$$\int_{\mathbb{C}^{2n}} |e^{-tH(\lambda)} f(z, w)|^2 e^{-\lambda \operatorname{Im}(z \cdot \bar{w})} p_{2t}^\lambda(2y, 2v) dz dw = \int_{\mathbb{R}^{2n}} |f *_\lambda p_t^\lambda(x, u)|^2 dx du.$$

Using the identity (3.7) once more we can rewrite the right hand side in terms of $f *_\lambda p_{2t}^\lambda$ leading to the interesting relation

$$\int_{\mathbb{C}^{2n}} |e^{-tH(\lambda)} f(z, w)|^2 W_t^{-\lambda}(z, w) dz dw = \int_{\mathbb{C}^{2n}} |e^{-2tL_\lambda} f(z, w)|^2 W_{2t}^\lambda(z, w) dz dw.$$

Remark 3.5. Let $\mathcal{S}_2(\Gamma_\lambda)$ be the image of $L^2(\mathbb{R}^{2n}, p_t^\lambda)$ under the Weyl transform. In other words, a densely defined linear operator T on $L^2(\mathbb{R}^n)$ belongs to $\mathcal{S}_2(\Gamma_\lambda)$ if $e^{-\frac{1}{2}H(\lambda)} T$ is Hilbert-Schmidt. Note that $B(L^2(\mathbb{R}^n)) \subset \mathcal{S}_2(\Gamma_\lambda)$ and G_λ initially defined on $B(L^2(\mathbb{R}^n))$ has a natural extension to $\mathcal{S}_2(\Gamma_\lambda)$. Thus if $T = \pi_\lambda(f)$ we have

$$G_\lambda(T)(z, w) = p_1^\lambda(z, w)^{-1} e^{-\frac{1}{2}H(\lambda)} f(z, w).$$

We denote the image of $\mathcal{S}_2(\Gamma_\lambda)$ under G_λ by $\mathcal{F}_0^\lambda(\mathbb{C}^{2n})$.

Remark 3.6. When $T = \pi_\lambda(f) \in B(L^2(\mathbb{R}^n))$ we have the estimate

$$\int_{\mathbb{C}^{2n}} |e^{-\frac{1}{2}H(\lambda)} f(z, w)|^2 W_t^\lambda(z, w) dzdw = \|Te^{-\frac{1}{2}H(\lambda)}\|_{HS}^2 \leq C\|T\|_{op}^2.$$

In particular, when $f \in L^p(\mathbb{R}^{2n}), 1 \leq p \leq 2, \|\pi_\lambda(f)\|_{op} \leq \|f\|_p$ and hence we have

$$\int_{\mathbb{C}^{2n}} |e^{-tH(\lambda)} f(z, w)|^2 W_t^\lambda(z, w) dzdw \leq C\|f\|_p^2.$$

3.5. Necessary and sufficient conditions for S_φ^λ to be bounded. From the above discussions, it is clear that a necessary and sufficient condition for S_φ^λ to be bounded on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ is that $\varphi(z, w) = p_1^\lambda(z, w)^{-1} e^{-\frac{1}{2}H(\lambda)} f(z, w)$ with $\pi_\lambda(f) \in B(L^2(\mathbb{R}^n))$. This in turn is equivalent to the statement that the operator $T_f : g \rightarrow g *_\lambda f$ initially defined on $\mathcal{S}(\mathbb{R}^{2n})$ extends to $L^2(\mathbb{R}^{2n})$ as a bounded operator if and only if $\pi_\lambda(f) \in B(L^2(\mathbb{R}^n))$. By using Gutzmer's formula for Hermite expansions, we can get a necessary condition on φ . In what follows we write $U_t^\lambda(\xi, \eta)$ in place of $U_t^\lambda(z, w)$ where $(z, w) = \zeta = \xi + i\eta$.

Proposition 3.7. *For $f \in L^2(\mathbb{R}^{2n}, p_{1/2}^\lambda)$ let $\varphi = G_\lambda(\pi_\lambda(f))$ and define $F = p_1^\lambda \varphi$. Then for any $0 < t < 1/2$, we have the identity*

$$\int_{\mathbb{R}^{4n}} |F(\xi + i\eta)|^2 U_t^\lambda(\xi, \eta) d\xi d\eta = c_n \sum_{k=0}^{\infty} \sum_{|\alpha|+|\beta|=k} |(\pi_\lambda(f) \Phi_\alpha^\lambda, \Phi_\beta^\lambda)|^2 e^{-(1-2t)(2k+2n)|\lambda|}.$$

Proof. Suppose $\varphi(z, w) = B_\lambda(p_{1/2}^\lambda *_\lambda f)(z, w)$ where $f \in L^2(\mathbb{R}^{2n}, p_{1/2}^\lambda)$. Then we have

$$F(z, w) = \varphi(z, w) p_1^\lambda(z, w) = e^{-\frac{1}{2}H(\lambda)} f(z, w).$$

Thus F is the holomorphic extension of the function $e^{-\frac{1}{2}H(\lambda)} f \in L^2(\mathbb{R}^{2n})$. By Gutzmer's formula stated in Theorem 3.2 we obtain

$$\begin{aligned} & \int_{U(2n)} \int_{\mathbb{R}^{2n}} |\pi_\lambda(i\sigma(\eta', \eta)) F(\xi)|^2 d\xi d\sigma \\ &= c_n \sum_{k=0}^{\infty} e^{-(2k+2n)|\lambda|} \frac{k!(2n-1)!}{(k+2n-1)!} \varphi_{k,\lambda}^{2n-1}(2i(\eta', \eta)) \|P_k(\lambda)f\|_2^2. \end{aligned}$$

As $H(\lambda)$ is a differential operator with real coefficients, we note that

$$H(\lambda) \bar{\Phi}_{\alpha\beta}^\lambda = (2|\alpha| + 2|\beta| + 2n) |\lambda| \bar{\Phi}_{\alpha\beta}^\lambda.$$

Hence $P_k(\lambda)f$ which is the projection of f into the k -th eigenspace corresponding to the eigenvalue $(2k+2n)|\lambda|$ is given by two expressions, viz.

$$P_k(\lambda)f = \sum_{|\alpha|+|\beta|=k} (f, \Phi_{\alpha\beta}^\lambda) \Phi_{\alpha\beta}^\lambda = \sum_{|\alpha|+|\beta|=k} (f, \bar{\Phi}_{\alpha\beta}^\lambda) \bar{\Phi}_{\alpha\beta}^\lambda.$$

We use the second expression to calculate $\|P_k(\lambda)f\|_2^2$. Recalling the definition of the special Hermite functions we see that

$$(f, \bar{\Phi}_{\alpha\beta}^\lambda) = (2\pi)^{-n/2} \int_{\mathbb{R}^{2n}} f(x, u) (\pi_\lambda(x, u) \Phi_\alpha^\lambda, \Phi_\beta^\lambda) dx du = (2\pi)^{-n/2} (\pi_\lambda(f) \Phi_\alpha^\lambda, \Phi_\beta^\lambda)$$

and hence we have

$$\|P_k(\lambda)f\|_2^2 = (2\pi)^{-n} \sum_{|\alpha|+|\beta|=k} |(\pi_\lambda(f)\Phi_\alpha^\lambda, \Phi_\beta^\lambda)|^2.$$

Therefore, we have proved the identity

$$\begin{aligned} & \int_{U(2n)} \int_{\mathbb{R}^{2n}} |\pi_\lambda(i\sigma(\eta', \eta))F(\xi)|^2 d\xi d\sigma \\ &= c_n \sum_{k=0}^{\infty} e^{-(2k+2n)|\lambda|} \frac{k!(2n-1)!}{(k+2n-1)!} \varphi_{k,\lambda}^{2n-1}(2i(\eta', \eta)) \sum_{|\alpha|+|\beta|=k} |(\pi_\lambda(f)\Phi_\alpha^\lambda, \Phi_\beta^\lambda)|^2. \end{aligned}$$

As $p_{2t}^\lambda(2(\eta', \eta))$ is invariant under the action of $U(2n)$ the above leads to the identity

$$\begin{aligned} & \int_{\mathbb{R}^{4n}} \left(\int_{\mathbb{R}^{2n}} |\pi_\lambda(i(\eta', \eta))F(\xi)|^2 d\xi \right) p_{2t}^\lambda(2(\eta', \eta)) d\eta d\eta' \\ &= \int_{\mathbb{R}^{2n}} \left(\int_{U(2n)} \int_{\mathbb{R}^{2n}} |\pi_\lambda(i\sigma(\eta', \eta))F(\xi)|^2 d\xi d\sigma \right) p_{2t}^\lambda(2(\eta', \eta)) d\eta d\eta'. \end{aligned}$$

The integral on the left can be simplified and easily seen to be

$$\int_{\mathbb{R}^{4n}} |F(\xi + i\eta)|^2 U_t^\lambda(\xi, \eta) d\xi d\eta$$

whereas in view of the identity proved above the integral on the right is given by

$$\sum_{k=0}^{\infty} \sum_{|\alpha|+|\beta|=k} |(\pi_\lambda(f)\Phi_\alpha^\lambda, \Phi_\beta^\lambda)|^2 c_{k,\lambda} e^{-(2k+2n)|\lambda|}$$

where we have written $c_{k,\lambda}$ for the constant defined by the following integral:

$$c_{k,\lambda} = \frac{k!(2n-1)!}{(k+2n-1)!} \left(\int_{\mathbb{R}^{4n}} \varphi_{k,\lambda}^{2n-1}(2i(\eta', \eta)) p_{2t}^\lambda(2(\eta', \eta)) d\eta d\eta' \right).$$

We now make use of the identity stated in Lemma 3.3 to conclude that

$$\int_{\mathbb{R}^{4n}} |F(\xi + i\eta)|^2 U_t^\lambda(\xi, \eta) d\xi d\eta = c_n \sum_{k=0}^{\infty} \sum_{|\alpha|+|\beta|=k} |(\pi_\lambda(f)\Phi_\alpha^\lambda, \Phi_\beta^\lambda)|^2 e^{-(1-2t)(2k+2n)|\lambda|}.$$

This completes the proof of the proposition. \square

Once we have the above proposition at our disposal, it is easy to get a necessary condition and a slightly stronger sufficient condition on φ so that S_φ^λ is bounded. With $F(z, w) = p_1^\lambda(z, w)\varphi(z, w)$ we can easily check that F is square integrable with respect to $U_t^\lambda(\xi, \eta)$ if and only if $\varphi(\xi + i\eta) = \varphi(z, w)$ is square integrable with respect to the weight function

$$U_t^\lambda(\xi, \eta) |p_1^\lambda(\xi + i\eta)|^2 = U_t^\lambda(\xi, \eta) e^{-\frac{\lambda}{2}(\coth \lambda)|\xi|^2} e^{\frac{\lambda}{2}(\coth \lambda)|\eta|^2}.$$

After simplification the above reduces to the weight function $w_t^\lambda(\xi, \eta)$ which appears in Theorem 1.3 and defined in (1.8). Therefore, Theorem 1.3 will follow once we prove the following result. Let us recall that $\mathcal{F}_0^\lambda(\mathbb{C}^{2n})$ stands for the image of $\mathcal{S}_2(\Gamma_\lambda)$ under G_λ .

Theorem 3.8. *Given $\varphi \in \mathcal{F}_0^\lambda(\mathbb{C}^{2n})$, let $F = p_1^\lambda \varphi$. A necessary condition for S_φ^λ to be bounded on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ is that*

$$\int_{\mathbb{R}^{4n}} |F(\xi + i\eta)|^2 U_t^\lambda(\xi, \eta) d\xi d\eta < \infty$$

for all $0 < t < 1/2$. Conversely, if the above condition holds for $t = 1/2$, then S_φ^λ is bounded.

Proof. Since $\varphi = G_\lambda(\pi_\lambda(f))$ for some $f \in L^2(\mathbb{R}^{2n}, p_{1/2}^\lambda)$ it follows that S_φ^λ is bounded on $\mathcal{F}^\lambda(\mathbb{C}^{2n})$ if and only if $\pi_\lambda(f)$ is bounded on $L^2(\mathbb{R}^n)$. When this happens

$$\sum_{|\alpha|+|\beta|=k} |(\pi_\lambda(f)\Phi_\alpha^\lambda, \Phi_\beta^\lambda)|^2 \leq C \frac{(k+2n-1)!}{k!(2n-1)!}$$

and consequently from the proposition proved above we get the estimate

$$\int_{\mathbb{R}^{4n}} |F(\xi + i\eta)|^2 U_t^\lambda(\xi, \eta) d\xi d\eta = C_n \sum_{k=0}^{\infty} \frac{(k+2n-1)!}{k!(2n-1)!} e^{-(1-2t)(2k+2n)|\lambda|}$$

which is finite as long as $0 < t < 1/2$. Conversely, if the condition holds for $t = 1/2$ we have

$$\sum_{k=0}^{\infty} \sum_{|\alpha|+|\beta|=k} |(\pi_\lambda(f)\Phi_\alpha^\lambda, \Phi_\beta^\lambda)|^2 < \infty.$$

But this simply means that $\pi_\lambda(f)$ is a Hilbert-Schmidt operator since

$$\|\pi_\lambda(f)\|_{HS}^2 = \sum_{\alpha \in \mathbb{N}^n} \|\pi_\lambda(f)\Phi_\alpha^\lambda\|_2^2 = \sum_{\alpha, \beta \in \mathbb{N}^n} |(\pi_\lambda(f)\Phi_\alpha^\lambda, \Phi_\beta^\lambda)|^2.$$

This completes the proof of the theorem and hence Theorem 1.3 is also proved. \square

ACKNOWLEDGEMENTS

This work was carried out when the author was visiting Harish-Chandra Research Institute as Infosys Visiting Professor. He thanks HRI for the facilities and warm hospitality. He was also partially supported by INSA.

REFERENCES

- [1] D.-W. Byun, *Inversions of Hermite semigroup*, Proc. Amer. Math. Soc. **118** (1993), 437–445..
- [2] S. Bais and V. N. Dogga, *\mathcal{L} -invariant radial singular integral operators on Fock spaces*, J. Pseudo-Differ. Oper. Appl. **14** (2023), no. 1, Paper No. 11, 35 pp.
- [3] G. Cao, L. He, J. Li, and M. Shen, *Boundedness criterion for integral operators on the fractional Fock-Sobolev spaces*, Math. Z. **301** (2022), no. 4, 3671–3693. MR 4449725
- [4] G. Cao, J. Li, M. Shen, B. D. Wick, and L. Yan, *A boundedness criterion for singular integral operators of convolution type on the Fock space*, Adv. Math. **363** (2020), 107001, 33. MR 4053517
- [5] G. B. Folland, *Harmonic analysis in phase space*, Annals of Mathematics Studies, vol. 122, Princeton University Press, Princeton, NJ, 1989. MR 983366
- [6] R. Garg and S. Thangavelu, *Boundedness of certain linear operators on twisted Fock spaces*, Math. Z. (2024) arXiv:2306.14188 (2023)
- [7] B. Krötz, S. Thangavelu, and Y. Xu, *The heat kernel transform for the Heisenberg group*, J. Funct. Anal. **225** (2005), no. 2, 301–336. MR 2152501
- [8] S. Thangavelu, *Lectures on Hermite and Laguerre expansions*, Mathematical Notes, vol. 42, Princeton University Press, Princeton, NJ, 1993, With a preface by Robert S. Strichartz. MR 1215939

- [9] S. Thangavelu, *An introduction to the uncertainty principle*, Progress in Mathematics, vol. 217, Birkhäuser Boston, Inc., Boston, MA, 2004, Hardy's theorem on Lie groups, With a foreword by Gerald B. Folland. MR 2008480
- [10] S. Thangavelu, *Gutzmer's formula and Poisson integrals on the Heisenberg group*, Pacific J. Math. **231** (2007), no.1, 217-237.
- [11] S. Thangavelu, *An analogue of Gutzmer's formula for Hermite expansions*, Studia Math. **185** (2008), 279-290.
- [12] S. Thangavelu, *Fourier multipliers and pseudo-differential operators on Fock-Sobolev spaces*, arXiv: 2304.01087 (2023).
- [13] S. Thangavelu, *Algebras of entire functions and representations of the twisted Heisenberg group*, Indian J. Pure and Appl. Math. (2024).
- [14] B. D. Wick and S. K. Wu, *Integral operators on Fock-Sobolev spaces via multipliers on Gauss-Sobolev spaces*, Integr. Equ. Oper. Theory **94** (2022), no. 2, Paper No. 22, 24 pp.

(S. Thangavelu) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE-560012, INDIA.

Email address: veluma@iisc.ac.in