

# THE DEGENERATION OF A FAMILY OF RATIONAL SURFACE AUTOMORPHISMS

QITONG JIANG

**ABSTRACT.** We consider a one-dimensional family of rational surfaces with automorphisms discussed in [7, 2, 3]. In a degeneration of this family, the limiting map is the identity map on a special fiber. We check that the map on the total space of the family has indeterminacy in the special fiber. However, we show that after blowing-up at an indeterminate curve, there is an induced birational map on the exceptional divisor over the indeterminate curve. Moreover, we show that this map has dynamical degree  $\lambda = 16$ .

## 0. INTRODUCTION

Given an automorphism  $f : X \rightarrow X$  of an algebraic variety  $X$ , if we blow-up  $X$  at a subvariety  $V$  with  $f(V) = V$ , then  $f$  induces an automorphism  $\tilde{f}|_E : E \rightarrow E$  of the exceptional divisor  $E$ . This map does not have any new dynamical properties, as it fibers over the original map  $f|_V : V \rightarrow V$ . On the other hand, if  $f(V) = V' \neq V$ , then the lift  $\tilde{f}$  on  $\text{Bl}_V X$  simply contracts  $E$ .

In this paper, we will be concerned with the situation when  $f : X \dashrightarrow X$  is a birational map and  $V$  is contained in  $\text{Ind}(f)$ . In this case, interesting dynamics is possible on  $E \subset \text{Bl}_V(X)$ . We particularly focus on cases related to degenerations of rational surface automorphisms. Let  $T$  be an elliptic curve and  $X = \mathbb{P}^2 \times T$ . Given  $t \in T$ , there is an involution  $\iota_t$  associated to  $t$  and  $T$  constructed in [7, 2, 3]. Fix a non-trivial 2-torsion point  $q \in T$ . We have the map  $\varphi : X \dashrightarrow X$  defined by

$$(p, t) \mapsto (\iota_q(p), t),$$

the map  $\psi : X \dashrightarrow X$  by

$$(p, t) \mapsto (\iota_t(p), t),$$

and the map  $\chi : X \dashrightarrow X$  by

$$(p, t) \mapsto (\iota_{-t}(p), t),$$

where  $-t$  is the inverse of  $t$ . The map  $F = \varphi \circ \chi \circ \varphi \circ \psi$  can be viewed as a family of birational maps of  $\mathbb{P}^2$ :

$$\{\iota_q \circ \iota_{-t} \circ \iota_q \circ \iota_t : t \in T\}.$$

For a general  $t \in T$ , the map  $F_t : X_t \dashrightarrow X_t$  can be regularized to an automorphism on some higher birational model of  $X$ , and the dynamical degree of  $F_t$  is  $\lambda(F_t) = 17 + 12\sqrt{2}$ ; whereas the dynamical degree of  $F_q$  is  $\lambda(F_q) = 1$  as  $F_q = (\iota_q)^4 = \text{id}_{\mathbb{P}^2}$ . As the family degenerates to the map on the fiber  $X_q = \mathbb{P}^2 \times \{q\}$ ,  $F$  develops indeterminate curves inside the fiber  $X_q$ . In fact, the interesting dynamics does not completely disappear as the maps degenerate, but becomes concentrated along the indeterminate curves. To see this, we blow-up  $X$  at the indeterminate curves and find that the induced map on the exceptional divisor has dynamical degree  $\lambda = 16$ .

---

*Date:* July 31, 2024.

Here is the outline of the discussion:

- In Section 2, we will exhibit families of rational surface birational maps whose general member can be regularized as automorphisms. We will study the indeterminate curves, that are in the fiber over  $q$ , of the corresponding threefold map by factoring the map as a sequence of automorphisms and Atiyah flops.
- In Section 3, we will find induced self-maps of the exceptional divisors over the identified indeterminate curves in the special fiber over  $q$ .
- In Section 4, we will study the dynamical properties of the induced self-maps of the exceptional divisors.

In particular, we show the following result.

**Theorem 1.** *There exists a curve  $q \subset X_q$  which is indeterminate for the map  $\varphi \circ \chi \circ \varphi \circ \psi$ . Let  $\text{Bl}_q X$  be the blow-up of  $X = \mathbb{P}^2 \times T$  at  $q$  with exceptional divisor  $E_q$ . There is an induced birational map on  $E_q$  by  $\varphi \circ \chi \circ \varphi \circ \psi$ , and the map has dynamical degree  $\lambda = 16$ .*

As a warm-up, let's consider several examples of birational self-maps of  $\mathbb{P}^3$  where the induced maps on exceptional divisors can be easily obtained.

**Example 1.** Given  $b \in \mathbb{C}$ , let  $f : \mathbb{P}^3 \rightarrow \mathbb{P}^3$  be the automorphism defined by the formula

$$[w, x, y, z] \mapsto [bw, x, y, z].$$

Let  $L$  be the line  $\{y = z = 0\}$  in  $\mathbb{P}^3$ . Then  $f(L) = L$ , and the exceptional divisor  $E$  over  $L$  is isomorphic to  $\mathbb{P}^1 \times \mathbb{P}^1$  on  $\text{Bl}_L \mathbb{P}^3$ . The induced map  $\tilde{f}|_E : E \rightarrow E$  can be written as

$$([u, v], [s, t]) \mapsto ([bu, v], [s, t]).$$

In the affine chart  $\mathbb{A}^1 \times \mathbb{A}^1$  where  $v, t \neq 0$ , this is a dilation in the  $u$ -direction:

$$(u, s) \mapsto (bu, s).$$

**Example 2.** For a slightly more complicated example, let  $f : \mathbb{P}^3 \dashrightarrow \mathbb{P}^3$  be the birational map given by

$$[w, x, y, z] \mapsto [x^2w, x^3, x^2y, x^2y - ayw^2 + x^2z],$$

for some  $a \in \mathbb{C}$ . Then  $f$  is defined on a dense open subset of the line  $L$ . The induced map  $\tilde{f}|_E : E \dashrightarrow E$  is birational in this case:

$$([u, v], [s, t]) \mapsto ([u, v], [v^2t - au^2t + v^2s, v^2t]).$$

We note that in the affine chart where  $v, t \neq 0$ , this is an “area-preserving” bend:

$$(u, s) \mapsto (u, 1 - au^2 + s).$$

**Example 3.** If  $f(V) \neq V$ , then the lifted map simply contracts  $E$ . Let  $f : \mathbb{P}^3 \rightarrow \mathbb{P}^3$  be the automorphism

$$[w, x, y, z] \mapsto [y, z, w, x].$$

The lifted map  $\tilde{f} : \text{Bl}_L \mathbb{P}^3 \dashrightarrow \text{Bl}_L \mathbb{P}^3$  will contract the exceptional surface  $E$  to the strict transform of the line  $\{w = x = 0\} \subset \mathbb{P}^3$ .

**Example 4.** We suppose that  $V \subset \text{Ind}(f)$ , and we blow-up  $X$  at  $V$ . We consider when there will be an induced birational map on the exceptional divisor. Let  $f : \mathbb{P}^3 \dashrightarrow \mathbb{P}^3$  be the composition of the automorphism

$$[w, x, y, z] \mapsto [y, z, w, x]$$

and the standard involution of  $\mathbb{P}^3$ . The line  $L = \{y = z = 0\}$  is contained in the indeterminacy locus of  $f$ . In this case,  $f$  induces a (regular) birational map  $f|_E : E \rightarrow E$ , whose formula can be written as

$$([u, v], [s, t]) \mapsto ([s, t], [u, v]).$$

We note that, in the affine chart where  $v, t \neq 0$ , this map is the reflection across the line  $u = s$ :

$$(u, s) \mapsto (s, u).$$

**Example 5.** We can actually obtain a Hénon map on the exceptional divisor by composing the maps introduced in Examples 1, 2 and 4. The composition  $f$  is undefined along the line  $L = \{y = z = 0\}$ , but the induced map on the exceptional divisor  $E$  over  $L$  is a Hénon map.

For our main discussion, we need to cover some background knowledge.

## 1. GROUNDWORK

We assume that the base field  $K$  is algebraically closed of characteristic  $\text{char } K = 0$ .

**1.1. Relative numerical equivalence.** Let  $X$  be a smooth variety of dimension  $d$  over  $K$ , and let  $p : X \rightarrow S$  be a proper morphism onto a variety  $S$ . We define  $Z_1(X/S)$  to be the free abelian group generated by reduced irreducible curves which are mapped to points by  $p$ . There is a numerical equivalence  $\equiv_S$  in  $\text{Pic}(X)$  defined by

$$D \equiv_S D' \quad \text{if and only if} \quad D \cdot C = D' \cdot C \quad \text{for all } C \in Z_1(X/S).$$

We write  $N^1(X/S) := (\text{Pic}(X)/\equiv_S) \otimes_{\mathbb{Z}} \mathbb{R}$ . When  $S = \text{Spec } K$ , we drop  $/S$  from the notation.

**1.2. Blow-up and exceptional locus.** The blow-up of a variety  $X$  at a subvariety  $V$  will be written as  $\text{Bl}_V X$ . Let  $\pi : \text{Bl}_V X \rightarrow X$  be the projection map. The exceptional divisor  $\pi^{-1}(V)$  is isomorphic to  $\mathbb{P}(N_{V/X})$ , where  $N_{V/X}$  is the normal bundle to  $V$  in  $X$ . We denote  $\pi^{-1}(V)$  by  $E_V$ . We can thus identify the points on  $E_V$  with elements in  $\mathbb{P}(N_{V/X})$ .

We will frequently consider sequences of blow-ups throughout the paper. To simplify notations, we use the same notation for a subvariety  $V$  and all of its strict transforms.

**1.3. Dynamical degree and algebraic stability.** Let  $X$  be a smooth projective variety of dimension  $k$ , and  $f : X \dashrightarrow X$  a dominant rational map. Let  $H \in \text{Div}(X)$  be a big and nef divisor. We define the *(H-)degree* of  $f$  to be

$$\deg_H(f) := H^{k-1} \cdot f^*(H).$$

The *(first) dynamical degree* of  $f$  is defined to be

$$\lambda(f) := \lim_{n \rightarrow \infty} (\deg_H(f^n))^{1/n}.$$

The fact that the above limit exists and is independent of the choices of  $H$  is proved in [5] [10].

Now, let  $S$  be a quasi-projective surface and  $f : S \dashrightarrow S$  a birational map. The map  $f$  is said to be *algebraically stable* if

$$(f^n)^* = (f^*)^n$$

for any integer  $n \geq 0$ . The following theorem is useful in practice.

**Theorem 2** (Theorem 1.14 in [6]). *Let  $S$  be a quasi-projective surface and  $f : S \dashrightarrow S$  a birational map.  $f$  is algebraically stable if and only if there is no curve  $C \subset S$  such that  $f^k(C) \subset \text{Ind}(f)$  for some integer  $k \geq 0$ .*

Moreover, we can use the following Theorem to compute the dynamical degree for an algebraically stable birational surface map.

**Theorem 3** (Corollary 1.19 in [6]). *Let  $S$  be a quasi-projective surface and  $f : S \dashrightarrow S$  an algebraically stable birational map. Then  $\lambda(f)$  coincides with the spectral radius of  $f_* : N^1(S) \rightarrow N^1(S)$ .*

**1.4. Atiyah flop.** Let  $X$  be a quasi-projective threefold, and let  $B$  and  $C$  be two curves on  $X$  intersecting only at a point  $p \in X$  with distinct tangent directions. Let

$$\begin{aligned}\tilde{X} &:= \mathrm{Bl}_C \mathrm{Bl}_B X \xrightarrow{\pi_2} \mathrm{Bl}_B X \xrightarrow{\pi_1} X \\ \tilde{X}^+ &:= \mathrm{Bl}_B \mathrm{Bl}_C X \xrightarrow{\pi_2^+} \mathrm{Bl}_C X \xrightarrow{\pi_1^+} X\end{aligned}$$

be the blow-ups of  $X$  at  $B$  and  $C$  in different orders. Finally, we take the preimage of the point  $p$  on  $\mathrm{Bl}_B X$ , which is a curve isomorphic to  $\mathbb{P}^1$ ; and we let  $\mathfrak{b}$  be the strict transform of the curve on  $\tilde{X}$ .

**Proposition 4.** *The normal bundle  $N_{\mathfrak{b}/\tilde{X}}$  is isomorphic to  $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$ , and the natural birational map  $\phi : \tilde{X} \dashrightarrow \tilde{X}^+$  is an Atiyah flop with the flopping curve  $\mathfrak{b}$ .*

In the case where a pseudo-automorphism  $f$  of a threefold can be factored into a sequence of Atiyah flops and automorphisms, the image of a divisor class can be computed recursively. We need the following lemma.

**Lemma 5** (cf. Lemma 8 in [9]). *Let  $\phi : X \dashrightarrow X^+$  be an Atiyah flop with the flopping curve  $c \subset X$ . Let  $c^+$  denote the flopping curve of  $\phi^{-1}$ . Suppose that  $D^+$  is a divisor on  $X^+$  and that  $D$  is the pullback of  $D^+$  under  $\phi$ . Then*

$$\mathrm{mult}_c(D) = \mathrm{mult}_{c^+}(D^+) + D^+ \cdot c^+.$$

Let  $p : X \rightarrow T$  be a proper and flat family of surfaces parametrized by a curve  $T$ . Let  $\phi : X \dashrightarrow X^+$  be an Atiyah flop with flopping curve  $c$  lying in the fiber  $F := X_q$  for some  $q \in T$ . Let  $c^+$  denote the flopped curve, and let  $F^+ := \phi(F) \subset X^+$ . For a divisor  $D^+ \in N^1(X^+/T)$ , we have

$$(1.1) \quad \phi^*(D^+)|_F = (\phi|_F)^*(D^+|_{F^+}) + (D^+ \cdot c^+)c.$$

Indeed, by Lemma 5, we have

$$\mathrm{mult}_c(\phi^*(D^+)|_F) = \mathrm{mult}_c(D) = \mathrm{mult}_{c^+}(D^+) + D^+ \cdot c^+ = \mathrm{mult}_{c^+}(D^+|_{F^+}) + D^+ \cdot c^+.$$

Since  $\phi|_F : F \rightarrow F^+$  is an isomorphism,

$$\mathrm{mult}_c((\phi|_F)^*(D^+|_{F^+})) = \mathrm{mult}_{c^+}(D^+|_{F^+}).$$

The equation (1.1) follows.

Inductively, we obtain the following result: Let  $X$  be a threefold and  $f : X \dashrightarrow X$  a pseudo-automorphism of  $X$  preserving the fibration  $p : X \rightarrow T$  with indeterminate curves lying inside the fiber  $F$ . Suppose  $f$  factors into a sequence of flops and isomorphisms

$$f : X = X_1 \xrightarrow{\phi_1} X_1^+ \xrightarrow{f_1} X_2 \xrightarrow{\phi_2} \cdots \xrightarrow{f_{n-1}} X_n = X,$$

where  $\phi_i$  are Atiyah flops, and  $f_i$  are the isomorphisms. Let  $c_i$  be the flopping curve of  $\phi_i$  with flopped curve  $c_i^+$ . We write

$$g_i := f_{n-1} \circ \cdots \circ f_{n-i} : X_{n-i}^+ \dashrightarrow X_n, i = 1, \dots, n-1;$$

and write

$$h_i := f_{n-i-1} \circ \cdots \circ \phi_1 : X_1 \dashrightarrow X_{n-i}, i = 1, \dots, n-2, \quad \text{and} \quad h_{n-1} := \text{id}_X.$$

**Proposition 6.** *For a divisor  $D \in N^1(X/T)$ ,*

$$f^*(D)|_F = (f|_F)^*(D|_F) + \sum_{i=1}^{n-1} (g_i^*(D) \cdot c_{n-i}^+) h_i^*(c_{n-i}).$$

In particular, the difference

$$(f^n)^*(D)|_F - (f^n|_F)^*(D|_F)$$

is an effective divisor supported on flopping curves and their pullbacks under  $f|_F$ .

Finally, we observe that there is a natural birational map between two birational models. Let  $B$  and  $C$  be two curves in a threefold  $X$  intersecting at a point  $p$  with distinct tangent directions. There are several ways to blow-up  $X$ :

- (1) The first sequence of blow-ups (at the obvious centers) is

$$\text{Bl}_C \text{Bl}_B \text{Bl}_p X \xrightarrow{\pi_C} \text{Bl}_B \text{Bl}_p X \xrightarrow{\pi_B} \text{Bl}_p X \xrightarrow{\pi} X$$

with exceptional divisors are  $E_p, E_B, E_C$  over  $p, B, C$ , respectively.

- (2) The second sequence of blow-ups is

$$\text{Bl}_C \text{Bl}_{f'} \text{Bl}_B X \xrightarrow{\pi'_C} \text{Bl}_{f'} \text{Bl}_B X \xrightarrow{\pi'} \text{Bl}_B X \xrightarrow{\pi'_B} X$$

with exceptional divisors are  $E'_B, E'_{f'}, E'_C$  over  $B, p, C$ , respectively. The blow-up center of  $\pi'$  is the fiber  $f'$  of  $E'_B$  over  $p$ .

- (3) The third sequence of blow-ups is

$$\text{Bl}_{f''} \text{Bl}_C \text{Bl}_B X \xrightarrow{\pi''} \text{Bl}_C \text{Bl}_B X \xrightarrow{\pi''_C} \text{Bl}_B X \xrightarrow{\pi''_B} X$$

with exceptional divisors are  $E''_B, E''_C, E''_{f''}$  over  $B, C, p$ , respectively. The blow-up center of  $\pi''$  is the flopping curve  $f''$  of the Atiyah flop  $\phi : \text{Bl}_C \text{Bl}_B X \dashrightarrow \text{Bl}_B \text{Bl}_C X$ .

**Proposition 7.** *There is an isomorphism (induced by the identity map)*

$$\text{Bl}_C \text{Bl}_B \text{Bl}_p X \xrightarrow{\cong} \text{Bl}_C \text{Bl}_{f'} \text{Bl}_B X,$$

which identifies  $E_p$  and  $E'_{f'}$ . Furthermore, there is an Atiyah flop (induced by the identity map)

$$\text{Bl}_C \text{Bl}_B \text{Bl}_p X \cong \text{Bl}_C \text{Bl}_{f'} \text{Bl}_B X \dashrightarrow \text{Bl}_{f''} \text{Bl}_C \text{Bl}_B X$$

that sends  $E_p \cong \text{Bl}_2 \mathbb{P}^2$  to  $E''_{f''} \cong \mathbb{P}^1 \times \mathbb{P}^1$ .

We have acquired all the necessary tools for further discussion.

## 2. FAMILY OF MAPS

We present families of rational surface birational self-maps in this section and examine their properties.

**2.1. Cubic involution of  $\mathbb{P}^2$ .** We recall the following construction in [7, 2, 3]: Let  $T$  be a smooth cubic curve in  $\mathbb{P}^2$ , and let  $q \in T$  be a point. For a general point  $p \in \mathbb{P}^2$ , the line  $L_{q,p}$  connecting  $q, p$  intersects  $T$  at three points  $q, a, b$ . There is a unique involution  $i_{q,p}$  of  $L_{q,p}$  with  $i_{q,p}(a) = b$  and  $i_{q,p}(b) = a$ . Therefore, given a smooth cubic curve  $T$  and  $q \in T$ , we can define a cubic involution  $\iota_q : \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$  by

$$p \mapsto \iota_q(p) := i_{q,p}(p).$$

We first note that the cubic involution fixes (a dense open subset of) the cubic curve pointwise by construction. Now, let  $q$  be a point in  $T$  such that there are four distinct lines  $l_1, l_2, l_3, l_4$  passing through  $q$  and being tangent to  $T$  at  $q_1, q_2, q_3, q_4$  respectively. The points  $q_1, q_2, q_3, q_4$  will be sometimes referred to as the (four) tangent points associated to  $q$ . The indeterminacy locus of  $\iota_q$  is

$$\text{Ind}(\iota_q) = \{q, q_1, q_2, q_3, q_4\}.$$

In fact, the cubic involution  $\iota_q$  contracts the conic  $\mathfrak{q}$  passing through  $q, q_1, q_2, q_3, q_4$  to  $q$  and contracts the tangent line  $l_i$  to  $q_i$  for each  $i$ . The (affine) picture of the cubic curve, the conic and the tangent lines is in Figure 1.




FIGURE 1. Curves contracted by  $\iota_q$

We will see that the sequence  $\{\deg((\iota_q \circ \iota_t)^n)\}$  grows quadratically for general  $t \neq q$ . So, by varying  $t \in T$ , we get a family of birational maps of  $\mathbb{P}^2$ , whose general member has quadratic degree growth, but the special map  $\iota_q \circ \iota_q$  is the identity of  $\mathbb{P}^2$ . Furthermore, the map  $\iota_q \circ \iota_{t'} \circ \iota_q \circ \iota_t$  has dynamical degree greater than 1 for general distinct  $t, t', q$ . We can obtain a family of birational maps of  $\mathbb{P}^2$ , whose general member has  $\lambda > 1$ , but again, the special member when  $t = t' = q$  is the identity.

**2.2. Involution of  $\mathbb{P}^2 \times T$ .** We define the families of maps formally as birational maps of a threefold. Let  $X$  be the threefold  $\mathbb{P}^2 \times T \subset \mathbb{P}^2 \times \mathbb{P}^2$ . Fix a point  $q \in T$  such that  $\#\text{Ind}(\iota_q) = 5$ . We define an involution  $\varphi : X \dashrightarrow X$  using  $\iota_q$  by

$$(p, t) \mapsto (\iota_q(p), t).$$

We note that  $\varphi$  can also be viewed as a family of birational involutions of  $\mathbb{P}^2$ , i.e.,  $\{\phi_t := \iota_q : t \in T\}$ . Next, we further let  $q$  be a nontrivial 2-torsion point with respect to the group law on  $T$ . We define

the involution  $\psi : X \dashrightarrow X$  by

$$(p, t) \mapsto (\iota_t(p), t)$$

and the involution  $\chi : X \dashrightarrow X$  by

$$(p, t) \mapsto (\iota_{-t}(p), t),$$

where  $-t$  is the inverse of  $t$  with respect to the group law on  $T$ . So, we have two other families of maps  $\{\psi_t := \iota_t : t \in T\}$  and  $\{\chi_t := \iota_{-t} : t \in T\}$ .

Let us take the open subset of  $T$  consisting of the point  $q$  and points  $t \in T$  such that

$$\#\text{Ind}(\iota_t) = \#\text{Ind}(\iota_{-t}) = 5,$$

and that

$$\text{Ind}(\iota_t) \cap \text{Ind}(\iota_q) = \text{Ind}(\iota_{-t}) \cap \text{Ind}(\iota_q) = \text{Ind}(\iota_t) \cap \text{Ind}(\iota_{-t}) = \emptyset;$$

and replace  $T$  with this open subset. We still keep the same notations for the threefold  $X = \mathbb{P}^2 \times T$  and restricted involutions  $\varphi, \psi, \chi$ . We are now ready to state the indeterminacy locus of the maps  $\varphi, \psi$  and  $\chi$ .

The indeterminacy locus of  $\varphi$  contains five curves

$$B = \{q\} \times T \quad \text{and} \quad B_i = \{q_i\} \times T$$

for  $i = 1, \dots, 4$ . The indeterminacy locus of  $\psi$  contains two irreducible curves, namely, the curve

$$C = \{(p, t) : p = t\}$$

and the curve  $C'$ . For  $t \in T$ , the restriction of the curve  $C'$  to the fiber  $\mathbb{P}^2 \times \{t\}$  consists of the tangent points  $t_1, t_2, t_3, t_4$  associated to  $t$ . The indeterminacy locus of  $\chi$  contains the curve

$$D = \{(p, t) : p = -t\}$$

and the curve  $D'$  whose restriction to  $\mathbb{P}^2 \times \{t\}$  are the four tangent points associated to  $-t$ . We note that

$$B \cap C \cap D = \{(q, q)\} \quad \text{and} \quad B_i \cap C' \cap D' = \{(q_i, q)\}$$

for  $i = 1, \dots, 4$ . These are the only intersection points of the indeterminate curves.

**Example 6.** Let  $T$  be an elliptic curve over a field  $K$  of characteristic  $\text{char } K \neq 2$  on  $\mathbb{A}^2$  given by

$$b^2 = a(a - \alpha_1)(a - \alpha_2),$$

where  $\alpha_1, \alpha_2$  are distinct nonzero elements of  $K$ , and  $q = (0, 0) \in T$ . We take the open subset of  $T$  as discussed above and still call it  $T$ . Let  $X = \mathbb{A}^2 \times T \subset \mathbb{A}^2 \times \mathbb{A}^2$  with coordinates  $((x, y), (a, b))$ .

The curve  $B$  is defined (in  $X$ ) by

$$x = y = 0.$$

The curve  $C$  is defined by

$$x - a = y - b = 0.$$

The curve  $D$  is defined by

$$x - a = y + b = 0.$$

For  $B_1, \dots, B_4, C'$ , and  $D'$ , we need explicit formulae for the tangent points associated to a given point on the elliptic curve, and we will follow the calculation in [1]. For example, for  $q = (0, 0)$ , the curves  $B_1$ , is defined in  $X$  by

$$x + \sqrt{-\alpha_1}\sqrt{-\alpha_2} = y - (-\sqrt{-\alpha_1} + \sqrt{-\alpha_2})x = 0.$$

The curves  $C'$  and  $D'$  can be obtained as the two irreducible components of the following reducible curve defined by

$$x^4 - 4ax^3 + 4(\alpha_1 + \alpha_2)ax^2 - 2\alpha_1\alpha_2x^2 - 4\alpha_1\alpha_2ax + \alpha_1^2\alpha_2^2 = y^2 - x(x - \alpha_1)(x - \alpha_2) = 0.$$

Using  $\varphi, \psi$  and  $\chi$ , we can define families of birational self-maps. We will first study the composition  $\varphi \circ \psi : X \dashrightarrow X$ , i.e., the family of maps  $\{\iota_q \circ \iota_t : t \in T\}$ . We will then take the composition  $\varphi \circ \chi \circ \varphi \circ \psi$ , which is the family of maps  $\{\iota_q \circ \iota_{-t} \circ \iota_q \circ \iota_t : t \in T\}$ .

**2.3. Indeterminate curves of  $\varphi \circ \psi$ .** Let us write  $\mathcal{B} := B \cup B_1 \cup B_2 \cup B_3 \cup B_4$ , and  $\mathcal{C} := C \cup C'$ . The map  $\varphi$  will be regularized on  $\text{Bl}_{\mathcal{B}} X$  and hence on  $\text{Bl}_{\mathcal{C}} \text{Bl}_{\mathcal{B}} X$  as  $\varphi$  fixes the curves in  $\mathcal{C}$  pointwise; and that the map  $\psi$  will be regularized on  $\text{Bl}_{\mathcal{C}} X$  and hence on  $\text{Bl}_{\mathcal{B}} \text{Bl}_{\mathcal{C}} X$  as  $\psi$  fixes the curves in  $\mathcal{B}$  pointwise (cf. [4]). Let  $\tilde{\varphi}$  and  $\tilde{\psi}$  be the lifts of  $\varphi$  and  $\psi$  on  $\text{Bl}_{\mathcal{C}} \text{Bl}_{\mathcal{B}} X$ . The map  $\tilde{\varphi} \circ \tilde{\psi}$  has no indeterminacy on each fiber over  $t$  for  $t \neq q$ . However, since  $\text{Bl}_{\mathcal{C}} \text{Bl}_{\mathcal{B}} X$  and  $\text{Bl}_{\mathcal{B}} \text{Bl}_{\mathcal{C}} X$  differ by a finite compositions of Atiyah flops with pairwise disjoint flopping curves in the fiber over  $q$ , we will see that  $\tilde{\varphi} \circ \tilde{\psi}$  will have indeterminate curves within the fiber over  $q$ .

Let us fix some notations for the rest of the discussion. Given a threefold  $X$  with a fibration  $X \rightarrow T$ , let  $X_t$  denote the fiber of  $X$  over  $t \in T$ . For a map  $f : X \dashrightarrow X$  preserving  $X \rightarrow T$ , let  $f_t$  denote the restriction  $f|_{X_t}$ .

*Blow-up, first round.* We denote by  $\text{Bl}_{\mathcal{B}} X$  the blow-up of  $X$  at  $\mathcal{B}$  with exceptional divisors  $E_{\mathcal{B}}$  and  $E_{B_i}$  over  $B$  and  $B_i$  respectively. For  $t \in T$ , let  $(\text{Bl}_{\mathcal{B}} X)_t$  be the fiber of  $\text{Bl}_{\mathcal{B}} X$  over  $t$ . Then  $(\text{Bl}_{\mathcal{B}} X)_t$  is the blow-up of  $X_t \cong \mathbb{P}^2$  at the five points

$$(t, t), (t_1, t), (t_2, t), (t_3, t), (t_4, t) \in X_t = \mathbb{P}^2 \times \{t\},$$

where  $t, t_1, t_2, t_3, t_4$  are the indeterminate points of  $\iota_t$  as a birational involution of  $\mathbb{P}^2$ . The involution  $\varphi$  lifts to a regular involution  $\hat{\varphi}$  on  $\text{Bl}_{\mathcal{B}} X$ . On each fiber  $(\text{Bl}_{\mathcal{B}} X)_t$ ,  $\hat{\varphi}|_{(\text{Bl}_{\mathcal{B}} X)_t} : (\text{Bl}_{\mathcal{B}} X)_t \rightarrow (\text{Bl}_{\mathcal{B}} X)_t$  swaps the exceptional curve over  $(q, t)$  and the strict transform of the conic  $\mathfrak{q} \times \{t\}$ , and exchanges the other four exceptional curves and the strict transforms of the four tangent lines respectively. In particular, on the fiber  $(\text{Bl}_{\mathcal{B}} X)_q$ , we have the five exceptional curves

$$\mathfrak{b} := E_{\mathcal{B}} \cap (\text{Bl}_{\mathcal{B}} X)_q \quad \text{and} \quad \mathfrak{b}_i := E_{B_i} \cap (\text{Bl}_{\mathcal{B}} X)_q$$

for  $i = 1, \dots, 4$ . Moreover, we still denote by  $\mathfrak{q}$  the strict transform of the curve  $\mathfrak{q} \times \{q\} \subset X_q$  on  $\text{Bl}_{\mathcal{B}} X$ , and by  $\mathfrak{l}_i$  the strict transforms of the curves  $\mathfrak{l}_i \times \{q\} \subset X_q$  on  $\text{Bl}_{\mathcal{B}} X$  for  $i = 1, \dots, 4$ . We have

$$\hat{\varphi}(\mathfrak{b}) = \mathfrak{q} \quad \text{and} \quad \hat{\varphi}(\mathfrak{b}_i) = \mathfrak{l}_i$$

for  $i = 1, \dots, 4$ .

Let  $\text{Bl}_{\mathcal{C}} X$  be the blow-up of  $X$  at  $\mathcal{C}$  with the exceptional divisors  $E_{\mathcal{C}}^+$  and  $E_{\mathcal{C}'}$  over  $\mathcal{C}$  and  $\mathcal{C}'$  respectively. The map  $\psi$  lifts to an automorphism  $\hat{\psi}^+$  on  $\text{Bl}_{\mathcal{C}} X$ . On the fiber  $(\text{Bl}_{\mathcal{C}} X)_q$  over  $q$ , there are the exceptional curves

$$\mathfrak{c}^+ = E_{\mathcal{C}}^+ \cap (\text{Bl}_{\mathcal{C}} X)_q \quad \text{and} \quad \mathfrak{c}_1^+ \sqcup \mathfrak{c}_2^+ \sqcup \mathfrak{c}_3^+ \sqcup \mathfrak{c}_4^+ = E_{\mathcal{C}'}, \cap (\text{Bl}_{\mathcal{C}} X)_q;$$

and there are the strict transform  $\mathfrak{q}^+$  of the conic and the strict transforms  $\mathfrak{l}_i^+$  of the tangent lines for  $i = 1, \dots, 4$ . We have

$$\hat{\psi}^+(\mathfrak{c}^+) = \mathfrak{q}^+ \quad \text{and} \quad \hat{\psi}^+(\mathfrak{c}_i^+) = \mathfrak{l}_i^+$$

for  $i = 1, \dots, 4$ .



*Blow-up, second round.* We want to relate  $\text{Bl}_{\mathcal{B}} X$  and  $\text{Bl}_{\mathcal{C}} X$ .

We can blow-up  $\text{Bl}_{\mathcal{B}} X$  at the strict transform of  $\mathcal{C}$  to get  $\text{Bl}_{\mathcal{C}} \text{Bl}_{\mathcal{B}} X$ . We still denote by  $E_B$  and  $E_{B_i}$  the exceptional divisors over  $B$  and  $B_i$ , respectively, on  $\text{Bl}_{\mathcal{C}} \text{Bl}_{\mathcal{B}} X$ ; and we denote by  $E_C$  and  $E_{C'}$  the exceptional divisors over  $C$  and  $C'$ , respectively, on  $\text{Bl}_{\mathcal{C}} \text{Bl}_{\mathcal{B}} X$ . We write  $\tilde{X} := \text{Bl}_{\mathcal{C}} \text{Bl}_{\mathcal{B}} X$ .

In the second round of blow-ups, we actually blow-up each fiber of  $\text{Bl}_{\mathcal{B}} X$  further at five additional points. For  $t \neq q$ , the fiber of  $\tilde{X}$  over  $t$  is isomorphic to the blow-up of  $X_t \cong \mathbb{P}^2$  at the ten distinct points  $(\mathcal{B} \cup \mathcal{C}) \cap X_t$ . However, since the curves  $\mathcal{B}$  and  $\mathcal{C}$  intersect in the fiber  $X_q$  at the five points

$$(q, q), (q_1, q), (q_2, q), (q_3, q), (q_4, q),$$

the fiber  $\tilde{X}_q$  is the blow-up of  $X_q \cong \mathbb{P}^2$  at five points and then at five infinitely near points. More specifically,  $(\text{Bl}_{\mathcal{B}} X)_q$  is the blow-up of  $X_q$  at  $(q, q), (q_1, q), (q_2, q), (q_3, q), (q_4, q)$  with exceptional curves  $\mathfrak{b}, \mathfrak{b}_1, \mathfrak{b}_2, \mathfrak{b}_3, \mathfrak{b}_4$ . The curves  $\mathfrak{b}, \mathfrak{b}_1, \mathfrak{b}_2, \mathfrak{b}_3, \mathfrak{b}_4$  intersect the strict transforms of  $C, C'$  transversally at the points  $(q', q), (q'_1, q), (q'_2, q), (q'_3, q), (q'_4, q)$  in  $(\text{Bl}_{\mathcal{B}} X)_q$  as shown in Figure 2. Therefore,  $\tilde{X}_q$  is the blow-up of  $(\text{Bl}_{\mathcal{B}} X)_q$  at such points.




FIGURE 2. Blow-ups of  $X_q$

Let

$$\mathfrak{c} := E_C \cap \tilde{X}_q \quad \text{and} \quad \mathfrak{c}_1 \sqcup \mathfrak{c}_2 \sqcup \mathfrak{c}_3 \sqcup \mathfrak{c}_4 := E_{C'} \cap \tilde{X}_q.$$

We still write  $\mathfrak{b}$  for the strict transform of  $\mathfrak{b}$  in  $\tilde{X}$ , and  $\mathfrak{b}_i$  for the strict transform of  $\mathfrak{b}_i$  in  $\tilde{X}$  for each  $i$ . Since the map  $\tilde{\varphi}$  fixes the strict transform  $C$  and  $C'$  pointwise, it lifts to an automorphism  $\tilde{\varphi}$  on  $\tilde{X}$ , and  $\tilde{\varphi}$  will preserve the new exceptional divisors  $E_C, E_{C'}$  as  $C$  and  $C'$  are fixed curves by  $\varphi$ . On  $\tilde{X}_q$ , we have

$$\tilde{\varphi}(\mathfrak{c}) = \mathfrak{c} \quad \text{and} \quad \tilde{\varphi}(\mathfrak{c}_i) = \mathfrak{c}_i$$

for  $i = 1, \dots, 4$ , and

$$\tilde{\varphi}(\mathfrak{b}) = \mathfrak{q} \quad \text{and} \quad \tilde{\varphi}(\mathfrak{b}_i) = \mathfrak{l}_i,$$

where we use the same notations for the strict transforms of the conic and tangent lines on  $\tilde{X}$ .

On the other hand, we can blow-up  $\text{Bl}_C X$  at the strict transform of  $\mathcal{B}$  to get  $\tilde{X}^+ := \text{Bl}_{\mathcal{B}} \text{Bl}_C X$ . The exceptional divisors over  $B$  and  $B_i$  in  $\tilde{X}^+$  are  $E_B^+$  and  $E_{B_i}^+$  respectively, and the exceptional divisors over  $C$  and  $C'$  in  $\tilde{X}^+$  are  $E_C^+$  and  $E_{C'}^+$  respectively. On the fiber  $(\tilde{X}^+)_q$ , we have

$$\mathfrak{b}^+ := E_B^+ \cap \tilde{X}_q^+ \quad \text{and} \quad \mathfrak{b}_i^+ := E_{B_i}^+ \cap \tilde{X}_q^+$$

for  $i = 1, \dots, 4$ ; and we again write  $\mathfrak{c}^+$  and  $\mathfrak{c}_i^+$  for the strict transforms of  $\mathfrak{c}^+$  and  $\mathfrak{c}_i^+$  in  $\tilde{X}^+$  for each  $i$ . The map  $\hat{\psi}^+$  lifts to an automorphism  $\tilde{\psi}^+$  on  $\tilde{X}^+$ , which preserves the new exceptional divisors  $E_B^+, E_{B_1}^+, E_{B_2}^+, E_{B_3}^+, E_{B_4}^+$ . In particular, we have

$$\tilde{\psi}^+(\mathfrak{b}^+) = \mathfrak{b}^+ \quad \text{and} \quad \tilde{\psi}^+(\mathfrak{b}_i^+) = \mathfrak{b}_i^+,$$

for  $i = 1, \dots, 4$ , and

$$\tilde{\psi}^+(\mathfrak{c}^+) = \mathfrak{q}^+ \quad \text{and} \quad \tilde{\psi}^+(\mathfrak{c}_i^+) = \mathfrak{l}_i^+,$$

for  $i = 1, \dots, 4$ .

*Factorization of  $\tilde{\varphi} \circ \tilde{\psi}$ .* Now the blow-ups  $\tilde{X}$  and  $\tilde{X}^+$  are related by a map  $\sigma : \tilde{X} \dashrightarrow \tilde{X}^+$ , which is a composition of five Atiyah flops. The five pairwise disjoint flopping curves for  $\sigma$  are  $\mathfrak{b}, \mathfrak{b}_1, \mathfrak{b}_2, \mathfrak{b}_3, \mathfrak{b}_4$ ; and the five pairwise disjoint flopped curves, which are the flopping curves for  $\sigma^{-1}$ , are  $\mathfrak{c}^+, \mathfrak{c}_1^+, \mathfrak{c}_2^+, \mathfrak{c}_3^+, \mathfrak{c}_4^+$ .

Let us denote by  $\tilde{\psi}$  the lift of  $\psi$  on  $\tilde{X}$ . We factorize the map  $\tilde{\varphi} \circ \tilde{\psi}$  in terms of flops and isomorphisms as follows:

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\tilde{\psi}} & \tilde{X} & \xrightarrow{\tilde{\varphi}} & \tilde{X} \\ \sigma \downarrow & & \uparrow \sigma^{-1} & & \\ \tilde{X}^+ & \xrightarrow{\tilde{\psi}^+} & \tilde{X}^+ & & \end{array} .$$

Although  $\tilde{\psi}^+$  is an automorphism on  $\tilde{X}^+$ ,  $\tilde{\psi}$  is a pseudo-automorphism on  $\tilde{X}$ , which is conjugate to  $\tilde{\psi}^+$  by  $\sigma$ . We note that  $\sigma$  is undefined at  $\mathfrak{b}, \mathfrak{b}_1, \mathfrak{b}_2, \mathfrak{b}_3, \mathfrak{b}_4$ , and  $\sigma^{-1}$  is undefined at  $\mathfrak{c}^+, \mathfrak{c}_1^+, \mathfrak{c}_2^+, \mathfrak{c}_3^+, \mathfrak{c}_4^+$ . So the composition  $\tilde{\psi} = \sigma^{-1} \circ \tilde{\psi}^+ \circ \sigma : \tilde{X} \dashrightarrow \tilde{X}$  is undefined along the ten curves

$$\mathfrak{b}, \mathfrak{b}_1, \mathfrak{b}_2, \mathfrak{b}_3, \mathfrak{b}_4, \mathfrak{q}, \mathfrak{l}_1, \mathfrak{l}_2, \mathfrak{l}_3, \mathfrak{l}_4.$$

Since  $\tilde{\varphi}$  is a regular involution exchanging  $\mathfrak{b}, \mathfrak{b}_1, \mathfrak{b}_2, \mathfrak{b}_3, \mathfrak{b}_4$  and  $\mathfrak{q}, \mathfrak{l}_1, \mathfrak{l}_2, \mathfrak{l}_3, \mathfrak{l}_4$ , the composition  $\tilde{\varphi} \circ \tilde{\psi} : \tilde{X} \dashrightarrow \tilde{X}$  is also a pseudo-automorphism that is undefined at the same ten curves.

**2.4. Dynamics on fibers.** For any divisor, we use the same notation to denote its class in the (relative) Néron-Severi group.

Let  $\pi_1 : \mathbb{P}^2 \times E \rightarrow \mathbb{P}^2$  be the first projection, and let  $\pi : \tilde{X} \rightarrow X$  be the blow-up map. For a hyperplane divisor  $h \in N^1(\mathbb{P}^2)$ , we write  $H := \pi^*(\pi_1^*(h)) \in N^1(\tilde{X}/T)$ . For any  $t \neq q$ , we write  $h_t$  for  $H|_{\tilde{X}_t}$ . To distinguish from the exceptional curves in the fiber over  $q$ , we denote by  $b_t := E_B|_{\tilde{X}_t}$ ,  $c_t := E_C|_{\tilde{X}_t}$ ,  $b_{t,i} := E_{B_i}|_{\tilde{X}_t}$ ,  $c_{t,i} := E_{C_i}|_{\tilde{X}_t}$ ,  $i = 1, \dots, 4$ , the classes of the exceptional curves of  $\tilde{X}_t$ . Together with  $h_t$ , they form a set of basis elements for  $N^1(\tilde{X}_t)$ :

$$\{h_t, b_t, b_{t,1}, b_{t,2}, b_{t,3}, b_{t,4}, c_t, c_{t,1}, c_{t,2}, c_{t,3}, c_{t,4}\}.$$

We note that  $\tilde{\varphi}_t = \tilde{\varphi}|_{\tilde{X}_t}$  and  $\tilde{\psi}_t = \tilde{\psi}|_{\tilde{X}_t}$  are the lifts of  $\iota_q$  and  $\iota_t$ , respectively, on the blow-up of  $\mathbb{P}^2$  at the ten indeterminate points. These maps are both regular, and as computed in [4], we have

**Proposition 8.** For  $t \neq q$ , the action of  $\tilde{\varphi}_t$  on  $N^1(\tilde{X}_t)$  is given by

$$M_\varphi := \begin{pmatrix} 3 & 2 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ -2 & -1 & -1 & -1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

and the action of  $\tilde{\psi}_t$  on  $N^1(\tilde{X}_t)$  is given by

$$M_\psi := \begin{pmatrix} 3 & 0 & 0 & 0 & 0 & 0 & 2 & 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ -2 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & -1 & -1 & -1 \\ -1 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & -1 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

**Corollary 9.** The product of  $M_\psi$  and  $M_\varphi$  has a  $3 \times 3$  Jordan block associated with the eigenvalue  $\lambda = 1$ . Hence, the sequence  $\{\deg((\tilde{\varphi} \circ \tilde{\psi})_t^n)\}_n$  grows quadratically for  $t \neq q$ .

On the other hand, the map  $(\tilde{\varphi} \circ \tilde{\psi})_q$  is the identity map of the fiber  $\tilde{X}_q$ . Write  $\mathfrak{h} := H|_{\tilde{X}_q}$ . The difference (Proposition 6)

$$(\tilde{\varphi} \circ \tilde{\psi})^*(H)|_{\tilde{X}_q} - ((\tilde{\varphi} \circ \tilde{\psi})_q)^*(\mathfrak{h}) = 2\mathfrak{q} + \sum \mathfrak{l}_i + 4\mathfrak{b} + 2 \sum \mathfrak{b}_i$$

is a sum of the flopping curves (indeterminate curves); the indeterminate curves contribute the difference between

$$\deg_{\mathfrak{h}}((\tilde{\varphi} \circ \tilde{\psi})_q^n) \quad \text{and} \quad \deg_{\mathfrak{h}_t}((\tilde{\varphi} \circ \tilde{\psi})_t^n)$$

for  $t \neq q$ . In the rest of the paper, we will show that the dynamical complexity persists but is concentrated around the indeterminacy locus. Once we blow-up at the indeterminate curves and then look at the induced maps on the exceptional divisors, the interesting dynamics will be seen again.

As for the composition of four involutions  $\varphi \circ \chi \circ \varphi \circ \psi$ , i.e., the family  $\{\iota_q \circ \iota_{-t} \circ \iota_q \circ \iota_t : t \in T\}$ , the map  $\iota_q \circ \iota_{-t} \circ \iota_q \circ \iota_t$  can be regularized and has dynamical degree  $17 + 12\sqrt{2}$  for  $t \neq q$ , while the limiting map is again the identity map when  $t = q$ . The fact that dynamical degree drops on the special fiber in a family of birational maps is consistent with the lower semicontinuity theorem on dynamical degree by [8]. We will also blow-up at the indeterminate curves of  $\varphi \circ \chi \circ \varphi \circ \psi$  in the fiber over  $q$ . The interesting dynamics will be recovered on the exceptional divisors too.

## 3. INDUCED MAP ON DIVISORS

In this section we aim to blow-up  $\tilde{X}$  at  $\mathfrak{q}$  and obtain (a map which is conjugate to) the birational self-map  $E_{\mathfrak{q}} \dashrightarrow E_{\mathfrak{q}}$  induced by  $\tilde{\varphi} \circ \tilde{\psi}$ .

**3.1. Some reductions.** To look at the exceptional divisor  $E_{\mathfrak{q}}$ , it suffices to blow-up  $X$  at  $\mathfrak{q} = \mathfrak{q} \times \{q\} \subset X$  and find the map induced by  $\varphi \circ \psi$  on  $E_{\mathfrak{q}} \subset \text{Bl}_{\mathfrak{q}} X$ .

Next, we note that the induced map can be broken down into the composition of maps

$$E_{\mathfrak{q}} \xrightarrow{\tilde{\psi}^{-1}} E_{\mathfrak{q}} \xrightarrow{\tilde{\varphi}} E_{\mathfrak{q}}$$

induced by  $\varphi$  and  $\psi$ , where  $E_{\mathfrak{q}}$  is the exceptional divisor of the blow-up of  $X$  at the point  $(q, q) \in X$ . In fact, by Proposition 7, there is a birational map from  $E_{\mathfrak{q}}$  to  $E_{\mathfrak{b}}$  (the exceptional divisor over the flopping curve  $\mathfrak{b}$ ) induced by the identity map.

Finally, to see the dynamical property of the induced map  $\tilde{\varphi} \circ \tilde{\psi}^{-1}$ , we will analyze the map  $\Phi := \tilde{\psi}^{-1} \circ \tilde{\varphi} : E_{\mathfrak{q}} \dashrightarrow E_{\mathfrak{q}}$ , which is birationally conjugate to  $\tilde{\varphi} \circ \tilde{\psi}^{-1} : E_{\mathfrak{q}} \dashrightarrow E_{\mathfrak{q}}$  by  $\tilde{\varphi}$ . As  $E_{\mathfrak{q}}$  is isomorphic to  $\mathbb{P}^2$  and  $E_{\mathfrak{q}}$  is isomorphic to  $\mathfrak{q} \times \mathbb{P}^1$ , it is simpler to study the map  $\tilde{\psi}^{-1} \circ \tilde{\varphi}$  than the map  $\tilde{\varphi} \circ \tilde{\psi}^{-1}$ .

The map  $\varphi \circ \chi \circ \varphi \circ \psi$  induces a birational self-map on  $E_{\mathfrak{q}} \subset \text{Bl}_{\mathfrak{q}} X$  as well, and it can be broken down into the composition

$$E_{\mathfrak{q}} \xrightarrow{\tilde{\psi}^{-1}} E_{\mathfrak{q}} \xrightarrow{\tilde{\varphi}} E_{\mathfrak{q}} \xrightarrow{\tilde{\chi}^{-1}} E_{\mathfrak{q}} \xrightarrow{\tilde{\varphi}} E_{\mathfrak{q}}.$$

Again, we will study the map  $\Psi := \tilde{\chi}^{-1} \circ \tilde{\varphi}$  which is birationally conjugate to  $\tilde{\varphi} \circ \tilde{\chi}^{-1}$  by  $\tilde{\varphi}$ . Therefore, the map  $\Psi \circ \Phi$  is birationally conjugate to  $\tilde{\varphi} \circ \tilde{\chi}^{-1} \circ \tilde{\varphi} \circ \tilde{\psi}^{-1}$  by  $\tilde{\varphi}$ .

**3.2. Explicit formulae for cubic involution.** In order to obtain simpler formulae, we now fix the base field  $K = \mathbb{Q}$ , a specific cubic curve  $T$  given by

$$y^2 z = x^3 + 5x^2 z + 4xz^2$$

in  $\mathbb{P}^2$  and the point  $q = [0, 0, 1] \in T$ . We note that this is an elliptic curve of the form

$$y^2 z = x(x - \alpha_1 z)(x - \alpha_2 z)$$

with  $\alpha_1 = -1$  and  $\alpha_2 = -4$  in Example 6. One of the affine pieces of  $T$  is depicted in Figure 1.

The cubic involution associated to  $T$  and  $q$  is  $\iota_q : \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$  defined by

$$[x, y, z] \mapsto [5x^3 - xy^2 + 8x^2 z, 5x^2 y - y^3 + 8xyz, -2x^3 - 5x^2 z + y^2 z].$$

The involution  $\iota_q$  contracts the four tangent lines  $l_1, \dots, l_4$  through  $q$  to the points  $q_1, \dots, q_4$  respectively. The conic through  $q, q_1, \dots, q_4$  is

$$\mathfrak{q} : y^2 = 5x^2 + 8xz,$$

and  $\iota_q$  contracts  $\mathfrak{q}$  to  $q$ .

Now let  $\mathbb{P}^2 \times \mathbb{P}^2$  be the product of two projective planes with coordinates  $([x, y, z], [a, b, c])$  and  $X = \mathbb{P}^2 \times T$  the threefold defined by the equation

$$b^2 c = a^3 + 5a^2 c + 4ac^2$$

in  $\mathbb{P}^2 \times \mathbb{P}^2$ . We define the involution  $\varphi : X \dashrightarrow X$  by

$$([x, y, z], [a, b, c]) \mapsto (\iota_q([x, y, z]), [a, b, c]).$$

We will use the map  $\varphi$  to explain how to find induced maps.

**3.3. Procedure for finding induced maps.** It suffices to make local calculations for finding birational maps. Let  $\mathbb{A}^2 \times \mathbb{A}^2$  be the affine space with coordinates  $((x, y), (a, b))$ . One of the affine pieces of the above threefold  $X$  is defined by the equation

$$b^2 = a^3 + 5a^2 + 4a$$

in  $\mathbb{A}^2 \times \mathbb{A}^2$ . We still use  $X$  to denote this affine piece. We also keep the same notation for the affine counterpart of each of the aforementioned curves and points correspondingly. Moreover, we keep writing  $\varphi$  for the involution of the affine threefold  $X$ .

*Blow-up of  $X$  at conic.* We explicitly define (one of the affine pieces of) the blow-up  $\text{Bl}_q X$  of  $X$  at  $\mathfrak{q} = \mathfrak{q} \times \{q\}$  in  $\mathbb{A}^2 \times \mathbb{A}^2 \times \mathbb{A}^2$  with coordinates  $((u, v), (c, d), (r, s))$ , and write the projection map  $\pi : \text{Bl}_q X \rightarrow X$  and inverse of the projection maps  $\pi^{-1} : X \dashrightarrow \text{Bl}_q X$  in coordinates. The exceptional divisor over  $\mathfrak{q}$  is denoted by  $E_q$ . We note in particular that (one of the affine pieces of)  $E_q$  is subvariety

$$\{(u, v, c, d, r, s) : v^2 = 5u^2 + 8u, c = d = 0, s = 0\} \subset (\mathbb{A}^2)^3.$$

So there is an isomorphism

$$E_q \cong \mathfrak{q} \times \mathbb{A}^1 = \{(u, v, r) : v^2 = 5u^2 + 8u\} \subset \mathbb{A}^2 \times \mathbb{A}^1.$$

*Blow-up of  $X$  at point.* Let  $\text{Bl}_q X$  be the blow-up of  $X$  at  $(q, q)$ . We want to represent the exceptional divisor  $E_q$  over  $(q, q)$  using curves, which can be considered as elements in  $\mathbb{P}(N_{(q,q)}/X)$ . We take parametrized curves through the point  $(q, q) = ((0, 0), (0, 0)) \in X$  in different tangent directions. For example, given  $(z, w) \in \mathbb{A}^2$ , the parametrized curve  $\gamma_{(z,w)} : T \rightarrow X$  with  $T := \{\beta^2 = \alpha^3 + 5\alpha^2 + 4\alpha\} \subset \mathbb{A}_{\alpha,\beta}^2$  given by

$$(\alpha, \beta) \mapsto ((\beta z, \beta w), (\alpha, \beta)).$$

has tangent vector  $((z, w), (0, 1))$  at  $((0, 0), (0, 0))$ . In this way, the curve  $\gamma_{(z,w)}$  corresponds to a point  $p_{(z,w)} \in E_q$ . Different curves  $\gamma_{(z,w)}, \gamma_{(z',w')}$  have different tangent vectors at  $(q, q)$  and therefore correspond to different points  $p_{(z,w)}, p_{(z',w')}$  in  $E_q$ . Taking all different points  $(z, w) \in \mathbb{A}^2$ , we obtain a family of parametrized curves  $\Gamma : \mathbb{A}^2 \times T \rightarrow X$  given by

$$((z, w), (\alpha, \beta)) \mapsto ((\beta z, \beta w), (\alpha, \beta)),$$

which correspond to a dense open subset of  $E_q$ . We identify the  $\mathbb{A}^2$  with (a dense open subset of)  $E_q$ , and now  $\Gamma$  has parameter domain  $E_q \times T$ .

*Induced map.* The composition

$$E_q \times T \xrightarrow{\Gamma} X \xrightarrow{\varphi} X \xrightarrow{\pi^{-1}} \text{Bl}_q X$$

is a family (parametrized by  $E_q$ ) of parameterizations (all parameterized by  $T$ ) of curves, each of them passing through a point on the exceptional divisor  $E_q$ .

Regard  $E_q$  as  $\mathfrak{q} \times \mathbb{A}^1$  with coordinates  $((u, v), r)$ . The correspondence between a pair  $(z, w)$  and a point on  $E_q$ , hence the induced map  $\bar{\varphi} : E_q \dashrightarrow E_q$ , is given by

$$(z, w) \mapsto \left( \left( \frac{8z^2}{-5z^2 + w^2}, \frac{8zw}{-5z^2 + w^2} \right), \frac{25z^4 - 10w^2z^2 + w^4}{72z^5 - 80w^2z^3 + 8w^4z} \right).$$

The inverse  $\bar{\varphi}^{-1} : E_q \rightarrow E_q$  is given by

$$((u, v), r) \mapsto \left( \frac{(25u^4 - 10u^2v^2 + v^4)u}{8u(9u^4 - 10u^2v^2 + v^4)r}, \frac{(25u^4 - 10u^2v^2 + v^4)v}{8u(9u^4 - 10u^2v^2 + v^4)r} \right).$$

**3.4. Projectivized induced maps.** Given the threefold  $X = \mathbb{P}^2 \times T$  defined by  $b^2c = a^3 + 5a^2c + 4ac^2$  in  $\mathbb{P}^2 \times \mathbb{P}^2$ , we have the involution  $\varphi : X \dashrightarrow X$  by

$$([x, y, z], [a, b, c]) \mapsto ([5x^3 - xy^2 + 8x^2z, 5x^2y - y^3 + 8xyz, -2x^3 - 5x^2z + y^2z], [a, b, c]).$$

We can also explicitly spell out the defining equations of  $\psi : X \dashrightarrow X$ :

$$([x, y, z], [a, b, c]) \mapsto ([P_1(x, y, z; a, b, c), P_2(x, y, z; a, b, c), P_3(x, y, z; a, b, c)], [a, b, c]),$$

where

$$P_1 = (x^3 + 2y^2z - 4xz^2)a + (-2xyz)b + (5x^3 - xy^2 + 8x^2z)c,$$

$$P_2 = (3x^2y + 10xyz + 4yz^2)a + (-2x^3 - 10x^2z - 8xz^2)b + (5x^2y - y^3 + 8xyz)c,$$

$$P_3 = (3x^2z + 10xz^2 + 4z^3)a + (-2yz^2)b + (-2x^3 - 5x^2z + y^2z)c.$$

For the elliptic curve we have chosen, the inverse of the point  $[a, b, c]$  has coordinates  $[a, -b, c]$ . Therefore, the involution  $\chi : X \dashrightarrow X$  is defined by

$$([x, y, z], [a, b, c]) \mapsto ([P_1(x, y, z; a, -b, c), P_2(x, y, z; a, -b, c), P_3(x, y, z; a, -b, c)], [a, b, c]).$$

The process in Section ?? above yields three induced maps  $\bar{\varphi}, \bar{\psi}, \bar{\chi} : E_q \dashrightarrow E_q$  with inverses  $\bar{\varphi}^{-1}, \bar{\psi}^{-1}, \bar{\chi}^{-1} : E_q \dashrightarrow E_q$ . To analyze the dynamical property of these maps, we projectivize the exceptional divisors and homogenize the defining equations. Now we have

$$E_q \cong \mathbb{P}^2 \quad \text{and} \quad E_q \cong \mathfrak{q} \times \mathbb{P}^1,$$

and, for example,  $\bar{\varphi} : E_q \dashrightarrow E_q$  is given by

$$[x, y, z] \mapsto ([8x^2, 8xy, (-5x^2 + y^2)], [(5x^2 - y^2)^2z, 8x(x - y)(x + y)(3x - y)(3x + y)]).$$

#### 4. DYNAMICS OF INDUCED MAPS

In the following discussion, if no confusion arises, we always use the same notation for a point and the corresponding point on any blow-up; and we always use the same notation for a subvariety and all of its strict transforms.

**4.1. The map induced by  $\varphi \circ \psi$ .** Recall that the map that is conjugate to the map induced by  $\varphi \circ \psi$  is  $\Phi := \bar{\psi}^{-1} \circ \bar{\varphi} : E_q \cong \mathbb{P}^2 \dashrightarrow \mathbb{P}^2 \cong E_q$ , and it is defined by

$$[x, y, z] \mapsto \begin{bmatrix} 9x^5 - 10x^3y^2 + xy^4 + 10x^3yz - 2xy^3z, \\ 9x^4y - 10x^2y^3 + y^5 + 9x^4z - y^4z, \\ 9x^4z - 10x^2y^2z + y^4z \end{bmatrix}^T.$$

The inverse of  $\Phi$  is  $\Phi^{-1} := \bar{\varphi}^{-1} \circ \bar{\psi} : E_q \dashrightarrow E_q$ . We will show that the sequence  $\{\deg(\Phi^n)\}$  grows quadratically.

A direct computation shows that the indeterminate points of  $\Phi$  are

$$p := [0, 0, 1], p_1 := [1, 1, 0], p_2 := [1, -1, 0], p_3 := [1, 3, 0], p_4 := [1, -3, 0].$$

The indeterminate points of  $\Phi^{-1}$  are

$$p' := [0, 1, 1], p_1 := [1, 1, 0], p_2 := [1, -1, 0], p_3 := [1, 3, 0], p_4 := [1, -3, 0].$$

We write  $P := \{p, p_1, p_2, p_3, p_4, p'\}$ .

Let  $\text{Bl}_P E_q$  be the blow-up of  $E_q$  at  $P$  with exceptional divisors  $e_p, e_{p_1}, e_{p_2}, e_{p_3}, e_{p_4}, e_{p'}$ . Both the lift of  $\Phi$  and the lift of  $\Phi^{-1}$  on  $\text{Bl}_P E_q$  will still have four indeterminate points  $p'_1, p'_2, p'_3, p'_4$ ,

one on each of the exceptional divisors  $e_{p_1}, e_{p_2}, e_{p_3}, e_{p_4}$ . We write  $P' := \{p'_1, p'_2, p'_3, p'_4\}$ . In fact, the pullback of a general line in  $E_q \cong \mathbb{P}^2$  under  $\Phi$  passes through the points  $p_1, p_2, p_3, p_4$  with tangent lines

$$\begin{aligned} l_1 &: 2x - 2y + z = 0, \\ l_2 &: 2x + 2y - z = 0, \\ l_3 &: 6x - 2y + z = 0, \\ l_4 &: 6x + 2y - z = 0 \end{aligned}$$

at the points respectively. The same claim holds for  $\Phi^{-1}$ .

In fact, the map  $\Phi$  preserves a fibration on  $E_q \cong \mathbb{P}^2$  of singular quartic curves with two nodal points at  $p, p'$  and with tangent lines  $l_1, \dots, l_4$  at  $p_1, \dots, p_4$ , respectively. Now we blow-up  $\text{Bl}_P E_q$  at  $P'$  to get  $\text{Bl}_{P'} \text{Bl}_P E_q$  with exceptional divisors  $e_{p'_1}, e_{p'_2}, e_{p'_3}, e_{p'_4}$  at the corresponding points. Both  $\Phi$  and  $\Phi^{-1}$  lift to an automorphism on  $\text{Bl}_{P'} \text{Bl}_P E_q$ .

**4.2. Quadratic degree growth of  $\Phi$ .** Given the sequence of blow-ups

$$\text{Bl}_{P'} \text{Bl}_P E_q \rightarrow \text{Bl}_P E_q \rightarrow E_q,$$

we write  $\hat{\Phi}$  for the lift of  $\Phi$  on  $\text{Bl}_P E_q$ , and write  $\tilde{\Phi}$  for the lift of  $\Phi$  on  $\text{Bl}_{P'} \text{Bl}_P E_q$ . To see the degree growth of  $\Phi$ , we examine the action of  $\tilde{\Phi}$  on  $N^1(\text{Bl}_{P'} \text{Bl}_P E_q)$ . Let  $h$  be a hyperplane class. A set of (ordered) basis elements for  $N^1(\text{Bl}_{P'} \text{Bl}_P E_q)$  is

$$\{h, e_p, e_{p_1}, e_{p_2}, e_{p_3}, e_{p_4}, e_{p'}, e_{p'_1}, e_{p'_2}, e_{p'_3}, e_{p'_4}\}.$$

**Proposition 10.** *The action of  $\tilde{\Phi}$  on  $N^1(\text{Bl}_{P'} \text{Bl}_P E_q)$  is given by*

$$M := \begin{pmatrix} 5 & 4 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & -1 & 1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ -1 & -1 & 0 & 1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ -1 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & -1 & 0 \\ -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & -1 \\ -4 & -3 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & -1 & -1 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

*Proof.* The image of a general hyperplane under  $\Phi$  is a degree-5 curve passing through  $p' = [0, 1, 1]$  with multiplicity 4 and  $p_1, p_2, p_3, p_4$  with tangent lines  $l_1, l_2, l_3, l_4$  respectively. On  $\text{Bl}_P E_q$ , we have

$$\hat{\Phi}_*(h) = 5h - 4e_{p'} - \sum e_{p_i}.$$

On  $\text{Bl}_{P'} \text{Bl}_P E_q$ , we have

$$\begin{aligned} \tilde{\Phi}_*(h) &= 5h - 4e_{p'} - \sum (e_{p_i} + e_{p'_i}) - \sum e_{p'_i} \\ &= 5h - 4e_{p'} - \sum e_{p_i} - 2 \sum e_{p'_i} \end{aligned}$$

$\Phi^{-1}$  contracts a degree-4 curve to  $p = [0, 0, 1]$ ; moreover, the curve passes through the point  $p' = [0, 1, 1]$  with multiplicity 3 and through  $p_1, p_2, p_3, p_4$  with tangent lines  $l_1, l_2, l_3, l_4$  respectively. We have

$$\hat{\Phi}_*(e_p) = 4h - 3e_{p'} - \sum e_{p_i},$$

and so

$$\begin{aligned} \tilde{\Phi}_*(e_p) &= 4h - 3e_{p'} - \sum (e_{p_i} + e_{p'_i}) - \sum e_{p'_i} \\ &= 4h - 3e_{p'} - \sum e_{p_i} - 2 \sum e_{p'_i}. \end{aligned}$$

We note that  $\Phi(0, 1, 1) = [0, 0, 1]$ , and it can be verified that

$$\tilde{\Phi}_*(e_{p'}) = e_p.$$

Next, we observe that the image of a general line passing through  $p_1 = [1, 1, 0]$  is a degree-4 curve passing through the point  $p' = [0, 1, 1]$  with multiplicity 3. The image also passes through the point  $p_1$  with tangent line not equal to  $l_1 : 2x - 2y + z = 0$  and through  $p_2, p_3, p_4$  with tangent lines  $l_2, l_3, l_4$  respectively. Thus,

$$\hat{\Phi}_*(h - e_{p_1}) = 4h - 3e_{p'} - \sum e_{p_i},$$

and hence

$$(4.1) \quad \tilde{\Phi}_*(h - e_{p_1} - e_{p'_1}) = 4h - 3e_{p'} - \sum e_{p_i} - 2 \sum e_{p'_i} + e_{p'_1}.$$

In particular, the image of the line  $l_1 : 2x - 2y + z = 0$  is a degree-3 curve passing through  $p' = [0, 1, 1]$  with multiplicity 2 and  $p_2, p_3, p_4$  with tangent lines  $l_2, l_3, l_4$  respectively. So

$$(4.2) \quad \tilde{\Phi}_*(h - e_{p_1} - 2e_{p'_1}) = 3h - 2e_{p'} - \sum e_{p_i} - 2 \sum e_{p'_i} + e_{p_1} + 2e_{p'_1}.$$

Combining (4.1) and (4.2) we get

$$(4.3) \quad \tilde{\Phi}_*(e_{p'_1}) = h - e_{p'} - e_{p_1} - e_{p'_1}.$$

Similarly,

$$\tilde{\Phi}_*(e_{p'_i}) = h - e_{p'} - e_{p_i} - e_{p'_i},$$

for  $i = 2, 3, 4$ .

Finally, substituting (4.3) back into (4.1) and rearranging terms, we get

$$\begin{aligned} \tilde{\Phi}_*(e_{p_1}) &= \tilde{\Phi}_*(h) - \tilde{\Phi}_*(e_p) - \tilde{\Phi}_*(e_{p'_1}) - e_{p'_1} \\ &= h - e_{p'} - e_{p'_1} - (h - e_{p'} - e_{p_1} - e_{p'_1}) \\ &= e_{p_1}. \end{aligned}$$

Similarly, we calculate that

$$\tilde{\Phi}_*(e_{p_i}) = e_{p_i},$$

for  $i = 2, 3, 4$ . □

**Corollary 11.**  *$M$  has a  $3 \times 3$  Jordan block associated with the eigenvalue 1. Hence the sequence  $\{\deg(\Phi^n)\}_n$  grows quadratically.*



Hence, the map  $E_{\mathfrak{q}} \dashrightarrow E_{\mathfrak{q}}$  induced by  $\varphi \circ \psi$  on the exceptional divisor over the curve  $\mathfrak{q}$  in the fiber over  $q$  has a quadratic degree growth. Recall that the map  $(\varphi \circ \psi)_t$  on a general fiber has a quadratic degree growth, but the map  $(\varphi \circ \psi)_q$  is the identity map. In this way, we found the interesting dynamics on the exceptional divisor  $E_{\mathfrak{q}}$ .

Moreover, we recall that the map  $(\varphi \circ \chi \circ \varphi \circ \psi)_t$  on a general fiber has dynamical degree  $> 1$ . We expect the induced map to have  $\lambda > 1$  as well.

**4.3. The map induced by  $\varphi \circ \chi \circ \varphi \circ \psi$ .** Using the maps  $\bar{\chi}$  and  $\bar{\varphi}$  (and their inverses), we obtain another map  $\Psi := \bar{\chi}^{-1} \circ \bar{\varphi} : E_q \dashrightarrow E_q$  with the inverse map  $\Psi^{-1} := \bar{\varphi}^{-1} \circ \bar{\chi}^{-1} : E_q \dashrightarrow E_q$ .  $\Psi \circ \Phi$  is conjugate to the map induced by  $\varphi \circ \chi \circ \varphi \circ \psi$ . We will see that  $\lambda(\Psi \circ \Phi) = 16$ , but we need to understand the map  $\Psi$  at first.

The map  $\Psi$  behaves almost the same as  $\Phi$ . The indeterminate points of  $\Psi$  are

$$p = [0, 0, 1], p_1 = [1, 1, 0], p_2 = [1, -1, 0], p_3 = [1, 3, 0], p_4 = [1, -3, 0].$$

Moreover, the indeterminate points of  $\Psi^{-1}$  are

$$p'' := [0, -1, 1], p_1 = [1, 1, 0], p_2 = [1, -1, 0], p_3 = [1, 3, 0], p_4 = [1, -3, 0].$$

The pullback of a general line in  $E_q$  under both  $\Psi$  passes through the points  $p_1, p_2, p_3, p_4$  with tangent lines

$$\begin{aligned} 2x - 2y - z &= 0 \\ 2x + 2y + z &= 0 \\ 6x - 2y - z &= 0 \\ 6x + 2y + z &= 0 \end{aligned}$$

at  $p_1, p_2, p_3, p_4$ , respectively. The same claim holds for  $\Psi^{-1}$ . Therefore, if we blow-up  $E_q$  at  $Q := \{p, p'', p_1, p_2, p_3, p_4\}$ , then both the lifts  $\hat{\Psi}$  and  $\hat{\Psi}^{-1}$  on  $\text{Bl}_Q E_q$  will have an indeterminate point  $p'_i$  at the exceptional divisors  $e_{p_i}$  for each  $i = 1, \dots, 4$ . If we blow-up again at  $Q' := \{p'_1, p'_2, p'_3, p'_4\}$ , then  $\hat{\Psi}$  and  $\hat{\Psi}^{-1}$  will be regularized on  $\text{Bl}_{Q'} \text{Bl}_Q E_q$  and lift to  $\tilde{\Psi}$  and  $\tilde{\Psi}^{-1}$ . The action of  $\tilde{\Psi}$  on  $N^1(\text{Bl}_{Q'} \text{Bl}_Q E_q)$  can be represented by the same matrix (with respect to a different basis, of course).

Now we are ready to compute the dynamical degree of  $\Psi \circ \Phi$ . Let us blow-up  $E_q$  at

$$R := \{p, p', p'', p_1, p_2, p_3, p_4\}$$

to get  $\text{Bl}_R \mathbb{P}^2$  and then at

$$R' := \{p'_1, p'_2, p'_3, p'_4, p''_1, p''_2, p''_3, p''_4\}$$

to get  $\text{Bl}_{R'} \text{Bl}_R E_q$ . Then  $N^1(\text{Bl}_{R'} \text{Bl}_R E_q)$  has rank 16 with an ordered basis

$$\{h, e_p, e_{p'}, e_{p''}, e_{p_1}, e_{p_2}, e_{p_3}, e_{p_4}, e_{p'_1}, e_{p'_2}, e_{p'_3}, e_{p'_4}, e_{p''_1}, e_{p''_2}, e_{p''_3}, e_{p''_4}\}.$$

Let  $\tilde{\Phi}$  and  $\tilde{\Psi}$  denote the lifts of  $\Phi$  and  $\Psi$  on  $\text{Bl}_{R'} \text{Bl}_R E_q$ . The maps  $\tilde{\Phi}$  (resp.  $\tilde{\Psi}$ ) is an isomorphism (which contracts no curves) on  $\text{Bl}_{P'} \text{Bl}_P E_q$  (resp.  $\text{Bl}_{Q'} \text{Bl}_Q E_q$ ). Nevertheless, some curves on  $\text{Bl}_{R'} \text{Bl}_R E_q$  will be contracted by  $\tilde{\Phi}$  and  $\tilde{\Psi}$ . The map  $\tilde{\Phi}$  contracts the curves  $e_{p''}, e_{p'_1}, e_{p'_2}, e_{p'_3}, e_{p'_4}$ , and the map  $\tilde{\Psi}$  contracts the curves  $e_{p'}, e_{p'_1}, e_{p'_2}, e_{p'_3}, e_{p'_4}$ .

**Proposition 12.** *For any curve  $C$  contracted by  $\tilde{\Phi}$ , we have  $\tilde{\Phi}(C) \notin \text{Ind}(\tilde{\Psi})$ . We conclude that*

$$(\tilde{\Psi} \circ \tilde{\Phi})_* = \tilde{\Psi}_* \circ \tilde{\Phi}_*.$$

*Proof.* Since  $\Phi(p'') = \Phi(0, -1, 1) = [0, -2, 1] =: p_{-2}$ ,  $\tilde{\Phi}(e_{p''}) = p_{-2}$  on  $\text{Bl}_{R'} \text{Bl}_R E_q$ . Moreover,  $\tilde{\Psi}$  is defined at  $p_{-2}$ , as  $\Psi$  is defined at  $[0, -2, 1]$ .

What happens at  $e_{p'_i}$ , for  $i = 1, \dots, 4$ , is similar, and we present only the local computations around  $p_1$ . The blow-up  $\text{Bl}_{p_1} E_q$  of  $E_q$  at  $p_1 = [1, 1, 0]$  can be defined in  $E_q \times \mathbb{P}^1 \cong \mathbb{P}^2 \times \mathbb{P}^1$  (with coordinates  $([x, y, z], [s, t])$ ) by the equation

$$(x - y)s = zt$$

with the exceptional divisor

$$e_{p_1} = \{([1, 1, 0], [s, t])\} \subset \text{Bl}_{p_1} E_q.$$

In terms of the coordinates on  $e_{p_1}$ ,  $p''_1 = ([1, 1, 0], [2, 1])$ . Let  $\Phi_1$  and  $\Psi_1$  be the lifts of  $\Phi$  and  $\Psi$  on  $\text{Bl}_{p_1} E_q$ . The restriction  $\Phi_1|_{e_{p_1}} : e_{p_1} \rightarrow e_{p_1}$  is given by

$$([1, 1, 0], [s, t]) \mapsto ([1, 1, 0], [2s, 2t - s]).$$

So  $\Phi_1([1, 1, 0], [2, 1]) = ([1, 1, 0], [1, 0])$ . This implies that  $\tilde{\Phi}$  contracts  $e_{p''_1}$  to the point corresponding to  $([1, 1, 0], [1, 0])$ . Moreover,  $\Psi_1$  is defined at  $([1, 1, 0], [1, 0])$ , so  $\tilde{\Psi}$  is defined at the point corresponding to  $([1, 1, 0], [1, 0])$ .  $\square$

Consequently,  $\tilde{\Psi} \circ \tilde{\Phi}$  contracts the curves

$$e_{p''}, e_{p''_1}, e_{p''_2}, e_{p''_3}, e_{p''_4}, (\tilde{\Phi}^{-1})(e_{p'}), (\tilde{\Phi}^{-1})(e_{p'_1}), (\tilde{\Phi}^{-1})(e_{p'_2}), (\tilde{\Phi}^{-1})(e_{p'_3}), (\tilde{\Phi}^{-1})(e_{p'_4}).$$

on  $\text{Bl}_{R'} \text{Bl}_R E_q$ , where  $(\tilde{\Phi}^{-1})(C)$  means the proper transform of a curve  $C$  under the inverse of  $\tilde{\Phi}$ .

**4.4. Algebraic stability of  $\tilde{\Psi} \circ \tilde{\Phi}$ .** We prove the algebraic stability of  $\tilde{\Psi} \circ \tilde{\Phi}$  on  $\text{Bl}_{R'} \text{Bl}_R E_q$ .

**Proposition 13.** *The map  $\tilde{\Psi} \circ \tilde{\Phi}$  is algebraically stable on  $\text{Bl}_{R'} \text{Bl}_R E_q$ .*

*Proof.* By Theorem 2, it suffices to show that for any curve  $C$  contracted by  $\tilde{\Psi} \circ \tilde{\Phi}$ ,  $(\tilde{\Psi} \circ \tilde{\Phi})(C)$  is a fixed point of  $\tilde{\Psi} \circ \tilde{\Phi}$ . We first deal with the curves  $e_{p''}, e_{p''_1}, e_{p''_2}, e_{p''_3}, e_{p''_4}$ .

For  $e_{p''}$ , we note that  $(\Psi \circ \Phi)(p'') = p''$ . So  $(\tilde{\Psi} \circ \tilde{\Phi})(e_{p''}) \subset e_{p''}$ . We can check that there is a curve (other than  $e_{p''}$ ) whose image under  $\tilde{\Psi} \circ \tilde{\Phi}$  is  $e_{p''}$ . So we cannot have  $(\tilde{\Psi} \circ \tilde{\Phi})(e_{p''}) = e_{p''}$  as  $\tilde{\Psi} \circ \tilde{\Phi}$  is birational. Thus  $(\tilde{\Psi} \circ \tilde{\Phi})(e_{p''})$  is a point on  $e_{p''}$ .

To see the points  $p''_i$ , we need to blow-up  $\mathbb{P}^2$  at  $p_i$  for  $i = 1, \dots, 4$ . Let  $\hat{\Phi}$  and  $\hat{\Psi}$  (resp.  $\hat{\Phi}^{-1}$  and  $\hat{\Psi}^{-1}$ ) be the lifts of  $\Psi$  and  $\Phi$  (resp.  $\hat{\Phi}^{-1}$  and  $\hat{\Psi}^{-1}$ ) on  $\text{Bl}_R E_q$ . We first claim that  $(\hat{\Psi} \circ \hat{\Phi})(p''_i) = p''_i$ . Again, we exhibit the calculation around  $p_1$  and omit the similar local calculations at  $p_2, p_3, p_4$ . Recall that in the proof of Proposition 12, we have  $p'_1 = ([1, 1, 0], [-2, 1])$  in terms of the coordinates on  $e_{p_1}$  and the restriction  $\Phi_1|_{e_{p_1}} : e_{p_1} \rightarrow e_{p_1}$  is given by

$$([1, 1, 0], [s, t]) \mapsto ([1, 1, 0], [2s, 2t - s]).$$

Moreover, we have  $p''_1 = ([1, 1, 0], [2, 1])$  and the restriction  $\Psi_1|_{e_{p_1}} : e_{p_1} \rightarrow e_{p_1}$  is given by

$$([1, 1, 0], [s, t]) \mapsto ([1, 1, 0], [2s, 2t + s]).$$

Therefore,  $\Psi_1 \circ \Phi_1$  fixes each point on  $e_{p_1}$  as long as it is defined at the point. In particular,  $(\Psi_1 \circ \Phi_1)(p''_1) = p''_1$ .

Now, for each  $i = 1, \dots, 4$ , we have  $(\tilde{\Psi} \circ \tilde{\Phi})(e_{p''_i}) \subset e_{p''_i}$ . There is a curve (other than  $e_{p''_i}$ ) whose image under  $(\tilde{\Psi} \circ \tilde{\Phi})$  is  $e_{p''_i}$ . So  $(\tilde{\Psi} \circ \tilde{\Phi})(e_{p''_i})$  is a point on  $e_{p''_i}$ .

We next consider the curves  $(\tilde{\Phi}^{-1})(e_{p'}), (\tilde{\Phi}^{-1})(e_{p'_1}), (\tilde{\Phi}^{-1})(e_{p'_2}), (\tilde{\Phi}^{-1})(e_{p'_3}), (\tilde{\Phi}^{-1})(e_{p'_4})$ . We note that  $\tilde{\Psi} \circ \tilde{\Phi}$  fixes the strict transform of the line  $\{x = 0\}$  and exceptional divisors  $e_{p'_1}, e_{p'_2}, e_{p'_3}, e_{p'_4}$

pointwise (wherever  $\tilde{\Psi} \circ \tilde{\Phi}$  is defined). Moreover,  $\tilde{\Psi} \circ \tilde{\Phi}$  contracts  $(\tilde{\Phi}^{-1})(e_{p'})$  to a point on the strict transform of  $\{x = 0\}$  and  $(\tilde{\Phi}^{-1})(e_{p'_i})$  to a point on  $e_{p'_i}$  for each  $i = 1, \dots, 4$ .  $\square$

**4.5. Dynamical degree of  $\Psi \circ \Phi$ .** The blow-up  $\text{Bl}_{R'} \text{Bl}_R E_q$  is a birational model on which  $\tilde{\Psi} \circ \tilde{\Phi}$  is algebraically stable. In order to see the action of  $\tilde{\Psi} \circ \tilde{\Phi}$  on  $N^1(\text{Bl}_{R'} \text{Bl}_R E_q)$ , we compute the matrices that represent the action of  $\tilde{\Phi}$  and  $\tilde{\Psi}$  and then take their product.

**Proposition 14.** *The action of  $\tilde{\Phi}$  on  $N^1(\text{Bl}_{R'} \text{Bl}_R E_q)$  is given by*

$$M_{\Phi} := \begin{pmatrix} 5 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ -4 & -3 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

and the action of  $\tilde{\Psi}$  on  $N^1(\text{Bl}_{R'} \text{Bl}_R E_q)$  is given by

$$M_{\Psi} := \begin{pmatrix} 5 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ -1 & -1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ -1 & -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ -1 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ -1 & -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ -1 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ -4 & -3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & -1 & -1 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ -2 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

*Proof.* We will show the action of  $\tilde{\Phi}$  is given by  $M_{\Phi}$ , and the proof for  $\tilde{\Psi}$  is similar.

First, we note that

$$\tilde{\Phi}_*(e_{p''}) = \tilde{\Phi}_*(e_{p'_i}) = 0$$

for  $i = 1, \dots, 4$ .

Continuing the arguments in the proof of Proposition 10, we have

$$\begin{aligned}\tilde{\Phi}_*(h) &= 5h - 4e_{p'} - \sum e_{p_i} - 2 \sum e_{p'_i} - \sum e_{p''_i} \\ \tilde{\Phi}_*(e_p) &= 4h - 3e_{p'} - \sum e_{p_i} - 2 \sum e_{p'_i} - \sum e_{p''_i} \\ \tilde{\Phi}_*(e_{p'}) &= e_p \\ \tilde{\Phi}_*(e_{p_i} + e_{p''_i}) &= e_{p_i} + e_{p''_i} \\ \tilde{\Phi}_*(e_{p'_i}) &= h - e_{p'} - e_{p_i} - e_{p'_i} - e_{p''_i}.\end{aligned}$$

Also

$$\tilde{\Phi}_*(e_{p_i}) = e_{p_i} + e_{p''_i} - \tilde{\Phi}_*(e_{p''_i}) = e_{p_i} + e_{p''_i}.$$

□

Proposition 12 suggests that we can use the product of  $M_\Psi$  and  $M_\Phi$  to represent the action of  $\tilde{\Psi} \circ \tilde{\Phi}$  on  $N^1(\text{Bl}_{R'} \text{Bl}_R E_q)$ . The spectral radius of  $M_\Psi M_\Phi$  is 16. By Theorem 3, we obtain the following result:

**Proposition 15.** *The induced map  $\Psi \circ \Phi$  on the exceptional divisor by  $\chi \circ \varphi \circ \psi \circ \varphi$  has  $\lambda(\Psi \circ \Phi) = 16$ . In particular,  $\Psi \circ \Phi$  is not regularizable.*

Thus, the conjugate map  $E_q \dashrightarrow E_q$  induced by  $\varphi \circ \chi \circ \varphi \circ \psi$  has  $\lambda = 16 > 1$ . This proves the Theorem 1, and we again found the interesting dynamics on the exceptional divisor  $E_q$  in the fiber over  $q$ .

In the attached SageMath code, we compute an explicit equation for the map  $\Psi \circ \Phi$ . We show that the dynamical degree  $\lambda = 16$  given here is consistent with the direct approximations of the  $\lambda$  given by assuming the Kawaguchi-Silverman conjecture and computing the height growth along a few orbits.

## REFERENCES

1. B. M. Bekker and Yu G. Zarhin, *Division by 2 of rational points on elliptic curves*, St. Petersburg mathematical journal **29** (2018), no. 4, 683–713 (English).
2. Jérémy Blanc, *On the inertia group of elliptic curves in the Cremona group of the plane*, Michigan Mathematical Journal **56** (2008), no. 2, 315 – 330.
3. Jérémy Blanc, *Dynamical degrees of (pseudo)-automorphisms fixing cubic hypersurfaces*, Indiana University Mathematics Journal **62** (2013), no. 4, 1143–1164.
4. Anna Bot, *On automorphisms of rational surfaces*, Master’s thesis, ETH Zürich, Zürich, Switzerland, 2021.
5. Nguyen-Bac Dang, *Degrees of iterates of rational maps on normal projective varieties*, Proceedings of the London Mathematical Society **121** (2020), no. 5, 1268–1310.
6. Jeffery Diller and Charles Favre, *Dynamics of bimeromorphic maps of surfaces*, American Journal of Mathematics **123** (2001), 1135 – 1169.
7. M. H. Gizatullin, *The decomposition, inertia and ramification groups in birational geometry*, Vieweg+Teubner Verlag, Wiesbaden, 1994.
8. Xie Junyi, *Periodic points of birational transformations on projective surfaces*, Duke Mathematical Journal **164** (2015), no. 5, 903 – 932.
9. John Lesieutre, *A pathology of asymptotic multiplicity in the relative setting*, Mathematical Research Letters **23** (2016), no. 5, 1433–1451.
10. Tuyen Trung Truong, *Relative dynamical degrees of correspondences over a field of arbitrary characteristic*, Journal für die reine und angewandte Mathematik (Crelles Journal) **2020** (2020), no. 758, 139–182.

## ACKNOWLEDGEMENT

I extend my sincere gratitude to my advisor, John Lesieutre, for proposing this problem and for his invaluable discussions and guidance throughout this work. I also thank Anna Bot, Junyi Xie, Luen-Chau Li, Tatiana Bandman, and Yuri Zarhin for their valuable discussions. This work was partially supported by NSF-DMS grant 2142966.

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16801, USA  
*Email address:* `qvj5031@psu.edu`