ON DEGENERATE WHITTAKER SPACE FOR $GL_4(\mathfrak{o}_2)$

ANKITA PARASHAR AND SHIV PRAKASH PATEL

ABSTRACT. Let \mathfrak{o}_2 be a finite principal ideal local ring of length 2. For a representation π of $GL_4(\mathfrak{o}_2)$, the degenerate Whittaker space $\pi_{N,\psi}$ is a representation of $GL_2(\mathfrak{o}_2)$. We describe $\pi_{N,\psi}$ explicitly for an irreducible strongly cuspidal representation π of $GL_4(\mathfrak{o}_2)$. This description verifies a special case of a conjecture of Prasad. We also prove that $\pi_{N,\psi}$ is a multiplicity free representation.

1. INTRODUCTION

Let *F* be a finite unramified extension of \mathbb{Q}_p or $\mathbb{F}_p((t))$. Let \mathfrak{o} be the ring of integers of *F* with a uniformizer \mathfrak{O} . For any positive integer *l*, let $\mathfrak{o}_l := \mathfrak{o}/(\mathfrak{O}^l)$. Then \mathfrak{o}_l is a finite principal ideal local ring of length *l*. Note that $\mathfrak{o}/(\mathfrak{O}) \cong \mathbb{F}_q$ is a finite field of order $q = p^f$ of characteristic p > 0. If the characteristic of *F* is zero then \mathfrak{o}_l is the the ring of Witt vectors over \mathbb{F}_q of length *l* and, if the characteristic of *F* is *p* then \mathfrak{o}_l is isomorphic to $\mathbb{F}_q[t]/(t^l)$. We will be working with both the rings simultaneously and our proofs are uniform for both the cases. We assume that $p \neq 2$.

Let $GL_{2n}(\mathfrak{o}_l)$ be the group of invertible $2n \times 2n$ matrices over the ring \mathfrak{o}_l . Let

$$P_{n,n} = \left\{ \begin{pmatrix} g & X \\ 0 & g' \end{pmatrix} : g, g' \in \operatorname{GL}_n(\mathfrak{o}_l), X \in M_n(\mathfrak{o}_l) \right\} \text{ and}$$
$$N = \left\{ \begin{pmatrix} I & X \\ 0 & I \end{pmatrix} : X \in M_n(\mathfrak{o}_l) \right\} \cong M_n(\mathfrak{o}_l).$$

Let $\psi_0 : \mathfrak{o}_l \to \mathbb{C}^{\times}$ be a non-trivial additive character which is also non-trivial when restricted to the subgroup $\varpi^{l-1}\mathfrak{o}_l$ for l > 1. Let $\psi : M_n(\mathfrak{o}_l) \to \mathbb{C}^{\times}$ be the additive character defined by $\psi(X) = \psi_0(\operatorname{tr}(X))$ for all $X \in M_n(\mathfrak{o}_l)$, where $\operatorname{tr}(X)$ denotes the trace of X. Since $M_n(\mathfrak{o}_l) \cong N$, we treat ψ as a character of N as well. Let (π, V) be an irreducible representation of $\operatorname{GL}_{2n}(\mathfrak{o}_l)$. Define $V_{N,\psi} := \{v \in V : \pi(X)v = \psi(X)v \; \forall X \in$ $N\}$ which is the subspace of V on which N operates by the character ψ . The space $V_{N,\psi}$ is called the (N,ψ) -Whittaker space of π , which we call degenerate Whittaker space of π . Since $\operatorname{tr}(gXg^{-1}) = \operatorname{tr}(X)$ for all $g \in \operatorname{GL}_n(\mathfrak{o}_l)$ and $X \in M_n(\mathfrak{o}_l)$, it follows that $\operatorname{GL}_n(\mathfrak{o}_l)$ sitting inside the subgroup $P_{n,n}$ as $g \mapsto \begin{pmatrix} g & 0 \\ 0 & g \end{pmatrix}$ stabilizes the character ψ of N. Therefore, $\operatorname{GL}_n(\mathfrak{o}_l)$ operates on the space $V_{N,\psi}$ giving rise to a representation of $\operatorname{GL}_n(\mathfrak{o}_l)$ which we denote by $\pi_{N,\psi}$. For a representation (π, V) , we often omit V and write only π . In this paper, we will be concerned with the following question.

Question 1.1. Given an irreducible representation π of $GL_{2n}(\mathfrak{o}_l)$, what is the degenerate Whittaker space $\pi_{N,\psi}$ as a representation of $GL_n(\mathfrak{o}_l)$?

²⁰²⁰ Mathematics Subject Classification. Primary: 20G25, Secondary: 20G05, 20C15.

Key words and phrases. Degenerate Whittaker space, Prasad's conjecture, Regular representations.

The above question is a variant of the question considered by Prasad [16] for l = 1. Prasad's work has inspired several related works, including ours. For example, when l = 1 Balasubramanian-Khurana [3, 4] consider a variant by taking different characters ψ for n = 2, 3 and Balasubramanian-Dangodara-Khurana [2] consider analogous variant for general n. Another variant, when the ring o_l is replaced by a non-Archimedean local field, this question has been considered by Prasad [17] for GL₄ and more recently Pandey-Venketasubramanian [12] for Sp_4 .

Now we briefly describe the work of Prasad [16] for l = 1. Note that $\mathfrak{o}_1 \cong \mathbb{F}_q$. In this case, (N, ψ) -degenerate Whittaker space $\pi_{N,\psi}$ is a representation of $\operatorname{GL}_n(\mathbb{F}_q)$. Recall that for any positive integer n, we have $\mathbb{F}_{q^n}^{\times} \hookrightarrow \operatorname{GL}_n(\mathbb{F}_q)$ and the embedding is unique up to conjugation. By the work of Green [8], it is well known that every irreducible cuspidal representation of $\operatorname{GL}_n(\mathbb{F}_q)$ is associated with a primitive character of $\mathbb{F}_{q^n}^{\times}$. Here is the theorem of Prasad [16, Theorem 1].

Theorem 1.2. Let π be an irreducible cuspidal representation of $\operatorname{GL}_{2n}(\mathbb{F}_q)$ obtained from a primitive character $\theta : \mathbb{F}_{q^{2n}}^{\times} \to \mathbb{C}^{\times}$. Then, as a representation of $\operatorname{GL}_n(\mathbb{F}_q)$,

$$\pi_{N,\psi} \cong \operatorname{Ind}_{\mathbb{F}_{q^n}^{\times}}^{\operatorname{GL}_n(\mathbb{F}_q)}(\theta|_{\mathbb{F}_{q^n}^{\times}}).$$

For n = 1 and π a regular representation of $GL_2(\mathfrak{o}_2)$, the Question 1.1 is answered in a work of the second author with P. Singla [14, 15]. In this case, $\pi_{N,\psi}$ is a representation of $GL_1(\mathfrak{o}_l)$ which is isomorphic to the center of $GL_2(\mathfrak{o}_l)$. More precisely, we have the following result [15, Theorem 1.1].

Theorem 1.3. Let π be an irreducible regular representation of $GL_2(\mathfrak{o}_l)$ with central character ω_{π} , then $\pi_{N,\psi}$ is one dimensional and $\pi_{N,\psi} \cong \omega_{\pi}$.

A more general version of Theorem 1.2 has been conjectured by Prasad which we state below. Let *L* (respectively, *E*) be an unramified extensions of *F* of degree *n* (respectively, 2*n*) and \mathfrak{O} (respectively, \mathcal{O}) the ring of integers of *L* (respectively, *E*). Write $\mathfrak{O}_l = \mathfrak{O}/(\mathfrak{O}^l)$ and $\mathcal{O}_l = \mathcal{O}/(\mathfrak{O}^l)$ then $\mathfrak{O}_l^{\times} \hookrightarrow \operatorname{GL}_n(\mathfrak{o}_l)$ and $\mathcal{O}_l^{\times} \hookrightarrow \operatorname{GL}_{2n}(\mathfrak{o}_l)$. We also have $\mathfrak{o}_l \subset \mathfrak{O}_l \subset \mathcal{O}_l$. It is known from Lusztig [11] and Aubert-Onn-Prasad [1] that a strongly cuspidal representation of $\operatorname{GL}_{2n}(\mathfrak{o}_l)$ is obtained from a strongly primitive character $\theta : \mathcal{O}_l^{\times} \to \mathbb{C}^{\times}$ (see Section 2 for more details).

Conjecture 1.4 (Prasad). Let π be an irreducible strongly cuspidal representation of $\operatorname{GL}_{2n}(\mathfrak{o}_l)$ associated to a strongly primitive character $\theta : \mathcal{O}_l^{\times} \to \mathbb{C}^{\times}$. Then as a representation of $\operatorname{GL}_n(\mathfrak{o}_l)$, we have

$$\pi_{N,\psi} \cong \operatorname{Ind}_{\mathfrak{O}_l^{\times}}^{\operatorname{GL}_n(\mathfrak{o}_l)}(\theta|_{\mathfrak{O}_l^{\times}}).$$

We prove the above conjecture of Prasad for n = 2, l = 2, which is one of the main results of this paper.

Theorem 1.5. *The Conjecture* 1.4 *is true for* n = 2 *and* l = 2*.*

We would like to mention that a few steps, but not all, in the proof of this theorem also work for general n and l = 2. Since, we could not achieve all the steps to prove the above conjecture of Prasad for general n (even for l = 2), we restrict ourselves to the case of n = 2 and l = 2.

In addition, we prove the following multiplicity one result.

Theorem 1.6. Let π be a strongly cuspidal representation of $GL_4(\mathfrak{o}_2)$. Then $\pi_{N,\psi}$ is multiplicity free representation of $GL_2(\mathfrak{o}_2)$, *i.e.* for any irreducible representation σ of $GL_2(\mathfrak{o}_2)$ we have

$$\dim \left(\operatorname{Hom}_{\operatorname{GL}_2(\mathfrak{o}_2)} \left(\pi_{N, \psi}, \sigma \right) \right) \leq 1.$$

The above theorem can be considered as a variant of a theorem of Rallis for *p*-adic groups [17, Theorem 1]. The theorem of Rallis is indeed a Bessel model case of Gan-Gross-Prasad (GGP) conjecture [6]. More precisely, it is the case of the GGP conjectures when one considers the restriction of representations of the group SO(6), which is closely related to GL₄ to the subgroup SO(3) which is closely related to GL₂. The analogous multiplicity one theorem is also true in Prasad's case for n = 2, which is a consequence of Theorem 1.2. Our proof of Theorem 1.5 uses the above multiplicity one result.

We now give an outline of the paper. Our proofs depend on the construction of the regular representations of $GL_n(\mathfrak{o}_l)$ as given by Hill [9] and Krakovski-Onn-Singla [10] which we recall in Section 2. In particular, an irreducible strongly cuspidal representation of $GL_n(\mathfrak{o}_2)$ is induced from certain subgroup, say *T*. Our strategy here is to restrict the representation π to the $P_{2,2}$ subgroup which we can describe by Mackey theory. We write an exhaustive set Ω of representatives for the double cosets in $T \setminus GL_4(\mathfrak{o}_2)/P_{2,2}$ in Section 4. We use these representatives for many calculations. In fact, $\pi|_{P_{2,2}} \cong \bigoplus_{s} \pi^{\delta}$ where δ varies over a set of representatives of the double cosets

mentioned above. Then, for every $\delta \in \Omega$ we describe the representation $\pi_{N,\psi}^{\delta}$ as a representation of $\operatorname{GL}_2(\mathfrak{o}_2)$. In Section 4, we compute the dimension of $\pi_{N,\psi}$ by computing the dimension of every $\pi_{N,\psi}^{\delta}$. In Section 5, we write down the character of $\pi_{N,\psi}^{\delta}$ for every δ . In Section 6, we describe all the characters of $I + \omega M_2(\mathfrak{o}_2)$ appearing in $\pi_{N,\psi}^{\delta}$ when restricted to $I + \omega M_2(\mathfrak{o}_2)$. There are three types of regular representations of $\operatorname{GL}_2(\mathfrak{o}_2)$ as mentioned in Table 1 which helps us to analyse their occurrence in $\pi_{N,\psi}^{\delta}$ and therefore in $\pi_{N,\psi}$ one by one. In fact, we describe all the irreducible representations of $\operatorname{GL}_2(\mathfrak{o}_2)$ which appear in $\pi_{N,\psi}$ without appealing to the induced representation $\Pi := \operatorname{Ind}_{\mathfrak{D}_2^{\times}}^{\operatorname{GL}(\mathfrak{o}_2)}(\theta|_{\mathfrak{D}_2^{\times}})$. In this section, we also prove the multiplicity one result mentioned above in Theorem 1.6. In Section 7, we describe all the irreducible appear in $\pi_{N,\psi}$ with the same multiplicity (in fact, multiplicity one), which in turn proves Theorem 1.5.

It should be noted that the proof of Prasad's theorem in [16] is indirect in the sense that there is no explicit isomorphism between $\pi_{N,\psi}$ and $\operatorname{Ind}_{\mathbb{F}_{q^n}}^{\operatorname{GL}_n(\mathbb{F}_q)}(\theta|_{\mathbb{F}_{q^n}})$, instead it is proved that the characters of these two representations coincide. We also prove Theorem 1.5 indirectly in the similar sense that we do not give any explicit isomorphism between $\pi_{N,\psi}$ and Π . The proof in [16] uses character theory, while our proof is a consequence of Mackey theory and Clifford theory since the construction of π is totally explicit.

2. Preliminaries

Let *F* be a finite unramified extension of \mathbb{Q}_p or $\mathbb{F}_p((t))$, and let \mathfrak{o} be its ring of integers with a uniformizer ω . For any positive integer *l*, let $\mathfrak{o}_l = \mathfrak{o}/(\omega^l)$. Note that

 $\mathfrak{o}/(\varpi) \cong \mathbb{F}_q$ is a finite field of order $q = p^f$ of characteristic p > 0. We assume $p \neq 2$. For any integer m with $1 \leq m < l$, we have the natural quotient map of rings $\mathfrak{o}_l \to \mathfrak{o}_m$. This quotient map induces a surjective group homomorphism $\operatorname{GL}_n(\mathfrak{o}_l) \to \operatorname{GL}_n(\mathfrak{o}_m)$ for any positive integer n. For any $g \in \operatorname{GL}_n(\mathfrak{o}_l)$, $h \in \operatorname{GL}_n(\mathfrak{o}_m)$ we write $\overline{g} \in \operatorname{GL}_n(\mathfrak{o}_m)$ for the image of g under the natural map and $\tilde{h} \in \operatorname{GL}_n(\mathfrak{o}_l)$ for any element in the inverse image of h under the natural map. Define K_l^m to be the kernel of the natural map $\operatorname{GL}_n(\mathfrak{o}_l) \to \operatorname{GL}_n(\mathfrak{o}_m)$ which is also called the m-th principal congruence subgroup of $\operatorname{GL}_n(\mathfrak{o}_l)$. Note that K_l^m are normal subgroups and, for $m \geq l/2$ the subgroups K_l^m are abelian. Thus we have the following natural filtration of subgroups

$$\{I\} \subset K_l^{l-1} \subset \cdots \subset K_l^{m-1} \subset K_l^m \subset \cdots \subset K_l^1 \subset \operatorname{GL}_n(\mathfrak{o}_l).$$

For $1 \le m < l$, we have $K_l^m / K_l^{m+1} \cong (M_n(\mathbb{F}_q), +)$. Let $M_n(\mathfrak{o}_l)$ denote the Pontryagin dual group of the group $(M_n(\mathfrak{o}_l), +)$. Fix a non-trivial additive character $\psi_0 : \mathfrak{o}_l \to \mathbb{C}^{\times}$ such that $\psi_0|_{\omega^{l-1}\mathfrak{o}_l} \ne 1$. For every $A \in M_n(\mathfrak{o}_l)$, define $\phi_A \in \widehat{M_n(\mathfrak{o}_l)}$ by $\phi_A(B) = \psi_0(\operatorname{tr}(AB))$, where tr denote the trace. The map $A \mapsto \phi_A$ defines a group isomorphism between $M_n(\mathfrak{o}_l)$ and $\widehat{M_n(\mathfrak{o}_l)}$. Note that this isomorphism depends on the choice of the character ψ_0 .

For any $1 \leq i < l$, $\mathfrak{o}_{l-i} \cong \omega^i \mathfrak{o}_l \subset \mathfrak{o}_l$ and there is a natural surjection $\mathfrak{o}_l \to \mathfrak{o}_{l-i}$ written as $x \mapsto \bar{x}$. Given a character $\psi_0 : \mathfrak{o}_l \to \mathbb{C}^{\times}$, its restriction to the subgroup $\omega^i \mathfrak{o}_l$ can be seen as character of \mathfrak{o}_{l-i} again denoted by ψ_0 satisfying the following property $\psi_0(\omega^i a) = \psi_0(\bar{a})$ for all $a \in \mathfrak{o}_l$.

2.1. Regular representations of $GL_n(\mathfrak{o}_l)$.

- **Definition 2.1.** (a) Let $x \in M_n(\mathbb{F}_q)$. Then x is called regular if its characteristic polynomial is the same as its minimal polynomial. Moreover, x is called regular elliptic if its characteristic polynomial is irreducible.
 - (b) For $x \in M_n(\mathbb{F}_q)$, the character $\phi_x : M_n(\mathbb{F}_q) \to \mathbb{C}^{\times}$ is called regular (respectively, regular elliptic) if x is a regular (respectively, regular elliptic) element.
 - (c) An irreducible representation of $\operatorname{GL}_n(\mathfrak{o}_l)$ is called regular (respectively, strongly cuspidal) if its restriction to $K_l^{l-1} \cong M_n(\mathbb{F}_q)$ contains a character which is regular (respectively, regular elliptic).

In the study of representations of $GL_n(\mathfrak{o}_l)$ the regular representations turn out to be basic objects. The regular representations of $GL_n(\mathfrak{o}_l)$ can be constructed using Clifford theory, see Gérardin [7], Hill [9], Krakovski-Onn-Singla [10], Stasinski [18], Stasinski-Stevens [19]. This construction is more intricate when l is odd, see [10]. Now, we briefly describe the construction of regular representations of $GL_n(\mathfrak{o}_l)$ for even l.

Theorem 2.2. Consider the group $\operatorname{GL}_n(\mathfrak{o}_l)$ for an even integer l > 1 and set m = l/2. Let $\rho : K_l^m \to \mathbb{C}^{\times}$ be a character such that $\rho|_{K_l^{l-1}}$ is a regular character and, let the inertia group of ρ be $I(\rho) := \{g \in \operatorname{GL}_n(\mathfrak{o}_l) : \rho(g^{-1}xg) = \rho(x) \forall x \in K_l^m\}$. Then the following hold. (1) The character ρ extends to $I(\rho)$ and let $\tilde{\rho}$ be an extension of ρ to $I(\rho)$. Then $\operatorname{Ind}_{I(\rho)}^{\operatorname{GL}_n(\mathfrak{o}_l)}\tilde{\rho}$ is

an irreducible regular representation of $GL_n(\mathfrak{o}_l)$.

(2) Suppose π is a regular representation of $\operatorname{GL}_n(\mathfrak{o}_l)$ such that ρ appears in $\pi|_{K_l^m}$. Then, there exists a character $\tilde{\rho}$ of $I(\rho)$ which extends ρ such that $\pi = \operatorname{Ind}_{I(\rho)}^{\operatorname{GL}_n(\mathfrak{o}_l)} \tilde{\rho}$.

Remark 2.3. Let l be even and l = 2m. Then, $K_l^m \cong M_n(\mathfrak{o}_m)$ and any character of $M_n(\mathfrak{o}_m)$ is ϕ_B for some $B \in M_n(\mathfrak{o}_m)$. Let π be an irreducible regular representation $\operatorname{GL}_n(\mathfrak{o}_l)$ such that ϕ_B appears in $\pi|_{K_l^m}$. It can be seen by Clifford theory that $\phi_{B'}$ appears in $\pi|_{K_l^m}$ if and only if $B' = gBg^{-1}$ for some $g \in \operatorname{GL}_n(\mathfrak{o}_m)$.

2.2. **Representations of** $GL_2(\mathfrak{o}_2)$. We summarise all the irreducible representations of $GL_2(\mathfrak{o}_2)$ below depending on their restriction to the first congruence subgroup K_2^1 which is isomorphic to $M_2(\mathbb{F}_q)$. Note that a regular matrix in $M_2(\mathbb{F}_q)$ is either non-split semisimple or split non-semisimple or split semisimple. It follows that the irreducible regular representation of $GL_2(\mathfrak{o}_2)$ can be classified into three classes described below. If an irreducible regular representation σ is such that $\sigma|_{K_2^1}$ contains ϕ_B with *B* non-split semisimple (respectively, split non-semisimple, split semisimple) then we say the type of σ is non-split semisimple (respectively, split non-semisimple, split semisimple). In $M_2(\mathbb{F}_q)$, for a given trace, there is a unique class of scalar matrix, a unique class of split non-semisimple matrices, $\frac{q-1}{2}$ distinct classes of split semisimple matrices and $\frac{q-1}{2}$ distinct classes of non-split semisimple matrices.

Fix a regular character ϕ_B of $K_2^1 \cong M_2(\mathbb{F}_q)$ and a character ω of the center Z of $GL_2(\mathfrak{o}_2)$. The character ϕ_B can be extended to the inertia group $I(\phi_B)$. Observe that $Z \cdot K_2^1 \subset I(\phi_B)$. The following table describes the number of irreducible regular representation of different types when its restriction to $Z \cdot K_2^1$ contains the character $\omega \cdot \phi_B$.

T	1
IARIE	
INDLL	т.

S.No.	Type of σ	dimension of σ	No. of σ with $\omega \cdot \phi_B \subset \sigma _{Z \cdot K_2^1}$
1.	non-split semisimple	q(q-1)	q+1
2.	split non-semisimple	$q^2 - 1$	9
3.	split semisimple	q(q+1)	q-1

The irreducible representations of $GL_2(\mathfrak{o}_2)$, which are not regular, are of the type $\sigma \otimes (\chi \circ \det)$, where σ is an irreducible representation of $GL_2(\mathbb{F}_q)$ considered as a representation of $GL_2(\mathfrak{o}_2)$ using the natural map $GL_2(\mathfrak{o}_2) \to GL_2(\mathbb{F}_q)$ and $\chi : \mathfrak{o}_2^{\times} \to \mathbb{C}^{\times}$ is a character.

2.3. **Strongly primitive characters.** Let *E* be an unramified extension of *F* of degree *n*. Let \mathcal{O} be the integral closure of \mathfrak{o} in *E* with maximal ideal $\mathcal{P} = \mathfrak{O}\mathcal{O}$. Let $\mathcal{O}_l = \mathcal{O}/\mathcal{P}^l$, then \mathcal{O}_l is isomorphic to a free \mathfrak{o}_l -module of rank *n*. Therefore, $\operatorname{GL}_n(\mathfrak{o}_l)$ can be identified with the group of automorphisms of \mathfrak{o}_l -module \mathcal{O}_l . This identification is determined up to an inner automorphism of $\operatorname{GL}_n(\mathfrak{o}_l)$. Clearly, for $a \in \mathcal{O}_l, x \mapsto ax$ is an \mathfrak{o}_l -module endomorphism of \mathcal{O}_l and we get an embedding of the ring \mathcal{O}_l to the ring of endomorphisms of \mathfrak{o}_l -module \mathcal{O}_l . Therefore, \mathcal{O}_l^{\times} becomes a subgroup of $\operatorname{GL}_n(\mathfrak{o}_l)$ under the same embedding.

- **Definition 2.4.** (a) A character $\phi : \mathcal{O}_1 \cong \mathbb{F}_{q^n} \to \mathbb{C}^{\times}$ is called primitive if there does not exist a proper subfield \mathbb{F}_{q^d} of \mathbb{F}_{q^n} and a character ϕ_0 of \mathbb{F}_{q^d} such that $\phi(x) = \phi_0(\operatorname{tr}_{\mathbb{F}_{q^n}|\mathbb{F}_{q^d}}(x))$ for all $x \in \mathbb{F}_{q^n}$, where $\operatorname{tr}_{\mathbb{F}_{q^n}|\mathbb{F}_{q^d}}$ denote the trace map for the field extension $\mathbb{F}_{q^n}/\mathbb{F}_{q^d}$.
 - (b) For l > 1, a character $\theta : \mathcal{O}_l^{\times} \to \mathbb{C}^{\times}$ is called strongly primitive if its restriction to the kernel of the map $\mathcal{O}_l^{\times} \to \mathcal{O}_{l-1}^{\times}$, which is isomorphic to \mathcal{O}_1 , is a primitive character.

Strongly cuspidal representations of $\operatorname{GL}_n(\mathfrak{o}_l)$ are associated to strongly primitive characters of \mathcal{O}_l^{\times} , which we explain for even l = 2m. Let $x \in M_n(\mathbb{F}_q)$ be a regular elliptic element and $\tilde{x} \in M_n(\mathfrak{o}_m)$ be such that it maps to x under the natural quotient $M_n(\mathfrak{o}_m) \to M_n(\mathbb{F}_q)$. Since $K_l^m \cong M_n(\mathfrak{o}_m)$, $\phi_{\tilde{x}}$ is a character of K_l^m . Then the inertia group $I(\phi_{\tilde{x}}) \cong \mathcal{O}_l^{\times} K_l^m$. Let $\tilde{\phi}_{\tilde{x}}$ be an extension of $\phi_{\tilde{x}}$ to $I(\phi_{\tilde{x}})$. Write $\theta = \tilde{\phi}_{\tilde{x}}|_{\mathcal{O}_l^{\times}}$. Then θ is a strongly primitive character of \mathcal{O}_l^{\times} associated to the strongly cuspidal representation $\pi = \operatorname{Ind}_{I(\tilde{\phi}_{\tilde{x}})}^{\operatorname{GL}_n(\mathfrak{o}_l)} \tilde{\phi}_{\tilde{x}}$, see Aubert-Onn-Prasad [1], Lusztig [11]. More precisely, we have the following result [1, Theorem B].

Theorem 2.5. There is a canonical bijective correspondence between strongly cuspidal representations of $GL_n(\mathfrak{o}_l)$ and Gal(E/F) orbits of strongly primitive characters of \mathcal{O}_l^{\times} .

- **Notation 2.6.** (a) Let L be a degree 2 unramified extension of F and \mathfrak{O} the integral closure of \mathfrak{o} in L. Let E be a degree 4 unramified extension of F containing L with \mathcal{O} the integral closure of \mathfrak{o} in E. Throughout the paper, we denote \mathfrak{o}_2 , \mathfrak{O}_2 and \mathcal{O}_2 for $\mathfrak{o}/(\varpi^2)$, $\mathfrak{O}/(\varpi^2)$ and $\mathcal{O}/(\varpi^2)$ respectively.
 - (b) We denote the subgroup $I + \varpi M_2(\mathfrak{o}_2)$ of $\operatorname{GL}_2(\mathfrak{o}_2)$ by J_2^1 and the subgroup $I + \varpi M_4(\mathfrak{o}_2)$ of $\operatorname{GL}_4(\mathfrak{o}_2)$ by K_2^1 .

3. Double cosets $T \setminus G/P$

For a representation π of a finite group H and an automorphism τ of H, we define a representation π^{τ} given by $\pi^{\tau} := \pi \circ \tau$. In particular, if $H \subset \tilde{H}$ is a subgroup and $x \in \tilde{H}$ normalises H, then we get an automorphism τ_x of H given by $\tau_x(y) = x^{-1}yx$. In this case, we write π^x instead of π^{τ_x} .

Throughout this section, $G = \operatorname{GL}_4(\mathfrak{o}_2)$ and $P = P_{2,2} \subset G$. We fix an irreducible strongly cuspidal representation $\pi = \operatorname{Ind}_{I(\phi_x)}^{\operatorname{GL}_4(\mathfrak{o}_2)}(\tilde{\phi}_x)$ where x is a regular elliptic element of $M_4(\mathbb{F}_q)$. We will choose an embedding $\mathbb{F}_{q^4}^{\times} \hookrightarrow \operatorname{GL}_4(\mathbb{F}_q)$ and take $x \in \mathbb{F}_{q^4}^{\times}$. We will also fix an embedding of $\mathcal{O}_2^{\times} \hookrightarrow \operatorname{GL}_4(\mathfrak{o}_2)$ such that \mathcal{O}_2^{\times} maps onto $\mathbb{F}_{q^4}^{\times}$ under the quotient map $\operatorname{GL}_4(\mathfrak{o}_2) \to \operatorname{GL}_4(\mathbb{F}_q)$. Since x is a regular elliptic element, we have $I(\phi_x) = \mathcal{O}_2^{\times} \cdot K_2^1$ and $I(\phi_x)/K_2^1 \cong \mathbb{F}_{q^4}^{\times}$. For convenience, we write $T = I(\phi_x)$ and then $\pi = \operatorname{Ind}_T^G \tilde{\phi}_x$. Therefore, $\pi_{N,\psi} = (\operatorname{Res}_P^G \pi)_{N,\psi} = (\operatorname{Res}_P^G \operatorname{Ind}_T^G \tilde{\phi}_x)_{N,\psi}$. By Mackey theory $\operatorname{Res}_P^G \operatorname{Ind}_T^G \tilde{\phi}_x \cong \bigoplus_{\delta \in T \setminus G/P} \operatorname{Ind}_{\delta^{-1}T\delta \cap P}^P \tilde{\phi}_x^{\delta^{-1}}$.

For $\delta \in T \setminus G/P$, we will write $\pi^{\delta} = \operatorname{Ind}_{\delta^{-1}T\delta \cap P}^{P} \tilde{\phi}_{x}^{\delta^{-1}}$. Now we describe the double cosets $T \setminus G/P$. We write \bar{P} for the image of P in $\operatorname{GL}_{4}(\mathbb{F}_{q})$ and $\bar{T} = T/K_{2}^{1} \cong \mathbb{F}_{q^{4}}^{\times}$. Since K_{2}^{1} is the kernel of the quotient map $\operatorname{GL}_{4}(\mathfrak{o}_{2}) \to \operatorname{GL}_{4}(\mathbb{F}_{q})$ and $K_{2}^{1} \subset T$, the quotient map induces a bijection between set of double cosets $T \setminus \operatorname{GL}_{4}(\mathfrak{o}_{2})/P$ and $\bar{T} \setminus \operatorname{GL}_{4}(\mathbb{F}_{q})/\bar{P}$.

Lemma 3.1. The number of distinct cosets $\overline{T} \setminus GL_4(\mathbb{F}_q) / \overline{P}$ is q + 1.

Proof. We know that $\operatorname{GL}_4(\mathbb{F}_q)/\bar{P} = \operatorname{Gr}(4,2)$ the Grassmannian, i.e. the set of all 2dimensional subspaces in a 4-dimensional vector space over \mathbb{F}_q . For a group *G* acting on a set *X* we write, $G \curvearrowright X$. Naturally, $\operatorname{GL}_4(\mathbb{F}_q) \curvearrowright \operatorname{Gr}(4,2)$ transitively and distinct orbits of the restricted action to the subgroup $\bar{T} \hookrightarrow \operatorname{GL}_4(\mathbb{F}_q)$ gives the distinct double cosets $\bar{T} \setminus \bar{G}/\bar{P}$. Now, consider \mathbb{F}_{q^4} as a 4-dimensional vector space over \mathbb{F}_q . A general element of $\operatorname{Gr}(4,2)$ can be written as $\operatorname{Span}\{x,y\}$ the span of vectors $x, y \in \mathbb{F}_{q^4}$ which are linearly independent over \mathbb{F}_q . Further, a representative of any orbit under the action $\mathbb{F}_{q^4}^{\times} \cap$ $\operatorname{Gr}(4,2)$ can be taken to be $\operatorname{Span}\{1,y\}$ where $y \in \mathbb{F}_{q^4} \setminus \mathbb{F}_q$. It can be easily verified that $\operatorname{Span}\{1,y\}$ and $\operatorname{Span}\{1,y'\}$ are in the same orbit if and only if $y' = \frac{py+q}{ry+s}$ for some $p, q, r, s \in \mathbb{F}_q$ where $\{py+q, ry+s\}$ is a linearly independent set of vectors of \mathbb{F}_{q^4} , which is equivalent to say that $\exists \begin{pmatrix} p & q \\ r & s \end{pmatrix} \in \operatorname{GL}_2(\mathbb{F}_q)$ such that $y' = \frac{py+q}{ry+s}$. Now consider the action $\operatorname{GL}_2(\mathbb{F}_q) \cap \mathbb{F}_{q^4} \setminus \mathbb{F}_q$ given by

$$\begin{pmatrix} p & q \\ r & s \end{pmatrix} \cdot y = \frac{py+q}{ry+s}, \text{ where } \begin{pmatrix} p & q \\ r & s \end{pmatrix} \in \operatorname{GL}_2(\mathbb{F}_q) \text{ and } y \in \mathbb{F}_{q^4} \setminus \mathbb{F}_q.$$
(3.1)

Thus, it is clear that there is a bijection between the orbits in the action $\mathbb{F}_{q^4}^{\times} \curvearrowright \operatorname{Gr}(2,4)$ and the action $\operatorname{GL}_2(\mathbb{F}_q) \curvearrowright \mathbb{F}_{q^4} \setminus \mathbb{F}_q$. We use the action $\operatorname{GL}_2(\mathbb{F}_q) \curvearrowright \mathbb{F}_{q^4} \setminus \mathbb{F}_q$ to compute the number of distinct orbits.

- If $y \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, the stabiliser of Span $\{1, y\}$ is $\mathbb{F}_{q^2}^{\times}$ and therefore the orbit size of Span $\{1, y\}$ is $q^2 q$.
- If $y \in \mathbb{F}_{q^4} \setminus \mathbb{F}_{q^2}$, the stabiliser of Span $\{1, y\}$ is \mathbb{F}_q^{\times} and therefore the orbit size of Span $\{1, y\}$ is $q(q^2 1)$.

Thus, there is a single orbit with stabiliser $\mathbb{F}_{a^2}^{\times}$ and the number of orbits with stabiliser

$$\mathbb{F}_q^{\times}$$
 is $\frac{\left|\mathbb{F}_{q^4} \setminus \mathbb{F}_{q^2}\right|}{q(q^2 - 1)} = q$. Hence, the number of distinct double cosets is $q + 1$.

Now we write down an exhaustive, although not exclusive, set of representatives for these double cosets for which we need to fix an embedding of $\mathbb{F}_{q^4}^{\times}$ inside $GL_4(\mathbb{F}_q)$. We fix an embedding of \mathbb{F}_{q^4} in $M_4(\mathbb{F}_q)$ as described below.

Let $\alpha \in \mathbb{F}_q^{\times} \setminus \mathbb{F}_q^{\times 2}$ and then $\mathbb{F}_{q^2} = \mathbb{F}_q(\sqrt{\alpha})$. Let $a, b \in \mathbb{F}_q$ be such that $a + b\sqrt{\alpha} \in \mathbb{F}_{q^2} \setminus \mathbb{F}_{q^2}^{\times 2} \setminus \mathbb{F}_{q^2}^{\times 2}$ and

Norm<sub>**F**_{*q*²}|**F**_{*q*}(
$$a + b\sqrt{\alpha}$$
) = $a^2 - b^2\alpha$ is a square free element in **F**_{*q*}, (3.2)</sub>

where Norm_{$\mathbb{F}_{q^2}|\mathbb{F}_q}$ denotes the norm map for the field extension $\mathbb{F}_{q^2}/\mathbb{F}_q$. Write $\beta = \sqrt{a + b\sqrt{\alpha}}$. Then $\mathbb{F}_{q^4} = \mathbb{F}_q(\beta)$. Note that $\mathbb{F}_q(\beta^2) = \mathbb{F}_{q^2}$. For $x \in \mathbb{F}_{q^4}$, the map $\mathbb{F}_{q^4} \to \mathbb{F}_{q^4}$ given by $y \mapsto xy$ is \mathbb{F}_q -linear and thus gives rise to an embedding $\mathbb{F}_{q^4} \hookrightarrow M_4(\mathbb{F}_q)$. Similarly, there is an embedding $\mathbb{F}_{q^2} \hookrightarrow M_2(\mathbb{F}_q)$. By fixing the ordered basis $\{1, \beta^2, \beta, \beta^3\}$ of \mathbb{F}_{q^4} as a \mathbb{F}_q -vector space, we write an explicit embedding which is given by}

$$x = a_0 + a_1 \beta^2 + a_2 \beta + a_3 \beta^3 \mapsto \begin{pmatrix} X_1 & X_2 \\ X_3 & X_1 \end{pmatrix},$$
 (3.3)

where $X_1, X_2, X_3 \in M_2(\mathbb{F}_q)$ are as follows:

$$X_{1} = \begin{pmatrix} a_{0} & -a_{1}(a^{2} - b^{2}\alpha) \\ a_{1} & a_{0} + 2aa_{1} \end{pmatrix}, \quad X_{2} = \begin{pmatrix} -a_{3}(a^{2} - b^{2}\alpha) & -(a_{2} + 2aa_{3})(a^{2} - b^{2}\alpha) \\ a_{2} + 2aa_{3} & a_{3}(3a^{2} + b^{2}\alpha) + 2aa_{2} \end{pmatrix}, \quad X_{3} = \begin{pmatrix} -a_{3}(a^{2} - b^{2}\alpha) & -(a_{2} + 2aa_{3})(a^{2} - b^{2}\alpha) \\ a_{2} + 2aa_{3} & a_{3}(3a^{2} + b^{2}\alpha) + 2aa_{2} \end{pmatrix}$$

 $\begin{pmatrix} a_2 & -a_3(a^2 - b^2\alpha) \\ a_3 & a_2 + 2aa_3 \end{pmatrix}$. Note that X_1 and X_3 are the images of $a_0 + a_1\beta^2$ and $a_2 + a_3\beta^2$ under the fixed embedding $\mathbb{F}_{q^2} \hookrightarrow M_2(\mathbb{F}_q)$ obtained by the ordered basis $\{1, \beta^2\}$ of \mathbb{F}_{q^2} as \mathbb{F}_q -vector space. More explicitly,

$$a_0 + a_1 \beta^2 \mapsto \begin{pmatrix} a_0 & -a_1(a^2 - b^2 \alpha) \\ a_1 & a_0 + 2aa_1 \end{pmatrix}.$$
 (3.4)

From now onwards, we identify $\mathbb{F}_{q^4}^{\times}$ as a subgroup of $GL_4(\mathbb{F}_q)$ and $\mathbb{F}_{q^2}^{\times}$ as a subgroup of $GL_2(\mathbb{F}_q)$ via the above mentioned embedding.

Definition 3.2. For
$$u, v, w \in \mathbb{F}_q$$
, define $A_{u,v} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & u & 1 & 0 \\ 0 & v & 0 & 1 \end{pmatrix}$ and $A_w = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & w & 0 & 1 \end{pmatrix}$.

Lemma 3.3. By taking the embedding $\mathbb{F}_{q^4}^{\times} \hookrightarrow \operatorname{GL}_4(\mathbb{F}_q)$ obtained from Equation (3.3), an exhaustive, but not exclusive, set of coset representatives of $\overline{T} \setminus \overline{G} / \overline{P}$ is given by

$$\{A_{u,v}, A_w : u, v, w \in \mathbb{F}_q\}$$

Proof. Recall that $\operatorname{GL}_4(\mathbb{F}_q)/\overline{P} \cong \operatorname{Gr}(4,2)$ where \overline{P} is the stabiliser of the subspace $W_0 = \operatorname{Span}\{1,\beta^2\} \cong \mathbb{F}_{q^2}$, therefore transitivity of the action $\operatorname{GL}_4(\mathbb{F}_q) \curvearrowright \operatorname{Gr}(4,2)$ implies given any $W \in \operatorname{Gr}(4,2)$, we get an element $g \in \operatorname{GL}_4(\mathbb{F}_q)$ such that $g \cdot W_0 = W$. Note that a representative of an orbit for the action $\mathbb{F}_{q^4}^{\times} \curvearrowright \operatorname{Gr}(4,2)$ can be taken to be $W = \operatorname{Span}\{1,y\}$ for $y \in \mathbb{F}_{q^4} \setminus \mathbb{F}_q$. Write $y = b_0 + b_1\beta^2 + b_2\beta + b_3\beta^3$ for some $b_0, b_1, b_2, b_3 \in \mathbb{F}_q$ then,

$$W = \text{Span}\{1, b_1\beta^2 + b_2\beta + b_3\beta^3\}.$$

- Case 1: If $b_1 \neq 0$, we can take $b_1 = 1$ and get $W = \text{Span}\{1, \beta^2 + b_2\beta + b_3\beta^3\}$ where $b_2, b_3 \in \mathbb{F}_q$. We define a linear map on \mathbb{F}_{q^4} which maps W_0 to W in such a way that $1 \mapsto 1, \beta^2 \mapsto \beta^2 + b_2\beta + b_3\beta^3, \beta \mapsto \beta, \beta^3 \mapsto \beta^3$. Then a representative for the double coset corresponding to W can be taken to be A_{b_2,b_3} .
- Case 2: If $b_1 = 0$, $b_2 \neq 0$ we can take $b_2 = 1$ and get $W = \text{Span}\{1, \beta + b_3\beta^3\}$ where $b_3 \in \mathbb{F}_q$. Similar to the Case 1, a representative for the double coset corresponding to W can be taken to be A_{b_3} .
- Case 3: If $b_1 = b_2 = 0$, $b_3 \neq 0$ we can take $b_3 = 1$ and get $W = \text{Span}\{1, \beta^3\}$. For this subspace, we claim that its representative can be taken to be either $A_{u,v}$ for some $u, v \in \mathbb{F}_q$ or A_w for some $w \in \mathbb{F}_q$. Note that β^{-3} action takes $\text{Span}\{1, \beta^3\}$ to $\text{Span}\{1, \beta^{-3}\}$. Write $\beta^{-3} = b_0 + b_1\beta^2 + b_2\beta + b_3\beta^3$ for some b_0, b_1, b_2, b_3 . We claim that either $b_1 \neq 0$ or $b_2 \neq 0$ and then the proof follows from the earlier two cases. If $b_1 = b_2 = 0$, then, $\beta^{-3} = b_0 + b_3\beta^3$ which gives $1 = b_0\beta^3 + b_3\beta^6 = b_0\beta^3 (2ab_3)(a^2 b^2\alpha) + (3a^2 + b^2\alpha)b_3\beta^2$. Comparing the coefficients both sides, we get, $b_0 = 0$ and $-2ab_3(a^2 b^2\alpha) = 1$ and $(3a^2 + b^2\alpha)b_3 = 0$ which implies $3a^2 + b^2\alpha = 0 \implies \text{Norm}_{\mathbb{F}_{q^2}|\mathbb{F}_q}(a + b\sqrt{\alpha}) = 4a^2$, which contradicts our assumption (3.2).

Lemma 3.4. For every $w \in \mathbb{F}_q$ there exists $u, v \in \mathbb{F}_q$ such that $\overline{T}A_w\overline{P} = \overline{T}A_{u,v}\overline{P}$.

Proof. It can be verified that $\overline{T}A_w\overline{P} = \overline{T}A_{u,v}\overline{P}$ if and only if

$$u^{2}[w(1+2aw)] - uv[1 - w^{2}(3a^{2} + b^{2}\alpha)] - v^{2}[2a + w(3a^{2} + b^{2}\alpha)] + (1 + 2aw + (a^{2} - b^{2}\alpha)w^{2}) = 0.$$
(3.5)

Therefore, it is enough to prove that for any $w \in \mathbb{F}_q$, there always exists $u, v \in \mathbb{F}_q$ such that Equation (3.5) holds. For a given $w \in \mathbb{F}_q$, we consider the above equation as a quadratic equation in *u*. If the coefficient of u^2 is 0, i.e. w(1 + 2aw) = 0, then the Equation (3.5) is linear and one can easily verify that there exists $u, v \in \mathbb{F}_q$ satisfying the equation. Now we assume that the coefficient of u^2 is not zero. Then Equation (3.5) has a solution $u \in \mathbb{F}_q$ if and only if the discriminant is a square element in \mathbb{F}_q . The discriminant D(v) is given by

$$D(v) = v^2 \left((1 + 2aw)^2 - (a^2 - b^2\alpha)w^2 \right) - 4w(1 + 2aw) \left((1 + aw)^2 - (bw)^2\alpha \right).$$

Clearly, the coefficient of v^2 is not zero, otherwise $a^2 - b^2 \alpha \in \mathbb{F}_a^{\times 2}$. Then,

$$\left|\left\{D(v): v \in \mathbb{F}_q\right\}\right| = \frac{q+1}{2}$$

Since the number of non-square elements in \mathbb{F}_q is $\frac{q-1}{2}$, there exists $v \in \mathbb{F}_q$ such that $D(v) \in \mathbb{F}_q^{\times 2}$. Hence we are through.

By using Lemma 3.3 and Lemma 3.4 we conclude the following.

Proposition 3.5. The set $\Omega := \{A_{u,v} : u, v \in \mathbb{F}_q\}$ is an exhaustive, but not exclusive, set of representatives for $\overline{T} \setminus \overline{G} / \overline{P}$.

We state the following lemma without proof which provides an equivalent condition for two elements from Ω representing the same double coset for $\overline{T} \setminus G/\overline{P}$.

Lemma 3.6. For $\delta = A_{u,v}$, $\delta' = A_{u',v'} \in \Omega$, let $C(\delta, \delta') = u^2 - {u'}^2 + uv{u'}^2 - u^2u'v' + 2a(v^2{u'}^2 - u^2{v'}^2) + 2a(uv - u'v')$ + $(3a^2 + b^2\alpha)(u'v'v^2 - uvv'^2) + (v^2 - v'^2)(a^2 - b^2\alpha).$

Then, $T\delta P = T\delta'P$ if and only if $C(\delta, \delta') = 0$. In particular, $T\delta P = TIP$ if and only if $\delta = I$, where I is the 4×4 identity matrix.

Here is a list of notations which we will use throughout this paper.

- (a) We denote the identity matrix of a suitable order by I, which will be Notation 3.7. clear from the context.
 - (b) Let $\Omega_0 \subset \Omega$ be such that every double coset in $\overline{T} \setminus \overline{G} / \overline{P}$ has a representative in Ω_0 and distinct elements in Ω_0 represent distinct double cosets in $\overline{T} \setminus \overline{G} / \overline{P}$.
 - (c) Since the quotient map $G \to \overline{G}$ induces a bijection between $T \setminus G / P$ and $\overline{T} \setminus \overline{G} / \overline{P}$, for a representative $\delta \in \overline{G}$ of a double coset in $\overline{T} \setminus \overline{G} / \overline{P}$ we write its lift to G by the same *letter* δ *for the corresponding representative of the double coset in* $T \setminus G/P$ *.*
 - (d) For $\delta = A_{u,v} = \begin{pmatrix} I & 0 \\ U & I \end{pmatrix}$ where $U = \begin{pmatrix} 0 & u \\ 0 & v \end{pmatrix}$, write $L_{\delta} = X_1 U - U X_1 + X_3 - U X_2 U$. In particular, $L_I = X_3$.
 - (e) If L_{δ} is invertible, then write $\mathcal{B}_{\delta} = L_{\delta}(X_1 + X_2 U)L_{\delta}^{-1} + X_1 UX_2$. In particular, $\mathcal{B}_{I} = 2X_{1} \text{ and } \operatorname{tr}(\mathcal{B}_{\delta}) = \operatorname{tr}(2X_{1}).$ (f) Write $X = a_{1}^{2} - a_{2}a_{3} - 2aa_{3}^{2} \text{ and } Y = a_{2}^{2} + a_{3}^{2}(a^{2} - b^{2}\alpha) + 2aa_{2}a_{3}.$

Observe that $Y \neq 0$ otherwise $a_2 = a_3 = 0$ and then *x* will not be a regular elliptic element of $M_4(\mathbb{F}_q)$. It can be checked that

$$\det(L_{\delta}) = -u^{2}X - uv\left(2aX + Y\right) - v^{2}\left(2aY + X(a^{2} - b^{2}\alpha)\right) + Y.$$
(3.6)

4. The dimension of $\pi_{N,\psi}$

For a finite dimensional representation ρ of a finite group H, we write Θ_{ρ} for the character of ρ . Suppose ρ' be an irreducible representation of the group H. The multiplicity of ρ' in ρ is defined to be the dimension of $\operatorname{Hom}_{H}(\rho, \rho')$. We say that ρ is multiplicity free if the multiplicity of any irreducible representation in ρ is ≤ 1 . If we define $\left\langle \Theta_{\rho}, \Theta_{\rho'} \right\rangle_{H} := \frac{1}{|H|} \sum_{g \in H} \Theta_{\rho}(g) \overline{\Theta}_{\rho'}(g)$, then by [5, Corollary 2.16] we have $\dim(\operatorname{Hom}_{H}(V, V')) = \left\langle \Theta_{\rho}, \Theta_{\rho'} \right\rangle_{H}$.

We fix $x \in \mathbb{F}_{q^4}^{\times}$ as in Equation (3.3) and consider a strongly cuspidal representation $\pi = \operatorname{Ind}_T^G \tilde{\phi}_x$ of $\operatorname{GL}_4(\mathfrak{o}_2)$ (see Theorem 2.2). If we write $\pi^{\delta} = \operatorname{Ind}_{\delta^{-1}T\delta\cap P}^P \tilde{\phi}_x^{\delta^{-1}}$, then by Mackey theory,

$$\pi_{N,\psi} = \bigoplus_{\delta \in \Omega_0} \pi_{N,\psi}^{\delta}.$$

Therefore, $\dim(\pi_{N,\psi}) = \sum_{\delta \in \Omega_0} \dim(\pi_{N,\psi}^{\delta})$. In what follows we compute the dimension of $\pi_{N,\psi}^{\delta}$ for all $\delta \in \Omega$ with the help of following series of lemmas.

Lemma 4.1. Write
$$\omega N = \left\{ \begin{pmatrix} I & \omega X \\ 0 & I \end{pmatrix} : X \in M_2(\mathfrak{o}_2) \right\}$$
. Then, for every $\delta \in \Omega$, we have
$$\pi_{N,\psi}^{\delta} \cong \bigoplus_{\gamma \in (\delta^{-1}T\delta \cap P) \setminus P/N} \operatorname{Hom}_{\omega N} \left(\tilde{\phi}_{\delta^{-1}x\delta}^{\gamma^{-1}}, \psi \right).$$

Proof. By Mackey theory, $\pi_{N,\psi}^{\delta} \cong \bigoplus_{\gamma \in (\delta^{-1}T\delta \cap P) \setminus P/N} \operatorname{Hom}_{\gamma^{-1}(\delta^{-1}T\delta \cap P)\gamma \cap N} \left(\tilde{\phi}_{\delta^{-1}x\delta}^{\gamma^{-1}} \psi \right).$

The Lemma follows from the fact that the subgroup $\gamma^{-1}(\delta^{-1}T\delta \cap P)\gamma \cap N = \varpi N$ for all $\gamma \in (\delta^{-1}T\delta \cap P) \setminus P/N$, which we prove now. For any $\gamma \in P$, we have

$$\gamma^{-1}(\delta^{-1}T\delta \cap P)\gamma \cap N = \gamma^{-1}(\delta^{-1}T\delta \cap P \cap N)\gamma = \gamma^{-1}(\delta^{-1}T\delta \cap N)\gamma.$$

Since $\delta^{-1}T\delta \pmod{K_2^1} \cong \mathbb{F}_{q^4}^{\times}$ we get, $(\delta^{-1}T\delta \cap N) \pmod{K_2^1} = \{I\}$. Then, $\delta^{-1}T\delta \cap N = \omega N$. Also, ωN is a normal subgroup of *P*, therefore $\gamma^{-1}(\delta^{-1}T\delta \cap N)\gamma = \omega N$.

Note that $(\delta^{-1}T\delta \cap P) \setminus P/N$ is in bijection with $(\delta^{-1}\mathbb{F}_{q^4}^{\times}\delta \cap \overline{P}) \setminus \overline{P}/\overline{N}$.

Lemma 4.2. A set of representatives of the double coset $(\delta^{-1}\mathbb{F}_{q^4}^{\times}\delta \cap \bar{P})\setminus \bar{P}/\bar{N}$ is given as

(a) For $\delta = I$, $\left\{ \begin{pmatrix} g_1 & 0 \\ 0 & g_2 \end{pmatrix} : g_1 \in \mathbb{F}_{q^2}^{\times} \setminus \operatorname{GL}_2(\mathbb{F}_q), g_2 \in \operatorname{GL}_2(\mathbb{F}_q) \right\}$ is a set of distinct representatives for $(\mathbb{F}_{q^4}^{\times} \cap \bar{P}) \setminus \bar{P} / \bar{N}$.

(b) For
$$\delta \neq I$$
, $\left\{ \begin{pmatrix} g_1 & 0 \\ 0 & g_2 \end{pmatrix} : g_1 \in \mathbb{F}_q^{\times} \setminus \operatorname{GL}_2(\mathbb{F}_q), g_2 \in \operatorname{GL}_2(\mathbb{F}_q) \right\}$ is a set of distinct representatives for $(\delta^{-1}\mathbb{F}_{q^4}^{\times}\delta \cap \bar{P}) \setminus \bar{P}/\bar{N}$.

Proof. Clearly, $\bar{P}/\bar{N} \cong \operatorname{GL}_2(\mathbb{F}_q) \times \operatorname{GL}_2(\mathbb{F}_q)$. Then (a) follows by noting that $\mathbb{F}_{q^4}^{\times} \cap \bar{P} \cong \mathbb{F}_{q^2}^{\times}$ and, (b) follows by noting that $\delta^{-1}\mathbb{F}_{q^4}^{\times}\delta \cap \bar{P} \cong \mathbb{F}_q^{\times}$ for $\delta \neq I$.

Lemma 4.3. (a) $\dim(\pi_{N,\psi}^{I}) = q(q-1)$. (b) Let $\delta \neq I$. If $\pi_{N,\psi}^{\delta} \neq 0$ then, $\dim(\pi_{N,\psi}^{\delta}) = q(q^{2}-1)$.

Proof. For $\gamma \in P$ we have

$$\left\langle \tilde{\phi}_{\delta^{-1}x\delta}^{\gamma^{-1}},\psi\right\rangle_{\varpi N} = 1 \text{ if } \tilde{\phi}_{\delta^{-1}x\delta}^{\gamma^{-1}}|_{\varpi N} = \psi|_{\varpi N} \text{ and } 0 \text{ otherwise.}$$

Write $\gamma = \begin{pmatrix} \tilde{g}_1 & 0 \\ 0 & \tilde{g}_2 \end{pmatrix}$ and using the definition of ϕ_x on ϖN we get, $\tilde{\phi}_{\delta^{-1}x\delta}^{\gamma^{-1}}|_{\varpi N} = \psi|_{\varpi N}$ if and only if the character $X \mapsto \psi_0(\operatorname{tr}((\tilde{g}_2^{-1}\tilde{L}_\delta \tilde{g}_1 - I)X))$ of ϖN is trivial (recall L_δ as defined in Notation 3.7(e)), which is equivalent to $L_\delta g_1 = g_2$.

(a) Let $\delta = I$. Note that $L_I = X_3$. By Lemma 4.2 (a), we get

$$\dim(\pi_{N,\psi}^{I}) = \left| \left\{ (g_1, g_2) \in \left(\mathbb{F}_{q^2}^{\times} \backslash \mathrm{GL}_2(\mathbb{F}_q) \right) \times \mathrm{GL}_2(\mathbb{F}_q) : g_2 = X_3 g_1 \right\} \right|$$
$$= q(q-1).$$

(b) Let $\delta \neq I$. By Lemma 4.2 (b), we get

$$\dim(\pi_{N,\psi}^{\delta}) = \frac{1}{|\varpi M_2(\mathfrak{o}_2)|} \sum_{\substack{\gamma \in (\delta^{-1}T\delta \cap P) \setminus P/N \\ X \in \varpi M_2(\mathfrak{o}_2)}} \psi_0(\operatorname{tr}((\tilde{g}_2^{-1}\tilde{L}_{\delta}\tilde{g}_1 - I)X))$$
$$= \left| \left\{ (g_1, g_2) \in \left(\mathbb{F}_q^{\times} \setminus \operatorname{GL}_2(\mathbb{F}_q) \right) \times \operatorname{GL}_2(\mathbb{F}_q) : g_2 = L_{\delta}g_1 \right\} \right|$$
$$= \left\{ \begin{array}{l} q(q^2 - 1) & \text{if } L_{\delta} \text{ is invertible,} \\ 0 & \text{otherwise.} \end{array} \right.$$

Lemma 4.4. There exists exactly one $\delta \in \Omega_0$ such that $\pi_{N,\psi}^{\delta} = 0$.

Proof. First, we prove that there exists a $\delta = A_{u,v} \in \Omega$ for some $u, v \in \mathbb{F}_q$ such that $\pi_{N,\psi}^{\delta} = 0$. From the proof of Lemma 4.3 it follows that $\pi_{N,\psi}^{\delta} = 0$ if and only if L_{δ} is not invertible, i.e. det $(L_{\delta}) = 0$.

Recall that $det(L_{\delta}) = -(u^2X + uv(2aX + Y) + v^2(2aY + X(a^2 - b^2\alpha)) - Y)$ where $Y \neq 0$, see (3.6). We claim that there exists $u, v \in \mathbb{F}_q$ such that $det(L_{\delta}) = 0$.

If X = 0, then det $(L_{\delta}) = 0 \Rightarrow (1 - uv - 2av^2)Y = 0$ and it can be easily verified that there exist $u, v \in F_q$ satisfying this condition. Now we assume $X \neq 0$ and consider the expression for det (L_{δ}) as a quadratic polynomial in variable u, whose discriminant is $(Y^2 - 4aXY + 4X^2b^2\alpha)v^2 + 4XY$. Note that if for some $v \in \mathbb{F}_q$, the discriminant is a square in \mathbb{F}_q then there exists a $u \in \mathbb{F}_q$ such that det $(L_{\delta}) = 0$. Write the discriminant as $C_1v^2 + C_2$ for $C_1 = Y^2 - 4aXY + 4X^2b^2\alpha$ and $C_2 = 4XY$. Note that $C_1 \neq 0$ otherwise,

$$Y^{2} - 4aXY + 4X^{2}b^{2}\alpha = (Y - 2aX)^{2} - 4X^{2}(a^{2} - b^{2}\alpha) = 0$$

i.e. $a^2 - b^2 \alpha$ is a square element in \mathbb{F}_q , which contradicts our assumption in (3.2). Since $C_1 \neq 0$, $|\{C_1v^2 + C_2 : v \in F_q\}| = \frac{q+1}{2}$ but the number of non-square elements in

 \mathbb{F}_q is $\frac{q-1}{2}$. Therefore, there exists a $v \in \mathbb{F}_q$ such that $C_1v^2 + C_2$ is a square in \mathbb{F}_q . This completes the proof of the existence of $\delta \in \Omega$ such that $\det(L_{\delta}) = 0$.

Now we prove the uniqueness, i.e. if $\pi_{N,\psi}^{\delta} = 0$ and $\pi_{N,\psi}^{\delta'} = 0$ then $T\delta P = T\delta' P$. Write $\delta = A_{u,v}$ and $\delta' = A_{u',v'}$ then

$$\det(L_{\delta}) = 0 \text{ and } \det(L_{\delta'}) = 0. \tag{4.1}$$

If X = 0, then (4.1) gives $1 - uv - 2av^2 = 0$ and $1 - u'v' - 2av'^2 = 0$. It can be easily verified that u, v and u', v' satisfy the condition in Lemma 3.6 giving $T\delta P = T\delta' P$. Now assume that $X \neq 0$. By eliminating Y from the two equations in (4.1) we get

$$X \cdot C(\delta, \delta') = 0.$$

Since $X \neq 0$, using Lemma 3.6, we get $T\delta P = T\delta' P$.

Proposition 4.5. Let π be a strongly cuspidal representation of $GL_4(\mathfrak{o}_2)$ then

$$\dim(\pi_{N,\psi}) = q^3(q-1).$$

Proof. Note that $\dim(\pi_{N,\psi}) = \sum_{\delta \in \Omega_0} \dim(\pi_{N,\psi}^{\delta})$. Recall that $|\Omega_0| = q + 1$ and by Lemma 4.4, there exists a unique $\delta \in \Omega_0$ such that $\delta \neq I$ and $\pi_{N,\psi}^{\delta} = 0$. Using Lemma 4.3, we get

$$\dim(\pi_{N,\psi}) = q(q-1) + (q-1) \cdot q(q^2 - 1) = q^3(q-1).$$

One of the main theorems in this paper is that $\pi_{N,\psi}$ is isomorphic to $\operatorname{Ind}_{\mathfrak{G}_2^{\times}}^{\operatorname{GL}_2(\mathfrak{o}_2)}(\theta|_{\mathfrak{G}_2^{\times}})$ where π is a strongly cuspidal representation of $\operatorname{GL}_4(\mathfrak{o}_2)$ associated to a strongly primitive character θ of \mathcal{O}_2^{\times} . Proposition 4.5 proves that these two representations have the same dimension.

5. Characters of $\pi_{N,\psi}^{\delta}$

For a finite dimensional representation ρ of a group we write ω_{ρ} for its central character, if it exists.

Lemma 5.1. Let \mathcal{M}, \mathcal{N} be subgroups of a finite group \mathcal{P} such that \mathcal{N} is a normal subgroup and $\mathcal{P} = \mathcal{M} \ltimes \mathcal{N}$. Let $\psi : \mathcal{N} \to \mathbb{C}^{\times}$ be a character of \mathcal{N} and $\mathcal{M}_{\psi} = \{m \in \mathcal{M} : \psi(mnm^{-1}) = \psi(n) \text{ for all } n \in \mathcal{N}\}$. Let (ρ, V) be a finite dimensional representation of \mathcal{P} and $V_{\mathcal{N},\psi} = \{v \in V : \rho(n)v = \psi(n)v \text{ for all } n \in \mathcal{N}\}$ which is a representation of \mathcal{M}_{ψ} denoted by $(\rho_{\mathcal{N},\psi}, V_{\mathcal{N},\psi})$. The character of $\rho_{\mathcal{N},\psi}$ is given by

$$\Theta_{\rho_{\mathcal{N},\psi}}(m) = \frac{1}{|\mathcal{N}|} \sum_{n \in \mathcal{N}} \Theta_{\rho}(mn) \overline{\psi}(n).$$

Proof. The projection map $\Pr: V \to V_{\mathcal{N},\psi}$ is given by $\Pr(v) = \frac{1}{|\mathcal{N}|} \sum_{n \in \mathcal{N}} \rho(n) \overline{\psi}(n) v$. Then,

$$\begin{split} \Theta_{\rho_{\mathcal{N},\psi}}(m) &= \operatorname{tr}(\rho_{\mathcal{N},\psi}(m)) = \operatorname{tr}(\rho(m)|_{V_{\mathcal{N},\psi}}) = \operatorname{tr}(\rho \circ \operatorname{Pr}) \\ &= \frac{1}{|\mathcal{N}|} \sum_{n \in \mathcal{N}} \operatorname{tr}(\rho(mn)) \overline{\psi}(n) = \frac{1}{|\mathcal{N}|} \sum_{n \in \mathcal{N}} \Theta_{\rho}(mn) \overline{\psi}(n). \end{split}$$

The main application of the above lemma is the following.

Corollary 5.2. Let ρ be a finite dimensional representation of $P_{n,n} = M \ltimes N$ where $\mathcal{M} \cong$ $\operatorname{GL}_n(\mathfrak{o}_l) \times \operatorname{GL}_n(\mathfrak{o}_l)$ and $N \cong M_n(\mathfrak{o}_l)$. Then, $g \in \operatorname{GL}_n(\mathfrak{o}_l)$ we have

$$\Theta_{\rho_{N,\psi}}(g) = \frac{1}{|N|} \sum_{X \in N} \Theta_{\rho} \begin{pmatrix} g & X \\ 0 & g \end{pmatrix} \overline{\psi}(Xg^{-1}).$$

We first calculate $\Theta_{\pi_{N,\psi}^{\delta}}$ on $J_2^1 = I + \omega M_2(\mathfrak{o}_2)$, for which we need $\Theta_{\pi^{\delta}}$. By using the formula for the character of an induced representation, the value $\Theta_{\pi^{\delta}} \begin{pmatrix} I + \omega A & X \\ 0 & I + \omega A \end{pmatrix}$ equals

$$\frac{1}{|\delta^{-1}T\delta\cap P|}\sum_{\substack{S,R\in\mathrm{GL}_{2}(\mathfrak{o}_{2})\\Q\in M_{2}(\mathfrak{o}_{2})}}\tilde{\phi}_{x}^{\delta^{-1}}\left(\begin{pmatrix}S&Q\\0&R\end{pmatrix}^{-1}\begin{pmatrix}I+\varpi A&X\\0&I+\varpi A\end{pmatrix}\begin{pmatrix}S&Q\\0&R\end{pmatrix}\right)$$

It is clear that this value is zero if $X \notin \omega M_2(\mathfrak{o}_2)$. Then by using Corollary 5.2, $\Theta_{\pi_{N,\psi}^{\delta}}(I + \omega A)$ is given by

$$\frac{1}{q^{8}|\delta^{-1}T\delta\cap P|} \sum_{\substack{S,R\in\operatorname{GL}_{2}(\mathfrak{o}_{2})\\X\in\omega M_{2}(\mathfrak{o}_{2})\\Q\in M_{2}(\mathfrak{o}_{2})}} \tilde{\phi}_{x}^{\delta^{-1}} \begin{pmatrix} I+\omega S^{-1}AS & 0\\ 0 & I+\omega R^{-1}AR \end{pmatrix} \tilde{\phi}_{x}^{\delta^{-1}} \begin{pmatrix} I & \omega (S^{-1}AQ-S^{-1}QR^{-1}AR)\\ 0 & I \end{pmatrix} \begin{pmatrix} I & \omega (S^{-1}AQ-S^{-1}QR^{-1}AR)\\ 0 & I \end{pmatrix} \tilde{\phi}_{x}^{\delta^{-1}} \begin{pmatrix} I & S^{-1}XR\\ 0 & I \end{pmatrix} \overline{\psi}(X)$$

$$(5.1)$$

We compute this by first summing over *X*, then over *Q*, and then over (S, R).

Lemma 5.3. For $\delta \in \Omega$, let $\pi_{N,\psi}^{\delta} \neq 0$ and recall L_{δ} and \mathcal{B}_{δ} from Notation 3.7 (d), (e). (a) For any $S, R \in \operatorname{GL}_2(\mathfrak{o}_2)$ we have

$$\sum_{X \in \omega M_2(\mathfrak{o}_2)} \tilde{\phi}_x^{\delta^{-1}} \begin{pmatrix} I & S^{-1}XR \\ 0 & I \end{pmatrix} \overline{\psi}(X) = \begin{cases} q^4 & \text{if } S = R\tilde{L}_\delta, \\ 0 & \text{otherwise.} \end{cases}$$

(b) For any δ and $S, R \in GL_2(\mathfrak{o}_2)$ with $S = R\tilde{L}_{\delta}$, we have

$$\sum_{Q \in M_2(\mathfrak{o}_2)} \tilde{\phi}_x^{\delta^{-1}} \begin{pmatrix} I & \mathcal{O}\left(S^{-1}AQ - S^{-1}QR^{-1}AR\right) \\ 0 & I \end{pmatrix} = q^8$$

(c) For $S = R\tilde{L}_{\delta}$, we have

$$\tilde{\phi}_{x}^{\delta^{-1}} \begin{pmatrix} I + \varpi S^{-1}AS & 0\\ 0 & I + \varpi R^{-1}AR \end{pmatrix} = \psi_{0}(\varpi \operatorname{tr}(R\tilde{\mathcal{B}}_{\delta}R^{-1}A)).$$
(5.2)

Proof. (a) Recall that $\phi_x(I + \omega A) = \psi_0(\omega \operatorname{tr}(\tilde{x}A))$. Since $X \in \omega M_2(\mathfrak{o}_2)$, for any $\delta \in \Omega$, we have

$$\sum_{X \in \omega M_2(\mathfrak{o}_2)} \tilde{\phi}_X^{\delta^{-1}} \begin{pmatrix} I & S^{-1}XR \\ 0 & I \end{pmatrix} \overline{\psi}(X) = \sum_{X \in \omega M_2(\mathfrak{o}_2)} \psi_0 \left(\omega \operatorname{tr}(R\tilde{L}_{\delta}S^{-1} - I)X \right)$$
$$= \begin{cases} |\omega M_2(\mathfrak{o}_2)| & \text{if } S = R\tilde{L}_{\delta}, \\ 0 & \text{otherwise.} \end{cases}$$

(b) Assume $S = R\tilde{L}_{\delta}$. The statement follows if we prove that the character

$$Q \mapsto \tilde{\phi}_{x}^{\delta^{-1}} \begin{pmatrix} I & \varpi \left(S^{-1}AQ - S^{-1}QR^{-1}AR \right) \\ 0 & I \end{pmatrix}$$

is the trivial character of
$$M_2(\mathfrak{o}_2)$$
. Since $\tilde{\phi}_x^{\delta^{-1}}|_{J_2^1} = \phi_{\delta^{-1}x\delta}|_{J_2^1}$, we get
 $\tilde{\phi}_x^{\delta^{-1}} \begin{pmatrix} I & \varpi(S^{-1}AQ - S^{-1}QR^{-1}AR) \\ 0 & I \end{pmatrix} = \psi_0 \left(\varpi \operatorname{tr}(\tilde{L}_{\delta}(S^{-1}AQ - S^{-1}QR^{-1}AR)) \right)$
 $= \psi_0 \left(\varpi \operatorname{tr}(R^{-1}AQ - R^{-1}QR^{-1}AR) \right)$
 $= 1.$

(c) Since $S = R\tilde{L}_{\delta}$, we get,

$$\begin{split} \tilde{\phi}_{x}^{\delta^{-1}} \begin{pmatrix} I + \varpi S^{-1}AS & 0\\ 0 & I + \varpi R^{-1}AR \end{pmatrix} &= \tilde{\phi}_{x}^{\delta^{-1}} \begin{pmatrix} I + \varpi \tilde{L}_{\delta}^{-1}R^{-1}AR\tilde{L}_{\delta} & 0\\ 0 & I + \varpi R^{-1}AR \end{pmatrix} \\ &= \psi_{0}(\varpi \operatorname{tr}(R\tilde{\mathcal{B}}_{\delta}R^{-1}A)). \end{split}$$

The last equality follows from the definition of ϕ_x .

Proposition 5.4. For $\delta \in \Omega$ with $\pi_{N,\psi}^{\delta} \neq 0$ and $A \in M_2(\mathfrak{o}_2)$, we have

$$\Theta_{\pi_{N,\psi}^{\delta}}(I+\varpi A) = \frac{q^{12}}{|\delta^{-1}T\delta \cap P|} \sum_{R \in \operatorname{GL}_2(\mathbb{F}_q)} \psi_0(\operatorname{tr}(R\mathcal{B}_{\delta}R^{-1}\bar{A})).$$

Proof. Using the sum over X, over Q and over (S, R) from Lemma 5.3 in Equation (5.1), we get

$$\Theta_{\pi^{\delta}_{N,\psi}}(I+\varpi A) = \frac{q^{12}}{q^8|\delta^{-1}T\delta\cap P|} \sum_{\substack{S,R\in\operatorname{GL}_2(\mathfrak{o}_2)\\S=R\tilde{L}_{\delta}}} \psi_0(\operatorname{\varpi tr}(R\tilde{\mathcal{B}}_{\delta}R^{-1}A)).$$

The above sum does not change if we replace *R* by *R'* and *S* by *S'* such that $R = R' \pmod{\varpi}$ and $S = S' \pmod{\varpi}$. Therefore, we get

$$\Theta_{\pi^{\delta}_{N,\psi}}(I + \varpi A) = \frac{q^{12} \cdot q^8}{q^8 |\delta^{-1}T\delta \cap P|} \sum_{R \in \operatorname{GL}_2(\mathbb{F}_q)} \psi_0(\operatorname{tr}(R\mathcal{B}_{\delta}R^{-1}\bar{A})).$$

Remark 5.5. Note that $|\delta^{-1}T\delta \cap P| = \begin{cases} (q^2 - 1)q^{12} & \text{if } \delta = I, \\ (q - 1)q^{12} & \text{if } \delta \neq I. \end{cases}$

We denote $A \sim B$ if the two matrices A and B are conjugate. Recall that any character of $M_2(\mathbb{F}_q)$ is of the form ϕ_B for $B \in M_2(\mathbb{F}_q)$. Since $M_2(\mathbb{F}_q) \cong J_2^1$ we write character of J_2^1 also as ϕ_B .

Corollary 5.6. Consider $\delta \in \Omega$ such that $\pi_{N,\psi}^{\delta} \neq 0$.

- (a) ϕ_B appears in $\pi^{\delta}_{N,\psi}|_{J^1_2}$ if and only if $B \sim \mathcal{B}_{\delta}$.
- (b) The multiplicity of $\bar{\phi}_{\mathcal{B}_{\delta}}$ in $\pi_{N,\psi}^{\delta}$ is given by

$$\left\langle \Theta_{\pi_{N,\psi}^{\delta}}, \phi_{\mathcal{B}_{\delta}} \right\rangle_{J_{2}^{1}} = \begin{cases} \frac{1}{q^{2}-1} \cdot |\operatorname{Stab}(2X_{1})| & \text{if } \delta = I, \\ \frac{1}{q-1} \cdot |\operatorname{Stab}(\mathcal{B}_{\delta})| & \text{if } \delta \neq I \end{cases}$$

where Stab(B) denotes the stabiliser of B under the conjugation action of $\text{GL}_2(\mathbb{F}_q)$ on $M_2(\mathbb{F}_q)$.

Proof. Using Proposition 5.4, we get

$$\langle \Theta_{\pi_{N,\psi}^{\delta}}, \phi_B \rangle_{J_2^1} = \frac{q^{12}}{|\delta^{-1}T\delta \cap P|} \cdot \frac{1}{q^4} \sum_{\substack{R \in \operatorname{GL}_2(\mathbb{F}_q)\\A \in M_2(\mathbb{F}_q)}} \psi_0(\operatorname{tr}(R\mathcal{B}_{\delta}R^{-1} - B)A).$$
(5.3)

Part (*a*) follows by observing that

$$\frac{1}{q^4} \cdot \sum_{A \in M_2(\mathbb{F}_q)} \psi_0(\operatorname{tr}(R\mathcal{B}_{\delta}R^{-1} - B)A) = \begin{cases} 0 & \text{if } R\mathcal{B}_{\delta}R^{-1} \neq B, \\ 1 & \text{if } R\mathcal{B}_{\delta}R^{-1} = B. \end{cases} \square$$

Part (*b*) follows from Remark 5.5 and Equation (5.3).

- **Remark 5.7.** (a) Under the considered embedding $GL_2(\mathfrak{o}_2) \hookrightarrow GL_4(\mathfrak{o}_2)$ the image of the center of $GL_2(\mathfrak{o}_2)$ is equal to the center of $GL_4(\mathfrak{o}_2)$ which are isomorphic to \mathfrak{o}_2^{\times} . We write Z for both the center of $GL_2(\mathfrak{o}_2)$ as well as that of $GL_4(\mathfrak{o}_2)$.
 - (b) Let the central character of representation π be ω_{π} i.e. $\Theta_{\pi}(z) = \omega_{\pi}(z) \dim(\pi) = \tilde{\phi}_{x}(z) \dim(\pi)$ for $z \in Z$. It can easily be seen that for $\delta \in \Omega$, if $\pi_{N,\psi}^{\delta} \neq 0$ then it has a central character which is equal to ω_{π} .

Proposition 5.4 describes the character of $\pi_{N,\psi}^{\delta}$ at elements of J_2^1 and now we describe the character of $\pi_{N,\psi}^{\delta}$ on remaining elements of $GL_2(\mathfrak{o}_2)$.

Theorem 5.8. Let $\delta \in \Omega$ be such that $\pi_{N,\psi}^{\delta} \neq 0$. Then,

(a) For $z \in Z$ and $k \in J_2^1$, we have $\Theta_{\pi_{N,\psi}^{\delta}}(z \cdot k) = \omega_{\pi}(z)\Theta_{\pi_{N,\psi}^{\delta}}(k)$. In particular, $\Theta_{\pi_{N,\psi}^{\delta}}(z) = \omega_{\pi}(z) \cdot \dim(\pi_{N,\psi}^{\delta}).$

(b) Let
$$\delta \neq I$$
. For $g \notin Z \cdot J_2^1$, we have $\Theta_{\pi_{M,\mu}^{\delta}}(g) = 0$.

- (c) If g is not conjugate to any element in $\mathfrak{D}_2^{\times} \cdot J_2^1$, then $\Theta_{\pi_{N,h}^I}(g) = 0$.
- (d) For $g \in (\mathfrak{O}_2^{\times} \cdot J_2^1) \setminus (Z \cdot J_2^1)$, we have

$$\Theta_{\pi_{N,\psi}^{I}}(g) = \frac{1}{q^{4}(q^{2}-1)} \left[\sum_{R \in \mathbb{N}(\mathfrak{O}_{2}^{\times} \cdot J_{2}^{1})} \tilde{\phi}_{x} \begin{pmatrix} R^{-1}gR & 0\\ 0 & R^{-1}gR \end{pmatrix} \right]$$

where $\mathsf{N}(\mathfrak{O}_2^{\times} \cdot J_2^1)$ is the normaliser of $\mathfrak{O}_2^{\times} \cdot J_2^1$ in $\mathrm{GL}_2(\mathfrak{o}_2)$.

Proof. Since $\pi^{\delta} = \operatorname{Ind}_{\delta^{-1}T\delta \cap P}^{P} \tilde{\phi}_{x}^{\delta^{-1}}$, we get

$$\Theta_{\pi^{\delta}}\begin{pmatrix}g & X\\ 0 & g\end{pmatrix} = \frac{1}{|\delta^{-1}T\delta \cap P|} \sum_{\gamma \in P} \tilde{\phi}_{x}^{\delta^{-1}} \left(\gamma^{-1}\begin{pmatrix}g & X\\ 0 & g\end{pmatrix}\gamma\right)$$
(5.4)

where the summation is over $\gamma = \begin{pmatrix} S & Q \\ 0 & R \end{pmatrix} \in P$ is such that

$$\gamma^{-1}\begin{pmatrix}g & X\\0 & g\end{pmatrix}\gamma = \begin{pmatrix}S^{-1}gS & S^{-1}gQ + S^{-1}XR - S^{-1}QR^{-1}gR\\0 & R^{-1}gR\end{pmatrix} \in \delta^{-1}T\delta \cap P.$$
(5.5)

- (a) It follows from the Remark 5.7 (b).
- (b) For $\delta \neq I$, $\delta^{-1}T\delta \cap P = Z \cdot (K_2^1 \cap P)$. By Equation (5.5), both $S^{-1}gS$ and $R^{-1}gR \in Z \cdot J_2^1$. Therefore, if *g* is not conjugate to $Z \cdot J_2^1$ (equivalently, $g \notin Z \cdot J_2^1$)

then there does not exist any $\gamma \in P$ such that $\gamma^{-1}\begin{pmatrix} g & X \\ 0 & g \end{pmatrix} \gamma \in (\delta^{-1}T\delta \cap P)$ and hence $\Theta_{\pi^{\delta}}\begin{pmatrix} g & X \\ 0 & g \end{pmatrix} = 0$. By Corollary 5.2, we conclude $\Theta_{\pi^{\delta}_{N,\psi}}(g) = 0$. (c) For $\delta = I$, $T \cap P = \Delta(\mathfrak{O}_2^{\times}) \cdot (K_2^1 \cap P)$ where $\Delta(\mathfrak{O}_2^{\times})$ is the diagonal embedding of \mathfrak{O}_2^{\times} in $GL_4(\mathfrak{o}_2)$. By Equation (5.5), both $S^{-1}gS$ and $R^{-1}gR \in \mathfrak{O}_2^{\times} \cdot J_2^1$. Therefore, if g is not conjugate to any element in $\mathfrak{O}_2^{\times} \cdot J_2^1$, then $\Theta_{\pi^I}\begin{pmatrix} g & X \\ 0 & g \end{pmatrix} = 0$. By Corollary 5.2, we conclude $\Theta_{\pi_{N,\psi}^{I}}(g) = 0.$

(d) Using Equation (5.5), if
$$\gamma^{-1}\begin{pmatrix}g&X\\0&g\end{pmatrix}\gamma\in\delta^{-1}T\delta\cap P$$
 then,
 $(S^{-1}gQ+S^{-1}XR-S^{-1}QR^{-1}gR)\in \mathscr{O}M_2(\mathfrak{o}_2).$

For $\delta = I$ and $g \in (\mathfrak{O}_2^{\times} \cdot J_2^1) \setminus (Z \cdot J_2^1)$, by Corollary 5.2, $\Theta_{\pi_{N,\psi}^I}(g)$ equals

$$\frac{1}{|N| \cdot |T \cap P|} \sum_{(S,R,Q,X) \in \mathcal{J}(g)} \tilde{\phi}_x \begin{pmatrix} S^{-1}gS & S^{-1}gQ + S^{-1}XR - S^{-1}QR^{-1}gR \\ 0 & R^{-1}gR \end{pmatrix} \overline{\psi}(Xg^{-1})$$
(5.6)

where

$$\mathcal{J}(g) = \left\{ \begin{array}{ll} S, R \in \operatorname{GL}_2(\mathfrak{o}_2) \text{ and } Q, X \in M_2(\mathfrak{o}_2), \\ (S, R, Q, X) : & S^{-1}gS, \ R^{-1}gR \in \mathfrak{O}_2^{\times} \cdot J_2^1 \text{ with } S^{-1}gS = R^{-1}gR \pmod{\omega}, \\ & S^{-1}gQ + S^{-1}XR - S^{-1}QR^{-1}gR \in \omega M_2(\mathfrak{o}_2) \end{array} \right\}.$$

The set $\mathcal{J}(g)$ describes the conditions on γ and X such that the matrix in (5.5) belongs to $T \cap P$.

Note that

$$S^{-1}gQ + S^{-1}XR - S^{-1}QR^{-1}gR = 0 \pmod{\omega}$$

$$\iff gQR^{-1} + X - QR^{-1}g = 0 \pmod{\omega}$$

$$\iff X = Q'g - gQ' \pmod{\omega},$$
(5.7)

where $Q' = QR^{-1}$. If Q'_1 and Q'_2 are two solutions of $X = Q'g - gQ' \pmod{\omega}$, then $(Q'_1 - Q'_2) \pmod{\omega}$ belongs to the centraliser of $g \pmod{\omega} \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. This gives $\mathcal{J}(g)$ consists of tuples (S, R, Q, X) such that

(i) $S, R \in \mathbb{N}(\mathfrak{D}_2^{\times} \cdot J_2^1)$ with $RS^{-1} \in \mathfrak{D}_2^{\times} \cdot J_2^1$. (ii) Q' belongs to the homogeneous space of centraliser of $g \pmod{\omega}$.

(iii) X (mod ω) belongs to $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \cdot \mathbb{F}_{q^2}$.

Hence,

$$|\{Q \in M_2(\mathfrak{o}_2) : (S, R, Q, X) \in \mathcal{J}(g)\}| = q^6 = |\{X \in M_2(\mathfrak{o}_2) : (S, R, Q, X) \in \mathcal{J}(g)\}|.$$
(5.8)
From Equation (5.6), $\Theta_{\pi^I_{N,\psi}}(g)$ is given by

$$\frac{1}{|N||T\cap P|} \sum_{(S,R,Q,X)\in\mathcal{J}(g)} \tilde{\varphi}_x \begin{pmatrix} S^{-1}gS & 0\\ 0 & R^{-1}gR \end{pmatrix} \tilde{\varphi}_x \begin{pmatrix} I & S^{-1}Q + S^{-1}g^{-1}XR - S^{-1}g^{-1}QR^{-1}gR \\ 0 & I \end{pmatrix} \overline{\psi}(Xg^{-1}).$$

The theorem follows easily from the following two lemmas, which can be verified by using the non-degeneracy of the trace pairing and using $R^{-1}gR(\text{mod } \omega)$, $RS^{-1}(\text{mod } \omega)$ and $X_3 \in \mathbb{F}_{q^2}^{\times}$. For more details, look at the PhD Thesis of first author [13].

Lemma 5.9. For fixed $R, S \in GL_2(\mathfrak{o}_2)$ such that $RS^{-1} \in \mathfrak{O}_2^{\times} \cdot J_2^1$, we have

$$\sum_{\substack{X,Q \in M_2(\mathfrak{o}_2) \text{ s.t.} \\ (S,R,Q,X) \in \mathcal{J}(g)}} \tilde{\phi}_x \begin{pmatrix} I & S^{-1}Q + S^{-1}g^{-1}XR - S^{-1}g^{-1}QR^{-1}gR \\ 0 & I \end{pmatrix} \overline{\psi}(Xg^{-1}) = q^{12}.$$

Lemma 5.10. Let $S, R \in \mathsf{N}(\mathfrak{D}_2^{\times} \cdot J_2^1)$ such that $SR^{-1} \in \mathfrak{D}_2^{\times} \cdot J_2^1$ then, for any $g \in \mathfrak{D}_2^{\times} \cdot J_2^1$ we have

$$\tilde{\phi}_x \begin{pmatrix} S^{-1}gS & 0\\ 0 & R^{-1}gR \end{pmatrix} = \tilde{\phi}_x \begin{pmatrix} R^{-1}gR & 0\\ 0 & R^{-1}gR \end{pmatrix}.$$

6. A description of $\pi_{N,\psi}$

Since $\pi_{N,\psi} = \bigoplus_{\delta \in \Omega_0} \pi_{N,\psi}^{\delta}$, we describe all the irreducible representations of $\operatorname{GL}_2(\mathfrak{o}_2)$ appearing in $\pi_{N,\psi}^{\delta}$ for all $\delta \in \Omega$.

Proposition 6.1. Let π be a strongly cuspidal representation of $GL_4(\mathfrak{o}_2)$. For $\delta \in \Omega$ with $\delta \neq I$ and an irreducible representation σ of $GL_2(\mathfrak{o}_2)$, if σ appears in $\pi_{N,\psi}^{\delta}$ then σ is a regular representation.

Proof. Let $B \in M_2(\mathbb{F}_q)$ be such that $\langle \Theta_{\sigma}, \phi_B \rangle_{J_2^1} \neq 0$. Since σ appears in $\pi_{N,\psi}^{\delta}$ then ϕ_B appears in $\pi_{N,\psi}^{\delta}|_{J_2^1}$. By Corollary 5.6, $B \sim \mathcal{B}_{\delta}$ and, therefore it is enough to prove that \mathcal{B}_{δ} is not a scalar matrix.

Now we assume that \mathcal{B}_{δ} is a scalar matrix and deduce a contradiction. Recall that $\operatorname{tr}(\mathcal{B}_{\delta}) = \operatorname{tr}(2X_1)$, then $\mathcal{B}_{\delta} = \begin{pmatrix} 2(a_0 + aa_1) & 0\\ 0 & 2(a_0 + aa_1) \end{pmatrix}$. Write $\mathcal{B}_{\delta} = L_{\delta}(X_1 + X_2Y)L_{\delta}^{-1} + X_1 - YX_2 = L_{\delta}\left(X_1 + X_2Y + L_{\delta}^{-1}(X_1 - YX_2)L_{\delta}\right)L_{\delta}^{-1}$.

This gives $X_1 + X_2Y + L_{\delta}^{-1}(X_1 - YX_2)L_{\delta}$ is diagonalizable and both the columns of L_{δ} are its eigenvectors with eigenvalue $2(a_0 + aa_1)$. Write

$$L_{\delta} = \begin{pmatrix} L_{\delta}^1 & L_{\delta}^2 \end{pmatrix}$$

where L^1_{δ} and L^1_{δ} are the columns of the matrix L_{δ} . Therefore,

$$\left(X_1 + X_2Y + L_{\delta}^{-1}(X_1 - YX_2)L_{\delta}\right) \cdot L_{\delta}^1 = 2(a_0 + aa_1) \cdot L_{\delta}^1.$$
(6.1)

$$\left(X_1 + X_2Y + L_{\delta}^{-1}(X_1 - YX_2)L_{\delta}\right) \cdot L_{\delta}^2 = 2(a_0 + aa_1) \cdot L_{\delta}^2.$$
(6.2)

Equation (6.1) gives

$$u = \frac{-2a_1(a_3(a^2 - b^2\alpha) + aa_2)}{a_2^2 + (3a^2 + b^2\alpha)a_3^2 + 4aa_2a_3} \text{ and } v = \frac{2a_1(a_2 + aa_3)}{a_2^2 + (3a^2 + b^2\alpha)a_3^2 + 4aa_2a_3}.$$
 (6.3)

Equation (6.2) gives two quadratic equations

$$Q_1(u,v) = 0 \text{ and } Q_2(u,v) = 0.$$
 (6.4)

If $a_1 = 0$, then Equation (6.3) gives u = v = 0 and hence $\delta = I$, which contradicts our assumption $\delta \neq I$.

Now we assume that $a_1 \neq 0$. Substituting the values of u, v from Equation (6.3) in Equations (6.4), one arrives at a contradiction to our assumption $x \in \mathbb{F}_{q^4} \setminus \mathbb{F}_{q^2}$, for which we skip the details.

Proposition 6.2. Let π be a strongly cuspidal representation of $GL_4(\mathfrak{o}_2)$. For $\delta \in \Omega$ and $\delta \neq I$, the representation $\pi_{N,\psi}^{\delta}$ is multiplicity free. In other words, if σ is a regular representation of $GL_2(\mathfrak{o}_2)$ such that $\omega_{\pi} = \omega_{\sigma}$ and $\phi_{\mathcal{B}_{\delta}}$ appears in $\sigma|_{J_2^1}$, then σ appears in $\pi_{N,\psi}^{\delta}$ with multiplicity one.

Proof. For $\delta \in \Omega$ and $\delta \neq I$, by Theorem 5.8 (b), we have $\Theta_{\pi_{N,\psi}^{\delta}}(g) = 0$ if $g \notin Z \cdot J_2^1$. Using $\omega_{\pi} = \omega_{\sigma}$ we get,

$$\begin{split} \left\langle \Theta_{\pi_{N,\psi}^{\delta}}, \Theta_{\sigma} \right\rangle_{\mathrm{GL}_{2}(\mathfrak{o}_{2})} &= \frac{1}{|\mathrm{GL}_{2}(\mathfrak{o}_{2})|} \sum_{g=z \cdot k \in Z \cdot J_{2}^{1}} \Theta_{\pi_{N,\psi}^{\delta}}(z \cdot k) \overline{\Theta}_{\sigma}(z \cdot k) \\ &= \frac{|Z/(J_{2}^{1} \cap Z)|}{|\mathrm{GL}_{2}(\mathfrak{o}_{2})|} \sum_{k \in J_{2}^{1}} \Theta_{\pi_{N,\psi}^{\delta}}(k) \overline{\Theta}_{\sigma}(k) \\ &= \frac{|Z/(J_{2}^{1} \cap Z)| \cdot |J_{2}^{1}|}{|\mathrm{GL}_{2}(\mathfrak{o}_{2})|} \left\langle \Theta_{\pi_{N,\psi}^{\delta}}, \Theta_{\sigma} \right\rangle_{J_{2}^{1}}. \end{split}$$

By Corollary 5.6, we have $\pi_{N,\psi}^{\delta}|_{J_2^1} = \bigoplus_{B \sim \mathcal{B}_{\delta}} \left(\frac{|\operatorname{Stab}(B)|}{q-1}\right) \phi_B$ and by Remark 2.3, it follows that $\sigma|_{J_2^1} = \bigoplus_{B \sim \mathcal{B}_{\delta}} \phi_B$. Therefore,

$$\left\langle \Theta_{\pi_{N,\psi}^{\delta}}, \Theta_{\sigma} \right\rangle_{J_{2}^{1}} = \left(\frac{|\operatorname{Stab}(\mathcal{B}_{\delta})|}{q-1} \right) \cdot |\operatorname{Conjugacy \ class \ of \ } \mathcal{B}_{\delta}|,$$

which gives $\left\langle \Theta_{\pi_{N,\psi}^{\delta}}, \Theta_{\sigma} \right\rangle_{\operatorname{GL}_{2}(\mathfrak{o}_{2})} = 1.$

Remark 6.3. Note that for $\delta \neq I$, the number of σ satisfying the property in Proposition 6.2 is $\frac{|\operatorname{Stab}(\mathcal{B}_{\delta})|}{q-1}$ and each having dimension $\frac{|\operatorname{GL}_2(\mathbb{F}_q)|}{|\operatorname{Stab}(\mathcal{B}_{\delta})|}$. This gives $\dim(\pi_{N,\psi}^{\delta}) = q(q^2 - 1) = \left(\frac{|\operatorname{Stab}(\mathcal{B}_{\delta})|}{q-1}\right) \cdot \left(\frac{|\operatorname{GL}_2(\mathbb{F}_q)|}{|\operatorname{Stab}(\mathcal{B}_{\delta})|}\right)$. Therefore, an irreducible representation σ appears in $\pi_{N,\psi}^{\delta}$ if and only if σ satisfies the properties mentioned in Proposition 6.2.

Lemma 6.4. The representation $\pi_{N,\psi}^{I} \cong \operatorname{Ind}_{\mathfrak{O}_{2}^{\times} \cdot J_{2}^{1}}^{\operatorname{GL}_{2}(\mathfrak{o}_{2})} \left(\tilde{\phi}_{x}|_{\mathfrak{O}_{2}^{\times} \cdot J_{2}^{1}} \right).$

Proof. It is enough to verify that the characters of the two representations are the same. The character of $\pi_{N,\psi}^{I}$ is given in Theorem 5.8. The character of $\operatorname{Ind}_{\mathfrak{D}_{2}^{\times} \cdot J_{2}^{1}}^{\operatorname{GL}_{2}(\mathfrak{o}_{2})} \left(\tilde{\phi}_{x} |_{\mathfrak{D}_{2}^{\times} \cdot J_{2}^{1}} \right)$ can be calculated using the formula for the character of an induced representation. We leave the verification to the reader that these two characters are the same.

Corollary 6.5. The representation $\pi_{N,\psi}^{I}$ is multiplicity free.

Proof. First consider $X_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. Then $\pi_{N,\psi}^I$ is an irreducible regular representation of non-split semisimple type and hence nothing to prove.

Now consider $X_1 \in \mathbb{F}_q$. By Clifford theory, there exists a character $\zeta : \operatorname{GL}_2(\mathfrak{o}_2) \to \mathbb{C}^{\times}$ such that $\zeta|_{J_2^1} = \phi_{-(2X_1)}$. Then $\theta' := (\theta \cdot \phi_{2X_1}) \cdot \zeta|_{\mathfrak{O}_2^{\times} \cdot J_2^1}$ is trivial on J_2^1 and therefore θ' factors through $(\mathfrak{O}_2^{\times} \cdot J_2^1)/J_2^1 \cong \mathbb{F}_{q^2}^{\times}$. Since $\pi_{N,\psi}^I \cong \operatorname{Ind}_{\mathfrak{O}_2^{\times} \cdot J_2^1}^{\operatorname{GL}(\mathfrak{o}_2)}(\theta \cdot \phi_{2X_1})$, we get $\pi_{N,\psi}^I \otimes \zeta \cong \operatorname{Ind}_{\mathfrak{O}_2^{\times} \cdot J_2^1}^{\operatorname{GL}(\mathfrak{o}_2)}(\theta')$ (see [5, Exercise 3.16]). It follows that $\pi_{N,\psi}^I \otimes \zeta$ is trivial on J_2^1 and hence factors through $\operatorname{GL}_2(\mathfrak{o}_2)/J_2^1 \cong \operatorname{GL}_2(\mathbb{F}_q)$. It is enough to prove that $\pi_{N,\psi}^I \otimes \zeta \cong \operatorname{Ind}_{\mathbb{F}_{q^2}^{\times}}^{\operatorname{GL}_2(\mathbb{F}_q)}(\theta')$ is multiplicity free representation as a representation of $\operatorname{GL}_2(\mathbb{F}_q)$. It follows from the fact that $(\operatorname{GL}_2(\mathbb{F}_q), \mathbb{F}_{q^2}^{\times})$ is a strong Gelfand pair which can be easily verified by looking at the character table of $\operatorname{GL}_2(\mathbb{F}_q)$. \Box

Theorem 6.6. For $\delta, \delta' \in \Omega$, $\mathcal{B}_{\delta} \sim \mathcal{B}_{\delta'}$ if and only if $T\delta P = T\delta' P$.

Proof. First, assume that $T\delta P = T\delta'P$. Then $\pi_{N,\psi}^{\delta} = \pi_{N,\psi}^{\delta'}$ and by Corollary 5.6, we get $\mathcal{B}_{\delta} \sim \mathcal{B}_{\delta'}$. For the converse, assume that $\mathcal{B}_{\delta} \sim \mathcal{B}_{\delta'}$. Then, $\det(\mathcal{B}_{\delta}) = \det(\mathcal{B}_{\delta'})$. It can be checked that for $\delta = A_{u,v}$, the determinant of \mathcal{B}_{δ} is given by

$$\det(\mathcal{B}_{\delta}) = 4(a_0 + aa_1)^2 - \left[\frac{4a_1^2(A) - ((u + av)^2 - (bv)^2\alpha) \cdot (a_2^2 + a_3^2(3a^2 + b^2\alpha) + 4aa_2a_3)^2}{\det(L_{\delta})}\right]$$

where A = A(u, v) is a quadratic polynomial in u and v.

Moreover, for $\delta = A_{u,v}$ and $\delta' = A_{u',v'}$ it can be deduced that the conditions for $\det(\mathcal{B}_{\delta}) = \det(\mathcal{B}_{\delta'})$ comes out to be the following for which we refer the reader to [13] and omit the details

$$M \cdot (a^2 a_3^2 + 2aa_2 a_3 + a_2^2 - a_3^2 b^2 \alpha) \cdot C(\delta, \delta') = 0,$$
(6.5)

where

$$M = 4a_1^4b^2\alpha - 4a_1^2(a^3a_3^2 + 2a^2a_2a_3 + aa_2^2 + 3aa_3^2b^2\alpha + 2a_2a_3b^2\alpha) + (a_2^2 + a_3^2(3a^2 + b^2\alpha) + 4aa_2a_3)^2.$$

Observe that the second factor in (6.5) is not zero, since

$$a^{2}a_{3}^{2} + 2aa_{2}a_{3} + a_{2}^{2} - a_{3}^{2}b^{2}\alpha = (a_{2} + aa_{3})^{2} - (a_{3}b)^{2}\alpha = 0 \implies a_{2} = 0, a_{3} = 0$$

and then $x \in \mathbb{F}_{q^2}$, a contradiction.

Now we claim that $M \neq 0$. Note that M is a quadratic polynomial in a_1^2 . If M = 0, then the discriminant of this quadratic polynomial is a square in \mathbb{F}_q . But, the discriminant is given by

$$16(a^2 - b^2\alpha) \left((a_2 + aa_3)^2 - (a_3b)^2\alpha \right)^2$$

which is not a square in \mathbb{F}_q (see the assumption (3.2)). This proves $M \neq 0$. Therefore, by Equation (6.5) we get $C(\delta, \delta') = 0$ and Lemma 3.6 gives $T\delta P = T\delta' P$. \Box

We have described each $\pi_{N,\psi}^{\delta}$ and proved that for distinct $\delta, \delta' \in \Omega_0$ the restriction of $\pi_{N,\psi}^{\delta}$ and $\pi_{N,\psi}^{\delta'}$ to J_2^1 are distinct, which gives the following result.

Corollary 6.7. For $\delta, \delta' \in \Omega_0$ with $\delta \neq \delta'$ the representations $\pi_{N,\psi}^{\delta}$ and $\pi_{N,\psi}^{\delta'}$ have no irreducible representation in common.

Theorem 6.8. Let π be a strongly cuspidal representation of $GL_4(\mathfrak{o}_2)$. Then $\pi_{N,\psi}$ is multiplicity free.

Proof. For $\delta, \delta' \in \Omega_0$ such that $\delta \neq \delta'$, by Corollary 6.7, $\pi_{N,\psi}^{\delta}$ and $\pi_{N,\psi}^{\delta'}$ have no irreducible representation in common. By Proposition 6.2 and Corollary 6.5 $\pi_{N,\psi}^{\delta}$ is multiplicity free for all $\delta \in \Omega_0$ and hence $\pi_{N,\psi}$ is multiplicity free.

Recall that $|\Omega_0| = q + 1$ (Lemma 3.1) and $|\{\delta \in \Omega_0 : \pi_{N,\psi}^{\delta} \neq 0\}| = q$ (Lemma 4.4). Therefore, there are exactly q conjugacy classes represented by \mathcal{B}_{δ} such that $\phi_{\mathcal{B}_{\delta}}$ appears in $\pi_{N,\psi}$ restricted to K_2^1 . Moreover, all the \mathcal{B}_{δ} 's have distinct determinant (Theorem 6.6) and then,

$$\{\det(\mathcal{B}_{\delta}): \delta \in \Omega_0\} = \mathbb{F}_q.$$

Note that the trace of each \mathcal{B}_{δ} is the same as tr(2 X_1). We conclude the following : (1) all the q-1 semisimple (both split and non-split) conjugacy classes with trace tr(2 X_1) are represented by \mathcal{B}_{δ} for some $\delta \in \Omega_0$.

(2) The determinant of the scalar matrix and the split non-semisimple matrices with trace $tr(2X_1)$ are the same and exactly one of these two conjugacy classes will be represented by \mathcal{B}_{δ} for some $\delta \in \Omega_0$.

We summarise the results discussed in Section 6 in the theorem below.

Theorem 6.9. Let $\pi = \operatorname{Ind}_{I(\phi_x)}^{\operatorname{GL}_4(\mathfrak{o}_2)} \tilde{\phi}_x$ be a strongly cuspidal representation of $\operatorname{GL}_4(\mathfrak{o}_2)$.

- (a) Let $X_1 \in \mathbb{F}_q$, then $\pi_{N,\psi}$ consists of the following representations of $GL_2(\mathfrak{o}_2)$.
 - (i) The representation $\pi_{N,\psi}^{I} = \operatorname{Ind}_{\mathfrak{O}_{2}^{\times} \cdot J_{2}^{1}}^{\operatorname{GL}_{2}(\mathfrak{o}_{2})}(\tilde{\phi}_{2X_{1}})$, which is not a regular representation and it is also not irreducible.
 - (ii) An irreducible regular representation σ of $GL_2(\mathfrak{o}_2)$ with multiplicity one such that $\omega_{\sigma} = \omega_{\pi}$ and $\sigma|_{J_2^1}$ contains the character ϕ_B with B a semisimple matrix of trace tr(2X₁).
- (b) Let $X_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_{q}$, then $\pi_{N,\psi}$ consists of the following representations of $\operatorname{GL}_2(\mathfrak{o}_2)$.
 - (i) The irreducible regular representation $\pi_{N,\psi}^{I} = \operatorname{Ind}_{\mathfrak{O}_{2}^{\times} \cdot J_{2}^{1}}^{\operatorname{GL}_{2}(\mathfrak{o}_{2})}(\tilde{\phi}_{2X_{1}})$ with multiplicity one.
 - (ii) An irreducible regular representation σ of $GL_2(\mathfrak{o}_2)$ with multiplicity one such that $\omega_{\sigma} = \omega_{\pi}$ and $\sigma|_{J_2^1}$ contains the character ϕ_B with *B* a regular matrix of trace $\operatorname{tr}(2X_1)$ and $B \not\sim 2X_1$.

7. A description of
$$\Pi = \operatorname{Ind}_{\mathfrak{G}_2^{\times}}^{\operatorname{GL}_2(\mathfrak{o}_2)}(\theta|_{\mathfrak{G}_2^{\times}})$$

Recall from Section 3 that we have fixed $a, b, \alpha \in \mathbb{F}_q$ such that α is a square free element in \mathbb{F}_q and $a + b\sqrt{\alpha}$ is a square free element in $\mathbb{F}_{q^2} = \mathbb{F}_q(\sqrt{\alpha})$. We have also fixed an embedding $\mathbb{F}_{q^2} \hookrightarrow M_2(\mathbb{F}_q)$ as given in (3.4). We state the following easy lemma without proof. It provides a set of representatives for various double cosets, which will be useful later in this section.

Lemma 7.1. For
$$y \in \mathbb{F}_q$$
, $z \in \mathbb{F}_q^{\times}$ write $\gamma(y, z) = \begin{pmatrix} 1 & y \\ 0 & z \end{pmatrix}$ and

$$\Gamma = \left\{ \gamma(y, z) : y \in \mathbb{F}_q, z \in \mathbb{F}_q^{\times} \right\}.$$

(a) The set Γ is a set of distinct representatives of $\mathbb{F}_{a^2}^{\times} \setminus \mathrm{GL}_2(\mathbb{F}_q)$.

(b) For
$$\gamma(y,z)$$
 and $\gamma(y',z') \in \Gamma$, $\mathbb{F}_{q^2}^{\times} \gamma(y,z) \mathbb{F}_{q^2}^{\times} = \mathbb{F}_{q^2}^{\times} \gamma(y',z') \mathbb{F}_{q^2}^{\times}$ if and only if

$$\frac{y^2 + 2ay(z-1) + (1+z^2)(a^2 - b^2\alpha)}{z} = \frac{y'^2 + 2ay'(z'-1) + (1+z'^2)(a^2 - b^2\alpha)}{z'}$$

We fix a subset $\Gamma_0 \subset \Gamma$ *representing distinct double cosets for* $\mathbb{F}_{q^2}^{\times} \setminus \mathrm{GL}_2(\mathbb{F}_q) / \mathbb{F}_{q^2}^{\times}$.

Remark 7.2. Let H, K be subgroups of $G = GL_2(\mathfrak{o}_2)$. Assume that the quotient map $G \to \overline{G} = GL_2(\mathbb{F}_q)$ induces a bijection between the double cosets $K \setminus G/H$ and $\overline{K} \setminus \overline{G}/\overline{H}$, where $\overline{H}, \overline{K}$ denote the images of H, K in \overline{G} . Then, for a representative $\gamma \in \overline{G}$ of a double coset in $\overline{K} \setminus \overline{G}/\overline{H}$, we write its lift to G by the same letter γ as a representative of the corresponding double coset in $K \setminus G/H$.

The following proposition describes the characters ϕ_B of $J_2^1 \cong M_2(\mathbb{F}_q)$ which appear in the restriction of Π to J_2^1 .

Proposition 7.3. For $B \in M_2(\mathbb{F}_q)$, ϕ_B appears in $\Pi|_{I_2^1}$ if and only if $\exists m, n \in \mathbb{F}_q$ such that

$$B \sim 2X_1 - \begin{pmatrix} m & n(a^2 - b^2 \alpha) + 2am \\ n & -m \end{pmatrix}.$$

Proof. Recall that the character $\phi_B : M_2(\mathbb{F}_q) \to \mathbb{C}^{\times}$ is given by $\phi_B(A) = \psi_0(\operatorname{tr}(BA))$. Since Π is an induced representation, by Mackey theory we have

$$\Pi|_{J_2^1} = \bigoplus_{\gamma \in \mathfrak{O}_2^{\times} \setminus \operatorname{GL}_2(\mathfrak{o}_2)/J_2^1} \operatorname{Ind}_{\gamma^{-1}\mathfrak{O}_2^{\times}\gamma \cap J_2^1}^{J_2^1} \left(\theta^{\gamma^{-1}}\right)$$

Using Frobenius reciprocity, we get

$$\left\langle \Theta_{\Pi}, \phi_{B} \right\rangle_{J_{2}^{1}} = \sum_{\gamma \in \mathfrak{O}_{2}^{\times} \setminus \operatorname{GL}_{2}(\mathfrak{o}_{2})/J_{2}^{1}} \left\langle \theta^{\gamma^{-1}}, \phi_{B} \right\rangle_{\gamma^{-1}(\mathfrak{O}_{2}^{\times})\gamma \cap J_{2}^{1}}.$$
(7.1)

Since J_2^1 is a normal subgroup, $\gamma^{-1}(\mathfrak{O}_2^{\times})\gamma \cap J_2^1 = \gamma^{-1}(\mathfrak{O}_2^{\times} \cap J_2^1)\gamma = \gamma^{-1}K\gamma$ where $K = \mathfrak{O}_2^{\times} \cap J_2^1$. Note that there is a bijection between $\mathfrak{O}_2^{\times} \backslash \mathrm{GL}_2(\mathfrak{o}_2)/J_2^1$ and $\mathbb{F}_{q^2}^{\times} \backslash \mathrm{GL}_2(\mathbb{F}_q)$. Using the Lemma 7.1 (a) we get,

$$\left\langle \Theta_{\Pi}, \phi_B \right\rangle_{J_2^1} = \sum_{\gamma \in \Gamma} \left\langle \theta^{\gamma^{-1}}, \phi_B \right\rangle_{\gamma^{-1}(K)\gamma} = \sum_{\gamma \in \Gamma} \left\langle \theta, \phi_B^{\gamma} \right\rangle_K.$$
(7.2)

Note that $\theta|_{K} = \phi_{2X_1}|_{K}$, which gives

$$\langle \theta, \phi_B^{\gamma} \rangle_K = \langle \phi_{2X_1}, \phi_B^{\gamma} \rangle_K = \frac{1}{|K|} \sum_{A \in K} \psi_0(\operatorname{tr}(2X_1A)) \overline{\psi_0}(\operatorname{tr}(\gamma B \gamma^{-1}A))$$
$$= \frac{1}{|K|} \sum_{A \in K} \psi_0(\operatorname{tr}(2X_1 - \gamma B \gamma^{-1})A).$$
(7.3)

We know that tr : $M_2(\mathbb{F}_q) \times M_2(\mathbb{F}_q) \to \mathbb{F}_q$ is a non-degenerate bilinear map. Using the isomorphism $J_2^1 \cong M_2(\mathbb{F}_q)$, *K* is identified with $\left\{ \begin{pmatrix} b_0 & -b_1(a^2 - b^2\alpha) \\ b_1 & b_0 + 2ab_1 \end{pmatrix} : b_0, b_1 \in \mathbb{F}_q \right\}$. It can be checked that

$$\left\{C \in M_2(\mathbb{F}_q) : \operatorname{tr}(CD) = 0 \ \forall \ D \in K\right\} = \left\{ \begin{pmatrix} m & n(a^2 - b^2\alpha) + 2am \\ n & -m \end{pmatrix} : m, n \in \mathbb{F}_q \right\}.$$

Note that $\langle \theta, \phi_B^{\gamma} \rangle_K$ is either 0 or 1. Moreover, $\langle \theta, \phi_B^{\gamma} \rangle_K = 1$ if and only if

$$\exists m, n \in \mathbb{F}_q \text{ such that } 2X_1 - \gamma B \gamma^{-1} = \begin{pmatrix} m & n(a^2 - b^2 \alpha) + 2am \\ n & -m \end{pmatrix}.$$
(7.4)

Therefore, ϕ_B appears in $\Pi|_{J_2^1}$ if and only if *B* satisfies (7.4) for some $\gamma \in \Gamma$ and $m, n \in \mathbb{F}_q$.

Corollary 7.4. If $X_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, then ϕ_{2X_1} appears in $\Pi|_{J_2^1}$ with multiplicity one.

Proof. Using Equation (7.2), (7.3) and (7.4) for $B = 2X_1$, we have

$$\begin{split} \left\langle \Theta_{\Pi}, \phi_{2X_{1}} \right\rangle_{J_{2}^{1}} &= \frac{1}{|K|} \sum_{\gamma \in \Gamma, \ A \in K} \psi_{0}(\operatorname{tr}(2X_{1} - \gamma 2X_{1}\gamma^{-1})A) \\ &= \left| \left\{ \gamma \in \Gamma : \ 2X_{1} - \gamma 2X_{1}\gamma^{-1} \in \left\{ \begin{pmatrix} m & n(a^{2} - b^{2}\alpha) + 2am \\ n & -m \end{pmatrix} : m, n \in \mathbb{F}_{q} \right\} \right\} \right| \\ &= \left| \left\{ \gamma(y, z) \in \Gamma : \frac{y^{2} + 2ay(z - 1) + (1 + z^{2})(a^{2} - b^{2}\alpha)}{z} = 2(a^{2} - b^{2}\alpha) \right\} \right| \\ &= 1. \end{split}$$

Lemma 7.5. Let $B = 2X_1 - \begin{pmatrix} m & n(a^2 - b^2\alpha) + 2am \\ n & -m \end{pmatrix}$ for some $m, n \in \mathbb{F}_q$.

(a) If $X_1 \in \mathbb{F}_q$, then either $B = 2X_1$ or B is a semisimple matrix with trace tr $(2X_1)$.

(b) If $X_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, then B is a regular matrix with trace $\operatorname{tr}(2X_1)$.

Proof. It is clear that $tr(B) = tr(2X_1)$. The characteristic polynomial of *B* is given by

$$(\lambda - 2(a_0 + aa_1))^2 - \left(4a_1^2b^2\alpha + (m + an)^2 - b^2n^2\alpha\right)$$

Clearly, if $D := 4a_1^2b^2\alpha + (m + an)^2 - b^2n^2\alpha \neq 0$, then *B* will be semisimple. If D = 0, then *B* is either scalar or split non-semisimple with trace tr(2*X*₁).

- (a) Let $X_1 \in \mathbb{F}_q$ i.e. $a_1 = 0$. Then, $D = 0 \implies m = 0, n = 0 \implies B = 2X_1$ which is scalar.
- (b) Let $X_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, i.e. $a_1 \neq 0$. If D = 0 and B is the scalar matrix with trace tr($2X_1$) then by comparing the entries we get $m = -2aa_1, n = 2a_1$ and $4a_1b^2\alpha = 0$. Since $a_1 \neq 0$, $4a_1b^2\alpha = 0$ is not possible, a contradiction. In fact, the choice $m = -2aa_1, n = 2a_1$ gives a split non-semisimple regular matrix. \Box

Let ϕ_B be a character of J_2^1 and $\sigma = \operatorname{Ind}_{I(\phi_B)}^{\operatorname{GL}_2(\mathfrak{o}_2)} \tilde{\phi}_B$. By Mackey theory, it follows that

$$\langle \Theta_{\Pi}, \Theta_{\sigma} \rangle_{\mathrm{GL}_{2}(\mathfrak{o}_{2})} = \langle \theta, \Theta_{\sigma} \rangle_{\mathfrak{G}_{2}^{\times}} = \sum_{\gamma \in \mathfrak{G}_{2}^{\times} \backslash \mathrm{GL}_{2}(\mathfrak{o}_{2})/I(\phi_{B})} \langle \theta, \tilde{\phi}_{B}^{\gamma} \rangle_{\gamma(I(\phi_{B}))\gamma^{-1} \cap \mathfrak{G}_{2}^{\times}}.$$
(7.5)

Lemma 7.6. Let $B \in M_2(\mathbb{F}_q)$ be a non-split semisimple matrix with trace $\operatorname{tr}(2X_1)$. Let $\tilde{\phi}_B : I(\phi_B) \to \mathbb{C}^{\times}$ be an extension of ϕ_B and $\sigma = \operatorname{Ind}_{I(\phi_B)}^{\operatorname{GL}_2(\mathfrak{o}_2)} \tilde{\phi}_B$. Then,

$$\langle \Theta_{\Pi}, \Theta_{\sigma} \rangle_{\mathrm{GL}_{2}(\mathfrak{o}_{2})} = \langle \theta, \tilde{\phi}_{B} \rangle_{\mathfrak{H}_{2}^{\times}} + \langle \theta, \tilde{\phi}_{B}^{\gamma_{0}} \rangle_{\mathfrak{H}_{2}^{\times}} + \sum_{\substack{\gamma \in \Gamma_{0} \text{ such that} \\ \gamma \notin \{I, \gamma_{0}\}}} \langle \tilde{\phi}_{x}, \tilde{\phi}_{B}^{\gamma} \rangle_{Z \cdot K}.$$
(7.6)

where $\gamma_0 \in \Gamma_0$ is a non-trivial element in $N_{GL_2(\mathbb{F}_q)}(\mathbb{F}_{q^2}^{\times})$ the normalizer of $\mathbb{F}_{q^2}^{\times}$, and $K = I + \omega \mathfrak{O}_2 \cong \mathbb{F}_{q^2}$.

Proof. We use Equation (7.5) to prove the lemma. Note that $I(\phi_B) = \mathfrak{O}_2^{\times} \cdot J_2^1$ and if we write $K = I + \mathfrak{O}\mathfrak{O}_2$, then

$$\gamma \left(\mathfrak{O}_{2}^{\times} \cdot J_{2}^{1}\right) \gamma^{-1} \cap \mathfrak{O}_{2}^{\times} = \begin{cases} \mathfrak{O}_{2}^{\times} & \text{if } \gamma \in \mathsf{N}_{\mathrm{GL}_{2}(\mathfrak{o}_{2})}(\mathfrak{O}_{2}^{\times} \cdot J_{2}^{1}) \\ Z \cdot K & \text{if } \gamma \notin \mathsf{N}_{\mathrm{GL}_{2}(\mathfrak{o}_{2})}(\mathfrak{O}_{2}^{\times} \cdot J_{2}^{1}). \end{cases}$$

Note that there is a bijection between $\mathfrak{D}_2^{\times} \backslash \mathrm{GL}_2(\mathfrak{o}_2)/\mathfrak{D}_2^{\times} \cdot J_2^1$ and $\mathbb{F}_{q^2}^{\times} \backslash \mathrm{GL}_2(\mathbb{F}_q)/\mathbb{F}_{q^2}^{\times}$ for which a set of representatives Γ_0 are given in Lemma 7.1. Since the index of $\mathbb{F}_{q^2}^{\times}$ in $\mathsf{N}_{\mathrm{GL}_2(\mathbb{F}_q)}(\mathbb{F}_{q^2}^{\times})$ is 2, we get the index of \mathfrak{D}_2^{\times} in $\mathsf{N}_{\mathrm{GL}_2(\mathfrak{o}_2)}(\mathfrak{D}_2^{\times})$ is 2. Now, the lemma follows by observing that $\theta|_{Z \cdot K} = \tilde{\phi}_x|_{Z \cdot K}$.

Corollary 7.7. Let $\tilde{\phi}_{2X_1} : I(\phi_{2X_1}) \to \mathbb{C}^{\times}$ be given by $\tilde{\phi}_{2X_1} = \theta \cdot \phi_{2X_1}$ and $\sigma = \operatorname{Ind}_{\mathfrak{G}_2^{\times} J_2^1}^{\operatorname{GL}_2(\mathfrak{o}_2)}(\theta \cdot \phi_{2X_1})$, then σ is a subrepresentation of Π . Moreover, if $X_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, then σ is an irreducible representation of $\operatorname{GL}_2(\mathfrak{o}_2)$ and σ appears in Π with multiplicity one.

Proof. First part follows easily from the transitivity of induction. For $X_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_{q}$, $B = 2X_1$ is non-split semisimple, we use Equation (7.6) to compute $\langle \Theta_{\Pi}, \Theta_{\sigma} \rangle_{\mathrm{GL}_2(\mathfrak{o}_2)}$. Since $\tilde{\phi}_{2X_1} = \theta \cdot \phi_{2X_1}$ and θ is a strongly primitive character of \mathfrak{O}_2^{\times} , we get

$$\langle \theta, \tilde{\phi}_{2X_1} \rangle_{\mathfrak{G}_2^{\times}} = 1 \text{ and } \left\langle \theta, \tilde{\phi}_{2X_1}^{\gamma_0^{-1}} \right\rangle_{\mathfrak{G}_2^{\times}} = \left\langle \theta^{\gamma_0}, \tilde{\phi}_{2X_1} \right\rangle_{\mathfrak{G}_2^{\times}} = \langle \theta^{\gamma_0}, \theta \rangle_{\mathfrak{G}_2^{\times}} = 0$$

It remains to prove that $\langle \theta, \tilde{\phi}_{2X_1}^{\gamma^{-1}} \rangle_{Z \cdot K} = 0$ for $\gamma \notin \{I, \gamma_0\}$, which can be easily verified using Equation (7.4) and we skip the details.

Corollary 7.8. Let $B \in M_2(\mathbb{F}_q)$ be a non-split semisimple matrix with trace $\operatorname{tr}(2X_1)$ and $B \nsim 2X_1$. Let $\tilde{\phi}_B$ be an extension of ϕ_B to its inertia group $I(\phi_B)$ such that $\tilde{\phi}_B|_Z = \omega_{\Pi}$ and $\sigma = \operatorname{Ind}_{I(\phi_B)}^{\operatorname{GL}_2(\mathfrak{o}_2)} \tilde{\phi}_B$. Then, σ appears in Π with multiplicity one.

Proof. Since *B* is non-split semisimple matrix, we can take that $B \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. We use Equation (7.6) to prove $\langle \Theta_{\Pi}, \Theta_{\sigma} \rangle_{\mathrm{GL}_2(\mathfrak{o}_2)} = 1$. We first claim that $\langle \theta, \tilde{\phi}_B \rangle_{\mathfrak{O}_2^{\times}} = 0$,

 $\left\langle \theta, \tilde{\phi}_B^{\gamma_0^{-1}} \right\rangle_{\mathfrak{D}_2^{\times}} = 0.$ Since $1 + \mathfrak{O}\mathfrak{O}_2 \subset \mathfrak{O}_2^{\times}$, it is enough to prove that $\langle \theta, \phi_B \rangle_{1+\mathfrak{O}\mathfrak{O}_2} = 0$ and $\left\langle \theta, \phi_B^{\gamma_0^{-1}} \right\rangle_{1+\mathfrak{O}\mathfrak{O}_2} = 0.$ We identify $1 + \mathfrak{O}\mathfrak{O}_2$ with \mathbb{F}_{q^2} under the isomorphism $J_2^1 \cong M_2(\mathbb{F}_q)$. Recall that tr : $M_2(\mathbb{F}_q) \times M_2(\mathbb{F}_q) \to \mathbb{F}_q$ is a non-degenerate bilinear map and so is its restriction to \mathbb{F}_{q^2} . Therefore, $B_1, B_2 \in \mathbb{F}_{q^2}$ satisfy the following

$$\phi_{B_1}|_{\mathbb{F}_{q^2}} = \phi_{B_2}|_{\mathbb{F}_{q^2}}$$
 if and only if $B_1 = B_2$.

Note that $\theta|_{1+\mathfrak{OD}_2} = \phi_{2X_1}|_{1+\mathfrak{OD}_2}$. Since $B \approx 2X_1$, we get $\phi_B|_{1+\mathfrak{OD}_2} \neq \phi_{2X_1}|_{1+\mathfrak{OD}_2}$ and $\phi_B^{\gamma_0^{-1}}|_{1+\mathfrak{OD}_2} \neq \phi_{2X_1}|_{1+\mathfrak{OD}_2}$. Hence the claim follows.

One can easily verify that there exists unique $\gamma \in \Gamma_0 \setminus \{I, \gamma_0\}$ with $\left\langle \tilde{\phi}_x, \tilde{\phi}_B^{\gamma^{-1}} \right\rangle_{Z \cdot K} = 1$, for which we skip the details.

Lemma 7.9. Let $B = \begin{pmatrix} m_1 & 0 \\ 0 & 4(a_0 + aa_1) - m_1 \end{pmatrix}$ be a split semisimple matrix such that $m_1 \neq 2(a_0 + aa_1)$. Let $\tilde{\phi}_B$ be an extension of ϕ_B to its inertia group $I(\phi_B)$ such that $\tilde{\phi}_B|_Z = \omega_{\Pi}$ and $\sigma = \operatorname{Ind}_{I(\phi_B)}^{\operatorname{GL}_2(\mathfrak{o}_2)} \tilde{\phi}_B$. Then, σ appears in Π with multiplicity one.

Proof. We use Equation (7.5) by noting that for the given *B*, the inertia group $I(\phi_B)$ is such that $I(\phi_B)/J_2^1$ is the diagonal torus T_1 in $\operatorname{GL}_2(\mathbb{F}_q)$. Moreover, there is a bijection between $\mathfrak{O}_2^{\times} \setminus \operatorname{GL}_2(\mathfrak{o}_2)/I(\phi_B)$ and $\mathbb{F}_{q^2}^{\times} \setminus \operatorname{GL}_2(\mathbb{F}_q)/T_1$ for which a set of representatives Γ_1 is known from Lemma 7.1 (c). One can easily verify that that there exists a unique $\gamma \in \Gamma_1$ satisfying $\langle \theta, \tilde{\phi}_B^{\gamma} \rangle_{\gamma I(\phi_B)\gamma^{-1} \cap \mathfrak{O}_2^{\times}} = 1$.

Lemma 7.10. Let $X_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ and $B = \begin{pmatrix} 2(a_0 + aa_1) & 1 \\ 0 & 2(a_0 + aa_1) \end{pmatrix}$. Let $\tilde{\phi}_B$ be an extension of ϕ_B to its inertia group $I(\phi_B)$ such that $\tilde{\phi}_B|_Z = \omega_{\Pi}$ and $\sigma = \operatorname{Ind}_{I(\phi_B)}^{\operatorname{GL}_2(\mathfrak{o}_2)} \tilde{\phi}_B$. Then, σ appears in Π with multiplicity one.

Proof. For the given *B* the inertia group $I(\phi_B)$ is such that $I(\phi_B)/J_2^1 = T_2$ as defined in Lemma 7.1. The rest of the proof is similar to that of Lemma 7.9.

We summarise the results from Section 7 in the following theorem.

Theorem 7.11. Let $\Pi = \operatorname{Ind}_{\mathfrak{D}_2^{\times}}^{\operatorname{GL}_2(\mathfrak{o}_2)}(\theta|_{\mathfrak{D}_2^{\times}})$. Recall that $\theta = \phi_{2X_1}$ when restricted to $\mathfrak{D}_2^{\times} \cap J_2^1 \cong 1 + \omega \mathfrak{D}_2$.

- (a) Let $X_1 \in \mathbb{F}_q$, then Π consists of the following representations of $GL_2(\mathfrak{o}_2)$.
 - (*i*) The representation $\operatorname{Ind}_{\mathfrak{O}_{2}^{\times},J_{2}^{1}}^{\operatorname{GL}_{2}(\mathfrak{o}_{2})}(\tilde{\phi}_{2X_{1}})$, which is not a regular representation and it is also not irreducible.
 - (ii) An irreducible regular representation σ of $GL_2(\mathfrak{o}_2)$ with multiplicity one such that $\omega_{\sigma} = \omega_{\Pi}$ and $\sigma|_{J_2^1}$ contains the character ϕ_B where B is a semisimple matrix of trace tr(2X₁).
- (b) Let $X_1 \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, then Π consists of the following representations of $GL_2(\mathfrak{o}_2)$.
 - (*i*) The irreducible regular representation $\operatorname{Ind}_{\mathfrak{O}_{2}^{\times} \cdot J_{2}^{1}}^{\operatorname{GL}(\mathfrak{o}_{2})}(\tilde{\phi}_{2X_{1}})$ with multiplicity one.
 - (ii) An irreducible regular representation σ of $GL_2(\mathfrak{o}_2)$ with multiplicity one such that $\omega_{\sigma} = \omega_{\Pi}$ and $\sigma|_{J_2^1}$ contains the character ϕ_B where B is a regular matrix of trace tr(2X₁) and $B \approx 2X_1$.

Using Theorem 6.9 and Theorem 7.11, we obtain our main result verifying the Conjecture 1.4 of Prasad for the case n = 2, l = 2.

Theorem 7.12. Let π be an irreducible strongly cuspidal representation of $GL_4(\mathfrak{o}_2)$ associated to a strongly primitive character $\theta : \mathcal{O}_2^{\times} \to \mathbb{C}^{\times}$. Then as a representation of $GL_2(\mathfrak{o}_2)$ we have

$$\pi_{N,\psi} \cong \operatorname{Ind}_{\mathfrak{G}_2^{\times}}^{\operatorname{GL}_2(\mathfrak{o}_2)}(\theta|_{\mathfrak{G}_2^{\times}}).$$

Acknowledgement

The authors thank Dipendra Prasad for several helpful conversations and insightful comments. This work is a part of AP's Thesis at IIT Delhi and she acknowledges the support of PMRF fellowship (1400772). SPP acknowledges the support of the SERB MATRICS grant (MTR/2022/000782).

References

- [1] A. M. Aubert, U. Onn, A. Prasad, and A. Stasinski. On cuspidal representations of general linear groups over discrete valuation rings. *Israel Journal of Mathematics*, 175:391–420, 2010.
- [2] K. Balasubramanian, A. Dangodara, and H. Khurana. On a twisted jacquet module of GL(2n) over a finite field. 2022. arXiv : 2206.03024.
- [3] K. Balasubramanian and H. Khurana. A certain twisted Jacquet module of GL(4) over a finite field. *J. Pure Appl. Algebra*, 226(5):Paper No. 106932, 16, 2022.
- [4] K. Balasubramanian and H. Khurana. A certain twisted Jacquet module of GL(6) over a finite field: the rank 2 case. *J. Pure Appl. Algebra*, 228(6):14–26, 2024.
- [5] W. Fulton and J. Harris. Representation theory a first course. *Springer New York*, *NY*, 129, 1991.
- [6] W. T. Gan, B. H. Gross, and D. Prasad. Symplectic local root numbers, central critical *L* values, and restriction problems in the representation theory of classical groups. *Astérisque*, (346):1–109, 2012.
- [7] P. Gérardin. *Construction de séries discrètes p-adiques,* volume Vol. 462 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1975.
- [8] J. A. Green. The characters of the finite general linear groups. *Trans. Amer. Math. Soc.*, 80:402–447, 1955.
- [9] G. Hill. Regular elements and regular characters of $GL_n(\mathfrak{o})$. *Journal of Algebra*, 174:610–635, 1995.
- [10] R. Krakovski, U. Onn, and P. Singla. Regular characters of groups of type *A_n* over discrete valuation rings. *Journal of Algebra*, 496:116–137, 2018.
- [11] G. Lusztig. Representations of reductive groups over finite rings. *Representation Theory. An Electronic Journal of the American Mathematical Society*, 8:1–14, 2004.
- [12] S. K. Pandey and C. G. Venketasubramanian. Structure of twisted jacquet modules of principal series representations of $sp_4(f)$. arXiv : 2401.16933, 2024.
- [13] A. Parashar. On degenerate Whittaker space for $GL_4(\mathfrak{o}_2)$. PhD thesis, IIT Delhi, In preparation.
- [14] S. P. Patel and P. Singla. Gelfand criterion and multiplicity one results for GL_n over finite chain rings. *Indian J. Pure Appl. Math.*, 52(3):761–772, 2021.
- [15] S. P. Patel and P. Singla. A multiplicity one theorem for groups of type *A_n* over discrete valuation rings. *Proc. Amer. Math. Soc.*, 150(6):2309–2322, 2022.
- [16] D. Prasad. The space of degenerate whittaker models for general linear groups over a finite field. *International Mathematics Research Notices*, 11:579–594, 2000.
- [17] D. Prasad. The space of degenerate Whittaker models for GL(4) over *p*-adic fields. In *Cohomology of arithmetic groups, L-functions and automorphic forms (Mumbai, 1998/1999)*, pages 103–115. Tata Inst. Fund. Res., Bombay, 2001.
- [18] A. Stasinski. Representations of GL_N over finite local principal ideal rings: an overview. In *Around Langlands correspondences*, volume 691 of *Contemp. Math.*, pages 337–358. Amer. Math. Soc., Providence, RI, 2017.
- [19] A. Stasinski and S. Stevens. The regular representations of GL_n over finite local principal ideal rings. *Bulletin of the London Mathematical Society*, pages 1066–1084, 2017.

Department of Mathematics, IIT Delhi, Hauz Khas, New Delhi - 110016 Email address: ankitaparashar216@gmail.com

Department of Mathematics, IIT Delhi, Hauz Khas, New Delhi - 110016

DEPARTMENT OF MATHEMATICS, IIT DHARWAD, CHIKKAMALLIGAWAD VILLAGE, DHARWAD - 580007 Email address: shivprakashpatel@gmail.com