

A framework for semi-universality: Semi-universality of 3-qudit $SU(d)$ -invariant gates

Austin Hulse^{1,2,*} Hanqing Liu^{3,†} and Iman Marvian^{1,2,4,‡}

¹*Duke Quantum Center, Duke University, Durham, NC 27708, USA*

²*Department of Physics, Duke University, Durham, NC 27708, USA*

³*Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

⁴*Department of Electrical and Computer Engineering, Duke University, Durham, NC 27708, USA*

Quantum circuits with symmetry-respecting gates have attracted broad interest in quantum information science. While recent work has developed a theory for circuits with Abelian symmetries, revealing important distinctions between Abelian and non-Abelian cases, a comprehensive framework for non-Abelian symmetries has been lacking. In this work, we develop novel techniques and a powerful framework that is particularly useful for understanding circuits with non-Abelian symmetries. Using this framework we settle an open question on quantum circuits with $SU(d)$ symmetry. We show that 3-qudit $SU(d)$ -invariant gates are semi-universal, i.e., generate all $SU(d)$ -invariant unitaries, up to certain constraints on the relative phases between sectors with inequivalent representation of symmetry. Furthermore, we prove that these gates achieve full universality when supplemented with 3 ancilla qudits. Interestingly, we find that studying circuits with 3-qudit gates is also useful for a better understanding of circuits with 2-qudit gates. In particular, we establish that even though 2-qudit $SU(d)$ -invariant gates are not themselves semi-universal, they become universal with at most 11 ancilla qudits. Additionally, we investigate the statistical properties of circuits composed of random $SU(d)$ -invariant gates. Our findings reveal that while circuits with 2-qudit gates do not form a 2-design for the Haar measure over $SU(d)$ -invariant unitaries, circuits with 3-qudit gates generate a t -design, with t that is quadratic in the number of qudits.

I. INTRODUCTION

The universality of 2-qudit quantum gates is a celebrated result in the fields of quantum computing and control theory [1–4]. According to this result, any unitary transformation on any finite number of qudits can be realized with a finite sequence of two qudit gates. However, in the presence of a global symmetry, this universality fails: generic unitaries that respect a global continuous symmetry cannot be realized, even approximately, using k -qudit gates that respect the same symmetry, with any fixed k [5–8]. In general, the locality of gates imposes various types of restrictions on the set of realizable unitaries. For instance, it restricts the possible relative phases between sectors with inequivalent irreducible representations (charges) of the symmetry. To distinguish these more common types of restrictions from other types, Ref. [8] proposes the notion of semi-universality, a weakening of the notion of universality. This concept, which is the main focus of the present paper, is defined below (see Sec. II A for the formal definition).

Recall that under the action of a symmetry group G , the total Hilbert space \mathcal{H} decomposes into subspaces (charge sectors), $\mathcal{H} = \bigoplus_{\lambda} \mathcal{H}_{\lambda}$, corresponding to inequivalent irreducible representations of G . A set of gates respecting this symmetry is called *semi-universal*, if for any unitary V that respects the symmetry, there exists a set of phases $\{\theta_{\lambda}\}$ such that $V \sum_{\lambda} e^{i\theta_{\lambda}} \Pi_{\lambda}$ can be realized as a sequence of gates in that set, where Π_{λ} is the orthogonal projector

to the charge sector \mathcal{H}_{λ} .

One may hope that even though 2-qudit gates are not generally universal, at least they might be semi-universal. Previous works have shown that this is indeed the case for qubits with $U(1)$ and $SU(2)$ symmetry [7, 8]. In particular, in the case of $SU(2)$ symmetry, 2-qubit gates realized by the Heisenberg exchange interaction, which is $SU(2)$ -invariant, are semi-universal. On the other hand, surprisingly, it turns out that for $d \geq 3$, 2-qudit $SU(d)$ -invariant gates on qudit systems are not semi-universal. Indeed, Ref. [6] identifies new conservation laws that restrict the time evolution of the system,¹ even when the state is restricted to one $SU(d)$ charge sector. Since 2-qudit $SU(d)$ -invariant gates are not semi-universal in general, it is natural to ask whether k -local gates are; and, in particular, what is the minimum locality that achieves semi-universality.

A. Summary of results

In this paper, we develop new powerful tools and a framework for understanding semi-universality in quantum circuits with arbitrary symmetries. While recent work [8] has found a simple characterization of circuits with Abelian symmetries, it is known that circuits with non-Abelian symmetries can show significantly more complicated behaviors. For instance, in the presence of non-Abelian symmetries, the unitaries realized in one charge

* austin.hulse@duke.edu

† hanqing.liu@lanl.gov

‡ iman.marvian@duke.edu

¹ In particular, in [6], it is shown that in certain subspaces, the dynamics of qudits under 2-qudit $SU(d)$ -invariant unitaries can be mapped to the dynamics of a fermionic system evolving under a free (non-interacting) Hamiltonian.

Ancillae	Local dimension d	Semi-universality	Universality
\times	2	$k = 2$	$k = 2 \lfloor \frac{n}{2} \rfloor$
	$d \geq 3$	$k = 3$	$k = n$
\checkmark	2	$k = 2$ with no ancillae	$k = 2$ and 2 ancillae
	$d \geq 3$	$k = 2$ and ≤ 8 ancillae	$k = 2$ and ≤ 11 ancillae, or $k = 3$ and ≤ 3 ancillae

TABLE I. **Locality of gates needed to achieve (semi-)universality.** This table lists the minimum k , such that k -qudit $SU(d)$ -invariant unitaries achieve semi-universality and universality on $n \geq 3$ qudits, with or without ancillae. The results for $SU(2)$ symmetry were shown in [7], and the results on $SU(d)$ symmetry with $d \geq 3$ are established in this work. Note that for $d = 2$, 2 ancilla qudits are needed to achieve universality, even if one is allowed to use k -qubit gates, with $k < 2 \lfloor \frac{n}{2} \rfloor$.

sector \mathcal{H}_λ , may dictate the unitaries in other (possibly multiple) sectors, whereas this cannot happen in the case of Abelian symmetries [6, 8].

Applying these tools to the important example of $SU(d)$ symmetry, we settle an open question about circuits with this symmetry. It was recently shown that 2-qudit $SU(d)$ -invariant gates are not semi-universal when $d > 2$, while they are for $d = 2$ [6, 7]. Furthermore, using properties of the Young-Jucys-Murphy elements and Okounkov-Vershik's approach to the representation theory of the symmetric group [9], Ref. [10] argues that 4-local $SU(d)$ -invariant unitaries are semi-universal. However, prior to the present work, it was not known if semi-universality can be achieved with 3-qudit unitaries or not.

Here, we settle this open question and prove that 3-qudit unitaries that respect a global $SU(d)$ symmetry on d -dimensional qudits are indeed semi-universal. It is worth emphasizing that our proof of the semi-universality of 3-qudit gates is elementary and based on the tools developed in this paper, which are applicable to other symmetries (see Sec. II, and in particular, Lemmas 1 and 2).

In our construction, the generating gate set includes all 2-qudit $SU(d)$ -invariant gates

$$\exp(i\theta \mathbf{P}_{ij}) : \quad \theta \in [0, 2\pi) , \quad (1)$$

where \mathbf{P}_{ij} is the SWAP operator on qudits² i and j , and a single 3-qudit gate, e.g., one of the unitaries

$$R_+ = \exp(i\pi \Pi_{\square\square}) , \quad \text{or} \quad R_- = \exp(i\pi \Pi_{\square}) . \quad (2)$$

Here, $\Pi_{\square\square}$ and Π_{\square} are, respectively, the Hermitian projectors to the symmetric and anti-symmetric subspaces of the three-qudit Hilbert space $(\mathbb{C}^d)^{\otimes 3}$, which means $R_+ = \mathbb{I} - 2\Pi_{\square}$ and $R_- = \mathbb{I} - 2\Pi_{\square\square}$ are reflection unitaries, and are also permutationally invariant gates (see Fig. 1). Here we emphasize that R_\pm are not special: almost any single 3-qudit $SU(d)$ -invariant unitary together with 2-qudit unitaries in Eq. (1) are semi-universal. Proposition 1




FIG. 1. **Semi-universality with 3-qudit gates.** Any set of $SU(d)$ -invariant gates is called *semi-universal*, if they generate all $SU(d)$ -invariant unitaries, up to possible constraints on the relative phases between sectors with inequivalent irreps of $SU(d)$. While 2-qudit $SU(d)$ -invariant gates are not semi-universal for $d \geq 3$, we show that amending them with any single generic 3-qudit unitary makes them semi-universal. In this schematic circuit, 2-qudit gates are in the form of Eq. (1) for arbitrary θ , and R_+ is the 3-qudit reflection unitary defined in Eq. (2).

provides a simple characterization of 3-qudit gates that are capable of achieving semi-universality. In particular, we find that any 3-qudit gate that cannot be realized with 2-qudit gates is sufficient to make them universal.

It is worth noting that when restricted to a 3-qudit system, 2-qudit $SU(d)$ -invariant gates already achieve semi-universality, and the gate R_+ is just a relative phase between the $SU(d)$ charge sectors. However, interestingly, when acting on 3 qudits in a system with $n > 3$ qudits, such gates can drastically change the set of realizable unitaries and make them semi-universal.

Table I summarizes the results on (semi-)universality from the current paper and [6, 7]. As listed in this table, in addition to the semi-universality of 3-qudit gates, which is shown in Secs. IV and V, in this work, we also prove that

1. Without ancilla qudits, for $d > 2$, $(n - 1)$ -qudit $SU(d)$ -invariant gates are still not sufficient to achieve universality on n qudits (see Sec. VI).
2. With at most 8 ancilla qudits, semi-universality can be achieved with 2-qudit $SU(d)$ -invariant gates when $d > 2$ (see Lemma 6).

² We label the qudits as if they were in a chain for convenience: this geometry plays no role in the proof of semi-universality.

3. With at most 11 ancilla qudits, universality can be achieved with 2-qudit $SU(d)$ -invariant gates when $d > 2$ (see Corollary 3).

Therefore, a remarkable (and perhaps, unexpected) corollary of our study of 3-qudit $SU(d)$ -invariant gates is a significantly simpler understanding of the computational universality of 2-qudit gates using ancilla qudits! This has been recently established in [11] by applying an advanced result in the mathematical Lie-algebraic literature by Marin [12]. (This work characterizes the Lie algebra generated by transpositions as a subalgebra of the group algebra of permutations.)

The semi-universality of 3-qudit gates allows us to characterize the group generated by k -qudit $SU(d)$ -invariant unitaries on n qudits, denoted by $\mathcal{V}_k^{(n)}$ for $k \geq 3$ (Note that we have suppressed the d dependence to simplify the notation). Recall that according to the general results of [5], $\mathcal{V}_k^{(n)}$ is a compact connected Lie group. Then, as we discuss in Sec. VI, in the regime $n \geq k \gg d \geq 2$, the difference between the dimensions of this group and the subgroup generated by 3-qudit $SU(d)$ -invariant gates is approximately

$$\dim \mathcal{V}_k^{(n)} - \dim \mathcal{V}_3^{(n)} \approx \frac{k^{d-1}}{d!(d-1)!} + \mathcal{O}(k^{d-2}). \quad (3)$$

In Fig. 2 we plot

$$\rho_{k,d}^{(n)} := \frac{\dim \mathcal{V}_k^{(n)} - \dim \mathcal{V}_3^{(n)}}{\dim \mathcal{V}^{(n)} - \dim \mathcal{V}_3^{(n)}} \approx \left(\frac{k}{n}\right)^{d-1}, \quad (4)$$

for various d together with their asymptotic behavior, where $\mathcal{V}^{(n)} = \mathcal{V}_n^{(n)}$ denotes the group of all $SU(d)$ -invariant unitaries on n qudits. This ratio determines how the dimension of the Lie group of realizable unitaries grows with k . In particular, $\rho_{k,d}^{(n)} = 1$ means that universality is achieved. Interestingly, when $d \geq 3$ this happens only if $k = n$, i.e., gates act on all qudits in the system.³ Note that the right-hand side of Eq. (3) is independent of n the number of qudits (indeed, according to the general results of [5], this is a consequence of the fact that the symmetry group $G = SU(d)$ is connected).

In Sec. VIII we also discuss the implications of this result on the statistical properties of random quantum circuits with $SU(d)$ -invariant gates. As was noted in [6], the additional conservation laws, that restrict unitaries realized by 2-qudit circuits, imply that the distribution of unitaries generated by such random circuits is not a 2-design for the Haar distribution over the group of $SU(d)$ -invariant unitaries $\mathcal{V}^{(n)}$ (see Sec. VIII for the definition of t -designs). As we show in Sec. VIII, a corollary of the




FIG. 2. **Convergence to universality as a function of the locality of gates.** According to the general no-go theorem of [5], in symmetric quantum circuits with continuous symmetries, without ancilla qudits, universality cannot be achieved with k -qudit gates with a fixed k . On the other hand, we show that in the case of $SU(d)$ symmetry, semi-universality is achieved with 3-qudit gates, which means for $k \geq 3$, the only constraints on the realizable unitaries are on the relative phases between sectors with inequivalent irreducible representations of $SU(d)$. Roughly speaking, the ratio $\rho_{k,d}^{(n)}$ defined in Eq. (4), describes the fraction of these constraints that vanish with k -qudit gates with $k \geq 3$. Universality is achieved when $\rho_{k,d}^{(n)} = 1$. Here, we plot this ratio for a system with $n = 10^4$ qudits for different values of d , as specified in the plot. The dots are $\rho_{k,d}^{(n)}$ and the lines are its asymptotic expression $(k/n)^{d-1}$. The denominators of the ratio in Eq. (4) are 4998, 8.3×10^6 , 7.0×10^9 , 3.5×10^{12} and 7.9×10^{23} for $d = 2, 3, 4, 5, 10$, respectively.

semi-universality of 3-qudit gates is that, assuming the number of qudits is $n > d$, the distribution of unitaries generated by random circuits formed from such gates is a t -design up to $t \approx n^2/2$.

Finally, we note that while the main emphasis of this paper is on semi-universality in the context of symmetric quantum circuits, many of the ideas and techniques developed here are broadly useful in the context of quantum computing and control theory. In particular, Lemma 1, which provides the necessary and sufficient conditions for semi-universality, and Lemma 2, which is used for extending controllability from a subspace to the full space, are of independent interest.

B. Outline

In Sec. II we formally define semi-universality for an arbitrary unitary symmetry group, and present a number of generally applicable tools. In particular, we provide a necessary and sufficient condition for semi-universality to hold in Lemma 1, and we describe how using ancillae, semi-universality can be promoted to universality.

³ On the other hand, in the case of $d = 2$, i.e., qubits with $SU(2)$ symmetry, when n is odd, the universality can be achieved with gates acting on $n - 1$ qubits.

In Sec. III we specify to the case of $SU(d)$ symmetry on d -dimensional qudits and state Theorem 1, the semi-universality of 3-qudit $SU(d)$ -invariant unitaries. We also describe one of the main tools used in this paper, namely Schur-Weyl duality.

To explain the applications of the tools developed in Sec. II, in Sec. IV we present a detailed discussion of the examples of $n = 3$ and $n = 4$ qudits. In particular, we describe in detail how semi-universality fails for 2-qudit gates and prove that it holds for 3-qudit gates using Lemma 1. This serves as the base case of the induction argument for Theorem 1, which is proven in Sec. V.

In Sec. VI, we show that, when $d \geq 3$ and without using ancilla, universality on n qudits cannot be achieved without full nonlocal control, i.e. n -local gates are required. We consider the use of ancilla qudits for achieving (semi-)universality in Sec. VII. The statistical properties of circuits generated from $SU(d)$ -invariant 3-local gates are studied in Sec. VIII.

Finally, in Sec. IX, we prove Lemma 1. We also describe more general scenarios in which semi-universality does not hold, even when there is subsystem universality on all charge sectors. In particular, in Lemma 8 we present a characterization of G -invariant groups that are subsystem universal.

TABLE II. Table of notations.

Notation	Definition
\mathcal{V}^G	G -invariant unitaries
$\mathcal{S}\mathcal{V}^G$	commutator subgroup of \mathcal{V}^G (G -invariant unitaries w/o relative phases)
$\mathcal{V}_k^{(n)}$	group generated by k -qudit $SU(d)$ -invariant unitaries on n qudits
$\mathcal{V}^{(n)} = \mathcal{V}_n^{(n)}$	all $SU(d)$ -invariant unitaries on n qudits
$\mathcal{S}\mathcal{V}^{(n)}$	commutator subgroup of $\mathcal{V}^{(n)}$ ($SU(d)$ -invariant unitaries w/o relative phases)
\mathbb{S}_n	symmetric group on n objects
\mathcal{M}_λ	multiplicity subsystem (irrep space of \mathbb{S}_n in the case of $SU(d)$ symmetry)
$\Lambda_{n,d}$	irreps of \mathbb{S}_n on n qudits

II. SEMI-UNIVERSALITY

In this section, we define semi-universality rigorously for arbitrary symmetry groups, and introduce three powerful lemmas that are later used in Sec. V to prove the semi-universality of 3-qudit gates. We anticipate that these lemmas will find other applications beyond this result.

A. Definition

Let \mathcal{V}^G be the set of all G -invariant unitaries; that is, $V \in \mathcal{V}^G$ if and only if $[V, U(g)] = 0$ for all $g \in G$, where

$U(g)$ is the unitary representation of a finite or compact Lie group G . Then, \mathcal{V}^G itself is a compact, connected Lie group [5].

In the introduction we (informally) defined the notion of semi-universality. As is explained below, an equivalent and useful definition of semi-universality is the following: a set of G -invariant unitaries is called *semi-universal* for \mathcal{V}^G , if the subgroup \mathcal{T} generated by them contains the commutator subgroup of \mathcal{V}^G , denoted by $\mathcal{S}\mathcal{V}^G$. That is,

$$\mathcal{T} \supseteq \mathcal{S}\mathcal{V}^G := [\mathcal{V}^G, \mathcal{V}^G], \quad (5)$$

where for any group \mathcal{S} the commutator subgroup $[\mathcal{S}, \mathcal{S}]$ is generated by the group commutators $WVW^{-1}V^{-1} : W, V \in \mathcal{S}$. As it is more apparent from Eq. (10) below, from a Lie-algebraic perspective, this condition implies that the semi-simple parts of the Lie algebras associated with \mathcal{V}^G and \mathcal{T} are identical, whereas their center can be different (this motivates the name ‘‘semi-universality’’).

A useful characterization of (semi-)universality can be obtained by considering the isotypic decomposition of representation $\{U(g) : g \in G\}$, namely

$$\mathcal{H} = \bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda = \bigoplus_{\lambda \in \Lambda} \mathcal{Q}_\lambda \otimes \mathcal{M}_\lambda, \quad (6)$$

where the sum is over a set Λ of inequivalent irreducible representations (irreps) of G , \mathcal{Q}_λ is a space carrying the irrep λ , and \mathcal{M}_λ corresponds to the multiplicity of λ , i.e., $\dim \mathcal{M}_\lambda > 0$ is the multiplicity of λ in \mathcal{H} . Applying Schur’s lemma, we find that, with respect to the decomposition in Eq. (6), any symmetric unitary $V \in \mathcal{V}^G$ is block-diagonal and takes the form

$$V = \bigoplus_{\lambda \in \Lambda} V_\lambda = \bigoplus_{\lambda \in \Lambda} (\mathbb{I}_{\mathcal{Q}_\lambda} \otimes v_\lambda), \quad (7)$$

where $\mathbb{I}_{\mathcal{Q}_\lambda}$ is the identity operator on \mathcal{Q}_λ and $v_\lambda \in U(\mathcal{M}_\lambda)$ is a unitary acting only on the multiplicity degrees of freedom. Here, $U(\mathcal{M}_\lambda)$ is the group of unitaries on the Hilbert space \mathcal{M}_λ , and $SU(\mathcal{M}_\lambda)$ is its subgroup with determinant one.

It is useful to consider the homomorphisms $\pi_\lambda : \lambda \in \Lambda$ from \mathcal{V}^G to $U(\mathcal{M}_\lambda)$, defined by $\pi_\lambda(V) = v_\lambda$. Then, we denote the collection of these unitaries $v_\lambda : \lambda \in \Lambda$, as $\pi_\Lambda(V) = (v_\lambda)_{\lambda \in \Lambda}$. This defines an isomorphism

$$\pi_\Lambda : \mathcal{V}^G \rightarrow \prod_{\lambda \in \Lambda} U(\mathcal{M}_\lambda), \quad (8)$$

from the group of all G -invariant unitaries to the Cartesian product of unitary groups on the multiplicity spaces. This means the unitary V is uniquely determined by the set $\pi_\Lambda(V)$, via $V = \bigoplus_{\lambda \in \Lambda} \mathbb{I}_{\mathcal{Q}_\lambda} \otimes \pi_\lambda(V)$.

Recall that $SU(\mathcal{M}_\lambda)$ is a perfect group, i.e., it is equal to its commutator subgroup, which means $SU(\mathcal{M}_\lambda) = [U(\mathcal{M}_\lambda), U(\mathcal{M}_\lambda)]$. This, in turn, implies

$$\mathcal{S}\mathcal{V}^G \cong \pi_\Lambda([\mathcal{V}^G, \mathcal{V}^G]) = \prod_{\lambda \in \Lambda} SU(\mathcal{M}_\lambda). \quad (9)$$

In particular, this means for any G -invariant unitary $V \in \mathcal{V}^G$, there exists a set of phases $\theta_\lambda \in [0, 2\pi)$ and $\tilde{V} \in \mathcal{SV}^G$, such that $V = \tilde{V} \sum_\lambda e^{i\theta_\lambda} \Pi_\lambda$, demonstrating the equivalence of the above definition with the definition presented in the introduction.

Therefore, the definition in Eq. (5) means that group \mathcal{T} is semi-universal if, and only if, its commutator subgroup is equal to \mathcal{SV}^G , i.e.,

$$[\mathcal{T}, \mathcal{T}] = [\mathcal{V}^G, \mathcal{V}^G] = \mathcal{SV}^G. \quad (10)$$

If the group $\mathcal{T} \subseteq \mathcal{V}^G$ is connected and each $\pi_\lambda(\mathcal{T}) = \{\pi_\lambda(V) : V \in \mathcal{T}\}$ acts irreducibly on \mathcal{M}_λ , then it follows that the connected component of the identity of $\mathcal{T} \cap \mathcal{SV}^G$ is equal to $[\mathcal{T}, \mathcal{T}]$. See Appendix B 6 for a proof of this statement.

Finally, it is worth noting that a subgroup of G -invariant unitaries $\mathcal{T} \subseteq \mathcal{V}^G$ contains \mathcal{SV}^G , if and only if it contains the one-parameter family of unitaries $\exp(iHt) : t \in \mathbb{R}$ for all centerless G -invariant Hamiltonians, where we say G -invariant Hamiltonian H is *centerless* if $\text{Tr}(\Pi_\lambda H) = 0$ for all $\lambda \in \Lambda$, or equivalently, if $\text{Tr}(U(g)H) = 0$ for $g \in G$.

B. Lemma 1: A simple characterization of semi-universality

The following lemma is one of our main new tools for studying semi-universality and can potentially have broad applications beyond the context of quantum circuits.

Lemma 1. *A subgroup $\mathcal{T} \subseteq \mathcal{V}^G$ of G -invariant unitaries contains the commutator subgroup of all G -invariant unitaries $\mathcal{SV}^G = [\mathcal{V}^G, \mathcal{V}^G]$ if, and only if, the following two conditions hold:*

A (Subsystem universality in all sectors): *For any irrep $\lambda \in \Lambda$ the action of \mathcal{T} on the corresponding multiplicity subsystem \mathcal{M}_λ contains $\text{SU}(\mathcal{M}_\lambda)$, i.e., $\text{SU}(\mathcal{M}_\lambda) \subseteq \pi_\lambda(\mathcal{T}) = \{\pi_\lambda(V) : V \in \mathcal{T}\}$.*

B (Pairwise independence): *For any pair of distinct irreps $\lambda_1, \lambda_2 \in \Lambda$, if $\dim(\mathcal{M}_{\lambda_1}) = \dim(\mathcal{M}_{\lambda_2}) \geq 2$, then there exists a unitary $V \in \mathcal{T}$ such that*

$$|\text{Tr}(\pi_{\lambda_1}(V))| \neq |\text{Tr}(\pi_{\lambda_2}(V))|. \quad (11)$$

As we show in Sec. IX E, this lemma can be established using Goursat's and Serre's lemmas (see Lemmas 9 and 10, respectively). Additionally, in Lemma 8 we present a variant of Lemma 1, which does not assume condition B holds. In particular, we find the most general form of the subgroups of G -invariant unitaries that respect condition A, subsystem universality in all sectors.

In words, condition A means that inside the subspace associated with any irrep λ , all G -invariant unitaries are realizable up to a phase. When irrep λ of group G is itself 1D, this means all unitaries inside the subspace \mathcal{H}_λ are realizable, a condition that is sometimes called "subspace controllability" in control theory. However, in

general, λ is not a 1D irrep, and therefore following the standard terminology in quantum information, we refer to this condition as "subsystem universality". Condition B, on the other hand, guarantees that for any pair of irreps, the realized unitaries in the corresponding subspaces are independent of each other.

When condition A holds, condition B is equivalent to the following, which can therefore replace it:

B': For any pair of distinct irreps λ_1, λ_2 , if $\dim(\mathcal{M}_{\lambda_1}) = \dim(\mathcal{M}_{\lambda_2}) \geq 2$, then there exists a unitary $V \in \mathcal{T}$ that acts as the identity operator on one of \mathcal{M}_{λ_1} or \mathcal{M}_{λ_2} , and is not proportional to the identity on the other, e.g.,

$$\pi_{\lambda_1}(V) = \mathbb{I}_{\mathcal{M}_{\lambda_1}}, \quad \pi_{\lambda_2}(V) \neq e^{i\theta} \mathbb{I}_{\mathcal{M}_{\lambda_2}}, \quad (12)$$

for any $e^{i\theta}$, where $\mathbb{I}_{\mathcal{M}_{\lambda_1}}$ and $\mathbb{I}_{\mathcal{M}_{\lambda_2}}$ are the identity operators on \mathcal{M}_{λ_1} and \mathcal{M}_{λ_2} , respectively.

Note that if a unitary V satisfies the condition in Eq. (12), then it also satisfies the condition in Eq. (11). This can be seen by noting that $|\text{Tr}(\pi_\lambda(V))| = \text{Tr}(\pi_\lambda(\mathbb{I}_{\mathcal{M}_\lambda}))$ if and only if all eigenvalues of $\pi_\lambda(V)$ have the same phase, which means in the complex plane they are aligned in the same direction.

If both conditions A and B (or, equivalently, conditions A and B') are satisfied for a pair of irreps λ_1 and λ_2 , then the joint projection of \mathcal{T} to \mathcal{M}_{λ_1} and \mathcal{M}_{λ_2} contains $\text{SU}(\mathcal{M}_{\lambda_1}) \times \text{SU}(\mathcal{M}_{\lambda_2})$. That is,

$$\pi_{\lambda_1, \lambda_2}(\mathcal{T}) = \{(v_{\lambda_1}, v_{\lambda_2}) : V \in \mathcal{T}\} \supseteq \text{SU}(\mathcal{M}_{\lambda_1}) \times \text{SU}(\mathcal{M}_{\lambda_2}). \quad (13)$$

This, in particular, means that the unitary realized in one sector is not dictated by the other, up to possible constraints on the global phases. In Sec. IX E we argue that together with Serre's Lemma 10, Eq. (13) implies independence in all sectors, as claimed in Lemma 1.

Pairwise independence implies full independence

While the necessity of both conditions A and B is trivial, their sufficiency is far from obvious. Indeed, it is remarkable that according to this lemma, to demonstrate semi-universality one needs to check the independence of the realized unitaries only among pairs of sectors and not, e.g., among 3-tuples of sectors. To see an example of such dependencies, consider the subgroup of $\text{U}(1)^3$ corresponding to a 3-tuple of phases

$$(e^{i\theta_1}, e^{i\theta_2}, e^{i(\theta_1+\theta_2)}) \quad : \theta_1, \theta_2 \in [0, 2\pi). \quad (14)$$

Then, any pair of these 3 phases are fully independent of each other. That is, for any 2-tuples the projection of this group is $\text{U}(1)^2$, and yet the overall group is isomorphic to $\text{U}(1)^2$, rather than $\text{U}(1)^3$. However, Lemma 1 implies that this situation cannot happen in the context of semi-universality. As we further explain in Sec. IX, this is a consequence of the fact that the group $\text{SU}(m)$ is perfect, while $\text{U}(1)$ in the above example is not.

Failure of semi-universality

As mentioned before, when semi-universality holds, the only constraints on realizable unitaries are constraints on the relative phases between sectors with different charges (irreps) of symmetry. On the other hand, failure of semi-universality can be due to various kinds of constraints. It is useful to recall the numbering system of [8] for all possible constraints on the universality of \mathcal{T} . Namely,

Type I: Constraints on the relative phases between charge sectors. Note that this is a failure of universality rather than semi-universality.

Types II and III: Charge sectors which are not subsystem universal, either because **(II)** the action of \mathcal{T} on this sector is not irreducible, or **(III)** the action is irreducible, but only a proper subset of determinant-one unitaries can be achieved.

Type IV: “Correlations” between the determinant-one unitaries in distinct charge sectors. That is, the unitary realized in one sector determines the unitary realized in the other sector (up to a possible global phase). Such correlations can arise independent of type **II** and **III** constraints.

Lemma 8 characterizes type **IV** constraints assuming type **II** and **III** do not exist. It is worth noting that constraints of type **I** on a smaller system can result in constraints of other types in a system with more qudits. Indeed, this is exactly what happens in the case of 2-qudit $SU(d)$ -invariant gates. As we show in Sec. IV A, when restricted to $n = 3$ qudits, the 2-qudit $SU(d)$ -invariant gates are semi-universal, and therefore the only constraints on the realizable unitaries are type **I** constraints. However, as soon as we go to $n = 4$ and more qudits, these constraints on the relative phases will also cause type **III** and **IV** constraints. See [6] for further discussions on restrictions on circuits with 2-qudit $SU(d)$ -invariant gates.

It is also worth noting that according to the result of [8], when the symmetry group G is Abelian and has on-site representation (see Sec. III A), semi-universality holds if and only if there are no type **II** restrictions. That is, types **III** and **IV** restrictions do not exist for Abelian symmetries.

C. Lemma 2: Extending controllability from a 3D subspace to the full space

In Sec. V we show how Lemma 1 can be applied recursively via an induction argument to establish the semi-universality of 3-qudit $SU(d)$ -invariant gates. To apply such recursion, we need one more tool, which is discussed in this subsection.

In Appendix F we present a series of results, which under different assumptions allow us to extend controllability from a subspace to the full space (see also [6] for

previous examples of such results). Using these results, in Appendix B 1 we prove the following lemma, which is one of the tools needed to establish the semi-universality of $SU(d)$ -invariant 3-qudit gates.

Lemma 2. *Let \mathcal{H} be a finite-dimensional Hilbert space with a subspace $\mathcal{H}_1 \subset \mathcal{H}$ with dimension $\dim(\mathcal{H}_1) \geq 3$. Let A_i , $i = 1, \dots, k$, be traceless anti-Hermitian operators on \mathcal{H} and consider the one-parameter groups $\mathcal{A}_i = \{e^{tA_i} : t \in \mathbb{R}\}$. If the group*

$$\mathcal{W} = \langle \mathcal{A}_i, SU(\mathcal{H}_1) : i = 1, \dots, k \rangle \quad (15)$$

acts irreducibly on \mathcal{H} , then $\mathcal{W} = SU(\mathcal{H})$.

Remark 1. Indeed, a similar result holds when $\dim(\mathcal{H}_1) = 2$ and $\dim(\mathcal{H})$ is odd. We present the proof of this for the special case of $\dim(\mathcal{H}) = 3$ in Appendix B 2, and postpone the proof of the general case to [13].

Furthermore, in [13], we also show that a variant of this theorem applies to the case when $\dim(\mathcal{H}_1) = 2$ and $\dim \mathcal{H}$ is even. However, in that case, the realized group \mathcal{W} can be either the full $SU(\mathcal{H})$ or a subgroup isomorphic to the symplectic group $Sp(\mathcal{H})$ (see [13] for further discussions).

D. Lemma 3: From semi-universality to universality with 3 ancillae

So far, we have not made any assumptions about the structure of the Hilbert space \mathcal{H} and the unitary representation of symmetry G on this space. For many applications in physics and quantum computing, we are interested in the scenarios where \mathcal{H} is the Hilbert space of n identical qudits, i.e., $\mathcal{H} = (\mathbb{C}^d)^{\otimes n}$. We are also often interested in the case where $U(g)$ is an “on-site” representation of the symmetry, such that $U(g) = u(g)^{\otimes n}$, where $u(g) : g \in G$ acts on a single qudit \mathbb{C}^d (this is often called “global rotations” on the system). However, it is worth noting that the above lemmas apply to other cases, e.g., when G is the permutation group \mathbb{S}_n .

Then, under the assumption that the representation of symmetry is “on-site”, it turns out that one can use ancilla qudits to elevate semi-universality to universality, as defined in the following. Consider a subset of G -invariant realizable unitaries \mathcal{T} on $n + c$ qudits and a fixed state $|\eta\rangle \in (\mathbb{C}^d)^{\otimes c}$ of c ancilla qudits. Then, we say unitary $V \in U((\mathbb{C}^d)^{\otimes n})$ on n qudits is realizable with c ancilla qudits, if there exists a unitary $\tilde{V} \in \mathcal{T}$ such that

$$\tilde{V}(|\psi\rangle \otimes |\eta\rangle) = (V|\psi\rangle) \otimes |\eta\rangle \quad (16)$$

for all $|\psi\rangle \in (\mathbb{C}^d)^{\otimes n}$.

Recent work in [8] shows that in the case of Abelian groups, such as $U(1)$, universality can be achieved with a single ancilla qudit. Furthermore, in [7] we showed that in the case of qubits with $SU(2)$ symmetry, universality can be achieved using 2 ancilla qubits, whereas 1 ancilla qubit is not sufficient in that case. Based on these previous

observations, one may expect that the number of required ancillae may grow with the size of the group (e.g., with d in the case of $SU(d)$ symmetry). However, as we show in Sec. VII, this is not the case: 3 ancilla qudits are sufficient to achieve universality, provided that semi-universality holds.

Lemma 3. *Let G be an arbitrary symmetry group with on-site representation on a system with $n + 3$ qudits (i.e., it acts via $g \mapsto u(g)^{\otimes(n+3)}$). Suppose a group \mathcal{T} is semi-universal, i.e., it is a subgroup of G -invariant unitaries that contains $\mathcal{S}\mathcal{V}^G$. Then, using 3 ancilla qudits, we can realize any G -invariant unitary on n qudits using unitaries in \mathcal{T} , as defined in Eq. (16).*

For instance, as we further explain in Sec. VII A, the state $|\eta\rangle$ of ancilla can be chosen to be either of

$$\begin{aligned} |\eta_1\rangle &= \frac{1}{2\sqrt{3}}(2|100\rangle + (\sqrt{3} - 1)|010\rangle - (1 + \sqrt{3})|001\rangle), \\ |\eta_2\rangle &= \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle) \otimes |0\rangle, \end{aligned} \quad (17)$$

where $|0\rangle$ and $|1\rangle$ are orthonormal states.

III. SEMI-UNIVERSALITY FOR $SU(d)$ SYMMETRY

Next, we apply the tools discussed in the previous section to the important case $G = SU(d)$, with the on-site representation of this symmetry on qudits.

A. Global $SU(d)$ symmetry and $SU(d)$ -invariant unitaries

Consider a quantum system composed of n qudits \mathbb{C}^d , with local dimension $d \geq 2$. There is a natural representation of $SU(d)$ on $(\mathbb{C}^d)^{\otimes n}$, where the single-qudit unitary $u \in SU(d)$ acts simultaneously on each qudit, $u \mapsto u^{\otimes n}$, corresponding to a global ‘‘rotation’’ of the system. In this case, the isotypic decomposition in Eq. (6) takes the form

$$(\mathbb{C}^d)^{\otimes n} \cong \bigoplus_{\lambda \in \Lambda_{n,d}} \mathcal{H}_\lambda = \bigoplus_{\lambda \in \Lambda_{n,d}} \mathcal{Q}_\lambda \otimes \mathcal{M}_\lambda, \quad (18)$$

where $\Lambda_{n,d}$ labels the inequivalent irreps of $SU(d)$ which show up in $(\mathbb{C}^d)^{\otimes n}$.

The elements of the set $\Lambda_{n,d}$ can be labeled by Young diagrams. Namely, elements of $\Lambda_{n,d}$ are in one-to-one correspondence with Young diagrams with n boxes and $\leq d$ rows, satisfying the property that the number of boxes in each row is non-increasing. However, our proof of semi-universality of symmetric 3-qudit gates does not require the manipulation of Young diagrams as long as one accepts certain basic facts about the representation theory of the symmetric group, which are reviewed in

Sec. V A. Any reader unfamiliar with Young diagrams may consider them an elaborate labeling scheme, and nothing more, for the purposes of this proof (we provide some details for the interested reader in Fig. 3).

Let $\mathcal{V}_k^{(n)} \subset U((\mathbb{C}^d)^{\otimes n})$ be the group generated by k -local $SU(d)$ -invariant unitaries, i.e., $SU(d)$ -invariant unitaries that can be written as $V' \otimes \mathbb{I}^{\otimes(n-k)}$ up to a permutation of qudits, where V' acts on k qudits. For each k , this is a compact, connected Lie group [5]. With the notation of Sec. II A, we have the equality

$$\mathcal{V}_n^{(n)} = \mathcal{V}^{SU(d)} \cong \prod_{\lambda \in \Lambda_{n,d}} U(\mathcal{M}_\lambda). \quad (19)$$

Then, the commutator subgroup of $\mathcal{V}_n^{(n)}$, denoted as $\mathcal{S}\mathcal{V}_n^{(n)}$, is isomorphic to $\prod_{\lambda \in \Lambda_{n,d}} SU(\mathcal{M}_\lambda)$. We will sometimes use the notation $\mathcal{V}^{(n)} = \mathcal{V}_n^{(n)}$, and $\mathcal{S}\mathcal{V}^{(n)} = \mathcal{S}\mathcal{V}_n^{(n)}$. Similarly, for k -qudit gates we define

$$\mathcal{S}\mathcal{V}_k^{(n)} = [\mathcal{V}_k^{(n)}, \mathcal{V}_k^{(n)}]. \quad (20)$$

Using these definitions, we can now present the formal statement of our result on the semi-universality of 3-qudit gates.

Theorem 1. *3-qudit $SU(d)$ -invariant gates are semi-universal on arbitrary n qudits. That is,*

$$[\mathcal{V}_3^{(n)}, \mathcal{V}_3^{(n)}] = \mathcal{S}\mathcal{V}^{(n)} \cong \prod_{\lambda \in \Lambda_{n,d}} SU(\mathcal{M}_\lambda). \quad (21)$$

Specifically, 2-qudit $SU(d)$ -invariant gates together with any single 3-qudit gate that is not realizable with 2-qudit gates, such as gates R_+ or R_- , defined in Eq. (2), are semi-universal.

See Proposition 1 for a characterization of the 3-qudit gates that cannot be realized with 2-qudit $SU(d)$ -invariant unitaries.

In Sec. IV A, we show that in the special case of $d = 2$, i.e., qubits with $SU(2)$ symmetry, all 3-qudit $SU(2)$ -invariant unitaries can be realized with 2-qudit $SU(2)$ -invariant unitaries (which is not true for $d \geq 3$). Therefore, we have the following immediate corollary of this result, which was also previously established in [6].

Corollary 1. *For qubit systems with $SU(2)$ symmetry, 2-qubit $SU(2)$ -invariant unitaries are semi-universal.*

Before presenting the proof of this theorem, we briefly review a powerful tool for understanding the properties of $SU(d)$ -invariant unitaries, which plays a crucial role in our arguments: Schur-Weyl duality.

B. Symmetric group \mathbb{S}_n and Schur-Weyl duality

An important class of $SU(d)$ -invariant unitaries are permutations. The symmetric group on n objects \mathbb{S}_n has

a natural representation on $(\mathbb{C}^d)^{\otimes n}$. In particular, for any permutation $\sigma \in \mathbb{S}_n$, let $\mathbf{P}(\sigma)$ denote the unitary operator on $(\mathbb{C}^d)^{\otimes n}$ which permutes the qudits according to σ . Occasionally, we will also use the notation \mathbf{P}_σ .

Since permutations are $SU(d)$ -invariant, they are block-diagonal with respect to the decomposition in Eq. (18). Furthermore, according to the Schur-Weyl duality, $\Lambda_{n,d}$ also labels inequivalent irreps of \mathbb{S}_n , i.e., for all $\sigma \in \mathbb{S}_n$

$$\mathbf{P}(\sigma) = \bigoplus_{\lambda \in \Lambda_{n,d}} \mathbb{I}_{\mathcal{Q}_\lambda} \otimes \mathbf{p}_\lambda(\sigma), \quad (22)$$

where $\mathbb{I}_{\mathcal{Q}_\lambda}$ is the identity operator on \mathcal{Q}_λ , and $\mathbf{p}_\lambda(\sigma) \in U(\mathcal{M}_\lambda)$, defines an irrep of \mathbb{S}_n on \mathcal{M}_λ (see, e.g., [14, 15]). This group of unitaries is generated by the transposition (swap) of qubits i and j , denoted by $\mathbf{P}(ij)$, or \mathbf{P}_{ij} for simplicity. It follows that all permutations $\mathbf{P}(\sigma)$ belong to $\mathcal{V}_2^{(n)}$. We conclude that for $k \geq 2$, the group $\mathcal{V}_k^{(n)}$, as well as its commutator subgroup $\mathcal{SV}_k^{(n)}$, acts irreducibly on \mathcal{M}_λ for all $\lambda \in \Lambda_{n,d}$.

Schur-Weyl duality implies that on any pair of distinct qudits i and j , any 2-qudit $SU(d)$ -invariant unitary can be written as a global phase $e^{i\phi} \mathbb{I}$ times

$$\exp(i\theta \mathbf{P}_{ij}) = \cos \theta \mathbb{I} + i \sin \theta \mathbf{P}_{ij}, \quad (23)$$

for $\phi, \theta \in [0, 2\pi)$, where \mathbb{I} denotes the identity operator on \mathcal{H} . Therefore, $\mathcal{V}_2^{(n)}$ is the group generated by

$$\mathcal{V}_2^{(n)} = \langle \exp(i\theta \mathbf{P}_{ij}), \exp(i\phi) \mathbb{I} : i \neq j, \theta, \phi \in [0, 2\pi) \rangle. \quad (24)$$

See [6] for further discussions on properties of group $\mathcal{V}_2^{(n)}$.

IV. ILLUSTRATIVE EXAMPLE: $n = 4$ QUDITS

As an illustrative example, in this section, we discuss the case of $n = 4$ qudits in depth and show that, 3-qudit $SU(d)$ -invariant unitaries are semi-universal in this case. This also provides the base case of an inductive proof that 3-qudit $SU(d)$ -invariant unitaries are semi-universal on arbitrary $n > 4$ qudits, discussed in Sec. V. Also, it demonstrates all the main techniques that are applied for the general proof. It is helpful to first consider 2-qudit gates—in doing so, the failure of 2-qudit $SU(d)$ -invariant gates to be semi-universal when $d > 2$ is illustrated.

It is worth noting that while 3-qudit $SU(d)$ -invariant gates are semi-universal on $n = 4$ qudits, they are not universal. In particular, applying the results of [5], reviewed in Eq. (59), we find that the difference between the dimensions of the Lie group of all $SU(d)$ -invariant unitaries on $n = 4$ qudits, and the subgroup generated by 3-qudit $SU(d)$ -invariant gates is equal to

$$\dim \mathcal{V}^{(4)} - \dim \mathcal{V}_3^{(4)} = |\Lambda_{4,d}| - |\Lambda_{3,d}| = \begin{cases} 1 & : d = 3 \\ 2 & : d \geq 4 \end{cases} \quad (25)$$

where $|\Lambda_{n,d}|$ is the number of inequivalent irreps of $SU(d)$, (or equivalently, \mathbb{S}_n) on n qudits.

A. 2-qudit gates on $n = 3$ qudits

First, we consider $n = 3$ qudits. In this case, for general $d \geq 3$, the decomposition in Eq. (18) contains 3 inequivalent irreps of \mathbb{S}_3 , which are labeled by the Young diagrams

$$\Lambda_3 = \left\{ \begin{array}{c} \square\square\square \\ \square\square \\ \square \end{array} \right\}, \quad (26)$$

where $\square\square\square$ and $\begin{array}{c} \square \\ \square \end{array}$ correspond to 1D irreps of \mathbb{S}_3 , namely the trivial representation and the sign representation⁴ $\mathbf{p}_{\begin{array}{c} \square \\ \square \end{array}}(\sigma) = \text{sgn}(\sigma) : \sigma \in \mathbb{S}_3$ (Note that the irrep $\begin{array}{c} \square \\ \square \end{array}$ appears only for qudits with $d > 2$). Diagram $\begin{array}{c} \square\square \\ \square \end{array}$ corresponds to the 2D irrep of \mathbb{S}_3 . In particular, in Young's orthonormal basis for $\mathcal{M}_{\begin{array}{c} \square\square \\ \square \end{array}}$, we obtain

$$[\mathbf{P}_{\begin{array}{c} \square\square \\ \square \end{array}}(12)] = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad [\mathbf{P}_{\begin{array}{c} \square\square \\ \square \end{array}}(23)] = \frac{1}{2} \begin{pmatrix} -1 & \sqrt{3} \\ \sqrt{3} & 1 \end{pmatrix}, \quad (27)$$

where we use the notation $[A]$ to denote the matrix representation of an operator A . Therefore, when projected to $\begin{array}{c} \square\square \\ \square \end{array}$, unitaries $\exp(i\theta \mathbf{P}(12))$ and $\exp(i\theta \mathbf{P}(23))$ correspond to $SU(2)$ rotations in two non-parallel directions, which means together they generate the full $SU(2)$ group inside $\mathcal{M}_{\begin{array}{c} \square\square \\ \square \end{array}}$. In general, the projection of such unitaries inside $\mathcal{M}_{\begin{array}{c} \square\square\square \\ \square \end{array}}$ and $\mathcal{M}_{\begin{array}{c} \square \\ \square \end{array}}$ are non-trivial phases. However, such phases, which are allowed in $\mathcal{V}^{(3)}$ but not $\mathcal{SV}^{(3)}$, can be canceled by noting that $SU(2)$ is equal to its commutator subgroup (In particular, any $U \in SU(2)$ can be decomposed as $U = U_2^\dagger U_1^\dagger U_2 U_1$ for some $U_1, U_2 \in SU(2)$ [16]).

We conclude that 2-qudit $SU(d)$ -invariant gates are semi-universal on $n = 3$ qudits. That is

$$\mathcal{V}_2^{(3)} \supseteq \mathcal{SV}^{(3)} \cong SU(2). \quad (28)$$

However, whether they are universal or not depends on d . In particular, the irrep $\begin{array}{c} \square \\ \square \end{array}$, whose charge sector $\mathcal{H}_{\begin{array}{c} \square \\ \square \end{array}}$ is the subspace of states of three qudits which are totally antisymmetric, shows up in the decomposition in Eq. (18) only if $d \geq 3$. In this case, there is a one-parameter family of relative phases between charge sectors that cannot be generated by 2-local $SU(d)$ -invariant gates. In particular, as we show in Appendix A 1,

Proposition 1 (Characterization of $\mathcal{V}_2^{(3)}$). *For a system with $n = 3$ qudits, the family of unitary evolutions $\exp(-iHt) : t \in \mathbb{R}$ is realizable with 2-qudit $SU(d)$ -invariant unitaries, i.e., $\exp(-iHt) \in \mathcal{V}_2^{(3)}$ if, and only if $\text{Tr}(HC) = 0$, where*

$$\begin{aligned} C &= 2(d-1)(d-2)\Pi_{\begin{array}{c} \square\square\square \\ \square \end{array}} - (d+2)(d-2)\Pi_{\begin{array}{c} \square\square \\ \square \end{array}} \\ &\quad + 2(d+2)(d+1)\Pi_{\begin{array}{c} \square \\ \square \end{array}} \\ &= d^2(\mathbf{P}_{(123)} + \mathbf{P}_{(132)}) - 2d(\mathbf{P}_{12} + \mathbf{P}_{13} + \mathbf{P}_{23}) + 4\mathbb{I}. \end{aligned} \quad (29)$$

⁴ Recall that $\text{sgn}(\sigma) = \pm 1$ depending on whether the number of transpositions needed to realize σ is odd or even.

Furthermore, when $d \geq 3$, the unitary $V \in \mathcal{V}^{(3)}$ is realizable by 2-qudit $SU(d)$ -invariant unitaries, i.e. $V \in \mathcal{V}_2^{(3)}$, if and only if

$$\det v_{\boxplus} = (\det v_{\boxminus})(\det v_{\boxtimes}). \quad (30)$$

Remark 2. In the special case of $d = 2$, operator $C = 0$, and therefore the condition holds trivially, which means 2-qudit $SU(2)$ -invariant unitaries are indeed *universal* on $n = 3$ qubits, i.e., $\mathcal{V}_2^{(3)} = \mathcal{V}^{(3)}$. Furthermore, this implies that in this case, $\mathcal{V}_2^{(n)} = \mathcal{V}_3^{(n)}$ for all $n \geq 3$.

The constraint imposed by Eq. (30) implies that when $d \geq 3$, the difference between the dimensions of the Lie group of all $SU(d)$ -invariant unitaries on 3 qudits, and its subgroup generated by 2-qudits gates is

$$\dim \mathcal{V}^{(3)} - \dim \mathcal{V}_2^{(3)} = 1, \quad (31)$$

saturating a general lower bound previously established in [5]. Therefore, a generic 3-qudit $SU(d)$ -invariant unitary cannot be generated by 2-qudit ones, since $\mathcal{V}_2^{(3)}$ forms a set of measure zero.

B. 2-qudit gates on $n = 4$ qudits

Next, we consider $n = 4$ qudits. The group \mathbb{S}_4 has 5 inequivalent irreps, which are labeled by the Young diagrams

$$\Lambda_4 = \left\{ \begin{array}{c} \square\square\square\square \\ \square\square\square \\ \square\square \\ \square \\ \square \end{array} \right\}. \quad (32)$$

For $d \geq 4$, all 5 diagrams appear in the decomposition in Eq. (18), whereas for $d = 3$ the last diagram does not appear. Again, the diagrams $\square\square\square\square$ and \square correspond to the 1D irreps, namely the trivial and sign representations of \mathbb{S}_4 , respectively.

To determine which unitary transformations can be realized with 2-qudit $SU(d)$ -invariant gates, we consider the unitaries in the form $V \otimes \mathbb{I} : V \in \mathcal{S}\mathcal{V}^{(3)}$, which according to Eq. (28) form a group isomorphic to $SU(2)$, and they can all be realized by 2-qudit $SU(d)$ -invariant gates that act trivially on qudit 4. Fortunately, to understand how these unitaries act on the multiplicity spaces \mathcal{M}_{\boxplus} , \mathcal{M}_{\boxminus} , and \mathcal{M}_{\boxtimes} , it suffices to understand the action of the \mathbb{S}_3 subgroup of $\mathcal{V}_2^{(3)} \otimes \mathbb{I}$ corresponding to permutations of the first 3 qudits. This follows from the fact that

$$\{\mathbf{P}(\sigma) : \sigma \in \mathbb{S}_3\} \subset \mathcal{V}_2^{(3)} \subset \text{span}_{\mathbb{C}}\{\mathbf{P}(\sigma) : \sigma \in \mathbb{S}_3\}. \quad (33)$$

Using the standard facts about the representation theory of \mathbb{S}_4 (see Sec. V A), it can be easily shown that a copy of the 2D irrep \boxplus of \mathbb{S}_3 appear inside each of the multiplicity spaces \mathcal{M}_{\boxplus} , \mathcal{M}_{\boxminus} , and \mathcal{M}_{\boxtimes} . In particular, under the

$[\mathbf{p}_{\boxplus}(\sigma)] = \left(\begin{array}{c c} [\mathbf{p}_{\boxplus}(\sigma)] & \\ \hline & \mathbf{p}_{\boxplus}(\sigma) \end{array} \right) = \left(\begin{array}{c c} [\mathbf{p}_{\boxplus}(\sigma)] & \\ \hline & 1 \end{array} \right)$
$[\mathbf{p}_{\boxminus}(\sigma)] = \left(\begin{array}{c c} [\mathbf{p}_{\boxminus}(\sigma)] & \\ \hline & \mathbf{p}_{\boxminus}(\sigma) \end{array} \right) = \left(\begin{array}{c c} [\mathbf{p}_{\boxminus}(\sigma)] & \\ \hline & \text{sgn}(\sigma) \end{array} \right)$
$[J] = \left(\begin{array}{c c} 1 & \\ \hline -1 & \\ \hline & 1 \end{array} \right) \quad [\mathbf{p}_{\boxtimes}(34)] = \left(\begin{array}{c c} \frac{1}{3} & \frac{2\sqrt{2}}{3} \\ \hline & 1 \\ \hline \frac{2\sqrt{2}}{3} & -\frac{1}{3} \end{array} \right)$

TABLE III. In the first two rows σ is in \mathbb{S}_3 subgroup of \mathbb{S}_4 that acts trivially on qudit 4, and $[\mathbf{p}_{\boxplus}(\sigma)]$ is given by Eq. (27). Matrix $[J]$ satisfies Eq. (39).

restriction to \mathbb{S}_3 subgroup that acts on the first three qudits, the irreps of \mathbb{S}_4 decompose as

$$\begin{aligned} \mathcal{M}_{\boxplus} &\cong \mathcal{M}_{\boxplus} \cong \mathbb{C}^2 \\ \mathcal{M}_{\boxminus} &\cong \mathcal{M}_{\boxplus} \oplus \mathcal{M}_{\boxtimes} \cong \mathbb{C}^2 \oplus \mathbb{C} \\ \mathcal{M}_{\boxtimes} &\cong \mathcal{M}_{\boxplus} \oplus \mathcal{M}_{\boxtimes} \cong \mathbb{C}^2 \oplus \mathbb{C}. \end{aligned} \quad (34)$$

The first two rows of Table III give the explicit matrix representations of this \mathbb{S}_3 subgroup in irreps \boxplus and \boxtimes .

Eq. (34) immediately implies that the projection of $\mathcal{S}\mathcal{V}_2^{(3)} \otimes \mathbb{I}$ to \mathcal{M}_{\boxplus} is isomorphic to $SU(2)$, i.e.,

$$\pi_{\boxplus}(\mathcal{S}\mathcal{V}_2^{(3)} \otimes \mathbb{I}) = SU(\mathcal{M}_{\boxplus}) \cong SU(2). \quad (35)$$

In other words, condition **A** in Lemma 1, namely, subsystem universality, is satisfied in the sector with irrep \boxplus .

Next, we consider the projection of this group to \mathcal{M}_{\boxminus} and \mathcal{M}_{\boxtimes} . Relative to the decomposition in Eq. (34), this group of unitaries will be in the form

$$\left(\begin{array}{c|c} SU(2) & \\ \hline & 1 \end{array} \right). \quad (36)$$

Note that this block-diagonal form is a consequence of the fact that under the operators that act on the first 3 qudits, the $SU(d)$ charge of qudit 4 is conserved. Now suppose we include 2-qudit $SU(d)$ -invariant unitaries $\exp(i\theta\mathbf{P}(34)) : \theta \in [0, 2\pi)$ that act on qudits 3 and 4, which allows qudit 4 to interact with the rest of the system. Since the transposition $\mathbf{P}(34)$ together with the aforementioned \mathbb{S}_3 subgroup generate \mathbb{S}_4 and \mathcal{M}_{\boxplus} and \mathcal{M}_{\boxtimes} are both irreps of \mathbb{S}_4 , the projection of these unitaries inside \mathcal{M}_{\boxplus} and \mathcal{M}_{\boxtimes} cannot be block-diagonal with respect to the decompositions in Eq. (34).

In summary, inside both subspaces \mathcal{M}_{\boxplus} and \mathcal{M}_{\boxtimes} 2-qudit unitaries acting on qudits 1, 2 and 3 generate the block-diagonal unitaries in the form Eq. (36), and

$\exp(i\theta\mathbf{P}(34)) : \theta \in [0, 2\pi)$ is a one-parameter family of unitaries that are not block diagonal with respect to this decomposition. According to Remark 1, adding any such one-parameter family of unitaries to unitaries in Eq. (36), generates a group that contains $SU(3)$ (see also Lemma 5 of [6]).

Therefore, similar to irrep \boxplus , condition **A** in Lemma 1 is also satisfied in irreps $\boxplus\boxplus$ and $\boxplus\boxminus$. Since $\mathcal{M}_{\boxplus\boxplus}$ and $\mathcal{M}_{\boxplus\boxminus}$ are 1D, this condition is trivially satisfied in those irreps. We conclude that condition **A** is fully satisfied, i.e., for all $\lambda \in \Lambda_4$, the projection of $\mathcal{V}_2^{(4)}$ to \mathcal{M}_λ is equal to the projection of $\mathcal{V}^{(4)}$ to \mathcal{M}_λ :

$$\forall \lambda \in \Lambda_4 : \quad \pi_\lambda(\mathcal{SV}_2^{(4)}) = \pi_\lambda(\mathcal{SV}^{(4)}) = SU(\mathcal{M}_\lambda) . \quad (37)$$

However, simply realizing all possible unitaries in the multiplicity spaces does not preclude the possibility of correlation between sectors. That is, condition **B** may not be satisfied, and in fact, it turns out that this is the case! The unitaries realized in $\mathcal{M}_{\boxplus\boxplus}$ and $\mathcal{M}_{\boxplus\boxminus}$ uniquely determine each other, up to a global phase. This is a consequence of a standard representation isomorphism between pairs of irreps of S_4 that are related by the sign representation, namely

$$\boxplus\boxplus \cong \text{sgn} \otimes \boxplus\boxminus , \quad \boxplus\boxminus \cong \text{sgn} \otimes \boxplus\boxplus . \quad (38)$$

That is, there exists a unitary operator $J : \mathcal{M}_{\boxplus\boxplus} \rightarrow \mathcal{M}_{\boxplus\boxminus}$ such that

$$\sigma \in S_4 : \quad J\mathbf{p}_{\boxplus\boxplus}(\sigma)J^\dagger = \text{sgn}(\sigma)\mathbf{p}_{\boxplus\boxminus}(\sigma) . \quad (39)$$

As presented in Table III, we can pick Young's basis where permutations have real matrix representation, denoted as $[\mathbf{p}_{\boxplus\boxplus}(\sigma)]$ and $[\mathbf{p}_{\boxplus\boxminus}(\sigma)]$. Then, relative to this basis, J also becomes a real orthogonal matrix $[J]$. Since for any transposition (SWAP) (ij) the parity $\text{sgn}((ij)) = -1$, this in turn implies

$$[J][\exp(i\theta\mathbf{p}_{\boxplus\boxplus}(ij))][J]^T = [\exp(i\theta\mathbf{p}_{\boxplus\boxminus}(ij))]^* , \quad (40)$$

where $[J]^T$ is the transpose of matrix $[J]$ in the aforementioned basis. Recall that any element of $\mathcal{V}_2^{(4)}$ can be decomposed to a sequence of 2-qudit gates $\exp(i\theta\mathbf{P}_{ij})$ and a global phase. Because products of unitaries that satisfy the above constraint also satisfy this constraint, it follows that for any $V \in \mathcal{V}_2^{(4)}$

$$\forall V \in \mathcal{V}_2^{(4)} : \quad [J][v_{\boxplus\boxplus}][J]^T = e^{i\phi} [v_{\boxplus\boxminus}]^* , \quad (41)$$

where $v_{\boxplus\boxplus} = \pi_{\boxplus\boxplus}(V)$, $v_{\boxplus\boxminus} = \pi_{\boxplus\boxminus}(V)$ are the components of V in $\mathcal{M}_{\boxplus\boxplus}$ and $\mathcal{M}_{\boxplus\boxminus}$ respectively, $[v_{\boxplus\boxminus}]^*$ is the complex conjugate of $[v_{\boxplus\boxminus}]$, and $e^{i\phi}$ is an unspecified phase that depends on V .

The relation in Eq. (41) means that the joint projection of $\mathcal{SV}_2^{(4)}$ to multiplicity spaces $\mathcal{M}_{\boxplus\boxplus}$ and $\mathcal{M}_{\boxplus\boxminus}$ does not

contain $SU(3) \times SU(3)$, as required by semi-universality. Rather, it satisfies

$$\begin{aligned} \pi_{\boxplus\boxplus, \boxplus\boxminus}(\mathcal{SV}_2^{(4)}) &:= \{(v_{\boxplus\boxplus}, v_{\boxplus\boxminus}) : V \in \mathcal{SV}_2^{(4)}\} \\ &\cong SU(\mathcal{M}_{\boxplus\boxplus}) \cong SU(\mathcal{M}_{\boxplus\boxminus}) \cong SU(3) , \end{aligned} \quad (42)$$

and the explicit form of the isomorphism is given by Eq. (41). In this situation we say the unitaries realized in $\boxplus\boxplus$ and $\boxplus\boxminus$ are ‘‘correlated’’.

On the other hand, since $\mathcal{M}_{\boxplus\boxplus}$ has a different dimension from $\mathcal{M}_{\boxplus\boxminus}$ and $\mathcal{M}_{\boxplus\boxminus}$, condition **B** is automatically satisfied for the pair \boxplus and $\boxplus\boxplus$, and the pair \boxplus and $\boxplus\boxminus$. In particular, according to Eq. (13), for both $\lambda = \boxplus\boxplus$ and $\lambda = \boxplus\boxminus$ the joint projection of $\mathcal{SV}_2^{(4)}$ to λ and \boxplus is isomorphic to

$$\begin{aligned} \pi_{\lambda, \boxplus}(\mathcal{SV}_2^{(4)}) &:= \{(v_\lambda, v_{\boxplus}) : V \in \mathcal{SV}_2^{(4)}\} \\ &\cong SU(\mathcal{M}_\lambda) \times SU(\mathcal{M}_{\boxplus}) \cong SU(3) \times SU(2) . \end{aligned} \quad (43)$$

In summary, the commutator subgroup of $\mathcal{V}_2^{(4)}$ is

$$\mathcal{SV}_2^{(4)} = [\mathcal{V}_2^{(4)}, \mathcal{V}_2^{(4)}] \cong SU(2) \times SU(3) ,$$

whereas the commutator subgroup of the group of all $SU(d)$ -invariant unitaries is

$$\mathcal{SV}^{(4)} = [\mathcal{V}^{(4)}, \mathcal{V}^{(4)}] \cong SU(2) \times SU(3) \times SU(3) ,$$

which means 2-qudit $SU(d)$ -invariant gates are not semi-universal. We revisit this example in Sec. IX C, and show that while $\mathcal{V}_2^{(4)}$ does not satisfy condition **B** of Lemma 1, because it still satisfies condition **A** (subsystem universality), it can be characterized via Lemma 8. (This lemma is an extension of Lemma 1 that does not assume condition **B**.)

Here, we saw the failure of semi-universality and constraints imposed by the locality of interactions in the case of a system with $n = 4$ qudits, as a consequence of Eq. (41). Indeed, this relation can be understood in terms of an anti-unitary transformation. See [6] for further details, where these facts can be interpreted both in terms of conservation of an observable K defined on two copies of the system, and in the language of a freely evolving fermionic system. This phenomenon, i.e., that the dynamics of 2-qudit symmetric interactions in distinct charge sectors determine each other, exists for general n when $d > 2$.⁵ For sufficiently many qudits n , as the dimension of the Hilbert space of each qudit d grows, there will be more such constraints among different sectors [6]. Furthermore, in addition to such correlations,

⁵ The reason that it does not exist for $d = 2$, i.e., that 2-qudit $SU(2)$ -invariant unitaries are semi-universal on qubits, is related to the fact that in this case there is no representation associated with the diagram \boxplus .

which correspond to type **IV** constraints, we also see type **III** constraints. For instance, for $n = 5$ qudits with $d \geq 3$, type **III** constraints appear: in the charge sector $\square\square\square$ we get the group $\text{SO}(6)$ rather than $\text{SU}(6)$ [6], which is required by semi-universality. In addition, the sectors $\square\square\square$ and $\square\square$ are correlated with each other, and similarly, the three sectors $\square\square\square$, $\square\square$, and \square are all correlated. It is worth noting again that these phenomena, i.e., correlations between unitaries in sectors with inequivalent irreps of $G = \text{SU}(d)$ (type **IV**) and appearance of irreducible subgroups of the realizable unitaries in one sector (type **III**), cannot occur in the case of Abelian symmetry groups G [8].

C. Generic 3-qudit gates are semi-universal

Next, we show that 2-qudit $\text{SU}(d)$ -invariant gates together with any $\text{SU}(d)$ -invariant unitary $Y \in \mathcal{V}^{(4)}$ that breaks the constraint in Eq. (41) become semi-universal on $n = 4$ qudits. Furthermore, such Y can be chosen as a 3-qudit gate that acts on 3 out of 4 qudits in the system (tensor product with the identity operator on the 4th qudit).

Proposition 2. *Any 4-qudit $\text{SU}(d)$ -invariant unitary Y together with 2-qudit $\text{SU}(d)$ -invariant unitaries are semi-universal on $n = 4$ qudits, i.e., $\mathcal{SV}^{(4)} \subset \langle Y, V : V \in \mathcal{V}_2^{(4)} \rangle$ if, and only if,*

$$[J][\pi_{\square\square\square}(Y)][J]^T \neq e^{i\phi}[\pi_{\square\square}(Y)]^* , \quad (44)$$

for any phase $e^{i\phi}$, where $[\pi_{\square\square\square}(Y)]$ and $[\pi_{\square\square}(Y)]$ are the matrix representations of the components of Y in $\mathcal{M}_{\square\square\square}$ and $\mathcal{M}_{\square\square}$, in the basis defined in the Table III where matrix $[J]$ is also defined.

This is proven in Appendix A 2. Note that if the unitary $Y \in \mathcal{V}^{(4)}$ satisfies

$$|\text{Tr}(\pi_{\square\square\square}(Y))| \neq |\text{Tr}(\pi_{\square\square}(Y))| , \quad (45)$$

then Eq. (44) cannot hold as equality for any phase ϕ and unitary J , which means, together with 2-qudit unitaries, Y is semi-universal. This allows using the characterization of elements of $\mathcal{V}_2^{(3)}$ in Proposition 1 to demonstrate that a generic 3-qudit $\text{SU}(d)$ -invariant unitary, together with 2-qudit ones, is semi-universal.

Proposition 3 (Generic 3-qudit gates are semi-universal on $n = 4$ qudits). *For any 3-qudit unitary $S \in \mathcal{V}^{(3)}$ that is not realizable with 2-qudit $\text{SU}(d)$ -invariant gates, i.e., $S \in \mathcal{V}^{(3)} \setminus \mathcal{V}_2^{(3)}$, the 4-qudit unitary $Y = S \otimes \mathbb{I}$ satisfies the requirement in Eq. (44) of the above proposition, and therefore the 3-qudit gate S together with 2-qudit unitaries $\exp(i\theta \mathbf{P}_{ij}) : \theta \in [0, 2\pi)$ are semi-universal, i.e., $\mathcal{SV}^{(4)} \subset \langle S \otimes \mathbb{I}, V : V \in \mathcal{V}_2^{(4)} \rangle$.*

Together with the results of Sec. V, in particular Eq. (53), this implies that any 3-qudit gate $S \in \mathcal{V}^{(3)} \setminus \mathcal{V}_2^{(3)}$ together with 2-qudit $\text{SU}(d)$ -invariant gates is semi-universal on n qudits, for all $n \geq 3$.

Proposition 3 is proven in Appendix A 2. Next, we present an example of a unitary Y satisfying the constraint in the above lemma.

Example: Semi-universality with 3-qudit reflections

Recall the three-qudit reflection unitaries $R_+ = \exp(i\pi\Pi_{\square\square\square})$ and $R_- = \exp(i\pi\Pi_{\square\square})$ in Eq. (2). Suppose in addition to the 2-qudit $\text{SU}(d)$ -invariant gates, one can also use one of the gates R_+ or R_- . Is this set semi-universal on $n = 4$ qudits?

It is easy to verify that $R_{\pm} \notin \mathcal{V}_2^{(3)}$ when $d \geq 3$ using Proposition 1. For instance, rewriting as

$$R_+ = -\Pi_{\square\square\square} + \Pi_{\square\square} + \Pi_{\square} , \quad (46)$$

we find that Eq. (30) does not hold, i.e.,

$$1 = \det \pi_{\square\square}(R_+) \neq (\det \pi_{\square\square\square}(R_+))(\det \pi_{\square}(R_+)) = -1. \quad (47)$$

Equivalently, we can see this using the criterion in Proposition 2 or trace condition in Eq. (45). In particular, it can be easily seen that $R_{\pm} \otimes \mathbb{I}$ acts trivially on one of $\mathcal{M}_{\square\square}$ and $\mathcal{M}_{\square\square\square}$ and non-trivially in the other (depending on the sign \pm). For instance, while $R_+ \otimes \mathbb{I}$ acts trivially in $\mathcal{M}_{\square\square}$, inside $\mathcal{M}_{\square\square\square}$ it has the matrix representation


$$[\pi_{\square\square\square}(R_+ \otimes \mathbb{I})] = \left(\begin{array}{c|c} 1 & \\ \hline & -1 \end{array} \right) , \quad (48)$$

which can be seen from Eq. (46) together with Table III. This means Eq. (44) cannot hold as equality for any phase $e^{i\phi}$. In particular,

$$\begin{aligned} |\text{Tr}(\pi_{\square\square\square}(R_+))| &= 1 \\ |\text{Tr}(\pi_{\square\square}(R_+))| &= 3 , \end{aligned} \quad (49)$$

so the condition in Eq. (45) holds.

It is also worth noting that for generic values of $\theta \in [0, 2\pi)$, the unitary realized by the following circuit is in $\mathcal{SV}^{(4)}$ but not in $\mathcal{SV}_2^{(4)}$.



Indeed, since R_{\pm} acts trivially in $\mathcal{M}_{\square\square}$ and one of $\mathcal{M}_{\square\square}$ or $\mathcal{M}_{\square\square\square}$, the unitary realized by the above circuit acts

trivially in all sectors except \mathcal{M}_{\boxplus} , or \mathcal{M}_{\boxminus} , depending on the sign \pm of R_{\pm} . Furthermore, in this sector, the realized unitary does not act as a global phase, which can be seen by noting that it is not permutationally invariant. Therefore, Proposition 2 implies that each one of R_+ or R_- , together with 2-qudit $SU(d)$ -invariant gates are semi-universal on $n = 4$ qudits.

It is worth noting that the reflection unitary for the charge sector \boxplus , i.e., $\exp(i\pi\Pi_{\boxplus})$, is already contained in $\mathcal{SV}^{(3)} \subseteq \mathcal{V}_2^{(3)}$, since $\det \pi_{\lambda}(\exp(i\pi\Pi_{\boxplus})) = 1$ for all $\lambda \in \Lambda_{3,d}$.

V. SEMI-UNIVERSALITY OF 3-QUDIT GATES ON ARBITRARY NUMBER OF QUDITS

Next, we prove the semi-universality of 3-qudit gates for systems with $n \geq 4$, i.e., Theorem 1. To establish this result, we use induction. Applying Lemma 1, we show that by composing m -qudit unitaries in $\mathcal{SV}^{(m)}$, one can obtain any element of $\mathcal{SV}^{(m+1)}$ on $m+1$ qudits. More precisely, let

$$\mathcal{W}^{(m+1)} = \langle \mathbf{P}_{ij}(\mathcal{SV}^{(m)} \otimes \mathbb{I})\mathbf{P}_{ij} : i \neq j \rangle \quad (50)$$

denote the group generated by m -local unitaries in $\mathcal{SV}^{(m)}$ acting on $m+1$ qudits. It is worth noting that $\mathcal{W}^{(m+1)}$ is generated by those unitaries restricted to only acting on the first m or the last m qudits, i.e.⁶

$$\mathcal{W}^{(m+1)} = \langle \mathcal{SV}^{(m)} \otimes \mathbb{I}, \mathbb{I} \otimes \mathcal{SV}^{(m)} \rangle. \quad (51)$$

Then, we show that

$$\mathcal{W}^{(m+1)} = \mathcal{SV}^{(m+1)} \quad \text{for } m \geq 4. \quad (52)$$

This means that for any $n \geq 4$, the group generated by the permuted versions of $\mathcal{SV}^{(4)}$ is equal to $\mathcal{SV}^{(n)}$, i.e.,

$$\langle \mathbf{P}_{ij}(\mathcal{SV}^{(4)} \otimes \mathbb{I}^{\otimes(n-4)})\mathbf{P}_{ij} : i \neq j \rangle = \mathcal{SV}^{(n)}. \quad (53)$$

As we showed in the previous section for 4 qudits, $\mathcal{SV}^{(4)}$ can be generated by 3-qudit $SU(d)$ -invariant unitaries. Combining these results, we conclude that 3-qudit $SU(d)$ -invariant gates are semi-universal on n qudits systems, i.e., they generate $\mathcal{SV}^{(n)}$, which proves Theorem 1.

To prove Eq. (52), first we note that $\mathcal{W}^{(m+1)} \subseteq \mathcal{SV}^{(m+1)}$, which follows from the fact that

$$\mathcal{SV}^{(m)} \otimes \mathbb{I} \subseteq \mathcal{SV}^{(m+1)}. \quad (54)$$

This can be seen, for instance, by noting that $\mathcal{SV}^{(m)} \otimes \mathbb{I} \subseteq \mathcal{V}^{(m+1)}$, together with $\mathcal{SV}^{(m)} = [\mathcal{SV}^{(m)}, \mathcal{SV}^{(m)}]$.

Therefore, to prove Eq. (52) we need to show that $\mathcal{W}^{(m+1)}$ contains all elements of $\mathcal{SV}^{(m+1)}$. The example case $n = 4$, discussed in Sec. IV, introduces most of the main ideas that are needed in this proof. The only new ingredients are a few useful facts about the representation theory of \mathbb{S}_n that are reviewed in the next section, and the following lemma which is of independent interest.

Lemma 4. *On a system with n qudits, consider the group generated by one-parameter families $e^{it(\mathbf{P}_{ij}-\mathbf{P}_{kl})}$ acting on qudits $i \neq j$ and $k \neq l$. This group is equal to the commutator subgroup of $\mathcal{V}_2^{(n)}$, i.e.*

$$\langle e^{it(\mathbf{P}_{ij}-\mathbf{P}_{kl})} : t \in \mathbb{R}, i \neq j, k \neq l \rangle = \mathcal{SV}_2^{(n)}. \quad (55)$$

In particular, for all irreps $\lambda \in \Lambda_{n,d}$, this group acts irreducibly on \mathcal{M}_{λ} .

This is proven in Appendix B6 using the fact that any swap $i\mathbf{P}_{ij}$ can be written as a linear combination of the differences $i(\mathbf{P}_{ij}-\mathbf{P}_{kl}) : i \neq j, k \neq l$ and the permutationally-invariant operator $iB = i\sum_{i \neq j} \mathbf{P}_{ij}$, which is in the center of the Lie algebra generated by transpositions.

A. Useful facts about the symmetric group \mathbb{S}_n

Recall the isotypic decomposition in Eq. (18),

$$(\mathbb{C}^d)^{\otimes(m+1)} = \bigoplus_{\lambda \in \Lambda_{m+1,d}} \mathcal{Q}_{\lambda} \otimes \mathcal{M}_{\lambda},$$

where $\Lambda_{m+1,d}$ is the set of Young diagrams with $m+1$ boxes and $\leq d$ rows, and \mathcal{M}_{λ} is the irrep of \mathbb{S}_{m+1} labeled with the Young diagram λ . Consider $\mathbb{S}_m \subset \mathbb{S}_{m+1}$ the subgroup which permutes the first m qudits. Then, the branching rule of \mathbb{S}_m says that, for each $\lambda \in \Lambda_{m+1,d}$, there is $\Gamma_{\lambda} \subseteq \Lambda_{m,d}$ such that

$$\mathcal{M}_{\lambda} = \bigoplus_{\gamma \in \Gamma_{\lambda}} \mathcal{M}_{\gamma}, \quad (56)$$

where each irrep in $\gamma \in \Gamma_{\lambda}$ of \mathbb{S}_m shows up with multiplicity one and the subspace \mathcal{M}_{γ} carries that irrep [9, 17]. For the proof of Theorem 1, in addition to this fact, we will also use the following facts.

Fact. Whenever $m \geq 4$ and $\lambda \in \Lambda_{m+1,d}$ is not one-dimensional (i.e., it is neither the trivial nor sign irrep of \mathbb{S}_{m+1}),

1. If λ has $\leq d$ rows then each $\gamma \in \Gamma_{\lambda}$ has $\leq d$ rows.
2. At least one irrep in the branching, $\gamma \in \Gamma_{\lambda} \subseteq \Lambda_{m,d}$, has $\dim \mathcal{M}_{\gamma} \geq 3$.
3. For any distinct $\lambda \neq \lambda' \in \Lambda_{m+1,d}$, there is some $\gamma \in \Lambda_{m,d}$ with $\dim \mathcal{M}_{\gamma} \geq 2$ which is in one of, but not both, Γ_{λ} or $\Gamma_{\lambda'}$.

⁶ This follows because every even permutation, i.e. element of the alternating subgroup $\sigma \in \mathbb{A}_m \subseteq \mathbb{S}_m$, is contained in $\mathcal{SV}^{(m)}$, as $\det \mathbf{p}_{\lambda}(\sigma) = 1$ for all $\lambda \in \Lambda_n$. Thus the group on the right-hand side of Eq. (51) is invariant under conjugation by $\mathbf{P}(\sigma) : \sigma \in \mathbb{A}_{m+1}$ and by \mathbf{P}_{23} , which together generate all of \mathbb{S}_{m+1} .




FIG. 3. The beginning of Young's lattice, from which the branching rule may be read: the representations that show up in the branching Eq. (56) of the Young diagram λ are precisely the subdiagrams, which are connected by an edge in the lattice. The last layer corresponds to $n = 5$ qudits. One can easily check Fact 1, Fact 2, and Fact 3. For instance, Eq. (56) implies that except for the two Young diagrams at the two ends of each layer, namely single-row and single-column Young diagrams, the rest of the diagrams have dimensions larger than 1, and indeed for diagrams with $n = 5$, they all have dimensions 3 or larger. We can explicitly check each fact for $\lambda \in \Lambda_{5,d}$. According to Fact 1, for any irrep $\lambda \in \Lambda_{5,d}$, the corresponding irreps in their branching denoted as Γ_λ , which are the diagrams connected to λ in the previous level, have less or equal number of rows. Furthermore, as stated in Fact 2, all the irreps $\lambda \in \Lambda_{5,d}$ that are not one-dimensional, namely everything except the diagrams at the left and right corners, have either $\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}$ or $\begin{smallmatrix} \square & \square \\ & \square \end{smallmatrix}$ in their branching, both of which are three-dimensional. Also, as stated in Fact 3, for any distinct irreps $\lambda \neq \lambda' \in \Lambda_{5,d}$ which are not one-dimensional, at least one of $\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}$, $\begin{smallmatrix} \square & \square \\ & \square \end{smallmatrix}$, or $\begin{smallmatrix} \square & \square \\ & \square \\ & \square \end{smallmatrix}$ is in the branching of λ but not λ' .

These facts can be deduced from Young's lattice in Fig. 3 (in the caption we explain these three facts for the example of $n = 5$). Note that Fact 2 and Fact 3 do not hold for $n = 4$, and this is another way to understand why 2-qudit gates are not semi-universal on $n = 4$ qudits (or, more precisely, why the induction starts at $m = 4$ rather than $m = 3$).

B. Extending semi-universality from m to $m + 1$ qudits (Proof of Eq. (52))

Here, we prove Eq. (52). Similar to the proof in the special case of $n = 4$, the proof is in 2 steps, namely we establish conditions **A** and **B** of Lemma 1.

- **Condition A:** We show that for each irrep $\lambda \in \Lambda_{m+1,d}$, the projection of $\mathcal{W}^{(m+1)}$ to irrep \mathcal{M}_λ is equal to

$$\pi_\lambda(\mathcal{W}^{(m+1)}) = \pi_\lambda(\mathcal{S}\mathcal{V}^{(m+1)}) = \text{SU}(\mathcal{M}_\lambda). \quad (57)$$

Since this condition is trivially satisfied when $\dim(\mathcal{M}_\lambda) = 1$, in the following, without loss of generality, we assume

$\dim(\mathcal{M}_\lambda) > 1$. Similar to the case of $n = 4$ qudits, first we consider elements of $\mathcal{W}^{(m+1)}$ that act trivially on qudit $m + 1$, i.e., unitaries in the form $(V \otimes \mathbb{I}) : V \in \mathcal{S}\mathcal{V}^{(m)}$. Relative to the decomposition in Eq. (56), the projection of this subgroup of $\mathcal{W}^{(m+1)}$ to \mathcal{M}_λ is equal to

$$\{\pi_\lambda(V \otimes \mathbb{I}) : V \in \mathcal{S}\mathcal{V}^{(m)}\} = \bigoplus_{\gamma \in \Gamma_\lambda} \text{SU}(\mathcal{M}_\gamma). \quad (58)$$

Again, this block-diagonal form is a consequence of the fact that qudit $m + 1$ does not interact with the rest of qudits. Then, we argue that as soon as we include interactions with this qudit we get the entire $\text{SU}(\mathcal{M}_\lambda)$. To achieve this we apply Lemma 2, whose assumptions are verified in the following:

1. In decomposition $\mathcal{M}_\lambda = \bigoplus_{\gamma \in \Gamma_\lambda} \mathcal{M}_\gamma$, for at least one irrep $\gamma^* \in \Gamma_\lambda$, $\dim \mathcal{M}_{\gamma^*} \geq 3$. This follows immediately from Fact 2, which holds because $m \geq 4$ and λ is not a 1D irrep of \mathbb{S}_{m+1} .
2. The one-parameter families $\exp(it(\mathbf{P}_{ij} - \mathbf{P}_{kl})) : t \in \mathbb{R}$ for all $i \neq j, k \neq l \in \{1, \dots, m + 1\}$ are inside $\mathcal{W}^{(m+1)}$, and according to Lemma 4, the group generated by them acts irreducibly on \mathcal{M}_λ for all $\lambda \in \Lambda_{m+1,d}$. To see why these families are inside $\mathcal{W}^{(m+1)}$, note that according to Lemma 4, $\exp(it(\mathbf{P}_{ij} - \mathbf{P}_{kl}))$ is in the group $\mathcal{S}\mathcal{V}^{(m)}$ defined on any m out of $m + 1$ qudits that contain qudits i, j, k, l . Then, together with Eq. (54), this implies that on $m + 1$ qudits, it is inside $\mathcal{W}^{(m+1)}$.

Therefore, both assumptions of Lemma 2 are satisfied for $\pi_\lambda(\mathcal{W}^{(m+1)})$, which implies $\pi_\lambda(\mathcal{W}^{(m+1)}) = \text{SU}(\mathcal{M}_\lambda)$. In summary, for all $\lambda \in \Lambda_{m+1,d}$, the subsystem universality holds, which means condition **A** of Lemma 1 is satisfied.

- **Condition B:** Consider an arbitrary pair of distinct irreps $\lambda, \lambda' \in \Lambda_{m+1,d}$. If one or both of λ and λ' are 1D irreps of \mathbb{S}_{m+1} , then condition **B** is satisfied for that pair. Therefore, without loss of generality, we assume they both have dimensions larger than one. Then, using Fact 3 we know that the branching sets $\Gamma_\lambda \neq \Gamma_{\lambda'}$, with some differing irrep having dimension greater than two. Without loss of generality, let $\gamma \in \Gamma_\lambda$ with $\gamma \notin \Gamma_{\lambda'}$ and $\dim \mathcal{M}_\gamma \geq 2$. Consider the subgroup of $\mathcal{S}\mathcal{V}^{(m)}$ isomorphic to $\text{SU}(\mathcal{M}_\gamma)$ formed from m -qudit $\text{SU}(d)$ -invariant unitaries that act trivially in all sectors except \mathcal{M}_γ . Then, for any unitary V in this subgroup $V \otimes \mathbb{I}$ acts trivially in $\mathcal{M}_{\lambda'}$ and non-trivially in \mathcal{M}_λ . That is, $\pi_{\lambda'}(V \otimes \mathbb{I}) = \mathbb{I}_{\lambda'}$ and $\pi_\lambda(V \otimes \mathbb{I}) = \pi_\gamma(V) \oplus \mathbb{I}_\perp$, where \mathbb{I}_\perp is the identity operator on the orthogonal complement of \mathcal{M}_γ in \mathcal{M}_λ . We conclude that condition **B'** of Lemma 1, and thus, condition **B** of this lemma are also satisfied.

In conclusion, the subgroup $\mathcal{W}^{(m+1)} \subset \mathcal{SV}^{(m+1)}$ satisfies both assumptions **A** and **B** of Lemma 1, and therefore, this lemma implies that $\mathcal{W}^{(m+1)} = \mathcal{SV}^{(m+1)}$. This completes the proof of semi-universality of 3-qudit $\text{SU}(d)$ -invariant gates.

VI. IMPOSSIBILITY OF ACHIEVING UNIVERSALITY WITH $(n-1)$ -QUDIT GATES

We saw that semi-universality can be achieved with 3-qudit gates. The next natural question is what is the minimum locality of gates that is needed to achieve universality. More precisely, assuming we cannot use any ancilla qudits, what is the minimum k for which k -qudit $\text{SU}(d)$ -invariant gates become universal on n qudits, such that $\mathcal{V}_k^{(n)} = \mathcal{V}_n^{(n)} = \mathcal{V}^{(n)}$?

For qubit systems with $\text{SU}(2)$ symmetry, which corresponds to the special case of $d=2$ in the present paper, it was shown in [7] that universality is achieved with $k=2\lfloor n/2 \rfloor$. This means that there is an even-odd effect, which was shown in [7] to be related to time-reversal symmetry. Specifically, when n is odd, $(n-1)$ -local symmetric gates are universal, but they are not when n is even. Here we show that, for $d \geq 3$, there is no such effect: without using ancillary qudits, $(n-1)$ -qudit $\text{SU}(d)$ -invariant gates are not universal. This proves Corollary 2.

According to the general results of [5], which apply to arbitrary symmetry groups, the semi-universality of k -qudit $\text{SU}(d)$ -invariant gates for $k \geq 3$ together with the fact $\text{SU}(d)$ is a connected Lie group imply that the difference between the dimensions of the Lie group of all $\text{SU}(d)$ -invariant unitaries on n qudits, and the subgroup generated by k -qudit $\text{SU}(d)$ -invariant gates is

$$\dim(\mathcal{V}_n^{(n)}) - \dim(\mathcal{V}_k^{(n)}) = |\Lambda_{n,d}| - |\Lambda_{k,d}|. \quad (59)$$

Here, $|\Lambda_{n,d}|$ is the size of $\Lambda_{n,d}$, the set of inequivalent irreps of $\text{SU}(d)$, or equivalently \mathbb{S}_n , that appear on n qudits, which can be labeled by Young diagrams with n boxes and $\leq d$ rows. In particular, in the case of $\text{SU}(d)$ symmetry, the arguments of [5] imply that, because $\text{SU}(d)$ is a connected Lie group, the center of the Lie algebra associated to $\mathcal{V}_k^{(n)}$ has dimension $|\Lambda_{k,d}|$, whereas the center of the Lie algebra associated to $\mathcal{V}_n^{(n)} = \mathcal{V}^{(n)}$ has dimension $|\Lambda_{n,d}|$, and because of semi-universality this is exactly the difference between dimensions of $\mathcal{V}_n^{(n)}$ and $\mathcal{V}_k^{(n)}$. In Appendix C, we show that

Lemma 5 (Strict monotonicity of the number of irreps, for $d \geq 3$). *Let $|\Lambda_{k,d}|$ be the number of inequivalent irreps of $\text{SU}(d)$, or equivalently \mathbb{S}_k , on k qudits with the total Hilbert space $(\mathbb{C}^d)^{\otimes k}$. Then, for $d \geq 3$, $|\Lambda_{k,d}| > |\Lambda_{k-1,d}|$ for all k .*

In summary, for $d \geq 3$, we have

$$|\Lambda_{n,d}| = |\Lambda_{k,d}| \iff n = k.$$

Then, together with Eq. (59), this proves the non-universality of k -qudit gates with $k < n$, and we have the following corollary,

Corollary 2. *Consider a system of n qudits with $d \geq 3$. Then, k -local $\text{SU}(d)$ -invariant unitaries are universal, if and only if $k = n$. In other words, if $k < n$, then $\mathcal{V}_k^{(n)} \subsetneq \mathcal{V}_n^{(n)}$ is a proper subgroup.*

We also note that while there is no simple general formula for $|\Lambda_{k,d}|$, it can be obtained [18] using its generating function

$$\sum_{k=0}^{\infty} |\Lambda_{k,d}| x^k = \prod_{l=1}^d \frac{1}{1-x^l}. \quad (60)$$

Using this method, in Fig. 4 we plot $|\Lambda_{k,d}|$ as a function of k . We see that $|\Lambda_{k,d}|$ is strictly monotonic in k for $d \geq 3$, while for $d=2$ it grows only every other step in k .




FIG. 4. The vertical axis is $|\Lambda_{k,d}|$, the number of inequivalent irreps of $\text{SU}(d)$ on k qudits, which according to the results of [5], is the the dimension of the center of the Lie algebra associated to $\mathcal{V}_k^{(n)}$, the group generated by k -qudit $\text{SU}(d)$ -invariant gates. When $d \geq 3$, $|\Lambda_{k,d}|$ is strictly monotonic in k . When $d=2$, $|\Lambda_{k,d}|$ increases only every other step in k .

Finally, we discuss the asymptotic behavior of $|\Lambda_{k,d}|$ in the regime $k \gg d$, which implies Eq. (3) and explain the plot in Fig. 2. Recall that $|\Lambda_{k,d}|$ is the number of partitions of k into $i \leq d$ positive integers, i.e., $k_1 \geq \dots \geq k_i$ and $k = k_1 + k_2 + \dots + k_i$.

In the limit of large $k \gg d$, the number of such partitions for each $i \leq d$ can be calculated through the number of the composition of k into exactly i parts, namely $\binom{k-1}{i-1}$. However, the order does not matter in integer partition, whereas it matters in integer composition, which should be corrected. When $k \gg i$, for almost all partitions, k_1, \dots, k_i take distinct values, and we can correct it ap-

proximately by a factor of $1/i!$. This implies that⁷

$$|\Lambda_{k,d}| \approx \sum_{i=1}^d \frac{1}{i!} \binom{k-1}{i-1} = \frac{k^{d-1}}{d!(d-1)!} + O(k^{d-2}). \quad (61)$$

Putting this into Eq. (59), we arrive at

$$\begin{aligned} \dim \mathcal{V}_k^{(n)} - \dim \mathcal{V}_3^{(n)} &= |\Lambda_{k,d}| - |\Lambda_{3,d}| = |\Lambda_{k,d}| - 3 \\ &\approx \frac{k^{d-1}}{d!(d-1)!} + O(k^{d-2}), \end{aligned}$$

where we have used the fact that $\Lambda_{3,d}$, given in Eq. (26), has 3 elements. This proves Eq. (3).

VII. THE USE OF ANCILLA QUDITS FOR ACHIEVING (SEMI-)UNIVERSALITY

In this section, we consider the use of ancillae for achieving both universality and semi-universality and prove various results in this context, including those presented in Table I. First, we consider achieving universality from semi-universality for arbitrary symmetry group G with on-site representation, and prove Lemma 3. That is, we prove that universality can be achieved on n qudits if semi-universality holds on $n+3$ qudits, and the 3 qudits are used as ancillae. We note that in general, one might be able to achieve universality with fewer ancilla qudits (for instance, 1 qudit in the case of Abelian symmetries [8], and 2 qubits in the case of $SU(2)$ symmetry with qubit systems [6]). Then, we focus on the case of 2-qudit gates with $SU(d)$ symmetry. Even though 2-qudit gates are not semi-universal for $n > 3$ qudits, by studying a system with 11 qudits with the total Hilbert space $(\mathbb{C}^d)^{\otimes 11}$, we show that 8 qudits can be used as ancillae to implement any $SU(d)$ -invariant unitary on the remaining 3 qudits. In the special case of $d = 3$, this can be achieved with only 6 ancilla qudits.

Since the state of ancillae remains unchanged and uncorrelated with the rest of qudits, we can reuse it to realize arbitrary 3-qudit gates on a system with arbitrary $n > 3$ qudits. We have previously shown in Theorem 1 that 3-qudit gates are semi-universal on a system with arbitrary n . Combining these we conclude that

Corollary 3. *Using 2-qudit $SU(d)$ -invariant unitaries, and 8 ancilla qudits we can realize any unitary in $\mathcal{S}\mathcal{V}^{(n)}$, and using 11 ancilla qudits we can realize any unitary in $\mathcal{V}^{(n)}$. In the special case of $d = 3$, these can be achieved with 6 and 9 ancilla qudits, respectively.*

A. From semi-universality to universality with three ancilla qudits (Proof of Proposition 3)

Consider an arbitrary symmetry group G with the on-site representation, as defined in Lemma 3. Consider 3 ancillary qudits, with the total Hilbert space

$$(\mathbb{C}^d)^{\otimes 3} = \mathcal{Q}_{\square\square\square} \oplus (\mathcal{Q}_{\square\square} \otimes \mathcal{M}_{\square}) \oplus \mathcal{Q}_{\square}, \quad (62)$$

where we have decomposed this Hilbert space according to the irreps of the permutation group \mathbb{S}_3 , and used the notation introduced in Eq. (18). Since it is permutationally-invariant, $u(g)^{\otimes 3}$ acts on $\mathcal{Q}_{\square\square} \otimes \mathcal{M}_{\square}$ as $u_{\square\square}(g) \otimes \mathbb{I}_{\mathcal{M}_{\square}}$, where $u_{\square\square}$ is a possibly reducible representation of G on $\mathcal{Q}_{\square\square}$.

Let $Z_{\square\square}$ be a nonzero Hermitian operator on $\mathcal{M}_{\square\square} \cong \mathbb{C}^2$ with $\text{Tr} Z_{\square\square} = 0$. Then, for any Hermitian G -invariant operator H on $(\mathbb{C}^d)^{\otimes n}$, we consider operator \tilde{H} on $n+3$ qudits, defined by

$$H \mapsto \tilde{H} = H \otimes (\mathbb{I}_{\mathcal{Q}_{\square\square}} \otimes Z_{\square\square}). \quad (63)$$

Then, \tilde{H} is also Hermitian, G -invariant, and in addition, it is centerless, because

$$\text{Tr}(u(g)^{\otimes(n+3)} \tilde{H}) = \text{Tr}(u(g)^{\otimes n} H) \text{Tr}(u_{\square\square}(g)) \text{Tr}(Z_{\square\square}) = 0, \quad (64)$$

for all $g \in G$, which follows from $\text{Tr} Z_{\square\square} = 0$. We conclude that the family of unitaries $\exp(i\tilde{H}t) : t \in \mathbb{R}$ are in the commutator subgroup of G -invariant unitaries on $n+3$ qudits, and therefore they can be realized by any set of gates that are semi-universal on $n+3$ qudits. For instance, in the case of $SU(d)$ symmetry discussed in the previous sections, $\exp(i\tilde{H}t)$ is in $\mathcal{S}\mathcal{V}^{(n+3)}$, which means it is realizable with 3-qudit $SU(d)$ -invariant unitaries.

In general, \tilde{H} is an entangling Hamiltonian. However, by preparing the 3 ancilla qudit in an eigenstate $|\eta\rangle \in (\mathbb{C}^d)^{\otimes 3}$ of $\mathbb{I}_{\mathcal{Q}_{\square\square}} \otimes Z_{\square\square}$, they remain unentangled with the system. That is,

$$\exp(i\tilde{H}t)(|\psi\rangle \otimes |\eta\rangle) = (\exp(iH\alpha t)|\psi\rangle) \otimes |\eta\rangle, \quad (65)$$

where α is the eigenvalue of $\mathbb{I}_{\mathcal{Q}_{\square\square}} \otimes Z_{\square\square}$ for eigenvector $|\eta\rangle$. Some examples of choices of $|\eta\rangle$ and $Z_{\square\square}$ include

- $|\eta_1\rangle = \frac{1}{2\sqrt{3}}(2|100\rangle + (\sqrt{3}-1)|010\rangle - (1+\sqrt{3})|001\rangle)$ and $\mathbb{I}_{\mathcal{Q}_{\square\square}} \otimes Z_{\square\square} = \frac{1}{\sqrt{3}}(\mathbf{P}_{12} - \mathbf{P}_{13})$.
- $|\eta_2\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle) \otimes |0\rangle$ and $\mathbb{I}_{\mathcal{Q}_{\square\square}} \otimes Z_{\square\square} = \frac{1}{3}(-2\mathbf{P}_{12} + 2\mathbf{P}_{23} + \mathbf{P}_{(123)} - \mathbf{P}_{(132)})$.

Here, $Z_{\square\square}$ is normalized so that its eigenvalues are ± 1 , and the given state is an eigenvector with eigenvalue $\alpha = 1$. Note that the above particular linear combination of permutations is restricted to irrep $\square\square$, such that

$$(\mathbb{I}_{\mathcal{Q}_{\square\square}} \otimes Z_{\square\square})\Pi_{\lambda} = 0 : \quad \lambda = \square\square\square, \square\square, \quad (66)$$

which can be seen using the fact that $\mathbf{P}(\sigma)\Pi_{\square\square} = \Pi_{\square\square}$ and $\mathbf{P}(\sigma)\Pi_{\square} = \text{sgn}(\sigma)\Pi_{\square}$.

⁷ We note that this formula can also be obtained using Sylvester's denominator formula [19] or the asymptotic formula from [18].

B. (Semi-)universality with 2-qudit $SU(d)$ -invariant gates and ancillae (Proof of Corollary 3)

Theorem 1 shows that 3-qudit $SU(d)$ -invariant gates are semi-universal on arbitrary n qudits. In this section we show how general 3-qudit $SU(d)$ -invariant gates can be realized with 2-qudit gates, provided that one can use 8 (or 6, in the special case of $d = 3$) ancilla qudits. Since the state of the ancillae remains unchanged, we can reuse it arbitrarily many times to implement all 3-qudit gates in the circuit. Therefore, this together with Theorem 1, proves Corollary 3.

As we proved in Sec. IV A, on $n = 3$ qudits, 2-qudit gates are semi-universal, i.e., $\mathcal{SV}^{(3)} \subset \mathcal{V}_2^{(3)}$. Therefore, in this case, the only constraints that the locality of gates imposes on the realizable unitaries are on the relative phases between sectors with different irreps of symmetry, namely type **I** constraints. Hence, to achieve universality it suffices to amend $\mathcal{SV}^{(3)}$ with relative phases between different sectors, i.e., unitaries in the form

$$\sum_{\lambda \in \Lambda_3} e^{i\theta_\lambda} \Pi_\lambda : \quad \theta_\lambda \in [0, 2\pi), \quad (67)$$

where $\Lambda_3 = \{\square\square\square, \square\square, \square\}$ (we note that a 2D Lie subgroup of this group is already included in $\mathcal{V}_2^{(3)}$). This may sound similar to the problem we studied in the previous section, where we used 3 ancilla qudits to elevate semi-universality to universality. However, there is an important difference: to apply that technique we need semi-universality on $n + 3$ qudits, i.e., the main qudits and the ancillae. Therefore, in the context of the problem at hand, this will require semi-universality on 6 qudits. However, as we discussed before, for $n > 3$ qudits 2-qudit gates are not semi-universal. Therefore, the technique of the previous section is not applicable here, and we need to develop a new scheme that works even in the absence of semi-universality.

We have found a solution to this problem that uses 8 ancilla qudits. This solution is based on the following fact about systems with 11 qudits. Recall that the wedge product of states is defined as

$$|\psi_1\rangle \wedge \dots \wedge |\psi_m\rangle := \frac{1}{\sqrt{m!}} \sum_{\sigma \in \mathcal{S}_m} \text{sgn}(\sigma) \mathbf{P}(\sigma)(|\psi_1\rangle \otimes \dots \otimes |\psi_m\rangle). \quad (68)$$

Lemma 6 (Centerless Hamiltonians on 11 qudits). *For any centerless $SU(d)$ -invariant Hamiltonian H on $(\mathbb{C}^d)^{\otimes 11}$ (i.e., a Hamiltonian satisfying $\text{Tr}(Hu(g)^{\otimes 11}) = 0$ for all $g \in SU(d)$) there exists a Hamiltonian \tilde{H} that is realizable with 2-qudit $SU(d)$ -invariant Hamiltonians, i.e., $\exp(it\tilde{H}) \in \mathcal{V}_2 : t \in \mathbb{R}$, which satisfies*

$$\exp(it\tilde{H})(|\psi\rangle \otimes |\eta\rangle) = \exp(itH)(|\psi\rangle \otimes |\eta\rangle), \quad (69)$$

for all $|\psi\rangle \in (\mathbb{C}^d)^{\otimes 3}$, where

$$|\eta\rangle = (|0\rangle \wedge |1\rangle \wedge |2\rangle \wedge |3\rangle)^{\otimes 2} \in (\mathbb{C}^d)^{\otimes 8}, \quad (70)$$

is an 8-qudit state. Furthermore, in the special case of $d = 3$, the same fact holds for any centerless Hamiltonian on 9 qudits, and state

$$|\eta'\rangle = (|0\rangle \wedge |1\rangle)^{\otimes 2} \otimes |00\rangle \in (\mathbb{C}^d)^{\otimes 6}. \quad (71)$$

This lemma can be shown directly by studying the Lie algebra generated by SWAPs on 11 qudits, or, as we do in Appendix D, by applying the seminal result of Marin [12], which characterizes the Lie algebra generated by transpositions (SWAPs) in terms of its simple factors.⁸

Next, we apply this lemma to a system containing 3 qudits plus 8 ancilla qudits, and show how we can realize unitaries in the form of Eq. (67). More precisely, we show that Hamiltonian Π_λ^{123} for all $\lambda \in \Lambda_3$ can be realized with 8 ancilla qudits.

Since Π_λ^{123} is not centerless, first we construct a centerless Hamiltonian, namely we consider $H = \Pi_\lambda^{123} - \Pi_\lambda^{456}$, where we have suppressed tensor product with the identity operators on the rest of qudits. This Hamiltonian is clearly centerless. In fact, any operator of the form $A - P_\sigma A P_\sigma^{-1}$ is centerless, because

$$\text{Tr}(\Pi(A - P_\sigma A P_\sigma^{-1})) = \text{Tr}(\Pi(A - A)) = 0, \quad (72)$$

where in the last step we used the cyclic property of the trace and Π is an arbitrary element in the center. Therefore, according to the above lemma, there exists an $SU(d)$ -invariant Hamiltonian \tilde{H} , such that Eq. (69) holds for all $t \in \mathbb{R}$ and all $|\psi\rangle \in (\mathbb{C}^d)^{\otimes 3}$.

Now for state $|\eta\rangle$ in Eq. (70) the reduced state of qudits 456 is restricted to the totally anti-symmetric subspace, corresponding to $\lambda = \square$, which means

$$\Pi_\lambda^{456}|\eta\rangle = 0 \quad : \quad \lambda = \square\square\square, \square\square, \quad (73a)$$

$$\Pi_\lambda^{456}|\eta\rangle = |\eta\rangle \quad : \quad \lambda = \square, \quad (73b)$$

where we have suppressed the tensor product with the identity operator on the rest of the qudits. Therefore, for Hamiltonian $H = \Pi_\lambda^{123} - \Pi_\lambda^{456}$, Eq. (69) becomes

$$e^{it\tilde{H}}(|\psi\rangle \otimes |\eta\rangle) = \begin{cases} (e^{it\Pi_\lambda^{123}}|\psi\rangle) \otimes |\eta\rangle & \lambda = \square\square\square, \square\square \\ (e^{it(\Pi_\lambda^{123} - \mathbb{I})}|\psi\rangle) \otimes |\eta\rangle & \lambda = \square \end{cases} \quad (74)$$

Note that in the second case, we get an extra global phase, which is not physically relevant (e.g., it can be removed by shifting \tilde{H} with a multiple of the identity operator).

We conclude that for all $\lambda \in \Lambda_3$ we can implement the Hamiltonian Π_λ on 3 qudits, using 8 qudits as ancilla and 2-qudit $SU(d)$ -invariant gates. Since the state of ancilla remains unchanged we can reuse it to implement all unitaries in the form of Eq. (67).

⁸ Note that this is the only result in the current paper depending on Marin's results in [12].

Now consider the special case of $d = 3$. Although in this case Eq. (73) does not hold, because both states $(|0\rangle \wedge |1\rangle) \otimes |0\rangle$ and $(|0\rangle \wedge |1\rangle) \otimes |1\rangle$ live in $\mathcal{Q}_{\mathbb{F}} \otimes \mathcal{M}_{\mathbb{F}}$, we have

$$\Pi_{\lambda}^{456} |\eta'\rangle = 0 \quad : \quad \lambda = \square\square\square, \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad (75a)$$

$$\Pi_{\lambda}^{456} |\eta'\rangle = |\eta'\rangle \quad : \quad \lambda = \begin{array}{|c|} \hline \square \\ \hline \end{array}. \quad (75b)$$

Therefore, in this case for Hamiltonian $H = \Pi_{\lambda}^{123} - \Pi_{\lambda}^{456}$ Eq. (69) simplifies to

$$e^{it\tilde{H}}(|\psi\rangle \otimes |\eta'\rangle) = \begin{cases} (e^{it(\Pi_{\lambda}^{123})} |\psi\rangle) \otimes |\eta'\rangle & \lambda = \square\square\square, \begin{array}{|c|} \hline \square \\ \hline \end{array} \\ (e^{it(\Pi_{\lambda}^{123}-1)} |\psi\rangle) \otimes |\eta'\rangle & \lambda = \begin{array}{|c|} \hline \square \\ \hline \end{array}, \end{cases} \quad (76)$$

and therefore the same results can be established with 6 qudit state $|\eta'\rangle$.

VIII. STATISTICAL PROPERTIES OF RANDOM $SU(d)$ -INVARIANT CIRCUITS: t -DESIGNS

In this section, we briefly discuss the statistical properties of random 3-qudit circuits and explain a corollary of semi-universality of 3-qudit gates.

First, recall that the no-go theorem of [5] puts strong constraints on the statistical properties of random symmetric circuits for general symmetries. According to the general results of [5], for any symmetry group G , the group of unitaries generated by k -local G -invariant unitaries is a compact connected Lie group, and therefore it has a unique notion of invariant (Haar) measure. Furthermore, when the symmetry group is a continuous group, such as $SU(d)$, [5] shows that for any fixed k , the dimension of this compact Lie group is strictly less than the compact Lie group of all G -invariant unitaries on n qudits. Clearly, the uniform distribution over a compact manifold cannot be fully mimicked by distributions restricted to compact submanifolds with lower dimensions. This can be more formally stated in terms of moments of the distributions.

Corollary 4 (corollary of [5]). *Consider the uniform distribution over the compact group \mathcal{V}_k^G of n -qudit unitaries realized with k -qudit G -invariant gates. Then, all moments of this distribution are equal to the corresponding moments of the uniform distribution over the group of all G -invariant unitaries, such that*

$$\mathbb{E}_{V \in \mathcal{V}^G} [V^{\otimes t} \otimes V^{*\otimes t}] = \mathbb{E}_{V \in \mathcal{V}_k^G} [V^{\otimes t} \otimes V^{*\otimes t}], \quad (77)$$

for all integer t , only if $|\text{Irreps}_G(k)| = |\text{Irreps}_G(n)|$, where $|\text{Irreps}_G(k)|$ is the number of inequivalent irreps of group G appearing on k qudits, i.e., in the representation $g \mapsto u(g)^{\otimes k} : g \in G$. Furthermore, assuming semi-universality holds for \mathcal{V}_k^G , and the group G is connected, then this condition becomes necessary and sufficient.

In the case of continuous symmetries such as $SU(d)$, there is no finite k such that $|\text{Irreps}_G(k)| = |\text{Irreps}_G(n)|$ for all $n > k$. Therefore, for any fixed k , the above equation cannot hold for arbitrarily large t and n .

For instance, in the case of $SU(d)$ symmetry discussed in this paper, the strict monotonicity of $|\Lambda_{n,d}|$ in Lemma 5, implies that for $d \geq 3$, unless $k = n$, certain moments of the uniform distribution over $\mathcal{V}_k^{(n)}$ will be different from the corresponding moment for the Haar distribution over the group of all $SU(d)$ -invariant unitaries.

However, it is still possible that some moments of these distributions match. This can be formulated based on the notion of t -designs [20–23]. We say the Haar distribution over the group $\mathcal{V}_k^{(n)}$ is a t -design for the Haar distribution over the group of all $SU(d)$ -invariant unitaries, if Eq. (77) holds.

Our previous result in [6] reveals that in the case of circuits with 2-qudit $SU(d)$ -invariant gates, for $d \geq 3$ this equation holds for $t = 1$, but not $t = 2$, which means the distribution generated by such random circuits converges to a 1-design but not a 2-design for the Haar distribution. Indeed, as it was noted in [6], the results of [24, 25] imply that whenever semi-universality does not hold then the distribution generated by the random circuits cannot be a 2-design for the Haar distribution over all symmetric unitaries. More recently, Ref. [26] shows that the distribution generated by circuits with 4-qudit $SU(d)$ -invariant gates is a t -design with t quadratic in n .

In this paper, we find that a similar quadratic scaling can be achieved with 3-qudit gates in the case of $SU(d)$ with $d \geq 3$, and 2-qudit gates in the case of $SU(2)$.

Proposition 4 (Quadratic scaling of t -designs with 3- or 2-qudit gates). *For systems with $n \geq 9$ qudits with $d < n - 1$, the Haar distribution over the group generated by 3-qudit $SU(d)$ -invariant gates, $\mathcal{V}_3^{(n)}$ is an exact t -design for the Haar distribution over the group of all $SU(d)$ -invariant unitaries $\mathcal{V}^{(n)}$ with $t < \frac{1}{2}n(n-3)$. In the case of $d = 2$, i.e., qubits with $SU(2)$ symmetry, $\mathcal{V}_3^{(n)} = \mathcal{V}_2^{(n)}$, which means the same quadratic scaling can be obtained with 2-qudit gates.*

As we explain in Appendix E, this proposition follows from the semi-universality of 3-qudit gates, together with the following fact

$$\pi_{\mu_0, \mu_1}(\mathcal{V}_3^{(n)}) = \pi_{\mu_0, \mu_1}(\mathcal{V}^{(n)}) \cong U(\mathcal{M}_{\mu_0}) \times U(\mathcal{M}_{\mu_1}), \quad (78)$$

where μ_0 and μ_1 are, respectively the symmetric and the $(n-1)$ -dimensional standard irrep of \mathbb{S}_n . In words, this means that when restricted to these two sectors, the group $\mathcal{V}_3^{(n)}$ and $\mathcal{V}^{(n)}$ are identical, which in turn implies that the design properties of $\mathcal{V}_3^{(n)}$ is determined by the smallest dimension of \mathcal{M}_{λ} , for $\lambda \in \Lambda_{n,d}$ other than μ_0, μ_1 . This dimension is equal to $n(n-3)/2$, when $n \geq 9$ and $d < n - 1$ [27]. We also use the standard techniques in the context of t -designs [20–23], which are used previously, e.g., in Ref. [26] in the context of $SU(d)$ -invariant circuits, and in [28] in the context of $U(1)$ -invariant circuits.

IX. FROM SUBSYSTEM UNIVERSALITY TO SEMI-UNIVERSALITY

In this section, we introduce a characterization of subsystem universality and various other techniques that are generally useful for understanding semi-universality (and its failure) for subgroups of G -invariant unitaries, for general symmetries and representations. We start by presenting a generalization of Lemma 1, namely Lemma 8, which is applicable even when condition **B** of Lemma 1, i.e., pairwise independence, does not hold. Then, we show how both lemmas follow from a combination of Goursat's and Serre's lemmas (Lemma 9 and Lemma 10). The ideas and techniques discussed in this section and in Appendix B are more broadly useful in the context of quantum computing and control theory.

A. Subdirect products

Before going through this section, we recall the concept of subdirect products, which is useful for understanding semi-universality and also appears in the statements of Goursat's and Serre's lemmas.

Definition 1 (Subdirect product). A *subdirect product* of the groups G_1, \dots, G_r is a subgroup of the direct product, $H \subseteq G_1 \times \dots \times G_r$, such that $\pi_i(H) = G_i$ for each factor, $i = 1, \dots, r$, where $\pi_i : G_1 \times \dots \times G_r \rightarrow G_i$ is the projection homomorphism.

It is useful to consider different subdirect products as exhibiting different types of *correlations*. In particular, if for any pair i and j of groups G_i and G_j , the projection of H to that pair is not surjective, i.e., $\pi_{ij}(H) \neq G_i \times G_j$, we say they are *correlated* in H , and otherwise, we say they are *independent* in H . This can be generalized to more factors, and in general, the correlations can be quite complicated (see, e.g., the example in Sec. II B of a subdirect product of three U(1) subgroups which are pairwise independent but still exhibit a tripartite correlation).

As we explain below, the notion of subdirect products appears naturally in the context of G -invariant unitaries.

B. Subsystem universality (condition A)

Consider any subgroup $\mathcal{T} \subset \mathcal{V}^G$ of G -invariant unitaries satisfying condition **A** in Lemma 1, namely subsystem universality, which means that for any irrep $\lambda \in \Lambda$ the projection of \mathcal{T} to \mathcal{M}_λ contains $\text{SU}(\mathcal{M}_\lambda)$, i.e.

$$\pi_\lambda(\mathcal{T}) = \{\pi_\lambda(V) : V \in \mathcal{T}\} \supseteq \text{SU}(\mathcal{M}_\lambda). \quad (79)$$

Then under this assumption, the commutator subgroup of \mathcal{T} , denoted as $[\mathcal{T}, \mathcal{T}]$, is a subdirect product of the groups $\text{SU}(\mathcal{M}_\lambda) : \lambda \in \Lambda$. Furthermore, semi-universality

means that this subdirect product is indeed the Cartesian product $\mathcal{SV}^G \cong \prod_{\lambda \in \Lambda} \text{SU}(\mathcal{M}_\lambda)$.

It is also worth noting that subsystem universality implies that any unitary $V \in \mathcal{T}$ can be decomposed as the product of an element of $[\mathcal{T}, \mathcal{T}]$ and a unitary in the subgroup of relative phases

$$\mathcal{P} = \left\{ \sum_{\lambda \in \Lambda} e^{i\theta_\lambda} \Pi_\lambda : \theta_\lambda \in [0, 2\pi) \right\}, \quad (80)$$

which is the center of \mathcal{V}^G . More precisely,

Lemma 7. *A subgroup of G -invariant unitaries $\mathcal{T} \subset \mathcal{V}^G$ contains \mathcal{SV}^G if, and only if, its commutator subgroup $[\mathcal{T}, \mathcal{T}] = \mathcal{SV}^G$. More generally, if \mathcal{T} satisfies the subsystem universality condition in Eq. (79) for all $\lambda \in \Lambda$, then any element of \mathcal{T} can be decomposed as a product of an element of $[\mathcal{T}, \mathcal{T}]$ and a unitary in the group of relative phases \mathcal{P} . That is, $\mathcal{T} \subseteq \mathcal{P}[\mathcal{T}, \mathcal{T}]$.*

Proof. Note that $\mathcal{V}^G = \mathcal{SV}^G \mathcal{P}$. If $\mathcal{SV}^G \subseteq \mathcal{T}$ then also $\mathcal{V}^G = \mathcal{P}\mathcal{T}$. Taking the commutator subgroup, since \mathcal{P} is the center, i.e. commutes with everything else, it follows that $\mathcal{SV}^G = [\mathcal{T}, \mathcal{T}]$. The proof of the second part is presented in Sec. IX E, by applying Lemma 8. \square

Therefore, in the following, we often consider the properties of the commutator subgroup $[\mathcal{T}, \mathcal{T}]$ and, assuming subsystem universality holds, interpret it as a subdirect product.

C. A characterization of subsystem universality

Here, we present a useful characterization of subgroups of G -invariant unitaries \mathcal{V}^G that satisfy the subsystem-universality condition in all charge sectors, i.e., condition **A** of Lemma 1, but do not necessarily satisfy the condition **B**, pairwise independence (According to the numbering of constraints in Sec. II B, this means that except type **I**, all constraints are of type **IV**, i.e., correlations between different sectors).

Lemma 8 (Extension of Lemma 1). *Consider a subgroup $\mathcal{T} \subseteq \mathcal{V}^G$ of G -invariant unitaries satisfying condition **A** in Lemma 1. That is, for any irrep $\lambda \in \Lambda$ the projection of \mathcal{T} to \mathcal{M}_λ contains $\text{SU}(\mathcal{M}_\lambda)$, as stated in Eq. (79). Then, the set of irreps Λ is partitioned into non-overlapping subsets $\Delta_1, \dots, \Delta_s$, such that for all pairs λ, λ' that belong to the same subset Δ_r , $\dim(\mathcal{M}_\lambda) = \dim(\mathcal{M}_{\lambda'}) =: m_r$, and that the commutator subgroup of \mathcal{T} is isomorphic to*

$$[\mathcal{T}, \mathcal{T}] \cong \prod_{r=1}^s \text{SU}(m_r). \quad (81)$$

Furthermore, for any pair of irreps $\lambda, \lambda' \in \Delta_r$, and fixed orthonormal bases for \mathcal{M}_λ and $\mathcal{M}_{\lambda'}$, there exists an $m_r \times$

m_r unitary matrix $W \in \text{SU}(m_r)$, such that for all $V \in [\mathcal{T}, \mathcal{T}]$ one of the followings holds

$$[\pi_{\lambda'}(V)] = W[\pi_{\lambda}(V)]W^\dagger, \text{ or} \quad (82a)$$

$$[\pi_{\lambda'}(V)] = W[\pi_{\lambda}(V)]^*W^\dagger, \quad (82b)$$

where $[\pi_{\lambda}(V)]^*$ is the complex conjugate of $\pi_{\lambda}(V)$ relative to this basis.

In words, this lemma means that for charge sectors that belong to the same part Δ , the unitary realized in one sector uniquely determines the unitaries realized in all the other sectors. Namely, they are all equal, up to a change of basis, a possible complex conjugation, and relative phases between sectors (i.e. an element of the group \mathcal{P} defined in Eq. (80)). Note that, as stated in Lemma 7, when condition **A** holds for all sectors, the commutator subgroup $[\mathcal{T}, \mathcal{T}]$ determines \mathcal{T} , up to additional freedoms on the relative phases between different charge sectors. Therefore, this lemma characterizes group \mathcal{T} , up to this freedom.

We next mention an example that has been already discussed in detail in Sec. IV B.

Example: Revisiting group $\mathcal{V}_2^{(4)}$

In the special case of $G = \text{SU}(d)$ symmetry and $n = 4$ qudits, in Sec. IV B we characterized the group $\mathcal{V}_2^{(4)}$ generated by 2-qudit $\text{SU}(d)$ -invariant gates. In the context of the above lemma, $\mathcal{V}_2^{(4)}$ corresponds to the group \mathcal{T} , which is a subgroup of $\mathcal{V}^{(4)}$ the group of all $\text{SU}(d)$ -invariant unitaries. Then, in the language of the above lemma, we found that relative to this group, the set of irreps Λ_4 is partitioned into 3 sets, namely

$$\Delta_1 = \left\{ \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right\}, \Delta_2 = \left\{ \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \end{array} \right\}, \Delta_3 = \left\{ \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \end{array} \right\}, \quad (83)$$

with corresponding dimensions $m_1 = 2$, $m_2 = 3$, and $m_3 = 1$. Furthermore, we showed while the commutator subgroup of all $\text{SU}(d)$ -invariant unitaries, denoted as $\mathcal{SV}^{(4)}$, is isomorphic to $\text{SU}(2) \times \text{SU}(3) \times \text{SU}(3)$, the commutator subgroup of $\mathcal{V}_2^{(4)}$, denoted as $\mathcal{SV}_2^{(4)}$ is isomorphic to

$$\text{SU}(m_1) \times \text{SU}(m_2) \times \text{SU}(m_3) = \text{SU}(2) \times \text{SU}(3). \quad (84)$$

Moreover, in Eq. (41) we saw that for any $V \in \mathcal{V}_2^{(4)}$, the unitaries realized in $\mathcal{M}_{\square\square}$ and \mathcal{M}_{\square} are equal, up to a complex conjugation and a change of basis, as described by Lemma 8.

Next, we explain the tools that are needed to prove Lemmas 1 and 8.

D. Two useful lemmas

1. Goursat's lemma: A characterization of pairwise correlations

Goursat's lemma [29] classifies all possible subdirect products $H \subseteq G_1 \times G_2$ of arbitrary groups G_1 and G_2 . We postpone the general form of this lemma, which applies to arbitrary groups G_1 and G_2 to Lemma 13 in Appendix B 3. Instead, here we discuss its specialized form applied to special unitary groups, which are the most relevant groups for applications in quantum computing and control.

Lemma 9 (Goursat's lemma for special unitary groups). *Let $l, l' \geq 2$ and let $H \subseteq \text{SU}(l) \times \text{SU}(l')$ be a subdirect product. There are two possibilities:*

$$(i) \ H = \text{SU}(l) \times \text{SU}(l').$$

$$(ii) \ l = l' \text{ and } H \cong \text{SU}(l) \times \mathbb{Z}_q, \text{ where } q \text{ divides } l.$$

Furthermore, in the second case, there exists an isomorphism $\Phi : \text{SU}(l) \rightarrow \text{SU}(l)$ such that

$$H = \{(U, e^{i\theta} \Phi(U)) : U \in \text{SU}(l) \text{ and } e^{i\theta} \in \mathbb{Z}_q \subset \text{SU}(l)\}. \quad (85)$$

Remark 3. When interpreted as explicit $l \times l$ determinant-one unitary matrices, then the above isomorphism in case (ii) must be in one of the following forms:

$$\Phi(U) = WUW^\dagger \quad (86a)$$

$$\Phi(U) = WU^*W^\dagger, \quad (86b)$$

where W is a unitary matrix and U^* is the component-wise complex conjugate of matrix U . For $l > 2$, these two possibilities correspond to two inequivalent representations of the group $\text{SU}(l)$,⁹ and these are the only two non-trivial representations of dimension l [17].¹⁰

Note that, in the second case of Lemma 9, upon taking the commutator subgroup, the discrete phase vanishes, i.e. $[H, H] \cong \text{SU}(l)$. In fact, this holds even if we consider H as a subgroup of $\text{U}(l) \times \text{U}(l)$. Goursat's lemma and the following corollary are proved in Appendix B 3.

Corollary 5. *Suppose that $H \subseteq \text{U}(l) \times \text{U}(l')$ is a subgroup with $[H, H] \subseteq \text{SU}(l) \times \text{SU}(l')$ a subdirect product. Then either $[H, H] \cong \text{SU}(l) \times \text{SU}(l)$, or $l = l'$ and there is an isomorphism $\Phi : \text{SU}(l) \rightarrow \text{SU}(l)$ such that $[H, H] = \{(U, \Phi(U)) : U \in \text{SU}(l)\} \cong \text{SU}(l)$. Furthermore, in the second case every element of H is of the form $(e^{i\theta} U, e^{i\phi} \Phi(U))$, for some $\theta, \phi \in [0, 2\pi)$.*

⁹ For $l = 2$, these are not distinct possibilities since the defining representation is self-dual: with Pauli Y defined in the chosen basis, we have $U^* = YUY^\dagger$ for all $U \in \text{SU}(2)$.

¹⁰ Note that this fact follows from ‘‘automatic continuity’’. See Appendix B 4 for further discussion.

2. *Serre's lemma: Pairwise independence implies full independence*

Serre's lemma characterizes when the subdirect product of perfect groups is actually a direct product, and roughly states that, if perfect groups are pairwise independent, then they are fully independent.¹¹ Recall that a group G is called *perfect* if it is equal to its commutator subgroup, i.e., $[G, G] = G$. For example, the unitary group $U(d)$ is not perfect, but the special unitary group $SU(d)$ is perfect (in fact, all finite-dimensional semi-simple Lie groups are perfect).

Lemma 10 (Serre's lemma [30]). *Let $H \subseteq G_1 \times \dots \times G_r$ be a subgroup such that $\pi_{ij}(H) = G_i \times G_j$ for all pairs $1 \leq i < j \leq r$, where $\pi_{ij} : G_1 \times \dots \times G_r \rightarrow G_i \times G_j$ is the projection homomorphism. If each G_i is perfect, then $H = G_1 \times \dots \times G_r$.*

We present the proof of this lemma in Appendix B 5.

E. Characterizations of semi-universality and subsystem universality: Proofs of Lemmas 1 and 8

Finally, we show how Goursat's and Serre's lemmas can be applied to the case of G -invariant unitaries, and prove Lemma 1 and Lemma 8. We start by Lemma 1.

Given that $\pi_\Lambda(\mathcal{SV}^G) = \prod_\lambda SU(\mathcal{M}_\lambda)$, it is clear that if $\mathcal{T} \supseteq \mathcal{SV}^G$, then \mathcal{T} satisfies both conditions **A** and **B** (see Sec. II B). Therefore, to prove the lemma, it suffices to prove the converse direction.

First, we focus only on a pair of irreps $\lambda_1, \lambda_2 \in \Lambda$, and discuss the implications of conditions **A** and **B** for this pair. Suppose the subgroup $\mathcal{T} \subset \mathcal{V}^G$ satisfies condition **A** for $\lambda_1, \lambda_2 \in \Lambda$. That is, Eq. (79) holds for $\lambda = \lambda_1, \lambda_2$. Then, case (i) of Goursat's lemma immediately implies that if $\dim \mathcal{M}_{\lambda_1} \neq \dim \mathcal{M}_{\lambda_2}$, the joint projection of $[\mathcal{T}, \mathcal{T}]$ to λ_1 and λ_2 , denoted as $\pi_{\lambda_1 \lambda_2}([\mathcal{T}, \mathcal{T}])$, is isomorphic to the Cartesian product $SU(\mathcal{M}_{\lambda_1}) \times SU(\mathcal{M}_{\lambda_2})$. However, if $m := \dim \mathcal{M}_{\lambda_1} = \dim \mathcal{M}_{\lambda_2} \geq 2$, then there exists a second possibility, namely case (ii) of Goursat's lemma. In particular, according to Remark 3 and Corollary 5, for any fixed orthonormal bases on \mathcal{M}_{λ_1} and \mathcal{M}_{λ_2} , there exists an $m \times m$ unitary matrix W such that, for all $V \in [\mathcal{T}, \mathcal{T}]$, one of the following holds

$$[\pi_{\lambda_2}(V)] = W[\pi_{\lambda_1}(V)]W^\dagger \quad (87a)$$

$$[\pi_{\lambda_2}(V)] = W[\pi_{\lambda_1}(V)]^*W^\dagger. \quad (87b)$$

Furthermore, the second part of Corollary 5 implies that, for all $V \in \mathcal{T}$,

$$[\pi_{\lambda_2}(V)] = e^{i\phi} W[\pi_{\lambda_1}(V)]W^\dagger \quad (88a)$$

$$[\pi_{\lambda_2}(V)] = e^{i\phi} W[\pi_{\lambda_1}(V)]^*W^\dagger, \quad (88b)$$

for some phase depending on V .

Now to prove Lemma 1, we note that in both cases in Eq. (88), $|\text{Tr}(\pi_{\lambda_1}(V))| = |\text{Tr}(\pi_{\lambda_2}(V))|$. Therefore, if condition **B** of Lemma 1 holds, i.e., there exists a unitary $V \in \mathcal{T}$ such that $|\text{Tr}(\pi_{\lambda_1}(V))| \neq |\text{Tr}(\pi_{\lambda_2}(V))|$ then the case (ii) of Goursat's lemma is ruled out, which means

$$SU(\mathcal{M}_{\lambda_1}) \times SU(\mathcal{M}_{\lambda_2}) = \pi_{\lambda_1 \lambda_2}([\mathcal{T}, \mathcal{T}]) \subseteq \pi_{\lambda_1 \lambda_2}(\mathcal{T}). \quad (89)$$

Similarly, if condition **B'** holds, i.e., there exists $V \in \mathcal{T}$, such that $\pi_{\lambda_1}(V) = \mathbb{I}_{\mathcal{M}_{\lambda_1}}$ and $\pi_{\lambda_2}(V) \neq e^{i\alpha} \mathbb{I}_{\mathcal{M}_{\lambda_2}}$ for any phase α , then, again Eq. (88) cannot hold, which implies the same result.

This proves Lemma 1 in the special case where Λ contains only two irreps. Finally, we note that because the special unitary group $SU(\mathcal{M}_\lambda)$ is perfect, and Eq. (89) applies to all pairs $\lambda_1, \lambda_2 \in \Lambda$, applying Serre's lemma to the subdirect product $\pi_\Lambda([\mathcal{T}, \mathcal{T}]) \subseteq \prod_{\lambda \in \Lambda} SU(\mathcal{M}_\lambda)$, we conclude that $\prod_{\lambda \in \Lambda} SU(\mathcal{M}_\lambda) \cong \mathcal{SV}^G \subset \mathcal{T}$, which completes the proof of Lemma 1.

Next, we prove Lemma 8, which does not assume condition **B**. Then, in this case, the case (ii) in Goursat's lemma, i.e., correlations between different sectors, cannot be ruled out. This means that it is possible that for any $V \in \mathcal{T}$ its projection to two sectors λ_1 and λ_2 , namely $\pi_{\lambda_1}(V)$ and $\pi_{\lambda_2}(V)$, are related via one of the two possibilities in Eq. (88) where the unitary realized in λ_1 uniquely determines the unitary in λ_2 , up to a possible phase.

As we explain in the following, based on the two possibilities in Goursat's lemma, i.e., case (i) and case (ii), we can partition all irreps in Λ into equivalency classes $\Delta_1, \dots, \Delta_s$. Namely, for any $\lambda_1, \lambda_2 \in \Lambda$ if case (i) of Goursat's lemma applies to $\pi_{\lambda_1 \lambda_2}([\mathcal{T}, \mathcal{T}])$, then they are in different parts and if case (ii) applies they are in the same part. Note that Eq. (87) guarantees that this rule defines equivalency classes and can be applied consistently. In particular, if for $\lambda_1, \lambda_2, \lambda_3 \in \Lambda$, the case (ii) applies to the pair λ_1, λ_2 and also to the pair λ_2, λ_3 , then it also applies to the pair λ_1, λ_3 , i.e., one of the two possibilities in Eq. (87) is applicable to $\pi_{\lambda_1}(V)$ and $\pi_{\lambda_3}(V)$, and therefore they are in the same part as well. Also, note that the case (ii) of Goursat's lemma applies to λ_1 and λ_2 only if $\dim \mathcal{M}_{\lambda_1} = \dim \mathcal{M}_{\lambda_2}$, which means for all irreps in the same part this dimension is identical. We label these parts as $\Delta_1, \dots, \Delta_s$, and define the corresponding dimensions as $m_r = \dim \mathcal{M}_\lambda$ for any $\lambda \in \Delta_r$.

Next, we pick a representative λ from each part Δ_r , denoted as λ_r , which together define $\Lambda_{\text{rep}} = \{\lambda_r : r = 1, \dots, s\}$. Then, we apply Serre's lemma to the subdirect product

$$\prod_{\lambda \in \Lambda_{\text{rep}}} \pi_\lambda([\mathcal{T}, \mathcal{T}]) \subseteq \prod_{r=1}^s SU(m_r). \quad (90)$$

¹¹ Serre's lemma for finite groups was first stated in a paper by Kenneth Ribet [30], where the proof is attributed to Jean-Pierre Serre. Later Terence Tao restated the lemma in a slightly different way and named it Serre's lemma [31]. Our statement and proof follow Ribet [30].

By the above construction, for any pair of parts Δ_r and $\Delta_{r'}$ their corresponding representatives λ_r and $\lambda_{r'}$ satisfy case (i) of Goursat's lemma, which means $\pi_{\lambda_r, \lambda_{r'}}([\mathcal{T}, \mathcal{T}]) = \text{SU}(m_r) \times \text{SU}(m_{r'})$. Therefore, by applying Serre's lemma, we conclude that Eq. (90) holds as equality, i.e., the subdirect product is indeed a Cartesian product. This completes the proof of Lemma 8.

Now we finish the proof of Lemma 7. Let $V \in \mathcal{T}$ be an arbitrary element. We show that there exists phases $e^{i\phi_\lambda}$: $\lambda \in \Lambda$ and $V' \in [\mathcal{T}, \mathcal{T}]$ such that $V = V' \sum_{\lambda \in \Lambda} e^{i\phi_\lambda} \Pi_\lambda$.

By Eq. (90) there is an element $U \in [\mathcal{T}, \mathcal{T}]$ such that, for all $\lambda_r \in \Lambda_{\text{rep}}$, $\pi_{\lambda_r}(UV) = e^{i\phi_r} \Pi_{\lambda_r}$ for some phase. By construction, every other $\lambda' \in \Lambda \setminus \Lambda_{\text{rep}}$ is contained in some equivalence class of irreps, say $\lambda' \in \Delta_r$. According to Eq. (88), if $\pi_{\lambda_r}(UV)$ is proportional to the identity, $\pi_{\lambda'}(UV)$ must be as well. But this holds for all $\lambda' \in \Lambda$, hence $UV \in \mathcal{P}$ is in the subgroup of relative phases. In other words, $V = U^{-1}(UV)$ is of the desired form.

X. CONCLUSION

A series of recent works has shown that in the presence of symmetries, the locality of gates can severely restrict the set of realizable unitaries [5–8, 32]. Interestingly, the type of restrictions significantly depends on the properties of the symmetry. In particular, in the case of Abelian symmetries, recent work [8] has revealed a simple characterization of the group of all realizable unitaries with k -qudit gates, showing that some restrictions that appear in the case of non-Abelian symmetries, such as $\text{SU}(d)$ symmetry, cannot appear in the case of Abelian symmetries. In general, understanding the restrictions in the case of non-Abelian symmetries is significantly more complicated. Prior to the present work, a general formal framework or techniques for understanding such circuits did not exist.

In this work, we developed novel tools and a new framework for understanding semi-universality and universality in symmetric quantum circuits, which are particularly useful in the case of non-Abelian symmetries where the simpler characterization of [8] is not applicable. We anticipate that, beyond the theory of symmetric quantum circuits, this framework and the new tools, such as our characterization of semi-universality in Lemma 1, characterization of subsystem universality in Lemma 8, and our Lemma 2 on extending controllability, will be more broadly useful in Quantum Control theory, Quantum Many-body Physics, and Quantum Thermodynamics. For instance, these new techniques could be useful for understanding the effect of non-Abelian conserved charges on the thermalization of quantum systems [33, 34].

As an example of applications, we applied these new tools to settle a fundamental question in the context of $\text{SU}(d)$ -invariant circuits with $d \geq 3$. Namely, we showed the semi-universality of 3-qudit gates. We also discussed

two corollaries of this result. Firstly, by studying 3-qudit circuits, we found a significantly simpler proof of the universality of 2-qudit gates, which was recently shown in a PhD thesis [11], based on the advanced results in Mathematical literature, namely, Marin's characterization of the Lie algebra generated by transpositions. Secondly, using this result, we showed that the distribution generated by random circuits with 3-qudit $\text{SU}(d)$ -invariant gates is a t -design, with t growing quadratically with the number of qudits. Such quadratic scaling has been recently shown in [12], albeit using 4-qudit gates (We also emphasize that circuits with 2-qudit gates are not 2-design [6]).

$\text{SU}(d)$ -invariant circuits, and more generally symmetric quantum circuits, have various applications in the context of quantum computation and control. In particular, they are useful for protecting information, e.g., via decoherence-free subspaces and noiseless subsystems [35–40]. More broadly, the notion of charge-conserving unitaries appears in various areas of quantum information science, and it is crucial to understand how such unitaries can be realized. This is particularly relevant in the context of quantum thermodynamics [41–48], the resource theory of asymmetry [49–51], and other related areas such as Quantum reference frames [52, 53], and covariant error-correcting codes [54–56]. Furthermore, such circuits have been found useful in other contexts including variational quantum machine learning [57–60], and variational quantum eigensolvers for quantum chemistry [61–66]. Besides such applications, the framework of symmetric quantum circuits has now become a standard tool in the area of many-body Physics, for modeling various physical phenomena, from quantum thermalization and quantum chaos [67] to symmetry-protected topological order [68, 69]. Therefore, we anticipate that the framework developed in this paper find applications beyond quantum computation.

ACKNOWLEDGEMENTS

This work is supported by a collaboration between the US DOE and other Agencies. This material is based upon work supported by the U.S. Department of Energy, Office of Science, National Quantum Information Science Research Centers, Quantum Systems Accelerator. Additional support is acknowledged from Army Research Office (W911NF-21-1-0005), NSF Phy-2046195, and NSF QLCI grant OMA-2120757. H.L. was supported by the Quantum Science Center (QSC), a National Quantum Information Science Research Center of the U.S. Department of Energy (DOE) and by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics (NP) contract DE-AC52-06NA25396.

- [1] D. P. DiVincenzo, Two-bit gates are universal for quantum computation, *Physical Review A* **51**, 1015 (1995).
- [2] S. Lloyd, Almost any quantum logic gate is universal, *Physical Review Letters* **75**, 346 (1995).
- [3] D. E. Deutsch, A. Barenco, and A. Ekert, Universality in quantum computation, *Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences* **449**, 669 (1995).
- [4] J.-L. Brylinski and R. Brylinski, Universal quantum gates, *Mathematics of quantum computation* **79** (2002).
- [5] I. Marvian, Restrictions on realizable unitary operations imposed by symmetry and locality, *Nature Physics* **18**, 283–289 (2022).
- [6] I. Marvian, H. Liu, and A. Hulse, Qudit circuits with $su(d)$ symmetry: Locality imposes additional conservation laws (2022), [arXiv:2105.12877 \[quant-ph\]](https://arxiv.org/abs/2105.12877).
- [7] I. Marvian, H. Liu, and A. Hulse, Rotationally invariant circuits: Universality with the exchange interaction and two ancilla qubits, *Physical Review Letters* **132**, 10.1103/physrevlett.132.130201 (2024).
- [8] I. Marvian, Theory of quantum circuits with abelian symmetries (2024), [arXiv:2302.12466 \[quant-ph\]](https://arxiv.org/abs/2302.12466).
- [9] A. Okounkov and A. Vershik, A new approach to representation theory of symmetric groups, *Selecta Mathematica* **2**, 581 (2005).
- [10] H. Zheng, Z. Li, J. Liu, S. Strelchuk, and R. Kondor, Speeding up learning quantum states through group equivariant convolutional quantum ansätze, *PRX Quantum* **4**, 10.1103/prxquantum.4.020327 (2023).
- [11] J. R. van Meter, Universality of swap for qudits: a representation theory approach (2021), [arXiv:2103.12303 \[quant-ph\]](https://arxiv.org/abs/2103.12303).
- [12] I. Marin, L’algèbre de lie des transpositions, *Journal of Algebra* **310**, 742 (2007).
- [13] H. Liu, A. Hulse, and I. Marvian, Under preparation.
- [14] R. Goodman and N. R. Wallach, *Symmetry, representations, and invariants*, Vol. 255 (Springer, 2009).
- [15] A. W. Harrow, Applications of coherent classical communication and the schur transform to quantum information theory, [arXiv preprint quant-ph/0512255](https://arxiv.org/abs/0512255) (2005).
- [16] M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information*, Cambridge Series on Information and the Natural Sciences (Cambridge University Press, 2000).
- [17] W. Fulton and J. Harris, *Representation theory: a first course*, Vol. 129 (Springer Science & Business Media, 2013).
- [18] P. Erdős and J. Lehner, The distribution of the number of summands in the partitions of a positive integer, *Duke Mathematical Journal* **8**, 335 (1941).
- [19] J. Alfonsin, *The Diophantine Frobenius Problem*, Oxford Lecture Series in Mathematics and Its Applications (OUP Oxford, 2005).
- [20] A. W. Harrow and R. A. Low, Random quantum circuits are approximate 2-designs, *Communications in Mathematical Physics* **291**, 257 (2009).
- [21] J. Emerson, E. Livine, and S. Lloyd, Convergence conditions for random quantum circuits, *Physical Review A* **72**, 060302 (2005).
- [22] F. G. Brandao, A. W. Harrow, and M. Horodecki, Efficient quantum pseudorandomness, *Physical review letters* **116**, 170502 (2016).
- [23] F. G. Brandao, A. W. Harrow, and M. Horodecki, Local random quantum circuits are approximate polynomial-designs, *Communications in Mathematical Physics* **346**, 397 (2016).
- [24] R. Zeier and Z. Zimborás, On squares of representations of compact Lie algebras, *Journal of Mathematical Physics* **56**, 081702 (2015), https://pubs.aip.org/aip/jmp/article-pdf/doi/10.1063/1.4928410/15804949/081702_1_online.pdf.
- [25] Z. Zimborás, R. Zeier, T. Schulte-Herbrüggen, and D. Burgarth, Symmetry criteria for quantum simulability of effective interactions, *Physical Review A* **92**, 042309 (2015).
- [26] Z. Li, H. Zheng, J. Liu, L. Jiang, and Z.-W. Liu, Designs from Local Random Quantum Circuits with $SU(d)$ Symmetry, (2023), [arXiv:2309.08155 \[quant-ph\]](https://arxiv.org/abs/2309.08155).
- [27] R. Rasala, On the minimal degrees of characters of sn , *Journal of Algebra* **45**, 132 (1977).
- [28] S. N. Hearth, M. O. Flynn, A. Chandran, and C. R. Laumann, Unitary k -designs from random number-conserving quantum circuits, [arXiv preprint arXiv:2306.01035](https://arxiv.org/abs/2306.01035) (2023).
- [29] S. Lang, *Algebra*, Graduate Texts in Mathematics (Springer New York, 2005).
- [30] K. A. Ribet, On ℓ -adic representations attached to modular forms, *Invent. Math.* **28**, 245–275 (1975).
- [31] T. Tao, *Goursat and Furstenberg-Weiss type lemmas* (2021).
- [32] S. Kazi, M. Larocca, and M. Cerezo, On the universality of sn -equivariant k -body gates, *New Journal of Physics* **26**, 053030 (2024).
- [33] N. Y. Halpern, M. E. Beverland, and A. Kalev, Noncommuting conserved charges in quantum many-body thermalization, *Physical Review E* **101**, 042117 (2020).
- [34] S. Majidy, W. F. Braasch Jr, A. Lasek, T. Upadhyaya, A. Kalev, and N. Yunger Halpern, Noncommuting conserved charges in quantum thermodynamics and beyond, *Nature Reviews Physics* **5**, 689 (2023).
- [35] D. A. Lidar, I. L. Chuang, and K. B. Whaley, Decoherence-free subspaces for quantum computation, *Physical Review Letters* **81**, 2594 (1998).
- [36] D. Bacon, J. Kempe, D. A. Lidar, and K. B. Whaley, Universal fault-tolerant quantum computation on decoherence-free subspaces, *Phys. Rev. Lett.* **85**, 1758 (2000).
- [37] D. P. DiVincenzo, D. Bacon, J. Kempe, G. Burkard, and K. B. Whaley, Universal quantum computation with the exchange interaction, *nature* **408**, 339 (2000).
- [38] P. Zanardi and M. Rasetti, Noiseless quantum codes, *Phys. Rev. Lett.* **79**, 3306 (1997).
- [39] J. Kempe and K. B. Whaley, Exact gate sequences for universal quantum computation using the xy interaction alone, *Physical Review A* **65**, 052330 (2002).
- [40] D. J. Brod and A. M. Childs, The computational power of matchgates and the xy interaction on arbitrary graphs, [arXiv preprint arXiv:1308.1463](https://arxiv.org/abs/1308.1463) (2013).
- [41] D. Janzing, P. Wocjan, R. Zeier, R. Geiss, and T. Beth, Thermodynamic cost of reliability and low temperatures: tightening Landauer’s principle and the Second Law, *Int. J. Theor. Phys.* **39**, 2717 (2000).
- [42] M. Horodecki and J. Oppenheim, Fundamental limitations for quantum and nanoscale thermodynamics, *Nat.*

- Commun. **4**, 1 (2013).
- [43] F. G. Brandao, M. Horodecki, J. Oppenheim, J. M. Renes, and R. W. Spekkens, Resource theory of quantum states out of thermal equilibrium, *Physical review letters* **111**, 250404 (2013).
- [44] Y. Guryanova, S. Popescu, A. J. Short, R. Silva, and P. Skrzypczyk, Thermodynamics of quantum systems with multiple conserved quantities, *Nature communications* **7**, ncomms12049 (2016).
- [45] M. Lostaglio, K. Korzekwa, D. Jennings, and T. Rudolph, Quantum coherence, time-translation symmetry, and thermodynamics, *Physical Review X* **5**, 021001 (2015).
- [46] N. Y. Halpern, P. Faist, J. Oppenheim, and A. Winter, Microcanonical and resource-theoretic derivations of the thermal state of a quantum system with noncommuting charges, *Nature communications* **7**, 12051 (2016).
- [47] N. Y. Halpern and J. M. Renes, Beyond heat baths: Generalized resource theories for small-scale thermodynamics, *Physical Review E* **93**, 022126 (2016).
- [48] M. Lostaglio, D. Jennings, and T. Rudolph, Thermodynamic resource theories, non-commutativity and maximum entropy principles, *New Journal of Physics* **19**, 043008 (2017).
- [49] G. Gour and R. W. Spekkens, The resource theory of quantum reference frames: manipulations and monotones, *New Journal of Physics* **10**, 033023 (2008).
- [50] I. Marvian, *Symmetry, Asymmetry and Quantum Information*, *PhD thesis*, Ph.D. thesis, University of Waterloo, <https://uwspace.uwaterloo.ca/handle/10012/7088> (2012).
- [51] I. Marvian and R. W. Spekkens, The theory of manipulations of pure state asymmetry: I. basic tools, equivalence classes and single copy transformations, *New Journal of Physics* **15**, 033001 (2013).
- [52] S. D. Bartlett, T. Rudolph, and R. W. Spekkens, Reference frames, superselection rules, and quantum information, *Reviews of Modern Physics* **79**, 555 (2007).
- [53] I. Marvian and R. Mann, Building all time evolutions with rotationally invariant hamiltonians, *Physical Review A* **78**, 022304 (2008).
- [54] P. Faist, S. Nezami, V. V. Albert, G. Salton, F. Pastawski, P. Hayden, and J. Preskill, Continuous symmetries and approximate quantum error correction, *Physical Review X* **10**, 041018 (2020).
- [55] P. Hayden, S. Nezami, S. Popescu, and G. Salton, Error correction of quantum reference frame information, *PRX Quantum* **2**, 010326 (2021).
- [56] M. P. Woods and Á. M. Alhambra, Continuous groups of transversal gates for quantum error correcting codes from finite clock reference frames, *Quantum* **4**, 245 (2020).
- [57] J. J. Meyer, M. Mularski, E. Gil-Fuster, A. A. Mele, F. Arzani, A. Wilms, and J. Eisert, Exploiting symmetry in variational quantum machine learning, *PRX Quantum* **4**, 010328 (2023).
- [58] Q. T. Nguyen, L. Schatzki, P. Braccia, M. Ragone, P. J. Coles, F. Sauvage, M. Larocca, and M. Cerezo, Theory for equivariant quantum neural networks, arXiv preprint arXiv:2210.08566 (2022).
- [59] F. Sauvage, M. Larocca, P. J. Coles, and M. Cerezo, Building spatial symmetries into parameterized quantum circuits for faster training, arXiv preprint arXiv:2207.14413 (2022).
- [60] H. Zheng, Z. Li, J. Liu, S. Strelchuk, and R. Kondor, Speeding up learning quantum states through group equivariant convolutional quantum ansätze, *PRX Quantum* **4**, 020327 (2023).
- [61] G. S. Barron, B. T. Gard, O. J. Altman, N. J. Mayhall, E. Barnes, and S. E. Economou, Preserving symmetries for variational quantum eigensolvers in the presence of noise, *Physical Review Applied* **16**, 034003 (2021).
- [62] V. Shkolnikov, N. J. Mayhall, S. E. Economou, and E. Barnes, Avoiding symmetry roadblocks and minimizing the measurement overhead of adaptive variational quantum eigensolvers. arxiv, arXiv preprint arXiv:2109.05340 (2021).
- [63] B. T. Gard, L. Zhu, G. S. Barron, N. J. Mayhall, S. E. Economou, and E. Barnes, Efficient symmetry-preserving state preparation circuits for the variational quantum eigensolver algorithm, *npj Quantum Information* **6**, 10 (2020).
- [64] M. Streif, M. Leib, F. Wudarski, E. Rieffel, and Z. Wang, Quantum algorithms with local particle-number conservation: Noise effects and error correction, *Physical Review A* **103**, 042412 (2021).
- [65] Z. Wang, N. C. Rubin, J. M. Dominy, and E. G. Rieffel, X y mixers: Analytical and numerical results for the quantum alternating operator ansatz, *Physical Review A* **101**, 012320 (2020).
- [66] P. K. Barkoutsos, J. F. Gonthier, I. Sokolov, N. Moll, G. Salis, A. Fuhrer, M. Ganzhorn, D. J. Egger, M. Troyer, A. Mezzacapo, *et al.*, Quantum algorithms for electronic structure calculations: Particle-hole hamiltonian and optimized wave-function expansions, *Physical Review A* **98**, 022322 (2018).
- [67] V. Khemani, A. Vishwanath, and D. A. Huse, Operator spreading and the emergence of dissipative hydrodynamics under unitary evolution with conservation laws, *Physical Review X* **8**, 031057 (2018).
- [68] X. Chen, Z.-C. Gu, and X.-G. Wen, Local unitary transformation, long-range quantum entanglement, wave function renormalization, and topological order, *Physical review b* **82**, 155138 (2010).
- [69] X. Chen, Z.-C. Gu, and X.-G. Wen, Classification of gapped symmetric phases in one-dimensional spin systems, *Physical review b* **83**, 035107 (2011).
- [70] V. Jurdjevic and H. J. Sussmann, Control systems on lie groups, *Journal of Differential equations* **12**, 313 (1972).
- [71] K. Hofmann and S. Morris, *The Structure of Compact Groups: A Primer for the Student – A Handbook for the Expert*, De Gruyter Studies in Mathematics (De Gruyter, 2020).
- [72] H. Georgi, *Lie Algebras In Particle Physics: from Isospin To Unified Theories* (CRC Press, 2018).

APPENDIX

- **Appendix A 3-qudit gates**25
 In this section, the group $\mathcal{V}_2^{(3)}$ of 3-qudit $SU(d)$ -invariant unitaries generated by 2-qudit symmetric gates is characterized. It is also shown that generic 3-qudit $SU(d)$ -invariant gates are semiuniversal together with 2-qudit ones.
- **Appendix B Tools for semiuniversality**29
 In this section, the remaining details of the proof of semi-universality of 3-qudit $SU(d)$ -invariant gates are given. In particular, Lemma 2 is proven, as well as Goursat's and Serre's lemmas.
- **Appendix C For $d \geq 3$, universality requires n -qudit interactions**37
 In this section, it is proven that, when $d \geq 3$, full n -local control is required for universality.
- **Appendix D Semi-universality on $n = 3$ qudits using 8 ancilla qudits**38
 Here the result that semi-universality can be achieved with 2-local $SU(d)$ -invariant gates and 8 ancilla qudits is proven.
- **Appendix E t -designs: Proof of Proposition 4**39
 In this section it is proven that 3-qudit $SU(d)$ -invariant gates are a t -design for all symmetric gates, with $t \approx n^2/2$.
- **Appendix F Techniques for controllability**40
 A number of strategies useful for extending controllability are considered, including the idea of isolating Lie algebra elements to blocks.

Appendix A: 3-qudit gates

In this section, we consider 2-qudit and 3-qudit $SU(d)$ -invariant gates. In Appendix A 1 we characterize the 3-qudit gates that can be generated by 2-qudit ones, and we prove in Appendix A 2 that the ones which cannot be generated, together with 2-qudit gates are semiuniversal. Together with the aforementioned characterization, this implies that generic $SU(d)$ -invariant 3-qudit gates are semiuniversal (Proposition 3).

1. Which 3-qudit gates are realizable with 2-qudit gates: A characterization of $\mathcal{V}_2^{(3)}$

In this section, we characterize the 3-qudit $SU(d)$ -invariant gates that can be generated by 2-qudit ones, i.e. the elements of $\mathcal{V}_2^{(3)}$. In particular, because 2-qudit gates are semi-universal, i.e. $\mathcal{S}\mathcal{V}^{(3)} \subseteq \mathcal{V}_2^{(3)}$, the only elements that cannot be generated are central and therefore act as relative phases between charge sectors.

On

$$(\mathbb{C}^d)^{\otimes 3} = \mathcal{H}_{\square\square\square} \oplus \mathcal{H}_{\square\Box} \oplus \mathcal{H}_{\Box\square} \quad (\text{A1})$$

consider the Hermitian operators

$$\begin{aligned} B_1 &= \mathbb{I} \\ B_2 &= \mathbf{P}(12) + \mathbf{P}(13) + \mathbf{P}(23) \\ B_3 &= \mathbf{P}(123) + \mathbf{P}(132). \end{aligned} \quad (\text{A2})$$

These are both $SU(d)$ - and permutationally-invariant and therefore commute with each other and each element of $\mathcal{V}^{(3)}$. Furthermore, when $d \geq 3$, they are linearly independent. Thus, the group of unitaries that they generate, is equal to the center of $\mathcal{V}^{(3)}$,

$$\langle e^{i\alpha_1 B_1}, e^{i\alpha_2 B_2}, e^{i\alpha_3 B_3} : \alpha_i \in \mathbb{R} \rangle = \{e^{i\phi_{\square\square\square}} \Pi_{\square\square\square} + e^{i\phi_{\square\Box}} \Pi_{\square\Box} + e^{i\phi_{\Box\square}} \Pi_{\Box\square} : \phi_\lambda \in [0, 2\pi)\}. \quad (\text{A3})$$

Using Table III, or using a character table for \mathbb{S}_3 , it can be determined that

$$\begin{aligned} B_1 &= \Pi_{\square\square\square} + \Pi_{\square\Box} + \Pi_{\Box\square} \\ B_2 &= 3\Pi_{\square\square\square} - 3\Pi_{\Box\square} \\ B_3 &= 2\Pi_{\square\square\square} - \Pi_{\square\Box} + 2\Pi_{\Box\square}. \end{aligned} \quad (\text{A4})$$

Note that, when $d = 2$, $\Pi_{\Box\square} = 0$, and so these three operators are linearly dependent. On the other hand, when $d \geq 3$, iB_3 does not satisfy the \mathbb{Z}_2 condition of [6], so the group generated by B_3 is not achievable with 2-local $SU(d)$ -invariant unitaries (one can also verify by hand that the Lie algebra generated by transpositions $i\mathbf{P}(jk)$ does not contain iB_3). To fully characterize $\mathcal{V}_2^{(3)}$, we find the linear combination of projectors to charge sectors which is orthogonal to both B_1 and B_2 [5].

Proposition (re 1). *For a system with $n = 3$ qudits, the family of unitary evolutions $\exp(-iHt) : t \in \mathbb{R}$ is realizable with 2-qudit $SU(d)$ -invariant unitaries, i.e., $\exp(-iHt) \in \mathcal{V}_2^{(3)}$, if and only if $\text{Tr}(HC) = 0$, where*

$$\begin{aligned} C &= 2(d-1)(d-2)\Pi_{\square\square\square} - (d+2)(d-2)\Pi_{\square\Box} + 2(d+2)(d+1)\Pi_{\Box\square} \\ &= d^2(\mathbf{P}_{(123)} + \mathbf{P}_{(132)}) - 2d(\mathbf{P}_{12} + \mathbf{P}_{13} + \mathbf{P}_{23}) + 4\mathbb{I}. \end{aligned} \quad (\text{A5})$$

Furthermore, when $d \geq 3$, the unitary $V \in \mathcal{V}^{(3)}$ is realizable by 2-qudit $SU(d)$ -invariant unitaries, i.e. $V \in \mathcal{V}_2^{(3)}$, if and only if

$$\det v_{\square\Box} = (\det v_{\square\square\square})(\det v_{\Box\square}). \quad (\text{A6})$$

Proof. Write $C = c_{\square\square\square}\Pi_{\square\square\square} + c_{\square\Box}\Pi_{\square\Box} + c_{\Box\square}\Pi_{\Box\square}$. We find c_λ , up to an overall normalization, using the conditions $\text{Tr } C = \text{Tr } B_1 C = 0$ and $\text{Tr } B_2 C = 0$. Recall the decomposition into isotypic components, $\mathcal{H} = \bigoplus_\lambda \mathcal{Q}_\lambda \otimes \mathcal{M}_\lambda$. Note that $\text{Tr } \Pi_\lambda = d_\lambda m_\lambda$ where $d_\lambda = \dim \mathcal{Q}_\lambda$ and $m_\lambda = \dim \mathcal{M}_\lambda$.

First find that

$$0 = \text{Tr } B_2 C = 3c_{\square\square}d_{\square\square} - 3c_{\square\boxplus}d_{\square\boxplus}, \quad (\text{A7})$$

so $c_{\square\square}d_{\square\square} = c_{\square\boxplus}d_{\square\boxplus}$. Using this in

$$\begin{aligned} 0 = \text{Tr } C &= c_{\square\square}d_{\square\square} + 2c_{\square\boxplus}d_{\square\boxplus} + c_{\square\boxminus}d_{\square\boxminus} \\ &= 2c_{\square\square}d_{\square\square} + 2c_{\square\boxplus}d_{\square\boxplus}, \end{aligned} \quad (\text{A8})$$

we find $c_{\square\boxplus}d_{\square\boxplus} = -c_{\square\square}d_{\square\square}$.

Finally, with the formulas for the dimensions,

$$\begin{aligned} d_{\square\square} &= \binom{d+2}{3} \\ d_{\square\boxplus} &= 2\binom{d+1}{3} \\ d_{\square\boxminus} &= \binom{d}{3}, \end{aligned} \quad (\text{A9})$$

the first part of Eq. (A5) is verified. The second part can be checked with Eqs. (A2) and (A4).

The Hamiltonians $\mathbb{I} = B_1$ and B_2 of Eq. (A2) generate the center of $\mathcal{V}_2^{(3)}$, since B_2 is a sum of 2-local terms. According to Eq. (A4),

$$e^{i\alpha_1\mathbb{I}} e^{i\alpha_2 B_2/3} = e^{i(\alpha_1+\alpha_2)} \Pi_{\square\square} + e^{i\alpha_1} \Pi_{\square\boxplus} + e^{i(\alpha_1-\alpha_2)} \Pi_{\square\boxminus}. \quad (\text{A10})$$

Each choice of values $\alpha_1, \alpha_2 \in [0, 2\pi)$ corresponds to a unique operator in $\mathcal{V}_2^{(3)}$, i.e. this parameterization is one-to-one, and every element of the center of $\mathcal{V}_2^{(3)}$ is of the form Eq. (A10) for some α_1, α_2 . Note that, when $\alpha_1 = \alpha_2 = \pi$, Eq. (A10) is actually inside $\mathcal{S}\mathcal{V}^{(3)}$, since $-\mathbb{I}_{\mathcal{M}_{\square\boxplus}} \in \text{SU}(\mathcal{M}_{\square\boxplus})$. This is the only point of intersection, since any other choice of phases will not have determinant one on all multiplicity subsystems \mathcal{M}_λ .

Conversely, given an arbitrary unitary acting as relative phases, where $\phi_\lambda \in [0, 2\pi)$,

$$e^{i\phi_{\square\square}} \Pi_{\square\square} + e^{i\phi_{\square\boxplus}} \Pi_{\square\boxplus} + e^{i\phi_{\square\boxminus}} \Pi_{\square\boxminus}, \quad (\text{A11})$$

it is of the form Eq. (A10) if and only if there are $\alpha_1, \alpha_2 \in [0, 2\pi)$ such that

$$\begin{aligned} \alpha_1 + \alpha_2 &= \phi_{\square\square} \pmod{2\pi} \\ \alpha_1 &= \phi_{\square\boxplus} \\ \alpha_1 - \alpha_2 &= \phi_{\square\boxminus} \pmod{2\pi}. \end{aligned} \quad (\text{A12})$$

Setting $\alpha_1 = \phi_{\square\boxplus}$, a phase $\alpha_2 \in [0, 2\pi)$ satisfying these equations exists if and only if $\phi_{\square\square} - \phi_{\square\boxplus} = \phi_{\square\boxplus} - \phi_{\square\boxminus} \pmod{2\pi}$, or, in other words,

$$e^{i2\phi_{\square\boxplus}} = e^{i\phi_{\square\square}} e^{i\phi_{\square\boxminus}}, \quad (\text{A13})$$

i.e. $2\phi_{\square\boxplus} = \phi_{\square\square} + \phi_{\square\boxminus} \pmod{2\pi}$.

Now consider an arbitrary element of $\mathcal{V}^{(3)}$:

$$V = (\mathbb{I}_{\mathcal{Q}_{\square\square}} \otimes v_{\square\square}) \oplus (\mathbb{I}_{\mathcal{Q}_{\square\boxplus}} \otimes v_{\square\boxplus}) \oplus (\mathbb{I}_{\mathcal{Q}_{\square\boxminus}} \otimes v_{\square\boxminus}). \quad (\text{A14})$$

How can we check if $V \in \mathcal{V}_2^{(3)}$? The first thing to note is that, since $\mathcal{V}_2^{(3)}$ is semiuniversal, for any $V \in \mathcal{V}^{(3)}$ there is some $\tilde{V} \in \mathcal{S}\mathcal{V}^{(3)} \subseteq \mathcal{V}_2^{(3)}$ such that $V\tilde{V}$ acts as phases,

$$V\tilde{V} = e^{i\phi_{\square\square}} \Pi_{\square\square} + e^{i\phi_{\square\boxplus}} \Pi_{\square\boxplus} + e^{i\phi_{\square\boxminus}} \Pi_{\square\boxminus}. \quad (\text{A15})$$

Since $\tilde{V} \in \mathcal{V}_2^{(3)}$, $V \in \mathcal{V}_2^{(3)}$ if and only if $V\tilde{V} \in \mathcal{V}_2^{(3)}$. Furthermore, since $\det \tilde{v}_\lambda = 1$ for each $\lambda = \square\square, \square\boxplus, \square\boxminus$, it also holds that

$$\det v_\lambda = \det v_\lambda \tilde{v}_\lambda = e^{i\phi_\lambda m_\lambda}, \quad (\text{A16})$$

where $m_\lambda = \dim \mathcal{M}_\lambda$. Since $V\tilde{V} \in \mathcal{V}_2^{(3)}$ if and only if $2\phi_{\square\square} = \phi_{\square\square} + \phi_{\square\square} \pmod{2\pi}$, it follows that $V \in \mathcal{V}_2^{(3)}$ if and only if

$$\det v_{\square\square} = (\det v_{\square\square})(\det v_{\square\square}). \quad (\text{A17})$$

□

Note that, since $\mathcal{V}_2^{(3)}$ is semiuniversal, i.e. it contains $\mathcal{SV}^{(3)} \cong \text{SU}(2)$ which acts nontrivially only in the 2D multiplicity subsystem $\mathcal{M}_{\square\square}$ (the other two are 1D), the discussion after Eq. (A10) implies

$$\mathcal{V}_2^{(3)} \cong \text{U}(1) \times \text{U}(2) \quad (\text{A18})$$

since the $\text{U}(1) \times \text{U}(1)$ subgroup parameterized by Eq. (A10) intersects $\mathcal{SV}^{(3)}$ only in the element which acts on $\mathcal{M}_{\square\square}$ as $-\mathbb{I}$. (When $d = 2$ this isomorphism is obvious since then $\mathcal{V}_2^{(3)} \cong \mathcal{V}^{(3)}$.)

2. Generic 3-qudit gates are semi-universal

Here we show Proposition 2, that any 4-qudit gate which does not satisfy the condition

$$[J][\pi_{\square\square}(Y)][J]^T \neq e^{i\phi}[\pi_{\square\square}(Y)]^*,$$

is enough to achieve semi-universality with, along with 2-qudit gates. Note that this condition is related to the \mathbb{Z}_2 condition of [6].

Proof of Proposition 2. One can easily see the necessity of this condition: All unitaries in $\mathcal{V}_2^{(4)}$ satisfy the constraint in Eq. (41). If Y also satisfies this constraint, then the group generated by them does as well, which means it cannot contain $\mathcal{SV}^{(4)}$ (Note that if unitaries V_1 and V_2 satisfy Eq. (41), then their inverses and their product also satisfies this constraint, i.e., this condition defines a subgroup).

Next, to see the sufficiency of the condition in Eq. (44), first recall that the projection of $\mathcal{V}_2^{(4)}$ to each of the multiplicity spaces $\mathcal{M}_{\square\square\square}$, $\mathcal{M}_{\square\square}$, and $\mathcal{M}_{\square\square}$, is equal to the full unitary group in these spaces, which means condition **A** of Lemma 1 is satisfied. In this case condition **B** is only relevant for the pair $\lambda = \square\square$ and $\lambda = \square\square$, which both have dimension 3.

Furthermore, for all such unitaries the components in $\mathcal{M}_{\square\square\square}$ and $\mathcal{M}_{\square\square}$ satisfy the constraint in Eq. (41). Therefore, there exists a unitary $\tilde{Y} \in \mathcal{V}_2^{(4)}$, such that $\pi_{\square\square}(\tilde{Y}) = \pi_{\square\square}(Y)$. Thus $T = Y\tilde{Y}^\dagger$ acts trivially on $\square\square$, and inside $\square\square$ acts non-trivially, i.e., it is not proportional to a global phase, such that $\pi_{\square\square}(Y\tilde{Y}^\dagger) \neq e^{i\phi} \mathbb{I}_{\square\square}$ for all phases $e^{i\phi}$. This together with Eq. (43) implies that condition **B'** in Lemma 1 is also satisfied. Therefore, Lemma 1 implies that $\mathcal{SV}^{(4)} \subset \langle Y, V : V \in \mathcal{V}_2^{(4)} \rangle$. □

We now prove Proposition 3, that for any 3-qudit $\text{SU}(d)$ -invariant unitary V which cannot be generated by 2-qudit ones, the unitary $Y = V \otimes \mathbb{I}$ satisfies the condition of Proposition 2. In particular, due to Proposition 1, generic 3-qudit $\text{SU}(d)$ -invariant unitaries are semi-universal when combined with 2-qudit ones.

Proof of Proposition 3. To see this first we consider unitaries of the form

$$\tilde{S} = e^{i\phi_{\square\square}} \Pi_{\square\square} + e^{i\phi_{\square\square}} \Pi_{\square\square} + e^{i\phi_{\square\square}} \Pi_{\square\square}. \quad (\text{A19})$$

Then, for $Y = \tilde{S} \otimes \mathbb{I}$, Eq. (45) is equivalent to

$$\left| \exp i(\phi_{\square\square} - \phi_{\square\square}) + \frac{1}{2} \right| \neq \left| \exp i(\phi_{\square\square} - \phi_{\square\square}) + \frac{1}{2} \right|. \quad (\text{A20})$$

This can only hold as equality if $\exp i(\phi_{\square\square} - \phi_{\square\square})$ and $\exp i(\phi_{\square\square} - \phi_{\square\square})$ are either equal or related by complex conjugate (see the diagram Fig. 5 for a proof). As seen in Eq. (A13), a unitary of the form \tilde{S} is an element of $\mathcal{V}_2^{(3)}$ if and only if they are complex conjugates, i.e.,

$$\exp i(\phi_{\square\square} - \phi_{\square\square}) = \exp i(\phi_{\square\square} - \phi_{\square\square})^*. \quad (\text{A21})$$

Now recall that, for any $V \in \mathcal{V}^{(3)}$ there exists $\tilde{V} \in \mathcal{SV}^{(3)} \subseteq \mathcal{V}_2^{(3)}$ such that $V\tilde{V}$ has the form Eq. (A19) and

$$\begin{aligned}\phi_{\square\square} &= \arg \det v_{\square\square} \\ \phi_{\square\boxplus} &= \arg \det v_{\square\boxplus} + b\pi \pmod{2\pi} \\ \phi_{\boxplus\boxplus} &= \arg \det v_{\boxplus\boxplus}\end{aligned}\tag{A22}$$

where $b = 0, 1$ is not specified by V . But, since $e^{i\pi} = e^{-i\pi}$, the value of b does not affect the validity of Eq. (A21). In other words, the condition Eq. (A21) for $V\tilde{V}$ is determined by

$$\det v_{\square\boxplus} \stackrel{?}{=} (\det v_{\square\square})(\det v_{\boxplus\boxplus}),\tag{A23}$$

which is precisely the condition Eq. (30) for $V \in \mathcal{V}_2^{(3)}$. \square




FIG. 5. A geometric description of the condition in Eq. (A20). This equation holds as equality if, and only if, $\exp i(\phi_{\square\boxplus} - \phi_{\square\square})$ and $\exp i(\phi_{\square\boxplus} - \phi_{\boxplus\boxplus})$ are either equal or complex conjugates. Both $\exp i(\phi_{\square\boxplus} - \phi_{\square\square}) + 1/2$ and $\exp i(\phi_{\square\boxplus} - \phi_{\boxplus\boxplus}) + 1/2$ sit on the translated unit circle centered at $1/2$. Equation (A20) holds as equality only when these points are equidistant from the origin, i.e. lie on the same circle centered at the origin. The intersection of any two such circles consists of at most two distinct points, which are complex conjugates of each other. Thus $|\exp i(\phi_{\square\boxplus} - \phi_{\square\square}) + 1/2| = |\exp i(\phi_{\square\boxplus} - \phi_{\boxplus\boxplus}) + 1/2|$ if and only if they are either equal or complex conjugates, and since $1/2$ is real, the claim holds.

Appendix B: Tools for semiuniversality

In this section, we prove Lemmas 2 and 12, which provide guarantees on controllability, and we prove Goursat's lemma for special unitary groups, Lemma 9, as a corollary of the asymmetric version of the original Goursat's lemma, Lemma 13. We also provide a proof of Serre's lemma, Lemma 10. Together with the argument in Sec. V, this completes the proof of semi-universality of 3-qudit $SU(d)$ -invariant unitaries.

1. Extending controllability from a single block

In this section, we prove theorems that provide guarantees on controllability. In particular, we demonstrate that irreducibility resulting from continuous families of unitaries together with particular subgroups of unitaries acting only on subspaces of the total Hilbert space \mathcal{H} , is enough to achieve full controllability, i.e. all unitaries up to a global phase, $SU(\mathcal{H})$.

In the following, we will often consider a Hilbert space with orthogonal decomposition $\mathcal{H} = \bigoplus_{i=1}^m \mathcal{H}_i$. With respect to such a decomposition, we consider $U(\mathcal{H}_i) \subseteq U(\mathcal{H})$ the subgroup of unitaries which act as the identity on \mathcal{H}_j for $j \neq i$. We also use the fact that, for *any* matrix group $\mathcal{W} \subseteq GL(\mathcal{H})$, the set of linear operators

$$\mathfrak{w} = \{X \in \mathcal{L}(\mathcal{H}) : \forall t \in \mathbb{R}, e^{tX} \in \mathcal{W}\} \quad (\text{B1})$$

is a Lie algebra [70], closed under real linear combination and commutator.¹²

Lemma (re 2). *Let \mathcal{H} be a finite-dimensional Hilbert space with a subspace $\mathcal{H}_1 \subset \mathcal{H}$ such that $\dim \mathcal{H}_1 \geq 3$. Let A_i , $i = 1, \dots, k$, be traceless anti-Hermitian operators on \mathcal{H} and consider the one-parameter groups $\mathcal{A}_i = \{e^{tA_i} : t \in \mathbb{R}\}$. If the group*

$$\mathcal{W} = \langle \mathcal{A}_i, SU(\mathcal{H}_1) : i = 1, \dots, k \rangle \quad (\text{B2})$$

acts irreducibly on \mathcal{H} , then $\mathcal{W} = SU(\mathcal{H})$.

Remark 4. Let $\mathfrak{g} \subseteq \mathfrak{su}(\mathcal{H})$ be a matrix Lie algebra acting irreducibly on a finite-dimensional Hilbert space \mathcal{H} . Lemma 2 is equivalent to the statement that, if there is a subspace $\mathcal{H}_1 \subseteq \mathcal{H}$ such that $\mathfrak{su}(\mathcal{H}_1) \subseteq \mathfrak{g}$ and $\dim \mathcal{H}_1 \geq 3$, then $\mathfrak{g} = \mathfrak{su}(\mathcal{H})$.

Remark 5. The assumption that $\dim \mathcal{H}_1 \geq 3$ can be relaxed to $\dim \mathcal{H}_1 \geq 2$ if the possibility that $\mathcal{W} = \text{Sp}(\mathcal{H})$ is allowed, when $\dim \mathcal{H}$ is even (see [13] for further details).

For the proof of Lemma 2, we will also need Lemmas 11 and 16: the first, which constructs an off-block-diagonal Lie algebra element, is proven in a number of ways Appendix F 3; the second, which describes a generating set for the Lie algebra of determinant-one unitaries, is postponed until after the proof of Lemma 2.

Proof of Lemma 2. Suppose that $\mathcal{H}' \subset \mathcal{H}$ is any proper subspace of \mathcal{H} which contains $\mathcal{H}_1 \subseteq \mathcal{H}'$ and such that $SU(\mathcal{H}') \subseteq \mathcal{W}$. We prove that there is a subspace $\mathcal{H}'' \subseteq \mathcal{H}$ properly containing $\mathcal{H}' \subset \mathcal{H}''$ and such that $SU(\mathcal{H}'') \subseteq \mathcal{W}$. In summary,

$$\mathcal{H}_1 \subseteq \mathcal{H}' \subset \mathcal{H}'' \subseteq \mathcal{H} . \quad (\text{B3})$$

Let Π be the Hermitian projector to \mathcal{H}' and Π_\perp the projector to its orthogonal complement. There necessarily exists some A_i , $i = 1, \dots, k$, such that $\Pi_\perp A_i \Pi \neq 0$, since otherwise \mathcal{W} would act block-diagonally between \mathcal{H}' and its orthogonal complement, i.e. it would not act irreducibly on \mathcal{H} . In the following we denote this operator A_i with A .

Consider the non-zero subspace $\mathcal{F} = \Pi_\perp A \mathcal{H}'$, that is the image of \mathcal{H}' under A , projected to Π_\perp . Let

$$\mathcal{H}'' = \mathcal{H}' \oplus \mathcal{F} = \mathcal{H}' \oplus \Pi_\perp A \mathcal{H}' . \quad (\text{B4})$$

In the following, we will show that for any arbitrary pair of vectors $|\phi\rangle \in \mathcal{H}'$ and $|\psi\rangle \in \mathcal{F}$, the Lie algebra \mathfrak{w} contains $|\psi\rangle\langle\phi| - |\phi\rangle\langle\psi|$.

¹² While \mathcal{W} may not be a Lie group itself, it at least contains a (possibly trivial) connected Lie group, namely the group generated by exponentials of its Lie algebra, $\langle e^{\mathfrak{w}} \rangle \subseteq \mathcal{W}$. In any case, we call \mathfrak{w} the Lie algebra of \mathcal{W} .

First, note that by definition, the linear map $\tilde{A} = \Pi_{\perp} A \Pi$ has full support on subspace $\mathcal{F} = \Pi_{\perp} A \mathcal{H}'$, which means for any vector $|\psi\rangle \in \mathcal{F}$, there exists a vector $|\psi'\rangle$ in \mathcal{H}' , such that

$$|\psi\rangle = \tilde{A}|\psi'\rangle = \Pi_{\perp} A |\psi'\rangle . \quad (\text{B5})$$

Next, we use Lemma 16. This lemma implies that, because $\dim(\mathcal{H}') \geq 3$ and $\text{SU}(\mathcal{H}') \subseteq \mathcal{W}$, then for any $B \in \mathcal{L}(\mathcal{H}')$ (i.e. $B = \Pi B \Pi$), $B^{\dagger} A \Pi_{\perp} + \Pi_{\perp} A B \in \mathfrak{w}$.

Suppose we choose $B = |\psi'\rangle\langle\phi|$, where $|\phi\rangle$ is an arbitrary vector in \mathcal{H}' . In this case, we have

$$\Pi_{\perp} A B = \Pi_{\perp} A |\psi'\rangle\langle\phi| = |\psi\rangle\langle\phi| . \quad (\text{B6})$$

Since A is anti-Hermitian $B^{\dagger} A \Pi_{\perp} = -(\Pi_{\perp} A B)^{\dagger} = -|\phi\rangle\langle\psi|$. Therefore, Lemma 16 implies

$$B^{\dagger} A \Pi_{\perp} + \Pi_{\perp} A B = |\psi\rangle\langle\phi| - |\phi\rangle\langle\psi| \in \mathfrak{w} . \quad (\text{B7})$$

Since this holds for arbitrary $|\phi\rangle \in \mathcal{H}'$ and $|\psi\rangle \in \mathcal{F}$, then applying Lemma 11 to $\mathcal{H}'' = \mathcal{H}' \oplus \mathcal{F}$, we conclude that $\mathfrak{su}(\mathcal{H}'') \subseteq \mathfrak{w}$, and, by exponentiation, $\text{SU}(\mathcal{H}'') \subseteq \mathcal{W}$.

Proceeding inductively, because \mathcal{H} is finite-dimensional and $\text{SU}(\mathcal{H}') \subseteq \mathcal{W}$, it follows that $\text{SU}(\mathcal{H}) \subseteq \mathcal{W}$. \square

Now we state and prove Lemma 11.

Lemma 11. *Consider a finite-dimensional Hilbert space with orthogonal decomposition $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$ and respective dimensions $d = d_1 + d_2$, with $d_1 \geq d_2 \geq 1$. The (real) Lie algebra \mathfrak{g} generated by all $|\psi_1\rangle\langle\psi_2| - |\psi_2\rangle\langle\psi_1|$, where $|\psi_i\rangle \in \mathcal{H}_i$, is equal to $\mathfrak{su}(\mathcal{H})$.*

Proof. Let $|1\rangle, \dots, |d\rangle$ be an orthonormal basis for \mathcal{H} with $|1\rangle, \dots, |d_1\rangle \in \mathcal{H}_1$ and $|d_1 + 1\rangle, \dots, |d_1 + d_2\rangle \in \mathcal{H}_2$. To prove the claim it suffices to show that for any $j \neq k$, both

$$Y_{jk} = |j\rangle\langle k| - |k\rangle\langle j| \quad (\text{B8})$$

and

$$X_{jk} = i(|j\rangle\langle k| + |k\rangle\langle j|) \quad (\text{B9})$$

are in the Lie algebra \mathfrak{g} , which implies

$$Z_{jk} = i(|j\rangle\langle j| - |k\rangle\langle k|) = \frac{1}{2}[X_{jk}, Y_{jk}] \quad (\text{B10})$$

is also in the Lie algebra. Since operators $X_{jk}, Y_{jk}, Z_{jk} : 1 \leq j < k \leq d$ form a basis for $\mathfrak{su}(\mathcal{H})$, this proves the claim.

First, note that for any $1 \leq j \leq d_1$ and $d_1 + 1 \leq k \leq d$, the assumption of lemma implies that $Y_{jk} \in \mathfrak{g}$. Furthermore, choosing $|\psi_1\rangle = i|j\rangle$ and $|\psi_2\rangle = |k\rangle$, the assumption also implies $X_{jk} \in \mathfrak{g}$. This way we obtain all the elements of the above basis that have support in both subspaces \mathcal{H}_1 and \mathcal{H}_2 . Next, we show how we can obtain elements with support restricted to one of these subspaces. We use the fact that for any 3 distinct $j, k, k' \in \{1, \dots, d\}$, we have the commutation relations

$$[Y_{jk}, Y_{kk'}] = Y_{jj'} , \quad (\text{B11})$$

and

$$[X_{jk}, Y_{kk'}] = X_{jj'} . \quad (\text{B12})$$

This completes the proof. \square

2. Two blocks

Lemma 2 proves that irreducibility allows for extending controllability from three-dimensional blocks to the entire Hilbert space. For the proof of Theorem 1 it is also required that controllability can be extended from a two-dimensional block to the entire Hilbert space when $\dim \mathcal{H} = 3$ (note that this holds more generally when $\dim \mathcal{H}$ is odd, see [13]). That is, we need to extend

$$\left(\begin{array}{c|c} \text{SU}(2) & \\ \hline & 1 \end{array} \right) . \quad (\text{B13})$$

to the entire $SU(3)$. Note that here we do not need to explicitly assume irreducibility on the entire Hilbert space: instead, the fact that the one-parameter group is not block-diagonal immediately implies it. (Note that the following lemma is actually more general than described above, but in the case of $\dim \mathcal{H}_\alpha = 2$ and $\dim \mathcal{H}_\beta = 1$ it is equivalent). This lemma is also previously presented in [6].¹³

Lemma 12. ([6]) *Consider a finite-dimensional Hilbert space with an orthogonal decomposition $\mathcal{H} = \mathcal{H}_\alpha \oplus \mathcal{H}_\beta$ such that neither $\dim \mathcal{H}_\alpha = \dim \mathcal{H}_\beta = 2$ or $\dim \mathcal{H}_\alpha = \dim \mathcal{H}_\beta = 1$. The subgroup $SU(\mathcal{H}_\alpha) \subseteq SU(\mathcal{H})$ acts as the identity on \mathcal{H}_β , and vice-versa. Let A be a Hermitian operator on \mathcal{H} which is not block-diagonal with respect to the above decomposition. Then the group*

$$\mathcal{W} = \langle e^{itA}, SU(\mathcal{H}_\alpha), SU(\mathcal{H}_\beta) : t \in \mathbb{R} \rangle, \quad (\text{B14})$$

generated by the unitaries e^{itA} and each $SU(\mathcal{H}_i)$ contains all determinant-one unitaries, $\mathcal{W} \supseteq SU(\mathcal{H})$.

Remark 6. In the case of $\dim \mathcal{H}_\alpha = \dim \mathcal{H}_\beta = 1$, if we instead consider the subgroups of relative phases, i.e. $U(\mathcal{H}_i)$, the theorem holds.

Remark 7. In the case of $\dim \mathcal{H}_\alpha = \dim \mathcal{H}_\beta = 2$, this result can fail in an interesting way. Choose a basis $\{|m\rangle \otimes |n\rangle : m, n = 0, 1\}$ for this space so that $\mathcal{H}_\alpha = \text{span}_{\mathbb{C}}\{|00\rangle, |01\rangle\}$ and $\mathcal{H}_\beta = \text{span}_{\mathbb{C}}\{|10\rangle, |11\rangle\}$. Then, for example, $A = \sigma_x \otimes \mathbb{I}$ is not block-diagonal with respect to the decomposition $\mathcal{H}_\alpha \oplus \mathcal{H}_\beta$. However, $\{e^{iAt} : t \in \mathbb{R}\}$ together with $SU(\mathcal{H}_i)$ generate a subgroup $\text{Sp}(2) \subseteq SU(4)$.

Note the change in notation from Lemma 2 to Lemma 12: A here is Hermitian rather than anti-Hermitian. Let $D_i = \dim \mathcal{H}_i$ and $\mathfrak{g} = \mathfrak{su}(\mathcal{H}_\alpha) \oplus \mathfrak{su}(\mathcal{H}_\beta)$.

Proof. First, we prove the result in the special case where the Hermitian operator A is in the form

$$A = |\Theta_\alpha\rangle\langle\Theta_\beta| + |\Theta_\beta\rangle\langle\Theta_\alpha|, \quad (\text{B15})$$

where $|\Theta_{\alpha,\beta}\rangle$ are normalized vectors in $\mathcal{H}_{\alpha,\beta}$, and then explain how the proof can be generalized.

Since \mathcal{H}_α and \mathcal{H}_β play equivalent roles in the proof, without loss of generality we assume $D_\alpha \geq D_\beta$. In particular, the assumption of the lemma is that we cannot have $D_\alpha = D_\beta = 1$ or $D_\alpha = D_\beta = 2$, and so $D_\alpha \geq 2$, and when $D_\beta = 2$ we also have $D_\alpha \geq 3$. Let $\{|l, \alpha\rangle : l = 1, \dots, D_\alpha\}$ be an orthonormal basis for \mathcal{H}_α and $\{|m, \beta\rangle : m = 1, \dots, D_\beta\}$ be an orthonormal basis for \mathcal{H}_β , with the property that they contain $|\Theta_{\alpha,\beta}\rangle$, so that there exist l_0, m_0 such that $|l_0, \alpha\rangle = |\Theta_\alpha\rangle$ and $|m_0, \beta\rangle = |\Theta_\beta\rangle$, i.e.,

$$\begin{aligned} iA &= i(|\Theta_\alpha\rangle\langle\Theta_\beta| + |\Theta_\beta\rangle\langle\Theta_\alpha|) \\ &= i(|l_0, \alpha\rangle\langle m_0, \beta| + |m_0, \beta\rangle\langle l_0, \alpha|) \\ &\equiv X_{l_0 m_0}. \end{aligned} \quad (\text{B16})$$

It can be easily shown that the operator $X_{l_0 m_0} = iA$ together with \mathfrak{g} generates the full $\mathfrak{su}(D)$. In particular, note that for any $l = 1, \dots, D_\alpha$ with $l \neq l_0$, the commutator of $X_{l_0 m_0} = iA$ with the traceless skew-Hermitian operator $|l_0, \alpha\rangle\langle l, \alpha| - |l, \alpha\rangle\langle l_0, \alpha| \in \mathfrak{g}$, is

$$\begin{aligned} X_{l m_0} &= [X_{l_0 m_0}, |l_0, \alpha\rangle\langle l, \alpha| - |l, \alpha\rangle\langle l_0, \alpha|] \\ &= i(|l, \alpha\rangle\langle m_0, \beta| + |m_0, \beta\rangle\langle l, \alpha|). \end{aligned} \quad (\text{B17})$$

Furthermore, the commutator of this operator with the traceless skew Hermitian operator $i(|l_0, \alpha\rangle\langle l_0, \alpha| - |l, \alpha\rangle\langle l, \alpha|) \in \mathfrak{g}$ is equal to

$$\begin{aligned} Y_{l m_0} &\equiv [X_{l m_0}, i(|l_0, \alpha\rangle\langle l_0, \alpha| - |l, \alpha\rangle\langle l, \alpha|)] \\ &= |m_0, \beta\rangle\langle l, \alpha| - |l, \alpha\rangle\langle m_0, \beta|. \end{aligned} \quad (\text{B18})$$

And,

$$Z_{l m_0} \equiv -\frac{1}{2}[X_{l m_0}, Y_{l m_0}] = i(|m_0, \beta\rangle\langle m_0, \beta| - |l, \alpha\rangle\langle l, \alpha|). \quad (\text{B19})$$

¹³ The proof presented below was included in the first arXiv version of [6].

Therefore, for all $l = 1, \dots, D_\alpha$, operators $X_{lm_0}, Y_{lm_0}, Z_{lm_0}$ are in the Lie algebra $\langle \mathfrak{g}, iA \rangle$. If $D_\beta = 1$, then the linear combinations of these operators with traceless skew-Hermitian operators with support restricted to \mathcal{H}_α yield all traceless skew Hermitian operators on $\mathcal{H}_\alpha \oplus \mathcal{H}_\beta$. Therefore, in this special case we conclude $\mathfrak{su}(D) \subseteq \langle \mathfrak{g}, \{iA\} \rangle$, which proves the lemma.

On the other hand, if $D_\beta > 1$, then for any arbitrary $m \in \{1, \dots, D_\beta\}$ with $m \neq m_0$, we define

$$\begin{aligned} Y_{lm} &= [Y_{lm_0}, |m, \beta\rangle\langle m_0, \beta| - |m_0, \beta\rangle\langle m, \beta|] \\ &= |l, \alpha\rangle\langle m, \beta| - |m, \beta\rangle\langle l, \alpha|. \end{aligned} \quad (\text{B20})$$

Similarly, considering the commutator of Y_{lm} with operator with $i(|m, \beta\rangle\langle m, \beta| - |m_0, \beta\rangle\langle m_0, \beta|) \in \mathfrak{g}$ we obtain operators $X_{lm} = i(|l, \alpha\rangle\langle m, \beta| + |m, \beta\rangle\langle l, \alpha|)$, and from the commutator of X_{lm} with Y_{lm} we obtain $Z_{lm} = \frac{1}{2}[X_{lm}, Y_{lm}] = i(|l, \alpha\rangle\langle l, \alpha| - |m, \beta\rangle\langle m, \beta|)$. Then, the linear combination of operators

$$X_{lm}, Y_{lm}, Z_{lm} : \quad m = 1, \dots, D_\beta ; \quad l = 1, \dots, D_\alpha , \quad (\text{B21})$$

together with operators in \mathfrak{g} contains all skew-Hermitian traceless operators, which implies $\mathfrak{su}(D) \subseteq \langle \{iA\}, \mathfrak{g} \rangle$.

The above argument proves the lemma in the special case where $A = |\Theta_\alpha\rangle\langle\Theta_\beta| + |\Theta_\beta\rangle\langle\Theta_\alpha|$. To complete the proof, next we show that if A is not block-diagonal with respect to $\mathcal{H}_\alpha \oplus \mathcal{H}_\beta$ then the Lie algebra generated by \mathfrak{g} and $\{iA\}$ contains an operator in the form $i(|\Theta_\alpha\rangle\langle\Theta_\beta| + |\Theta_\beta\rangle\langle\Theta_\alpha|)$. Therefore, by applying the above argument this proves the lemma in the general case.

To prove this, we consider the cases of $D_\beta = 1$ and $D_\beta \geq 2$ separately.

The case of $D_\beta = 1$: Let $|\Theta_\beta\rangle \in \mathcal{H}_\beta$ be a normalized vector. Any Hermitian operator A can be written as

$$A = A_\alpha + a|\Theta_\beta\rangle\langle\Theta_\beta| + b|\Gamma\rangle\langle\Theta_\beta| + b^*|\Theta_\beta\rangle\langle\Gamma| . \quad (\text{B22})$$

where $A_\alpha = A_\alpha^\dagger$ has support restricted to \mathcal{H}_α , a is real, $|\Gamma\rangle \in \mathcal{H}_\alpha$ is a normalized vector and the assumption that A is not block-diagonal implies $b \neq 0$.

Let Π_α be the projector to \mathcal{H}_α and define the operator

$$E \equiv |\Gamma\rangle\langle\Gamma| - D_\alpha^{-1}\Pi_\alpha , \quad (\text{B23})$$

which is nonzero because $D_\alpha > 1$. Note that iE is an element of \mathfrak{g} . Next, note that

$$\begin{aligned} [iA, iE] &= (EA_\alpha - A_\alpha E) + b E|\Gamma\rangle\langle\Theta_\beta| - b^* |\Theta_\beta\rangle\langle\Gamma|E \\ &= (EA_\alpha - A_\alpha E) \\ &\quad + [1 - D_\alpha^{-1}](b |\Gamma\rangle\langle\Theta_\beta| - b^* |\Theta_\beta\rangle\langle\Gamma|) . \end{aligned} \quad (\text{B24})$$

The first term $EA_\alpha - A_\alpha E$ is a traceless skew-Hermitian operator with support restricted to \mathcal{H}_α . Therefore $EA_\alpha - A_\alpha E$ is an element of \mathfrak{g} . This means the second term, i.e., $[1 - D_\alpha^{-1}](b |\Gamma\rangle\langle\Theta_\beta| - b^* |\Theta_\beta\rangle\langle\Gamma|)$ is in the Lie algebra generated by \mathfrak{g} and $\{iA\}$. Furthermore, since $b \neq 0$ and $1 - D_\alpha^{-1} \neq 0$, we conclude that

$$i(|\Theta_\alpha\rangle\langle\Theta_\beta| + |\Theta_\beta\rangle\langle\Theta_\alpha|) \in \langle \{iA\}, \mathfrak{g} \rangle , \quad (\text{B25})$$

where

$$|\Theta_\alpha\rangle = -i \frac{b}{|b|} |\Gamma\rangle , \quad (\text{B26})$$

is a normalized state. Since this operator is in the form given in Eq. (B16) then we can proceed with the rest of the proof of the lemma, as presented above.

The case of $D_\beta \geq 2$: The fact that A is hermitian means that it can be written

$$A = A_\alpha + A_{\alpha\beta} + A_{\beta\alpha} + A_\beta , \quad (\text{B27})$$

where $A_\alpha = A_\alpha^\dagger$ has support restricted to \mathcal{H}_α (and likewise for $A_\beta = A_\beta^\dagger$) and $A_{\alpha\beta} = \Pi_\alpha A \Pi_\beta$ satisfies $A_{\alpha\beta} = A_{\beta\alpha}^\dagger$. The assumption that A is not block-diagonal means that there exists a normalized state $|\Gamma\rangle \in \mathcal{H}_\alpha$ so that $0 \neq A_{\beta\alpha}|\Gamma\rangle \in \mathcal{H}_\beta$. Consider again the operator E from Eq. (B23) and the commutator

$$[iA, iE] = (EA_\alpha - A_\alpha E) + EA_{\alpha\beta} - A_{\beta\alpha}E . \quad (\text{B28})$$

Similar to before, $EA_\alpha - A_\alpha E$ is in \mathfrak{g} . Thus $B_1 \equiv EA_{\alpha\beta} - A_{\beta\alpha}E \in \langle \{iA\}, \mathfrak{g} \rangle$. Consider further the commutator

$$B_2 \equiv [[B_1, iE], iE] = A_{\beta\alpha}E^3 - E^3A_{\alpha\beta}. \quad (\text{B29})$$

Then

$$B_1 + D_\alpha^2 B_2 = [-2 + 3D_\alpha - D_\alpha^2](|\Gamma\rangle\langle\Gamma|A_{\alpha\beta} - A_{\beta\alpha}|\Gamma\rangle\langle\Gamma|), \quad (\text{B30})$$

which is not zero because it is assumed that $D_\alpha > 2$ and so

$$-2 + 3D_\alpha - D_\alpha^2 < 0. \quad (\text{B31})$$

(Interestingly, the fact that this is zero when $D_\alpha = 2$ opens up the possibility for the proper subgroup $\text{Sp}(2) \subset \text{SU}(4)$ to appear when $D_\beta = 2$ also. See Appendix F 2 for more discussion.)

With $|\Theta_\alpha\rangle \equiv |\Gamma\rangle$ and

$$|\Theta_\beta\rangle \equiv \frac{i}{\langle\Gamma|A_{\alpha\beta}A_{\beta\alpha}|\Gamma\rangle} A_{\beta\alpha}|\Gamma\rangle, \quad (\text{B32})$$

it is apparent that $i(|\Theta_\alpha\rangle\langle\Theta_\beta| + |\Theta_\beta\rangle\langle\Theta_\alpha|) \in \langle \{iA\}, \mathfrak{g} \rangle$ is in the form Eq. (B16), and so the proof can proceed as before. \square

3. Goursat's lemma for the special unitary group

Recall Definition 1 of a subdirect product: a subgroup of the direct product whose projection to each factor is surjective. Goursat's lemma characterizes all subdirect products of a pair of groups G_1 and G_2 .

Lemma 13 (Asymmetric Goursat's lemma). *The set of all subdirect products $H \subseteq G_1 \times G_2$ is in bijection with the set of all surjective homomorphisms $G_1 \rightarrow G_2/N_2$ where $N_2 \trianglelefteq G_2$ is a normal subgroup.*

Proof. First, suppose that $H \subseteq G_1 \times G_2$ is a subdirect product and consider the set

$$N_2 = \{g_2 \in G_2 : (e_1, g_2) \in H\}. \quad (\text{B33})$$

This is a normal subgroup of G_2 : if $g'_2 \in G_2$ then, since H is a subdirect product, there exists $g_1 \in G_1$ such that $(g_1, g'_2) \in H$, and

$$H \ni (g_1, g'_2)(e_1, g_2)(g_1, g'_2)^{-1} = (e_1, g'_2 g_2 g_2'^{-1}). \quad (\text{B34})$$

For any $g_1 \in G_1$, assign the set of elements of G_2 that show up with g_1 in H ,

$$g_1 \mapsto \tilde{\Phi}_{g_1} = \{g_2 \in G_2 : (g_1, g_2) \in H\}. \quad (\text{B35})$$

If $g_2, g'_2 \in \tilde{\Phi}_{g_1}$, then $(g_1, g_2)^{-1}(g_1, g'_2) = (e_1, g_2^{-1}g'_2) \in H$, so $g_2^{-1}g'_2 \in N_2$. In other words, the cosets $g_2N_2 = g'_2N_2$, and in fact $\tilde{\Phi}_{g_1} = g_2N_2$ since if $(e_1, n_2) \in H$ then $(g_1, g_2n_2) \in H$. It follows that $\tilde{\Phi} : G_1 \rightarrow G_2/N_2$ is surjective, since H is subdirect, and it is a homomorphism since if $(g_1, g_2), (g'_1, g'_2) \in H$ then $g_2g'_2 \in \tilde{\Phi}_{g_1g'_1}$.

Conversely, given a surjective homomorphism $\tilde{\Phi} : G_1 \rightarrow G_2/N_2$ for some normal subgroup $N_2 \trianglelefteq G_2$, define the subset

$$H = \{(g_1, g_2) \in G_1 \times G_2 : \tilde{\Phi}_{g_1} = g_2N_2\}. \quad (\text{B36})$$

It is easily verified that this is a subdirect product: it's a subgroup since $\tilde{\Phi}$ is a homomorphism and the projections $\pi_i(H) = G_i$ since $\tilde{\Phi}$ is surjective (every element $g_2 \in G_2$ shows up in some coset, namely g_2N_2). \square

In applying Goursat's lemma to the special unitary groups, we take advantage of two facts: that special unitary groups are simply connected simple Lie groups, and that they enjoy an "automatic continuity" property for their homomorphisms to compact groups (see Theorem 2). Note that simple Lie groups are examples of so-called "quasisimple" groups: in particular, every proper normal subgroup of $\text{SU}(l)$ is contained in its center \mathbb{Z}_l .

Lemma (re 9). *Let $l, l' \geq 2$ and let $G \subseteq \text{SU}(l) \times \text{SU}(l')$ be a subdirect product. There are two possibilities:*

(i) $G = \text{SU}(l) \times \text{SU}(l')$.

(ii) $l = l'$ and $G \cong \text{SU}(l) \times \mathbb{Z}_q$, where q divides l .

Furthermore, in the second case, there exists an isomorphism $\Phi : \text{SU}(l) \rightarrow \text{SU}(l)$ such that

$$G = \{(U, e^{i\theta} \Phi(U)) : U \in \text{SU}(l) \text{ and } e^{i\theta} \in \mathbb{Z}_q \subset \text{SU}(l)\}. \quad (\text{B37})$$

Proof. Consider the proof of the forward direction of Lemma 13, and let

$$U \mapsto \tilde{\Phi}_U = \{U' \in \text{SU}(l') : (U, U') \in G\} \quad (\text{B38})$$

define the surjective homomorphism $\tilde{\Phi} : \text{SU}(l) \rightarrow \text{SU}(l')/N'$, where $N' \trianglelefteq \text{SU}(l')$ is the set of elements that show up with the identity $\mathbb{I} \in \text{SU}(l)$. Since $\text{SU}(l')$ is quasisimple, either $N' = \text{SU}(l')$ or $N' \subseteq \mathbb{Z}_{l'} \subseteq \text{SU}(l')$. In the first case, it immediately follows that $G = \text{SU}(l) \times \text{SU}(l')$. So suppose that $N' \triangleleft \text{SU}(l')$ is a proper subgroup, which, as a subgroup of $\mathbb{Z}_{l'}$, is equal to a cyclic subgroup, $N' = \mathbb{Z}_{q'}$ for some q' that divides l' .

By Theorem 2, since $\text{SU}(l)$ and $\text{SU}(l')/\mathbb{Z}_{q'}$ are both compact Lie groups, it follows that $\tilde{\Phi}$ is continuous. Therefore, since $\text{SU}(l)$ is simply connected, there is a (continuous) surjective homomorphism $\Phi : \text{SU}(l) \rightarrow \text{SU}(l')$ which lifts $\tilde{\Phi}$.¹⁴ The kernel of any homomorphism should be a proper normal subgroup of $\text{SU}(l)$, namely in the center of $\text{SU}(l)$, which is the discrete \mathbb{Z}_l subgroup of phases $e^{i2\pi k/l} \mathbb{I} : k = 0, \dots, l-1$. Since Φ is continuous this implies that $\text{SU}(l')$ has the same dimension as $\text{SU}(l)$ as a Lie group (i.e. manifold), which means $l = l'$ and therefore Φ is an isomorphism.

Setting $q = q'$, it is clear that G consists of the elements of the form Eq. (85). \square

From this result, it immediately follows that the commutator subgroup $[G, G] \subseteq \text{SU}(l') \times \text{SU}(l')$ is perfect and connected, equal to either $\text{SU}(l) \times \text{SU}(l')$ or $\{(U, \Phi(U)) : U \in \text{SU}(l)\} \cong \text{SU}(l)$. Using the following description of normal subgroups of $\text{U}(l)$, we extend this result to subgroups of $\text{U}(l) \times \text{U}(l')$ in Corollary 5. Let $\mathcal{P} = \{e^{i\theta} \mathbb{I} : \theta \in [0, 2\pi)\} \cong \text{U}(1)$ denote the subgroup of overall phases and let $\mathbb{Z}_l = \mathcal{P} \cap \text{SU}(l)$ denote the subgroup with determinant one, isomorphic to the cyclic group of order l .

Lemma 14. *Let $N \leq \text{U}(l)$ be a normal subgroup. Then either $N \subseteq \mathcal{P}$ or $N \supseteq \text{SU}(l)$.*

Proof. Every element of $\text{U}(l)$, and, in particular every element of N , can be written $e^{i\theta} U$ for some $e^{i\theta} \in \mathcal{P}$ and some $U \in \text{SU}(l)$. Thus, $U \in \text{SU}(l) \cap (\mathcal{P}N)$. Furthermore, since the product and intersection of normal subgroups are normal, it follows that

$$\text{SU}(l) \cap (\mathcal{P}N) \trianglelefteq \text{SU}(l). \quad (\text{B39})$$

Since $\text{SU}(l)$ is quasisimple, (see the discussion before Lemma 9), either $\text{SU}(l) \cap (\mathcal{P}N) = \text{SU}(l)$, in which case $\text{SU}(l) \subseteq \mathcal{P}N$, or $\text{SU}(l) \cap (\mathcal{P}N) \subseteq \mathbb{Z}_l$. In the former case, taking commutators gives, since $\text{SU}(l)$ is perfect, $\text{SU}(l) \subseteq [N, N]$. In the latter case, every $U \in \text{SU}(l) \cap (\mathcal{P}N)$ is a phase, so N consists entirely of phases, i.e. $N \subseteq \mathcal{P}$. This completes the proof. \square

Corollary (re 5). *Suppose that $H \subseteq \text{U}(l) \times \text{U}(l')$ is a subgroup with $[H, H] \subseteq \text{SU}(l) \times \text{SU}(l')$ a subdirect product. Then either $[H, H] \cong \text{SU}(l) \times \text{SU}(l)$, or $l = l'$ and there is an isomorphism $\Phi : \text{SU}(l) \rightarrow \text{SU}(l)$ such that $[H, H] = \{(U, \Phi(U)) : U \in \text{SU}(l)\} \cong \text{SU}(l)$. Furthermore, in the second case every element of H is of the form $(e^{i\theta} U, e^{i\phi} \Phi(U))$.*

Proof. Since $[H, H] \subseteq \text{SU}(l) \times \text{SU}(l')$ is a subdirect product, Lemma 9 implies there are two possibilities: either it is the direct product, or $l = l'$ and $[H, H]$ consists entirely of elements of the form $(U, e^{i\theta} \Phi(U))$ where $\Phi : \text{SU}(l) \rightarrow \text{SU}(l)$ is an isomorphism and $e^{i\theta} \in \mathbb{Z}_q \subseteq \mathbb{Z}_l$ for some q that divides l .

Suppose the second case holds and let $(e^{i\theta_1} U_1, e^{i\theta_2} U_2) \in H$ be an arbitrary element, where $U_1, U_2 \in \text{SU}(l)$. Then also $(e^{i\theta_1} \mathbb{I}, e^{i\theta_2} \Phi(U_1)^{-1} U_2) \in H$. But the group $\langle W_2 \in \text{U}(l) : (e^{i\theta_1} \mathbb{I}, W_2) \in H \rangle$ is a normal subgroup (note that it is sufficient to check only that this set is invariant under conjugation by $W_2' \in \text{SU}(l)$ since overall phases vanish). Thus by Lemma 14 it either consists entirely of phases or it contains all $\text{SU}(l)$ elements.

But if for every $W_2 \in \text{SU}(l)$ it held that $(e^{i\phi} \mathbb{I}, W_2) \in H$ for some phase $e^{i\phi}$, then taking commutators with elements $(U, \Phi(U)) \in [H, H]$ would give $(\mathbb{I}, V_2) \in [H, H]$ for every commutator $V_2 = W_2 \Phi(U) W_2^\dagger \Phi(U)^\dagger$, which is in contradiction with the conclusion from Lemma 9 that $(\mathbb{I}, V_2) \in [H, H]$ if and only if $V_2 \in \mathbb{Z}_q$. Thus, it must hold that $\Phi(U_1)^{-1} U_2$ is a phase, i.e. $U_2 = e^{i\theta} \Phi(U_1)$. This proves that all elements of H are of the form $(e^{i\phi_1} U, e^{i\phi_2} \Phi(U))$. Upon taking commutators, we find that the phases vanish, i.e. $[H, H] = \{(U, \Phi(U))\}$. \square

¹⁴ That is, for each $U \in \text{SU}(l)$, some $U' \in \tilde{\Phi}_U$ can be chosen so that the assignment $\Phi : U \mapsto U'$ is a proper homomorphism.

4. Automatic continuity

There is a subtlety in the argument for Lemma 9, namely that in order to lift the projective homomorphism, continuity is required. This is guaranteed by the following result, which is an instance of a more general property known as “automatic continuity”.

Theorem 2 (Theorem 5.64 of [71]). *Assume that $f : G \rightarrow H$ is a group homomorphism where H is a compact group and G is a linear Lie group such that $\mathfrak{g} = [\mathfrak{g}, \mathfrak{g}]$, i.e. $G_0 = [G_0, G_0]$ (where \mathfrak{g} is the Lie algebra of G and $G_0 \subseteq G$ is the connected component of the identity). Then f is continuous.*

Note that “linear Lie group” in [71] is more general than the typical meaning of “matrix Lie group”, so indeed all compact Lie groups are linear Lie groups.

The usual classification of irreps of $SU(l)$ goes through its Lie algebra, and thus requires an assumption of continuity. The above result implies that all representations are automatically continuous. This ensures, for instance, that all isomorphisms $\Phi : SU(l) \rightarrow SU(l)$ are of the form $U \mapsto WUW^\dagger$ or $U \mapsto WU^*W^\dagger$, as in Eq. (86).

5. Serre’s lemma

Now we consider a subdirect product $H \subseteq G_1 \times \cdots \times G_r$ which is pairwise independent. Recall that a group is perfect if it is equal to its commutator subgroup, $[G_i, G_i] = G_i$.

Lemma (Serre’s lemma [30], re 10). *Let $H \subseteq G_1 \times \cdots \times G_r$ be a subgroup such that $\pi_{ij}(H) = G_i \times G_j$ for all pairs $1 \leq i < j \leq r$, where $\pi_{ij} : G_1 \times \cdots \times G_r \rightarrow G_i \times G_j$ is the projection homomorphism. If each G_i is perfect, then $H = G_1 \times \cdots \times G_r$.*

Proof. Without loss of generality, it suffices to prove that, for all $g_1 \in G_1$,

$$(g_1, e_2, \dots, e_r) \in H, \tag{B40}$$

where $e_i \in G_i$ is the identity element. In particular, if this is true for G_1 then it is symmetrically true for all factors G_i , and altogether these elements generate the entire direct product.

We proceed by induction; the base case of $r = 2$ holds from the assumption of the lemma. The induction hypothesis applied to $G_1 \times G_3 \times \cdots \times G_r$ and $G_1 \times G_2 \times G_4 \times \cdots \times G_r$, respectively, implies that, for all $g_1, g'_1 \in G_1$, there exist $g_2 \in G_2$ and $g_3 \in G_3$ such that

$$h_1 = (g_1, g_2, e_3, \dots, e_r) \in H \tag{B41a}$$

$$h_2 = (g'_1, e_2, g_3, e_4, \dots, e_r) \in H. \tag{B41b}$$

Therefore,

$$[h_1, h_2] = ([g_1, g'_1], e_2, \dots, e_r) \in H, \tag{B42}$$

where $[h_1, h_2] = h_1 h_2 h_1^{-1} h_2^{-1}$ is the group commutator. Since G_1 is assumed perfect, elements of the form $[g_1, g'_1]$ generate G_1 , and so Eq. (B40) holds. \square

6. Commutator subgroup is connected component of identity

Here we verify the statement that,

Lemma 15. *Let $\mathcal{T} \subseteq \mathcal{V}^G$ be a subgroup such that \mathcal{T} acts irreducibly on each multiplicity subsystem (i.e. $\pi_\lambda(\mathcal{T}) \subseteq U(\mathcal{M}_\lambda)$ is irreducible for each $\lambda \in \Lambda$). Then the connected component of the identity of $\mathcal{T} \cap \mathcal{SV}^G$ is equal to the connected component of the identity of the commutator subgroup $[\mathcal{T}, \mathcal{T}]$.*

In particular, when $[\mathcal{T}, \mathcal{T}]$ is connected (which is always the case when \mathcal{T} is connected), it is equal to the identity component of $\mathcal{T} \cap \mathcal{SV}^G$. We note that when the group \mathcal{T} is generated by k -local G -invariant gates, it is always compact and connected [5]. Furthermore, in the case of $G = SU(d)$ on d -dimensional qudits, each of $\mathcal{V}_k^{(n)}$ acts irreducibly on the multiplicity subsystems since they contain $\mathbf{P}(\sigma) : \sigma \in \mathbb{S}_n$.

Proof. Let \mathfrak{v} and \mathfrak{t} be the Lie algebras of \mathcal{V}^G and \mathcal{T} , respectively (see Eq. (B1)). Both are groups of unitaries, hence their Lie algebras are reductive [24, 25], meaning that they split into a direct sum of their commutator subalgebra and center,

$$\begin{aligned}\mathfrak{v} &= \mathfrak{sv} \oplus \mathfrak{p} = [\mathfrak{v}, \mathfrak{v}] \oplus \mathfrak{p} \\ \mathfrak{t} &= \mathfrak{s} \oplus \mathfrak{z} = [\mathfrak{t}, \mathfrak{t}] \oplus \mathfrak{z},\end{aligned}\tag{B43}$$

where \mathfrak{sv} and \mathfrak{s} are semisimple and \mathfrak{p} and \mathfrak{z} are commutative (the respective centers of \mathfrak{v} and \mathfrak{t}). Since $\mathfrak{t} \subseteq \mathfrak{v}$, it follows that $\mathfrak{s} \subseteq \mathfrak{sv}$. By Schur's lemma and the fact that each $\pi_\lambda(\mathcal{T})$ acts irreducibly, elements of the center \mathfrak{z} must be real linear combinations of the projectors $i\Pi_\lambda : \lambda \in \Lambda$ to charge sectors, which span \mathfrak{p} . Thus $\mathfrak{z} \subseteq \mathfrak{p}$.

The Lie subalgebra $\mathfrak{s} = [\mathfrak{t}, \mathfrak{t}]$ is also the Lie algebra of the commutator subgroup $[\mathcal{T}, \mathcal{T}]$, meaning that its exponentials generate its connected component of the identity. Because its Lie algebra $\mathfrak{s} \subseteq \mathfrak{sv}$, it follows that \mathfrak{s} is also the Lie algebra of the intersection $\mathcal{T} \cap \mathcal{SV}^G$, i.e. the identity components coincide. \square

It is worth noting that the above argument implies that every element of the connected component of the identity of \mathcal{T} can be written as a product of an element of $[\mathcal{T}, \mathcal{T}]$ and a relative phase in the form e^{iP} with $P \in \mathfrak{z}$.

We now prove Lemma 4.

Proof. Recall that $\mathcal{V}_2^{(n)} = \langle e^{it\mathbb{I}}, e^{it\mathbf{P}_{ij}} : t \in \mathbb{R}, i \neq j \rangle$ is a compact connected Lie group [5], which means any unitary $V \in \mathcal{V}_2^{(n)}$ can be written as exponential $V = \exp(A)$, where A is in the real Lie algebra generated by operators $i\mathbf{P}_{ij} : i \neq j$ and the identity $i\mathbb{I}$. The Lie subalgebra generated by transpositions is equal to the real Lie algebra generated by operators $i(\mathbf{P}_{ij} - \mathbf{P}_{kl}) : i \neq j, k \neq l$ and the permutationally-invariant operator $iB = i\sum_{i \neq j} \mathbf{P}_{ij}$ (This can be seen by noting that the linear spans of the generators of the two Lie algebras are equal). Since B is permutationally-invariant, it commutes with all $i(\mathbf{P}_{ij} - \mathbf{P}_{kl}) : i \neq j, k \neq l$. Therefore any unitary $V \in \mathcal{V}_2^{(n)}$ can be decomposed as $e^{i\theta} \tilde{V} \exp(iBs) = e^{i\theta} \exp(iBs) \tilde{V}$, where \tilde{V} is in the group on the left-hand side of Eq. (55), denoted as \mathcal{X} , and $s \in \mathbb{R}$. It follows that \mathcal{X} contains the commutator subgroup of $\mathcal{V}_2^{(n)}$, namely $\mathcal{SV}_2^{(n)}$. On the other hand, \mathcal{X} itself is also contained in $\mathcal{SV}_2^{(n)}$. This can be seen, for instance, by noting that Hamiltonians $\mathbf{P}_{ij} - \mathbf{P}_{kl}$ are centerless, i.e., $\text{Tr}(u^{\otimes n}[\mathbf{P}_{ij} - \mathbf{P}_{kl}]) = 0$, which means the unitaries realized by them are in $\mathcal{SV}^{(n)} \cap \mathcal{V}_2^{(n)}$, hence by Lemma 15 they are in $\mathcal{SV}_2^{(n)}$. This completes the proof of the first part of the lemma. The second part of the lemma follows from the fact that $\mathcal{V}_2^{(n)}$ contains $\mathbf{P}(\sigma) : \sigma \in \mathbb{S}_n$, which acts irreducibly on \mathcal{M}_λ . This in turn implies its commutator subgroup $\mathcal{SV}_2^{(n)}$ also acts irreducibly on \mathcal{M}_λ . \square

We note that, with Lemma 15, we could alternatively prove this by showing that $i(\mathbf{P}_{ij} - \mathbf{P}_{kl})$ generate the semisimple part of the Lie algebra of $\mathcal{V}_2^{(n)}$, for instance using the commutator identity

$$i(\mathbf{P}_{12} - \mathbf{P}_{23}) = \frac{1}{2} [[i\mathbf{P}_{12}, i\mathbf{P}_{23}], i\mathbf{P}_{31}].\tag{B44}$$

Appendix C: For $d \geq 3$, universality requires n -qudit interactions

In this appendix, we prove that, without ancillae, and when $d \geq 3$, $(n-1)$ -local $SU(d)$ -invariant unitaries are not universal. As discussed before Eq. (59), applying the results of [5], we find that

$$\dim \mathcal{V}_n^{(n)} - \dim \mathcal{V}_k^{(n)} = |\Lambda_{n,d}| - |\Lambda_{k,d}| ,$$

where $|\Lambda_{k,d}|$ is the number of inequivalent irreps of $SU(d)$, or, equivalently \mathbb{S}_n that appear on k qudits.

Lemma 5 (Strict monotonicity of the number of irreps, for $d \geq 3$). *Let $|\Lambda_{k,d}|$ be the number of inequivalent irreps of $SU(d)$, or equivalently \mathbb{S}_k , on k qudits with the total Hilbert space $(\mathbb{C}^d)^{\otimes k}$. Then, for $d \geq 3$, $|\Lambda_{k,d}| > |\Lambda_{k-1,d}|$ for all k .*

Proof. We prove this lemma by defining an injective map $i : \Lambda_{n-1,d} \rightarrow \Lambda_{n,d}$, which implies $|\Lambda_{n,d}| \geq |\Lambda_{n-1,d}|$. Then, we show $\Lambda_{n,d}$ contains elements that are not in the image of i .

This map i is defined as follows. Let $\lambda \in \Lambda_{n-1,d}$, which can be labeled by a Young diagram with $n-1$ boxes. Then $i(\lambda)$ is defined by adding a box to the first row of λ . Clearly, this map is injective, because if $i(\lambda_1) = i(\lambda_2)$, then by removing a box from the first row, we must have $\lambda_1 = \lambda_2$.

It is easy to see that for any $i(\lambda)$, the number of boxes in the first row is always greater than the number of boxes in the second row. Then as long as there exists a $\mu \in \Lambda_{n,d}$, such that the first two rows of μ have the same length, then $\mu \notin i(\Lambda_{n-1,d})$. When $d \geq 3$, such μ always exists: if n is even, we can choose $\mu = \begin{array}{c} \square \cdots \square \\ \square \cdots \square \end{array}$; if n is odd, we can choose $\mu = \begin{array}{c} \square \cdots \square \\ \square \end{array}$. \square

Note that when n is odd and $d = 2$, the above μ does not exist. Indeed, in this case we have $|\Lambda_{n,2}| = |\Lambda_{n-1,2}|$ when n is odd.

Appendix D: Semi-universality on $n = 3$ qudits using 8 ancilla qudits

In this appendix, we consider unitaries that are realizable on 11 qudits using 2-qudit $SU(d)$ -invariant unitaries. More precisely, we focus on the behavior of the group $\mathcal{SV}_2^{(11)}$ in certain irreps that are relevant for the use of ancilla qudits. To understand the properties of this group, we use Marin's characterization of the Lie algebra generated by transpositions (SWAPs) as a subalgebra of the group algebra [12]. Roughly speaking, according to this characterization, for any system with an arbitrary number of qudits, there are two sources of constraints: a relation between charge sectors labeled by L-shape diagrams, and a relation between λ and λ' , where λ' is the transpose (sometimes called conjugate) Young diagram (which may or may not be the same as λ). In particular, this result implies

Proposition 5 (Marin [12]). *Two-qudit $SU(d)$ symmetric Hamiltonians are semi-universal in a subset of irreps labeled by Λ , provided that Λ satisfies the following two criteria:*

1. Λ does not contain L-shape Young diagrams;
2. All $\lambda \in \Lambda$ satisfies $\lambda' \notin \Lambda$.

The following lemma does not actually depend on Proposition 5 in its full strength, which applies to a general number of qudits. In fact, we only need Proposition 5 applied to 11 qudits.

Recall that an operator H is called centerless if $\text{Tr } H\Pi_\lambda = 0$ for all $\lambda \in \Lambda_{n,d}$.

Lemma (re 6). *For any centerless $SU(d)$ -invariant Hamiltonian H , there exists a Hamiltonian \tilde{H} that is realizable with 2-qudit $SU(d)$ -invariant Hamiltonians, i.e., $e^{it\tilde{H}} \in \mathcal{V}_2 : t \in \mathbb{R}$ such that*

$$e^{itH}(|\psi\rangle \otimes |\eta\rangle) = e^{it\tilde{H}}(|\psi\rangle \otimes |\eta\rangle), \quad (\text{D1})$$

for all $|\psi\rangle \in (\mathbb{C}^d)^{\otimes 3}$, where for $d \geq 4$,

$$|\eta\rangle = (|0\rangle \wedge |1\rangle \wedge |2\rangle \wedge |3\rangle)^{\otimes 2} \in (\mathbb{C}^d)^{\otimes 8} \quad (\text{D2})$$

is a 8-qudit state, and for $d = 3$,

$$|\eta'\rangle = (|0\rangle \wedge |1\rangle)^{\otimes 2} \otimes |00\rangle \in (\mathbb{C}^3)^{\otimes 6}, \quad (\text{D3})$$

is a 6-qudit state.

In addition to Proposition 5, the following proof of this lemma requires some elementary knowledge of manipulating Young diagrams, which can be found, for example, in [72].

Proof. When $d \geq 4$, the 8-qudit state $|\eta\rangle$ is within the irrep labeled by the Young diagram $\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}$. Then, one can show that the 11-qudit state $|\psi\rangle \otimes |\eta\rangle$ only has components in the following 9 Young diagrams, which are obtained by adding 3 boxes to $\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}$,

$$\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \quad (\text{D4})$$

These Young diagrams will be denoted as Λ . Here we assume $d \geq 7$. When $d < 7$, we just have to remove all the diagrams with more than d rows. It can be easily checked that Λ satisfies the above two criteria in Proposition 5. Therefore, 2-qudit Hamiltonians are semi-universal in Λ . In other words, for any centerless $SU(d)$ -invariant Hamiltonian H , there exists a Hamiltonian \tilde{H} that is realizable with 2-qudit $SU(d)$ -invariant Hamiltonians, such that

$$e^{itH}(|\psi\rangle \otimes |\eta\rangle) = e^{it\tilde{H}}(|\psi\rangle \otimes |\eta\rangle). \quad (\text{D5})$$

Next, we consider the 6-qudit state $|\eta'\rangle \in (\mathbb{C}^3)^{\otimes 6}$. This state is within the irrep labeled by the Young diagram $\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}$. By tensoring it with the 3 qudits in arbitrary states, we find that the resulting set of Young diagrams is

$$\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \quad (\text{D6})$$

which still satisfies the above two criteria in Proposition 5. By a similar argument, we know that H is realizable using 2-qudit symmetric Hamiltonians. \square

Appendix E: t -designs: Proof of Proposition 4

In this section, we prove Proposition 4. To prove this result, in addition to the semi-universality of 3-qudit gates, we also use the following fact: Let μ_0 and μ_1 be, respectively, the symmetric $\square\square\square\square\square\cdots$ and the standard irrep $\square\square\square\square\cdots$ of \mathbb{S}_n . Then, the projection of $\mathcal{V}_3^{(n)}$ to these two sectors is equal to

$$\pi_{\mu_0, \mu_1}(\mathcal{V}_3^{(n)}) = \pi_{\mu_0, \mu_1}(\mathcal{V}^{(n)}) \cong \mathrm{U}(\mathcal{M}_{\mu_0}) \times \mathrm{U}(\mathcal{M}_{\mu_1}). \quad (\text{E1})$$

To see this note that $\mathcal{V}_3^{(n)}$ contains the subgroup $\exp(i\phi_0 A_0) \exp(i\phi_1 A_1) : \phi_0, \phi_1 \in [0, 2\pi)$, where

$$A_0 = \frac{1}{n} B_2 - \frac{n-3}{2} \mathbb{I}, \quad A_1 = \frac{n-1}{2} \mathbb{I} - \frac{1}{n} B_2, \quad (\text{E2})$$

and $B_2 = \frac{1}{2} \sum_{i,j} \mathbf{P}_{ij}$. It can be easily seen that $A_i \Pi_{\mu_j} = \delta_{i,j} \Pi_{\mu_i}$, for $i, j \in \{0, 1\}$, which means when projected to the sectors μ_0 and μ_1 , $\pi_{\mu_0, \mu_1} \{ \exp(i\phi_0 A_0) \exp(i\phi_1 A_1) : \phi_0, \phi_1 \in [0, 2\pi) \} \cong \mathrm{U}(1) \times \mathrm{U}(1)$. Together with $\mathcal{SV}^{(n)} \subset \mathcal{V}_3^{(n)}$ proves Eq. (E1).

Recall the decomposition $V = \bigoplus_{\lambda} (\mathbb{I}_{\mathcal{Q}_{\lambda}} \otimes v_{\lambda})$. Applying this decomposition to both sides of Eq. (77), we see that this equation holds, if and only if

$$\mathbb{E}_{V \in \mathcal{V}_3^{(n)}} \left[\bigotimes_{i=1}^t v_{\lambda_i} \otimes v_{\lambda'_i}^* \right] = \mathbb{E}_{V \in \mathcal{V}^{(n)}} \left[\bigotimes_{i=1}^t v_{\lambda_i} \otimes v_{\lambda'_i}^* \right], \quad (\text{E3})$$

for all $\lambda_1, \dots, \lambda_t, \lambda'_1, \dots, \lambda'_t \in \Lambda_{n,d}$. To analyze these expectation values, we use the following standard fact: For the expectation value with respect to the Haar measure over $\mathrm{U}(m)$, unless $r = r'$, $\mathbb{E}_{U} (U^{\otimes r} \otimes U^{*\otimes r'}) = 0$. This means that unless $\lambda'_1, \dots, \lambda'_t$ is a permutation of $\lambda_1, \dots, \lambda_t$ the right-hand side of Eq. (E3) vanishes. On the other hand, when they are permutations of each other, phases do not contribute to the expectation value, so semi-universality implies both sides are equal. Therefore, to guarantee Eq. (E3), we only need to consider cases where $\lambda'_1, \dots, \lambda'_t$ is not a permutation of $\lambda_1, \dots, \lambda_t$, and make sure the left-hand side of Eq. (E3) vanishes.

Next, we use the result of [27] which implies that when $n \geq 9$ and $d < n - 1$, the three irreps of \mathbb{S}_n with the lowest dimensions are (i) the symmetric irrep $\mu_0 = \square\square\square\square\square\cdots$ with dimension 1, (ii) the standard irrep μ_1 with dimension $n - 1$, and (iii) the irrep $\square\square\square\cdots$ (two-row Young diagrams with two boxes in the second row) with dimension $\frac{1}{2}n(n - 3)$.

Let Δ be the set of irreps in $\Lambda_{n,d}$ for which the number of their occurrences in $\{\lambda_1, \dots, \lambda_t\}$ and $\{\lambda'_1, \dots, \lambda'_t\}$ are not equal. There are two cases depending on whether Δ contains an element of $\Lambda_{n,d} \setminus \{\mu_0, \mu_1\}$. First, assume it does contain such elements, denoted as δ . Then, for all $t < \frac{1}{2}n(n - 3)$,

$$\mathbb{E}_{V \in \mathcal{SV}^{(n)}} \left[\bigotimes_{i=1}^t v_{\lambda_i} \otimes v_{\lambda'_i}^* \right] \cong \mathbb{E}_{V \in \mathcal{SV}^{(n)}} [v_{\delta}^{\otimes n_{\delta}} \otimes v_{\delta}^{*\otimes n'_{\delta}} \otimes V_{\mu} \otimes V_{\nu}^*] = 0, \quad (\text{E4})$$

where in the second expression we have separated the occurrences of δ from all other irreps, denoted $V_{\mu} \otimes V_{\nu}^*$, and n_{δ} and n'_{δ} are the numbers of times δ shows up in $\lambda_1, \dots, \lambda_t$ and $\lambda'_1, \dots, \lambda'_t$, respectively. To show that this expectation value vanishes, it suffices to show that $\mathbb{E}_{v_{\delta} \in \mathrm{SU}(\mathcal{M}_{\delta})} v_{\delta}^{\otimes n_{\delta}} \otimes (v_{\delta})^{*\otimes n'_{\delta}} = 0$. Since $\dim(\mathcal{M}_{\delta}) \geq \frac{1}{2}n(n - 3)$, we have $\dim(\mathcal{M}_{\delta}) > t \geq n_{\delta}, n'_{\delta}$. This, in turns, implies $|n_{\delta} - n'_{\delta}| < \dim(\mathcal{M}_{\delta})$. We also know that because $\delta \in \Delta$, $n_{\delta} \neq n'_{\delta}$. Recall that with the respect to the Haar measure over $\mathrm{SU}(m)$, we have $\mathbb{E}_{U \in \mathrm{SU}(m)} (U^{\otimes r} \otimes U^{*\otimes r'}) \neq 0$, if and only if $r = r' \pmod{m}$. We conclude that $\mathbb{E}_{v_{\delta} \in \mathrm{SU}(\mathcal{M}_{\delta})} v_{\delta}^{\otimes n_{\delta}} \otimes (v_{\delta})^{*\otimes n'_{\delta}} = 0$, which in turn proves the above equality. Finally, we note that because $\mathcal{SV}^{(n)}$ is a subgroup of $\mathcal{V}_3^{(n)}$ (semi-universality), the above identity implies that

$$\mathbb{E}_{V \in \mathcal{V}_3^{(n)}} \left[\bigotimes_{i=1}^t v_{\lambda_i} \otimes v_{\lambda'_i}^* \right] = \mathbb{E}_{V \in \mathcal{SV}^{(n)}} \left[\bigotimes_{i=1}^t v_{\lambda_i} \otimes v_{\lambda'_i}^* \right] = 0, \quad (\text{E5})$$

Next, we focus on the second case, i.e., when Δ does not have any element if $\Lambda_{n,k} \setminus \{\mu_0, \mu_1\}$, which means $\Delta \subseteq \{\mu_0, \mu_1\}$. In this case there exists $\delta \in \{\mu_0, \mu_1\}$ such that the number of its occurrence in $\{\lambda_1, \dots, \lambda_t\}$ and $\{\lambda'_1, \dots, \lambda'_t\}$, denoted as n_{δ} and n'_{δ} are not equal. Then, in the case applying Eq. (E1) we have

$$\mathbb{E}_{V \in \mathcal{V}_3^{(n)}} v_{\delta}^{\otimes n_{\delta}} \otimes (v_{\delta})^{*\otimes n'_{\delta}} = \mathbb{E}_{V \in \mathcal{V}^{(n)}} v_{\delta}^{\otimes n_{\delta}} \otimes (v_{\delta})^{*\otimes n'_{\delta}} = \mathbb{E}_{v_{\delta} \in \mathrm{U}(\mathcal{M}_{\delta})} v_{\delta}^{\otimes n_{\delta}} \otimes (v_{\delta})^{*\otimes n'_{\delta}} = 0. \quad (\text{E6})$$

Therefore, we conclude that for all $t < n(n - 3)/2$, Eq. (E3) holds for all $\{\lambda_1, \dots, \lambda_t\}$ and $\{\lambda'_1, \dots, \lambda'_t\}$. This completes the proof of Proposition 4.

Appendix F: Techniques for controllability

In this appendix, we describe techniques for controllability using the notion of isolation to subspaces of a Hilbert space. In Appendix F 1 we define “isolation” and we discuss some of its uses. In Appendix F 2 we show that one of the techniques described in Appendix F 1 can fail when the subspaces are two-dimensional. Appendix F 3 provides strategies for generating isolated elements of Lie algebras.

1. Using isolation

In this section we describe some uses of isolating Lie algebra elements to off-diagonal blocks. First, we discuss what we mean by “isolation”. Consider a finite-dimensional Hilbert space with an orthogonal decomposition $\mathcal{H} = \bigoplus_{i=1}^m \mathcal{H}_i$ and a group of unitaries $\mathcal{W} \subseteq \mathbf{U}(\mathcal{H})$. A nonzero element $X \in \mathfrak{w}$ of the Lie algebra of \mathcal{W} ,

$$\mathfrak{w} = \{X \in \mathcal{L}(\mathcal{H}) : \forall t \in \mathbb{R}, e^{tX} \in \mathcal{W}\}, \quad (\text{re B1})$$

is called (off-diagonal) *isolated* to block i if

$$X = \Pi_i X + X \Pi_i. \quad (\text{F1})$$

Note that this implies that $\Pi_i X \Pi_i = 0$ by applying Π_i on the right and left to both sides of Eq. (F1). We also consider isolation to two blocks, say i and j : $X \neq 0$ is *isolated* to these if

$$X = \Pi_i X \Pi_j + \Pi_j X \Pi_i. \quad (\text{F2})$$

Isolated elements in the Lie algebra are particularly helpful when there are nontrivial elements $U_i \in \mathcal{W}$ which act as the identity on \mathcal{H}_j for $j \neq i$. Let $\mathcal{W}_i \subseteq \mathcal{W}$ be the subgroup which acts nontrivially only on the i th block, defined via

$$\mathcal{W}_i = \{U_i \in \mathcal{W} : \Pi_j U_i = U_i \Pi_j = U_i^{\delta_{ij}} \Pi_j\}, \quad (\text{F3})$$

where δ_{ij} is the Kronecker delta, i.e.,

$$U_i^{\delta_{ij}} \Pi_j := \begin{cases} U_i \Pi_i & j = i \\ \Pi_j & j \neq i. \end{cases} \quad (\text{F4})$$

Note that if $W \in \mathcal{W}$ and $X \in \mathfrak{w}$ then $WXW^\dagger \in \mathfrak{w}$ since for all $t \in \mathbb{R}$

$$e^{tWXW^\dagger} = W e^{tX} W^\dagger \in \mathcal{W}. \quad (\text{F5})$$

If $X \in \mathfrak{w}$ is isolated to the i th block (i.e. it satisfies Eq. (F1)), then $\Pi_i X U_i = \Pi_i X$, which, in turn, implies

$$U_i X U_i^\dagger = U_i \Pi_i X + X \Pi_i U_i^\dagger \in \mathfrak{w}, \quad (\text{F6})$$

Furthermore, \mathfrak{w} is closed under real linear combinations, so for all

$$B \in \text{span}_{\mathbb{R}}\{U_i \Pi_i : U_i \in \mathcal{W}_i\} = \text{span}_{\mathbb{R}} \mathcal{W}_i \Pi_i, \quad (\text{F7})$$

it holds that $BX + XB^\dagger \in \mathfrak{w}$. In other words,

$$\{BX + XB^\dagger : B \in \text{span}_{\mathbb{R}} \mathcal{W}_i \Pi_i\} = \text{span}_{\mathbb{R}}\{U_i X U_i^\dagger : U_i \in \mathcal{W}_i\} \subseteq \mathfrak{w}. \quad (\text{F8})$$

As long as there is some $U_i \in \mathcal{W}_i$ such that $U_i X U_i^\dagger \neq X$, this can be used to generate new elements of \mathcal{W} which act only in the subspace $\mathcal{H}_i \oplus X\mathcal{H}_i$ (where $X\mathcal{H}_i$ is the image of $X\Pi_i$; note that $X\mathcal{H}_i$ consists entirely of vectors orthogonal to \mathcal{H}_i since $\Pi_i X \Pi_i = 0$).

For instance, suppose that

$$\text{span}_{\mathbb{R}}\{\Pi_i U_i : U_i \in \mathcal{W}_i\} = \text{span}_{\mathbb{C}}\{\Pi_i U_i : U_i \in \mathcal{W}_i\} \quad (\text{F9})$$

(this implies that \mathcal{W}_i does not act on \mathcal{H}_i by orthogonal or symplectic matrices), and that the action of \mathcal{W}_i on \mathcal{H}_i is irreducible, so that

$$\text{span}_{\mathbb{C}}\{\Pi_i U_i : U_i \in \mathcal{W}_i\} = \mathcal{L}(\mathcal{H}_i) = \{B \in \mathcal{L}(\mathcal{H}) : B = \Pi_i B \Pi_i\}. \quad (\text{F10})$$

i.e. the span of the unitaries in \mathcal{W}_i , restricted to \mathcal{H}_i , consists of all linear operators on \mathcal{H}_i . Then the real span includes all rank-one projectors on \mathcal{H}_i , and so Eq. (F8) includes all operators of the form $|\psi\rangle\langle\varphi| - |\varphi\rangle\langle\psi|$ where $|\psi\rangle \in \mathcal{H}_i$ and $|\varphi\rangle \in X\mathcal{H}_i$. Then, since this is a generating set for $\mathfrak{su}(\mathcal{H}_i \oplus X\mathcal{H}_i) \subseteq \mathfrak{w}$ (see Lemma 11),

$$\mathrm{SU}(\mathcal{H}_i \oplus X\mathcal{H}_i) \subseteq \mathcal{W}. \quad (\text{F11})$$

Applying this technique in the section we prove the following lemma, which is used in Appendix B 1 for the proof of Lemma 2.

Lemma 16. *Let $\mathcal{W} \subseteq \mathrm{U}(\mathcal{H})$ be a group of unitaries on $\mathcal{H} = \bigoplus_{i=1}^m \mathcal{H}_i$. Suppose that, for some $i = 1, \dots, m$ the dimension $\dim \mathcal{H}_i \geq 3$ and $\mathrm{SU}(\mathcal{H}_i) \subseteq \mathcal{W}_i$, where $\mathcal{W}_i \subseteq \mathcal{W}$ is the subgroup acting nontrivially only on \mathcal{H}_i , defined in Eq. (F3). Let Π_i be the Hermitian projector to \mathcal{H}_i and Π_\perp the projector to its orthogonal complement. If A is an (anti-Hermitian) operator such that $e^{tA} \in \mathcal{W}$ for all $t \in \mathbb{R}$, then for all $B = \Pi_i B \Pi_i$ and $t \in \mathbb{R}$,*

$$\exp t(BA\Pi_\perp + \Pi_\perp AB^\dagger) \in \mathcal{W}. \quad (\text{F12})$$

After discussing isolation techniques, we prove this lemma in Appendix F 4. In the next subsection, we discuss why the assumption that $\dim \mathcal{H}_i \geq 3$ is necessary.

It is worth noting that, if \mathcal{W}_i has a nontrivial Lie algebra, denoted \mathfrak{w}_i , then an isolated $X \in \mathfrak{w}$ can possibly (as long as it does not commute with \mathfrak{w}_i) be used to generate a larger subalgebra $\langle \mathfrak{w}_i, X \rangle \subseteq \mathfrak{w}$.

2. Why two-dimensional blocks are special

The assumption that $\mathrm{span}_{\mathbb{R}} \mathcal{W}_i \Pi_i = \mathrm{span}_{\mathbb{C}} \mathcal{W}_i \Pi_i$, discussed above, is not satisfied for a particularly important group: when $\mathcal{W}_i = \mathrm{SU}(\mathcal{H}_i)$ and $\dim \mathcal{H}_i = 2$. This is related to the exceptional isomorphism $\mathrm{SU}(2) \cong \mathrm{Sp}(1)$ with the compact symplectic group [14].

One way of seeing this is the following. Every element $U \in \mathrm{SU}(2)$ can be written

$$U = e^{i\theta(\hat{\mathbf{n}} \cdot \boldsymbol{\sigma})} = \cos \theta \mathbb{I} + i \sin \theta (\hat{\mathbf{n}} \cdot \boldsymbol{\sigma}), \quad (\text{F13})$$

where $\boldsymbol{\sigma}$ is the Pauli vector operator, $\hat{\mathbf{n}} \in \mathbb{R}^3$ is a unit vector, and $\theta \in [0, 2\pi)$. This implies

$$\mathrm{span}_{\mathbb{R}} \mathrm{SU}(2) = \mathrm{span}_{\mathbb{R}} \{\mathbb{I}, i\sigma_x, i\sigma_y, i\sigma_z\}. \quad (\text{F14})$$

Notably, the traceless part of any $B \in \mathrm{span}_{\mathbb{R}} \mathrm{SU}(2)$ is anti-Hermitian. Hence $\mathrm{span}_{\mathbb{R}} \mathrm{SU}(2)$ is clearly a proper *real* subspace of $\mathcal{L}(\mathbb{C}^2)$, the space of linear operators over \mathbb{C}^2 :

$$\mathrm{span}_{\mathbb{R}} \mathrm{SU}(2) \neq \mathrm{span}_{\mathbb{C}} \mathrm{SU}(2) = \mathcal{L}(\mathbb{C}^2). \quad (\text{F15a})$$

In fact, with the identity $(\hat{\mathbf{n}} \cdot \boldsymbol{\sigma})(\hat{\mathbf{n}}' \cdot \boldsymbol{\sigma}) = \hat{\mathbf{n}} \cdot \hat{\mathbf{n}}' \mathbb{I} + i(\hat{\mathbf{n}} \times \hat{\mathbf{n}}') \cdot \boldsymbol{\sigma}$, it can be seen that $\mathrm{span}_{\mathbb{R}} \mathrm{SU}(2) \cong \mathbb{H}$ where \mathbb{H} is the associative algebra of quaternions.

On the other hand, for $\mathrm{SU}(d)$ with $d > 2$, the element $e^{i2\pi/d} \mathbb{I} \in \mathrm{SU}(d)$. It follows that

$$d \geq 3 \implies \mathrm{span}_{\mathbb{R}} \mathrm{SU}(d) = \mathrm{span}_{\mathbb{C}} \mathrm{SU}(d), \quad (\text{F15b})$$

since any complex scalar is a real number plus a real multiple of the root of unity $e^{i2\pi/d}$. Similarly,

$$\mathrm{span}_{\mathbb{R}} \mathrm{U}(2) = \mathrm{span}_{\mathbb{C}} \mathrm{U}(2), \quad (\text{F15c})$$

so this phenomenon does not occur for $\mathrm{U}(2)$ subgroups in the diagonal blocks of \mathcal{W} , or, more generally, for any subgroups that contain phases that are not real.

3. Strategies for isolation

In this section we consider a number of strategies for isolating Lie algebra elements to off-diagonal blocks of $\mathcal{H} = \bigoplus_{i=1}^m \mathcal{H}_i$. The strategies we discuss are summarized in Table IV. For the following, fix a group of unitaries $\mathcal{W} \subseteq \mathrm{U}(\mathcal{H})$ and let $\mathcal{W}_i \subseteq \mathcal{W}$ be the subgroups defined in Eq. (F3), which necessarily act as the identity on \mathcal{H}_j for $j \neq i$.

$\int_{\mathcal{V}_i} dU_i f(U_i) U_i X U_i^\dagger$		assumptions	
$D_{U_i}(X) = U_i X U_i^\dagger - X$	$\begin{pmatrix} 0 & & D_{U_i}(X_{1i}) & \\ & \ddots & \vdots & \\ D_{U_i}(X_{i1}) & \cdots & D_{U_i}(X_{ii}) & \cdots \\ & & \vdots & \ddots \end{pmatrix}$	<ul style="list-style-type: none"> \mathcal{V}_i nontrivial 	Equation (F19)
$(\mathcal{I} - \mathcal{E}_i)(X)$	$\begin{pmatrix} 0 & & X_{1i} & \\ & \ddots & \vdots & \\ X_{i1} & \cdots & X_{ii} - \mathcal{E}_i(X_{ii}) & \cdots \\ & & \vdots & \ddots \end{pmatrix}$	<ul style="list-style-type: none"> \mathcal{V}_i is compact, with no nonzero invariant vectors in \mathcal{H}_i \mathcal{E}_i is uniform Haar integral over \mathcal{V}_i 	Equation (F25)
$f(U_i) = \text{tr}(B^\dagger U_i + B U_i^\dagger)$	$\begin{pmatrix} 0 & & X_{1i} B^\dagger & \\ & \ddots & \vdots & \\ B X_{i1} & \cdots & 0 & \cdots \\ & & \vdots & \ddots \end{pmatrix}$	<ul style="list-style-type: none"> \mathcal{V}_i is compact, acts nontrivially and irreducibly on \mathcal{H}_i $\mathcal{H}_i \otimes \mathcal{H}_i$ does not contain subrepresentation isomorphic to trivial or \mathcal{H}_i $B = \Pi_i B \Pi_i$ 	Equation (F37)
$(D_{U_j} \circ D_{U_i})(X)$	$\begin{pmatrix} \ddots & & & & \\ & 0 & & \tilde{X}_{ij} & \\ & & \ddots & & \\ \tilde{X}_{ji} & & & 0 & \\ & & & & \ddots \end{pmatrix}$	<ul style="list-style-type: none"> \mathcal{V}_i and \mathcal{V}_j nontrivial $\tilde{X}_{ij} = (\mathbb{I} - U_i) X_{ij} (\mathbb{I} - U_j^\dagger)$ and $\tilde{X}_{ji} = -\tilde{X}_{ij}^\dagger$ 	Equation (F20)
$((\mathcal{I} - \mathcal{E}_j) \circ (\mathcal{I} - \mathcal{E}_i))(X)$	$\begin{pmatrix} \ddots & & & & \\ & 0 & & X_{ij} & \\ & & \ddots & & \\ X_{ji} & & & 0 & \\ & & & & \ddots \end{pmatrix}$	<ul style="list-style-type: none"> \mathcal{V}_i and \mathcal{V}_j are compact with no nonzero invariant vectors in \mathcal{H}_i and \mathcal{H}_j, respectively \mathcal{E}_i and \mathcal{E}_j are uniform Haar integrals over \mathcal{V}_i and \mathcal{V}_j, respectively 	Equation (F27)

TABLE IV. This table shows various isolation strategies for block i of $\mathcal{H} = \bigoplus_{j=1}^m \mathcal{H}_j$. The first three rows can be understood as an integral weighted by a real-valued function f over the group $\mathcal{V}_i \subseteq \mathcal{W}_i \subseteq \text{U}(\mathcal{H}_i)$ which acts nontrivially on \mathcal{H}_i and trivially on \mathcal{H}_j for $j \neq i$ (really f is to be understood as a distribution: for instance the Dirac delta distribution δ_{U_i} , satisfying $\int dU'_i \delta_{U'_i} K(U'_i) = K(U_i)$, is used in the first row). The last two rows have two such integrals, over \mathcal{V}_i and \mathcal{V}_j . The color scheme for blocks has (1) white background for unchanged blocks; (2) red background for blocks which necessarily become 0; and (3) blue background for blocks which may be changed but are not necessarily 0 (the exact value depending on X and f).

a. Using one nontrivial group element: D_{U_i}

For $X \in \mathfrak{w}$, let $X_{jk} = \Pi_j X \Pi_k$. Then for $U_i \in \mathcal{W}_i$,

$$U_i X U_i^\dagger = \sum_{j,k} U_i X_{jk} U_i^\dagger = \sum_{j,k} (U_i)^{\delta_{ij}} X_{jk} (U_i^\dagger)^{\delta_{ik}} \in \mathfrak{w}, \quad (\text{F16})$$

where $X_{ij} = \Pi_i X \Pi_j$ and δ_{ij} is the Kronecker delta. Written in block-matrix form,

$$U_i X U_i^\dagger = \begin{pmatrix} X_{11} & & X_{1i} U_i^\dagger & \\ & \ddots & \vdots & \\ U_i X_{i1} & \cdots & U_i X_{ii} U_i^\dagger & \cdots \\ & & \vdots & \ddots \end{pmatrix} \in \mathfrak{w}, \quad (\text{F17})$$

where the blue-shaded region denotes all the blocks which are possibly distinct from X , namely those in the i th row and i th column. This immediately implies that subtracting off X will set all non-shaded blocks to zero. Suggested by this, for any unitary U and operator X , define

$$D_U(X) = U X U^\dagger - X. \quad (\text{F18})$$

To first order, the superoperator D_U can be understood as a derivation: if $U = e^{\varepsilon B}$ then expanding in ε gives $D_U(X) = \varepsilon[B, X] + O(\varepsilon^2)$.

If $X \in \mathfrak{w}$ and $U \in \mathcal{W}$, then $D_U(X) \in \mathfrak{w}$ since it is a sum of terms of \mathfrak{w} with real coefficients. Hence D_U can possibly be used as a tool for isolating to particular blocks. To reiterate, $D_{U_i}(X)$ has a large number of blocks which are guaranteed to be zero. In block-matrix form, $D_{U_i}(X)$ can be nonzero only in the blue-shaded region of

$$D_{U_i}(X) = \begin{pmatrix} 0 & & D_{U_i}(X_{1i}) & \\ & \ddots & \vdots & \\ D_{U_i}(X_{i1}) & \cdots & D_{U_i}(X_{ii}) & \cdots \\ & & \vdots & \ddots \end{pmatrix} \in \mathfrak{w}. \quad (\text{F19})$$

Furthermore, when $j \neq i$, $D_{U_i}(X_{ij}) = (U_i - \mathbb{1})X_{ij}$, and $D_{U_i}(X_{ji}) = -D_{U_i}(X_{ij})$ since $X \in \mathfrak{w}$ is anti-Hermitian. If $X_{ij} \neq 0$ and if \mathcal{H}_i does not contain any non-zero invariant vector under \mathcal{W}_i , which is always the case if \mathcal{W}_i acts nontrivially and irreducibly on \mathcal{H}_i , then it is always possible to find $U_i \in \mathcal{W}_i$ such that $U_i X_{ij} \neq X_{ij}$, so that $D_{U_i}(X_{ij}) \neq 0$. (Recall that if \mathcal{W}_i acts nontrivially and irreducibly on \mathcal{H}_i , for any vector $0 \neq |\psi\rangle \in \mathcal{H}_i$, there exists $U_i \in \mathcal{W}_i$, such that $U_i |\psi\rangle \neq |\psi\rangle$. This, in turn, implies that for any nonzero operator $X_{ij} = \Pi_i X_{ij}$, there exists $U_i \in \mathcal{W}_i$ such that $(U_i - \mathbb{1})X_{ij} \neq 0$.)

The utility of D_{U_i} comes in two forms:

- i) First, the possibly-remaining block-diagonal element $D_{U_i}(X_{ii}) \in \mathfrak{su}(\mathcal{H}_i)$ since it is traceless and anti-Hermitian, although any of the off-diagonal blocks in the i th row or i th column may be zero or nonzero. If $\dim \mathcal{H}_i = 1$, then, if nonzero, this implies $D_{U_i}(X) \in \mathfrak{w}$ is isolated to block i . Otherwise, if $\mathfrak{su}(\mathcal{H}_i) \subset \mathfrak{w}$ then $D_{U_i}(X) - D_{U_i}(X_{ii}) \in \mathfrak{w}$ is purely off-diagonal (i.e. isolated).
- ii) Second, by sequentially applying $D_{U_j} \circ D_{U_i}$ for distinct blocks $i \neq j$, we obtain an operator which, when nonzero, is guaranteed to be purely off-diagonal and isolated to just these two blocks. (It is worth noting that, since $[U_i, U_j] = 0$, also $[D_{U_i}, D_{U_j}] = 0$, i.e. it does not matter in which order they are applied.)

To clarify the second point, note that the only surviving terms have to be in the overlap of the blue-shaded regions corresponding to D_{U_i} and D_{U_j} ,

$$D_{U_j} \circ D_{U_i}(X) = \begin{pmatrix} \ddots & & & & \\ & 0 & & \tilde{X}_{ij} & \\ & & \ddots & & \\ & \tilde{X}_{ji} & & 0 & \\ & & & & \ddots \end{pmatrix} \in \mathfrak{w}. \quad (\text{F20})$$

That is, only $D_{U_j} \circ D_{U_i}(X_{ij})$ and $D_{U_j} \circ D_{U_i}(X_{ji})$ can possibly be nonzero. Once again, if $X_{ij} \neq 0$ and U_i and U_j come from nontrivial groups which do not have nonzero invariant vectors in their respective subspaces, the group elements can always be chosen so that the result is nonzero.

b. Integration over the uniform Haar measure

The scheme using D_{U_i} ensures that the only nonzero blocks are in the i th row and i th column; however, any of the blocks may be changed (possibly becoming zero). We next discuss a technique using the Haar measure which does the same, except that it leaves all the off-diagonal blocks invariant. First, in full generality, we notice that integrating the adjoint action (Eq. (F5)) of \mathcal{W} on its Lie algebra \mathfrak{w} over group elements with real coefficients stays in the Lie algebra, since it is a closed (real) subspace of the set of all operators.

Proposition 6. *Let \mathcal{W} be a matrix Lie group and $\mathcal{S} \subseteq \mathcal{W}$ a subset with a measure ds (e.g. a finite set with the counting measure), and let $f(s)$ be a real-valued integrable function (or, more generally, a real-valued distribution) on \mathcal{S} . For any matrix A , if $e^{tA} \in \mathcal{W}$ for all $t \in \mathbb{R}$, then $e^{t\tilde{A}} \in \mathcal{W}$ for all $t \in \mathbb{R}$, where*

$$\tilde{A} = \int ds f(s) s A s^{-1}. \quad (\text{F21})$$

In particular, assuming that there is a nontrivial compact subgroup $\mathcal{V}_i \subseteq \mathcal{W}_i \subseteq \mathcal{W}$ which does not have any nonzero invariant vectors on \mathcal{H}_i , we will apply this to

$$\mathcal{E}_i(X) = \int_{\mathcal{V}_i} dU_i U_i X U_i^\dagger \in \mathfrak{w}. \quad (\text{F22})$$

Then $\mathcal{E}_i(X)$ is zero in the red-shaded region and the same as X everywhere else except for the i th diagonal block:

$$\mathcal{E}_i(X) = \begin{pmatrix} X_{11} & & 0 & & \\ & \ddots & \vdots & & \\ 0 & \cdots & \mathcal{E}_i(X_{ii}) & \cdots & \\ & & \vdots & & \ddots \end{pmatrix} \in \mathfrak{w}. \quad (\text{F23})$$

This follows because

$$\int_{\mathcal{V}_i} dU_i U_i \Pi_i = \int_{\mathcal{V}_i} dU_i U_i^\dagger \Pi_i = 0, \quad (\text{F24})$$

is the projector to the subspace of invariant vectors of \mathcal{H}_i , which consists only of the zero vector. Letting \mathcal{I} be the identity superoperator, $\mathcal{I}(X) = X$ for all X , it follows that

$$(\mathcal{I} - \mathcal{E}_i)(X) = \begin{pmatrix} 0 & & X_{1i} & & \\ & \ddots & \vdots & & \\ X_{i1} & \cdots & X_{ii} - \mathcal{E}_i(X_{ii}) & \cdots & \\ & & \vdots & & \ddots \end{pmatrix} \in \mathfrak{w}. \quad (\text{F25})$$

Note that $\text{Tr } X_{ii} = \text{Tr } \mathcal{E}_i(X_{ii})$. Thus $X_{ii} - \mathcal{E}_i(X_{ii})$ is traceless and anti-Hermitian, hence an element of $\mathfrak{su}(\mathcal{H}_i)$. If $\mathfrak{su}(\mathcal{H}_i) \subseteq \mathfrak{w}$, then

$$(\mathcal{I} - \mathcal{E}_i)(X - X_{ii}) = (\mathcal{I} - \mathcal{E}_i)(X) - (X_{ii} - \mathcal{E}_i(X_{ii})) \in \mathfrak{w} \quad (\text{F26})$$

is an element isolated to the i th block. Regardless, if a compact subgroup $\mathcal{V}_j \subseteq \mathcal{W}_j$ also does not have invariant nonzero vectors on \mathcal{H}_j for some $j \neq i$, then

$$(\mathcal{I} - \mathcal{E}_j) \circ (\mathcal{I} - \mathcal{E}_i)(X) = \begin{pmatrix} \ddots & & & & \\ & 0 & & X_{ij} & \\ & & \ddots & & \\ & X_{ji} & & 0 & \\ & & & & \ddots \end{pmatrix} \in \mathfrak{w} \quad (\text{F27})$$

is isolated to blocks i and j .

Another use-case involves using the Haar measure on multiple blocks to set many off-diagonal blocks to 0, when there are many compact $\mathcal{V}_i \subseteq \mathcal{W}_i$ which do not have nonzero invariant vectors. This can be helpful if, for instance, there is a block j such that \mathcal{V}_j is trivial, but all other blocks are irreducible (for instance, as an alternative way to prove Lemma 12 in the case needed for semi-universality, i.e. when one of the blocks is one-dimensional). As a specific example, suppose that $\text{SU}(\mathcal{H}_j) \subseteq \mathcal{W}$ for $j \geq 2$, or, in other words, $\mathfrak{su}(\mathcal{H}_j) \subseteq \mathfrak{w}$. Then, by integrating over each of these,

$$(\mathcal{E}_m \circ \dots \circ \mathcal{E}_2)(X) = \begin{pmatrix} X_{11} & 0 & & 0 \\ 0 & \mathcal{E}_2(X_{22}) & & 0 \\ & & \ddots & \\ 0 & 0 & & \mathcal{E}_m(X_{mm}) \end{pmatrix} \in \mathfrak{w}. \quad (\text{F28})$$

But then, since $X_{ii} - \mathcal{E}_i(X_{ii}) \in \mathfrak{su}(\mathcal{H}_i) \subseteq \mathfrak{w}$ for $i \geq 2$, it follows that

$$\sum_{i=1}^m X_{ii} = \mathcal{E}_m \circ \dots \circ \mathcal{E}_2(X) + \sum_{i=2}^m (X_{ii} - \frac{\text{Tr } X_{ii}}{\dim \mathcal{H}_i} \Pi_i) \in \mathfrak{w}. \quad (\text{F29})$$

And therefore,

$$\tilde{X} = X - \sum_{i=1}^m X_{ii} = \begin{pmatrix} 0 & X_{12} & \dots & X_{1m} \\ X_{21} & 0 & \ddots & X_{2m} \\ \vdots & \ddots & \ddots & \\ X_{m1} & X_{m2} & & 0 \end{pmatrix} \in \mathfrak{w}. \quad (\text{F30})$$

Finally, we may isolate to, e.g., blocks 1 and 2:

$$\mathcal{E}_m \circ \dots \circ \mathcal{E}_3(\tilde{X}) = \begin{pmatrix} 0 & X_{12} & \\ X_{21} & 0 & \\ & & \ddots \end{pmatrix} \in \mathfrak{w}. \quad (\text{F31})$$

c. Weighting with characteristic function

Suppose that $\mathcal{V}_i \subseteq \mathcal{W}_i$ is compact and acts nontrivially and irreducibly. If we assume also that, as a representation of \mathcal{V}_i , $\mathcal{H}_i \otimes \mathcal{H}_i$ does not contain a subrepresentation isomorphic to \mathcal{H}_i or the trivial representation, then we can isolate without requiring any of the other subgroups \mathcal{V}_j . Note that the latter assumption is not satisfied for $\mathcal{V}_i \cong \text{SU}(2)$, since $\mathcal{H}_i \otimes \mathcal{H}_i$ contains a trivial subrepresentation for all of its irreps. More generally, this condition is not satisfied if the group \mathcal{V}_i is a subgroup of the orthogonal or symplectic unitary group.

Recall the notion of the characteristic function $\text{Tr } BU(g)$ of the operator B (see e.g. [50]). We weight the Haar measure by the real part of the characteristic function in the following construction.

Lemma 17. *Consider a Hilbert space with orthogonal decomposition $\mathcal{H} = \mathcal{H}_\lambda \oplus \mathcal{H}_\perp$ and associated Hermitian projectors Π_λ and Π_\perp , where \mathcal{H}_λ carries a nontrivial irreducible representation $U_\lambda(g)$ of a compact group G . Write $U(g) = U_\lambda(g) \oplus \mathbb{I}_\perp \in \text{U}(\mathcal{H})$ for the action on \mathcal{H} . If the representation $U_\lambda(g) \otimes U_\lambda(g)$ on $\mathcal{H}_\lambda \otimes \mathcal{H}_\lambda$ does not contain the trivial representation as a subrepresentation, or a subrepresentation isomorphic to \mathcal{H}_λ , then for any operators $A \in \mathcal{L}(\mathcal{H})$ and $B = \Pi_\lambda B \Pi_\lambda \in \mathcal{L}(\mathcal{H})$,*

$$\dim \mathcal{H}_\lambda \int dg \text{Tr}(BU(g)^\dagger)U(g)AU(g)^\dagger = B A \Pi_\perp \quad (\text{F32a})$$

$$\dim \mathcal{H}_\lambda \int dg \text{Tr}(BU(g))U(g)AU(g)^\dagger = \Pi_\perp A B, \quad (\text{F32b})$$

where dg is the normalized Haar measure on G .

Remark 8. If the irreducible group of unitaries $U_\lambda(G) = \{U_\lambda(g) : g \in G\}$ has a nontrivial center, then it always satisfies the condition that $\mathcal{H}_\lambda \otimes \mathcal{H}_\lambda$ does not contain a subrepresentation isomorphic to \mathcal{H}_λ . To see this, note that irreducibility implies that the center of this group should be in the form of phases $e^{i\theta} \mathbb{I}_\lambda \in U_\lambda(G)$. For two copies of nontrivial phases, i.e., the unitary $e^{i\theta} \mathbb{I}_\lambda \otimes e^{i\theta} \mathbb{I}_\lambda$, all eigenvalues are $e^{i2\theta}$, and therefore, there is no subspace of $\mathcal{H}_\lambda \otimes \mathcal{H}_\lambda$ in which this unitary acts as $e^{i\theta}(\mathbb{I}_\lambda \otimes \mathbb{I}_\lambda)$.

Proof. Write $A_{ij} = \Pi_i A \Pi_j$ for $i, j = \lambda, \perp$ and expand

$$U(g)AU(g)^\dagger = \sum_{i,j} U(g)A_{ij}U(g)^\dagger = U_\lambda(g)A_{\lambda\lambda}U_\lambda(g)^\dagger + U_\lambda(g)A_{\lambda\perp} + A_{\perp\lambda}U_\lambda(g)^\dagger + A_{\perp\perp}. \quad (\text{F33})$$

According to the Schur orthogonality theorem, for any pair of irreps λ and λ' of a compact group G

$$\dim \mathcal{H}_\lambda \int dg U_{\lambda'}(g) \otimes U_\lambda(g)^\dagger = \delta_{\lambda,\lambda'} \text{SWAP}_\lambda, \quad (\text{F34})$$

where $\delta_{\lambda,\lambda'}$ is 0 if λ and λ' are inequivalent representations, and is equal to 1 if $\lambda = \lambda'$, and where SWAP_λ is the linear operator on $\mathcal{H}_\lambda \otimes \mathcal{H}_\lambda$ which swaps product states, $|\psi\rangle \otimes |\varphi\rangle \mapsto |\varphi\rangle \otimes |\psi\rangle$. Schur orthogonality plus the assumption that $\mathcal{H}_\lambda \otimes \mathcal{H}_\lambda$ does not contain \mathcal{H}_λ or the trivial representation as subrepresentations implies

$$\int dg U_\lambda(g) \otimes U_\lambda(g) = 0 \quad (\text{F35a})$$

$$\int dg U_\lambda(g) \otimes U_\lambda(g) \otimes U_\lambda(g)^\dagger = 0. \quad (\text{F35b})$$

Similarly, switching $U_\lambda(g) \leftrightarrow U_\lambda(g)^\dagger$ in these expressions also obtains vanishing integrals. Finally, since $U_\lambda(g)$ is assumed nontrivial, $\int dg U_\lambda(g) = 0$. Thus the only term from Eq. (F33) that survives in the following has exactly one $U_\lambda(g)$ and one $U_\lambda(g)^\dagger$ in the integral:

$$\begin{aligned} \dim \mathcal{H}_\lambda \int dg \text{Tr}(BU(g))U(g)AU(g)^\dagger &= \dim \mathcal{H}_\lambda \int dg \text{Tr}(BU_\lambda(g))A_{\perp\lambda}U_\lambda(g)^\dagger \\ &= \dim \mathcal{H}_\lambda \text{Tr}_{(1)} \int dg (B \otimes A_{\perp\lambda})(U_\lambda(g) \otimes U_\lambda(g)^\dagger) \\ &= \text{Tr}_{(1)}(B \otimes A_{\perp\lambda}) \text{SWAP}_\lambda \\ &= A_{\perp\lambda}B = \Pi_\perp AB. \end{aligned} \quad (\text{F36})$$

where $\text{Tr}_{(1)}$ is the partial trace¹⁵. The other integral is performed similarly. \square

Supposing that the conditions of Lemma 17 are satisfied for the action of \mathcal{V}_i on \mathcal{H}_i , it follows that, for any $B = \Pi_i B \Pi_i \in \mathcal{L}(\mathcal{H})$,

$$\int_{\mathcal{V}_i} dU_i \text{Tr}(B^\dagger U_i + BU_i^\dagger) U_i X U_i^\dagger = \begin{pmatrix} 0 & & X_{1i} B^\dagger & \\ & \ddots & \vdots & \\ BX_{i1} & \cdots & 0 & \cdots \\ & & \vdots & \ddots \end{pmatrix} \in \mathfrak{w}, \quad (\text{F37})$$

since the coefficients $\text{Tr}(B^\dagger U_i + BU_i^\dagger)$ are real.

4. Application of strategies: 3 different proofs of Lemma 16

Proof of Lemma 16. If $\Pi_i A = 0$ then this is just the statement that $\mathbb{I} \in \mathcal{W}$. So suppose that $\Pi_i A \neq 0$. Any of the first three rows of Table IV may be used. In particular, the first two directly use Eq. (F8) and the fact that $\dim \mathcal{H}_i \geq 3$, so that $\text{span}_{\mathbb{R}} \text{SU}(\mathcal{H}_i)\Pi_i = \text{span}_{\mathbb{C}} \text{SU}(\mathcal{H}_i)\Pi_i = \mathcal{L}(\mathcal{H}_i)$ (see Eq. (F15b)).

¹⁵ This can also be shown using $\dim \mathcal{H}_\lambda \int dg \text{Tr}(BU_\lambda(g))U_\lambda(g)^\dagger = B$.

1. Recall the definition $D_U(X) = UXU^\dagger - X$. Choose $U_i \in \text{SU}(\mathcal{H}_i)$ such that $U_i - \Pi_i$ is invertible. Then $(U_i - \Pi_i)A \neq 0$. Then $D_{U_i}(A_{ii}) \in \mathfrak{su}(\mathcal{H}_i)$ and therefore $\tilde{A} := D_{U_i}(A) - D_{U_i}(A_{ii}) \in \mathfrak{w}$ is isolated to block i (see also Eq. (F19)). Recall

$$\{BX + XB^\dagger : B \in \text{span}_{\mathbb{R}} \mathcal{W}_i \Pi_i\} = \text{span}_{\mathbb{R}} \{U_i X U_i^\dagger : U_i \in \mathcal{W}_i\} \subseteq \mathfrak{w}. \quad (\text{re F8})$$

which holds under the assumption that X is anti-Hermitian, and inside \mathfrak{w} , and $\Pi_i X + X \Pi_i = X$, i.e., is isolated. Since $\dim \mathcal{H} \geq 3$, $\text{span}_{\mathbb{R}} \text{SU}(\mathcal{H}_i) \Pi_i = \mathcal{L}(\mathcal{H}_i)$, which means B can be arbitrary operator. Choosing $X = \tilde{A}$ we conclude that for any $\tilde{B} \in \mathcal{L}(\mathcal{H}_i)$,

$$\tilde{B} \tilde{A} + \tilde{A} \tilde{B}^\dagger = \tilde{B}(U_i - \Pi_i) A \Pi_\perp + \Pi_\perp A (U_i^\dagger - \Pi_i) \tilde{B}^\dagger \in \mathfrak{w}. \quad (\text{F38})$$

In particular, for any $B \in \mathcal{L}(\mathcal{H}_i)$ choose $\tilde{B} = B(U_i - \Pi_i)^{-1}$ to obtain Eq. (F12).

2. **Integration over uniform Haar measure:** By Eq. (F25), $(\mathcal{I} - \mathcal{E}_i)(A) \in \mathfrak{w}$. But note that $\Pi_i(\mathcal{I} - \mathcal{E}_i)(A)\Pi_i = A_{ii} - \mathcal{E}_i(A_{ii}) \in \mathfrak{su}(\mathcal{H}_i)$ because it is anti-Hermitian and traceless, so

$$\Pi_i A \Pi_\perp + \Pi_\perp A \Pi_i = \Pi_i A + A \Pi_i - 2\Pi_i A \Pi_i = (\mathcal{I} - \mathcal{E}_i)(A) - (A_{ii} - \mathcal{E}_i(A_{ii})) \in \mathfrak{w}. \quad (\text{F39})$$

Thus Eq. (F8) proves the claim.

3. **Weighting with characteristic function:** By Remark 8, since $\text{SU}(\mathcal{H}_i)$ has a nontrivial center, $\mathcal{H}_i \otimes \mathcal{H}_i$ does not contain a subrepresentation isomorphic to \mathcal{H}_i . Furthermore, $\dim \mathcal{H}_i \geq 3$ implies that $\mathcal{H}_i \otimes \mathcal{H}_i$ does not contain a trivial subrepresentation, since it is not self-dual as a representation of $\text{SU}(\mathcal{H}_i)$. Thus, since $\dim \mathcal{H}_i \geq 3$, $\text{SU}(\mathcal{H}_i)$ satisfies all the assumptions of Lemma 17, Eq. (F37) applies, and it follows immediately.

□