Tractable and Provably Efficient Distributional Reinforcement Learning with General Value Function Approximation

Taehyun Cho¹, Seungyub Han¹, Kyungjae Lee², Seokhun Ju¹, Dohyeong Kim¹, Jungwoo Lee^{1*}

Abstract

Distributional reinforcement learning improves performance by effectively capturing environmental stochasticity, but a comprehensive theoretical understanding of its effectiveness remains elusive. In this paper, we present a regret analysis for distributional reinforcement learning with general value function approximation in a finite episodic Markov decision process setting. We first introduce a key notion of *Bellman unbiasedness* for a tractable and exactly learnable update via *statistical functional dynamic programming*. Our theoretical results show that approximating the infinite-dimensional return distribution with a finite number of moment functionals is the only method to learn the statistical information unbiasedly, including nonlinear statistical functionals. Second, we propose a provably efficient algorithm, SF-LSVI, achieving a regret bound of $\tilde{O}(d_E H^{\frac{3}{2}}\sqrt{K})$ where H is the horizon, K is the number of episodes, and d_E is the eluder dimension of a function class.

1 Introduction

Distributional reinforcement learning (DistRL) [7, 13, 30, 41] is an advanced approach to reinforcement learning (RL) that focuses on the entire probability distribution of returns rather than solely on the expected return. By considering the full distribution of returns, distRL provides deeper insight into the uncertainty of each action, such as the mode or median. This framework enables us to make safer and more effective decisions that account for various risks [14, 22, 29, 46], particularly in complex real-world situations, such as robotic manipulation [10], neural response [36], stratospheric balloon navigation [8], algorithm discovery [19], and several game benchmarks [6, 34].

Although distRL has a solid theoretical basis and demonstrates effective empirical performance, its efficiency and tractability are still not fully understood. In practice, distributions contain an infinite amount of information, and we should resort to approximations using a finite number of parameters or statistical functionals, such as categorical [7] and quantile representations [16]. However, not all statistical functionals can be learned through the Bellman operator, as the meaning of statistical functionals is not typically preserved after updates. For example, the median is not preserved under the Bellman updates, as the median of a mixture of two distributions does not necessarily equate to the mixture of their medians. Hence, a distinct analysis is required to determine whether there exists a corresponding Bellman operator for each statistical functional that ensures commutativity. Rowland et al. [41] defined *Bellman closedness* as a property of statistical functionals that can be exactly learned by the existence of a corresponding Bellman operator.

^{*}Corresponding author

Preprint. Under review.



Figure 1: Venn-Diagram of Statistical Functional Classes. The diagram shows the category of statistical functional. (Yellow) Within the linear statistical functional class, Rowland et al. [41] demonstrated that the solution of Bellman closedness is uniquely represented by polynomial exponential functionals. (Red) We extend this concept by introducing Bellman unbiasedness, which not only covers moment functionals but also central moment functionals from the broader class including nonlinear statistical functionals. (A) Maximum and minimum functionals are Bellman closed, while those are not unbiasedly estimatable. (B) Median and quantile functionals are neither Bellman closed and unbiased, highlighting that they are not proper to encode the distribution.

At this point, we take a closer look at the distributional Bellman update and revisit what additional properties of statistical functionals beyond Bellman closedness are needed to construct tractable algorithms. In doing so, we identify and provide the following challenges that arise when statistical functionals are used to express the update process:

- Representing a mixture distribution with a finite, fixed number of parameters inevitably leads to approximation errors during the update process. For example, when expressing the mixture of two distributions, each represented by N parameters, compressing 2N into N parameters in the mixture results in inevitable information loss.
- Due to the unknown nature of the transition probability $\mathbb{P}(\cdot|s, a)$, the target distribution is estimated by sampling the next state s'. For a tractable algorithm, it is crucial that the statistical functionals of the target distribution can be unbiasedly estimated using the statistical functionals from the sampled distribution.

In this paper, we describe the *statistical functional* and *sketch* of the distribution, and introduce the additional approximation errors that can occur with the distRL method. We define the notion of *Bellman unbiasedness*, a property for representing the distribution tractably and prove that the moment functional is the only solution that satisfies this property, including *nonlinear* statistical functionals. Based on this property, we propose a provably efficient statistical functional reinforcement learning algorithm with general value function approximation, denoted as SF-LSVI. We will discuss the inherent issues of *distributional Bellman completeness*, a structural assumption previously defined in the literature [11, 49], and investigate the benefits of redesigning this concept using a sketch.

In summary, our main contributions are as follows:

- Introduce a key property of Bellman unbiasedness for collections of statistical functionals (sketch) and show that the moment functionals are the unique structure that can be unbiasedly estimated and exactly learned from distributions, including nonlinear statistical functionals.
- Revisit the existing concept of distributional Bellman Completeness through a statistical functional lens. We enhance the tractability by using moment functional, that the agent can only access a finite-dimensional embedding space. This removes the misspecification errors commonly encountered when discretizing infinite-dimensional distributions.
- Propose a tractable and provably efficient distributional RL algorithm called SF-LSVI, achieving a tightest regret upper bound $\tilde{O}(d_E H^{\frac{3}{2}}\sqrt{K})^{-1}$ in distributional framework. Our

¹We ignore poly-log terms in H, S, A, K in the $\tilde{O}(\cdot)$ notation.

Table 1: Comparison for different methods under distributional RL framework. \mathcal{H} represents a subspace of infinite-dimensional space \mathcal{F}^{∞} . To bound the eluder dimesion d_E , Wang et al. [49] and Chen et al. [11] assumed the discretized reward MDP.

Algorithm	Regret	Eluder dimension d_E	Bellman Completeness	MDP assumption	Tractability of Distribution Estimation
0-DISCO [49]	$\tilde{\mathcal{O}}(\mathrm{poly}(d_E H)\sqrt{K})$	$\dim_E(\mathcal{H},\epsilon)$	distributional BC	discretized reward, small-loss bound	×
V-EST-LSR [11]	$\tilde{\mathcal{O}}(d_E H^2 \sqrt{K})$ ²	$\dim_E(\mathcal{H},\epsilon)$	distributional BC	discretized reward, lipschitz continuity	×
SF-LSVI [Ours]	$\tilde{\mathcal{O}}(d_E H^{\frac{3}{2}}\sqrt{K})$	$\dim_E(\mathcal{F}^N,\epsilon)$	statistical functional BC	none	1

framework yields a better regret bound with a weaker structural assumption compared to prior results.

2 Related Work

Distributional RL. In classical RL, the Bellman equation, which is based on expected returns, has a closed-form expression. However, it is questionable whether any statistical functionals of return distribution always have their corresponding closed-form expressions. Rowland et al. [41] introduced the notion of *Bellman Closedness* for collections of statistical functionals that can be updated in a closed form via Bellman update. They showed that the only Bellman-closed statistical functionals in the discounted setting are the moments $\mathbb{E}_{Z \sim \eta}[Z^k]$. More recently, Marthe et al. [35] proposed a general framework for distRL, where the agent plans to maximize their own utility functionals instead of expected return and formalized the property of *Bellman Optimizability*. They proved that the only W_1 -continuous and linear Bellman Optimizable statistical functionals are shown to be exponential utilities $\frac{1}{\lambda} \log \mathbb{E}_{Z \sim \eta} [\exp(\lambda Z)]$ in the undiscounted setting.

In practice, C51 [7] and QR-DQN [16] are notable distributional RL algorithms where the convergence guarantees of sampled-based algorithms are proved [40, 42]. Dabney et al. [15] expanded the class of policies on arbitrary distortion risk measures by taking the based distribution non-uniformly and improve the sample efficiency from their implicit representation of the return distribution. Cho et al. [12] highlighted the drawbacks of optimistic exploration in distRL, introducing a randomized exploration method that perturbs the return distribution when the agent selects their next action.

RL with General Value Function Approximation. Regret bounds have been studied for a long time in RL, across various domains such as bandit [1, 32, 43], tabular RL [3, 26, 28, 38, 39], and linear function approximation [27, 51, 52]. In recent years, deep RL has shown significant performance using deep neural networks as function approximators, and attempts have been made to analyze whether it is efficient in terms of general function approximation. Wang et al. [50] established a provably efficient RL algorithm with general value function approximation based on the eluder dimension d_E [43] and achieves a regret upper bound of $\tilde{O}(\text{poly}(d_EH)\sqrt{K})$. To circumvent the intractability from computing the upper confidence bound, Ishfaq et al. [25] injected the stochasticity on the training data and get the optimistic value function instead of upper confidence bound, enhancing computationally efficiency. Beyond risk-neutral setting, several prior works have shown regret bounds under risk-sensitive objectives (e.g., entropic risk [20, 33], CVaR [5]), which align with our approach in that they are built on a distribution framework. Liang and Luo [33] achieved the regret upper bound of $\tilde{O}(\exp(H)\sqrt{|\mathcal{S}|^2|\mathcal{A}|H^2K})$ and the lower bound of $\Omega(\exp(H)\sqrt{|\mathcal{S}||\mathcal{A}|HK})$ in tabular setting.

DistRL with General Value Function Approximation. Recently, only few efforts have aimed to bridge the gap between two fields. Wang et al. [49] proposed a distributional RL algorithm, 0-DISCO, which enjoys small-loss bound by using a log-likelihood objective. Similarly, Chen et al. [11] provided a risk-sensitive reinforcement learning framework with static lipschitz risk measure. While these studies analyze within a distributional framework, they do not address the intractability

²In Chen et al. [11], the regret bound is written as $\tilde{O}(d_E L_{\infty}(\rho) H \sqrt{K})$, where $L_{\infty}(\rho)$ represents the lipschitz constant of the risk measure ρ , i.e., $|\rho(Z) - \rho(Z')| \le L_{\infty}(\rho) ||F_Z - F_{Z'}||_{\infty}$. Since $L_{\infty}(\rho) \ge H$ in risk-neutral setting, we translate the regret bound into $\tilde{O}(d_E H^2 \sqrt{K})$.

of implementation in infinite-dimensional space of distributions. In contrast, our approach focuses on a statistical functional framework, providing a detailed comparison with other distRL methods as shown in Table 1.

3 Preliminaries

Episodic MDP. We consider a episodic Markov decision process which is defined as a $\mathcal{M} = (S, \mathcal{A}, H, \mathbb{P}, r)$ characterized by state space S, action space \mathcal{A} , horizon length H, transition kernels $\mathbb{P} = \{\mathbb{P}_h\}_{h \in [H]}$, and reward $r = \{r_h\}_{h \in [H]}$ at step $h \in [H]$. The agent interacts with the environment across K episodes. For each $k \in [K]$ and $h \in [H]$, $\mathbb{H}_h^k = (s_1^1, a_1^1, \ldots, s_H^1, a_H^1, \ldots, s_h^k, a_h^k)$ represents the history up to step h at episode k. We assume the reward is bounded by [0, 1] and the agent always transit to terminal state s_{end} at step H + 1 with $r_{H+1} = 0$.

Policy and Value Functions. A (deterministic) policy π is a collection of H functions $\{\pi_h : S \to \mathcal{A}\}_{h=1}^H$. Given a policy π , a step $h \in [H]$, and a state-action pair $(s, a) \in S \times \mathcal{A}$, the Q and V-function are defined as $Q_h^{\pi}(s, a)(: S \times \mathcal{A} \to \mathbb{R}) := \mathbb{E}_{\pi} \left[\sum_{h'=h}^{H} r_{h'}(s_{h'}, a_{h'}) \mid s_h = s, a_h = a \right]$ and $V_h^{\pi}(s)(: S \to \mathbb{R}) := \mathbb{E}_{\pi} \left[\sum_{h'=h}^{H} r_{h'}(s_{h'}, a_{h'}) \mid s_h = s \right]$.

Random Variables and Distributions. For a sample space Ω , we extend the definition of the Q-function into a random variable and its distribution,

$$Z_h^{\pi}(s,a)(:\mathcal{S}\times\mathcal{A}\times\Omega\to\mathbb{R}):=\sum_{h'=h}^{H}r_{h'}(s_{h'},a_{h'})\mid s_h=s, a_h=a, a_{h'}=\pi_{h'}(s_{h'}),$$
$$\eta_h^{\pi}(s,a)(:\mathcal{S}\times\mathcal{A}\to\mathscr{P}(\mathbb{R})):=\operatorname{law}(Z_h^{\pi}(s,a)).$$

Analogously, we extend the definition of V-function by introducing a bar notation.

$$\bar{Z}_{h}^{\pi}(s)(:\mathcal{S}\times\Omega\to\mathbb{R})\coloneqq\sum_{h'=h}^{H}r_{h'}(s_{h'},a_{h'})\mid s_{h}=s,a_{h'}=\pi_{h'}(s_{h'}),$$
$$\bar{\eta}_{h}^{\pi}(s)(:\mathcal{S}\to\mathscr{P}(\mathbb{R}))\coloneqq\operatorname{law}(\bar{Z}_{h}^{\pi}(s)).$$

Note that $\bar{Z}_{h}^{\pi}(s) = Z_{h}^{\pi}(s,\pi(s))$ and $\bar{\eta}_{h}^{\pi}(s) = \eta_{h}^{\pi}(s,\pi(s))$. We use π^{\star} to denote an optimal policy and denote $V_{h}^{\star}(s) = V_{h}^{\pi^{\star}}(s), Q_{h}^{\star}(s,a) = Q_{h}^{\pi^{\star}}(s,a), \eta_{h}^{\star}(s,a) = \eta_{h}^{\pi^{\star}}(s,a),$ and $\bar{\eta}_{h}^{\star}(s) = \bar{\eta}_{h}^{\pi^{\star}}(s)$. For notational simplicity, we denote the expectation over transition, $[\mathbb{P}_{h}V_{h+1}^{\pi}](s,a) = \mathbb{E}_{s'\sim\mathbb{P}_{h}(\cdot|s,a)}V_{h+1}^{\pi}(s'), [\mathbb{P}_{h}\bar{Z}_{h+1}^{\pi}](s,a) = \mathbb{E}_{s'\sim\mathbb{P}_{h}(\cdot|s,a)}\bar{Z}_{h+1}^{\pi}(s'),$ and $[\mathbb{P}_{h}\bar{\eta}_{h+1}^{\pi}](s,a) = \mathbb{E}_{s'\sim\mathbb{P}_{h}(\cdot|s,a)}\bar{\eta}_{h+1}^{\pi}(s').$ ³ For brevity, we refer to $\bar{\eta}^{\pi}$ simply as $\bar{\eta}$.

In the episodic MDP, the agent aims to learn the optimal policy through a fixed number of interactions with the environment across a number of episodes. At the beginning of each episode $k (\in [K])$, the agent starts at the initial state s_1^k and choose a policy π^k . In step $h (\in [H])$, the agent observes $s_h^k (\in S)$, takes an action $a_h^k (\in A) \sim \pi_h^k (\cdot | s_h^k)$, receives a reward $r_h(s_h^k, a_h^k)$, and the environment transits to the next state $s_{h+1}^k \sim \mathbb{P}_h(\cdot | s_h^k, a_h^k)$. Finally, we measure the suboptimality of an agent by its regret, which is the accumulated difference between the ground truth optimal and the return received from the interaction. The regret after K episodes is defined as $\text{Reg}(K) = \sum_{k=1}^{K} V_1^{\star}(s_1^k) - V_1^{\pi^k}(s_1^k)$.

Distributional Bellman Optimality Equation. Recall that η_h^* satisfies the distributional Bellman optimality equation:

$$\eta_{h}^{\star}(s,a) = (\mathcal{T}_{h}\eta_{h+1}^{\star})(s,a) = \mathbb{E}_{s' \sim \mathbb{P}_{h}(\cdot|s,a),a' \sim \pi_{h}^{\star}(\cdot|s')}[(\mathcal{B}_{r_{h}})_{\#}\eta_{h+1}^{\star}(s',a')] \\ = (\mathcal{B}_{r_{h}})_{\#}[\mathbb{P}_{h}\eta_{h+1}^{\star}](s,a)$$

where $\mathcal{B}_r : \mathbb{R} \to \mathbb{R}$ is defined by $\mathcal{B}_r(x) = r + x$, and $g_{\#}\eta \in \mathscr{P}(\mathbb{R})$ is the pushforward of the distribution η through g, i.e., $g_{\#}\eta(A) = \eta(g^{-1}(A))$ for any Borel set $A \subseteq \mathbb{R}$.

³Note that $\mathbb{E}_{s' \sim \mathbb{P}_h(\cdot|s,a)} \bar{\eta}_{h+1}^{\pi}(s')$ is a mixture distribution.



Figure 2: Illustrative representation of sketch-based Bellman updates for a mixture distribution. Instead of updating the distributions directly, each sampled distribution is embedded through a sketch ψ (e.g., mean μ , quantile q_i). The transformation ϕ_{ψ} aims to compress the mixture distribution into the same number of parameters, ensuring unbiasedness to prevent information loss.

Additional Notations. For a given N, we denote an N-dimensional function class $\mathcal{F}^N := \mathcal{F}^{(1)} \times \cdots \times \mathcal{F}^{(N)} \subseteq \left\{ f = [f^{(1)}, \cdots, f^{(N)}] : \mathcal{S} \times \mathcal{A} \to \mathbb{R}^N \right\}$. Given a dataset $\mathcal{D} = \{(s_t, a_t, [z_t^{(1)}, \dots, z_t^{(N)}])\}_{t=1}^{|\mathcal{D}|} \subseteq \mathcal{S} \times \mathcal{A} \times \mathbb{R}^N$, a set of state-action pairs $\mathcal{Z} = \{(s_t, a_t)\}_{t=1}^{|\mathcal{Z}|} \subseteq \mathcal{S} \times \mathcal{A}$ and for a function $f : \mathcal{S} \times \mathcal{A} \to \mathbb{R}^N$, we define the norm $\|f^{(n)}\|_{\infty}, \|f\|_{\infty,1}, \|f\|_{\mathcal{D}}, \|f\|_{\mathcal{Z}}$ as written in Appendix A. For a set of (vector-valued) functions $\mathcal{F}^N \subseteq \{f : \mathcal{S} \times \mathcal{A} \to \mathbb{R}^N\}$, the width function of (s, a) is defined as $w^{(n)}(\mathcal{F}^N, s, a) \coloneqq \max_{f, q \in \mathcal{F}^N} |f^{(n)}(s, a) - g^{(n)}(s, a)|$.

4 Statistical Functionals in Distributional RL

4.1 Bellman Unbiasedness

We revisit the definitions of statistical functionals and illustrate the property of *Bellman closedness* described in Bellemare et al. [9]. Then we introduce another pivotal property, *Bellman unbiasedness*, required for learning in terms of statistical functionals, rather than approximated distribution.

Definition 4.1 (Statistical functionals, Sketch; [9]). A statistical functional is a mapping from a probability distribution to a real value $\psi : \mathscr{P}(\mathbb{R}) \to \mathbb{R}$. A sketch is a vector-valued function $\psi_{1:N} : \mathscr{P}(\mathbb{R}) \to \mathbb{R}^N$ specified by an N-tuple where each component is a statistical functional,

$$\psi_{1:N}(\cdot) = (\psi_1(\cdot), \cdots, \psi_N(\cdot))$$

We denote the domain of sketch as $\mathscr{P}_{\psi_{1:N}}(\mathbb{R})$ and its image as $I_{\psi_{1:N}} = \{\psi(\bar{\eta}) : \bar{\eta} \in \mathscr{P}_{\psi_{1:N}}(\mathbb{R})\}$. We further extend to state return distribution functions $\psi_{1:N}(\bar{\eta}) = (\psi_{1:N}(\bar{\eta}(s)) : s \in \mathcal{S})$.

Definition 4.2 (Bellman closedness; [41]). A sketch $\psi_{1:N}$ is **Bellman closed** if there exists an operator $\mathcal{T}_{\psi_{1:N}} : I_{\psi_{1:N}}^{S} \to I_{\psi_{1:N}}^{S}$ such that

$$\psi_{1:N}(\mathcal{T}\bar{\eta}) = \mathcal{T}_{\psi_{1:N}}\psi_{1:N}(\bar{\eta}) \quad \text{for all } \bar{\eta} \in \mathscr{P}(\mathbb{R})^{\mathcal{S}}$$

which is closed under the distributional Bellman operator $\mathcal{T}: \mathscr{P}(\mathbb{R})^{\mathcal{S}} \to \mathscr{P}(\mathbb{R})^{\mathcal{S}}$.

Bellman closedness is the property that a sketch are exactly learnable when updates are performed from the infinite-dimensional distribution space to the finite-dimensional embedding space. Notably, Rowland et al. [41] showed that the only finite linear statistical functionals that are Bellman closed are given by the collections of statistical functionals where its linear span is equal to the set of exponential-polynomial functionals where ψ_0 is the constant functional equal to 1.

Remark 4.3. *Rowland et al.* [41] discussed only linear statistical functionals when defining a sketch, leaving open questions about the Bellman closedness of nonlinear statistical functionals such as variance, central moments, or quantiles. However, it is noteworthy that nonlinear statistical functionals, such as maximum or minimum, can also be Bellman closed. In their proof, there might be an ambiguity regarding the assumption that the corresponding sketch Bellman operator for quantile is linear. In this paper, we show the non-existence of sketch Bellman operator for quantile and discuss in detail in Appendix B.1.

$$\begin{array}{cccc} \eta & \longrightarrow & \mathcal{T}\eta & \longrightarrow & \psi(\mathcal{T}\eta) & & \eta & \longrightarrow & \mathcal{T}\eta & \longrightarrow & \psi(\mathcal{T}\eta) \\ \downarrow & & & \downarrow & & \uparrow \\ \psi(\eta) & \longrightarrow & \mathcal{T}_{\psi}\psi(\eta) & & & \psi(\eta) & & & \psi(\eta) & & \\ & & & \psi(\eta) & \xrightarrow{\text{sample}} & \hat{\mathcal{T}}_{\psi}\psi(\eta) & \xrightarrow{\text{unbiasedly}} & \mathbb{E}_{\mathbb{P}}[\phi_{\psi}\circ\hat{\mathcal{T}}_{\psi}\psi(\eta)] \\ & & & \text{estimate} \end{array}$$

Figure 3: (Left) Bellman Closedness. (**Right**) Bellman Unbiasedness. The above path represents an ideal distributional Bellman update. Due to the infinite-dimensionality, the update process should be represented by using a finite-dimensional embedding (sketch) ψ . Since the transition kernel \mathbb{P} is unknown, the below path describes that the implementation should sample the next state and update by using $\hat{\mathcal{T}}_{\psi}$ with the empirical transition kernel $\hat{\mathbb{P}}$. A sketch ψ is Bellman unbiased if $\hat{\mathcal{T}}_{\psi} \circ \psi$ can unbiasedly estimate $\psi \circ \mathcal{T}$ through some transformation ϕ_{ψ} .

Theorem 4.4. Quantile functional cannot be Bellman closed under any additional sketch.

We investigate whether there exists another sketch that is both nonlinear and Bellman closed, and introduce additional conditions under which the Bellman closed property might be missing. During the implementation phase, it is important to note that the agent does not have access to the transition kernel. Specifically, the agent can only access the empirical transition kernel $\hat{\mathbb{P}}$ which is derived from a finite number of sample estimates. This limitation implies that the operator should be treated as an empirical operator $\hat{\mathcal{T}}_{\psi}$, rather than \mathcal{T}_{ψ} . Therefore, we introduce a new notion of *Bellman unbiasedness* that allows us to unbiasedly estimate the expected distribution $(\mathcal{B}_r)_{\#}\mathbb{E}_{s' \sim \mathbb{P}(\cdot|s,a)}[\bar{\eta}(s')]$, which is a mixture by transitions, from the sample distribution $(\mathcal{B}_r)_{\#}\bar{\eta}(s')$.

Definition 4.5 (Bellman unbiasedness). A sketch ψ is **Bellman unbiased** if there exists an integer kand a corresponding vector-valued estimator $\phi_{\psi} = \phi_{\psi}(\psi(x_1), \dots, \psi(x_k)) : (I_{\psi}^{S})^k \to I_{\psi}^{S}$ where the sketch of expected distribution $(\mathcal{B}_r)_{\#} \mathbb{E}_{s' \sim \mathbb{P}(\cdot|s,a)}[\bar{\eta}(s')]$ can be unbiasedly estimated by ϕ_{ψ} using the k samples from the sketch of the sample distribution $(\mathcal{B}_r)_{\#}\bar{\eta}(s')$, i.e.,

$$\mathbb{E}_{s_1',\cdots,s_k'\sim\mathbb{P}(\cdot|s,a)}\left[\phi_{\psi}\left(\underbrace{\psi\Big((\mathcal{B}_r)_{\#}\bar{\eta}(s_1')}_{\hat{\mathcal{T}}_{\psi}\psi(\eta)}\right),\cdots,\psi\Big((\mathcal{B}_r)_{\#}\bar{\eta}(s_k')\Big)\right)\right]=\psi\Big((\mathcal{B}_r)_{\#}\mathbb{E}_{s'\sim\mathbb{P}(\cdot|s,a)}[\bar{\eta}(s')]\Big).$$

Bellman unbiasedness is another natural definition, similar to Bellman closedness, which takes into account a finite number of samples for the transition. As a simple example, median functional is not Bellman unbiased as the sample median is not an unbiased estimate of the median functional. In general, there is no unbiased estimator for the median, implying that the median functional cannot be unbiasedly estimated within an embedding space represented by a finite number of statistical functionals. On the other hand, variance functional is a nonlinear statistical functional but can be unbiasedly estimated. Then, the following question naturally arises;

"Which sketches are unbiasedly estimatable under the sketch-based Bellman update?"

The following lemma answers this question.

Lemma 4.6. Let $F_{\bar{\eta}}$ be a CDF of the probability distribution $\bar{\eta} \in \mathscr{P}_{\psi}(\mathbb{R})^{S}$. Then a sketch is Bellman unbiased if and only if the sketch is homogeneous over $\mathscr{P}_{\psi}(\mathbb{R})^{S}$ of degree k, i.e., there exists some vector-valued function $h = h(x_1, \dots, x_k) : \mathcal{X}^k \to \mathbb{R}^N$ such that

$$\psi(\bar{\eta}) = \int \cdots \int h(x_1, \cdots, x_k) dF_{\bar{\eta}}(x_1) \cdots dF_{\bar{\eta}}(x_k).$$

Lemma 4.6 states that in statistical functional dynamic programming, the unbiasedly estimatable embedding of a distribution can only be structured in the form of functions that are homogeneous of finite degree. (e.g., variance functional is a homogeneous of degree 2 [23]). Taking this concept further and combining it with the results on Bellman closedness, we prove that even when including a nonlinear statistical functional, the only sketch that can be unbiasedly estimated in a finite-dimensional embedding space is the moment functional.

Theorem 4.7. The only finite statistical functionals that are both Bellman unbiased and closed are given by the collections of ψ_1, \ldots, ψ_N where its linear span $\{\sum_{n=0}^N \alpha_n \psi_n | \alpha_n \in \mathbb{R}, \forall N\}$ is equal to the set of exponential polynomial functionals $\{\eta \to \mathbb{E}_{Z \sim \eta}[Z^l \exp(\lambda Z)] | l = 0, 1, \ldots, L, \lambda \in \mathbb{R}\}$, where ψ_0 is the constant functional equal to 1. In discount setting, it is equal to the linear span of the set of moment functionals $\{\eta \to \mathbb{E}_{Z \sim \eta}[Z^l] | l = 0, 1, \ldots, L, \lambda \in \mathbb{R}\}$,

Compared to Rowland et al. [41], our results extend beyond linear statistical functionals to include unbiasedly estimatable nonlinear statistical functionals, showing the uniqueness of the moment functional. Our theoretical results not only show that high-order central moments such as variance or skewness are exactly learnable, but also reveal that other nonlinear statistical functionals like median or quantiles inevitably involve approximation errors due to biased estimations. In Section 4.2, we emphasize that unbiasedly estimable sketches play an important role in mitigating misspecification errors, thereby enabling us to achieve sublinear regret.

4.2 Statistical Functional Bellman Completeness

We consider distributional reinforcement learning with value function approximation. For successful TD learning, reinforcement learning with function approximation commonly requires the assumption, *Bellman Completeness*, that after applying Bellman operator, the output lies in the function class \mathcal{F} [4, 25, 50]. On the other hand, our approach receives a tuple of function class $\mathcal{F}^N \subseteq \{f : S \times \mathcal{A} \rightarrow \mathbb{R}^N\}$ as input to represent *N*-moment of distribution. Building on this, we assume that for any $\bar{\eta} : S \rightarrow \mathcal{P}([0, H])$, the sketch of target function that results from applying sketch Bellman operator lies in the function class \mathcal{F}^N .

In the seminal works, Wang et al. [49] and Chen et al. [11] assumed that the function class $\mathcal{H} \subseteq \{\eta : S \times \mathcal{A} \to \mathscr{P}([0, H])\}$ follows the *distributional Bellman Completeness* assumption, i.e., if $\eta \in \mathcal{H}$ for all $\pi, h \in [H]$, then $\mathcal{T}_h^{\pi} \eta \in \mathcal{H}$. However, since the distributional Bellman operator mixes distributions for the next state transition, it implies that a function class \mathcal{H} must be closed under mixture. Due to this closedness under mixture, both previous studies assumed a discretized reward MDP, which requires the prior knowledge that outcomes of return distribution are additively closed. In general, representing a distribution with a fixed number of parameters inevitably increases the required number of parameters due to the emergence of a mixture distribution. Consequently, trying to represent it with a limited number of parameters leads to approximating the learning outcome, thereby causing *model misspecification* which generally introduces an additional linear regret. Precisely, a regret lower bound with model misspecification error ζ is known as $\Omega(\zeta K)$ in a linear bandit setting [52]. Therefore, assuming Bellman Completeness on the distribution space is invalid as it significantly constrains the MDP structure or causes linear regret.

To circumvent this issue, we revisit the assumption of distributional Bellman completeness through the statistical functional lens. We propose a new framework that matches a finite number of all statistical functionals to the target function, rather than the entire distribution itself.

Assumption 4.8 (Statistical Functional Bellman Completeness). For any distribution $\bar{\eta} : S \to \mathscr{P}([0, H])$ and $h \in [H]$, there exists $f_{\bar{\eta}} \in \mathcal{F}^N$ which satisfies

$$f_{\bar{\eta}}(s,a) = \psi_{1:N} \big((\mathcal{B}_{r_h})_{\#} [\mathbb{P}_h \bar{\eta}](s,a) \big) \quad \forall (s,a) \in \mathcal{S} \times \mathcal{A}$$

Assumption 4.9 (Model Misspecification). There exists a set of function \mathcal{F}^N and a real number $\zeta > 0$, such that for any $\bar{\eta} : S \to \mathscr{P}([0, H])$ and $h \in [H]$, there exists $f_{\bar{\eta}} \in \mathcal{F}^N$ which satisfies

$$\max_{(s,a)\in\mathcal{S}\times\mathcal{A}}\left\|f_{\bar{\eta}}(s,a)-\psi_{1:N}((\mathcal{B}_{r_h})_{\#}[\mathbb{P}_h\bar{\eta}](s,a))\right\|\leq \zeta\quad\forall (s,a)\in\mathcal{S}\times\mathcal{A}.$$

We call ζ the **misspecification error**.

5 Algorithms

In this section, we propose SF-LSVI for distributional RL framework with general value function approximation. Leveraging the result from Theorem 4.7, we introduce a moment least square regression. This allows us to capture a finite set of moment information from the distribution, which can be unbiasedly estimated, thereby leading to the *truncated moment problem*. Unlike previous work [11, 49] that estimates in infinite-dimensional distribution spaces, our method enables tractable distribution estimation in finite-dimensional embedding spaces without approximation error.

Algorithm 1 Statistical Functional Least Square Value Iteration (SF-LSVI(δ))

Input: failure probability $\delta \in (0, 1)$ and the number of episodes K 1: for episode k = 1, 2, ..., K do 2: Receive initial state s_1^k Initialize $\psi_{1:N}(\bar{\eta}_{H+1}^k(\cdot)) \leftarrow \mathbf{0}^N$ for step $h = H, H-1, \dots, 1$ do 3: 4: $\mathcal{D}_{h}^{k} \leftarrow \left\{ s_{h'}^{\tau}, a_{h'}^{\tau}, \psi_{1:N} \left((\mathcal{B}_{r_{h'}}^{\tau})_{\#} \bar{\eta}_{h+1}^{k} (s_{h'+1}^{\tau}) \right) \right\}_{(\tau,h') \in [k-1] \times [H]}$ // Data collection 5:
$$\begin{split} \tilde{f}_{h,\bar{\eta}}^k &\leftarrow \arg\min_{f \in \mathcal{F}^N} \|f\|_{\mathcal{D}_h^k} \\ b_h^k(\cdot,\cdot) &\leftarrow w^{(1)}((\mathcal{F}^N)_h^k,\cdot,\cdot) \end{split}$$
// Distribution Estimation 6: 7: $\begin{aligned} b_h(\cdot, \cdot) &\leftarrow \operatorname{wid}(U \to f_h, \cdot, \cdot) \\ Q_h^k(\cdot, \cdot) &\leftarrow \min\{(\tilde{f}_{h,\bar{\eta}}^k)^{(1)}(\cdot, \cdot) + b_h^k(\cdot, \cdot), H\} \\ \pi_h^k(\cdot) &= \arg\max_{a \in \mathcal{A}} Q_h^k(\cdot, a), V_h^k(\cdot) = Q_h^k(\cdot, \pi_h^k(\cdot)) \qquad // \operatorname{Optimisti} \\ \psi_1\Big(\eta_h^k(\cdot, \cdot)\Big) &\leftarrow Q_h^k(\cdot, \cdot), \ \psi_{2:N}\Big(\eta_h^k(\cdot, \cdot)\Big) \leftarrow \Big(\min\{(\tilde{f}_{h,\bar{\eta}}^k)^{(n)}(\cdot, \cdot), H\}\Big)_{n \in [2:N]} \end{aligned}$ 8: // Optimistic planning 9: 10: $\psi_1\left(\bar{\eta}_h^k(\cdot,\cdot)\right) \leftarrow V_h^k(\cdot), \ \psi_{2:N}\left(\bar{\eta}_h^k(\cdot)\right) \leftarrow \psi_{1:N}\left(\eta_h^k(\cdot,\pi_h^k(\cdot))\right)_{n \in [2:N]}$ 11: $\begin{array}{l} \text{for } h=1,2,\ldots,H \text{ do} \\ \text{Take action } a_h^k \leftarrow \pi_h^k(s_h^k) \end{array}$ 12: 13: Observe reward $r_h^k(s_h^k, a_h^k)$ and get next state s_{h+1}^k . 14:

Overview. At the beginning of episode $k \in [K]$, we maintain all previous samples $\{(s_{h'}^{\tau}, a_{h'}^{\tau}, r_{h'}^{\tau})\}_{(\tau,h')\in[k-1]\times[H]}$ and initialize a sketch $\psi_{1:N}(\bar{\eta}_{H+1}^k(\cdot)) = \mathbf{0}^N$. For each step $h = H, \ldots, 1$, we compute the normalized sample moments of target distribution $\{(\mathcal{B}_{r_{h'}^{\tau}})_{\#}\bar{\eta}_{h+1}^k(s_{h'+1}^{\tau})\}_{h'\in[H]}$ with the help of binomial theorem,

$$\psi_n\Big((\mathcal{B}_{r_{h'}})_{\#}\bar{\eta}_h(s_{h'+1}^{\intercal})\Big) = \mathbb{E}[(\bar{Z}_{h+1}^k(s_{h'+1}^{\intercal}) + r_{h'}^{\intercal})^n]/H^{n-1}$$
$$= \sum_{n'=0}^n H^{n'}\psi_{n'}\Big(\bar{\eta}_h(s_{h'+1}^{\intercal})\Big)(r_{h'}^{\intercal})^{n-n'}/H^{n-1}$$

and iteratively solve the N-moment least square regression

$$\tilde{f}_{h,\bar{\eta}}^{k} \leftarrow \arg\min_{f \in \mathcal{F}} \sum_{\tau=1}^{k-1} \sum_{h'=1}^{H} \left(\sum_{n=1}^{N} f^{(n)}(s_{h'}^{\tau}, a_{h'}^{\tau}) - \psi_n \left((\mathcal{B}_{r_{h'}^{\tau}})_{\#} \bar{\eta}_{h+1}^{k}(s_{h'+1}^{\tau}) \right) \right)^2$$

based on the dataset \mathcal{D}_{h}^{k} which contains the sketch of temporal target $\psi_{1:N}\left((\mathcal{B}_{r_{h}^{\tau}})_{\#}\bar{\eta}_{h+1}^{k}(s_{h'+1}^{\tau})\right)$. Then we define $Q_{h}^{k}(\cdot, \cdot) = \min\{(\tilde{f}_{h,\bar{\eta}}^{k})^{(1)}(\cdot, \cdot) + b_{h}^{k}(\cdot, \cdot), H\}$ and choose the greedy policy $\pi_{h}^{k}(\cdot)$ with respect to Q_{h}^{k} . We repeat the procedure until all the K episodes are completed.

Remark 5.1. For an optimistic planning, we define the bonus function as the width function $b_h^k(s, a) \coloneqq w_h^k((\mathcal{F}^N)_h^k, s, a)$ where $(\mathcal{F}^N)_h^k$ denotes a confidence region at step h, episode k. When \mathcal{F} is a linear function class, the width function can be evaluated by simply computing the maximal distance of weight vector. For a general function class \mathcal{F} , computing the width function requires to solve a set-constrained optimization problem, which is known as NP-hard [17]. However, a width function is computed simply for optimistic exploration, and approximation errors are known to have a small effect on regret [1].

6 Theoretical Analysis

In this section, we provide the theoretical guarantees for SF-LSVI under assumption 4.8. We apply proof techniques from Wang et al. [50] and extend the result to a statistical functional lens. First, we generalize the concept of eluder dimension [43] to the vector-valued function, which has been widely used in RL literatures [4, 27, 50] to measure the complexity of learning with the function approximators.

Definition 6.1 (ϵ -dependent, ϵ -independent, Eluder dimension for vector-valued function). Let $\epsilon \ge 0$ and $\mathcal{Z} = \{(s_i, a_i)\}_{i=1}^n \subseteq S \times A$ be a sequence of state-action pairs.

- A state-action pair $(s, a) \in S \times A$ is ϵ -dependent on Z with respect to \mathcal{F}^N if $||f g||_{\mathcal{Z}} \leq \epsilon$ for any vector-valued function $f, g \in \mathcal{F}^N$, then $|f^{(1)}(s, a) g^{(1)}(s, a)| \leq \epsilon$.
- An (s, a) is ϵ -independent on \mathcal{Z} with respect to \mathcal{F}^N if (s, a) is not ϵ -dependent on \mathcal{Z} .
- The ϵ -eluder dimension $\dim_E(\mathcal{F}^N, \epsilon)$ of a vector-valued function class \mathcal{F}^N is the length of the longest sequence of elements in $\mathcal{S} \times \mathcal{A}$ such that, for some $\epsilon' \geq \epsilon$, every element is ϵ' -independent on its predecessors.

We assume that the function class \mathcal{F}^N and state-action space $\mathcal{S} \times \mathcal{A}$ have bounded covering numbers. Assumption 6.2 (Covering number). For any $\epsilon > 0$, the following holds:

- there exists an ϵ -cover $\mathcal{C}(\mathcal{F}^N, \epsilon) \subseteq \mathcal{F}^N$ with size $|\mathcal{C}(\mathcal{F}^N, \epsilon)| \leq \mathcal{N}(\mathcal{F}^N, \epsilon)$, such that for any $g \in \mathcal{F}^N$, there exists $g' \in \mathcal{C}(\mathcal{F}^N, \epsilon)$ with $\|g g'\|_{\infty, 1} \leq \epsilon$.
- there exists an ϵ -cover $\mathcal{C}(\mathcal{S} \times \mathcal{A}, \epsilon)$ with size $|\mathcal{C}(\mathcal{S} \times \mathcal{A}, \epsilon)| \leq \mathcal{N}(\mathcal{S} \times \mathcal{A}, \epsilon)$, such that for any $(s, a) \in \mathcal{S} \times \mathcal{A}$, there exists $(s', a') \in \mathcal{C}(\mathcal{S} \times \mathcal{A}, \epsilon)$ with $\max_{f \in \mathcal{F}} |f(s, a) f(s', a')| \leq \epsilon$

The following two lemmas give confidence bounds on the sum of the l_2 norms of all normalized moments.

Lemma 6.3 (Single Step Optimization Error). Consider a fixed $k \in [K]$ and a fixed $h \in [H]$. Let $\mathcal{Z}_{h}^{k} = \{(s_{h}^{\tau}, a_{h}^{\tau})\}_{\tau \in [k-1]}$ and $\mathcal{D}_{h,\bar{\eta}}^{k} = \left\{\left(s_{h}^{\tau}, a_{h}^{\tau}, \psi_{1:N}\left((\mathcal{B}_{r_{h'}}^{\tau})_{\#}\bar{\eta}(s_{h'+1}^{\tau})\right)\right)\right\}_{\tau \in [k-1]}$ for any $\bar{\eta}$: $\mathcal{S} \to \mathscr{P}([0, H])$. Define $\tilde{f}_{h,\bar{\eta}}^{k} = \arg\min_{f \in \mathcal{F}^{N}} \|f\|_{\mathcal{D}_{h,\bar{\eta}}^{k}}^{2}$. For any $\bar{\eta}$ and $\delta \in (0, 1)$, there is an event $\mathcal{E}(\bar{\eta}, \delta)$ such that conditioned on $\mathcal{E}(\bar{\eta}, \delta)$, with probability at least $1 - \delta$, for any $\bar{\eta}' : \mathcal{S} \to \mathscr{P}([0, H])$ with $\|\psi_{1:N}(\bar{\eta}') - \psi_{1:N}(\bar{\eta})\|_{\infty, 1} \leq 1/T$, we have

$$\left\| \tilde{f}_{h,\bar{\eta}'}(\cdot,\cdot) - \psi_{1:N} \left((\mathcal{B}_{r(\cdot,\cdot)})_{\#} [\mathbb{P}\bar{\eta}'](\cdot,\cdot) \right) \right\|_{\mathcal{Z}_{h}^{k}} \leq c' \left(N^{\frac{1}{2}} H \sqrt{\log(1/\delta) + \log \mathcal{N}(\mathcal{F}^{N},1/T)} \right)^{k}$$

for some constant c' > 0.

We remark that using moment functional as a sketch is a unique structure that leads to a concentration results for the difference between the target and sampled sketches $\psi\left((\mathcal{B}_r)_{\#}[\mathbb{P}\bar{\eta}]\right) - \psi\left((\mathcal{B}_r)_{\#}\bar{\eta}\right)$, as the tranformation ϕ_{ψ} of moment functional is an identity. Furthermore, using normalized moments minimizes the size of the confidence region from $O(H^N)$ to $O(\sqrt{N})$. This adjustment is akin to scaling the optimization function in multi-objective optimization to treat each objective equally, which effectively prevents the model from favoring objectives with larger scales.

Lemma 6.4 (Confidence Region). Let $(\mathcal{F}^N)_h^k = \{f \in \mathcal{F}^N | \|f - \tilde{f}_{h,\bar{\eta}}^k\|_{\mathcal{Z}_h^k}^2 \leq \beta(\mathcal{F}^N, \delta)\},$ where

$$\beta(\mathcal{F}^N, \delta) \ge c' \cdot NH^2(\log(T/\delta) + \log \mathcal{N}(\mathcal{F}^N, 1/T))$$

for some constant c' > 0. Then with probability at least $1 - \delta/2$, for all $k, h \in [K] \times [H]$, we have

$$\psi_n\Big((\mathcal{B}_{r_h(\cdot,\cdot)})_{\#}[\mathbb{P}_h\bar{\eta}_{h+1}^k](\cdot,\cdot)\Big)\in(\mathcal{F}^N)_h^k$$

Lemma 6.4 guarantees that the sequence of moments from the target distribution $\psi_{1:N}((\mathcal{B}_{r_h(\cdot,\cdot)})_{\#}[\mathbb{P}_h\bar{\eta}_{h+1}^k](\cdot,\cdot))$ lies in the confidence region $(\mathcal{F}^N)_h^k$ with high probability. Supported by the aforementioned lemma, we can further guarantee that all *Q*-functions are optimistically estimated with high probability and derive our final result.

Theorem 6.5. Under Assumption 4.8, with probability at least $1 - \delta$, SF-LSVI achieves a regret bound of

$$\operatorname{Reg}(K) \le 2H \operatorname{dim}_E(\mathcal{F}^N, 1/T) + 4H\sqrt{KH}\log(1/\delta)$$

Under weaker structural assumptions, we show that SF-LSVI enjoys sublinear regret of order $\tilde{O}(d_E H^{\frac{3}{2}}\sqrt{K})$, which is \sqrt{H} better than the state-of-the-art distributional framework algorithm V-EST-LSR [11]. For the linear MDP setting, we have $d_E = \tilde{O}(d)$ and thus the regret bound of SF-LSVI is $\tilde{O}(\sqrt{d^2H^3K})$ which matches a lower bound $\Omega(\sqrt{d^2H^3K})$ [53]. In our analysis, we highlight two main technical novelties which significantly reduces the complexity of regret;

- 1. We refine previous lemma of Osband et al. [39] and Wang et al. [50] to remove the dependency of $\beta(\mathcal{F}^N, 1/\delta)$ (See Appendix C.4), ensuring that regret bound depends only on the defined function class, not on the number of moment extracted.
- 2. As shown in Table 1, we define the eluder dimension d_E in a finite-dimensional embedding space \mathcal{F}^N , while other methods rely on an infinite-dimensional distribution space $\mathcal{H} \subseteq \mathcal{F}^{\infty}$.

7 Conclusions

We describe the sources of approximation error inherent in distribution-based updates and introduce a pivotal concept of Bellman unbiasedness, which enables to exactly learn the information of distribution. We also present a provably efficient distRL algorithm, SF-LSVI, with general value function approximation. Notably, our algorithm achieves a near-optimal regret bound of $\tilde{O}(d_E H^{\frac{3}{2}}\sqrt{K})$, matching the tightest upper bound achieved by non-distributional framework [24, 53]. One interesting future direction would be to reformulate the definition of regret as discrepencies in moments rather than the expected return, and to show the sample-efficiency of distRL. We hope that our work sheds some light on future research in analyzing the provable efficiency of distRL.

References

- [1] Yasin Abbasi-Yadkori, Dávid Pál, and Csaba Szepesvári. Improved algorithms for linear stochastic bandits. *Advances in neural information processing systems*, 24, 2011.
- [2] Naum Ilich Akhiezer. *The classical moment problem and some related questions in analysis*. SIAM, 2020.
- [3] Peter Auer, Thomas Jaksch, and Ronald Ortner. Near-optimal regret bounds for reinforcement learning. *Advances in neural information processing systems*, 21, 2008.
- [4] Alex Ayoub, Zeyu Jia, Csaba Szepesvari, Mengdi Wang, and Lin Yang. Model-based reinforcement learning with value-targeted regression. In *International Conference on Machine Learning*, pages 463–474. PMLR, 2020.
- [5] Osbert Bastani, Jason Yecheng Ma, Estelle Shen, and Wanqiao Xu. Regret bounds for risksensitive reinforcement learning. Advances in Neural Information Processing Systems, 35: 36259–36269, 2022.
- [6] Marc G Bellemare, Yavar Naddaf, Joel Veness, and Michael Bowling. The arcade learning environment: An evaluation platform for general agents. *Journal of Artificial Intelligence Research*, 47:253–279, 2013.
- [7] Marc G Bellemare, Will Dabney, and Rémi Munos. A distributional perspective on reinforcement learning. In *International Conference on Machine Learning*, pages 449–458. PMLR, 2017.
- [8] Marc G Bellemare, Salvatore Candido, Pablo Samuel Castro, Jun Gong, Marlos C Machado, Subhodeep Moitra, Sameera S Ponda, and Ziyu Wang. Autonomous navigation of stratospheric balloons using reinforcement learning. *Nature*, 588(7836):77–82, 2020.
- [9] Marc G Bellemare, Will Dabney, and Mark Rowland. *Distributional reinforcement learning*. MIT Press, 2023.
- [10] Cristian Bodnar, Adrian Li, Karol Hausman, Peter Pastor, and Mrinal Kalakrishnan. Quantile qt-opt for risk-aware vision-based robotic grasping. arXiv preprint arXiv:1910.02787, 2019.
- [11] Yu Chen, Xiangcheng Zhang, Siwei Wang, and Longbo Huang. Provable risk-sensitive distributional reinforcement learning with general function approximation. *arXiv preprint arXiv:2402.18159*, 2024.
- [12] Taehyun Cho, Seungyub Han, Heesoo Lee, Kyungjae Lee, and Jungwoo Lee. Pitfall of optimism: Distributional reinforcement learning by randomizing risk criterion. arXiv preprint arXiv:2310.16546, 2023.
- [13] Yunho Choi, Kyungjae Lee, and Songhwai Oh. Distributional deep reinforcement learning with a mixture of gaussians. In 2019 International conference on robotics and automation (ICRA), pages 9791–9797. IEEE, 2019.
- [14] Yinlam Chow, Aviv Tamar, Shie Mannor, and Marco Pavone. Risk-sensitive and robust decisionmaking: a cvar optimization approach. Advances in neural information processing systems, 28, 2015.
- [15] Will Dabney, Georg Ostrovski, David Silver, and Rémi Munos. Implicit quantile networks for distributional reinforcement learning. In *International conference on machine learning*, pages 1096–1105. PMLR, 2018.
- [16] Will Dabney, Mark Rowland, Marc Bellemare, and Rémi Munos. Distributional reinforcement learning with quantile regression. In *Proceedings of the AAAI Conference on Artificial Intelligence*, 2018.
- [17] Christoph Dann, Nan Jiang, Akshay Krishnamurthy, Alekh Agarwal, John Langford, and Robert E Schapire. On oracle-efficient pac rl with rich observations. *Advances in neural information processing systems*, 31, 2018.

- [18] Martin Engert. Finite dimensional translation invariant subspaces. *Pacific Journal of Mathematics*, 32(2):333–343, 1970.
- [19] Alhussein Fawzi, Matej Balog, Aja Huang, Thomas Hubert, Bernardino Romera-Paredes, Mohammadamin Barekatain, Alexander Novikov, Francisco J R Ruiz, Julian Schrittwieser, Grzegorz Swirszcz, et al. Discovering faster matrix multiplication algorithms with reinforcement learning. *Nature*, 610(7930):47–53, 2022.
- [20] Yingjie Fei, Zhuoran Yang, and Zhaoran Wang. Risk-sensitive reinforcement learning with function approximation: A debiasing approach. In *International Conference on Machine Learning*, pages 3198–3207. PMLR, 2021.
- [21] Fei Feng, Wotao Yin, Alekh Agarwal, and Lin Yang. Provably correct optimization and exploration with non-linear policies. In *International Conference on Machine Learning*, pages 3263–3273. PMLR, 2021.
- [22] Ido Greenberg, Yinlam Chow, Mohammad Ghavamzadeh, and Shie Mannor. Efficient riskaverse reinforcement learning. Advances in Neural Information Processing Systems, 35:32639– 32652, 2022.
- [23] Paul R Halmos. The theory of unbiased estimation. *The Annals of Mathematical Statistics*, 17 (1):34–43, 1946.
- [24] Jiafan He, Heyang Zhao, Dongruo Zhou, and Quanquan Gu. Nearly minimax optimal reinforcement learning for linear markov decision processes. In *International Conference on Machine Learning*, pages 12790–12822. PMLR, 2023.
- [25] Haque Ishfaq, Qiwen Cui, Viet Nguyen, Alex Ayoub, Zhuoran Yang, Zhaoran Wang, Doina Precup, and Lin Yang. Randomized exploration in reinforcement learning with general value function approximation. In *International Conference on Machine Learning*, pages 4607–4616. PMLR, 2021.
- [26] Chi Jin, Zeyuan Allen-Zhu, Sebastien Bubeck, and Michael I Jordan. Is q-learning provably efficient? *Advances in neural information processing systems*, 31, 2018.
- [27] Chi Jin, Zhuoran Yang, Zhaoran Wang, and Michael I Jordan. Provably efficient reinforcement learning with linear function approximation. In *Conference on Learning Theory*, pages 2137– 2143. PMLR, 2020.
- [28] Sham Machandranath Kakade. On the sample complexity of reinforcement learning. University of London, University College London (United Kingdom), 2003.
- [29] Dohyeong Kim, Kyungjae Lee, and Songhwai Oh. Trust region-based safe distributional reinforcement learning for multiple constraints. In *Thirty-seventh Conference on Neural Information Processing Systems*, 2023.
- [30] Dohyeong Kim, Kyungjae Lee, and Songhwai Oh. Trust region-based safe distributional reinforcement learning for multiple constraints. *Advances in neural information processing* systems, 36, 2024.
- [31] Mark Grigorevich Kreuin, AA Nudel, and D Louvish. *The Markov moment problem and extremal problems*. American Mathematical Society, 1977.
- [32] Tor Lattimore and Csaba Szepesvári. Bandit algorithms. Cambridge University Press, 2020.
- [33] Hao Liang and Zhi-Quan Luo. Bridging distributional and risk-sensitive reinforcement learning with provable regret bounds. arXiv preprint arXiv:2210.14051, 2022.
- [34] Marlos C Machado, Marc G Bellemare, Erik Talvitie, Joel Veness, Matthew Hausknecht, and Michael Bowling. Revisiting the arcade learning environment: Evaluation protocols and open problems for general agents. *Journal of Artificial Intelligence Research*, 61:523–562, 2018.
- [35] Alexandre Marthe, Aurélien Garivier, and Claire Vernade. Beyond average return in markov decision processes. arXiv preprint arXiv:2310.20266, 2023.

- [36] Timothy H Muller, James L Butler, Sebastijan Veselic, Bruno Miranda, Joni D Wallis, Peter Dayan, Timothy EJ Behrens, Zeb Kurth-Nelson, and Steven W Kennerley. Distributional reinforcement learning in prefrontal cortex. *Nature Neuroscience*, pages 1–6, 2024.
- [37] Thanh Nguyen-Tang, Sunil Gupta, and Svetha Venkatesh. Distributional reinforcement learning via moment matching. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 35, pages 9144–9152, 2021.
- [38] Ian Osband and Benjamin Van Roy. On lower bounds for regret in reinforcement learning. arXiv preprint arXiv:1608.02732, 2016.
- [39] Ian Osband, Benjamin Van Roy, Daniel J Russo, Zheng Wen, et al. Deep exploration via randomized value functions. J. Mach. Learn. Res., 20(124):1–62, 2019.
- [40] Mark Rowland, Marc Bellemare, Will Dabney, Rémi Munos, and Yee Whye Teh. An analysis of categorical distributional reinforcement learning. In *International Conference on Artificial Intelligence and Statistics*, pages 29–37. PMLR, 2018.
- [41] Mark Rowland, Robert Dadashi, Saurabh Kumar, Rémi Munos, Marc G Bellemare, and Will Dabney. Statistics and samples in distributional reinforcement learning. In *International Conference on Machine Learning*, pages 5528–5536. PMLR, 2019.
- [42] Mark Rowland, Rémi Munos, Mohammad Gheshlaghi Azar, Yunhao Tang, Georg Ostrovski, Anna Harutyunyan, Karl Tuyls, Marc G Bellemare, and Will Dabney. An analysis of quantile temporal-difference learning. *arXiv preprint arXiv:2301.04462*, 2023.
- [43] Daniel Russo and Benjamin Van Roy. Eluder dimension and the sample complexity of optimistic exploration. Advances in Neural Information Processing Systems, 26, 2013.
- [44] K Schmüdgen and PJ di Dio. "truncated moment problem: set of atoms and carathéodory numbers. Oberwolfach Rep., 14:77–79, 2017.
- [45] Matthew J Sobel. The variance of discounted markov decision processes. *Journal of Applied Probability*, 19(4):794–802, 1982.
- [46] Kyunghwan Son, Junsu Kim, Yung Yi, and Jinwoo Shin. Disentangling sources of risk for distributional multi-agent reinforcement learning. 2021.
- [47] Aviv Tamar, Dotan Di Castro, and Shie Mannor. Learning the variance of the reward-to-go. *Journal of Machine Learning Research*, 17(13):1–36, 2016.
- [48] Henry Teicher. On the mixture of distributions. *The Annals of Mathematical Statistics*, 31(1): 55–73, 1960.
- [49] Kaiwen Wang, Kevin Zhou, Runzhe Wu, Nathan Kallus, and Wen Sun. The benefits of being distributional: Small-loss bounds for reinforcement learning. arXiv preprint arXiv:2305.15703, 2023.
- [50] Ruosong Wang, Russ R Salakhutdinov, and Lin Yang. Reinforcement learning with general value function approximation: Provably efficient approach via bounded eluder dimension. *Advances in Neural Information Processing Systems*, 33:6123–6135, 2020.
- [51] Yining Wang, Ruosong Wang, Simon S Du, and Akshay Krishnamurthy. Optimism in reinforcement learning with generalized linear function approximation. arXiv preprint arXiv:1912.04136, 2019.
- [52] Andrea Zanette, Alessandro Lazaric, Mykel Kochenderfer, and Emma Brunskill. Learning near optimal policies with low inherent bellman error. In *International Conference on Machine Learning*, pages 10978–10989. PMLR, 2020.
- [53] Dongruo Zhou, Quanquan Gu, and Csaba Szepesvari. Nearly minimax optimal reinforcement learning for linear mixture markov decision processes. In *Conference on Learning Theory*, pages 4532–4576. PMLR, 2021.

Appendix

A Notation

Notation	Description					
S	state space of size S					
\mathcal{A}	action space of size A					
Н	horizon length of one episode					
Т	number of episodes					
$r_h(s,a)$	reward of (s, a) at step h					
$\mathbb{P}_h(s' s,a)$	probability transition of (s, a) to s' at step h					
\mathbb{H}_{h}^{k}	history up to step h , episode k					
N	number of statistical functionals					
$Q_h^{\pi}(s, a)$	Q-function of a given policy π at step h					
$V_h^{\pi}(s)$	V-function of a given policy π at step h					
$Z_h^{\pi}(s,a)$	random variable of Q-function					
$\bar{Z}_h^{\pi}(s)$	random variable of V-function					
$\eta_h^{\pi}(s, a)$	probability distribution of Q-function					
$\bar{\eta}_h^{\pi}(s)$	probability distribution of V-function					
$[\mathbb{P}_h(\cdot)]$	expectation over transition $[\mathbb{P}_h(\cdot)] = \mathbb{E}_{s' \sim \mathbb{P}_h}(\cdot)$					
$(\mathcal{B}_r)_{\#}$	pushforward of the distribution through $\mathcal{B}_r(x) \coloneqq r + x$					
$f^{(n)}$	n-th element of N -dimensional vector f					
$\ f\ _{\infty}$	max norm of $f: X \to \mathbb{R}$ defined as $\ f\ _{\infty} \coloneqq \max_{x \in X} f^{(n)}(x) $					
$ f _{\infty,1}$	l_1 -norm of max norm of $f: X \to \mathbb{R}$ defined as $ f _{\infty,1} \coloneqq \sum_{n=1}^N \max_{x \in X} f^{(n)}(x) $					
\mathcal{F}^N	a function class of N -dimensional embedding space					
Z	a set of state-action pairs $\mathcal{Z} \coloneqq \{(s_t, a_t)\}_{t=1}^{ \mathcal{Z} }$					
\mathcal{D}	a dataset $\mathcal{D} \coloneqq \{(s_t, a_t, [d_t^{(1)}, \cdots, d_t^{(N)}])\}_{t=1}^{ \mathcal{D} }$					
$\ f\ _{\mathcal{Z}}^2$	for $f: \mathcal{S} \times \mathcal{A} \to \mathbb{R}$, define $\ f\ _{\mathcal{Z}}^2 \coloneqq \sum_{n=1}^N \sum_{(s,a) \in \mathcal{Z}} (f^{(n)}(s_t, a_t))^2$					
$\ f\ _{\mathcal{D}}^2$	for $f: S \times A \to \mathbb{R}$, define $ f _{\mathcal{D}}^2 \coloneqq \sum_{n=1}^N \sum_{t=1}^{\mathcal{D}} (f^{(n)}(s_t, a_t) - d_t^{(n)})^2$					
$w^{(n)}(\mathcal{F}^N, s, a)$	width function of (s, a) defined as $w^{(n)}(\mathcal{F}^N, s, a) \coloneqq \max_{f,g \in \mathcal{F}^N} f^{(n)}(s, a) - g^{(n)}(s, a) $					
$\tilde{f}^k_{h,\bar{\eta}}$	a solution of moment least squre regression, defined as $\tilde{f}^k_{h,\bar{\eta}} \coloneqq \arg\min_{f \in \mathcal{F}^N} \ f\ _{\mathcal{D}^k_h}$					
$f_{ar\eta}$	a target sketch of distribution $\bar{\eta}$, defined as $f_{\bar{\eta}} \coloneqq \psi_{1:N}((\mathcal{B}_r)_{\#}[\mathbb{P}_h\bar{\eta}])$					
$(\mathcal{F}^N)_h^k$	a confidence region at step h , episode k , defined as $(\mathcal{F}^N)_h^k \coloneqq \{f \in \mathcal{F}^N \ f - \tilde{f}_{h,\bar{\eta}}^k\ _{\mathcal{Z}_h^k}^2 \leq \beta(\mathcal{F}^N, \delta)$					
$\psi(\bar{\eta})$	a statistical functional $\mathscr{P}_{\psi}(\mathbb{R})^{\mathcal{S}} \to \mathbb{R}^{\mathcal{S}}$					
$\psi_{1:N}(\bar{\eta})$	a <i>N</i> -collection of statistical functional $\mathscr{P}_{\psi_{1:N}}(\mathbb{R})^{\mathcal{S}} \to \mathbb{R}^{N \times S}$					
$\mathscr{P}_{\psi_{1:N}}(\mathbb{R})$	a domain of sketch $\psi_{1:N}$					
$I_{\psi_{1:N}}$	an image of sketch $\psi_{1:N}$					
${\mathcal T}$	distributional Bellman operator, defined as $\mathcal{T}\bar\eta\coloneqq(\mathcal{B}_r)_\#[\mathbb{P}\bar\eta]$					
\mathcal{T}_ψ	sketch Bellman operator w.r.t ψ , defined as $\mathcal{T}_{\psi}\psi(ar{\eta})\coloneqq\psiig((\mathcal{B}_r)_{\#}[\mathbb{P}ar{\eta}]ig)$					
$\hat{\mathcal{T}}_\psi$	empirical sketch Bellman operator w.r.t ψ , defined as $\hat{\mathcal{T}}_{\psi}\psi(ar{\eta}) \coloneqq \psi(\check{\mathcal{B}}_r)_{\#}[\hat{\mathbb{P}}ar{\eta}]$					
$\mathcal{N}(\mathcal{F}^N,\epsilon)$	covering number of \mathcal{F}^N w.r.t the ϵ -ball					
$\dim_E(\mathcal{F}^N,\epsilon)$	eluder dimension of \mathcal{F}^N w.r.t ϵ					

Table 2: Table of notation

B Related Work and Discussion

B.1 Technical Clarifications on Linearity Assumption in Existing Results

Bellman Closedness and Linearity. Rowland et al. [41] proved that quantile functional is not Bellman closed by providing a specific counterexample. However, their discussion based on counterexamples can be generalized as it assumes that the sketch Bellman operator for the quantile functional needs to be linear.

They consider an discounted MDP with initial state s_0 with single action a, which transits to one of two terminal states s_1, s_2 with equal probability. Letting no reward at state s_0 , Unif([0,1]) at state s_1 , and Unif([1/K, 1+1/K]) at state s_2 , the return distribution at state s_0 is computed as mixture $\frac{1}{2}\text{Unif}([0,\gamma]) + \frac{1}{2}\text{Unif}([\gamma/K, \gamma + \gamma/K])$. Then the $\frac{1}{2K}$ -quantile at state s_0 is $\frac{\gamma}{K}$. They proposed a counterexample where each quantile distribution of state s_1, s_2 is represented as $\frac{1}{K}\sum_{k=1}^{K} \delta_{\frac{2k-1}{K}}$ and

 $\frac{1}{K}\sum_{k=1}^{K} \delta_{\frac{2k+1}{K}} \text{ respectively, the } \frac{1}{2K} - \text{quantile of state } s_0 \text{ is } \psi_{q_{2K}} \left(\frac{1}{2K} \sum_{k=1}^{K} \delta_{\frac{\gamma(2k-1)}{K}} + \delta_{\frac{\gamma(2k+1)}{K}} \right) = \frac{3\gamma}{2K}.$ However, this example does not consider that the mixture of quantiles is not a quantile of the mixture distribution (i.e., $\psi_q(\lambda\eta_1 + (1-\lambda)\eta_2) \neq \lambda\psi_q(\eta_1) + (1-\lambda)\psi_q(\eta_2)$), due to the nonlinearity of the quantile functional. Therefore, this does not present a valid counterexample to prove that quantile functionals are not Bellman closed.

Bellman Optimizability and Linearity. Marthe et al. [35] proposed the notion of Bellman optimizable statistical functional which redefine the Bellman update by planning with respect to statistical functionals rather than expected returns. They proved that W_1 -continuous Bellman Optimizable statistical functionals are characterized by exponential utilities $\frac{1}{\lambda} \log \mathbb{E}_{Z \sim \eta} [\exp(\lambda Z)]$. However, their proof requires some technical clarification regarding the assumption that such statistical functionals are linear.

To illustrate, they define a statistical functional ψ_f and consider two probability distributions $\eta_1 = \frac{1}{2}(\delta_0 + \delta_h)$ and $\eta_2 = \delta_{\phi(h)}$ where $\phi(h) = f^{-1}(\frac{1}{2}(f(0) + f(h)))$. Using the translation property, they lead $\psi_f(\eta_1) = \psi_f(\eta_2)$ to $\frac{1}{2}(f(x) + f(x+h)) = f(x + \phi(h))$ for all $x \in \mathbb{R}$. However, this equality $\psi_f(\frac{1}{2}(\delta_x + \delta_{x+h})) = \frac{1}{2}(f(x) + f(x+h))$ holds only if ψ_f is linear, which is not necessarily a valid assumption for all statistical functionals.

B.2 Existence of Nonlinear Bellman Closed Sketch.

The previous two examples may not have considered the possibility that the sketch Bellman operator might not necessarily be linear. However, some statistical functionals are Bellman-closed even if they are nonlinear, so it is open question whether there is a nonlinear sketch Bellman operator that makes the quantile functional Bellman-closed. In this section, we present examples of maximum and minimum functionals that are Bellman-closed, despite being nonlinear.

In a nutshell, consider the maximum of return distribution at state s_1, s_2 is $\gamma, \gamma + \gamma/K$ respectively. Beyond linearity, the maximum of return distribution at state s_0 can be computed by taking the maximum of these values;

$$\max(\max(\bar{\eta}(s_1)), \max(\bar{\eta}(s_2))) = \max(\gamma, \gamma + \gamma/K) = \gamma + \gamma/K$$

which produces the desired result. This implies the existence of a nonlinear sketch that is Bellman closed. More precisely, by defining $\max_{s' \sim \mathbb{P}(\cdot|s,a)}$ and $\min_{s' \sim \mathbb{P}(\cdot|s,a)}$ as the maximum and minimum of the sampled sketch $\psi((\mathcal{B}_r)_{\#}\bar{\eta}(s'))$ with the distribution $\mathbb{P}(\cdot|s,a)$, we can derive the sketch Bellman operator for maximum and minimum functionals as follows;

•
$$\mathcal{T}_{\psi_{\max}}\left(\psi_{\max}(\bar{\eta}(s))\right) = \max_{s' \sim \mathbb{P}(\cdot|s,a)}\left(\psi_{\max}\left((\mathcal{B}_r)_{\#}\bar{\eta}(s')\right)\right)$$

• $\mathcal{T}_{\psi_{\min}}\left(\psi_{\min}(\bar{\eta}(s))\right) = \min_{s' \sim \mathbb{P}(\cdot|s,a)}\left(\psi_{\min}\left((\mathcal{B}_r)_{\#}\bar{\eta}(s')\right)\right).$

B.3 Non-existence of sketch Bellman operator for quantile functional

In this section, we prove that quantile functional cannot be Bellman closed under any additional sketch. First we introduce the definition of *mixture-consistent*, which is the property that the sketch of a mixture can be computed using only the sketch of the distribution of each component.

Definition B.1 (mixture-consistent). A sketch ψ is mixture-consistent if for any $\lambda \in [0, 1]$ and any distributions $\eta_1, \eta_2 \in \mathscr{P}_{\psi}(\mathbb{R})$, there exists a corresponding function h_{ψ} such that

$$\psi(\lambda\eta_1 + (1-\lambda)\eta_2) = h_{\psi}(\psi(\eta_1), \psi(\eta_2), \lambda).$$

Next, we will provide some examples of determining whether a sketch is mixture-consistent or not.

Example 1. Every moment or exponential polynomial functional is mixture-consistent.

Proof. For any $n \in [N]$ and $\lambda \in \mathbb{C}$, $\mathbb{E}_{Z \sim \lambda \eta_1 + (1-\lambda)\eta_2}[Z^n \exp(\lambda Z)] = \lambda \mathbb{E}_{Z \sim \eta_1}[Z^n \exp(\lambda Z)] + (1-\lambda)\mathbb{E}_{Z \sim \eta_2}[Z^n \exp(\lambda Z)].$

Example 2. Variance functional is not mixture-consistent.

Proof. Let $\lambda = \frac{1}{2}$ and Z, Y be the random variables where $Z \sim \frac{1}{2}\delta_0 + \frac{1}{2}\delta_2$ and $Y \sim \frac{1}{2}\delta_k + \frac{1}{2}\delta_{k+2}$. Then, $\operatorname{Var}(Z) = \operatorname{Var}(Y) = 1$. While RHS is constant for any k, LHS is not a constant for any k, i.e.,

$$\operatorname{Var}_{X \sim \frac{1}{2}(\frac{1}{2}\delta_0 + \frac{1}{2}\delta_2) + \frac{1}{2}(\frac{1}{2}\delta_k + \frac{1}{2}\delta_{k+2})}(X) = \frac{1}{4}(k^2 + 5)$$

While variance functional is not mixture consistent by itself, it can be mixture consistent with another statistical functional, the mean.

Example 3. Variance functional is mixture-consistent under mean functional.

Proof. Notice that mean functional is mixture-consistent. We need to show that variance functional is mixture-consistent under mean functional.

$$\begin{aligned} \operatorname{Var}_{Z \sim \lambda \eta_1 + (1 - \lambda)\eta_2}[Z] \\ &= \mathbb{E}_{Z \sim \lambda \eta_1 + (1 - \lambda)\eta_2}[Z^2] - (\mathbb{E}_{Z \sim \lambda \eta_1 + (1 - \lambda)\eta_2}[Z])^2 \\ &= \lambda \mathbb{E}_{Z \sim \eta_1}[Z^2] + (1 - \lambda)\mathbb{E}_{Z \sim \eta_2}[Z^2] - (\lambda \mathbb{E}_{Z \sim \eta_1}[Z] + (1 - \lambda)\mathbb{E}_{Z \sim \eta_2}[Z])^2 \\ &= \lambda (\operatorname{Var}_{Z \sim \eta_1}[Z] + (\mathbb{E}_{Z \sim \eta_1}[Z])^2) + (1 - \lambda)(\operatorname{Var}_{Z \sim \eta_2}[Z] + (\mathbb{E}_{Z \sim \eta_2}[Z])^2) \\ &- (\lambda \mathbb{E}_{Z \sim \eta_1}[Z] + (1 - \lambda)\mathbb{E}_{Z \sim \eta_2}[Z])^2. \end{aligned}$$

This means that to determine whether it is mixture-consistent or not, we should check it on a per-sketch basis, rather than on a per-statistical functional basis.

Example 4. Maximum and minimum functional are both mixture-consistent.

Proof.

$$\max_{Z \sim \lambda \eta_1 + (1-\lambda)\eta_2} [Z] = \max(\max_{Z \sim \eta_1} [Z], \max_{Z \sim \eta_2} [Z])$$

and

$$\min_{Z \sim \lambda \eta_1 + (1-\lambda)\eta_2} [Z] = \min(\min_{Z \sim \eta_1} [Z], \min_{Z \sim \eta_2} [Z])$$

Since maximum and minimum functionals are mixture consistent, we can construct a nonlinear sketch bellman operator like the one in section B.2. This is possible because there is a nonlinear function h_{ψ} that ensures the sketch is closed under mixture.

Before demonstrating that a quantile sketch cannot be mixture consistent under any additional sketch, we will first illustrate with the example of a median functional that is not mixture consistent.

Example 5. Median sketch is not mixture-consistent.

Proof. Let $\lambda = \frac{1}{2}$ and Z, Y be the random variables where $Z \sim 0.2\delta_0 + 0.8\delta_1$ and $Y \sim 0.6\delta_0 + 0.4\delta_k$ for some 0 < k < 1. Then $\psi_{\text{med}}(Z) = 1$ and $\psi_{\text{med}}(Y) = 0$. However,

$$\operatorname{med}_{X=\frac{Z+Y}{2}}[X] = \psi_{\operatorname{med}}(0.4\delta_0 + 0.2\delta_k + 0.4\delta_1) = k$$

which is dependent in k.

Lemma B.2. Quantile sketch cannot be mixture-consistent, under any additional sketch.

Proof. For a given integer N > 0 and a quantile level $\alpha \in [0, 1]$, let $\lambda = \frac{1}{2}$ and a random variable $Y \sim p_{y_0}\delta_0 + p_{y_1}\delta_{y_1} + \dots + p_{y_N}\delta_{y_N} (0 < y_1 < \dots < y_N < 1)$ where $p_{y_0} > \alpha$ so that $\psi_{q_\alpha}(Y) = 0$. Consider another random variable $Z \sim p_{z_0}\delta_0 + p_{z_1}\delta_1$ where $p_{z_0} < \alpha$ so that $\psi_{q_\alpha}(Z) = 1$. Then the α -quantile of the mixture $X = \frac{Y+Z}{2}$ is

$$\psi_{q_{\alpha}}[X] = y_n \text{ where } n = \min\left\{ n \le N \Big| \ \frac{1}{2} \sum_{n'=0}^n p_{y_{n'}} + \frac{1}{2} p_{z_0} > \alpha
ight\}.$$

Letting $p_{z_0} = 2\alpha - \sum_{n'=0}^{n} p_{y_{n'}}$, we can manipulate $\psi_{q_\alpha}[X]$ to be any value of y_n . Hence, $\psi(q_\alpha)[X]$ is a function of all possible outcomes of Y.

If there exists a finite number of statistical functionals which make quantile sketch mixture-consistent, then such sketch would uniquely determine the distribution for any N. This results in a contradiction that infinite-dimensional distribution space can be represented by a finite number of statistical functional.

Lemma B.3. If a sketch ψ is Bellman closed, then it is mixture-consistent.

Proof. Consider an MDP where initial state s_0 has no reward and transits to two state s_1, s_2 with probability $\lambda, 1 - \lambda$ and reward distribution $\bar{\eta}_1, \bar{\eta}_2$. Since ψ is Bellman closed, $\psi(\bar{\eta}(s_0))$ is a function of $\psi(\bar{\eta}(s_1))$ and $\psi(\bar{\eta}(s_2))$, (i.e., $\psi(\bar{\eta}(s_0)) = g_{\psi}(\psi(\bar{\eta}(s_1)), \psi(\bar{\eta}(s_2)))$ for some g_{ψ}). Since $\psi(\bar{\eta}(s_0)) = \psi(\lambda \bar{\eta}(s_1) + (1 - \lambda) \bar{\eta}(s_2))$, it implies that ψ is mixture-consistent.

Combining the results of Lemma B.2 and Lemma B.3, we prove that a quantile sketch cannot be Bellman closed, no matter what additional sketches are provided.

C Proof

Theorem (4.4). *Quantile functional cannot be Bellman closed under any additional sketch.*

Proof. See Lemma B.2 and Lemma B.3.

Lemma (4.6). Let $F_{\bar{\eta}}$ be a CDF of the probability distribution $\bar{\eta} \in \mathscr{P}(\mathbb{R})^{S}$. Then a sketch is Bellman unbiased if and only if the sketch is a homogeneous of degree k, i.e., there exists some vector-valued function $h = h(x_1, \dots, x_k) : \mathcal{X}^k \to \mathbb{R}^N$ such that

$$\psi(\bar{\eta}) = \int \cdots \int h(x_1, \cdots, x_k) dF_{\bar{\eta}}(x_1) \cdots dF_{\bar{\eta}}(x_k).$$

Proof. (\Rightarrow) Consider an two-stage MDP with a single action a, and an initial state s_0 which transits to one of terminal state $\{s_1, \dots, s_K\}$ with transition kernel $\mathbb{P}(\cdot|s_0, a)$. Assume that the reward $r(s_0) = 0$. Then $\bar{\eta}(s_0) = \sum_{k=1}^K \mathbb{P}(s_k) \delta_{r(s_k)}$. Note that s'_1, \dots, s'_k are independent and identically distributed random variable in distribution $\mathbb{P}(\cdot|s, a)$.

$$\begin{split} & \mathbb{E}_{s' \sim \mathbb{P}(\cdot|s_0,a)} \left[\phi_{\psi} \left(\psi \Big((\mathcal{B}_r)_{\#} \bar{\eta}(s_1') \Big), \cdots, \psi \Big((\mathcal{B}_r)_{\#} \bar{\eta}(s_k') \Big) \right) \right] = \psi_{1:N} \Big((\mathcal{B}_r)_{\#} \mathbb{E}_{s' \sim \mathbb{P}(\cdot|s_0,a)} [\bar{\eta}(s')] \Big) \\ & \Longrightarrow \mathbb{E}_{s' \sim \mathbb{P}(\cdot|s_0,a)} \left[\phi_{\psi} \left(\psi \Big(\delta_{r(s_1')} \Big), \cdots, \psi \Big(\delta_{r(s_k')} \Big) \Big) \right] = \psi \Big(\mathbb{E}_{s' \sim \mathbb{P}(\cdot|s_0,a)} [\delta_{r(s')}] \Big) \\ & \Longrightarrow \mathbb{E}_{s' \sim \mathbb{P}(\cdot|s_0,a)} \left[\phi_{\psi} \Big(g(s_1'), \cdots, g(s_k') \Big) \right] = \psi \Big(\bar{\eta}(s_0) \Big) \\ & \Longrightarrow \int \cdots \int h(s_1', \cdots, s_k') dF_{\bar{\eta}}(s_1') \cdots dF_{\bar{\eta}}(s_k') = \psi \Big(\bar{\eta}(s_0) \Big). \end{split}$$

(⇐)

$$\begin{split} \psi\Big((\mathcal{B}_{r})_{\#}\mathbb{E}_{s'\sim\mathbb{P}(\cdot|s,a)}[\bar{\eta}(s')]\Big) \\ &= \int \cdots \int h(x_{1},\cdots,x_{k})dF_{(\mathcal{B}_{r})_{\#}\mathbb{E}_{s'\sim\mathbb{P}(\cdot|s,a)}[\bar{\eta}(s')]}(x_{1}),\cdots,dF_{(\mathcal{B}_{r})_{\#}\mathbb{E}_{s'\sim\mathbb{P}(\cdot|s,a)}[\bar{\eta}(s')](x_{k})} \\ &= \int \cdots \int h(x_{1}+r,\cdots,x_{k}+r)d\Big(\mathbb{E}_{s'\sim\mathbb{P}(\cdot|s,a)}F_{\bar{\eta}(s')}(x_{1})\Big),\cdots,d\Big(\mathbb{E}_{s'\sim\mathbb{P}(\cdot|s,a)}F_{\bar{\eta}(s')}(x_{k})\Big) \\ &= \mathbb{E}_{s'\sim\mathbb{P}(\cdot|s,a)}\left[\int \cdots \int h(x_{1}+r,\cdots,x_{k}+r)dF_{\bar{\eta}(s')}(x_{1})\cdots dF_{\bar{\eta}(s')}(x_{k})\right] \\ &= \mathbb{E}_{s'\sim\mathbb{P}(\cdot|s,a)}\left[\psi\Big((\mathcal{B}_{r})_{\#}[\bar{\eta}(s')]\Big)\right] \end{split}$$

Theorem (4.7). The only finite statistical functionals that are Bellman unbiased and closed are given by the collections of ψ_1, \ldots, ψ_N where its linear span $\{\sum_{n=0}^N \alpha_n \psi_n | \alpha_n \in \mathbb{R}, \forall N\}$ is equal to the set of exponential polynomial functionals $\{\eta \to \mathbb{E}_{Z \sim \eta}[Z^l \exp(\lambda Z)] | l = 0, 1, \ldots, L, \lambda \in \mathbb{R}\}$, where ψ_0 is the constant functional equal to 1. In discount setting, it is equal to the linear span of the set of moment functionals $\{\eta \to \mathbb{E}_{Z \sim \eta}[Z^l] | l = 0, 1, \ldots, L\}$ for some $L \leq N$.

Proof. Our proof is mainly based on the proof techniques of Rowland et al. [41] and we describe in an extended form. Since their proof also considers the discounted setting, we will define $\mathcal{B}_{r,\gamma}(x) = r + \gamma x$ for discount factor $\gamma \in [0, 1)$. By assumption of Bellman closedness, $\psi_n((\mathcal{B}_{r,\gamma})_{\#}\bar{\eta}(s'))$ will

be written as $g(r, \gamma, \psi_{1:N}(\bar{\eta}(s')))$ for some g. By assumption of Bellman unbiasedness and Lemma 4.6, both $\psi_{1:N}(\bar{\eta}(s'))$ and $\psi_n((\mathcal{B}_{r,\gamma})_{\#}\bar{\eta}(s'))$ are affine as functions of the distribution $\bar{\eta}(s')$,

$$\begin{split} \psi_{1:N}(\alpha \bar{\eta}_1(s') + (1-\alpha) \bar{\eta}_2(s')) \\ &= \mathbb{E}_{Z_i \sim \alpha \bar{\eta}_1(s') + (1-\alpha) \bar{\eta}_2(s')} [h_{1:N}(\bar{Z}_1, \cdots, \bar{Z}_k)] \\ &= \alpha \mathbb{E}_{\bar{Z}_i \sim \bar{\eta}_1(s')} [h_{1:N}(\bar{Z}_1, \cdots, \bar{Z}_k)] + (1-\alpha) \mathbb{E}_{\bar{Z}_i \sim \bar{\eta}_2(s')} [h_{1:N}(\bar{Z}_1, \cdots, \bar{Z}_k)] \\ &= \alpha \psi_{1:N}(\bar{\eta}_1(s')) + (1-\alpha) \psi_{1:N}(\bar{\eta}_2(s')) \end{split}$$

and

$$\begin{split} \psi_n \Big((\mathcal{B}_{r,\gamma})_{\#} (\alpha \bar{\eta}_1(s') + (1-\alpha) \bar{\eta}_2(s')) \Big) \\ &= \mathbb{E}_{Z_i \sim \alpha \bar{\eta}_1(s') + (1-\alpha) \bar{\eta}_2(s')} [h_n(r+\gamma \bar{Z}_1, \cdots, r+\gamma \bar{Z}_k)] \\ &= \alpha \mathbb{E}_{\bar{Z}_i \sim \bar{\eta}_1(s')} [h_n(r+\gamma \bar{Z}_1, \cdots, r+\gamma \bar{Z}_k)] + (1-\alpha) \mathbb{E}_{\bar{Z}_i \sim \bar{\eta}_2(s')} [h_n(r+\gamma \bar{Z}_1, \cdots, r+\gamma \bar{Z}_k)] \\ &= \alpha \psi_n \Big((\mathcal{B}_{r,\gamma})_{\#} \bar{\eta}_1(s') \Big) + (1-\alpha) \psi_n \Big((\mathcal{B}_{r,\gamma})_{\#} \bar{\eta}_2(s') \Big) \end{split}$$

Therefore, $g(r, \gamma, \cdot)$ is also affine on the convex codomain of $\psi_{1:N}$. Thus, we have

$$\mathbb{E}_{\bar{Z}_i \sim \bar{\eta}}[\phi_{\psi_n}(r + \gamma \bar{Z}_1, \cdots, r + \gamma \bar{Z}_k)] = a_0(r, \gamma) + \sum_{n'=1}^N a_{n'}(r, \gamma) \mathbb{E}_{\bar{Z}_i \sim \bar{\eta}}[\phi_{\psi_{n'}}(\bar{Z}_1, \cdots, \bar{Z}_k)]$$

for some function $a_{0:N} : \mathbb{R} \times [0,1] \to \mathbb{R}$. By taking $\bar{\eta}(s') = \delta_x$, we obtain

$$\phi_{\psi_n}(r+\gamma x,\cdots,r+\gamma x) = a_0(r,\gamma) + \sum_{n'=1}^N a_{n'}(r,\gamma)\phi_{\psi_{n'}}(x,\cdots,x)$$

According to Engert [18], for any translation invariant finite-dimensional space is spanned by a set of function of the form

$$\{x \mapsto x^l \exp(\lambda_j x) | j \in [J], 0 \le l \le L\}$$

for some finite subset $\{\lambda_1, \dots, \lambda_J\}$ of \mathbb{C} . Hence, each function $x \mapsto \phi_{\psi_n}(x, \dots, x)$ is expressed as linear combination of exponential polynomial functions. In addition, the linear combination of ϕ_{ψ_n} should be closed under composition with for any discount factor $\gamma \in [0, 1]$, all λ_j should be zero. Hence, the linear combination of $\phi_{\psi_1}, \dots, \phi_{\psi_N}$ must be equal to the span of $\{x \mapsto x^l \mid 0 \le l \le L\}$ for some $L \in \mathbb{N}$.

Lemma (6.3). Consider a fixed $k \in [K]$ and a fixed $h \in [H]$. Let $\mathcal{Z}_{h}^{k} = \{(s_{h}^{\tau}, a_{h}^{\tau})\}_{\tau \in [k-1]}$ and $\mathcal{D}_{h,\bar{\eta}}^{k} = \{(s_{h}^{\tau}, a_{h}^{\tau}), \psi_{1:N}((\mathcal{B}_{r_{h'}}^{\tau}) \# \bar{\eta}(s_{h'+1}^{\tau}))\}_{\tau \in [k-1]}$ for any $\bar{\eta} : S \to \mathcal{P}([0, H])$. Define $\tilde{f}_{h,\bar{\eta}}^{k} = \arg\min_{f \in \mathcal{F}^{N}} \|f\|_{\mathcal{D}_{h,\bar{\eta}}^{k}}^{2}$. For any $\bar{\eta}$ and $\delta \in (0, 1)$, there is an event $\mathcal{E}(\bar{\eta}, \delta)$ such that conditioned on $\mathcal{E}(\bar{\eta}, \delta)$, with probability at least $1-\delta$, for any $\bar{\eta}' : S \to \mathcal{P}([0, H])$ with $\|\psi_{1:N}(\bar{\eta}') - \psi_{1:N}(\bar{\eta})\|_{\infty,1} \leq 1/T$ or $\sum_{n=1}^{N} \|\psi_{n}(\bar{\eta}') - \psi_{n}(\bar{\eta})\|_{\infty} \leq 1/T$, we have

$$\left\|\tilde{f}_{h,\bar{\eta}'}(\cdot,\cdot) - \psi_{1:N}\left((\mathcal{B}_{r(\cdot,\cdot)})_{\#}[\mathbb{P}\bar{\eta}'](\cdot,\cdot) \right) \right\|_{\mathcal{Z}_{h}^{k}} \leq c' \left(N^{\frac{1}{2}} H \sqrt{\log(1/\delta) + \log \mathcal{N}(\mathcal{F}^{N}, 1/T)} \right)$$

for some constant c' > 0.

Proof. Define the sketch of target $f_{\bar{\eta}} : S \times A \to \mathbb{R}^N$,

$$f_{\bar{\eta}}(\cdot,\cdot) \coloneqq \psi_{1:N}\Big((\mathcal{B}_{r(\cdot,\cdot)})_{\#}[\mathbb{P}\bar{\eta}](\cdot,\cdot)\Big)$$

for all $i \in [N]$.

$$\begin{split} \|f\|_{\mathcal{D}_{h,\eta'}^{k}}^{2} &- \|f_{\bar{\eta}'}\|_{\mathcal{D}_{h,\eta'}^{k}}^{2} \\ &= \sum_{n=1}^{N} \sum_{s_{h}^{\tau}, a_{h}^{\tau} \in \mathcal{Z}_{h,\bar{\eta}'}^{k}} \left(f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n} \Big((\mathcal{B}_{r_{h}^{\tau}})_{\#} \bar{\eta}'(s_{h+1}^{\tau}) \Big) \Big)^{2} - \Big(f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n} \Big((\mathcal{B}_{r_{h}^{\tau}})_{\#} \bar{\eta}'(s_{h+1}^{\tau}) \Big) \Big)^{2} \\ &= \sum_{n=1}^{N} \sum_{s_{h}^{\tau}, a_{h}^{\tau} \in \mathcal{Z}_{h,\bar{\eta}'}^{k}} (f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}))^{2} \\ &+ 2(f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau})) \Big(f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n} \Big((\mathcal{B}_{r_{h}^{\tau}})_{\#} \bar{\eta}'(s_{h+1}^{\tau}) \Big) \Big) \\ &\geq \|f - f_{\bar{\eta}'}\|_{\mathcal{Z}_{h}^{k}}^{2} - 4 \sum_{n=1}^{N} \|f_{\bar{\eta}}^{(n)} - f_{\bar{\eta}'}^{(n)}\|_{\infty} (H+1) |\mathcal{Z}_{h}^{k}| \\ &+ \sum_{n=1}^{N} \sum_{s_{h}^{\tau}, a_{h}^{\tau} \in \mathcal{Z}_{h,\bar{\eta}'}^{k}} \Big[\underbrace{ 2(f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n} \Big((\mathcal{B}_{r_{h}^{\tau}})_{\#} \bar{\eta}(s_{h+1}^{\tau}) \Big) \Big) \Big] \\ &\geq \|f - f_{\bar{\eta}'}\|_{\mathcal{Z}_{h}^{k}}^{2} - 4N(H+1) - \Big| \sum_{n=1}^{N} \sum_{s_{h}^{\tau}, a_{h}^{\tau} \in \mathcal{Z}_{h,\bar{\eta}'}^{k}} \chi_{h}^{\tau}(f^{(n)}) \Big|. \end{split}$$

For the first inequality, we change the second term from $\bar{\eta}'$ to $\bar{\eta}$ which are the ϵ -covers. Notice that $AC - BC' \ge -|AC - BC'| \ge -|(A - B)C| - |(A - B)C'| \ge -2|A - B||\max(C, C')|$.

$$\begin{split} (f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau})) \Big(f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n}\Big((\mathcal{B}_{r_{h}^{\tau}})_{\#}\bar{\eta}'(s_{h+1}^{\tau}) \Big) \Big) \\ &- (f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau})) \Big(f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n}\Big((\mathcal{B}_{r_{h}^{\tau}})_{\#}\bar{\eta}(s_{h+1}^{\tau}) \Big) \Big) \\ &\geq -2 \| f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) \| \\ &\times \max\Big(\Big| f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n}\Big((\mathcal{B}_{r_{h}^{\tau}})_{\#}\bar{\eta}'(s_{h+1}^{\tau}) \Big) \Big|, \Big| f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n}\Big((\mathcal{B}_{r_{h}^{\tau}})_{\#}\bar{\eta}(s_{h+1}^{\tau}) \Big) \Big| \Big) \\ &\geq -2 \| f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) \| (H+1) \end{split}$$

For the second inequality, consider $\bar{\eta}' : S \to \mathscr{P}([0, H])$ with $\sum_{n=1}^{N} \|\psi_n(\bar{\eta}') - \psi_n(\bar{\eta})\|_{\infty} \leq 1/T$. We have

$$\begin{split} \|f_{\bar{\eta}}^{(n)} - f_{\bar{\eta}'}^{(n)}\|_{\infty} &= \max_{s,a} \Big| \sum_{n'=1}^{n} H^{n'}[\psi_{n'}([\mathbb{P}\bar{\eta}](s,a)) - \psi_{n'}([\mathbb{P}\bar{\eta}'](s,a))]r^{n-n'}/H^{n-1} \Big| \\ &\leq \sum_{n'=1}^{n} \max_{s'} \Big| \psi_{n'}(\bar{\eta}(s')) - \psi_{n'}(\bar{\eta}'(s')) \Big| \\ &\leq 1/T. \end{split}$$

Defining \mathbb{F}_h^k as the filtration induced by the sequence $\{(s_{h'}^{\tau}, a_{h'}^{\tau})\}_{\tau, h' \in [k-1] \times [H]} \cup \{(s_1^k, a_1^k), (s_2^k, a_2^k), \dots, (s_h^k, a_h^k)\}$, notice that

$$\begin{split} & \mathbb{E}\Big[\sum_{n=1}^{N} \chi_{h}^{\tau}(f^{(n)}) \Big| \mathbb{F}_{h}^{\tau} \Big] \\ &= \sum_{n=1}^{N} 2(f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}))(f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \mathbb{E}\Big[\psi_{n}\Big((\mathcal{B}_{r_{h}^{\tau}})_{\#}\bar{\eta}(s_{h+1}^{\tau})\Big) \Big| \mathbb{F}_{h}^{\tau} \Big] \big) \\ &= \sum_{n=1}^{N} 2(f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}))(f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \mathbb{E}_{s_{h+1}^{\tau} \sim \mathbb{P}_{h}(\cdot|s_{h}^{\tau}, a_{h}^{\tau})}\Big[\psi_{n}\Big((\mathcal{B}_{r_{h}^{\tau}})_{\#}\bar{\eta}(s_{h+1}^{\tau})\Big)\Big] \big) \\ &= \sum_{n=1}^{N} 2(f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}))(f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n}\Big((\mathcal{B}_{r_{h}^{\tau}})_{\#}\mathbb{E}_{s_{h+1}^{\tau} \sim \mathbb{P}_{h}(\cdot|s_{h}^{\tau}, a_{h}^{\tau})}[\bar{\eta}(s_{h+1}^{\tau})]\Big) \big) \\ &= 0 \end{split}$$

and

$$\begin{split} \left| \sum_{n=1}^{N} \chi_{h}^{\tau}(f^{(n)}) \right| &= \left| \sum_{n=1}^{N} 2(f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}))(f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n} \Big((\mathcal{B}_{r_{h}^{\tau}})_{\#} \bar{\eta}(s_{h+1}^{\tau}) \Big)) \right| \\ &\leq \max_{n \in [N]} \Big\{ 2(f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - \psi_{n} \Big((\mathcal{B}_{r_{h}^{\tau}})_{\#} \bar{\eta}(s_{h+1}^{\tau}) \Big)) \Big\} \sum_{n=1}^{N} \left| f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) \Big| \\ &\leq 2(H+1) \sum_{n=1}^{N} \left| f^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) - f_{\bar{\eta}}^{(n)}(s_{h}^{\tau}, a_{h}^{\tau}) \right| \end{split}$$

In third equality, we emphasize that only Bellman unbiased sketch can derive the martingale difference sequence which induce the concentration result. Since every moment functional is commutable with mixing operation, the transformation ϕ_{ψ_n} in Definition 4.5 is identity for all $n \in [N]$. Hence, we choose the sketch as moment which already knows ϕ_{ψ} .

By Azuma-Hoeffding inequality,

$$\mathbb{P}\Big[\Big|\sum_{(\tau,h)\in[k-1]\times[H]}\sum_{n=1}^{N}\chi_{h}^{\tau}(f^{(n)})\Big| \ge \epsilon\Big] \le 2\exp\Big(-\frac{\epsilon^{2}}{2(2(H+1))^{2}\sum_{(\tau,h)\in[k-1]\times[H]}\Big(\sum_{n=1}^{N}|f^{(n)}-f_{\bar{\eta}}^{(n)}|\Big)^{2}}\Big)$$
$$\le 2\exp\Big(-\frac{\epsilon^{2}}{2(2(H+1))^{2}\sum_{(\tau,h)\in[k-1]\times[H]}\Big(N\sum_{n=1}^{N}|f^{(n)}-f_{\bar{\eta}}^{(n)}|^{2}\Big)}\Big)$$
$$= 2\exp\Big(-\frac{\epsilon^{2}}{2N(2(H+1))^{2}\|f-f_{\bar{\eta}}\|_{\mathcal{Z}_{h}^{k}}^{2}}\Big)$$

where the second inequality follows from the Cauchy-Schwartz inequality. We set

$$\epsilon = \sqrt{8N(H+1)^2 \|f - f_{\bar{\eta}}\|_{\mathcal{Z}_h^k}^2 \log\left(\frac{\mathcal{N}(\mathcal{F}^N, 1/T)}{\delta}\right)}$$

With union bound for all $f \in C(\mathcal{F}^N, 1/T)$, with probability at least $1 - \delta$,

$$\Big|\sum_{(\tau,h)\in[k-1]\times[H]}\sum_{n=1}^{N}\chi_{h}^{\tau}(f^{(n)})\Big| \leq c'N^{\frac{1}{2}}(H+1)\|f-f_{\bar{\eta}}\|_{\mathcal{Z}_{h}^{k}}\sqrt{\log\left(\frac{\mathcal{N}(\mathcal{F}^{N},1/T)}{\delta}\right)}$$

for some constant c' > 0.

For all $f \in \mathcal{F}^N$, there exists $g \in \mathcal{C}(\mathcal{F}^N, 1/T)$, such that $||f - g||_{\infty,1} \leq 1/T$ or $\sum_{n=1}^N ||f^{(n)} - g^{(n)}||_{\infty} \leq 1/T$ for all $n \in [N]$,

$$\begin{split} \Big| \sum_{(\tau,h)\in[k-1]\times[H]} \sum_{n=1}^{N} \chi_{h}^{\tau}(f^{(n)}) \Big| &\leq \Big| \sum_{(\tau,h)\in[k-1]\times[H]} \sum_{n=1}^{N} \chi_{h}^{\tau}(g^{(n)}) \Big| + 2(H+1) |\mathcal{Z}_{h}^{k}| \sum_{n=1}^{N} \frac{1}{T} \\ &\leq c' N^{\frac{1}{2}} (H+1) |\|g - f_{\bar{\eta}}\|_{\mathcal{Z}_{h}^{k}} \sqrt{\log\left(\frac{\mathcal{N}(\mathcal{F}^{N}, 1/T)}{\delta}\right)} + 2N(H+1) \\ &\leq c' N^{\frac{1}{2}} (H+1) (\|f - f_{\bar{\eta}}\|_{\mathcal{Z}_{h}^{k}} + 1) \sqrt{\log\left(\frac{\mathcal{N}(\mathcal{F}^{N}, 1/T)}{\delta}\right)} + 2N(H+1) \\ &\leq c' N^{\frac{1}{2}} (H+1) (\|f - f_{\bar{\eta}'}\|_{\mathcal{Z}_{h}^{k}} + 2) \sqrt{\log\left(\frac{\mathcal{N}(\mathcal{F}^{N}, 1/T)}{\delta}\right)} + 2N(H+1) \end{split}$$

where the third inequality follows from,

$$\begin{split} \|f - g\|_{\mathcal{Z}_h^k}^2 &\leq \sum_{n=1}^N \sum_{(\tau,h) \in [k-1] \times [H]} |f^{(n)}(s_h^{\tau}, a_h^{\tau}) - g^{(n)}(s_h^{\tau}, a_h^{\tau})|^2 \\ &\leq NT \Big(\frac{1}{T}\Big)^2 \\ &\leq 1. \end{split}$$

Recall that $\tilde{f}_{h,\eta'}^k = \arg\min_{f\in\mathcal{F}} \|f\|_{\mathcal{D}_{h,\eta'}^k}^2$. We have $\|\tilde{f}_{h,\eta'}^k\|_{\mathcal{D}_{h,\eta'}^k}^2 - \|f_{\bar{\eta}'}\|_{\mathcal{D}_{h,\eta'}^k}^2 \leq 0$, which implies,

$$\begin{split} 0 &\geq \|\tilde{f}_{h,\bar{\eta}'}^{k}\|_{\mathcal{D}_{h,\bar{\eta}'}^{k}}^{2} - \|f_{\bar{\eta}'}\|_{\mathcal{D}_{h,\bar{\eta}'}^{k}}^{2} \\ &= \|\tilde{f}_{h,\bar{\eta}'}^{k} - f_{\bar{\eta}'}\|_{\mathcal{Z}_{h}^{k}}^{2} \\ &+ 2\sum_{n=1}^{N}\sum_{(\tau,h)\in[k-1]\times[H]} \left[((\tilde{f}_{h,\bar{\eta}'}^{k})^{(n)}(s_{h}^{\tau},a_{h}^{\tau}) - f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau},a_{h}^{\tau}))((f_{\bar{\eta}'}^{(n)}(s_{h}^{\tau},a_{h}^{\tau}) - \psi_{n}\Big((\mathcal{B}_{r_{h}^{\tau}})_{\#}\bar{\eta}'(s_{h+1}^{\tau})\Big)) \right] \\ &\geq \|\tilde{f}_{h,\bar{\eta}'}^{k} - f_{\bar{\eta}'}\|_{\mathcal{Z}_{h}^{k}}^{2} - c'N^{\frac{1}{2}}(H+1)(\|\hat{f}_{h,\bar{\eta}'}^{k} - f_{\bar{\eta}'}\|_{\mathcal{Z}_{h}^{k}}^{2} + 2)\sqrt{\log(2/\delta) + \log\mathcal{N}(\mathcal{F}^{N}, 1/T)} - 6N(H+1). \end{split}$$

Recall that if $x^2 - 2ax - b \le 0$ holds for constant a, b > 0, then $x \le a + \sqrt{a^2 + b} \le c' \cdot a$ for some constant c' > 0.

Hence,

$$\|\tilde{f}_{h,\eta'}^k - f_{\bar{\eta}'}\|_{\mathcal{Z}_h^k} \le c'(N^{\frac{1}{2}}H\sqrt{\log(1/\delta)} + \log\mathcal{N}(\mathcal{F}^N, 1/T))$$

for some constant c' > 0.

Lemma (6.4). Let $(\mathcal{F}^N)_h^k = \{f \in \mathcal{F}^N | \|f - \tilde{f}_{h,\bar{\eta}}^k\|_{\mathcal{Z}_h^k}^2 \leq \beta(\mathcal{F}^N, \delta)\}$, where $\beta(\mathcal{F}^N, \delta) \geq c' \cdot NH^2(\log(T/\delta) + \log \mathcal{N}(\mathcal{F}^N, 1/T))$

for some constant c' > 0. Then with probability at least $1 - \delta/2$, for all $k, h \in [K] \times [H]$, we have

$$\psi_n\Big((\mathcal{B}_{r_h(\cdot,\cdot)})_{\#}[\mathbb{P}_h\bar{\eta}_{h+1}^k](\cdot,\cdot)\Big)\in(\mathcal{F}^N)_h^k$$

Proof. For all $(k, h) \in [K] \times [H]$,

$$\mathbf{S} \coloneqq \begin{cases} \left\{ \left(\min\{f^{(1)}(\cdot, \cdot) + b_{h+1}^k(\cdot, \cdot), H\} \right) \middle| f \in \mathcal{C}(\mathcal{F}^N, 1/T) \right\} \cup \left\{ 0 \right\} & n = 1 \\ \left\{ \left(\min\{f^{(n)}(\cdot, \cdot), H\} \right) \middle| f \in \mathcal{C}(\mathcal{F}^N, 1/T) \right\} \cup \left\{ 0 \right\} & 2 \le n \le N \end{cases}$$

is a (1/T)-cover of $\psi_{1:N}(\eta_{h+1}^k(\cdot, \cdot))$ where

$$\psi_{1:N}(\eta_{h+1}^k(\cdot,\cdot)) = \begin{cases} \min\{(f_{h+1}^k)^{(1)}(\cdot,\cdot) + b_{h+1}^k(\cdot,\cdot),H\} & n = 1 \text{ and } h < H \\ \min\{(f_{h+1}^k)^{(n)}(\cdot,\cdot),H\} & 2 \le n \le N \text{ and } h < H \\ \mathbf{0}^N & h = H \end{cases}$$

i.e., there exists $\psi_{1:N}(\eta) \in \mathbf{S}$ such that $\|\psi_{1:N}(\eta) - \psi_{1:N}(\eta_{h+1}^k)\|_{\infty,1} \leq 1/T$. This implies

$$\bar{\mathbf{S}} \coloneqq \left\{ \psi_{1:N} \Big(\eta(\cdot, \arg\max_{a \in \mathcal{A}} \psi_1(\eta(\cdot, a))) \Big) \mid \psi_{1:N}(\eta) \in \mathbf{S} \right\}$$

is a (1/T)-cover of $\psi_{1:N}(\bar{\eta}_{h+1}^k)$ with $\log(|\bar{\mathbf{S}}|) \leq \log \mathcal{N}(\mathcal{F}^N, 1/T)$.

For each $\psi_{1:N}(\bar{\eta}) \in \bar{\mathbf{S}}$, let $\mathcal{E}(\bar{\eta}, \delta/2|\bar{\mathbf{S}}|T)$ be the event defined in Lemma 6.3. By union bound for all $\psi_{1:N}(\bar{\eta}) \in \bar{\mathbf{S}}$, we have $\Pr[\bigcap_{\psi_{1:N}(\bar{\eta})\in \bar{\mathbf{S}}} \mathcal{E}(\bar{\eta}, \delta/2|\bar{\mathbf{S}}|T)] \ge 1 - \delta/2T$.

Let $\psi_{1:N}(\bar{\eta}) \in \bar{\mathbf{S}}$ such that $\|\psi_{1:N}(\bar{\eta}) - \psi_{1:N}(\bar{\eta}_{h+1}^k)\|_{\infty,1} \leq 1/T$. Conditioned on $\bigcap_{s_N(\bar{\eta})\in\bar{\mathbf{S}}} \mathcal{E}(\bar{\eta}, \delta/2|\bar{\mathbf{S}}|T)$ and by Lemma 6.3, we have

$$\left\| \tilde{f}_{h,\bar{\eta}}^{k}(\cdot,\cdot) - \psi_{1:N} \Big((\mathcal{B}_{r_{h}(\cdot,\cdot)})_{\#} [\mathbb{P}_{h} \bar{\eta}_{h+1}^{k}](\cdot,\cdot) \Big) \right\|_{\mathcal{Z}_{h}^{k}}^{2} \leq c' \Big(NH^{2}(\log(T/\delta) + \log\mathcal{N}(\mathcal{F}^{N}, 1/T)) \Big)$$

for some constant c' > 0.

By union bound for all $(k,h) \in [K] \times [H]$, we have $\psi_{1:N} \left((\mathcal{B}_{r_h(\cdot,\cdot)})_{\#} [\mathbb{P}_h \bar{\eta}_{h+1}^k](\cdot,\cdot) \right) \in (\mathcal{F}^N)_h^k$ with probability $1 - \delta/2$.

Lemma C.1. Let $Q_h^k(s, a) \coloneqq \min\{H, \tilde{f}_h^k(s, a) + b_h^k(s, a)\}$ for some bonus function $b_h^k(s, a)$ for all $(s, a) \in S \times A$. If $b_h^k(s, a) \ge w^{(1)}((\mathcal{F}^N)_h^k, s, a)$, then with probability at least $1 - \delta/2$,

 $Q_h^*(s,a) \leq Q_h^k(s,a) \text{ and } V_h^*(s) \leq V_h^k(s)$

for all $(k,h) \in [K] \times [H]$, for all $(s,a) \in \mathcal{S} \times \mathcal{A}$.

Proof. We use induction on h from h = H to 1 to prove the statement. Let \mathcal{E} be the event that for $(k,h) \in [K] \times [H], \psi_{1:N} \left((\mathcal{B}_{r_h(\cdot,\cdot)})_{\#} [\mathbb{P}_h \bar{\eta}_{h+1}^k](\cdot,\cdot) \right) \in (\mathcal{F}^N)_h^k$. By Lemma 6.4, $\Pr|\mathcal{E}| \ge 1 - \delta/2$. In the rest of the proof, we condition on \mathcal{E} .

When h = H + 1, the desired inequality holds as $Q_{H+1}^*(s, a) = V_{H+1}^*(s) = Q_{H+1}^k(s, a) = V_{H+1}^k(s) = 0$. Now, assume that $Q_{h+1}^*(s, a) \leq Q_{h+1}^k(s, a)$ and $V_{h+1}^*(s) \leq V_{h+1}^k(s)$ for some $h \in [H]$. Then, for all $(s, a) \in \mathcal{S} \times \mathcal{A}$,

$$\begin{aligned} Q_h^*(s, a) &= \min\{H, r_h(s, a) + [\mathbb{P}_h V_{h+1}^*](s, a)\} \\ &\leq \min\{H, r_h(s, a) + [\mathbb{P}_h V_{h+1}^k](s, a)\} \\ &\leq \min\{H, \tilde{f}_h^k(s, a) + w^{(1)}(\mathcal{F}_h^k, s, a)\} \\ &= \min\{H, Q_h^k(s, a) - b_h^k(s, a) + w^{(1)}(\mathcal{F}_h^k, s, a)\} \\ &\leq Q_h^k(s, a) \end{aligned}$$

Lemma C.2 (Regret decomposition). With probability at least $1 - \delta/4$, we have

$$Reg(K) \le \sum_{k=1}^{K} \sum_{h=1}^{H} (2b_h^k(s_h^k, a_h^k) + \xi_h^k),$$

where $\xi_h^k = [\mathbb{P}_h(V_{h+1}^k - V_{h+1}^{\pi^k})](s_h^k, a_h^k) - (V_{h+1}^k(s_{h+1}^k) - V_{h+1}^{\pi^k}(s_{h+1}^k))$ is a martingale difference sequence with respect to the filtration \mathbb{F}_h^k induced by the history \mathbb{H}_h^k .

Proof. We condition on the above event \mathcal{E} in the rest of the proof. For all $(k,h) \in [K] \times [H]$, we have

$$\left\| \tilde{f}_{h,\bar{\eta}}^{k}(\cdot,\cdot) - \psi_{1:N} \left((\mathcal{B}_{r_{h}(\cdot,\cdot)})_{\#} [\mathbb{P}_{h}\bar{\eta}_{h+1}^{k}](\cdot,\cdot) \right) \right\|_{\mathcal{Z}_{h}^{k}}^{2} \leq \beta(\mathcal{F}^{N},\delta).$$

Recall that $(\mathcal{F}^N)_h^k = \{f \in \mathcal{F}^N \mid ||f - \tilde{f}_{h,\bar{\eta}}^k||_{\mathcal{Z}_h^k}^2 \leq \beta(\mathcal{F}^N, \delta)\}$ is the confidence region. Since $\psi_{1:N}\Big((\mathcal{B}_{r_h(\cdot,\cdot)})_{\#}[\mathbb{P}_h\bar{\eta}_{h+1}^k](\cdot,\cdot)\Big) \in (\mathcal{F}^N)_h^k$, then by the definition of width function $w^{(1)}(\mathcal{F}_h^k, s, a)$, for $(k,h) \in [K] \times [H]$, we have

$$w^{(1)}(\mathcal{F}_{h}^{k}, s, a) \geq \left| \psi_{1} \Big((\mathcal{B}_{r_{h}(s,a)})_{\#} [\mathbb{P}_{h} \bar{\eta}_{h+1}^{k}](s,a) \Big) - (\tilde{f}_{h,\bar{\eta}}^{k})^{(1)}(s,a) \right| \\ = \left| r_{h}(s,a) + [\mathbb{P}_{h} V_{h+1}^{k}](s,a) - (\tilde{f}_{h,\bar{\eta}}^{k})^{(1)}(s,a) \right|.$$

Recall that $Q_h^*(\cdot, \cdot) \leq Q_h^k(\cdot, \cdot)$.

$$\begin{split} \operatorname{Reg}(K) &= \sum_{k=1}^{K} V_1^*(s_1^k) - V_1^{\pi^k}(s_1^k) \\ &\leq \sum_{k=1}^{K} V_1^k(s_1^k) - V_1^{\pi^k}(s_1^k) \\ &= \sum_{k=1}^{K} Q_1^k(s_1^k, a_1^k) - Q_1^{\pi^k}(s_1^k, a_1^k) \\ &= \sum_{k=1}^{K} Q_1^k(s_1^k, a_1^k) - (r_1(s_1^k, a_1^k) + [\mathbb{P}_1 V_2^k](s_1^k, a_1^k)) + (r_1(s_1^k, a_1^k) + [\mathbb{P}_1 V_2^k](s_1^k, a_1^k)) \\ &- Q_1^{\pi^k}(s_1^k, a_1^k) \\ &\leq \sum_{k=1}^{K} w^{(1)}((\mathcal{F}^N)_1^k, s_1^k, a_1^k) + b_1^k(s_1^k, a_1^k) + [\mathbb{P}_1(V_2^k - V_2^{\pi^k})](s_1^k, a_1^k) \\ &\leq \sum_{k=1}^{K} w^{(1)}((\mathcal{F}^N)_1^k, s_1^k, a_1^k) + b_1^k(s_1^k, a_1^k) + (V_2^k(s_2^k) - V_2^{\pi^k}(s_2^k)) + \xi_1^k \\ &\vdots \\ &\leq \sum_{k=1}^{K} \sum_{h=1}^{H} (w^{(1)}((\mathcal{F}^N)_h^k, s_h^k, a_h^k) + b_h^k(s_h^k, a_h^k) + \xi_h^k) \\ &\leq \sum_{k=1}^{K} \sum_{h=1}^{H} (2b_h^k(s_h^k, a_h^k) + \xi_h^k) \end{split}$$

It remains to bound $\sum_{k=1}^{K} \sum_{h=1}^{H} b_h^k(s_h^k, a_h^k)$, for which we will exploit fact that \mathcal{F}^N has bounded eluder dimension.

Lemma C.3. If $b_h^k(s, a) \ge w^{(1)}((\mathcal{F}^N)_h^k, s, a)$ for all $(s, a) \in \mathcal{S} \times \mathcal{A}$ and $k \in [K]$ where $(\mathcal{F}^N)_h^k = \{f \in \mathcal{F}^N | \|f - \tilde{f}_{h,\bar{\eta}}^k\|_{\mathcal{Z}_h^k}^2 \le \beta(\mathcal{F}^N, \delta)\},$

then

$$\sum_{k=1}^{K} \sum_{h=1}^{H} \mathbf{1}\{b_h^k(s_h^k, a_h^k) > \epsilon\} \le \left(\frac{4\beta(\mathcal{F}^N, \delta)}{\epsilon^2} + 1\right) \dim_E(\mathcal{F}^N, \epsilon)$$

for some constant c > 0.

Proof. We first want to show that for any sequence $\{(s_1, a_1), \ldots, (s_{\kappa}, a_{\kappa})\} \subseteq S \times A$, there exists $j \in [\kappa]$ such that (s_j, a_j) is ϵ -dependent on at least $L = \lceil (\kappa - 1)/\dim_E(\mathcal{F}^N, \epsilon) \rceil$ disjoint subsequences in $\{(s_1, a_1), \ldots, (s_{j-1}, a_{j-1})\}$ with respect to \mathcal{F}^N . We demonstrate this by using the following procedure. Start with L disjoint subsequences of $\{(s_1, a_1), \ldots, (s_{j-1}, a_{j-1})\}$, $\mathcal{B}_1, \mathcal{B}_2, \ldots, \mathcal{B}_L$, which are initially empty. For each j, if (s_j, a_j) is ϵ -dependent on every $\mathcal{B}_1, \ldots, \mathcal{B}_L$, we achieve our goal so we stop the process. Else, we choose $i \in [L]$ such that (s_j, a_j) is ϵ -independent on \mathcal{B}_i and update $\mathcal{B}_i \leftarrow \mathcal{B}_i \cup \{(s_j, a_j)\}, j \leftarrow j + 1$. Since every element of \mathcal{B}_i is ϵ -independent on its predecessors, $|\mathcal{B}_i|$ cannot get bigger than $\dim_E(\mathcal{F}^N, \epsilon)$ at any point in this process. Therefore, the process stops at most step $j = L\dim_E(\mathcal{F}^N, \epsilon) + 1 \leq \kappa$.

Now we want to show that if for some $j \in [\kappa]$ such that $b_h^k(s_j, a_j) > \epsilon$, then (s_j, a_j) is ϵ -dependent on at most $4\beta(\mathcal{F}^N, \delta)/\epsilon^2$ disjoint subsequences in $\{(s_1, a_1), \ldots, (s_{j-1}, a_{j-1})\}$ with respect to \mathcal{F}^N . If $b_h^k(s_j, a_j) > \epsilon$ and (s_j, a_j) is ϵ -dependent on a subsequence of $\{(s'_1, a'_1), \ldots, (s'_l, a'_l)\} \subseteq$ $\{(s_1, a_1), \ldots, (s_\kappa, a_\kappa)\}$, it implies that there exists $f, g \in \mathcal{F}^N$ with $||f - \tilde{f}_{h,\bar{\eta}}^k||_{\mathcal{Z}_h^k}^2 \leq \beta(\mathcal{F}^N, \delta)$ and $||g - \tilde{f}_{h,\bar{\eta}}^k||_{\mathcal{Z}_h^k}^2 \leq \beta(\mathcal{F}^N, \delta)$ such that $f^{(1)}(s'_t, a'_t) - g^{(1)}(s'_t, a'_t) \geq \epsilon$. By triangle inequality, $||f - g||_{\mathcal{Z}_h^k}^2 \leq 4\beta(\mathcal{F}^N, \delta)$. On the other hand, if (s_j, a_j) is ϵ -dependent on L disjoint subsequences in $\{(s_1, a_1), \ldots, (s_\kappa, a_\kappa)\}$, then

$$4\beta(\mathcal{F}^{N},\delta) \geq \|f - g\|_{\mathcal{Z}^{k}}^{2} \geq \|f^{(1)} - g^{(1)}\|_{\mathcal{Z}^{k}}^{2} \geq L\epsilon^{2}$$

resulting in $L \leq 4\beta(\mathcal{F}^N, \delta)/\epsilon^2$. Therefore, we have $(\kappa/\dim_E(\mathcal{F}^N, \epsilon)) - 1 \leq 4\beta(\mathcal{F}^N, \delta)/\epsilon^2$ which results in

$$\kappa \leq \left(\frac{4\beta(\mathcal{F},\delta)}{\epsilon^2} + 1\right) \dim_E(\mathcal{F}^N,\epsilon)$$

Lemma C.4 (Refined version of Lemma 10 in Wang et al. [50]). If $b_h^k(s, a) \ge w^{(1)}((\mathcal{F}^N)_h^k, s, a)$ for all $(s, a) \in \mathcal{S} \times \mathcal{A}$ and $k \in [K]$, then

$$\sum_{k=1}^K \sum_{h=1}^H b_h^k(s_h^k, a_h^k) \leq H \mathrm{dim}_E(\mathcal{F}^N, 1/T).$$

Proof. We first sort the sequence $\{b_h^k(s_h^k, a_h^k)\}_{(k,h)\in[K]\times[H]}$ in a decreasing order and denote it by $\{e_1, \ldots, e_T\}(e_1 \ge e_2 \ge \cdots \ge e_T)$. By Lemma C.3, for any constant M > 0 and $e_t \ge 1/\sqrt{MT}$, we have

$$t \leq \left(\frac{4\beta(\mathcal{F}^{N},\delta)}{Me_{t}^{2}} + 1\right) \dim_{E}(\mathcal{F}^{N},\sqrt{M}e_{t}) \leq \left(\frac{4\beta(\mathcal{F}^{N},\delta)}{Me_{t}^{2}} + 1\right) \dim_{E}(\mathcal{F}^{N},1/T)$$

which implies

$$e_t \leq \left(\frac{t}{\dim_E(\mathcal{F}^N, 1/T)} - 1\right)^{-1/2} \sqrt{\frac{4\beta(\mathcal{F}^N, \delta)}{M}},$$

for $t \geq \dim_E(\mathcal{F}^N, 1/T)$. Since we have $e_t \leq H$,

$$\begin{split} \sum_{t=1}^{T} e_t &= \sum_{t=1}^{T} e_t \mathbf{1} \{ e_t < 1/\sqrt{M}T \} + \sum_{t=1}^{T} e_t \mathbf{1} \{ e_t \ge 1/\sqrt{M}T, t < \dim_E(\mathcal{F}^N, 1/T) \} \\ &+ \sum_{t=1}^{T} e_t \mathbf{1} \{ e_t \ge 1/\sqrt{M}T, t \ge \dim_E(\mathcal{F}^N, 1/T) \} \\ &\leq \frac{1}{\sqrt{M}} + H \dim_E(\mathcal{F}^N, 1/T) + \sum_{\dim_E(\mathcal{F}^N, 1/T) \le t \le T} \left(\frac{t}{\dim_E(\mathcal{F}^N, 1/T)} - 1 \right)^{-1/2} \sqrt{\frac{4\beta(\mathcal{F}^N, \delta)}{M}} \\ &\leq \frac{1}{\sqrt{M}} + H \dim_E(\mathcal{F}^N, 1/T) + 2 \left(\frac{T}{\dim_E(\mathcal{F}^N, 1/T)} - 1 \right)^{1/2} \dim_E(\mathcal{F}^N, 1/T) \sqrt{\frac{4\beta(\mathcal{F}^N, \delta)}{M}} \\ &= \frac{1}{\sqrt{M}} + H \dim_E(\mathcal{F}^N, 1/T) + \sqrt{16 \cdot \dim_E(\mathcal{F}^N, 1/T) \cdot T \cdot \beta(\mathcal{F}^N, \delta)/M}. \end{split}$$

Taking $M \to \infty$,

$$\sum_{k=1}^{K} \sum_{h=1}^{H} b_h^k(s_h^k, a_h^k) \le H \dim_E(\mathcal{F}^N, 1/T).$$

Theorem (6.5). Under Assumption 4.8, with probability at least $1 - \delta$, *SF-LSVI* achieves a regret bound of

 $\operatorname{Reg}(K) \leq 2H \operatorname{dim}_E(\mathcal{F}^N, 1/T) + 4H \sqrt{KH \log(2/\delta)}.$

Proof. Recall that $\xi_h^k = [\mathbb{P}_h(V_{h+1}^k - V_{h+1}^{\pi^k})](s_h^k, a_h^k) - (V_{h+1}^k(s_{h+1}^k) - V_{h+1}^{\pi^k}(s_{h+1}^k))$ is a martingale difference sequence where $\mathbb{E}[\xi_h^k]\mathbb{F}_h^k] = 0$ and $|\xi_h^k| \leq 2H$. By Azuma-Hoeffding's inequality, with probability at least $1 - \delta/2$,

$$\sum_{k=1}^{K} \sum_{h=1}^{H} \xi_h^k \le 4H\sqrt{KH\log(2/\delta)}.$$

Conditioning on the above event and Lemma C.4, we have

$$\begin{split} & \operatorname{Reg}(K) \leq 2 \sum_{k=1}^{K} \sum_{h=1}^{H} b_{h}^{k}(s_{h}^{k}, a_{h}^{k}) + \sum_{k=1}^{K} \sum_{h=1}^{H} \xi_{h}^{k} \\ & \leq 2H \operatorname{dim}_{E}(\mathcal{F}^{N}, 1/T) + 4H \sqrt{KH \log(2/\delta)} \end{split}$$

		L
		L