

# Considerations on measuring spin-spin correlation of hyperons in heavy-ion experiments

Diyu Shen,<sup>1,\*</sup> Jinhui Chen,<sup>1</sup> and Aihong Tang<sup>2</sup>

<sup>1</sup>*Key Laboratory of Nuclear Physics and Ion-beam Application (MOE),  
Institute of Modern Physics, Fudan University, Shanghai 200433, China*

<sup>2</sup>*Brookhaven National Laboratory, Upton, New York 11973*

(Dated: July 2024)

The significant global spin alignment observed for  $\phi$  mesons in heavy-ion collisions has sparked intense discussions about its origin and implications. One explanation suggests that fluctuations in the strong force field may introduce strong spin correlations between strange ( $s$ ) and anti-strange ( $\bar{s}$ ) quarks, leading to the global spin alignment of  $\phi$  mesons. Extending this line of research, the theoretical community has proposed studying the spin correlation between  $\Lambda$  and  $\bar{\Lambda}$  hyperons. In this paper, we construct experimental observables and make connections between them and the theoretical proposed quantities.

The concept of global polarization in heavy-ion collisions was first proposed in Ref.[1]. In off-central heavy-ion collisions, quarks are predicted to be globally polarized along the direction of angular momentum due to spin-orbital coupling in strong interactions. This effect has recently been confirmed experimentally through measurements of  $\Lambda$  global polarization [2–6]. However, another related phenomenon, vector meson spin alignment, cannot be explained solely by vorticity or other mechanisms, most of which are conventional [7–13]. The measured  $\rho_{00}$ , the 00-component of the spin density matrix (with 1/3 indicating no spin alignment), of  $\phi$  mesons is significantly above 1/3 [14], while predictions from the quark coalescence model, considering  $\Lambda$  data, suggest [7, 9]  $\rho_{00}$  below 1/3 with a much smaller deviation from 1/3. This discrepancy between hyperon polarization and vector meson spin alignment implies richer spin physics beyond global quark polarization. It was later realized that spin correlation between quarks and anti-quarks plays a crucial role in determining vector meson spin [15]. For instance, short-range correlations between  $s$  and  $\bar{s}$  quarks due to strong-force fields successfully explain the observed  $\phi$  meson spin alignment [10, 16–19].

The spin-spin correlation of spin 1/2 particles was usually studied in helicity-frame in  $e^+e^-$  collisions or  $p + p$  collisions [20, 21], and the correlation function is defined as

$$c_{ij} = \frac{f_{\uparrow\uparrow} + f_{\downarrow\downarrow} - f_{\downarrow\uparrow} - f_{\uparrow\downarrow}}{f_{\uparrow\uparrow} + f_{\downarrow\downarrow} + f_{\downarrow\uparrow} + f_{\uparrow\downarrow}}, \quad (1)$$

where  $f_{m_i m_j} = \langle m_i m_j | \hat{\rho} | m_i m_j \rangle$  denotes the fraction of fermion pairs in the spin state of  $|m_i m_j\rangle$  along spin quantization direction, with  $\hat{\rho}$  being the spin density matrix. In general, the quantization directions of particle  $i$  and  $j$  differ in helicity-frame. However, in heavy-ion collisions, we are interested in global spin-spin correlations, meaning the quantization directions of particle  $i$  and  $j$  are fixed. In this case, Eq. 1 is modified as [15]:

$$c'_{ij} = c_{ij} - P_i P_j, \quad (2)$$

where  $P_i$  and  $P_j$  are the polarization of particle  $i$  and  $j$  respectively. It is clear that the  $c'_{ij}$  represents  $\langle \hat{\sigma}_{m_i} \hat{\sigma}_{m_j} \rangle - \langle \hat{\sigma}_{m_i} \rangle \langle \hat{\sigma}_{m_j} \rangle$ , where  $\hat{\sigma}_m$  is the Pauli operator along spin quantization direction.

The left-hand side of Eq. 2 represents the global spin correlations in heavy-ion collisions and can be calculated by measuring the  $c_{ij}$  and global polarization. In the following, we demonstrate how to measure  $c_{ij}$  using the example of  $\Lambda$ - $\Lambda$  spin correlation. This method can be easily extended to other hyperons.

We begin with the proton distribution from  $\Lambda$  weak decay,

$$\frac{dN}{d \cos \theta^*} = \frac{1}{2} (1 + \alpha_\Lambda P_\Lambda \cos \theta^*), \quad (3)$$

where  $\theta^*$  is the angle, in  $\Lambda$ 's rest frame, between proton momentum and  $\Lambda$  spin quantization direction.  $P_\Lambda$  defines  $\Lambda$  polarization and  $\alpha_\Lambda$  is a decay constant. For  $\Lambda$  pairs in the spin state  $|\uparrow\uparrow\rangle$  (both are fully polarized thus  $P_\Lambda = 1$ ), the distribution of proton-pairs can be written as

$$\frac{dN_{\uparrow\uparrow}}{d \cos \theta_i^* d \cos \theta_j^*} = \frac{1}{4} [1 + \alpha_\Lambda (\cos \theta_i^* + \cos \theta_j^*) + \alpha_\Lambda^2 \cos \theta_i^* \cos \theta_j^*], \quad (4)$$

---

\* dyshen@fudan.edu.cn

where  $N_{\uparrow\uparrow}$  denotes proton pairs from  $\Lambda$  pairs in  $|\uparrow\uparrow\rangle$  state. The footnote  $i$  and  $j$  denote the first proton and the second proton, respectively. Similarly, for  $\Lambda$  pairs in the spin state of  $|\downarrow\downarrow\rangle$ ,  $|\uparrow\downarrow\rangle$ , and  $|\downarrow\uparrow\rangle$ , the proton pairs are described by

$$\frac{dN_{\downarrow\downarrow}}{d\cos\theta_i^* d\cos\theta_j^*} = \frac{1}{4} [1 - \alpha_\Lambda (\cos\theta_i^* + \cos\theta_j^*) + \alpha_\Lambda^2 \cos\theta_i^* \cos\theta_j^*], \quad (5)$$

$$\frac{dN_{\uparrow\downarrow}}{d\cos\theta_i^* d\cos\theta_j^*} = \frac{1}{4} [1 + \alpha_\Lambda (\cos\theta_i^* - \cos\theta_j^*) - \alpha_\Lambda^2 \cos\theta_i^* \cos\theta_j^*], \quad (6)$$

$$\frac{dN_{\downarrow\uparrow}}{d\cos\theta_i^* d\cos\theta_j^*} = \frac{1}{4} [1 - \alpha_\Lambda (\cos\theta_i^* - \cos\theta_j^*) - \alpha_\Lambda^2 \cos\theta_i^* \cos\theta_j^*]. \quad (7)$$

In reality,  $\Lambda$  pairs are mixture of the above spin states, with the fractions given by  $f_{m_i m_j}$ . Therefore, the general distribution of daughter proton pairs can be expressed as follows :

$$\begin{aligned} \frac{dN}{d\cos\theta_i^* d\cos\theta_j^*} &= f_{\uparrow\uparrow} \frac{dN_{\uparrow\uparrow}}{d\cos\theta_i^* d\cos\theta_j^*} + f_{\downarrow\downarrow} \frac{dN_{\downarrow\downarrow}}{d\cos\theta_i^* d\cos\theta_j^*} + f_{\uparrow\downarrow} \frac{dN_{\uparrow\downarrow}}{d\cos\theta_i^* d\cos\theta_j^*} + f_{\downarrow\uparrow} \frac{dN_{\downarrow\uparrow}}{d\cos\theta_i^* d\cos\theta_j^*} \\ &= \frac{1}{4} [1 + A\alpha_\Lambda \cos\theta_i^* + B\alpha_\Lambda \cos\theta_j^* + C\alpha_\Lambda^2 \cos\theta_i^* \cos\theta_j^*], \end{aligned} \quad (8)$$

where

$$\begin{aligned} A &= \frac{f_{\uparrow\uparrow} - f_{\downarrow\downarrow} + f_{\uparrow\downarrow} - f_{\downarrow\uparrow}}{f_{\uparrow\uparrow} + f_{\downarrow\downarrow} + f_{\uparrow\downarrow} + f_{\downarrow\uparrow}}, \\ B &= \frac{f_{\uparrow\uparrow} - f_{\downarrow\downarrow} - f_{\uparrow\downarrow} + f_{\downarrow\uparrow}}{f_{\uparrow\uparrow} + f_{\downarrow\downarrow} + f_{\uparrow\downarrow} + f_{\downarrow\uparrow}}, \\ C &= \frac{f_{\uparrow\uparrow} + f_{\downarrow\downarrow} - f_{\uparrow\downarrow} - f_{\downarrow\uparrow}}{f_{\uparrow\uparrow} + f_{\downarrow\downarrow} + f_{\uparrow\downarrow} + f_{\downarrow\uparrow}}. \end{aligned} \quad (9)$$

The parameter  $C$  is the  $c_{ij}$  defined in Eq. 1. Indeed, the parameter  $A$  could be reduced to  $(f_\uparrow - f_\downarrow)/(f_\uparrow + f_\downarrow)$  which represents the polarization of the first  $\Lambda$  by definition, where  $f_\uparrow = f_{\uparrow\uparrow} + f_{\uparrow\downarrow}$  ( $f_\downarrow = f_{\downarrow\uparrow} + f_{\downarrow\downarrow}$ ) is the fraction of it in the spin state  $|\uparrow\rangle$  ( $|\downarrow\rangle$ ). Similarly, the parameter  $B$  represents the polarization of the second  $\Lambda$ . It is reasonable to expect that parameter  $C$  reflects the spin correlation between the first and second  $\Lambda$ . Mathematically, the  $c_{ij}$  can be obtained by the integral of  $\cos\theta_i^* \cos\theta_j^*$ , it is equivalent to calculate the average  $\langle \cos\theta_i^* \cos\theta_j^* \rangle$  in the experiment.

Based on Eq. 2, we establish the connection between experimental observable and the  $\Lambda$  spin correlation as used in the context of theoretical work:

$$c'_{\Lambda\Lambda} = \frac{9}{\alpha_\Lambda^2} \langle \cos\theta_i^* \cos\theta_j^* \rangle - P_\Lambda^2. \quad (10)$$

In experiment, the spin quantization direction (orbital angular momentum direction) is estimated from the event plane, which has a finite resolution. To correct for event plane resolution, similar to Ref. [22], we project Eq. 8 onto transverse plane, resulting in :

$$\frac{dN}{d\Delta\phi_i^* d\Delta\phi_j^*} = \frac{1}{16\pi} (1 + A\alpha_\Lambda \sin\Delta\phi_i^* + B\alpha_\Lambda \sin\Delta\phi_j^* + \frac{C\alpha_\Lambda^2 \pi}{4} \sin\Delta\phi_i^* \sin\Delta\phi_j^*), \quad (11)$$

and  $c'_{\Lambda\Lambda}$  can be calculated by

$$c'_{\Lambda\Lambda} = \frac{64}{\pi^2 \alpha_\Lambda^2} \langle \sin\Delta\phi_i^* \sin\Delta\phi_j^* \rangle - P_\Lambda^2, \quad (12)$$

where  $\Delta\phi_i^*$  ( $\Delta\phi_j^*$ ) is the azimuthal angle relative to the reaction plane for proton  $i$  ( $j$ ) in first (second)  $\Lambda$  rest frame. Considering a finite event plane resolution, Eq. 12 can be expressed as

$$c'_{\Lambda\Lambda} = \frac{64}{\pi^2 \alpha_\Lambda^2} \frac{\langle \sin(\phi_i^* - \Psi_{EP}) \sin(\phi_j^* - \Psi_{EP}) \rangle}{\langle \cos^2(\Psi_{EP} - \Psi_{RP}) \rangle} - P_\Lambda^2, \quad (13)$$

where  $\Psi_{EP}$  and  $\Psi_{RP}$  denote the event plane and the reaction plane, respectively. Note that

$$\langle \cos^2(\Psi_{EP} - \Psi_{RP}) \rangle = \frac{1 + \langle \cos(2(\Psi_{EP} - \Psi_{RP})) \rangle}{2}, \quad (14)$$

and  $\langle \cos(2(\Psi_{EP} - \Psi_{RP})) \rangle$  can be determined from sub-event method in experiment [23].

Building on the structure of balance function [24, 25], we propose a spin balance function to detect the difference of spin correlation between “same sign” pairs ( $\Lambda$ - $\Lambda$  or  $\bar{\Lambda}$ - $\bar{\Lambda}$ ) and “opposite sign” pairs ( $\Lambda$ - $\bar{\Lambda}$ ),

$$B(c'_{OS}, c'_{SS}) = \frac{1}{2} \left( \frac{c'_{\Lambda\bar{\Lambda}} - c'_{\Lambda\Lambda}}{c'_{\Lambda\bar{\Lambda}} + c'_{\Lambda\Lambda}} + \frac{c'_{\bar{\Lambda}\Lambda} - c'_{\bar{\Lambda}\bar{\Lambda}}}{c'_{\bar{\Lambda}\Lambda} + c'_{\bar{\Lambda}\bar{\Lambda}}} \right). \quad (15)$$

The advantage of the above equation is that the detector effect and flavor independent correlation will be canceled, only additional correlation between  $s$  and  $\bar{s}$  (in  $\Lambda$ - $\bar{\Lambda}$  pairs) remains.

To summarize, in this paper, we investigated the correlation function used in a theoretical context, identified its connection to experimental observables, and outlined the procedure to measure it in experiments. Additionally, we proposed a method for measuring the correlation using a spin balance function, which minimize the impact of detector and other trivial effects. This work will be instrumental in studying hyperon spin correlations and facilitating the understanding of spin-spin correlations and their implications for fluctuations in the strong force field.

### ACKNOWLEDGMENT

This work was inspired by the theoretical study [15] and benefited from discussions with X.-N. Wang, Z.-T. Liang, Q. Wang, X.-L. Sheng., and Z. Tu.

- 
- [1] Z.-T. Liang and X.-N. Wang, Globally polarized quark-gluon plasma in non-central A+A collisions, *Phys. Rev. Lett.* **94**, 102301 (2005), [Erratum: *Phys.Rev.Lett.* 96, 039901 (2006)], arXiv:nucl-th/0410079.
  - [2] L. Adamczyk et al. (STAR), Global  $\Lambda$  hyperon polarization in nuclear collisions: evidence for the most vortical fluid, *Nature* **548**, 62 (2017), arXiv:1701.06657 [nucl-ex].
  - [3] J. Adam et al. (STAR), Global polarization of  $\Lambda$  hyperons in Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV, *Phys. Rev. C* **98**, 014910 (2018), arXiv:1805.04400 [nucl-ex].
  - [4] S. Acharya et al. (ALICE), Global polarization of  $\Lambda\bar{\Lambda}$  hyperons in Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  and 5.02 TeV, *Phys. Rev. C* **101**, 044611 (2020), arXiv:1909.01281 [nucl-ex].
  - [5] M. S. Abdallah et al. (STAR), Global  $\Lambda$ -hyperon polarization in Au+Au collisions at  $\sqrt{s_{NN}}=3$  GeV, *Phys. Rev. C* **104**, L061901 (2021), arXiv:2108.00044 [nucl-ex].
  - [6] M. I. Abdulhamid et al. (STAR), Global polarization of  $\Lambda$  and  $\Lambda^-$  hyperons in Au+Au collisions at sNN=19.6 and 27 GeV, *Phys. Rev. C* **108**, 014910 (2023), arXiv:2305.08705 [nucl-ex].
  - [7] Z.-T. Liang and X.-N. Wang, Spin alignment of vector mesons in non-central A+A collisions, *Phys. Lett. B* **629**, 20 (2005), arXiv:nucl-th/0411101.
  - [8] F. Becattini, L. Csernai, and D. J. Wang,  $\Lambda$  polarization in peripheral heavy ion collisions, *Phys. Rev. C* **88**, 034905 (2013), [Erratum: *Phys.Rev.C* 93, 069901 (2016)], arXiv:1304.4427 [nucl-th].
  - [9] Y.-G. Yang, R.-H. Fang, Q. Wang, and X.-N. Wang, Quark coalescence model for polarized vector mesons and baryons, *Phys. Rev. C* **97**, 034917 (2018), arXiv:1711.06008 [nucl-th].
  - [10] X.-L. Sheng, L. Oliva, and Q. Wang, What can we learn from the global spin alignment of  $\phi$  mesons in heavy-ion collisions?, *Phys. Rev. D* **101**, 096005 (2020), [Erratum: *Phys.Rev.D* 105, 099903 (2022)], arXiv:1910.13684 [nucl-th].
  - [11] X.-L. Xia, H. Li, X.-G. Huang, and H. Zhong Huang, Local spin alignment of vector mesons in relativistic heavy-ion collisions, *Phys. Lett. B* **817**, 136325 (2021), arXiv:2010.01474 [nucl-th].
  - [12] J.-H. Gao, Helicity polarization in relativistic heavy ion collisions, *Phys. Rev. D* **104**, 076016 (2021), arXiv:2105.08293 [hep-ph].
  - [13] B. Müller and D.-L. Yang, Anomalous spin polarization from turbulent color fields, *Phys. Rev. D* **105**, L011901 (2022), [Erratum: *Phys.Rev.D* 106, 039904 (2022)], arXiv:2110.15630 [nucl-th].
  - [14] M. S. Abdallah et al. (STAR), Pattern of global spin alignment of  $\phi$  and  $K^{*0}$  mesons in heavy-ion collisions, *Nature* **614**, 244 (2023), arXiv:2204.02302 [hep-ph].
  - [15] J.-p. Lv, Z.-h. Yu, Z.-t. Liang, Q. Wang, and X.-N. Wang, Global quark spin correlations in relativistic heavy ion collisions, *Phys. Rev. D* **109**, 114003 (2024), arXiv:2402.13721 [hep-ph].
  - [16] X.-L. Sheng, L. Oliva, Z.-T. Liang, Q. Wang, and X.-N. Wang, Spin Alignment of Vector Mesons in Heavy-Ion Collisions, *Phys. Rev. Lett.* **131**, 042304 (2023), arXiv:2205.15689 [nucl-th].
  - [17] X.-L. Sheng, Q. Wang, and X.-N. Wang, Improved quark coalescence model for spin alignment and polarization of hadrons, *Phys. Rev. D* **102**, 056013 (2020), arXiv:2007.05106 [nucl-th].
  - [18] X.-L. Sheng, L. Oliva, Z.-T. Liang, Q. Wang, and X.-N. Wang, Relativistic spin dynamics for vector mesons, *Phys. Rev. D* **109**, 036004 (2024), arXiv:2206.05868 [hep-ph].

- [19] A. Kumar, B. Müller, and D.-L. Yang, Spin alignment of vector mesons by glasma fields, *Phys. Rev. D* **108**, 016020 (2023), arXiv:2304.04181 [nucl-th].
- [20] K.-b. Chen, W.-h. Yang, Y.-j. Zhou, and Z.-t. Liang, Energy dependence of hadron polarization in  $e^+e^- \rightarrow hX$  at high energies, *Phys. Rev. D* **95**, 034009 (2017), arXiv:1609.07001 [hep-ph].
- [21] X. Li, Z.-X. Chen, S. Cao, and S.-Y. Wei, Correlations of dihadron polarization in central, peripheral, and ultraperipheral heavy-ion collisions, *Phys. Rev. D* **109**, 014035 (2024), arXiv:2309.09487 [hep-ph].
- [22] B. I. Abelev et al. (STAR), Global polarization measurement in Au+Au collisions, *Phys. Rev. C* **76**, 024915 (2007), [Erratum: *Phys.Rev.C* 95, 039906 (2017)], arXiv:0705.1691 [nucl-ex].
- [23] A. M. Poskanzer and S. A. Voloshin, Methods for analyzing anisotropic flow in relativistic nuclear collisions, *Phys. Rev. C* **58**, 1671 (1998), arXiv:nucl-ex/9805001.
- [24] J. Adams et al. (STAR), Narrowing of the balance function with centrality in au + au collisions at  $(S(NN))^{1/2} = 130$ -GeV, *Phys. Rev. Lett.* **90**, 172301 (2003), arXiv:nucl-ex/0301014.
- [25] S. A. Bass, P. Danielewicz, and S. Pratt, Clocking hadronization in relativistic heavy ion collisions with balance functions, *Phys. Rev. Lett.* **85**, 2689 (2000), arXiv:nucl-th/0005044.