On the mean-field limit of the Cucker-Smale model with Random Batch Method

Yuelin Wang *a and Yiwen Lin $^{\dagger a}$

^aSchool of Mathematical Sciences, MOE-LSC, Shanghai Jiao Tong University, Shanghai, 200240, P.R.China

Abstract

In this work, we focus on the mean-field limit of the Random Batch Method (RBM) for the Cucker-Smale model. Different from the classical mean-field limit analysis, the chaos in this model is imposed at discrete time and is propagated to discrete time flux. We approach separately the limits of the number of particles $N \to \infty$ and the discrete time interval $\tau \to 0$ with respect to the RBM, by using the flocking property of the Cucker-Smale model and the observation in combinatorics. The Wasserstein distance is used to quantify the difference between the approximation limit and the original mean-field limit. Also, we combine the RBM with generalized Polynomial Chaos (gPC) expansion and proposed the RBM-gPC method to approximate stochastic mean-field equations, which conserves positivity and momentum of the mean-field limit with random inputs.

 ${\bf Keywords:} \ {\rm Random \ Batch \ Method, \ mean-field \ limit, \ Cucker-Samle \ model, \ stochastic \ Galerkin.}$

1 Introduction

Collective behaviors of many-body systems are ubiquitous in the real world, like flocking of birds [18, 19, 10, 28], swarming of fishes [32], synchronicity of fireflies [26, 5] and pacemaker cells [30]. We use the jargon "flocking" to describe the phenomenon in which self-propelled particles organize into an ordered motion using only limited environmental information and simple rules [32]. Since there is a great deal of literature on collective behaviors and related models, we refer the readers to [12, 25, 1, 2, 3, 35, 17, 13, 33] and the references therein.

The Cucker-Smale model, proposed by Cucker and Smale [10], is a famous model of collective behaviors that models phenomenologically the flocking phenomenon. It is in the form of an N-body second-order system of ordinary differential equations or particle systems for position and velocity, which resemble Newton's equations.

In addition, the mean-field limit, as the number of particles $N \to \infty$, is also a prevalent tool for simplifying large particle systems, which makes use of continuum models derived within the framework of mathematical kinetic theory, to approximate collective behaviors. In these years, there has been extensive research on the mean-field limit of particle systems and we refer to [20, 8, 9] for survey articles.

As a classical type of N-body systems, in the Cucker-Smale model, each particle interacts with N-1 particles, and thus the computational cost to solve it is $\mathcal{O}(N^2)$ per time step. An efficient algorithm to reduce computational complexity to $\mathcal{O}(N)$ is the Random Batch Method (RBM) proposed by Jin et al. [22] in 2020. The RBM indeed constructs a randomly decoupled system with subsystems of interaction between p particles, where the constant $p \ll N$. In this system, at each time step, the interaction only occurs within small batches

^{*}sjtu_wyl@sjtu.edu.cn

[†]linyiwen@sjtu.edu.cn

with p particles. The choice of batches is random and thus the time-average effect makes it a good approximation of the original system [22, 15, 23].

An interesting problem is to understand the mean-field limit of the RBM. In 2022, Jin and Li considered the mean-field limit of the RBM on first-order systems with Gaussian noise in [21]. Inspired by [21], in this paper, we investigate the mean-field limit of the RBM on the Cucker-Smale model as the number of particles tends to infinity. Unlike the first-order system considered in [21], the Cucker-Smale model is a second-order system with interaction coupled with both velocity and position. Technically, moment control needs to be derived by the asymptotic flock estimate instead of using the contraction property in first-order systems in [22, 23]. Specifically, we mainly analyze the path from (b) to (c), and briefly discuss the path from (a) to (d) in the following figure (1.1). On the upper left corner, Ha et al. give an estimate on (a) in [15] and a proof of the existence of mean-field limit on (d) in [16]. Their analysis on (d) is uniform in time but implicit in N. In later work [29], an explicit estimate on N is given but only works on finite time. The four models in (1.1) are represented in Section 2.

Cucker-Smale system (2.1)
$$\xrightarrow{(d) N \to \infty} \text{ mean-field limit (2.6)}$$

$$\stackrel{(a) \tau \to 0}{\uparrow} \qquad \qquad \uparrow^{(c) \tau \to 0}$$
dom Batch Cucker-Smale model (2.10)
$$\xrightarrow{(b) N \to \infty} \text{ mean-field limit of RBM (2.11)}$$

$$(1.1)$$

With the justification of (b) and (c), we employ the RBM, which is an efficient particlebased numerical scheme for the approximation of stochastic mean-field equations of the Cucker-Smale model, inspired by [7]. We consider such an application in the stochastic setting, where the interacting force depends on a random variable modeling uncertainties. This is an important problem in Uncertainty Quantification [24]. The method combines the RBM with a generalized Polynomial Chaos (gPC) expansion in the random space and thus we call it the RBM-gPC. It avoids loss of positivity like MCgPC in [7], and moreover preserves the mean velocity during evolution.

The rest of the paper is organized as follows. In Section 2, we give a concise introduction to the Cucker-Smale model including notations, assumptions and prerequisites. Section 3 presents our main results on the analysis of limits (b) and (c) in (1.1) with helpful discussions on the limits from (a) to (d). Section 4 and Section 5 provide the proof details. In Section 6, we show some numerical experiments with the RBM-gPC. Finally, Section 7 is devoted to a summary of our main results and some remaining issues to be explored in the future.

2 Preliminaries

Ran

In this section, we introduce the main models in (1.1) with assumptions and notations. Also, some basic properties of the models are given.

2.1 The Cucker-Smale model

Let X_i and V_i be the position and velocity of the *i*-th particle with unit mass, and $\psi(|X_j - X_i|)$ be the communication weight between the *j*-th and *i*-th particles. The Cucker-Smale model reads as the following

$$\begin{cases} \frac{d}{dt}X_{i}(t) = V_{i}(t), & i = 1, \cdots, N, \\ \frac{d}{dt}V_{i}(t) = \frac{\kappa}{N-1}\sum_{j=1}^{N}\psi(|X_{j}(t) - X_{i}(t)|)(V_{j}(t) - V_{i}(t)), \end{cases}$$
(2.1)

where κ is the nonnegative coupling strength and ψ satisfies positivity, boundedness, Lipschitz continuity and mononticity conditions, i.e., there exist positive constants $\psi_0, \psi_M > 0$ such that

$$0 < \psi_0 \le \psi(r) \le \psi_M, \, \forall r \ge 0; \quad \|\psi\|_{\text{Lip}} < \infty; \quad (\psi(r_1) - \psi(r_2))(r_1 - r_2) \le 0, \, r_1, r_2 \in \mathbb{R}_+.$$
(2.2)

Without loss of generality, we set $\psi_M = 1$ in this paper for convenience. Under the condition (2.2) of ψ , it is observed in [10, 18] that the total momentum is conserved as a constant and the total energy is nonincreasing along the Cucker-Smale flow. Actually, supposing that $\{(X_i, V_i)\}$ is the solution of system (2.1), then for any t > 0, one has

$$\frac{d}{dt}\sum_{i=1}^{N}V_i(t) = 0, \quad \frac{d}{dt}\sum_{i=1}^{N}|V_i(t)|^2 = -\frac{\kappa}{N-1}\sum_{i,j}\psi(|X_j(t) - X_i(t)|)|V_j(t) - V_i(t)|^2.$$
(2.3)

A key observation of the Cucker-Smale model, given by Ha et al. in [18], is that the standard deviations of particle phase-space positions are dominated by SDDI (the system of dissipative differential inequalities):

$$\left|\frac{d\mathcal{X}}{dt}\right| \le \mathcal{V}, \quad \frac{d\mathcal{V}}{dt} \le -\phi(\mathcal{X})\mathcal{V}, \tag{2.4}$$

where $(\mathcal{X}, \mathcal{V})$ are nonnegative functions and ϕ is a nonnegtive measurable function. This is a useful tool to analyze the Cucker-Smale model. In this paper, we use a variant of (2.4) (Lemma 1) in the proof of our main result. One can find its proof in Lemma 3.1 in [16].

Lemma 1. Suppose that two nonnegative Lipschitz functions X and V satisfy the coupled differential inequalities:

$$\left|\frac{d\mathcal{X}}{dt}\right| \le \mathcal{V}, \quad \frac{d\mathcal{V}}{dt} \le -\alpha \mathcal{V} + \gamma e^{-\alpha t} \mathcal{X}, \quad a.e. \, t > 0, \tag{2.5}$$

where α and γ are positive constants. Then, \mathcal{X} and \mathcal{V} satisfy the uniform bound and decay estimates:

$$\mathcal{X}(t) \le \frac{2M}{\alpha}(\mathcal{X}(0) + \mathcal{V}(0)), \quad \mathcal{V}(t) \le M(\mathcal{X}(0) + \mathcal{V}(0))e^{-\frac{\alpha t}{2}}, \quad t \ge 0,$$

where M is given by

$$M := \max\left\{1, \frac{2\gamma}{\alpha e}\right\} + \frac{8\gamma}{\alpha^3 e^3}.$$

2.2 The mean-field limit of the Cucker-Smale model

The corresponding evolution for the distribution function $\tilde{f}(x, v, t)$ was first derived by Tadmor, etc. in [19] by the BBGKY hierarchy (e.g. [4, 12]), which follows

$$\partial_t \tilde{f} + v \cdot \nabla_x \tilde{f} + \nabla_v \cdot [\xi[\tilde{f}]\tilde{f}] = 0, \qquad (2.6)$$

where the operator ξ is defined by

$$\xi[f](x,v,t) = \kappa \int \int \psi(|x-y|)(w-v)f(y,w,t) \, dw dy.$$

It is clear that

$$\frac{\partial}{\partial t} \int \int v \tilde{f} dx dv = 0.$$
(2.7)

Similar with energy-decreasing in (2.3) in the microscopic scale, the mean-field limit has the following property:

Lemma 2 ([31], Theorem 3.1). Let f_0 be compactly supported in \mathbb{R}^{2d} with $\bar{x} = \int x f_0(x, v) dx dv$ and $\bar{v} = \int v f_0(x, v) dx dv$. Define $C_{x0} := \inf\{X \ge 0 \mid supp(f_0) \subset B(\bar{x}, X) \times \mathbb{R}^d\}$ and $C_v := \inf\{V \ge 0 | supp(f_0) \subset \mathbb{R}^d \times B(\bar{v}, V)\}, \text{ where } B(\bar{x}, X) \text{ designates the ball of center } \bar{x} \text{ and radius } X \text{ in } \mathbb{R}^d.$ Let \tilde{f}_t be the unique solution of (2.6) with $\tilde{f}(0) = f_0$. If the initial data satisfies

$$C_v < \kappa \int_{C_{x0}}^{\infty} \psi(2s) ds, \qquad (2.8)$$

1

then there exists $C_x > 0$ such that

$$supp[\tilde{f}_t] \subset B(\bar{x} + t\bar{v}, C_x) \times B(\bar{v}, C_v e^{-\kappa \psi(2C_x)t}).$$

$$(2.9)$$

Remark 1. Under assumption (2.2) of ψ , one has $\psi(2C_x) \ge \psi_0$. Actually, the assumption of the lower bound ψ_0 is for the convenience of the proof and can be relaxed if (2.8) is satisfied by both discrete and continuous systems, guaranteeing the flocking in the Cucker-Smale model.

2.3 The Wasserstein distance

To introduce the topology of the space of probability measures, here we recall the definitions of the Wasserstein space [36]. For a metric space (\mathbb{R}^d, ρ) and $q \ge 1$, define the Wasserstein space of order q by

$$\mathcal{P}_{q,\rho}(\mathbb{R}^d) := \left\{ \mu \in \mathcal{P}(\mathbb{R}); \int_{\mathbb{R}^d} \rho(x,0)^q \nu(dx) < \infty \right\},$$

endowed with the Wasserstein metric

$$W_{q,\rho}(\mu,\nu) := \left(\inf_{\pi \in \Pi(\mu,\nu)} \left\{ \int_{\mathbb{R}^d \times \mathbb{R}^d} \rho(x,y)^q \pi(dx,dy) \right\} \right)^{\frac{1}{q}},$$

where $\mu, \nu \in \mathcal{P}_q(\mathbb{R}^d)$ and $\Pi(\mu, \nu)$ is the set of probability measures on $\mathbb{R}^d \times \mathbb{R}^d$ with marginals μ and ν respectively. The W_q metric, or distance, induces a kind of weak topology and measures the closeness between distributions. We use the notation $W_{q,p}$ when $\rho(x, y) := \|x - y\|_p$ and typically abbreviate $W_q = W_{q,2}$. For convenience in notation, we simplify $\|\cdot\|$ as $|\cdot|$ without ambiguity.

Recall the Jordan decomposition for a signed measure $|\mu| := \mu^+ + \mu^-$. One can control the W_q distance by the weighted total variation by the following lemma (see e.g. Theorem 6.15 in [36]), which helps the proof of our main theorems, i.e., Theorem 1 and Theorem 2, in Section 3.2.

Lemma 3 (The Wasserstein distance is controlled by the weighted total variation). Let μ and ν be two probability measures on a Polish space (\mathfrak{X}, d) . Let $q \in [1, \infty)$ and $x_0 \in \mathfrak{X}$. Then

$$W_q(\mu,\nu) \le 2^{1-\frac{1}{q}} \left(\int d(x_0,x)^q d|\mu-\nu|(x) \right)^{\frac{1}{q}}$$

2.4 The Random Batch method and its mean-field dynamics

In each sub-time interval $[t_{k-1}, t_k)$, we set $[i]_k$ as the batch containing the particle *i*. The RBM-(approximated) system of (2.1) becomes:

$$\begin{cases} \partial_t X_i^R(t) = V_i^R(t), \quad i = 1, \cdots, N, \\ \partial_t V_i^R(t) = \frac{\kappa}{p-1} \sum_{j \in [i]_k} \psi(|X_j^R(t) - X_i^R(t)|) (V_j^R(t) - V_i^R(t)), \end{cases}$$
(2.10)

with initial data $(X_i^R(0), V_i^R(0)) = (X_i^{in}, V_i^{in}), i = 1, \dots, N$. See Algorithm 1.

In the $N \to \infty$ limit, the N-particle system is then reduced to a p-particle subsystem described by the following system for $t \in [t_k, t_{k+1})$:

$$\begin{cases} \partial_t X_i(t) = V_i(t), & i = 1, \cdots, N, \\ \partial_t V_i(t) = \frac{\kappa}{p-1} \sum_{j=1}^p \psi(|X_j(t) - X_i(t)|) (V_j(t) - V_i(t)), \end{cases}$$
(2.11)

Algorithm 1: The RBM for (2.1)

with $Z_i := (X_i, V_i)$ and $Z_i(t_k)$ being *i.i.d* drawn from $f(\cdot; t_k)$. Here $f(\cdot; t_k)$ is the law of $Z_1(t_k^-)$ for $k \ge 1$ and $f(\cdot; 0)$ equals the initial distribution f_0 . We impose $Z_1(t_k^-) = Z_1(t_k^+)$. Except the individual particle 1, the rest p-1 particles are sampled from an infinite pool of independent particles from particle 1 at each time step t_k . This process becomes the mean-field limit model of the RBM system (2.10), and one may write the following mean-field limit for the RBM in terms of the probability distribution as shown in Algorithm 2, while (2.11) becomes its microscopic description.

Algorithm 2: Mean-field Dynamics of the RBM (2.11)

- 1 From t_k to t_{k+1} the distribution function f_{t_k} will be transformed into
- $f_{t_{k+1}} := \mathcal{Q}_{\infty}(f_{t_k})$ as follows;
- **2** Let $f^{(p)}(\cdots;t_k) := f_{t_k}^{\otimes p}$ be a probability measure on \mathbb{R}^{2dp} ;
- **3** Evolve $f^{(p)}$ by time τ according to

$$\partial_t f^{(p)} + \sum_{i=1}^p \nabla_{x_i} \cdot (v_i f^{(p)}) + \sum_{i=1}^p \nabla_{v_i} \cdot \left(\xi_i f^{(p)}\right) = 0, \qquad (2.12)$$

where

$$\xi_i := \frac{\kappa}{p-1} \sum_{j=1}^p \psi(|x_j - x_i|)(v_j - v_i).$$

Set $\mathcal{Q}_{\infty}(f_{t_k}) := \int_{\mathbb{R}^{2d(p-1)}} f^{(p)}(\cdots; t_{k+1}^-) dx_2 \cdots dx_p dv_2 \cdots dv_p.$

3 The Main Result

In this paper, we show the conservation law and the mean-field limit of the Random-Batch Cucker-Smale model. First, we show the conservation of momentum and dissipation of kinetic energy of the mean-field limit system of RBM in Section 3.1. Then, we analyze the procedure (b) and (c) in Section 3.2.

Assume that the initial data of Algorithm 1 and Algorithm 2 obey the same law $f_0(x, v)dxdv$. For notational convenience, recall $\bar{x} := \int x f_0(x, v)$ and $\bar{v} := \int v f_0(x, v)$; and set $z_i := (x_i, v_i)$ for each i and

$$f(x,v;t) := \int_{\mathbb{R}^{2d(p-1)}} f^{(p)}(\cdots;t^{-}) dx_2 \cdots dx_p dv_2 \cdots dv_p = f_t(x,v).$$

Note that f is defined on \mathbb{R}^{2d} in Algorithm 2 and $f^{(p)}$ is defined on \mathbb{R}^{2dp} . Also, the definition of f is consistent with that in Algorithm 2 at $t = t_k$.

In the following, we suppose that f_0 is compactly supported and then there exists a constant R > 0 such that $\operatorname{supp}[f_0] \subset B(0, R)$, where B(0, R) is a ball with radius R centered in the origin. Without loss of generality, we set $\overline{v} = 0$ as a standardization of the initial momentum.

3.1 Basic characteristics of the mean-field limit system of RBM

Here, we show that the two basic characteristics: conservation of momentum and dissipation of kinetic energy, as observed in the Cucker-Smale model (2.1) and the RBM system (2.10), also holds for the RBM mean-field system described in Algorithm 2.

Proposition 1 (Conservation of momentum). For $t \in (t_k, t_{k+1})$, it holds

$$\frac{\partial}{\partial t} \int_{\mathbb{R}^{2d}} v f(x,v;t) dx dv = 0.$$

In particular, since $\bar{v} = 0$, $\int v f(x, v; t) = 0$ for any $t \in [0, T]$.

Proof. By the definition of f, one has

$$\begin{aligned} \frac{\partial}{\partial t} \int_{\mathbb{R}^{2d}} vf(x,v;t) dx dv &= \int_{\mathbb{R}^{2d}} \int_{\mathbb{R}^{2d(p-1)}} v \frac{\partial}{\partial t} f^{(p)}(x,\cdot,v,\cdot;t^{-}) dx dx_2 \cdots dx_p dv dv_2 \cdots dv_p \\ &= \int_{\mathbb{R}^{2dp}} v_1 \left[-\sum_{i=1}^p \nabla_{x_i} \cdot v_i f^{(p)}(\cdots;t) - \sum_{i=1}^p \nabla_{v_i} \cdot \xi_i f^{(p)}(\cdots;t) \right] dz_1 \cdots dz_p \\ &= \frac{\kappa}{p-1} \sum_{j=2}^p \int_{\mathbb{R}^{2dp}} \psi(|x_j - x_1|) (v_j - v_1) f_t^{(p)}(z_1,\cdots,z_p) dz_1 \cdots dz_p \\ &= 0. \end{aligned}$$

The last equality is derived from the fact that $\psi(|x_j - x_1|)(v_j - v_1)f_t^{(p)}$ is an odd function on (z_j, z_1) . Note that $f(x, v; t_k^-) = f(x, v; t_k) = f(x, v; t_k^+)$. Then by induction one has $\int vf(x, v; t) = \bar{v}$.

Proposition 2 (Dissipation of kinetic energy). For any $t \in (t_k, t_{k+1})$,

$$\frac{\partial}{\partial t} \int_{\mathbb{R}^{2d}} |v|^2 f(x,v;t) dx dv \le 0.$$

In particular, $M_2^v f_t \leq M_2^v f_0$, where $M_2^v f_t := \int_{\mathbb{R}^{2d}} |v|^2 f(x,v;t) dx dv$. Moreover, since $0 < \psi_0 \leq \psi$, it holds

$$M_2^v f_{k+1} \le M_2^v f_0 \left(e^{-2\kappa\psi_0\tau} + \frac{8\kappa^2\psi_0}{p-1}\tau^2 \right)^k$$

Proof. For any $t \in (t_k, t_{k+1})$,

$$\begin{aligned} \frac{\partial}{\partial t} M_2^v f_t &= \int_{\mathbb{R}^{2dp}} |v_1|^2 \left(-\sum_{i=1}^p \nabla_{x_i} \cdot (v_i f_t^{(p)}) - \sum_{i=1}^p \nabla_{v_i} \cdot (\xi_i f_t^{(p)}) \right) \\ &= \int_{\mathbb{R}^{2dp}} \frac{\kappa}{p-1} \sum_{j=1}^p \psi(|x_1 - x_j|) (v_j - v_1) \cdot 2v_1 f_t^{(p)} \\ &= -\int_{\mathbb{R}^{2dp}} \frac{\kappa}{p-1} \sum_{j=1}^p \psi(|x_1 - x_j|) |v_j - v_1|^2 f_t^{(p)} \\ &\leq 0. \end{aligned}$$

The last equality is derived by the symmetry of ψ and $f_t^{(p)}$. By the continuity of f, one has $M_2^v f_t \leq M_2^v f_0$. Since ψ has a positive lower bound ψ_0 , furthermore, it holds

$$\frac{\partial}{\partial t} M_2^v f_t \leq -\int_{\mathbb{R}^{2d_p}} \frac{\kappa}{p-1} \sum_{j=1}^p \psi_0 |v_j - v_1|^2 f_t^{(p)}
= -2\kappa \psi_0 \int_{\mathbb{R}^{2d}} |v|^2 f_t(x, v) dx dv + 2\kappa \psi_0 \int_{\mathbb{R}^{2d_p}} v_1 \cdot v_2 f_t^{(p)},$$
(3.1)

by the symmetry of $f_t^{(p)}$. For the second term in (3.1), one has

$$\frac{\partial}{\partial t} \int_{\mathbb{R}^{2dp}} v_1 \cdot v_2 f_t^{(p)} \\
= \kappa \int_{\mathbb{R}^{2dp}} \frac{v_2}{p-1} \cdot \sum_{k \neq 1}^p \psi(|x_1 - x_k|) (v_k - v_1) f_t^{(p)} + \frac{v_1}{p-1} \cdot \sum_{k \neq 2}^p \psi(|x_2 - x_k|) (v_k - v_2) f_t^{(p)} \\
= 2\kappa \int_{\mathbb{R}^{2dp}} \frac{v_2}{p-1} \cdot \sum_{k \neq 1}^p \psi(|x_1 - x_k|) (v_k - v_1) f_t^{(p)}.$$
(3.2)

For the last integration in (3.2), by the indistinguishability of the particles, one has

$$\begin{split} &\int_{\mathbb{R}^{2dp}} v_2 \cdot \sum_{k \neq 1}^p \psi(|x_1 - x_k|)(v_k - v_1) f_t^{(p)} \\ &= \int_{\mathbb{R}^{2dp}} v_2 \cdot \sum_{k \neq 1,2}^p \psi(|x_1 - x_k|)(v_1 - v_k) f_t^{(p)} + \int_{\mathbb{R}^{2dp}} v_2 \cdot \psi(|x_1 - x_2|)(v_2 - v_1) f_t^{(p)} \\ &= \int_{\mathbb{R}^{2dp}} v_2 \cdot \sum_{k \neq 1}^p \psi(|x_1 - x_k|)(v_1 - v_k) f_t^{(p)} + \int_{\mathbb{R}^{2dp}} v_2 \cdot \psi(|x_1 - x_2|)(v_2 - v_1) f_t^{(p)} \\ &+ \int_{\mathbb{R}^{2dp}} v_2 \cdot \psi(|x_1 - x_2|)(v_2 - v_1) f_t^{(p)} \\ &= -\int_{\mathbb{R}^{2dp}} v_2 \cdot \sum_{k \neq 1}^p \psi(|x_1 - x_k|)(v_k - v_1) f_t^{(p)} + \int_{\mathbb{R}^{2dp}} |v_2 - v_1|^2 \psi(|x_1 - x_2|) f_t^{(p)}. \end{split}$$

Hence, recalling $\psi \leq 1$, it holds

$$\int_{\mathbb{R}^{2dp}} v_2 \cdot \sum_{k \neq 1}^p \psi(|x_1 - x_k|)(v_k - v_1) f_t^{(p)} = \frac{1}{2} \int_{\mathbb{R}^{2dp}} |v_2 - v_1|^2 \psi(|x_1 - x_2|) f_t^{(p)} \le 2M_2^v f_t.$$

Then it yields from (3.2) that

$$\frac{\partial}{\partial t} \int_{\mathbb{R}^{2dp}} v_1 \cdot v_2 f_t^{(p)} \le \frac{4\kappa}{p-1} M_2^v f_{t_{k-1}}.$$
(3.3)

By combining (3.2) and (3.3), it holds

$$\begin{split} \frac{\partial}{\partial t} M_2^v f_t &\leq -2\kappa \psi_0 M_2^v f_t + 2\kappa \psi_0 \frac{4\kappa\tau}{p-1} M_2^v f_{t_{k-1}} + 2\kappa \psi_0 \int v_1 \cdot v_2 f_{t_{k-1}}^{(p)} \\ &= -2\kappa \psi_0 M_2^v f_t + 2\kappa \psi_0 \frac{4\kappa\tau}{p-1} M_2^v f_{t_{k-1}} \\ &\leq -2\kappa \psi_0 M_2^v f_t + \frac{8\kappa^2 \psi_0}{p-1} \tau M_2^v f_{t_{k-1}}. \end{split}$$

By Grönwall's inequality, one has

$$M_2^v f_k \le M_2^v f_{t_{k-1}} \left(e^{-2\kappa\psi_0\tau} + \frac{8\kappa^2\psi_0}{p-1}\tau^2 \right) \le M_2^v f_0 \left(e^{-2\kappa\psi_0\tau} + \frac{8\kappa^2\psi_0}{p-1}\tau^2 \right)^k.$$

3.2 The main theorems

First we discuss the Wasserstein distance between the distribution of the RBM of the N-particle system (2.10) and the distribution of the $N \to \infty$ system (2.11).

Define the operator $\mathcal{Q}_N^{(k)}$ on the probability measure space $\mathcal{P}(\mathbb{R}^{2d})$ as follows. Let $Z_i^R(0)$ be *i.i.d* drawn from f_0 . Corresponding to Algorithm 1, we define

$$\mathcal{Q}_N^{(k)}(f_0) := \operatorname{Law}(Z_1^R(t_k)),$$

where $\text{Law}(Z_1^R)$ means the law of Z_1^R . Conditioning on a specific sequence of random batches, the particles are not exchangeable. However, when considering the mixture of all possible sequences of random batches, the laws of $Z_i^R(t_k)$ are identical.

In addition, we define \mathcal{Q}_{∞} as one time step evolution in Algorithm 2. Thus one arrives at

$$\mathcal{Q}^k_{\infty}(f_0) = \mathcal{Q}_{\infty} \circ \cdots \circ \mathcal{Q}_{\infty}(f_0), \ (k \text{ copies}),$$

which is expected to be the mean field limit of the RBM after k steps. Note that the \mathcal{Q}_{∞} dynamics can fully determine the probability transition, knowing the marginal distribution of Z_1 , while knowing only the marginal distribution is not enough for the dynamics of $\mathcal{Q}_N^{(k)}$. The joint distribution must be known in the latter case.

Theorem 1 gives an estimate between $\mathcal{Q}_{\infty}^{k}(f_{0})$ and $\mathcal{Q}_{N}^{(k)}(f_{0})$ on the W_{q} distance. In this theorem, we introduce a quantity $\epsilon_{k} \in \mathbb{R}$, which was first introduced in [21]. Plainly speaking, the ϵ_{k} means the probability that the particle *i* is not "clean", where a particle *i* is clean at t_{k}^{-} means its batchmates at $t \leq t_{k}$ were mutually independent and independent to the particle *i* when they interacted. For better organization, we give the definition of "clean" and more details in Section 4.1.

Theorem 1. For f_0 with the compact support $supp[f_0]$, it holds that

$$W_q(\mathcal{Q}^k_{\infty}(f_0), \mathcal{Q}^{(k)}_N(f_0)) \le C(1+t_k)\epsilon_k^{1/q},$$
(3.4)

with $q \in [1, \infty)$ and $C = C(\kappa, \psi_0, R)$. For fixed k, it holds

$$\lim_{N \to \infty} \epsilon_k = 0, \tag{3.5}$$

with $\epsilon_k \leq \mathcal{O}(N^{-1})$.

Then, we consider the limit dynamics given by the operator \mathcal{Q}_{∞} and how $\mathcal{Q}_{\infty}^{n}(f_{0})$ approximates the dynamics of the Fokker-Planck equation (2.6), i.e. the procedure (c) in (1.1). We prove that the W_{q} distance can be controlled by τ with a smooth initial distribution f_{0} with a compact support uniformly in time.

Theorem 2. Let \tilde{f} be a solution to the Fokker-Planck equation (2.6) with initial condition $\tilde{f}(0) = f_0$. It holds the uniform-in-time estimate

$$\sup W_q(\mathcal{Q}^n_\infty(f_0), f(n\tau)) \le C\tau, \tag{3.6}$$

with $q \in [1, \infty)$ and $C = C(\kappa, \psi_0, R)$.

3.3 Some helpful discussions

In addition, we give some discussions on Lemma 12 that will be used for the proof of Theorem 2, and on the commutation of (1.1).

3.3.1 Approximation using PDE analysis

Similar with [[21], Lemma 4.4], one can obtain an $\mathcal{O}(\tau^{\frac{2}{q}})$ bound on the W_q distance between $\mathcal{Q}_{\infty}(\tilde{f}_{t_k})$ and $\tilde{S}(\tilde{f}_{t_k})$ in Proposition 3. Here $\tilde{S}(\tilde{f}_{t_k})$ denotes the law of (5.14) in Section 5.2.

Proposition 3. Let \tilde{f} be a solution to the Fokker-Planck equation (2.6) with initial condition $\tilde{f}(0) = f_0 \in (C^2 \cap W^{2,\infty})(\mathbb{R}^{2d})$. Suppose that the interaction kernel ψ satisfies

$$\sup_{r \ge 0} (|\psi'(r)| + |\psi''(r)|) \le \epsilon_{\psi}$$

with a constant ϵ_{ψ} . It holds

$$W_q(\mathcal{Q}_{\infty}(\tilde{f}_{t_k}), \tilde{S}(\tilde{f}_{t_k})) \le C(\kappa, \epsilon_{\psi}, R, d)\tau^{\frac{2}{q}}, \quad q \in [1, \infty).$$
(3.7)

The proof relies on the PDE analysis on the evolution equations on $\mathcal{Q}_{\infty}(\tilde{f}_{t_k})$ and $\tilde{S}(\tilde{f}_{t_k})$, using the corresponding operator groups. There are two limitations on this method. One is that more regular assumptions on both the interaction kernel and initial data are needed to guarantee the existence of a unique classical solution $\tilde{f}_t \in C^2([0,T] \times \mathbb{R}^{2d})$ for (2.6). In addition, if one uses the estimate obtained by Lemma 3 with the total variance bound, the order of τ is $\frac{2}{a}$. A detailed proof of Proposition 3 is given in Appendix B.

3.3.2 The mean-field limit of the Cucker-Smale system

Up to now, we have given a brief analysis of the path ((b),(c)) in the lower right corner of (1.1). Here, we discuss the upper left corner path in the sense of the Wasserstein distance as well. Denote by

$$\mu_t^N := \sum_{i=1}^N \frac{1}{N} \delta_{z_i(t)}$$

the empirical measure and μ^R_t the empirical measure derived by the RBM with $\mu^R_0=\mu^N_0.$

Proposition 4. For the path from (a) to (d) in (1.1), it holds

$$\mathbb{E}W_2(\tilde{f}_t, \mu_t^R) \le C(\kappa, R, \psi) \sqrt{\frac{\tau}{p-1}} + C(R) \cdot \begin{cases} N^{-\frac{1}{4}} & \text{if } d < 4, \\ N^{-\frac{1}{4}} \sqrt{\log(1+N)} & \text{if } d = 4, \\ N^{-\frac{1}{d}} & \text{if } d > 4. \end{cases}$$
(3.8)

Proof. We first consider the procedure (a). For the first $\tau \to 0$ limit, by the definition of the W_q distance, one has

$$W_2^2(\mu_t^R, \mu_t^N) \le \frac{1}{N} \sum_{i=1}^N (|X_i^R - X_i|^2 + |V_i^R - V_i|^2).$$

Based on [[15], Theorem 3.2], one has

$$\mathbb{E}\left(\frac{1}{N}\sum_{i=1}^{N}|V_{i}^{R}(t)-V_{i}(t)|^{2}\right) \leq C\left(\frac{1}{p-1}-\frac{1}{N-1}\right)\tau + C\tau^{2} + C(1+\tau)\exp\left(-\kappa\psi_{0}t\right),$$

for some constant C depending on κ , R and ψ . Note that

$$\frac{d}{dt}\sum_{i=1}^{N}|X_{i}^{R}-X_{i}|^{2} \leq 2\sqrt{\sum_{i=1}^{N}|X_{i}^{R}-X_{i}|^{2}}\sqrt{\sum_{i=1}^{N}|V_{i}^{R}-V_{i}|^{2}},$$

by the Cauchy-Schwarz inequality and

$$\sqrt{\sum_{i=1}^{N} |X_i^R - X_i|^2} \le \sqrt{\sum_{i=1}^{N} |V_i^R - V_i|^2} T.$$

It thus holds

$$\mathbb{E}W_2(\mu_t^N, \mu_t^R) \le \mathbb{E}\frac{1}{\sqrt{N}} \sqrt{\sum_{i=1}^N |X_i^R - X_i|^2 + \sum_{i=1}^N |V_i^R - V_i|^2} \le \sqrt{C\tau \left(\frac{1}{p-1} - \frac{1}{N-1}\right)} (1+T).$$

Next, for the procedure (d), by the estimate of [[16], Corollary 1], one has $W_q(\mu_t^N, \tilde{f}_t) \leq C(\kappa, R)W_q(\mu_0^N, \tilde{f}_0)$. In addition, according to the convergence rate of the empirical measure in [[14], Theorem 1], with f_0 compactly supported, one has

$$\mathbb{E}W_{q}^{q}(f_{0}, \mu_{0}^{N}) \leq C(R) \cdot \begin{cases} N^{-\frac{1}{2}} & \text{if } q > \frac{d}{2}, \\ N^{-\frac{1}{2}} \log(1+N) & \text{if } q = \frac{d}{2}, \\ N^{-\frac{q}{d}} & \text{if } q < \frac{d}{2}. \end{cases}$$

Therefore, under the same assumption of Theorem 1 and Theorem 2, for the path from (a) to (d), one has

$$\mathbb{E}W_{2}(\tilde{f}_{t},\mu_{t}^{R}) \leq \mathbb{E}\left[W_{2}(\mu_{t}^{N},\mu_{t}^{R}) + W_{2}(\tilde{f}_{t},\mu_{t}^{N})\right] \leq C(\kappa,R,\psi)\sqrt{\frac{\tau}{p-1}} + C(\kappa,R)\mathbb{E}W_{2}(f_{0},\mu_{0}^{N})$$
$$\leq C(\kappa,R,\psi)\sqrt{\frac{\tau}{p-1}} + C(\kappa,R)\left(\mathbb{E}W_{2}^{2}(f_{0},\mu_{0}^{N})\right)^{\frac{1}{2}}.$$

4 Proof of Theorem 1

In this section, we prove Theorem 1 by an observation in combinatorics. Recalling the critical quantity ϵ_k we mentioned above Theorem 1, we first give the Lemmas 4-5, for the proof of Theorem 1, and show the mathematical definition of ϵ_k in Section 4.1. The proof of Lemmas 4-5 are shown in Sections 4.2-4.3 respectively. Finally, Theorem 1 will be proved in Section 4.4 by combining Lemma 4, Lemma 5 and an estimate on the diameters of the compact support in space and velocity, shown in Lemma 7.

Lemma 4. Under the assumption and notation of Theorem 1, it holds

$$|\mathcal{Q}_{\infty}^{k}(f_{0}) - \mathcal{Q}_{N}^{(k)}(f_{0})|_{TV} \leq 2\epsilon_{k},$$

where ϵ_k is a positive constant with respect to N.

Lemma 5 ([21], Theorem 3.2). For any fixed k, with the assumption and notation of Theorem 1, it holds that $\epsilon_k \leq \mathcal{O}(N^{-1})$.

4.1 Prerequisites

To give a detailed explanation of ϵ_k in Theorem 1, we introduce the concept "clean" in Definition 1. First, for each particle *i*, we define a sequence of lists $\{L_i^{(k)}\}$ associated with *i*, given as follows:

1. $L_i^{(0)} = \{i\};$

2. For $k \geq 1$, let $\mathcal{C}_q^{(k-1)}$ be the batch that particle *i* stays in for $t \in [t_{k-1}, t_k)$, then

$$L_i^{(k)} = \bigcup_{j \in \mathcal{C}_q^{(k-1)}} L_j^{(k-1)}$$

Here, $L_i^{(k)}$ can be viewed as the particles that have impacted i for $t < t_k$. Clearly, a particle $i_1 \in L_i^{(k)}$ might not have been a batchmate of i. It could have been a batchmate of i_2 , and then i_2 was a batchmate of i at some time. The important observation is that if $L_i^{(k)}$ and $L_j^{(k)}$ do not intersect for a given sequence of random batches, then particles i and j are independent at t_k^- . Note that we are not claiming all particles in $L_j^{(k)}$ are independent of those in $L_j^{(k)}$ at t_k^- . In fact, it is possible that some $i_1 \in L_i^{(k)}$ and $j_1 \in L_j^{(k)}$ are in the same batch on $[t_{k-1}, t_k)$. However, i_1 and j_1 must be independent at the times when they were added to the batches that eventually impact i, j at t_k^- . Then we are motivated to define the following.

Definition 1. We say the particle *i* is clean on $[t_k, t_{k+1})$ if the batch $C_q^{(k)}$ that contains *i* at t_k^+ satisfies the following:

- (1) any $j \in C_q^{(k)}$ is clean at t_k^- ;
- (2) any $j, \ell \in \mathcal{C}_q^{(k)}$ with $j \neq \ell$, $L_j^{(k)}$ and $L_\ell^{(k)}$ do not intersect.

In other words, a particle *i* is "clean" at t_k^- if its batchmates at $t < t_k$ were mutually independent and independent to *i* when they interacted.

Let A_k denote the set of particles that are clean at t_k^- . Then

$$A_0 = \{1, \cdots, N\}.$$

For $k \geq 1$, one has

$$A_{k} = \left\{ i \in A_{k-1} : i \in \mathcal{C}_{q}^{(k-1)}, \, \forall j, \ell \in \mathcal{C}_{q}^{(k-1)}, \, j \neq \ell, \\ j \in A_{k-1}, \, \ell \in A_{k-1}, \, L_{j}^{(k-1)} \cap L_{\ell}^{(k-1)} = \emptyset \right\}.$$
(4.1)

Now we define

$$\epsilon_k := \mathbb{P}(1 \notin A_k). \tag{4.2}$$

Note that by symmetry, ϵ_k is also the probability that particle *i* is not clean.

4.2 Proof of Lemma 4

We first introduce Lemma 6.

Lemma 6. Consider a fixed sequence of divisions of random batches $\{C^{(\ell)}\}_{\ell \leq k-1}$.

• It holds that

 $|L_i^{(k)}| \le p^k,$

and the particle i is clean at t_k^- if and only if the equality holds.

• The distribution of Z_i (Recall $Z_i := (X_i, V_i)$.) for a clean particle i at t_k^- is $\mathcal{Q}^k_{\infty}(f_0)$.

Here the symbol |A| means the cardinality of a set A.

Proof. The proof is a straightforward induction. Here, let us just briefly mention the proof of the second claim. For k = 0, the statement is trivial. Now, suppose the statement is true for all $k \leq m - 1$. We now consider k = m.

For the given sequence of random batches $\{\mathcal{C}^{(\ell)}\}_{\ell \leq k-1}$, that a particle *i* is clean at $t_m^$ means that on $[t_{m-1}, t_m)$, the particles in the batch for *i* are independent of *i* at t_{m-1} . By the induction assumption, the distribution of one particle at t_{m-1} is given by $\mathcal{Q}_{\infty}^{m-1}(f_0)$. By the independence, the joint distribution of them at t_{m-1} is therefore

$$f^{(p)}(\cdots,t_{m-1}) = \mathcal{Q}_{\infty}^{m-1}(f_0)^{\otimes p}.$$

From t_{m-1} to t_m , the evolution of the joint distribution obeys equation (2.12). Hence, at t_m^- , the distribution of particle *i* is given by $\mathcal{Q}_{\infty}^m(f_0)$.

Now we begin to prove Lemma 4.

Proof of Lemma 4. By Lemma 6, one has

$$\mathcal{Q}_N^{(k)}(f_0) = \mathbb{P}(1 \in A_k) \mathcal{Q}_\infty^k(f_0) + \mathbb{P}(1 \notin A_k) \nu_k, \tag{4.3}$$

for some probability measure ν_k . To see this, consider all possible sequences of random batches. Only the first k divisions of batches (i.e. ones at t_0, \dots, t_{k-1}) will affect the distribution at t_k . This subsequence (the first k divisions) can take only finitely many values, and let $\{c^\ell\}_{\ell \leq k-1}$ be such values. Then, for any $E \subset \mathbb{R}^d$ that is Borel measurable,

$$\mathcal{Q}_{N}^{(k)}(f_{0})[E] = \sum_{\{\mathcal{C}^{\ell} = c^{\ell}, \ell \leq k-1\}} \mathbb{P}(\mathcal{C}^{\ell} = c^{\ell}, \ell \leq k-1) \mathbb{P}(Z_{1} \in E \mid \mathcal{C}^{\ell} = c^{\ell}, \ell \leq k-1)$$

Lemma 6 tells us if $\{c^{\ell}\}_{\ell \leq k-1}$ is a value such that particle 1 is clean, then

$$\mathbb{P}(Z_1 \in E \mid \mathcal{C}^{\ell} = c^{\ell}, \, \ell \leq k-1) = \mathcal{Q}_{\infty}^k(f_0)[E].$$

Hence,

$$\mathcal{Q}_{N}^{(k)}(f_{0})[E] = \mathbb{P}(1 \in A_{k})\mathcal{Q}_{\infty}^{k}(f_{0})[E] + \sum_{\{\mathcal{C}^{\ell} = c^{\ell}, \ell \leq k-1\}, \ 1 \notin A_{k}} \mathbb{P}(\mathcal{C}^{\ell} = c^{\ell}, \ \ell \leq k-1)\mathbb{P}(Z_{1} \in E \ | \ \mathcal{C}^{\ell} = c^{\ell}, \ \ell \leq k-1) = \mathbb{P}(1 \in A_{k})\mathcal{Q}_{\infty}^{k}(f_{0})[E] + \mathbb{P}(1 \notin A_{k})\nu_{k}(E),$$

with

$$\nu_k(E) = \sum_{\{\mathcal{C}^{\ell} = c^{\ell}, \ell \leq k-1\}, 1 \notin A_k} \frac{\mathbb{P}(\mathcal{C}^{\ell} = c^{\ell}, \ell \leq k-1)}{\mathbb{P}(1 \notin A_k)} \mathbb{P}(Z_1 \in E \,|\, \mathcal{C}^{\ell} = c^{\ell}, \, \ell \leq k-1).$$

Clearly, ν_k is a convex combination of some conditional marginal distributions of Z_1 , each being Law (Z_1) conditioning on a particular sequence of batches for $\{1 \notin A_k\}$. Hence ν_k is a probability measure.

By (4.3), it holds that

$$|\mathcal{Q}_{\infty}^{k}(f_{0}) - \mathcal{Q}_{N}^{(k)}(f_{0})| \le (1 - \mathbb{P}(1 \in A_{k}))\mathcal{Q}_{\infty}^{k}(f_{0}) + \mathbb{P}(1 \notin A_{k})\nu_{k} = \epsilon_{k}(\mathcal{Q}_{\infty}^{k}(f_{0}) + \nu_{k}).$$

Therefore, the total variation distance between the two measures is controlled by

$$|\mathcal{Q}_{\infty}^{k}(f_{0}) - \mathcal{Q}_{N}^{(k)}(f_{0})|_{TV} \leq (1 - \mathbb{P}(1 \in A_{k})) + \mathbb{P}(1 \notin A_{k}) = 2\epsilon_{k}.$$

4.3 Proof of Lemma 5

Furthermore, we provide the proof of Lemma 5 which is a simple variant of [[21], Theorem 3.2].

Proof of Lemma 5. First of all, clearly, one has

$$\epsilon_0 = 1 - 1 = 0.$$

Now, one can do induction on k. Assume

$$\lim_{N \to \infty} \epsilon_k = 0,$$

and then consider the batches for $t_k \to t_{k+1}^-$. Assume the batch for particle 1 is $\mathcal{C}_q^{(k)}$. Denote

$$B_k = \left\{ \forall j, \ell \in \mathcal{C}_q^{(k)}, \, j \neq \ell, \, L_j^{(k)} \cap L_\ell^{(k)} = \emptyset \right\}.$$

Let $\mathcal{B} := \mathcal{C}_q^{(k)} \setminus \{1\}$ be the set of other particles that share the same batch with particle 1. Then, by definition of A_{k+1} , it holds

$$\mathbb{P}(1 \in A_{k+1}) = \sum_{j_1, \cdots, j_{p-1}} \mathbb{P}(\mathcal{B} = \{j_1, \cdots, j_{p-1}\}) \times \mathbb{P}(B_k \cap \{1 \in A_k\} \cap_{\ell=1}^{p-1} \{j_\ell \in A_k\} \mid \mathcal{B} = \{j_1, \cdots, j_{p-1}\}).$$
(4.4)

Denote $E := \{ \mathcal{B} = \{j_1, \dots, j_{p-1}\} \}$, where we omit the dependence in $j_{\ell}, 1 \leq \ell \leq p-1$ for notational convenience. Conditioning on $B_k \cap E$ (i.e., provided that the event $B_k \cap E$ happens), whether the particles are clean or not is independent. Hence,

$$\mathbb{P}(B_k \cap \{1 \in A_k\} \cap_{\ell=1}^{p-1} \{j_\ell \in A_k\} | E) = \mathbb{P}(B_k | E))\mathbb{P}(\{1 \in A_k\} \cap_{\ell=1}^{p-1} \{j_\ell \in A_k\} | E, B_k) = \mathbb{P}(B_k | E)\Pi_{\ell=1}^p \mathbb{P}(j_\ell \in A_k | E, B_k),$$

where one has set $j_p = 1$. Moreover,

$$\mathbb{P}(j_{\ell} \in A_k | E, B_k) = \frac{\mathbb{P}(\{j_{\ell} \in A_k\} \cap E \cap B_k)}{\mathbb{P}(E \cap B_k)} = \frac{\mathbb{P}(\{1 \in A_k\} \cap E \cap B_k)}{\mathbb{P}(E \cap B_k)} = \frac{\mathbb{P}(\{1 \in A_k\} \cap B_k)}{\mathbb{P}(B_k)}$$

The second and the last equalities are due to symmetry. For the last equality, $\mathbb{P}(\{\mathcal{B} = \{j_1 \cdots, j_{p-1}\}\} \cap B_k)$ should be equal for all possible j_1, \cdots, j_{p-1} , and the same is true for the numerator. This actually is a kind of independence. Hence, eventually due to the fact

$$\sum_{j_1,\cdots,j_{p-1}} \mathbb{P}(\mathcal{B} = \{j_1,\cdots,j_{p-1}\}) \mathbb{P}(B_k | \mathcal{B} = \{j_1,\cdots,j_{p-1}\}) = \mathbb{P}(B_k)$$

one has

$$1 - \epsilon_{k+1} = \mathbb{P}(1 \in A_{k+1}) \ge \mathbb{P}(B_k)(1 - \epsilon_k)^p.$$

Hence, it suffices to show

$$\lim_{N \to \infty} \mathbb{P}(B_k) = 1.$$

To get an estimate for this, we consider the following equivalent way to construct $L_i^{(k)}$: one starts with $L_i \leftarrow \{i\}$ and repeat the following k times:

- 1. Set $L_{tmp} \leftarrow L_i$ and $A = \emptyset$.
- 2. Loop the following until L_{tmp} is empty.
 - (a) Pick a particle $i_1 \in L_{tmp}$, then choose p-1 particles from $\{1, \dots, N\} \setminus A \cup \{i_1\}$ denoted by $\{i_2, \dots, i_p\}$.
 - (b) Set $L_i \leftarrow L_i \cup \{i_2, \cdots, i_p\}$.
 - (c) Set $A \leftarrow A \cup \{i_1, i_2, \cdots, i_p\}$.
 - (d) Set $L_{tmp} \leftarrow L_{tmp} \setminus \{i_1, i_2, \cdots, i_p\}$.

In the above, we are actually looking back from t_{k-1} . In the *j*-th iteration, we are constructing batches at t_{k-j} . Hence, this is an equivalent way to construct $L_i^{(k)}$.

Now, we estimate $\mathbb{P}(B_k)$ by constructing the lists $L_{j_\ell}^{(k)} : 1 \leq \ell \leq p$ for $j_\ell \in \mathcal{C}_q^{(k)}$ using the above procedure. Consider that the lists for $j_1, \dots, j_{\ell-1}$ have been constructed, which have included at most $(\ell-1)p^k$ particles. Now, for $L_{j_\ell}^{(k)}$ not to intersect with the previous lists, one has to choose particles from $\{1, \dots, N\} \setminus [\bigcup_{z=1}^{\ell-1} L_{j_z}^{(k)} \cup A \cup \{i_1\}]$ in 2.(a) step. Conditioning on the specific choices of $L_{j_\ell}^{(k)} : 1 \leq \ell \leq p$ and A with

$$N_1 := |L_{j_\ell}^{(k)} \cup A \cup \{i_1\}|, N_2 := |A|,$$

this probability is controlled from below by

$$\frac{\binom{N-N_1}{p-1}}{\binom{N-1-N_2}{p-1}} \ge \frac{\binom{N-1-\ell p^k}{p-1}}{\binom{N-1}{p-1}}.$$

Hence, as $N \to \infty$,

$$\mathbb{P}(B_k) \ge \prod_{\ell=1}^p \left[\frac{\binom{N-1-\ell p^k}{p-1}}{\binom{N-1}{p-1}} \right]^k = 1 - \mathcal{O}(N^{-1}).$$

Hence, $\lim_{N \to \infty} \epsilon_{k+1} = 0$ and the claim follows.

4.4 Proof of Theorem 1

Now we can prove Theorem 1. Recall that the TV distance is defined by

$$|\mu - \nu|_{TV} := \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)|,$$

for two probability measures μ , ν defined on certain sample space Ω with events \mathcal{F} .

We denote the diameters of the compact support in the spatial and velocity variables at time t by $\mathcal{D}_X(t)$ and $\mathcal{D}_V(t)$, respectively, i.e.

$$\mathcal{D}_X(t) := \max_{i,j} |X_i - X_j|, \quad \mathcal{D}_V(t) := \max_{i,j} |V_i - V_j|.$$
(4.5)

For the RBM system (2.10), Ha et al. [15] gave an estimate on \mathcal{D}_{X^R} and \mathcal{D}_{V^R} shown in Lemma 7. This estimate works on any fixed sequence of batches and we provide the proof in Appendix A for completeness.

Lemma 7 ([15], Lemma 2.4). Let X^R, V^R be a solution of the RBM model (2.10) with initial data f_0 satisfying

$$\mathcal{D}_X(0) + \mathcal{D}_V(0) < \infty.$$

Then one has

$$\mathcal{D}_{V^R}(t) \le \mathcal{D}_V(0), \quad \mathcal{D}_{X^R}(t) \le \mathcal{D}_X(0) + \mathcal{D}_V(0)t, \quad \forall t \ge 0$$

Proof of Theorem 1. Denote $z := (x, v), Z_i^R := (X_i^R, V_i^R)$, and $|z|^q := |x|^q + |v|^q$. By Lemma 3, taking $\mu = \mathcal{Q}_{\infty}^k(f_0)$ and $\nu = \mathcal{Q}_N^{(k)}(f_0)$, one has

$$W_q(\mathcal{Q}^k_{\infty}(f_0), \mathcal{Q}^{(k)}_N(f_0)) \le 2^{1-\frac{1}{q}} \left(\int_{\mathbb{R}^{2d}} |z|^q d |\mathcal{Q}^k_{\infty}(f_0) - \mathcal{Q}^{(k)}_N(f_0)|(z) \right)^{\frac{1}{q}}.$$

Note that Lemma 7 degenerates into the original Cucker-Smale model when p = N. Hence for $t \in [0, t_1)$, any $\{Z_i^{(0)}(t)\}_{i=1}^p$ satisfying (2.11), one obtains

$$\mathcal{D}_{V^{(0)}}(t) \le \mathcal{D}_{V}(0), \quad \mathcal{D}_{X^{(0)}}(t) \le \mathcal{D}_{X}(0) + \mathcal{D}_{V}(0)t.$$

Also, by the assumption of f_0 , one has $\bar{v} = 0$, and therefore $\bar{x} = 0$. Since (2.11) holds independent of the choice Z_i , defining

$$\begin{aligned} \mathsf{X}(t) &:= \max\{|x| \mid \exists v, \, s.t. \, f(x,v;t) > 0\}, \\ \mathsf{V}(t) &:= \max\{|v| \mid \exists x, \, s.t. \, f(x,v;t) > 0\}, \end{aligned}$$

one has for $t \in [0, t_1]$,

$$\mathsf{V}(t) \le C, \quad \mathsf{X}(t) \le C(1+t), \tag{4.6}$$

where C is a constant depending on R. Then for $t \in [t_1, t_2]$, for the new subsystem $\{Z_i^{(1)}\}_{i=1}^p$ with initial distribution f_{t_1} , one can use Lemma 7 again to get

$$\begin{aligned} \mathcal{D}_{V^{(1)}}(t) &\leq \mathcal{D}_{V^{(1)}}(0) \leq 2\mathsf{V}(t_1) \leq C, \\ \mathcal{D}_{X^{(1)}}(t) &\leq \mathcal{D}_{X^{(1)}}(0) + \mathcal{D}_{V^{(1)}}(t-\tau) \leq 2\mathsf{X}(t_1) + 2\mathsf{V}(t_1)(t-\tau) \leq C(1+t), \end{aligned}$$

and (4.6) holds for $t \in [t_1, t_2]$. By induction, for any t, (4.6) holds. Therefore $|X(t_k)|^q + |V(t_k)|^q \leq C(1+t_k)^q$.

Also, by Lemma 7, it holds $|Z_i^R|^q \leq C(1+t_k)^q$. Then for any $z \in \operatorname{supp}[|\mathcal{Q}_{\infty}^k(f_0) - \mathcal{Q}_N^{(k)}(f_0)|], |z|^q \leq C(1+t_k)^q$. Therefore,

$$\int_{\mathbb{R}^{2d}} |z|^q d |\mathcal{Q}_{\infty}^k(f_0) - \mathcal{Q}_N^{(k)}(f_0)|(z) \le C(1+t_k)^q |\mathcal{Q}_{\infty}^k(f_0) - \mathcal{Q}_N^{(k)}(f_0)|_{TV}.$$

By Lemma 4, one has

$$\int_{\mathbb{R}^{2d}} |z|^q d |\mathcal{Q}^k_{\infty}(f_0) - \mathcal{Q}^{(k)}_N(f_0)|(z) \le C(1+t_k)^q \epsilon_k.$$

Hence $W_q(\mathcal{Q}^k_{\infty}(f_0), \mathcal{Q}^{(k)}_N(f_0)) \leq C(1+t_k)\epsilon_k^{\frac{1}{q}}$. Then by Lemma 5, we finish the proof. \Box

Remark 2. The above analysis is not enough to get the mean-field limit independent of τ and for fixed N, $\epsilon_k \to 1$ as $k \to \infty$. This is a limitation of the technique in the paper.

5 Proof of Theorem 2

Here gives an outline of the proof of Theorem 2. First, we show the stability of \mathcal{Q}_{∞} in Section 5.1. Second, an auxiliary system is introduced in Section 5.2 as a bridge to estimate the W_q distance between $\mathcal{Q}_{\infty}(\tilde{f}_{t_k})$ and $\tilde{f}_{t_{k+1}}$. Last, the proof of Theorem 2 is given in Section 5.3.

5.1 Stability of Q_{∞}

In Lemma 8 and Lemma 9, we give the stability analysis on the operator \mathcal{Q}_{∞} .

Lemma 8. Suppose two probability measures μ_1, μ_2 satisfying $\int v d\mu_1 = \int v d\mu_2$, with compact supports $supp[\mu_1]$, $supp[\mu_2]$ respectively, where $supp[\mu_i] \subset \mathbb{R}^d \times B(0, R_\mu)$, i = 1, 2, and the constant $R_\mu > 0$. Then it holds

$$W_q(\mathcal{Q}_{\infty}(\mu_1), \mathcal{Q}_{\infty}(\mu_2)) \le CW_q(\mu_1, \mu_2), \quad q = [1, \infty),$$
 (5.1)

where $C(\kappa, \psi, R_{\mu})$ is a positive constant.

Proof. Set $(X_i, V_i)(0) \sim \mu_1$, and $(X_i, V_i)(0) \sim \mu_2$, for $i = 1, \dots, p$. The key to the proof is to make use of Lemma 1 to get the contraction property of the velocity difference. The two steps are as follows.

Step 1. First, we show an auxiliary estimate on $\mathcal{D}_V(t)$. By following the formulation of [[16], Lemma 2.2], denote

$$\Psi_{ij} := \frac{\psi(|X_i - X_j|)}{p} + \left(1 - \frac{\sum\limits_{j=1}^p \psi(|X_i - X_j|)}{p}\right) \delta_{ij},$$

where δ_{ij} is the Kronecker delta. Then Ψ_{ij} is nonnegative and satisfies

$$\Psi_{ij} \ge \frac{\psi(|X_i - X_j|)}{p}, \quad \sum_{j=1}^p \Psi_{ij} = 1, \quad \sum_{j=1}^p \Psi_{ij}(V_j - V_i) = \sum_{j=1}^p \frac{\psi(|X_i - X_j|)}{p}(V_j - V_i).$$

We denote $V_i = (V_i^k)$ and obtain the following equality

$$\frac{1}{2} \frac{d}{dt} |V_i - V_j|^2 = \frac{1}{2} \frac{d}{dt} \sum_{k=1}^d |V_i^k - V_j^k|^2$$

$$= \sum_{k=1}^d (V_i^k - V_j^k) (\dot{V}_i^k - \dot{V}_j^k)$$

$$= \sum_{k=1}^d \frac{\kappa p}{p-1} (V_i^k - V_j^k) \left[\sum_{m=1}^p \Psi_{im} (V_m^k - V_i^k) - \sum_{m=1}^p \Psi_{jm} (V_m^k - V_j^k) \right]$$

$$= -\sum_{k=1}^d \frac{\kappa p}{p-1} (V_i^k - V_j^k)^2 + \sum_{k=1}^d \kappa (V_i^k - V_j^k) \left(\sum_{m=1}^p (\Psi_{im} - \Psi_{jm}) V_m^k \right)$$

$$= -\sum_{k=1}^d \frac{\kappa p}{p-1} (V_i^k - V_j^k)^2$$

$$+ \sum_{k=1}^d \frac{\kappa p}{p-1} (V_i^k - V_j^k) \left[\sum_{m=1}^p (\Psi_{im} - \min\{\Psi_{im}, \Psi_{jm}\} + \min\{\Psi_{im}, \Psi_{jm}\} - \Psi_{jm}) V_m^k \right].$$
(5.2)

For a given time t, one can choose indices i_t and j_t satisfying the relation

$$\mathcal{D}_V(t) = |V_{i_t}(t) - V_{j_t}(t)|,$$

and thus for each $m = 1, 2, \cdots, p$,

$$\sum_{k=1}^{d} (V_{i_t}^k - V_{j_t}^k) V_{j_t}^k \le \sum_{k=1}^{d} (V_{i_t}^k - V_{j_t}^k) V_m^k \le \sum_{k=1}^{d} (V_{i_t}^k - V_{j_t}^k) V_{i_t}^k.$$
(5.3)

Combining (5.2) and (5.3), one obtains

$$\frac{1}{2} \frac{d}{dt} |V_{i_t} - V_{j_t}|^2 \leq -\sum_{k=1}^d \frac{\kappa p}{p-1} (V_{i_t}^k - V_{j_t}^k)^2$$

$$+ \sum_{k=1}^d \frac{\kappa p}{p-1} (V_{i_t}^k - V_{j_t}^k) V_{i_t}^k \sum_{m=1}^p (\Psi_{im} - \min\{\Psi_{im}, \Psi_{jm}\})$$

$$- \sum_{k=1}^d \frac{\kappa p}{p-1} (V_{i_t}^k - V_{j_t}^k) V_{j_t}^k \sum_{m=1}^p (\Psi_{jm} - \min\{\Psi_{im}, \Psi_{jm}\})$$

$$= -\sum_{m=1}^p \frac{\kappa p}{p-1} \min\{\Psi_{im}, \Psi_{jm}\} |V_{i_t} - V_{j_t}|^2$$

$$\leq -\frac{\kappa p}{p-1} \psi_0 |V_{i_t} - V_{j_t}|^2.$$
(5.4)

Note that \mathcal{D}_V is Lipschitz continuous with respect to t and thus it is differentiable almost everywhere. Consider the instant t at which \mathcal{D}_V is differentiable. Then,

$$\frac{d}{dt}\mathcal{D}_V^2 \le \frac{d}{dt}|V_{i_t} - V_{j_t}|^2.$$
(5.6)

Combining with (5.4) and (5.6), one obtains

$$\left|\frac{d}{dt}\mathcal{D}_X\right| \leq \mathcal{D}_V, \quad \frac{d}{dt}\mathcal{D}_V \leq -\frac{\kappa p}{p-1}\psi_0\mathcal{D}_V,$$

which is an SDDI. By Lemma 1, there exists a positive constant R_{μ} such that

$$\mathcal{D}_V(t) \le 2e^{-\kappa\psi_0 t} R_\mu. \tag{5.7}$$

Step 2. It can be obtained that

$$\frac{d}{dt}\mathbb{E}|V_1 - \bar{V}_1|^q = \frac{1}{p}\frac{d}{dt}\mathbb{E}\sum_{i=1}^p \left(\sum_{k=1}^d |V_i^k - \bar{V}_i^k|^2\right)^{\frac{d}{2}},\tag{5.8}$$

with

$$\frac{d}{dt} \mathbb{E} \left(\sum_{k=1}^{d} |V_i^k - \bar{V}_i^k|^2 \right)^{\frac{q}{2}} = \mathbb{E} \left[\frac{q}{2} \left(\sum_{k=1}^{d} |V_i^k - \bar{V}_i^k|^2 \right)^{\frac{q}{2}-1} \frac{d}{dt} \sum_{k=1}^{d} |V_i^k - \bar{V}_i^k|^2 \right] \\
= \frac{q\kappa}{p-1} \mathbb{E} \left[|V_i - \bar{V}_i|^{q-2} \sum_{j=1}^{p} \psi(|X_j - X_i|) \sum_{k=1}^{d} (V_j^k - V_i^k - \bar{V}_j^k + \bar{V}_i^k) (V_i^k - \bar{V}_i^k) \right. \\
\left. + \sum_{j=1}^{p} |V_i - \bar{V}_i|^{q-2} \left(\psi(|X_j - X_i|) - \psi(|\bar{X}_j - \bar{X}_i|) \right) \sum_{k=1}^{d} (\bar{V}_j^k - \bar{V}_i^k) (V_i^k - \bar{V}_i^k) \right].$$

Rewrite the right-hand side of (5.8) as

$$\frac{d}{dt}\mathbb{E}|V_1 - \bar{V}_1|^q := I_{21} + I_{22},$$

where

$$I_{21} := \frac{q\kappa}{p-1} \mathbb{E} \sum_{i,j=1}^{p} \sum_{k=1}^{d} \psi(|X_j - X_i|) |V_i - \bar{V}_i|^{q-2} (V_j^k - V_i^k - \bar{V}_j^k + \bar{V}_i^k) (V_i^k - \bar{V}_i^k),$$

and

$$I_{22} := \frac{q\kappa}{p-1} \mathbb{E} \sum_{i,j=1}^{p} \sum_{k=1}^{d} \left(\psi(|X_j - X_i|) - \psi(|\bar{X}_j - \bar{X}_i|) \right) \\ \cdot |V_i - \bar{V}_i|^{q-2} (V_i^k - \bar{V}_i^k) (\bar{V}_j^k - \bar{V}_i^k).$$

First estimate ${\cal I}_{21}$ using the index changing trick:

$$\begin{split} I_{21} &= \frac{q\kappa}{p-1} \mathbb{E} \sum_{i,j=1}^{p} \sum_{k=1}^{d} \psi(|X_j - X_i|) |V_i - \bar{V}_i|^{q-2} [(V_j^k - \bar{V}_j^k) - (V_i^k - \bar{V}_i^k)] (V_i^k - \bar{V}_i^k) \\ &= \frac{q\kappa}{p-1} \mathbb{E} \left[\sum_{i,j=1}^{p} \sum_{k=1}^{d} \psi(|X_j - X_i|) |V_i - \bar{V}_i|^{q-2} (V_j^k - \bar{V}_j^k) (V_i^k - \bar{V}_i^k) \\ &- \sum_{i,j=1}^{p} \sum_{k=1}^{d} \psi(|X_j - X_i|) |V_i - \bar{V}_i|^{q-2} (V_i^k - \bar{V}_i^k)^2 \right] \\ &= \frac{q\kappa}{p-1} \mathbb{E} \left[\sum_{i,j=1}^{p} \sum_{k=1}^{d} \psi(|X_j - X_i|) |V_j - \bar{V}_j|^{q-2} (V_j^k - \bar{V}_j^k) (V_i^k - \bar{V}_i^k) \\ &- \sum_{i,j=1}^{p} \sum_{k=1}^{d} \psi(|X_j - X_i|) |V_i - \bar{V}_i|^{q-2} (V_i^k - \bar{V}_i^k)^2 \right] \\ &= -\frac{q\kappa}{p-1} \mathbb{E} \left[\sum_{i,j=1}^{p} \sum_{k=1}^{d} \psi(|X_j - X_i|) \\ &\cdot \left[|V_i - \bar{V}_i|^{q-2} (V_i^k - \bar{V}_i^k) - |V_j - \bar{V}_j|^{q-2} (V_j^k - \bar{V}_j^k) \right] (V_i^k - \bar{V}_i^k) \right] \\ &= \frac{q\kappa}{2(p-1)} \mathbb{E} \left[\sum_{i,j=1}^{p} \sum_{k=1}^{d} \psi(|X_j - X_i|) \\ &\cdot \left[|V_i - \bar{V}_i|^{q-2} (V_i^k - \bar{V}_i^k) - |V_j - \bar{V}_j|^{q-2} (V_j^k - \bar{V}_j^k) \right] \left[(V_j^k - \bar{V}_j^k) - (V_i^k - \bar{V}_i^k) \right] \right]. \end{split}$$

One can use the relation

$$\sum_{k=1}^{d} \left[|V_i - \bar{V}_i|^{q-2} (V_i^k - \bar{V}_i^k) - |V_j - \bar{V}_j|^{q-2} (V_j^k - \bar{V}_j^k) \right] \cdot \left[(V_j^k - \bar{V}_j^k) - (V_i^k - \bar{V}_i^k) \right] \le 0,$$

and the condition (2.2) of ψ to obtain

$$\begin{split} I_{21} &\leq \frac{q\kappa}{2(p-1)} \psi_0 \mathbb{E} \Biggl\{ \sum_{i,j=1}^p \sum_{k=1}^d \left[|V_i - \bar{V}_i|^{q-2} (V_i^k - \bar{V}_i^k) - |V_j - \bar{V}_j|^{q-2} (V_j^k - \bar{V}_j^k) \right] \\ &\quad \cdot \left[(V_j^k - \bar{V}_j^k) - (V_i^k - \bar{V}_i^k) \right] \Biggr\} \\ &= \frac{q\kappa}{2(p-1)} \psi_0 \mathbb{E} \left[\sum_{i=1}^p \sum_{k=1}^d |V_i - \bar{V}_i|^{q-2} (V_i^k - \bar{V}_i^k) \sum_{j=1}^p (V_j^k - \bar{V}_j^k) \\ &\quad - \sum_{i=1}^p \sum_{k=1}^d p |V_i - \bar{V}_i|^{q-2} (V_i^k - \bar{V}_i^k)^2 - \sum_{j=1}^p \sum_{k=1}^d p |V_j - \bar{V}_j|^{q-2} (V_j^k - \bar{V}_j^k)^2 \\ &\quad + \sum_{j=1}^p \sum_{k=1}^d |V_j - \bar{V}_j|^{q-2} (V_j^k - \bar{V}_j^k) \sum_{i=1}^p (V_i^k - \bar{V}_i^k) \Biggr] \,. \end{split}$$

By the momentum condition $\mathbb{E}\sum_{i=1}^{p}(V_{i}^{k}-\bar{V}_{i}^{k})=0$, it holds

$$I_{21} \le -\frac{p^2 q \kappa}{p-1} \psi_0 \mathbb{E} |V_1 - \bar{V}_1|^q.$$
(5.9)

For I_{22} , one has

$$I_{22} \leq \frac{q\kappa}{p-1} \|\psi\|_{\text{Lip}} \mathbb{E}\left[\sum_{i,j=1}^{p} \sum_{k=1}^{d} \left| |X_j - X_i| - |\bar{X}_j - \bar{X}_i| |V_i - \bar{V}_i|^{q-2} |V_i^k - \bar{V}_i^k| |\bar{V}_j^k - \bar{V}_i^k| \right] \\ \leq \frac{q\kappa}{p-1} \|\psi\|_{\text{Lip}} \mathbb{E}\left[\sum_{i,j=1}^{p} \left| |X_j - X_i| - |\bar{X}_j - \bar{X}_i| |V_i - \bar{V}_i|^{q-1} |\bar{V}_j - \bar{V}_i| \right],$$

by the Cauchy inequality. Then by the inequality $||a - b| - |c - d|| \le |a - c| + |b - d|$, one has

$$I_{22} \leq \frac{q\kappa}{p-1} \|\psi\|_{\mathrm{Lip}} \mathbb{E} \left[\sum_{i,j=1}^{p} \left| |X_i - \bar{X}_i| + |X_j - \bar{X}_j| \right| |V_i - \bar{V}_i|^{q-1} |\bar{V}_j - \bar{V}_i| \right] \\ \leq \frac{q\kappa}{p-1} \|\psi\|_{\mathrm{Lip}} \mathcal{D}_{\bar{V}}(t) \mathbb{E} \left[p \sum_{i=1}^{p} |X_i - \bar{X}_i| |V_i - \bar{V}_i|^{q-1} + \sum_{i,j=1}^{p} |X_j - \bar{X}_j| |V_i - \bar{V}_i|^{q-1} \right] \\ \leq \frac{2qp^2\kappa}{p-1} \|\psi\|_{\mathrm{Lip}} \mathcal{D}_{\bar{V}}(t) \mathbb{E} \left[|X_1 - \bar{X}_1| |V_1 - \bar{V}_1|^{q-1} \right],$$
(5.10)

by Hölder's inequality. Now by combining (5.8) and (5.9) and (5.10), one has

$$\frac{d}{dt}\mathbb{E}|V_{1}-\bar{V}_{1}|^{q} \leq -\frac{p}{p-1}q\kappa\psi_{0}\mathbb{E}|V_{1}-\bar{V}_{1}|^{q} \\
+\frac{2p}{p-1}q\kappa\|\psi\|_{\mathrm{Lip}}\mathcal{D}_{\bar{V}}(t)\mathbb{E}[|X_{1}-\bar{X}_{1}||V_{1}-\bar{V}_{1}|^{q-1}] \\
\leq -\frac{p}{p-1}q\kappa\psi_{0}\mathbb{E}|V_{1}-\bar{V}_{1}|^{q} \\
+\frac{2p}{p-1}q\kappa\|\psi\|_{\mathrm{Lip}}\mathcal{D}_{\bar{V}}(t)(\mathbb{E}[|X_{1}-\bar{X}_{1}|^{q}])^{\frac{1}{q}}(\mathbb{E}|V_{1}-\bar{V}_{1}|^{q})^{1-\frac{1}{q}},$$

similarly by Hölder's inequality. Since

$$\frac{d}{dt} (\mathbb{E}|V_1 - \bar{V}_1|^q)^{\frac{1}{q}} = \frac{1}{q} (\mathbb{E}|V_1 - \bar{V}_1|^q)^{\frac{1}{q} - 1} \frac{d}{dt} \mathbb{E}|V_1 - \bar{V}_1|^q
\leq -\frac{p\kappa}{p - 1} \psi_0 (\mathbb{E}[|V_1 - \bar{V}_1|^q_q])^{\frac{1}{q}} + \frac{2p\kappa}{p - 1} \|\psi\|_{\operatorname{Lip}} \mathcal{D}_{\bar{V}}(t) (\mathbb{E}[|X_1 - \bar{X}_1|^q])^{\frac{1}{q}}.$$
(5.11)

and

$$\frac{d}{dt}\mathbb{E}|X_1 - \bar{X}_1|^q = \mathbb{E}\frac{q}{2} \left(\sum_{k=1}^d |X_1^k - \bar{X}_1^k|^2 \right)^{\frac{q}{2}-1} \sum_{k=1}^d (X_1^k - \bar{X}_1^k) \cdot 2(V_1^k - \bar{V}_1^k) \\
\leq \mathbb{E}q|X_1 - X_1|^{q-1}|V_1 - \bar{V}_1| \\
\leq q(\mathbb{E}|X_1 - \bar{X}_1|^q)^{1-\frac{1}{q}} (\mathbb{E}|V_1 - \bar{V}_1|^q)^{\frac{1}{q}},$$

thus it holds

$$\frac{d}{dt} (\mathbb{E}|X_1 - \bar{X}_1|^q)^{\frac{1}{q}} = \frac{1}{q} (\mathbb{E}|X_1 - \bar{X}_1|^q)^{\frac{1}{q} - 1} \frac{d}{dt} \mathbb{E}|X_1 - \bar{X}_1|^q \le (\mathbb{E}|V_1 - \bar{V}_1|^q)^{\frac{1}{q}}.$$
 (5.12)

Note that $\mathcal{D}_{\bar{V}} \leq 2e^{-\kappa\psi_0 t}R_{\mu}$ by Step 1. Now one knows that both $(\mathbb{E}|X_1 - \bar{X}_1|^q)^{\frac{1}{q}}$ and $(\mathbb{E}|V_1 - \bar{V}_1|^q)^{\frac{1}{q}}$ satisfy Lemma 1 and then there exists a constant C such that

$$\left(\mathbb{E}|X_1 - \bar{X}_1|^q\right)^{\frac{1}{q}}(t) + \left(\mathbb{E}|V_1 - \bar{V}_1||^q\right)^{\frac{1}{q}}(t) \le C\left[\left(\mathbb{E}|X_1 - \bar{X}_1|^q\right)^{\frac{1}{q}}(0) + \left(\mathbb{E}|V_1 - \bar{V}_1|^q\right)^{\frac{1}{q}}(0)\right].$$

Then,

$$W_{q}(\mathcal{Q}_{\infty}(\mu_{1}),\mathcal{Q}_{\infty}(\mu_{2})) \leq \left(\mathbb{E}|X_{1}-\bar{X}_{1}|^{q}(t)+\mathbb{E}|V_{1}-\bar{V}_{1}|^{q}(t)\right)^{\frac{1}{q}} \\ \leq \left(\mathbb{E}|X_{1}-\bar{X}_{1}|^{q}(t)\right)^{\frac{1}{q}}+\left(\mathbb{E}|V_{1}-\bar{V}_{1}|^{q}(t)\right)^{\frac{1}{q}} \\ \leq C\left[\left(\mathbb{E}|X_{1}-\bar{X}_{1}|^{q}\right)^{\frac{1}{q}}(0)+\left(\mathbb{E}|V_{1}-\bar{V}_{1}|^{q}\right)^{\frac{1}{q}}(0)\right] \\ \leq 2C\left[\mathbb{E}|X_{1}-\bar{X}_{1}|^{q}(0)+\mathbb{E}|V_{1}-\bar{V}_{1}|^{q}(0)\right]^{\frac{1}{q}},$$

since $a^{\frac{1}{q}} + b^{\frac{1}{q}} \le 2^{1-\frac{1}{q}}(a+b)^{\frac{1}{q}}$ for $a \ge 0$, $b \ge 0$ with $q \ge 1$. For any $\epsilon > 0$, the initial data can be chosen by a coupling such that

$$\left[\mathbb{E}|X_1 - \bar{X}_1|^q(0) + \mathbb{E}|V_1 - \bar{V}_1|^q(0)\right]^{\frac{1}{q}} \le \epsilon + W_1(\mu_1, \mu_2).$$

Hence by the arbitrariness of ϵ , it holds $W_q(\mathcal{Q}_{\infty}(\mu_1), \mathcal{Q}_{\infty}(\mu_2)) \leq CW_q(\mu_1, \mu_2)$,

Lemma 9. For \tilde{f} defined in (2.6) with $R < \infty$, one has

$$W_q(\mathcal{Q}^{n-m+1}_{\infty}(\tilde{f}_{m-1}), \mathcal{Q}^{n-m}_{\infty}(\tilde{f}_m)) \le CW_q(\mathcal{Q}_{\infty}(\tilde{f}_{m-1}), \tilde{f}_m), \quad q = [1, \infty),$$

where C is relevant to ψ, κ, R and d.

Proof. By the flocking property shown in Step 1 in Lemma 8, one has $\operatorname{supp}^{v}[\mathcal{Q}_{\infty}(\tilde{f}_{m-1})] \subset$ supp^v[\tilde{f}_{m-1}]. Also, supp^v[\tilde{f}_{m-1}] \subset supp^v[\tilde{f}_0] by Lemma 2. Furthermore, for $n - m - 1 \ge k \ge 0$, $\mathcal{Q}_{\infty}^{k+1}(\tilde{f}_{m-1})$ and $\mathcal{Q}_{\infty}^k(\tilde{f}_m)$ have compact supports, where $\mathcal{Q}_{\infty}^0(\tilde{f}_m) := \tilde{f}_m$. Similarly with the proof of Proposition 1, note that $(X_1(k\tau), V_1(k\tau) \text{ and } (\bar{X}_1(k\tau), \bar{V}_1(k\tau)))$ obey $\mathcal{Q}_{\infty}^{k+1}(\tilde{f}_{m-1})$ and $\mathcal{Q}_{\infty}^k(\tilde{f}_m)$ respectively, which satisfy the momentum equality assumption \tilde{f}_m .

tion of Lemma 8.

Denote

$$|\Delta X_1| := (\mathbb{E}|X_1 - \bar{X}_1|^q)^{\frac{1}{q}}, \quad |\Delta V_1| := (\mathbb{E}|V_1 - \bar{V}_1|^q)^{\frac{1}{q}}, \quad |\Delta Z_1| := |\Delta X_1| + |\Delta V_1|,$$

for $q \ge 1$. Hence, by the analysis in Lemma 8, (5.12) and (5.11) still hold for $t \in [0, (n-m)\tau]$. By Lemma 9, one obtains

$$|\Delta Z_1((n-m)\tau)| \le C |\Delta Z_1(0)|.$$

By choosing a suitable coupling of the initial data, the proof is finished.

 \square

5.2An auxiliary system

We introduce an auxiliary system (5.14) with law $\tilde{S}(\tilde{f}_{t_k})$ to get the estimate on

$$W_q(\mathcal{Q}_{\infty}(\tilde{f}_{t_k}), \tilde{f}_{t_{k+1}}) \le W_q(\tilde{S}(\tilde{f}_{t_k}), \tilde{f}_{t_{k+1}}) + W_q(\tilde{S}(\tilde{f}_{t_k}), \mathcal{Q}_{\infty}(\tilde{f}_{t_k})).$$

Rewrite (2.6) into the following particle system

$$\begin{cases} \frac{d}{dt}\tilde{X}_{i}(t) = \tilde{V}_{i}(t), \ i = 1, \cdots, N, \\ \frac{d}{dt}\tilde{V}_{i}(t) = \int \kappa \psi(|x - \tilde{X}_{i}|)(v - \tilde{V}_{i})\tilde{f}_{t}(x, v)dxdv. \end{cases}$$
(5.13)

In order to show the law of (\tilde{X}, \tilde{V}) at time $n\tau$ is close to \mathcal{Q}_{∞}^{n} , we consider the transient system 1

$$\begin{cases} \frac{d}{dt}\hat{X}_{i}(t) = \hat{V}_{i}(t), \ i = 1, \cdots, N, \\ \frac{d}{dt}\hat{V}_{i}(t) = \int \kappa \psi(|x - \hat{X}_{i}|)(v - \hat{V}_{i})\tilde{f}_{t_{k}}(x, v)dxdv, \end{cases}$$

$$(5.14)$$

with $(\hat{X}(0), \hat{V}(0)) \sim \tilde{f}_{t_k}$. Denote $\tilde{S}(\tilde{f}_{t_k}) := \operatorname{Law}(\hat{X}(\tau), \hat{V}(\tau))$.

Lemma 10. Under the same assumption of Lemma 9, it holds

$$supp[\tilde{S}(\tilde{f}_{t_k})] \subset B(0,C) \times B(0,Ce^{-\kappa\psi_0 t_{k+1}}).$$
(5.15)

Proof. For $\tilde{S}(\tilde{f}_{t_k})$, define

$$\begin{split} \mathsf{X}_{k}(t) &:= \max\{ |x(t, x^{k}, v^{k})| \mid (x^{k}, v^{k}) \in \mathrm{supp}[\tilde{f}_{t_{k}}] \}, \\ \mathsf{V}_{k}(t) &:= \max\{ |v(t, x^{k}, v^{k})| \mid (x^{k}, v^{k}) \in \mathrm{supp}[\tilde{f}_{t_{k}}] \}, \end{split}$$

where the pair $(x(t, x^k, v^k), v(t, x^k, v^k))$ denotes (x(t), v(t)) being the solution of (5.14) with $(x(t_k), v(t_k)) = (x^k, v^k)$. Note that the maximum is reached because it is assumed that $\operatorname{supp}[\tilde{f}_{t_k}]$ is compact. Denote $K_t^x \subset \operatorname{supp}^x[\tilde{f}_{t_k}]$ (resp. $K_t^v \subset \operatorname{supp}^v[\tilde{f}_{t_k}]$) the set of points (x^k, v^k) such that the maximum is reached in $X_k(t)$ (resp. in $V_k(t)$). By definition, one has $|X_k(t)|^2 = |x(t, x^k, v^k)|^2$ for every $(x^k, v^k) \in K_t^x$. It follows from Danskin's theorem [31, 11] and from the fact that $\partial_t x(t, x^k, v^k) = v(t, x^k, v^k)$ that

$$\mathsf{X}_k(t) \cdot \dot{\mathsf{X}}_k(t) = \max\{x(t, x^k, v^k) \cdot v(t, x^k, v^k) \mid (x^k, v^k) \in K_t^x\}.$$

Using the Cauchy-Schwarz inequality, one obtains that

$$\dot{\mathsf{X}}_k(t) \leq \mathsf{V}_k(t).$$

Similarly, by Danskin's theorem again, one has

$$\mathsf{V}_k(t) \cdot \dot{\mathsf{V}}_k(t) = \max\{v(t, x^k, v^k) \cdot \int \kappa \psi(|x - x(t, x^k, v^k)|)(v - v(t, x^k, v^k))\tilde{f}_{t_k} dx dv$$
$$|(x^k, v^k) \in K_t^v\}.$$

Note that by the definition of K_t^v , $v(t, x^k, v^k) \cdot (v - v(t, x^k, v^k)) \leq 0$. It holds that

$$\mathsf{V}_k(t) \cdot \dot{\mathsf{V}}_k(t) \le -\kappa \psi_0 \int v(t, x^k, v^k) \cdot v(t, x^k, v^k) \tilde{f}_{t_k} dx dv, \quad (x^k, v^k) \in K_t^v,$$

since $\int v \tilde{f}_{t_k} dx dv = 0$ and $\int \tilde{f}_{t_k} dx dv = 1$. Then one has

$$\dot{\mathsf{V}}_k(t) \leq -\kappa \psi_0 \mathsf{V}_k(t)$$

and thus (5.15) holds.

Lemma 11. Under the same assumption of Lemma 9, there exists a positive constant α such that

$$W_q(\tilde{S}(\tilde{f}_{t_k}), \tilde{f}_{t_{k+1}}) \le Ce^{-\kappa\psi_0 t_k} \tau^2, \quad q \in [1, \infty),$$
 (5.16)

with $C = C(\kappa, \psi_0, \|\psi\|_{Lip}, R).$

Proof. Let $\mathcal{R}(x, v, t) = \int \psi(|y - x|)(w - v)(\tilde{f}_t(y, w) - \tilde{f}_{t_k}(y, w))dydw$. For $t \in [t_k, t_{k+1})$, by straightforward calculus, it holds

$$\begin{split} \left| \frac{\partial \mathcal{R}}{\partial t}(x,v,t) \right| &= \left| \int \psi(|y-x|)(w-v) \frac{\partial \tilde{f}_t}{\partial t}(y,w) dy dw \right| \\ &= \left| \int \psi(|y-x|)(w-v) \left[-w \cdot \nabla_y \tilde{f}_t(y,w) + \nabla_w \cdot \left(\tilde{f}_t(y,w) \kappa \int \psi(|\bar{y}-y|)(\bar{w}-w) \tilde{f}_t(\bar{y},\bar{w}) d\bar{y} d\bar{w} \right) \right] dy dw \right| \\ &+ \nabla_w \cdot \left(\tilde{f}_t(y,w) \kappa \int \psi(|\bar{y}-y|)(\bar{w}-w) \tilde{f}_t(\bar{y},\bar{w}) d\bar{y} d\bar{w} \right) \right] dy dw \\ &= \left| \int \left[w \cdot \nabla_y \psi(|y-x|)(w-v) \tilde{f}_t(y,w) - d\kappa \psi(|y-x|) \tilde{f}_t(y,w) \int \psi(|\bar{y}-y|)(\bar{w}-w) \tilde{f}_t(\bar{y},\bar{w}) d\bar{y} d\bar{w} \right] dy dw \right| \\ &\leq ||\psi||_{\text{Lip}} \int |w||v-w| \tilde{f}_t(y,w) dy dw + d\kappa \int |\bar{w}-w| \tilde{f}_t(y,w) \tilde{f}_t(\bar{y},\bar{w}) d\bar{y} d\bar{w} dy dw \\ &\leq ||\psi||_{\text{Lip}} \int (|w||v|+|w|^2) \tilde{f}_t(y,w) dy dw + d\kappa \int 2|w| \tilde{f}_t(y,w) dy dw. \end{split}$$

By Lemma 2,

$$\left|\frac{\partial \mathcal{R}}{\partial t}(\tilde{X}_1, \tilde{V}_1, t)\right| \le C e^{-\kappa \psi_0 t}.$$

We first consider q to be an even integer. For $(\tilde{X}_1, \tilde{V}_1)$ obeying (5.13) and (\hat{X}_1, \hat{V}_1) obeying (5.14), with $(\tilde{X}_1, \tilde{V}_1)(t_k) = (\hat{X}_1, \hat{V}_1)(t_k) \sim \tilde{f}_{t_k}$, it holds

$$\begin{split} &\frac{d}{dt} |\tilde{V}_{1} - \hat{V}_{1}|^{2} = 2 \sum_{k=1}^{d} (\tilde{V}_{1}^{k} - \hat{V}_{1}^{k}) \frac{d}{dt} (\tilde{V}_{1}^{k} - \hat{V}_{1}^{k}) \\ &= 2\kappa \sum_{k=1}^{d} (\tilde{V}_{1}^{k} - \hat{V}_{1}^{k}) \Big[\int \psi(|x - \tilde{X}_{1}|)(v - \tilde{V}_{1}) \tilde{f}_{t_{k}}(x, v) dx dv \\ &- \int \psi(|x - \hat{X}_{1}|)(v - \hat{V}_{1}) \tilde{f}_{t_{k}}(x, v) dx dv + \mathcal{R}(\tilde{X}_{1}, \tilde{V}_{1}) \Big]^{(k)} \\ &= 2\kappa \sum_{k=1}^{d} (\tilde{V}_{1}^{k} - \hat{V}_{1}^{k}) \left(\int \psi(|x - \tilde{X}_{1}|)[(v - \tilde{V}_{1}) - (v - \hat{V}_{1})] \tilde{f}_{t_{k}}(x, v) dx dv \right)^{(k)} \\ &+ 2\kappa \sum_{k=1}^{d} (\tilde{V}_{1}^{k} - \hat{V}_{1}^{k}) \left(\int [\psi(|x - \hat{X}_{1}|) - \psi(|x - \tilde{X}_{1}|)](v - \hat{V}_{1}) \tilde{f}_{t_{k}}(x, v) dx dv \right)^{(k)} \\ &+ 2\kappa \sum_{k=1}^{d} (\tilde{V}_{1}^{k} - \hat{V}_{1}^{k}) \mathcal{R}(\tilde{X}_{1}, \tilde{V}_{1}) | \\ &\leq -q\psi_{0} |\tilde{V}_{1} - \hat{V}_{1}|^{2} + 2\kappa ||\psi||_{\mathrm{Lip}} |\tilde{X}_{1} - \hat{X}_{1}|| \sum_{k=1}^{d} (\tilde{V}_{1}^{k} - \hat{V}_{1}^{k}) \left(\int (v - \hat{V}_{1}) \tilde{f}_{t_{k}} \right)^{(k)} \\ &\leq -2\kappa\psi_{0} |\tilde{V}_{1} - \hat{V}_{1}|^{2} + 2\kappa ||\psi||_{\mathrm{Lip}} |\tilde{X}_{1} - \hat{X}_{1}|| \tilde{V}_{1} - \hat{V}_{1}|| \hat{V}_{1} \\ &\leq -2\kappa\psi_{0} |\tilde{V}_{1} - \hat{V}_{1}|^{2} + 2\kappa ||\psi||_{\mathrm{Lip}} |\tilde{X}_{1} - \hat{X}_{1}|| \tilde{V}_{1} - \hat{V}_{1}|| \hat{V}_{1}| \\ &+ 2\kappa |\tilde{V}_{1} - \hat{V}_{1}||\mathcal{R}(\tilde{X}_{1}, \tilde{V}_{1})|, \end{split}$$

where the superscript (k) means the k-th dimension of vectors. Since $\int v \tilde{f}_t dx dv = \bar{v} = 0$ by (2.7), and $|\hat{V}_1| \leq C e^{-\kappa \psi_0 t_k}$ by Lemma 2 and Lemma 10, it holds

$$\frac{d}{dt} |\tilde{V}_1 - \hat{V}_1|^2 \leq -2\kappa\psi_0 |\tilde{V}_1 - \hat{V}_1|^2 + C\kappa ||\psi||_{\text{Lip}} e^{-\kappa\psi_0 t_k} |\tilde{X}_1 - \hat{X}_1||\tilde{V}_1 - \hat{V}_1| + 2\kappa |\tilde{V}_1 - \hat{V}_1||\mathcal{R}(\tilde{X}_1, \tilde{V}_1)|, \quad (5.17)$$

and

$$\frac{d}{dt}|\tilde{V}_1 - \hat{V}_1| \le -\psi_0 \kappa |\tilde{V}_1 - \hat{V}_1| + C\kappa \|\psi\|_{\operatorname{Lip}} e^{-\kappa\psi_0 t_k} |\tilde{X}_1 - \hat{X}_1| + \kappa |\mathcal{R}(\tilde{X}_1, \tilde{V}_1)|.$$

Also, with

$$\frac{d}{dt}|\tilde{X}_1 - \hat{X}_1| \le |\tilde{V}_1 - \hat{V}_1|,$$

one has

$$\frac{d}{dt}\left(|\tilde{X}_1 - \hat{X}_1| + |\tilde{V}_1 - \hat{V}_1|\right) \le C\left(|\tilde{X}_1 - \hat{X}_1| + |\tilde{V}_1 - \hat{V}_1|\right) + C|\mathcal{R}|.$$

By Grönwall's inequality, it holds

1

$$|\hat{X}_1 - \tilde{X}_1| + |\hat{V}_1 - \tilde{V}_1| \le C e^{-\kappa \psi_0 t_k} \tau^2$$

Then

$$\left[\mathbb{E}\left[|\hat{X}_{1}-\tilde{X}_{1}|^{q}+|\hat{V}_{1}-\tilde{V}_{1}|^{q}\right]\right]^{\frac{1}{q}} \leq \left[\mathbb{E}\left[|\hat{X}_{1}-\tilde{X}_{1}|+|\hat{V}_{1}-\tilde{V}_{1}|\right]^{q}\right]^{\frac{1}{q}} \leq Ce^{-\kappa\psi_{0}t_{k}}\tau^{2},$$
implies

which implies

$$W_q(S(f_{t_k}), f_{t_{k+1}}) \le C(\kappa, \psi_0, \|\psi\|_{\operatorname{Lip}}, R) e^{-\kappa \psi_0 t_k} \tau^2.$$

~

The estimate on $W_q(\tilde{S}(\tilde{f}_{t_k}), \mathcal{Q}_{\infty}(\tilde{f}_{t_k}))$ is given by Lemma 12.

Lemma 12. Under the same assumption of Theorem 2, it holds

$$W_q(\mathcal{Q}_{\infty}(\tilde{f}_{t_k}), \tilde{S}(\tilde{f}_{t_k})) \le C(\kappa, R)\tau^2, \quad q \in [1, \infty).$$
(5.18)

Proof. Set (\hat{X}_1, \hat{V}_1) and $\{(X_i, V_i)\}$ to be the solutions of (5.14) and (2.11) respectively. The initial data $(\hat{X}_1(0), \hat{V}_1(0))$ and $\{(X_i(0), V_i(0))\}$ follow \tilde{f}_{t_k} . Take the coupling $(\hat{X}_1(0), \hat{V}_1(0)) = (X_1(0), V_1(0))$. We introduce $\{(X_i^E, V_i^E)\}$ as the approximation of $\{(X_i, V_i)\}$ by the forward Euler method with time step τ . Then, taking the expectation of the selection of $\{(X_j(0), V_j(0)\}_{j \neq 1}$, one has

$$\begin{split} \frac{d}{dt} \frac{1}{2} \mathbb{E} |V_1^E - \hat{V}_1|^2 &= \int \kappa \psi(|x - X_1(0)|) (v - V_1(0)) \cdot \mathbb{E}(V_1^E(t) - \hat{V}_1(t)) \tilde{f}_{t_k} \\ &- \int \kappa \psi(|x - \hat{X}_1(t)|) (v - \hat{V}_1(t)) \cdot \mathbb{E}(V_1^E(t) - \hat{V}_1(t)) \tilde{f}_{t_k} \\ &\leq \kappa \int \psi(|x - X_1(0)|) (\hat{V}_1(t) - V_1(0)) \cdot \mathbb{E}(V_1^E(t) - \hat{V}_1(t)) \tilde{f}_{t_k} \\ &+ \kappa \int \|\psi\|_{\mathrm{Lip}} |X_1(0) - \hat{X}_1(t)| (v - \hat{V}_1(t)) \cdot \mathbb{E}(V_1^E(t) - \hat{V}_1(t)) \tilde{f}_{t_k}. \end{split}$$

By Lemma 2 and Lemma 10, one has

$$|v - \hat{V}_1(t)| \le C e^{-\kappa \psi_0 t_k}.$$

Note that

$$|X_1(0) - \hat{X}_1(t)| \le C e^{-\kappa \psi_0 t_k} \tau$$

and

$$|\hat{V}_1(t) - V_1(0)| \le C e^{-\kappa \psi_0 t_k} \tau.$$

Take the expectation of $(X_1(0), V_1(0))$, one has

$$\frac{d}{dt}\mathbb{E}|V_1^E - \hat{V}_1|^2 \le Ce^{-\kappa\psi_0 t_k}\tau\sqrt{\mathbb{E}|V_1^E - \hat{V}_1|^2}.$$

Hence

$$\mathbb{E}|V_1^E - \hat{V}_1|^2 \le Ce^{-2\kappa\psi_0 t_k}\tau^4.$$

Similar with (5.12), one has

$$\mathbb{E}|X_1^E - \hat{X}_1|^2 \le Ce^{-2\kappa\psi_0 t_k}\tau^6.$$

Now consider

$$\frac{d}{dt} \frac{1}{2} \mathbb{E} |V_1^E(t) - V_1(t)|^2 = \mathbb{E} (V_1^E(t) - V_1(t)) \frac{\kappa}{p-1} \sum_{j=1}^p \left[\psi(X_j(0) - X_1(0))(V_j(0) - V_1(0)) - \psi(X_j(t) - X_1(t))(V_j(t) - V_1(t)) \right] \\
= \psi(X_j(t) - X_1(t))(V_j(t) - V_1(t)) \left[\|\psi\|_{\text{Lip}} |X_2(0) - X_1(0) - X_2(t) + X_1(t))(V_2(0) - V_1(0)) + \psi(X_2(t) - X_1(t))(V_2(0) - V_1(0) - V_2(t) + V_1(t)) \right].$$

By Lemma 2 and Lemma 7, one knows that $|V_j(t)| \leq e^{-\kappa \psi_0 t_k}$, for any j and $t \in [0, \tau]$. Hence it's easy to obtain

$$\frac{d}{dt}\mathbb{E}|V_1^E - V_1|^2 \le Ce^{-\kappa\psi_0 t_k}\tau\sqrt{\mathbb{E}|V_1^E - V_1|^2},$$

and

$$\mathbb{E}|V_1^E - V_1|^2 \le Ce^{-2\kappa\psi_0 t_k}\tau^4, \quad \mathbb{E}|X_1^E - \hat{X}_1|^2 \le Ce^{-2\kappa\psi_0 t_k}\tau^6.$$

Then, by the triangular inequality of Wasserstein distance, one has

$$W_2(\mathcal{Q}_{\infty}(\tilde{f}_{t_k}), \tilde{S}(\tilde{f}_{t_k})) \le C(\kappa, R)e^{-\kappa\psi_0 t_k}\tau^2.$$

The estimate of W_q is similar by calculating

$$\frac{d}{dt} (\mathbb{E}|V_1^E - \hat{V}_1|^q)^{\frac{1}{q}} = \frac{1}{q} (\mathbb{E}|V_1^E - \hat{V}_1|^q)^{\frac{1}{q} - 1} \frac{d}{dt} \mathbb{E}|V_1^E - \hat{V}_1|^q \le C(\kappa, R) e^{-\kappa\psi_0 t_k} \tau.$$

5.3 Proof of Theorem 2

Now we prove Theorem 2 by combining the above lemmas.

Proof of Theorem 2. By the triangular inequality, it holds

$$W_q(\mathcal{Q}^n_{\infty}(f_0), \tilde{f}(n\tau)) \le \sum_{m=1}^n W_q(\mathcal{Q}^{n-m+1}_{\infty}(\tilde{f}_{m-1}), \mathcal{Q}^{n-m}_{\infty}(\tilde{f}_m)).$$

And then, by the stability result in Lemma 9, it holds

$$\sum_{m=1}^{n} W_q(\mathcal{Q}_{\infty}^{n-m+1}(\tilde{f}_{m-1}), \mathcal{Q}_{\infty}^{n-m}(\tilde{f}_m)) \le \sum_{m=1}^{n} CW_q(\mathcal{Q}_{\infty}(\tilde{f}_{m-1}), \tilde{f}_m).$$

Again by the triangular inequality, one has

$$W_q(\mathcal{Q}_{\infty}(\tilde{f}_{m-1}), \tilde{f}_m) \le W_q(\mathcal{Q}_{\infty}(\tilde{f}_{m-1}), \tilde{S}(\tilde{f}_{m-1})) + W_q(\tilde{S}(\tilde{f}_{m-1}), \tilde{f}_m).$$

Then, by Lemma 11 and Lemma 12, we have

$$\sum_{m=1}^{n} W_q(\mathcal{Q}_{\infty}^{n-m+1}(\tilde{f}_{m-1}), \mathcal{Q}_{\infty}^{n-m}(\tilde{f}_m)) \le \sum_{m=1}^{n} C e^{-\kappa \psi_0(m-1)\tau} \tau^2 \le C\tau,$$
(5.19)

where the constant C is independent of n. One completes the proof.

Remark 3. Although our analysis is based on $W_{q,2}$ distance, the main results in Section 3 also work on $W_{q,p}$, $p \in [1, \infty)$, by noting that

$$||x||_p \le d^{1 - \frac{1}{q} + \frac{1}{p}} ||x||_q, \quad x \in \mathbb{R}^d, \quad 1 < p, q < \infty.$$

6 Application: the RBM-gPC

In this section, we show a simple application for the mean-field limit analysis for the Cucker-Smale model combined with the RBM. Based on the above analysis, and inspired by Carrillo's work [7], we can construct the numerical scheme for the approximation of stochastic mean-field models of swarming which preserve the nonnegativity of the solution.

Let us consider the stochastic mean-field equation for the distribution function $f(\theta, x, v; t)$:

$$\partial_t f + v \cdot \nabla_x f = \nabla_v \cdot [\mathcal{H}[f]f], \tag{6.1}$$

where

$$\mathcal{H}[f](\theta, x, v; t) = \kappa \int \int \psi(|x - y|, \theta)(v - w)f(\theta, y, w; t)dwdy,$$

with ψ satisfying positivity, boundedness, Lipschitz continuity and mononticity conditions: there exists a positive constant $\psi_M > 0$ such that

$$0 \le \psi(r, \theta) \le \psi_M, \, \forall r \ge 0, \quad \|\psi\|_{\operatorname{Lip}} < \infty,$$

$$(\psi(r_1,\theta) - \psi(r_2,\theta))(r_1 - r_2) \le 0, \quad r_1, r_2 \in \mathbb{R}_+.$$

Here, θ is a random variable from a measurable space (Ω, \mathcal{F}) to $(I_{\Theta}, \mathcal{B}_{\mathbb{R}})$, with a probability distribution $\pi(\theta)$, with $I_{\Theta} \subset \mathbb{R}$ and $\mathcal{B}_{\mathbb{R}}$ the Borel set.

It is easy to see that the total mass and momentum of (6.1) is conserved in time. Note that if we directly apply the classical stochastic Galerkin method to (6.1), the solution might lose its positivity. However, we can make use of the microscopic dynamics to approximate f. Our method is to consider the corresponding Cucker-Smale model with random input θ on the interaction kernel ψ :

$$\begin{cases} \partial_t x_i(t,\theta) = v_i(t,\theta), \ i = 1, \cdots, N, \\ \partial_t v_i(t,\theta) = \frac{\kappa}{N-1} \sum_{j=1}^N \psi(|x_j(t,\theta) - x_i(t,\theta)|, \theta)(v_j(t,\theta) - v_i(t,\theta)). \end{cases}$$

Denote $\{\Phi_k(\theta)\}_{j=0}^{\infty}$ to be the gPC basis with respect to π , which means that Φ_k is a polynomial of degree k, and these polynomials form an orthonormal basis of $L^2(\Omega, \mathcal{F}, \pi)$. We approximate the position and the velocity of *i*-th agent as follows:

$$x_i(\theta, t) \approx x_i^K = \sum_{k=0}^K {}^k \hat{x}_i(t) \Phi_k(\theta), \quad v_i(\theta, t) \approx v_i^K = \sum_{k=0}^K {}^k \hat{v}_i(t) \Phi_k(\theta),$$

where

$${}^k\hat{x}_i = \int x_i(\theta, t)\Phi_k(\theta)d\pi, \quad {}^k\hat{v}_i = \int v_i(\theta, t)\Phi_k(\theta)d\pi$$

We obtain the following system

$$\begin{cases} \frac{d}{dt}^{\ell} \hat{x}_{i}(t) = {}^{\ell} \hat{v}_{i}(t), \\ \frac{d}{dt}^{\ell} \hat{v}_{i}(t) = \frac{\kappa}{N-1} \sum_{j=1}^{N} \sum_{k=0}^{K} e_{\ell k}^{ij} ({}^{k} \hat{v}_{j}(t) - {}^{k} \hat{v}_{i}(t)), \end{cases} \quad l = 0, 1, \dots, K.$$
(6.2)

where

$$e_{\ell k}^{ij} = \frac{1}{|\Phi_{\ell}|^2} \int \psi(|x_j - x_i|, \theta) \Phi_k(\theta) \Phi_{\ell}(\theta) d\pi.$$

Then, we design Algorithm 3, on the generalized Polynomial Chaos expansion in the random space based on the RBM:

Algorithm 3: RBM-gPC

- Consider N samples (x_i, v_i) with i = 1, · · · , N from the initial f₀(x, v), and fix the batch size p;
 Apply the RBM on (6.2): for k = 1 to T/τ do
- **3** Divide $\{1, \dots, N\}$ into n = N/p batches randomly;
- 4 for each batch C_q do
- 5 Update the position and velocity change by $\begin{cases}
 \frac{d}{dt}^{\ell} \hat{x}_{i}(t) =^{\ell} \hat{v}_{i}(t), \\
 \frac{d}{dt}^{\ell} \hat{v}_{i}(t) = \frac{\kappa}{p-1} \sum_{j \in C_{q}} \sum_{k=0}^{K} e_{\ell k}^{ij}(^{k} \hat{v}_{j}(t) ^{k} \hat{v}_{i}(t)).
 \end{cases}$ 6 end
 7 end
 8 Reconstruct $\mathbb{E}_{\theta}[f(\theta, x, v; T)].$

Remark 4. Although being derived from different numerical methods (the RBM and the Monte Carlo method), Algorithm 3 is similar with the Monte Carlo gPC (MCgPC) proposed in [7] at first glance. Both can reduce the total cost of (6.2) to $\mathcal{O}(KN)$. The main difference between them is that the RBM-gPC preserves the momentum during evolution. Actually, since the RBM updates information in the unit of batches, (instead, the MCgPC updates information of each particle independently at per time step.) for any $t \in [t_m, t_{m+1}]$, the total sum of velocities is conserved in each batch by calculus:

$$\frac{d}{dt} \sum_{i \in \mathcal{C}_q} {}^{\ell} \hat{v}_i(t) = \frac{\kappa}{p-1} \sum_{i,j \in \mathcal{C}_q} \sum_{k=0}^K e^{ij}_{\ell k} ({}^k \hat{v}_j(t) - {}^k \hat{v}_i(t)) = 0,$$

for $l = 0, \dots, K$. Furthermore, it holds

$$\sum_{i=1}^{N} {}^{\ell} \hat{v}_i(t) = \sum_{q=1}^{N/p} \sum_{i \in \mathcal{C}_q} {}^{\ell} \hat{v}_i(t) = \sum_{q=1}^{N/p} \sum_{i \in \mathcal{C}_q} {}^{\ell} \hat{v}_i(t_m) = \sum_{q=1}^{N/p} \sum_{i \in \mathcal{C}_q} {}^{\ell} \hat{v}_i(t_0) = \sum_{i=1}^{N} {}^{\ell} \hat{v}_i(0).$$

Remark 5. We use

$$x_{i}^{K,R}(t,\theta) := \sum_{k=0}^{K} {}^{k} \hat{x}_{i}(t) \Phi_{k}(\theta), \quad v_{i}^{K,R}(t,\theta) := \sum_{k=0}^{K} {}^{k} \hat{v}_{i}(t) \Phi_{k}(\theta)$$

to denote the approximation of position and velocity obtained by Algorithm 3. The kinetic distribution $f_{N,K}^{R}(\theta, x, v, t)$ is recovered from the empirical density function

$$f_{N,K}^R(\theta, x, v, t) = \frac{1}{N} \sum_{i=1}^N \delta(x - x_i^{K,R}(t,\theta)) \otimes \delta(v - v_i^{K,R}(t,\theta)).$$

Common approximations methods for Dirac delta distributions can be used when needed. For example, we fix an upper and lower bound for the (x, v)-space, to restrict our computational domain in a cube space and discretize it in uniform cells $\{C_l\}_{l=1}^{N_l}$ of size $|C_l| = h^{2d}$, where h is the edge length of each cell. One can reconstruct the distribution $f_{N,K}^R$ by histograms counting the number of particles belonging to each cell. Then the discrete distribution function $\bar{f}_{N,K}^R$ reads

$$\bar{f}_{N,K}^{R}(\theta, x, v, t) = \frac{1}{|C_l|} \left(\frac{1}{N} \sum_{i=1}^{N} \mathbb{I}(x_i^{K,R}(t,\theta), v_i^{K,R}(t,\theta) \in C_l) \right) \cdot \left(\sum_{l=1}^{N_l} \mathbb{I}((x,v) \in C_l) \right), \quad (6.3)$$

where $\mathbb{I}(\cdot)$ is the indicator function. Clearly, the positivity of the distribution function is preserved. One can also refer to a brief review, by Lee et al. [27], on the numerical methods for regularized Dirac delta functions.

Remark 6. With respect to the stochastic variable θ , the dynamics of the N-particle system achieves spectral accuracy as we adopt the generalized polynomial chaos (gPC) basis [37], provided that we have a smooth dependence of the particle solution from the random field.

For any given θ and K, denote the distribution function of the N-particle system (6.2) by $f_{N,K}$. If one uses the reconstruction (6.3), it holds that

$$\mathbb{E}W_2(f_{N,K}^R, \bar{f}_{N,K}^R) \le \left(\mathbb{E}\frac{1}{N}\sum_{i=1}^N |(x_i^R, v_i^R) - c_{l(i)}|^2\right)^{\frac{1}{2}},$$

where $c_{l(i)}$ is the center of the cell C_l containing (x_i^R, v_i^R) . Since h is the uniform edge length, $|(x_i^R, v_i^R) - c_{l(i)}|^2 \leq \sum_{i=1}^d (h/2)^2$ for any i. Combining with the estimate of $\mathbb{E}W_2(f_{N,K}, f_{N,K}^R)$ in Section 3.3.2, it holds that

$$\mathbb{E}W_{2}(f_{N,K}, \bar{f}_{N,K}^{R}) \leq \mathcal{O}(\sqrt{\frac{\tau}{p-1}}) + \begin{cases} \mathcal{O}(N^{-\frac{1}{4}}) & \text{if } d < 2, \\ \mathcal{O}(N^{-\frac{1}{4}}\sqrt{\log(1+N)}) & \text{if } d = 2, + \mathcal{O}(\sqrt{d}h), \\ \mathcal{O}(N^{-\frac{1}{d}}) & \text{if } d > 2, \end{cases}$$

6.1 Numerical experiments

Here we present some numerical tests for the RBM-gPC. In the following, we take the Legendre gPC base.

The space homogeneous case We consider the space-independent case in the onedimensional setting:

$$\partial_t f(\theta, v; t) = \partial_v \left[K(\theta)(v - u) f(\theta, v; t) \right], \tag{6.4}$$

with $u = \int vf(v)dv$ and $K(\theta) = 0.5 + 0.01\theta$, $\theta \sim \mathcal{U}(0, 1)$. The long time solution is a Dirac delta $\delta(v - u)$ provided $K(\theta) > 0$, see [34, 6]. We take the Legendre polynomial basis. The gPC approximation is given for $h = 0, \dots, M$ by

$$\partial_t \hat{f}_h(v;t) = \frac{1}{|\Phi_h|^2} \partial_v \left[\sum_{k=0}^M (v-u) \mathcal{H}_{hk} \hat{f}_{t_k}(v;t) \right], \tag{6.5}$$

where

$$\mathcal{H}_{hk} = \int K(\theta) \Phi_h(\theta) \Phi_k(\theta) dg(\theta), \quad \hat{f}_h(v;t) = \int f(\theta,v;t) \Phi_h(\theta) dg(\theta).$$

We consider an initial density function $f_0(v)$ by

$$f_0(v) = \beta \left[\exp\left(-\frac{(v-\mu)^2}{2\sigma^2}\right) + \exp\left(-\frac{(v+\mu)^2}{2\sigma^2}\right) \right], \quad \sigma^2 = 0.1, \ \mu = 0.5.$$

with $\beta > 0$ a normalization constant. Discrete samples of the initial data of the ODE system are obtained from $f_0(v)$. Denote the expected temperature of the system \mathcal{T} by

$$\mathcal{T}(f) = \int (v-u)^2 f(\theta, v; t) dv dg(\theta).$$

In Figure 1, we show the evolution of the expected density toward the Dirac delta function $\delta(v-u)$, u = 0 for the RBM-gPC scheme at different times with p = 2 and $N = 10^4$. In



Figure 1: Evolution of the expected density $\delta(v-u)$, u = 0 for the RBM-gPC scheme.

Figure 2, we take T = 0.5, K = 3, $dt = 10^{-2}$ and $dv = 10^{-2}$. Time integration is performed through a 4th order Runge-Kutta method. In Figure 2(a), we evaluate the mean square error (MSE) for the expected temperature

$$\mathrm{MSE}_{\mathrm{T}} := \frac{1}{n_m} \sum_{i=1}^{n_m} \left| \mathcal{T}(f_{\mathrm{ref}}) - \mathcal{T}(f_{\mathrm{rbm}}^{(i)}) \right|^2,$$

as N grows with p = 2, where $\mathcal{T}(f_{\text{ref}})$ denotes the reference expected temperature obtained by (6.5) through a central difference scheme, and $\mathcal{T}(f_{\text{rbm}}^{(i)})$ denotes the expected temperature obtained by the RBM-gPC in the *i*-th turn. We have applied $n_m = 100$ for random initialization and batch splits. Figure 2(b) shows the difference of TV distance between the expected reference distribution and the expected distribution by the RBM-gPC:

$$\operatorname{Err}_{\mathrm{TV}} := \left| \mathbb{E}_{\theta} f_{\mathrm{ref}} - \mathbb{E}_{\theta} \frac{1}{n_m} \sum_{i=1}^{n_m} f_{\mathrm{rbm}}^{(i)} \right|_{L^1}$$

with $n_m = 100$ as well.



Figure 2: (a): The MSE_T of expected temperature between reference solution and the RBMgPC solution. (b): The Err_{TV} between the expected reference distribution and the expected distribution by the RBM-gPC. Both are obtained at T = 0.5, with $dt = dv = 10^{-2}$, p = 2.

Also, Figure 3(a) and Figure 3(b) show the decay of MSE_T as the batch size p and the time step dt, respectively. The reference distribution of Figure 3(a) is obtained by the RBM-gPC with $N = 2^8$, $p = 2^8$, and $dt = 10^{-2}$, and of Figure 3(b) is obtained with $N = 2^8$, p = 2, and $dt = 10^{-2}$. We still take $n_m = 100$.



Figure 3: (a):The MSE_T of expected temperature of the RBM-gPC solution, with $dt = 10^{-2}$ and batch size p varying. (b): The MSE_T of expected temperature of the RBM-gPC solution, with p = 2 and time step dt varying. Both are obtained at T = 0.5 with $N = 2^8$.

Stochastic 1D Cucker-Smale dynamics In this test, we consider the 1D Cucker-Smale dynamics. Let us consider the initial distribution as the following bivariate and bimodal

distribution

$$f_0(x,v) = \frac{1}{2\pi\sigma_x\sigma_v} \exp\left(-\frac{x^2}{2\sigma_v^2}\right) \left[\exp\left(-\frac{(v+1)^2}{2\sigma_x^2}\right) + \exp\left(-\frac{(v-1)^2}{2\sigma_x^2}\right)\right],$$

with $\sigma_x^2 = 0.5$, $\sigma_v^2 = 0.2$. Our initial data for particle positions and velocities are sampled from f_0 . The stochastic interactions are given by

$$H(\theta; |x_i - x_j|) = \frac{1}{(1 + |x_i - x_j|^2)^{\gamma(\theta)}},$$
(6.6)

with $\gamma(\theta) = 0.1 + 0.05\theta$, $\theta \sim \mathcal{U}[-1, 1]$. The results are obtained through the RBM-gPC scheme with $dt = 10^{-2}$, $N = 10^4$, batch size p = 2, and K = 3. The expected distribution is reconstructed by ksdensity in Matlab in the domain $[-3, 3] \times [-3, 3]$ with 100 gridpoints in both space and velocity. Figure 4 presents the evolution over the time interval $t \in [0, 4]$ of the expected distribution.



Figure 4: The expected distribution of the stochastic 1D Cucker-Smale dynamics with $N = 10^4$ agents.

Stochastic 2D Cucker-Smale dynamics Figure 5 shows the evolution over the time interval $t \in [0, 4]$ of the 2D Cucker-Smale nonhomogeneous mean-field model. For the initial distribution, we consider uniformly distributed $N = 10^4$ particles on a 2D annulus with a circular counterclockwise motion

$$f_0(x,v) = \frac{1}{|C|} \chi(x \in C) \delta\left(v - \frac{k \wedge x}{|x|}\right),$$

being $C := \{x \in \mathbb{R}^2 : 0.5 \le |x_1 - x_2| \le 1\}$ and k the fundamental unit vector of the z-axis, where |C| means the cardinality of set C. The interaction kernel shares the same form with (6.6). The evolution shows the flocking phenomenon in the top row and that the mean velocity is conserved in the bottom row of Figure 5. The results are obtained through the RBM-gPC scheme with $dt = 10^{-2}$, $N = 10^4$, batch size p = 2, and K = 3. The reconstruction follows the same method as in the 1D case.

7 Conclusion

In conclusion, we give an analysis on the mean-field limit of the RBM for the Cucker-Smale model and obtain a long-time approximation. In addition, as a simple application, we propose a generalized Polynomial Chaos based Random Batch Method (RBM-gPC) as a numerical method to simulate the mean-field limit dynamics of the Cucker-Smale model with random inputs.

This paper focuses on the separate limits of the number of particles N and the time step τ but not a uniform limit of (N, τ) . For future work, it will be interesting to investigate the mixture of particles as time evolves, and to study on the uniform limit. Also, numerically, an error estimate on this model applied with the particle method (corresponding to a new model) can be an interesting topic to study.



Figure 5: The expected distribution of the stochastic 2D Cucker-Smale dynamics with $N = 10^4$ agents.

Acknowledgement

We thank Prof. Shi Jin and Prof. Lei Li for detailed and valuable comments and advice on the manuscript. This work is supported by the NSFC grant (No. 12031013, 12201404), the Shanghai Municipal Science and Technology Key Project (No. 22JC1402300), and Project supported by the National Science Foundation for International Senior Scientists (No. 12350710181).

A Proof of Lemma 7

The proof is given by Lemma 2.4 in [15].

Proof of Lemma 7. For the first assertion, we claim that the relative velocities are nonincreasing in time. Let $t \in [t_{m-1}, t_m)$ be given. Then one can choose time-dependent indices k and ℓ such that

$$|V_k^R(t) - V_\ell^R(t)| = \mathcal{D}_{V^R}(t).$$

Then for such k and ℓ , one has

$$\begin{aligned} \frac{d}{dt} |V_k^R(t) - V_\ell^R(t)|^2 &= 2(V_k^R(t) - V_\ell^R(t)) \cdot \frac{d}{dt} (V_k^R(t) - V_\ell^R(t)) \\ &= \frac{2\kappa}{p-1} \sum_{j \in [k]_m} \psi(|X_j^R - X_k^R|) (V_j^R - V_k^R) \cdot (V_k^R - V_\ell^R) \\ &- \frac{2\kappa}{p-1} \sum_{j \in [i]_m} \psi(|X_j^R - X_\ell^R|) (V_j^R - V_\ell^R) \cdot (V_k^R - V_\ell^R). \end{aligned}$$
(A.1)

In order to show that the right-hand side of (A.1) is not positive, we use the maximality of $|V_k^R(t) - V_\ell^R(t)|$ at time t. Since

$$|V_k^R(t) - V_\ell^R(t)| \ge |V_j^R(t) - V_\ell^R(t)|, \quad j = 1, \cdots, N_\ell$$

one has

$$\begin{split} (V_j^R(t) - V_k^R(t)) \cdot (V_k^R(t) - V_\ell^R(t)) &= -((V_k^R - V_\ell^R) - (V_j^R - V_\ell^R)) \cdot (V_k^R - V_\ell^R) \\ &\leq -|V_k^R(t) - V_\ell^R(t)|^2 + |V_j^R(t) - V_\ell^R(t)||V_k^R(t) - V_\ell^R(t)| \leq 0, \quad j = 1, \cdots, N. \end{split}$$

Similarly, one has

$$(V_j^R(t) - V_\ell^R(t)) \cdot (V_k^R(t) - V_\ell^R(t)) \ge 0, \quad j = 1, \cdots, N.$$

Hence, $|V_k^R(t) - V_\ell^R(t)|^2$ is non-increasing in time. Then by the definition of \mathcal{D}_{X^R} and \mathcal{D}_{V^R} , one has

$$\frac{d}{dt}\mathcal{D}_{X^R}(t) \le \mathcal{D}_{V^R}(t) \le \mathcal{D}_{V^R}(0), \quad a.e.\, t > 0.$$

This yields the second assertion on \mathcal{D}_{X^R} .

Β Details on Proposition 3

Fixed-time regularity of \tilde{f} . First we check the regularity of \tilde{f} by Lemma 13, which is needed in the proof of Proposition 3.

Lemma 13. For the flocking kinetic model (2.6), suppose that the initial probability measure $f_0 \in (C^2 \cap W^{2,\infty})(\mathbb{R}^{2d})$ is compactly supported, and the interaction kernel ψ satisfies

$$\sup_{r\geq 0}(|\psi'(r)|+|\psi''(r)|)\leq \epsilon_{\psi}$$

with a constant ϵ_{ψ} . Then, for any $T \in (0, \infty)$, there exists a unique classical solution $\tilde{f}_t \in$ $C^{2}([0,T) \times \mathbb{R}^{2d})$ for (2.6).

The proof stem of Lemma 13 is generally a smoothness check for \tilde{f} inspired by in [[19], Theorem 3.1]. The main difference with [19] is that here we need better regularity for the operator $(\bar{\mathcal{L}}^*)^2$ and $(\tilde{\mathcal{L}}^*_k)^2$ in Lemma 12.

Proof of Lemma 13. Local existence follows from [[19] Theorem 3.1] by the standard fixed point argument. One only needs to obtain a C^2 bound of f to conclude the global existence of classical solutions. One can express the non-conservative kinetic model (2.6) in terms of the non-linear transport operator $\mathcal{T} := \partial_t + v \cdot \nabla_x + \xi[\tilde{f}] \cdot \nabla_v$,

$$\mathcal{T}\tilde{f} = -\tilde{f}\nabla_v \cdot \xi[\tilde{f}].$$

We claim that there exists a positive constant $C(d, \kappa, \epsilon_{\psi}, R)$ such that

$$\left|\mathcal{T}\tilde{f}\right| \le C|\tilde{f}|,\tag{B.1}$$

$$\left| \mathcal{T}(\partial_{x_i} \tilde{f}) \right| \le C(|\tilde{f}| + |\nabla_v \tilde{f}| + |\partial_{x_i} \tilde{f})|, \tag{B.2}$$

$$\left| \mathcal{T}(\partial_{v_i} \tilde{f}) \right| \le C(|\nabla_v \tilde{f}| + |\partial_{x_i} \tilde{f}|), \tag{B.3}$$

$$\left| \mathcal{T}(\partial_{x_i x_j} \tilde{f}) \right| \le C(\left| \partial_{x_i} \tilde{f} \right| + \left| \partial_{x_j} \tilde{f} \right| + \left| \partial_{x_i x_j} \tilde{f} \right| + \left| \nabla_v \tilde{f} \right| + \left| \nabla_v \partial_{x_i} \tilde{f} \right| + \left| \nabla_v \partial_{x_j} \tilde{f} \right|), \tag{B.4}$$

$$\left| \mathcal{T}(\partial_{x_i v_j} \tilde{f}) \right| \le C(\left| \partial_{x_i x_j} \tilde{f} \right| + \left| \nabla_v \tilde{f} \right| + \left| \partial_{v_j} \partial_{x_i} \tilde{f} \right| + \left| \nabla_v \partial_{v_j} \tilde{f} \right| + \left| \nabla_v \partial_{x_i} \tilde{f} \right|), \tag{B.5}$$

$$\left| \mathcal{T}(\partial_{v_i v_j} \tilde{f}) \right| \le C(\left| \partial_{x_i v_j} \tilde{f} \right| + \left| \partial_{x_j} \partial_{v_i} \tilde{f} \right| + \left| \partial_{v_j} \partial_{v_i} \tilde{f} \right|).$$
(B.6)

To verify these inequalities, note that $-\nabla_v \cdot \xi[\tilde{f}] = \kappa d \int \psi(|y-x|)\tilde{f}(y,w,t)dydw$ and thus $|\nabla_v \cdot \xi[\tilde{f}]| \leq \kappa d$. By direct calculus, one obtains the following estimates:

For (B.1),

$$\left|\mathcal{T}\tilde{f}\right| \leq \left|-\kappa\tilde{f}\nabla_{v}\cdot\int\psi(|y-x|)(w-v)\tilde{f}(y,w,t)dydw\right| \leq \kappa d|\tilde{f}|;$$

for (B.2),

$$\begin{aligned} \left| \mathcal{T}(\partial_{x_i}\tilde{f}) \right| &\leq \left| -(\partial_{x_i}\xi[\tilde{f}]) \cdot \nabla_v \tilde{f} - (\partial_{x_i}\nabla_v \cdot \xi[\tilde{f}])\tilde{f} - (\nabla_v \cdot \xi[\tilde{f}])\partial_{x_i}\tilde{f} \right| \\ &\leq \kappa \epsilon_\psi (\sqrt{M_2^v f_0} + 2R) |\nabla_v \tilde{f}| + \kappa d\epsilon_\psi |\tilde{f}| + \kappa d|\partial_{x_i}\tilde{f}|) \\ &\leq C(|\tilde{f}| + |\nabla_v \tilde{f}| + |\partial_{x_i}\tilde{f})|; \end{aligned}$$

For (B.3),

$$|\mathcal{T}\partial_{v_i}\tilde{f}| \le |-\partial_{x_i}\tilde{f}| + \kappa |\nabla_v\tilde{f}| + \kappa d|\partial_{v_i}\tilde{f}| \le C(|\nabla_v\tilde{f}| + |\partial_{x_i}\tilde{f}|).$$

The estimate on (B.4)-(B.6) is similar and we omit it for brevity.

Now, let F(t) measure the $W^{2,\infty}$ -norm of \tilde{f} ,

$$F := \|\tilde{f}(\cdot, t)\|_{W^{2,\infty}}$$

The above inequalities imply

$$\frac{d}{dt}F(t) \le CF(t).$$

Then we end up with the energy bound by Grönwall's inequality $F(t) \leq F(0)e^{Ct}$. Equipped with this a priori $W^{2,\infty}$ estimate, standard continuation principle yields a global extension of local classical solutions.

The following is a simple prerequisite for Proposition 3.

Lemma 14. For any $g_1 \in L^1(\mathbb{R}^{2d})$, $g_2 \in L^1(\mathbb{R}^{2dp})$, the operators $\tilde{\mathcal{L}}_k^*$ and $\bar{\mathcal{L}}^*$ defined by (B.9) and (B.10) respectively, it holds

$$\int |e^{\tau \tilde{\mathcal{L}}_k^*} g_1| dz \le \int |g_1| dz, \tag{B.7}$$

$$\int \left| e^{\tau \bar{\mathcal{L}}^*} g_2 \right| dz_1 \cdots dz_p \le \int \left| g_2 \right| dz_1 \cdots dz_p.$$
(B.8)

Proof. One only needs to consider $g_1 \ge 0$ since there is composition $g_1 = g_1^+ - g_1^-$. Also, since $e^{\tau \mathcal{L}_k^*} cg_1 = c e^{\tau \mathcal{L}_k^*} g_1$ and $e^{\tau \bar{\mathcal{L}}_k^*} cg_2 = c e^{\tau \bar{\mathcal{L}}_k^*} g_2$ for any constant c, one can set g_1 be a probability density function. Then,

$$\int |e^{\tau \tilde{\mathcal{L}}_k^*} g_1| dz = \int e^{\tau \tilde{\mathcal{L}}_k^*} g_1 dz,$$

since $e^{\tau \tilde{\mathcal{L}}_k^*} g_1$ is another probability density function derived from g_1 by the dynamics (5.14). It's clear that

$$\int e^{\tau \tilde{\mathcal{L}}_k^*} g_1 dz = \int \int_0^\tau \frac{\partial}{\partial t} \tilde{g}_1(t) dt dz + \int g_1 dz = \int g_1 dz,$$

where $\tilde{g}_1(t)$ denotes the solution of (2.6) with initial datum g_1 . For the general g_1 ,

$$\int |e^{\tau \tilde{\mathcal{L}}_{k}^{*}} g_{1}| dz \leq \int |e^{\tau \tilde{\mathcal{L}}_{k}^{*}} g_{1}^{+}| dz + \int |e^{\tau \tilde{\mathcal{L}}_{k}^{*}} g_{1}^{-}| dz = \int |g_{1}^{+}| dz + \int |g_{1}^{-}| dz = \int |g_{1}| dz.$$

The proof of (B.8) is similar.

Proof of Proposition 3. It's clear that $\tilde{S} = e^{\tau \tilde{\mathcal{L}}_k^*} \tilde{f}_{t_k}$, where

$$\tilde{\mathcal{L}}_{k}^{*} := -(v \cdot \nabla_{x}) \cdot -\nabla_{v} \cdot \left(\int \kappa \psi(|y-x|)(w-v)f_{t_{k}}(y,w)dydw \cdot \right).$$
(B.9)

Define the operator of (2.12) as

$$\bar{\mathcal{L}}^* := -\sum_{i=1}^p (v_i \nabla_{x_i}) \cdot -\sum_{i=1}^p \nabla_{v_i} \cdot (\xi_i \cdot).$$
(B.10)

Then one has

$$\begin{aligned} \mathcal{Q}_{\infty}(\tilde{f}_{t_k}) &= \int e^{\tau \bar{\mathcal{L}}^*} \Pi_{i=1}^p \tilde{f}_{t_k}(x_i, v_i) dx_2 \cdots dx_p dv_2 \cdots dv_p \\ &= \tilde{f}_{t_k}(x_1, v_1) + \tau \int \bar{\mathcal{L}}^* \Pi_{i=1}^p \tilde{f}_{t_k}(x_i, v_i) dx_2 \cdots dx_p dv_2 \cdots dv_p \\ &+ \int_0^\tau (\tau - s) \int (\bar{\mathcal{L}}^*)^2 e^{s \bar{\mathcal{L}}^*} \Pi_{i=1}^p \tilde{f}_{t_k}(x_i, v_i) dx_2 \cdots dx_p dv_2 \cdots dv_p ds, \end{aligned}$$

where

$$\int \bar{\mathcal{L}}^* \Pi_{i=1}^p \tilde{f}_{t_k}(z_i) dz_2 \cdots dz_p = -\sum_{i=1}^p \int (v_i \nabla_{x_i}) \cdot \Pi_{j=1}^p \tilde{f}_{t_k}(z_j) - \nabla_{v_i} \cdot (\xi_i \Pi_{j=1}^p) \tilde{f}_{t_k}(z_j) dz_2 \cdots dz_p)$$

= $-(v_1 \nabla_{x_1}) \cdot \tilde{f}_{t_k}(z_1) - \nabla_{v_1} \cdot \int \kappa \psi(|y-x_1|) (w-v_1) \tilde{f}_{t_k}(y,w) dy dw.$

Note that $\tilde{S}(\tilde{f}_{t_k}) = \tilde{f}_{t_k} + \tau \tilde{\mathcal{L}}_k^* \tilde{f}_{t_k} + \int_0^\tau (\tau - s) (\tilde{\mathcal{L}}_k^*)^2 e^{s \tilde{\mathcal{L}}_k^*} \tilde{f}_{t_k} ds$, then $\left|\tilde{S}(\tilde{f}_{t_k}) - \mathcal{Q}_{\infty}(\tilde{f}_{t_k})\right|(z_1) \leq \int_0^\tau (\tau - s) \left[\left| (\tilde{\mathcal{L}}_k^*)^2 e^{s\tilde{\mathcal{L}}_k^*} \tilde{f}_{t_k}(z_1) \right| + \left| \int (\bar{\mathcal{L}}^*)^2 e^{s\bar{\mathcal{L}}^*} \Pi_{i=1}^p \tilde{f}_{t_k}(z_i) dz_2 \cdots dz_p \right| \right] ds.$

As shown in Lemma 13, $\tilde{f}_{t_k} \in C^2(z)$, hence it holds

$$\int |(\tilde{\mathcal{L}}_k^*)^2 \tilde{f}_{t_k}| dz \le C, \quad \int \left| (\bar{\mathcal{L}}^*)^2 \Pi_{i=1}^p \tilde{f}_{t_k}(z_i) \right| dz_2 \cdots dz_p \le C.$$

where $C = C(\kappa, \epsilon_{\psi}, R, d)$. By Lemma 14, taking $g_1 = (\tilde{\mathcal{L}}_k^*)^2 \tilde{f}_{t_k}$ and $g_2 = (\bar{\mathcal{L}}^*)^2 \Pi_{i=1}^p \tilde{f}_{t_k}(z_i)$, one has

$$\int |(\tilde{\mathcal{L}}_k^*)^2 e^{\tau \tilde{\mathcal{L}}_k^*} \tilde{f}_{t_k} | dz \le \int |(\tilde{\mathcal{L}}_k^*)^2 \tilde{f}_{t_k} | dz$$

and

$$\int \left| (\bar{\mathcal{L}}^*)^2 e^{\tau \bar{\mathcal{L}}^*} \Pi_{i=1}^p \tilde{f}_{t_k}(z_i) \right| dz_1 \cdots dz_p \leq \int \left| (\bar{\mathcal{L}}^*)^2 \Pi_{i=1}^p \tilde{f}_{t_k}(z_i) \right| dz_1 \cdots dz_p$$

Using the exchangability of \mathcal{L}^* and $e^{\tau \mathcal{L}^*}$, with $\mathcal{L}^* = \tilde{\mathcal{L}}_k^*$ or $\bar{\mathcal{L}}^*$. Then one has $|\tilde{S}(\tilde{f}_{t_k}) - \tilde{\mathcal{L}}_k^*|$ $\begin{array}{l} \mathcal{Q}_{\infty}(\tilde{f}_{t_k})|_{TV} \leq C\tau^2.\\ \text{For }\tilde{S}(\tilde{f}_{t_k}), \text{ by Lemma 10, it holds} \end{array}$

$$\operatorname{supp}[\tilde{S}_{\infty}(\tilde{f}_{t_k})] \subset B(0,C) \times B(0,Ce^{-\kappa\psi_0 t}).$$

For the support of $\mathcal{Q}_{\infty}(\tilde{f}_{t_k})$, by Lemma 2, one knows that

$$\operatorname{supp}[\mathcal{Q}_{\infty}(\tilde{f}_{t_k})] \subset B(0,C) \times B(0,Ce^{-\kappa\psi_0 t_k}).$$

Hence for any $z \in \text{supp}|\tilde{S}(\tilde{f}_{t_k}) - \tilde{f}_{t_k}|$, it holds $|z| \leq C$. Then, by Lemma 3 and applying similar analysis on Lemma 8 Step 1, one has

$$W_q(\mathcal{Q}_{\infty}(\tilde{f}_{t_k}), \tilde{S}(\tilde{f}_{t_k})) \le C(\kappa, \epsilon_{\psi}, f_0, d) \tau^{\frac{2}{q}}.$$

References

- Giacomo Albi, Nicola Bellomo, Luisa Fermo, Seung-Yeal Ha, Jeongho Kim, Lorenzo Pareschi, David Poyato, and Juan Soler. Vehicular traffic, crowds, and swarms: From kinetic theory and multiscale methods to applications and research perspectives. *Mathematical Models and Methods in Applied Sciences*, 29(10):1901–2005, 2019.
- [2] Nicola Bellomo and Seung-Yeal Ha. A quest toward a mathematical theory of the dynamics of swarms. Mathematical Models and Methods in Applied Sciences, 27(04):745– 770, 2017.
- [3] Nicola Bellomo and J Soler. On the mathematical theory of the dynamics of swarms viewed as complex systems. Mathematical Models and Methods in Applied Sciences, 22(supp01):1140006, 2012.
- [4] Dario Benedetto, Emanuele Caglioti, and Mario Pulvirenti. A kinetic equation for granular media. ESAIM: Mathematical Modelling and Numerical Analysis, 31(5):615– 641, 1997.
- [5] John Buck and Elisabeth Buck. Biology of synchronous flashing of fireflies, 1966.
- [6] José A Carrillo, Massimo Fornasier, Jesús Rosado, and Giuseppe Toscani. Asymptotic flocking dynamics for the kinetic Cucker–Smale model. SIAM Journal on Mathematical Analysis, 42(1):218–236, 2010.
- [7] José A Carrillo, Lorenzo Pareschi, Mattia Zanella, et al. Particle based gpc methods for mean-field models of swarming with uncertainty. *Comunications in Computational Physics*, 25(2):508–531, 2019.
- [8] Louis-Pierre Chaintron and Antoine Diez. Propagation of chaos: A review of models, methods and applications. i. models and methods. *Kinetic and Related Models*, 15(6):895–1015, 2022.
- [9] Louis-Pierre Chaintron and Antoine Diez. Propagation of chaos: A review of models, methods and applications. ii. applications. *Kinetic and Related Models*, 15(6):1017– 1173, 2022.
- [10] Felipe Cucker and Steve Smale. Emergent behavior in flocks. *IEEE Transactions on automatic control*, 52(5):852–862, 2007.
- John M Danskin. The theory of max-min, with applications. SIAM Journal on Applied Mathematics, 14(4):641–664, 1966.
- [12] Pierre Degond and Sébastien Motsch. Continuum limit of self-driven particles with orientation interaction. Mathematical Models and Methods in Applied Sciences, 18(supp01):1193–1215, 2008.
- [13] Florian Dörfler and Francesco Bullo. Synchronization in complex networks of phase oscillators: A survey. Automatica, 50(6):1539–1564, 2014.
- [14] Nicolas Fournier and Arnaud Guillin. On the rate of convergence in wasserstein distance of the empirical measure. Probability theory and related fields, 162(3-4):707-738, 2015.
- [15] Seung-Yeal Ha, Shi Jin, Doheon Kim, and Dongnam Ko. Uniform-in-time error estimate of the random batch method for the Cucker–Smale model. *Mathematical Models and Methods in Applied Sciences*, 31(06):1099–1135, 2021.
- [16] Seung-Yeal Ha, Jeongho Kim, and Xiongtao Zhang. Uniform stability of the Cucker-Smale model and its application to the mean-field limit. *Kinetic and Related Models*, 11(5):1157–1181, 2018.

- [17] Seung-Yeal Ha, Dongnam Ko, Jinyeong Park, and Xiongtao Zhang. Collective synchronization of classical and quantum oscillators. *EMS Surveys in Mathematical Sciences*, 3(2):209–267, 2016.
- [18] Seung-Yeal Ha and Jian-Guo Liu. A simple proof of the Cucker-Smale flocking dynamics and mean-field limit. *Communications in Mathematical Sciences*, 7(2):297–325, 2009.
- [19] Seung-Yeal Ha and Eitan Tadmor. From particle to kinetic and hydrodynamic descriptions of flocking. *Kinetic and Related Models*, 1(3):415–435, 2008.
- [20] Pierre-Emmanuel Jabin. A review of the mean field limits for vlasov equations. *Kinetic and Related models*, 7(4):661–711, 2014.
- [21] Shi Jin and Lei Li. On the mean field limit of the Random Batch Method for interacting particle systems. *Science China Mathematics*, pages 1–34, 2022.
- [22] Shi Jin, Lei Li, and Jian-Guo Liu. Random batch methods (rbm) for interacting particle systems. Journal of Computational Physics, 400:108877, 2020.
- [23] Shi Jin, Lei Li, and Yiqun Sun. On the Random Batch Method for second order interacting particle systems. *Multiscale Modeling & Simulation*, 20(2):741–768, 2022.
- [24] Shi Jin and Lorenzo Pareschi. Uncertainty quantification for hyperbolic and kinetic equations. Springer, 2017.
- [25] Eric W Justh and Perinkulam S Krishnaprasad. A simple control law for uav formation flying. Technical report, Technical Report 2002-38, Institute for Systems Research, 2002.
- [26] Yoshiki Kuramoto. Self-entrainment of a population of coupled non-linear oscillators. In International Symposium on Mathematical Problems in Theoretical Physics: January 23–29, 1975, Kyoto University, Kyoto/Japan, pages 420–422. Springer, 1975.
- [27] Hyun Geun Lee and Junseok Kim. Regularized dirac delta functions for phase field models. International journal for numerical methods in engineering, 91(3):269–288, 2012.
- [28] Sebastien Motsch and Eitan Tadmor. A new model for self-organized dynamics and its flocking behavior. *Journal of Statistical Physics*, 144:923–947, 2011.
- [29] Roberto Natalini and Thierry Paul. On the mean field limit for Cucker-Smale models. arXiv preprint arXiv:2011.12584, 2020.
- [30] Charles S Peskin. Mathematical aspects of heart physiology. Courant Inst. Math, 1975.
- [31] Benedetto Piccoli, Francesco Rossi, and Emmanuel Trélat. Control to flocking of the kinetic Cucker–Smale model. SIAM Journal on Mathematical Analysis, 47(6):4685– 4719, 2015.
- [32] John Toner and Yuhai Tu. Flocks, herds, and schools: A quantitative theory of flocking. *Physical review E*, 58(4):4828, 1998.
- [33] Chad M Topaz and Andrea L Bertozzi. Swarming patterns in a two-dimensional kinematic model for biological groups. SIAM Journal on Applied Mathematics, 65(1):152– 174, 2004.
- [34] Andrea Tosin and Mattia Zanella. Boltzmann-type models with uncertain binary interactions. arXiv preprint arXiv:1709.02353, 2017.
- [35] Tamás Vicsek, András Czirók, Eshel Ben-Jacob, Inon Cohen, and Ofer Shochet. Novel type of phase transition in a system of self-driven particles. *Physical review letters*, 75(6):1226, 1995.

- [36] Cédric Villani et al. Optimal transport: old and new, volume 338. Springer, 2009.
- [37] Dongbin Xiu and George Em Karniadakis. The wiener–askey polynomial chaos for stochastic differential equations. SIAM journal on scientific computing, 24(2):619–644, 2002.