

WHITNEY EXTENSION THEOREMS ON SYMMETRIC SPACES, AN EXAMPLE.

BIRGIT SPEH  AND PETER VANG UTTENTHAL 

ABSTRACT. H. Whitney introduced in 1934 the problem of extending a function f on a set of points in \mathbb{R}^n to an analytic function on the ambient space. In this article we prove Whitney type extension theorems for data on some homogeneous spaces. We use harmonic analysis on the homogeneous spaces and representation theory of compact as well as noncompact reductive groups.

CONTENTS

List of Figures	2
1. Introduction	2
1.1. Main results	3
1.2. Related work	4
1.3. Contents	4
1.4. Acknowledgements	6
2. Low dimensional examples	6
2.1. The sphere S^2	6
2.2. Preliminaries	7
2.3. The first interpolation theorem	7
2.4. Harmonic polynomials on \mathbb{R}^3	8
2.5. An open subset $\mathbb{R}^2 \setminus \overline{B}^1$ of the plane	9
3. The symmetric space $X(2, 2)_+$	10
3.1. Preliminaries	10
3.2. From rational data to Whitney functions. The first case	12
3.3. Cayley's rational parametrization of $\mathrm{SO}(n)$	14
3.4. From rational data sets to Whitney functions: the general case	16
3.5. From Whitney functions to Whitney representations	17
3.6. An exact Whitney theorem	19
4. The higher dimensional symmetric spaces $X_{\pm}(p, 2)$	20
4.1. Generalities	20
4.2. Whitney functions on $X(p, 2)_- = \mathrm{SO}(p, 2)^\circ / \mathrm{SO}(p, 1)^\circ$	22
4.3. Whitney representations	23
4.4. Data in orbits of orthogonal subgroups	23
5. Whitney representations in elementary particle physics	23
References	24

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NY 14853-4201, USA

DEPARTMENT OF MATHEMATICS, AARHUS UNIVERSITY, 8000 AARHUS C, DENMARK

E-mail addresses: speh@math.cornell.edu, petervang@math.au.dk.

Date: August 1, 2024.

LIST OF FIGURES

1	Anti de Sitter space from a physicist's perspective [18, J. Maldacena, <i>The Illusion of Gravity</i> , Scientific American (2005) no. 293]	3
2	Data on orbits in a symmetric space	13
3	Data in general position on a symmetric space	13

1. INTRODUCTION

In 1934, H. Whitney introduced the problem of extending a function f on a set of points in \mathbb{R}^n to an analytic function on the ambient space. In the 2000s, C. Fefferman took up Whitney's work and proved a sharp form of the Whitney extension theorem [5]. In particular, if the set of points in Euclidean space is finite of cardinality n , there is an extension which is continuously differentiable of some degree depending on n .

In this paper, we study a version of Whitney's extension problem where the points have an inherent symmetry. More precisely, we consider points $x^{(i)}$ in a symmetric homogeneous space $X = G/H$ for reductive Lie groups G, H and a nonconstant function f on these points with values $f(x^{(i)}) = y^{(i)}$ in the complex numbers \mathbb{C} . We are especially interested not just in proving existence of an analytic function extending f , but to obtain a formula for this function which can be effectively computed by an algorithm. Initially, we weaken the assumption

$$f(x^{(i)}) = y^{(i)}, \quad 1 \leq i \leq n.$$

Instead, we find for all $\varepsilon > 0$ a complex valued function $f_n : X \rightarrow \mathbb{C}$ so that

$$|f_n(x^{(i)}) - y^{(i)}| < \varepsilon, \quad 1 \leq i \leq n.$$

Observe that the function f_n depends on ε .

Furthermore, we show how to modify the proof to obtain an *exact* Whitney extension theorem. The function f_n generates a representation Π_n of G , which we call the Whitney representation of the data. By acting with the maximal compact subgroup K in G , we modify the Whitney function without changing the K -types occurring in its Whitney representation to obtain an analytic, square integrable function, still denoted f_n , such that $f_n(x^{(i)}) = y^{(i)}$ for all $i \leq n$.


We consider symmetric spaces of the form G/H for certain orthogonal Lie groups G and H . The analysis on symmetric spaces was intensively studied starting with the work of M. Flensted-Jensen in the late 1970s and culminating in the Plancherel theorem around 2005 by H. Schlichtkrull and E. van den Ban. In this article, we use information about the representation of the symmetry group G on the square integrable functions on G/H and in particular about the irreducible subrepresentations. We also use results about the restriction of these representations to orthogonal subgroups.

The group of symmetries of the quadrics

$$X(p, q)_{\pm} = \{(x_i) \in \mathbb{R}^{p+q} | x_1^2 + \dots + x_p^2 - x_{p+1}^2 \dots - x_{p+q}^2 = \pm 1\}$$

is the noncompact Lie group $G = \mathrm{SO}(p, q)^{\circ}$, where $p \geq 2$ and the superscript denotes the connected component.

This class of homogeneous spaces contains the spaces known in physics as



Anti de Sitter space from a physicist's perspective [18, J. Maldacena, *The Illusion of Gravity*, Scientific American (2005) no. 293]

de Sitter spaces (dS)

respectively as

anti de Sitter spaces (AdS)

[30, Chapters X and XI]. The anti de Sitter space is considered in physics as a model of space time with negative curvature.

1.1. Main results. We prove

Theorem 1.1. *Let $p \geq 2$. Let $x^{(i)}$, $1 \leq i \leq n$, be a set of points in $X(p, 2)_-$ and let $f(x^{(i)}) = y^{(i)}$ be a nonconstant, complex-valued function. For all $\varepsilon > 0$, there exists an analytic, square integrable function $f_n : X(p, 2)_- \rightarrow \mathbb{C}$ such that*

$$|f_n(x^{(i)}) - y^{(i)}| < \varepsilon.$$

We call f_n the Whitney function of the data points.

We approach the extension problem by realizing space $X(p, 2)_-$ as the homogeneous space

$$X(p, 2)_- = \mathrm{SO}(p, 2)^\circ / \mathrm{SO}(p, 1)^\circ.$$

This allows us to obtain an explicit formula for the Whitney function f_n using the representation theory of noncompact Lie groups. In particular, we build on the work of M. Flensted-Jensen [8], who published in 1980 an article in which he constructs a family of square integrable functions on symmetric spaces which are eigenfunctions of all invariant differential operators. We refer to these functions as Flensted-Jensen functions.

Theorem 1.2. *Let $x^{(i)}$, $1 \leq i \leq n$, be a set of points in $X(p, 2)_-$ and let $f(x^{(i)}) = y^{(i)}$ be a nonconstant complex-valued function. In the notation of Theorem 1.1, the Whitney function f_n can be chosen as a \mathbb{C} -linear combination of Flensted-Jensen functions*

$$\varphi_\ell(\xi', t) = (\xi'_1 - \sqrt{-1}\xi'_2)^\ell \cosh(t)^{-\ell},$$

for integers $\ell \geq \ell_0$ for some lower bound ℓ_0 depending on p . Here, (ξ, ξ') = $(\xi_1, \dots, \xi_p, \xi'_1, \xi'_2)$ is an angular variable and t a radial variable such that

$$(\xi_1 \sinh(t), \xi_2 \sinh(t), \dots, \xi_p \sinh(t), \xi'_1 \cosh(t), \xi'_2 \cosh(t)) \in X(p, 2)_-.$$

In applications of the theorem to a list of observable data, we may assume that the results of all measurements are rational. In the case of the three-dimensional symmetric space $X(2, 2)_+ = \mathrm{SO}(2, 2)^\circ / \mathrm{SO}(1, 2)^\circ$, we prove therefore also a rational version of our Whitney extension theorem starting from the assumption that all the data points are rational.

Theorem 1.3. *Let $x^{(i)} \in X(2, 2)_+(\mathbb{Q})$, $1 \leq i \leq n$, be a list of \mathbb{Q} -valued points in the three-dimensional symmetric space $X(2, 2)_+$ realized as the quadric surface*

$$X(2, 2)_+ = \{x \in \mathbb{R}^4 : x_1^2 + x_2^2 - x_3^2 - x_4^2 = 1\}$$

and suppose f is a nonconstant function with $f(x^{(i)}) = y^{(i)} \in \mathbb{Q}$. Let $\varepsilon > 0$. We can choose coefficients a_ℓ in the number field $\mathbb{Q}(\sqrt{-1})$ such that

$$f_n = \sum_{2 \leq \ell \leq n+1} a_\ell \varphi_\ell$$

is a Whitney function of the points: f_n is analytic, square integrable and K -finite and $|f_n(x^{(i)}) - y^{(i)}| < \varepsilon$ for all i . Here, the Flensted-Jensen functions φ_ℓ are given by the formula

$$\varphi_\ell(x) = (x_1 + \sqrt{-1}x_2)^{-\ell}, \quad x \in X(2, 2)_+.$$

1.2. Related work. In the 2000s, C. Fefferman proved a sharp form of the Whitney extension theorem [5]. In particular, if the set of points in Euclidean space is finite of cardinality n , he showed that there is an extension which is continuously differentiable of some degree depending on n . In this paper we start with a set of points of finite cardinality n , and using additional assumptions we prove the existence of a Whitney function which is infinitely often differentiable. The reason we are able to get an extension of such high regularity is that we take advantage of the fact that the points live in a subset of Euclidean space obeying a high amount of symmetries.

C. Fefferman et al. [7] have also studied Whitney extension problems in situations where the points live on a smooth manifold X . The Whitney extension theorems proved in that context, however, are existence theorems. In contrast, the results given in this article are effective. We give an algorithm computing the extension with an explicit formula. This is possible because we build the extension from functions in the discrete spectrum of the symmetric space G/H . Such Flensted-Jensen functions are solutions to a system of differential equations and hence satisfy a high amount symmetries.

Whitney extension theorems have found a range of applications in data science [6]. There is a growing literature on equivariant machine learning, where the data are invariant with respect to a group of symmetries [2]. Most results in this direction, however, are only available when the symmetry group of the points is either finite or a *compact* Lie group [2], [17], [4]. However, many groups of importance in physics are noncompact, such as the Lorentz group $\mathrm{SO}(3, 1)^\circ$, the symmetry group $\mathrm{SO}(p, 1)^\circ$ of $X(p, 1)_\pm$, and the symmetry group $\mathrm{SO}(p, 2)^\circ$ of $X(p, 2)_\pm$. One motivation for this work is to generalize the results in harmonic analysis applicable to Lie group equivariant machine learning to include *noncompact* Lie groups.

1.3. Contents. The paper has four parts. In this section, we summarize the contents of each part.

1.3.1. *Data on a sphere.* In the first part, we prove Whitney extension theorems for points on a sphere S^2 in \mathbb{R}^3 , whose group of symmetries is the compact group $\text{SO}(3)$. The proof is instructive for the noncompact case, so we give it in details. To do so, we use spherical harmonics, which will be used in the later parts of the article as well. Indeed, spherical harmonics also play a role in the representation theory of non compact groups and semisimple symmetric spaces, since Flensted-Jensen functions can be written as products of a Jacobi polynomial, which is the restriction of a harmonic polynomial, with a hyperbolic function.

1.3.2. *Data in the plane.* In this part, we prove a Whitney extension theorem for functions on a set of points in the Euclidean 2-space \mathbb{R}^2 . In contrast to the results by H. Whitney and C. Fefferman, we do not have any limitations on how differentiable the extension can be: We can compute a Whitney extension which is analytic right away. Notice that in this case, we do not need the points to lie in a symmetric space; all we need is that they are contained in the complement of the unit disk, in order to make certain functions square integrable. We include these results here to illustrate our techniques.

The result we prove is as follows.

Theorem 1.4. *Let $x^{(i)}$, $1 \leq i \leq n$, be a set of points in the plane \mathbb{R}^2 . Assume $x^{(i)}$ is not contained in the unit disk, and consider a nonconstant complex-valued function $f(x^{(i)}) = y^{(i)}$. There exists an analytic, square integrable function on \mathbb{R}^2 with the closed unit disk removed which extends this function.*

We prove this theorem by observing that it falls out, automatically, of the proof of the Whitney extension theorem for the three-dimensional symmetric space $X(2, 2)_+$, given in the next part.

1.3.3. *Data on the symmetric space $X(2, 2)_+ = \text{SO}(2, 2)^\circ / \text{SO}(1, 2)^\circ$.* This part forms the core of the paper. We discuss the Whitney extension problem for the three-dimensional manifold $X(2, 2)_+ = \text{SO}(2, 2)^\circ / \text{SO}(1, 2)^\circ$. We are especially interested not just in proving the existence, but also to obtain a formula for this function which can be easily computed. More precisely, we consider the question:

Question 1.1. *Does there exist a smooth, square integrable function $f \in C^\infty(X(2, 2)_+)$ such that $f(x^{(i)}) = y^{(i)}$ for all $i \leq n$? Can we obtain a closed formula for f ?*

Note that the manifold is a fibration of Minkowski space over the circle, $\text{SO}(2)$. In particular, we have no well-defined global time direction. We make the following assumptions:

- (1) We consider only finitely many points $x^{(i)}$, $1 \leq i \leq n$, although the number may be large.
- (2) We assume that $x^{(i)} \in \mathbb{Q}^4$ for all $i \leq n$ and consider a nonconstant function f on these points with values $f(x^{(i)}) = y^{(i)}$ in \mathbb{Q} .
- (3) We assume that the points are in general position, in the sense that they are not contained in the orbit of a proper subgroup of the symmetry group of the ambient symmetric space (cf. 3.2).

Not only do we find a smooth, square integrable function extending the function on the points, but we find a constructive way to obtain a formula for this function using the representation theory of noncompact Lie groups and symmetric spaces. We show that

Theorem 1.5. *Under these assumptions, for all $\varepsilon > 0$, there exists an analytic, square integrable function $f \in C^\infty(X(2, 2)_+)$ such that*

$$|f(x^{(i)}) - y^{(i)}| < \varepsilon$$

for all $i \leq n$.

Under the above assumptions, we consider the representation of G generated by f_n in the representation space $L^2(X(2, 2)_+)$. We show that to each Whitney function, we can associate a reducible, unitary representation, Π_n , which we call the Whitney representation of the data.

We then consider the problem: If we enlarge the set of points from cardinality n to $n + 1$, can we choose a smooth, square integrable, K -finite function $f_{n+1} \in L^2(X(2, 2)_+)$ such that $f_{n+1}(x^{(i)}) = y^{(i)}$ for all i and such that Π_n is a subrepresentation of Π_{n+1} ? We give an analytic proof that we can always act on the points by the maximal compact subgroup $\mathrm{SO}(2) \times \mathrm{SO}(2)$ to arrange that all irreducible subrepresentations below some fixed degree will occur in the Whitney representation.

1.3.4. *Data on symmetric spaces $X(p, 2)_- = \mathrm{SO}(p, 2)^\circ / \mathrm{SO}(p, 1)^\circ$.* We show that our results extend to the symmetric spaces

$$X(p, 2)_- = \mathrm{SO}(p, 2)^\circ / \mathrm{SO}(p, 1)^\circ$$

for all $p \geq 2$. The essential feature is that the homogeneous space $X(p, 2)_-$ has a holomorphic structure and holomorphic discrete series representations.

1.4. Acknowledgements. These results were first presented in a lecture at the AIM virtual seminar “Representation Theory and Noncommutative Geometry”. We are grateful to Bent Ørsted for helpful conversations. We would like to thank David Vogan and Robert Yuncken for insightful questions and comments related to this work.

The first author was supported by grant 89086 from the Simons Foundation.

2. LOW DIMENSIONAL EXAMPLES

2.1. **The sphere S^2 .** In this subsection we consider a set of points

$$x^{(i)} \in S^2 = \{x \in \mathbb{R}^3 : x_1^2 + x_2^2 + x_3^2 = 1\}, \quad 1 \leq i \leq n,$$

and a function $f(x^{(i)}) = y^{(i)}$ ($1 \leq i \leq n$) which we assume is not constant. We think of this as an observable set of data with an inherent spherical symmetry. From this perspective, it is natural to assume

$$x^{(i)} \in \mathbb{Q}^3, \quad y^{(i)} \in \mathbb{Q}$$

for all i , and we will impose this assumption throughout this section. Our goal is to find an explicit function interpolating the points. Since the symmetry group of a sphere is $\mathrm{SO}(3)$, it is natural to solve this problem using the representation theory of $\mathrm{SO}(3)$.

2.2. Preliminaries. There is a natural action of $\mathrm{SO}(3)$ on S^2 . We endow it with the surface area measure, μ , which is invariant under the action of $\mathrm{SO}(3)$, and normalized such that $\mu(S^2) = 1$. We consider the Hilbert space $L^2(S^2, \mu)$ of square integrable functions with respect to μ . Then $\mathrm{SO}(3)$ acts on $L^2(S^2)$ as

$$(g \cdot \psi)(x) = \psi(g^{-1}x), \quad g \in \mathrm{SO}(3), \psi \in L^2(S^2), \quad x \in S^2.$$

Let $\Delta = \sum_{i=1}^3 \partial^2 / \partial x_i^2$ be the Laplacian on \mathbb{R}^3 . Recall that a harmonic polynomial on \mathbb{R}^3 is a polynomial $p = p(x_1, x_2, x_3) \in \mathbb{C}[x_1, x_2, x_3]$ in three variables with complex coefficients satisfying

$$\Delta p = 0.$$

Example: Let $\ell \in \mathbb{Z}_{\geq 0}$. The polynomial

$$p(x_1, x_2, x_3) = (x_1 + \sqrt{-1}x_2)^\ell$$

is a harmonic of degree ℓ . Note that p is independent of x_3 and is holomorphic as a function of the complex variable $z = x_1 + \sqrt{-1}x_2$. By [1, 4.3], it follows that p is a harmonic function of (x_1, x_2) in the sense that

$$\left(\frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} \right) p = 0.$$

In particular, $\Delta p = 0$. The space of spherical harmonics of degree ℓ ($\ell \in \mathbb{Z}_{\geq 0}$) is the subspace V_ℓ of $L^2(S^2)$ consisting of restrictions to S^2 of harmonic polynomials on \mathbb{R}^3 that are homogeneous of degree ℓ . Recall that the space V_ℓ of spherical harmonics of degree ℓ is an irreducible representation of $\mathrm{SO}(3)$ of degree $2\ell + 1$. If V is any irreducible finite-dimensional representation of $\mathrm{SO}(3)$, then there is a unique integer ℓ such that V is isomorphic to V_ℓ . The space $L^2(S^2)$ decomposes as a Hilbert space direct sum

$$L^2(S^2) \simeq \hat{\bigoplus}_{\ell \in \mathbb{Z}_{\geq 0}} V_\ell.$$

In the physics literature [9, Definition 17.5], ℓ is called the spin of the representation V . In Lie theoretic language, the irreducible representations of $\mathrm{SO}(3)$ are classified by their highest weight [10], and ℓ is the highest weight of V .

2.3. The first interpolation theorem.

Theorem 2.1. *Consider a set of points $x^{(i)} \in S^2 \cap \mathbb{Q}^3$, $1 \leq i \leq n$, and a function f on the points with values $f(x^{(i)}) = y^{(i)} \in \mathbb{Q}$. Suppose $x^{(i)} \neq (0, 0, 1)$ and $(x_1^{(j)}, x_2^{(j)}) \neq (x_1^{(i)}, x_2^{(i)})$ whenever $i \neq j$. There are a unique coefficients $a_1, \dots, a_\ell \in \mathbb{Q}(\sqrt{-1})$ such that the harmonic polynomial*

$$f(x_1, x_2, x_3) = \sum_{\ell=1}^n a_\ell (x_1 + \sqrt{-1}x_2)^\ell$$

satisfies $f(x^{(i)}) = y^{(i)}$ for all i .

Proof. Let $p_\ell(x) = (x_1 + \sqrt{-1}x_2)^\ell$ be the spherical harmonic of degree ℓ . The $n \times n$ matrix $(p_\ell(x^{(i)}))_{i,\ell}$ is the product of the diagonal matrix

$$\begin{pmatrix} (x_1^{(1)} + \sqrt{-1}x_2^{(1)}) & & & \\ & \ddots & & \\ & & & (x_1^{(n)} + \sqrt{-1}x_2^{(n)}) \end{pmatrix}$$

with the Vandermonde matrix

$$\begin{pmatrix} 1 & (x_1^{(1)} + \sqrt{-1}x_2^{(1)}) & (x_1^{(1)} + \sqrt{-1}x_2^{(1)})^2 & \cdots & (x_1^{(1)} + \sqrt{-1}x_2^{(1)})^{n-1} \\ 1 & (x_1^{(2)} + \sqrt{-1}x_2^{(2)}) & (x_1^{(2)} + \sqrt{-1}x_2^{(2)})^2 & \cdots & (x_1^{(2)} + \sqrt{-1}x_2^{(2)})^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & (x_1^{(n)} + \sqrt{-1}x_2^{(n)}) & (x_1^{(n)} + \sqrt{-1}x_2^{(n)})^2 & \cdots & (x_1^{(n)} + \sqrt{-1}x_2^{(n)})^{n-1} \end{pmatrix}$$

By the formula for a Vandermonde determinant (which holds over any commutative ring, cf. [19, p. 42]),

$$\det(p_\ell(x^{(i)}))_{i,\ell} = \prod_{i=1}^n (x_1^{(i)} + \sqrt{-1}x_2^{(i)}) \prod_{1 \leq i < j \leq n} \left((x_1^{(j)} + \sqrt{-1}x_2^{(j)}) - (x_1^{(i)} + \sqrt{-1}x_2^{(i)}) \right)$$

Since $x^{(i)} \neq (0, 0, 1)$ and $(x_1^{(j)}, x_2^{(j)}) \neq (x_1^{(i)}, x_2^{(i)})$ for all $i \neq j$, this determinant is nonzero. In particular, the system

$$\begin{pmatrix} y^{(1)} \\ \vdots \\ y^{(n)} \end{pmatrix} = \begin{pmatrix} p_1(x^{(1)}) & p_2(x^{(1)}) & \cdots & p_n(x^{(1)}) \\ \vdots & \vdots & \ddots & \vdots \\ p_1(x^{(n)}) & p_2(x^{(n)}) & \cdots & p_n(x^{(n)}) \end{pmatrix} \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix}$$

has a unique solution in $a_\ell \in \mathbb{Q}(\sqrt{-1})$. \square

2.4. Harmonic polynomials on \mathbb{R}^3 . Let \mathcal{P}_ℓ be the complex vector space of homogeneous polynomials on \mathbb{R}^3 with complex coefficients. Recall that the restriction map

$$\mathcal{P}_\ell \longrightarrow L^2(S^2)$$

is injective [14], so we may identify V_ℓ with a space of functions on \mathbb{R}^3 . We endow \mathbb{R}^3 with the product measure

$$\mu \otimes 4\pi r^2 dr$$

of the surface area measure μ on S^2 with the measure $4\pi r^2 dr$ on $(0, \infty)$. The measure $\mu \otimes 4\pi r^2 dr$ is invariant with respect to the natural action of $\text{SO}(3)$ on \mathbb{R}^3 , and we define $L^2(\mathbb{R}^3)$ with respect to this measure. Then $L^2(\mathbb{R}^3)$ is a unitary representation of $\text{SO}(3)$.

Lemma 2.1. *There is a canonical isomorphism of $L^2(\mathbb{R}^3)$ with the Hilbert tensor product*

$$L^2(\mathbb{R}^3) \simeq L^2(S^2) \hat{\otimes} L^2((0, \infty), 4\pi r^2 dr).$$

Under this isomorphism, a vector $f \otimes g$ in $L^2(S^2) \hat{\otimes} L^2(0, \infty)$ corresponds to the function $f(\theta, \phi)g(r)$ in $L^2(\mathbb{R}^3)$ written in polar coordinates.

Lemma 2.2. *Let $\ell \in \mathbb{Z}_{\geq 0}$, and let f be a measurable function on $(0, \infty)$ such that $f(r)r^\ell \in L^2(0, \infty)$, i.e.*

$$\int_{(0, \infty)} |f(r)|^2 r^{2\ell} 4\pi r^2 dr < \infty.$$

If $p \in V_\ell$, then $p(x)f(|x|) \in L^2(\mathbb{R}^3)$. Define $V_{\ell, f}$ to be the subspace of $L^2(\mathbb{R}^3)$ of functions of the form

$$\varphi(x) = p(x)f(\|x\|), \quad p \in V_\ell.$$

Then $V_{\ell, f}$ is invariant and irreducible under the action of $\text{SO}(3)$.

Proof. Note that if $\varphi \in V_{\ell, f}$, then under the isomorphism in Lemma 2.1,

$$\varphi(x) = p(x/\|x\|)\|x\|^\ell f(\|x\|) \simeq p \otimes g,$$

where $g(r) = r^\ell f(r)$ and $p \in V_\ell$. \square

Lemma 2.3. *Let V be any irreducible representation of $\mathrm{SO}(3)$ in $L^2(\mathbb{R}^3)$. Then V is of the form $V_{\ell, f}$ for some ℓ and f satisfying the condition in Lemma 2.2.*

Proof. See [9, Prop. 17.19]. \square

2.5. An open subset $\mathbb{R}^2 \setminus \overline{B}^1$ of the plane. Let $B^1 = \{x \in \mathbb{R}^2 : x_1^2 + x_2^2 < 1\}$ be an open ball of radius 1, and let $\overline{B}^1 = \{x \in \mathbb{R}^2 : x_1^2 + x_2^2 \leq 1\}$ be its closure. In this subsection we consider the domain $\mathbb{R}^2 \setminus \overline{B}^1$ and a nonconstant function on the set of points $x^{(i)} \in \mathbb{R}^2 \setminus \overline{B}^1$ for $i \leq n$. Assume $x^{(i)} \in \mathbb{Q}^2$ and that the function maps any $x^{(i)}$ to a rational point $y^{(i)} \in \mathbb{Q}$. We equip $\mathbb{R}^2 \setminus \overline{B}^1$ with the measure

$$\mu = d\theta \otimes r \, dr,$$

which is Lebesgue measure in polar coordinates.

Lemma 2.4. *For any integer $\ell \geq 2$, the functions*

$$\varphi_\ell^\pm(x) = (x_1 \pm \sqrt{-1}x_2)^{-\ell}, \quad x \in \mathbb{R}^2 \setminus \overline{B}^1$$

are in $L^2(\mathbb{R}^2 \setminus \overline{B}^1, d\theta \otimes r \, dr)$.

Proof.

$$\begin{aligned} \int_{\mathbb{R}^2 \setminus \overline{B}^1} |\varphi_\ell^\pm(x)|^2 d\theta \otimes r \, dr &= \int_{-\pi}^{\pi} \int_1^{\infty} |r \cos \theta \pm \sqrt{-1}r \sin \theta|^{-2\ell} d\theta \otimes r \, dr \\ &= 2\pi \int_1^{\infty} r^{-2\ell} r \, dr \end{aligned}$$

which is finite if and only if $2\ell - 1 > 1 \iff \ell \geq 2$. \square

Lemma 2.5. *For any $\ell \geq 2$, let*

$$\psi_\ell^+(x) = (x_1 + \sqrt{-1}x_2)^{-\ell}, \quad \psi_\ell^-(x) = (x_1 - \sqrt{-1}x_2)^{-\ell}, \quad x = (x_1, x_2) \in \mathbb{R}^2 \setminus \{(0, 0)\},$$

and let $x^{(i)}$ ($1 \leq i \leq n$) be a sequence of points with $x^{(i)} = (x_1^{(i)}, x_2^{(i)}) \in \mathbb{Q}^2 \setminus \{(0, 0)\}$ for all $i \leq n$. Up to multiplication by a diagonal matrix, $(\psi_\ell^+(x^{(i)}))_{i, \ell}$ and $(\psi_\ell^-(x^{(i)}))_{i, \ell}$ where $1 \leq i \leq n$ and $2 \leq \ell \leq n + 1$, are Vandermonde matrices. In particular, $(\psi_\ell^+(x^{(i)}))_{i, \ell}$ is invertible if and only if the points $x^{(i)} = (x_1^{(i)}, x_2^{(i)})$ have the properties

$$(x_1^{(i)}, x_2^{(i)}) \neq (x_1^{(j)}, x_2^{(j)}), \quad \text{whenever } i < j, \quad i, j \leq n.$$

Proof. Define

$$\alpha_i = (x_1^{(i)} + \sqrt{-1}x_2^{(i)})^{-1} \in \mathbb{Q}(\sqrt{-1}), \quad 1 \leq i \leq n,$$

then

$$(\psi_\ell^+(x^{(i)}))_{i, \ell} = \begin{pmatrix} \alpha_1^2 & & & \\ & \ddots & & \\ & & \alpha_n^2 & \\ & & & \alpha_n^2 \end{pmatrix} \begin{pmatrix} 1 & \alpha_1 & \alpha_1^2 & \cdots & \alpha_1^{n-1} \\ 1 & \alpha_2 & \alpha_2^2 & \cdots & \alpha_2^{n-1} \\ \vdots & & & & \\ 1 & \alpha_n & \alpha_n^2 & \cdots & \alpha_n^{n-1} \end{pmatrix}.$$

By the formula for a Vandermonde determinant [19, p. 42],

$$\det(\psi_\ell^+(x^{(i)}))_{i,\ell} = \prod_{i=1}^n \alpha_i^2 \prod_{1 \leq i < j \leq n} (\alpha_j - \alpha_i)$$

□

Lemma 2.6. *For any $\ell \geq 2$, let*

$$\psi_\ell^\pm(x) = (x_1 \pm \sqrt{-1}x_2)^{-\ell}, \quad x = (x_1, x_2) \in \mathbb{R}^2 \setminus \overline{B}^1.$$

Let $x^{(i)} = (x_1^{(i)}, x_2^{(i)}) \in \mathbb{Q}^2 \setminus \{(0, 0)\}$ be a sequence of points on which a function f is defined such that $f(x^{(i)}) = y^{(i)} \in \mathbb{Q}$ for all $i \leq n$. Suppose that

$$x^{(i)} \neq x^{(j)}$$

whenever $i < j$. There exist unique coefficients $a_\ell \in \mathbb{Q}(\sqrt{-1})$ such that if we define the function

$$f_n = \sum_{l=2}^{n+1} a_l \psi_l^+ \in L^2(\mathbb{R}^2 \setminus \overline{B}^1, d\theta \otimes r dr)$$

then $f_n(x^{(i)}) = y^{(i)}$ for all $i \leq n$.

Proof. By Lemma 2.5, the column vector $(a_\ell)_\ell$ can be computed in terms of the points $(x^{(i)}, y^{(i)})$ as

$$(a_\ell)_\ell = \left(\psi_\ell^+(x^{(i)}) \right)_{i,\ell}^{-1} (y^{(i)})_i.$$

□

3. THE SYMMETRIC SPACE $X(2, 2)_+$

3.1. Preliminaries. Let $p, q \geq 1$ be integers. Let $(\cdot, \cdot)_{p,q}$ be the symmetric bilinear form on \mathbb{R}^{p+q} of signature (p, q) defined by

$$(x, y)_{p,q} = \sum_{k=1}^p x_k y_k - \sum_{k=p+1}^{p+q} x_k y_k, \quad x, y \in \mathbb{R}^{p+q}.$$

The generalized orthogonal group of signature (p, q) is the group

$$\mathrm{O}(p, q) = \{g \in \mathrm{GL}(p+q, \mathbb{R}) : (gx, gy)_{p,q} = (x, y)_{p,q} \forall x, y \in \mathbb{R}^{p+q}\},$$

and $\mathrm{SO}(p, q) = \{g \in \mathrm{O}(p, q) : \det g = 1\}$. Note that if $g \in \mathrm{GL}(p+q, \mathbb{R})$, then $g \in \mathrm{O}(p, q)$ if and only if

$$\begin{aligned} (ge_j, ge_k)_{p,q} &= 0, & j &\neq k, \\ (ge_j, ge_j)_{p,q} &= 1, & j &\leq p, \\ (ge_j, ge_j)_{p,q} &= -1, & p+1 &\leq j \leq p+q. \end{aligned}$$

Lemma 3.1. *Let*

$$Q(x) = x_1^2 + x_2^2 - x_3^2 - x_4^2, \quad x \in \mathbb{R}^4.$$

Then $X = \{x \in \mathbb{R}^4 : Q(x) = 1\}$ is a manifold endowed with a transitive Lie group action by $G = \mathrm{SO}(2, 2)$, and there is a G -equivariant diffeomorphism

$$X \simeq \mathrm{SO}(2, 2) / \mathrm{SO}(1, 2).$$

Proof. There is a smooth, injective group homomorphism $\varphi : \mathrm{SO}(1, 2) \longrightarrow \mathrm{SO}(2, 2)$ of the form

$$\varphi : g \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & g_{11} & g_{12} & g_{13} \\ 0 & g_{21} & g_{22} & g_{23} \\ 0 & g_{31} & g_{32} & g_{33} \end{pmatrix}, \quad g \in \mathrm{SO}(1, 2)$$

since for $x, y \in \mathbb{R}^4$ and $g \in \mathrm{SO}(1, 2)$,

$$\begin{aligned} \left(\begin{pmatrix} 1 & 0 \\ 0 & g \end{pmatrix} x, \begin{pmatrix} 1 & 0 \\ 0 & g \end{pmatrix} y \right)_{2,2} &= x_1 y_1 + \left(g \begin{pmatrix} x_2 \\ x_3 \\ x_4 \end{pmatrix}, g \begin{pmatrix} y_2 \\ y_3 \\ y_4 \end{pmatrix} \right)_{1,2} \\ &= x_1 y_1 + \left(\begin{pmatrix} x_2 \\ x_3 \\ x_4 \end{pmatrix}, \begin{pmatrix} y_2 \\ y_3 \\ y_4 \end{pmatrix} \right)_{1,2} = (x, y)_{2,2}. \end{aligned}$$

Let $G = \mathrm{SO}(2, 2)$. Let $e_1 = (1, 0, 0, 0) \in \mathbb{R}^4$. The stabilizer in G at e_1 is

$$G_{e_1} = \{g \in G : ge_1 = e_1\} = \left\{ g \in G : g = \begin{pmatrix} 1 & 0 \\ 0 & \tilde{g} \end{pmatrix} \right\} = \mathrm{SO}(1, 2).$$

By definition, $Q(x) = (x, x)_{2,2}$, and the G -orbit of e_1 is

$$Ge_1 = \{x \in \mathbb{R}^4 : (x, x)_{2,2} = 1\} = \{x \in \mathbb{R}^4 : Q(x) = 1\}.$$

In particular, the orbit Ge_1 is a manifold, and there is a diffeomorphism

$$\mathrm{SO}(2, 2)/\mathrm{SO}(1, 2) \longrightarrow Ge_1 = \{x \in \mathbb{R}^4 : Q(x) = 1\}.$$

□

We follow the conventions in the literature and work with the connected components, so we consider

$$X(2, 2)_+ = \mathrm{SO}(2, 2)^\circ / \mathrm{SO}(1, 2)^\circ$$

respectively

$$X(2, 2)_- = \mathrm{SO}(2, 2)^\circ / \mathrm{SO}(2, 1)^\circ.$$

Lemma 3.2. *Let $\mathfrak{so}(2, 2)$ be the Lie algebra of $\mathrm{SO}(2, 2)$, and let $\mathfrak{so}(2, 2; \mathbb{C})$ be its complexification. There is an isomorphism of Lie algebras*

$$\varphi : \mathfrak{so}(4; \mathbb{C}) \longrightarrow \mathfrak{so}(2, 2; \mathbb{C})$$

In particular, if $\mathfrak{g}_1 \simeq \mathfrak{sl}(2, \mathbb{C})$, then

$$\mathfrak{so}(2, 2; \mathbb{C}) \simeq \mathfrak{g}_1 \oplus \overline{\mathfrak{g}_1}$$

and

$$\mathfrak{so}(2, 2) \simeq \mathfrak{sl}(2; \mathbb{C})_{\mathbb{R}}.$$

Hence $\mathfrak{so}(2, 2)$ is simple as a real Lie algebra, while $\mathfrak{so}(2, 2; \mathbb{C})$ is not simple (but semisimple.)

Proof. Let

$$g = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix}.$$

Let $M_4(\mathbb{R})$ be the space of 4×4 real matrices, and for any matrix X , let tX be its transpose. We have $g = g^{-1}$ and

$$\mathfrak{so}(2, 2) \simeq \{X \in M_2(\mathbb{R}) : g({}^tX)g^{-1} = -X\}.$$

Note that g has a complex square root:

$$g^{1/2} = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & i & \\ & & & i \end{pmatrix} \in \mathrm{GL}(4, \mathbb{C}).$$

In particular,

$${}^tX = -X \iff g \left({}^t(g^{1/2}Xg^{-1/2}) \right) g^{-1} = -g^{1/2}Xg^{-1/2}.$$

It follows that there is a Lie algebra isomorphism

$$\mathrm{Ad}_{g^{1/2}} : \mathfrak{so}(4; \mathbb{C}) \longrightarrow \mathfrak{so}(2, 2; \mathbb{C})$$

By [10, Section 7.7.2 and 7.8 Exercise 4],

$$\mathfrak{so}(4; \mathbb{C}) \simeq \mathfrak{sl}(2; \mathbb{C}) \oplus \mathfrak{sl}(2; \mathbb{C}).$$

As in the proof of [10, Theorem 7.32], these two copies of $\mathfrak{sl}(2; \mathbb{C})$ are complex conjugate. If \mathfrak{g}_1 denotes the first copy in the decomposition, then $\mathfrak{so}(2, 2; \mathbb{C}) = \mathfrak{g}_1 \oplus \overline{\mathfrak{g}_1}$. The map $\varphi : \mathfrak{g}_1 \rightarrow \mathfrak{so}(2, 2; \mathbb{C})$

$$\varphi : X \mapsto X + \overline{X}$$

is an \mathbb{R} -linear isomorphism and respects Lie brackets. We conclude that

$$\mathfrak{so}(2, 2) \simeq \mathfrak{sl}(2; \mathbb{C})_{\mathbb{R}}.$$

□

3.2. From rational data to Whitney functions. The first case. The functions constructed by M. Flensted-Jensen [8] will in our considerations play the role for the noncompact group $\mathrm{SO}(2, 2)^\circ$ analogous to the role played by spherical harmonics for the compact group $\mathrm{SO}(3)$.


Proposition 3.1. *Let $X = \{x \in \mathbb{R}^4 : x_1^2 + x_2^2 - x_3^2 - x_4^2 = 1\}$. For any integer $\ell \geq 2$, define $\psi_\ell^+ : X \rightarrow \mathbb{C}$ and $\psi_\ell^- : X \rightarrow \mathbb{C}$ by*

$$\psi_\ell^+(x) = (x_1 + \sqrt{-1}x_2)^{-\ell}, \quad \psi_\ell^-(x) = (x_1 - \sqrt{-1}x_2)^{-\ell}.$$


Then ψ_ℓ^\pm are Flensted-Jensen functions for the parameter $\lambda = 2\ell - 2$, and the (\mathfrak{g}, K) -modules generated by ψ_ℓ^+ and ψ_ℓ^- form a complete set of discrete series representations for X .

Proof. Cf. [12] Part 4, section 9, p. 215. □

We say that the points $x^{(i)}$ ($1 \leq i \leq n$) are *in general position* if not all points in the sequence have the same coordinate equal to zero, i.e. not all points $x^{(i)} \in G/H$ have representatives in $G = \mathrm{SO}(2, 2)^\circ$ contained in the same subgroup G' isomorphic to one of the groups $\mathrm{SO}(1, 2)^\circ$, $\mathrm{SO}(2, 1)^\circ$, $\mathrm{SO}(1, 1)^\circ$, $\mathrm{SO}(2, 0)$, or $\mathrm{SO}(0, 2)$. In this section we assume that the points $x^{(i)}$ are in general position.



Data on orbits in a symmetric space



Data in general position on a symmetric space

Let $x^{(i)}$ ($1 \leq i \leq n$) be a sequence of points of $X = G/H$ in general position, and assume $x^{(i)} \in \mathbb{Q}^4$ for all i . Let f be a nonconstant function on the points with $f(x^{(i)}) = y^{(i)} \in \mathbb{Q}$. We would like to find coefficients a_ℓ such that

$$y^{(i)} = \sum_{\ell \geq 2} a_\ell \psi_\ell^\pm(x^{(i)})$$

for all $i \leq n$, and where $a_\ell \neq 0$ for finitely many $\ell \geq 2$. Then

$$f = \sum_{\ell \geq 2} a_\ell \psi_\ell^\pm \in L^2(X)$$

is a K -finite vector in the representation space $L^2(X)$ with the property that $f(x^{(i)}) = y^{(i)}$ for all $i \leq n$. We find (a_ℓ) by solving a linear system of equations

$$\begin{pmatrix} y^{(1)} \\ \vdots \\ y^{(n)} \end{pmatrix} = \begin{pmatrix} \psi_2^\pm(x^{(1)}) & \psi_3^\pm(x^{(1)}) & \cdots & \psi_m^\pm(x^{(1)}) \\ \vdots \\ \psi_2^\pm(x^{(n)}) & \psi_3^\pm(x^{(n)}) & \cdots & \psi_m^\pm(x^{(n)}) \end{pmatrix} \begin{pmatrix} a_1 \\ \vdots \\ a_m \end{pmatrix}$$

for a choice of $m \geq n$.

Proposition 3.2. *The matrix $(\psi_\ell^+(x^{(i)}))_{i,\ell}$ where $1 \leq i \leq n$, $2 \leq \ell \leq n+1$ is Vandermonde (up to multiplication by a diagonal matrix). Furthermore, $(\psi_\ell^+(x^{(i)}))_{i,\ell}$ is invertible if and only if the points $x^{(i)} = (x_1^{(i)}, x_2^{(i)}, x_3^{(i)}, x_4^{(i)})$ in X have the property*

$$(x_1^{(i)}, x_2^{(i)}) \neq (x_1^{(j)}, x_2^{(j)}), \quad \text{whenever } i < j, \quad i, j \leq n.$$

In this case, $a_\ell \in \mathbb{Q}(\sqrt{-1})$ for all $2 \leq \ell \leq n+1$, and the column vector $(a_\ell)_\ell$ can be computed in terms of the points $(x^{(i)}, y^{(i)})$ as

$$(a_\ell)_\ell = \left(\psi_\ell^+(x^{(i)}) \right)_{i,\ell}^{-1} (y^{(i)})_i.$$

Proof. Follows from Lemma 2.5. \square

3.3. Cayley's rational parametrization of $\mathrm{SO}(n)$. A matrix A is called non-exceptional if $\det(I+A) \neq 0$, where I is the identity matrix, and otherwise A is called exceptional.

Lemma 3.3. *Let A be a non-exceptional matrix. Then the matrix S defined by the equation*

$$I+S = 2(I+A)^{-1}$$

is also non-exceptional, $(I+A) = 2(I+S)^{-1}$, and A and S are mutually related by the equations

$$S = (I-A)(I+A)^{-1} = (I+A)^{-1}(I-A), \quad A = (I-S)(I+S)^{-1} = (I+S)^{-1}(I-S).$$

Since the factors commute, we will follow Weyl [28] and use the rational notation

$$A = \frac{I-S}{I+S}.$$

For a matrix $R = (r_{ij})$, the bilinear form $\sum_{ij} r_{ij} x_i y_j$, is A -invariant if and only if

$${}^t A R A = R,$$

where ${}^t A$ is the transpose of A .

Lemma 3.4. *Let A and S be a pair of non-exceptional matrices related as in Lemma 3.3, and let R be any matrix. Then ${}^t A R A = R$ if and only if*

$${}^t S R + R S = 0.$$

In particular, if $R = I$ and A belongs to the orthogonal group $\mathrm{O}(n)$, then $A \in \mathrm{SO}(n)$ and S is skew-symmetric: ${}^t S + S = 0$. Conversely, if S is skew-symmetric matrix, then $A \in \mathrm{SO}(n)$.

Lemma 3.5. *Let $A \in \mathrm{SO}(n)$. Then $A = A_1 A_2$ for two non-exceptional matrices*

$$A_1 = \frac{I-S_1}{I+S_1}, \quad A_2 = \frac{I-S_2}{I+S_2} \in \mathrm{SO}(n).$$

If $\det(I+S_1 S_2) \neq 0$, then A is itself non-exceptional and

$$A = \frac{I-S}{I+S}$$

where

$$S = \frac{S_1 + S_2}{I + S_1 S_1}.$$

The following theorem implies that there are plenty of rational points in the space X .

Proposition 3.3. *Let*

$$f = a_1x_1^2 + a_2x_2^2 + a_3x_3^2 + a_4x_4^2 + a_5x_5^2, \quad a_i \in \mathbb{Q}$$

be a quadratic form over \mathbb{Q} of rank 5. Then f represents zero in \mathbb{Q} if and only if it represents zero in \mathbb{R} . In particular, there are coordinates $x_i \in \mathbb{Q}$ such that

$$0 = x_1^2 + x_2^2 - x_3^2 - x_4^2 - x_5^2,$$

$x_5 \neq 0$, and the point $x' = (x_1/x_5, x_2/x_5, x_3/x_5, x_4/x_5) \in X(2, 2)_+ \cap \mathbb{Q}^4$, where

$$X(2, 2)_+ = \{x \in \mathbb{R}^4 : x_1^2 + x_2^2 - x_3^2 - x_4^2 = 1\}.$$

For any proper orthogonal matrix $A \in \text{SO}(2, \mathbb{Q})$, the point

$$\begin{pmatrix} A & \\ & I \end{pmatrix} x' \in X(2, 2)_+ \cap \mathbb{Q}^4.$$

In particular, there are infinitely many rational points on $X(2, 2)_+$.

Proof. See [24] p. 43. □

Lemma 3.6. *Let $x = (x_1, x_2, x_3, x_4)$ and $x' = (x'_1, x'_2, x'_3, x'_4)$ be rational points in $X(2, 2)_+$. Suppose*

$$x_1 = x'_1, \quad x_2 = x'_2.$$

For any matrix $A \in \text{SO}(2, \mathbb{Q})$, let

$$\begin{pmatrix} x_1^\circ \\ x_2^\circ \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x'_1 \\ x'_2 \end{pmatrix} = \begin{pmatrix} ax'_1 + bx'_2 \\ cx'_1 + dx'_2 \end{pmatrix}$$

and let $x_3^\circ = x'_3$ and $x_4^\circ = x'_4$. Then the perturbed point $x^\circ = (x_1^\circ, x_2^\circ, x'_3, x'_4)$ is a rational point on $X(2, 2)_+$.

Proof. By assumption,

$$(x'_1)^2 + (x'_2)^2 = 1 + (x'_3)^2 + (x'_4)^2,$$

so (x'_1, x'_2) lies on a circle of radius $1 + (x'_3)^2 + (x'_4)^2$, and $\text{SO}(2)$ acts on this circle. □

Lemma 3.7. *Let $\varepsilon \in \mathbb{Q}, \varepsilon > 0$, and define the skew-symmetric matrix*

$$S = \begin{pmatrix} 0 & \varepsilon \\ -\varepsilon & 0 \end{pmatrix}.$$

Then $\det(I + S) = 1 + \varepsilon^2 > 0$, so S is non-exceptional, and

$$A = \frac{I - S}{I + S} = \begin{pmatrix} 1 & \varepsilon \\ -\varepsilon & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 & -\varepsilon \\ \varepsilon & 1 \end{pmatrix} = \frac{1}{1 + \varepsilon^2} \begin{pmatrix} 1 - \varepsilon^2 & -2\varepsilon \\ 2\varepsilon & 1 - \varepsilon^2 \end{pmatrix}$$

is a non-exceptional matrix in $\text{SO}(2, \mathbb{Q})$. Also, the distance between A and the identity transformation I is

$$\|A - I\|^2 = \frac{8\varepsilon^2}{1 + \varepsilon^2},$$

which tends to 0 as ε tends to zero.

Lemma 3.8. *Let x and \tilde{x} be rational points in $X(2, 2)_+$ whose first and second coordinates coincide: $x_1 = \tilde{x}_1$ and $x_2 = \tilde{x}_2$. Then the point*

$$\tilde{x}^{(\varepsilon)} = \begin{pmatrix} \frac{1-\varepsilon^2}{1+\varepsilon^2}x_1 - \frac{2\varepsilon}{1+\varepsilon^2}x_2 \\ \frac{2\varepsilon}{1+\varepsilon^2}x_1 + \frac{1-\varepsilon^2}{1+\varepsilon^2}x_2 \\ \tilde{x}_3 \\ \tilde{x}_4 \end{pmatrix}$$

is a rational point on $X(2, 2)_+$. Clearly, the points \tilde{x} and $\tilde{x}^{(\varepsilon)}$ are equidistant from x , i.e.

$$\|x - \tilde{x}\|_{\mathbb{R}^4} = \|x - \tilde{x}^{(\varepsilon)}\|_{\mathbb{R}^4},$$

while the distance between \tilde{x} and the perturbed point $\tilde{x}^{(\varepsilon)}$ is given by

$$\|\tilde{x} - \tilde{x}^{(\varepsilon)}\|_{\mathbb{R}^4}^2 = \frac{4\varepsilon^2}{(1+\varepsilon^2)^2} \left((x_2 - \varepsilon x_1)^2 + (x_1 - \varepsilon x_2)^2 \right),$$

which tends to zero as ε tends to zero: Indeed, if $\varepsilon \leq 1$, and if $C = \max_{i,j} |x_j^{(i)}|$ then $\|\tilde{x} - \tilde{x}^{(\varepsilon)}\|_{\mathbb{R}^4} \leq 2^{5/2}C\varepsilon$. Finally,

$$x_1 \neq \tilde{x}_1^{(\varepsilon)}, \quad x_2 \neq \tilde{x}_2^{(\varepsilon)}.$$

3.4. From rational data sets to Whitney functions: the general case.

Consider a set of points $x^{(i)}$ ($1 \leq i \leq n$) and a nonconstant function $f(x^{(i)}) = y^{(i)}$ where $x^{(i)} \in X(2, 2)_+$, $x_j^{(i)} \in \mathbb{Q}$ for all $j \leq 4$, and $y^{(i)} \in \mathbb{Q}$ for all $i \leq n$.

- (1) Define a small error tolerance $\varepsilon \in \mathbb{Q}$, $\varepsilon > 0$.
- (2) Loop through indices i with $2 \leq i \leq n$. Let such an i be fixed. By induction, it holds for all $j, k < i$ ($j \neq k$) that

$$(x_1^{(j)}, x_2^{(j)}) \neq (x_1^{(k)}, x_2^{(k)}).$$

Suppose there is some $k < i$ such that

$$(x_1^{(k)}, x_2^{(k)}) = (x_1^{(i)}, x_2^{(i)}).$$

In particular, $k < i$ is the only index with this property. Apply a non-exceptional proper orthogonal transformation to the point $x^{(i)}$ to obtain the point $x^{(i,\varepsilon)}$ for $\varepsilon \in \mathbb{Q}_{>0}$ as in Lemma 3.8 with the property that $(x_1^{(i,\varepsilon)}, x_2^{(i,\varepsilon)})$ is *distinct from all the points in the finite list*

$$(x_1^{(j)}, x_2^{(j)}), \quad j < i.$$

Replace the point $x^{(i)}$ with the point $x^{(i,\varepsilon)}$.

- (3) After having checked all points and replaced some with small perturbations if necessary, all the assumptions of Proposition 3.2 are in place. Compute the column vector

$$(a_\ell)_{\ell=2}^{n+1} = \left((x_1^{(i,\varepsilon)} + \sqrt{-1}x_2^{(i,\varepsilon)})^{-\ell} \right)_{1 \leq i \leq n, 2 \leq \ell \leq n+1}^{-1} (y^{(i)})_{i=1}^n,$$

where $a_\ell \in \mathbb{Q}(\sqrt{-1})$ for all ℓ . Then the function

$$f_n = \sum_{\ell=2}^{n+1} a_\ell \psi_\ell^+$$

is a linear combination of n linearly independent Flensted-Jensen functions with the property that

$$y_i = f_n(x^{(i,\varepsilon)})$$

for all $i \leq n$.

Let $(x^{(i)})$ be a data set of n points in G/H in general position and let $f(x^{(i)}) = y^{(i)} (1 \leq i \leq n)$ be a nonconstant function. After fixing some $\varepsilon > 0$, we associate an analytic, square integrable, K -finite function f_n to the data according to the algorithm in this section 3.4. We will refer to f_n as a *Whitney function of the data*.

3.5. From Whitney functions to Whitney representations. Let $G = \mathrm{SO}(2, 2)^\circ$ and $H = \mathrm{SO}(1, 2)^\circ$ and let f_n be a Whitney function of a data set in G/H . Let V_{f_n} be the (\mathfrak{g}, K) -module in $L^2(G/H)$ generated by f_n . This is the vector space of K -finite functions in $L^2(G/H)$ obtained by acting on f_n with the universal enveloping algebra $U(\mathfrak{g}_\mathbb{C})$. Let Π_n be the representation of G on the completion of V_{f_n} in $L^2(G/H)$ and let (π_ℓ, V_ℓ) ($\ell \geq 2$) be the Flensted-Jensen representation of G in $L^2(G/H)$ generated by φ_ℓ^+ . The results of M. Flensted-Jensen imply in the special case of $X(2, 2)_+$ that if $\ell, \ell' \geq 2$ are distinct indices, then the representations π_ℓ and $\pi_{\ell'}$ are inequivalent representations of G . Evidently, for a Whitney function $f_n = \sum_{\ell \leq n+1} a_\ell \varphi_\ell^+$,

$$\Pi_n = \bigoplus_{\ell \leq n+1: a_\ell \neq 0} \pi_\ell.$$

Note that if $a_\ell = 0$ then the representation π_ℓ is not a direct summand.

Lemma 3.9. *Define a hyperplane H_k in \mathbb{C}^n as*

$$H_k = \{z \in \mathbb{C}^n : z_k = 0\}, \quad 1 \leq k \leq n.$$

The union $\cup_{k \leq n} H_k$ is a subset of \mathbb{C}^n of Lebesgue measure zero. Let $x^{(i)} (1 \leq i \leq n)$ be a data set in G/H in general position, and let $f(x^{(i)}) = y^{(i)}$ be a nonconstant function. Fix $\varepsilon > 0$ and choose modified points $\tilde{x}^{(i)} = x^{(i,\varepsilon)} = k_\varepsilon \cdot x^{(i)}$ such that the matrix

$$T = \left(\varphi_\ell^+(\tilde{x}^{(i)}) \right)_{i,\ell} \in \mathrm{GL}(n, \mathbb{C}).$$

Let T^{-1} be the inverse of T in $\mathrm{GL}(n, \mathbb{C})$. Suppose

$$(y^{(i)})_{i \leq n} \in \mathbb{C}^n \setminus T(\cup_{k \leq n} H_k).$$

Then the Whitney function $f_n = \sum a_\ell \varphi_\ell^+$ of the data has the property that $a_\ell \neq 0$ for all ℓ . In particular, the representation Π_n which it generates contains a maximal number of summands:

$$\Pi_n = \bigoplus_{2 \leq \ell \leq n+1} \pi_\ell.$$

Proof. If the vector $(y^{(i)})_i \in \mathbb{C}^n$ lives in the set $\mathbb{C}^n \setminus T \cup_{k \leq n} H_k$ (which has full Lebesgue measure in \mathbb{C}^n) then

$$(a_\ell)_\ell = T^{-1}(y^{(i)})_i \notin \cup_{k \leq n} H_k$$

which means that $a_\ell \neq 0$ for all ℓ . □

Proposition 3.4. *Let $x^{(i)} (1 \leq i \leq n)$ be a set of points in G/H in general position, and let $f(x^{(i)}) = y^{(i)}$ be a nonconstant function. Fix $\varepsilon > 0$ and suppose there is a Whitney function*

$$f_n = \sum_{2 \leq \ell \leq n+1} a_\ell \varphi_\ell^+$$

such that $|f_n(x^{(i)}) - y^{(i)}| < \varepsilon$ for all i . There exists $\delta \in (-1/m, 1/m)$ for some $m \in \mathbb{Z}_{\geq 1}$ and a Whitney function

$$f_n^{(\delta)} = \sum_{\ell} a_\ell^{(\delta)} \varphi_\ell^+$$

such that

$$|f_n^{(\delta)}(x^{(i)}) - y^{(i)}| < \varepsilon \quad \forall i \leq n,$$

and such that $a_\ell^{(\delta)} \neq 0$ for all $\ell \leq n+1$. In particular the Whitney representation $\Pi_n^{(\delta)}$ generated by $f_n^{(\delta)}$, is of the form

$$\Pi_n^{(\delta)} = \bigoplus_{2 \leq \ell \leq n+1} \pi_\ell.$$

Proof. The proof relies on the observation that set of regular points

$$(\mathbb{C}^n)^{\text{reg}} := \{z \in \mathbb{C}^n : z_i \neq 0 \forall i \leq n\}$$

is dense in \mathbb{C}^n . First, we verify that the function

$$W : z = (z^{(i)})_i \mapsto (a_\ell(z)_\ell) \in \mathbb{C}^n, \quad (z^{(i)})_i \in (G/H)^n$$

is an open mapping. The map $\varphi : (G/H)^n \rightarrow \mathbb{C}^{n \times n}$

$$z = (z^{(i)})_i \mapsto \varphi(z) = ((z_1^{(i)} + \sqrt{-1}z_2^{(i)})^{-\ell})_{i,\ell}$$

is an analytic, open mapping, and so is the restriction $\varphi|_{\varphi^{-1}(\text{GL}(n, \mathbb{C}))}$ to the open set $\varphi^{-1}(\text{GL}(n, \mathbb{C}))$. Likewise, the evaluation map $e_y : \text{GL}(n, \mathbb{C}) \rightarrow \mathbb{C}^n$ defined by $e_y : T \mapsto T(y^{(i)})_i$ for $T \in \text{GL}(n, \mathbb{C})$ and the inversion map $\iota : \text{GL}(n, \mathbb{C}) \rightarrow \text{GL}(n, \mathbb{C})$ are both open mappings. By definition,

$$W = e_y \circ \iota \circ \varphi|_{\varphi^{-1}(\text{GL}(n, \mathbb{C}))}$$

so W is also an open mapping from an open subset of \mathbb{C}^n into \mathbb{C}^n .

Let

$$S_\delta = \begin{pmatrix} 0 & \delta \\ -\delta & 0 \end{pmatrix}$$

be the non-exceptional skew symmetric matrix whose Cayley transform $A(S_\delta) = (I - S_\delta)/(I + S_\delta) \in \text{SO}(2)$. Define

$$k_\delta = \begin{pmatrix} \frac{I - S_\delta}{I + S_\delta} & \\ & I \end{pmatrix} \in K = \text{SO}(2) \times \text{SO}(2).$$

Note that $x = (x^{(i)})_i \in \varphi^{-1}(\text{GL}(n, \mathbb{C}))$, so there is some $m \in \mathbb{Z}_{\geq 1}$ such that $k_\delta \cdot x \in \varphi^{-1}(\text{GL}(n, \mathbb{C}))$ for all $\delta \in (-1/m, 1/m)$. The open set

$$U_{m,\varepsilon} = \left\{ W(k_\delta \cdot x) : \delta \in (-1/m, 1/m), \left| \sum_{2 \leq \ell \leq n+1} a_\ell(k_\delta \cdot x) \varphi_\ell^+(x^{(i)}) - y^{(i)} \right| < \varepsilon \right\}$$

is an open neighborhood of $(a_\ell(x))_\ell$ in \mathbb{C}^n , and therefore it meets the dense set $(\mathbb{C}^n)^{\text{reg}}$. Choose some δ such that $(a_\ell(k_\delta \cdot x))_\ell \in U_{m,\varepsilon} \cap (\mathbb{C}^n)^{\text{reg}}$. Then the function

$$f_n^{(\delta)} = \sum_{2 \leq \ell \leq n+1} a_\ell(k_\delta \cdot x) \varphi_\ell^+$$

has the desired properties. \square

We call the representation $\Pi_n^{(\delta)}$ the *Whitney representation of the data set*. This answers the question in the introduction and shows

Theorem 3.1. *We fix $\varepsilon > 0$. If we enlarge the set of points from cardinality n to $n+1$, we can choose a small real number δ and a smooth, square integrable, K -finite function $f_{n+1}^{(\delta)} \in L^2(X(2,2)_+)$ such that $|f_{n+1}^{(\delta)}(x^{(i)}) - y^{(i)}| < \varepsilon$ for all $i \leq n+1$ and such that the representation Π_n generated by f_n is a subrepresentation of $\Pi_{n+1}^{(\delta)}$.*

3.6. An exact Whitney theorem. In this subsection, we obtain an exact Whitney theorem. Let

$$\kappa_t = \begin{pmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{pmatrix}.$$

By direct calculation,

$$\begin{aligned} \varphi_\ell^+(\kappa_t \cdot x) &= (x_1 \cos(t) + x_2 \sin(t) + \sqrt{-1}(x_1(-\sin(t)) + x_2 \cos(t)))^{-\ell} \\ &= (x_1(\cos(t) - \sqrt{-1} \sin(t)) + x_2(\sin(t) + \sqrt{-1} \cos(t)))^{-\ell} \\ &= (x_1(\cos(t) - \sqrt{-1} \sin(t)) + \sqrt{-1}x_2(\cos(t) - \sqrt{-1} \sin(t)))^{-\ell} \\ &= e^{\sqrt{-1}(-t)(-\ell)} (x_1 + \sqrt{-1}x_2)^{-\ell} \\ &= e^{\sqrt{-1}t\ell} \varphi_\ell^+(x). \end{aligned}$$

As a consequence of the algorithm 3.4, there exists a real number t_n and a Whitney function f_n such that

$$f_n(\kappa_{t_n} \cdot x^{(i)}) = y^{(i)}$$

for all $i \leq n$. Then

$$\begin{aligned} y^{(i)} &= f_n(\kappa_{t_n} \cdot x^{(i)}) \\ &= \sum_{\ell \leq n+1} a_\ell \varphi_\ell^+(k_{t_n} \cdot x^{(i)}) \\ &= \sum_{\ell \leq n+1} a_\ell e^{\sqrt{-1}t_n \ell} \varphi_\ell^+(x^{(i)}). \end{aligned}$$

In particular, if

$$a'_\ell = e^{\sqrt{-1}t_n \ell} a_\ell, \quad \ell \leq n+1,$$

then the linear combination $f'_n = \sum a'_\ell \varphi_\ell^+$ of Flensted-Jensen functions is an exact Whitney function for the data.

4. THE HIGHER DIMENSIONAL SYMMETRIC SPACES $X_{\pm}(p, 2)$

In this section, we sketch the generalization of the Whitney extension theorems to the higher dimensional symmetric spaces

$$X(p, 2)_{-} = \mathrm{SO}(p, 2)^{\circ} / \mathrm{SO}(p, 1)^{\circ} \text{ and } X(p, 2)_{+} = \mathrm{SO}(p, 2)^{\circ} / \mathrm{SO}(p-1, 2)^{\circ}.$$

4.1. Generalities. First, we clarify the definitions of certain semisimple symmetric spaces that we will study in the sequel. We refer to [15, Section 2.5] for further details. Define

$$X(p, q)_{+} = \{(x, y) \in \mathbb{R}^{p+q} : |x|^2 - |y|^2 = 1\}$$

and

$$X(p, q)_{-} = \{(x, y) \in \mathbb{R}^{p+q} : |x|^2 - |y|^2 = -1\}.$$

There is a diffeomorphism

$$X(p, q)_{-} \simeq X(q, p)_{+}$$

given by $(x, y) \mapsto (y, x)$. The Lie group $G = \mathrm{SO}(p, q)^{\circ}$ acts transitively on $X(p, q)_{+}$ and on $X(p, q)_{-}$, and there are G -diffeomorphisms

$$X(p, q)_{+} \simeq \mathrm{SO}(p, q)^{\circ} / \mathrm{SO}(p-1, q)^{\circ},$$

and

$$X(p, q)_{-} \simeq \mathrm{SO}(p, q)^{\circ} / \mathrm{SO}(p, q-1)^{\circ}.$$

In the remaining part of the paper, we will generalize our results from the space $X(2, 2)_{-}$ to the spaces $X(p, 2)_{-}$ for all $p \geq 2$. We keep $q = 2$ because we want the maximal compact subgroup K of the symmetry group $\mathrm{SO}(p, 2)^{\circ}$ to have an abelian factor which is diffeomorphic to the circle group $\mathrm{SO}(2)$, so that our space has a complex structure and we can speak about holomorphic and antiholomorphic discrete series representations.

Let $p \geq 2$. Next, we parametrize the symmetric space

$$X(p, 2)_{-} = \{(x, y) \in \mathbb{R}^{p+2} : |x|^2 - |y|^2 = -1\}$$

in terms of hyperbolic coordinates. The lemma below, which gives the KJH -decomposition of the symmetric space $X(p, q)_{-}$, motivates this reparametrization.

Lemma 4.1. *Let $p, q \geq 2$ and let $G = \mathrm{SO}(p, q)^{\circ}$. There is a decomposition*

$$G = KJH,$$

where

$$H = \begin{pmatrix} g & \\ & 1 \end{pmatrix}, \quad g \in \mathrm{SO}(p, q-1)^{\circ}, \quad K = \begin{pmatrix} k_1 & \\ & k_2 \end{pmatrix}, \quad k_1 \in \mathrm{SO}(p), k_2 \in \mathrm{SO}(q)$$

and

$$J = \exp \begin{pmatrix} & t \\ t & \end{pmatrix} = \begin{pmatrix} \cosh(t) & \sinh(t) \\ -\sinh(t) & \cosh(t) \end{pmatrix}, \quad t \in \mathbb{R}.$$

In particular, $G/H \simeq X(p, q)_{-}$ is a symmetric space of rank one.

Proof. Let $G = \mathrm{SO}(p, q)^{\circ}$ and let H be the image in G of $\mathrm{SO}(p, q-1)^{\circ}$ under the embedding

$$g \mapsto \begin{pmatrix} g & \\ & 1 \end{pmatrix}, \quad g \in \mathrm{SO}(p, q-1)^{\circ}.$$

Let

$$J/H \subseteq G/H$$

be a maximal, split, abelian subspace of G/H . We may decompose G as

$$G = KJH,$$

where

$$K = \begin{pmatrix} k_1 & \\ & k_2 \end{pmatrix}, \quad k_1 \in \mathrm{SO}(p), k_2 \in \mathrm{SO}(q)$$

is the maximal compact subgroup of G . In particular,

$$\mathrm{rank} G/H = \dim_{\mathbb{R}} J = 1.$$

□

We introduce the coordinates

$$\Phi(\xi, \xi', t) = (\xi_1 \sinh(t), \dots, \xi_p \sinh(t), \xi'_1 \cosh(t), \xi'_2 \cosh(t))$$

where $\sum_{j=1}^p \xi_j^2 = 1$ and $(\xi'_1)^2 + (\xi'_2)^2 = 1$.

4.1.1. *Data points.* Let $G = \mathrm{SO}(p, 2)^\circ$ and $H = \mathrm{SO}(p, 1)^\circ$. Recall that we say that the points $x^{(i)} \in G/H$ are *in general position* if not all points are contained in an orbit $G'/G' \cap H$ in G/H of a proper reductive subgroup G' of G isomorphic to one of the orthogonal groups $\mathrm{SO}(p', 2)^\circ$, $\mathrm{SO}(p', 1)^\circ$, or $\mathrm{SO}(p', 0)$ for some $p' \leq p$.

4.1.2. *Flensted-Jensen functions.* If $p = 2$, the Flensted-Jensen functions φ_ℓ^+ on the space

$$\begin{aligned} X &= X(2, 2)_+ = \{x \in \mathbb{R}^4 : x_1^2 + x_2^2 - x_3^2 - x_4^2 = 1\} \\ &\simeq X(2, 2)_- = \{(x_3, x_4, x_1, x_2) \in \mathbb{R}^4 : x_3^2 + x_4^2 - x_1^2 - x_2^2 = -1\} \end{aligned}$$

can be expressed in the new coordinates as follows:

$$\begin{aligned} \varphi_\ell^+(x) &= (x_1 + \sqrt{-1}x_2)^{-\ell} = (x_1 - \sqrt{-1}x_2)^\ell \frac{1}{(x_1^2 + x_2^2)^\ell} \\ &= (\xi'_1 - \sqrt{-1}\xi'_2)^\ell \cosh(t)^\ell \frac{1}{((\xi'_1)^2 + (\xi'_2)^2)^\ell \cosh(t)^{2\ell}} \\ &= (\xi'_1 - \sqrt{-1}\xi'_2)^\ell \cosh(t)^{-\ell}, \quad \ell \geq 2. \end{aligned}$$

We see that φ_ℓ^+ is a product of a spherical harmonic polynomial

$$p_\ell(\xi'_1, \xi'_2) = (\xi'_1 - \sqrt{-1}\xi'_2)^\ell$$

of degree ℓ , and a negative power of a hyperbolic cosine. This form of the Flensted-Jensen functions generalizes to symmetric spaces $X(p, q)_- = \mathrm{SO}(p, q)^\circ / \mathrm{SO}(p, q - 1)^\circ$ of arbitrary signature (p, q) [25, p. 8].

Lemma 4.2. *Consider the symmetric space $X(p, 2)_- = \mathrm{SO}(p, 2)^\circ / \mathrm{SO}(p, 1)^\circ$.*

The functions

$$\varphi_\ell(\xi', t) = (\xi'_1 - \sqrt{-1}\xi'_2)^\ell \cosh(t)^{-\ell}, \quad \ell \geq \frac{p+2}{2}, \ell \in \mathbb{Z}$$

are Flensted-Jensen functions for $X(p, 2)_-$. Any Flensted-Jensen function takes the form

$$p_\ell(\xi'_1, \xi'_2) \cosh(t)^{-\ell}, \quad (\xi'_1)^2 + (\xi'_2)^2 = 1,$$

for a harmonic polynomial p_ℓ of degree ℓ .

4.2. Whitney functions on $X(p, 2)_- = \mathrm{SO}(p, 2)^\circ / \mathrm{SO}(p, 1)^\circ$.

Theorem 4.1. *Let $p \geq 2$, and let $K = \mathrm{SO}(p) \times \mathrm{SO}(2)$ be the maximal compact subgroup of $G = \mathrm{SO}(p, 2)^\circ$. Let $H = \mathrm{SO}(p, 1)^\circ$ and consider the symmetric space $X(p, 2)_- \simeq G/H$. Let $x^{(i)} \in X(p, 2)_-, 1 \leq i \leq n$, be points in general position and consider a nonconstant function on these points with values $y^{(i)} \in \mathbb{C}, i \leq n$. Let $\varepsilon > 0$. There exists an analytic, K -finite, square-integrable function f_n on the symmetric space $X(p, 2)_-$ such that*

$$|f_n(x^{(i)}) - y^{(i)}| < \varepsilon, \quad \forall i \leq n.$$

The function f_n may be chosen as a \mathbb{C} -linear combination

$$f_n(\xi', t) = \sum_{\ell} a_{\ell} \varphi_{\ell}(\xi', t)$$

of Flensted-Jensen functions $\varphi_{\ell}(\xi', t) = (\xi'_1 - \sqrt{-1}\xi'_2)^{\ell} \cosh(t)^{-\ell}$.

Proof. For each $x^{(i)} \in G/H$, let $(\xi^{(i)}, \xi'^{(i)}, t^{(i)})$ be such that

$$x^{(i)} = (\xi_1 \sinh(t^{(i)}), \dots, \xi_p \sinh(t^{(i)}), \xi'_1 \cosh(t^{(i)}), \xi'_2 \cosh(t^{(i)})).$$

Let ℓ_0 be the smallest integer occurring as the weight of a Flensted-Jensen function $\varphi_{\ell}(\xi', t) = (\xi'_1 - \sqrt{-1}\xi'_2)^{\ell} \cosh(t)^{-\ell}$ (note that ℓ_0 depends on p). Let T be the complex $n \times n$ matrix

$$T = \left(\left((\xi'_1^{(i)} - \sqrt{-1}\xi'_2^{(i)}) \cosh(t^{(i)})^{-1} \right)^{\ell} \right)_{1 \leq i \leq n, \ell_0 \leq \ell \leq \ell_0 + n - 1}$$

and consider the linear system $(a_{\ell})_{\ell_0 \leq \ell \leq \ell_0 + n - 1} = T(y^{(i)})_i$. Since T is the product of the diagonal matrix

$$T = \left(\begin{array}{ccc} \left((\xi'_1^{(1)} - \sqrt{-1}\xi'_2^{(1)}) \cosh(t^{(1)})^{-1} \right)^{\ell_0} & & \\ & \ddots & \\ & & \left((\xi'_1^{(n)} - \sqrt{-1}\xi'_2^{(n)}) \cosh(t^{(n)})^{-1} \right)^{\ell_0} \end{array} \right)$$

with the Vandermonde matrix,

$$\left(\left((\xi'_1^{(i)} - \sqrt{-1}\xi'_2^{(i)}) \cosh(t^{(i)})^{-1} \right)^{\ell} \right)_{1 \leq i \leq n, 0 \leq \ell \leq n - 1}$$

the statement follows once the points are sufficiently distinct in $X(p, 2)_-$. This can be arranged as in the algorithm 3.4: After specifying $\varepsilon > 0$, define the skew symmetric matrix $S_{\varepsilon} = \begin{pmatrix} & \varepsilon \\ -\varepsilon & \end{pmatrix}$ with Cayley transform $k_{\varepsilon} = (I - S_{\varepsilon}) / (I + S_{\varepsilon}) \in \mathrm{SO}(2)$ and move the points $x^{(i)}$ slightly within their K -orbit in G/H by acting with

$$\begin{pmatrix} I_{p \times p} & \\ & k_{\varepsilon} \end{pmatrix} \in K$$

where $I_{p \times p}$ is the identity matrix in $\mathrm{SO}(p)$. \square

Remark: Holomorphic discrete series representations on symmetric spaces of Hermitian type were studied in [20], [21]. The Flensted-Jensen functions on these Hermitian spaces admit closed formulas involving consecutive powers of hyperbolic cosine, indexed by weights [21, Theorem 5.2], consistent with the formulas presented here. Consequently, our Whitney type extension theorems can be adapted (at least) to Hermitian spaces whose symmetry group is $\mathrm{SU}(1, 1)$.

4.3. Whitney representations. As in the classical cases of the sphere in section 2.1 and of anti de Sitter space in section 3 we associate a unitary representation of the symmetry group $\mathrm{SO}(p, 2)^\circ$ to a data set in the symmetric space $\mathrm{SO}(p, 2)^\circ / \mathrm{SO}(p, 1)^\circ$ in general position on which a nonconstant function f is defined.

Let f_n be the Whitney function associated to this data, and let V_{f_n} be the (\mathfrak{g}, K) -module generated by f_n . It defines a unitary admissible representation which we call again the Whitney representation of the data set. It is a direct sum of holomorphic discrete series representations of $\mathrm{SO}(p, 2)^\circ$.

4.4. Data in orbits of orthogonal subgroups. Assume first that the orbits have a holomorphic structure compatible with the holomorphic structure on $X(p, 2)_-$. There is a nested sequence of orthogonal symmetric spaces

$$X(2, 2)_- \subset X(3, 2)_- \subset \cdots \subset X(p-1, 2)_- \subset X(p, 2)_-.$$

Let p_0 be the smallest integer less than or equal to p such that all the points belong to $X(p_0, 2)_-$ and such that the data is in general position for the space $X(p_0, 2)_-$. We may then observe that the functions φ_ℓ^+ for

$$\ell = (p+2)/2 + \delta, (p+2)/2 + \delta + 1, \dots, (p+2)/2 + \delta + (n-1),$$

where $\delta = 1/2$ if p is odd and $\delta = 0$ if p is even, form a list of n independent Flensted-Jensen functions for all the spaces $X(p', 2)_-$ where $p_0 \leq p' \leq p$. In particular, there exists a Whitney function

$$f_n = \sum_{(p+1)/2 + \delta \leq \ell \leq (p+1)/2 + \delta + n - 1} a_\ell \varphi_\ell^+$$

for the data which is a Flensted-Jensen function for the ambient space $X(p, 2)_-$: f_n is analytic and square integrable on $X(p, 2)_-$ and K -finite for $K = \mathrm{SO}(p) \times \mathrm{SO}(2)$ and $f_n(x^{(i)}) = y^{(i)}$ for all i . Furthermore, f_n has the property that when restricted to $X(p_0, 2)$, f_n can be regarded as a Whitney function for the space $X(p_0, 2)$.

Remark: Suppose that the data are in an orbit of an orthogonal subgroup isomorphic to $\mathrm{SO}(p', 1)^\circ$ for $p' < p$. We conjecture that it is best to proceed in two steps:

- (1) In the first step, we use our algorithm to obtain a Whitney function for the proper orthogonal subspace $X(p', 1)_-$ of $X(p, 2)_-$.
- (2) In the second step, we use the results on holography by Kobayashi and Pevzner [16] to obtain an extension to $X(p, 2)_-$ and the group $\mathrm{SO}(p, 2)^\circ$. The restriction of the Whitney representations to orthogonal subgroups are known by the work of T. Kobayashi [15] and we may use this to understand the extension of a spherical Whitney function from a subgroup to the larger group.

5. WHITNEY REPRESENTATIONS IN ELEMENTARY PARTICLE PHYSICS

The Whitney representations of the generalized orthogonal groups introduced in 4.3 may be of interest in physics [3]. For the sphere in the compact case, the representations are used to describe the hydrogen atom [9]. In the work of I. E. Segal our Whitney representations are used to describe electrons [23]; see also S. Sternberg [26]. The proceedings of the biannual conferences in Varna (Bulgaria) organized

by V. Dobrev discuss more recent developments in applications of symmetries to physics [3].

REFERENCES

1. C. Berg, *Complex analysis*, 2012, noter.math.ku.dk/koman-12.pdf.
2. B. Blum-Smith and S. Villar, *Machine learning and invariant theory*, 2023, [arXiv:2209.14991](https://arxiv.org/abs/2209.14991).
3. V. Dobrev (ed.), *Quantum theory and symmetries with Lie theory and its applications in physics*, vol. 1 and 2, Springer Proceedings in Mathematics and Statistics, June 2017.
4. Ennes, H. and Tinarrage, R., *LieDetect: Detection of representation orbits of compact Lie groups from point clouds*, 2023, [arXiv:2309.03086v1](https://arxiv.org/abs/2309.03086v1).
5. C. Fefferman, *A sharp form of Whitney's extension theorem*, Ann. of Math. (2) **161** (2005), no. 1, 509–577.
6. ———, *Whitney's extension problems and interpolation of data*, Bull. Amer. Math. Soc. (N.S.) **46** (2009), no. 2, 207–220.
7. C. Fefferman, S. Ivanov, Y. Kurylev, M. Lassas, and H. Narayanan, *Reconstruction and interpolation of manifolds. I: The geometric Whitney problem*, Found. Comput. Math. **20** (2020), no. 5, 1035–1133.
8. M. Flensted-Jensen, *Discrete series for semisimple symmetric spaces*, Annals of Mathematics (1980), no. 111, 253–311.
9. B. C. Hall, *Quantum Theory for Mathematicians*, Graduate Texts in Mathematics 267, Springer, 2013.
10. ———, *Lie Groups, Lie Algebras, and Representations*, 2nd ed., Graduate Texts in Mathematics 222, Springer, 2015.
11. F. Kassel and T. Kobayashi, *Stable spectrum for pseudo-Riemannian locally symmetric spaces*, C. R. Acad. Sci. Paris (2011), 29–33.
12. ———, *Poincaré series for non-Riemannian locally symmetric spaces*, Advances in Mathematics (2016), no. 287, 123–236.
13. ———, *Spectral analysis on standard locally homogeneous spaces*, (2019), [arXiv:1912.12601](https://arxiv.org/abs/1912.12601).
14. A.W. Knap, *Lie groups beyond an introduction*, second ed., Progress in mathematics, vol. 140, Birkhauser, 2002.
15. T. Kobayashi, *Branching laws of unitary representations associated to minimal elliptic orbits for indefinite orthogonal group $O(p, q)$* , Adv. Math. **388** (2021).
16. T. Kobayashi and M. Pevzner, *Inversion of Rankin-Cohen operators via holographic transform*, Ann. Inst. Fourier (Grenoble) **70** (2020), no. 5, 2131–2190.
17. C. Kondor and S. Trivedi, *On the generalization of equivariance and convolution in neural networks to the action of compact groups*, Proceedings of the 35th International Conference on Machine Learning (2018).
18. J. Maldacena, *The illusion of gravity*, Scientific American (2005), no. 293, 56–63.
19. D. Marcus, *Number Fields*, Springer, 1977.
20. B. Ørsted and G. Olafsson, *The holomorphic discrete series for affine symmetric spaces, I*, Journal of Functional Analysis **81** (1988), 126–159.
21. ———, *The holomorphic discrete series of an affine symmetric space and representations with reproducing kernels*, Transactions of the American Mathematical Society **326** (1991), no. 1, 385–405.
22. H. Schlichtkrull, *The Langlands parameters of Flensted-Jensen's discrete series for semisimple symmetric spaces*, J. Functional Analysis **50** (1983), no. 2, 133–150.
23. I. E. Segal, H. P. Jakobsen, B. Ørsted, S. M. Paneitz, and B. Speh, *Covariant chronogeometry and extreme distances: elementary particles*, Proc. Nat. Acad. Sci. U.S.A. **78** (1981), no. 9, 5261–5265.
24. J.-P. Serre, *A course in Arithmetic*, Springer, 1973.
25. B. Speh and B. Ørsted, *Branching laws for discrete series of some affine symmetric spaces*, 2019, [arXiv:1907.07544](https://arxiv.org/abs/1907.07544).
26. S. Sternberg, *Group theory and physics*, Cambridge University Press, Cambridge, 1994.
27. E. P. van den Ban and H. Schlichtkrull, *The Plancherel decomposition for a reductive symmetric space. II. Representation theory*, Invent. Math. **161** (2005), no. 3, 567–628.
28. H. Weyl, *Classical Groups*, second ed., Princeton University Press, 1946.

29. H. Whitney, *Analytic extensions of differentiable functions defined in closed sets*, Trans. Amer. Math. Soc. **36** (1934), no. 1, 63–89.
30. A. Zee, *Einstein gravity in a nutshell*, Princeton University Press, 2013.