

# ON $\mu$ -INVARIANTS AND ISOGENIES FOR ABELIAN VARIETIES OVER FUNCTION FIELDS

SOHAN GHOSH, JISHNU RAY, AND TAKASHI SUZUKI

ABSTRACT. We give several formulas for how Iwasawa  $\mu$ -invariants of abelian varieties over unramified  $\mathbb{Z}_p$ -extensions of function fields change under isogeny. These are analogues of Schneider’s formula in the number field setting. We also prove that the validity of the Birch–Swinnerton-Dyer conjecture (including the leading coefficient formula) over function fields is invariant under isogeny, without using the result of Kato–Trihan.

## CONTENTS

1. Introduction	1
2. Notation and conventions	3
3. The ind-rational pro-étale site	4
4. Conductors of isogenies in the local case	6
5. Conductors of isogenies in the global case	8
6. Change in $\mu$ by flat Euler characteristics: general base field case	9
7. Change in $\mu$ by flat Euler characteristics: finite base field case	11
8. Lie complexes	12
9. Duality of Euler characteristics	13
10. Change in $\mu$ by degrees of Lie complexes	14
11. Isogeny invariance of the BSD formula	15
12. Base change stable invariants	16
13. Change in $\mu$ with base change conductors	17
14. Another proof of Theorem 10.3 by the $\mu$ -invariant formula	18
References	19

## 1. INTRODUCTION

Schneider [Sch87] studies how Iwasawa  $\mu$ -invariants of dual Selmer groups of abelian varieties over cyclotomic  $\mathbb{Z}_p$ -extensions of number fields change under isogenies of abelian varieties. On the other hand, Iwasawa theory for abelian varieties over unramified  $\mathbb{Z}_p$ -extensions of function fields in one variable of characteristic  $p$  is initiated by Ochiai–Trihan [OT09]. In this paper, we will carry out an analogue of Schneider’s studies over unramified  $\mathbb{Z}_p$ -extensions of function fields.

---

*Date:* July 31, 2024.

*2020 Mathematics Subject Classification.* 11R23 (Primary) 11G10, 11R58, 11G40, 14F20 (Secondary).

*Key words and phrases.* Iwasawa invariants; abelian varieties; function fields; Birch–Swinnerton-Dyer conjectures.

Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over a function field  $K$  of characteristic  $p > 0$ . Let  $k$  be the constant field of  $K$ . Let  $K_\infty$  be the (unique) unramified  $\mathbb{Z}_p$ -extension of  $K$ . For  $i = 1, 2$ , let  $\mu_{A_i/K}$  be  $1/[k : \mathbb{F}_p]$  times the  $\mu$ -invariant of the dual Selmer group of  $A_i$  over  $K_\infty$  as a  $\mathbb{Z}_p[[\text{Gal}(K_\infty/K)]]$ -module, which is finite by [OT09, Theorem 1.7]. We will give three different formulas for  $\mu_{A_2/K} - \mu_{A_1/K}$ .

Let  $X$  be the proper smooth curve with function field  $K$ . Let  $\mathcal{A}_i$  be the Néron model over  $X$  of  $A_i$ . Let  $\mathcal{N}_f \subset \mathcal{A}_1$  be the schematic closure of the kernel of  $f: A_1 \rightarrow A_2$  in  $\mathcal{A}_1$ , which is a quasi-finite flat separated group scheme over  $X$ . In Definition 5.1, we will define a certain non-negative integer  $c(f)$ , which we call the *conductor* of  $f$ . It is a locally defined quantity that measures the potential failure of  $f$  to extend to an isogeny on the Néron models.

For a non-zero rational number  $r$ , let  $\text{ord}_q(r)$  be the  $q$ -valuation of  $r$ , which is the  $p$ -adic valuation divided by  $[k : \mathbb{F}_p]$  (so  $\text{ord}_q(q) = 1$ ). Our first formula is:

**Theorem 1.1** (Proposition 7.1, Theorem 7.2).

- (1) *For any quasi-finite flat separated group scheme  $\mathcal{N}$  over  $X$ , the flat cohomology group  $H^i(X, \mathcal{N})$  is finite for all  $i$  and zero for almost all  $i$ . This defines a well-defined number*

$$\chi(X, \mathcal{N}) := \prod_i (\#H^i(X, \mathcal{N}))^{(-1)^i}.$$

- (2) *We have*

$$\mu_{A_2/K} - \mu_{A_1/K} = \text{ord}_q(\chi(X, \mathcal{N}_f)) - c(f).$$

*In particular, if  $f$  extends to an isogeny on the Néron models, then*

$$\mu_{A_2/K} - \mu_{A_1/K} = \text{ord}_q(\chi(X, \mathcal{N}_f)).$$

We can rewrite the term  $\text{ord}_q(\chi(X, \mathcal{N}_f))$  with a certain linear or coherent object. For any quasi-finite flat separated group scheme  $\mathcal{N}$  over  $X$ , let  $R\text{Lie}\mathcal{N} := L_{\mathcal{N}}^\vee$  be the Lie complex as defined by Illusie [Ill72, Chapter VII, Section 3.1.1]. It is a perfect complex of  $\mathcal{O}_X$ -modules whose zero-th cohomology is the usual Lie algebra  $\text{Lie}\mathcal{N}$ . Let  $\text{deg } R\text{Lie}\mathcal{N}$  be its degree with respect to the base field  $k$ .

**Theorem 1.2** (Theorem 10.2). *For any quasi-finite flat separated group scheme  $\mathcal{N}$  over  $X$ , we have*

$$\chi(X, \mathcal{N}) = q^{\text{deg } R\text{Lie}\mathcal{N}}.$$

This solves the problem of finding a formula for  $\chi(X, \mathcal{N})$ , which is originally posed by Milne [Mil06, Chapter III, Problem 8.10]. It implies our second formula, which is a function field version of the ‘‘Isogeny formula (First form)’’ at the end of [Sch87]:

**Theorem 1.3** (Theorem 10.3). *We have*

$$\mu_{A_2/K} - \mu_{A_1/K} = \text{deg } R\text{Lie}\mathcal{N}_f - c(f).$$

*In particular, if  $f$  extends to an isogeny on the Néron models, then*

$$\mu_{A_2/K} - \mu_{A_1/K} = \text{deg } R\text{Lie}\mathcal{N}_f.$$

The term  $\deg R\mathrm{Lie}\mathcal{N}_f$  changes in a complicated manner if  $K$  is replaced by a finite extension. We can replace it by a more stable invariant, at the expense of adding and subtracting Chai’s base change conductors [Cha00] of the abelian varieties.

Let  $c(A_i)$  be the base change conductor of  $A_i$ . Let  $L/K$  be a finite separable extension over which  $A_i$  has semistable reduction everywhere. Let  $\mathcal{N}_{f,L}$  be the object  $\mathcal{N}_f$ , this time defined from the base change  $f: A_1 \times_K L \rightarrow A_2 \times_K L$  of  $f$  to  $L$ . The ratio  $\deg R\mathrm{Lie}\mathcal{N}_{f,L}/[L:K]$  is independent of the choice of  $L$ . Denote it by  $\deg_X R\mathrm{Lie}\mathcal{N}_{f,K^{\mathrm{sep}}}$ . Our third formula is:

**Theorem 1.4** (Theorem 13.1). *We have*

$$\mu_{A_2/K} - \mu_{A_1/K} = \deg_X R\mathrm{Lie}\mathcal{N}_{f,K^{\mathrm{sep}}} + c(A_2) - c(A_1).$$

Our results and methods are also useful to study the behavior of the Birch–Swinnerton-Dyer conjecture over function fields under isogeny, including the formula for the leading coefficient of the  $L$ -function. This formula (under the assumption of the finiteness of  $\mathrm{III}(A)$ ) is proved by Kato–Trihan [KT03]. Here we prove, without using the result of [KT03], that the validity of the leading coefficient formula is invariant under arbitrary isogeny on  $A$ , whose degree may be divisible by the characteristic  $p > 0$ . Proving this isogeny invariance is a problem posed by Milne [Mil06, Chapter III, Problem 9.10].

**Theorem 1.5** (Theorem 11.1). *The Birch–Swinnerton-Dyer formula holds for  $A_1$  if and only if it does for  $A_2$ .*

Kato–Trihan’s proof of the BSD formula uses heavy integral  $p$ -adic cohomology arguments, while our simple proof of isogeny invariance does not.

We will also give another simpler proof of Theorem 1.3 without using Theorem 1.1 or 1.2. It instead uses the  $\mu$ -invariant formula given by the third author’s article with Lai–Longhi–Tan–Trihan [LLS<sup>+</sup>21]. This formula is proved in many cases, but not in full expected generality. Therefore the proof based on this  $\mu$ -invariant formula, though simpler, is conditional.

Actually a certain version of  $\mu$ -invariant can be geometrically defined even if the constant field  $k$  is not finite but perfect of positive characteristic more generally. It is the dimension of the “Tate–Shafarevich scheme” constructed by the third author [Suz20a] based on “the ind-rational pro-étale site of  $k$ ”  $\mathrm{Spec} k_{\mathrm{proet}}^{\mathrm{indrat}}$ . That it agrees with the  $\mu$ -invariant when  $k$  is finite is proved in [LLS<sup>+</sup>21]. In this paper, we will review some notation about  $\mathrm{Spec} k_{\mathrm{proet}}^{\mathrm{indrat}}$  and then formulate and prove geometric versions of Theorems 1.1, 1.2, 1.3 and 1.4 for general perfect  $k$ . These versions immediately imply the original versions above.

*Acknowledgments.* The second author thanks Aditya Karnataki for inviting him to Chennai Mathematical Institute in July 2023. He also thanks Anwesh Ray for discussing similar questions dealt in this article in the number field setup for non-commutative Lie extensions which inspired him to ask these questions in the function field case. He also gratefully acknowledges support from the Inspire research grant. The first author acknowledges the support received from the HRI postdoctoral fellowship.

## 2. NOTATION AND CONVENTIONS

All groups and group schemes are assumed to be commutative.

Let  $\text{Ab}$  be the category of abelian groups. For an abelian category  $\mathcal{A}$ , its derived category is denoted by  $D(\mathcal{A})$  and its bounded derived category by  $D^b(\mathcal{A})$ . The mapping cone of a morphism  $A \rightarrow B$  between complexes in  $\mathcal{A}$  is denoted by  $[A \rightarrow B]$  (hence its negative shift  $[A \rightarrow B][-1]$  is the mapping fiber). If we say  $A \rightarrow B \rightarrow C$  is a distinguished triangle in  $D(\mathcal{A})$ , we implicitly assume that a morphism  $C \rightarrow A[1]$  to the shift of  $A$  is given and the resulting triangle is  $A \rightarrow B \rightarrow C \rightarrow A[1]$  is distinguished. If  $F: \mathcal{A} \rightarrow \mathcal{B}$  is an additive functor from an abelian category with enough K-injectives to another abelian category, then its right derived functor is denoted by  $RF: D(\mathcal{A}) \rightarrow D(\mathcal{B})$ .

We denote the category of sheaves of abelian groups on a Grothendieck site  $S$  by  $\text{Ab}(S)$ . Denote  $D^*(S) = D^*(\text{Ab}(S))$  with  $*$  = (blank) or  $b$ .

For an irreducible quasi-compact regular scheme  $X$  of dimension 1 with perfect residue fields at closed points, let  $X_0$  be the set of closed points of  $X$ . For  $v \in X_0$ , let  $\mathcal{O}_v$  be the completed local ring of  $X$  at  $v$ , with fraction field  $K_v$  and residue field  $k(v)$ .

From Section 6 onward, we take  $X$  to be a proper smooth geometrically connected curve over a perfect field  $k$  of characteristic  $p > 0$ , with function field  $K$ .

For a power  $q = p^e$  of a prime  $p$  and a non-zero rational number  $r$ , let  $\text{ord}_q(r)$  be the  $q$ -valuation of  $r$ , which is the  $p$ -adic valuation divided by  $e$  (so  $\text{ord}_q(q) = 1$ ).

### 3. THE IND-RATIONAL PRO-ÉTALE SITE

Let  $k$  be a perfect field of characteristic  $p > 0$ . We recall some notation about perfection of algebraic groups and the ind-rational pro-étale site  $\text{Spec } k_{\text{proet}}^{\text{indrat}}$  from [Suz20b] and [Suz20a]. See also [OS23, Section 3].

The perfection of a  $k$ -algebra  $R$  (respectively a  $k$ -scheme  $X$ ) is the direct limit  $R \rightarrow R \rightarrow \cdots$  (respectively the inverse limit  $\cdots \rightarrow X \rightarrow X$ ) along the Frobenius morphisms, equipped with the natural  $k$ -algebra (respectively  $k$ -scheme) structure.

Let  $\text{LAlg}/k$  be the category of perfection of smooth group schemes over  $k$  with group scheme morphisms. Let  $\text{Alg}/k \subset \text{LAlg}/k$  be the full (abelian) subcategory of quasi-compact groups. Let  $\text{LAlg}_{\text{uc}}/k \subset \text{LAlg}/k$  be the full subcategory of groups with unipotent identity component. Let  $\text{Alg}_{\text{uc}}/k = \text{Alg}/k \cap \text{LAlg}_{\text{uc}}/k$ .

A  $k$ -algebra  $k'$  is said to be *rational* if it can be written as  $k'_1 \times \cdots \times k'_n$ , where each  $k'_i$  is the perfection of a finitely generated extension field over  $k$ . A  $k$ -algebra is said to be *ind-rational* if it can be written as a filtered direct limit of rational  $k$ -algebras. Let  $\text{Spec } k_{\text{proet}}^{\text{indrat}}$  be (the opposite of) the category of ind-rational  $k$ -algebras with  $k$ -algebra homomorphisms equipped with the pro-étale topology. We denote  $\text{Ab}(\text{Spec } k_{\text{proet}}^{\text{indrat}}) = \text{Ab}(k_{\text{proet}}^{\text{indrat}})$  and  $D^*(\text{Spec } k_{\text{proet}}^{\text{indrat}}) = D^*(k_{\text{proet}}^{\text{indrat}})$  with  $*$  = (blank) or  $b$ . The cohomology functor  $R\Gamma(\text{Spec } k_{\text{proet}}^{\text{indrat}}, \cdot)$  on this site is simply denoted by  $R\Gamma(k, \cdot)$ . The natural Yoneda functor  $\text{Alg}/k \rightarrow \text{Ab}(k_{\text{proet}}^{\text{indrat}})$  is fully faithful exact and induces a fully faithful embedding  $D^b(\text{Alg}/k) \hookrightarrow D(k_{\text{proet}}^{\text{indrat}})$  ([Suz20b, Proposition 2.3.4]). The (bounded) derived category of étale group schemes over  $k$  is also a full subcategory of  $D(k_{\text{proet}}^{\text{indrat}})$  ([Suz20b, Proposition 2.3.4]). In particular, the natural Yoneda functor  $\text{LAlg}/k \rightarrow \text{Ab}(k_{\text{proet}}^{\text{indrat}})$  is fully faithful exact.

Let  $X$  be a proper smooth geometrically connected curve over  $k$ . Let  $U \subset X$  be a dense open subscheme. Let  $U_{\text{fppf}}$  be the fppf site of  $U$ . Let

$$\Gamma(U, \cdot): \text{Ab}(U_{\text{fppf}}) \rightarrow \text{Ab}(k_{\text{proet}}^{\text{indrat}})$$

be the left exact functor defined in [Suz20a, Section 2.7]. Denote its  $i$ -th right derived functor by  $\mathbf{H}^i(U, \cdot)$ . If  $G$  is a group scheme locally of finite type over  $U$  (or a bounded complex thereof), then we have a canonical isomorphism

$$R\Gamma(k, R\Gamma(U, G)) \cong R\Gamma(U, G)$$

([Suz20a, Proposition 2.7.8] and the paragraph thereafter), where the right-hand side is the usual fppf cohomology.

Let  $v \in X_0$ . Let  $\text{Ab}(\mathcal{O}_{v, \text{fppf}})$  be the category of sheaves of abelian groups on the fppf site  $\text{Spec } \mathcal{O}_{v, \text{fppf}}$  of  $\mathcal{O}_v$ . Let

$$\Gamma(\mathcal{O}_v, \cdot): \text{Ab}(\mathcal{O}_{v, \text{fppf}}) \rightarrow \text{Ab}(k(v)_{\text{proet}}^{\text{indrat}})$$

be the left exact functor defined in [Suz20a, Section 2.5]. Denote its  $i$ -th right derived functor by  $\mathbf{H}^i(\mathcal{O}_v, \cdot)$ . For any  $G \in D(\mathcal{O}_{v, \text{fppf}})$ , there is a canonical morphism

$$R\Gamma(\mathcal{O}_v, G) \rightarrow R\Gamma(k, R\Gamma(\mathcal{O}_v, G))$$

in  $D(\text{Ab})$ . It is an isomorphism if  $G$  is a smooth group scheme or a finite flat group scheme (the paragraph before [Suz20a, Proposition 2.5.2]). It is also an isomorphism if  $G$  is a quasi-finite flat separated group scheme since such a group is an extension of an étale (hence smooth) group scheme by a finite flat group scheme (the finite part of  $G$ ; see the paragraph before [BLR90, Section 7.3, Proposition 3]).

Let  $f: \text{Spec } k(v) \rightarrow \text{Spec } k$  be the natural morphism. Let  $f: \text{Spec } k(v)_{\text{proet}}^{\text{indrat}} \rightarrow \text{Spec } k_{\text{proet}}^{\text{indrat}}$  be the induced morphism of sites (defined by the functor  $k' \otimes_k k(v) \leftarrow k'$  on the underlying categories). Set

$$\Gamma(\mathcal{O}_v/k, \cdot) := f_*\Gamma(\mathcal{O}_v, \cdot): \text{Ab}(\mathcal{O}_{v, \text{fppf}}) \rightarrow \text{Ab}(k_{\text{proet}}^{\text{indrat}})$$

as in the end of [Suz20a, Section 2.5]. Denote its  $i$ -th right derived functor by  $\mathbf{H}^i(\mathcal{O}_v/k, \cdot)$ .

Let

$$R\Gamma_c(U, \cdot): D(U_{\text{fppf}}) \rightarrow D(k_{\text{proet}}^{\text{indrat}})$$

be the triangulated functor defined in [Suz20a, Section 2.7]. Denote  $\mathbf{H}_c^i(U, \cdot) = H^i R\Gamma_c(U, \cdot)$ . If  $G$  is a smooth group scheme or a finite flat group scheme (or a bounded complex thereof) over  $U$ , then there exists a canonical distinguished triangle

$$(3.1) \quad R\Gamma_c(U, G) \rightarrow R\Gamma(X, G) \rightarrow \bigoplus_{v \in X \setminus U} R\Gamma(\mathcal{O}_v/k, G)$$

in  $D(k_{\text{proet}}^{\text{indrat}})$  functorial in  $G$  ([Suz20a, Propositions 2.7.4 and 2.7.5]). This also applies to  $G$  being a quasi-finite flat separated group scheme since the required condition (the ‘‘cohomological approximation’’ property) is étale local and  $G$  is étale locally an extension of an étale group scheme by a finite flat group scheme. Applying  $R\Gamma(k, \cdot)$  recovers the compact support fppf cohomology distinguished triangle

$$R\Gamma_c(U, G) \rightarrow R\Gamma(X, G) \rightarrow \bigoplus_{v \in X \setminus U} R\Gamma(\mathcal{O}_v, G)$$

in  $D(\text{Ab})$  given in [Mil06, Chapter III, Proposition 0.4 (c), Remark 0.6 (b)] (see also [DH19, Proposition 2.1 (3)]).

We will need the following result to pass from the general base field case to the finite base field case.

**Proposition 3.1.** *Assume that  $k = \mathbb{F}_q$  is finite.*

- (1) *Let  $G \in D^b(\text{Alg}/k)$ . Then  $H^i(k, G)$  is finite for all  $i$  and zero for almost all  $i$ .*  
(2) *Let  $G \in D^b(\text{Alg}_{\text{uc}}/k)$ . Then*

$$\sum_i (-1)^i \text{ord}_q(\#H^i(k, G)) = \sum_i (-1)^i \dim H^i(G)$$

(where  $H^i(G)$  is the  $i$ -th cohomology of  $G$  as a complex).

*Proof.* (1) We may assume  $G \in \text{Alg}/k$ . Then  $H^0(k, G) = G(k)$  and  $H^1(k, G) = H^1(k, \pi_0(G))$  are finite and  $H^i(k, G) = 0$  for  $i \geq 2$  by Lang's theorem.

(2) We may assume that  $G \in \text{Alg}_{\text{uc}}/k$ . If  $G$  is finite étale, then the exact sequence

$$0 \rightarrow H^0(k, G) \rightarrow G(\bar{k}) \xrightarrow{F_k^{-1}} G(\bar{k}) \rightarrow H^1(k, G) \rightarrow 0$$

shows that  $H^0(k, G)$  and  $H^1(k, G)$  have the same cardinality, where  $F_k$  denotes the  $q$ -th power Frobenius over  $k$ . With  $H^i(k, G) = 0$  for  $i \neq 0, 1$ , we get the result in this case.

If  $G$  is connected (and hence unipotent), we may assume  $G = \mathbf{G}_a$ . Then  $\#H^0(k, G) = \#k = q$  and  $H^i(k, G) = 0$  for  $i \neq 0$ , which give the result.  $\square$

#### 4. CONDUCTORS OF ISOGENIES IN THE LOCAL CASE

In this section, we will define a number that we call the “conductor”,  $c(f)$ , for an isogeny  $f$  between abelian varieties over local fields. Such an isogeny does not extend to an isogeny on the Néron models (in the sense of having finite kernel on the special fibers) in general. Our conductor measures this defect. The kernel of the morphism induced on the Néron models is not finite, not even quasi-finite, in general. We first extract a certain quasi-finite group scheme  $\mathcal{N}_f$  from this non-quasi-finite group scheme. With this and the Liu–Lorenzini–Raynaud constructions [LLR04, Theorem 2.1 (b)], we define  $c(f)$ . Both  $\mathcal{N}_f$  and  $c(f)$  contribute to the change in  $\mu$ -invariants in the global situation.

Let  $K$  be a complete discrete valuation field with ring of integers  $\mathcal{O}_K$  and perfect residue field  $k$ . Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over  $K$  with kernel  $N_f$ . We have an exact sequence

$$0 \rightarrow N_f \rightarrow A_1 \rightarrow A_2 \rightarrow 0.$$

For  $i = 1, 2$ , let  $\mathcal{A}_i$  be the Néron model over  $\mathcal{O}_K$  of  $A_i$ . Then  $f$  induces a morphism  $\tilde{f}: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  of group schemes over  $\mathcal{O}_K$ . Let  $\mathcal{N}_f \subset \mathcal{A}_1$  be the schematic closure of  $N_f$  in  $\mathcal{A}_1$ , which is a quasi-finite flat separated closed group subscheme over  $\mathcal{O}_K$ . The quotient  $\mathcal{A}'_2 := \mathcal{A}_1/\mathcal{N}_f$  as an fppf sheaf over  $\mathcal{O}_K$  is representable by a smooth separated group scheme of finite type by [Ana73, Theorem 4.C]. Since  $\tilde{f}: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  annihilates  $\mathcal{N}_f$ , we have an induced morphism  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$ :

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{N}_f & \longrightarrow & \mathcal{A}_1 & \longrightarrow & \mathcal{A}'_2 & \longrightarrow & 0 \\ & & & & & & \downarrow & & \\ & & & & & & \mathcal{A}_2 & & \end{array}$$

The morphism  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$  is an isomorphism on the generic fibers. Hence by [LLR04, Theorem 2.1 (b)] (see also [OS23, Proposition 2.4 and Remark 2.5]), there exists a canonical smooth group scheme  $D = D_f$  of finite type over  $k$  such that  $D_f(\bar{k})$

is canonically isomorphic to the cokernel of  $\mathcal{A}'_2(\hat{\mathcal{O}}_K^{\text{ur}}) \hookrightarrow \mathcal{A}_2(\hat{\mathcal{O}}_K^{\text{ur}}) = \mathcal{A}_2(\hat{K}^{\text{ur}})$  as  $\text{Gal}(\bar{k}/k)$ -modules and  $\dim D_f$  is equal to the  $\mathcal{O}_K$ -length of the cokernel of  $\text{Lie } \mathcal{A}'_2 \hookrightarrow \text{Lie } \mathcal{A}_2$ , where  $\hat{\mathcal{O}}_K^{\text{ur}}$  is the completed maximal unramified extension of  $\mathcal{O}_K$  and  $\hat{K}^{\text{ur}}$  its fraction field. More explicitly,  $D_f$  is given by the cokernel of the morphism  $\text{Gr}_{\mathcal{O}_K} \mathcal{A}'_2 \rightarrow \text{Gr}_{\mathcal{O}_K} \mathcal{A}_2$  induced on the infinite-level Greenberg transforms of  $\mathcal{A}'_2$  and  $\mathcal{A}_2$  ([BGA18, Section 14], [OS23, Remark 2.5]):

$$(4.1) \quad \text{Gr}_{\mathcal{O}_K} \mathcal{A}'_2 \rightarrow \text{Gr}_{\mathcal{O}_K} \mathcal{A}_2 \rightarrow D_f \rightarrow 0.$$

The kernel of the first morphism  $\text{Gr}_{\mathcal{O}_K} \mathcal{A}'_2 \rightarrow \text{Gr}_{\mathcal{O}_K} \mathcal{A}_2$  is pro-infinitesimal (since it is injective on geometric points). This kernel is zero if  $K$  has equal characteristic by the explicit description of Greenberg transforms as Weil restrictions in this case. In general,  $D_f$  has unipotent identity component since  $\mathcal{A}_1 \rightarrow \mathcal{A}'_2 \rightarrow \mathcal{A}_2$  are isomorphisms up to bounded torsion.

**Definition 4.1.** *Define  $c(f)$  to be the dimension of  $D_f$  and call it the conductor of the isogeny  $f$ .*

Recall from [BLR90, Section 7.3, Definition 4] that  $\tilde{f}: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  is called an isogeny if the induced morphism  $\mathcal{A}_{1,k} \rightarrow \mathcal{A}_{2,k}$  on the special fibers is an isogeny (namely, has finite kernel). The number  $c(f)$  measures the defect of  $\tilde{f}$  being an isogeny:

**Proposition 4.2.** *We have  $c(f) = 0$  if and only if  $\tilde{f}: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  is an isogeny, in which case its kernel is  $\mathcal{N}_f$  and the morphism  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$  is an open immersion.*

*Proof.* If  $\tilde{f}: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  is an isogeny, then  $\text{Ker}(\tilde{f})$  is quasi-finite flat separated. Such a closed subgroup scheme of  $\mathcal{A}_1$  is necessarily the schematic closure of its generic fiber. Hence  $\text{Ker}(f) = \mathcal{N}_f$ . Therefore  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$  is a monomorphism. As it is a morphism of smooth group schemes and an isomorphism on the generic fibers, it follows that  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$  is an open immersion. Hence  $c(f) = 0$ .

Conversely, if  $c(f) = 0$ , then  $\text{Lie } \mathcal{A}'_2 \xrightarrow{\sim} \text{Lie } \mathcal{A}_2$ , so  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$  is an open immersion. Hence  $\mathcal{A}_{1,k} \rightarrow \mathcal{A}_{2,k}$  is surjective on the identity components and hence has finite kernel. Thus  $\tilde{f}$  is an isogeny.  $\square$

Note that  $\tilde{f}: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  is an isogeny if  $\mathcal{A}_1$  has semistable reduction or the degree of  $f: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  is prime to the characteristic of  $k$  ([BLR90, Section 7.3, Proposition 6]). The converse holds if  $f$  is multiplication by a natural number on  $\mathcal{A}_1 = \mathcal{A}_2$  ([Mil06, Chapter III, Corollary C.9]). This gives examples with  $c(f) \neq 0$ .

We also recall Chai's base change conductors. Let  $A$  be an abelian variety over  $K$ . Let  $L$  be a finite extension of  $K$  with ring of integers  $\mathcal{O}_L$  and ramification index  $e_{L/K}$  over which  $A$  has semistable reduction. Let  $\mathcal{A}$  be the Néron model over  $\mathcal{O}_K$  of  $A$  and  $\mathcal{A}_L$  the Néron model over  $\mathcal{O}_L$  of  $A \times_K L$ .

**Definition 4.3** ([Cha00]). *Define a rational number  $c(A)$  by*

$$c(A) = \frac{1}{e_{L/K}} \text{length}_{\mathcal{O}_L} \frac{\text{Lie } \mathcal{A}_L}{\text{Lie } \mathcal{A} \otimes_{\mathcal{O}_K} \mathcal{O}_L},$$

where  $\text{length}_{\mathcal{O}_L}$  denotes the length of  $\mathcal{O}_L$ -modules, and call it the base change conductor of  $A$ .

This is independent of the choice of  $L$ .

## 5. CONDUCTORS OF ISOGENIES IN THE GLOBAL CASE

We will globalize the constructions of the previous section.

Let  $X$  be an irreducible quasi-compact regular scheme of dimension 1 with perfect residue fields at closed points. Let  $K$  be its function field. Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over  $K$ . For any  $v \in X_0$ , it gives an isogeny  $f: A_1 \times_K K_v \rightarrow A_2 \times_K K_v$  over the local field  $K_v$  at  $v$ . Applying the constructions in the previous section, we have the corresponding group scheme  $D_{f,v}$  and the conductor  $c_v(f) = \dim D_{f,v}$ . Since  $A_1$  has semistable reduction almost everywhere, we know that  $c_v(f) = 0$  for almost all  $v$  by Proposition 4.2.

**Definition 5.1.**

- (1) Define a divisor on  $X$  by  $\mathfrak{c}(f) = \sum_{v \in X_0} c_v(f)v$  and call it the conductor divisor of  $f$ .
- (2) When  $X$  is a proper smooth geometrically connected curve over a perfect field  $k$ , define  $c(f)$  to be the degree of  $\mathfrak{c}(f)$  (with respect to  $k$ ) and call it the conductor of  $f$ .
- (3) When  $X$  is the Spec of the ring of integers of a number field, define  $c(f)$  to be the norm of  $\mathfrak{c}(f)$  (with respect to  $\mathbb{Q}$ ) and call it the conductor of  $f$ .

We do not really treat the number field case in this paper. The number field definition is included here just for completeness.

Let  $N_f$  be the kernel of  $f$ . Let  $\mathcal{A}_i$  be the Néron model over  $X$  of  $A_i$ . Let  $\tilde{f}: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  be the induced morphism. Let  $\mathcal{N}_f \subset \mathcal{A}_1$  be the schematic closure of  $N_f$ , which is a quasi-finite flat separated closed group subscheme over  $X$ . The quotient  $\mathcal{A}'_2 := \mathcal{A}_1/\mathcal{N}_f$  as an fppf sheaf over  $\mathcal{O}_K$  is representable by a smooth separated group scheme of finite type by [Ana73, Theorem 4.C]. We have an induced morphism  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$ :

$$(5.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{N}_f & \longrightarrow & \mathcal{A}_1 & \longrightarrow & \mathcal{A}'_2 & \longrightarrow & 0 \\ & & & & & & \downarrow & & \\ & & & & & & \mathcal{A}_2 & & \end{array}$$

**Proposition 5.2.** *We have  $\mathfrak{c}(f) = 0$  if and only if  $\tilde{f}: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  is an isogeny, in which case its kernel is  $\mathcal{N}_f$  and the morphism  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$  is an open immersion.*

*Proof.* This follows from the corresponding local statement, Proposition 4.2.  $\square$

We also globalize base change conductors. Let  $A$  be an abelian variety over  $K$ . For each  $v \in X_0$ , we have the base change conductor  $c_v(A)$  of  $A \times_K K_v$ .

**Definition 5.3.**

- (1) Define a  $\mathbb{Q}$ -divisor on  $X$  by  $\mathfrak{c}(A) = \sum_{v \in X_0} c_v(A)v$  and call it the base change conductor divisor of  $A$ .
- (2) When  $X$  is a proper smooth geometrically connected curve over a perfect field  $k$ , define  $c(A)$  to be the degree of  $\mathfrak{c}(A)$  (with respect to  $k$ ) and call it the base change conductor of  $A$ .
- (3) When  $X$  is the Spec of the ring of integers of a number field, define  $c(A)$  to be the norm of  $\mathfrak{c}(A)$  (with respect to  $\mathbb{Q}$ ) and call it the base change conductor of  $A$ .



*Remark 5.4.* The basic idea in the constructions in this and the previous sections is to imagine to have a kind of “exact sequence”

$$“ 0 \rightarrow \mathcal{N}_f \rightarrow \mathcal{A}_1 \rightarrow \mathcal{A}_2 \rightarrow \bigoplus_{v \in X_0} i_{v,*} D_{f,v} \rightarrow 0 ”,$$

where  $i_v: \text{Spec } k(v) \hookrightarrow X$  is the inclusion. It is not really clear in what category this supposed exact sequence actually lives, since  $D_{f,v}$  is not an étale scheme and  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$  is not injective on the special fibers in general. This sequence is trying to decompose the information about  $\mathcal{A}_1 \rightarrow \mathcal{A}_2$  into the information about  $\mathcal{N}_f$  and  $D_{f,v}$ . This idea works out in practice, as we will see in the rest of the paper.

## 6. CHANGE IN $\mu$ BY FLAT EULER CHARACTERISTICS: GENERAL BASE FIELD CASE

From now on throughout the paper, let  $X$  be a proper smooth geometrically connected curve over a perfect field  $k$  of characteristic  $p > 0$ . Let  $K$  be its function field. In this and the next sections, we will formulate and prove a formula for the change in  $\mu$ -invariants under isogeny in terms of certain Euler characteristics in the fppf topology. This section treats the version of the statement for general, not necessarily finite base field  $k$ .

For an fppf sheaf  $\mathcal{F}$  on  $X$ , if  $\mathbf{H}^i(X, \mathcal{F}) \in \text{LAlg}/k$  for all  $i$  and  $\mathbf{H}^i(X, \mathcal{F}) = 0$  for almost all  $i$ , then we denote

$$\chi^0(X, \mathcal{F}) = \sum_i (-1)^i \dim \mathbf{H}^i(X, \mathcal{F}).$$

(The notation 0 indicates that only the identity components matter.) For an abelian variety  $A$  over  $K$  with Néron model  $\mathcal{A}$  over  $X$ , we have  $\mathbf{H}^i(X, \mathcal{A}) \in \text{LAlg}/k$  for all  $i$  and  $\mathbf{H}^i(X, \mathcal{A}) = 0$  for  $i \neq 0, 1, 2$  by [Suz20a, Theorem 3.4.1 (1)]. Define

$$(6.1) \quad \mu_{A/K} := \dim \mathbf{H}^1(X, \mathcal{A})$$

as in [LLS<sup>+</sup>21, Section 6.2].

**Proposition 6.1.** *For any quasi-finite flat separated group scheme  $\mathcal{N}$  over  $X$ , we have  $\mathbf{H}^i(X, \mathcal{N}) \in \text{Alg}_{\text{uc}}/k$  for all  $i$  and zero for almost all  $i$ .*

*Proof.* This is [Suz20a, Theorem 3.1.3] if  $\mathcal{N}$  is finite over  $X$ . In general, let  $N$  be the generic fiber of  $\mathcal{N}$ . Let  $N^0 \subset N$  be the identity component of  $N$ . Let  $\mathcal{N}^0 \subset \mathcal{N}$  be the schematic closure of  $N^0$  in  $\mathcal{N}$ . Then  $\mathcal{N}^0$  is finite over  $X$  since  $N^0$  is infinitesimal. Hence the finite case implies that we may assume  $N$  to be étale. Let  $U \subset X$  be a dense open subscheme where  $\mathcal{N}$  is étale. We have a distinguished triangle

$$R\Gamma_c(U, \mathcal{N}) \rightarrow R\Gamma(X, \mathcal{N}) \rightarrow \bigoplus_{v \in X \setminus U} R\Gamma(\mathcal{O}_v/k, \mathcal{N})$$

in  $D(k_{\text{proet}}^{\text{indrat}})$  by (3.1). The term  $R\Gamma_c(U, \mathcal{N})$  agrees with  $R\pi_! \mathcal{N}$  (where  $\pi: U_{\text{et}} \rightarrow \text{Spec } k_{\text{et}}$  is the structure morphism) and hence is a bounded complex of finite étale groups over  $k$  by a standard result on étale cohomology [Mil80, Chapter VI, Theorem 3.2 (d)]. For the third term for each  $v \in X \setminus U$ , let  $\mathcal{M} \subset \mathcal{N}$  be the finite part of  $\mathcal{N}$  over  $\mathcal{O}_v$ . Then we have

$$R\Gamma(\mathcal{O}_v/k, \mathcal{M}) \xrightarrow{\sim} R\Gamma(\mathcal{O}_v/k, \mathcal{N})$$

by [Suz20b, Proposition 5.2.3.5]. Since the generic fiber of  $\mathcal{M}$  is étale, we have  $R\Gamma(\mathcal{O}_v/k, \mathcal{M}) \in D^b(\text{Alg}_{\text{uc}}/k)$  by [OS23, Proposition 3.7].  $\square$

Here is the promised formula in this section:

**Theorem 6.2.** *Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over  $K$ . Then we have*

$$\mu_{A_2/K} - \mu_{A_1/K} = \chi^0(X, \mathcal{N}_f) - c(f).$$

*In particular, if  $f$  extends to an isogeny on the Néron models, then*

$$\mu_{A_2/K} - \mu_{A_1/K} = \chi^0(X, \mathcal{N}_f).$$

*Proof.* For each  $j = 1, 2$ , let  $\mathcal{A}_j$  be the Néron model of  $A_j$ . We first show that

$$(6.2) \quad \mu_{A_2/K} - \mu_{A_1/K} = \chi^0(X, \mathcal{A}_1) - \chi^0(X, \mathcal{A}_2).$$

The object  $\mathbf{H}^i(X, \mathcal{A}_j)$  is étale for  $i \neq 0, 1$  by [Suz20a, Theorem 3.4.1 (1) and (2)]. The object  $\mathbf{H}^0(X, \mathcal{A}_j)$  is an extension of an étale group with finitely generated group of geometric points by (the perfection of) an abelian variety over  $k$  by [Suz20a, Theorem 3.4.1 (2)]. Since  $A_1$  and  $A_2$  are isogenous, the objects  $\mathbf{H}^0(X, \mathcal{A}_1)$  and  $\mathbf{H}^0(X, \mathcal{A}_2)$  are isomorphic up to bounded torsion. Hence their abelian variety parts are isogenous. Combining these, we get (6.2).

Let  $\mathcal{A}'_2 = \mathcal{A}_1/\mathcal{N}_f$ . Then we have the diagram (5.1). Let  $\mathcal{D}_f = [\mathcal{A}'_2 \rightarrow \mathcal{A}_2]$  be the mapping cone of the morphism  $\mathcal{A}'_2 \rightarrow \mathcal{A}_2$ . We have distinguished triangles

$$(6.3) \quad R\Gamma(X, \mathcal{N}_f) \rightarrow R\Gamma(X, \mathcal{A}_1) \rightarrow R\Gamma(X, \mathcal{A}'_2),$$

$$(6.4) \quad R\Gamma(X, \mathcal{A}'_2) \rightarrow R\Gamma(X, \mathcal{A}_2) \rightarrow R\Gamma(X, \mathcal{D}_f)$$

in  $D(k_{\text{proet}}^{\text{indrat}})$ .

We will calculate  $R\Gamma(X, \mathcal{D}_f)$ . Let  $v \in X_0$ . By [Suz20b, Proposition 3.4.2 (a)] and its proof, we know that  $\mathbf{H}^i(\mathcal{O}_v, \mathcal{A}'_2) = 0$  for  $i \neq 0$  and  $\mathbf{H}^0(\mathcal{O}_v, \mathcal{A}'_2)$  is (the perfection of) the Greenberg transform  $\text{Gr}_{\mathcal{O}_v} \mathcal{A}'_2$  of  $\mathcal{A}'_2 \times_X \text{Spec } \mathcal{O}_v$ . The same is true with  $\mathcal{A}'_2$  replaced by  $\mathcal{A}_2$ . Hence the distinguished triangle

$$R\Gamma(\mathcal{O}_v/k, \mathcal{A}'_2) \rightarrow R\Gamma(\mathcal{O}_v/k, \mathcal{A}_2) \rightarrow R\Gamma(\mathcal{O}_v/k, \mathcal{D}_f)$$

reduces to the short exact sequence

$$0 \rightarrow \text{Res}_{k(v)/k} \text{Gr}_{\mathcal{O}_v} \mathcal{A}'_2 \rightarrow \text{Res}_{k(v)/k} \text{Gr}_{\mathcal{O}_v} \mathcal{A}_2 \rightarrow \text{Res}_{k(v)/k} \mathcal{D}_f \rightarrow 0$$

coming from (4.1), where  $\text{Res}_{k(v)/k}$  is the Weil restriction functor. In particular,  $\mathbf{H}^0(\mathcal{O}_v/k, \mathcal{D}_f) \in \text{Alg}_{\text{uc}}/k$  and

$$(6.5) \quad \dim \mathbf{H}^0(\mathcal{O}_v/k, \mathcal{D}_f) = [k(v) : k] \dim \mathcal{D}_f.$$

Let  $U \subset X$  be a dense open subscheme where  $\tilde{f}: \mathcal{A}_1 \rightarrow \mathcal{A}_2$  is an isogeny. Then the restriction of  $\mathcal{D}_f$  to  $U$  is zero by Proposition 5.2. Hence by (3.1), we have a commutative diagram of distinguished triangles

$$\begin{array}{ccccc} R\Gamma_c(U, \mathcal{A}'_2) & \xrightarrow{\sim} & R\Gamma_c(U, \mathcal{A}_2) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ R\Gamma(X, \mathcal{A}'_2) & \longrightarrow & R\Gamma(X, \mathcal{A}_2) & \longrightarrow & R\Gamma(X, \mathcal{D}_f) \\ \downarrow & & \downarrow & & \downarrow \wr \\ \bigoplus_{v \in X \setminus U} R\Gamma(\mathcal{O}_v/k, \mathcal{A}'_2) & \longrightarrow & \bigoplus_{v \in X \setminus U} R\Gamma(\mathcal{O}_v/k, \mathcal{A}_2) & \longrightarrow & \bigoplus_{v \in X \setminus U} R\Gamma(\mathcal{O}_v/k, \mathcal{D}_f). \end{array}$$

Hence  $R\Gamma(X, \mathcal{D}_f)$  has cohomology only in degree zero that is an object of  $\text{Alg}_{\text{uc}}/k$  and

$$(6.6) \quad \dim \mathbf{H}^0(X, \mathcal{D}_f) = \sum_{v \in X \setminus U} \dim \mathbf{H}^0(\mathcal{O}_v/k, \mathcal{D}_f) = c(f)$$

by (6.5).

With the properties of  $\mathbf{H}^i(X, \mathcal{A}_2)$  noted before Proposition 6.1, we know that  $\mathbf{H}^i(X, \mathcal{A}'_2)$  is in  $\text{LAlg}/k$  for all  $i$  and zero for almost all  $i$ . The distinguished triangle (6.4) and the equality (6.6) then imply

$$(6.7) \quad \chi^0(X, \mathcal{A}_2) = \chi^0(X, \mathcal{A}'_2) + \dim \mathbf{H}^0(X, \mathcal{D}_f) = \chi^0(X, \mathcal{A}'_2) + c(f).$$

The distinguished triangle (6.3) implies

$$(6.8) \quad \chi^0(X, \mathcal{A}_1) = \chi^0(X, \mathcal{N}_f) + \chi^0(X, \mathcal{A}'_2).$$

Combining (6.2), (6.7) and (6.8), we get the result.  $\square$

## 7. CHANGE IN $\mu$ BY FLAT EULER CHARACTERISTICS: FINITE BASE FIELD CASE

Assume that  $k = \mathbb{F}_q$  is a finite field in this section. We will deduce a formula from Theorem 6.2 specific to the finite base field.

Let  $A$  be an abelian variety over  $K$ . Let  $\mu_{A/K}$  be the  $\mu$ -invariant of the Pontryagin dual of the Selmer group of  $A$  over the unramified  $\mathbb{Z}_p$ -extension of  $K$  as defined in [LLS<sup>+</sup>21, Definition 2.0.1] (which is the  $\mu$ -invariant in the usual sense divided by  $[k : \mathbb{F}_p]$ ). This is consistent with the previous definition of  $\mu_{A/K}$  in (6.1) by [LLS<sup>+</sup>21, Theorem 2.4.3].

For an fppf sheaf  $\mathcal{F}$  on  $X$ , let

$$\chi(X, \mathcal{F}) = \prod_i (\#H^i(X, \mathcal{F}))^{(-1)^i}$$

if  $H^i(X, \mathcal{F})$  is finite for all  $i$  and zero for almost all  $i$ . The following proposition generalizes [Mil06, Chapter III, Lemma 8.9] on finite flat group schemes to quasi-finite ones:

**Proposition 7.1.** *For any quasi-finite flat separated group scheme  $\mathcal{N}$  over  $X$ , the groups  $H^i(X, \mathcal{N})$  are finite for all  $i$  and zero for almost all  $i$ .*

*Proof.* This follows from Proposition 6.1 by applying Proposition 3.1 (1) to  $G = R\Gamma(X, \mathcal{N})$ .  $\square$

**Theorem 7.2.** *Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over  $K$ . Then we have*

$$\mu_{A_2/K} - \mu_{A_1/K} = \text{ord}_q(\chi(X, \mathcal{N}_f)) - c(f).$$

*In particular, if  $f$  extends to an isogeny on the Néron models, then*

$$\mu_{A_2/K} - \mu_{A_1/K} = \text{ord}_q(\chi(X, \mathcal{N}_f)).$$

*Proof.* This follows from Theorem 6.2 by applying Proposition 3.1 (2) to  $G = R\Gamma(X, \mathcal{N}_f)$ .  $\square$

## 8. LIE COMPLEXES

Our next task is to deduce “linear” (or coherent) versions of Theorems 6.2 and 7.2, which will be completed in Section 10. As preparations, in this section, we will define the linearized terms needed for this, which are derived versions of Lie algebras (relative tangent spaces at the zero section).

Let  $U \subset X$  be a dense open subscheme. For a flat group scheme  $G$  locally of finite type over  $U$ , we denote the Lie complex  $l_G^\vee$  ([Ill72, Chapter VII, Section 3.1.1]) of  $G$  by  $R\mathrm{Lie} G$ :

$$R\mathrm{Lie} G := l_G^\vee \in D^b(\mathcal{O}_U).$$

It is (represented by) a perfect complex of  $\mathcal{O}_U$ -modules with tor-amplitude in  $[0, 1]$  ([Ill72, Chapter VII, (3.1.1.3)'] whose zero-th cohomology is the usual Lie algebra  $\mathrm{Lie} G$  of  $G$ . If  $0 \rightarrow G' \rightarrow G \rightarrow G'' \rightarrow 0$  is an exact sequence of flat group schemes locally of finite type over  $U$ , then we have a canonical distinguished triangle

$$R\mathrm{Lie} G' \rightarrow R\mathrm{Lie} G \rightarrow R\mathrm{Lie} G''$$

by [Ill72, Chapter VII, Proposition 3.1.1.5]. If  $G$  is smooth over  $U$ , then  $R\mathrm{Lie} G$  is the usual Lie algebra  $\mathrm{Lie} G$  of  $G$ . Hence if a flat group scheme  $N$  locally of finite type over  $U$  can be embedded into a smooth group scheme  $G$  as a closed group subscheme with quotient  $H = G/N$ , then

$$(8.1) \quad R\mathrm{Lie} N \cong [\mathrm{Lie} G \rightarrow \mathrm{Lie} H][-1].$$

Such an embedding always exists if we restrict all the objects to  $K$  by [DG70, Chapter III, Section 5, Lemma 6.5] or if  $N$  is finite by [Mil06, Chapter III, Theorem A.5]. Hence in these case, one may take this formula as the definition of  $R\mathrm{Lie} N$ .

This object  $R\mathrm{Lie} N$  is particularly simple if  $N$  has étale generic fiber:

**Proposition 8.1.** *Let  $\mathcal{N}$  be a quasi-finite flat separated group scheme over  $U$  with étale generic fiber. Let  $e: U \rightarrow \mathcal{N}$  be its zero section. Then the  $\mathcal{O}_U$ -dual of  $R\mathrm{Lie} \mathcal{N}$  is isomorphic to  $e^* \Omega_{\mathcal{N}/U}^1$  placed in degree zero.*

*Proof.* The  $H^{-1}$  of the  $\mathcal{O}_U$ -dual of  $R\mathrm{Lie} \mathcal{N}$  is torsion-free by the tor-amplitude condition. It is torsion by the étaleness assumption. Hence this  $H^{-1}$  is zero. The description of  $H^0$  directly follows from the definition.  $\square$

For a perfect complex of  $\mathcal{O}_U$ -modules  $\mathcal{F}$ , let  $\det \mathcal{F}$  be its determinant invertible sheaf ([Sta24, Tag 0FJW]). When  $U = X$ , let  $\deg \mathcal{F} := \deg \det \mathcal{F}$ , where  $\deg$  on the right-hand side is with respect to  $k$ . Let  $\chi(X, \mathcal{F})$  be the coherent Euler characteristic with respect to the base field  $k$ . Let  $g: \mathrm{Spec} K \rightarrow X$  be the natural morphism. Then

$$\chi(X, \mathcal{F}) = \chi(K, g^* \mathcal{F}) \chi(X, \mathcal{O}_X) + \deg \mathcal{F}$$

by the Riemann–Roch theorem, where  $\chi(K, \cdot)$  is with respect to  $K$ .

**Proposition 8.2.** *Let  $\mathcal{N}$  be a quasi-finite flat separated group scheme over  $X$ . Then*

$$\chi(X, R\mathrm{Lie} \mathcal{N}) = \deg R\mathrm{Lie} \mathcal{N}.$$

*Proof.* Let  $N$  be the generic fiber of  $\mathcal{N}$ . It is enough to show that  $\chi(K, R\mathrm{Lie} N) = 0$ . This follows from (8.1) since  $\dim G = \dim H$  in this case.  $\square$

9. DUALITY OF EULER CHARACTERISTICS

We need duality results for  $\deg R\mathrm{Lie} \mathcal{N}$  and  $\chi^0(X, \mathcal{N})$ .

**Proposition 9.1.** *Let  $U \subset X$  be a dense open subscheme. Let  $N$  and  $M$  be finite flat group schemes over  $U$  Cartier dual to each other. Then there exists a canonical isomorphism  $\det R\mathrm{Lie} N \cong (\det R\mathrm{Lie} M)^{\otimes -1}$  of  $\mathcal{O}_U$ -modules.*

*Proof.* First assume that  $N$  can be embedded into an abelian scheme as a closed group subscheme. Then we have a resolution  $0 \rightarrow N \rightarrow A_1 \rightarrow A_2 \rightarrow 0$  by abelian schemes over  $U$ . Denote its dual ([Oor66, Theorem (19.1)]) by  $0 \rightarrow M \rightarrow B_2 \rightarrow B_1 \rightarrow 0$ . Let  $d$  be the relative dimension of  $A_1$  (or  $A_2$ ). Let  $\pi_i: A_i \rightarrow U$  be the structure morphism. Since  $\mathrm{Lie} B_i \cong R^1\pi_{i,*}\mathcal{O}_{A_i}$  and by [Ser88, Chapter VII, Section 4.21, Theorem 10], we have

$$\det \mathrm{Lie} B_i \cong \det R^1\pi_{i,*}\mathcal{O}_{A_i} \cong R^d\pi_{i,*}\mathcal{O}_{A_i},$$

which is dual to  $\pi_{i,*}\Omega_{A_i}^d$  by Serre duality. This sheaf  $\pi_{i,*}\Omega_{A_i}^d$  is dual to  $\det \mathrm{Lie} A_i$ . Hence  $\det \mathrm{Lie} B_i \cong \det \mathrm{Lie} A_i$  canonically. Therefore

$$(9.1) \quad \begin{aligned} \det R\mathrm{Lie} N &\cong \det \mathrm{Lie} A_1 \otimes (\det \mathrm{Lie} A_2)^{\otimes -1} \\ &\cong \det \mathrm{Lie} B_1 \otimes (\det \mathrm{Lie} B_2)^{\otimes -1} \cong (\det R\mathrm{Lie} M)^{\otimes -1}. \end{aligned}$$

If we have another resolution  $0 \rightarrow N \rightarrow A'_1 \rightarrow A'_2 \rightarrow 0$  and a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & N & \longrightarrow & A'_1 & \longrightarrow & A'_2 & \longrightarrow & 0 \\ & & & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & N & \longrightarrow & A_1 & \longrightarrow & A_2 & \longrightarrow & 0 \end{array}$$

with  $A'_1 \rightarrow A_1$  smooth with connected fibers, then the kernel of  $A'_1 \rightarrow A_1$  and  $A'_2 \rightarrow A_2$  are isomorphic. Hence the resulting isomorphism

$$(9.2) \quad \begin{aligned} \det R\mathrm{Lie} N &\cong \det \mathrm{Lie} A'_1 \otimes (\det \mathrm{Lie} A'_2)^{\otimes -1} \\ &\cong \det \mathrm{Lie} B'_1 \otimes (\det \mathrm{Lie} B'_2)^{\otimes -1} \cong (\det R\mathrm{Lie} M)^{\otimes -1}. \end{aligned}$$

is compatible with (9.1). If we just have another resolution  $0 \rightarrow N \rightarrow A'_1 \rightarrow A'_2 \rightarrow 0$  (no commutative diagram), then take  $A'_1 = A_1 \times A'_1$  and the diagonal embedding  $N \hookrightarrow A'_1$ , defining  $A'_2 = A'_1/N$ . Then we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & N & \longrightarrow & A'_1 & \longrightarrow & A'_2 & \longrightarrow & 0 \\ & & & & \parallel & & \uparrow & & \uparrow \\ 0 & \longrightarrow & N & \longrightarrow & A''_1 & \longrightarrow & A''_2 & \longrightarrow & 0 \\ & & & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & N & \longrightarrow & A_1 & \longrightarrow & A_2 & \longrightarrow & 0 \end{array}$$

Hence (9.2) is still compatible with (9.1). This means that we have the required canonical isomorphism in this (embeddable) case.

In general, such an embedding exists Zariski locally on  $U$  by [BBM82, Theorem 3.1.1]. Now the canonicity of the isomorphism in the embeddable case proved right above allows us to patch the local isomorphisms.  $\square$

**Proposition 9.2.** *Let  $N$  and  $M$  be finite flat group schemes over  $X$  Cartier dual to each other. Then*

$$\deg R\mathrm{Lie} N = -\deg R\mathrm{Lie} M.$$

*Proof.* This follows from Propositions 9.1 and 8.2.  $\square$

**Proposition 9.3.** *Let  $N$  and  $M$  be finite flat group schemes over  $X$  Cartier dual to each other. Then*

$$\chi^0(X, N) = -\chi^0(X, M).$$

*Proof.* This follows from [Suz20a, Theorem 3.1.3].  $\square$

## 10. CHANGE IN $\mu$ BY DEGREES OF LIE COMPLEXES

Now we can interpret  $\chi(X, \mathcal{N})$  in terms of  $\deg R\mathrm{Lie} \mathcal{N}$ , which gives a linearized version Theorems 6.2 and 7.2. This is a function field version of Schneider's formula. This interpretation also answers Milne's question.

**Theorem 10.1.** *Let  $\mathcal{N}$  be a quasi-finite flat separated group scheme over  $X$ . Then*

$$\chi^0(X, \mathcal{N}) = \deg R\mathrm{Lie} \mathcal{N}.$$

*Proof.* We may assume  $\mathcal{N}$  is killed by a power of  $p$ . For a vector bundle  $G$  on  $X$  (viewed as a vector group), we have

$$R\Gamma(X, G) \cong R\Gamma(X, G) \otimes_k \mathbf{G}_a$$

with respect to the natural  $k$ -linear structures on  $G$  and  $\mathbf{G}_a$ . Hence

$$(10.1) \quad \chi^0(X, G) = \chi(X, \mathrm{Lie} G).$$

Assume that  $\mathcal{N}$  is finite flat with height one Cartier dual. Then we have an exact sequence  $0 \rightarrow \mathcal{N} \rightarrow G \rightarrow H \rightarrow 0$  over  $X$  with  $G$  and  $H$  vector groups by [Mil06, Chapter III, Section 6]. Hence (10.1) and Proposition 8.2 give the result in this case.

Assume next that  $\mathcal{N}$  has étale generic fiber. Let  $U \subset X$  be dense open where  $\mathcal{N}$  is étale. Then  $R\Gamma_c(U, \mathcal{N})$  has finite étale cohomologies as in the proof of Proposition 6.1 and  $R\Gamma_c(U, R\mathrm{Lie} \mathcal{N}) = 0$ . Hence it is enough to show that

$$\chi^0(\mathcal{O}_v, \mathcal{N}) = \chi(\mathcal{O}_v, R\mathrm{Lie} \mathcal{N})$$

for each  $v \in X \setminus U$  (where the left-hand side is the alternating sum of the dimensions of  $\mathbf{H}^i(\mathcal{O}_v, \mathcal{N})$  and the right-hand side is the coherent Euler characteristic with respect to  $k(v)$ ). This equality follows from

$$\chi^0(\mathcal{O}_v, \mathcal{N}) = -\dim \mathbf{H}^1(\mathcal{O}_v, \mathcal{N}) = -\mathrm{length}_{\mathcal{O}_v} e^* \Omega_{\mathcal{N}/\mathcal{O}_v}^1 = \chi(\mathcal{O}_v, R\mathrm{Lie} \mathcal{N})$$

by [OS23, Proposition 3.7] and Proposition 8.1, where  $e$  denotes the zero section for  $\mathcal{N}$ .

In general, let  $N$  be the generic fiber of  $\mathcal{N}$ . Let

$$0 \subset N^m \subset N^0 \subset N$$

be the multiplicative part and the identity component, respectively. Let

$$0 \subset \mathcal{N}^m \subset \mathcal{N}^0 \subset \mathcal{N}$$

be their schematic closures in  $\mathcal{N}$ . Then  $\mathcal{N}^m$  and  $\mathcal{N}^0$  are finite flat, and the subquotients of this filtration are all quasi-finite flat separated. We will check the statement for each subquotient. The statement for  $\mathcal{N}^m$  reduces to the statement

for its Cartier dual by Propositions 9.2 and 9.3, which has étale generic fiber. Hence this case is done. For the statement for  $\mathcal{N}^0/\mathcal{N}^m$ , its Cartier dual is infinitesimal, so it has a filtration with height one subquotients, which is done. The quotient  $\mathcal{N}/\mathcal{N}^0$  has étale generic fiber, so it is done.  $\square$

The following answers [Mil06, Chapter III, Problem 8.10].

**Theorem 10.2.** *Assume  $k = \mathbb{F}_q$  is finite. Let  $\mathcal{N}$  be a quasi-finite flat separated group scheme over  $X$ . Then*

$$\chi(X, \mathcal{N}) = q^{\deg R \operatorname{Lie} \mathcal{N}}.$$

*Proof.* This follows from Theorem 10.1 and Proposition 3.1 (2).  $\square$

The following gives a function field version of the ‘‘Isogeny formula (First form)’’ at the end of [Sch87].

**Theorem 10.3.** *Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over  $K$ . Then we have*

$$\mu_{A_2/K} - \mu_{A_1/K} = \deg R \operatorname{Lie} \mathcal{N}_f - c(f).$$

*In particular, if  $f$  extends to an isogeny on the Néron models, then*

$$\mu_{A_2/K} - \mu_{A_1/K} = \deg R \operatorname{Lie} \mathcal{N}_f.$$

*Proof.* This follows from Theorems 10.1 and 6.2.  $\square$

## 11. ISOGENY INVARIANCE OF THE BSD FORMULA

As an application of our methods and Theorem 10.2, we will prove that the validity of the Birch–Swinnerton-Dyer conjecture for abelian varieties over function fields is invariant under arbitrary isogeny, whose degree may be divisible by the characteristic  $p$ .

Assume that  $k = \mathbb{F}_q$  is finite in this section. Let  $A/K$  be an abelian variety with Néron model  $\mathcal{A}/X$ . Let  $r$  be the rank of  $A(K)$ . Recall the Weil-étale BSD formula from [GS20, Theorem 1.1, Proposition 8.4]:

$$(11.1) \quad \lim_{t \rightarrow q^{-1}} \frac{L(A, t)}{(1 - qt)^r} = \frac{q^{\chi(X, \operatorname{Lie} \mathcal{A})}}{\chi_W(X, \mathcal{A})},$$

where  $L(A, t)$  is the Hasse–Weil  $L$ -function of  $A/K$  as a rational function of  $t$ , and we assume that  $\text{III}(A)$  is finite so that the Weil-étale Euler characteristic  $\chi_W(X, \mathcal{A})$  is finite. This formula (under the assumption of the finiteness of  $\text{III}(A)$ ) is proved by [KT03]. Here we prove, without using the result of [KT03], that the validity of this formula is invariant under arbitrary isogeny on  $A$ . This is [Mil06, Chapter III, Problem 9.10].

**Theorem 11.1.** *Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over  $K$ . Assume that  $\text{III}(A_1)$  and  $\text{III}(A_2)$  are finite. Then the formula (11.1) holds for  $A_1$  if and only if the formula (11.1) holds for  $A_2$ .*

*Proof.* We need to show that

$$\frac{\chi_W(X, \mathcal{A}_2)}{\chi_W(X, \mathcal{A}_1)} = q^{\chi(X, \operatorname{Lie} \mathcal{A}_2) - \chi(X, \operatorname{Lie} \mathcal{A}_1)}.$$

Let  $\mathcal{N}_f$  and  $D_f$  as before. Then

$$\chi_W(X, \mathcal{A}_1) = \chi(X, \mathcal{N}_f) \chi_W(X, \mathcal{A}'_2).$$

Applying the Weil-étale cohomology functor  $R\Gamma_W(k, \cdot)$  to (6.4), we have a distinguished triangle

$$R\Gamma_W(X, \mathcal{A}'_2) \rightarrow R\Gamma_W(X, \mathcal{A}_2) \rightarrow \bigoplus_v R\Gamma(k(v), D_f).$$

Hence

$$\chi_W(X, \mathcal{A}_2) = \chi_W(X, \mathcal{A}'_2) \prod_v \chi(k(v), D_f).$$

Arguing similarly, we have

$$(11.2) \quad \chi(X, \text{Lie } \mathcal{A}_1) = \chi(X, R\text{Lie } \mathcal{N}_f) + \chi(X, \text{Lie } \mathcal{A}'_2),$$

$$(11.3) \quad \chi(X, \text{Lie } \mathcal{A}_2) = \chi(X, \text{Lie } \mathcal{A}'_2) + \sum_v [k(v) : k] \chi(k(v), \text{Lie } D_f),$$

where  $\chi(k(v), \cdot)$  is with respect to  $k(v)$ . We have

$$\chi(k(v), D_f) = q_v^{\chi(k(v), \text{Lie } D_f)}$$

for each  $v$  by Proposition 3.1 (2) (where  $q_v = \#k(v)$ ). We have

$$\chi(X, \mathcal{N}_f) = q^{\deg R\text{Lie } \mathcal{N}_f} = q^{\chi(X, R\text{Lie } \mathcal{N}_f)}$$

by Theorem 10.2 and Proposition 8.2.  $\square$

## 12. BASE CHANGE STABLE INVARIANTS

The term  $\deg R\text{Lie } \mathcal{N}_f$  in Theorem 10.3 changes in a complicated manner if  $K$  is replaced by a finite extension. We can replace it by a more stable invariant at the expense of adding and subtracting base change conductors of the abelian varieties (Theorem 13.1). We first need to define this stable invariant in this section.

Let  $X^{\text{sep}}$  be the normalization of  $X$  in the separable closure  $K^{\text{sep}}$  of  $K$ . Let  $\mathcal{F}_{K^{\text{sep}}} \in D^b(\mathcal{O}_{X^{\text{sep}}})$  be (represented by) a perfect complex of  $\mathcal{O}_{X^{\text{sep}}}$ -modules. Choose a finite Galois extension  $L/K$  such that  $\mathcal{F}_{K^{\text{sep}}}$  can be obtained as a derived pullback of a perfect complex  $\mathcal{F}_L$  of  $\mathcal{O}_Y$ -modules, where  $Y$  is the normalization of  $X$  in  $L$ .

**Definition 12.1.** *Define a rational number by*

$$\deg_X \mathcal{F}_{K^{\text{sep}}} := \frac{\deg \mathcal{F}_L}{[L : K]},$$

where the numerator on the right-hand side is the degree with respect to  $k$ .

**Proposition 12.2.** *This definition is independent of the choice of  $L$  or  $\mathcal{F}_L$ .*

*Proof.* This reduces to the equality

$$\deg \mathcal{O}_X(v) = \frac{\deg \mathcal{O}_Y(f^{-1}v)}{[L : K]} \quad (= [k(v) : k]),$$

where  $v$  is a closed point of  $X$  and  $f: Y \rightarrow X$  is the structure morphism.  $\square$

On the other hand, if  $\mathcal{F}$  is a perfect complex of  $\mathcal{O}_X$ -modules, then we denote its base change to  $X^{\text{sep}}$  by  $\mathcal{F}_{X^{\text{sep}}}$ . We have  $\deg \mathcal{F} = \deg_X \mathcal{F}_{X^{\text{sep}}}$  by Proposition 12.2.

We can rewrite base change conductors using this  $\deg_X$  construction. Let  $A$  be an abelian variety over  $K$  with Néron model  $\mathcal{A}$  over  $X$ . If  $A$  has semistable



reduction over a finite separable extension  $L/K$  and  $\mathcal{A}_L$  denotes the Néron model of  $A \times_K L$ , then  $\mathrm{Lie} \mathcal{A}_L$  commutes with base change:

$$g^* \mathrm{Lie} \mathcal{A}_L \xrightarrow{\sim} \mathrm{Lie} \mathcal{A}_{L'},$$

where  $L'/L$  is a finite separable extension and  $g: Y' \rightarrow Y$  the morphism on the normalizations of  $X$  in  $L$  and  $L'$ . Hence we may define  $\mathrm{Lie} \mathcal{A}_{K^{\mathrm{sep}}}$  to be the base change of  $\mathrm{Lie} \mathcal{A}_L$  to  $X^{\mathrm{sep}}$  for any choice of such  $L$ . It is a locally free sheaf on  $X^{\mathrm{sep}}$  of finite rank.

On the other hand, we have the base change  $\mathrm{Lie} \mathcal{A}_{X^{\mathrm{sep}}}$  of  $\mathrm{Lie} \mathcal{A}$  to  $X^{\mathrm{sep}}$ .

**Proposition 12.3.** *We have*

$$c(A) = \deg_X \mathrm{Lie} \mathcal{A}_{K^{\mathrm{sep}}} - \deg_X \mathrm{Lie} \mathcal{A}_{X^{\mathrm{sep}}}.$$

*Proof.* Let  $L/K$  be a finite Galois extension over which  $A$  has semistable reduction everywhere. Let  $Y$  be the normalization of  $X$  in  $L$ . Let  $k_L$  be the constant field of  $Y$ . We have

$$\begin{aligned} & \deg_X \mathrm{Lie} \mathcal{A}_{K^{\mathrm{sep}}} - \deg_X \mathrm{Lie} \mathcal{A}_{X^{\mathrm{sep}}}. \\ &= \sum_{w \in Y_0} \frac{[k_L(w) : k]}{[L : K]} \mathrm{length}_{\mathcal{O}_w} \frac{\mathrm{Lie} \mathcal{A}_L}{\mathrm{Lie} \mathcal{A} \otimes_{\mathcal{O}_X} \mathcal{O}_Y} \\ &= \sum_{v \in X_0} \frac{[k(v) : k]}{e_v} \mathrm{length}_{\mathcal{O}_w} \frac{\mathrm{Lie} \mathcal{A}_L}{\mathrm{Lie} \mathcal{A} \otimes_{\mathcal{O}_X} \mathcal{O}_Y}, \end{aligned}$$

where, in the third line,  $w$  is any point of  $Y$  above  $v$  that we fix for each  $v$  and  $e_v$  is the ramification index of  $L/K$  at  $v$ . The quantity in the third line is equal to the definition of  $c(A)$ .  $\square$

We stabilize  $R\mathrm{Lie} \mathcal{N}_f$  as follows. Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over  $K$ . Let  $L$  be a finite separable extension of  $K$  over which  $A_1$  has semistable reduction. Let  $\mathcal{A}_{i,L}$  be the Néron model of  $A_i \times_K L$ . Then the induced morphism  $\tilde{f}_L: \mathcal{A}_{1,L} \rightarrow \mathcal{A}_{2,L}$  is an isogeny. Let  $\mathcal{N}_{f,L}$  be its kernel. Then  $R\mathrm{Lie} \mathcal{N}_{f,L}$  commutes with base change:

$$g^* R\mathrm{Lie} \mathcal{N}_{f,L} \xrightarrow{\sim} R\mathrm{Lie} \mathcal{N}_{f,L'},$$

where  $L'/L$  is a finite separable extension and  $g: Y' \rightarrow Y$  the morphism on the normalizations of  $X$  in  $L$  and  $L'$ . Hence we may define  $R\mathrm{Lie} \mathcal{N}_{f,K^{\mathrm{sep}}}$  to be the base change of  $R\mathrm{Lie} \mathcal{N}_{f,L}$  to  $X^{\mathrm{sep}}$  for any choice of such  $L$ . It is a perfect complex of  $\mathcal{O}_{X^{\mathrm{sep}}}$ -modules.

### 13. CHANGE IN $\mu$ WITH BASE CHANGE CONDUCTORS

With these stabilized objects, we can give a stabilized version of Theorem 10.3:

**Theorem 13.1.** *Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over  $K$ . Then*

$$\mu_{A_2/K} - \mu_{A_1/K} = \deg_X R\mathrm{Lie} \mathcal{N}_{f,K^{\mathrm{sep}}} + c(A_2) - c(A_1).$$

*Proof.* By Theorem 10.3, it is enough to show that

$$\deg R\mathrm{Lie} \mathcal{N}_f - c(f) = \deg_X R\mathrm{Lie} \mathcal{N}_{f,K^{\mathrm{sep}}} + c(A_2) - c(A_1).$$

We have a morphism of distinguished triangles

$$\begin{array}{ccccc} R\mathrm{Lie}\mathcal{N}_{f,X^{\mathrm{sep}}} & \longrightarrow & \mathrm{Lie}\mathcal{A}_{1,X^{\mathrm{sep}}} & \longrightarrow & \mathrm{Lie}\mathcal{A}'_{2,X^{\mathrm{sep}}} \\ \downarrow & & \downarrow & & \downarrow \\ R\mathrm{Lie}\mathcal{N}_{f,K^{\mathrm{sep}}} & \longrightarrow & \mathrm{Lie}\mathcal{A}_{1,K^{\mathrm{sep}}} & \longrightarrow & \mathrm{Lie}\mathcal{A}_{2,K^{\mathrm{sep}}} \end{array}$$

of perfect complexes of  $\mathcal{O}_{X^{\mathrm{sep}}}$ -modules. Taking  $\deg_X$ , we have

$$\begin{aligned} & \deg_X R\mathrm{Lie}\mathcal{N}_{f,K^{\mathrm{sep}}} - \deg_X R\mathrm{Lie}\mathcal{N}_{f,X^{\mathrm{sep}}} \\ &= \deg_X \mathrm{Lie}\mathcal{A}_{1,K^{\mathrm{sep}}} - \deg_X \mathrm{Lie}\mathcal{A}_{1,X^{\mathrm{sep}}} - \deg_X \mathrm{Lie}\mathcal{A}_{2,K^{\mathrm{sep}}} + \deg_X \mathrm{Lie}\mathcal{A}'_{2,X^{\mathrm{sep}}}. \end{aligned}$$

We have

$$\deg_X \mathrm{Lie}\mathcal{A}_{1,K^{\mathrm{sep}}} - \deg_X \mathrm{Lie}\mathcal{A}_{1,X^{\mathrm{sep}}} = c(A_1)$$

by Proposition 12.3. On the other hand,

$$\begin{aligned} & \deg_X \mathrm{Lie}\mathcal{A}_{2,K^{\mathrm{sep}}} - \deg_X \mathrm{Lie}\mathcal{A}'_{2,X^{\mathrm{sep}}} \\ &= c(A_2) + \deg_X \mathrm{Lie}\mathcal{A}_{2,X^{\mathrm{sep}}} - \deg_X \mathrm{Lie}\mathcal{A}'_{2,X^{\mathrm{sep}}}. \\ &= c(A_2) + c(f). \end{aligned}$$

Combining these, we get the result.  $\square$

#### 14. ANOTHER PROOF OF THEOREM 10.3 BY THE $\mu$ -INVARIANT FORMULA

In this section, we will give another proof of Theorem 10.3 using the  $\mu$ -invariant formula in [LLS<sup>+</sup>21]. This is a simpler but conditional proof.

Assume  $k = \mathbb{F}_q$  is finite in this section. Let  $A$  be an abelian variety over  $K$ . Write the  $L$ -function  $L(A, t)$  of  $A$  as

$$L(A, t) = \frac{P_1(t)}{P_0(t)P_2(t)},$$

where  $P_i(t) \in \mathbb{Z}[t]$  is a polynomial in  $t$  with constant term 1 whose reciprocal roots are Weil  $q$ -numbers of weight  $i + 1$ . See [LLS<sup>+</sup>21, Section 2.2]. Define  $\theta_A$  to be the non-negative rational number such that  $q^{\theta_A} P_1(t/q)$  is  $p$ -primitive (namely, all the coefficients are  $p$ -adically integral and some coefficients are  $p$ -adic units). (It is actually an integer by [LLS<sup>+</sup>21, Proposition A.3].) Recall the  $\mu$ -invariant formula of [LLS<sup>+</sup>21] in the form stated in [LLS<sup>+</sup>21, Remark 2.5.5]:

$$(14.1) \quad \theta_A = \chi^0(X, \mathcal{A}) - \chi(X, \mathrm{Lie}\mathcal{A}).$$

This is expected to be true for all  $A$ . So far it is verified in the following cases:

- $\mathrm{III}(A/K\mathbb{F}_{q^{p^n}})$  is finite for all  $n$  ([LLS<sup>+</sup>21, Corollary 2.5.1]).
- $A$  is the Jacobian of a proper smooth geometrically connected curve over  $K$  ([LLS<sup>+</sup>21, Corollary 3.4.1]).
- $A$  has semistable reduction everywhere ([LLS<sup>+</sup>21, Theorem 4.1.1]).

Let  $f: A_1 \rightarrow A_2$  be an isogeny of abelian varieties over  $K$ . Assume that  $A_1$  and  $A_2$  satisfy the formula (14.1). We will give a simpler proof of Theorem 10.3 under this assumption.

*Another proof of Theorem 10.3.* We have  $\theta_{A_1} = \theta_{A_2}$  since  $L$ -functions are isogeny invariant. By (6.2) and (14.1), we are reduced to showing that

$$\chi(X, \mathrm{Lie}\mathcal{A}_1) - \chi(X, \mathrm{Lie}\mathcal{A}_2) = \deg R\mathrm{Lie}\mathcal{N}_f - c(f).$$

But this follows from (11.2) and (11.3).  $\square$

*Remark 14.1.* Without assuming  $k$  to be finite, we can still define the number  $\theta_A$  to be the sum of slopes less than one of the  $F$ -isocrystal  $\mathcal{H}_{\mathbb{Q}_p}^1(-1)$  over  $k$  given in [LLS<sup>+</sup>21, Appendix] by [LLS<sup>+</sup>21, Proposition A.1]. The formula (14.1) is expected to be true for general  $k$  with this definition of  $\theta_A$ . It implies Theorem 10.3 by the same proof, with “ $L$ -functions” replaced by “ $F$ -isocrystals”.

## REFERENCES

- [Ana73] Sivaramakrishna Anantharaman. Schémas en groupes, espaces homogènes et espaces algébriques sur une base de dimension 1. In *Sur les groupes algébriques*, Supplément au Bull. Soc. Math. France, Tome 101, pages 5–79. Soc. Math. France, Paris, 1973.
- [BBM82] Pierre Berthelot, Lawrence Breen, and William Messing. *Théorie de Dieudonné cristalline. II*, volume 930 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1982.
- [BGA18] Alessandra Bertapelle and Cristian D. González-Avilés. The Greenberg functor revisited. *Eur. J. Math.*, 4(4):1340–1389, 2018.
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud. *Néron models*, volume 21 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1990.
- [Cha00] Ching-Li Chai. Néron models for semiabelian varieties: congruence and change of base field. *Asian J. Math.*, 4(4):715–736, 2000.
- [DG70] Michel Demazure and Pierre Gabriel. *Groupes algébriques. Tome I: Géométrie algébrique, généralités, groupes commutatifs*. Masson & Cie, Éditeur, Paris; North-Holland Publishing Co., Amsterdam, 1970. Avec un appendice *Corps de classes local* par Michiel Hazewinkel.
- [DH19] Cyril Demarche and David Harari. Artin-Mazur-Milne duality for fppf cohomology. *Algebra Number Theory*, 13(10):2323–2357, 2019.
- [GS20] Thomas H. Geisser and Takashi Suzuki. A Weil-étale version of the Birch and Swinnerton-Dyer formula over function fields. *J. Number Theory*, 208:367–389, 2020.
- [Ill72] Luc Illusie. *Complexe cotangent et déformations. II*. Lecture Notes in Mathematics, Vol. 283. Springer-Verlag, Berlin-New York, 1972.
- [KT03] Kazuya Kato and Fabien Trihan. On the conjectures of Birch and Swinnerton-Dyer in characteristic  $p > 0$ . *Invent. Math.*, 153(3):537–592, 2003.
- [LLR04] Qing Liu, Dino Lorenzini, and Michel Raynaud. Néron models, Lie algebras, and reduction of curves of genus one. *Invent. Math.*, 157(3):455–518, 2004.
- [LLS<sup>+</sup>21] King-Fai Lai, Ignazio Longhi, Takashi Suzuki, Ki-Seng Tan, and Fabien Trihan. On the  $\mu$ -invariants of abelian varieties over function fields of positive characteristic. *Algebra Number Theory*, 15(4):863–907, 2021.
- [Mil80] James S. Milne. *Étale cohomology*, volume 33 of *Princeton Mathematical Series*. Princeton University Press, Princeton, N.J., 1980.
- [Mil06] James S. Milne. *Arithmetic duality theorems*. BookSurge, LLC, Charleston, SC, second edition, 2006.
- [Oor66] Frans Oort. *Commutative group schemes*, volume 15 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1966.
- [OS23] Otto Overkamp and Takashi Suzuki. Chai’s conjectures on base change conductors. Preprint: arXiv:2310.01289v1, 2023.
- [OT09] Tadashi Ochiai and Fabien Trihan. On the Selmer groups of abelian varieties over function fields of characteristic  $p > 0$ . *Math. Proc. Cambridge Philos. Soc.*, 146(1):23–43, 2009.
- [Sch87] Peter Schneider. The  $\mu$ -invariant of isogenies. *J. Indian Math. Soc. (N.S.)*, 52:159–170 (1988), 1987.
- [Ser88] Jean-Pierre Serre. *Algebraic groups and class fields*, volume 117 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1988. Translated from the French.
- [Sta24] The Stacks Project Authors. *Stacks Project*. <http://stacks.math.columbia.edu>, 2024.
- [Suz20a] Takashi Suzuki. Duality for cohomology of curves with coefficients in abelian varieties. *Nagoya Math. J.*, 240:42–149, 2020.

[Suz20b] Takashi Suzuki. Grothendieck's pairing on Néron component groups: Galois descent from the semistable case. *Kyoto J. Math.*, 60(2):593–716, 2020.

HARISH CHANDRA RESEARCH INSTITUTE, A CI OF HOMI BHABHA NATIONAL INSTITUTE, CHHATNAG ROAD, JHUNSI, PRAYAGRAJ (ALLAHABAD) 211 019 INDIA  
*Email address:* [ghoshsohan4@gmail.com](mailto:ghoshsohan4@gmail.com)

HARISH CHANDRA RESEARCH INSTITUTE, A CI OF HOMI BHABHA NATIONAL INSTITUTE, CHHATNAG ROAD, JHUNSI, PRAYAGRAJ (ALLAHABAD) 211 019 INDIA  
*Email address:* [jishnuray@hri.res.in](mailto:jishnuray@hri.res.in); [jishnuray1992@gmail.com](mailto:jishnuray1992@gmail.com)

DEPARTMENT OF MATHEMATICS, CHUO UNIVERSITY, 1-13-27 KASUGA, BUNKYO-KU, TOKYO 112-8551, JAPAN  
*Email address:* [tsuzuki@gug.math.chuo-u.ac.jp](mailto:tsuzuki@gug.math.chuo-u.ac.jp)