Quantum Thermodynamics of Open Quantum Systems: Nature of Thermal Fluctuations

Neha Pathania,^{*} Devvrat Tiwari,[†] and Subhashish Banerjee[‡] Department of Physics, Indian Institute of Technology, Jodhpur, India

We investigate the thermodynamic behavior of open quantum systems through the Hamiltonian of Mean Force, focusing on two models: a two-qubit system interacting with a thermal bath and a Jaynes-Cummings Model without the rotating wave approximation. By analyzing both weak and strong coupling regimes, we uncover the impact of environmental interactions on quantum thermodynamic quantities, including specific heat capacity, internal energy, and entropy. Further, the ergotropy and entropy production are computed. We also explore the thermodynamic uncertainty relation, which sets an upper bound on the signal-to-noise ratio.

I. INTRODUCTION

The study of thermodynamics within the realm of open quantum systems is a burgeoning area that seeks to understand how classical thermodynamic principles apply when quantum effects are significant [1-4]. This field is particularly relevant for systems interacting with their surroundings, leading to complex behaviors and phenomena not observed in isolated systems. A fundamental development in the theory of open quantum systems is the Gorini-Kossakowski-Lindblad-Sudarshan (GKLS) master equation [5-10], which offers a framework for describing the time evolution of a system S under the influence of weak interactions with a large external environment. Advances in theoretical and experimental domains implore one to go beyond the scope of GKLS dynamics, specifically where strong-coupling non-Markovian effects are prominent [11].

Open quantum systems that are non-Markovian demonstrate the intricate connection between a quantum system and its surroundings. Unlike their Markovian counterparts, non-Markovian systems retain a memory of their past, resulting in complex dynamical behavior [11– 21]. A prominent feature of non-Markovian evolution is the resurgence of quantum properties [12], the examination of which is crucial for comprehending the system's dynamics under varying levels of coupling with the environment. Studying the dynamics of thermodynamic quantities when the system has strong coupling with the environment in the non-Markovian regime presents a formidable challenge [22–25]. Numerous techniques have been developed recently to address the dynamics of such systems. These encompass the Hamiltonian of mean force (HMF) [26–31], the method of hierarchical equations of motion, a numerical method utilizing the influence-functional formalism [32, 33], the reaction coordinate method, which explores strong coupling effects through mapping the Hamiltonian with a reaction coordinate [34–36] and pseudomodes technique [37].

Here, we make use of the Hamiltonian of Mean Force to investigate the strong-coupling non-Markovian effects. This concept of HMF has been previously studied in the Caldeira-Leggett Model [38], Spin-Boson Model [39], Quantum Brownian Motion [40] to ascertain the system's effective dynamics and thermodynamics and also in molecular systems [41] to study solvation dynamics and chemical reaction rates in complex environments. It has been observed that HMF provides consistent generalizations of the laws of thermodynamics and fluctuation relations [26, 42].

Recent advances in the study of open quantum systems have vielded significant insights into the behavior of quantum thermodynamic quantities [43, 44], and with technological advancements, this is being leveraged towards quantum thermal devices [45–48]. The specific heat capacity of the systems coupled to thermal reservoirs demonstrates non-trivial temperature dependencies influenced by quantum effects [26, 27]. Along with ubiquitous entropy, another quantity prominent in quantum thermodynamics is entropy production, which, along with system dynamics, also depends on global state evolution [49–51]. The Fluctuation Dissipation Relation (FDR), which connects response functions to correlation functions, is fundamental in understanding dissipation and fluctuations [52]. It turns out that in the strong coupling regime of thermodynamics, FDR has modifications coming from the specific heat capacity and another term that is an essential ingredient in computing the system's internal energy [1, 42]. Additionally, the concept of ergotropy, representing the maximum extractable work from a quantum system via unitary transformations, was introduced [53-55] and is a crucial ingredient in characterizing the thermodynamic behavior of a system. These studies collectively advance our understanding of the complex behavior and properties inherent to open quantum systems.

Another development in this field is a bound on signalto-noise ratio for estimation of temperature [42]. This involves tools of quantum metrology [56, 57] in the form of the quantum Cramer-Rao inequality for an unbiased estimation of the temperature of the system. As is well known, the quantum Cramer-Rao inequality is bounded by the Fisher information [58, 59]. The other ingredient

^{*} neha@ctp-jamia.res.in

[†] devvrat.1@iitj.ac.in

[‡] subhashish@iitj.ac.in

in this bound is the modified fluctuation-dissipation relation involving the specific heat capacity of the system as well as the Wigner-Yanase-Dyson skew information and another term that is pertinent to computing the system's internal energy [1, 42].

Dynamics of two-qubit systems interacting with a squeezed thermal bath has revealed significant insights into entanglement dynamics, coherence, and quantum correlations [60, 61]. The Jaynes-Cummings model [62, 63]—which describes a single two-level system interacting with a single-mode resonator—has seen significant progress in improving quantum measurement techniques specifically in quantum sensing and metrology and also in understanding the interplay between coherence and quantum thermal states [64-67]. In this article, we discuss the quantum thermodynamics of the following open quantum systems models: (i) a two-qubit model interacting with a thermal bath and (ii) the Jaynes-Cummings Model without the rotating wave approximation, using the Hamiltonian of Mean force across all coupling regimes.

The structure of the paper is as follows. The preliminary section (Sec. II) includes an introduction to the Hamiltonian of Mean Force, certain thermodynamical potentials, thermodynamic quantifiers like ergotropy and entropy production, and a brief explanation of quantum information theoretic quantities such as WYD skewinformation and Quantum Fisher information. In Sec. III, we have briefly introduced the models under study and discussed their quantum thermodynamic properties using the Hamiltonian of Mean Force. This is followed by the conclusion in Sec. IV.

II. PRELIMINARIES

This section is divided into two categories. The first category focuses on Quantum Information-theoretic quantities, where we define quantum uncertainty, skew information, and Fisher information. The second category pertains to quantum statistical mechanics, introducing the Hamiltonian of Mean Force and thermodynamic potentials such as entropy and specific heat capacity. Additionally, we provide a brief introduction to entropy production and ergotropy.

A. Quantum Information-Theoretic Quantities

1. Quantum Uncertainty and Wigner-Yanase-Dyson Skew Information

Skew information [68], quantifies the quantum uncertainty in a state relative to an observable defined by $I(\rho, Y) = -\frac{1}{2} \text{Tr}([\sqrt{\rho}, Y]^2)$. It measures the non-commutativity between the state and the observable and

has been extensively studied and generalized. The concept of skew information was extended through the definition $I_{\eta}(\rho, Y) = -\frac{1}{2} \operatorname{Tr}([\rho^{\eta}, Y][\rho^{1-\eta}, Y])$, commonly referred to as WYD entropy [69], where $0 < \eta < 1$. When $\eta = 1/2$, it simplifies to the well-known skew information. This generalized form can be expressed as:

$$I_{\eta}(\rho, Y) = \operatorname{Tr}(\rho Y^{2}) - \operatorname{Tr}(\rho^{\eta} Y \rho^{1-\eta} Y).$$
(1)

In comparison, the usual variance of an observable Y for the quantum state ρ is given by [70],

$$\operatorname{Var}(\rho, Y) = \operatorname{Tr}(\rho Y^2) - (\operatorname{Tr}(\rho Y))^2 = Q[\hat{\rho}, \hat{Y}] + K[\hat{\rho}, \hat{Y}].$$
(2)

The variance of the observable Y is influenced by both quantum and classical uncertainties inherent in the system. The measures of quantum and classical uncertainties, $Q[\hat{\rho}, \hat{Y}]$ and $K[\hat{\rho}, \hat{Y}]$, respectively, are defined as follows [42]

$$Q[\hat{\rho}, \hat{Y}] = \int_0^1 Q_a[\hat{\rho}, \hat{Y}] \, da, \qquad (3)$$

where $Q_a[\hat{\rho}, \hat{Y}] = -\frac{1}{2} \operatorname{Tr} \left([\hat{Y}, \hat{\rho}^a] [\hat{Y}, \hat{\rho}^{1-a}] \right)$ with $a \in (0, 1)$. This measure $Q[\hat{\rho}, \hat{Y}]$ quantifies the quantum uncertainty associated with the observable \hat{Y} in the state $\hat{\rho}$. The negative sign reflects the non-commutativity of $\hat{\rho}$ and \hat{Y} , capturing the essence of quantum fluctuations. Similarly,

$$K[\hat{\rho}, \hat{Y}] := \int_0^1 K_a[\hat{\rho}, \hat{Y}] \, da, \tag{4}$$

where $K_a[\hat{\rho}, \hat{Y}] := \operatorname{Tr}\left(\hat{\rho}^a \delta \hat{Y} \hat{\rho}^{1-a} \delta \hat{Y}\right)$ with $a \in (0, 1)$. Here, $\delta \hat{Y} = \hat{Y} - \langle \hat{Y} \rangle$ represents the deviation of \hat{Y} from its expectation value. The measure $K[\hat{\rho}, \hat{Y}]$ captures the classical uncertainty, reflecting the statistical spread of outcomes due to classical probabilities. The parameter aindicates that there is no unique way of decomposing the variance into quantum and classical components. These quantum and classical uncertainty parameters play a very important role in the thermodynamics of quantum systems which is explored in the later sections.

2. Quantum Fisher Information

Quantum Fisher Information (QFI) is a crucial concept in quantum metrology and information theory, representing the sensitivity of a quantum state to changes in a parameter. It is a fundamental tool for determining the ultimate precision limits in parameter estimation tasks, such as those encountered in quantum thermometry. Mathematically, QFI is defined through the symmetric logarithmic derivative (SLD), Λ_{θ}

$$F(\theta) = \text{Tr}[\rho(\theta)\Lambda_{\theta}^2], \qquad (5)$$

where $\rho(\theta)$ is the density matrix of the quantum state dependent on the parameter θ . This quantity is pivotal as it establishes the quantum Cramér-Rao bound (QCRB), expressed as

$$\Delta \theta \ge \frac{1}{\sqrt{NF(\theta)}},\tag{6}$$

where N is the number of measurements, dictating that the variance of any unbiased estimator θ is inversely proportional to the QFI. It has been recently proved [42] that for an exponential state $\hat{\rho}_{\theta} = \exp\left(-\hat{\Phi}_{\theta}\right)/\mathcal{Z}_{\theta}$ the QFI concerning the parameter θ is bounded by,

$$F(\theta) \le K[\hat{\rho}_{\theta}, \hat{Y}_{\theta}],\tag{7}$$

where, \hat{Y}_{θ} is the thermodynamic observable defined as $\hat{Y}_{\theta} = \partial_{\theta} \hat{\Phi}_{\theta}$, and $K[\hat{\rho}_{\theta}, \hat{Y}_{\theta}]$ is the classical uncertainty defined in Eq. (4). The spectral decomposition of the state $\hat{\rho}_{\theta} = \exp(-\hat{\Phi}_{\theta})/\mathcal{Z}_{\theta}$ is given by $\hat{\rho}_{\theta} = \sum_{n} p_{n} |\psi_{n}\rangle \langle \psi_{n}|$ where the eigenstates satisfy $\hat{\Phi}_{\theta} |\psi_{n}\rangle = \lambda_{n} |\psi_{n}\rangle$. From this spectral decomposition, the QFI can be expressed as [42],

$$F(\theta) = 2\sum_{n,m} \frac{\left|\langle \psi_n | \partial_\theta \hat{\rho}_\theta | \psi_m \rangle\right|^2}{p_n + p_m}.$$
(8)

In this work, we have considered this unbiased estimator parameter θ as Temperature. For a quantum thermal state, the Fisher Information for the temperature is proportional to the specific heat capacity of the system as $F(T) = C_S/T^2$ [58].

B. Quantum Statistical Mechanics Related Quantities

1. Hamiltonian of Mean Force

A quantum system S interacting with a thermal bath \mathcal{B} is described by the Hamiltonian,

$$\hat{\mathcal{H}}_{\mathcal{T}} = \hat{H}_S \otimes \hat{\mathbb{I}}_B + \hat{\mathbb{I}}_S \otimes \hat{H}_B + \hat{\mathcal{V}}_{SB}, \tag{9}$$

where \hat{H}_S and \hat{H}_B are the hamiltonian of system and bath respectively, while $\hat{\mathcal{V}}_{SB}$ is an interaction term of arbitrary strength. Here, we will consider situations where the environment is huge compared to the system, i.e. the operator norms fulfill $||\hat{H}_B|| \gg ||\hat{H}_S||$, $||\hat{\mathcal{V}}_{SB}||$. The Hamiltonian of mean force takes into account the average effects of the environment on the subsystem. The global equilibrium state of the system and the reservoir at temperature T is denoted by $\hat{\zeta}_{SB}$, which is of Gibbs form,

$$\hat{\zeta}_{SB} := \frac{\exp\left(-\beta \hat{\mathcal{H}}_{\mathcal{T}}\right)}{\mathcal{Z}_{SB}},\tag{10}$$

where, $\mathcal{Z}_{SB} = \operatorname{Tr}_{SB} \left[\exp\left(-\beta \hat{\mathcal{H}}_{\mathcal{T}}\right) \right]$ is the partition function for SB. Also, $\beta = (k_b T)^{-1}$, the Boltzmann constant k_b is set to unity throughout. The interaction term causes the reduced state of the system S, $\hat{\zeta}_S(T) = \operatorname{Tr}_B \left[\hat{\zeta}_{SB}(T) \right]$ to deviate from the thermal state unless the coupling is very weak (the magnitude of interaction term $\hat{\mathcal{V}}_{SB}$ is much lesser than the magnitude of the bare system hamiltonian H_S). Thus, for the purpose of calculating internal energy in the strong coupling regime, the system's bare Hamiltonian would not be useful. This issue can be resolved by rewriting the state of the system S as an effective Gibbs state $\hat{\zeta}_S(T) = \frac{\exp(-\beta \hat{\mathcal{H}}_S^*(T))}{Z_S^*}$, where the partition function for the system S can be expressed as the ratio

$$\mathcal{Z}_S^* = \mathcal{Z}_{SB} / \mathcal{Z}_B, \tag{11}$$

where $\mathcal{Z}_{SB} = \text{Tr}_{SB}[e^{-\beta \hat{H}_{SB}}]$ and $\mathcal{Z}_B = \text{Tr}_B[e^{-\beta \hat{H}_B}]$ and,

$$\hat{\mathcal{H}}_{\mathcal{S}}^{*}(\mathcal{T}) := -\frac{1}{\beta} \ln \left(\frac{\operatorname{Tr}_{B} \left[\exp\left(-\beta \hat{\mathcal{H}}_{\mathcal{T}}\right) \right]}{\operatorname{Tr}_{B} \left[\exp\left(-\beta \hat{\mathcal{H}}_{\mathcal{B}}\right) \right]} \right), \qquad (12)$$

is the Hamiltonian of Mean Force [1, 71, 72]. This operator can be interpreted as an effective Hamiltonian describing S, and unlike the bare Hamiltonian \hat{H}_S , it implicitly depends upon both the temperature T and the interaction $\hat{\mathcal{V}}_{SB}$. In the weak coupling regime, this reduces to the bare system Hamiltonian. HMF is essential for understanding thermodynamic properties such as free energy, entropy, and heat capacities in systems where direct interactions with an environment cannot be ignored. It allows for the formulation of generalized thermodynamic potentials and facilitates the study of equilibrium and non-equilibrium processes, providing insights into phenomena such as quantum dissipation and decoherence.

2. Thermodynamic Potentials

The partition function in statistical mechanics contains information regarding the distribution of occupation probabilities among the various microstates of the system S. It enables the calculation of important thermodynamic potentials like free energy, internal energy, and entropy. The internal energy of the system S can be written as [1],

$$U_{S}(T) = U_{SB}(T) - U'_{B}(T).$$
(13)

This $U_S(T)$ is simply the difference between the total energy $U_{SB}(T) = -\partial_\beta \ln \mathcal{Z}_{SB}$ and the energy of the reservoir, $U'_B(T) = -\partial_\beta \ln \mathcal{Z}_B$. $U_S(T)$ can be expressed as an expectation value [1, 73],

$$U_{S}(T) = \left\langle \hat{E}_{S}^{*}(T) \right\rangle = \operatorname{Tr}\left[\hat{E}_{S}^{*}(T).\hat{\zeta}_{S}\right]$$
$$= \operatorname{Tr}\left[\partial_{\beta}(\beta\hat{\mathcal{H}}_{\mathcal{S}}^{*}(\mathcal{T})).\hat{\zeta}_{S}\right], \qquad (14)$$

of the observable,

$$\hat{E}_{S}^{*}(T) = \partial_{\beta}[\beta\hat{\mathcal{H}}_{S}^{*}(\mathcal{T})] = \hat{H}_{S} + \partial_{\beta}[\beta(\hat{\mathcal{H}}_{S}^{*}(\mathcal{T}) - \hat{H}_{S})].$$
(15)

This $\hat{E}_{S}^{*}(T)$ can be interpreted as the effective energy operator describing the system whose eigenstates are referred to as the system's energy eigenstates. With the inclusion of this operator, the fluctuations in the internal energy are considered as:

$$\Delta U_S = \sqrt{\operatorname{Var}[\hat{\zeta}_S, \hat{E}_S^*]}.$$
 (16)

It is significant to remember that $\hat{E}_{S}^{*}(T)$ depends on both the temperature T and the system-environment coupling $\hat{\mathcal{V}}_{SB}$. This $\hat{E}_{S}^{*}(T)$ differs from both the bare system hamiltonian \hat{H}_{S} and hamiltonian of mean force $\hat{H}_{S}^{*}(T)$ [42]. From Eq. (15) we can observe that only in those cases where $\hat{H}_{S}^{*}(T)$ becomes temperature independent will $\hat{H}_{S}^{*}(T)$ be almost equivalent to $\hat{E}_{S}^{*}(T)$. One example is the bilinear coupling model for Quantum Brownian motion [42].

The entropy of the system is calculated as

$$\mathcal{S}_S = -\mathrm{Tr}[(\ln \hat{\zeta}_S).\hat{\zeta}_S] + \beta^2 \mathrm{Tr}[(\partial_\beta \hat{H}_S^*).\hat{\zeta}_S].$$
(17)

Here, we recognize the first contribution as the von Neumann entropy for the system, while the second term stems from the temperature dependence of the HMF. This implies that the thermodynamic entropy is generally not equivalent to the information in the equilibrium state when interactions are present.

In the weak-coupling limit, it is well established that the specific heat capacity C_S is proportional to the variance in internal energy, expressed as [1]

$$C_S = \beta^2 \Delta U_S^2. \tag{18}$$

This is a standard result of the fluctuation-dissipation relation, which connects the rate of change of the system's energy with temperature to the energy fluctuations at equilibrium. Recent research has indicated that for open quantum systems of the form given in Eq. (9), the heat capacity can become negative at low temperatures, suggesting that it is not generally proportional to a positive variance. Further, in the presence of strong coupling, the fluctuation-dissipation relation (FDR) includes two additional contributions [1, 42],

$$C_S(\beta) = \beta^2 \Delta U_S^2 - \beta^2 Q[\hat{\zeta}_S, \hat{E}_S^*] - \frac{1}{\beta^2} \langle \partial_\beta \hat{E}_S^* \rangle, \quad (19)$$

4

where all the terms are as defined above. This equation implies that $C_S(\beta)$ can be less than $\beta^2 \Delta U_S^2$ and even negative. The first correction term arises from quantum fluctuations in energy, represented by the average WYD information related to the observable \hat{E}_S^* . The second correction term $\langle \partial_\beta \hat{E}_S^* \rangle$ accounts for the temperature dependence of \hat{E}_S^* and remains in the classical limit. In the weak coupling limit, where $\hat{E}_S^* = \hat{\mathcal{H}}_S^*$, these correction terms disappear, making the heat capacity proportional to the energy variance. When considering temperature as the unbiased parameter θ , the Cramer-Rao bound inequality in Eq. (6) can be written as $\Delta\beta \geq \frac{1}{\sqrt{F(\beta)}}$ for a single measurement N = 1. Thus, Eq. (7) can be reformulated as $F(\beta) \leq K[\hat{\zeta}_S, \hat{E}_S^*]$. From this, a modified thermodynamic uncertainty relation was obtained as [42],

$$\Delta \beta \ge \frac{1}{\sqrt{\Delta U_S^2 - Q[\hat{\zeta}_S, \hat{E}_S^*]}}.$$
(20)

Comparing Eqs. (19) and (20), the uncertainty in the temperature in the strong coupling regime can be expressed as

$$\Delta T \ge \frac{T}{\sqrt{C_S(T) - \langle \partial_T \hat{E}_S^* \rangle}}.$$
(21)

This expression sets a lower bound on the thermal fluctuations ΔT . In the scenario of strong coupling, the best possible signal-to-noise ratio $\frac{T}{\Delta T}$ for determining the temperature of the system S is constrained by both the specific heat capacity and an additional dissipation term $\langle \partial_T \hat{E}_S^* \rangle$, which can be negative or positive. In the case of weak coupling, this additional dissipation term is zero. Therefore, in that case, the optimal signal-to-noise ratio $\frac{T}{\Delta T}$ depends solely on the specific heat capacity, indicating that accurate temperature measurement requires a large heat capacity.

C. Quantum thermodynamic quantifiers

Here, we briefly discuss quantum thermodynamic quantifiers, particularly ergotropy and entropy production.

1. Ergotropy

Ergotropy is the maximum work that can be extracted from a quantum system [53, 54]. Consider a quantum state ρ , whose spectral decomposition is given by

$$\rho = \sum_{i} r_{i} \left| r_{i} \right\rangle \left\langle r_{i} \right|, \qquad (22)$$

where r_i 's (in the order $r_1 > r_2 > r_3 > ...$) are the eigenvalues of ρ , and the spectral decomposition of the system's Hamiltonian H_S is given by

$$H_S = \sum_j e_j |e_j\rangle \langle e_j|, \qquad (23)$$

where e_j (in the order $e_1 < e_2 < e_3...$) are eigenvalues of the system's Hamiltonian. The ergotropy for this system is given by

$$\mathcal{W}[\rho(t)] = \sum_{j,i} r_i e_j \left(|\langle r_i | e_j \rangle|^2 - \delta_{ij} \right).$$
(24)

2. Entropy production

The transfers undergone by the system during its evolution are alluded to as entropy production. It accounts for the irreversibility that arises due to discarding the bath degrees of freedom. Consider a system bath (S-B)evolution of an arbitrary initial state $\rho_{SB}(0)$ dictated by the unitary operator U, such that $\rho_{SB}(t) = U\rho_{SB}(t)U^{\dagger}$. Entropy production [49, 50] is given by

$$\Sigma = \mathcal{I}(S:B) + S\left[\rho_B(t)||\rho_B(0)\right], \qquad (25)$$

where $\mathcal{I}(S:B)$ is the mutual information given by the von Neumann entropies $(S(\rho) = -\text{Tr}(\rho \ln \rho))$ of the subsystems S and B and the joint system SB, that is,

$$I(S:B) = S(\rho_S) + S(\rho_B) - S(\rho_{SB}).$$
 (26)

 $S(\rho||\sigma) = \text{Tr}(\rho \ln \rho - \rho \ln \sigma)$ is the quantum relative entropy. The above relation can further be simplified as

$$\Sigma = S\left[\rho_{SB}(t)||\rho_S(t) \otimes \rho_B(0)\right],\tag{27}$$

which is the Kullback-Leibler divergence between the joint system-bath evolved state and the state $\rho_S(t) \otimes \rho_B(0)$, where $\rho_S(t) = \text{Tr}_B \left(U \rho_{SB}(t) U^{\dagger} \right)$.

III. ILLUSTRATIVE OPEN QUANTUM SYSTEM MODELS

Here, we briefly discuss the models under consideration in this work. This is followed by a study of their corresponding quantum thermodynamic properties.

A. Two-qubit system interacting with a thermal bath

We start with a two-qubit system interacting with a thermal bath via a dissipative interaction. The dissipative interaction between 2-qubits (a two-level atomic system) and the bath (represented as a three-dimensional electromagnetic field (EMF)) via the dipole interaction [74] is described by the Hamiltonian,

$$H = H_{S} + H_{B} + H_{SB}$$

= $\sum_{j=1}^{2} \hbar \omega_{j} S_{j}^{z} + \sum_{\vec{k}s} \hbar \omega_{k} \left(\alpha_{\vec{k}s}^{\dagger} \alpha_{\vec{k}s} + \frac{1}{2} \right)$
 $- i\hbar \sum_{\vec{k}s} \sum_{j=1}^{2} [\vec{\mu}_{j} \cdot \vec{g}_{\vec{k}s} (\vec{r}_{j}) (S_{j}^{+} + S_{j}^{-}) \alpha_{\vec{k}s} - \text{h.c.}]$
(28)

The two-qubits are modeled as two-level systems with excited state $|e_j\rangle$, ground state $|g_j\rangle$, transition frequency ω_j , and transition dipole moments μ_j . We assume that the qubits are located at different atomic positions $\vec{r_j}$. The transition dipole moments are dependent on $\vec{r_j}$. S_j⁺ = $|e_j\rangle\langle g_j|$ and $S_j^- = |g_j\rangle\langle e_j|$ are the dipole raising and lowering operators, respectively, satisfying the commutation relation and $S_j^z = \frac{1}{2}(|e_j\rangle\langle e_j| - |g_j\rangle\langle g_j|)$ is the energy operator of the jth-atom. $\alpha_{\vec{ks}}^{\dagger}$, $\alpha_{\vec{ks}}$ are the creation and annihilation operators of the field mode \vec{ks} with wave vector \vec{k} , frequency ω_k and the polarization index s. The system-reservoir coupling constant is

$$\vec{g}_{\vec{k}s}(\vec{r}_j) = \left(\frac{\omega_j}{2\epsilon\hbar V}\right)^{1/2} \vec{e}_{\vec{k}s} e^{i\vec{k}.\vec{r}_j}.$$
(29)

V is the normalization volume, and $\vec{e}_{\vec{k}s}$ is the unit polarization of the field. This equation implies that the system-reservoir coupling constant depends on the atomic position r_j . This leads to two regimes: one where the inter-qubit spacing is less than the length scale set by the bath, that is, the collective regime, and the other where the inter-qubit spacing is comparable with the length scale set by the bath, the independent regime [61]. In this work, the collective regime is considered for further analysis.

For simplification, we assume that both the qubits have the same transition frequency and the first qubit is at the origin while the distance between them is ξ . Further, \vec{k} is taken to be along \vec{r}_{12} . The dipole moments $\vec{\mu}_j$'s are taken to be equal for both qubits ($\mu_1 = \mu_2 = \mu$) and along the direction of a polarization vector, such that $\vec{\mu}_j \cdot \vec{e}_{\vec{k}s} =$ μ . Incorporating these quantities, the Hamiltonian in Eq. (28) (for $\hbar = c = 1$) can be rewritten as

$$H = \omega_0 \left(S_1^z + S_2^z \right) + \sum_{\vec{k}} \omega_k \left(\alpha_k^{\dagger} \alpha_k + \frac{1}{2} \right)$$
$$- i \sum_k \lambda \sqrt{\omega_k} \left[\left(S_1^+ + S_1^- \right) \left(\alpha_k - \alpha_k^{\dagger} \right) + \left(S_2^+ + S_2^- \right) \left(\alpha_k e^{i\omega_k \xi} - \alpha_k^{\dagger} e^{-i\omega_k \xi} \right) \right], \qquad (30)$$

where $\lambda = \left(\frac{\mu}{\sqrt{2\epsilon_0 V}}\right)$. The reduced density matrix of the two-qubit system in the interaction picture and the stan-



FIG. 1. Variation of (a) specific heat capacity $C_S(T)$, (b) fluctuations in the internal energy $\Delta U_S(T)$, (c) quantum uncertainty Q(T), and (d) $\langle \partial_T \hat{E}_S^* \rangle$ with temperature T. The parameters are $\omega_0 = 2.0, \xi = 0.05$. In strong-coupling $\lambda \sqrt{\omega_k} \sim \omega_0$, while in weak coupling limit, it is $\sim 10^{-3}\omega_0$.

dard Born-Markov rotating wave approximation (RWA) was obtained in [61, 74] by assuming separable initial system-bath correlation and taking a trace over the bath. When $\lambda \sqrt{\omega_k} \sim \omega_0$, the interaction strength between the system and the bath is comparable to the system's natural frequency ω_0 . This characterizes a strong coupling regime, wherein the system and the bath can efficiently exchange energy, resulting in significant mutual influence on each other's dynamics. Conversely, when $\lambda \sqrt{\omega_k} \ll \omega_0$, the interaction strength is considerably smaller than the system's natural frequency. This corresponds to a weak coupling regime, where the interaction between the system and the bath is minimal. Consequently, the system's dynamics are less perturbed by the bath, and energy exchange between the two is limited.

HMF for this model is calculated using Eqs. (12) and (30), where $\mathcal{H}_{\mathcal{T}}$ is the total Hamiltonian of the system and bath H. The numerical form of the HMF for this model is given by

$$H_S^* = \begin{pmatrix} a & 0 & 0 & e \\ 0 & b & f & 0 \\ 0 & g & c & 0 \\ h & 0 & 0 & d \end{pmatrix}.$$
 (31)

It is observed that in the weak coupling limit, the offdiagonal elements e, f, g, h tend to become zero, and the above Hamiltonian becomes equal to the bare Hamiltonian of the system H_S . The specific heat capacity and the corresponding variables in its expression are plotted in Fig. 1. As can be observed, the quantum uncertainty and the quantity pertaining to E_S^* are zero in the weak coupling regime but non-zero in the strong coupling regime. This is consistent with the generalized fluctuation-dissipation relation, Eq. (19).

An upper bound on the signal-to-noise ratio, obtained



FIG. 2. Variation of the upper bound $C_S(T) - \langle \partial_T \hat{E}^* \rangle$ on the signal-to-noise ratio $\left(\frac{T}{\Delta T_S}\right)^2$, and of $\left(\frac{T}{\Delta T_S}\right)^2_{\text{opt}} = T^2 F_S(T)$ as a function of temperature T. The parameters are $\omega_0 = 2.0, \xi = 0.05$. In strong-coupling $\lambda \sqrt{\omega_k} \sim \omega_0$, while in weak coupling limit, it is $\sim 10^{-3}\omega_0$.



FIG. 3. Plot for the entropy [Eq. (17)] as a function of temperature for the two-qubit decoherence model. The parameters are chosen to be $\omega_0 = 2.0, \xi = 0.05$. In strong-coupling $\lambda \sqrt{\omega_k} \sim \omega_0$, while in weak coupling limit, it is $\sim 10^{-3} \omega_0$.

from the modified thermodynamic uncertainty relation, Eq. (21), is depicted in Fig. 2. The bounds are well satisfied for both the weak and strong coupling regimes, with the weak coupling regime depicting the tighter bound, in fact, saturating the bound.

The entropy of the reduced dynamics of the system is plotted in Fig. 3. The weak coupling behavior of the entropy is along the expected line, whereas for strong coupling, the entropy is negative at lower temperatures. However, it approaches the entropy in the weak-coupling regime as the temperature is raised.

The ergotropy is calculated using the state $\zeta_S(T) =$



FIG. 4. Variation of Ergotropy $\mathcal{W}(\hat{\zeta}_S)$ with Temperature T in the strong, moderate, and weak coupling limit for a twoqubit squeezed thermal bath model. The parameters are chosen to be $\omega_0 = 2.0, \xi = 0.05$. In strong-coupling $\lambda \sqrt{\omega_k} \sim \omega_0$, moderate-coupling is when the coupling constant is kept half of the strong-coupling constant value, while in weak coupling limit, it is $\sim 10^{-3}\omega_0$.

 $e^{-\beta H_S^*}/Z^*$, and is plotted in Fig. 4. From this figure, one can infer that in the weak coupling regime, the HMF coincides with the bare Hamiltonian of the system, and the ergotropy becomes zero, as the state $\zeta_S(T)$ becomes the Gibbs state for the bare Hamiltonian, which is not the case in the moderate and strong coupling regimes. Further, ergotropy is composed of coherent and incoherent parts. It is observed that for the state $\zeta_S(T)$ of the twoqubit model, the incoherent contribution is zero. The non-zero ergotropy comes from the coherent part. This is due to the fact that the state $\zeta_S(T)$ has coherence terms present in it at stronger system-bath couplings, which are absent in the weak coupling regime.

Considering the single-mode resonator approximation of the above mode (for single k in Eq. (30)), we calculate the entropy production, Eq. (27), using the Bell state $\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ as the initial state of the two-qubit system, and $\rho_B(0) = \frac{e^{-\beta H_B}}{\text{Tr}(e^{-\beta H_B})}$ as the initial state of the bath. The dynamics of the total and the reduced systems are obtained using $U\{\rho_S(0) \otimes \rho_B(0)\} U^{\dagger}$ and $\text{Tr}_B \left[U\{\rho_S(0) \otimes \rho_B(0)\} U^{\dagger}\right]$, respectively, where $U = e^{-iHt}$. The variation of entropy production Σ as a function of temperature, at time t = 1, is plotted in Fig. 5. As can be observed from Fig. 5, in this case, the hierarchy of the entropy production with respect to the coupling strength is similar to that of the ergotropy.



FIG. 5. Variation of the entropy production Σ as a function of temperature at time t = 1 for the two-qubit model interacting with single bath resonator. The strong, moderate, and weak couplings are taken as discussed in the above figures. The parameters are taken to be: $\omega_0 = 2.0, \omega = 1, \xi = 0.05$.

B. A single two-level system interacting with a single-mode resonator

Here, we take an example of a two-level system interacting with a single-mode field, that is, the Jaynes-Cummings (JC) model without rotating wave approximation [63, 75]. The Hamiltonian (for $\hbar = 1$) of the total system is given by

$$H = H_S + H_B + H_{SB}$$

= $\frac{\omega_0}{2} \sigma^z + \omega_c \hat{a}^{\dagger} \hat{a} + \lambda \sigma^x \left(\hat{a} + \hat{a}^{\dagger} \right).$ (32)

We find out the HMF numerically for this model using Eq. (12). It is found to be diagonal in form, though not equal to H_S in the strong coupling regime. It is, however, the same as the bare Hamiltonian in the weak coupling regime. The specific heat capacity $C_S(T)$ and related quantities as a function of temperature are plotted in Fig. 6. It is observed that the quantum uncertainty is zero in both weak and strong coupling limits. The term $\langle \partial_T \hat{E}_S^* \rangle$ is non-zero for strong coupling but vanishes in the weak coupling limit. The generalized fluctuation-dissipation relation, Eq. (19), is satisfied. Further, $C_S(T)$ is negative in the strong-coupling regime, which was also observed in [76].

The upper bound on the signal-to-noise ratio is depicted in Fig. 7 and is observed to be tight both for the weak and strong coupling regimes. From this, one could infer the local temperature attained by the system S. The entropy of the system using HMF is plotted in Fig. 8 and is along expected lines.

The ergotropy is calculated using the state $\hat{\zeta}_S(T) = e^{-\beta H_S^*}/Z_S^*$, which is zero due to the absence of coherence, and the diagonal terms having $\text{Tr}[\sigma^z \hat{\zeta}_S(T)] < 0$,



FIG. 6. Variation of (a) specific heat capacity $C_S(T)$, (b) fluctuations in the internal energy $\Delta U_S(T)$, (c) quantum uncertainty Q(T), and (d) $\langle \partial_T \hat{E}_S^* \rangle$ with temperature T for the JC model. The parameters are: $\omega_0 = 2.0, \omega_c = 1.0$. The weak coupling limit corresponds to $\lambda \ll \omega_0$, and $\lambda \sim \omega_0$ corresponds to the strong coupling limit.



FIG. 7. Variation of the the upper bound on the signal-tonoise ratio $(C_S(T) - \langle \partial_T \hat{E}^* \rangle)$ for the JC model using Eq. (19) as a function of temperature. The parameters are: $\omega_0 = 2.0, \omega_c = 1.0$.

that is, the population of the excited state is always lower than the ground state of the system in the state $\zeta_S(T)$ [77]. The entropy production for the JC model is calculated using the initial states $|\psi\rangle_S = \frac{1}{\sqrt{2}} (|0\rangle + |1\rangle)$ and $\rho_B(0) = e^{-\beta H_B}/\text{Tr} \left[e^{-\beta H_B}\right]$ of the system and the bath, using Eq. (27), where the unitary evolution is given by $U = e^{-iHt}$. The variation of entropy production as a function of temperature is shown in Fig. 9 and exhibits the same pattern as in the earlier model.



FIG. 8. Variation of entropy [Eq. (17)] as a function of temperature for the JC model. Here, $\omega_0 = 2.0, \omega_c = 1.0$.



FIG. 9. Variation of the entropy production Σ as a function of temperature at time t = 0.5 for the JC model. The parameters are $\omega_0 = 2.0, \omega_c = 1.0$.

IV. CONCLUSIONS

In this work, we studied the quantum thermodynamic behavior of open quantum system models. To understand the effect of weak and strong coupling, use was made of the Hamiltonian of mean force, which could be interpreted as an effective Hamiltonian of the system instead of the system's bare Hamiltonian in the strong coupling regime. The thermodynamic potentials, particularly specific heat capacity, internal energy, and entropy, were studied using this Hamiltonian of mean force. Further, we also discussed the thermodynamic uncertainty relation that provides an upper-bound to signal-to-noise ratio. This upper bound was computed using the specific heat capacity of the system of interest and a quantity used for computing the internal energy of the system. The behavior of the above thermodynamic quantities was illustrated in two models. First, we considered a twoqubit model interacting with an electromagnetic field, and next the Jaynes-Cumming model without the rotating wave approximation. The generalized fluctuationdissipation relation was satisfied for both models for all coupling limits, ranging from weak to strong coupling. The upper bound on the signal-to-noise ratio was tight for the two-qubit model in the strong coupling regime. In the weak coupling regime of the two-qubit model and

- H. J. D. Miller, Hamiltonian of Mean Force for Strongly-Coupled Systems (Springer International Publishing, Cham, 2018), pp. 531–549, ISBN 978-3-319-99046-0, URL https://doi.org/10.1007/978-3-319-99046-0_ 22.
- [2] T. Jahnke, S. Lanéry, and G. Mahler, Physical Review E—Statistical, Nonlinear, and Soft Matter Physics 83, 011109 (2011).
- [3] R. Alicki, Journal of Physics A: Mathematical and General 12, L103 (1979).
- [4] R. Kosloff, Entropy 15, 2100 (2013), ISSN 1099-4300, URL https://www.mdpi.com/1099-4300/15/6/2100.
- [5] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, Journal of Mathematical Physics 17, 821 (1976).
- [6] A. Kossakowski, Reports on Mathematical Physics 3, 247 (1972).
- [7] H.-P. Breuer and F. Petruccione, *The theory of open quantum systems* (Oxford University Press, USA, 2002).
- [8] G. Lindblad, Communications in mathematical physics 48, 119 (1976).
- [9] A. Rivas and S. F. Huelga, Open quantum systems, vol. 10 (Springer, 2012).
- S. Banerjee, Open Quantum Systems: Dynamics of Nonclassical Evolution (Springer Singapore, 2018), ISBN 9789811331824, URL http://dx.doi.org/10. 1007/978-981-13-3182-4.
- [11] I. de Vega and D. Alonso, Rev. Mod. Phys. 89, 015001 (2017), URL https://link.aps.org/doi/10. 1103/RevModPhys.89.015001.
- [12] H.-P. Breuer, E.-M. Laine, and J. Piilo, Phys. Rev. Lett. 103, 210401 (2009), URL https://link.aps.org/doi/ 10.1103/PhysRevLett.103.210401.
- [13] A. Rivas, S. F. Huelga, and M. B. Plenio, Phys. Rev. Lett. **105**, 050403 (2010), URL https://link.aps.org/ doi/10.1103/PhysRevLett.105.050403.
- [14] R. Vasile, S. Maniscalco, M. G. A. Paris, H.-P. Breuer, and J. Piilo, Phys. Rev. A 84, 052118 (2011), URL https://link.aps.org/doi/10.1103/PhysRevA. 84.052118.
- [15] N. P. Kumar, S. Banerjee, R. Srikanth, V. Jagadish, and F. Petruccione, Open Systems & Information Dynamics 25, 1850014 (2018), https://doi.org/10.1142/S1230161218500142, URL https://doi.org/10.1142/S1230161218500142.
- [16] H.-P. Breuer, E.-M. Laine, J. Piilo, and B. Vacchini, Reviews of Modern Physics 88, 021002 (2016).

for the single-qubit model for all coupling strengths, the inequality was saturated. Further, in the case of the twoqubit model, the Gibbs state found using the Hamiltonian of mean force had a non-zero capacity for doing work, as its coherent ergotropy was non-zero in strong and moderate coupling limits due to the presence of coherence in this state. The entropy production for both systems was greater in the strong coupling regime, reducing with decrease in the coupling strength. Our results provide insights into the complex behavior of open quantum systems in the context of quantum thermodynamics.

- [17] W.-M. Zhang, P.-Y. Lo, H.-N. Xiong, M. W.-Y. Tu, and F. Nori, Physical review letters **109**, 170402 (2012).
- [18] B.-H. Liu, L. Li, Y.-F. Huang, C.-F. Li, G.-C. Guo, E.-M. Laine, H.-P. Breuer, and J. Piilo, Nature Physics 7, 931 (2011).
- [19] S. Utagi, R. Srikanth, and S. Banerjee, Scientific Reports 10 (2020), ISSN 2045-2322, URL http://dx.doi.org/ 10.1038/s41598-020-72211-3.
- [20] J. Naikoo, S. Dutta, and S. Banerjee, Phys. Rev. A 99, 042128 (2019), URL https://link.aps.org/doi/ 10.1103/PhysRevA.99.042128.
- [21] N. P. Kumar, S. Banerjee, and C. M. Chandrashekar, Scientific Reports 8 (2018), ISSN 2045-2322, URL http: //dx.doi.org/10.1038/s41598-018-27132-7.
- [22] P. Strasberg, G. Schaller, N. Lambert, and T. Brandes, New Journal of Physics 18, 073007 (2016).
- [23] G. Thomas, N. Siddharth, S. Banerjee, and S. Ghosh, Physical Review E 97, 062108 (2018).
- [24] W.-M. Huang and W.-M. Zhang, Phys. Rev. A 106, 032607 (2022), URL https://link.aps.org/doi/10. 1103/PhysRevA.106.032607.
- [25] D. Tiwari, B. Bose, and S. Banerjee, arXiv preprint arXiv:2404.15915 (2024), URL https://arxiv.org/ abs/2404.15915.
- [26] M. Campisi, P. Talkner, and P. Hänggi, Phys. Rev. Lett. 102, 210401 (2009), URL https://link.aps.org/doi/ 10.1103/PhysRevLett.102.210401.
- [27] P. Hänggi, G.-L. Ingold, and P. Talkner, New Journal of Physics 10, 115008 (2008).
- [28] M. Campisi, P. Hänggi, and P. Talkner, Reviews of Modern Physics 83, 771 (2011).
- [29] P. Talkner and P. Hänggi, Reviews of Modern Physics 92, 041002 (2020).
- [30] S. Hilt, B. Thomas, and E. Lutz, Physical Review E—Statistical, Nonlinear, and Soft Matter Physics 84, 031110 (2011).
- [31] P. Strasberg and M. Esposito, Physical Review E 101, 050101 (2020).
- [32] Y. Tanimura, The Journal of chemical physics 153 (2020).
- [33] J. Bätge, Y. Ke, C. Kaspar, and M. Thoss, Physical Review B 103, 235413 (2021).
- [34] A. Nazir and G. Schaller, Thermodynamics in the Quantum Regime: Fundamental Aspects and New Directions pp. 551–577 (2018).
- [35] J. Iles-Smith, N. Lambert, and A. Nazir, Phys. Rev. A 90, 032114 (2014), URL https://link.aps.org/doi/

10.1103/PhysRevA.90.032114.

- [36] N. Anto-Sztrikacs and D. Segal, Physical Review A 104, 052617 (2021).
- [37] G. Pleasance, B. M. Garraway, and F. Petruccione, Physical Review Research 2, 043058 (2020).
- [38] A. O. Caldeira and A. J. Leggett, Physica A: Statistical mechanics and its Applications 121, 587 (1983).
- [39] U. Weiss, Quantum dissipative systems (World Scientific, 2012).
- [40] P. Hänggi and G.-L. Ingold, Chaos: An Interdisciplinary Journal of Nonlinear Science 15 (2005).
- [41] R. Kapral and G. Ciccotti, The Journal of chemical physics 110, 8919 (1999).
- [42] H. J. Miller and J. Anders, Nature communications 9, 2203 (2018).
- [43] S. Lahiri, S. Banerjee, and A. M. Jayannavar, Quantum Information Processing 20 (2021), ISSN 1573-1332, URL http://dx.doi.org/10.1007/s11128-021-03260-4.
- [44] S. Banerjee, S. Choudhury, S. Chowdhury, J. Knaute, S. Panda, and K. Shirish, Nuclear Physics B 996, 116368 (2023), ISSN 0550-3213, URL http://dx.doi.org/10. 1016/j.nuclphysb.2023.116368.
- [45] S. Vinjanampathy and J. Anders, Contemporary Physics 57, 545 (2016).
- [46] A. Kumar, S. Lahiri, T. Bagarti, and S. Banerjee, Physica A: Statistical Mechanics and its Applications 623, 128832 (2023), ISSN 0378-4371, URL http://dx.doi. org/10.1016/j.physa.2023.128832.
- [47] G. Bhanja, D. Tiwari, and S. Banerjee, Phys. Rev. A 109, 012224 (2024), URL https://link.aps.org/doi/ 10.1103/PhysRevA.109.012224.
- [48] G. Manzano, R. Sánchez, R. Silva, G. Haack, J. B. Brask, N. Brunner, and P. P. Potts, Phys. Rev. Res. 2, 043302 (2020), URL https://link.aps.org/doi/10. 1103/PhysRevResearch.2.043302.
- [49] G. T. Landi and M. Paternostro, Rev. Mod. Phys. 93, 035008 (2021), URL https://link.aps.org/doi/ 10.1103/RevModPhys.93.035008.
- [50] M. Esposito, K. Lindenberg, and C. Van den Broeck, New Journal of Physics 12, 013013 (2010).
- [51] S. Deffner and E. Lutz, Physical review letters 107, 140404 (2011).
- [52] R. Kubo, Reports on progress in physics **29**, 255 (1966).
- [53] A. E. Allahverdyan and T. M. Nieuwenhuizen, Phys. Rev. Lett. 85, 1799 (2000), URL https://link.aps. org/doi/10.1103/PhysRevLett.85.1799.
- [54] A. E. Allahverdyan, R. Balian, and T. M. Nieuwenhuizen, Europhysics Letters 67, 565 (2004).
- [55] R. Alicki, M. Horodecki, P. Horodecki, and R. Horodecki, Open Systems & Information Dynamics 11, 205 (2004).

- [56] V. Giovannetti, S. Lloyd, and L. Maccone, Nature photonics 5, 222 (2011).
- [57] M. G. Paris, International Journal of Quantum Information 7, 125 (2009).
- [58] J. Liu, H. Yuan, X.-M. Lu, and X. Wang, Journal of Physics A: Mathematical and Theoretical 53, 023001 (2020).
- [59] C. W. Helstrom, Physics letters A 25, 101 (1967).
- [60] S. Banerjee and R. Ghosh, Journal of Physics A: Mathematical and Theoretical 40, 13735 (2007).
- [61] S. Banerjee, V. Ravishankar, and R. Srikanth, Annals of Physics 325, 816 (2010).
- [62] E. T. Jaynes and F. W. Cummings, Proceedings of the IEEE 51, 89 (1963).
- [63] J. Larson and T. Mavrogordatos, The Jaynes-Cummings model and its descendants: modern research directions (IoP Publishing, 2021).
- [64] S. Deffner and E. Lutz, Physical review letters 111, 010402 (2013).
- [65] J. Casanova, G. Romero, I. Lizuain, J. J. García-Ripoll, and E. Solano, Physical review letters 105, 263603 (2010).
- [66] K. Xu, W. Han, Y.-J. Zhang, and H. Fan, Chinese Physics B 27, 010302 (2018).
- [67] A. Messinger, A. Ritboon, F. Crimin, S. Croke, and S. M. Barnett, New Journal of Physics 22, 043008 (2020).
- [68] E. P. Wigner and M. M. Yanase, Proceedings of the National Academy of Sciences 49, 910 (1963).
- [69] E. H. Lieb, Les rencontres physiciens-mathématiciens de Strasbourg-RCP25 19, 0 (1973).
- [70] X. Li, D. Li, H. Huang, X. Li, and L. Kwek, The European Physical Journal D 64, 147 (2011).
- [71] J. G. Kirkwood, The Journal of chemical physics 3, 300 (1935).
- [72] G. Timofeev and A. Trushechkin, International Journal of Modern Physics A 37, 2243021 (2022).
- [73] U. Seifert, Physical review letters **116**, 020601 (2016).
- [74] Z. Ficek and R. Tanaś, Physics Reports 372, 369 (2002).
- [75] A. Smirne and B. Vacchini, Phys. Rev. A 82, 022110 (2010), URL https://link.aps.org/doi/10. 1103/PhysRevA.82.022110.
- [76] M. Campisi, P. Talkner, and P. Hänggi, Journal of Physics A: Mathematical and Theoretical 42, 392002 (2009), ISSN 1751-8121, URL http://dx.doi.org/10. 1088/1751-8113/42/39/392002.
- [77] D. Tiwari and S. Banerjee, Frontiers in Quantum Science and Technology 2 (2023), ISSN 2813-2181, URL http: //dx.doi.org/10.3389/frqst.2023.1207552.