

A-DAVIS-WIELANDT RADIUS BOUNDS OF SEMI-HILBERTIAN SPACE OPERATORS

MESSAOUD GUESBA, SOMDATTA BARIK, PINTU BHUNIA, KALLOL PAUL

ABSTRACT. Consider \mathcal{H} is a complex Hilbert space and A is a positive operator on \mathcal{H} . The mapping $\langle \cdot, \cdot \rangle_A : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$, defined as $\langle y, z \rangle_A = \langle Ay, z \rangle$ for all $y, z \in \mathcal{H}$, induces a seminorm $\|\cdot\|_A$. The A -Davis-Wielandt radius of an operator S on \mathcal{H} is defined as $d\omega_A(S) = \sup \left\{ \sqrt{|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4} : \|z\|_A = 1 \right\}$. We investigate some new bounds for $d\omega_A(S)$ which refine the existing bounds. We also give some bounds for the 2×2 off-diagonal block matrices.

1. Introduction and preliminaries

Here we collect all the technical ingredients that are necessary to read the paper. Throughout, \mathcal{H} is a complex Hilbert space and $\mathcal{B}(\mathcal{H})$ is the \mathbb{C}^* -algebra of bounded operators on \mathcal{H} . For $S \in \mathcal{B}(\mathcal{H})$, the null space, range space and closure of the range space of S are respectively denoted as $\mathcal{N}(S)$, $\mathcal{R}(S)$ and $\overline{\mathcal{R}(S)}$. A positive operator $A \in \mathcal{B}(\mathcal{H})$ defines a sesquilinear form $\langle \cdot, \cdot \rangle_A : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$ defined as $\langle x, z \rangle_A = \langle Ax, z \rangle$. This gives a seminorm $\|\cdot\|_A$ defined by $\|z\|_A = \|A^{\frac{1}{2}}z\|$. Clearly, $\|z\|_A = 0$ iff $z \in \mathcal{N}(A)$. Therefore, $\|\cdot\|_A$ is a norm on \mathcal{H} iff A is one-one. Suppose $\|S\|_A = \sup_{\substack{z \in \overline{\mathcal{R}(A)} \\ z \neq 0}} \frac{\|Sz\|_A}{\|z\|_A} = \sup_{\substack{z \in \overline{\mathcal{R}(A)} \\ \|z\|_A=1}} \|Sz\|_A$ is the A -operator seminorm of $S \in \mathcal{B}(\mathcal{H})$. If there is a scalar $c > 0$ such that $\|Sz\|_A \leq c\|z\|_A$ for all $z \in \overline{\mathcal{R}(A)}$, then $\|S\|_A < \infty$. If $\|S\|_A < \infty$, then $\|S\|_A = \sup \left\{ |\langle Sx, z \rangle_A| : x, z \in \overline{\mathcal{R}(A)}, \|x\|_A = \|z\|_A = 1 \right\}$. An operator $R \in \mathcal{B}(\mathcal{H})$ is A -adjoint of $S \in \mathcal{B}(\mathcal{H})$ if $\langle Sx, z \rangle_A = \langle x, Rz \rangle_A$ for all $x, z \in \mathcal{H}$,

2010 *Mathematics Subject Classification.* 47A05, 47A12, 47A30, 47B15.

Key words and phrases. Positive operator, semi-inner product, A -Davis-Wielandt radius, seminorm.

The authors would like to acknowledge the referee for his/her valuable comments towards improve the article. Miss Somdatta Barik would like to thank UGC, Govt. of India for the financial support in the form of JRF under the mentorship of Prof. Kallol Paul. Dr. Pintu Bhunia would like to thank SERB, Govt. of India for the financial support in the form of National Post Doctoral Fellowship (N-PDF, File No. PDF/2022/000325) under the mentorship of Prof. Apoorva Khare.

or equivalently, $AR = S^*A$. If $AS = S^*A$ then S is A -selfadjoint and if AS is positive then S is A -positive and we write $S \geq_A 0$. Consider $\mathcal{B}_A(\mathcal{H})$ is the collection of operators that admit A -adjoints. From Douglas result [11], we have $\mathcal{B}_A(\mathcal{H}) = \{S \in \mathcal{B}(\mathcal{H}) : \mathcal{R}(S^*A) \subseteq \mathcal{R}(A)\} = \{S \in \mathcal{B}(\mathcal{H}) : \exists c > 0 : \|ASz\| \leq c\|Az\|, \forall z \in \mathcal{H}\}$. For $S \in \mathcal{B}_A(\mathcal{H})$, S admits A -adjoint. Furthermore, there exists a distinguished A -adjoint of S , namely, the reduced solution of $AZ = S^*A$, i.e., $S^{\sharp A} = A^\dagger S^*A$, where A^\dagger is the Moore-Penrose inverse of A . The operator $S^{\sharp A}$ satisfies $AS^{\sharp A} = S^*A$, $\mathcal{R}(S^{\sharp A}) \subseteq \overline{\mathcal{R}(A)}$, $\mathcal{N}(S^{\sharp A}) = \mathcal{N}(S^*A)$. Again, by Douglas result, we can see that $\mathcal{B}_{A^{1/2}}(\mathcal{H}) = \{S \in \mathcal{B}(\mathcal{H}) : \exists c > 0 : \|Sz\|_A \leq c\|z\|_A, \forall z \in \mathcal{H}\}$. If $S \in \mathcal{B}_{A^{1/2}}(\mathcal{H})$, $\|S\|_A = \sup_{\substack{z \notin \mathcal{N}(A) \\ \|z\|_A=1}} \frac{\|Sz\|_A}{\|z\|_A} = \sup_{z \in \mathcal{H}} \|Sz\|_A$, see [2]. If $S \in \mathcal{B}_{A^{1/2}}(\mathcal{H})$ then $S(\mathcal{N}(A)) \subseteq \mathcal{N}(A)$ and $\|Sz\|_A \leq \|S\|_A \|z\|_A$ for all $z \in \mathcal{H}$. The sets $\mathcal{B}_A(\mathcal{H})$ and $\mathcal{B}_{A^{1/2}}(\mathcal{H})$ are subalgebras (see [2]) and $\mathcal{B}_A(\mathcal{H}) \subseteq \mathcal{B}_{A^{1/2}}(\mathcal{H})$. Note that S is A -selfadjoint, but $S = S^{\sharp A}$ may not be true. For example, consider $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \in \mathcal{M}_2(\mathbb{C})^+$ and $S = \begin{pmatrix} 2 & 2 \\ 0 & 0 \end{pmatrix}$. Here S is A -selfadjoint, however $S^{\sharp A} = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \neq S$. Note that, $S = S^{\sharp A}$ iff $\mathcal{R}(S^{\sharp A}) \subseteq \mathcal{R}(A)$ and S is A -selfadjoint. An operator S in $\mathcal{B}_A(\mathcal{H})$ is A -unitary if $\|Sx\|_A = \|S^{\sharp A}x\|_A = \|x\|_A$, $\forall x \in \mathcal{H}$. For more we refer to [1, 2]. If $S \in \mathcal{B}_A(\mathcal{H})$ then the following holds:

- (1) If $AS = SA$, then $S^{\sharp A} = P_{\overline{\mathcal{R}(A)}}S^*$, where $P_{\overline{\mathcal{R}(A)}}$ denote the orthogonal projection on $\overline{\mathcal{R}(A)}$.
- (2) $S^{\sharp A}S$, $SS^{\sharp A}$ are A -selfadjoint and are A -positive.
- (3) $S^{\sharp A} \in \mathcal{B}_A(\mathcal{H})$, $(S^{\sharp A})^{\sharp A} = P_{\overline{\mathcal{R}(A)}}SP_{\overline{\mathcal{R}(A)}}$, $\left((S^{\sharp A})^{\sharp A}\right)^{\sharp A} = S^{\sharp A}$.
- (4) If $R \in \mathcal{B}_A(\mathcal{H})$ then $SR \in \mathcal{B}_A(\mathcal{H})$ and $(SR)^{\sharp A} = R^{\sharp A}S^{\sharp A}$. Furthermore, for all $S, R \in \mathcal{B}_{A^{1/2}}(\mathcal{H})$, we have

$$\|SR\|_A \leq \|S\|_A \|R\|_A. \quad (1.1)$$

$$(5) \|S^{\sharp A}\|_A = \|S\|_A = \|S^{\sharp A}S\|_A^{\frac{1}{2}} = \|SS^{\sharp A}\|_A^{\frac{1}{2}}.$$

To simplify we write P instead of $P_{\overline{\mathcal{R}(A)}}$. An operator R in $\mathcal{B}_A(\mathcal{H})$ can be written as $R = \Re_A(R) + i\Im_A(R)$, where $\Re_A(R) = \frac{R+R^{\sharp A}}{2}$ and $\Im_A(R) = \frac{R-R^{\sharp A}}{2i}$. Recall that, $\omega(S) := \sup\{|\langle Sz, z \rangle| : z \in \mathcal{H}, \|z\| = 1\}$ is the numerical radius of $S \in \mathcal{B}(\mathcal{H})$, see [18]. For more facts, see [12, 16, 20, 28] and references therein. One of the generalization of the numerical radius is A -numerical radius of $S \in \mathcal{B}_A(\mathcal{H})$, it is defined by $\omega_A(S) := \sup\{|\langle Sz, z \rangle_A| : z \in \mathcal{H}, \|z\|_A = 1\}$. Obviously, for $A = I$ we obtain the (classical) numerical radius which received considerable attention in recent years by many researchers. We refer to [7, 8, 15, 17] and references therein. Clearly, $\omega_A(S) = \omega_A(S^{\sharp A})$ for any $S \in \mathcal{B}_A(\mathcal{H})$. Also, the A -numerical radius satisfies the power

inequality (see [24, 30]): for $S \in \mathcal{B}_{A^{1/2}}(\mathcal{H})$, $\omega_A(S^n) \leq \omega_A^n(S)$, $n = 1, 2, \dots$. It defines a seminorm on $\mathcal{B}_{A^{1/2}}(\mathcal{H})$, and it satisfies

$$\frac{1}{2} \|S\|_A \leq \omega_A(S) \leq \|S\|_A \quad \text{for every } S \in \mathcal{B}_{A^{1/2}}(\mathcal{H}). \quad (1.2)$$

Here $\omega_A(S) = \|S\|_A$ if S is A -selfadjoint. Recall that, the A -Crawford number of $S \in \mathcal{B}_{A^{1/2}}(\mathcal{H})$ is defined by $c_A(S) = \inf \{|\langle Sz, z \rangle_A| : z \in \mathcal{H}, \|z\|_A = 1\}$. For more basic facts, we refer to [14, 27, 30]. The Davis-Wielandt radius of $S \in \mathcal{B}(\mathcal{H})$ is defined as $d\omega(S) = \sup \{(|\langle Sz, z \rangle|^2 + \|Sz\|^4)^{1/2} : z \in \mathcal{H}, \|z\| = 1\}$. For more about Davis-Wielandt radius we refer to [5, 6, 9, 23, 29, 31]. The A -Davis-Wielandt radius of $S \in \mathcal{B}_{A^{1/2}}(\mathcal{H})$ is defined (see [3]) as $d\omega_A(S) = \sup \{(|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4)^{1/2} : z \in \mathcal{H}, \|z\|_A = 1\}$. It satisfies $\max \{\omega_A(S), \|S\|_A^2\} \leq d\omega_A(S) \leq \sqrt{\omega_A^2(S) + \|S\|_A^4}$.

Very recently some results on the A -Davis-Wielandt radius are studied (see [3]). In this manuscript we continue the study in this direction and we provide some new bounds of the A -Davis-Wielandt radius. Further, we give some new bounds for the Davis-Wielandt radius of 2×2 off-diagonal block matrices.

2. BOUNDS FOR THE A -DAVIS-WIELANDT RADIUS OF OPERATORS

To develop our first bound of A -Davis-Wielandt radius, the following inner-product inequality is necessary.

Lemma 2.1. [4, 21] *If $a, b, c \in \mathcal{H}$ and $\|c\|_A = 1$ and $0 \neq \alpha \in \mathbb{C}$, then*

$$|\langle a, c \rangle_A \langle c, b \rangle_A| \leq \frac{1}{|\alpha|} \left(\max\{1, |\alpha - 1|\} \|a\|_A \|b\|_A + |\langle a, b \rangle_A| \right).$$

In particular, for $\alpha = 2$,

$$|\langle a, c \rangle_A \langle c, b \rangle_A| \leq \frac{1}{2} \left(\|a\|_A \|b\|_A + |\langle a, b \rangle_A| \right).$$

Theorem 2.2. *Let $S \in \mathcal{B}_A(\mathcal{H})$ and $0 \neq \alpha \in \mathbb{C}$. Then*

$$d\omega_A^2(S) \leq \omega_A^2(S^{\sharp_A} S + S) + \frac{2}{|\alpha|} \omega_A(S^{\sharp_A} S^2) + \frac{\max\{1, |\alpha - 1|\}}{|\alpha|} \left\| (S^{\sharp_A} S)^2 + S^{\sharp_A} S \right\|_A.$$

Proof. Take $z \in \mathcal{H}$ and $\|z\|_A = 1$. We obtain

$$\begin{aligned} & |\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \\ &= |\langle Sz, z \rangle_A + \langle Sz, Sz \rangle_A|^2 - 2\Re_A(\langle Sz, Sz \rangle_A \langle Sz, z \rangle_A) \\ &\leq |\langle (S + S^{\sharp_A} S)z, z \rangle_A|^2 + 2 |\langle S^{\sharp_A} Sz, z \rangle_A \langle z, Sz \rangle_A| \\ &\leq |\langle (S + S^{\sharp_A} S)z, z \rangle_A|^2 + \frac{2}{|\alpha|} \left[\max\{1, |\alpha - 1|\} \|S^{\sharp_A} Sz\|_A \|Sz\|_A + |\langle S^{\sharp_A} Sz, Sz \rangle_A| \right] \\ &\quad (\text{by Lemma 2.1}) \end{aligned}$$

$$\begin{aligned}
&\leq |\langle (S + S^{\sharp_A} S) z, z \rangle_A|^2 + \frac{\max\{1, |\alpha - 1|\}}{|\alpha|} \left(\|S^{\sharp_A} S z\|_A^2 + \|S z\|_A^2 \right) + \frac{2}{|\alpha|} |\langle S^{\sharp_A} S z, S z \rangle_A| \\
&\quad (\text{by the AM-GM inequality}) \\
&= |\langle (S + S^{\sharp_A} S) z, z \rangle_A|^2 + \frac{\max\{1, |\alpha - 1|\}}{|\alpha|} (\langle S^{\sharp_A} S z, S^{\sharp_A} S z \rangle_A + \langle S^{\sharp_A} S z, z \rangle_A) \\
&\quad + \frac{2}{|\alpha|} |\langle z, S^{\sharp_A} S^2 z \rangle_A| \\
&= |\langle (S + S^{\sharp_A} S) z, z \rangle_A|^2 + \frac{\max\{1, |\alpha - 1|\}}{|\alpha|} \left(\langle (S^{\sharp_A} S)^2 z, z \rangle_A + \langle S^{\sharp_A} S z, z \rangle_A \right) \\
&\quad + \frac{2}{|\alpha|} |\langle z, S^{\sharp_A} S^2 z \rangle_A| \\
&\leq \omega_A^2(S^{\sharp_A} S + S) + \frac{2}{|\alpha|} \omega_A(S^{\sharp_A} S^2) + \frac{\max\{1, |\alpha - 1|\}}{|\alpha|} \left\| (S^{\sharp_A} S)^2 + S^{\sharp_A} S \right\|_A.
\end{aligned}$$

This gives the bound as desired. \square

Now from Theorem 2.2 (for $\alpha = 2$), we deduce the bound:

Corollary 2.3. *If $S \in \mathcal{B}_A(\mathcal{H})$ then*

$$d\omega_A^2(S) \leq \omega_A^2(S^{\sharp_A} S + S) + \omega_A(S^{\sharp_A} S^2) + \frac{1}{2} \left\| (S^{\sharp_A} S)^2 + S^{\sharp_A} S \right\|_A.$$

Again, considering $\alpha = n$ and $n \rightarrow \infty$ in Theorem 2.2, we get the bound:

Corollary 2.4. *If $S \in \mathcal{B}_A(\mathcal{H})$ then*

$$d\omega_A^2(S) \leq \omega_A^2(S^{\sharp_A} S + S) + \left\| (S^{\sharp_A} S)^2 + S^{\sharp_A} S \right\|_A.$$

To get next result we need the following semi-inner product inequality.

Lemma 2.5. [22] *If $a, b, c \in \mathcal{H}$ and $\|c\|_A = 1$ then*

$$|\langle a, c \rangle_A \langle c, b \rangle_A| \leq \frac{1}{2} (\|a\|_A \|b\|_A + |\langle a, b \rangle_A|) - \delta(a, b, c),$$

$$\text{where } \delta(a, b, c) = \begin{cases} \frac{\|b\|_A}{\|a\|_A} \left(|\langle a, c \rangle_A| \inf_{\lambda \in \mathbb{C}} \|c - \lambda b\|_A - \frac{1}{2} \inf_{\mu \in \mathbb{C}} \|a - \mu b\|_A \right)^2 & \text{if } \|a\|_A \|b\|_A \neq 0 \\ 0 & \text{if } \|a\|_A \|b\|_A = 0. \end{cases}$$

Theorem 2.6. *If $S \in \mathcal{B}_A(\mathcal{H})$ then*

$$d\omega_A^2(S) \leq \omega_A^2(S^{\sharp_A} S + S) + \omega_A(S^{\sharp_A} S^2) + \frac{1}{2} \left\| (S^{\sharp_A} S)^2 + S^{\sharp_A} S \right\|_A - 2 \inf_{\|z\|_A=1} \delta(S^{\sharp_A} S z, S z, z),$$

$$\text{where } \delta(a, b, c) = \begin{cases} \frac{\|b\|_A}{\|a\|_A} \left(|\langle a, c \rangle_A| \inf_{\lambda \in \mathbb{C}} \|c - \lambda b\|_A - \frac{1}{2} \inf_{\mu \in \mathbb{C}} \|a - \mu b\|_A \right)^2 & \text{if } \|a\|_A \|b\|_A \neq 0 \\ 0 & \text{if } \|a\|_A \|b\|_A = 0. \end{cases}$$

Proof. Take $z \in \mathcal{H}$ and $\|z\|_A = 1$. Using Lemma 2.5, we obtain

$$\begin{aligned}
& |\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \\
= & |\langle Sz, z \rangle_A + \langle Sz, Sz \rangle_A|^2 - 2\Re_A(\langle Sz, Sz \rangle_A \langle Sz, z \rangle_A) \\
\leq & |\langle (S + S^{\sharp_A} S) z, z \rangle_A|^2 + 2 |\langle S^{\sharp_A} Sz, z \rangle_A \langle z, Sz \rangle_A| \\
\leq & |\langle (S + S^{\sharp_A} S) z, z \rangle_A|^2 + \|S^{\sharp_A} Sz\|_A \|Sz\|_A + |\langle S^{\sharp_A} Sz, Sz \rangle_A| - 2\delta(S^{\sharp_A} Sz, Sz, z) \\
\leq & |\langle (S + S^{\sharp_A} S) z, z \rangle_A|^2 + \frac{1}{2} \left(\langle (S^{\sharp_A} S)^2 z, z \rangle_A + \langle S^{\sharp_A} Sz, z \rangle_A \right) \\
& + |\langle z, S^{\sharp_A} S^2 z \rangle_A| - 2\delta(T^{\sharp_A} Sz, Sz, z) \\
\leq & \omega_A^2 (S + S^{\sharp_A} S) + \omega_A (S^{\sharp_A} S^2) + \frac{1}{2} \left\| (S^{\sharp_A} S)^2 + S^{\sharp_A} S \right\|_A - 2\delta(S^{\sharp_A} Sz, Sz, z),
\end{aligned}$$

which implies the desired bound. \square

We now obtain both upper and lower bounds.

Theorem 2.7. *If $S \in \mathcal{B}_A(\mathcal{H})$, then*

$$\begin{aligned}
& \max \{ \omega_A(\Re_A(S) + iS^{\sharp_A} S), \omega_A(\Im_A(S) + iS^{\sharp_A} S) \} \leq d\omega_A(S) \\
\leq & \min \left\{ \sqrt{\omega_A^2(\Re_A(S) + iS^{\sharp_A} S) + \|\Im_A(S)\|_A^2}, \sqrt{\omega_A^2(\Im_A(S) + iS^{\sharp_A} S) + \|\Re_A(S)\|_A^2} \right\}.
\end{aligned}$$

Proof. Take $z \in \mathcal{H}$ and $\|z\|_A = 1$. We have

$$\begin{aligned}
|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 &= |\langle (\Re_A(S) + i\Im_A(S))z, z \rangle_A|^2 + \langle S^{\sharp_A} Sz, z \rangle_A^2 \\
&= |\langle \Re_A(S)z, z \rangle_A|^2 + |\langle \Im_A(S)z, z \rangle_A|^2 + \langle S^{\sharp_A} Sz, z \rangle_A^2 \\
&= |\langle (\Re_A(S) + iS^{\sharp_A} S)z, z \rangle_A|^2 + |\langle \Im_A(S)z, z \rangle_A|^2 \quad (2.1) \\
&\leq \omega_A^2(\Re_A(S) + iS^{\sharp_A} S) + \|\Im_A(S)\|_A^2.
\end{aligned}$$

This implies

$$d\omega_A^2(S) \leq \omega_A^2(\Re_A(S) + iS^{\sharp_A} S) + \|\Im_A(S)\|_A^2. \quad (2.2)$$

Similarly, we obtain

$$d\omega_A^2(S) \leq \omega_A^2(\Im_A(S) + iS^{\sharp_A} S) + \|\Re_A(S)\|_A^2. \quad (2.3)$$

Now combining the inequalities (2.2) and (2.3), we get the desired upper bound. Again from (2.1), we get

$$|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \geq |\langle (\Re_A(S) + iS^{\sharp_A} S)z, z \rangle_A|^2 \quad (2.4)$$

and

$$|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \geq |\langle (\Im_A(S) + iS^{\sharp_A} S)z, z \rangle_A|^2. \quad (2.5)$$

Now first taking the supremum over $\|z\|_A = 1$ and then combining them we obtain the desired lower bound. \square

From Theorem 2.7, we now obtain the result which gives the equality for the A -numerical radius.

Corollary 2.8. *If $S \in \mathcal{B}_A(\mathcal{H})$ with $(\Re_A(S^{\#A}))^2 = \Im_A(S^{\#A})$ then*

$$\omega_A(S) = \|\Re_A(S)\|_A \sqrt{1 + \|\Re_A(S)\|_A^2}.$$

Proof. First we suppose $S \in \mathcal{B}_A(\mathcal{H})$ with $S = S^{\#A}$. Then S is A -selfadjoint and so $d\omega_A(S) = \sqrt{\|S\|_A^2 + \|S\|_A^4}$ (see [3]). From Theorem 2.7, it follows $d\omega_A(S) = \omega_A(S + iS^2)$. Hence, $\omega_A(S + iS^2) = \sqrt{\|S\|_A^2 + \|S\|_A^4}$. Now substitute S by $\Re_A(S^{\#A})$ for $S \in \mathcal{B}_A(\mathcal{H})$, the result derives from the facts $\omega_A(S) = \omega_A(S^{\#A})$ and $\|\Re_A(S)\|_A = \|\Re_A(S^{\#A})\|_A$. \square

Remark 2.9. Theorem 2.7 is better than the existing bound in [3], namely

$$d\omega_A^2(S) \geq 2 \max\{\omega_A(S)c_A(S^{\#A}S), c_A(S)\|S\|_A^2\} \quad (2.6)$$

for some operators. If we assume $A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ and $S = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ then (2.6) gives $d\omega_A(S) \geq 0$, but Theorem 2.7 gives $d\omega_A(S) \geq \sqrt{2}$.

A generalization of the Cauchy-Schwarz inequality is as the lemma.

Lemma 2.10. [25] *If $S \in \mathcal{B}_A(\mathcal{H})$ and $x, z \in \mathcal{H}$ with $\|z\|_A = \|x\|_A = 1$ then*

$$|\langle Sx, z \rangle_A|^2 \leq \langle S^{\#A}Sx, x \rangle_A^{\frac{1}{2}} \langle SS^{\#A}z, z \rangle_A^{\frac{1}{2}}.$$

We now obtain an upper bound.

Theorem 2.11. *If $S \in \mathcal{B}_A(\mathcal{H})$ then*

$$d\omega_A^2(S) \leq \|S\|_A^4 + 2\|S\|_A^2 - \sqrt{c_A(S^{\#A}S)c_A(SS^{\#A})}.$$

Proof. Take $z \in \mathcal{H}$ and $\|z\|_A = 1$. We obtain

$$\begin{aligned} & |\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \\ & \leq \langle S^{\#A}Sz, z \rangle_A^{\frac{1}{2}} \langle SS^{\#A}z, z \rangle_A^{\frac{1}{2}} + \|Sz\|_A^4 \\ & \quad (\text{by Lemma 2.10}) \\ & \leq \frac{1}{2} (\langle S^{\#A}Sz, z \rangle_A + \langle SS^{\#A}z, z \rangle_A) + \|Sz\|_A^4 \\ & \quad (\text{by the AM-GM inequality}) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \left(\langle S^{\sharp_A} Sz, z \rangle_A^{\frac{1}{2}} + \langle SS^{\sharp_A} z, z \rangle_A^{\frac{1}{2}} \right)^2 - \langle S^{\sharp_A} Sz, z \rangle_A^{\frac{1}{2}} \langle SS^{\sharp_A} z, z \rangle_A^{\frac{1}{2}} + \|Sz\|_A^4 \\
&\leq \frac{1}{2} \left(\|S^{\sharp_A} Sz\|_A^{\frac{1}{2}} + \|SS^{\sharp_A} z\|_A^{\frac{1}{2}} \right)^2 + \|Sz\|_A^4 - \langle T^{\sharp_A} Sz, z \rangle_A^{\frac{1}{2}} \langle SS^{\sharp_A} z, z \rangle_A^{\frac{1}{2}} \\
&\leq \frac{1}{2} \left(\|S^{\sharp_A} S\|_A^{\frac{1}{2}} + \|SS^{\sharp_A}\|_A^{\frac{1}{2}} \right)^2 + \|S\|_A^4 - \sqrt{c_A(S^{\sharp_A} S) c_A(SS^{\sharp_A})} \\
&= 2\|S\|_A^2 + \|S\|_A^4 - \sqrt{c_A(S^{\sharp_A} S) c_A(SS^{\sharp_A})},
\end{aligned}$$

which gives the desired bound. \square

Remark 2.12. Theorem 2.11 is better than the existing bound in [13], namely

$$d\omega_A^2(S) \leq \max\{\|S\|_A^2, \|S\|_A^4\} + \omega_A(S^{\sharp_A} S^2) \quad (2.7)$$

for some operators. For example, if we consider $A = S = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ then (2.7) gives $d\omega_A^2(S) \leq 24$, whereas Theorem 2.11 gives $d\omega_A^2(S) \leq 23$.

We next obtain a bound which is a refinement of the existing bound given in [13].

Theorem 2.13. If $S \in \mathcal{B}_A(\mathcal{H})$ then

$$d\omega_A^2(S) \leq \omega_A^2(e^{i\theta} S + S^{\sharp_A} S) + 2\|S\|_A^2 \|\Re_A(e^{i\theta} S)\|_A, \quad \forall \theta \in \mathbb{R}.$$

Proof. For any $z \in \mathcal{H}$ and $\|z\|_A = 1$, we obtain

$$\begin{aligned}
|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 &= |\langle (S + S^{\sharp_A} S)z, z \rangle_A|^2 - 2\|Sz\|_A^2 \langle \Re_A(S)z, z \rangle_A \\
&\leq |\langle (S + S^{\sharp_A} S)z, z \rangle_A|^2 + 2\|Sz\|_A^2 \langle \Re_A(S)z, z \rangle_A \\
&\leq \omega_A^2(S + S^{\sharp_A} S) + 2\|S\|_A^2 \|\Re_A(S)\|_A.
\end{aligned}$$

This implies

$$d\omega_A^2(S) \leq \omega_A^2(S + S^{\sharp_A} S) + 2\|S\|_A^2 \|\Re_A(S)\|_A. \quad (2.8)$$

Replacing S by $e^{i\theta} S$, we obtain the desired bound. \square

Remark 2.14. Following Theorem 2.13, we get

$$d\omega_A^2(S) \leq \omega_A^2(S \pm S^{\sharp_A} S) + 2\|S\|_A^2 \|\Re_A(S)\|_A. \quad (2.9)$$

Clearly, (2.9) is sharper than the existing bound in [13], namely

$$d\omega_A^2(S) \leq \omega_A^2(S^{\sharp_A} S - S) + 2\|S\|_A^2 \|\omega_A(S)\|_A. \quad (2.10)$$

We now obtain a lower bound.

Theorem 2.15. If $S \in \mathcal{B}_A(\mathcal{H})$ then

$$d\omega_A^2(S) + 2\|S\|_A^2\|\Re_A(S)\|_A \geq \max\{\omega_A^2(S + S^{\sharp_A}S), \omega_A^2(S - S^{\sharp_A}T)\}.$$

Proof. Take $z \in \mathcal{H}$, $\|z\|_A = 1$, we get

$$\begin{aligned} |\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 &\geq |\langle (S + S^{\sharp_A}S)z, z \rangle_A|^2 - 2\|Sz\|_A^2|\langle \Re_A(S)z, z \rangle_A| \\ &\geq |\langle (S + S^{\sharp_A}S)z, z \rangle_A|^2 - 2\|S\|_A^2\|\Re_A(S)\|_A. \end{aligned}$$

From this we get

$$d\omega_A^2(S) + 2\|S\|_A^2\|\Re_A(S)\|_A \geq \omega_A^2(S + S^{\sharp_A}S). \quad (2.11)$$

Replacing S by $-S$, we get

$$d\omega_A^2(S) + 2\|S\|_A^2\|\Re_A(S)\|_A \geq \omega_A^2(S - S^{\sharp_A}S). \quad (2.12)$$

Combining (2.11) and (2.12), we obtain the bound as desired. \square

Remark 2.16. Theorem 2.15 is better than the existing bound in [3], namely

$$d\omega_A^2(S) \geq 2 \max\{\omega_A(S)c_A(S^{\sharp_A}S), c_A(S)\|S\|_A^2\} \quad (2.13)$$

for some operators. If we consider the same example as in Remark 2.12 then (2.13) gives $d\omega_A^2(S) \geq 8$ but Theorem 2.15 gives $d\omega_A^2(S) \geq 20$.

Now the following lemmas are needed.

Lemma 2.17. [10] If $S \in \mathcal{B}(\mathcal{H})$ is A -positive, and $z \in \mathcal{H}$, $\|z\|_A = 1$ then

$$\langle Sz, z \rangle_A^n \leq \langle S^n z, z \rangle_A \text{ for all } n \in \mathbb{N}.$$

Lemma 2.18. [22] If $x, z \in \mathcal{H}$ and $\|x\|_A \neq 0$ then

$$|\langle x, z \rangle_A| \leq \|z\|_A \left(\|x\|_A - \frac{\inf_{\lambda \in \mathbb{C}} \|x - \lambda z\|_A^2}{2\|x\|_A} \right).$$

We now develop a bound.

Theorem 2.19. Suppose $S \in \mathcal{B}_A(\mathcal{H})$. Then for all $\alpha \in [0, 1]$,

$$d\omega_A^2(S) \leq \left\| \alpha S^{\sharp_A}S + (1 - \alpha)SS^{\sharp_A} + (S^{\sharp_A}S)^2 \right\|_A - (\alpha\mu + (1 - \alpha)\eta),$$

where

$$\mu = \inf_{\|z\|_A=1} \left[\inf_{\lambda \in \mathbb{C}} \|Sz - \lambda z\|_A^2 - \frac{\inf_{\lambda \in \mathbb{C}} \|Sz - \lambda z\|_A^4}{4\|Sz\|_A^2} \right]$$

and

$$\eta = \inf_{\|z\|_A=1} \left[\inf_{\lambda \in \mathbb{C}} \|S^{\sharp_A} z - \lambda z\|_A^2 - \frac{\inf_{\lambda \in \mathbb{C}} \|S^{\sharp_A} z - \lambda z\|_A^4}{4 \|S^{\sharp_A} z\|_A^2} \right].$$

Proof. Take $z \in \mathcal{H}$, $\|z\|_A = 1$ and $\alpha \in [0, 1]$. By using lemma 2.18, we obtain

$$\begin{aligned} |\langle Sz, z \rangle_A|^2 &= \alpha |\langle Sz, z \rangle_A|^2 + (1 - \alpha) |\langle z, S^{\sharp_A} z \rangle_A|^2 \\ &\leq \alpha \left(\|Sz\|_A - \frac{\inf_{\lambda \in \mathbb{C}} \|Sz - \lambda z\|_A^2}{2 \|Sz\|_A} \right)^2 \\ &\quad + (1 - \alpha) \left(\|S^{\sharp_A} z\|_A - \frac{\inf_{\lambda \in \mathbb{C}} \|S^{\sharp_A} z - \lambda z\|_A^2}{2 \|S^{\sharp_A} z\|_A} \right)^2 \\ &= \alpha \|Sz\|_A^2 + (1 - \alpha) \|S^{\sharp_A} z\|_A^2 - \alpha \left(\inf_{\lambda \in \mathbb{C}} \|Sz - \lambda z\|_A^2 - \frac{\inf_{\lambda \in \mathbb{C}} \|Sz - \lambda z\|_A^4}{4 \|Sz\|_A^2} \right) \\ &\quad - (1 - \alpha) \left(\inf_{\lambda \in \mathbb{C}} \|S^{\sharp_A} z - \lambda z\|_A^2 - \frac{\inf_{\lambda \in \mathbb{C}} \|S^{\sharp_A} z - \lambda z\|_A^4}{4 \|S^{\sharp_A} z\|_A^2} \right) \\ &\leq \langle (\alpha S^{\sharp_A} S + (1 - \alpha) SS^{\sharp_A}) z, z \rangle_A - (\alpha \mu + (1 - \alpha) \eta). \end{aligned}$$

Therefore, we have

$$\begin{aligned} &|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \\ &= \langle (\alpha S^{\sharp_A} S + (1 - \alpha) SS^{\sharp_A}) z, z \rangle_A + \langle S^{\sharp_A} S z, z \rangle_A^2 - (\alpha \mu + (1 - \alpha) \eta) \\ &\leq \langle (\alpha S^{\sharp_A} S + (1 - \alpha) SS^{\sharp_A}) z, z \rangle_A + \langle (S^{\sharp_A} S)^2 z, z \rangle_A - (\alpha \mu + (1 - \alpha) \eta) \\ &\quad (\text{by Lemma 2.17}) \\ &\leq \left\| \alpha S^{\sharp_A} S + (1 - \alpha) SS^{\sharp_A} + (S^{\sharp_A} S)^2 \right\|_A - (\alpha \mu + (1 - \alpha) \eta). \end{aligned}$$

This gives

$$d\omega_A^2(S) \leq \left\| \alpha S^{\sharp_A} S + (1 - \alpha) SS^{\sharp_A} + (S^{\sharp_A} S)^2 \right\|_A - (\alpha \mu + (1 - \alpha) \eta). \quad (2.14)$$

□

Remark 2.20. (i) Form Theorem 2.19 (for $\alpha = 0$) we obtain

$$d\omega_A^2(S) \leq \|SS^{\sharp_A} + (S^{\sharp_A} S)^2\|_A - \eta. \quad (2.15)$$

(ii) The bound (2.15) is better than the existing bound in [13], namely

$$d\omega_A^2(S) \leq \sqrt{\omega_A((S^{\sharp_A} S)^2 + (S^{\sharp_A} S)^4)} + 2\omega_A^2(S^{\sharp_A} S^2) \quad (2.16)$$

for some operators. For example, if we consider $A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ and $S = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ then (2.16) gives $d\omega_A^2(S) \leq \frac{\sqrt{5}}{4}$ but (2.15) gives $d\omega_A^2(S) \leq \frac{1}{2}$.

We now present another upper bound.

Theorem 2.21. *If $S \in \mathcal{B}_A(\mathcal{H})$ then $d\omega_A^2(S) \leq \min \{\beta, \gamma\}$, where*

$$\beta = \min_{\alpha \in [0,1]} \left\{ \left\| \frac{\alpha}{4} S^{\sharp_A} S + \left(1 - \frac{3\alpha}{4}\right) S S^{\sharp_A} + (S^{\sharp_A} S)^2 \right\|_A + \frac{\alpha}{2} \omega_A(S^2) - (1-\alpha)\eta \right\},$$

$$\gamma = \min_{\alpha \in [0,1]} \left\{ \left\| \left(1 - \frac{3\alpha}{4}\right) S^{\sharp_A} S + \frac{\alpha}{4} S S^{\sharp_A} + (S^{\sharp_A} S)^2 \right\|_A + \frac{\alpha}{2} \omega_A(S^2) - (1-\alpha)\mu \right\}$$

and μ, η are as in Theorem 2.19.

Proof. Suppose $z \in \mathcal{H}$, $\|z\|_A = 1$ and $\alpha \in [0,1]$. Then we obtain

$$\begin{aligned} |\langle Sz, z \rangle_A|^2 &= \alpha |\langle Sz, z \rangle_A|^2 + (1-\alpha) |\langle z, T^{\sharp_A} z \rangle_A|^2 \\ &\leq \alpha |\langle Sz, z \rangle_A|^2 + (1-\alpha) \left(\|S^{\sharp_A} z\|_A - \frac{\inf_{\lambda \in \mathbb{C}} \|S^{\sharp_A} z - \lambda z\|_A^2}{2 \|S^{\sharp_A} z\|_A} \right)^2 \\ &\quad (\text{by Lemma 2.18}) \\ &= \alpha |\langle Sz, z \rangle_A|^2 + (1-\alpha) \|S^{\sharp_A} z\|_A^2 \\ &\quad - (1-\alpha) \left(\inf_{\lambda \in \mathbb{C}} \|S^{\sharp_A} z - \lambda z\|_A^2 - \frac{\inf_{\lambda \in \mathbb{C}} \|S^{\sharp_A} z - \lambda z\|_A^4}{4 \|S^{\sharp_A} z\|_A^2} \right) \\ &\leq \alpha |\langle Sz, z \rangle_A|^2 + (1-\alpha) \|S^{\sharp_A} z\|_A^2 - (1-\alpha)\eta. \end{aligned}$$

So,

$$\begin{aligned} &|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \\ &\leq \alpha |\langle Sz, z \rangle_A|^2 + (1-\alpha) \langle SS^{\sharp_A} z, z \rangle_A + \langle S^{\sharp_A} S z, z \rangle_A^2 - (1-\alpha)\eta \\ &\leq \alpha |\langle Sz, z \rangle_A|^2 + (1-\alpha) \langle SS^{\sharp_A} z, z \rangle_A + \langle (S^{\sharp_A} S)^2 z, z \rangle_A - (1-\alpha)\eta \\ &\quad (\text{by Lemma 2.17}) \\ &\leq \frac{\alpha}{2} (\|Sz\|_A \|S^{\sharp_A} z\|_A + |\langle Sz, S^{\sharp_A} z \rangle_A|) + (1-\alpha) \langle SS^{\sharp_A} z, z \rangle_A + \langle (S^{\sharp_A} S)^2 z, z \rangle_A \\ &\quad - (1-\alpha)\eta \quad (\text{by Lemma 2.1}) \\ &\leq \frac{\alpha}{4} (\|Sz\|_A^2 + \|S^{\sharp_A} z\|_A^2) + \frac{\alpha}{2} |\langle S^2 z, z \rangle_A| + (1-\alpha) \langle SS^{\sharp_A} z, z \rangle_A + \langle (S^{\sharp_A} S)^2 z, z \rangle_A \\ &\quad - (1-\alpha)\eta \quad (\text{by the AM-GM inequality}) \end{aligned}$$

$$\begin{aligned}
&= \left\langle \left(\frac{\alpha}{4} S^{\sharp_A} S + \left(1 - \frac{3\alpha}{4}\right) S S^{\sharp_A} + (S^{\sharp_A} S)^2 \right) z, z \right\rangle_A + \frac{\alpha}{2} |\langle S^2 z, z \rangle_A| - (1 - \alpha)\eta \\
&\leq \left\| \frac{\alpha}{4} S^{\sharp_A} S + \left(1 - \frac{3\alpha}{4}\right) S S^{\sharp_A} + (S^{\sharp_A} S)^2 \right\|_A + \frac{\alpha}{2} \omega_A(S^2) - (1 - \alpha)\eta.
\end{aligned}$$

This gives

$$d\omega_A^2(S) \leq \left\| \frac{\alpha}{4} S^{\sharp_A} S + \left(1 - \frac{3\alpha}{4}\right) S S^{\sharp_A} + (S^{\sharp_A} S)^2 \right\|_A + \frac{\alpha}{2} \omega_A(S^2) - (1 - \alpha)\eta.$$

Considering the minimum over $\alpha \in [0, 1]$, we get

$$d\omega_A^2(S) \leq \min_{\alpha \in [0, 1]} \left\{ \left\| \frac{\alpha}{4} S^{\sharp_A} S + \left(1 - \frac{3\alpha}{4}\right) S S^{\sharp_A} + (S^{\sharp_A} S)^2 \right\|_A + \frac{\alpha}{2} \omega_A(S^2) - (1 - \alpha)\eta \right\}. \quad (2.17)$$

Similarly, we get

$$d\omega_A^2(S) \leq \min_{\alpha \in [0, 1]} \left\{ \left\| \left(1 - \frac{3\alpha}{4}\right) S^{\sharp_A} S + \frac{\alpha}{4} S S^{\sharp_A} + (S^{\sharp_A} S)^2 \right\|_A + \frac{\alpha}{2} \omega_A(S^2) - (1 - \alpha)\mu \right\}. \quad (2.18)$$

Combining (2.17) and (2.18) we get the result as desired. \square

From Theorem 2.21 (for $\alpha = 1$) we obtain the bound:

Corollary 2.22. *If $S \in \mathcal{B}_A(\mathcal{H})$ then*

$$d\omega_A^2(S) \leq \frac{1}{4} \left\| S^{\sharp_A} S + S S^{\sharp_A} + 4(S^{\sharp_A} S)^2 \right\|_A + \frac{1}{2} \omega_A(S^2). \quad (2.19)$$

Remark 2.23. The bound (2.19) is better than the following bound in [13], namely

$$d\omega_A^2(S) \leq \frac{1}{2} [\omega_A((S^{\sharp_A} S)^2 + S^{\sharp_A} S) + \omega_A((S^{\sharp_A} S)^2 - S^{\sharp_A} S)] + \omega_A(S^{\sharp_A} S^2) \quad (2.20)$$

for some operator. If we consider the same example as in Remark 2.12, the bound (2.20) gives $d\omega_A^2(S) \leq 24$ whereas (2.19) gives $d\omega_A^2(S) \leq 20$.

To achieve the next result we need the inequality.

Lemma 2.24. [19] *If $a, c > 0$ and $0 \leq \alpha \leq 1$ then*

$$a^\alpha c^{1-\alpha} \leq \alpha a + (1 - \alpha)c \leq (\alpha a^r + (1 - \alpha)c^r)^{\frac{1}{r}}, \quad \text{for all } r \geq 1.$$

Theorem 2.25. *If $S \in \mathcal{B}_A(\mathcal{H})$ then*

$$d\omega_A^{4n}(S) \leq 4^{n-1} \left\| (S^{\sharp_A} S)^n + (S^{\sharp_A} S)^{2n} \right\|_A \left\| (S S^{\sharp_A})^n + (S^{\sharp_A} S)^{2n} \right\|_A \quad \text{for each } n \in \mathbb{N}.$$

Proof. Take $z \in \mathcal{H}$ and $\|z\|_A = 1$, we get

$$\begin{aligned}
& |\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \\
& \leq 2 \left(\frac{|\langle Sz, z \rangle_A|^{2n} + \|Sz\|_A^{4n}}{2} \right)^{\frac{1}{n}} \quad (\text{by Lemma 2.24}) \\
& = 2^{1-\frac{1}{n}} \left(|\langle Sz, z \rangle_A|^{2n} + \langle T^{\sharp_A} Sz, z \rangle_A^{2n} \right)^{\frac{1}{n}} \\
& \leq 2^{1-\frac{1}{n}} \left(\langle S^{\sharp_A} Sz, z \rangle_A^{\frac{n}{2}} \langle SS^{\sharp_A} z, z \rangle_A^{\frac{n}{2}} + \langle S^{\sharp_A} Sz, z \rangle_A^{2n} \right)^{\frac{1}{n}} \quad (\text{by Lemma 2.10}) \\
& \leq 2^{1-\frac{1}{n}} \left[\left(\langle S^{\sharp_A} Sz, z \rangle_A^n + \langle S^{\sharp_A} Sz, z \rangle_A^{2n} \right) \left(\langle SS^{\sharp_A} z, z \rangle_A^n + \langle S^{\sharp_A} Sz, z \rangle_A^{2n} \right) \right]^{\frac{1}{2n}} \\
& \quad (\text{since } (ab + cd)^2 \leq (a^2 + c^2)(b^2 + d^2) \text{ with } a, b, c, d \geq 0) \\
& \leq 2^{1-\frac{1}{n}} \left(\langle (S^{\sharp_A} S)^n z, z \rangle_A + \langle (S^{\sharp_A} S)^{2n} z, z \rangle_A \right)^{\frac{1}{2n}} \\
& \quad \times \left(\langle (SS^{\sharp_A})^n z, z \rangle_A + \langle (SS^{\sharp_A})^{2n} z, z \rangle_A \right)^{\frac{1}{2n}} \quad (\text{by Lemma 2.17}) \\
& \leq 2^{1-\frac{1}{n}} \left(\|(S^{\sharp_A} S)^n + (S^{\sharp_A} S)^{2n}\|_A \|(SS^{\sharp_A})^n + (SS^{\sharp_A})^{2n}\|_A \right)^{\frac{1}{2n}}.
\end{aligned}$$

Thus,

$$d\omega_A^2(S) \leq 2^{1-\frac{1}{n}} \left(\|(S^{\sharp_A} S)^n + (S^{\sharp_A} S)^{2n}\|_A \|(SS^{\sharp_A})^n + (SS^{\sharp_A})^{2n}\|_A \right)^{\frac{1}{2n}}.$$

This gives the bound. \square

From Theorem 2.25 (for $n = 1$), we obtain the bound:

Corollary 2.26. *If $S \in \mathcal{B}_A(\mathcal{H})$ then*

$$d\omega_A^2(S) \leq \|SS^{\sharp_A} + (S^{\sharp_A} S)^2\|_A^{\frac{1}{2}} \|S^{\sharp_A} S + (S^{\sharp_A} S)^2\|_A^{\frac{1}{2}}. \quad (2.21)$$

Remark 2.27. The bound (2.21) is better than the existing bound in [3], namely

$$d\omega_A^2(S) \leq \|S^{\sharp_A} S + (S^{\sharp_A} S)^{\sharp_A} S^{\sharp_A} S\|_A \quad (2.22)$$

for some operators. For the same example as in Remark 2.20, the bound (2.22) gives $d\omega_A^2(S) \leq \frac{3}{4}$, but (2.21) gives $d\omega_A^2(S) \leq \frac{\sqrt{3}}{2\sqrt{2}}$.

We now obtain a lower bound.

Theorem 2.28. *If $S \in \mathcal{B}_A(\mathcal{H})$ then*

$$d\omega_A^2(S) \geq \max \{c_A^2(S)(1 + \|S\|_A^2), \omega_A^2(S)(1 + c_A^2(S^{\sharp_A} S))\}.$$

Proof. Suppose $z \in \mathcal{H}$ and $\|z\|_A = 1$. Then

$$|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \geq |\langle Sz, z \rangle_A|^2 + |\langle Sz, z \rangle_A|^2 \|Sz\|_A^2 \geq c_A^2(S) (1 + \|Sz\|_A^2).$$

This implies

$$d\omega_A^2(S) \geq c_A^2(S) (1 + \|S\|_A^2). \quad (2.23)$$

We also have

$$|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \geq |\langle Sz, z \rangle_A|^2 (1 + \|Sz\|_A^2) \geq |\langle Sz, z \rangle_A|^2 (1 + c_A^2(S^\#_A S)).$$

From this we get

$$d\omega_A^2(S) \geq \omega_A^2(S) (1 + c_A^2(S^\#_A S)). \quad (2.24)$$

Combining (2.23) and (2.24), we get

$$d\omega_A^2(S) \geq \max \{c_A^2(S) (1 + \|S\|_A^2), \omega_A^2(S) (1 + c_A^2(S^\#_A S))\}.$$

This completes the proof. \square

We conclude this section by giving another lower bound.

Theorem 2.29. Let $S \in \mathcal{B}_A(\mathcal{H})$ and $m_A(S) = \inf_{\|x\|_A=1} \|Sz\|_A$. Then

$$d\omega_A^2(S) \geq \max \{(1 + m_A^2(S)) \omega_A^2(S), (1 + \|S\|_A^2) c_A^2(S)\}.$$

Proof. Take $z \in \mathcal{H}$ and $\|z\|_A = 1$. We obtain

$$|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \geq |\langle Sz, z \rangle_A|^2 + |\langle Sz, z \rangle_A|^2 m_A^2(S) = (1 + m_A^2(S)) |\langle Sz, z \rangle_A|^2.$$

This gives

$$d\omega_A^2(S) \geq (1 + m_A^2(S)) \omega_A^2(S). \quad (2.25)$$

Moreover, we see

$$|\langle Sz, z \rangle_A|^2 + \|Sz\|_A^4 \geq (1 + \|Sz\|_A^2) |\langle Sz, z \rangle_A|^2 \geq (1 + \|Sz\|_A^2) c_A^2(S).$$

So

$$d\omega_A^2(S) \geq (1 + \|S\|_A^2) c_A^2(S). \quad (2.26)$$

Combining (2.25) and (2.26), we get the result as desired. \square

It should be mentioned here that the bound in Theorem 2.29 is also studied in [5] in the setting of Hilbert space operators.

3. BOUNDS FOR THE A_0 -DAVIS-WIELANDT RADIUS OF BLOCK MATRICES

Consider 2×2 block diagonal matrix $A_0 = \begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix}$. Clearly, $A_0 \in \mathcal{B}(\mathcal{H} \oplus \mathcal{H})$ is positive. Therefore, A_0 induces $\langle x, z \rangle_{A_0} = \langle A_0 x, z \rangle = \langle x_1, z_1 \rangle_A + \langle x_2, z_2 \rangle_A$, for all $x = (x_1, x_2)$ and $z = (z_1, z_2) \in \mathcal{H} \oplus \mathcal{H}$. If $S_{ij} \in \mathcal{B}_A(\mathcal{H})$ for $i, j = 1, 2$, then $\begin{bmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{bmatrix} \in \mathcal{B}_{A_0}(\mathcal{H} \oplus \mathcal{H})$ and $\begin{bmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{bmatrix}^{\sharp_{A_0}} = \begin{bmatrix} S_{11}^{\sharp_A} & S_{21}^{\sharp_A} \\ S_{12}^{\sharp_A} & S_{22}^{\sharp_A} \end{bmatrix}$, shown in [26]. To start our analysis we begin with the lemmas.

Lemma 3.1. [3] *If $S \in \mathcal{B}_A(\mathcal{H})$ and V is A -unitary, then $d\omega_A(V^{\sharp_A} SV) = d\omega_A(S)$.*

Lemma 3.2. [26] *If $S, T \in \mathcal{B}_A(\mathcal{H})$ then $\omega_{A_0}\left(\begin{bmatrix} S & 0 \\ 0 & T \end{bmatrix}\right) = \max\{\omega_A(S), \omega_A(T)\}$.*

Lemma 3.3. [26] *If $S, T \in \mathcal{B}_A(\mathcal{H})$ then $\omega_{A_0}\left(\begin{bmatrix} S & T \\ T & S \end{bmatrix}\right) = \max\{\omega_A(S+T), \omega_A(S-T)\}$.*

In particular, $\omega_{A_0}\left(\begin{bmatrix} 0 & S \\ S & 0 \end{bmatrix}\right) = \omega_A(S)$.

We now prove our results.

Proposition 3.4. *If $S, T \in \mathcal{B}_A(\mathcal{H})$ then $d\omega_{A_0}\left(\begin{bmatrix} S & 0 \\ 0 & T \end{bmatrix}\right) = \max\{d\omega_A(S), d\omega_A(T)\}$.*

Proof. Let $X = \begin{bmatrix} T & 0 \\ 0 & S \end{bmatrix}$. Suppose $u = (z, 0) \in \mathcal{H} \oplus \mathcal{H}$, $\|z\|_A = 1$. Clearly $\|u\|_{A_0} = 1$. Then $|\langle Tz, z \rangle_A|^2 + \|Tz\|_A^4 = |\langle Xu, u \rangle_{A_0}|^2 + \|Xu\|_{A_0}^4 \leq d\omega_{A_0}^2(X)$. Hence, we obtain $d\omega_A^2(T) \leq d\omega_{A_0}^2(X)$. Similarly, we can see that $d\omega_A^2(S) \leq d\omega_{A_0}^2(X)$. Therefore, we infer that $\max\{d\omega_A(T), d\omega_A(S)\} \leq d\omega_{A_0}(X)$.

Now, let $v = (z, y) \in \mathcal{H} \oplus \mathcal{H}$ and $\|v\|_{A_0} = 1$, then

$$\begin{aligned} \sqrt{|\langle Xv, v \rangle_{A_0}|^2 + \|Xv\|_{A_0}^4} &= \sqrt{|\langle Tz, z \rangle_A + \langle Sy, y \rangle_A|^2 + \|Tz\|_A^4 + \|Sy\|_A^4} \\ &\leq \sqrt{|\langle Tz, z \rangle_A|^2 + \|Tz\|_A^4} + \sqrt{|\langle Sy, y \rangle_A|^2 + \|Sy\|_A^4} \\ &\leq d\omega_A(T) \|z\|_A^2 + d\omega_A(S) \|y\|_A^2 \\ &\leq \max\{d\omega_A(T), d\omega_A(S)\}. \end{aligned}$$

This gives $d\omega_{A_0}(X) \leq \max\{d\omega_A(T), d\omega_A(S)\}$. \square

We now discuss the following equality.

Lemma 3.5. If $S, T \in \mathcal{B}_A(\mathcal{H})$ then

$$d\omega_{A_0} \left(\begin{bmatrix} S & T \\ T & S \end{bmatrix} \right) = d\omega_{A_0} \left(\begin{bmatrix} S-T & 0 \\ 0 & S+T \end{bmatrix} \right).$$

In particular,

$$d\omega_{A_0} \left(\begin{bmatrix} 0 & S \\ S & 0 \end{bmatrix} \right) = d\omega_A(S). \quad (3.1)$$

Proof. Let $U = \frac{1}{\sqrt{2}} \begin{bmatrix} I & I \\ -I & I \end{bmatrix}$ and $W = \begin{bmatrix} S & T \\ T & S \end{bmatrix}$. Then $\begin{bmatrix} P(S-T) & 0 \\ 0 & P(S+T) \end{bmatrix} = U^{\sharp_{A_0}} W U$. Using Lemma 3.1, we get $d\omega_{A_0}(U^{\sharp_{A_0}} W U) = d\omega_{A_0}(W)$. So, $d\omega_{A_0} \left(\begin{bmatrix} S & T \\ T & S \end{bmatrix} \right) = d\omega_{A_0} \left(\begin{bmatrix} P(S-T) & 0 \\ 0 & P(S+T) \end{bmatrix} \right)$. Taking $U = I$ in Lemma 3.1, we can observe that $d\omega_A(P\bar{S}) = d\omega_A(S)$. So,

$$d\omega_{A_0} \left(\begin{bmatrix} S & T \\ T & S \end{bmatrix} \right) = d\omega_{A_0} \left(\begin{bmatrix} P(S-T) & 0 \\ 0 & P(S+T) \end{bmatrix} \right) = d\omega_{A_0} \left(\begin{bmatrix} S-T & 0 \\ 0 & S+T \end{bmatrix} \right).$$

In particular, we get

$$d\omega_{A_0} \left(\begin{bmatrix} 0 & S \\ S & 0 \end{bmatrix} \right) = d\omega_{A_0} \left(\begin{bmatrix} -S & 0 \\ 0 & S \end{bmatrix} \right) = \max \{d\omega_A(-S), d\omega_A(S)\} = d\omega_A(S).$$

□

We now develop a bound of A_0 -Davis–Wielandt radius for $\begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix}$ in $\mathcal{B}_A(\mathcal{H} \oplus \mathcal{H})$.

Theorem 3.6. If $C, B \in \mathcal{B}_A(\mathcal{H})$ then

$$\begin{aligned} & d\omega_{A_0}^2 \left(\begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix} \right) \\ & \leq \frac{1}{4} \max \left\{ \omega_A \left(BB^{\sharp_A} + C^{\sharp_A}C + 4(C^{\sharp_A}C)^2 \right), \omega_A \left(B^{\sharp_A}B + CC^{\sharp_A} + 4(B^{\sharp_A}B)^2 \right) \right\} \\ & \quad + \frac{1}{2} \max \{ \omega_A(BC), \omega_A(CB) \}. \end{aligned}$$

Proof. Let $S = \begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix}$ and $z \in \mathcal{H} \oplus \mathcal{H}$ with $\|z\|_{A_0} = 1$. Then

$$\begin{aligned} & |\langle Sz, z \rangle_{A_0}|^2 + \|Sz\|_{A_0}^4 \\ & \leq \frac{1}{2} \left(\|Sz\|_{A_0} \|S^{\sharp_A}z\|_{A_0} + \left| \langle Sz, S^{\sharp_A}z \rangle_{A_0} \right| \right) + \|Sz\|_{A_0}^4 \quad (\text{by Lemma 2.1}) \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{4} \left(\|Sz\|_{A_0}^2 + \|S^{\sharp_A} z\|_{A_0}^2 \right) + \frac{1}{2} \left| \langle S^2 z, z \rangle_{A_0} \right| + \|Sz\|_{A_0}^4 \quad (\text{by AM-GM inequality}) \\
&\leq \frac{1}{4} \langle (S^{\sharp_A} S + SS^{\sharp_A}) z, z \rangle_{A_0} + \langle (S^{\sharp_A} S)^2 z, z \rangle_{A_0} + \frac{1}{2} \left| \langle S^2 z, z \rangle_{A_0} \right| \quad (\text{by Lemma 2.17}) \\
&= \frac{1}{4} \langle (S^{\sharp_A} S + SS^{\sharp_A} + 4(S^{\sharp_A} S)^2) z, z \rangle_{A_0} + \frac{1}{2} \left| \langle S^2 z, z \rangle_{A_0} \right| \\
&\leq \frac{1}{4} \omega_{A_0} (S^{\sharp_A} S + SS^{\sharp_A} + 4(S^{\sharp_A} S)^2) + \frac{1}{2} \omega_{A_0} (S^2) \\
&= \frac{1}{4} \omega_{A_0} \left(\begin{bmatrix} BB^{\sharp_A} + C^{\sharp_A} C + 4(C^{\sharp_A} C)^2 & 0 \\ 0 & B^{\sharp_A} B + CC^{\sharp_A} + 4(B^{\sharp_A} B)^2 \end{bmatrix} \right) \\
&\quad + \frac{1}{2} \omega_{A_0} \left(\begin{bmatrix} BC & 0 \\ 0 & CB \end{bmatrix} \right) \\
&= \frac{1}{4} \max \left\{ \omega_A (BB^{\sharp_A} + C^{\sharp_A} C + 4(C^{\sharp_A} C)^2), \omega_A (B^{\sharp_A} B + CC^{\sharp_A} + 4(B^{\sharp_A} B)^2) \right\} \\
&\quad + \frac{1}{2} \max \{ \omega_A (BC), \omega_A (CB) \} \quad (\text{by Lemma 3.2}).
\end{aligned}$$

This implies the bound as desired. \square

We now need the lemma.

Lemma 3.7. [13] If $x, u, v \in \mathcal{H}$ then

$$|\langle x, v \rangle_A|^2 + |\langle x, u \rangle_A|^2 \leq \|x\|_A^2 \left(\max \{ \|v\|_A^2, \|u\|_A^2 \} + |\langle v, u \rangle_A| \right).$$

We next obtain another upper bound of A_0 -Davis–Wielandt radius of $\begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix}$.

Theorem 3.8. If $C, B \in \mathcal{B}_A(\mathcal{H})$, then

$$\begin{aligned}
&d\omega_{A_0}^2 \left(\begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix} \right) \\
&\leq \frac{1}{2} \max \left\{ \omega_A ((C^{\sharp_A} C)^2 + C^{\sharp_A} C), \omega_A (B^{\sharp_A} B + (B^{\sharp_A} B)^2) \right\} \\
&\quad + \frac{1}{2} \max \left\{ \omega_A ((C^{\sharp_A} C)^2 - C^{\sharp_A} C), \omega_A (B^{\sharp_A} B - (B^{\sharp_A} B)^2) \right\} \\
&\quad + \omega_{A_0} \left(\begin{bmatrix} 0 & C^{\sharp_A} CB \\ B^{\sharp_A} BC & 0 \end{bmatrix} \right).
\end{aligned}$$

Proof. Let $S = \begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix}$ and $z \in \mathcal{H} \oplus \mathcal{H}$ with $\|z\|_{A_0} = 1$. Replacing $x = z$, $v = Sz$, $u = S^{\sharp_A}Sz$ in Lemma 3.7, we obtain

$$\begin{aligned}
& |\langle Sz, z \rangle_{A_0}|^2 + \|Sz\|_{A_0}^4 = |\langle z, Sz \rangle_{A_0}|^2 + \langle z, S^{\sharp_A}Sz \rangle_{A_0}^2 \\
& \leq \|z\|_{A_0}^2 \left(\max \left\{ \|Sz\|_{A_0}^2, \|S^{\sharp_A}Sz\|_{A_0}^2 \right\} + |\langle Sz, S^{\sharp_A}Sz \rangle_{A_0}| \right) \\
& = \frac{1}{2} \left(\|Sz\|_{A_0}^2 + \|S^{\sharp_A}Sz\|_{A_0}^2 + \left| \|Sz\|_{A_0}^2 - \|S^{\sharp_A}Sz\|_{A_0}^2 \right| \right) + |\langle S^2z, Sz \rangle_{A_0}| \\
& \quad (\text{since } \max\{a, b\} = \frac{1}{2}(a + b + |a - b|) \text{ for real scalars } a, b) \\
& = \frac{1}{2} \left(\left\langle \left(S^{\sharp_A}S + (S^{\sharp_A}S)^2 \right) z, z \right\rangle_{A_0} + \left| \left\langle \left(S^{\sharp_A}S - (S^{\sharp_A}S)^2 \right) z, z \right\rangle_{A_0} \right| \right) + |\langle S^{\sharp_A}S^2z, z \rangle_{A_0}| \\
& \leq \frac{1}{2} \left[\omega_{A_0} \left(S^{\sharp_A}S + (S^{\sharp_A}S)^2 \right) + \omega_{A_0} \left(S^{\sharp_A}S - (S^{\sharp_A}S)^2 \right) \right] + \omega_{A_0} (S^{\sharp_A}S^2) \\
& = \frac{1}{2} \omega_{A_0} \left(\begin{bmatrix} (C^{\sharp_A}C)^2 + C^{\sharp_A}C & 0 \\ 0 & B^{\sharp_A}B + (B^{\sharp_A}B)^2 \end{bmatrix} \right) \\
& \quad + \frac{1}{2} \omega_{A_0} \left(\begin{bmatrix} (C^{\sharp_A}C)^2 - C^{\sharp_A}C & 0 \\ 0 & (B^{\sharp_A}B)^2 - B^{\sharp_A}B \end{bmatrix} \right) + \omega_{A_0} \left(\begin{bmatrix} 0 & C^{\sharp_A}CB \\ B^{\sharp_A}BC & 0 \end{bmatrix} \right) \\
& = \frac{1}{2} \max \left\{ \omega_A \left((C^{\sharp_A}C)^2 + C^{\sharp_A}C \right), \omega_A \left(B^{\sharp_A}B + (B^{\sharp_A}B)^2 \right) \right\} \\
& \quad + \frac{1}{2} \max \left\{ \omega_A \left((C^{\sharp_A}C)^2 - C^{\sharp_A}C \right), \omega_A \left(B^{\sharp_A}B - (B^{\sharp_A}B)^2 \right) \right\} \\
& \quad + \omega_{A_0} \left(\begin{bmatrix} 0 & C^{\sharp_A}CB \\ B^{\sharp_A}BC & 0 \end{bmatrix} \right) \quad (\text{by Lemma 3.2}).
\end{aligned}$$

This implies the inequality as desired. \square

Remark 3.9. Considering $B = C$ in Theorem 3.8, and using (3.1) and Lemma 3.3, we derive the existing inequality (see [13, Corollary 2.10])

$$d\omega_A^2(B) \leq \frac{1}{2} \left[\omega_A \left(B^{\sharp_A}B + (B^{\sharp_A}B)^2 \right) + \omega_A \left(B^{\sharp_A}B - (B^{\sharp_A}B)^2 \right) \right] + \omega_A (B^{\sharp_A}B^2)$$

and so the inequality in Theorem 3.8 is a generalization of [13, Corollary 2.10].

To develop our final result we need the lemma.

Lemma 3.10. *If $x, u, v \in \mathcal{H}$ then*

$$|\langle x, u \rangle_A|^2 + |\langle x, v \rangle_A|^2 \leq \|x\|_A^2 \sqrt{|\langle u, u \rangle_A|^2 + 2|\langle u, v \rangle_A|^2 + |\langle v, v \rangle_A|^2}.$$

Theorem 3.11. *If $C, B \in \mathcal{B}_A(\mathcal{H})$ then*

$$\begin{aligned} d\omega_{A_0}^4 \left(\begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix} \right) &\leq \max \left\{ \omega_A \left((C^{\sharp_A} C)^2 + (C^{\sharp_A} C)^4 \right), \omega_A \left((B^{\sharp_A} B)^2 + (B^{\sharp_A} B)^4 \right) \right\} \\ &\quad + 2\omega_{A_0}^2 \left(\begin{bmatrix} 0 & C^{\sharp_A} CB \\ B^{\sharp_A} BC & 0 \end{bmatrix} \right). \end{aligned}$$

Proof. Let $S = \begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix}$ and $z \in \mathcal{H} \oplus \mathcal{H}$ with $\|z\|_{A_0} = 1$, then

$$\begin{aligned} &\left(|\langle Sz, z \rangle_{A_0}|^2 + \|Sz\|_{A_0}^4 \right)^2 = \left(|\langle z, Sz \rangle_{A_0}|^2 + \left| \langle z, S^{\sharp_A} Sz \rangle_{A_0} \right|^2 \right)^2 \\ &\leq \|z\|_{A_0}^4 \left(|\langle Sz, Sz \rangle_{A_0}|^2 + 2 \left| \langle Sz, S^{\sharp_A} Sz \rangle_{A_0} \right|^2 + \left| \langle S^{\sharp_A} Sz, S^{\sharp_A} Sz \rangle_{A_0} \right|^2 \right) \text{ (by Lemma 3.10)} \\ &= \left| \langle S^{\sharp_A} Sz, z \rangle_{A_0} \right|^2 + 2 \left| \langle S^{\sharp_A} S^2 z, z \rangle_{A_0} \right|^2 + \left| \langle (S^{\sharp_A} S)^2 z, z \rangle_{A_0} \right|^2 \\ &\leq \left\langle (S^{\sharp_A} S)^2 z, z \right\rangle_{A_0} + \left\langle (S^{\sharp_A} S)^4 z, z \right\rangle_{A_0} + 2 \left| \langle S^{\sharp_A} S^2 z, z \rangle_{A_0} \right|^2 \text{ (by Lemma 2.17)} \\ &\leq \omega_{A_0} \left((S^{\sharp_A} S)^2 + (S^{\sharp_A} S)^4 \right) + 2\omega_{A_0}^2 (S^{\sharp_A} S^2) \\ &= \omega_{A_0} \left(\begin{bmatrix} (C^{\sharp_A} C)^2 + (C^{\sharp_A} C)^4 & 0 \\ 0 & (B^{\sharp_A} B)^2 + (B^{\sharp_A} B)^4 \end{bmatrix} \right) + 2\omega_{A_0}^2 \left(\begin{bmatrix} 0 & C^{\sharp_A} CB \\ B^{\sharp_A} BC & 0 \end{bmatrix} \right) \\ &= \max \left\{ \omega_A \left((C^{\sharp_A} C)^2 + (C^{\sharp_A} C)^4 \right), \omega_A \left((B^{\sharp_A} B)^2 + (B^{\sharp_A} B)^4 \right) \right\} \\ &\quad + 2\omega_{A_0}^2 \left(\begin{bmatrix} 0 & C^{\sharp_A} CB \\ B^{\sharp_A} BC & 0 \end{bmatrix} \right) \text{ (by Lemma 3.2).} \end{aligned}$$

Therefore, this gives the inequality as desired. □

Remark 3.12. Considering $B = C$ in Theorem 3.11, and using (3.1) and Lemma 3.3, we derive the existing inequality (see [13, Corollary 2.7])

$$d\omega_A^4(B) \leq \omega_A \left((B^{\sharp_A} B)^2 + (B^{\sharp_A} B)^4 \right) + 2\omega_A^2 (B^{\sharp_A} B^2)$$

and so Theorem 3.11 is a generalization of [13, Corollary 2.7].

Declarations. All authors contributed to the preparation of the article. All authors read and approved the final manuscript.

Competing interests. Authors declare that there is no competing interest.

REFERENCES

- [1] M.L. Arias, G. Corach and M.C. Gonzalez, Metric properties of projections in semi-Hilbertian spaces, *Integr. Equ. Oper. Theory*, 62(1) (2008), 11-28.
- [2] M.L. Arias, G. Corach and M.C. Gonzalez, Partial isometries in semi-Hilbertian spaces, *Linear Algebra Appl.*, 428(7) (2008), 1460-1475.
- [3] A. Bhanja, P. Bhunia and K. Paul, On generalized Davis-Wielandt radius inequalities of semi-Hilbertian space operators, *Oper. Matrices*, 15(4) (2021), 1201-1225.
- [4] P. Bhunia, A. Sen, S. Barik and K. Paul, Berezin number and Berezin norm inequalities for operator matrices, *Linear Multilinear Algebra*, (2024). <https://doi.org/10.1080/03081087.2023.2299388>
- [5] P. Bhunia, A. Bhanja and K. Paul, New Inequalities for Davis-Wielandt Radius of Hilbert Space Operators, *Bull. Malays. Math. Sci. Soc.*, 44 (5) (2021), 3523-3539.
- [6] P. Bhunia, A. Bhanja, S. Bag and K. Paul, Bounds for the Davis-Wielandt radius of bounded linear operators, *Ann. Funct. Anal.*, 12 (1) (2021), 23 pp.
- [7] P. Bhunia, K. Paul and R.K. Nayak, On inequalities for A -numerical radius of operators, *Electron. J. Linear Algebra*, 36 (2020), 143-157.
- [8] P. Bhunia and R.K. Nayak, Refinements of A -numerical radius inequalities and their applications, *Adv. Oper. Theory*, 5 (4) (2020), 1498-1511.
- [9] P. Bhunia, D. Sain and K. Paul, On the Davis-Wielandt shell of an operator and the Davis-Wielandt index of a normed linear space, *Collect. Math.*, 73 (3) (2020), 521-533.
- [10] C. Conde and K. Feki, On some inequalities for the generalized joint numerical radius of semi-Hilbert space operators, *Ric. Mat.*, 73 (2024), no.2, 661-679.
- [11] R.G. Douglas, On majorization, factorization and range inclusion of operators in Hilbert space, *Proc. Amer. Math. Soc.*, 17 (1966), 413-416.
- [12] S.S. Dragomir, Some inequalities for the norm and the numerical radius of linear operator in Hilbert spaces, *Tamkang J. Math.*, 39(1) (2008), 1-7.
- [13] K. Feki, Inequalities for the A -joint numerical radius of two operators and their application, *Hacet. J. Math. Stat.*, 53 (2024), no.1, 22-39.
- [14] M. Guesba, A -numerical radius inequalities via Dragomir inequalities and related results, *Novi Sad J. Math.*, (2023). <https://doi.org/10.30755/NSJOM.15867>
- [15] M. Guesba, P. Bhunia and K. Paul, A -numerical radius inequalities and A -translatable radii of semiHilbert space operators, *Filomat*, 37 (11) (2023), 3443-3456.
- [16] M. Guesba, On some numerical radius inequalities for normal operators in Hilbert spaces, *J. Interdiscip. Math.*, 25 (2) (2022), 463-470.
- [17] M. Guesba, Some generalizations of A -numerical radius inequalities for semi-Hilbert space operators, *Boll. Unione Mat. Ital.*, 14 (4) (2021), 681-692.
- [18] K.E. Gustafson and D.K.M. Rao, *Numerical Range*, Springer-Verlag, New York, (1997).
- [19] G.H. Hardy, J.E. Littlewood and G. Polya, *Inequalities*, 2nd ed, Cambridge University Press, Cambridge, (1988).
- [20] F. Kittaneh, M.S. Moslehian and T. Yamazaki, Cartesian decomposition and numerical radius inequalities, *Linear Algebra Appl.*, 471 (2015), 46-53.

- [21] F. Kittaneh and A. Zamani, Bounds for A -numerical radius based on an extension of A-Buzano inequality, *J. Comput. Appl. Math.*, 426 (2023), Paper No. 115070, 14 pp.
- [22] F. Kittaneh and A. Zamani, A refinement of A-Buzano inequality and applications to A-numerical radius inequalities, *Linear Algebra Appl.*, (2023). <https://doi.org/10.1016/j.laa.2023.02.020>
- [23] C.K. Li, Y.T. Poon and N.S. Sze, Davis-Wielandt shells of operators, *Oper. Matrices*, 2 (3) (2008), 341-355.
- [24] M.S. Moslehian, Q. Xu and A. Zamani, Seminorm and numerical radius inequalities of operators in semi-Hilbertian spaces, *Linear Algebra Appl.*, 591 (2020), 299-321.
- [25] H. Qiao, G. Hai and E. Bai, A-numerical radius and A-norm inequalities for semi-Hilbertian space operators, *Linear Multilinear Algebra*, 70 (21) (2022), 6891-6907.
- [26] N.C. Rout, S. Sahoo and D. Mishra, On A -numerical radius inequalities for 2×2 operator matrices, *Linear Multilinear Algebra*, 70 (2022), no.14, 2672–2692.
- [27] A. Saddi, A -normal operators in semi-Hilbertian spaces, *Aust. J. Math. Anal. Appl.*, 9 (1) (2012), 1-12.
- [28] K. Shebrawi and H. Albadawi, Numerical Radius and Operator Norm Inequalities, *J. Inequal. Appl.*, (2009), Article ID 492154, 11 pp.
- [29] A. Zamani and K. Shebrawi, Some upper bounds for the Davis-Wielandt radius of Hilbert space operators, *Mediterr. J. Math.*, 17 (2020), no.1, Paper No. 25, 13 pp.
- [30] A. Zamani, A -numerical radius inequalities for semi-Hilbertian space operators, *Linear Algebra Appl.*, 578 (2019), 159-183.
- [31] A. Zamani, M.S. Moslehian, M.-T. Chien and H. Nakazato, Norm-parallelism and the Davis-Wielandt radius of Hilbert space operators, *Linear Multilinear Algebra*, 67(11) (2019), 2147-2158.

(Guesba) FACULTY OF EXACT SCIENCES, DEPARTMENT OF MATHEMATICS, EL OUED UNIVERSITY, 39000 ALGERIA

Email address: guesbamessaoud2@gmail.com

(Barik) DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, WEST BENGAL, INDIA

Email address: bariksomdatta97@gmail.com

(Bhunia) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BENGALURU 560012, KARNATAKA, INDIA

Email address: pintubhunia5206@gmail.com

(Paul) DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, WEST BENGAL, INDIA

Email address: kalloldada@gmail.com