PRIME IDEALS OF MOH AND THE CHARACTERISTIC OF THE FIELD

LAURA GONZÁLEZ AND FRANCESC PLANAS-VILANOVA

ABSTRACT. We reprove and generalize a result of Moh which bounds the minimal number of generators of an ideal in a power series ring in three variables over a field. As a consequence, we obtain minimal generating sets for the first prime of Moh, proving that the minimal number of generators might decrease depending on the field characteristic. This contradicts an statement of Sally. Finally, we show that these minimal generating sets are standard basis with the negative degree reverse lexicographic order.

1. Introduction

Let k be a field, $n \geq 3$ be an odd integer, m = (n+1)/2 and λ an integer greater than n(n+1)m, with $\gcd(\lambda,m)=1$. In his papers [7, 8], Moh considers the following family of k-algebra morphisms $\rho_n: k[[x,y,z]] \to k[[t]]$, where $\rho_n(x) = t^{nm} + t^{nm+\lambda}$, $\rho_n(y) = t^{(n+1)m}$ and $\rho_n(z) = t^{(n+2)m}$. Let $P_n:=\ker(\rho_n)$ and let $\mu(P_n)$ denote its minimal number of generators. Moh proves that, if k has characteristic zero, then $\mu(P_n) = n+1$. Shortly after, Sally gives in her book [9] an excellent overview of Moh's work. In a Remark in pages 61-62, she writes "Once Moh's primes are constructed in k[[x,y,z]], with k of characteristic 0, then one can show that they also exist in k[[x,y,z]], for k a field of any characteristic." In her discussion, she affirms that, if \overline{P}_n is the corresponding ideal in characteristic p>0, then $\mu(\overline{P}_n) \geq \mu(P_n)$. Much more recently, Mehta, Saha and Sengupta in [6] revisit Moh's primes focusing on given an iterative procedure for finding a set of polynomials that minimally generates P_n . It is worth to mention that in the papers of Moh, it is not mentioned, at least explicity, that the ideals P_n can be generated by polynomials. Mehta, Saha and Sengupta also give a Gröbner basis of the ideal $P_1 \cap k[x,y,z]$, revealing the difficulty of finding a Gröbner basis of $P_n \cap k[x,y,z]$ in general.

In the present paper we first study, what is to us, the key result of Moh in [7]. That result gives a lower bound on the minimal number of generators of an ideal Q of $\mathbb{k}[[x,y,z]]$ in terms of the dimension of some \mathbb{k} -vector spaces V_r associated to an order σ in $\mathbb{k}[[x,y,z]]$ and to the ideal Q (see [7, Theorem 4.3]). It is readily seen that that theorem holds true in any dimension and for any characteristic of \mathbb{k} . Our main contribution, besides giving a more detailed proof, is to show a somewhat more explicit method of finding (and extending to) minimal generating sets of Q, contained in the \mathbb{k} -linear space spanned by a fixed minimal generating set of Q, and satisfying certain conditions linked to the V_r and σ (see Theorem 3.1 for more details).

Secondly, by means of our extension of the result of Moh, we construct sets of polynomials that minimally generate $P:=P_3$, (n=3), so m=2, and for $\lambda=25$, but for any characteristic. We prove that $\mu(P)=4$, not only in characteristic 0, but also in characteristic $p\geq 5$ (see Theorem 4.7). Surprisingly enough, we see that $\mu(P)=2$ in characteristic 2 and $\mu(P)=3$ in characteristic 3, which contradicts the aforementioned inequality $\mu(\overline{P}_n)\geq \mu(P_n)$ in the book of Sally (Theorems 4.10 and 4.13). And what is more, it leaves as an open problem to find a subfamily of the family of primes of Moh, or an alternative family of primes in $\mathbb{k}[[x,y,z]]$, with an unbounded minimal number

Date: August 1, 2024.

²⁰²⁰ Mathematics Subject Classification. 13E05, 13H05, 13J05, 13P10.

Partially supported by Grant PID2019-103849GB-I00 funded by MICIU/AEI/10.13039/501100011033 and AGAUR project 2021 SGR 00603.

of generators, and where k is a field of characteristic other than 0. We hope to come back to this question in a near future.

Finally, we show that the minimal generating sets provided for P are, in fact, standard basis with the negative degree reverse lexicographic order $>_{ds}$ on the set of monomials.

For any unexplained notation we refer to the books [1, 4, 5].

2. Notations and preliminaries

This section is devoted to introduce some terminology and easy preliminary results.

Notation 2.1. We denote the set of non-negative integers by \mathbb{N} . Fix $d \in \mathbb{N}$, $d \geq 1$. Let $n_1, \ldots, n_d \in \mathbb{N}$, $n_i \geq 1$, with $\gcd(n_1, \ldots, n_d) = 1$. Let $\mathcal{S} = \langle n_1, \ldots, n_d \rangle$ be the numerical semigroup generated by n_1, \ldots, n_d . For $r \in \mathbb{N}$, let

$$F(r, S) = \{ \alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}^d \mid \alpha_1 n_1 + \dots + \alpha_d n_d = r \}$$

stand for the set of factorizations of r. Note that if $r \notin \mathcal{S}$, then $F(r,\mathcal{S}) = \emptyset$.

Notation 2.2. Let \mathbbm{k} be a field, $\underline{\mathbf{x}} = x_1, \dots, x_d$ a set of variables over \mathbbm{k} and $\mathrm{Mon}_d = \{\mathbf{x}^\alpha \mid \alpha \in \mathbb{N}^d\}$ the set of monomials $\mathbf{x}^\alpha = x_1^{\alpha_1} \cdots x_d^{\alpha_d}$ with multi-index exponent $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}^d$. The degree of \mathbf{x}^α is defined as $\deg(\mathbf{x}^\alpha) = |\alpha| = \alpha_1 + \dots + \alpha_d$. Let $\mathbbm{k}[\underline{\mathbf{x}}] = \mathbbm{k}[x_1, \dots, x_d]$ denote the power series ring with coefficients in \mathbbm{k} . Any power series can be written as $f = \sum_{\alpha \in \mathbb{N}^d} \lambda_\alpha \mathbf{x}^\alpha$, with $\lambda_\alpha \in \mathbbm{k}$. The support of $f, f \neq 0$, is $\mathrm{Supp}(f) = \{\alpha \in \mathbb{N}^d \mid \lambda_\alpha \neq 0\}$. The order of $f, f \neq 0$, is $\mathrm{ord}(f) = \min\{|\alpha| \in \mathbb{N} \mid \alpha \in \mathrm{Supp}(f)\}$. The leading form of $f, f \neq 0$, is $\mathrm{LF}(f) = \sum_{|\alpha| = \mathrm{ord}(f)} \lambda_\alpha \mathbf{x}^\alpha$; f is said to be homogeneous if $f = \mathrm{LF}(f)$. If f, g are homogeneous, both of order r, then f + g = 0 or f + g is homogeneous of order r. In this paper, whenever we work with power series rings in three variables or in one variable, they will be denoted x, y, z or t, respectively. For instance, if d = 3 and $f = z^3 - 2xy^3 + x^2yz - y^5z^4 \in \mathbb{k}[[x,y,z]]$, then $\mathrm{Supp}(f) = \{(0,0,3), (1,3,0), (2,1,1), (0,5,4)\}$, $\mathrm{ord}(f) = 3$ and $\mathrm{LF}(f) = z^3$.

Let $\sigma: \mathbb{k}[\![\underline{x}]\!] \to \mathbb{k}[\![\underline{x}]\!]$ be the \mathbb{k} -algebra morphism defined by $\sigma(x_i) = x_i^{n_i}$, for $i=1,\ldots,d$. Given $f=\sum_{\alpha\in\mathbb{N}^d}\lambda_\alpha x^\alpha\in\mathbb{k}[\![\underline{x}]\!]$, $f\neq 0$, the σ -order of f is defined as $\sigma\mathrm{ord}(f):=\mathrm{ord}(\sigma(f))$. Note that $\sigma\mathrm{ord}(f)\in\mathcal{S}$. The σ -leading form of f is $f^\sigma:=\sigma^{-1}(\mathrm{LF}(\sigma(f)))=\sum_{\alpha\in\mathrm{F}(r,\mathcal{S})}\lambda_\alpha x^\alpha,$ where $r=\sigma\mathrm{ord}(f)$; f is said to be σ -homogeneous if $\sigma(f)$ is homogeneous, that is, if $f=f^\sigma$. Let $f^\tau:=f-f^\sigma$, which will be called the tail of f. Following with the previous example, if $\mathcal{S}=\langle 3,4,5\rangle,$ and $\sigma(x)=t^3,$ $\sigma(y)=t^4$ and $\sigma(z)=t^5,$ then $\sigma(f)=z^{15}-2x^3y^{12}+x^6y^4z^5-y^{20}z^{20},$ $\sigma\mathrm{ord}(f)=15\in\mathcal{S},$ $\mathrm{LF}(\sigma(f))=z^{15}-2x^3y^{12}+x^6y^4z^5,$ $f^\sigma=z^3-2xy^3+x^2yz$ and $f^\tau=-y^5z^4.$

Given a subset \mathscr{B} of $\mathbb{k}[\![\underline{\mathbf{x}}]\!]$, let $\mathscr{B}^{\sigma} = \{f^{\sigma} \mid f \in \mathscr{B}, f \neq 0\}$ denote the set of its σ -leading forms. We understand that if $\mathscr{B} = \emptyset$ or $\mathscr{B} = \{0\}$, then $\mathscr{B}^{\sigma} = \emptyset$.

Remark 2.3. Let $r \in \mathbb{N}$. Set $W_r = \{g \in \mathbb{k}[\![\underline{\mathbf{x}}]\!] \mid g \neq 0, \ g = g^{\sigma}, \ \sigma \text{ord}(g) = r\} \cup \{0\}$. Then, $g \in W_r$ if and only if $g = \sum_{\alpha \in \mathcal{F}(r,\mathcal{S})} \lambda_{\alpha} \mathbf{x}^{\alpha}$. Since $\mathcal{F}(r,\mathcal{S})$ is finite, $W_r = \langle \mathbf{x}^{\alpha} \mid \alpha \in \mathcal{F}(r,\mathcal{S}) \rangle$ is the \mathbb{k} -vector space spanned by the monomials \mathbf{x}^{α} , with $\alpha \in \mathcal{F}(r,\mathcal{S})$. Thus, the elements of W_r are polynomials. If $r \in \mathcal{S}$, then $W_r \neq 0$ and dim $W_r \geq 1$. If r = 0, then $W_0 = \langle 1 \rangle$. If $r \notin \mathcal{S}$, we just define $W_r = \{0\}$. Observe that the definition of W_r depends on the chosen σ .

Let $h = \sum_{\alpha \in \mathbb{N}^d} \lambda_\alpha \mathbf{x}^\alpha \in \mathbb{k}[\underline{\mathbf{x}}]$. Given $r \in \mathcal{S}$, let $h_r = \sum_{\alpha \in \mathcal{F}(r,\mathcal{S})} \lambda_\alpha \mathbf{x}^\alpha \in W_r$ be the polynomial obtained as the summation of all the monomial terms of h of σ -order r. In particular, the summation $\sum_{r \in \mathcal{S}} h_r$ is well-defined and h can be written as $h = \sum_{r \in \mathcal{S}} h_r$.

Lemma 2.4. Let Q be a proper ideal of $\mathbb{k}[\underline{x}]$. Let $r \in \mathcal{S}$. Set

$$V_r = \{g \in W_r \mid g \neq 0, \text{ there exists } f \in Q \text{ such that } g = f^{\sigma}\} \cup \{0\}.$$

Then V_r is a k-vector subspace of W_r . Moreover, if $g \in W_r$, $g \neq 0$, then the following are equivalent:

- (a) $g \in V_r$;
- (b) $g \in Q$ or there exists $h \in \mathbb{K}[\underline{x}]$, $h \neq 0$, such that $\sigma \operatorname{ord}(h) > r$ and such that $f := g + h \in Q$.

In such a case, $q = f^{\sigma}$ and $h = f^{\tau}$.

Proof. If $g_1, g_2 \in V_r$, with $g_1 = -g_2$, then $g_1 + g_2 = 0$, and $g_1 + g_2 \in V_r$. If $g_1, g_2 \in V_r$, with $g_1, g_2 \neq 0$ and $g_1 \neq -g_2$, then $g_1, g_2 \in W_r$ and $g_1 + g_2 \in W_r$, because W_r is a vector space. Moreover, there are $f_i = \sum_{\alpha \in \mathbb{N}^d} \lambda_{i,\alpha} \mathbf{x}^\alpha \in Q$, $f_i \neq 0$, such that $g_i = f_i^\sigma = \sum_{\alpha \in \mathbf{F}(r,\mathcal{S})} \lambda_{i,\alpha} \mathbf{x}^\alpha$. Then, $f_1 + f_2 \in Q$, where $f_1 + f_2 = \sum_{\alpha \in \mathbb{N}^d} (\lambda_{1,\alpha} + \lambda_{2,\alpha}) \mathbf{x}^\alpha$, and $f_1 + f_2 \neq 0$, otherwise $g_1 = f_1^\sigma = -f_2^\sigma = -g_2$. Moreover, $g_1 + g_2 = f_1^\sigma + f_2^\sigma = \sum_{\alpha \in \mathbf{F}(r,\mathcal{S})} \lambda_{1,\alpha} \mathbf{x}^\alpha + \sum_{\alpha \in \mathbf{F}(r,\mathcal{S})} \lambda_{2,\alpha} \mathbf{x}^\alpha = \sum_{\alpha \in \mathbf{F}(r,\mathcal{S})} (\lambda_{1,\alpha} + \lambda_{2,\alpha}) \mathbf{x}^\alpha = (f_1 + f_2)^\sigma$. Therefore, $g_1 + g_2 \in V_r$. Similarly, if $g \in V_r$, $g \neq 0$, and $\nu \in \mathbb{K}$, $\nu \neq 0$, then $\nu g \in W_r$, because W_r is a vector space. Furthermore, there exists $f = \sum_{\alpha \in \mathbb{N}^d} \lambda_\alpha \mathbf{x}^\alpha \in Q$, $f \neq 0$, such that $g = f^\sigma = \sum_{\alpha \in \mathbf{F}(r,\mathcal{S})} \lambda_{\alpha} \mathbf{x}^\alpha$. Then, $\nu f \in Q$, where $\nu f = \sum_{\alpha \in \mathbb{N}^d} (\nu \lambda_\alpha) \mathbf{x}^\alpha$, and $\nu f \neq 0$. Moreover, $\nu g = \nu f^\sigma = \sum_{\alpha \in \mathbf{F}(r,\mathcal{S})} (\nu \lambda_\alpha) \mathbf{x}^\alpha = (\nu f)^\sigma$. Therefore, $\nu g \in V_r$. Thus, V_r is a \mathbb{k} -vector space.

Suppose that $g \in W_r$, $g \neq 0$, $g \in V_r$. By definition of V_r , there exists $f = \sum_{s \geq r} f_s$, where $f_s = \sum_{\alpha \in \mathcal{F}(s,S)} \lambda_{\alpha} \mathbf{x}^{\alpha}$, $f \in Q$, and such that $g = f^{\sigma} = f_r$. If $f_s = 0$, for all s > r, then $g = f_r = f$, so $g \in Q$. If $f_s \neq 0$, for some s > r, take $h := \sum_{s > r} f_s$. Then $h \neq 0$, $\operatorname{\sigmaord}(h) > r$, and $f = f_r + h = g + h$. Conversely, suppose that $g \in W_r$, $g \neq 0$, in particular, $g = g^{\sigma}$ and $\operatorname{\sigmaord}(g) = r$. If $g \in Q$, take f := g and we are done. On the other hand, if there exists $h \in \mathbb{k}[\![\mathbf{x}]\!]$, $h \neq 0$, such that $\operatorname{\sigmaord}(h) > r$ and such that $f := g + h \in Q$, then, clearly, $f^{\sigma} = g$ and $g \in V_r$.

Note that, since Q is proper, if r = 0, it follows that $V_0 = 0$. If $r \notin \mathcal{S}$, we just define $V_r = \{0\}$. Observe that V_r depends on σ and Q.

3. Revisiting a theorem of Moh

A careful reading of the proof of Theorem 4.3 of Moh in [7] reveals a very deep and rich result (see also the proof of [8, Theorem]). The purpose of this section is to bring to the surface the somehow hidden ideas in the proof while, at the same time, we reprove it more detailedly and generalize it.

To simplify the statement, we will understand that a null vector space has a unique basis, or a unique linearly independent subset, which is the empty set.

Theorem 3.1. Let Q be a proper ideal of $\mathbb{k}[\underline{x}]$ and let \mathscr{C} be a finite system of generators of Q. Let $\sigma : \mathbb{k}[\underline{x}] \to \mathbb{k}[\underline{x}]$ be the \mathbb{k} -algebra morphism defined by $\sigma(x_i) = x_i^{n_i}$, for $i = 1, \ldots, d$. Let V_r be associated to σ and Q. Let $\xi = \min(n_1, \ldots, n_d)$, $s = \min\{\sigma \operatorname{ord}(f) \mid f \in Q, f \neq 0\}$, with $\xi, s \geq 1$.

- (1) Then there exist $\mathscr{B}_0, \ldots, \mathscr{B}_{\xi-1}, \mathscr{C}_{\xi-1}$ included in $\langle \mathscr{C} \rangle_{\mathbb{k}}$, the \mathbb{k} -linear space spanned by \mathscr{C} , such that for each $i = 0, \ldots, \xi 1$,
 - \mathscr{B}_{i}^{σ} is a (possibly empty) basis of V_{s+i} .
 - $\mathscr{B}_0, \ldots, \mathscr{B}_{\xi-1}, \mathscr{C}_{\xi-1}$ are pairwise disjoint.
 - $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1}$ is a minimal generating set of Q.

In particular,

$$\mu(Q) \ge \sum_{i=0}^{\xi-1} \dim V_{s+i}$$
.

(2) Let $\mathscr{D}_0, \ldots, \mathscr{D}_{\xi-1} \subset Q$. Suppose that, for each $i = 0, \ldots, \xi - 1$, \mathscr{D}_i^{σ} is a (possibly empty) linearly independent subset of V_{s+i} . Then $\mathscr{D}_0 \cup \ldots \cup \mathscr{D}_{\xi-1}$ can be extended to a minimal generating set of Q with elements of $\langle \mathscr{C} \rangle_{\Bbbk}$.

Moreover, if the elements of \mathscr{C} are polynomials, then the elements of $\mathscr{B}_0, \ldots, \mathscr{B}_{\xi-1}, \mathscr{C}_{\xi-1}$ can be chosen to be polynomials and $\mathscr{D}_0 \cup \ldots \cup \mathscr{D}_{\xi-1}$ can be extended to a minimal generating set of Q with polynomials.

Before the proof, we want to emphasize some aspects of this result, by comparison to the Moh's original theorem. First observe that it holds in any dimension and in any characteristic of the field k. Moh focused on the lower bound of $\mu(Q)$. Instead, we focus on detailing the construction of

the minimal generating set of Q from where one can deduce that lower bound. Concretely: fixed a (minimal) system of generators \mathscr{C} of Q, we construct a (minimal) generating set $\mathscr{B}_0 \cup \ldots \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1}$ of Q, included in the \mathbb{k} -linear space spanned by \mathscr{C} , and such that each \mathscr{B}_i^{σ} is a basis of V_{s+i} . In particular, $\mu(Q) \geq \sum_{i=0}^{\xi-1} \dim V_{s+i}$ and, if Q is generated by polynomials, then the elements of \mathscr{B}_i and $\mathscr{C}_{\xi-1}$ can be taken polynomials. We have not been able to see in the papers of Moh a proof that the ideals P_n are generated by polinomials. This is not the case in the paper of Mehta, Saha and Sengupta, in characteristic zero, where they give an iterative method to construct minimal generating sets of P_n given by polynomials (see [6, Section 4]).

The second part of our theorem asserts that it is possible to extend $\mathscr{D}_0 \cup \ldots \cup \mathscr{D}_{\xi-1}$, where \mathscr{D}_i^{σ} is a linearly independent subset of V_{s+i} , to a minimal generating set of Q. We will use this second part to find the minimal generating sets of P_3 in any characteristic.

In the end, two are the main ingredients of the proof: the equality $\sigma \operatorname{ord}(fg) = \sigma \operatorname{ord}(f) + \sigma \operatorname{ord}(g)$, where $f, g \in \mathbb{k}[\![\underline{\mathbf{x}}]\!]$, $f, g \neq 0$, in the first part and the Steinitz exchange lemma, a ubiquitous procedure in linear algebra, in the second part.

Proof of the Theorem 3.1. For $r \geq 1$, set $d_r = \dim V_r$ and, for $i = 0, \ldots, \xi - 1$, set $e_i = d_s + \cdots + d_{s+i}$. Take $\mathscr{C} = \{f_1, \ldots, f_{\eta}\}$, a finite set of generators of Q of cardinality $\eta \geq 1$. Let us iteratively construct, for each $i = 0, \ldots, \xi - 1$, \mathscr{B}_i and \mathscr{C}_i satisfying the following list of conditions \mathbb{H}_i .

- \mathscr{B}_i and \mathscr{C}_i are subsets of $\langle \mathscr{C} \rangle_{\mathbb{k}}$.
 - If $V_{s+i} = 0$, then $\mathscr{B}_i = \emptyset$.
 - If $V_{s+i} \neq 0$, then $\mathscr{B}_i = \{f_{i,1}, \dots, f_{i,d_{s+i}}\}$, $\operatorname{card}(\mathscr{B}_i) = d_{s+i}$, and \mathscr{B}_i^{σ} is a basis of V_{s+i} .
- $\bullet \ \eta \ge e_i = d_s + \dots + d_{s+i}.$
 - If $\eta = e_i$, then $V_r = 0$, for all $s + i < r < s + \xi$ and $\mathscr{C}_i = \emptyset$.
 - If $\eta > e_i$, then $\mathscr{C}_i = \{g_{i,1}, \dots, g_{i,\eta-e_i}\} \subset \langle \mathscr{C} \rangle_{\mathbb{k}}$, $\operatorname{card}(\mathscr{C}_i) = \eta e_i$ and $\operatorname{\sigmaord}(g_{i,j}) > s + i$.
- In any case, $\mathscr{B}_0, \ldots, \mathscr{B}_i, \mathscr{C}_i$ are pairwise disjoint and $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_i \cup \mathscr{C}_i$ is a set of generators of Q of cardinality η .

In the end, once all the \mathscr{B}_i and \mathscr{C}_i are constructed satisfying the conditions \mathbb{H}_i , we deduce that $\eta \geq e_{\xi-1} = d_s + \cdots + d_{s+\xi-1} = \sum_{i=0}^{\xi-1} \dim V_{s+i}$, and that there exist $\mathscr{B}_0, \ldots, \mathscr{B}_{\xi-1}, \mathscr{C}_{\xi-1}$, a family of pairwise disjoint subsets of $\langle \mathscr{C} \rangle_{\mathbb{K}}$, $\mathscr{B}_i = \emptyset$, if $V_{s+i} = 0$, \mathscr{B}_i^{σ} a basis of V_{s+i} , if $V_{s+i} \neq 0$, and such that $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1}$ is a set of generators of Q of cardinality η . In particular, if \mathscr{C} is a minimal generating set of Q, since $\mathbb{K}[x_1, \ldots, x_d]$ is Noetherian local, one deduces that $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1}$ is a minimal generating set for Q too and so $\mu(Q) = \eta \geq \sum_{i=0}^{\xi-1} \dim V_{s+i}$. This will prove item (1).

Construction of \mathscr{B}_0 and \mathscr{C}_0 . By definition of $s, V_s \neq 0$. Take $\{v_{0,1}, \ldots, v_{0,d_s}\} \subset Q$ such that $\{v_{0,1}^{\sigma}, \ldots, v_{0,d_s}^{\sigma}\}$ is a basis of V_s . For each $1 \leq j \leq d_s$, since $v_{0,j} \in Q$ and $Q = (f_1, \ldots, f_\eta)$, then

$$v_{0,j} = \sum_{k=1}^{\eta} a_{0,j}^k f_k,$$

where $a_{0,j}^k \in \mathbb{k}[\underline{x}]$. Note that, if $a_{0,j}^k$ is not a unit and $a_{0,j}^k \neq 0$, then $\operatorname{\sigmaord}(a_{0,j}^k f_k) \geq \xi + s > s = \operatorname{\sigmaord}(v_{0,j})$. In particular, the subset $\Lambda_{0,j}$ of indices l of $\{1,\ldots,\eta\}$ such that $a_{0,j}^l$ is invertible, with term of order zero $\lambda_{0,j}^l \in \mathbb{k}$, $\lambda_{0,j}^l \neq 0$, and such that $\operatorname{\sigmaord}(f_l) = s$, is not empty. Taking σ -leading forms in both parts of the equality, we obtain

$$v_{0,j}^{\sigma} = \sum_{l \in \Lambda_{0,j}} \lambda_{0,j}^l f_l^{\sigma}.$$

Let $\Lambda_0 = \Lambda_{0,1} \cup \ldots \cup \Lambda_{0,d_s}$. Then

$$V_s = \langle v_{0,1}^{\sigma}, \dots, v_{0,d_s}^{\sigma} \rangle \subseteq \langle f_l^{\sigma} \mid l \in \Lambda_0 \rangle \subseteq V_s.$$

Thus $V_s = \langle f_l^{\sigma} \mid l \in \Lambda_0 \rangle$. Since V_s has dimension d_s and $\{f_l^{\sigma} \mid l \in \Lambda_0\}$ spans V_s , then $\operatorname{card}(\Lambda_0) \geq d_s$, and there exists a subset $\Gamma \subseteq \Lambda_0 \subseteq \{1, \ldots, \eta\}$, Γ of cardinality d_s , such that $\{f_l^{\sigma} \mid l \in \Gamma\}$ is a basis

of V_s . In particular $\eta \geq d_s$. Renaming the elements f_1, \ldots, f_{η} , we can suppose that $\{f_1^{\sigma}, \ldots, f_{d_s}^{\sigma}\}$ is a basis of V_s . For $l = 1, \ldots, d_s$, set $f_{0,l} := f_l$ and let $\mathscr{B}_0 = \{f_{0,1}, \ldots, f_{0,d_s}\}$. Note that $\mathscr{B}_0 \subseteq \mathscr{C}$, $\operatorname{card}(\mathscr{B}_0) = d_s$, \mathscr{B}_0^{σ} is a basis of V_s , and $\operatorname{card}(\mathscr{C} \setminus \mathscr{B}_0) = \eta - d_s = \eta - e_0 \geq 0$.

Suppose that $\eta = e_0 = d_s$. In particular, $\mathscr{B}_0 = \mathscr{C}$, which is a set of generators of Q of cardinality η . Let us prove that $V_r = 0$, for all $s < r < s + \xi$. Since $\eta = d_s$, f_1, \ldots, f_η have all σ -order equal to s and $\{f_1^{\sigma}, \ldots, f_{\eta}^{\sigma}\}$ is a basis of V_s . If $\xi = 1$, there is nothing to prove. Suppose that $\xi \geq 2$, and that there exists $g \in V_r$, $g \neq 0$, for some $s < r < s + \xi$. Then $g = f^{\sigma}$, with $f \in Q$, $\sigma \operatorname{ord}(f) = r$. Since $f \in Q = (f_1, \ldots, f_{\eta})$, then $f = \sum_{j=1}^{\eta} b_j f_j$, where $b_j \in \mathbb{k}[\![\underline{x}]\!]$. Let $\Lambda \subseteq \{1, \ldots, \eta\}$ denote the subset of indices l such that b_l is invertible, with term of order zero $\mu_l \in \mathbb{k}$, $\mu_l \neq 0$. Suppose that $\Lambda \neq \emptyset$. Since $\sigma \operatorname{ord}(f) = r > s$, then, necessarily, $\sum_{l \in \Lambda} \mu_l f_l^{\sigma} = 0$. Since $\{f_1^{\sigma}, \ldots, f_{\eta}^{\sigma}\}$ is a basis of V_s , this implies $\mu_l = 0$, a contradiction. Therefore $\Lambda = \emptyset$ and all the b_l are not units. But then $\sigma \operatorname{ord}(b_j f_j) \geq \xi + s > r$, whereas $\sigma \operatorname{ord}(f) = r$, a contradiction again. Therefore, f must be zero and $V_r = 0$, for all $s < r < s + \xi$.

Thus, when $\eta = e_0 = d_s$, let $\mathscr{C}_0 = \emptyset$, so that $\mathscr{B}_0 \cup \mathscr{C}_0$ is a set of generators of Q of cardinality η .

Suppose that $\eta > e_0 = d_s$. For any $g \in \mathcal{C} \setminus \mathcal{B}_0$, with $\sigma \operatorname{ord}(g) = s$, if any, then $g^{\sigma} = \sum_{l=1}^{d_s} \lambda_l f_{0,l}^{\sigma}$, for some $\lambda_l \in \mathbb{k}$. Moreover $g - \sum_{l=1}^{d_s} \lambda_l f_{0,l} \in Q$ and $\sigma \operatorname{ord}(g - \sum_{l=1}^{d_s} \lambda_l f_{0,l}) > s$. Take

$$\mathscr{C}_0 := \{g \in \mathscr{C} \setminus \mathscr{B}_0 \mid \operatorname{\sigmaord}(g) > s\} \bigcup \left\{g - \sum_{l=1}^{d_s} \lambda_l f_{0,l} \middle| g \in \mathscr{C} \setminus \mathscr{B}_0, \operatorname{\sigmaord}(g) = s\right\} \subset \langle \mathscr{C} \rangle_{\Bbbk},$$

where $\langle \mathcal{C} \rangle_{\mathbb{k}}$ stands for the \mathbb{k} -linear space spanned by \mathcal{C} . Since all the elements in \mathcal{C}_0 have σ -order bigger than s, it follows that \mathcal{B}_0 and \mathcal{C}_0 are disjoint. Since \mathcal{C} is a set of generators of Q and $\mathcal{B}_0 \subset \mathcal{C}$, one deduces that $\mathcal{B}_0 \cup \mathcal{C}_0$ is a set of generators of Q of cardinality η .

This proves that \mathcal{B}_0 and \mathcal{C}_0 fulfill the conditions \mathbb{H}_0 .

Construction of \mathcal{B}_i and \mathcal{C}_i , $1 \leq i \leq \xi - 1$. Note that, in particular, $\xi \geq i + 1$. By the step i-1, there exist a family of pairwise disjoint subsets $\mathcal{B}_0, \ldots, \mathcal{B}_{i-1}, \mathcal{C}_{i-1}$ of $\langle \mathcal{C} \rangle_{\mathbb{k}}$ satisfying the conditions \mathbb{H}_{i-1} . Therefore, $\mathcal{B}_k = \emptyset$, if $V_{s+k} = 0$, and $\mathcal{B}_k = \{f_{k,1}, \ldots, f_{k,d_{s+k}}\}$, with \mathcal{B}_k^{σ} a basis of V_{s+k} , if $V_{s+k} \neq 0$. Moreover, $\mathcal{B}_0 \cup \ldots \cup \mathcal{B}_{i-1} \cup \mathcal{C}_{i-1}$ is a set of generators of Q of cardinality η . Furthermore, $\eta \geq e_{i-1} = d_s + \cdots + d_{s+i-1}$ and, if $\eta = e_{i-1}$, then $V_r = 0$, for all $s+i-1 < r < s+\xi$ and $\mathcal{C}_{i-1} = \emptyset$. If $\eta > e_{i-1}$, then $\mathcal{C}_{i-1} = \{g_{i-1,1}, \ldots, g_{i-1,\eta-e_{i-1}}\} \subset \langle \mathcal{C} \rangle_{\mathbb{k}}$, $\operatorname{card}(\mathcal{C}_{i-1}) = \eta - e_{i-1}$ and $\operatorname{\sigmaord}(g_{i-1,j}) > s+i-1$.

Suppose that $\eta = e_{i-1}$. In this case, by the step i-1, $V_{s+i} = 0$ and $\mathscr{C}_{i-1} = \emptyset$. Take $\mathscr{B}_i = \emptyset$ and $\mathscr{C}_i = \emptyset$. Since $\mathscr{C}_{i-1} = \emptyset$, it follows that $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_i \cup \mathscr{C}_i = \mathscr{B}_0 \cup \ldots \cup \mathscr{B}_{i-1} = \mathscr{B}_0 \cup \ldots \cup \mathscr{B}_{i-1} \cup \mathscr{C}_{i-1}$, which is a set of generators of Q of cardinality η , by the step i-1.

Suppose that $\eta > e_{i-1}$ and $V_{s+i} = 0$. Then take $\mathscr{B}_i = \emptyset$ and $\mathscr{C}_i = \mathscr{C}_{i-1}$, so $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_i \cup \mathscr{C}_i = \mathscr{B}_0 \cup \ldots \cup \mathscr{B}_{i-1} \cup \mathscr{C}_{i-1}$, which, by the step i-1, is a set of generators of Q of cardinality η .

Suppose that $\eta > e_{i-1}$ and $V_{s+i} \neq 0$. Take $\{v_{i,1}, \ldots, v_{i,d_{s+i}}\} \subset Q$ such that $\{v_{i,1}^{\sigma}, \ldots, v_{i,d_{s+i}}^{\sigma}\}$ is a basis of V_{s+i} . Since $v_{i,j} \in Q$ and Q is generated by the elements of $\mathcal{B}_0 \cup \ldots \mathcal{B}_{i-1} \cup \mathcal{C}_{i-1}$, then, for each $1 \leq j \leq d_{s+i}$,

$$v_{i,j} = \sum_{k=0}^{i-1} \sum_{l=1}^{d_{s+k}} a_{i,j}^{k,l} f_{k,l} + \sum_{l=1}^{\eta - e_{i-1}} b_{i,j}^l g_{i-1,l}, \tag{1}$$

where $a_{i,j}^{k,l}, b_{i,j}^l \in \mathbb{k}[\underline{x}]$, $f_{k,l} \in \mathcal{B}_k$ and $g_{i-1,l} \in \mathcal{C}_{i-1}$. Note that $\operatorname{\sigmaord}(v_{i,j}) = s + i$ and $\operatorname{\sigmaord}(f_{k,l}) = s + k$, where $0 \le k \le i-1$ and $1 \le l \le d_{s+k}$. In addition, $\operatorname{\sigmaord}(g_{i-1,l}) \ge s + i$, for $1 \le l \le \eta - e_{i-1}$. If $a_{i,j}^{k,l}$ is not a unit and $a_{i,j}^{k,l} \ne 0$, then $\operatorname{\sigmaord}(a_{i,j}^{k,l}f_{k,l}) \ge \xi + s + k > s + i = \operatorname{\sigmaord}(v_{i,j})$, because $\xi \ge i+1$.

Let $\Lambda_{i,j}^0 \subseteq \{1,\ldots,d_s\}$ denote the subset of indices l such that $a_{i,j}^{0,l}$ is invertible, with term of order zero $\lambda_{i,j}^{0,l} \in \mathbb{R}$, $\lambda_{i,j}^{0,l} \neq 0$, and suppose that $\Lambda_{i,j}^0 \neq \emptyset$. Taking σ -leading forms in Equality (1), we deduce that $\sum_{l \in \Lambda_{i,j}^0} \lambda_{i,j}^{0,l} f_{0,l}^{\sigma} = 0$. Since $\{f_{0,1}^{\sigma},\ldots,f_{0,d_s}^{\sigma}\}$ is a basis of V_s , then $\lambda_{i,j}^{0,l} = 0$, a contradiction. Therefore, all the $a_{i,j}^{0,l}$ are not units. For all $k = 1,\ldots,i-1$, $\sigma \operatorname{ord}(f_{k,l}) = s + k < s + i = \sigma \operatorname{ord}(v_{i,j})$. Thus, similarly, and recursively on $k = 1,\ldots,i-1$, one can prove that $a_{i,j}^{k,l}$ are not invertible. Hence, $\sigma \operatorname{ord}(a_{i,j}^{k,l}f_{k,l}) > s + i = \sigma \operatorname{ord}(v_{i,j})$, for all $k = 0,\ldots,i-1$ and for all $l = 1,\ldots,d_{s+k}$. Let $\Lambda_{i,j} \subseteq \{1,\ldots,\eta-e_{i-1}\}$ be the set of indices l such that $b_{i,j}^l$ is invertible, with term of order zero $\mu_{i,j}^l \in \mathbb{R}$, $\mu_{i,j}^l \neq 0$, and such that $\sigma \operatorname{ord}(g_{i-1,l}) = s + i$. Note that $\Lambda_{i,j} \neq \emptyset$ for, if $b_{i,j}^l$ is not a unit, then $\sigma \operatorname{ord}(b_{i,j}^l g_{i-1,l}) \geq \xi + s + i > \sigma \operatorname{ord}(v_{i,j})$. Taking σ -leading forms in Equality (1), we deduce that

$$v_{i,j}^{\sigma} = \sum_{l \in \Lambda_{i,j}} \mu_{i,j}^l g_{i-1,l}^{\sigma}.$$

Let $\Lambda_i = \Lambda_{i,1} \cup \ldots \cup \Lambda_{i,d_{s+i}}$. Then

$$V_{s+i} = \langle v_{i,1}^{\sigma}, \dots, v_{i,d_{s+i}}^{\sigma} \rangle \subseteq \langle g_{i-1,l}^{\sigma} \mid l \in \Lambda_i \rangle \subseteq V_{s+i}.$$

So $V_{s+i} = \langle g_{i-1,l}^{\sigma} \mid l \in \Lambda_i \rangle$. Since dim $V_{s+i} = d_{s+i}$ and $\{g_{i-1,l} \mid l \in \Lambda_i\}$ spans V_{s+i} , then $\operatorname{card}(\Lambda_i) \geq d_{s+i}$, and there exists a subset $\Gamma \subseteq \Lambda_i \subseteq \{1, \ldots, \eta - e_{i-1}\}$, $\operatorname{card}(\Gamma) = d_{s+i}$, such that $\{g_{i-1,l}^{\sigma} \mid l \in \Gamma\}$ is a basis of V_{s+i} . In particular,

$$\eta - e_{i-1} \ge \operatorname{card}(\Lambda_i) \ge \operatorname{card}(\Gamma) = d_{s+i},$$

and $\eta \geq e_{i-1} + d_{s+i} = e_i$. Renaming the elements of $\mathscr{C}_{i-1} = \{g_{i-1,1}, \dots, g_{i-1,\eta-e_{i-1}}\}$, we can suppose that $\{g_{i-1,1}^{\sigma}, \dots, g_{i-1,d_{s+i}}^{\sigma}\}$ is a basis of V_{s+i} . For $l = 1, \dots, d_{s+i}$, set $f_{i,l} := g_{i-1,l}$ and let $\mathscr{B}_i = \{f_{i,1}, \dots, f_{i,d_{s+i}}\}$. Note that $\mathscr{B}_i \subseteq \mathscr{C}_{i-1} \subset \langle \mathscr{C} \rangle_{\mathbb{k}}$, $\operatorname{card}(\mathscr{B}_i) = d_{s+i}$ and $\operatorname{card}(\mathscr{C}_{i-1}) = \eta - e_{i-1}$.

Suppose that $\eta = e_i$. In particular, $\operatorname{card}(\mathcal{B}_i) = d_{s+i} = e_i - e_{i-1} = \eta - e_{i-1} = \operatorname{card}(\mathcal{C}_{i-1})$. Thus $\mathcal{B}_i = \mathcal{C}_{i-1}$ and $\mathcal{B}_0 \cup \ldots \cup \mathcal{B}_i = \mathcal{B}_0 \cup \ldots \mathcal{B}_{i-1} \cup \mathcal{C}_{i-1}$, which is a system of generators of Q of cardinality η , by the conditions \mathbb{H}_{i-1} of the former step. Let us show that $V_r = 0$, for all $s+i < r < s+\xi$. If $\xi = i+1$, there is nothing to prove. Suppose that $\xi \geq i+2$, and that there exists $g \in V_r$, $g \neq 0$, for some $s+i < r < s+\xi$. Then $g = f^{\sigma}$, with $f \in Q$, $\sigma \operatorname{ord}(f) = r$. Since $f \in Q = (\mathcal{B}_0 \cup \ldots \cup \mathcal{B}_i)$, then

$$f = \sum_{k=0}^{i} \sum_{l=1}^{d_{s+k}} a_{k,l} f_{k,l},$$

where $a_{k,l} \in \mathbb{k}[\![\underline{\mathbf{x}}]\!]$. Let $\Lambda_0 \subseteq \{1,\ldots,d_s\}$ denote the subset of indices l such that $a_{0,l}$ is invertible, with term of order zero $\lambda_{0,l} \in \mathbb{k}$, $\lambda_{0,l} \neq 0$. Suppose that $\Lambda_0 \neq \emptyset$. Since $\sigma \operatorname{ord}(f_{0,l}) = s < s + i < r = \sigma \operatorname{ord}(f)$, then, necessarily, $\sum_{l \in \Lambda_0} \lambda_{0,l} f_{0,l}^{\sigma} = 0$. Since $\{f_{0,1}^{\sigma},\ldots,f_{0,d_s}^{\sigma}\}$ is a basis of V_s , this implies $\lambda_{0,l} = 0$, a contradiction. Therefore $\Lambda_0 = \emptyset$ and all the $a_{0,l}$ are not units. For all $k = 1,\ldots,i$, $\sigma \operatorname{ord}(f_{k,l}) = s + k \leq s + i < r = \sigma \operatorname{ord}(f)$. So, in a similar vein, and recursively on k, one can prove that $a_{k,l}$ are not invertible for all $k = 1,\ldots,i$. But then, $\sigma \operatorname{ord}(a_{k,l}f_{k,l}) \geq \xi + s > r$, whereas $\sigma \operatorname{ord}(f) = r$, a contradiction again. Therefore, f must be zero and $V_r = 0$, for all $s + i < r < s + \xi$. Thus, when $n = e_i$, let $\mathscr{C}_i = \emptyset$. Since $\mathscr{D}_i = \mathscr{C}_i$, $1 \in \mathscr{C}(\mathscr{C})_k$, and by the former step, it follows

Thus, when $\eta = e_i$, let $\mathscr{C}_i = \emptyset$. Since $\mathscr{B}_i = \mathscr{C}_{i-1} \subset \langle \mathscr{C} \rangle_{\mathbb{k}}$, and by the former step, it follows that $\mathscr{B}_0, \ldots, \mathscr{B}_i, \mathscr{C}_i$ is a family of pairwise disjoint subsets of $\langle \mathscr{C} \rangle_{\mathbb{k}}$, such that $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_i \cup \mathscr{C}_i = \mathscr{B}_0 \cup \ldots \mathscr{B}_{i-1} \cup \mathscr{C}_{i-1}$ is a system of generators of Q of cardinality η .

Suppose that $\eta > e_i$. Then $\mathscr{B}_i \subsetneq \mathscr{C}_{i-1}$. Let $g \in \mathscr{C}_{i-1} \setminus \mathscr{B}_i$, in particular, $\sigma \operatorname{ord}(g) \geq s+i$. If $\sigma \operatorname{ord}(g) = s+i$, then $g^{\sigma} = \sum_{l=1}^{d_{s+i}} \lambda_l f_{i,l}^{\sigma}$, for some $\lambda_l \in \mathbb{k}$. Therefore, $g - \sum_{l=1}^{d_{s+i}} \lambda_l f_{i,l} \in Q$ and

 σ ord $(g - \sum_{l=1}^{d_{s+i}} \lambda_l f_{i,l}) > s+i$. Take

$$\mathscr{C}_i := \left\{ g \in \mathscr{C}_{i-1} \setminus \mathscr{B}_i \middle| \operatorname{\sigmaord}(g) > s+i \right\} \bigcup \left\{ g - \sum_{l=1}^{d_{s+i}} \lambda_l f_{i,l} \middle| g \in \mathscr{C}_{i-1} \setminus \mathscr{B}_i, \operatorname{\sigmaord}(g) = s+i \right\}.$$

Observe that $\mathscr{C}_i \subset \langle \mathscr{C} \rangle_{\mathbb{k}}$. Since all the elements in \mathscr{B}_i have σ -order equal to s+i and all the elements of \mathscr{C}_i have σ -order bigger than s+i, it follows that $\mathscr{B}_0, \ldots, \mathscr{B}_i, \mathscr{C}_i$ are pairwise disjoint. Since $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_{i-1} \cup \mathscr{C}_{i-1}$ is a set of generators of Q and $\mathscr{B}_i \subset \mathscr{C}_{i-1}$, one deduces that $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_i \cup \mathscr{C}_i$ is a set of generators of Q of cardinality η . This proves that $\mathscr{B}_i, \mathscr{C}_i$ fulfill the conditions \mathbb{H}_i and finishes the proof of (1).

PROOF OF (2). Let $\mathscr{D}_0, \ldots, \mathscr{D}_{\xi-1} \subset Q$ be as in the hypothesis (2), so that $\mathscr{D}_i = \emptyset$ or \mathscr{D}_i^{σ} is a linearly independent subset of V_{s+i} . In this last case, set $\mathscr{D}_i = \{u_{i,1}, \ldots, u_{i,\delta_i}\}$, where $\operatorname{card}(\mathscr{D}_i) = \delta_i$. Let $\mathscr{B}_0, \ldots, \mathscr{B}_{\xi-1}, \mathscr{C}_{\xi-1}$ be defined as in item (1). In particular, when they are not empty, set $\mathscr{B}_i = \{f_{i,1}, \ldots, f_{i,d_{s+i}}\}$ and $\mathscr{C}_{\xi-1} = \{g_{\xi-1,1}, \ldots, g_{\xi-1,\mu-e_{\xi-1}}\}$, where \mathscr{B}_i^{σ} is a basis of V_{s+i} , $\sigma \operatorname{ord}(g_{\xi-1,l}) > s + \xi - 1$, and $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1}$ is a minimal generating set for Q of cardinality μ . Let us iteratively construct, for each $i = 0, \ldots, \xi - 1$, \mathscr{B}_i' satisfying the following list of conditions \mathbb{L}_i .

- \mathscr{B}'_i is a subset of \mathscr{B}_i .
- $\mathscr{D}_{i}^{\sigma} \cup (\mathscr{B}_{i}')^{\sigma}$ is a basis of V_{s+i} , understanding that $\mathscr{D}_{i}^{\sigma} = \emptyset$ if $\mathscr{D}_{i} = \emptyset$, or $(\mathscr{B}_{i}')^{\sigma} = \emptyset$ if $\mathscr{B}_{i}' = \emptyset$.
- $\mathcal{D}_0, \mathcal{B}'_0, \dots, \mathcal{D}_i, \mathcal{B}'_i, \mathcal{B}_{i+1}, \dots, \mathcal{B}_{\xi-1}, \mathcal{C}_{\xi-1}$ are pairwise disjoint subsets of Q.
- $(\mathscr{D}_0 \cup \mathscr{B}'_0) \cup \ldots \cup (\mathscr{D}_i \cup \mathscr{B}'_i) \cup \mathscr{B}_{i+1} \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1}$ is a minimal generating set of Q.

In the end, once all the \mathscr{B}'_i are constructed satisfying the conditions \mathbb{L}_i , we deduce that

$$(\mathscr{D}_0 \cup \mathscr{B}'_0) \cup \ldots \cup (\mathscr{D}_{\xi-1} \cup \mathscr{B}'_{\xi-1}) \cup \mathscr{C}_{\xi-1}$$

is a minimal generating set of Q. In particular, $\mathscr{D}_0 \cup \ldots \cup \mathscr{D}_{\xi-1}$ can be extended to a minimal generating set of Q. The proof basically consists on applying repeatedly the Steinitz exchange lemma in linear algebra.

CONSTRUCTION OF \mathscr{B}'_0 . If $\mathscr{D}_0 = \emptyset$, let $\mathscr{B}'_0 = \mathscr{B}_0$, which clearly satisfies the conditions \mathbb{L}_0 . Suppose that $\mathscr{D}_0 \neq \emptyset$. Take $u_{0,1} \in \mathscr{D}_0 \subset Q$. Since $\mathscr{B}_0 \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1}$ is a generating set of the ideal Q, it follows that

$$u_{0,1} = \sum_{k=0}^{\xi-1} \sum_{l=1}^{d_{s+k}} a_{0,1}^{k,l} f_{k,l} + \sum_{l=1}^{\mu-e_{\xi-1}} b_{0,1}^{l} g_{\xi-1,l},$$

where $a_{0,1}^{k,l}, b_{0,1}^l \in \mathbb{k}[\![\underline{\mathbf{x}}]\!]$. Note that, if $a_{0,1}^{0,l}$ is not a unit and $a_{0,1}^{0,l} \neq 0$, then $\sigma \operatorname{ord}(a_{0,1}^{0,l}f_{0,l}) \geq \xi + s > s = \sigma \operatorname{ord}(u_{0,1})$. Moreover, $\sigma \operatorname{ord}(g_{\xi-1,l}) \geq s + \xi > s = \sigma \operatorname{ord}(u_{0,1})$. In particular, the subset $\Lambda_{0,1} \subseteq \{1,\ldots,d_s\}$ of indices l such that $a_{0,1}^{0,l}$ is invertible, with term of order zero $\lambda_{0,1}^{0,l} \in \mathbb{k}$, $\lambda_{0,1}^{0,l} \neq 0$, is not empty. Taking σ -leading forms in both parts of the equality, one deduces that $u_{0,1}^{\sigma} = \sum_{l \in \Lambda_{0,1}} \lambda_{0,1}^{0,l} f_{0,l}^{\sigma}$. Renaming the elements of \mathcal{B}_0 , we can suppose that $1 \in \Lambda_{0,1}$, so $\lambda_{0,1}^{0,1}$ and $a_{0,1}^{0,1}$ are invertible. Set $\mathcal{B}_{0,1} := \{u_{0,1}, f_{0,2}, \ldots, f_{0,d_s}\}$. Clearly, $f_{0,1}^{\sigma}$ is in the vector space spanned by $\mathcal{B}_{0,1}^{\sigma}$ and $\mathcal{B}_{0,1}^{\sigma}$ is a basis of V_s . Moreover, since $a_{0,1}^{0,1}$ is invertible, $f_{0,1}$ is in the ideal generated by $\mathcal{B}_{0,1} \cup \mathcal{B}_2 \cup \ldots \cup \mathcal{B}_{\xi-1} \cup \mathcal{C}_{\xi-1}$. Thus, $\mathcal{B}_{0,1} \cup \mathcal{B}_2 \cup \ldots \cup \mathcal{B}_{\xi-1} \cup \mathcal{C}_{\xi-1}$ is a minimal generating set of Q.

If $\delta_0 = 1$, i.e., $\mathscr{D}_0 = \{u_{0,1}\}$, take $\mathscr{B}_0' = \mathscr{B}_{0,1} \setminus \mathscr{D}_0$, which satisfies the conditions \mathbb{L}_0 . Suppose that $\delta_0 \geq 2$. So, let $u_{0,2} \in \mathscr{D}_0 \subset Q$. Then

$$u_{0,2} = a_{0,2}^{0,1} u_{0,1} + \sum_{l=2}^{d_s} a_{0,2}^{0,l} f_{0,l} + \sum_{k=1}^{\xi-1} \sum_{l=1}^{d_{s+k}} a_{0,2}^{k,l} f_{k,l} + \sum_{l=1}^{\mu-e_{\xi-1}} b_{0,2}^l g_{\xi-1,l},$$

where $a_{0,2}^{k,l}, b_{0,2}^l \in \mathbb{k}[\![\underline{x}]\!]$. Let $\lambda_{0,2}^{0,l} \in \mathbb{k}$ be the term of order zero of $a_{0,2}^{0,l}$. Note that, if $a_{0,2}^{0,l}$ is not a unit and $a_{0,2}^{0,l} \neq 0$, then $\sigma \operatorname{ord}(a_{0,2}^{0,l}f_{0,l}) \geq \xi + s > s = \sigma \operatorname{ord}(u_{0,2})$. Moreover, $\sigma \operatorname{ord}(g_{\xi-1,l}) \geq s + \xi > s = \sigma \operatorname{ord}(u_{0,1})$. Let $\Lambda_{0,2} \subseteq \{2,\ldots,d_s\}$ be the subset of indices l, $1 \leq l \leq l \leq l$, such that $a_{0,2}^{0,l}$ is invertible, so $a_{0,2}^{0,l} \neq 0$. If $a_{0,2}^{0,l} = l$, a contradiction with $a_{0,2}^{0,l} = l$ being a linearly independent subset of $a_{0,2}^{0,l} = l$. Thus $a_{0,2}^{0,l} \neq l$ and $a_{0,2}^{0,l} = l$, a contradiction with $a_{0,2}^{0,l} = l$. Renaming the elements of $a_{0,1}^{0,l} = l$, we can suppose that $a_{0,2}^{0,l} = l$, so $a_{0,2}^{0,l} = l$ and $a_{0,2}^{0,l} = l$. Renaming the elements of $a_{0,1}^{0,l} = l$. Clearly, $a_{0,2}^{0,l} = l$, so $a_{0,2}^{0,l} = l$ and $a_{0,2}^{0,l} = l$ and $a_{0,2}^{0,l} = l$ and $a_{0,2}^{0,l} = l$ and $a_{0,2}^{0,l} = l$ is in the vector space spanned by $a_{0,2}^{0,l} = l$ is a basis of $a_{0,2}^{0,l} = l$. Furthermore, since $a_{0,2}^{0,l} = l$ is invertible, it follows that $a_{0,2}^{0,l} = l$ is a minimal generating set of $a_{0,2}^{0,l} = l$. Thus, $a_{0,2}^{0,l} = l$ is a minimal generating set of $a_{0,2}^{0,l} = l$.

If $\delta_0=2$, i.e., $\mathscr{D}_0=\{u_{0,1},u_{0,2}\}$, take $\mathscr{B}_0'=\mathscr{B}_{0,2}\setminus\mathscr{D}_0$, which clearly satisfies the conditions \mathbb{L}_0 . If $\delta_0>2$, proceed recursively to substitute all the elements of \mathscr{D}_0 in \mathscr{B}_0 . After some renaming, one gets $\mathscr{B}_{0,\delta_0}:=\{u_{0,1},\ldots,u_{0,\delta_0},f_{0,\delta_0+1},\ldots,f_{0,d_s}\}$, such that $\mathscr{B}_{0,\delta_0}\cup\mathscr{B}_2\cup\ldots\cup\mathscr{B}_{\xi-1}\cup\mathscr{C}_{\xi-1}$ is a minimal generating set of Q. Then the set $\mathscr{B}_0':=\mathscr{B}_{0,\delta_0}\setminus\mathscr{D}_0$ satisfies the conditions \mathbb{L}_0 .

Construction of \mathcal{B}'_i , $1 \leq i \leq \xi - 1$. Note that, in particular, $\xi \geq i + 1$. By the step i - 1, there exist a family of subsets $\mathcal{B}'_0, \ldots, \mathcal{B}'_{i-1}$, each \mathcal{B}'_k contained in \mathcal{B}_k , satisfying the conditions of \mathbb{L}_{i-1} . That is, $\mathcal{D}^{\sigma}_k \cup (\mathcal{B}'_k)^{\sigma}$ is a (possibly empty) basis of V_{s+k} , for $k = 0, \ldots, i-1$; the family of subsets of Q, $\mathcal{D}_0, \mathcal{B}'_0, \ldots, \mathcal{D}_{i-1}, \mathcal{B}'_{i-1}, \mathcal{B}_i, \ldots, \mathcal{B}_{\xi-1}, \mathcal{E}_{\xi-1}$ are pairwise disjoint and their union $(\mathcal{D}_0 \cup \mathcal{B}'_0) \cup \ldots \cup (\mathcal{D}_{i-1} \cup \mathcal{B}'_{i-1}) \cup \mathcal{B}_i \cup \ldots \cup \mathcal{B}_{\xi-1} \cup \mathcal{E}_{\xi-1}$ is a minimal generating set of Q. For the sake of convenience, set $\mathcal{B}'_k = \{u_{k,\delta_k+1}, \ldots, u_{k,d_{s+k}}\}$.

If $\mathscr{D}_i = \emptyset$, let $\mathscr{B}'_i = \mathscr{B}_i$, which satisfies the conditions \mathbb{L}_i . Suppose that $\mathscr{D}_i \neq \emptyset$. Take $u_{i,1} \in \mathscr{D}_i \subset Q$. Since $(\mathscr{D}_0 \cup \mathscr{B}'_0) \cup \ldots \cup (\mathscr{D}_{i-1} \cup \mathscr{B}'_{i-1}) \cup \mathscr{B}_i \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1}$ is a minimal generating set of Q, then

$$u_{i,1} = \sum_{k=0}^{i-1} \sum_{l=1}^{d_{s+k}} a_{i,1}^{k,l} u_{k,l} + \sum_{k=i}^{\xi-1} \sum_{l=1}^{d_{s+k}} a_{i,1}^{k,l} f_{k,l} + \sum_{l=1}^{\mu-e_{\xi-1}} b_{i,1}^{l} g_{\xi-1,l},$$
(2)

where $a_{i,1}^{k,l}, b_{i,1}^l \in \mathbb{k}[\![\underline{\mathbf{x}}]\!]$, $f_{k,l} \in \mathcal{B}_k$, $g_{\xi-1,l} \in \mathscr{C}_{\xi-1}$. If $a_{i,1}^{0,l}$ is not a unit and $a_{i,1}^{0,l} \neq 0$, then $\sigma \operatorname{ord}(a_{i,1}^{0,l}u_{0,l}) \geq \xi + s > s + i = \sigma \operatorname{ord}(u_{i,1})$. Let $\Lambda_{i,1}^0 \subseteq \{1,\ldots,d_s\}$ denote the subset of indices l such that $a_{i,1}^{0,l}$ is invertible, with term of order zero $\lambda_{i,1}^{0,l} \in \mathbb{k}$, $\lambda_{i,1}^{0,l} \neq 0$. Suppose that $\Lambda_{i,1}^0 \neq \emptyset$. Taking σ -leading forms in Equality (2), we deduce that $\sum_{l \in \Lambda_{i,1}^0} \lambda_{i,1}^{0,l} u_{k,l}^{\sigma} = 0$. Since $\mathscr{D}_0^{\sigma} \cup (\mathscr{B}_0')^{\sigma}$ is a basis of V_s , this implies that $\lambda_{i,1}^{0,l} = 0$, which is a contradiction. Therefore, all the $a_{i,1}^{0,l}$ are not units for all $l = 1, \ldots, d_s$. Similarly, one proves that $a_{i,1}^{k,l}$ are not units for all $k = 1, \ldots, i-1$ and all $l = 1, \ldots, d_{s+k}$.

Let now $\Lambda_{i,1}^i \subseteq \{1,\ldots,d_{s+i}\}$ be the subet of indices l such that $a_{i,1}^{i,l}$ is invertible, with term of order zero $\lambda_{i,1}^{i,l}$, $\lambda_{i,1}^{i,l} \neq 0$. Taking σ -leading forms, one deduces that $u_{i,1}^{\sigma} = \sum_{l \in \Lambda_{i,1}^i} \lambda_{i,1}^{i,l} f_{i,1}^{\sigma}$. Since $\mathscr{B}_i^{\sigma} = \{f_{i,1}^{\sigma},\ldots,f_{i,d_{s+i}}^{\sigma}\}$ is a basis of V_{s+i} and $u_{i,1}^{\sigma} \in V_{s+i}$, $u_{i,1}^{\sigma} \neq 0$, renaming the elements of \mathscr{B}_i , one can suppose that $1 \in \Lambda_{i,1}^i$, so $\lambda_{i,1}^{i,1}$ and $a_{i,1}^{i,1}$ are invertible. Set $\mathscr{B}_{i,1} = \{u_{i,1},f_{i,2},\ldots,f_{i,d_{s+i}}\}$. Then, $f_{i,1}^{\sigma}$ is in the vector space spanned by $\mathscr{B}_{i,1}^{\sigma}$ and $\mathscr{B}_{i,1}^{\sigma}$ is a basis of V_{s+i} . Moreover, since $a_{i,1}^{i,1}$ is in the ideal generated by

$$(\mathscr{D}_0 \cup \mathscr{B}'_0) \cup \ldots \cup (\mathscr{D}_{i-1} \cup \mathscr{B}'_{i-1}) \cup \mathscr{B}_{i,1} \cup \mathscr{B}_{i+1} \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1},$$

which is a minimal generating set for Q.

If $\delta_i = 1$, i.e., $\mathscr{D}_i = \{u_{i,1}\}$, take $\mathscr{B}'_i = \mathscr{B}_{i,1} \setminus \mathscr{D}_0$, which satisfies \mathbb{L}_i . Suppose that $\delta_i \geq 2$. Let $u_{i,2} \in \mathcal{D}_0 \subset Q$. Then, similarly as before, write

$$u_{i,2} = \sum_{k=0}^{i-1} \sum_{l=1}^{d_{s+k}} a_{i,2}^{k,l} u_{k,l} + a_{i,2}^{i,1} u_{i,1} + \sum_{l=2}^{d_{s+i}} a_{i,2}^{i,l} f_{i,l} + \sum_{k=i+1}^{\xi-1} \sum_{l=1}^{d_{s+k}} a_{i,2}^{k,l} f_{k,l} + \sum_{l=1}^{\mu-e_{\xi-1}} b_{i,2}^{l} g_{\xi-1,l}.$$

As before, one first deduces that $a_{i,2}^{k,l}$ are not units, for $k=0,\ldots,i-1$, and $l=1,\ldots,d_{s+k}$. So, $\operatorname{\sigma ord}(a_{i,2}^{k,l}u_{k,l}) \geq \xi + s > s + i = \operatorname{\sigma ord}(u_{i,2})$. Let $\Lambda_{i,2} \subseteq \{2,\ldots,d_{s+i}\}$ denote the subset of indices $l \geq 2$ such that $a_{i,2}^{i,l}$ is invertible, with term of order zero $\lambda_{i,2}^{i,l} \in \mathbb{k}$, $\lambda_{i,2}^{i,l} \neq 0$. If $\Lambda_{i,2}$ were empty, on taking σ -leading forms in the equality above one would deduce that $u_{i,2}^{\sigma} = \lambda_{i,2}^{i,1} u_{i,1}^{\sigma}$, a contradiction with \mathscr{D}_i^{σ} being a set of linearly independent elements of V_{s+i} . Thus $\Lambda_{i,2} \neq \emptyset$ and $u_{i,2}^{\sigma} = \lambda_{i,2}^{i,1} u_{i,1}^{\sigma} + \sum_{l \in \Lambda_{i,2}} \lambda_{i,2}^{i,l} f_{i,l}^{\sigma}$. Renaming the elements of $\mathcal{B}_{i,1}$, we can suppose that $2 \in \Lambda_{i,2}$, so $\lambda_{i,2}^{i,2}$ and $a_{i,2}^{i,2}$ are invertible. Set $\mathcal{B}_{i,2} = \{u_{i,1}, u_{i,2}, f_{i,3}, \dots, f_{i,d_{s+i}}\}$. Then, $f_{i,2}^{\sigma}$ is in the vector space spanned by $\mathscr{B}_{i,2}^{\sigma}$ and $\mathscr{B}_{i,2}^{\sigma}$ is a basis of V_{s+i} . Furthermore, since $a_{i,2}^{i,2}$ is invertible, $f_{i,2}$ is in the ideal generated by

$$(\mathscr{D}_0 \cup \mathscr{B}'_0) \cup \ldots \cup (\mathscr{D}_{i-1} \cup \mathscr{B}'_{i-1}) \cup \mathscr{B}_{i,2} \cup \mathscr{B}_{i+1} \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1},$$

which is a minimal generating set for Q. If $\delta_i = 2$, i.e., $\mathcal{D}_i = \{u_{i,1}, u_{i,2}\}$, take $\mathcal{B}'_i = \mathcal{B}_{i,2} \setminus \mathcal{D}_i$, which clearly satisfies the conditions \mathbb{L}_i . Otherwise, proceed recursively to substitute all the elements of \mathcal{D}_i in \mathcal{B}_i . After some renaming, one gets $\mathcal{B}_{i,\delta_i} := \{u_{i,1}, \dots, u_{i,\delta_i}, f_{i,\delta_i+1}, \dots, f_{i,d_{s+i}}\}$, where

$$(\mathscr{D}_0 \cup \mathscr{B}'_0) \cup \ldots \cup (\mathscr{D}_{i-1} \cup \mathscr{B}'_{i-1}) \cup \mathscr{B}_{i,\delta_i} \cup \mathscr{B}_{i+1} \cup \ldots \cup \mathscr{B}_{\xi-1} \cup \mathscr{C}_{\xi-1}$$

is a minimal generating set of Q. Then, the set $\mathscr{B}'_i := \mathscr{B}_{i,\delta_i} \setminus \mathscr{D}_i$ satisfies the conditions of \mathbb{L}_i , which finishes the whole proof.

4. The example of Moh, for n=3, in any characteristic

In this section, \Bbbk is a field and $\rho: \Bbbk[\![x,y,z]\!] \to \Bbbk[\![t]\!]$ is the \Bbbk -algebra morphism defined by Moh in [7], with n=3, m=2 and $\lambda=25$, so $\rho(x)=t^6+t^{31}=t^6(1+t^{25})$, $\rho(y)=t^8$ and $\rho(z)=t^{10}$. Moreover $\sigma: \mathbb{k}[[x,y,z]] \to \mathbb{k}[[x,y,z]]$ is the k-algebra morphism defined by $\sigma(x) = x^3$, $\sigma(y) = y^4$ and $\sigma(z) = z^5$ and $S = \langle 3, 4, 5 \rangle$ is the numerical semigroup generated by 3, 4, 5. The purpose of this section is to find minimal generating sets for the first prime of Moh, $P = \ker(\rho)$, showing that its cardinality depend on the characteristic of the field k, contradicting a statement of Sally.

Remark 4.1. For two natural numbers m, n, let $b_{m,n} \in \mathbb{k}$ be the image of the binomial number $\binom{m}{n} \in \mathbb{N}$ through the ring homomorphism $\mathbb{Z} \to \mathbb{k}$. We understand that if $m \leq n$, then $\binom{m}{n} = 0$. The well-known Theorem of Lucas says that $\binom{m}{n} \equiv \prod_{i=0}^k \binom{m_i}{n_i} \mod p$, where $m = m_0 + m_1 \cdot p + \ldots + m_k \cdot p^k$ and $n = n_0 + n_1 \cdot p + \ldots + n_k \cdot p^k$, $0 \le m_i, n_i < p$. In particular, if char(\mathbb{k}) = p, then $b_{m,n} = 0$ if and only if $m_i < n_i$, for some i.

Lemma 4.2. Let $r \in \mathcal{S}$, $r \neq 0$. Let $\nu_r = \max\{\alpha_1 \mid \alpha \in F(r, \mathcal{S})\}$. The following hold.

- (1) If $\alpha \in F(r, \mathcal{S})$, then $\rho(\mathbf{x}^{\alpha}) = t^{2r}(1 + t^{25})^{\alpha_1} = t^{2r} \sum_{k=0}^{\alpha_1} b_{\alpha_1, k} t^{25k}$. In particular, $\rho(\mathbf{x}^{\alpha}) \neq 0$, $\{2r, 2r + 25\alpha_1\} \subseteq \operatorname{Supp}(\rho(\mathbf{x}^{\alpha})) \subseteq \{2r + 25k \mid 0 \leq k \leq \alpha_1\} \text{ and } \operatorname{ord}(\rho(\mathbf{x}^{\alpha})) = 2r$. (2) Let $g = \sum_{\alpha \in F(r, \mathcal{S})} \lambda_{\alpha} \mathbf{x}^{\alpha} \in W_r$. If $g \in V_r$, then $\sum_{\alpha \in F(r, \mathcal{S})} \lambda_{\alpha} = 0$ and $\sum_{\alpha \in F(r, \mathcal{S})} b_{\alpha_1, 1} \lambda_{\alpha} = 0$.
- Furthermore.
 - (3) If dim $W_r = 1$, then $V_r = 0$.
 - (4) If $W_r = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta} \rangle$, $\alpha \neq \beta$, then $V_r \subseteq \langle \mathbf{x}^{\alpha} \mathbf{x}^{\beta} \rangle$. Moreover, if $b_{\alpha_1, 1} \neq b_{\beta_1, 1}$, then $V_r = 0$.
 - (5) Suppose that $W_r = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta}, \mathbf{x}^{\gamma} \rangle$, with α, β, γ distinct, and

$$(b_{\gamma_1,1}-b_{\beta_1,1},b_{\alpha_1,1}-b_{\gamma_1,1},b_{\beta_1,1}-b_{\alpha_1,1})\neq (0,0,0).$$

Then
$$V_r \subseteq \langle (b_{\gamma_1,1} - b_{\beta_1,1}) \mathbf{x}^{\alpha} + (b_{\alpha_1,1} - b_{\gamma_1,1}) \mathbf{x}^{\beta} + (b_{\beta_1,1} - b_{\alpha_1,1}) \mathbf{x}^{\gamma} \rangle$$
.

Proof. If $r \in \mathcal{S}$ and $\alpha \in F(r, \mathcal{S})$, then $r = \alpha_1 \cdot 3 + \alpha_2 \cdot 4 + \alpha_3 \cdot 5$. Thus,

$$\rho(\mathbf{x}^{\alpha}) = (t^6 + t^{31})^{\alpha_1} t^{8\alpha_2} t^{10\alpha_3} = t^{2r} (1 + t^{25})^{\alpha_1} = t^{2r} \left(b_{\alpha_1,0} + b_{\alpha_1,1} t^{25} + \dots + b_{\alpha_1,\alpha_1} t^{25\alpha_1} \right).$$

Since $b_{\alpha_1,0} = b_{\alpha_1,\alpha_1} = 1$, $\rho(\mathbf{x}^{\alpha}) \neq 0$ and $\{2r, 2r + 25\alpha_1\} \subseteq \operatorname{Supp}(\rho(\mathbf{x}^{\alpha})) \subseteq \{2r + 25k \mid 0 \leq k \leq \alpha_1\}$. So $\sigma \operatorname{ord}(\rho(\mathbf{x}^{\alpha})) = 2r$. Note that, depending on the characteristic of \mathbb{k} , some $b_{\alpha_1,k}$ might be zero, which is the reason why the equality may fail. This proves (1).

Let $g = \sum_{\alpha \in F(r,\mathcal{S})} \lambda_{\alpha} \mathbf{x}^{\alpha} \in W_r$. By Item (1), and using that, for any $\alpha \in F(r,\mathcal{S})$, $\alpha_1 \leq \nu_r$, then

$$\rho(g) = \sum_{\alpha \in F(r,S)} \lambda_{\alpha} t^{2r} \left(b_{\alpha_{1},0} + b_{\alpha_{1},1} t^{25} + \dots + b_{\alpha_{1},\alpha_{1}} t^{25\alpha_{1}} \right) = \sum_{\alpha \in F(r,S)} \lambda_{\alpha} t^{2r} \left(b_{\alpha_{1},0} + b_{\alpha_{1},1} t^{25} + \dots + b_{\alpha_{1},\alpha_{1}} t^{25\alpha_{1}} + \dots + b_{\alpha_{1},\nu_{r}} t^{25\nu_{r}} \right) =$$

$$\left(\sum_{\alpha \in F(r,\mathcal{S})} b_{\alpha_1,0} \lambda_{\alpha}\right) t^{2r} + \left(\sum_{\alpha \in F(r,\mathcal{S})} b_{\alpha_1,1} \lambda_{\alpha}\right) t^{2r+25} + \dots + \left(\sum_{\alpha \in F(r,\mathcal{S})} b_{\alpha_1,\nu_r} \lambda_{\alpha}\right) t^{2r+25\nu_r},$$

where $b_{\alpha_1,k} = 0$, whenever $k > \alpha_1$. If $g \in \ker(\rho)$, then $\sum_{\alpha \in F(r,\mathcal{S})} b_{\alpha_1,k} \lambda_{\alpha} = 0$, for all $0 \le k \le \nu_r$. Suppose that $g \in V_r$, $g \notin \ker(\rho)$. By Lemma 2.4, there exists $h = \sum_{s>r} \sum_{\alpha \in F(s,\mathcal{S})} \mu_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{k}[[x,y,z]]$, $h \neq 0$, $\sigma \operatorname{ord}(h) > r$, such that $\rho(g) + \rho(h) = 0$. Since $g \notin \ker(\rho)$, $\rho(h) \neq 0$. As before, and letting $\nu_s = \max\{\alpha_1 \mid \alpha \in F(s,\mathcal{S})\}$, then $\rho(h)$ is equal to

$$\sum_{s>r} \left[\left(\sum_{\alpha \in \mathcal{F}(s,\mathcal{S})} b_{\alpha_1,0} \mu_{\alpha} \right) t^{2s} + \left(\sum_{\alpha \in \mathcal{F}(s,\mathcal{S})} b_{\alpha_1,1} \mu_{\alpha} \right) t^{2s+25} + \dots + \left(\sum_{\alpha \in \mathcal{F}(s,\mathcal{S})} b_{\alpha_1,\nu_s} \mu_{\alpha} \right) t^{2s+25\nu_s} \right].$$

From the equality $\rho(g) + \rho(h) = 0$ and since 2r < 2s, $2r + 25 \neq 2s$ and 2r + 25 < 2s + 25, it follows that $\sum_{\alpha \in F(r,\mathcal{S})} b_{\alpha_1,0} \lambda_{\alpha} = 0$ and $\sum_{\alpha \in F(r,\mathcal{S})} b_{\alpha_1,1} \lambda_{\alpha} = 0$. This proves (2).

If $W_r = \langle \mathbf{x}^{\alpha} \rangle$ and $g = \lambda \mathbf{x}^{\alpha} \in V_r$, then, by (2), $\lambda = 0$ and g = 0. This proves (3).

Suppose that $W_r = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta} \rangle$ and $g = \lambda \mathbf{x}^{\alpha} + \mu \mathbf{x}^{\beta} \in W_r$. By (2), $\lambda + \mu = 0$ and $b_{\alpha_1, 1}\lambda + b_{\beta_1, 1}\mu = 0$. Thus, $g = \lambda(\mathbf{x}^{\alpha} - \mathbf{x}^{\beta})$. Moreover, if $b_{\alpha_1, 1} \neq b_{\beta_1, 1}$, then $\lambda = 0$ and g = 0, which proves (4).

Suppose that $W_r = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta}, \mathbf{x}^{\gamma} \rangle$ and let $g = \lambda \mathbf{x}^{\alpha} + \mu \mathbf{x}^{\beta} + \eta \mathbf{x}^{\gamma} \in V_r$. By (2), $\lambda + \mu + \eta = 0$ and $b_{\alpha_1,1}\lambda + b_{\beta_1,1}\mu + b_{\gamma_1,1}\eta = 0$. By hypothesis,

$$\operatorname{rank}\left(\begin{array}{ccc} 1 & 1 & 1 \\ b_{\alpha_1,1} & b_{\beta_1,1} & b_{\gamma_1,1} \end{array}\right) = 2$$

and $(\lambda, \mu, \eta) \in \langle (b_{\gamma_1, 1} - b_{\beta_1, 1}, b_{\alpha_1, 1} - b_{\gamma_1, 1}, b_{\beta_1, 1} - b_{\alpha_1, 1}) \rangle$, which shows (5).

Convention 4.3. From now on, a subset of monomials will be ordered from biggest to smallest, according to the local negative degree reverse lexicographical order $>_{ds}$ on the set of monomials $Mon_3 = \{x^{\alpha} \mid \alpha \in \mathbb{N}^3\}$, that is: $x^{\alpha} >_{ds} x^{\beta} \Leftrightarrow \deg(x^{\alpha}) < \deg(x^{\beta})$ or $\deg(x^{\alpha}) = \deg(x^{\beta})$ and there exists $1 \leq i \leq d-1$ such that $\alpha_d = \beta_d, \ldots, \alpha_{i+1} = \beta_{i+1}$ and $\alpha_i < \beta_i$. For instance,

$$1 > x > y > z > x^2 > xy > y^2 > xz > yz > z^2 > x^3 > x^2y > xy^2 > y^3 > x^2z > xyz > \dots$$

A non-zero polynomial f will be written as $f = \sum_{\nu=0}^{n} a_{\nu} \mathbf{x}^{\alpha(\nu)}$, $a_{\nu} \in \mathbb{k} \setminus \{0\}$, with $\mathbf{x}^{\alpha(0)} > \ldots > \mathbf{x}^{\alpha(n)}$, so the leading monomial of f is $\mathrm{LM}(f) = \mathbf{x}^{\alpha(0)}$ (see Notation 5.1). A subset of polynomials f_1, \ldots, f_r will be listed according to $\mathrm{LM}(f_1) > \ldots > \mathrm{LM}(f_r)$. We will usually underline the σ -leading form of any polynomial with more than one term.

Remark 4.4. A simple calculation of F(r, S) gives rise to the following W_r , $r \in S$, $3 \le r \le 16$:

$$\begin{split} W_{3} &= \langle x \rangle, \ W_{4} = \langle y \rangle, \ W_{5} = \langle z \rangle, \ W_{6} = \langle x^{2} \rangle, \ W_{7} = \langle xy \rangle, \ W_{8} = \langle y^{2}, xz \rangle, \ W_{9} = \langle yz, x^{3} \rangle, \\ W_{10} &= \langle z^{2}, x^{2}y \rangle, \ W_{11} = \langle xy^{2}, x^{2}z \rangle, \ W_{12} = \langle y^{3}, xyz, x^{4} \rangle, \ W_{13} = \langle y^{2}z, xz^{2}, x^{3}y \rangle, \\ W_{14} &= \langle yz^{2}, x^{2}y^{2}, x^{3}z \rangle, \ W_{15} = \langle z^{3}, xy^{3}, x^{2}yz, x^{5} \rangle, \ W_{16} = \langle x^{4}y, y^{4}, xy^{2}z, x^{2}z^{2} \rangle. \end{split}$$

Lemma 4.5. Let $s = \min\{\sigma \operatorname{ord}(f) \mid f \in P, f \neq 0\}$. If $\operatorname{char}(\mathbb{k}) = 0$ or $p \geq 5$, then s = 12. If $\operatorname{char}(\mathbb{k}) = 2$, then s = 10. If $\operatorname{char}(\mathbb{k}) = 3$, then s = 9.

Proof. By Lemma 4.2, $V_r = 0$, for $3 \le r \le 7$. Suppose that $\operatorname{char}(\mathbb{k}) = 0$ or $p \ge 5$. By Lemma 4.2, we deduce that:

- $W_8 = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta} \rangle$, with $\alpha = (0, 2, 0), \beta = (1, 0, 1), b_{\alpha_1, 1} b_{\beta_1, 1} = -1$ and $V_8 = 0$;
- $W_9 = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta} \rangle$, with $\alpha = (0, 1, 1), \beta = (3, 0, 0), b_{\alpha_1, 1} b_{\beta_1, 1} = -3 \text{ and } V_9 = 0$;
- $W_{10} = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta} \rangle$, with $\alpha = (0, 0, 2)$, $\beta = (2, 1, 0)$, $b_{\alpha_1, 1} b_{\beta_1, 1} = -2$ and $V_{10} = 0$;
- $W_{11} = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta} \rangle$, with $\alpha = (1, 2, 0), \beta = (2, 0, 1), b_{\alpha_1, 1} b_{\beta_1, 1} = -1$ and $V_{11} = 0$;
- $W_{12} = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta}, \mathbf{x}^{\gamma} \rangle$, with $\alpha = (0, 3, 0)$, $\beta = (1, 1, 1)$, $\gamma = (4, 0, 0)$ and $(b_{\gamma_1, 1} b_{\beta_1, 1}, b_{\alpha_1, 1} b_{\gamma_1, 1}, b_{\beta_1, 1} b_{\alpha_1, 1}) = (3, -4, 1)$.

Let us find $f_1 \in P$, $f_1 \neq 0$, with $f_1^{\sigma} \in V_{12}$. By Lemma 4.2, $V_{12} \subseteq \langle 3y^3 - 4xyz + x^4 \rangle$, so pick $f_1^{\sigma} = 3y^3 - 4xyz + x^4$. Then, $\rho(f_1^{\sigma}) = 6t^{74} + 4t^{99} + t^{124}$. Now we want to construct the tail f_1^{τ} . As a general rule, we will try to find a σ -homogeneous tail whose monomials have exponents in x as smallest as possible. By Lemma 4.2, if $r \in \mathcal{S}$, $r \neq 0$, and $\alpha \in F(r, \mathcal{S})$, then $\operatorname{ord}(\rho(\mathbf{x}^{\alpha})) = 2r$. Thus, pick $f_1^{\tau} \in W_{37}$, where W_{37} is generated by the monomials

$$y^3z^5, xyz^6, y^8z, xy^6z^2, x^2y^4z^3, x^3y^2z^4, x^4z^5, x^3y^7, x^4y^5z, x^5y^3z^2, x^6yz^3, x^7y^4, x^8y^2z, x^9z^2, x^{11}y.$$

Note that the image by ρ of the monomials $y^{\alpha_2}z^{\alpha_3} \in W_{37}$ is just t^{74} . However, we need to get a t^{99} , which has an odd exponent. Thus, f_1^{τ} must contain at least a monomial multiple of x. On the other hand, to get the term t^{124} , we need a monomial multiple of x^2 . Considering that $\rho(xy^6z^2)=t^{74}+t^{99}$ and that $\rho(x^2y^4z^3)=t^{74}+2t^{99}+t^{124}$, we see that one can take $f_1^{\tau}=3y^3z^5-2xy^6z^2-x^2y^4z^3$. Therefore, let $f_1:=3y^3-4xyz+x^4-3y^3z^5-2xy^6z^2-x^2y^4z^3$. Then, $f_1^{\sigma}=3y^3-4xyz+x^4\in W_{12}$, $f_1\in\ker(\rho)=P, f_1^{\sigma}\in V_{12}$ and $s=\min\{\sigma\mathrm{ord}(f)\mid f\in P, f\neq 0\}=12$.

Suppose that $\operatorname{char}(\Bbbk)=2$. From the previous discussion we deduce that $V_r=0$, for $3 \leq r \leq 9$. Let us find $g_1 \in P$, $g_1 \neq 0$, with $g_1^{\sigma} \in V_{10}$. As before, by Lemma 4.2, g_1^{σ} must be of the form $g_1^{\sigma}=z^2+x^2y$. Then, $\rho(g_1^{\sigma})=t^{70}$. Take the monomial $y^5z^3 \in W_{35}$, which has no x and whose image is $\rho(y^5z^3)=t^{70}$. Let $g_1:=z^2+x^2y+y^5z^3$. Then, $g_1^{\sigma}=z^2+x^2y \in W_{10}$, $g_1 \in \ker(\rho)=P$, $g_1^{\sigma} \in V_{10}$ and $s=\min\{\sigma\operatorname{ord}(f) \mid f \in P, f \neq 0\}=10$.

Suppose that $\operatorname{char}(\mathbb{k})=3$. As before, we can affirm that $V_r=0$, for $3 \leq r \leq 8$. Let us find $h_1 \in P, \ h_1 \neq 0$, with $h_1^{\sigma} \in V_9$. By Lemma 4.2, $h_1^{\sigma} = yz - x^3$. Then, $\rho(h_1^{\sigma}) = -t^{93}$. To vanish this term, the tail must contain a monomial multiple of x, so we pick one of the form $xy^{\alpha_2}z^{\alpha_3}$. Then, $\rho(xy^{\alpha_2}z^{\alpha_3}) = t^{6+8\alpha_2+10\alpha_3} + t^{31+8\alpha_2+10\alpha_3}$. A possible solution of $31 + 8\alpha_2 + 10\alpha_3 = 93$ is $(\alpha_2, \alpha_3) = (4, 3)$. Note that $xy^4z^3 \in W_{34}$ and $\rho(xy^4z^3) = t^{68} + t^{93}$. To vanish t^{68} , we can choose y^6z^2 , which is in W_{34} . So, let $h_1 = \underline{yz - x^3} - y^6z^2 + xy^4z^3 \in \mathbb{k}[[x, y, z]]$. Then, $h_1^{\sigma} = yz - x^3 \in W_9$, $h_1 \in \ker(\rho) = P$, $h_1^{\sigma} \in V_9$ and $s = \min\{\sigma \operatorname{ord}(f) \mid f \in P, f \neq 0\} = 9$.

Discussion 4.6. Suppose that $\operatorname{char}(\mathbb{k}) = 0$ or p > 5. Let us find $f_1, f_2, f_3, f_4 \in P$, $f_i \neq 0$, such that $f_1^{\sigma} \in V_{12}, f_2^{\sigma} \in V_{13}, f_3^{\sigma} \in V_{14}$ and $f_4^{\sigma} \in V_{15}$. Observe that $s = \min\{\sigma\operatorname{ord}(f) \mid f \in P, f \neq 0\} = 12$ and $\xi = 3$. So, by Theorem 3.1, we are interested in finding elements whose σ -leading form are in V_{12}, V_{13} and V_{14} . Moreover, by [8, Theorem], we also have to consider V_{15} , where, for n = 3, $n^2 + 2n = 15$.

By Lemma 4.5, we can pick $f_1 = 3y^3 - 4xyz + x^4 - 3y^3z^5 - 2xy^6z^2 - x^2y^4z^3$. By Lemma 4.2, we can pick $f_2^{\sigma} = 2y^2 - 3xz^2 + x^3y \in V_{13}$. Then, $\rho(f_2^{\sigma}) = 3t^{76} + t^{101}$. Let us find

By Lemma 4.2, we can pick $f_2^{\sigma} = 2y^2 - 3xz^2 + x^3y \in V_{13}$. Then, $\rho(f_2^{\sigma}) = 3t^{76} + t^{101}$. Let us find $f_2^{\tau} \in W_{38}$ with monomials whose exponents in x be as smallest as possible. To vanish t^{101} , take $xy^{\alpha_2}z^{\alpha_3}$, so $\rho(xy^{\alpha_2}z^{\alpha_3}) = t^{6+8\alpha_2+10\alpha_3} + t^{31+8\alpha_2+10\alpha_3}$. To obtain $31 + 8\alpha_2 + 10\alpha_3 = 101$, choose $(\alpha_2, \alpha_3) = (5, 3)$. So, $xy^5z^3 \in W_{38}$ and $\rho(xy^5z^3) = t^{76} + t^{101}$. To vanish t^{76} , choose $y^7z^2 \in W_{38}$. So, we can pick the polynomial $f_2 = 2y^2 - 3xz^2 + x^3y - 2y^7z^2 - xy^5z^3$.

By Lemma 4.2, we can pick $f_3^{\sigma} = yz^2 - 3x^2y^2 + 2x^3z \in V_{14}$. Then, $\rho(f_3^{\sigma}) = 3t^{78} + 2t^{103}$. Let us find $f_3^{\tau} \in W_{39}$ with monomials whose exponents in x be as smallest as possible. To vanish t^{103} , take $xy^{\alpha_2}z^{\alpha_3}$ and solve $31 + 8\alpha_2 + 10\alpha_3 = 103$. Thus, we can choose $(\alpha_2, \alpha_3) = (4, 4)$. So, $xy^4z^4 \in W_{39}$

and $\rho(xy^4z^4)=t^{78}+t^{103}$. To vanish t^{78} , choose $y^6z^3\in W_{39}$. So, we can pick the polynomial $f_3=\underline{yz^2-3x^2y^2+2x^3z-y^6z^3-2xy^4z^4}$. Finally, take $f_4^\sigma=az^3+bxy^3+cx^2yz\in V_{15}$, omitting the x^5 for the sake of simplicity. Then,

$$\rho(f_4^{\sigma}) = (a+b+c)t^{30} + (b+2c)t^{55} + ct^{80}.$$

Choose (a,b,c)=(1,-2,1), so $f_4^{\sigma}=z^3-2xy^3+x^2yz$ and $\rho(f_4^{\sigma})=t^{80}$. To vanish t^{80} , choose $y^5z^4\in W_{40}$. Thus, we can take $f_4=\underline{z^3-2xy^3+x^2yz}-y^5z^4$.

Note that the σ -leading forms f_1^{σ} , $\overline{f_2^{\sigma}}$, f_3^{σ} , f_4^{σ} , coincide with the homonymous elements found in [6, Exemple 4.4], though in that example $\lambda = 27$.

In the next theorem we prove that the polynomials f_1, f_2, f_3, f_4 form a minimal generating set of P if char(\mathbb{k}) = 0 or $p \geq 5$. The demonstration consists in proving that the ideal $I := (f_1, f_2, f_3, f_4)$ is equal to P, where, by construction $I \subseteq P$. To prove the other inclusion we calculate the lengths of R/(I+yR) and R/(P+yR), where $R=\mathbb{k}[x,y,z]$, and see that they coincide. The advantage of this method is that it can be reproduced whatever the characteristic of k is.

Theorem 4.7. If $char(\mathbb{k}) = 0$ or $p \geq 5$, then P is minimally generated by

$$f_1 = \underbrace{3y^3 - 4xyz + x^4}_{f_3 = yz^2 - 3x^2y^2 + 2x^3z} - 2xy^6z^2 - x^2y^4z^3, \quad f_2 = \underbrace{2y^2z - 3xz^2 + x^3y}_{f_3 = yz^2 - 3x^2y^2 + 2x^3z} - y^6z^3 - 2xy^4z^4, \quad f_4 = z^3 - 2xy^3 + x^2yz - y^5z^4.$$

Proof. Let $R = \mathbb{k}[x, y, z]$, with maximal ideal $\mathfrak{m} = (x, y, z)$. Set $I = (f_1, f_2, f_3, f_4)$. By Discussion 4.6, $I \subseteq P$. For the sake of clarity, we divide the proof in several steps.

Step 1: If I = P, then $\mu(P) = 4$ and f_1, f_2, f_3, f_4 is a minimal system of generators of P.

By Lemma 4.5, Discussion 4.6 and Theorem 3.1, it follows that f_1, f_2, f_3 is part of a minimal system of generators of P, so $\mu(P) \geq 3$. Since $(f_1, f_2, f_3) \subset (x, y)$ and $f_4 = z^3 + g$, with $g \in (x, y)$, it follows that $f_4 \notin (f_1, f_2, f_3)$. If $\mu(P) = 3$, then $(f_1, f_2, f_3) = P$ and, since $f_4 \in P$, then $f_4 \in (f_1, f_2, f_3)$, a contradiction. Therefore, $\mu(P) > 3$. But, if I = P, then $\mu(P) \le 4$. So, $\mu(P) = 4$ and f_1, f_2, f_3, f_4 is a minimal system of generators of P.

Step 2: ht(I), ht(P) = 2 and $length_R(R/(I+yR))$, $length_R(R/(P+yR))$ are finite.

Clearly, $(f_1, f_4, y) = (x^4, z^3, y)$ and x^4, z^3, y is a regular sequence. By [1, Corollary 1.6.19.], $\operatorname{grade}(f_1, f_4, y) = \operatorname{grade}(x^4, z^3, y) = 3$ and f_1, f_4, y is a regular sequence. In particular, since $P \subseteq \mathfrak{m}$, $2 \leq \operatorname{grade}(I) = \operatorname{ht}(I) \leq \operatorname{ht}(P) \leq 2$ and $\operatorname{ht}(I) = \operatorname{ht}(P) = 2$. Since $(f_1, f_4, y) \subset I + yR \subseteq P + yR$, it follows that $\operatorname{grade}(I+yR)$, $\operatorname{grade}(P+yR) \geq 3$, so $\operatorname{ht}(I+yR) = 3$, $\operatorname{ht}(P+yR) = 3$ and R/(I+yR), R/(P+yR) are Artinian rings. Thus, R/(I+yR) and R/(P+yR) have finite length as R-modules.

STEP 3: length_R(R/(I+yR)) = 8.

Set $J = I + yR = (x^4, xz^2, x^3z, z^3, y)$. Let S = R/yR, which is a local ring of maximal $\mathfrak{m}_S =$ \mathfrak{m}/yR , say, with residue field $S/\mathfrak{m}_S = \mathbb{k}$. By abuse of notation, we still denote x, z to the classes of the variables in S. Let $\overline{J} = J/yR = (x^4, xz^2, x^3z, z^3)$. Since $R \to S$ is a surjective map and $S/\overline{J} = R/J$ is an S-module, it follows that $\operatorname{length}_R(R/(I+yR)) = \operatorname{length}_R(R/J) = \operatorname{length}_R(S/\overline{J}) = \operatorname{length}_R(S/\overline{J})$ $\operatorname{length}_S(S/\overline{J})$. For $0 \leq i \leq 2$, understanding $\mathfrak{m}_S^0 = S$ and noting that $\overline{J} \subset \mathfrak{m}_S^3$, we consider the short exact sequences:

$$0 \to \mathfrak{m}_S^{i+1}/\overline{J} \to \mathfrak{m}_S^i/\overline{J} \to \mathfrak{m}_S^i/\mathfrak{m}_S^{i+1} \to 0.$$

By the additivity of length, length_S $(S/\overline{J}) = 6 + \text{length}_S(\mathfrak{m}_S^3/\overline{J})$. Since $\mathfrak{m}_S(\mathfrak{m}_S^3/\overline{J}) = 0$, it follows that $\mathfrak{m}_S^3/\overline{J} = \langle x^3, x^2z \rangle$ is a two-dimensional S/\mathfrak{m}_S -vectorial space. Thus, length_S $(S/\overline{J}) = 8$.

Let D = R/P, which is a local Noetherian one-dimensional domain. Let $\mathfrak{m}_D = \mathfrak{m}/P$ be its maximal ideal and $k_D = \mathbb{k}$ its residue field. By abuse of notation, let x, y, z still denote the classes of the variables in D, so $\mathfrak{m}_D = (x, y, z)$. Since R is Cohen-Macaulay, $\dim(R/P) = \dim(R) - \operatorname{ht}(P) = 1$. Note that $D = R/P = \mathbb{k}[[x, y, z]] / \ker(\rho) \cong \operatorname{Im}(\rho) = \mathbb{k}[[t^6 + t^{31}, t^8, t^{10}]]$.

Step 4: The integral closure of $D = \mathbb{k}[[t^6 + t^{31}, t^8, t^{10}]]$ (in its field of fractions) is $\mathbb{k}[[t]]$.

Let $\mathbb{k}((t)) = \{\sum_{r=-n}^{\infty} \lambda_r t^r \mid \lambda_r \in \mathbb{k}, n \in \mathbb{N}\}$ be the ring of formal Laurent series over \mathbb{k} , which is well-known to be the field of fractions of $\mathbb{k}[[t]]$.

Let K(D) denote the field of fractions of D. Since $t^8t^{-10} = t^{-2} \in K(D)$, it follows that $t^2 \in K(D)$, so $(t^6 + t^{31} - (t^2)^3)(t^{-2})^{15} = t \in K(D)$ and $t^{-1} \in K(D)$. Since $t \in K(D)$ is integral over D, then $D \subset D[t]$ is an integral extension. Clearly, $D[t] = \mathbb{k}[[t^6 + t^{31}, t^8, t^{10}]][t]$ is a subring of $\mathbb{k}[[t]]$. Let us prove that $D[t] = \mathbb{k}[[t]]$. To do that, consider the numerical semigroup $S = \langle 4, 5 \rangle$, whose Frobenius number is F(S) = 11, and M the subsemigroup $M = 2S = \{2s \mid s \in S\} = \langle 8, 10 \rangle$, where 2F(S) = 22 is the last even element not in M.

Let $f \in \mathbb{k}[[t]]$. Write $f = \varphi_1 + \varphi_2 + \varphi_3$, where $\varphi_1 := \sum_{i=0}^{54} \lambda_i t^i$, $\varphi_2 := \sum_{i\geq 0} \lambda_{55+2i} t^{55+2i}$ and $\varphi_3 := \sum_{i\geq 0} \lambda_{56+2i} t^{56+2i}$. Clearly $\varphi_1 \in \mathbb{k}[t] \subset D[t]$. Note that $\varphi_2 = \sum_{i\geq 0} \lambda_{55+2i} ((t^6 + t^{31}) t^{24+2i} - t^{30+2i})$. Since 22 is the last even number not in $\langle 8, 10 \rangle$, then, for each $i \geq 0$, both t^{24+2i} and t^{30+2i} are in $\mathbb{k}[[t^8, t^{10}]] \subset D$. Thus, $\varphi_2 \in D$. Again, since 22 is the last even number not in $\langle 8, 10 \rangle$, then, for each $i \geq 0$, $t^{56+2i} \in \mathbb{k}[[t^8, t^{10}]] \subset D$. Therefore, $\varphi_3 \in D$. So, $f \in D[t]$ and $D[t] = \mathbb{k}[[t]]$.

As a consequence, we deduce that $K(D) = \mathbb{k}((t))$. Clearly, $K(D) \subset K(D[t]) = K(\mathbb{k}[[t]]) = \mathbb{k}((t))$. On the other hand, let $g \in \mathbb{k}((t)) = K(\mathbb{k}[[t]]) = K(D[t])$, so g = p(t)/q(t), where $p(t), q(t) \in D[t]$, $q(t) \neq 0$. Since $t \in K(D)$ and $D \subset K(D)$, it follows that p(t), q(t) and $g = p(t)/q(t) \in K(D)$.

Since $D \subset D[t] = \mathbb{k}[[t]]$ is an integral extension, $\mathbb{k}[[t]]$ is integrally closed over its field of fractions $\mathbb{k}((t))$, and $K(D) = \mathbb{k}((t))$, it follows that $\mathbb{k}[[t]]$ is the integral closure of D.

Step 5: length_R(R/(P+yR)) = 8.

Since $R \to D = D/P$ is a surjective map and D/yD = R/(P + yR) is a D-module, it follows that $\operatorname{length}_R(R/(P + yR)) = \operatorname{length}_R(D/yD) = \operatorname{length}_D(D/yD)$. Since y is a D-regular sequence in \mathfrak{m}_D , $1 \leq \operatorname{grade}(\mathfrak{m}_D, D) = \operatorname{depth}(D) \leq \dim(D) = 1$, so D is a Cohen-Macaulay ring. Since $f_1, f_4 \in I \subseteq P$, then $f_1 = f_4 = 0$ in D. Thus, $x^4 \in yD$ and $z^3 \in yD$. So, yD is an \mathfrak{m}_D -primary ideal and y is a system of parameters of D.

Let $V = \mathbb{k}[[t]]$, which is a DVR. Set \mathfrak{m}_V its maximal ideal and $k_V = \mathbb{k}$ its residue field. Let ν be the valuation in $K(V) = \mathbb{k}([t]) = K(D)$ corresponding to V. Through the isomorphism $D = R/P \cong \mathbb{k}[[t^6 + t^{31}, t^8, t^{10}]] \subset V$, then $y = t^8$ and $\nu(y) = \nu(t^8) = 8\nu(t) = 8$.

We have seen that $D \subset D[t] = \mathbb{k}[[t]] = V$, where t is integral over D. Thus, V is a finitely generated D-module. Moreover, V is a DVR, so a Cohen-Macaulay ring. Therefore, V is a finitely generated Cohen-Macaulay D-module.

Note that $K(D) = (D \setminus \{0\})^{-1}D \subseteq (D \setminus \{0\})^{-1}V \subseteq (D \setminus \{0\})^{-1}K(D) = K(D)$. Therefore, $V \otimes_D K(D) = (D \setminus \{0\})^{-1}V = K(D)$. So, $\operatorname{rank}_D(V) = 1$. Using that y is a system of parameters of D and [1, Corollary 4.6.11], we conclude that $\operatorname{length}_D(V/yV) = \operatorname{length}_D(D/yD) \cdot \operatorname{rank}_D(V) = \operatorname{length}_D(D/yD)$.

Let $0 = M_0 \subset M_1 \subset \ldots \subset M_{n-1} \subset M_n = V/yV$ be a chain of V-submodules of V/yV of maximal length, so $M_{i+1}/M_i \cong k_V$ and length_V(V/yV) = n. Applying the additivity of length to the exact sequences of D-modules $0 \to M_{i+1}/M_i \to M_n/M_i \to M_n/M_{i+1} \to 0$, $0 \le i \le n-1$, length_D $(V/yV) = \sum_{i=0}^{n-1} \operatorname{length}_D(M_{i+1}/M_i) = n \cdot \operatorname{length}_D(k_V)$. Since $k_V = \mathbb{k} = k_D$, it follows that length_D $(k_V) = 1$ and length_D $(V/yV) = n = \operatorname{length}_V(V/yV)$. Thus,

 $\operatorname{length}_R(R/(P+yR)) = \operatorname{length}_D(D/yD) = \operatorname{length}_D(V/yV) = \operatorname{length}_V(V/yV) = \nu(y) = 8.$

CONCLUSION.

Since $\rho(y) = t^8$, then $y \notin P$, $P \cap yR = yP$ and $\text{Tor}_1(R/P, R/yR) = 0$. Consider the short exact sequence, $0 \to P/I \to R/I \to R/P \to 0$. On tensoring by R/yR we obtain the short exact sequence

$$0 \to (P/I)/y(P/I) \to R/(I+yR) \to R/(P+yR) \to 0.$$

Since $\operatorname{length}_R(R/(I+yR)) = \operatorname{length}_R(R/(P+yR)) = 8$, it follows that $\operatorname{length}_R((P/I)/y(P/I)) = 0$ and P/I = y(P/I). By Nakayama's Lemma, I = P, as desired.

Remark 4.8. Sally's diagram (I). The following diagram, with natural vertical morphisms, is considered in the book of Sally [9, p. 61]. Now, let $\tilde{\rho}(x) = t^6 + t^{31}$, $\tilde{\rho}(y) = t^8$ and $\tilde{\rho}(z) = t^{10}$, and suppose that $\overline{\rho}$ and ρ are defined in an analogous way.

Let $P = \ker(\rho)$, $\tilde{P} = \ker(\tilde{\rho})$ and $\overline{P} = \ker(\bar{\rho})$. Theorem 4.7 says that P is minimally generated by the polynomials

$$f_1 = \underbrace{3y^3 - 4xyz + x^4 - 3y^3z^5 - 2xy^6z^2 - x^2y^4z^3}_{f_3 = yz^2 - 3x^2y^2 + 2x^3z - y^6z^3 - 2xy^4z^4}, \quad f_2 = \underbrace{2y^2z - 3xz^2 + x^3y}_{f_3 = yz^2 - 3x^2y^2 + 2x^3z - y^6z^3 - 2xy^4z^4}_{f_4 = z^3 - 2xy^3 + x^2yz - y^5z^4}.$$

Since the f's have integer coefficients, they can also be considered in $\mathbb{Z}_{(p)}[[x,y,z]]$ and clearly belong to \tilde{P} because ι is injective. Therefore, the classes $\overline{f_i}$ of the f_i in $(\mathbb{Z}/p\mathbb{Z})[[x,y,z]]$ belong to \tilde{P} . If p=2, then

$$\overline{f_1} = \underline{y^3 + x^4} + y^3 z^5 + x^2 y^4 z^3, \quad \overline{f_2} = \underline{xz^2 + x^3 y} + xy^5 z^3,
\overline{f_3} = \underline{yz^2 + x^2 y^2} + y^6 z^3, \quad \overline{f_4} = \underline{z^3 + x^2 yz} + y^5 z^4.$$

If p=3, then

$$\overline{f_1} = \underline{-xyz + x^4} + xy^6z^2 - x^2y^4z^3, \quad \overline{f_2} = \underline{-y^2z + x^3y} + y^7z^2 - xy^5z^3,$$

$$\overline{f_3} = yz^2 - x^3z - y^6z^3 + xy^4z^4, \quad \overline{f_4} = z^3 + xy^3 + x^2yz - y^5z^4.$$

Discussion 4.9. Suppose that $\operatorname{char}(\mathbb{k}) = 2$. Let us find $g_1, g_2 \in P$, $g_i \neq 0$, such that $g_1^{\sigma} \in V_{10}$ and $g_2^{\sigma} \in V_{12}$. Observe that now $s = \min\{\sigma\operatorname{ord}(f) \mid f \in P, f \neq 0\} = 10$ and $\xi = 3$. Theorem 3.1 says that, in order to find a minimal generating set of P, one can search for linearly independent subsets of V_{10} , V_{11} and V_{12} . However, $W_{11} = \langle xy^2, x^2z \rangle = \langle x^{\alpha}, x^{\beta} \rangle$, with $\alpha = (1, 2, 0)$ and $\beta = (2, 0, 1)$, and where $b_{\alpha_{1},1} = 1$ and $b_{\beta_{1},1} = 0$. Thus, by Lemma 4.2, $V_{11} = 0$.

where $b_{\alpha_1,1}=1$ and $b_{\beta_1,1}=0$. Thus, by Lemma 4.2, $V_{11}=0$. By Lemma 4.5, we can pick $g_1=\underline{z^2+x^2y+y^5z^3}$. With the notations as in Remark 4.8, $\overline{f_2}=xg_1$, $\overline{f_3}=yg_1$, $\overline{f_4}=zg_1$. Moreover, let g_2 be chosen as the polynomial corresponding to the element $\overline{f_1}$ in Remark 4.8. Concretely, let $g_2=\underline{y^3+x^4}+y^3z^5+x^2y^4z^3$. Then, $g_2^{\sigma}\in V_{12},\,g_2\in P$ and $\overline{f_1}=g_2$.

The proof of the next theorem is very similar to the proof of Theorem 4.7. Consequentely, in each step, we only give the demonstration of those details which differ from Theorem 4.7. We maintain the same notations unless stated otherwise.

Theorem 4.10. If $char(\mathbb{k}) = 2$, then P is minimally generated by

$$g_1 = \underline{z^2 + x^2y} + y^5z^3$$
, $g_2 = \underline{y^3 + x^4} + y^3z^5 + x^2y^4z^3$.

Proof. Set $I = (g_1, g_2)$. By Discussion 4.9, $I \subseteq P$.

STEP 1. By Lemma 4.5, s=10. By Theorem 3.1, since $\xi=3$, it follows that g_1, g_2 is part of a minimal system of generators of P, so $\mu(P) \geq 2$. But, if I=P, then $\mu(P) \leq 2$. So, $\mu(P)=2$ and g_1, g_2 is a minimal generating set of P.

STEP 2. Observe that $(g_1, g_1, y) = (z^2, x^4, y)$ and z^2, x^4, y is a regular sequence. The rest follows as in Theorem 4.7.

STEP 3. Note that $J = I + yR = (z^2, x^4, y)$ and $\overline{J} = J/yR = (z^2, x^4)$. For i = 0, 1, consider the exact sequences:

$$0 \to \mathfrak{m}_S^{i+1}/\overline{J} \to \mathfrak{m}_S^i/\overline{J} \to \mathfrak{m}_S^i/\mathfrak{m}_S^{i+1} \to 0.$$

By the additivity of length, length_S $(S/\overline{J}) = 3 + \text{length}_S(\mathfrak{m}_S^2/\overline{J})$. Let H be the ideal of S generated by $q_1 = x^3 z$, $q_2 = x^3$, $q_3 = x^2 z$, $q_4 = x^2$ and $q_5 = xz$. Let

$$\overline{J}_0 = \overline{J}$$
, $\overline{J}_i = \overline{J}_{i-1} + (q_i)$, for $1 \le i \le 5$.

Then, $\overline{J}_5 = \overline{J} + H = \mathfrak{m}_S^2$. Since $\mathfrak{m}_S \overline{J}_i \subseteq \overline{J}_{i-1}$, it follows that, for $i = 1, \ldots, 5$, $\overline{J}_i / \overline{J}_{i-1} = (\overline{q}_i)$ is a one-dimensional vector space, so a simple S-module. For $i = 1, \ldots, 5$, consider the following exact sequences of S-modules:

$$0 \to \overline{J}_{i-1}/\overline{J} \to \overline{J}_i/\overline{J} \to \overline{J}_i/\overline{J}_{i-1} \to 0.$$

Hence, $\operatorname{length}_S(\mathfrak{m}_S^2/\overline{J}) = \sum_{i=1}^5 \operatorname{length}_S(\overline{J}_i/\overline{J}_{i-1}) = 5$. Thus, $\operatorname{length}_S(S/yS) = 8$.

STEP 4 works as in Theorem 4.7. In STEP 5, note that $g_1 = g_2 = 0$ in D, so $z^2, x^4 \in yD$ and y is a system of parameters of D. Finally, the Conclusion works as in Theorem 4.7.

Remark 4.11. Sally's diagram (II). Theorem 4.10 says that \overline{P} is minimally generated by the polynomials g_1, g_2 . Moreover, we have shown that, $\overline{f_1} = g_2$, $\overline{f_2} = xg_1$, $\overline{f_3} = yg_1$ and $\overline{f_4} = zg_1$. Let us see that the representative $\varphi := z^2 + x^2y + y^5z^3$ of g_1 in $\mathbb{Z}_{(2)}[[x, y, z]]$ is not in the ideal $\tilde{P} + 2\mathbb{Z}_{(2)}[[x, y, z]]$. Indeed, if $\varphi \in \tilde{P} + 2\mathbb{Z}_{(2)}[[x, y, z]]$, then $\varphi = f - 2g$, where $f \in \tilde{P}$ and $g \in \mathbb{Z}_{(2)}[[x, y, z]]$. Hence, $2\tilde{\rho}(g) = \tilde{\rho}(f) - \tilde{\rho}(\varphi) = 2t^{20} + 2t^{45} + 2t^{70}$ and, since $\mathbb{Z}_{(2)}[[t]]$ is a domain, it follows that $g \in \tilde{\rho}^{-1}(t^{20} + t^{45} + t^{70})$. Let us see that such a g does not exist. Indeed, let $g = \sum_{s > r} g_s$, $g_s = \sum_{\alpha \in F(s,S)} \lambda_{\alpha} x^{\alpha}$, and suppose that $g \in \tilde{\rho}^{-1}(t^{20} + t^{45} + t^{70})$. Similarly as in Lemma 4.2,

$$\tilde{\rho}(g) = \sum_{s \ge r} \left[\left(\sum_{\alpha \in \mathcal{F}(s,\mathcal{S})} b_{\alpha_1,0} \lambda_{\alpha} \right) t^{2s} + \left(\sum_{\alpha \in \mathcal{F}(s,\mathcal{S})} b_{\alpha_1,1} \lambda_{\alpha} \right) t^{2s+25} + \cdots \right].$$

This forces 2r=20, so r=10 and $g^{\sigma} \in W_{10}=\langle z^2,x^2y\rangle$. Thus, $g^{\sigma}=\lambda z^2+\mu x^2y$, with $\lambda+\mu=1$ and $b_{0,1}\lambda + b_{2,1}\mu = 1$, so $2\mu = 1$, a contradiction because μ must be in $\mathbb{Z}_{(2)}$. Therefore,

$$\overline{P} \not\subset (\widetilde{P}, 2)\mathbb{Z}_{(2)}[[x, y, z]]/2\mathbb{Z}_{(2)}[[x, y, z]],$$

showing that the equality in [9, page 62, line 1] is not correct.

Observe that when passing to characteristic 2 it emerges the element $g_1 \in \overline{P}$, whose σ -order is smaller than the σ -order of $\overline{f_1}$, $\overline{f_2}$, $\overline{f_3}$, $\overline{f_4}$. The equalities $\overline{f_2} = xg_1$, $\overline{f_3} = yg_1$ and $\overline{f_4} = zg_1$, reveal that f_2 , f_3 and f_4 can be substituted by g_1 . "This" is the reason why the minimal number of generators of P might decrease when passing to characteristic p.

Discussion 4.12. Suppose that char(\mathbb{k}) = 3. Let us find $h_1, h_2, h_3 \in P$, $h_i \neq 0$, such that $h_1^{\sigma} \in V_9$, $h_2^{\sigma} \in V_{15}$ and $h_3^{\sigma} \in V_{16}$. Now $s = \min\{\sigma \operatorname{ord}(f) \mid f \in P, f \neq 0\} = 9$ and $\xi = 3$. To directly apply Theorem 3.1, one should find elements whose σ -leading form are in V_9 , V_{10} and V_{11} . However, $W_{10} = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta} \rangle$, with $b_{\alpha_1, 1} - b_{\beta_1, 1} = 1 \neq 0$, and $W_{11} = \langle \mathbf{x}^{\alpha}, \mathbf{x}^{\beta} \rangle$, with $b_{\alpha_1, 1} - b_{\beta_1, 1} = -1 \neq 0$. Therefore, by Lemma 4.2, $V_{10} = 0$ and $V_{11} = 0$. In other words, the only element that we can deduce from Theorem 3.1 that is part of a minimal generating set of P is h_1 . In Remark 4.14 we

see that V_{12} , V_{13} are non-zero, but they do not lead to a minimal generating set of P. By Lemma 4.5, we can pick $h_1 = \underline{yz - x^3} - y^6z^2 + xy^4z^3$. With the notations as in Remark 4.8, $\overline{f_1} = -xh_1, \overline{f_2} = -yh_1, \overline{f_3} = zh_1.$ Take $h_2^{\sigma} = az^3 + bxy^3 + cx^2yz \in W_{15}$, omitting again the x^5 for the sake of simplicity. Then,

$$\rho(h_2^{\sigma}) = (a+b+c)t^{30} + (b+2c)t^{55} + ct^{80}.$$

Choose (a,b,c)=(1,1,1), so $h_2^\sigma=z^3+xy^3+x^2yz$ and $\rho(h_2^\sigma)=t^{80}$. So we can pick $h_2^\tau=-y^5z^4$. Therefore, let $h_2=z^3+xy^3+x^2yz-y^5z^4$. Take $h_3^\sigma=ay^4+\overline{bxy^2z+cx^2z^2}\in W_{16}$, omitting the x^4y for the sake of simplicity. Then,

$$\rho(h_3^{\sigma}) = (a+b+c)t^{32} + (b+2c)t^{57} + ct^{82}.$$

Choose (a,b,c)=(1,1,1), so $h_3^{\sigma}=y^4+xy^2z+x^2z^2$ and $\rho(h_3^{\sigma})=t^{82}$. So we can pick $h_3^{\tau}=-y^4z^5$. Therefore, let $h_3=\underline{y^4+xy^2z+x^2z^2}-y^4z^5$.

Theorem 4.13. If char(k) = 3, then P is minimally generated by

$$h_1 = \underline{yz - x^3} - y^6z^2 + xy^4z^3, \quad h_2 = \underline{z^3 + xy^3 + x^2yz} - y^5z^4, \quad h_3 = \underline{y^4 + xy^2z + x^2z^2} - y^4z^5.$$

Proof. Set $I = (h_1, h_2, h_3)$. By Discussion 4.12, $I \subseteq P$.

Here STEP 1 is more laborious than in Theorem 4.10. By Lemma 4.5, s = 9. By Theorem 3.1, since $\xi = 3$, it follows that h_1 is part of a minimal system of generators of P and $\mu(P) \geq 1$. Since $h_1 \in (x,y)$ and $h_2 = z^3 + f$, with $f \in (x,y)$, it follows that $h_2 \notin (h_1)$. Similarly, since $h_1 \in (x,z)$ and $h_3 = y^4 + g$, with $g \in (x,z)$, it follows that $h_3 \notin (h_1)$. If $\mu(P) = 1$, then $P=(h_1)$ and, since $h_2,h_3\in P$, then $h_2,h_3\in (h_1)$, a contradiction. Therefore, $\mu(P)>1$. Suppose that $\mu(P) = 2$. Then there exist $f \in R$ such that h_1, f is a minimal system of generators of P. Since $h_2, h_3 \in P = (h_1, f)$, it follows that $h_2 = a_1 h_1 + a_2 f$, $h_3 = b_1 h_1 + b_2 f$, where $a_i, b_i \in R$, $1 \le i \le 2$. Since $h_2, h_3 \not\in (h_1)$, it follows that $a_2 \ne 0$ and $b_2 \ne 0$. Moreover, if I = P, then $f = r_1 h_1 + r_2 h_2 + r_3 h_3$, for some $r_i \in R$, $1 \le i \le 3$. Clearly, $(r_2, r_3) \ne (0, 0)$, otherwise $f \in (h_1)$, a contradiction. Substituting the expressions of h_2 and h_3 in the equality $f = r_1h_1 + r_2h_2 + r_3h_3$, we obtain $(1 - r_2a_2 - r_3b_2)f = (r_1 + r_2a_1 + r_3b_1)h_1$. If $1 - r_2a_2 - r_3b_2$ is a unit, then $f \in (h_1)$, a contradiction. Hence, $1-r_2a_2-r_3b_2$ is not a unit, i.e., $1-r_2a_2-r_3b_2 \in \mathfrak{m}$. Therefore, either $r_2a_2 \notin \mathfrak{m}$, or else $r_3b_2 \notin \mathfrak{m}$. In the first case, a_2 is a unit, $f \in (h_1, h_2), P = (h_1, f) \subseteq (h_1, h_2) \subseteq I = P$, so $h_3 \in I = (h_1, h_2) \subset (x, z)$, which is a contradiction, because $h_3 \notin (x, z)$. In the second case, b_2 is a unit, $f \in (h_1, h_3), P = (h_1, f) \subseteq (h_1, h_3) \subseteq I = P$, so $h_2 \in I = (h_1, h_3) \subset (x, y)$, which is a contradiction because $h_2 \notin (x,y)$. Thus, $\mu(P) > 2$. But, if I = P, then $\mu(P) \leq 3$. So, $\mu(P) = 3$ and h_1, h_2, h_3 is a minimal generating set of P.

STEP 2. Now $(h_1, h_2, h_3, y) = (x^3, z^3, x^2z^2, y)$ and x^3, z^3, y is a regular sequence. Then proceed as in Theorem 4.7.

STEP 3. Observe that $J = I + yR = (x^3, z^3, x^2z^2, y)$ and $\overline{J} = J/yR = (x^3, x^2z^2, z^3)$. Note that $\mathfrak{m}_S^4 \subset \overline{J} \subset \mathfrak{m}_S^3$, where S = R/yR and $\mathfrak{m}_S = \mathfrak{m}/yR$. For $0 \le i \le 2$, and understanding $\mathfrak{m}_S^0 = S$, consider the short exact sequences:

$$0 \to \mathfrak{m}_S^{i+1}/\overline{J} \to \mathfrak{m}_S^i/\overline{J} \to \mathfrak{m}_S^i/\mathfrak{m}_S^{i+1} \to 0.$$

By the additivity of the length, length_S $(S/\overline{J}) = 6 + \text{length}_S(\mathfrak{m}_S^3/\overline{J})$. Since $\mathfrak{m}_S^4 \subset \overline{J} \subset \mathfrak{m}_S^3$, it follows that $\mathfrak{m}_S^3/\overline{J} = \langle x^2z, xz^2 \rangle$ is a two-generated S/\mathfrak{m}_S -vectorial space. Thus, length $S/\overline{J} = 8$.

STEP 4 works as in Theorem 4.7. In STEP 5, note that $h_1 = h_2 = 0$ in D, so $x^3, z^3 \in yD$ and y is a system of parameters of D. Finally, the Conclusion works as in Theorem 4.7.

Remark 4.14. In Discussion 4.12, we could look for elements h_2 and h_3 , such that $h_2^{\sigma} \in V_{12}$ and $\hat{h}_3^{\sigma} \in V_{13}$. By Lemma 4.2, we could pick, for instance,

$$\hat{h}_2 = \underline{-xyz + x^4} - xy^6z^2 + y^3z^5 - y^3z^{10} \quad \text{and} \quad \hat{h}_3 = \underline{-y^2z + x^3y} + y^2z^6 - xz^7.$$

However, $(h_1, \hat{h}_2, \hat{h}_3) \subsetneq P = (h_1, h_2, h_3)$. Indeed, h_2 can be written as $h_2 = z^3 + g$, where $g \in (x, y)$ and $(h_1, \hat{h}_2, \hat{h}_3) \subset (x, y)$, thus $h_2 \notin (h_1, \hat{h}_2, \hat{h}_3)$, otherwise $z^3 \in (x, y)$.

Remark 4.15. Sally's diagram (III). Theorem 4.13 says that \overline{P} is minimally generated by the polynomials h_1, h_2, h_3 . Moreover, we have seen that $\overline{f_1} = -xh_1$, $\overline{f_2} = -yh_1$, $\overline{f_3} = zh_1$ and $\overline{f_4} = h_2$. Thus, $\overline{f_1}$, $\overline{f_2}$ and $\overline{f_3}$ can be substituted by h_1 . Nevertheless, h_1 and h_2 are not sufficient to generate P. We need an extra h_3 , which is not a divisor of any of the $\overline{f_i}$.

Remark 4.16. A run with SINGULAR [2] shows that, for the second prime of Moh, when n = 5, $\mu(\ker(\rho_5)) = 6$ if $\operatorname{char}(\mathbb{k}) = 0$ or $p \geq 7$, $\mu(\ker(\rho_5)) = 3$ if $\operatorname{char}(\mathbb{k}) = 2$, $\mu(\ker(\rho_5)) = 2$ if $\operatorname{char}(\mathbb{k}) = 3$, and $\mu(\ker(\rho_5)) = 5$ if $\operatorname{char}(\mathbb{k}) = 5$. To prove it, one could reproduce our proof given for the case n = 3. See the papers of Moh [7, 8] and Mehta, Saha and Sengupta [6] for alternative proofs. For n = 7 or higher, it seems that the computation with SINGULAR stops and does not provide an answer, at least in characteristic zero.

5. Standard Bases

The purpose of this section is to prove that the minimal generating sets of P constructed in Theorems 4.7, 4.10 and 4.13 are standard bases with respect to the negative degree reverse lexicographic order $>_{ds}$. Concretely, we display the Mora standard representation for the S-polynomials (see, e.g., [3, Definition 2.3.8 and Theorem 4.4.16] and [4, Definition 1.6.9 and Algorithm 1.7.6]). We start by recalling some definitions and results. We mainly follow [4, Sections 1.5 to 1.7 and 6.4] and [3, Section 4.4].

Notation 5.1. A monomial ordering > is a total ordering on $\operatorname{Mon}_d = \{\mathbf{x}^\alpha \mid \alpha \in \mathbb{N}^d\}$ satisfying that if $\mathbf{x}^\alpha > \mathbf{x}^\beta$, then $\mathbf{x}^\gamma \mathbf{x}^\alpha > \mathbf{x}^\gamma \mathbf{x}^\beta$, for all $\alpha, \beta, \gamma \in \mathbb{N}^d$. A monomial ordering > is said to be local if $\mathbf{x}^\alpha < 1$, for all $\alpha \neq (0, \dots, 0)$. A local monomial ordering > is a local degree ordering if $\deg(\mathbf{x}^\alpha) < \deg(\mathbf{x}^\beta)$ implies that $\mathbf{x}^\alpha > \mathbf{x}^\beta$. A non-zero polynomial f can be written $f = \sum_{\nu=0}^n a_\nu \mathbf{x}^{\alpha(\nu)}, \ a_\nu \in \mathbb{k} \setminus \{0\}$, where $\mathbf{x}^{\alpha(0)} > \dots > \mathbf{x}^{\alpha(n)}$. The leading term of f is $\operatorname{LT}(f) = a_0 \mathbf{x}^{\alpha(0)}$ and the leading monomial of f is $\operatorname{LM}(f) = \mathbf{x}^{\alpha(0)}$.

Let $\mathbb{k}[\underline{\mathbf{x}}]_{>} := \{f/u \mid f \in \mathbb{k}[\underline{\mathbf{x}}], u \in S_{>}\}$ be the localization of $\mathbb{k}[\underline{\mathbf{x}}]$ with respect to the multiplicative closed set $S_{>} := \{u \in \mathbb{k}[\underline{\mathbf{x}}] \setminus \{0\} \mid \mathrm{LM}(u) = 1\}$. If > is local, then $\mathbb{k}[\underline{\mathbf{x}}]_{>} = \mathbb{k}[\underline{\mathbf{x}}]_{(\underline{\mathbf{x}})}$. For $f \in \mathbb{k}[\underline{\mathbf{x}}]_{>}$, choose $u \in \mathbb{k}[\underline{\mathbf{x}}]$ such that $\mathrm{LT}(u) = 1$ and $uf \in \mathbb{k}[\underline{\mathbf{x}}]$. Then $\mathrm{LT}(f) := \mathrm{LT}(uf)$ and $\mathrm{LM}(f) := \mathrm{LM}(uf)$.

Given a subset $G \subset \mathbb{k}[\underline{x}]_>$, the *leading ideal* of G is defined as $L(G) = \langle \mathrm{LM}(g) \mid g \in G, g \neq 0 \rangle_{\mathbb{k}[\underline{x}]}$, which is seen as an ideal of $\mathbb{k}[\underline{x}]_>$. Let I be an ideal of $\mathbb{k}[\underline{x}]_>$. A finite subset set $G \subset \mathbb{k}[\underline{x}]_>$ is a standard basis of I if $G \subset I$ and L(I) = L(G). In the non-local case, a standard basis is also called a Gröbner basis.

Definition 5.2. Let > be a monomial ordering. Let \mathcal{G} be the set of all finite lists in $\mathbb{k}[\underline{x}]_{>}$. A weak normal form on $\mathbb{k}[\underline{x}]_{>}$ is a map NF: $\mathbb{k}[\underline{x}]_{>} \times \mathcal{G} \to \mathbb{k}[\underline{x}]_{>}$ satisfying:

- (0) NF(0 | G) = 0 for all $G \in \mathcal{G}$;
- (1) If NF($f \mid G$) $\neq 0$, then LM(NF($f \mid G$)) $\notin L(G)$, for all $f \in \mathbb{k}[\underline{x}]$ and all $G \in \mathcal{G}$;
- (2) For all $f \in \underline{\mathbb{k}[\underline{x}]}$ and all $G = g_1, \ldots, g_m, G \in \mathcal{G}$, there exist $u \in (\underline{\mathbb{k}[\underline{x}]})^*$ such that uf has a standard representation with respect to NF(- | G), that is, $uf NF(uf | G) = \sum_{i=1}^m a_i g_i$, with $a_i \in \underline{\mathbb{k}[\underline{x}]}_{(\underline{x})}$ satisfying that $LM(\sum_{i=1}^m a_i g_i) \geq LM(a_i g_i)$, for all i such that $a_i g_i \neq 0$.

A weak normal form is called *polynomial* if whenever $f \in \mathbb{k}[\underline{x}]$ and G is a list in $\mathbb{k}[\underline{x}]$, there exists a $u \in (\mathbb{k}[\underline{x}]_{>})^* \cap \mathbb{k}[\underline{x}]$ such that uf has a standard representation with $a_i \in \mathbb{k}[\underline{x}]$, for all $1 \leq i \leq m$.

Fixed a list G, a weak normal form with respect to G is a map NF: $\mathbb{k}[\underline{x}] \to \mathbb{k}[\underline{x}] \to \mathbb{k}[\underline{x}]$

Definition 5.3. Let > be any monomial ordering. Let $f \in \mathbb{k}[\underline{x}]$ and let $G = g_1, \ldots, g_m$ be a finite list of $\mathbb{k}[\underline{x}]$. The *Mora normal form algorithm* returns polynomials $u, a_i, h \in \mathbb{k}[\underline{x}]$, such that LM(u) = 1, $uf - h = \sum_{i=1}^m a_i g_i$, where $LM(f) \geq LM(a_i g_i)$ for all $a_i g_i \neq 0$ and, if $h \neq 0$, then LM(h) is not divisible by any $LM(g_i)$ (see, e.g., [3, Theorem 4.4.16] and [4, Algorithm 1.7.6]). The reminder h of this expression is called the *Mora Normal Form* of f with respect to G and will be denote it as NFMora $(f \mid G)$.

Remark 5.4. In fact, NFMora : $\mathbb{k}[\underline{x}] \to \mathbb{k}[\underline{x}] \to \mathbb{k}[\underline{x}]$

Remark 5.5. Let > be a local degree ordering on Mon_d . A non-zero element $f \in \mathbb{k}[\![\underline{x}]\!]$ can be written as $f = \sum_{\nu \geq 0}^{\infty} a_{\nu} \mathbf{x}^{\alpha(\nu)}$, $a_{\nu} \in \mathbb{k} \setminus \{0\}$, where $\mathbf{x}^{\alpha(\nu)} > \mathbf{x}^{\alpha(\nu+1)}$. The leading term of f is defined as $\operatorname{LT}(f) = a_0 \mathbf{x}^{\alpha(0)}$ and the leading monomial of f as $\operatorname{LM}(f) = \mathbf{x}^{\alpha(0)}$. Similarly as before, given a subset $G \subset \mathbb{k}[\![\underline{x}]\!]$, the leading ideal of G is defined as $L(G) = \langle G \rangle_{\mathbb{k}[\![\underline{x}]\!]}$, seen as an ideal in $\mathbb{k}[\![\underline{x}]\!]$. Furthermore, G is a standard basis of an ideal $I \subset \mathbb{k}[\![\underline{x}]\!]$ if $G \subset I$ and L(I) = L(G).

Proposition 5.6. Let $F = \{f_1, f_2, f_3, f_4\}$ be defined as in Theorem 4.7. If $\operatorname{char}(\mathbb{k}) = 0$ or if $\operatorname{char}(\mathbb{k}) \geq 5$, then F is a standard basis of P with respect to the negative degree reverse lexicographic order $>_{\operatorname{ds}}$.

Proof. Let J be the ideal of $\mathbb{k}[\underline{\mathbf{x}}]$ generated by $F = \{f_1, f_2, f_3, f_4\}$. By [4, Theorem 6.4.3], if F is a standard basis of J, then F is a standard basis of $J\mathbb{k}[[\underline{\mathbf{x}}]] = P$. Hence, it suffices to prove that F is a standard basis of J. So we need to check that NFMora $(S(f_i, f_j) \mid F) = 0$ for $1 \le i, j \le 4$ (see [4, Theorem 1.7.3 (Buchberger's criterion)]). Using the Mora division algorithm (see Remark below), we prove that:

$$\begin{split} u\,S(f_1,f_2) &= \left(-z^6 + y^5z^2 + (4/3)xy^3z^3 + (16/9)x^2yz^4 - (1/3)x^4y^2z^2 + (2/9)x^5z^3\right)f_1 \\ &- (22/9)x^3z^4f_2 + \left((1/2)x - (9/2)xz^5 - (1/3)x^2y^3z^2 - (11/18)x^3yz^3\right)f_3 + (5/3)x^4z^3f_4; \\ S(f_1,f_3) &= 3x^2yf_1 - 3x^3f_2 - (9/2)xzf_3 + zS(f_1,f_2); \\ S(f_1,f_4) &= 2xy^3f_1 - x^3zf_2 - \left(2xz^2 + (2/3)x^3y\right)f_3 + x^4f_4 + \left((4/9)z^2 + (4/3)x^2y\right)S(f_1,f_2) \\ &+ (5/9)zS(f_1,f_3); \\ S(f_2,f_3) &= -3xf_4; \\ S(f_2,f_4) &= \left((4/3)xy^2 - (8/9)x^2z\right)f_1 - 3xzf_4 + (8/9)x^2S(f_1,f_2); \\ S(f_3,f_4) &= (2/3)xyf_1 - (2/3)x^2f_2; \end{split}$$

where $u=1-z^5-(2/3)xy^3z^2-(11/9)x^2yz^3+(2/9)x^5z^2$. It follows that NFMora $(S(f_1,f_2)\mid F)=$ NFMora $(S(f_2,f_3)\mid F)=$ NFMora $(S(f_3,f_4)\mid F)=0$. By multiplying the second equality by u and substituting the value of $uS(f_1,f_2)$ from the first equality, we get NFMora $(S(f_1,f_3)\mid F)=0$. Similarly, one can check that NFMora $(S(f_1,f_4)\mid F)=$ NFMora $(S(f_2,f_4)\mid F)=0$.

Remark 5.7. To compute the explicit Mora standard representation for the S-polynomials, we have replicated the Mora's Division Algorithm ([4, Algorithm 1.7.6]) in Singular [2]. We use the library teachstd.lib which provides the computations for the S-polynomials and the minimal ecart. The input of the algorithm is a polynomial f and a finite list of polynomials I, and the output is the Mora normal form of f with respect to I denoted by h. In each iteration, we store the divisors in the list D and the quotients in the list Q. Since the Mora's Division Algorithm adds a new element in I in certain iterations, we keep track of it in the liest L. Let us see a more detailed explanation on how to use the code to recover the Mora standard representation for the $S(f_1, f_2)$.

Define the ring k[x, y, z]> where > is the ds ordering.
 LIB "teachstd.lib";
 ring R=0,(x,y,z),ds;

• Define the subset of polynomials $F = \{f_1, f_2, f_3, f_4\}$ as an ideal. poly f1=3y3-4xyz+x4-3y3z5-2xy6z2-x2y4z3; poly f2=2y2z-3xz2+x3y-2y7z2-xy5z3; poly f3=yz2-3x2y2+2x3z-y6z3-2xy4z4; poly f4=z3-2xy3+x2yz-y5z4; ideal F=f1,f2,f3,f4;

- Define the S-polynomial $S(f_1, f_2)$. polv s12=spolv(f1,f2);
- Define the empty lists D, Q and L, initialize h and run the algorithm code.

```
list D;
list Q;
list L;
poly h=s12;
while(h!=0 \&\& minEcart(F,h)!=0){
    poly d=minEcart(F,h);
    poly q=lead(h)/lead(minEcart(F,h));
    D=insert(D,d);
    Q=insert(Q,q);
    if(ecart(d)>ecart(h)){F=F+h;L=insert(L,h);}
        else{L=insert(L,0);}
    h=h-q*d;
};
```

- Run h; to print the Mora normal form for $S(f_1, f_2)$ with respect to F and check that is
- Run any of the three lists to see how many steps the algorithm needed. In this case, it needed 15 steps. The three lists are ordered such that the first element of the list is the last one to be inserted, and so on.
- Do the following iterative process step by step from k=1 to k=15.
 - Let $d_k = D[n-k+1]$ and $q_k = Q[n-k+1]$.
 - Let $h_k = S(f_1, f_2) \sum_{i=1}^{k} q_i d_i$.
 - If L[n-k+1]!=0, then the algorithm has added h_k as a new element of I.
 - Identify d_k as an element of I. Note that in previous steps, we could have added some
- new elements in I. Then, $d_k = f_j$ for some $1 \le j \le 5$ or $d_k = h_i$ for some $1 \le i < k$.

 In the last step, we get $h_n = S(f_1, f_2) \sum_{i=1}^n q_i d_i$ which is the Mora normal form of $S(f_1, f_2)$ and it is zero. Hence, $S(f_1, f_2) = \sum_{i=1}^n q_i d_i$. Note that the elements d_i are either f_i for some $1 \le i \le 4$ or h_k for some $1 \le k \le n$. Moreover, $h_1 \in (f_1, f_2, f_3, f_4, S(f_1, f_2)), h_2$ is in $(f_1, f_2, f_3, f_4, s_{1,2}, h_1) = (f_1, f_2, f_3, f_4, S(f_1, f_2)).$ Recursively, $h_k \in (f_1, f_2, f_3, f_4, S(f_1, f_2)).$ This means that $d_i \in (f_1, f_2, f_3, f_4, S(f_1, f_2))$ and so we can recover the Mora standard representation of $S(f_1, f_2)$ with respect to F. This is $uS(f_1, f_2) = \sum_{i=1}^s a_i f_i$ where $u, a_i \in$ $k[\underline{x}]$ for $1 \le i \le 4$ and LM(u) = 1.

After computing the Mora standard representation for $S(f_1, f_2)$, we repeat the process for $S(f_1, f_3)$, and so on, keeping the extended ideal F from the previous step. This provides a more efficient way of computing the Mora standard representation for all the S-polynomials.

Proposition 5.8. Let $G = \{g_1, g_2\}$ be defined as in Theorem 4.10. If $\operatorname{char}(\mathbb{k}) = 2$, then G is a standard basis of P with respect to the negative degree reverse lexicographic order $>_{ds}$.

Proof. Using the Mora division algorithm we obtain:

$$(1+y^5z+x^2yz^3)S(g_1,g_2)=(x^4+y^3z^5+y^8z)g_1+(x^2y+x^2yz^5)g_2.$$

Therefore, NFMora $(S(g_1, g_2) \mid G) = 0$.

Proposition 5.9. Let $H = \{h_1, h_2, h_3\}$ be defined as in Theorem 4.13. If $\operatorname{char}(\mathbb{k}) = 3$, then H is a standard basis of P with respect to the negative degree reverse lexicographic order $>_{ds}$.

Proof. Using the Mora division algorithm we obtain:

$$S(h_1, h_2) = -xh_3;$$

$$(1 - z^5) S(h_1, h_3) = (-xyz + xyz^6) h_1 + (-x^2 - xy^2z^4) h_2 + (z^6 - y^5z^2 + xy^3z^3 + x^2yz^4) h_3;$$

$$S(h_2, h_3) = xyz^3h_1 - xy^2zh_2 + (xy^3 + x^2yz) h_3 + z^2S(h_1, h_3).$$

So, NFMora $(S(h_1, h_2) \mid H) = \text{NFMora}(S(h_1, h_3) \mid H) = 0$. Multiplying the third equality by $v = 1 - z^5$ and substituting the value of $vS(h_1, h_3)$, we get NFMora $(S(h_2, h_3) \mid H) = 0$. The rest follows as in Proposition 5.6.

References

- W. Bruns, J. Herzog, Cohen-Macaulay rings. Cambridge Studies in Advanced Mathematics, 39. Cambridge University Press, Cambridge, 1993
- [2] W. Decker, G.-M. Greuel, G. Pfister, H. Schönemann, SINGULAR 4-3-0 A computer algebra system for polynomial computations. https://www.singular.uni-kl.de (2022).
- [3] W. Decker and F. Schreyer, Varieties, groebner bases, and algebraic curves. Manuscript. Available at: https://www.math.uni-sb.de/ag/schreyer/index.php/teaching/ss-18/128-algebraische-geometrie
- [4] G.-M. Greuel, G. Pfister, A Singular introduction to commutative algebra. Second, extended edition. With contributions by Olaf Bachmann, Christoph Lossen and Hans Schönemann. Springer-Verlag, Berlin, 2008.
- [5] H. Matsumura, Commutative ring theory. Translated from the Japanese by M. Reid. Second edition. Cambridge Studies in Advanced Mathematics, 8. Cambridge University Press, Cambridge, 1989
- [6] R. Mehta, J. Saha, I. Sengupta, Moh's example of algebroid space curves. J. Symbolic Comput. 104 (2021), 168–182.
- [7] T.T. Moh, On the unboundedness of generators of prime ideals in power series rings of three variables. J. Math. Soc. Japan 26 (1974), 722-734
- [8] T.T. Moh, On generators of ideals. Proc. Amer. Math. Soc. 77 (1979), no. 3, 309-312.
- [9] J. Sally, Numbers of generators of ideals in local rings. Marcel Dekker, Inc., New York-Basel, 1978.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT POLITÈCNICA DE CATALUNYA. DIAGONAL 647, ETSEIB, E-08028 BARCELONA.

 $Email\ address: \verb|laura.gonzalez.hernandez@upc.edu|$

Email address: francesc.planas@upc.edu